

The unstable homotopy groups of 2-cell complexes

Zhongjian Zhu

College of Mathematics and Physics, Wenzhou University, Wenzhou 325035, China

Abstract

In this paper, we develop the new method, initiated by B. Gray (1972), to compute the unstable homotopy groups of the mapping cone, especially for 2-cell complex $X = S^m \cup_{\alpha} e^n$. By Gray's work mentioned above or the traditional method given by I.M.James (1957) which were widely used in previous related work to compute $\pi_i(X)$, the dimension $i \leq 2n + m - 4$. By our method, we can compute $\pi_i(X)$ for $i > 2n + m - 4$. We use this different technique to generalize J.Wu's work, at Mem. of AMS, on homotopy groups of mod 2 Moore spaces to higher dimensional homotopy groups of mod 2^r Moore spaces $P^n(2^r)$ for all $r \geq 1$. This practice shows that the technique given here is a new general method to compute the unstable homotopy groups of CW complexes with higher dimension.

Keywords: Relative James construction, fibration sequence, homotopy group, Moore space

2000 MSC: 55P10, 55Q05, 55Q15, 55Q52

1. Introduction

Calculating the unstable homotopy groups of finite CW-complexes is a fundamental and difficult problem in algebraic topology. This problem is reduced to compute $\pi_i(Y^*)$, where $Y^* = Y \cup_f e^n$, f is the attaching map of the cell e^n . There are two traditional methods to compute the homotopy groups $\pi_i(Y^*)$ for $i \leq 2n + m - 4$ with $m - 1$ connected space Y . The first is applying James's method, i.e., the Theorem 2.1 of [20], to the homotopy exact sequences of a pair (Y^*, Y) . By this method, a lot of related work has been done on 2-cell complexes, such as mod 2^r Moore spaces when $r = 1, 2, 3$ and suspended complex projective spaces. Let $P^n(p^r)$ denote n -dimensional elementary Moore spaces, whose only nontrivial reduced homology is $\overline{H}_{n-1}(P^n(p^r)) = \mathbb{Z}_{p^r}$ ($\mathbb{Z}_k := \mathbb{Z}/k\mathbb{Z}$). J. Mukai computed $\pi_i(P^n(2))$ for some $i \leq 3n - 5$ and $3 \leq n \leq 7$ in [28, 29]. In 1999, J.Mukai and T.Shinpo computed the $\pi_i(P^n(4))$ in the range $i = 2n - 3, 2n - 4$ and $n \leq 24$ [27]; In 2007, X.G.Liu computed $\pi_i(P^n(8))$ in the range

Email address: 20160118@wzu.edu.cn (Zhongjian Zhu)

$i = 2n - 3$ for $2 \leq n \leq 20$ and $i = 2n - 2$ for $3 \leq n \leq 7$ [24]. Some homotopy groups of suspended complex projective plane $C_\eta^n := \Sigma^{n-4}\mathbb{C}P^2$, $n \geq 5$, was also computed in [31].

The second method is using the fibration sequence $\Omega\Sigma X \xrightarrow{\partial} F \rightarrow Y \cup_f CX \xrightarrow{\text{pinch}} \Sigma X$, where $Y \cup_f CX$ is the mapping cone of the map f , which is also denoted by C_f . In 1972, B.Gray constructed the relative James construction $J(X, A)$ for a CW-pair (X, A) and inclusion $A \xrightarrow{i} X$, which is filtrated by $J_r(X, A)$ ($r = 1, 2, \dots$) and proved that $J(X, A)$ is homotopy equivalent to the homotopy fiber of the pinch map $X \cup_i CA \rightarrow \Sigma A$ [13]. So the homotopy type of the above homotopy fiber F is homotopy equivalent to Gray's relative James construction $J(M_f, X)$ which is filtrated by a sequence of subspace $J_r(M_f, X)$, $r = 1, 2, \dots$, where M_f is the mapping cylinder of f . When $X = \Sigma X'$, $Y = \Sigma Y'$, the homotopy type of the second filtration $J_2(M_f, X)$ of $J(M_f, X)$ was also obtained by Gray in [13]. It has the homotopy type $Y \cup_{\gamma_2} C(Y \wedge X')$, where $\gamma = [id_Y, f]$ is the generalized Whitehead product. By the recently developed work of Yang, Mukai and Wu [45] on the homotopy analysis of the connection map $\Omega\Sigma X \xrightarrow{\partial} F$, Gray's method to compute homotopy groups of CW-complex becomes more operable. As applications, they compute homotopy groups of suspended quaternionic projective plane, i.e., $\pi_i(\Sigma^k \mathbb{H}P^2)$ for $i \leq k + 15$ and we (the author and Pan) compute the first two dimensional unstable homotopy groups of indecomposable \mathbf{A}_3^2 -complexes, i.e., 2-connected finite CW-complexes with dimensions of cells less than or equal to 5 [49].

So far, the only general way to compute $\pi_i(Y^*)$ for $i > 2n + m - 4$ is given by Wu in 2003 [43]. In it, a p -local functorial decomposition $\Omega\Sigma X \simeq A^{\min}(X) \times \Omega(\bigvee_{n=2}^{\infty} Q_n^{\max}(X))$ is given, which is helpful to calculate homotopy groups of ΣX . $Q_n^{\max}(X)$ is a wedge summand of $\Sigma X^{\wedge n}$, where $X^{\wedge n}$ is a n -fold self-smash product of X . It is an amazing method since it connects the algebraic representation theory and the homotopy theory, by which he got a lot of unstable homotopy groups $\pi_i(P^n(2))$ ($n \geq 3$) where i can be larger than $3n - 5$ (i.e., $2n + m - 4$ with $m = n - 1$). However it contains so high technique that the homotopy type of $A^{\min}(X)$ or skeletons of it is difficult to get [36]. $Q_n^{\max}(X)$ is also hard to be obtained for X with more than 2 cells since the homotopy wedge decomposition of $X^{\wedge n}$ is not easy [46, 47, 48].

Another way to compute $\pi_i(Y^*)$ for $i > 2n + m - 4$ is to get information of homotopy type of the filtration $J_r(M_f, X)$ of the Gray's relative James construction $J(M_f, X)$ for the pinch map $Y \cup_f CX \xrightarrow{\text{pinch}} \Sigma X$, where $X = \Sigma X'$, $Y = \Sigma Y'$ are suspensions. By our previous work we show that the attaching map $\gamma_r : \Sigma^{r-1} Y' \wedge X'^{\wedge(r-1)} \rightarrow J_{r-1}(M_f, X)$ is an element of set of r -th order Whitehead products defined by G.J.Porter [34], that is

Theorem 1.1 (Theorem 3.4 of [50]). *Let $X \xrightarrow{f} Y$ be a map of simply connected CW complexes, $X = \Sigma X'$, $Y = \Sigma Y'$. Then $J_n(M_f, X)$ has the homotopy type $Y \cup_{\gamma_2} C(\Sigma Y' \wedge X') \cup_{\gamma_3} \dots \cup_{\gamma_n} C(\Sigma^{n-1} Y' \wedge X'^{\wedge(n-1)})$, γ_r is an element of r -th order Whitehead products*

in $[j_Y^{r-1}, j_Y^{r-1}f, \dots, j_Y^{r-1}f]$ where $j_Y^{r-1} : Y \hookrightarrow Y \cup_{\gamma_2} C(\Sigma Y' \wedge X') \cup_{\gamma_3} \dots \cup_{\gamma_{r-1}} C(\Sigma^{r-2} Y' \wedge X'^{\wedge(r-2)})$ is the canonical inclusion for $r = 2, \dots, n$.

As an application, we used Theorem 1.1 to compute the homotopy groups $\pi_k(P^3(2^r))$ ($k = 5, 6$) for all positive integers r in [50]. However the above calculation relies on a result of one special third order Whitehead products computed in [35], previously we did not very sure that Theorem 1.1 had wide applications in computing high dimensional homotopy groups of mapping cones because of that $\gamma_r (r \geq 3)$ is an element of r -th order Whitehead products $[j_Y^{r-1}, j_Y^{r-1}f, \dots, j_Y^{r-1}f]$, which is a set that may contain more than one element so that it will not be enough to determine γ_r .

In this paper, we find that by combining Theorem 1.1 and some homotopy property, especially the naturality of γ_r , we can give the detailed description of γ_3 for a lot of 2-cell complexes. This enable us to develop the method of Gray to compute the $\pi_i(Y^*)$ for $i > 2n + m - 4$. We use this new method to generalize J.Wu's work [43] on homotopy groups of mod 2 Moore spaces to higher dimensional homotopy groups of mod 2^r Moore spaces $P^n(2^r)$ for all $r \geq 1$. The following is our main result.

Theorem 1.2. *The unstable homotopy groups of $\pi_i(P^n(2^r))$ for $n = 4, 5, 6$ and $2n-2 \leq i \leq 4n-6$ are given as follows.*

$i =$	6	7	8	9	10
$\pi_i(P^4(2^r))$	$(2)^{1+\theta_r+2^{r+1}}, r \leq 2$ $2+4+2^r, r \geq 3$	$(2)^{2+4\theta_r}$	$(2)^3, r=1$ $(2)^2+(4)^2, r=2$ $(2)^4+2^r, r \geq 3$	$(2)^{2+\theta_r+2^{m^3}}$	$(2)^{2+4}, r=1$ $(2)^2+2^{m^3}+2^r, r \geq 2$

Table 1

$i =$	8	9	10
$\pi_i(P^5(2^r))$	$(2)^{1+\theta_r+2^{m^3}}$	$(2)^{2+4\theta_r}, r \leq 2$ $(2)^4, r \geq 3$	$2+4, r=1$ $(2)^2+2^{m^3+1}, r \geq 2$
11	12	13	14
$2+2^{m^3+1}+2^r$	$(2)^3$	$(2)^2+(4)^2, r=1$ $(2)^6, r \geq 2$	$(2)^7+4, r=1$ $(2)^7+(4)^3, r=2$ $(2)^6+4+(8)^2+2^r, r \geq 3$

Table 2

$i =$	10	11	12	13	
$\pi_i(P^6(2^r))$	$8, r=1$ $2+2^{r+1}, r \geq 2$	$(2)^2$	$(2)^{2+\theta_r}$	$(2)^3, r=1$ $2+4+2^{m^3}, r \geq 2$	
14	15	16	17	18	
$(2)^4+4$ $(2)^4+2^{m^3}+2^r,$	$(2)^4+4,$ $(2)^5+2^{m^3}$	$(2)^6+4$ $(2)^6+(2^{m^3})^2$	$(2)^5+(8)^2$ $(2)^4+4+(2^{m^3})^2+2^{m^4}$	$(2)^5+8$ $(2)^4+2^r+2^{m^3}+2^{m^4}$	$r=1$ $r \geq 2$

Table 3

In the above tables, an integer k indicates a cyclic group $\mathbb{Z}_k := \mathbb{Z}/k\mathbb{Z}$ of order k , the symbol “+” the direct sum of the groups and $(2)^t$ indicates the direct sum of t -copies of \mathbb{Z}_2 . $m_a^b := \min\{a, b\}$ indicates the minimal integer of a and b . $\vartheta_1 = 0$ and $\vartheta_r = 1$ for $r \geq 2$.

Theorem 1.3. *The first 9 dimensional unstable homotopy groups of $\pi_i(P^n(2^r))$ for $n = 7, 8$ are given as follows.*

$i =$	11	12	13	14
$\pi_i(P^7(2^r))$	2^r	2	$(2)^2 + 2^{m_{r-1}^2}$	$(2)^2 + 2^{m_{r+1}^3} + 2^{m_r^3}$
15	16	17	18	19
$2 + (4)^2 + 8, r=1$ $(2)^2 + 2^{m_r^3}, r \geq 2$	$(2)^4 + 4, r=1$ $(2)^5 + 2^{m_r^3}, r \geq 2$	$2 + 4 + (2^{m_r^3})^3$	$(2)^3 + 2^{m_r^3} + 2^{m_r^4}$	$(2)^2 + 4, r=1$ $(2)^3 + 2^{m_r^4}, r \geq 2$

Table 4

$i =$	13	14	15	16
$\pi_i(P^8(2^r))$	2	$2 + 2^{r-1} + 2^{r+1}, r \leq 3$ $2 + 8 + 2^r, r \geq 4$	$(2)^{3+\vartheta_r} + 2^{m_r^3}$	$(2)^3 + (4)^2, r=1$ $(2)^7 + 4, r=2$ $(2)^9, r \geq 3$
17	18	19	20	21
$(2)^3 + (4)^2, r=1$ $(2)^6 + 2^{m_{r-1}^3} + 8, r \geq 2$	$(2)^2 + 2^{m_{r-1}^3}$ $+ 2^{m_r^3} + 2^{m_{r+1}^3}$	$2 + 2^{m_r^3}$	$(2)^{1+\vartheta_r} + 2^r$	$(2)^3 + 2^{m_r^2}$ $+ 2^{m_{r-1}^3} + 2^{m_{r+1}^4}$

Table 5

where for $r = 1$, $\mathbb{Z}_{2^{m_{r-1}^3}} = \mathbb{Z}_{2^{m_{r-1}^2}} = \mathbb{Z}_{2^0} = \mathbb{Z}_1 = 0$.

Remark 1.4.

- In Theorem 1.2, $\pi_i(P^4(2^r))$ for $i = 6, 7$ are given by [49]; $\pi_8(P^5(2^r))$ is given by [23].

- $2n - 3$ dimensional homotopy groups of $P^n(2^r), n = 4, 5, 6$ (which may be unstable) are not listed in the Tables of Theorem 1.2, since $\pi_5(P^4(2^r)) = \pi_5^s(P^4(2^r)) \cong \begin{cases} \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2, & r \geq 2 \end{cases}$ is given in [3, Part IV]; $\pi_7(P^5(2^r)) \cong \mathbb{Z}_{2^{m_{r-1}^2}} \oplus \mathbb{Z}_{2^{r+1}} \oplus \mathbb{Z}_2$ and $\pi_9(P^6(2^r)) \cong \mathbb{Z}_{2^{m_r^3}} \oplus \mathbb{Z}_2$ are given in [49] and [23] respectively.

- Some homotopy groups of mod 2 Moore spaces in the above Tables, that is $\pi_i(P^4(2))$ $i \leq 10$, $\pi_i(P^5(2))$ $i \leq 13$, $\pi_i(P^6(2))$ $i \leq 14$, $\pi_i(P^7(2))$ $i \leq 15$ and $\pi_i(P^8(2))$ $i \leq 16$ are also given by [43].

- We also generalize the results of $\pi_i(P^n(2))$ for $n = 9, 10, 11$ in [43] to $\pi_i(P^n(2^r))$ for any $r \geq 1$ in Theorem 7.1 without proof.

There are two advantages of the method given in this paper to compute the unstable homotopy groups. Firstly, it is easy to get the generators of homotopy groups by the calculation process. Secondly, we are able to calculate the homotopy groups of $P^n(2^r)$ for all $r \geq 1$ simultaneously.

Some new techniques are given here to determine the connection homomorphism $\partial_{k*} : \pi_{k+1}(X) \rightarrow \pi_k(F)$ and solve the extension problem of the short exact sequence of homotopy groups

$$0 \rightarrow \text{Coker}(\partial_{k*}) \rightarrow \pi_k(C_f) \rightarrow \text{Ker}(\partial_{k-1*}) \rightarrow 0$$

which is induced by fibration sequence $\Omega\Sigma X \xrightarrow{\partial} F \rightarrow Y \cup_f CX \xrightarrow{\text{pinch}} \Sigma X$, where $\text{Ker}(h)$ and $\text{Coker}(h)$ denote the Kernel and Cokernal of a homomorphism $h : G \rightarrow H$ of abelian groups respectively; f_{m*} denotes the induced homomorphism $f_* : \pi_m(X) \rightarrow \pi_m(Y)$ for a map $f : X \rightarrow Y$ of spaces. We also show that the homotopy type of $J_3(M_f, X)$ for some other spaces such as suspended complex or quaternionic projective plane can be obtained by our method. This enables us to calculate the homotopy groups of them with dimension larger than that in the previous work. This gives us reason to believe that the method given in this paper has wide applications in computing high dimensional homotopy groups of CW-complexes, at least for 2-cell complexes.

By the way, recently the research on the suspension homotopy of manifolds becomes a popular topic [11, 18, 19, 25, 37], however the homology groups of manifolds considered by them have no 2-torsion. And the known unstable homotopy groups of mod q^r (q is odd prime) Moore spaces are very important in their methods. So the above results of unstable homotopy groups of mod 2^r Moore spaces have a potential application, that is to classify the homotopy types of the suspension of non-simply connected manifolds whose homology groups are allowed to have 2-torsion.

The paper is arranged as follows. Section 2 introduces some concepts and properties of higher order Whitehead products and relative James construction; Section 3 gives some homotopy properties of the connection map $\Omega\Sigma X \xrightarrow{\partial} F_p$ and some q which are useful to solve the extension problem of the short exact sequence. Section 5 and Section 6 calculate the unstable homotopy groups of mod 2^r Moore spaces in Theorem 1.2 and Theorem 1.3 respectively. The last section, i.e., Section 7 gives some remarks to the method given in this paper.

2. Higher order Whitehead products and relative James construction

In this paper, all spaces and maps are in the category of pointed topological spaces and maps (i.e. continuous functions) preserving base point. And we always use $*$ and 0 to denote the basepoint and the constant map mapping to the basepoint respectively. Without special mention, all spaces are CW-complexes and all the space

pairs are CW-pairs. We denote $A \hookrightarrow X$ as an inclusion map. By abuse of notion, we will not distinguish the notions between a map and its homotopy class in many cases.

For the maps $f_i : X_i \rightarrow X_{i+1} (i = 1, 2, \dots, n)$, $f_n f_{n-1} \cdots f_1 : X_1 \rightarrow X_n$ denotes the composition of $f_i (i = 1, 2, \dots, n)$; Moreover, if $[X_1, X_n]$ is an abelian group, for example X_1 is a double suspension, then for $k \in \mathbb{Z}$, $k f_n f_{n-1} \cdots f_1 := k(f_n f_{n-1} \cdots f_1)$ denotes the k times of $f_n f_{n-1} \cdots f_1$;

2.1. Higher order Whitehead products

Let $T_r(X_1, X_2, \dots, X_n)$ be the subset of the Cartesian product $X_1 \times X_2 \times \cdots \times X_n$, consisting of those n -tuples with at least r co-ordinates at a base point. Thus $T_{n-1}(X_1, X_2, \dots, X_n) = X_1 \vee X_2 \vee \cdots \vee X_n$, $T_1(X_1, X_2, \dots, X_n)$ is the ‘‘fat wedge’’, and $X_1 \times X_2 \times \cdots \times X_n / T_1(X_1, X_2, \dots, X_n) = X_1 \wedge X_2 \wedge \cdots \wedge X_n$. From Theorem 1.2 and Theorem 2.1 of [34], there is a principle cofibration

$$\Sigma^{n-1} X_1 \wedge \cdots \wedge X_n \xrightarrow{W_n} T_1(\Sigma X_1, \dots, \Sigma X_n) \rightarrow \Sigma X_1 \times \cdots \times \Sigma X_n \quad (1)$$

where the map W_n is natural.

Given a map $f : T_1(\Sigma X_1, \dots, \Sigma X_n) \rightarrow X$, $n \geq 2$, define

$$W(f) := f_*(W_n) = f W_n \in [\Sigma^{n-1} X_1 \wedge \cdots \wedge X_n, X]$$

the n -th order Whitehead product, which depends only upon the homotopy class of f [34].

Let $k_j : \Sigma X_j \hookrightarrow T_1(\Sigma X_1, \dots, \Sigma X_n)$, $j = 1, 2, \dots, n$, be the canonical injections.

Definition 2.1. *The set of n -th order Whitehead products of $f_j : \Sigma X_j \rightarrow X$, $j = 1, \dots, n$, is*

$$[f_1, \dots, f_n] := \{W(f) \mid f : T_1(\Sigma X_1, \dots, \Sigma X_n) \rightarrow X, f k_j \simeq f_j, j = 1, \dots, n\}.$$

Remark 2.2. 1) $[f_1, \dots, f_n]$ is a subset of $[\Sigma^{n-1} X_1 \wedge \cdots \wedge X_n, X]$, and it is perhaps empty. We also have $[f_1, \dots, f_n] := \{W(f) := f W_n \mid f : T_1(\Sigma X_1, \dots, \Sigma X_n) \rightarrow X\}$ for all f extending the wedge sum map $(f_1, \dots, f_n) : \Sigma X_1 \vee \cdots \vee \Sigma X_n \rightarrow X$ up to homotopy.

2) $[f_1, \dots, f_n]$ depends only upon the homotopy classes of $f_i (i = 1, \dots, n)$;

3) Hardie[14] has also given the definition of $[f_1, \dots, f_n]$ when all X_i are spheres (called the n -th order spherical Whitehead product). $[f_1, f_2, f_3]$ is, in this case, the Zeeman product studied by Hardie[15]. When X_1, X_2 are arbitrary, $[f_1, f_2]$ is the ‘‘generalized Whitehead product’’ studied by Arkowitz[1].

4) From the Theorem 2.5 [34], if X is an H -space, then $[f_1, \dots, f_n] = \{0\}$; if Σ is the suspension homomorphism, then $\Sigma[f_1, \dots, f_n] = \{0\}$.

The following naturality of higher order Whitehead products comes from Theorem 2.1 of [34] and Corollary 2.4 of [50].

Proposition 2.3. (*Naturality*) Let $f_i : A_i \rightarrow B_i$, $h_i : \Sigma B_i \rightarrow X$, $i = 1, \dots, n$, $g : X \rightarrow Y$ and $\varphi : T_1(\Sigma B_1, \dots, \Sigma B_n) \rightarrow X$, then

$$(a) (\Sigma^{n-1}(f_1 \wedge \dots \wedge f_n))^*[h_1, \dots, h_n] \subset [h_1 \Sigma f_1, \dots, h_n \Sigma f_n]$$

$$(b) g*[h_1, \dots, h_n] \subset [gh_1, \dots, gh_n].$$

(c) If $B_j = \Sigma B'_j$ for some j , then $k[h_1, \dots, h_j, \dots, h_n] \subset [h_1, \dots, kh_j, \dots, h_n]$ for any integer k ;

If $f_i : A_i \rightarrow B_i$ and $g : X \rightarrow Y$ are homotopy equivalences, then

$$(d) (\Sigma^{n-1}(f_1 \wedge \dots \wedge f_n))^*[h_1, \dots, h_n] = [h_1 \Sigma f_1, \dots, h_n \Sigma f_n];$$

$$(e) g*[h_1, \dots, h_n] = [gh_1, \dots, gh_n].$$

The following lemma about the property of third order Whitehead product is from the Theorem 4.2 of [12], and the special case for spheres $X_i (i = 1, 2, 3)$ are given by Hardie [15].

Lemma 2.4. $[f_1, f_2, f_3]$ with $f_i : \Sigma X_i \rightarrow X (i = 1, 2, 3)$ is a coset of the following subgroup of group $[\Sigma^2 X_1 \wedge X_2 \wedge X_3, X]$

$$[[\Sigma^2 X_2 \wedge X_3, X], f_1] + [[\Sigma^2 X_1 \wedge X_3, X], f_2] + [[\Sigma^2 X_1 \wedge X_2, X], f_3].$$

2.2. Relative James construction

Let (X, A) be a pair of spaces with base point $* \in A$, and suppose that A is closed in X . In [13], B.Gray constructed a space $(X, A)_\infty$ analogous to the James construction, which is denoted by us as $J(X, A)$ in [49] to parallel with the the absolute James construction $J(X)$ [21]. In fact, $J(X, A)$ is the subspace of $J(X)$ of words for which letters after the first are in A . Especially, $J(X, X) = J(X)$. As parallel with the familiar symbol $J_r(X)$ which is the r -th filtration of $J(X)$, we denote the r -th filtration of $J(X, A)$ by $J_n(X, A) := J(X, A) \cap J_r(X)$, which is denoted by Gray as $(X, A)_r$ in [13]. For example,

$$J_1(X, A) = X, J_2(X, A) = (X \times A)/((a, *) \sim (*, a)),$$

$$J_3(X, A) = (X \times A \times A)/((x, *, a) \sim (x, a, *); (a, a', *) \sim (*, a, a') \sim (a, *, a')).$$

In fact there is a pushout diagram for $r \geq 2$:

$$\begin{array}{ccc} X \times A^{\wedge(n-1)} & \xrightarrow{\Pi_r} & J_r(X, A) \\ \uparrow & & \uparrow I_{r-1} \\ T_1(X, A, \dots, A) & \longrightarrow & J_{r-1}(X, A) \end{array}$$

where Π_r and I_{r-1} are the projection and the inclusion respectively and both of them are natural. Moreover, $J_n(X, A)/J_{n-1}(X, A)$ is naturally homeomorphic to $(X \times A^{n-1})/T_1(X, A, \dots, A) = X \wedge A^{\wedge(n-1)}$.

It is well known that there is a natural weak homotopy equivalence $\omega : J(X) \rightarrow \Omega\Sigma X$, which is a homotopy equivalence when X is a finite CW-complex, and satisfies

$$X \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow[\Omega\Sigma]{\quad} \\ \xrightarrow{\quad} \end{array} J(X) \xrightarrow{\omega} \Omega\Sigma X, \text{ where } X \xrightarrow{\Omega\Sigma} \Omega\Sigma X, \text{ also denoted by } E_X, \text{ is the inclusion}$$

$x \mapsto \psi$ where $\psi : S^1 \rightarrow S^1 \wedge X, t \mapsto t \wedge x$.

In the following text, the ‘‘commutative’’ means ‘‘homotopy commutative’’.

Let $X \xrightarrow{f} Y$ be a map. There is a cofibration sequence $X \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{p} \Sigma X$. We always use C_f, F_f and M_f to denote the mapping cone (or say, cofibre), homotopy fibre and mapping cylinder of f , $C_f \xrightarrow{p} \Sigma X$ the pinch map and $\Omega\Sigma X \xrightarrow{\partial} F_p \xrightarrow{\tau_p} C_f \xrightarrow{p} \Sigma X$ the homotopy fibration sequence induced by p respectively. Suppose that $X = \Sigma X', Y = \Sigma Y'$ are suspensions. Applying Theorems of [13] to (M_f, X) , we get

- (i) $F_p \simeq J(M_f, X)$;
- (ii) $\Sigma J(M_f, X) \simeq \bigvee_{k \geq 0} (\Sigma Y \wedge X^{\wedge k}); \Sigma J_k(M_f, X) \simeq \bigvee_{i=0}^{k-1} (\Sigma Y \wedge X^{\wedge i})$;
- (iii) $J_2(M_f, X) \simeq Y \cup_{\gamma_2} C(Y \wedge X')$, where $\gamma_2 = [id_Y, f]$ is the generalized Whitehead product.

We denote both the inclusion $Y \hookrightarrow J_2(M_f, X)$ and the composition of the inclusions $Y \hookrightarrow J_2(M_f, X) \hookrightarrow J(M_f, X) \simeq F_p$ by j_p without ambiguous. As pointed out in the proof of Lemma 4.1.of [13], j_p can be chosen naturally as the lift of the inclusion $Y \xrightarrow{i} C_f$, i.e.,

$$\tau_p j_p \simeq i \tag{2}$$

Moreover the relative James construction is natural in the sense of the following

Lemma 2.5 (Lemma 2.2 of [49]). *Suppose the left diagram is commutative*

$$\begin{array}{ccc} X \xrightarrow{f} Y & ; & F_p \simeq J(M_f, X) \longrightarrow M_f/X \simeq C_f \xrightarrow{p} \Sigma X \\ \downarrow \mu_X & \downarrow \mu_Y & \downarrow J(\widehat{\mu}, \mu_X) \\ X_1 \xrightarrow{f_1} Y_1 & & F_{p_1} \simeq J(M_{f_1}, X_1) \longrightarrow M_{f_1}/X_1 \simeq C_{f_1} \xrightarrow{p_1} \Sigma X' \end{array}$$

then it induces the right commutative diagrams on fibrations, where $\widehat{\mu}$ satisfies

$$Y \begin{array}{c} \xrightarrow{\mu_Y} \\ \xrightarrow[\widehat{\mu}]{\quad} \\ \xrightarrow{\quad} \end{array} M_f \xrightarrow{\quad} M_{f_1} \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow[\simeq]{\quad} \\ \xrightarrow{\quad} \end{array} Y_1 .$$

Let $J_r(M_f, X) \xrightarrow{J(\widehat{\mu}, \mu_X)|_{J_r(M_f, X)} = J_r(\widehat{\mu}, \mu_X)} J_r(M_{f_1}, X_1)$ ($r \geq 1$). Then we have the following homotopy commutative diagram

$$\begin{array}{ccccccc}
J_{n-1}(M_f, X) & \xrightarrow{I_{n-1}} & J_n(M_f, X) & \xrightarrow{proj.} & J_n(M_f, X)/J_{n-1}(M_f, X) & \xrightarrow{\cong} & Y \wedge X^{\wedge^{n-1}} \\
\downarrow J_{n-1}(\widehat{\mu}, \mu_X) & & \downarrow J_n(\widehat{\mu}, \mu_X) & & \downarrow \overline{J_n(\widehat{\mu}, \mu_X)} & & \downarrow \mu_Y \wedge \mu_X^{\wedge^{n-1}} \\
J_{n-1}(M_{f_1}, X_1) & \xrightarrow{I_{n-1}} & J_n(M_{f_1}, X_1) & \xrightarrow{proj.} & J_n(M_{f_1}, X_1)/J_{n-1}(M_{f_1}, X_1) & \xrightarrow{\cong} & Y_1 \wedge X_1^{\wedge^{n-1}}
\end{array}$$

2.3. homotopy of γ_n and Hopf invariant

Let $j_X^r : X = J_1(X, A) \hookrightarrow J_r(X, A)$ be the canonical inclusion. By Theorem 1.1, there is a cofibration sequence

$$\Sigma^{n-1}Y' \wedge X'^{\wedge(n-1)} \xrightarrow{\gamma_n} J_{n-1}(M_f, X) \xrightarrow{I_{n-1}} J_n(M_f, X)$$

where γ_n is an element of n -th order Whitehead products $[j_Y^{n-1}, j_Y^{n-1}f, \dots, j_Y^{n-1}f]$. Thus from Remark 2.2,

$$\Sigma\gamma_n = 0.$$

By [50, Remark 3.5] γ_n is natural in the sense of the following

Lemma 2.6. For $\mu_X = \Sigma\mu'_X : X = \Sigma X' \rightarrow X_1 = \Sigma X'_1$, $\mu_Y = \Sigma\mu'_Y : Y = \Sigma Y' \rightarrow Y_1 = \Sigma Y'_1$, there is the following commutative diagram

$$\begin{array}{ccccc}
\Sigma^{n-1}Y' \wedge X'^{\wedge(n-1)} & \xrightarrow{\gamma_n} & J_{n-1}(Y, X) & \xrightarrow{I_{n-1}} & J_n(Y, X) \\
\downarrow \Sigma^{n-1}\mu'_Y \wedge \mu'_X{}^{\wedge(n-1)} & & \downarrow J_{n-1}(\mu_Y, \mu_X) & & \downarrow J_n(\mu_Y, \mu_X) \\
\Sigma^{n-1}Y'_1 \wedge X'_1{}^{\wedge(n-1)} & \xrightarrow{\gamma_n} & J_{n-1}(Y_1, X_1) & \xrightarrow{I_{n-1}} & J_n(Y_1, X_1)
\end{array}$$

Define the n -th relative James-Hopf invariant

$$J(X, A) \xrightarrow{H_n} J(X \wedge A^{\wedge(n-1)}), x_1 x_2 \dots x_t \mapsto \prod_{1 \leq i_1 < i_2 < \dots < i_n \leq t} (x_{i_1} \wedge x_{i_2} \wedge \dots \wedge x_{i_n})$$

which are natural for pairs.

By abuse of notion, the second James-Hopf invariant H_2 also denotes the composition of the maps

$$\Omega\Sigma X \xrightarrow{\cong} J(X) = J(X, X) \xrightarrow{H_2} J(X \wedge X) \xrightarrow{\cong} \Omega\Sigma(X \wedge X) \quad (3)$$

where X is a CW-complex and let $H_2' : F_p \simeq J(M_f, X) \xrightarrow{H_2} J(M_f \wedge X) \xrightarrow{\cong} J(Y \wedge X) \simeq \Omega\Sigma(Y \wedge X)$. Then by Lemmas 2.5 and 2.7 of [49], we get

Lemma 2.7. *Let $X \xrightarrow{f} Y$ be a map. Then the following diagram is homotopy commutative*

$$\begin{array}{ccccc}
\Omega\Sigma X & \xrightarrow{\partial} & F_p \simeq J(M_f, X) & \xleftarrow{\quad} & J_2(M_f, X) \xrightarrow{\text{pinch}} J_2(M_f, X)/J_1(M_f, X) = M_f \wedge X \\
\downarrow H_2 & & \downarrow H'_2 & & \downarrow \\
\Omega\Sigma(X \wedge X) & \xrightarrow{\Omega\Sigma(f \wedge id_X)} & \Omega\Sigma(Y \wedge X) & \xleftarrow{\omega} & J(Y \wedge X) \simeq J(M_f \wedge X) .
\end{array}$$

Let X, Y be connected CW-complexes. Recall that the Hilton-Milnor theorem states that there is a homotopy equivalence

$$\Omega\Sigma(X \vee Y) \simeq \Omega\Sigma X \times \Omega\Sigma Y \times \Omega\Sigma\left(\bigvee_{i \geq 1} X^{\wedge i} \wedge Y\right).$$

Let $j_1: X \rightarrow X \vee Y$ and $j_2: Y \rightarrow X \vee Y$ be the canonical inclusion into wedge sum, respectively. Recall the following duality isomorphism

$$\Omega_0: [\Sigma X, Y] \rightarrow [X, \Omega Y], \quad \Omega_0 f = (\Omega f) \circ E_X.$$

For maps $f: \Sigma X \rightarrow Z, g: \Sigma Y \rightarrow Z$, the Whitehead product

$$[f, g]: \Sigma X \wedge Y \rightarrow Z$$

is the duality of the Samelson product

$$[\Omega_0 f, \Omega_0 g]^S: X \wedge Y \rightarrow \Omega Z;$$

that is, $\Omega_0[f, g] = [\Omega_0 f, \Omega_0 g]^S$. Let X be a connected CW co- H -space. Recall that the Hopf-Hilton invariant H ([5], Section 5.3 of [32]) is the composition

$$H: \Omega\Sigma X \xrightarrow{\Omega\Sigma(j_1+j_2)} \Omega\Sigma(X \vee X) \xrightarrow{p_{[j_1, j_2]}} \Omega\Sigma(X \wedge X),$$

where $p_{[j_1, j_2]}$ is the canonical projection after pre-composing with the homotopy equivalence given by the Hilton-Milnor theorem.

By Lemma 3.12 of [38], the Hopf-Hilton invariant H and the second James-Hopf invariant H_2 (absolute) are homotopic.

Lemma 2.8. *Let (X, A) be a CW-pair with $X = \Sigma X', A = \Sigma A'$. $i_A: A \hookrightarrow X$ is the inclusion map, then $p_{X \wedge A} \gamma_3 \simeq 0$, where $p_{X \wedge A}: J_2(X, A) \rightarrow J_2(X, A)/X \cong X \wedge A$ is the canonical projection. Moreover, if $[id_X, i_A] = 0$, then $J_2(X, A) \simeq X \vee X \wedge A$. $J_3(X, A) \simeq (X \vee X \wedge A) \cup_{\gamma_3} C(X' \wedge A \wedge A)$ such that the γ_3 also satisfies*

$$p_X \gamma_3 = [p_X, \mu p_{X \wedge A}] H_2(\gamma_3) \text{ for some } \mu \in [X \wedge A, X]$$

where $p_X: X \vee X \wedge A \rightarrow X$ is the canonical projection of the first wedge summand.

Proof. Let $j_X : X \hookrightarrow J_2(X, A)$ be the canonical inclusion. From Theorem 1.1, $\gamma_3 \in [j_X, j_X i_A, j_X i_A]$. Then from Proposition 2.3, $p_{X \wedge A} \gamma_3 \in p_{X \wedge A} [j_X, j_X i_A, j_X i_A] \subset [p_{X \wedge A} j_X, p_{X \wedge A} j_X i_A, p_{X \wedge A} j_X i_A] = [0, 0, 0]$. Since $0 \in [0, 0, 0]$ by [12, Corollary 2.6] and it is a coset of the subgroup $\{0\}$ by Lemma 2.4, $[0, 0, 0] = \{0\}$. Hence $p_{X \wedge A} \gamma_3 = 0$.

If $[id_X, i_A] = 0$, then $J_2(X, A) \simeq X \vee X \wedge A$. By Corollary 5.9 of [13], X is a wedge summand of $J(X, A)$, i.e., there is a projection $P_\infty : J(X, A) \rightarrow X$, such that $P_\infty I_X = id_X$ where $I_X : X = J_1(X, A) \hookrightarrow J(X, A)$ is the canonical inclusion which is also the following composition

$$\begin{array}{ccccccc} & & & I_X & & & \\ & & & \curvearrowright & & & \\ X & \xhookrightarrow{j_X} & J_2(X, A) & \xhookrightarrow{I_2} & J_3(X, A) & \xhookrightarrow{\quad} & J(X, A) \end{array}$$

Let $P_n : J_n(X, A) \hookrightarrow J(X, A) \xrightarrow{P_\infty} X$. $P_2 j_X = id_X$ implies that $P_2 = (id_X, \mu) = p_X + \mu p_{X \wedge A}$ for some $\mu \in [X \wedge A, X]$. By the following exact sequence

$$[J_3(X, A), X] \xrightarrow{I_2^*} [X \vee X \wedge A, X] \xrightarrow{\gamma_3^*} [X' \wedge A \wedge A, X]$$

we get $I_2^*(P_3) = P_2$. Thus $\gamma_3^*(P_2) = P_2 \gamma_3 = 0$, i.e., $(p_X + \mu p_{X \wedge A}) \gamma_3 = 0$. From the (4.11) of [5], we have $p_X \gamma_3 + \mu p_{X \wedge A} \gamma_3 + [\mu p_{X \wedge A}, p_X] H_2(\gamma_3) = 0$, i.e., $p_X \gamma_3 = [p_X, \mu p_{X \wedge A}] H_2(\gamma_3)$.

We complete the proof of this Lemma. \square

The above lemma easily gives the following corollary

Corollary 2.9. *Let $X \xrightarrow{f} Y$ be a map with $X = \Sigma X'$, $Y = \Sigma Y'$. Then $p_{Y \wedge X} \gamma_3 \simeq 0$, where $p_{Y \wedge X} : J_2(M_f, X) \rightarrow J_2(M_f, X)/Y \simeq Y \wedge X$ is the canonical projection. Moreover, if $[id_Y, f] = 0$, then $J_2(M_f, X) \simeq Y \vee Y \wedge X$. $J_3(M_f, X) \simeq (Y \vee Y \wedge X) \cup_{\gamma_3} C(Y' \wedge X \wedge X)$ such that the γ_3 also satisfies*

$$p_Y \gamma_3 = [p_Y, \mu p_{Y \wedge X}] H_2(\gamma_3) \text{ for some } \mu \in [Y \wedge X, Y]$$

where $p_Y : Y \vee Y \wedge X \rightarrow Y$ is the canonical projection of the first wedge summand.

Here we give the following conjecture which holds in all the special cases we have encountered and a lot of maps satisfy the condition of the conjecture, such as the attaching map of the top cell in the even dimensional mod 2 Moore space, suspended complex projective plane $\Sigma \mathbb{C}P^2$, quaternionic projective plane $\Sigma \mathbb{H}P^2$ and so on.

Conjecture 2.10. *Let $X \xrightarrow{f} Y$ be a map with $X = \Sigma X'$, $Y = \Sigma Y'$. If $[id_Y, f] = 0$, that is $J_2(M_f, A) \simeq Y \vee Y \wedge X$, then $\gamma_3 = [j_Y f, j_{Y \wedge X}] : Y' \wedge X \wedge X \rightarrow Y \vee Y \wedge X$, where j_Y and $j_{Y \wedge X}$ are canonical inclusions of the corresponding wedge summands of $J_2(M_f, X) \simeq X \vee X \wedge A$.*

Let $j_i : X_i \hookrightarrow \bigvee_{i=1}^n X_i$, $q_i : \bigvee_{i=1}^n X_i \twoheadrightarrow X_i$ (resp. $j_i : A_i \hookrightarrow \bigoplus_{i=1}^n A_i$, $q_i : \bigoplus_{i=1}^n A_i \twoheadrightarrow A_i$, A_i are abelian groups) are inclusion and projection of the i -th wedge summand X_i (rep. direct summand A_i). If X_i is a sphere S^m , denote j_i, q_i by j_i^m, q_i^m respectively. Let $(f_1, f_2, \dots, f_n) : \bigvee_{i=1}^n X_i \rightarrow Y$ be the map such that $(f_1, f_2, \dots, f_n)j_i \simeq f_i$ ($i = 1, 2, \dots, n$) and $g_1 \vee g_2 \vee \dots \vee g_n$ denote the map $(j_1 g_1, j_2 g_2, \dots, j_n g_n) : \bigvee_{i=1}^n X_i \rightarrow \bigvee_{i=1}^n Y_i$.

Consider the cofibration sequence $S^{n+k} \xrightarrow{\alpha} S^n \rightarrow C_\alpha \xrightarrow{p} S^{n+k+1}$. There is the cofibration sequence

$$\begin{aligned} S^{2n+k-1} &\xrightarrow{\gamma_2} S^n \xrightarrow{j_p} J_2(M_\alpha, S^{n+k}) \xrightarrow{\bar{p}} S^{2n+k}, \quad \gamma_2 = [\iota_n, \alpha]. \\ Sk_{5n+2k-3}(F_{\bar{p}}) &\simeq J_2(M_{\gamma_2}, S^{2n+k-1}) \simeq S^n \cup_{[\iota_n, [\iota_n, \alpha]]} e^{3n+k-1}. \end{aligned} \quad (4)$$

Moreover if $[\iota_n, [\iota_n, \alpha]] = 0$, then there is the inclusion of the wedge summand $\bar{\beta}_{3n+k-1} : S^{3n+k-1} \hookrightarrow J_2(M_{\gamma_2}, S^{2n+k-1}) \simeq S^n \vee S^{3n+k-1}$. Let β_{3n+k-1} be the composition of the following canonical inclusions

$$\beta_{3n+k-1} : S^{3n+k-1} \xrightarrow{\bar{\beta}_{3n+k-1}} J_2(M_{\gamma_2}, S^{2n+k-1}) \hookrightarrow F_{\bar{p}} \xrightarrow{\bar{\tau}} J_2(M_\alpha, S^{n+k}).$$

Lemma 2.11. *Let $S^{n+k} \xrightarrow{\alpha} S^n \rightarrow C_\alpha \xrightarrow{p} S^{n+k+1}$ be the cofibration sequence, with $k \geq 0, n \geq 2$ and $k < n - 1$.*

If $[\iota_n, [\iota_n, \alpha]] = 0$, i.e., $J_2(M_{\gamma_2}, S^{2n+k-1}) \simeq S^n \vee S^{3n+k-1}$, then

$$\gamma_3 = \beta_{3n+k-1}\theta + j_p\lambda : S^{3n+2k-1} \rightarrow J_2(M_\alpha, S^{n+k}), \theta \in \pi_{3n+2k-1}(S^{3n+k-1}), \lambda \in \pi_{3n+2k-1}(S^n).$$

Especially, if $[\iota_n, \alpha] = 0$, i.e., $J_2(M_\alpha, S^{n+k}) \simeq S^n \vee S^{2n+k}$, then $\gamma_3 \simeq [j_1^n, j_2^{2n+k}]\theta + j_1^n\lambda$.

Proof. The fibration sequence $\Omega S^{2n+k} \xrightarrow{\bar{\delta}} F_{\bar{p}} \xrightarrow{\bar{\tau}} J_2(M_\alpha, S^{n+k}) \xrightarrow{\bar{p}} S^{2n+k}$ induces the exact sequence with commutative square

$$\pi_{3n+2k}(S^{2n+k}) \xrightarrow{\bar{\delta}_*} \pi_{3n+2k-1}(F_{\bar{p}}) \xrightarrow{\bar{\tau}_*} \pi_{3n+2k-1}(J_2(M_\alpha, S^{n+k})) \xrightarrow{\bar{p}_*} \pi_{3n+k-1}(S^{2n+k})$$

By (4), $\pi_{3n+2k-1}(F_{\bar{p}}) = j_{\bar{p}*}\pi_{3n+2k-1}(S^n) \oplus \bar{\beta}_{3n+k-1*}\pi_{3n+2k-1}(S^{3n+k-1})$, $k < n - 1$, where $\bar{\tau}j_{\bar{p}} \simeq j_p$ by (2). From Corollary 2.9, $\bar{p}\gamma_3 \simeq 0$, that is,

$$\gamma_3 \in \bar{\tau}_*(\pi_{3n+2k-1}(F_{\bar{p}})) = j_p \circ \pi_{3n+2k-1}(S^n) + \bar{\beta}_{3n+k-1} \circ \pi_{3n+2k-1}(S^{3n+k-1}).$$

This implies the first conclusion of this lemma. \square

Remark 2.12. *For the case $[\iota_n, \alpha] \simeq 0$ in Lemma 2.11, if $\Sigma : \pi_{3n+2k-1}(S^n) \rightarrow \pi_{3n+2k}(S^{n+1})$ is an injection, then by $\Sigma\gamma_3 = 0$, we get $j_1^{n+1}\Sigma\lambda = 0$ which implies $\lambda = 0$. Hence $\gamma_3 = [j_1^n, j_2^{2n+k}]\theta$ for some $\theta \in \pi_{3n+2k-1}(S^{3n+k-1})$.*

For a map $f: X \rightarrow Y$, denote by f_E the composition

$$f_E = E_Y \circ f = (\Omega\Sigma f) \circ E_X = \Omega_0(\Sigma f): X \rightarrow \Omega\Sigma Y.$$

Lemma 2.13. *Let $H_*: \pi_{m+n}(\Omega\Sigma(S^m \vee S^n)) \rightarrow \pi_{m+n}(\Omega\Sigma(S^m \vee S^n)^{\wedge 2})$ be the homomorphism induced by the second James-Hopf invariant. Then*

$$H_*([(j_1^m)_E, (j_2^n)_E]^S) = (j_1^m \wedge j_2^n)_E - (j_2^n \wedge j_1^m)_E.$$

Thus we have

$$\begin{aligned} H_2([(j_1^{m+1}, j_2^{n+1})]) &= \Sigma j_1^m \wedge j_2^n - \Sigma j_2^n \wedge j_1^m, \\ \text{where } H_2: \pi_{n+m+1}(S^{n+1} \vee S^{m+1}) &\rightarrow \pi_{n+m+1}(\Sigma(S^m \vee S^n)^{\wedge 2}) \end{aligned}$$

is the induced by the second James-Hopf invariant.

Proof. There are isomorphisms

$$\begin{aligned} H_{m+n}(S^{m+n}) &\xrightarrow{\tau_{m,n}} H_{m+n}(S^m \wedge S^n) \cong H_m(S^m) \otimes H_n(S^n): \\ \iota_{m+n} &\mapsto (-1)^k \cdot \iota_m \otimes \iota_n, \quad k \in \mathbb{Z}. \end{aligned}$$

In the following we simplify j_1^m, j_2^n by j_1, j_2 respectively. By the Bott-Samelson theorem,

$$\begin{aligned} H_{m+n}(\Omega\Sigma(S^m \vee S^n)) &\cong T\langle (j_1)_{E*}(\iota_m), (j_2)_{E*}(\iota_n) \rangle, \\ H_{m+n}(\Omega\Sigma(S^m \vee S^n)^{\wedge 2}) &\cong T\langle (j_t \wedge j_s)_{E*}(\iota_m \otimes \iota_n) \mid t, s = 1, 2 \rangle, \end{aligned}$$

where $T\langle x_1, \dots, x_m \rangle$ is the free tensor algebra $T(V)$ and V is the free abelian group generated by x_1, \dots, x_m .

Let $h_X: \pi_{m+n}(\Omega\Sigma X) \rightarrow H_{m+n}(\Omega\Sigma X)$ be the Hurewicz homomorphism with $X = S^m \vee S^n$. We have

$$\begin{aligned} h_X([(j_1)_E, (j_2)_E]^S) &= (-1)^k [(j_1)_E, (j_2)_E]_*^S(\iota_m \otimes \iota_n) \\ &= (-1)^k ((j_1)_{E*}(\iota_m) \otimes (j_2)_{E*}(\iota_n) - (j_2)_{E*}(\iota_n) \otimes (j_1)_{E*}(\iota_m)), \\ h_{X^{\wedge 2}}((j_t \wedge j_s)_E) &= (-1)^k (j_t \wedge j_s)_{E*}(\iota_m \otimes \iota_n), \quad t, s = 1, 2. \end{aligned}$$

Set $H_*([(j_1)_E, (j_2)_E]^S) = \sum_{t=1, s=1}^{m,n} x_{ts} \cdot (j_t \wedge j_s)_E, x_{ts} \in \mathbb{Z}$. Then

$$\begin{aligned} h_{X^{\wedge 2}} H_*([(j_1)_E, (j_2)_E]^S) &= h_{X^{\wedge 2}} \left(\sum_{t,s} x_{ts} \cdot (j_t \wedge j_s)_E \right) \\ &= (-1)^k \sum_{t,s} x_{ts} \cdot (j_t \wedge j_s)_{E*}(\iota_m \otimes \iota_n), \\ H_* h_X([(j_1)_E, (j_2)_E]^S) &= (-1)^k H_*((j_1)_{E*}(\iota_m) \otimes (j_2)_{E*}(\iota_n) - (j_2)_{E*}(\iota_n) \otimes (j_1)_{E*}(\iota_m)) \\ &= (-1)^k \overline{((j_1)_{E*}(\iota_m) \otimes (j_2)_{E*}(\iota_n) - (j_2)_{E*}(\iota_n) \otimes (j_1)_{E*}(\iota_m))} \\ &= (-1)^k ((j_1 \wedge j_2)_{E*}(\iota_m \otimes \iota_n) - (j_2 \wedge j_1)_{E*}(\iota_n \otimes \iota_m)), \end{aligned}$$

where the second to last equality holds by [32, Lemma 5.2.5]. Thus from the equality

$$H_*h_X([(t_1)_E, (t_2)_E]^S) = h_{X \wedge 2}H_*([(t_1)_E, (t_2)_E]^S)$$

we get $x_{11} = x_{22} = 0$, $x_{12} = 1$, $x_{21} = -1$. \square

3. homotopy of $\Omega\Sigma X \xrightarrow{\partial} F_p$ and extension problem

In this section we give some lemmas for the homotopy properties of the connection map $\Omega\Sigma X \xrightarrow{\partial} F_p$ and for the extension problem of the short exact sequence $0 \rightarrow \text{Coker}(\partial_{k*}) \rightarrow \pi_k(C_f) \rightarrow \text{Ker}(\partial_{k-1*}) \rightarrow 0$.

The following lemma comes from [49, Lemma 2.3], which generalizes the Lemma 4.4.1 of [45].

Lemma 3.1. *Let $f : X \rightarrow Y$ be a map, then for the following fibration sequence,*

$$\Omega\Sigma X \xrightarrow{\partial} J(M_f, X) \simeq F_p \rightarrow C_f \xrightarrow{p} \Sigma X$$

there exists homotopy-commutative diagram,

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Omega\Sigma \downarrow & & \downarrow j_p \\ \Omega\Sigma X & \xrightarrow{\partial} & J(M_f, X) \simeq F_p \end{array}$$

Lemma 3.2. *Let $f : X \rightarrow Y$ be a map, $\beta \in \pi_n(S^m)$. Then $\partial_{n*}(\alpha\Sigma\beta) = \partial_{m*}(\alpha)\beta$ for any $\alpha \in \pi_{m+1}(\Sigma X)$, i.e., the following diagram commutes*

$$\begin{array}{ccc} \pi_{m+1}(\Sigma X) & \xrightarrow{\partial_{m*}} & \pi_m(F_p) \\ (\Sigma\beta)^* \downarrow & & \downarrow \beta_* \\ \pi_{n+1}(\Sigma X) & \xrightarrow{\partial_{n*}} & \pi_n(F_p) \end{array} .$$

Proof. The lemma is obtained by the following equations

$$\begin{aligned} \partial_{m*}(\alpha\Sigma\beta) &= \partial\Omega_0(\alpha\Sigma\beta) = \partial\Omega(\alpha\Sigma\beta)E_{S^n} \simeq \partial\Omega(\alpha)(\Omega\Sigma\beta)E_{S^n} \\ &= \partial\Omega(\alpha)E_{S^m}\beta = \partial\Omega_0(\alpha)\beta = \partial_{n*}(\alpha)\beta. \end{aligned}$$

\square

Lemma 3.3. *Let $S^m \xrightarrow{\alpha} S^n \rightarrow C_\alpha \xrightarrow{p} S^{m+1}$ be the cofibration sequence, with $m \geq n \geq 2$. Consider the fibration sequence $\Omega S^{m+k+1} \xrightarrow{\partial^k} F_{\Sigma^k p} \xrightarrow{i^k} \Sigma^k C_\alpha \xrightarrow{\Sigma^k p} S^{m+k+1}$. The map*

$E^k : F_p \rightarrow \Omega^k F_{\Sigma^k p}$ which is the restriction of $\Omega^k \Sigma^k : C_\alpha \rightarrow \Omega^k \Sigma^k C_\alpha$ on the fiber satisfies the following left homotopy commutative diagram of fibration sequences

$$\begin{array}{ccccccc} \Omega S^{m+1} & \xrightarrow{\partial} & F_p & \xrightarrow{i} & C_\alpha & \xrightarrow{p} & S^{m+1} \\ \downarrow \Omega(\Omega^k \Sigma^k) & & \downarrow E^k & & \downarrow \Omega^k \Sigma^k & & \downarrow \Omega^k \Sigma^k \\ \Omega^{k+1} S^{m+k+1} & \xrightarrow{\Omega^k \partial^k} & \Omega^k F_{\Sigma^k p} & \xrightarrow{\Omega^k i^k} & \Omega^k \Sigma^k C_\alpha & \xrightarrow{\Omega^k \Sigma^k p} & \Omega^k S^{m+k+1}, \end{array} \quad \begin{array}{ccc} S^n & \xrightarrow{j_p} & F_p \\ \downarrow a \Omega^k \Sigma^k & & \downarrow E^k \\ \Omega^k S^{n+k} & \xrightarrow{\Omega^k j_{\Sigma^k p}} & \Omega^k F_{\Sigma^k p} \end{array}$$

where $\Omega^k \Sigma^k$ represents the k -fold canonical inclusions. Then E^k satisfies the above right homotopy commutative diagram with $a = \pm 1$ for $m > n$ and $(a, t) = 1$ for $m = n$ and $\alpha = t\tau_n$.

Proof. Since $Sk_n(\Omega^k F_{\Sigma^k p}) = S^n \hookrightarrow \Omega^k F_{\Sigma^k p}$ is homotopic to the composition of the maps $S^n \xrightarrow{\Omega^k \Sigma^k} \Omega^k S^{n+k} \xrightarrow{\Omega^k j_{\Sigma^k p}} \Omega^k F_{\Sigma^k p}$, there is an integer a such that the right square of the lemma is homotopy commutative. Then the condition of a in the lemma is obtained by observing the commutative diagram of n -dimensional homotopy groups which is induced by the following homotopy commutative diagram

$$\begin{array}{ccccc} & & i & & \\ & & \curvearrowright & & \\ S^n & \xrightarrow{j_p} & F_p & \xrightarrow{i} & C_\alpha \\ \downarrow \Omega(\Omega^k \Sigma^k) & & \downarrow E^k & & \downarrow \Omega^k \Sigma^k \\ \Omega^k S^{n+k} & \xrightarrow{\Omega^k j_{\Sigma^k p}} & \Omega^k F_{\Sigma^k p} & \xrightarrow{\Omega^k i^k} & \Omega^k \Sigma^k C_\alpha. \\ & & \curvearrowleft & & \\ & & \Omega^k \Sigma^k i & & \end{array}$$

□

If k is large enough in above lemma, we get the following corollary

Corollary 3.4. *For $a \in \mathbb{Z}$ is given in Lemma 3.3, there is the following two commutative squares*

$$\begin{array}{ccc} \pi_{u+1}(S^{m+1}) & \xrightarrow{\partial_{u*}} & \pi_u(F_p) \\ \Sigma^\infty \downarrow & & E^\infty \downarrow \\ \pi_u^s(S^m) & \xrightarrow{\alpha_*} & \pi_u^s(S^n); \end{array} \quad \begin{array}{ccc} \pi_u(S^n) & \xrightarrow{j_{p*}} & \pi_u(F_p) \\ \searrow a \Sigma^\infty & & \downarrow E^\infty \\ & & \pi_u^s(S^n). \end{array}$$

Lemma 3.5. *Let $S^m \xrightarrow{f} C_\alpha \rightarrow C_f \xrightarrow{p} S^{m+1}$ be the cofibration sequence, where $C_\alpha = S^n \cup_\alpha e^{n+k+1}$ with $k \geq 1, n \geq 2, m > n, k + 2$. F_p is the homotopy fiber of map p . Then*

$$Sk_{n+m} F_p \simeq C_\alpha \cup_{[i_\alpha, f]} e^{n+m}$$

where $i_\alpha : S^n \hookrightarrow C_\alpha$ is the canonical inclusion.

Proof. $F_p \simeq J(M_f, S^m)$ where $J_1(M_f, S^m) = M_f \simeq C_\alpha$. $\Sigma J_2(M_f, S^m) \simeq \Sigma C_\alpha \vee \Sigma C_\alpha \wedge S^m = S^{n+1} \cup e^{n+k+2} \cup e^{n+m+1} \cup e^{n+k+m+2}$. By the natural inclusion $J_1(M_f, S^m) \hookrightarrow J_2(M_f, S^m)$, we have

$$\begin{aligned} Sk_{n+m-1}F_p &\simeq Sk_{n+m-1}J_2(M_f, S^m) \simeq J_1(M_f, S^m) \simeq C_\alpha; \\ Sk_{n+k+m}F_p &\simeq Sk_{n+m}J_2(M_f, S^m) \simeq C_\alpha \cup_\beta e^{n+m}, \text{ for some } \beta \in \pi_{n+m-1}(C_\alpha). \end{aligned}$$

We have the following two commutative diagrams for cofibration sequences left and fibration sequences right respectively.

$$\begin{array}{ccc} S^m & \xrightarrow{f} & C_\alpha \longrightarrow C_f \xrightarrow{p} S^{m+1}, & \Omega S^{m+1} & \longrightarrow & F_p & \longrightarrow & C_f \\ \parallel & & \uparrow (i_\alpha, f) & \bar{\omega} \uparrow & & \uparrow \omega & & \bar{\omega} \uparrow \\ S^m & \xrightarrow{j_2^m} & S^n \vee S^m \xrightarrow{\iota_n \vee i_C} S^n \vee C_{j_2^m} \xrightarrow{p_\vee} S^{m+1} & & \Omega S^{m+1} & \longrightarrow & F_{p_\vee} & \longrightarrow & S^n \vee C_{j_2^m} \end{array},$$

where the homotopy fiber $F_{p_\vee} \simeq J(M_{j_2^m}, S^m)$ and $J_2(M_{j_2^m}, S^m) \simeq (S^n \vee S^m) \cup_{[id, j_2^m]} C(S^n \vee S^m) \wedge S^{m-1} \simeq (S^n \vee S^m) \cup_{([j_1^n, j_2^m], [j_2^m, j_2^m])} C(S^{n+m-1} \vee S^{2m-1})$.

Consider the following homotopy commutative diagram of the cofibration sequences

$$\begin{array}{ccccc} J_1(M_f, S^m) \simeq C_\alpha & \xrightarrow{j_p} & J_2(M_f, S^m) & \longrightarrow & C_\alpha \wedge S^m \\ \uparrow \omega|_{J_1} = (i_\alpha, f) & & \uparrow \omega|_{J_2} & & \uparrow (i_\alpha, f) \wedge \iota_m \\ J_1(M_{j_2^m}, S^m) \simeq S^n \vee S^m & \xrightarrow{j_{p_\vee}} & J_2(M_{j_2^m}, S^m) & \longrightarrow & (S^n \vee S^m) \wedge S^m. \end{array}$$

It is easy to get that the induced homomorphism $H_{n+m}((i_\alpha, f) \wedge \iota_m)$ is isomorphic and so is $H_{n+m}(\omega|_{J_2})$.

We can decompose the following two inclusions j_p and j_{p_\vee} by

$$\begin{aligned} j_p &: C_\alpha \xrightarrow{j'_p} Sk_{n+m}J_2(M_f, S^m) \hookrightarrow J_2(M_f, S^m), \\ j_{p_\vee} &: S^n \vee S^m \xrightarrow{j'_{p_\vee}} Sk_{n+m}J_2(M_{j_2^m}, S^m) \hookrightarrow J_2(M_{j_2^m}, S^m) \end{aligned}$$

We have the following homotopy commutative diagram of the cofibration sequences

$$\begin{array}{ccccccc} S^{n+m-1} & \xrightarrow{\beta} & J_1(M_f, S^m) \simeq C_\alpha & \xrightarrow{j'_p} & Sk_{n+m}J_2(M_f, S^m) & \xrightarrow{proj} & S^{n+m} \\ \uparrow \theta & & \uparrow \omega|_{J_1} & & \uparrow (\omega|_{J_2})|_{Sk_{n+m}} & & \uparrow \Sigma\theta \\ S^{n+m-1} & \xrightarrow{[j_1^n, j_2^m]} & J_1(M_{j_2^m}, S^m) \simeq S^n \vee S^m & \xrightarrow{j'_{p_\vee}} & Sk_{n+m}J_2(M_{j_2^m}, S^m) & \xrightarrow{proj} & S^{n+m} \end{array}$$

where $(\omega|_{J_2})|_{Sk_{n+m}}$ is the restriction of $\omega|_{J_2}$ to the $(n+m)$ -skeleton and there is a θ making the left square homotopy commutative because the following sequence

$\pi_{n+m-1}(S^{n+m-1}) \xrightarrow{\beta_*} \pi_{n+m-1}(C_\alpha) \xrightarrow{j'_{p*}} \pi_{n+m-1}(Sk_{n+m}J_2(M_f, S^m))$ is exact by Theorem 1.16 of [7] and $j'_{p*}(\omega|_{J_1}[j_1^n, j_2^m]) = 0$.

The isomorphism $H_{n+m}(\omega|_{J_2})$ implies that $H_{n+m}((\omega|_{J_2})|_{Sk_{n+m}})$ is isomorphic. Thus we get $H_{n+m}(\Sigma\theta)$ is also isomorphic and so is the $H_{n+m-1}(\theta)$. Hence $\theta = \pm\iota_{n+m-1}$. Then

$$\beta = \pm(\omega|_{J_1})[j_1^n, j_2^m] = \pm(i_\alpha, f)[j_1^n, j_2^m] = \pm[i_\alpha, f].$$

Since the sign \pm in $\beta = \pm[i_\alpha, f]$ does not effect the homotopy type of $Sk_{n+m}J_2(M_f, S^m) \simeq C_\alpha \cup_\beta e^{n+m}$, we choose $\beta = [i_\alpha, f]$. It completes the proof of this lemma. \square

Note. In above lemma, C_α may not be a suspension, so we can not use the Gray's Theorem to get $J_2(M_f, S^m) \simeq C_\alpha \cup_{[id, f]} C(C_\alpha \wedge S^{m-1})$.

The following two lemmas will be used to solve the extension problem of the short exact sequence of the homotopy groups.

Lemma 3.6 ([23, Lemma 2.7]). Let $X \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{p} \Sigma X$ be a cofibration sequence of CW complexes. Consider the fibration sequence $\Omega\Sigma^{k+1}X \xrightarrow{\partial^k} F_{\Sigma^k p} \xrightarrow{i^k} \Sigma^k C_f \xrightarrow{\Sigma^k p} \Sigma^{k+1}X$. Assume that the following restriction of $\Sigma^k : \pi_n(\Sigma X) \rightarrow \pi_{n+k}(\Sigma^{k+1}X)$ is isomorphic

$$Ker(\partial_{n-1*} : \pi_n(\Sigma X) \rightarrow \pi_{n-1}(F_p)) \xrightarrow{\Sigma^k} Ker(\partial_{n+k-1*}^k : \pi_{n+k}(\Sigma^{k+1}X) \rightarrow \pi_{n+k-1}(F_{\Sigma^k p}))$$

If $0 \rightarrow Coker\partial_{n*} \rightarrow \pi_n(C_f) \rightarrow Ker\partial_{n-1*} \rightarrow 0$ is split, then so is the short exact sequence

$$0 \rightarrow Coker\partial_{n+k*}^k \rightarrow \pi_n(\Sigma^k C_f) \rightarrow Ker\partial_{n+k-1*}^k \rightarrow 0.$$

Lemma 3.7. For a 2-local suspension space X which is 2-cell complexes such that the reduced homology $\bar{H}_*(X, \mathbb{Z}_2)$ of dim 2 with generators u and v such that $|u| < |v|$, where $|x|$ denotes the degree of $x \in \bar{H}_*(X, \mathbb{Z}_2)$. Then $\pi_i(\Sigma^{1+|u|+|v|}X)$ is a direct summand of $\pi_i(\Sigma X)$ for any positive integer i .

Proof. By Proposition 3.1 of [6], $L_3(X) \simeq \Sigma^{|u|+|v|}X$, where $L_3(X)$ is defined in Notation 1.5 of [42]. Then by Corollary 4.1 of [42],

$$\Omega\Sigma X \simeq \Omega\Sigma^{1+|u|+|v|}X \times \text{other space.}$$

This implies the Lemma 3.7. \square

Lemma 3.8. Let $S^{n+k} \xrightarrow{\alpha} S^n \xrightarrow{i} C_\alpha \xrightarrow{p} S^{n+k+1}$ be the cofibration sequence, with $k \geq 0, n \geq 2$. For any $\Sigma\theta \in \pi_m(S^{n+k+1})$ with $\alpha\theta = 0$, there is an element $-\mathbf{x}$ in the Toda bracket $\{i, \alpha, \theta\} \subset \pi_m(C_\alpha)$, such that $p(\mathbf{x}) = \Sigma\theta$.

Proof. From Corollary 2.5.1 of [45], $\iota_{n+k+1} \in \{p, i, \alpha\}$. By Proposition 1.4 of [39], $-\Sigma\theta \in -\{p, i, \alpha\}\Sigma\theta = p\{i, \alpha, \theta\}$. This implies the existence of \mathbf{x} . \square

Lemma 3.9. *r is a positive integer. Assume $2^r \beta = 0$ for an $\beta \in \pi_{m+k}(S^m), k > 0$. Then for $\alpha \in \pi_{m+1}(S^{n_1}), \gamma \in \pi_{n_2}(S^{m+k})$,*

$$\{2^r \alpha, \Sigma \beta, 2^r \Sigma \gamma\} \ni \begin{cases} \alpha \Sigma \beta \eta_{m+k+1} \Sigma^2 \gamma, & r = 1; \\ 0, & r \geq 2. \end{cases}$$

Proof.

$$\begin{aligned} \{2^r \alpha, \Sigma \beta, 2^r \Sigma \gamma\} &= \{\alpha(2^r \iota_{m+1}), \Sigma \beta, (2^r \iota_{m+k+1}) \Sigma \gamma\} \supset \alpha \{2^r \iota_{m+1}, \Sigma \beta, 2^r \iota_{m+k+1}\} \Sigma^2 \gamma \\ &\supset \alpha \{2^r \iota_{m+1}, \Sigma \beta, 2^r \iota_{m+k+1}\}_1 \Sigma^2 \gamma \ni \begin{cases} \alpha \Sigma \beta \eta_{m+k+1} \Sigma^2 \gamma, & r = 1; \\ 0, & r \geq 2 \end{cases} \end{aligned}$$

where the above first “ \supset ” comes from [39, Proposition 1.2]; the second “ \supset ” comes from [39, (1.15)] and the last “ \ni ” comes from [39, Corollary 3.7]. \square

4. Skeletons of the homotopy fiber

For an abelian group A , $A^{\oplus k}$ denotes the direct sum of k -copies of A . $\langle \alpha_1, \alpha_2, \dots, \alpha_s \rangle$ denotes the subgroup of A which is generated by $\alpha_1, \alpha_2, \dots, \alpha_s \in A$. Let A_1 be a subgroup of A . Then “ $a \equiv b \pmod{A_1}$ ” means $a - b \in A_1$ and “ $a \in H \pmod{A_1}$ ” or “ $H \ni a \pmod{A_1}$ ” means H , which contains an element a , is a coset of subgroup A_1 . For a map $f : X \rightarrow Y$ and $\alpha \in [Z, Y]$, if there is an $\tilde{\alpha} \in [Z, X]$ such that $f_*(\tilde{\alpha}) = p\tilde{\alpha} = \alpha$, then $\tilde{\alpha}$ is called the lift of α by f .

- For $P^{k+1}(2^r)$, there is a canonical cofibration sequence

$$S^k \xrightarrow{2^r \iota_k} S^k \xrightarrow{i_k} P^{k+1}(2^r) \xrightarrow{p_{k+1}} S^{k+1} \quad (5)$$

Note that for $r = 0$, $P^{k+1}(2^0) = *$ is contractible. We simplify the notation $F_{p_{k+1}}$ by F_{k+1} , which is the homotopy fiber of the pinch map p_{k+1} above, by denoting $J_{i,k}^r := J_i(M_{2^r \iota_k}, S^k)$, we have

$$\begin{aligned} Sk_{3k-1}(F_{k+1}) &\simeq J_{2,k}^r \simeq S^k \cup_{\gamma_2^r} CS^{2k-1}, \quad \gamma_2^r = 2^r [\iota_k, \iota_k]; \\ Sk_{4k-1}(F_{k+1}) &\simeq J_{3,k}^r \simeq J_{2,k}^r \cup_{\gamma_3^r} CS^{3k-1} \end{aligned}$$

We have the natural inclusion $I_{i-1}^r : J_{i-1,k}^r \rightarrow J_{i,k}^r$. and denote the natural inclusion $J_{i,k}^r \rightarrow F_{k+1}$ by \check{I}_i^r . So we have $\check{I}_i^r I_{i-1}^r = \check{I}_{i-1}^r$.

Let $\beta_k = j_{p_{k+1}} : S^k \hookrightarrow J_{2,k}^r$ and $\bar{p}_{2k} : J_{2,k}^r \rightarrow S^{2k}$ be the canonical pinch map.

Moreover if $\gamma_2^r = 2^r [\iota_k, \iota_k] \simeq 0$, then there is a canonical inclusion $\beta_{2k} : S^{2k} \hookrightarrow J_{2,k}^r \simeq S^k \vee S^{2k}$ and in this case, $\beta_k = j_1^k, \bar{p}_{2k} = q_2^{2k}$.

Let $\Omega S^{k+1} \xrightarrow{\partial^{k+1,r}} F_{k+1} \xrightarrow{\tau_{k+1}} P^{k+1}(2^r) \xrightarrow{p_{k+1}} S^{k+1}$ be the homotopy fibration sequence which implies the following exact sequence with homotopy commutative squares by

Lemma 3.1

$$\begin{array}{ccccc}
\pi_{m+1}(S^{k+1}) & \xrightarrow{\partial_{m*}^{k+1,r}} & \pi_m(F_{k+1}) & \xrightarrow{\tau_{k+1*}} & \pi_m(P^{k+1}(2^r)) & \xrightarrow{p_{k+1*}} & \pi_m(S^{k+1}) & \xrightarrow{\partial_{m-1*}^{k+1,r}} & \pi_{m-1}(F_{k+1}) & (6) \\
\uparrow \Sigma & & \uparrow \beta_{k*} & & & & \uparrow \Sigma & & \uparrow \beta_{k*} \\
\pi_m(S^k) & \xrightarrow{(2^r \iota_k)_*} & \pi_m(S^k) & & & & \pi_{m-1}(S^k) & \xrightarrow{(2^r \iota_k)_*} & \pi_{m-1}(S^k)
\end{array}$$

For any $\alpha \in \text{Ker}(\partial_{m-1*}^{k+1,r})$, We always denote the $\tilde{\alpha} \in \pi_m(P^{k+1}(2^r))$ a lift of α by p_{k+1} , i.e., $p_{k+1*}(\tilde{\alpha}) = \alpha$.

- For $J_{2,k}^r$, there are the following cofibration and fibration sequences

$$S^{2k-1} \xrightarrow{\gamma_2^r} S^k \xrightarrow{\beta_k} J_{2,k}^r \xrightarrow{\bar{p}_{2k}} S^{2k}; \quad \Omega S^{2k} \xrightarrow{\bar{\partial}_{2k,r}^{2k}} F_{\bar{p}_{2k}} \xrightarrow{\bar{\tau}_{2k}} J_{2,k}^r \xrightarrow{\bar{p}_{2k}} S^{2k}.$$

If $[\iota_k, [\iota_k, \iota_k]] = 0$ (this holds under 2-localization or for odd k by [41, Theorem 8.8 in Chapter XI]), then $[\iota_k, \gamma_2] = 0$. Denoting $J_2(M_{\gamma_2}, S^{2k-1})$ by $J_{2,k}^{\gamma,r}$ and $I_2^{\gamma,r} : J_{2,k}^{\gamma,r} \rightarrow F_{\bar{p}_{2k}}$ the canonical inclusion.

$$Sk_{5k-3}(F_{\bar{p}_{2k}}) \simeq J_{2,k}^{\gamma,r} \simeq S^k \vee S^{3k-1}. \quad (7)$$

Let $\bar{\beta}_i : S^i \hookrightarrow S^k \vee S^{3k-1} \simeq J_{2,k}^{\gamma,r} \simeq Sk_{5k-3}(F_{\bar{p}_{2k}}) \hookrightarrow F_{\bar{p}_{2k}}$ ($i = k, 3k-1$), where the first map is the canonical inclusions of the corresponding wedge summands.

Note that $\bar{\tau}_{2k}\bar{\beta}_k \simeq \beta_k$. We also denote $\beta_{3k-1} := \bar{\tau}_{2k}\bar{\beta}_{3k-1}$

Similar as the definition of j_p in Section 2.2, we also denote the composition

$$S^i \xrightarrow{\beta_i} J_{2,k}^r \xrightarrow{\bar{i}_2^r} F_p \text{ by } \beta_i \text{ without ambiguous for } i = k, 2k, 3k-1.$$

Let $\text{Im}(h) := h(G)$ denote the Image of the homomorphism $h : G \rightarrow H$ of abelian groups.

Lemma 4.1. *Let $m \leq 4k-3$, there is a short exact sequence*

$$0 \rightarrow \frac{\bar{\beta}_{k*}(\pi_m(S^k))}{\bar{\beta}_{k*}\gamma_{2*}^r(\pi_m(S^{2k-1}))} \oplus G_m^k \xrightarrow{\bar{\tau}_{2k*}} \pi_m(J_{2,k}^r) \xrightarrow{\bar{p}_{2k*}} \Sigma(\text{Ker}(\gamma_{2*}^r)) \rightarrow 0$$

where the right γ_{2*}^r is the induced homomorphism on $\pi_{m-1}(-)$ by γ_2^r and $G_m^k = \begin{cases} 0, & m \leq 3k-2; \\ \bar{\beta}_{3k-1*}(\pi_m(S^{3k-1})), & 3k-2 \leq m \leq 4k-3. \end{cases}$

Proof. By Lemma 3.1, we have exact sequence with commutative squares

$$\begin{array}{ccccccc}
\pi_{m+1}(S^{2k}) & \xrightarrow{\bar{\partial}_{m*}^{2k,r}} & \pi_m(F_{\bar{p}_{2k}}) & \xrightarrow{\bar{\tau}_{2k*}} & \pi_m(J_{2,k}^r) & \xrightarrow{\bar{p}_{2k*}} & \pi_m(S^{2k}) & \xrightarrow{\bar{\partial}_{m-1*}^{2k,r}} & \pi_{m-1}(F_{\bar{p}_{2k}}) \\
\uparrow \Sigma & & \uparrow \bar{\beta}_{k*} & \nearrow \beta_{k*} & & & \uparrow \cong \Sigma & & \uparrow \bar{\beta}_{k*} \\
\pi_m(S^{2k-1}) & \xrightarrow{\gamma_{2*}^r} & \pi_m(S^k) & & & & \pi_{m-1}(S^{2k-1}) & \xrightarrow{\gamma_{2*}^r} & \pi_{m-1}(S^k)
\end{array}$$

where the suspension homomorphisms Σ in the left and right squares are epimorphism and isomorphism respectively by the Freudenthal Suspension Theorem. Note that

$$\pi_m(F_{\bar{p}_{2k}}) = \bar{\beta}_{k*}(\pi_m(S^k)) \oplus \bar{\beta}_{3k-1*}(\pi_m(S^{3k-1}))(m \leq 4k-3),$$

Thus we complete the proof of this lemma by the following

$$\begin{aligned} \text{Coker}(\bar{\delta}_{m*}^{2k,r}) &= \frac{\pi_m(F_{\bar{p}_{2k}})}{\text{Im}(\bar{\beta}_{k*}\gamma_{2*}^r)} = \frac{\bar{\beta}_{k*}(\pi_m(S^k))}{\bar{\beta}_{k*}\gamma_{2*}^r(\pi_m(S^{2k-1}))} \oplus \bar{\beta}_{3k-1*}(\pi_m(S^{3k-1})); \\ \text{Ker}(\bar{\delta}_{m-1*}^{2k,r}) &= \Sigma(\text{Ker}(\gamma_{2*}^r)). \end{aligned}$$

□

Remark 4.2. Take $m = 3k - 1$ in Lemma 4.1, we get short exact sequence

$$0 \rightarrow \frac{\bar{\beta}_{k*}(\pi_{3k-1}(S^k))}{\bar{\beta}_{k*}\gamma_{2*}^r(\pi_{3k-1}(S^{2k-1}))} \oplus \mathbb{Z}_{(2)}\{\bar{\beta}_{3k-1}\} \xrightarrow{\bar{\tau}_{2k*}} \pi_{3k-1}(J_{2,k}^r) \xrightarrow{\bar{p}_{2k*}} \Sigma(\text{Ker}(\gamma_{2*}^r)) \rightarrow 0$$

where the right γ_{2*}^r is the induced homomorphism on $\pi_{3k-2}(-)$ by γ_2^r .

Note that in the following text, for any $\alpha \in \Sigma(\text{Ker}(\gamma_{2*}^r))$, we always denote the $\widehat{\alpha} \in \pi_{3k-1}(J_2(2^r t_k))$ a lift of α by \bar{p}_{2k} , i.e., $\bar{p}_{2k*}(\widehat{\alpha}) = \alpha$.

- For $J_{3,k}^r$, there are the following cofibration and fibration sequences

$$S^{3k-1} \xrightarrow{\gamma_3^r} J_{2,k}^r \xrightarrow{I_2^r} J_{3,k}^r \xrightarrow{\check{p}_{3k}} S^{3k}; \quad \Omega S^{3k} \xrightarrow{\check{\delta}^{3k,r}} F_{\check{p}_{3k}} \xrightarrow{\check{\tau}_{3k}} J_{3,k}^r \xrightarrow{\check{p}_{3k}} S^{3k}.$$

We denote $I_2^r \beta_i$ by $\check{\beta}_i$ for $i = k, 2k, 3k - 1$.

Then from Lemma 3.5, we get

$$Sk_{4k-1}(F_{\check{p}_{3k}}) \simeq J_{2,k}^r \cup_{[\beta_k, \gamma_3^r]} e^{4k-1}. \quad (8)$$

Lemma 4.3. Let $m \leq 4k - 2$, there is a short exact sequence

$$0 \rightarrow \frac{\pi_m(J_{2,k}^r)}{\gamma_{3*}^r(\pi_m(S^{3k-1})) + L_m^k} \xrightarrow{I_{2*}^r} \pi_m(J_{3,k}^r) \xrightarrow{\check{p}_{3k*}} \Sigma(\text{Ker}(\gamma_{3*}^r)) \rightarrow 0$$

where the right γ_{3*} is the induced homomorphism on $\pi_{m-1}(-)$ by γ_3 and

$$L_m^k = \begin{cases} 0, & m \leq 4k-3; \\ \langle [\beta_k, \gamma_3^r] \rangle, & m = 4k-2. \end{cases}$$

Note that in the following text, for any $\alpha \in \Sigma_*(\text{Ker}(\gamma_{3*}^r))$, we always denote the $\widehat{\alpha} \in \pi_m(J_{3,k}^r)$ a lift of α by \check{p}_{3k} , i.e., $\check{p}_{3k*}(\widehat{\alpha}) = \alpha$.

Proof. From (8), we have the isomorphism

$$\pi_m(J_{2,k}^r)/L_m^k \xrightarrow{j\check{p}_{3k}^* \cong} \pi_m(F_{\check{p}_{3k}})(m \leq 4k - 2).$$

Then this lemma is obtained by the following exact sequence with commutative triangle and squares for $m \leq 4k - 2$

$$\begin{array}{ccccccc} \pi_{m+1}(S^{3k}) & \xrightarrow{\check{\delta}_{m^*}^{3k,r}} & \pi_m(F_{\check{p}_{3k}}) & \xrightarrow{\check{\tau}_{3k^*}} & \pi_m(J_{3,k}^r) & \xrightarrow{\check{p}_{3k^*}} & \pi_m(S^{3k}) & \xrightarrow{\check{\delta}_{m-1^*}^{2k,r}} & \pi_{m-1}(F_{\check{p}_{3k}}) \\ \uparrow \cong \Sigma & & \uparrow j\check{p}_{3k^*} & \nearrow I_{2^*} & & & \uparrow \cong \Sigma & & \uparrow \cong j\check{p}_{3k^*} \\ \pi_m(S^{3k-1}) & \xrightarrow{\gamma_{3^*}^r} & \pi_m(J_2(2^r \iota_k)) & & & & \pi_{m-1}(S^{3k-1}) & \xrightarrow{\gamma_{3^*}^r} & \pi_{m-1}(J_{2,k}^r) \end{array}$$

where the suspension homomorphisms Σ in squares are isomorphisms by the Freudenthal Suspension Theorem; $j\check{p}_{3k^*}$ are isomorphisms except the left one for $m = 4k - 2$ which is just an epimorphism. \square

If we need to emphasize the dependency of the above spaces or maps on the variable r , we will replace the $F_{k+1}, i_k, p_{k+1}, \tau_{k+1}, \beta_i (i = k, 2k, 3k - 1), \bar{p}_{2k}, \bar{\tau}_{2k}, \bar{\beta}_i (i = k, 3k - 1), \check{p}_{3k}, \check{\beta}_i (i = k, 2k, 3k - 1)$ by $F_{k+1}^r, i_k^r, p_{k+1}^r, \tau_{k+1}^r, \beta_i^r, \bar{p}_{2k}^r, \bar{\tau}_{2k}^r, \bar{\beta}_i^r, \check{p}_{3k}^r, \check{\beta}_i^r$ respectively.

We have the following homotopy commutative diagrams of cofibrations (left) and fibrations (right)

$$\begin{array}{ccccccc} S^k & \xrightarrow{2^s \iota_k} & S^k & \longrightarrow & P^{k+1}(2^s) & \xrightarrow{P_{k+1}^s} & S^{k+1} & , & \Omega S^{k+1} & \xrightarrow{\partial^{k+1,s}} & F_{k+1}^s & \xrightarrow{\tau_{k+1}^s} & P^{k+1}(2^s) \\ \downarrow 2^{M_{s-t}^0 \iota_k} & & \downarrow 2^{M_{t-s}^0 \iota_k} & & \downarrow \bar{\chi}_t^s & & \downarrow 2^{M_{s-t}^0 \iota_{k+1}} & & \downarrow \Omega(2^{M_{s-t}^0 \iota_{k+1}}) & & \downarrow \chi_t^s & & \downarrow \bar{\chi}_t^s \\ S^k & \xrightarrow{2^t \iota_k} & S^k & \longrightarrow & P^{k+1}(2^t) & \xrightarrow{P_{k+1}^t} & S^{k+1} & & \Omega S^{k+1} & \xrightarrow{\partial^{k+1,t}} & F_{k+1}^t & \xrightarrow{\tau_{k+1}^t} & P^{k+1}(2^t) \end{array} \quad (9)$$

where $M_a^b := \text{Max}\{r, s\}$ be the maximal number of integers a and b .

Let $\bar{g}_t^s := (\chi_t^s)|_{J_2}$, there is a map $g_t^s : F_{\bar{p}_{2k}^s} \rightarrow F_{\bar{p}_{2k}^t}$ such that the following diagrams homotopy commutes

$$\begin{array}{ccccccccccc} J_{2,k}^{\gamma,s} & \xrightarrow{I_2^{\gamma,s}} & F_{\bar{p}_{2k}^s} & \xrightarrow{\bar{\tau}_{2k}^s} & J_{2,k}^s & \xrightarrow{I_2^s} & J_{3,k}^s & \xrightarrow{\bar{I}_3^s} & F_{k+1}^s & \xrightarrow{\tau_{k+1}^s} & P^{k+1}(2^s) \\ \downarrow (g_t^s)|_{J_2} & & \downarrow g_t^s & & \downarrow \bar{g}_t^s & & \downarrow (\chi_t^s)|_{J_3} & & \downarrow \chi_t^s & & \downarrow \bar{\chi}_t^s \\ J_{2,k}^{\gamma,t} & \xrightarrow{I_2^{\gamma,t}} & F_{\bar{p}_{2k}^t} & \xrightarrow{\bar{\tau}_{2k}^t} & J_{2,k}^t & \xrightarrow{I_2^t} & J_{3,k}^t & \xrightarrow{\bar{I}_3^t} & F_{k+1}^t & \xrightarrow{\tau_{k+1}^t} & P^{k+1}(2^t) \end{array} \quad (10)$$

From Lemma 2.5 and 2.6

$$\begin{array}{ccc}
S^{2k-1} & \xrightarrow{\gamma_2^s} & S^k \xrightarrow{\beta_k^s} J_{2,k}^s \xrightarrow{\bar{p}_{2k}^s} S^{2k}, & S^{3k-1} & \xrightarrow{\gamma_3^s} & J_{2,k}^s \xrightarrow{I_2^s} J_{3,k}^r \xrightarrow{\check{p}_{3k}^s} S^{3k} \\
\downarrow 2^{[s-t]}\iota_{2k-1} & & \downarrow 2^{M_{t-s}^0}\iota_k \bar{g}_t^s & \downarrow 2^{[s-t]}\iota_{2k} & & \downarrow 2^{M_{2(s-t)}^s}\iota_{3k-1} \bar{g}_t^s & \downarrow (\chi_t^s)|_{J_3} \\
S^{2k-1} & \xrightarrow{\gamma_2^t} & S^k \xrightarrow{\beta_k^t} J_{2,k}^t \xrightarrow{\bar{p}_{2k}^t} S^{2k}, & S^{3k-1} & \xrightarrow{\gamma_3^t} & J_{2,k}^t \xrightarrow{I_2^t} J_{3,k}^t \xrightarrow{\check{p}_{3k}^t} S^{3k}
\end{array} \quad (11)$$

$$\text{So we have } \quad \bar{g}_t^s \beta_k^s \simeq 2^{M_{t-s}^0} \beta_k^t; \quad 2^{M_{2(s-t)}^s} \gamma_3^t \simeq \bar{g}_t^s \gamma_3^s. \quad (12)$$

$$\begin{array}{ccc}
\Omega S^{2k} & \xrightarrow{\bar{\partial}^{2k,s}} & F_{\bar{p}_{2k}^s} \xrightarrow{\bar{\tau}_{2k}^s} & J_{2,k}^s; & S^k & \xrightarrow{\bar{\beta}_k^s} & J_{2,k}^{\gamma,s} = S^k \vee S^{3k-1} \xrightarrow{q_2^{3k-1}} & S^{3k-1} \\
\downarrow \Omega(2^s \iota_{2k}) & & \downarrow g_t^s & \downarrow \bar{g}_t^s & \downarrow 2^{M_{t-s}^0} \iota_k & & \downarrow (g_t^s)|_{J_2} & \downarrow 2^{M_{2(s-t)}^s} \iota_{3k-1} \\
\Omega S^{2k} & \xrightarrow{\bar{\partial}^{2k,t}} & F_{\bar{p}_{2k}^t} \xrightarrow{\bar{\tau}_{2k}^t} & J_{2,k}^t & S^k & \xrightarrow{\bar{\beta}_k^t} & J_{2,k}^{\gamma,t} = S^k \vee S^{3k-1} \xrightarrow{q_2^{3k-1}} & S^{3k-1}
\end{array} \quad (13)$$

$$\bar{g}_t^s \circ \bar{\tau}_{2k}^s = \bar{\tau}_{2k}^t g_t^s; \quad (14)$$

$$(g_t^s)|_{J_2} = 2^{M_{t-s}^0} j_1^k q_1^k + 2^{M_{2(s-t)}^s} j_2^{3k-1} q_2^{3k-1} + j_1^k \theta q_2^{3k-1}, \quad \theta \in \pi_{3k-1}(S^k). \quad (15)$$

Lemma 4.4. For k is odd, $r \geq 1$, then

(i) $J_{2,k}^r \simeq S^k \vee S^{2k}$;

(ii) $\gamma_3^r = a_r [j_1^k, j_2^{2k}] + a_r j_1^k [\iota_k, \mu_r] \in \pi_{3k-1}(S^k \vee S^{2k})$ for some $a_r \in \mathbb{Z}$, $\mu_r \in \pi_{2k}(S^k)$;

(iii) Moreover, if $k = 3, 7$, then $\gamma_3^r = \pm 2^r [j_1^k, j_2^{2k}]$.

Proof. For k is odd, Theorem 8.8 of [41] implies $2[\iota_k, \iota_k] = 0$. $J_{2,k}^r \simeq S^k \vee S^{2k}$ ($r \geq 1$).

For(ii), there is an isomorphism

$$\pi_{3k-1}(S^{3k-1}) \oplus \pi_{3k-1}(S^k) \oplus \pi_{3k-1}(S^{2k}) \cong \pi_{3k-1}(S^k \vee S^{2k})$$

By the well known definition of the above isomorphism and from Lemma 2.11, we assume

$$\gamma_3^r = a_r [j_1^k, j_2^{2k}] + j_1^k \lambda_r, \quad a_r \in \mathbb{Z}, \lambda_r \in \pi_{3k-1}(S^k).$$

Moreover from Corollary 2.9, there is an element $\mu_r \in \pi_{2k}(S^k)$ such that

$$\begin{aligned}
\lambda_r &= q_1^k \gamma_3^r = [q_1^k, \mu_r q_2^{2k}] H_2(\gamma_3^r) \\
&= [\iota_k, \mu_r] (\Sigma q_1^{k-1} \wedge q_2^{2k-1}) (H_2(a_r [j_1^k, j_2^{2k}]) + H_2(j_1^k \lambda_r)) \\
&= a_r [\iota_k, \mu_r] (\Sigma q_1^{k-1} \wedge q_2^{2k-1}) (\Sigma j_1^{k-1} \wedge j_2^{2k-1} - \Sigma j_2^{2k-1} \wedge j_1^{k-1}) \quad (\text{by Lemma 2.13}) \\
&+ [\iota_k, \mu_r] (\Sigma q_1^{k-1} \wedge q_2^{2k-1}) (\Sigma j_1^{k-1} \wedge j_1^{k-1}) H_2(\lambda_r) \\
&= a_r [\iota_k, \mu_r].
\end{aligned}$$

This completes the proof of (ii) of this lemma.

For (iii), it is well known that $[\iota_k, \iota_k] = 0$ (odd $k \geq 2$) if and only if $k = 3, 7$. Thus $J_{2,k}^0 \simeq S^k \vee S^{2k}$ for $k = 3, 7$. From Proposition 5.9 and Theorem 7.7 of [39], $\Sigma : \pi_{3k-1}(S^k) \rightarrow \pi_{3k}(S^{k+1})$ are injections, for any $r \geq 0$, $S^{k_{4k-1}}(F_{k+1}) \simeq J_{3,k}^r \simeq (S^k \vee S^{2k}) \cup_{\gamma_3^r} CS^{3k-1}$ with $\gamma_3^r = a_r[j_1^k, j_2^{2k}] \in \pi_{3k-1}(S^k \vee S^{2k})$ for some $a_r \in \mathbb{Z}$ by Remark 2.12. So by the exact sequence (6) for $r = 0$, we get

$$\pi_{3k}(S^{k+1}) \cong \pi_{3k-1}(F_{k+1}^0) \cong \frac{\pi_{3k-1}(S^k) \oplus \pi_{3k-1}(S^{2k}) \oplus \mathbb{Z}\{[j_1^k, j_2^{2k}]\}}{\langle (0, 0, a_0[j_1^k, j_2^{2k}]) \rangle}. \quad (16)$$

By the Hopf bundles $S^k \rightarrow S^{2k+1} \rightarrow S^{k+1}$ ($k = 3, 7$), there is an isomorphism

$$\pi_{3k}(S^{k+1}) \cong \pi_{3k-1}(S^k) \oplus \pi_{3k}(S^{2k+1}) \cong \pi_{3k-1}(S^k) \oplus \pi_{3k-1}(S^{2k}). \quad (17)$$

Comparing (16) and (17), we get $a_0 = \pm 1$, i.e., $\gamma_3^0 = \pm[j_1^k, j_2^{2k}]$

We assume $\bar{g}_0^r = j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k} + j_1^k \theta q_2^{2k} \in [S^k \vee S^{2k}, S^k \vee S^{2k}]$ for some $\theta \in \pi_{2k}(S^k)$. By noting that $q_1^k \gamma_3^r \simeq 0$, $q_2^{2k} \gamma_3^r \simeq 0$, from (12),

$$\begin{aligned} 2^{2r} \gamma_3^0 &= \bar{g}_0^r \circ \gamma_3^r = (j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k} + j_1^k \lambda q_2^{2k}) \gamma_3^r \\ &= (j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k}) \gamma_3^r + j_1^k \theta q_2^{2k} \gamma_3^r \pm [j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k}, j_1^k \theta q_2^{2k}] H_2(\gamma_3^r) \\ (j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k}) \gamma_3^r &= \pm a_r [j_1^k q_1^k, 2^r j_2^{2k} q_2^{2k}] H_2([j_1^k, j_2^{2k}]) \\ &= \pm 2^r a_r [j_1^k, j_2^{2k}] (\Sigma q_1^{k-1} \wedge q_2^{2k-1}) (\Sigma j_1^{k-1} \wedge j_2^{2k-1} - \Sigma j_2^{2k-1} \wedge j_1^{k-1}) \\ &= \pm 2^r a_r [j_1^k, j_2^{2k}]. \\ \Rightarrow (2^{2r} \pm 2^r a_r) [j_1^k, j_2^{2k}] &= \pm [j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k}, j_1^k \theta q_2^{2k}] H_2(\gamma_3^r). \end{aligned} \quad (18)$$

Since θ is a torsion element, so is the right term of (18). However the left term of (18) is torsion free or zero. We get

$$(2^{2r} \pm 2^r a_r) [j_1^k, j_2^{2k}] = \pm [j_1^k q_1^k + 2^r j_2^{2k} q_2^{2k}, j_1^k \theta q_2^{2k}] H_2(\gamma_3^r) = 0$$

That is $a_r = \pm 2^r$, we get $\gamma_3^r = \pm 2^r [j_1^k, j_2^{2k}]$. \square

At the end of this section, we generalize the map $\bar{H}_2 : \Omega P^{n+1}(2) \rightarrow \Omega P^{2n+1}(2)$ in [43, section 4.4] to the mod 2^r Moore space as follows

$$\bar{H}_2 : \Omega P^{n+1}(2^r) \simeq J(P^n(2^r)) \xrightarrow{H_2} J(P^n(2^r) \wedge P^n(2^r)) \xrightarrow{J(q)} J(P^{2n}(2^r)) \simeq \Omega P^{2n+1}(2^r).$$

Thus we have the following homotopy commutative diagram of fibration sequence

$$\begin{array}{ccccccc} \Omega^2 S^{n+1} & \xrightarrow{\Omega \partial^{n+1,r}} & \Omega F_{n+1} & \xrightarrow{\Omega \tau_{n+1}} & \Omega P^{n+1}(2^r) & \xrightarrow{\Omega p_{n+1}} & \Omega S^{n+1} \\ \downarrow \Omega H_2 & & \downarrow \bar{H}_2 & & \downarrow \bar{H}_2 & & \downarrow H_2 \\ \Omega^2 S^{n+1} & \xrightarrow{\Omega \partial^{2n+1,r}} & \Omega F_{2n+1} & \xrightarrow{\Omega \tau_{2n+1}} & \Omega P^{2n+1}(2^r) & \xrightarrow{\Omega p_{2n+1}} & \Omega S^{2n+1} \end{array} \quad (19)$$

where the right vertical map H_2 is the second James Hopf invariant defined in (3); the homotopy commutativity of the right square is obtained by the naturality of James construction and the left vertical map \tilde{H}_2 is induced by the right homotopy commutative square.

5. Homotopy groups of 4, 5 and 6 dimensional mod 2^r Moore spaces

In this section, we compute homotopy groups of mod 2^r Moore spaces in Theorem 1.2. All the spaces and homotopy groups in the following are under 2-localization.

Firstly, we give some notations. For a homomorphism $f : G \rightarrow H$ of abelian groups, where G_1 is subgroups of G , then we denote $f|_{res} : G_1 \rightarrow H$ as the restriction of f on G_1 . For a monomorphism $f : G \hookrightarrow H$ (resp. epimorphism $g : H \twoheadrightarrow K$), if there is a map $f' : H \rightarrow G$ (resp. $g' : K \rightarrow H$) such that $f'f = id_G$ (resp. $gg' = id_K$), then f (resp. g) is called split into (resp. split onto). In this case, $f(G)$ (resp. $g'(K)$) is a direct summand of H and we say G (resp. K) splits out of H . $o(\alpha)$ denotes the order of α , where α is an element of an abelian group and the sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ always means a short exact sequence. All the following ladders, squares and triangles of abelian groups (CW-complexes) are commutative (homotopy commutative). In the following, i and p denote the $\Sigma^\infty i_k \in \pi_k^s(P^{k+1}(2^r))$ and $\Sigma^\infty p_{k+1} \in \Sigma^\infty [P^{k+1}(2^r), S^{k+1}]$.

5.1. Homotopy groups of $P^4(2^r)$

$$S^3 \xrightarrow{2^r i_3} S^3 \xrightarrow{i_3} P^4(2^r) \xrightarrow{p_4} S^4; \quad \Omega S^4 \xrightarrow{\partial^{4,r}} F_4 \xrightarrow{\tau_4} P^4(2^r) \xrightarrow{p_4} S^4$$

By (iii) of Lemma 4.4

$Sk_8(F_4) \simeq J_{2,3}^r \simeq S^3 \cup_{\gamma_2^r} CS^5 \simeq S^3 \vee S^6$; $Sk_{11}(F_5) \simeq J_{3,3}^r \simeq J_{2,3}^r \cup_{\gamma_3^r} CS^8$ where $\gamma_3^r = \pm 2^r [j_1^3, j_2^6]$.

Lemma 5.1. *For $r \geq 1$, $\pi_i(F_4) \cong \pi_i(J_{3,3}^r)$ ($i \leq 11$) and*

- (1) $\pi_8(J_{3,3}^r) = \mathbb{Z}_2\{\check{\beta}_3 \nu' \eta_6^2\} \oplus \mathbb{Z}_2\{\check{\beta}_6 \eta_6^2\} \oplus \mathbb{Z}_{2^r}\{[\check{\beta}_3, \check{\beta}_6]\}$.
- (2) $\pi_9(J_{3,3}^r) = \mathbb{Z}_8\{\check{\beta}_6 \nu_6\} \oplus \mathbb{Z}_2\{[\check{\beta}_3, \check{\beta}_6] \eta_8\}$.
- (3) $\pi_{10}(J_{3,3}^1) = \mathbb{Z}_2\{[\check{\beta}_3, [\check{\beta}_3, \check{\beta}_6]]\} \oplus \mathbb{Z}_4\{\widehat{\eta}_9\}$ with $[\check{\beta}_3, \check{\beta}_6] \eta_8 = 2\widehat{\eta}_9$;
 $\pi_{10}(J_{3,3}^r) = \mathbb{Z}_2\{[\check{\beta}_3, \check{\beta}_6] \eta_8^2\} \oplus \mathbb{Z}_{2^r}\{[\check{\beta}_3, [\check{\beta}_3, \check{\beta}_6]]\} \oplus \mathbb{Z}_2\{\widehat{\eta}_9\}$ ($r \geq 2$).

Proof. From Lemma 4.3 for $k = 3, m = 8, 9$, we get the following isomorphism

$$\pi_m(S^3 \vee S^6) / \gamma_{3*}^r(\pi_m(S^{3k-1})) \xrightarrow[\cong]{l_{2*}^r} \pi_m(J_{3,3}^r) \quad (m = 8, 9)$$

which implies (1) and (2) of Lemma 5.1.

For (3) of Lemma 5.1, there is an exact sequence

$$0 \rightarrow \text{Coker}(\check{\partial}_{10*}^{9,r}) \xrightarrow{I_{2*}^r} \pi_{10}(J_{3,3}^r) \xrightarrow{\check{p}_{9*}} \pi_{10}(S^9) = \mathbb{Z}_2\{\eta_9\} \rightarrow 0,$$

$$\text{Coker}(\check{\partial}_{10*}^{9,r}) = \pi_{10}(S^3 \vee S^6) / (\gamma_3^r \circ \pi_{10}(S^8) + \langle [j_1^3, \gamma_3^r] \rangle) = \mathbb{Z}_2\{[j_1^3, j_2^6]\eta_8^2\} \oplus \mathbb{Z}_{2^r}\{[j_1^3, [j_1^3, j_2^6]]\}.$$

For $r \geq 1$, by Lemma 3.8, there is an $-\mathbf{x} \in \{I_2^r, \gamma_3^r, \eta_8\} \subset \pi_{10}(J_{3,3}^r)$, such that $\check{p}_{9*}(\mathbf{x}) = \eta_9$. By Proposition 1.2 (IV) and Proposition 1.4 of [39],

$$2\mathbf{x} \in -\{I_2^r, \gamma_3^r, \eta_8\}(2\iota_{10}) = I_2^r\{\gamma_3^r, \eta_8, 2\iota_9\}.$$

$$\Rightarrow \{\gamma_3^r, \eta_8, 2\iota_9\} \ni \pm 2^{r-1}[j_1^3, j_2^6]\eta_8^2 \bmod \langle 2[j_1^3, [j_1^3, j_2^6]] \rangle \quad (\text{Lemma 2.4}).$$

Thus for $r = 1$, $2\mathbf{x} = I_2^1[j_1^3, j_2^6]\eta_8 = [\check{\beta}_3, \check{\beta}_6]\eta_8$, which implies that $o(\mathbf{x}) = 4$ and $\pi_{10}(J_{3,3}^1) = \mathbb{Z}_2\{[\check{\beta}_3, [\check{\beta}_3, \check{\beta}_6]]\} \oplus \mathbb{Z}_4\{\widehat{\eta}_9\}$ with $[\check{\beta}_3, \check{\beta}_6]\eta_8 = 2\widehat{\eta}_9$.

For $r \geq 2$, $2\mathbf{x} = I_2^r(2t[j_1^3, [j_1^3, j_2^6]]), t \in \mathbb{Z}$. Take $\mathbf{x}' = \mathbf{x} - I_2^r(t[j_1^3, [j_1^3, j_2^6]])$. We get $2\mathbf{x}' = 0$ and $\check{p}_{9*}(\mathbf{x}') = \eta_9$. So $\pi_{10}(J_{3,3}^r) \xrightarrow{\check{p}_{9*}} \pi_{10}(S^9)$ is split onto. which implies the (3) of Lemma 5.1 for $r \geq 2$. \square

- $\pi_8(P^4(2^r))$.

Consider diagram (6) for $k = 3, m = 8$, we get

$$\pi_9(S^4) \xrightarrow{\partial_{8*}^{4,r}} \pi_8(F_4) \xrightarrow{\tau_{4*}} \pi_8(P^4(2^r)) \xrightarrow{p_{4*}} \pi_8(S^4) \xrightarrow{\partial_{7*}^{4,r}} \pi_7(F_5)$$

we have $\partial_{7*}^{4,r}(\nu_4\eta_7) = (1 - \vartheta_r)\beta_3\nu'\eta_6$; $\partial_{7*}^{4,r}(\Sigma\nu'\eta_7) = 0$ ([49, (3.12),(3.13)]).

$$\Rightarrow \text{Ker}(\partial_{7*}^{4,r}) = \mathbb{Z}_2\{\Sigma\nu'\eta_7\} \oplus \vartheta_r\mathbb{Z}_2\{\nu_4\eta_7\}.$$

$$\partial_{8*}^{4,r}(\Sigma\nu'\eta_7^2) = \partial_{7*}^{4,r}(\Sigma\nu'\eta_7)\eta_7 = 0; \partial_{8*}^{4,r}(\nu_4\eta_7^2) = \partial_{7*}^{4,r}(\nu_4\eta_7)\eta_7 = (1 - \vartheta_r)\beta_3\nu'\eta_6^2; \quad (20)$$

$$\Rightarrow \text{Coker}(\partial_{8*}^{4,r}) = \vartheta_r\mathbb{Z}_2\{\beta_3\nu'\eta_6^2\} \oplus \mathbb{Z}_2\{\beta_6\eta_6^2\} \oplus \mathbb{Z}_{2^r}\{[\beta_3, \beta_6]\}. \quad (21)$$

Thus we have the following short exact sequence

$$0 \rightarrow \vartheta_r\mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_{2^r} \rightarrow \pi_8(P^4(2^r)) \rightarrow \vartheta_r\mathbb{Z}_2\{\nu_4\eta_7\} \oplus \mathbb{Z}_2\{\Sigma\nu'\eta_7\} \rightarrow 0 \quad (22)$$

By Lemma 3.8, there is an $-\mathbf{x} \in \{i_3, 2^r\iota_3, \nu'\eta_6\} \subset \pi_8(P^4(2^r))$, such that $p_{4*}(\mathbf{x}) = \Sigma\nu'\eta_7$. By Lemma 8.2 (i) and (20)

$$2\mathbf{x} \in -\{i_3, 2^r\iota_3, \nu'\eta_6\}(2\iota_8) = i_3\{2^r\iota_3, \nu'\eta_6, 2\iota_7\} = 2^{r-1}i_3\nu'\eta_6^2 = 0 \Rightarrow o(\mathbf{x}) = 2(r \geq 1).$$

Hence (22) splits for $r = 1$ and $\pi_8(P^4(2)) \cong \mathbb{Z}_2^{\oplus 3}$.

For $r \geq 2$, firstly we prove the following lemma

Lemma 5.2. For $r \geq 2$, there are $\tilde{\eta}_7 \in \pi_7(P^7(2^{r-1}))$ and $\tilde{v}_4 \in [P^7(2^{r-1}), P^4(2^r)]$ satisfying the following homotopy commutative diagram

$$\begin{array}{ccccc} S^8 & \xrightarrow{\tilde{\eta}_7} & P^7(2^{r-1}) & \xrightarrow{\tilde{v}_4} & P^4(2^r) \\ & \searrow \eta_7 & \downarrow p_7^{r-1} & & \downarrow p_4^r \\ & & S^7 & \xrightarrow{\nu_4} & S^4 \end{array} .$$

Moreover, the order of $\tilde{v}_4\tilde{\eta}_7$ is 4 and 2 respectively for $r = 2$ and $r \geq 3$.

Proof. $\tilde{\eta}_7$ comes from $\pi_8(P^7(2^{r-1})) = \begin{cases} \mathbb{Z}_4\{\tilde{\eta}_7\}, & r = 2; \\ \mathbb{Z}_2\{i_6\eta_6^2\} \oplus \mathbb{Z}_2\{\tilde{\eta}_7\}, & r \geq 3 \end{cases}$ by [4].

We have the following exact sequence

$$[P^7(2^{r-1}), P^4(2^r)] \xrightarrow{p_{4*}^r} [P^7(2^{r-1}), S^4] \xrightarrow{\partial_*^{4,r}} [P^6(2^{r-1}), F_4^r].$$

By noting $o(p_6^{r-1}) = 2^{r-1}$, from Lemma 3.2 and (3.5), (3.8) of [49],

$$\partial_*^{4,r}(\nu_4 p_7^{r-1}) = \partial_{6*}^{4,r}(\nu_4) p_6^{r-1} = (\pm 2^{r-1} \beta_3 \nu' + 2^r \beta_6) p_6^{r-1} = 0.$$

Thus \tilde{v}_4 exists. $\tilde{v}_4\tilde{\eta}_7$ is a lift of $\nu_4\eta_7$ by p_4^r implies $o(\tilde{v}_4\tilde{\eta}_7) \geq 2$. So $o(\tilde{v}_4\tilde{\eta}_7) = 2$ for $r \geq 3$, since $o(\tilde{\eta}_7) = 2$ for $r \geq 3$.

For $r = 2$, $o(\tilde{\eta}_7) = 4$ and $2\tilde{\eta}_7 = i_6\eta_6^2$. Hence, $2\tilde{v}_4\tilde{\eta}_7 = \tilde{v}_4 i_6\eta_6^2$.

Since $(4\iota_4) \circ \nu_4 = 16\nu_4 + 2\Sigma\nu' = (8\nu_4 + \Sigma\nu')(2\iota_7)$ by Lemma A.1 of [49], the following diagram is homotopy commutative

$$\begin{array}{ccccccc} P^7(2) & \xrightarrow{p_7^1} & S^7 & \xrightarrow{2\iota_7} & S^7 & \xrightarrow{i_7} & P^8(2) \\ \downarrow \tilde{v}_4 & & \downarrow \nu_4 & & \downarrow 8\nu_4 + \Sigma\nu' & & \downarrow \Sigma\tilde{v}_4 \\ P^4(4) & \xrightarrow{p_4^2} & S^4 & \xrightarrow{4\iota_4} & S^4 & \xrightarrow{i_4} & P^5(4) \end{array}$$

implies $\Sigma\tilde{v}_4 i_7 = i_4(8\nu_4 + \Sigma\nu')$.

$$\Sigma(\tilde{v}_4 i_6 \eta_6^2) = \Sigma\tilde{v}_4 i_7 \eta_7^2 = i_4(8\nu_4 + \Sigma\nu') \eta_7^2 = i_4 \Sigma\nu' \eta_7^2 \neq 0 \text{ by (29).}$$

Hence $\Sigma(\tilde{v}_4 i_6 \eta_6^2) = \Sigma(2\tilde{v}_4\tilde{\eta}_7) \neq 0 \Rightarrow 2\tilde{v}_4\tilde{\eta}_7 \neq 0 \Rightarrow o(\tilde{v}_4\tilde{\eta}_7) = 4$. \square

From above lemma and (22), we get

$$\begin{aligned} \pi_8(P^4(2^r)) &\cong \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_{2^r} \quad (r \geq 3); \\ \pi_8(P^4(4)) &\cong \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_4 \text{ or } \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_8 \text{ or } \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4^{\oplus 2}. \end{aligned}$$

From Lemma 3.7, $\mathbb{Z}_4 \cong \pi_8(P^9(4))$ is a summand of $\pi_8(P^4(4))$, implies that $\pi_8(P^9(4)) \not\cong \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_8$. Assume that there is an order 2 lift $\mathbf{y} \in \pi_8(P^4(4))$ of $\nu_4\eta_7$, then $\tilde{v}_4\tilde{\eta}_7 - \mathbf{y} \in \text{Ker}(p_{4*}^2) \Rightarrow \tilde{v}_4\tilde{\eta}_7 - \mathbf{y} = \tau_4^2[\beta_3, \beta_6] + \alpha$, with $o(\alpha) \leq 2$ (21). Then $2\tilde{v}_4\tilde{\eta}_7 = 2(\tilde{v}_4\tilde{\eta}_7 - \mathbf{y}) =$

$2\tau_4^2[\beta_3, \beta_6] \Rightarrow \Sigma(2\tilde{\nu}_4\tilde{\eta}_7) = 0$. This is a contradiction since we previously obtained $\Sigma(2\tilde{\nu}_4\tilde{\eta}_7) \neq 0$. Thus the short exact sequence (22) does not split for $r = 2$. This implies $\pi_8(P^4(4)) \not\cong \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_4$. So we get $\pi_8(P^4(4)) \cong \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4^{\oplus 2}$. Thus

$$\pi_8(P^4(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 3}, & r = 1; \\ \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4^{\oplus 2}, & r = 2; \\ \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_{2^r}, & r \geq 3. \end{cases}$$

- $\pi_9(P^4(2^r))$.

$$\mathbb{Z}_8\{v_4^2\} = \pi_{10}(S^4) \xrightarrow{\partial_{9*}^{4,r}} \pi_9(F_4) \xrightarrow{\tau_{4*}} \pi_9(P^4(2^r)) \xrightarrow{p_{4*}} \pi_9(S^4) \xrightarrow{\partial_{8*}^{4,r}} \pi_8(F_5)$$

$$\text{Ker}(\partial_{8*}^{4,r}) = \vartheta_r \mathbb{Z}_2\{v_4\eta_7^2\} \oplus \mathbb{Z}_2\{\Sigma v'\eta_7^2\}, \text{ by (20).}$$

$$\partial_{9*}^{4,r}(v_4^2) = \partial_{6*}^{4,r}(v_4)v_6 = (\pm 2^{r-1}\beta_3v' + 2^r\beta_6)v_6 = 2^r\beta_6v_6.$$

$$\Rightarrow \text{Coker}(\partial_{9*}^{4,r}) = \mathbb{Z}_{2m_r^3}\{\beta_6v_6\} \oplus \mathbb{Z}_2\{[\beta_3, \beta_6]\eta_8\}.$$

$$\Rightarrow 0 \rightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_{2m_r^3} \rightarrow \pi_9(P^4(2^r)) \xrightarrow{p_{4*}} \vartheta_r \mathbb{Z}_2\{v_4\eta_7^2\} \oplus \mathbb{Z}_2\{\Sigma v'\eta_7^2\} \rightarrow 0$$

Take $\mathbf{x}\eta_8 \in \pi_9(P^4(2^r))$ and $\tilde{\nu}_4\tilde{\eta}_7\eta_8 \in \pi_9(P^4(2^r))$ ($r \geq 2$), where \mathbf{x} and $\tilde{\nu}_4\tilde{\eta}_7 \in \pi_8(P^4(2^r))$ are lifts of $\Sigma v'\eta_7$ and $v_4\eta_7$ respectively. Thus $o(\mathbf{x}\eta_8) = o(\tilde{\nu}_4\tilde{\eta}_7\eta_8) = 2$ and $p_{4*}(\mathbf{x}\eta_8) = \Sigma v'\eta_7^2$, $p_{4*}(\tilde{\nu}_4\tilde{\eta}_7\eta_8) = v_4\eta_7^2$ ($r \geq 2$). Hence $\pi_9(P^4(2^r)) \xrightarrow{p_{4*}} \vartheta_r \mathbb{Z}_2\{v_4\eta_7^2\} \oplus \mathbb{Z}_2\{\Sigma v'\eta_7^2\}$ is split onto, which implies

$$\pi_9(P^4(2^r)) \cong \vartheta_r \mathbb{Z}_2 \oplus \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_{2m_r^3}, r \geq 1.$$

- $\pi_{10}(P^4(2^r))$.

By $\pi_{11}(S^4) = 0$ and $\partial_{9*}^{4,r}(v_4^2) = 2^r\beta_6v_6$ implies the short exact sequence

$$0 \longrightarrow \pi_{10}(F_4) \xrightarrow{\tau_{4*}} \pi_{10}(P^4(2^r)) \xrightarrow{p_{4*}} \mathbb{Z}_{2m_r^3}\{\delta_r v_4^2\} \longrightarrow 0$$

$$\pi_{10}(F_4^1) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_4; \quad \pi_{10}(F_4^r) \cong \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_{2^r} (r \geq 2)$$

where $\delta_r = 4, 2$ for $r = 1, 2$ and $\delta_r = 1$ for $r \geq 3$.

Lemma 5.3. *For any integer s , such that $0 \leq s < r$, $2^s \tau_4 \check{I}_3^r[\check{\beta}_3, [\check{\beta}_3, \check{\beta}_6]]$ is not divisible by 2^{s+1} in $\pi_{10}(P^4(2^r))$.*

Proof. Let $R = \mathbb{Z}_{2^r}$. Note that $H_*(\Omega P^4(2^r), R) \cong T(u, v)$, which is the tensor algebra over R generated by u, v with degrees 2 and 3 respectively.

$Sk_8 F_4 \simeq S^3 \vee S^6$ implies that $Sk_6(\Omega F_4) \simeq S^2 \vee S^4 \vee S^5 \vee S^6$.

Consider the R -homology the Serre Spectral Sequence for the fibration sequence

$$\Omega F_4 \xrightarrow{\Omega \tau_4} \Omega P^4(2^r) \xrightarrow{\Omega p_4} \Omega S^4$$

$R \cong H_5(\Omega F_4, R) \cong E_{0,5}^\infty = F_5^0 \subset F_5^5 = H_5(\Omega P^4(2^r), R) \cong R^{\oplus 2}$.

where the above inclusion is just the induced homomorphism

$$(\Omega\tau_4)_* : H_5(\Omega F_4, R) \rightarrow H_5(\Omega P^4(2^r), R)$$

by the first commutative diagram (the right commutative square) in page 22 of [16].

By Proposition 21.1 of [8], $[u, v] := uv - vu \in H_5(\Omega P^4(2^r), R)$ is spherical, i.e., it is the mod 2^r -reduction of Hurewicz image h_w of some element in $\pi_5(\Omega P^4(2^r))$, where $\Omega_0 : \pi_6(P^4(2^r)) \xrightarrow{\cong} \pi_5(\Omega P^4(2^r))$. By the calculation of $\pi_6(P^4(2^r))$ in [49, Section 3.1], we get

$$\pi_6(P^4(2^r)) = \begin{cases} \mathbb{Z}_4\{\tau_4\beta_6\} \oplus \mathbb{Z}_2\{\tilde{\eta}_4\eta_5\}, & r = 1; \\ \mathbb{Z}_8\{\tau_4\beta_6\} \oplus \mathbb{Z}_2\{2\tau_4\beta_6 + i_3\nu'\} \oplus \mathbb{Z}_2\{\tilde{\eta}_4\eta_5\}, & r = 2; \\ \mathbb{Z}_{2^r}\{\tau_4\beta_6\} \oplus \mathbb{Z}_4\{i_3\nu'\} \oplus \mathbb{Z}_2\{\tilde{\eta}_4\eta_5\}, & r \geq 3. \end{cases}$$

where $\tilde{\eta}_4 \in \pi_5(P^4(2^r))$ is a lift of η_4 by p_{4*} .

It is easy to see $h_w(\Omega_0(i_3\nu')) = 0$ and $h_w(\Omega_0(\tilde{\eta}_4\eta_5)) = 0$.

Hence $\bar{h}_w(\Omega_0(\tau_4\beta_6)) = l[u, v] \in H_5(\Omega P^4(2^r), R)$ where l is odd and \bar{h}_w is the mod 2^r -reduction of h_w .

Now we compute $\bar{h}_w(\Omega_0([i_3, [i_3, \tau_4\beta_6]]))$ by $(\Omega_0 i_3)_*(\iota_2) = u$, $(\Omega_0(\tau_4\beta_6))_*(\iota_5) = l[u, v]$

$$\begin{aligned} (\Omega_0[i_3, \tau_4\beta_6])_*(\iota_7) &= (-1)^{a_1}([\Omega_0 i_3, \Omega_0(\tau_4\beta_6)]^S)_*(\iota_2 \wedge \iota_5) \\ &= (-1)^{a_1}((\Omega_0 i_3)_*(\iota_2) \otimes (\Omega_0(\tau_4\beta_6))_*(\iota_5) - (\Omega_0(\tau_4\beta_6))_*(\iota_5) \otimes (\Omega_0 i_3)_*(\iota_2)) \\ &= (-1)^{a_1}l(u \otimes [u, v] - [u, v] \otimes u) = (-1)^{a_1}l[u, [u, v]]. \\ \bar{h}_w(\Omega_0([i_3, [i_3, \tau_4\beta_6]])) &= (-1)^{a_2}([\Omega_0 i_3, \Omega_0[i_3, \tau_4\beta_6]]^S)_*(\iota_2 \wedge \iota_7) \\ &= (-1)^{a_2}(u \otimes (\Omega_0[i_3, \tau_4\beta_6])_*(\iota_7) - (\Omega_0[i_3, \tau_4\beta_6])_*(\iota_7) \otimes u) \\ &= (-1)^{a_3}l[u, [u, [u, v]]], \quad a_1, a_2 \in \mathbb{Z}, a_3 = a_1 + a_2. \end{aligned}$$

This implies that $2^s \Omega_0([i_3, [i_3, \tau_4\beta_6]])$ is not divisible by 2^{s+1} in $\pi_9(\Omega P^4(2^r))$ for $0 \leq s < r$, so is $\tau_4 \check{I}_3^r[\check{\beta}_3, [\check{\beta}_3, \check{\beta}_6]] = [i_3, [i_3, \tau_4\beta_6]]$ in $\pi_{10}(P^4(2^r))$. \square

By Lemma 3.7, $\pi_{10}(P^9(2^r)) \cong \begin{cases} \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2^{\oplus 2}, & r \geq 2 \end{cases}$ is a direct summand of $\pi_{10}(P^4(2^r))$ and from Lemma 5.3, $\mathbb{Z}_{2^r}\{[\check{\beta}_3, [\check{\beta}_3, \check{\beta}_6]]\}$ in Lemma 5.1 splits out of $\pi_{10}(P^4(2^r))$. Hence

$$\pi_{10}(P^4(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_{2^r} \oplus \mathbb{Z}_{2^r m^3}, & r \geq 2. \end{cases}$$

5.2. Homotopy groups of $P^5(2^r)$

There are cofibration and fibration sequences

$$\begin{aligned} S^4 \xrightarrow{2^r \iota_4} S^4 \xrightarrow{i_4} P^5(2^r) \xrightarrow{p_5} S^5; \quad \Omega S^5 \xrightarrow{\partial^5} F_5 \xrightarrow{\tau_5} P^5(2^r) \xrightarrow{p_5} S^5 \\ Sk_{11}(F_5) \simeq J_{2,4}^r \simeq S^4 \cup_{\gamma_2^r} CS^7; \quad Sk_{15}(F_5) \simeq J_{3,4}^r \simeq J_{2,4}^r \cup_{\gamma_3^r} CS^{11}. \end{aligned}$$

where $\gamma_2^r = 2^r[\iota_4, \iota_4] = 2^{r+1}\nu_4 - 2^r\Sigma\nu'$.

Since $\pi_{11}(S^4) = 0$, from diagrams of (13), we get

$$\bar{g}_0^r \bar{\tau}_8^r = \bar{\tau}_8^0 g_0^r; g_0^r|_{J_2} \simeq \iota_4 \vee 2^r \iota_{11} \text{ which implies } g_0^r \bar{\beta}_4^r = \bar{\beta}_4^0 \text{ and } g_0^r \bar{\beta}_{11}^r = 2^r \bar{\beta}_{11}^0. \quad (23)$$

Lemma 5.4. For $r \geq 0$,

$$\pi_9(F_5^r) \cong \pi_9(J_{2,4}^r) = \begin{cases} \mathbb{Z}_2\{\beta_4\nu_4\eta_7^2\} \oplus \vartheta_r^0 \mathbb{Z}_4\{\widehat{\eta}_8\}, & r = 0, 1; \\ \mathbb{Z}_2\{\beta_4\nu_4\eta_7^2\} \oplus \mathbb{Z}_2\{\beta_4\Sigma\nu'\eta_7^2\} \oplus \mathbb{Z}_2\{\widehat{\eta}_8\}, & r \geq 2, \end{cases}$$

with $\beta_4^1 \Sigma\nu'\eta_7^2 = 2\widehat{\eta}_8$ for $r = 1$.

$$\pi_{10}(F_5^r) \cong \pi_{10}(J_{2,4}^r) = \mathbb{Z}_{2^{m_{r+1}}} \{\beta_4\nu_4^2\} \oplus \vartheta_r^0 \mathbb{Z}_2\{\eta_8^2\}.$$

$$\pi_i(J_{2,4}^r) = \begin{cases} \mathbb{Z}_{(2)}\{\beta_{11}\} \oplus \mathbb{Z}_{2^{m_{r+1}}} \{\widehat{\delta_{r+1}\nu_8}\}, & i = 11; \\ \mathbb{Z}_2\{\beta_4\varepsilon_4\} \oplus \mathbb{Z}_2\{\beta_{11}\eta_{11}\}, & i = 12; \\ \mathbb{Z}_2\{\beta_4\nu_4^3\} \oplus \mathbb{Z}_2\{\beta_4\mu_4\} \oplus \mathbb{Z}_2\{\beta_4\eta_4\varepsilon_5\} \oplus \mathbb{Z}_2\{\beta_{11}\eta_{11}^2\}, & i = 13. \end{cases}$$

where $\vartheta_r^0 = 0$ for $r = 0$ and $\vartheta_r^0 = 1$ for $r \geq 1$; $\widehat{\delta_{r+1}\nu_8}$ is a lift of $\delta_{r+1}\nu_8 \in \pi_{11}(S^8)$ by \bar{p}_8 ; the $\varepsilon_4, \mu_4, \eta_4\varepsilon_5$ comes from the generators of $\pi_{12}(S^4) = \mathbb{Z}_2\{\varepsilon_4\}$ and $\pi_{13}(S^4) = \mathbb{Z}_2\{\nu_4^3\} \oplus \mathbb{Z}_2\{\mu_4\} \oplus \mathbb{Z}_2\{\beta_{11}\eta_{11}^2\}$.

Proof.

- $\pi_9(J_{2,4}^r)$. Lemma 4.1 for $k = 4, m = 9$ implies the following short exact sequence

$$0 \rightarrow \mathbb{Z}_2\{\bar{\beta}_4\nu_4\eta_7^2\} \oplus \vartheta_r^0 \mathbb{Z}_2\{\bar{\beta}_4\Sigma\nu'\eta_7^2\} \xrightarrow{\bar{\tau}_{8*}} \pi_9(J_{2,4}^r) \xrightarrow{\bar{p}_{8*}} \vartheta_r^0 \mathbb{Z}_2\{\eta_8\} \rightarrow 0.$$

For $r \geq 1$, by Lemma 3.8, $\exists -\mathbf{x} \in \{\beta_4, \gamma_2^r, \eta_7\} \subset \pi_9(J_{2,4}^r)$, such that $\bar{p}_{8*}(\mathbf{x}) = \eta_8$.

$$2\mathbf{x} \in -\{\beta_4, \gamma_2^r, \eta_7\}(2\iota_9) = \beta_4\{\gamma_2^r, \eta_7, 2\iota_8\}.$$

$$\{\gamma_2^r, \eta_7, 2\iota_8\} \ni (2^{r-1}[\iota_4, \iota_4])\eta_8 \bmod \pi_9(S^4)(2\iota_9) + (\gamma_2^r)\pi_9(S^7) = \{0\}. \text{ (Lemma 3.9)}$$

Thus $2\mathbf{x} = 0$ for $r \geq 2$, implies that

$$\pi_9(J_{2,4}^r) = \mathbb{Z}_2\{\beta_4\nu_4\eta_7^2\} \oplus \mathbb{Z}_2\{\beta_4\Sigma\nu'\eta_7^2\} \oplus \mathbb{Z}_2\{\widehat{\eta}_8\} \quad (r \geq 2).$$

$2\mathbf{x} = \beta_4^1 \Sigma\nu'\eta_7^2$ for $r = 1$, implies that $o(\mathbf{x}) = 4$. Hence

$$\pi_9(J_2(2\iota_4)) = \mathbb{Z}_2\{\beta_4^1\nu_4\eta_7^2\} \oplus \mathbb{Z}_4\{\widehat{\eta}_8\} \text{ with } \beta_4^1 \Sigma\nu'\eta_7^2 = 2\widehat{\eta}_8.$$

- $\pi_{10}(J_{2,4}^r)$. Lemma 4.1 for $k = 4, m = 10$ implies

$$0 \rightarrow \mathbb{Z}_{2^{m_{r+1}}} \{\bar{\beta}_4\nu_4^2\} \xrightarrow{\bar{\tau}_{8*}} \pi_{10}(J_{2,4}^r) \xrightarrow{\bar{p}_{8*}} \vartheta_r^0 \mathbb{Z}_2\{\eta_8^2\} \rightarrow 0.$$

For $r \geq 2$, by Lemma 3.8, there is $-\mathbf{x} \in \{\beta_4, \gamma_2^r, \eta_7\} \subset \pi_9(J_{2,4}^r)$, such that $\bar{p}_{8*}(\mathbf{x}) = \eta_8$.

Then $\bar{p}_{8*}(\mathbf{x}\eta_9) = \eta_8^2$. Moreover, $2(\mathbf{x}\eta_9) = \mathbf{x}(2\eta_9) = 0$ implies $o(\mathbf{x}\eta_9) = 2$. Hence

$\pi_{10}(J_{2,4}^r) \xrightarrow{\bar{p}_{8*}} \vartheta_r^0 \mathbb{Z}_2\{\eta_8^2\}$ is split onto.

- $\pi_{11}(J_{2,4}^r)$. Since $\pi_{11}(S^4) = 0$, Remark 4.2 implies

$$0 \rightarrow \mathbb{Z}_{(2)}\{\bar{\beta}_{11}\} \xrightarrow{\bar{\tau}_{8*}} \pi_{11}(J_{2,4}^r) \xrightarrow{\bar{p}_{8*}} \mathbb{Z}_{2^{m_{r+1}^3}}\{\delta_{r+1}\nu_8\} \rightarrow 0.$$

The rest only requires proof of the split of above short exact sequence, which we only prove for $r = 1$ since the proof of other cases are just similar.

By Lemma 3.8, there is $-\mathbf{x} \in \{\beta_4^1, \gamma_2^1, 2\nu_7\} \subset \pi_{11}(J_2(2\iota_4))$, such that $\bar{p}_{8*}(\mathbf{x}) = 2\nu_8$. $o(2\nu_8) = 4$ implies that $o(\mathbf{x}) \geq 4$. On the other hand, $4\mathbf{x} \in -\{\beta_4^1, \gamma_2^1, 2\nu_7\}(4\iota_{11}) = \beta_4^1\{\gamma_2^1, 2\nu_7, 4\iota_{10}\} \subset \beta_{4*}^1(\pi_{11}(S^4)) = \{0\}$, which implies that $o(\mathbf{x}) \leq 4$. Hence $o(\mathbf{x}) = 4$.

So $\pi_{11}(J_{2,4}^r) \xrightarrow{\bar{p}_{8*}} \mathbb{Z}_{2^{m_{r+1}^3}}\{\delta_{r+1}\nu_8\}$ is split onto.

- $\pi_i(J_{2,4}^r)$ ($i = 12, 13$). These are easily obtained by the following isomorphisms for $i = 12, 13$ from Lemma 4.1

$$\frac{\bar{\beta}_{4*}(\pi_i(S^4))}{\bar{\beta}_{4*}\gamma_{2*}^r(\pi_i(S^7))} \oplus \bar{\beta}_{11*}(\pi_i(S^{11})) \xrightarrow{\bar{\tau}_{8*} \cong} \pi_i(J_{2,4}^r).$$

□

Lemma 5.5. For $r \geq 0$, $\gamma_3^r = 2^r a_0 \beta_{11} \in \pi_{11}(J_{2,4}^r)$, where a_0 is an odd integer in $\mathbb{Z}_{(2)}$.

Proof. From Lemma 5.4, we assume

$$\gamma_3^r = a_r \beta_{11}^r + b_r \widehat{\delta_{r+1}\nu_8} = a_r \bar{\tau}_8^r \bar{\beta}_{11}^r + b_r \widehat{\delta_{r+1}\nu_8}, a_r \in \mathbb{Z}_{(2)}, b_r \in \mathbb{Z}_{2^{m_{r+1}^3}}.$$

By Lemma 2.8 and $\bar{p}_8^r \bar{\tau}_8^r \simeq 0$, we get $0 = \bar{p}_8^r \gamma_3^r = \bar{p}_8^r (b_r \widehat{\delta_{r+1}\nu_8}) = b_r \delta_{r+1}\nu_8$. Hence $b_r = 0$, which implies $\gamma_3^r = a_r \beta_{11}^r$.

By exact sequence (6) for $k = 4, r = 0, m = 11$, we get

$$\begin{aligned} \mathbb{Z}_2 &\cong \pi_{12}(S^5) \cong \pi_{11}(F_5^0) \cong \pi_{11}(J_{3,4}^0) \cong \pi_{11}(J_{2,4}^0) / \text{Im}(\gamma_{3*}^0) \\ &= \frac{\mathbb{Z}_{(2)}\{\beta_{11}^0\} \oplus \mathbb{Z}_2\{\widehat{4\nu_8}\}}{\langle a_0 \beta_{11}^0 \rangle} \quad \Rightarrow \quad a_0 \text{ is an odd integer in } \mathbb{Z}_{(2)}. \end{aligned}$$

$$\begin{aligned} 2^{2r} \gamma_3^0 &= \bar{g}_0^r \gamma_3^r = a_r \bar{g}_0^r (\bar{\tau}_8^r \bar{\beta}_{11}^r) \quad (\text{by (12)}) \\ &= a_r \bar{\tau}_8^0 g_0^r \bar{\beta}_{11}^r = a_r \bar{\tau}_8^0 g_0^r I_2^{\gamma, r} j_2^{11} = a_r \bar{\tau}_8^0 I_2^{\gamma, 0} (g_0^r)|_{J_2} j_2^{11} \quad (\text{by (10) and (14)}) \\ &= a_r \bar{\tau}_8^0 I_2^{\gamma, 0} (j_1^4 q_1^4 + 2^r j_2^{11} q_2^{11}) j_2^{11} = 2^r a_r \bar{\tau}_8^0 \bar{\beta}_{11}^0 \quad (\text{by (15) and } \pi_{11}(S^4) = 0) \\ \text{i.e., } 2^{2r} a_0 \bar{\tau}_8^0 \bar{\beta}_{11}^0 &= 2^r a_r \bar{\tau}_8^0 \bar{\beta}_{11}^0 \quad \Rightarrow \quad a_r = 2^r a_0. \end{aligned}$$

□

Lemma 5.6. For $r \geq 0$, $\pi_i(F_5) \cong \pi_i(J_{3,4}^r)$ ($i \leq 14$) and

- (1) $\pi_{11}(J_{3,4}^r) = \mathbb{Z}_{2^r} \{\check{\beta}_{11}\} \oplus \mathbb{Z}_{2^{m_{r+1}^3}} \{\widehat{I_2^r \delta_{r+1} \nu_8}\}$.
- (2) $\pi_{12}(J_{3,4}^r) = \mathbb{Z}_2 \{\check{\beta}_4 \varepsilon_4\} \oplus \vartheta_r^0 \mathbb{Z}_2 \{\check{\beta}_{11} \eta_{11}\}$.
- (3) $\pi_{13}(J_{3,4}^0) = \mathbb{Z}_2 \{\check{\beta}_4 \nu_4^3\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \mu_4\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \eta_4 \varepsilon_5\}$.
 $\pi_{13}(J_{3,4}^1) = \mathbb{Z}_2 \{\check{\beta}_4 \nu_4^3\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \mu_4\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \eta_4 \varepsilon_5\} \oplus \mathbb{Z}_4 \{\widehat{\eta}_{12}\}$ with $\check{\beta}_{11} \eta_{11}^2 = 2\widehat{\eta}_{12}$.
 $\pi_{13}(J_{3,4}^r) = \mathbb{Z}_2 \{\check{\beta}_4 \nu_4^3\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \mu_4\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \eta_4 \varepsilon_5\} \oplus \mathbb{Z}_2 \{\check{\beta}_{11} \eta_{11}^2\} \oplus \mathbb{Z}_2 \{\widehat{\eta}_{12}\}$ ($r \geq 2$).

Proof. From Lemmas 5.4 and 5.5, (1) and (2) of this lemma are easily get by the following isomorphism

$$\frac{\pi_m(J_{2,4}^r)}{2^r a_0 \beta_{11*} (\pi_m(S^{11}))} \xrightarrow{I_{2*}^r} \pi_m(J_{3,4}^r).$$

For (3), the case $r = 0$ is easily obtained. For $r \geq 1$, we have the following short exact sequence by Lemma 5.4

$$0 \rightarrow \pi_{13}(J_{2,4}^r) \xrightarrow{I_{2*}^r} \pi_{13}(J_{3,4}^r) \xrightarrow{\check{p}_{12*}} \mathbb{Z}_2 \{\eta_{12}\} \rightarrow 0$$

For $r \geq 1$, by Lemma 3.8, there is $-\mathbf{x} \in \{I_2^r, \gamma_3^r, \eta_{11}\} \subset \pi_{13}(J_{3,4}^r)$, such that $\check{p}_{12*}(\mathbf{x}) = \eta_{12}$ and $2\mathbf{x} \in -\{I_2^r, \gamma_3^r, \eta_{11}\}(2\iota_{13}) = I_2^r \{\gamma_3^r, \eta_{11}, 2\iota_{12}\} = I_2^r \{2^r a_0 \beta_{11}, \eta_{11}, 2\iota_{12}\}$.

$$\{2^r a_0 \beta_{11}, \eta_{11}, 2\iota_{12}\} \ni 2^{r-1} a_0 \beta_{11} \eta_{11}^2 \bmod \pi_{13}(J_{2,4}^r) \circ (2\iota_{13}) + (2^r a_0 \beta_{11}) \circ \pi_{13}(S^{11}) = \{0\}$$

Thus $2\mathbf{x} = 0$ for $r \geq 2$, implies that

$$\pi_{13}(J_{3,4}^r) = \mathbb{Z}_2 \{\check{\beta}_4 \nu_4^3\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \mu_4\} \oplus \mathbb{Z}_2 \{\check{\beta}_4 \eta_4 \varepsilon_5\} \oplus \mathbb{Z}_2 \{\check{\beta}_{11} \eta_{11}^2\} \oplus \mathbb{Z}_2 \{\widehat{\eta}_{12}\}.$$

$2\mathbf{x} = I_2^r \beta_{11} \eta_{11}^2 = \check{\beta}_{11} \eta_{11}^2$ for $r = 1$, we get $o(\mathbf{x}) = 4$ and this implies the result of $\pi_{13}(J_{3,4}^1)$ in (3) of Lemma 5.6. \square

Next we compute $\pi_{14}(J_{2,4}^r)$ and $\pi_{14}(J_{3,4}^r)$. We have the following exact sequence with commutative diagrams.

$$\begin{array}{ccccccc} \pi_{15}(S^8) & \xrightarrow{\bar{\delta}_{14*}^8} & \pi_{14}(F_{\bar{p}_8}) & \xrightarrow{\bar{\tau}_{8*}} & \pi_{14}(J_{2,4}^r) & \xrightarrow{\bar{p}_{8*}} & \pi_{14}(S^8) \xrightarrow{\bar{\delta}_{13*}^8} \pi_{13}(F_{\bar{p}_8}) \\ \uparrow \Sigma & & \uparrow \bar{\beta}_{4*} & \nearrow \beta_{4*} & & & \uparrow \cong \Sigma \\ \pi_{14}(S^7) & \xrightarrow{\gamma_{2*}^r} & \pi_{14}(S^4) & & & & \pi_{13}(S^7) \xrightarrow{\gamma_{2*}^r} \pi_{13}(S^4) \\ & & & & & & \uparrow \bar{\beta}_{4*} \end{array} \quad (24)$$

where $\pi_{15}(S^8) = \mathbb{Z}_{(2)} \{\sigma_8\} \oplus \mathbb{Z}_8 \{\Sigma \sigma'\}$, $\pi_{14}(S^7) = \mathbb{Z}_8 \{\sigma'\}$,
 $\pi_{14}(F_{\bar{p}_8}) = \mathbb{Z}_8 \{\bar{\beta}_4 \nu_4 \sigma'\} \oplus \mathbb{Z}_4 \{\bar{\beta}_4 \Sigma \varepsilon'\} \oplus \mathbb{Z}_2 \{\bar{\beta}_4 \eta_4 \mu_5\} \oplus \mathbb{Z}_8 \{\bar{\beta}_{11} \nu_{11}\} \oplus \mathbb{Z}_{(2)} \{[\bar{\beta}_4, \bar{\beta}_{11}]\}$.

Lemma 5.7. For $r \geq 1$, we have the following split short exact sequence

$$0 \rightarrow \text{Coker}(\bar{\partial}_{14*}^{8,r}) \xrightarrow{\bar{\tau}_{8*}} \pi_{14}(J_{2,4}^r) \xrightarrow{\bar{p}_{8*}} \pi_{14}(S^8) = \mathbb{Z}_2\{v_8^2\} \rightarrow 0, \text{ i.e.,}$$

$$\pi_{14}(J_{2,4}^r) = \bar{\tau}_{8*}(\text{Coker}(\bar{\partial}_{14*}^{8,r})) \oplus \mathbb{Z}_2\{\widehat{v}_8^2\}, \text{ where}$$

$$\bar{\tau}_{8*}(\text{Coker}(\bar{\partial}_{14*}^{8,r})) = \frac{\mathbb{Z}_8\{\beta_4 v_4 \sigma'\} \oplus \mathbb{Z}_4\{\beta_4 \Sigma \varepsilon'\} \oplus \mathbb{Z}_2\{\beta_4 \eta_4 \mu_5\} \oplus \mathbb{Z}_8\{\beta_{11} v_{11}\} \oplus \mathbb{Z}_{(2)}\{[\beta_4, \beta_{11}]\}}{\langle 2^{r+1} \beta_4 v_4 \sigma', \pm 2^r \beta_4 v_4 \sigma' + 2^r \beta_4 \Sigma \varepsilon' + 2^r (1 - \vartheta_r) c_0 \beta_{11} v_{11} + 2^r d_0 [\beta_4, \beta_{11}] \rangle}$$

c_0 and d_0 are some integers.

Proof.

$$\begin{aligned} \gamma_{2*}^r(v_7^2) &= (2^{r+1}v_4 - 2^r\Sigma v')(v_7^2) = 0 \text{ by } v'_7 v_6 = 0 \text{ and } 2v_4^3 = 0, \\ \Rightarrow \text{Ker}(\bar{\partial}_{13*}^{8,r}) &= \pi_{14}(S^8) = \mathbb{Z}_2\{v_8^2\} (r \geq 0). \end{aligned} \quad (25)$$

$$(t\alpha)\sigma' = t\alpha\sigma', t \in \mathbb{Z}, \alpha \in \pi_7(X), \text{ from Lemma 4.5 of [39]}. \quad (26)$$

$$\begin{aligned} \gamma_{2*}^r(\sigma') &= (2^{r+1}v_4 - 2^r\Sigma v')(\sigma') \\ &= (2^{r+1}v_4)\sigma' - (2^r\Sigma v')\sigma' \pm [2^{r+1}v_4, 2^r\Sigma v']H_2(\sigma') \text{ (Proposition 2.10 of [33])} \\ &\quad [2^{r+1}v_4, 2^r\Sigma v']H_2(\sigma') = [v_4, \Sigma v'](2^{2r+1}H_2(\sigma')) = [v_4, \Sigma v'](2^{2r+1}\eta_{13}) = 0 \\ \Rightarrow \gamma_{2*}^r(\sigma') &= 2^{r+1}v_4\sigma' - 2^r\Sigma v'\sigma' = \begin{cases} 2^{r+1}v_4\sigma', & r \geq 1; \\ 2v_4\sigma' - 2\Sigma\varepsilon', & r = 0 \end{cases} \text{ (by Lemma 8.1 (i))} \\ \Rightarrow \bar{\partial}_{14*}^{8,r}(\Sigma\sigma') &= \begin{cases} 2^{r+1}\bar{\beta}_4^r v_4\sigma', & r \geq 1; \\ 2\bar{\beta}_4^0 v_4\sigma' - 2\bar{\beta}_4^0 \Sigma\varepsilon', & r = 0 \end{cases} \end{aligned} \quad (27)$$

Assume $\bar{\partial}_{14*}^{8,r}(\sigma_8) = x_r \bar{\beta}_4^r v_4 \sigma' + y_r \bar{\beta}_4^r \Sigma \varepsilon' + z_r \bar{\beta}_4^r \eta_4 \mu_5 + c_r \bar{\beta}_{11}^r v_{11} + d_r [\bar{\beta}_4^r, \bar{\beta}_{11}^r]$, where $x_r \in \mathbb{Z}_8, y_r \in \mathbb{Z}_4, z_r \in \mathbb{Z}_2, c_r \in \mathbb{Z}_8, d_r \in \mathbb{Z}_{(2)}, r \geq 0$.

Then by (23),

$$g_{0*}^r \bar{\partial}_{14*}^{8,r}(\sigma_8) = x_r \bar{\beta}_4^0 v_4 \sigma' + y_r \bar{\beta}_4^0 \Sigma \varepsilon' + z_r \bar{\beta}_4^0 \eta_4 \mu_5 + 2^r c_r \bar{\beta}_{11}^0 v_{11} + 2^r d_r [\bar{\beta}_4^0, \bar{\beta}_{11}^0].$$

On the other hand, from left commutative diagram (13) for $s = r, t = 0, k = 4$,

$$\begin{aligned} g_{0*}^r \bar{\partial}_{14*}^{8,r}(\sigma_8) &= \bar{\partial}_{14*}^{8,0}(2^r \iota_8)_*(\sigma_8) = 2^{2r} \bar{\partial}_{14*}^{8,0}(\sigma_8) - 2^{r-1}(2^r - 1) \bar{\partial}_{14*}^{8,0}(\Sigma\sigma') \\ &= 2^{2r} (x_0 \bar{\beta}_4^0 v_4 \sigma' + y_0 \bar{\beta}_4^0 \Sigma \varepsilon' + z_0 \bar{\beta}_4^0 \eta_4 \mu_5 + c_0 \bar{\beta}_{11}^0 v_{11} + d_0 [\bar{\beta}_4^0, \bar{\beta}_{11}^0]) \\ &\quad - 2^{r-1} (2^r - 1) (2 \bar{\beta}_4^0 v_4 \sigma' - 2 \bar{\beta}_4^0 \Sigma \varepsilon') \\ &= \begin{cases} (4x_0 - 2) \bar{\beta}_4^0 v_4 \sigma' + 2 \bar{\beta}_4^0 \Sigma \varepsilon' + 4c_0 \bar{\beta}_{11}^0 v_{11} + 4d_0 [\bar{\beta}_4^0, \bar{\beta}_{11}^0], & r = 1; \\ 4 \bar{\beta}_4^0 v_4 \sigma' + 2^4 d_0 [\bar{\beta}_4^r, \bar{\beta}_{11}^r], & r = 2; \\ 2^r d_0 [\bar{\beta}_4^r, \bar{\beta}_{11}^r], & r \geq 3. \end{cases} \end{aligned}$$

By comparing above two equations of $g_{0*}^r \bar{\partial}_{14*}^{8,r}(\sigma_8)$, we get

$$\begin{aligned} x_r &= \pm 2^r, \quad y_r = 2^r, \quad z_r = 0, \quad c_r = 2^r(1-\vartheta_r)c_0, \quad d_r = 2^r d_0 \quad (r \geq 1). \\ \bar{\partial}_{14*}^{8,r}(\sigma_8) &= \pm 2^r \bar{\beta}_4^r v_4 \sigma' + 2^r \bar{\beta}_4^r \Sigma \varepsilon' + 2^r(1-\vartheta_r)c_0 \bar{\beta}_{11}^r v_{11} + 2^r d_0 [\bar{\beta}_4^r, \bar{\beta}_{11}^r]. \end{aligned}$$

Together with (27), we have

$$\text{Coker}(\bar{\partial}_{14*}^{8,r}) = \frac{\mathbb{Z}_8\{\bar{\beta}_4^r v_4 \sigma'\} \oplus \mathbb{Z}_4\{\bar{\beta}_4^r \Sigma \varepsilon'\} \oplus \mathbb{Z}_2\{\bar{\beta}_4^r \eta_4 \mu_5\} \oplus \mathbb{Z}_8\{\bar{\beta}_{11}^r v_{11}\} \oplus \mathbb{Z}_{(2)}\{[\bar{\beta}_4^r, \bar{\beta}_{11}^r]\}}{\langle 2^{r+1} \bar{\beta}_4^r v_4 \sigma', \quad \pm 2^r \bar{\beta}_4^r v_4 \sigma' + 2^r \bar{\beta}_4^r \Sigma \varepsilon' + 2^r(1-\vartheta_r)c_0 \bar{\beta}_{11}^r v_{11} + 2^r d_0 [\bar{\beta}_4^r, \bar{\beta}_{11}^r] \rangle}$$

Now we get the short exact sequence of Lemma 5.7 by above equation and (25). It remains to show the split.

By Lemma 3.8, there is $-\mathbf{x} \in \{\beta_4^r, \gamma_2^r, \nu_7^2\} \subset \pi_{14}(J_{2,4}^r)$, such that $\bar{p}_{8*}(\mathbf{x}) = \nu_8^2$. $2\mathbf{x} \in -\{\beta_4^r, \gamma_2^r, \nu_7^2\}(2\iota_{14}) = \beta_4^r\{\gamma_2^r, \nu_7^2, 2\iota_{13}\}$.

$$\begin{aligned} \{\gamma_2^r, \nu_7^2, 2\iota_{13}\} \ni \gamma_2^{r-1} \nu_7^2 \eta_{13} &= 0 \pmod{\pi_{14}(S^4)(2\iota_{14}) + (2^r[\iota_4, \iota_4])\pi_{14}(S^7)} \\ (2^r[\iota_4, \iota_4])\sigma' &= (2^{r+1}v_4 - 2^r \Sigma v')\sigma' = 2^{r+1}v_4 \sigma' - 2^{r+1} \Sigma \varepsilon', \end{aligned}$$

we get $\{\gamma_2^r, \nu_7^2, 2\iota_{13}\} = 2\pi_{14}(S^4)$, $\Rightarrow 2\mathbf{x} = \beta_4^r(2\theta) = 2\beta_4^r \theta$, $\theta \in \pi_{14}(S^4)$.

Let $\mathbf{x}' = \mathbf{x} - \beta_4^r \theta$, we have $\bar{p}_{8*}(\mathbf{x}') = \nu_8^2$, $o(\mathbf{x}') = 2$.

So the split of the short exact sequence in Lemma 5.7 is obtained. \square

There is the following exact sequence with commutative squares

$$\begin{array}{ccccc} \mathbb{Z}_8\{v_{12}\} = \pi_{15}(S^{12}) & \xrightarrow{\check{\delta}_{14*}^{12,r}} & \pi_{14}(F\check{p}_{12}^r) & \xrightarrow{\check{\tau}_{3k*}} & \pi_{14}(J_{3,4}^r) & \xrightarrow{\check{p}_{12*}^r} & \pi_{14}(S^{12}) & \xrightarrow{\check{\delta}_{13*}^{12,r}} & \pi_{13}(F\check{p}_{12}^r) \\ \uparrow \cong \Sigma & & \uparrow \text{empi.} & \nearrow I_{4*}^r & & & \uparrow \cong \Sigma & & \uparrow \cong \\ \mathbb{Z}_8\{v_{11}\} = \pi_{14}(S^{11}) & \xrightarrow{\gamma_{3*}^r} & \pi_{14}(J_{2,4}^r) & & & & \pi_{13}(S^{11}) & \xrightarrow{\gamma_{3*}^r} & \pi_{13}(J_{2,4}^r) \end{array}$$

where $\pi_{14}(F\check{p}_{12}^r) \cong \frac{\pi_{14}(J_{2,4}^r)}{\langle 2^r a_0 [\beta_4^r, \beta_{11}^r] \rangle}$ by (8).

Lemma 5.8. For $r \geq 1$,

$$\pi_{14}(J_{3,4}^r) = \begin{cases} \mathbb{Z}_2\{\check{\beta}_4 v_4 \sigma' + \check{\beta}_4 \Sigma \varepsilon'\} \oplus H_{14}^1, & r = 1; \\ \mathbb{Z}_{2m_3^r}\{\check{\beta}_4 v_4 \sigma'\} \oplus H_{14}^r, & r \geq 2. \end{cases}$$

$$H_{14}^r = \mathbb{Z}_4\{\check{\beta}_4 \Sigma \varepsilon'\} \oplus \mathbb{Z}_2\{\check{\beta}_4 \eta_4 \mu_5\} \oplus \mathbb{Z}_{2m_3^r}\{\check{\beta}_{11} v_{11}\} \oplus \mathbb{Z}_{2^r}\{[\check{\beta}_4, \check{\beta}_{11}]\} \oplus \mathbb{Z}_2\{\widehat{I}_2^r \widehat{\nu}_8^2\} \oplus \mathbb{Z}_2\{\widehat{\eta}_{12} \eta_{13}\}.$$

Proof. From Lemma 5.5, we have the following short exact sequence

$$0 \rightarrow \text{Coker}(\check{\delta}_{14*}^{12,r}) \rightarrow \pi_{14}(J_{3,4}^r) \xrightarrow{\check{p}_{12*}^r} \pi_{14}(S^{12}) \rightarrow 0,$$

$$\text{where } \text{Coker}(\check{\delta}_{14*}^{12,r}) \cong \frac{\pi_{14}(J_{2,4}^r)}{\langle 2^r a_0 [\beta_4, \beta_{11}], 2^r a_0 \beta_{11} v_{11} \rangle}.$$

Moreover, the order 2 element $\widehat{\eta}_{12}\eta_{13}$ ($\widehat{\eta}_{12}$ is get in Lemma 5.6) satisfies $\check{p}_{12*}^r(\widehat{\eta}_{12}\eta_{13}) = \eta_{12}^2$, which implies $\pi_{14}(J_{3,4}^r) \xrightarrow{\check{p}_{12*}^r} \pi_{14}(S^{12})$ is split onto. Then this lemma is obtained by the following computation of $Coker(\check{\partial}_{14*}^{12,r})$ by Lemma 5.7, i.e., $Coker(\check{\partial}_{14*}^{12,r})$ is isomorphic to

$$\begin{aligned} & \frac{\mathbb{Z}_8\{\beta_4\nu_4\sigma'\} \oplus \mathbb{Z}_4\{\beta_4\Sigma\varepsilon'\} \oplus \mathbb{Z}_2\{\beta_4\eta_4\mu_5\} \oplus \mathbb{Z}_8\{\beta_{11}\nu_{11}\} \oplus \mathbb{Z}_{(2)}\{[\beta_4, \beta_{11}]\} \oplus \mathbb{Z}_2\{\widehat{\nu}_8^2\}}{\langle 2^{r+1}\beta_4\nu_4\sigma', \pm 2^r\beta_4\nu_4\sigma' + 2^r\beta_4\Sigma\varepsilon' + 2^r(1-\vartheta_r)c_0\beta_{11}\nu_{11} + 2^r d_0[\beta_4, \beta_{11}], 2^r a_0[\beta_4, \beta_{11}], 2^r a_0\beta_{11}\nu_{11} \rangle} \\ &= \frac{\mathbb{Z}_8\{\beta_4\nu_4\sigma'\} \oplus \mathbb{Z}_4\{\beta_4\Sigma\varepsilon'\} \oplus \mathbb{Z}_8\{\beta_{11}\nu_{11}\} \oplus \mathbb{Z}_{(2)}\{[\beta_4, \beta_{11}]\}}{\langle 2^{r+1}\beta_4\nu_4\sigma', \pm 2^r\beta_4\nu_4\sigma' + 2^r\beta_4\Sigma\varepsilon', 2^r a_0[\beta_4, \beta_{11}], 2^r a_0\beta_{11}\nu_{11} \rangle} \oplus \mathbb{Z}_2\{\beta_4\eta_4\mu_5\} \oplus \mathbb{Z}_2\{\widehat{\nu}_8^2\} \\ &= \frac{\mathbb{Z}_8\{\beta_4\nu_4\sigma'\} \oplus \mathbb{Z}_4\{\beta_4\Sigma\varepsilon'\}}{\langle 2^{r+1}\beta_4\nu_4\sigma', \pm 2^r\beta_4\nu_4\sigma' + 2^r\beta_4\Sigma\varepsilon' \rangle} \oplus \mathbb{Z}_2\{\beta_4\eta_4\mu_5\} \oplus \mathbb{Z}_{2m^3}\{\beta_{11}\nu_{11}\} \oplus \mathbb{Z}_{2^r}\{[\beta_4, \beta_{11}]\} \oplus \mathbb{Z}_2\{\widehat{\nu}_8^2\} \\ &= \begin{cases} \mathbb{Z}_2\{\beta_4\nu_4\sigma' + \beta_4\Sigma\varepsilon'\} \oplus \mathbb{Z}_4\{\beta_4\Sigma\varepsilon'\} \oplus \mathbb{Z}_2\{\beta_4\eta_4\mu_5\} \oplus \mathbb{Z}_2\{\beta_{11}\nu_{11}\} \oplus \mathbb{Z}_2\{[\beta_4, \beta_{11}]\} \oplus \mathbb{Z}_2\{\widehat{\nu}_8^2\}, & r = 1; \\ \mathbb{Z}_{2m^3}\{\beta_4\nu_4\sigma'\} \oplus \mathbb{Z}_4\{\beta_4\Sigma\varepsilon'\} \oplus \mathbb{Z}_2\{\beta_4\eta_4\mu_5\} \oplus \mathbb{Z}_{2m^3}\{\beta_{11}\nu_{11}\} \oplus \mathbb{Z}_{2^r}\{[\beta_4, \beta_{11}]\} \oplus \mathbb{Z}_2\{\widehat{\nu}_8^2\}, & r \geq 2. \end{cases} \end{aligned}$$

□

- $\pi_9(P^5(2^r))$.

Consider diagram (6) for $k = 4, m = 9$.

$$\partial_{8*}^{5,r}(\nu_5\eta_8) = \beta_{4*}(2^r\iota_4)_*(\nu_4\eta_7) = (2^{2r}\nu_4 - 2^{r-1}(2^r-1)\Sigma\nu')\eta_7 = \begin{cases} \beta_4\Sigma\nu'\eta_7, & r = 1; \\ 0, & r \geq 2. \end{cases}$$

$$\pi_8(F_5) \cong \pi_8(J_{2,4}^r) = \mathbb{Z}_2\{\beta_4\nu_4\eta_7\} \oplus \mathbb{Z}_2\{\beta_4\Sigma\nu'\eta_7\}, \text{ by (9) of [23]}$$

$$\Rightarrow Ker(\partial_{8*}^{5,r}) = \vartheta_r\mathbb{Z}_2\{\nu_5\eta_8\}.$$

$$\partial_{9*}^{5,r}(\nu_5\eta_8^2) = \beta_{4*}(2^r\iota_4)_*(\nu_4\eta_7^2) = \begin{cases} \beta_4\Sigma\nu'\eta_7^2, & r = 1; \\ 0, & r \geq 2. \end{cases} \quad (28)$$

$$\text{Lemma 5.4} \Rightarrow Coker(\partial_{9*}^{5,r}) = \mathbb{Z}_2\{\beta_4\nu_4\eta_7^2\} \oplus \vartheta_r\mathbb{Z}_2\{\beta_4\Sigma\nu'\eta_7^2\} \oplus \mathbb{Z}_2\{\widehat{\eta}_8\}. \quad (29)$$

Thus we have $\pi_9(P^5(2)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ and the short exact sequence

$$0 \rightarrow \mathbb{Z}_2^{\oplus 3} \cong Coker(\partial_{9*}^{5,r}) \rightarrow \pi_9(P^5(2^r)) \xrightarrow{p_{5*}} \mathbb{Z}_2\{\nu_5\eta_8\} \rightarrow 0 \quad (r \geq 2).$$

By Lemma 3.8, there is $-\mathbf{x}_r \in \{i_4, 2^r\iota_4, \nu_4\eta_7\}$, such that $p_{5*}(\mathbf{x}_r) = \nu_5\eta_8$.

$2\mathbf{x}_r \in -\{i_4, 2^r\iota_4, \nu_4\eta_7\}(2\iota_9) = i_4\{2^r\iota_4, \nu_4\eta_7, 2\iota_8\}$. Hence $2\mathbf{x}_r = i_4\Sigma\nu'\eta_7^2$ and 0 for $r = 2$ and $r \geq 3$ respectively by (ii) of Lemma 8.2, which implies $o(\mathbf{x}_2) = 4$ and $o(\mathbf{x}_r) = 2, r \geq 3$. So

$$\pi_9(P^5(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4, & r = 2; \\ \mathbb{Z}_2^{\oplus 4}, & r \geq 3. \end{cases}$$

- $\pi_{10}(P^5(2^r))$.

Consider diagram (6) for $k = 4, m = 10$.
 Computation of $\pi_9(P^5(2^r))$ implies $Ker(\partial_{9*}^{5,r}) = \vartheta_r \mathbb{Z}_2\{v_5\eta_8^2\}$.

$$\begin{aligned}\partial_{10*}^{5,r}(v_5^2) &= \beta_{4*}(2^r \iota_4)_*(v_4^2) = \beta_4(2^{2r} v_4 - 2^{r-1}(2^r - 1)\Sigma v')v_7 \\ &= 2^{2r} \beta_4 v_4^2 = 0, \text{ by Lemma 5.4.} \\ \Rightarrow Coker(\partial_{10*}^{5,r}) &= \mathbb{Z}_{2^{m_{r+1}^3}}\{\beta_4 v_4^2\} \oplus \mathbb{Z}_2\{\widehat{\eta_8^2}\}.\end{aligned}\tag{30}$$

Thus we have $\pi_{10}(P^5(2)) \cong \mathbb{Z}_4 \oplus \mathbb{Z}_2$ and the short exact sequence

$$0 \rightarrow \mathbb{Z}_{2^{m_{r+1}^3}} \oplus \mathbb{Z}_2 \cong Coker(\partial_{10*}^{5,r}) \rightarrow \pi_{10}(P^5(2^r)) \xrightarrow{p_{5*}^{5,r}} \mathbb{Z}_2\{v_5\eta_8^2\} \rightarrow 0 (r \geq 2)$$

Since there is an $\mathbf{x} \in \pi_9(P^5(2^r)) (r \geq 2)$ such that $p_{5*}(\mathbf{x}) = v_5\eta_8$, the order 2 element $\mathbf{x}\eta_9 \in \pi_{10}(P^5(2^r))$ satisfies $p_{5*}(\mathbf{x}\eta_9) = v_5\eta_8^2$, i.e., $\pi_{10}(P^5(2^r)) \xrightarrow{p_{5*}^{5,r}} \mathbb{Z}_2\{v_5\eta_8^2\}$ is split onto. We get

$$\pi_{10}(P^5(2^r)) \cong \mathbb{Z}_{2^{m_{r+1}^3}} \oplus \mathbb{Z}_2^{\oplus 2} (r \geq 2).$$

- $\pi_{11}(P^5(2^r))$.

We have the exact sequence with $\pi_{11}(F_5)$ obtained by Lemmas 5.5, 5.6

$$\begin{aligned}\pi_{12}(S^5) &\xrightarrow{\partial_{11*}^{5,r}} \pi_{11}(F_5) \xrightarrow{\tau_{5*}} \pi_{11}(P^5(2^r)) \xrightarrow{p_{5*}} \pi_{11}(S^5) \xrightarrow{\partial_{10*}^{5,r}=0} \pi_{10}(F_5) \\ \pi_{11}(F_5) &\cong \pi_{11}(J_{3,4}^r) \cong \mathbb{Z}_{2^r}\{\check{\beta}_{11}^r\} \oplus \mathbb{Z}_{2^{m_{r+1}^3}}\{\widehat{\delta_{r+1}v_8}\}.\end{aligned}$$

The right commutative diagram of (9) for $k = 4, s = 1, t = r \geq 2$ induces

$$\begin{array}{ccccccccc} \mathbb{Z}_2\{\sigma'''\} = \pi_{12}(S^5) & \xrightarrow{\partial_{11*}^{5,1}} & \pi_{11}(F_5^1) & \xrightarrow{\tau_{5*}^1} & \pi_{11}(P^5(2)) & \xrightarrow{p_{5*}^1} & \pi_{11}(S^5) & \longrightarrow & 0 \\ \parallel & & \downarrow \chi_{r*}^1 & & \downarrow \bar{\chi}_{r*}^1 & & \parallel & & \\ \pi_{12}(S^5) & \xrightarrow{\partial_{11*}^{5,r}} & \pi_{11}(F_5^r) & \xrightarrow{\tau_{5*}^r} & \pi_{11}(P^5(2^r)) & \xrightarrow{p_{5*}^r} & \pi_{11}(S^5) & \longrightarrow & 0 \end{array}$$

By Theorem 5.10 of [43], $\pi_{11}(P^5(2)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_4$. It implies $\partial_{11*}^{5,1}(\sigma''') = 0$, hence $\partial_{11*}^{5,r}(\sigma''') = \chi_{r*}^1 \partial_{11*}^{5,1}(\sigma''') = 0$. So we get the following short exact sequence which is split for $r = 1$.

$$0 \longrightarrow \mathbb{Z}_{2^r} \oplus \mathbb{Z}_{2^{m_{r+1}^3}} \cong Coker(\partial_{11*}^{5,r}) \longrightarrow \pi_{11}(P^5(2^r)) \xrightarrow{p_{5*}^r} \mathbb{Z}_2\{v_5^2\} \longrightarrow 0 \tag{31}$$

There is an order 2 element $\mathbf{x}_1 \in \pi_{11}(P^5(2))$ such that $p_{5*}^1(\mathbf{x}_1) = v_5^2$. Then the order 2 element $\mathbf{x}_r := \bar{\chi}_{r*}^1 \mathbf{x}_1 \in \pi_{11}(P^5(2^r)) (r \geq 2)$ makes $p_{5*}^r(\mathbf{x}_r) = v_5^2$. So above exact sequence is also split for $r \geq 2$. Hence

$$\pi_{11}(P^5(2^r)) \cong \mathbb{Z}_{2^r} \oplus \mathbb{Z}_{2^{m_{r+1}^3}} \oplus \mathbb{Z}_2, r \geq 1.$$

- $\pi_{12}(P^5(2^r))$.

Consider diagram (6) for $k = 4, m = 12$.

$$\begin{aligned} \mathbb{Z}_2\{\varepsilon_5\} &= \pi_{13}(S^5) \xrightarrow{\partial_{12*}^{5,r}} \pi_{12}(F_5) \xrightarrow{\tau_{5*}} \pi_{12}(P^5(2^r)) \xrightarrow{p_{5*}} \pi_{12}(S^5) \xrightarrow{\partial_{11*}^{5,r}=0} \pi_{11}(F_5) \\ \partial_{12*}^{5,r}(\varepsilon_5) &= \beta_{4*}(2^r \iota_4)_*(\varepsilon_4) = 0 \implies \text{Coker}(\partial_{12*}^{5,r}(\varepsilon_5)) = \pi_{12}(F_5) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2. \\ 0 &\longrightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2 \longrightarrow \pi_{12}(P^5(2^r)) \xrightarrow{p_{5*}} \mathbb{Z}_2\{\sigma'''\} \longrightarrow 0. \end{aligned} \quad (32)$$

Since $\pi_{12}(P^5(2)) \cong \mathbb{Z}_2^{\oplus 3}$ by Theorem 5.10 of [43], the exact sequence (32) splits for $r = 1$. By the same proof as that of the split of short exact sequence (31), we get (32) splits for any $r \geq 1$. Hence

$$\pi_{12}(P^5(2^r)) \cong \mathbb{Z}_2^{\oplus 3}.$$

- $\pi_{13}(P^5(2^r))$.

Consider diagram (6) for $k = 4, m = 13$.

$$\begin{aligned} \pi_{14}(S^5) &= \mathbb{Z}_2\{\nu_5^3\} \oplus \mathbb{Z}_2\{\mu_5\} \oplus \mathbb{Z}_2\{\eta_5 \varepsilon_6\}. \\ \partial_{13*}^{5,r}(\nu_5^3) &= \beta_{4*}(2^r \iota_4)_*(\nu_4^3) = \beta_4(2^{2r} \nu_4 - 2^{r-1}(2^r - 1)\Sigma \nu') \nu_7^2 = 0. \\ \text{Similarly, } \partial_{13*}^{5,r}(\mu_5) &= \partial_{13*}^{5,r}(\eta_5 \varepsilon_6) = 0. \text{ We have} \\ 0 &\longrightarrow \pi_{13}(F_5) \cong \pi_{13}(J_{3,4}^r) \longrightarrow \pi_{13}(P^5(2^r)) \xrightarrow{p_{5*}} \mathbb{Z}_2\{\varepsilon_5\} \longrightarrow 0. \end{aligned}$$

By Lemma 3.8, $\exists -\mathbf{x}_r \in \{i_4, 2^r \iota_4, \varepsilon_4\}$ such that $p_{5*}(\mathbf{x}) = \varepsilon_5$ and $2\mathbf{x} \in i_4\{2^r \iota_4, \varepsilon_4, 2\iota_{12}\}$. By Lemma 8.2 (iii), $2\mathbf{x}_1 = i_4 \eta_4 \varepsilon_5$ and $2\mathbf{x}_r = 0$ for $r \geq 2$. Hence

$$\pi_{13}(P^5(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4^{\oplus 2}, & r = 2; \\ \mathbb{Z}_2^{\oplus 6}, & r \geq 3. \end{cases}$$

- $\pi_{14}(P^5(2^r))$.

Consider diagram (6) for $k = 4, m = 14$. $\pi_{14}(S^5) = \mathbb{Z}_2\{\nu_5^3\} \oplus \mathbb{Z}_2\{\mu_5\} \oplus \mathbb{Z}_2\{\eta_5 \varepsilon_6\}$.

$$\pi_{15}(S^5) \xrightarrow{\partial_{14*}^{5,r}} \pi_{14}(F_5) \cong \pi_{14}(J_{3,4}^r) \longrightarrow \pi_{14}(P^5(2^r)) \xrightarrow{p_{5*}} \pi_{14}(S^5) \longrightarrow 0$$

Lemma 5.9. *The short exact sequence $0 \rightarrow \text{Coker}(\partial_{14*}^{5,r}) \rightarrow \pi_{14}(P^5(2^r)) \rightarrow \pi_{14}(S^5) \rightarrow 0$ does not split for $r = 1$ and splits for $r \geq 2$.*

Proof. We have the order 2 element $\mathbf{x}_r \in \pi_{11}(P^5(2^r))$ which is obtained in the proof of the split of the exact sequence (31) satisfies $p_{5*}(\mathbf{x}_r) = \nu_5^2$. This implies the order 2 element $\mathbf{x}_r \nu_{11} \in \pi_{14}(P^5(2^r))$ satisfies $p_{5*}(\mathbf{x}_r \nu_{11}) = \nu_5^3$. Since $\pi_6(P^5(2^r)) \xrightarrow{p_{5*}} \pi_6(S^5)$ is epimorphism, let $\mathbf{y}_r \in \pi_6(P^5(2^r))$ such that $p_{5*}(\mathbf{y}_r) = \eta_5$. Then the order 2 element

$\mathbf{y}_r \varepsilon_6 \in \pi_{13}(P^5(2^r))$ satisfies $p_{5*}(\mathbf{y}_r \varepsilon_6) = \eta_5 \varepsilon_6$. For $\mu_5 \in \pi_{14}(S^5)$, there is a $-\mathbf{z}_r \in \{i_4, 2^r \iota_4, \mu_4\}$ such that $p_{5*}(\mathbf{z}_r) = \mu_5$ with $2\mathbf{z}_r \in \{i_4, 2^r \iota_4, \mu_4\}(2\iota_{14}) = i_4\{2^r \iota_4, \mu_4, 2\iota_{13}\}$. By Lemma 8.2 (iv), $o(\mathbf{z}_1) \geq 4$ and $2\mathbf{z}_r = i_4(2\rho)$, $\rho \in \pi_{14}(S^4)$ for $r \geq 2$. Thus the short exact sequence in the lemma dose not split for $r = 1$. For $r \geq 2$, take $\mathbf{z}'_r = \mathbf{z}_r - i_4(\rho)$. Then we get $p_{5*}(\mathbf{z}'_r) = \mu_5$ with $o(\mathbf{z}'_r) = 2$, which implies the short exact sequence in the lemma splits for $r \geq 2$. \square

$\pi_{15}(S^5) = \mathbb{Z}_8\{v_5\sigma_8\} \oplus \mathbb{Z}_2\{\eta_5\mu_6\}$ and $\partial_{14*}^{5,r}(\eta_5\mu_6) = \check{\beta}_{4*}(2^r \iota_4)_*(\eta_4\mu_5) = 0$ (Here we calculate the image of $\partial_{14*}^{5,r}$ in $\pi_{14}(J_{3,4}^r)$ under isomorphism $\pi_{14}(F_5) \cong \pi_{14}(J_{3,4}^r)$).

Case : r = 1 Since $2v_5\sigma_8 = \pm\Sigma^2\varepsilon'$ [39, (7.10)],

$$2\partial_{14*}^{5,1}(v_5\sigma_8) = \partial_{14*}^{5,1}(2v_5\sigma_8) = \pm\check{\beta}_{4*}(2\iota_4)_*(\Sigma\varepsilon') = 2\check{\beta}_4\Sigma\varepsilon' \neq 0,$$

implies $\partial_{14*}^{5,1}(v_5\sigma_8) = \pm\check{\beta}_4\Sigma\varepsilon' + x_1(\check{\beta}_4v_4\sigma' + \check{\beta}_4^1\Sigma\varepsilon') + y_1\check{\beta}_4\eta_4\mu_5 + z_1\check{\beta}_{11}v_{11}$
 $+w_1[\check{\beta}_4, \check{\beta}_{11}] + u_1I_2^1v_8^2 + v_1\widehat{\eta}_{12}\eta_{13}$, $x_1, y_1, z_1, w_1, u_1, v_1 \in \mathbb{Z}_2$ (Lemma 5.8).
 $\Rightarrow \text{Coker}(\partial_{14*}^{5,1}) = (\mathbb{Z}_2\{\check{\beta}_4v_4\sigma' + \check{\beta}_4\Sigma\varepsilon'\} \oplus H_{14}^1) / \langle \partial_{14*}^{5,1}(v_5\sigma_8) \rangle$
 $= \mathbb{Z}_2\{\check{\beta}_4v_4\sigma' + \check{\beta}_4\Sigma\varepsilon'\} \oplus (H_{14}^1/\mathbb{Z}_4\{\check{\beta}_4\Sigma\varepsilon'\}) \cong \mathbb{Z}_2^{\oplus 6}$.
 $\Rightarrow \pi_{14}(P^5(2)) \cong \mathbb{Z}_2^{\oplus 7} \oplus \mathbb{Z}_4$ by Lemma 5.9.

Case : r ≥ 2 We have the following left and right homotopy commutative diagrams of cofibration sequences respectively for $r = 2$ and $r \geq 3$

$$\begin{array}{ccccccc} S^7 & \xrightarrow{2\iota_7} & S^7 & \longrightarrow & P^8(2) & \xrightarrow{p_8^1} & S^7, & S^7 & \xrightarrow{2^{2r}\iota_7} & S^7 & \longrightarrow & P^8(2^{2r}) & \xrightarrow{p_8^{2r}} & S^7 \\ \downarrow v_4 & & \downarrow 8v_4 + \Sigma v' & & \downarrow \phi_2^1 & & \downarrow v_5 & \downarrow v_4 & \downarrow v_4 & & \downarrow \bar{\phi}_r^{2r} & & \downarrow v_5 & \\ S^4 & \xrightarrow{4\iota_4} & S^4 & \longrightarrow & P^5(4) & \xrightarrow{p_5^2} & S^5 & S^4 & \xrightarrow{2^r\iota_4} & S^4 & \longrightarrow & P^5(2^r) & \xrightarrow{p_5^r} & S^5 \end{array} \quad (33)$$

There are maps $\phi_r^s : F_8^s \rightarrow F_5^r$ making the following diagrams of exact sequences commutative where $s = 2$ and $2r$ repectively for $r = 2$ and $r \geq 3$

$$\begin{array}{ccccccccc} \pi_{15}(S^8) & \xrightarrow{\partial_{14*}^{8,s}} & \pi_{14}(F_8^s) & \xrightarrow{\tau_{8*}^s} & \pi_{14}(P^8(2^s)) & \xrightarrow{p_{8*}^s} & \pi_{14}(S^8) & \xrightarrow{\partial_{13*}^{8,s}} & 0 \\ \downarrow v_{5*} & & \downarrow \phi_{r*}^s & & \downarrow \bar{\phi}_{r*}^s & & \downarrow v_{5*} & & \\ \pi_{15}(S^5) & \xrightarrow{\partial_{14*}^{5,r}} & \pi_{14}(F_5^r) & \xrightarrow{\tau_{5*}^r} & \pi_{14}(P^5(2^r)) & \xrightarrow{p_{5*}^r} & \pi_{14}(S^5) & \longrightarrow & 0 \end{array} \quad (34)$$

where $\partial_{13*}^{8,s} = 0$ in the top row since $\partial_{13*}^{8,s}(v_8^2) = \beta_{7*}(2^s \iota_7)_*(v_7^2) = 0$; $\pi_{14}(F_8^s) = \mathbb{Z}_{(2)}\{\beta_{14}^s\} \oplus \mathbb{Z}_8\{\beta_7^s\sigma'\}$; $\phi_2^1\beta_7^1 = \beta_4^2(8v_4 + \Sigma v')$ and $\bar{\phi}_r^{2r}\beta_7^{2r} = \beta_4^r v_4$ by Lemma 2.5.

Lemma 5.10. $\partial_{14*}^{8,s}(\sigma_8) = 2^s\beta_{14} + y\beta_{14}\sigma'$, where

$$\mathbb{Z}_8 \ni y = \begin{array}{|c|c|c|c|} \hline s = 1 & s = 2 & s = 3 & s \geq 4 \\ \hline odd & \pm 2 & 4 & 0 \\ \hline \end{array}$$

Proof. Lemma 2.7 for map $S^7 \xrightarrow{2^s \iota_7} S^7$ gives the following commutative diagram

$$\begin{array}{ccccc}
\pi_{15}(S^8) & \xrightarrow{\partial_{14*}^{8,s}} & \pi_{14}(F_8^s) & \xleftarrow{\cong} & \pi_{14}(S^7 \vee S^{14}) & \xrightarrow{Proj} & \pi_{14}(S^{14}) \\
H_2 \downarrow & & H_2' \downarrow & & & & \Sigma_* \cong \downarrow \\
\pi_{15}(S^{15}) & \xrightarrow{(2^s \iota_{15})_*} & \pi_{15}(S^{15}) & \xlongequal{\quad\quad\quad} & \pi_{15}(S^{15}) & & \pi_{15}(S^{15})
\end{array}$$

By the right commutative square, $H_2'(\beta_{14}) = \iota_{15}$. By the left commutative square, we get $H_2' \partial_{14*}^{8,s}(\sigma_8) = 2^s H_2(\sigma_8) = 2^s \iota_{15}$. This implies that

$$\partial_{14*}^{8,s}(\sigma_8) = 2^s \beta_{14} + y \beta_{14} \sigma', \quad y \in \mathbb{Z}_8.$$

From Lemma 2.4 of [49], there is a map $\phi : F_8^s \rightarrow \Omega S^8$ such that the following left diagram is homotopy commutative and it induces the right commutative diagram

$$\begin{array}{ccc}
\begin{array}{ccc}
\Omega S^8 & \xrightarrow{\partial^s} & F_8^s \\
\parallel & & \downarrow \beta_7 \\
\Omega S^8 & \xrightarrow{\Omega(-2^s \iota_8)} & \Omega S^8 \\
& & \downarrow \phi \\
& & J(P^8(2^s), S^7)
\end{array} & \longrightarrow & P^8(2^s) \\
& & \downarrow \\
& & J(P^8(2^s), S^7)
\end{array}
\quad
\begin{array}{ccc}
\pi_{15}(S^8) & \xrightarrow{\partial_{14*}^8} & \pi_{14}(F_8) \\
& \searrow P_1(-2^s \iota_8)_* & \downarrow P_1 \phi_* \\
& & \mathbb{Z}_8\{\Sigma \sigma'\}
\end{array}
\quad
\begin{array}{c}
\mathbb{Z}_{(2)}\{\beta_{14}\} \oplus \mathbb{Z}_8\{\beta_7 \sigma'\} \\
\parallel \\
\mathbb{Z}_8\{\Sigma \sigma'\}
\end{array}
\quad (35)$$

$P_1 : \pi_{15}(S^8) = \mathbb{Z}_{(2)}\{\sigma_8\} \oplus \mathbb{Z}_8\{\Sigma \sigma'\} \rightarrow \mathbb{Z}_8\{\Sigma \sigma'\}$ is the canonical projection.

By comparing the integral Homology $H_7(-)$, we get

$$\phi \beta_7 \simeq h \Omega \Sigma : S^7 \rightarrow \Omega S^8 = \Omega \Sigma S^7, \quad h \text{ is odd integer.}$$

By the same proof as that of Lemma A.1 of [49], $(-2^s \iota_8)_*(\sigma_8) = 2^{2s} \sigma_8 - 2^{s-1} (2^s + 1) \Sigma \sigma'$.

$$\begin{aligned}
P_1 \phi_* \partial_{14*}^8(\sigma_8) &= P_1(y \phi_*(\beta_7 \sigma') + 2^s \phi_*(\beta_{14})) = hy \Sigma \sigma' + 2^s P_1 \phi_*(\beta_{14}) \\
&= (hy + 2^s l) \Sigma \sigma' \quad (\text{for some integer } l).
\end{aligned}$$

$$P_1(-2^s \iota_8)_*(\sigma_4) = P_1(2^{2s} \sigma_8 - 2^{s-1} (2^s + 1) \Sigma \sigma') = -2^{s-1} (2^s + 1) \Sigma \sigma'.$$

From (35), we get $(hy + 2^s l) \Sigma \sigma' = -2^{s-1} (2^s + 1) \Sigma \sigma'$, thus we get the value of y in Lemma 5.10. \square

Now by commutative diagram (34) and Lemma 5.8, Lemma 5.10

$$\begin{aligned}
\partial_{14*}^{5,r}(\nu_5 \sigma_8) &= \phi_{r*}^s \partial_{14*}^{8,s}(\sigma_8) = \phi_{r*}^s(2^s \beta_{14} + y \beta_7 \sigma') = 2^s \phi_{r*}^s(\beta_{14}) + y \phi_{r*}^s \beta_7 \sigma' \\
&= \begin{cases} 2\phi_{2*}^1(\beta_{14}) + y \beta_4(8\nu_4 + \Sigma \nu') \sigma' = 2(\phi_{2*}^1(\beta_{14}) + \beta_4 \Sigma \varepsilon'), & r = 2; \\ 0, & r \geq 3. \end{cases}
\end{aligned}$$

Thus for $r \geq 3$, $\text{Coker}(\partial_{14*}^{5,r}) \cong \pi_{14}(J_{3,4}^r) \cong \mathbb{Z}_{2^r} \oplus \mathbb{Z}_8^{\oplus 2} \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_2^{\oplus 3}$.

For $r = 2$, we have $x_2, y_2, z_2, w_2 \in \mathbb{Z}_4$ such that

$$\partial_{14*}^{5,2}(\nu_5\sigma_8) = 2x_2\tilde{\beta}_4^2\nu_4\sigma' + 2y_2\tilde{\beta}_4^2\Sigma\varepsilon' + 2z_2\tilde{\beta}_{11}^2\nu_{11} + 2w_2[\tilde{\beta}_4^2, \tilde{\beta}_{11}^2], \quad (36)$$

Consider commutative diagrams in (9) for $s = 2, t = 1$,
 $\chi_{1*}^2(\partial_{14*}^{5,2}(\nu_5\sigma_8)) = \partial_{14*}^{5,1}(2t_5)_*(\nu_5\sigma_8) = \partial_{14*}^{5,1}(2\nu_5\sigma_8) \neq 0$, where the second equation holds since $\Sigma((2t_5)(\nu_5\sigma_8)) = \Sigma(2\nu_5\sigma_8)$ and $\Sigma : \pi_{15}(S^8) \rightarrow \pi_{16}(S^9)$ is a monomorphism ([39, page 69]). Hence $\partial_{14*}^{5,2}(\nu_5\sigma_8) \neq 0$, i.e., one of x_2, y_2, z_2, w_2 must be nonzero in (36). So

$$\text{Coker}(\partial_{14*}^{5,2}) \cong \mathbb{Z}_4^{\oplus 3} \oplus \mathbb{Z}_2^{\oplus 4}$$

which is obtained by reducing one of \mathbb{Z}_4 -summands of $\pi_{14}(F_5^2) \cong \pi_{14}(J_{3,4}^2)$ to \mathbb{Z}_2 . Now from Lemma 5.9,

$$\pi_{14}(P^5(2^r)) \cong \begin{cases} \mathbb{Z}_4^{\oplus 3} \oplus \mathbb{Z}_2^{\oplus 7}, & r = 2; \\ \mathbb{Z}_{2^r} \oplus \mathbb{Z}_8^{\oplus 2} \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_2^{\oplus 6}, & r \geq 3. \end{cases}$$

5.3. Homotopy groups of $P^6(2^r)$

There are cofibration and fibration sequences

$$\begin{aligned} S^5 \xrightarrow{2^r t_5} S^5 \xrightarrow{i_5} P^6(2^r) \xrightarrow{p_6} S^6; \quad \Omega S^6 \xrightarrow{\partial^6} F_6 \xrightarrow{\tau_6} P^6(2^r) \xrightarrow{p_6} S^6 \\ Sk_{14}(F_6) \simeq J_{2,5}^r \simeq S^5 \vee S^{10} (r \geq 1); \quad Sk_{19}(F_6) \simeq J_{3,5}^r \simeq J_{2,5}^r \cup_{\gamma_3^r} CS^{14}. \end{aligned}$$

Here $\beta_5 = j_1^5, \beta_{10} = j_2^{10}$.

Lemma 5.11. For $r \geq 1$, $\gamma_3^r = 2^r a_0 [j_1^5, j_2^{10}] \in \pi_{14}(J_{2,5}^r)$, where a_0 is an odd integer in $\mathbb{Z}_{(2)}$.

Proof. $J_{2,5}^0 \simeq S^5 \cup_{[t_5, t_5]} e^{10} \simeq S^5 \cup_{\nu_5 \eta_8} e^{10}$. It is easy to see from (7) that

$$\begin{aligned} \pi_{14}(J_{2,5}^0) &\cong \pi_{14}(F_{\bar{p}_{10}}^0) \cong \pi_{14}(S^5 \vee S^{14}) \\ \Rightarrow \pi_{14}(J_{2,5}^0) &= \mathbb{Z}_2\{\beta_5^0 \nu_5^3\} \oplus \mathbb{Z}_2\{\beta_5^0 \mu_5\} \oplus \mathbb{Z}_2\{\beta_5^0 \eta_5 \varepsilon_6\} \oplus \mathbb{Z}_{(2)}\{\beta_{14}^0\}. \end{aligned}$$

Assume $\gamma_3^0 = \beta_5^0 \theta_0 + a_0 \beta_{14}^0$, $\theta_0 \in \pi_{14}(S^5)$, $a_0 \in \mathbb{Z}_{(2)}$.

$$\text{By } \mathbb{Z}_2^{\oplus 3} \cong \pi_{15}(S^6) \xrightarrow[\cong]{\partial_{14*}^{6,0}} \pi_{14}(F_6^0) \cong \pi_{14}(J_{3,5}^0) \cong \pi_{14}(J_{2,5}^0) / \langle \gamma_3^0 \rangle$$

$\Rightarrow a_0$ is an odd integer in $\mathbb{Z}_{(2)}$.

From lemma 4.4, it is easy to get

$$\gamma_3^r = a_r [j_1^5, j_2^{10}], a_r \in \mathbb{Z}$$

Since the composition of the maps $S^5 \vee S^{14} \simeq J_{2,5}^{\gamma,r} \hookrightarrow F_{\bar{p}_{10}^r} \xrightarrow{\bar{\tau}_{10}^r} J_{2,5}^r \simeq S^5 \vee S^{10}$ induces the isomorphism $\pi_{14}(S^5 \vee S^{14}) \xrightarrow{\cong} \pi_{14}(S^5 \vee S^{10})$. We get

$$\bar{\tau}_{10}^r \bar{\beta}_{14}^r = \pm [j_1^5, j_2^{10}] + j_1^5 \theta_1, \theta_1 \in \pi_{14}(S^5).$$

$$\begin{aligned} 2^{2r} \gamma_3^0 &= \bar{g}_0^r \gamma_3^r = \pm a_r \bar{g}_0^r (\bar{\tau}_{10}^r \bar{\beta}_{14}^r - j_1^5 \theta_1) = \pm a_r \bar{g}_0^r (\bar{\tau}_{10}^r \bar{\beta}_{14}^r - \bar{\tau}_{10}^r \bar{\beta}_5^r \theta_1) \quad (\text{by (12)}) \\ &= \pm a_r \bar{\tau}_{10}^0 g_0^r (\bar{\beta}_{14}^r - \bar{\beta}_5^r \theta_1) = \pm 2^r a_r \bar{\tau}_{10}^0 \bar{\beta}_{14}^0 + \bar{\tau}_{10}^0 \bar{\beta}_5^0 \theta', \quad (\text{by (14) and (15)}) \\ &= \pm 2^r a_r \beta_{14}^0 + \beta_5^0 \theta', \quad \theta' \in \pi_{14}(S^5) \\ \text{i.e., } 2^{2r} (a_0 \beta_{14}^0 + \beta_5^0 \theta_0) &= \pm 2^r a_r \beta_{14}^0 + \beta_5^0 \theta' \Rightarrow a_r = \pm 2^r a_0. \end{aligned}$$

Thus we obtain this lemma by replacing $\pm a_0$ by a_0 in equation $a_r = \pm 2^r a_0$. \square

Lemma 2.7 gives the following commutative diagram

$$\begin{array}{ccccc} \pi_{m+1}(S^6) & \xrightarrow{\partial_{m*}^{6,r}} & \pi_m(F_6^r) & \xleftarrow{\bar{I}_{2*}^r} & \pi_m(S^5 \vee S^{10}) & \xrightarrow{Proj} & \pi_m(S^{10}) & (37) \\ H_2 \downarrow & & H_2' \downarrow & & & & \Sigma_* \downarrow & \\ \pi_{m+1}(S^{11}) & \xrightarrow{(2^r \iota_{11})_*} & \pi_{m+1}(S^{11}) & \xlongequal{\quad\quad\quad} & \pi_{m+1}(S^{11}) & & \pi_{m+1}(S^{11}) \end{array}$$

where \bar{I}_{2*}^r is isomorphic for $m \leq 13$.

Lemma 5.12. $\pi_i(F_6) \cong \pi_i(J_{3,5}^r), (i \leq 18)$ and

- (1) $\pi_{14}(J_{3,5}^r) = \mathbb{Z}_2\{\check{\beta}_5 \nu_5^3\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \mu_5\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \eta_5 \varepsilon_6\} \oplus \mathbb{Z}_{2^r}\{[\check{\beta}_5, \check{\beta}_{10}]\}$.
- (2) $\pi_{15}(J_{3,5}^r) = \mathbb{Z}_8\{\check{\beta}_5 \nu_5 \sigma_8\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \eta_5 \mu_6\} \oplus \mathbb{Z}_2\{[\check{\beta}_5, \check{\beta}_{10}] \eta_{14}\}$.
- (3) $\pi_{16}(J_{3,5}^1) = \mathbb{Z}_8\{\check{\beta}_5 \zeta_5\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \bar{\nu}_8\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \varepsilon_8\} \oplus \mathbb{Z}_2\{\check{\beta}_{10} \nu_{10}^2\} \oplus \mathbb{Z}_4\{\widehat{\eta}_{15}\},$
with $[\check{\beta}_5, \check{\beta}_{10}] \eta_{14}^2 = 2\widehat{\eta}_{15}$.
- $\pi_{16}(J_{3,5}^r) = \mathbb{Z}_8\{\check{\beta}_5 \zeta_5\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \bar{\nu}_8\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \varepsilon_8\} \oplus \mathbb{Z}_2\{\check{\beta}_{10} \nu_{10}^2\}$
 $\oplus \mathbb{Z}_2\{[\check{\beta}_5, \check{\beta}_{10}] \eta_{14}^2\} \oplus \mathbb{Z}_2\{\widehat{\eta}_{15}\}, (r \geq 2).$
- (4) $\pi_{17}(J_{3,5}^r) = \mathbb{Z}_2\{\check{\beta}_5 \nu_5^4\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \bar{\mu}_8\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \eta_8 \varepsilon_9\} \oplus \mathbb{Z}_{16}\{\check{\beta}_{10} \sigma_{10}\}$
 $\oplus \mathbb{Z}_{2m^3}\{[\check{\beta}_5, \check{\beta}_{10}] \nu_{14}\} \oplus \mathbb{Z}_2\{\widehat{\eta}_{15} \eta_{16}\}.$
- (5) $\pi_{18}(J_{3,5}^r) = \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \sigma_8 \nu_{15}\} \oplus \mathbb{Z}_2\{\check{\beta}_5 \nu_5 \eta_8 \mu_9\} \oplus \mathbb{Z}_2\{\check{\beta}_{10} \bar{\nu}_{10}\} \oplus \mathbb{Z}_2\{\check{\beta}_{10} \varepsilon_{10}\}$
 $\oplus \mathbb{Z}_{2^r}\{[\check{\beta}_5, [\check{\beta}_5, \check{\beta}_{10}]]\} \oplus \mathbb{Z}_{2m^3}\{\widehat{\delta}_r \nu_{15}\}.$

Proof. (1) of this lemma is easily obtained by Lemma 5.11.

Consider the Lemma 4.3 for $k = 5$.

(2) is obtained by the isomorphism $\pi_{15}(S^5 \vee S^{10}) \xrightarrow[\cong]{I_{2*}^r} \pi_{15}(J_{3,5}^r).$

For (3), we have the short exact sequence

$$0 \rightarrow \pi_{16}(S^5 \vee S^{10}) \xrightarrow{I_{2*}^r} \pi_{16}(J_{3,5}^r) \xrightarrow{\check{p}_{15*}} \pi_{16}(S^{15}) = \mathbb{Z}_2\{\eta_{15}\} \rightarrow 0,$$

$$\pi_{16}(S^5 \vee S^{10}) = \mathbb{Z}_8\{j_1^5 \zeta_5\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \bar{\nu}_8\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \varepsilon_8\} \oplus \mathbb{Z}_2\{j_2^{10} \nu_{10}^2\} \oplus \mathbb{Z}_2\{[j_1^5, j_2^{10}] \eta_{14}^2\}.$$

By Lemma 3.8, there is an $-\mathbf{x}_r \in \{I_2^r, \gamma_3^r, \eta_{14}\} \subset \pi_{16}(J_{3,5}^r)$, such that $\check{p}_{15*}(\mathbf{x}_r) = \eta_{15}$ and $2\mathbf{x}_r \in I_2^r\{2^r a_0[j_1^5, j_2^{10}], \eta_{14}, 2\iota_{15}\}$. By Lemma 3.9

$$\{2^r a_0[j_1^5, j_2^{10}], \eta_{14}, 2\iota_{15}\} \ni 2^{r-1}[j_1^5, j_2^{10}] \eta_{14}^2$$

$$\text{mod } (2^r a_0[j_1^5, j_2^{10}] \circ \pi_{16}(S^{14}) + \pi_{16}(S^5 \vee S^{10}) \circ (2\iota_{16})) = \langle 2j_1^5 \zeta_5 \rangle.$$

For $r = 1$, $2\mathbf{x}_1 = [\check{\beta}_5, \check{\beta}_{10}] \eta_{14}^2 + 2u_1 \check{\beta}_5 \zeta_5, u_1 \in \mathbb{Z}_4$. Let $\mathbf{x}'_1 = \mathbf{x}_1 - u_1 \check{\beta}_5 \zeta_5$, then $o(\mathbf{x}'_1) = 4$ and $\check{p}_{15*}(\mathbf{x}'_1) = \eta_{15}$ with $2\mathbf{x}'_1 = [\check{\beta}_5, \check{\beta}_{10}] \eta_{14}^2$. Thus we get $\pi_{16}(J_{3,5}^1)$.

For $r \geq 2$, $2\mathbf{x}_r = 2u_2 \check{\beta}_5 \zeta_5, u_2 \in \mathbb{Z}_4$. Let $\mathbf{x}'_r = \mathbf{x}_r - u_2 \check{\beta}_5 \zeta_5$, then $o(\mathbf{x}'_r) = 2$ and $\check{p}_{15*}(\mathbf{x}'_r) = \eta_{15}$. Thus $\pi_{16}(J_{3,5}^r) \xrightarrow{\check{p}_{15*}} \mathbb{Z}_2\{\eta_{15}\}$ is split onto. Hence we get $\pi_{16}(J_{3,5}^r), r \geq 2$ in (3) of this lemma.

For (4), we have the exact sequence

$$0 \rightarrow \frac{\pi_{17}(S^5 \vee S^{10})}{\langle \gamma_3^r \nu_{14} \rangle} \xrightarrow{I_{2*}^r} \pi_{17}(J_{3,5}^r) \xrightarrow{\check{p}_{15*}} \pi_{17}(S^{15}) = \mathbb{Z}_2\{\eta_{15}^2\} \rightarrow 0,$$

$$\frac{\pi_{17}(S^5 \vee S^{10})}{\langle \gamma_3^r \nu_{14} \rangle} = \mathbb{Z}_2\{j_1^5 \nu_5^4\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \mu_8\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \eta_8 \varepsilon_9\} \oplus \mathbb{Z}_{16}\{j_1^{10} \sigma_{10}\} \oplus \mathbb{Z}_{2m_r^3}\{[j_1^5, j_2^{10}] \nu_{14}\}.$$

Note that the order 2 element $\widehat{\eta}_{15} \eta_{16}$ satisfies $\check{p}_{15*}(\widehat{\eta}_{15} \eta_{16}) = \eta_{15}^2$. Hence $\pi_{17}(J_{3,5}^r) \xrightarrow{\check{p}_{15*}} \mathbb{Z}_2\{\eta_{15}^2\}$ is split onto. We get the $\pi_{17}(J_{3,5}^r)$ of this lemma.

For (5), we have the exact sequence

$$0 \rightarrow \frac{\pi_{18}(S^5 \vee S^{10})}{\langle 2^r a_0[j_1^5, [j_1^5, j_2^{10}]] \rangle} \xrightarrow{I_{2*}^r} \pi_{18}(J_{3,5}^r) \xrightarrow{\check{p}_{15*}} \mathbb{Z}_{2m_r^3}\{\delta_r \nu_{15}\} \rightarrow 0, \quad \frac{\pi_{18}(S^5 \vee S^{10})}{\langle 2^r a_0[j_1^5, [j_1^5, j_2^{10}]] \rangle} =$$

$$\mathbb{Z}_2\{j_1^5 \nu_5 \sigma_8 \nu_{15}\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \eta_8 \mu_9\} \oplus \mathbb{Z}_2\{j_2^{10} \bar{\nu}_{10}\} \oplus \mathbb{Z}_2\{j_2^{10} \varepsilon_{10}\} \oplus \mathbb{Z}_{2^r}\{[j_1^5, [j_1^5, j_2^{10}]]\}.$$

There is $-\mathbf{x}_r \in \{I_2^r, \gamma_3^r, \delta_r \nu_{14}\} \subset \pi_{18}(J_{3,5}^r)$, such that $\check{p}_{15*}(\mathbf{x}_r) = \delta_r \nu_{15}$ and $2^{m_r^3} \mathbf{x}_r \in I_2^r\{\gamma_3^r, \delta_r \nu_{14}, 2^{m_r^3} \iota_{17}\}$. $\{\gamma_3^r, \delta_r \nu_{14}, 2^{m_r^3} \iota_{17}\}$ is a coset of subgroup $\gamma_3^r \circ \pi_{18}(S^{14}) + \pi_{18}(S^5 \vee S^{10}) \circ (2^{m_r^3} \iota_{18}) = 2^{m_r^3} \pi_{18}(S^5 \vee S^{10})$. Moreover, by $\pi_{18}(S^{14}) = 0$, $\{\gamma_3^r, \delta_r \nu_{14}, 2^{m_r^3} \iota_{17}\} \supset (a_0[j_1^5, j_2^{10}] \circ \{2^r \iota_{14}, \delta_r \nu_{14}, 2^{m_r^3} \iota_{17}\}) = \{0\}$. Thus $\{\gamma_3^r, \delta_r \nu_{14}, 2^{m_r^3} \iota_{17}\} = 2^{m_r^3} \pi_{18}(S^5 \vee S^{10})$ which implies $2^{m_r^3} \mathbf{x}_r = 2^{m_r^3} I_2^r \theta$ for some $\theta \in \pi_{18}(S^5 \vee S^{10})$. Take $\mathbf{x}'_r = \mathbf{x}_r - I_2^r \theta$, then $o(\mathbf{x}'_r) = 2^{m_r^3}$ and $\check{p}_{15*}(\mathbf{x}'_r) = \delta_r \nu_{15}$. Hence we finish the proof of (5) of this lemma. \square

• $\pi_{10}(P^6(2^r))$.

There is the exact sequence

$$\mathbb{Z}_{(2)}\{\Delta(\iota_{13})\} = \pi_{11}(S^6) \xrightarrow{\partial_{10*}^{6,r}} \pi_{10}(F_6) = \mathbb{Z}_2\{\beta_5 \nu_5 \eta_8^2\} \oplus \mathbb{Z}_{(2)}\{\beta_{10}\} \xrightarrow{\tau_{6*}} \pi_{10}(P^6(2^r)) \rightarrow 0.$$

Assume $\partial_{10*}^{6,r}(\Delta(\iota_{13})) = x_r \beta_5 \nu_5 \eta_8^2 + y_r \beta_{10}, x_r \in \mathbb{Z}_2, y_r \in \mathbb{Z}$.

By the right commutative square of (37) for $m = 10$, $H'_2(\beta_{10}) = \iota_{11}$. Since $H_2(\Delta(\iota_{13})) = \pm\iota_{11}$ ([39, Proposition 2.7]), by the left commutative square of (37) for $m = 10$, we get $H'_2\partial_{10*}^{6,r}(\Delta(\iota_{13})) = 2^r H_2(\Delta(\iota_{13})) = 2^{r+1}\iota_{11}$. This implies that $y_r = \pm 2^{r+1}$.

For $r = 1$, by [10, Theorem 2.2], there is an element in $\pi_{10}(P^6(2)) \cong \frac{\mathbb{Z}_2\{\beta_5\nu_5\eta_8^2\} \oplus \mathbb{Z}_{(2)}\{\beta_{10}\}}{\langle \pm 4\beta_{10} + x_1\beta_5\nu_5\eta_8^2 \rangle}$ with order 8. So $\pi_{10}(P^6(2)) \cong \mathbb{Z}_8$ and $x_1 = 1$.

For $r \geq 2$, consider the diagram (9) for $s = 1, t = r$, we get

$$\begin{aligned} \partial_{10*}^{6,r}(\Delta(\iota_{13})) &= \chi_{r*}^1 \partial_{10*}^{6,1}(\Delta(\iota_{13})) = \chi_{r*}^1(4\beta_{10}^1 + \beta_5^1\nu_5\eta_8^2) \\ &= 4\chi_r^1\beta_{10}^1 + 2^{r-1}\beta_5^r\nu_5\eta_8^2 = 4\chi_r^1\beta_{10}^1, \text{ by first equation of (12)}. \end{aligned}$$

This implies that $x_r = 0$ for $r \geq 2$. Hence

$$\pi_{10}(P^6(2^r)) \cong \frac{\mathbb{Z}_2\{\beta_5\nu_5\eta_8^2\} \oplus \mathbb{Z}_{(2)}\{\beta_{10}\}}{\langle \pm 2^{r+1}\beta_{10} \rangle} \cong \mathbb{Z}_2 \oplus \mathbb{Z}_{2^{r+1}}, r \geq 2.$$

- $\pi_{11}(P^6(2^r))$.

Consider diagram (6) for $k = 5, m = 11, r \geq 1$, we get $\partial_{10*}^{6,r}(\nu_6^2) = 0$. Since $\partial_{10*}^{6,r}$ is a monomorphism, $\text{Ker}(\partial_{10*}^{6,r}) = 0$. So we have

$$\pi_{11}(P^6(2^r)) \cong \pi_{11}(F_6) \cong \pi_{11}(S^5 \vee S^{10}) \cong \mathbb{Z}_2^{\oplus 2}.$$

- $\pi_{12}(P^6(2^r))$.

There is the exact sequence

$$\mathbb{Z}_4\{\sigma''\} = \pi_{13}(S^6) \xrightarrow{\partial_{12*}^{6,r}} \pi_{12}(F_6) \xrightarrow{\tau_{6*}} \pi_{12}(P^6(2^r)) \xrightarrow{p_{6*}} \mathbb{Z}_2\{\nu_6^2\} \rightarrow 0.$$

$$\text{Assume } \partial_{12*}^{6,r}(\sigma'') = x_r\beta_5\sigma''' + y_r\beta_{10}\eta_{10}^2 \in \pi_{12}(F_6) = \mathbb{Z}_2\{\beta_5\sigma'''\} \oplus \mathbb{Z}_2\{\beta_{10}\eta_{10}^2\}.$$

Consider the commutative diagram (37) for $m = 12, r \geq 1$. We get

$$H'_2\partial_{12*}^{6,r}(\sigma'') = H'_2(x_r\beta_5\sigma''' + y_r\beta_{10}\eta_{10}^2) = y_r\eta_{11}^2 = 0.$$

This implies that $y_r = 0$.

For $r = 1$,

$$\chi_{0*}^1 \partial_{12*}^{6,1}(\sigma'') = \partial_{12*}^{6,0}(2\iota_6)_*(\sigma'') = \partial_{12*}^{6,0}(2\sigma'') = 2\partial_{12*}^{6,0}(\sigma'') \neq 0$$

where the first equation is from the homotopy commutative diagrams (9) for $s = 1, r = 0, k = 5$; the second equation holds since $(2\iota_6)\sigma'' = 2\sigma'' \pm [\iota_6, \iota_6]H_2(\sigma'') = 2\sigma''$ ($[\iota_6, \iota_6]H_2(\sigma'') = 0$ because $\Sigma([\iota_6, \iota_6]H_2(\sigma'')) = 0$ and $\pi_{13}(S^6) \xrightarrow{\Sigma} \pi_{14}(S^7)$ is injective);

the last inequality holds since $\pi_{13}(S^6) \xrightarrow{\partial_{12*}^{6,0}} \pi_{12}(F_6^0) \cong \mathbb{Z}_4$ is isomorphic. This implies $x_1 = 0$, i.e.,

$$\partial_{12*}^{6,1}(\sigma'') = \beta_5^1\sigma'''.$$

For $r \geq 2$,

$$\partial_{12*}^{6,r}(\sigma'') = \chi_{r*}^1 \partial_{12*}^{6,1}(\sigma'') = \chi_r^1 \beta_5^1 \sigma''' = (2^{r-1} \beta_5^r) \sigma''' = \beta_5^r (2^{r-1} \iota_5) \sigma''' = \beta_5^r (2^{r-1} \sigma''') = 0$$

where the first equation is induced by (9); the third equation is from the first equation of (12); the last equation holds since $(2^{r-1} \iota_5) \sigma''' = 2^{r-1} \sigma''' \pm \binom{2^{r-1}}{2} [\iota_5, \iota_5] H_2(\sigma''') = 2^{r-1} \sigma''' \pm \binom{2^{r-1}}{2} (\nu_5 \eta_8) (4\nu_4) = 2^{r-1} \sigma'''$ by [33, Proposition 2.10] and [39, Lemma 5.13].

Now we have the following short exact sequence

$$0 \rightarrow \vartheta_r \mathbb{Z}_2 \oplus \mathbb{Z}_2 \rightarrow \pi_{12}(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_2\{\nu_6^2\} \rightarrow 0.$$

since short exact sequence (31) splits, so is the above sequence by Lemma 3.6. Hence

$$\pi_{12}(P^6(2^r)) \cong \vartheta_r \mathbb{Z}_2 \oplus \mathbb{Z}_2^{\oplus 2}.$$

- $\pi_{13}(P^6(2^r))$.

We have the following exact sequence

$$\mathbb{Z}_8\{\bar{\nu}_6\} \oplus \mathbb{Z}_2\{\varepsilon_6\} = \pi_{14}(S^6) \xrightarrow{\partial_{13*}^{6,r}} \pi_{13}(F_6) \xrightarrow{\tau_{6*}} \pi_{13}(P^6(2^r)) \xrightarrow{P_{6*}} \text{Ker}(\partial_{12*}^{6,r}) \rightarrow 0.$$

$$\text{Ker}(\partial_{12*}^{6,1}) = \mathbb{Z}_2\{2\sigma''\}; \quad \text{Ker}(\partial_{12*}^{6,r}) = \mathbb{Z}_4\{\sigma''\} r \geq 2.$$

$$\pi_{13}(F_6) = \mathbb{Z}_2\{\beta_5 \varepsilon_5\} \oplus \mathbb{Z}_8\{\beta_{10} \nu_{10}\}; \quad \partial_{13*}^{6,r}(\varepsilon_6) = \beta_{5*} (2^r \iota_5)_*(\varepsilon_5) = 0.$$

$$\text{Assume } \partial_{13*}^{6,r}(\bar{\nu}_6) = x_r \beta_5 \varepsilon_5 + y_r \beta_{10} \nu_{10}, x_r \in \mathbb{Z}_2, y_r \in \mathbb{Z}_8.$$

Consider the commutative diagram (37) for $m = 13, r \geq 1$. By $H_2(\bar{\nu}_6) = l\nu_{11}, l$ is odd [39, Lemma 6.2], we get

$$H_2' \partial_{13*}^{6,r}(\bar{\nu}_6) = H_2'(x_r \beta_5 \varepsilon_5 + y_r \beta_{10} \nu_{10}) = y_r \nu_{11} = 2^r l \nu_{11}.$$

This implies that $y_r = 2^r l$.

From left commutative square of Corollary 3.4 for $m = n = 5, u = 13$,

$$\begin{array}{ccc} \mathbb{Z}_8\{\bar{\nu}_6\} \oplus \mathbb{Z}_2\{\varepsilon_6\} = \pi_{14}(S^6) & \xrightarrow{\partial_{13*}^{6,r}} & \pi_{13}(F_5) \\ \Sigma^\infty \downarrow & & \downarrow E^\infty \\ \mathbb{Z}_2\{\bar{\nu}\} \oplus \mathbb{Z}_2\{\varepsilon\} = \pi_{13}^s(S^5) & \xrightarrow{(2^r \iota)_* = 0} & \pi_{13}^s(S^5) \end{array}$$

$$\begin{aligned} \Rightarrow 0 &= E^\infty \partial_{13*}^{6,r}(\bar{\nu}_6) = E^\infty(x_r \beta_5 \varepsilon_5 + 2^r l \beta_{10} \nu_{10}) = x_r E^\infty(\beta_5 \varepsilon_5) + 2^r l E^\infty(\beta_{10} \nu_{10}) \\ &= x_r \varepsilon, \text{ by right commutative triangle of Corollary 3.4.} \end{aligned}$$

So $x_r = 0 \Rightarrow \text{Coker}(\partial_{13*}^{6,r}) = \mathbb{Z}_2\{\beta_5 \varepsilon_5\} \oplus \mathbb{Z}_{2m^3}\{\beta_{10} \nu_{10}\}$.

From (19) we have the following commutative diagram

$$\begin{array}{ccc}
\mathbb{Z}_2\{\beta_5\varepsilon_5\} \oplus \mathbb{Z}_{2m_r^3}\{\beta_{10}\nu_{10}\} = \text{Coker}\partial_{13*}^{6,r} & \xrightarrow{\tau_{6*}} & \pi_{13}(P^6(2^r)) \\
\downarrow \bar{H}_{2*} & & \downarrow \bar{H}_{2*} \\
\text{Coker}(\partial_{13*}^{11,r}) \cong \mathbb{Z}_{2m_r^3}\{\nu_{10}\} & \xrightarrow{i_{10*}} & \pi_{13}(P^{11}(2^r))
\end{array} \quad (38)$$

Lemma 5.13. *In the diagram (38), $\bar{H}_2(\beta_{10}\nu_{10}) = t\nu_{10}$, t is odd.*

Proof. By the same proof as that of Lemma 5.3, we get

$$\bar{h}_w(\Omega_0(\tau_6\beta_{10})) = [u, v] \in H_9(\Omega P^6(2^r), \mathbb{Z}_2) \quad (39)$$

where \bar{h}_w is the mod 2-reduction of Hurewicz homomorphism, i.e., the composition $\pi_9(\Omega P^6(2^r)) \xrightarrow{h_w} H_9(\Omega P^6(2^r)) \xrightarrow{\text{reduce}} H_9(\Omega P^6(2^r); \mathbb{Z}_2) \subset T(u, v)$, where $T(u, v)$ the tensor algebra over \mathbb{Z}_2 generated by u, v with degrees 4 and 5 respectively.

$\bar{H}_{2*} : H_*(\Omega P^6(2^r), \mathbb{Z}_2) \xrightarrow{H_{2*}} H_*(\Omega \Sigma P^5(2^r) \wedge P^5(2^r), \mathbb{Z}_2) \xrightarrow{(\Omega \Sigma q)_*} H_*(\Omega P^{11}(2^r), \mathbb{Z}_2)$. where $H_*(\Omega P^{11}(2^r), \mathbb{Z}_2) = T(u', v')$ and the degrees of u' and v' are 9 and 10 respectively; q is the pinch map $P^5(2^r) \wedge P^5(2^r) \rightarrow P^5(2^r) \wedge P^5(2^r)/(S^4 \wedge P^5(2^r)) = P^{10}(2^r)$. By [9, Proposition 5.3] we can get $\bar{H}_{2*}([u, v]) = u'$ and $\bar{H}_{2*}(v \otimes v) = v'$. Hence $\bar{H}_{2*}\bar{h}_w(\Omega_0(\tau_6\beta_{10})) = u'$. This implies the following composition is homotopy equivalent to $t\Omega_0(i_{10}) : S^9 \rightarrow \Omega P^{10}(2^r)$ with odd t

$$S^9 \xrightarrow{\Omega \Sigma} \Omega S^{10} \xrightarrow{\Omega \beta_{10}} \Omega F_6 \xrightarrow{\Omega \tau_6} \Omega P^6(2^r) \xrightarrow{\bar{H}_2} \Omega P^{10}(2^r) .$$

$$\xrightarrow[\Omega_0(\tau_6\beta_{10})]{\Omega_0(\tau_6\beta_{10})}$$

$$\begin{aligned}
i_{10*}\bar{H}_{2*}(\beta_{10}\nu_{10}) &= \bar{H}_{2*}(\tau_6\beta_{10}\nu_{10}) \quad (\text{by (37)}) \\
&= \Omega_0^{-1}(\bar{H}_2\Omega_0(\tau_6\beta_{10}\nu_{10})) \quad (\text{by definition of right map } \bar{H}_{2*} \text{ in (37)}) \\
&= \Omega_0^{-1}(\bar{H}_2\Omega(\tau_6\beta_{10}\nu_{10})E_{S^{12}}) = \Omega_0^{-1}(\bar{H}_2\Omega(\tau_6\beta_{10})(\Omega\nu_{10})E_{S^{12}}) \\
&= \Omega_0^{-1}(\bar{H}_2\Omega(\tau_6\beta_{10})E_{S^9\nu_9}) = \Omega_0^{-1}(\bar{H}_2\Omega_0(\tau_6\beta_{10})\nu_9) \\
&= \Omega_0^{-1}(t\Omega_0(i_{10})\nu_9) = \Omega_0^{-1}(t\Omega(i_{10})E_{S^9\nu_9}) = \Omega_0^{-1}(t\Omega(i_{10})\Omega(\nu_{10})E_{S^{12}}) \\
&= \Omega_0^{-1}(t\Omega(i_{10}\nu_{10})E_{S^{12}}) = \Omega_0^{-1}(t\Omega_0(i_{10}\nu_{10})) = i_{10*}(t\nu_{10}). \\
&\Rightarrow \bar{H}_{2*}(\beta_{10}\nu_{10}) = t\nu_{10}, \text{ since } i_{10*} \text{ is an injection.}
\end{aligned}$$

□

From above lemma, the composition $\mathbb{Z}_{2m_r^3}\{\beta_{10}\nu_{10}\} \xrightarrow{j_2} \text{Coker}\partial_{13*}^{6,r} \xrightarrow{\bar{H}_{2*}} \mathbb{Z}_{2m_r^3}\{\nu_{10}\}$ is isomorphic. Since the bottom injection $i_{10*} : \mathbb{Z}_{2m_r^3}\{\nu_{10}\} \rightarrow \pi_{13}(P^{11}(2^r))$ of (37) is split into. So is the map $\mathbb{Z}_{2m_r^3}\{\beta_{10}\nu_{10}\} \xrightarrow{j_2} \text{Coker}\partial_{13*}^{6,r} \xrightarrow{\tau_{6*}} \pi_{13}(P^6(2^r))$.

For $r = 1$, $\text{Ker}(\partial_{12*}^{6,1}) = \mathbb{Z}_2\{2\sigma''\}$. By $2\sigma'' = \Sigma\sigma'''$ [39, Lemma 5.14], there is $-\mathbf{x}_1 \in \{i_5, 2\iota_5, \sigma'''\}$, such that $p_{6*}(\mathbf{x}_1) = 2\sigma''$. $2\mathbf{x}_1 \in -\{i_5, 2\iota_5, \sigma'''\}(2\iota_{13}) = i_5\{2\iota_5, \sigma''', 2\iota_{12}\} = \{0\}$ by Lemma 8.2 (v). Hence $2\mathbf{x}_1 = 0$ implies that $\pi_{13}(P^6(2)) \xrightarrow{P_{6*}} \mathbb{Z}_2\{2\sigma''\}$ is split onto. We get

$$\pi_{13}(P^6(2)) \cong \mathbb{Z}_2^{\oplus 3}.$$

By the diagram (9) for $s = r \geq 2, t = 1, k = 5$, we have the commutative diagram

$$\begin{array}{ccccc} \mathbb{Z}_2\{\beta_5^r \varepsilon_5\} & \xrightarrow{j_1} & \pi_{13}(F_6^r) & \xrightarrow{\tau_{6*}^r} & \pi_{13}(P^6(2^r)) \\ \parallel & & \downarrow \chi_{1*}^r & & \downarrow \bar{\chi}_{1*}^r \\ \mathbb{Z}_2\{\beta_5^1 \varepsilon_5\} & \xrightarrow{j_1} & \pi_{13}(F_6^1) & \xrightarrow{\tau_{6*}^1} & \pi_{13}(P^6(2)) \end{array}$$

The composition of the bottom maps $\tau_{6*}^1 j_1$ is split into, so is the composition of the top maps $\tau_{6*}^r j_1$. Hence

$$\pi_{13}(P^6(2^r)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_{2m_r^3} \oplus \mathbb{Z}_4(r \geq 2).$$

- $\pi_{14}(P^6(2^r))$.

Commutative diagram (6) for $k = 5, m = 14, r \geq 1$ gives the following short exact sequence

$$0 \rightarrow \pi_{14}(J_{3,5}^r) \cong \pi_{14}(F_6) \xrightarrow{\tau_{6*}} \pi_{14}(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_{2m_r^3}\{\delta_r \bar{\nu}_6\} \oplus \mathbb{Z}_2\{\varepsilon_6\} \rightarrow 0.$$

Consider the commutative diagram of exact sequences

$$\begin{array}{ccccccc} \pi_{15}(S^6) & \xrightarrow{\partial_{14*}^{6,r}} & \pi_{14}(F_6^r) & \xrightarrow{\tau_{6*}} & \pi_{14}(P^6(2^r)) & \xrightarrow{P_{6*}} & \mathbb{Z}_{2m_r^3}\{\delta_r \bar{\nu}_6\} \oplus \mathbb{Z}_2\{\varepsilon_6\} \longrightarrow 0 \\ \Sigma^\infty \downarrow & & \downarrow E^\infty & & \Sigma^\infty \downarrow & & \downarrow \Sigma^\infty \\ \pi_{14}^s(S^5) & \xrightarrow{(2^r \iota)_* = 0} & \pi_{14}^s(S^5) & \xrightarrow{i_*} & \pi_{14}^s(P^6(2^r)) & \xrightarrow{P_*} & \mathbb{Z}_2\{\bar{\nu}\} \oplus \mathbb{Z}_2\{\varepsilon\} \longrightarrow 0 \end{array}$$

There is an $\alpha_r \in \pi_{14}(P^6(2^r))$, such that $p_{6*}(\alpha_r) = \delta_r \bar{\nu}_6$.

For $r = 1$, $p_* \Sigma^\infty(\alpha_1) = \Sigma^\infty p_{6*}(\alpha_1) = \Sigma^\infty(4\bar{\nu}_6) = 4\Sigma^\infty(\bar{\nu}_6) = 4\bar{\nu} = 0$. Hence $\Sigma^\infty(\alpha_1) \in i_*(\pi_{14}^s(S^5))$, which implies that $\Sigma^\infty(2\alpha_1) = 0$. For $r \geq 2$, $\Sigma^\infty(2^{m_r^3} \alpha_r) = 2^{m_r^3} \Sigma^\infty(\alpha_r) = 0$ since $4\pi_{14}^s(P^6(2^r)) = 0$.

From Corollary 3.4, we get

$$2^{m_r^3} \alpha_r = b_r \tau_6 \check{I}_2[\check{\beta}_5, \check{\beta}_{10}] = b_r [i_5, \tau_6 \beta_{10}], b_r \in \mathbb{Z}_{2^r}.$$

By the same proof as that of Lemma 5.3, we get $2^s [i_5, \tau_6 \beta_{10}]$ is not divisible by 2^{s+1} in $\pi_{14}(P^6(2^r))$ for any $0 \leq s \leq r - 1$. Hence $b_r = 0 \in \mathbb{Z}_{2^r}$. This implies that the order of α_r is $2^{m_r^3}$.

There is also an element $-\mathbf{x}_r^\varepsilon \in \{i_5, 2^r \iota_5, \varepsilon_5\} \subset \pi_{14}(P^6(2^r))$, such that $p_{6*}(\mathbf{x}_r^\varepsilon) = \varepsilon_6$. $2\mathbf{x}_r^\varepsilon \in i_{5\circ}\{2^r \iota_5, \varepsilon_5, 2\iota_{13}\}$. By Lemma 8.2 (vi),

$$\begin{aligned} \text{for } r = 1, \quad 2\mathbf{x}_1^\varepsilon = i_5\eta_6\varepsilon_5 &\Rightarrow \pi_{14}(P^6(2)) \cong \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_4; \\ \text{for } r \geq 2, \quad 2\mathbf{x}_r^\varepsilon = 0 &\Rightarrow \pi_{14}(P^6(2^r)) \cong \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_{2^r} \oplus \mathbb{Z}_{2m^3}. \end{aligned}$$

- $\pi_{15}(P^6(2^r))$.

Consider diagram (6) for $k = 5, m = 15, r \geq 1$. $\partial_{15*}^{6,r}(v_6\sigma_9) = \beta_5(2^r \iota_5)(v_5\sigma_8) = \beta_5 v_5(2^r \iota_8)\sigma_8 = \beta_5 v_5(2^{2r}\sigma_8 - 2^{r-1}(2^r - 1)\Sigma\sigma) = 2^r \beta_5 v_5 \sigma_8$. $\partial_{15*}^{6,r}(\eta_6\mu_7) = \beta_5(2^r \iota_6)(\eta_5\mu_6) = 0$. There is the short exact sequence

$$\begin{aligned} 0 \rightarrow \text{Coker}(\partial_{15*}^{6,r}) \xrightarrow{\tau_{6*}} \pi_{15}(P^6(2^r)) \xrightarrow{P_{6*}} \pi_{15}(S^6) = \mathbb{Z}_2\{v_6^3\} \oplus \mathbb{Z}_2\{\mu_6\} \oplus \mathbb{Z}_2\{\eta_6\varepsilon_7\} \rightarrow 0, \\ \text{Coker}(\partial_{15*}^{6,r}) = \mathbb{Z}_{2m^3}\{\beta_5 v_5 \sigma_8\} \oplus \mathbb{Z}_2\{\beta_5 \eta_5 \mu_6\} \oplus \mathbb{Z}_2\{[\beta_5, \beta_{10}]\eta_{14}\}. \end{aligned}$$

Let $x_1 = v_5^3; x_2 = \mu_5; x_3 = \eta_5\varepsilon_6$. For $i = 1, 2, 3$, there is an $-\mathbf{x}_r^i \in \{i_5, 2^r \iota_5, x_i\}$ such that $p_{6*}(-\mathbf{x}_r^i) = \Sigma x_i$ and $2\mathbf{x}_r^i \in i_{5\circ}\{2^r \iota_5, x_i, 2\iota_{14}\}$. By Lemma 3.9,

$$\begin{aligned} \{2^r \iota_5, x_i, 2\iota_{14}\} \ni 2^{r-1}x_i\eta_{14} \bmod (2^r \iota_5) \circ \pi_{15}(S^5) + \pi_{15}(S^5) \circ (2\iota_{15}) &= \langle 2v_5\sigma_8 \rangle. \\ x_1\eta_{14} = v_5^2(v_{11}\eta_{14}) = 0, &\Rightarrow 2\mathbf{x}_r^1 = 2a_1i_5v_5\sigma_8, \\ x_2\eta_{14} = \mu_5\eta_{14} = \eta_5\mu_6 \text{ [39, page 66]}, &\Rightarrow 2\mathbf{x}_r^2 = 2^{r-1}i_5\eta_5\mu_6 + 2a_2i_5v_5\sigma_8, \\ x_3\eta_{14} = \eta_5\varepsilon_6\eta_{14} = \eta_5^2\varepsilon_7 = 2\Sigma^2\varepsilon' = 4v_5\sigma_8 \text{ [39, page 67, (7.5),(7.10)]}, & \\ \Rightarrow 2\mathbf{x}_r^3 = 2a_3i_5v_5\sigma_8, \text{ where } a_1, a_2, a_3 \in \mathbb{Z}. & \end{aligned}$$

Let $\mathbf{x}_r^i = \mathbf{x}_r^i - a_i i_5 v_5 \sigma_8 (i = 1, 2, 3)$. Then $p_{6*}(\mathbf{x}_r^i) = \Sigma x_i$ with $o(\mathbf{x}_r^i) = 2$ for $(r, i) \neq (1, 2)$ and $o(\mathbf{x}_2^1) = 4$. Thus

$$\pi_{15}(P^6(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2^{\oplus 5} \oplus \mathbb{Z}_{2m^3}, & r \geq 2. \end{cases}$$

- $\pi_{16}(P^6(2^r))$.

Consider diagram (6) for $k = 5, m = 16, r \geq 1$. By [33, Proposition 2.10] and $H_2(\zeta_5) = 8\sigma_9$ [39, Lemma 6.7]

$$\begin{aligned} \partial_{16*}^{6,r}(\zeta_6) = \beta_5(2^r \iota_5)(\zeta_5) = \beta_5(2^r \zeta_5 \pm \binom{2^r}{2}[\iota_5, \iota_5]H_2(\zeta_5)) &= 2^r \beta_5 \zeta_5; \\ \partial_{16*}^{6,r}(\bar{v}_6 v_{14}) = \partial_{13*}^{6,r}(\bar{v}_6) v_{13} = 2^r l v_{11} v_{13} = 0 \text{ (} l \text{ is odd)}. & \\ \Rightarrow 0 \rightarrow \text{Coker}(\partial_{16*}^{6,r}) \xrightarrow{\tau_{6*}} \pi_{16}(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_{2m^3}\{\delta_r v_6 \sigma_9\} \oplus \mathbb{Z}_2\{\eta_6 \mu_7\} \rightarrow 0, & \\ \text{Coker}(\partial_{16*}^{6,r}) = \mathbb{Z}_{2m^3}\{\beta_5 \zeta_5\} \oplus \mathbb{Z}_2\{\beta_5 v_5 \bar{v}_8\} \oplus \mathbb{Z}_2\{\beta_5 v_5 \varepsilon_8\} \oplus \mathbb{Z}_2\{\beta_{10} v_{10}^2\} \oplus A_r & \\ \text{where } A_1 = \mathbb{Z}_4\{\tilde{I}_3^1 \widehat{\eta}_{15}\}, \quad A_r = \mathbb{Z}_2\{[\beta_5, \beta_{10}]\eta_{14}^2\} \oplus \mathbb{Z}_2\{\tilde{I}_3^r \widehat{\eta}_{15}\}, r \geq 2. & \end{aligned}$$

By Lemma 3.2 of [23], $\pi_9(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_{2^{m_r^3}}\{\delta_r \nu_6\}$ is split onto, i.e. there is an order $2^{m_r^3}$ element $\widetilde{\delta_r \nu_6} \in \pi_9(P^6(2^r))$ such that $p_{6*}(\widetilde{\delta_r \nu_6}) = \delta_r \nu_6$. Then $\widetilde{\delta_r \nu_6} \sigma_9$ is the order $2^{m_r^3}$ element such that $p_{6*}(\widetilde{\delta_r \nu_6} \sigma_9) = \delta_r \nu_6 \sigma_9$. There is an element $\widetilde{\eta}_6 \in \pi_7(P^6(2^r))$ such that $p_{6*}(\widetilde{\eta}_6) = \eta_6$. Then the order 2 element $\widetilde{\eta}_6 \mu_7$ satisfies $p_{6*}(\widetilde{\eta}_6 \mu_7) = \eta_6 \mu_7$. So $\pi_{16}(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_{2^{m_r^3}}\{\delta_r \nu_6 \sigma_9\} \oplus \mathbb{Z}_2\{\eta_6 \mu_7\}$ is split onto. Hence

$$\pi_{16}(P^6(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 6} \oplus \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2^{\oplus 6} \oplus \mathbb{Z}_{2^{m_r^3}}^{\oplus 2}, & r \geq 2. \end{cases}$$

- $\pi_{17}(P^6(2^r))$.

By [39, Proposition 2.5], $\Delta(\sigma_{13}) = l\Delta(\iota_{13})\sigma_{11}$, l is odd. Hence by the previous calculation of $\partial_{10*}^{6,r}$ and Lemma 8.1

$$\partial_{17*}^{6,r}(\Delta(\sigma_{13})) = l\partial_{10*}^{6,r}(\Delta(\iota_{13}))\sigma_{10} = \begin{cases} \beta_5 \nu_5^4 + \beta_5 \nu_5 \eta_8 \varepsilon_9 \pm 4\beta_{10}\sigma_{10}, & r = 1; \\ \pm 2^{r+1}\beta_{10}\sigma_{10}, & r \geq 2. \end{cases} \quad \text{There is}$$

the short exact sequence

$$0 \rightarrow \text{Coker}(\partial_{17*}^{6,r}) \xrightarrow{\tau_{6*}} \pi_{17}(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_{2^{m_r^3}}\{\delta_r \zeta_6\} \oplus \mathbb{Z}_4\{\bar{\nu}_6 \nu_{14}\} \rightarrow 0,$$

$$\text{Coker}(\partial_{17*}^{6,1}) = \mathbb{Z}_2\{\beta_5 \nu_5 \bar{\mu}_8\} \oplus \mathbb{Z}_2\{\beta_5 \nu_5 \eta_8 \varepsilon_9\} \oplus \mathbb{Z}_8\{\beta_{10}\sigma_{10}\} \oplus \mathbb{Z}_2\{[\beta_5, \beta_{10}]\nu_{14}\} \oplus \mathbb{Z}_2\{\tilde{I}_3^1 \widetilde{\eta_{15}\eta_{16}}\}$$

$$\text{Coker}(\partial_{17*}^{6,r}) = \mathbb{Z}_2\{\beta_5 \nu_5^4\} \oplus \mathbb{Z}_2\{\beta_5 \nu_5 \bar{\mu}_8\} \oplus \mathbb{Z}_2\{\beta_5 \nu_5 \eta_8 \varepsilon_9\} \oplus \mathbb{Z}_{2^{m_{r+1}^4}}\{\beta_{10}\sigma_{10}\}$$

$$\oplus \mathbb{Z}_{2^{m_r^3}}\{[\beta_5, \beta_{10}]\nu_{14}\} \oplus \mathbb{Z}_2\{\tilde{I}_3^r \widetilde{\eta_{15}\eta_{16}}\}, r \geq 2.$$

There is $-\mathbf{x}_r \in \{i_5, 2^r \iota_5, \delta_r \zeta_5\}$ such that $2^{m_r^3} \mathbf{x}_r \in i_{5\circ}\{2^r \iota_5, \delta_r \zeta_5, 2^{m_r^3} \iota_{16}\}$ and $\{2^r \iota_5, \delta_r \zeta_5, 2^{m_r^3} \iota_{16}\}$ is a coset of $\{0\}$. For $r = 1$, by $4\zeta_5 = \eta_5^2 \mu_7 = \Sigma(\eta_4^2 \mu_6)$ [39, (7.14)], $\{2\iota_5, 4\zeta_5, 2\iota_{16}\} \ni \eta_5^2 \mu_7 \eta_{16} = \eta_5^3 \mu_8 = 4\nu_5 \mu_8 = 0$ (Lemma 8.1). For $r \geq 2$, $\{2^r \iota_5, \delta_r \zeta_5, 2^{m_r^3} \iota_{16}\} \subset \{2\iota_5, (2^{r-1} \iota_5) \delta_r \zeta_5 (2^{m_r^3-1}), 2\iota_{16}\} = \{2\iota_5, 0, 2\iota_{16}\} = \{0\}$. Thus $\{2^r \iota_5, \delta_r \zeta_5, 2^{m_r^3} \iota_{16}\} = \{0\}$, $\Rightarrow 2^{m_r^3} \mathbf{x}_r = 0$, which implies that $\pi_{17}(P^6(2^r)) \xrightarrow{q_{17} P_{6*}} \mathbb{Z}_{2^{m_r^3}}\{\delta_r \zeta_6\}$ is split onto.

There is $-\mathbf{y}_r \in \{i_5, 2^r \iota_5, \nu_5 \varepsilon_8\}$ such that $p_{6*}(\mathbf{y}_r) = \nu_6 \varepsilon_9 = 2\bar{\nu}_6 \nu_{14}$ [39, (7.18)] and $2\mathbf{y}_r \in i_{5\circ}\{2^r \iota_5, \nu_5 \varepsilon_8, 2\iota_{16}\}$. By Lemma 3.9

$$\{2^r \iota_5, \nu_5 \varepsilon_8, 2\iota_{16}\} \ni 2^{r-1} \nu_5 \varepsilon_8 \eta_9 \text{ mod } \{0\}.$$

Hence $2\mathbf{y}_r = (1 - \vartheta_r) i_5 \nu_5 \varepsilon_8 \eta_9$.

Claim 1: There is an element $\mathbf{z}_r \in \pi_{17}(P^6(2^r))$ with $o(\mathbf{z}_r) \leq 8$ such that $p_{6*}(\mathbf{z}_r) = \bar{\nu}_6 \nu_{14}$.

Proof of the Claim 1. Consider the cofibration sequence $S^{k+3} \xrightarrow{2\nu_k} S^k \xrightarrow{i_k^{2\nu}} C_{2\nu_k} \xrightarrow{p_{k+4}^{2\nu}} S^{k+4}$ ($k \geq 5$). For $k = 10$, it induces exact sequence

$$[C_{2\nu_{10}}, P^6(2^r)] \xrightarrow{P_{6*}} [C_{2\nu_{10}}, S^6] \xrightarrow{\partial_*^{6,r}} [C_{2\nu_{10}}, F_6],$$

with $\partial_*^{6,r}(\bar{\nu}_6 p_{14}^{2\nu}) = \partial_{13*}^{6,r}(\bar{\nu}_6) p_{13}^{2\nu} = 2^r l \nu_{10} p_{13}^{2\nu} = 0$ ($r \geq 1, l$ is odd).

Thus there is a $f_r \in [C_{2\nu_{10}}, P^6(2^r)]$, such that $p_6 f_r = \bar{v}_6 p_{14}^{2\nu}$. This implies the following commutative diagram with the top row which is exact sequence

$$\begin{array}{ccccccc} \pi_{17}(S^{10}) & \longrightarrow & \pi_{17}(C_{2\nu_{10}}) & \xrightarrow{p_{14*}^{2\nu}} & \pi_{17}(S^{14}) = \mathbb{Z}_8\{\nu_{14}\} & \longrightarrow & 0 \\ & & \downarrow f_{r*} & & \downarrow \bar{v}_{6*} & & \\ & & \pi_{17}(P^6(2^r)) & \xrightarrow{p_{6*}} & \pi_{17}(S^6) & & \end{array}$$

The following left homotopy commutative diagram induces the right commutative one

$$\begin{array}{ccccccc} S^{13} & \xrightarrow{2\nu_{10}} & S^{10} & \xrightarrow{i_{10}^{2\nu}} & C_{2\nu_{10}} & ; & 0 \longrightarrow \pi_{17}(S^{10}) \xrightarrow{i_{10*}^{2\nu}} \pi_{17}(C_{2\nu_{10}}) \longrightarrow \mathbb{Z}_8\{\nu_{14}\} \longrightarrow 0 \\ \downarrow 2\iota_{13} & & \parallel & & \downarrow \chi_{\nu}^{2\nu} & & \downarrow \times 2 \\ S^{13} & \xrightarrow{\nu_{10}} & S^{10} & \xrightarrow{i_{10}^{\nu}} & C_{\nu_{10}} & & 0 \longrightarrow \pi_{17}(S^{10}) \xrightarrow{i_{10*}^{\nu}} \pi_{17}(C_{\nu_{10}}) \longrightarrow \mathbb{Z}_4\{2\nu_{14}\} \longrightarrow 0. \end{array}$$

From Theorem 1 of [45], $\pi_{17}(C_{\nu_{10}}) \cong \mathbb{Z}_{16} \oplus \mathbb{Z}_4$ implies that $\pi_{17}(S^{10}) \xrightarrow{i_{10*}^{\nu}} \pi_{17}(C_{\nu_{10}})$ is split into, so is $\pi_{17}(S^{10}) \xrightarrow{i_{10*}^{2\nu}} \pi_{17}(C_{2\nu_{10}})$. Hence $\pi_{17}(C_{2\nu_{10}}) \xrightarrow{p_{14*}^{2\nu}} \mathbb{Z}_8\{\nu_{14}\}$ is split onto, i.e., there exists an order 8 element $\mathbf{z}_r^{2\nu} \in \pi_{17}(C_{2\nu_{10}})$, such that $p_{14*}^{2\nu}(\mathbf{z}_r^{2\nu}) = \nu_{14}$. Let $\mathbf{z}_r := f_r \mathbf{z}_r^{2\nu} \in \pi_{17}(P^6(2^r))$. We get $o(\mathbf{z}_r) \leq 8$ and $p_{6*}(\mathbf{z}_r) = \bar{v}_6 \nu_{14}$. We finish the proof of **Claim 1**. \square

Claim 2: The monomorphism $\mathbb{Z}_{2m^3}\{[\beta_5, \beta_{10}]\nu_{14}\} \xrightarrow{\tau_{6*}} \pi_{17}(P^6(2^r))$ is split into.

Proof of the Claim 2. From [42], $\Omega\Sigma L_3(P^5(2^r)) \simeq \Omega\Sigma P^{14}(2^r)$ is the Cartesian product factor of $\Omega\Sigma P^5(2^r)$ with the inclusion $\Omega\Sigma L_3(P^5(2^r)) \simeq \Omega\Sigma P^{14}(2^r) \xrightarrow{\mathcal{I}} \Omega\Sigma P^5(2^r)$, where \mathcal{I} satisfies following commutative diagram

$$\begin{array}{ccccccc} S^4 \wedge S^9 \simeq S^{13} & \xrightarrow{i_{13}} & P^{14}(2^r) \simeq L_3(P^5(2^r)) & \xrightarrow{\Omega\Sigma} & \Omega\Sigma L_3(P^5(2^r)) & \xrightarrow{\mathcal{I}} & \Omega\Sigma P^5(2^r) \\ & \searrow i_4 \wedge \alpha & \uparrow \text{proj.} & & \nearrow [E, [E, E]^S]^S & & \\ & & (P^5(2^r))^{\wedge 3} & & & & \end{array}$$

where $[E, [E, E]^S]^S$ ($E = E_{P^5(2^r)}$) is the 3-fold Samelson product; the map α and the left commutative triangle come from Proposition 2.1 and Proposition 3.1 of [6]; the right commutative triangle comes from the proof of Theorem 1.6 of [42].

By the result of $\pi_{10}(P^6(2^r))$, assume the composition $S^9 \xrightarrow{\alpha} P^5(2^r) \wedge P^5(2^r) \xrightarrow{[E, E]^S} \Omega\Sigma P^5(2^r)$ to be $\Omega_0(ai_5\nu_5\eta_8^2 + b\tau_6\beta_{10})$. Since by Proposition 2.1 of [6]

$$H_9(S^9, \mathbb{Z}_2) = \mathbb{Z}_2\{[\iota_9]\} \xrightarrow{([E, E]^S \alpha)_*} H_*(\Omega\Sigma P^5(2^r); \mathbb{Z}_2) = T(u, v), ([E, E]^S \alpha)_*([\iota_9]) = [u, v]$$

where $[\iota_9]$ is the homology class represented by ι_9 , we have b is odd by (39).

Note that $\mathbb{Z}_{2m^3}\{v_{14}\} \xrightarrow{i_{14}^*} \pi_{17}(P^{15}(2^r))$ is split into. Consider the following composition of maps

$$\mathbb{Z}_{2m^3}\{v_{14}\} \xrightarrow{i_{14}^*} \pi_{17}(P^{15}(2^r)) \xrightarrow[\cong]{\Omega_0} \pi_{16}(\Omega P^{15}(2^r)) \xrightarrow{I_*} \pi_{16}(\Omega P^6(2^r)) \xrightarrow[\cong]{\Omega_0^{-1}} \pi_{17}(P^6(2^r)).$$

$$\begin{aligned} I_*\Omega_0 i_{14}^*(v_{14}) &= I\Omega_0(i_{14}v_{14}) = I\Omega\Sigma(i_{13}v_{13})E_{S^{16}} = I\Omega\Sigma(i_{13})E_{S^{13}}v_{13} \\ &= [E, [E, E]^S]^S(i_4 \wedge \alpha)v_{13} = [\Omega_0(i_5), \Omega_0(ai_5v_5\eta_8^2 + b\tau_6\beta_{10})]^S v_{13} \\ &= (\Omega_0[i_5, ai_5v_5\eta_8^2 + b\tau_6\beta_{10}])v_{13} = b(\Omega_0[i_5, \tau_6\beta_{10}])v_{13}. \end{aligned}$$

where the last equation holds because $[i_5, ai_5v_5\eta_8^2] = ai_5[\iota_5, \iota_5]v_9\eta_{12}^2 = 0$.

$$\Omega_0^{-1}I_*\Omega_0 i_{14}^*(v_{14}) = \Omega_0^{-1}(b(\Omega_0[i_5, \tau_6\beta_{10}])v_{13}) = b[i_5, \tau_6\beta_{10}]v_{14}.$$

Since $\Omega_0^{-1}I_*\Omega_0 i_{14}^* : \mathbb{Z}_{2m^3}\{v_{14}\} \hookrightarrow \pi_{17}(P^6(2^r))$ is split into, so is the monomorphism in **Claim 2**. \square

For \mathbf{z}_r in **Claim 1**, $p_{6^*}(2\mathbf{z}_r - \mathbf{y}_r) = 0$, i.e. $2\mathbf{z}_r - \mathbf{y}_r \in \tau_{6^*}(\text{Coker}(\partial_{17^*}^{6,r}))$, which implies $4\mathbf{z}_r - \epsilon_r i_5 v_5 \epsilon_8 \eta_9 = 2a_r \tau_6 \beta_{10} \sigma_{10} + 2b_r \tau_6 [\beta_5, \beta_{10}] v_{14}$, $a_1 \in \mathbb{Z}_8$, $a_r \in \mathbb{Z}_{2^{m^4}} (r \geq 2)$, $b_r \in \mathbb{Z}_{2m^3}$.

For $r = 1$, $4\mathbf{z}_1 = i_5 v_5 \epsilon_8 \eta_9 + 2a_1 \tau_6 \beta_{10} \sigma_{10}$. Since $o(\mathbf{z}_1) \leq 8$, $2a_1 = 4a'_1$ for some $a'_1 \in \mathbb{Z}$. Let $\mathbf{z}'_1 = \mathbf{z}_1 - a'_1 \tau_6 \beta_{10} \sigma_{10}$, then $o(\mathbf{z}'_1) = 8$ and $4\mathbf{z}'_1 = i_5 v_5 \epsilon_8 \eta_9$. Hence

$$\begin{aligned} \pi_{17}(P^6(2)) &= \mathbb{Z}_2\{i_5 v_5^4\} \oplus \mathbb{Z}_2\{i_5 v_5 \bar{\mu}_8\} \oplus \mathbb{Z}_8\{\tau_6 \beta_{10} \sigma_{10}\} \oplus \mathbb{Z}_2\{[i_5, \tau_6 \beta_{10}]v_{14}\} \\ &\quad \oplus \mathbb{Z}_2\{\tau_6 \check{I}_3 \widehat{\eta_{15} \eta_{16}}\} \oplus \mathbb{Z}_2\{\widehat{4\zeta_6}\} \oplus \mathbb{Z}_8\{\mathbf{z}'_1\}, \quad 4\mathbf{z}'_1 = i_5 v_5 \epsilon_8 \eta_9. \\ \Rightarrow \pi_{17}(P^6(2)) &\cong \mathbb{Z}_2^{\oplus 5} \oplus \mathbb{Z}_8^{\oplus 2}. \end{aligned}$$

For $r \geq 2$, $2\mathbf{y}_r = 0$. $4\mathbf{z}_r = 2a_r \tau_6 \beta_{10} \sigma_{10} + 2b_r \tau_6 [\beta_5, \beta_{10}] v_{14}$. $o(\mathbf{z}_r) \leq 8$ implies $a_r = 2a'_r$, $a'_r \in \mathbb{Z}$. Then $2b_r \tau_6 [\beta_5, \beta_{10}] v_{14} = 4(\mathbf{z}_r - a'_r \tau_6 \beta_{10} \sigma_{10})$. By **Claim 2**, $b_r = 2b'_r$, $b'_r \in \mathbb{Z}$. Take $\mathbf{z}'_r = \mathbf{z}_r - a'_r \tau_6 \beta_{10} \sigma_{10} - b'_r \tau_6 [\beta_5, \beta_{10}] v_{14}$, which is an order 4 element such that $p_{6^*}(\mathbf{z}'_r) = \bar{v}_6 v_{14}$. So

$$\pi_{17}(P^6(2^r)) \cong \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_{2m^3}^{\oplus 2} \oplus \mathbb{Z}_{2^{m^4}}.$$

- $\pi_{18}(P^6(2^r))$.

Consider the commutative diagram (6) for $k = 5, m = 18, r \geq 1$.

$$\partial_{18^*}^{6,r}(v_6 \sigma_9 v_{16}) = \beta_{5^*}(2^r \iota_6)_*(v_5 \sigma_8 v_{15}) = 0.$$

We have the short exact sequence

$$\begin{aligned} 0 \rightarrow \text{Coker}(\partial_{18^*}^{6,r}) \xrightarrow{\tau_{6^*}} \pi_{18}(P^6(2^r)) \xrightarrow{p_{6^*}} \mathbb{Z}_{2^{m^4}}\{\delta_r \Delta(\sigma_{13})\} \rightarrow 0, \quad (40) \\ \text{Coker}(\partial_{18^*}^{6,r}) = \mathbb{Z}_2\{\beta_5 v_5 \sigma_8 v_{15}\} \oplus \mathbb{Z}_2\{\beta_5 v_5 \eta_8 \mu_9\} \oplus \mathbb{Z}_2\{\beta_{10} \bar{v}_{10}\} \oplus \mathbb{Z}_2\{\beta_{10} \epsilon_{10}\} \\ \oplus \mathbb{Z}_{2^r}\{[\beta_5, [\beta_5, \beta_{10}]]\} \oplus \mathbb{Z}_{2m^3}\{\check{I}_3 \widehat{\delta_r v_{15}}\}. \end{aligned}$$

For $r = 1$, by Lemma 3.8, there is $-\mathbf{x}_1 \in \{i_5, 2\iota_5, \nu_5\mu_8\}$ such that $p_{6*}(\mathbf{x}_1) = \nu_6\mu_9 = 8\Delta(\nu_{13})$ [39, (7.25)] and $2\mathbf{x}_1 \in i_{5\circ}\{2\iota_5, \nu_5\mu_8, 2\iota_{17}\}$. $\{2\iota_5, \nu_5\mu_8, 2\iota_{17}\} \ni \nu_5\mu_8\eta_{17} = \nu_5\eta_8\mu_9 \pmod{\{0\}}$ by Lemma 8.1 (ii) and [39, Corollary 3.7]. Hence $2\mathbf{x}_1 = i_5\nu_5\eta_8\mu_9 \neq 0$, which implies that $o(\mathbf{x}_1) = 4$. Assume that the exact sequence (40) is split, then $\pi_{18}(P^6(2)) \cong \mathbb{Z}_2^{\oplus 6} \oplus \mathbb{Z}_4$ has no order 4 element taken to $8\Delta(\sigma_{13}) \in \mathbb{Z}_4\{4\Delta(\sigma_{13})\}$ by p_{6*} . This contradicts to existence of \mathbf{x}_1 . So (40) is not split for $r = 1$. We have

$$\pi_{18}(P^6(2)) \cong \mathbb{Z}_2^{\oplus 5} \oplus \mathbb{Z}_8.$$

For $r \geq 2$, consider the following exact sequence

$$[P^{11}(2^r), P^6(2^r)] \xrightarrow{p_{6*}} [P^{11}(2^r), S^6] \xrightarrow{\partial_*^{6,r}} [P^{11}(2^r), F_6],$$

with $\partial_*^{6,r}(l\Delta(\iota_{13})p_{11}) = l\partial_{10*}^{6,r}(\Delta(\iota_{13}))p_{10} = 2^{r+1}l\beta_{10}p_{10} = l\beta_{10}(2^{r+1}p_{10}) = 0$,

where the odd integer l is given by $\Delta(\sigma_{13}) = l\Delta(\iota_{13})\sigma_{11}$.

Thus there is an $f_r \in [P^{11}(2^r), P^6(2^r)]$, such that $p_6 f_r = l\Delta(\iota_{13})p_{11}$. This implies the following commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}_2^{\oplus 2} \cong \pi_{18}(S^{10}) & \longrightarrow & \pi_{18}(P^{11}(2^r)) & \xrightarrow{p_{11*}} & \mathbb{Z}_{2^{m_r^4}}\{\delta_r^{16}\sigma_{11}\} \longrightarrow 0 \\ & & \downarrow & & \downarrow f_{r*} & & (l\Delta(\iota_{13}))_* \downarrow \\ 0 & \longrightarrow & \text{Coker}(\partial_{18*}^{6,r}) & \longrightarrow & \pi_{18}(P^6(2^r)) & \xrightarrow{p_{6*}} & \mathbb{Z}_{2^{m_{r+1}^4}}\{\delta_r\Delta(\nu_{13})\} \longrightarrow 0 \end{array}$$

where $\delta_r^{16} = 2\delta_r$ for $r \leq 3$ and $\delta_r^{16} = 1$ for $r \geq 4$.

There is $-\mathbf{x}_r \in \{i_{10}, 2^r\iota_{10}, \delta_r^{16}\sigma_{10}\}$ such that $p_{11*}(\mathbf{x}_r) = \delta_r^{16}\sigma_{11}$ and $2^{m_r^4}\mathbf{x}_r \in i_{10}\{2^r\iota_{10}, \delta_r^{16}\sigma_{10}, 2^{m_r^4}\iota_{17}\}$.

$$\{2^r\iota_{10}, \delta_r^{16}\sigma_{10}, 2^{m_r^4}\iota_{17}\} \subset \{2\iota_{10}, (2^{r-1}\iota_{10})(\delta_r^{16}\sigma_{10})(2^{m_r^4-1}\iota_{17}), 2\iota_{17}\} = \{2\iota_{10}, 0, 2\iota_{17}\} = \{0\}.$$

Thus $2^{m_r^4}\mathbf{x}_r = 0$. We get that the top short exact sequence in the above diagram is split, so is the bottom one for $2 \leq r \leq 4$ by $2^{m_r^4} \cdot \text{Coker}(\partial_{18*}^{6,r}) = 0$ with [49, Lemma 2.9] and for $r \geq 4$ by the isomorphism of right map $(l\Delta(\iota_{13}))_*$. Thus

$$\pi_{18}(P^6(2^r)) \cong \mathbb{Z}_2^4 \oplus \mathbb{Z}_{2^r} \oplus \mathbb{Z}_{2^{m_r^3}} \oplus \mathbb{Z}_{2^{m_{r+1}^4}}, r \geq 2.$$

6. Homotopy groups of 7 and 8 dimensional mod 2^r Moore spaces

In this section we will give the unstable homotopy groups $\pi_i(P^7(2^r))$ for $i \leq 19$ and $\pi_i(P^8(2^r))$ for $i \leq 21$.

6.1. Homotopy groups of $P^7(2^r)$

There are cofibration and fibration sequences

$$\begin{aligned} S^6 \xrightarrow{2^r \iota_6} S^6 \xrightarrow{i_6} P^7(2^r) \xrightarrow{p_7} S^7; \quad \Omega S^7 \xrightarrow{\partial^7} F_7 \xrightarrow{\tau_7} P^7(2^r) \xrightarrow{p_7} S^7 \\ Sk_{17}(F_7) \simeq J_{2,6}^r \simeq S^6 \cup_{\gamma_2^r = 2^r \Delta(\iota_{13})} CS^{11} (r \geq 1); \quad Sk_{23}(F_7) \simeq J_{3,6}^r \simeq J_{2,6}^r \cup_{\gamma_3^r} CS^{17}. \end{aligned}$$

Lemma 6.1. *For $r \geq 1$ and $i \leq 16$, $\pi_i(F_7) \cong \pi_i(J_{2,6}^r)$ and*

- (1) $\pi_{11}(J_{2,6}^r) = \mathbb{Z}_{2^r} \{\beta_6 \Delta(\iota_{13})\}$.
- (2) $\pi_{12}(J_{2,6}^r) = \mathbb{Z}_2 \{\beta_6 \nu_6^2\}$.
- (3) $\pi_{13}(J_{2,6}^r) = \mathbb{Z}_4 \{\beta_6 \sigma''\} \oplus \mathbb{Z}_2 \{\widehat{\eta}_{12}\}$.
- (4) $\pi_{14}(J_{2,6}^r) = \mathbb{Z}_{2^{m_{r+1}}} \{\beta_6 \bar{\nu}_6\} \oplus \mathbb{Z}_2 \{\beta_6 \varepsilon_6\} \oplus \mathbb{Z}_2 \{\widehat{\eta}_{12} \eta_{13}\}$.
- (5) $\pi_{15}(J_{2,6}^r) = \mathbb{Z}_2 \{\beta_6 \nu_6^3\} \oplus \mathbb{Z}_2 \{\beta_6 \mu_6\} \oplus \mathbb{Z}_2 \{\beta_6 \eta_6 \varepsilon_7\} \oplus \mathbb{Z}_{2^{m_{r+1}}} \{\widehat{\delta_{r+1} \nu_{12}}\}$.
- (6) $\pi_{16}(J_{2,6}^r) = \mathbb{Z}_8 \{\beta_6 \nu_6 \sigma_9\} \oplus \mathbb{Z}_2 \{\beta_6 \eta_6 \mu_7\}$.
- (7) $\pi_{17}(J_{2,6}^r) = \mathbb{Z}_8 \{\beta_6 \zeta_6\} \oplus \mathbb{Z}_4 \{\beta_6 \bar{\nu}_6 \nu_{14}\} \oplus \mathbb{Z}_{(2)} \{\beta_{17}\}$
 $\pi_{17}(J_{2,6}^0) = \mathbb{Z}_8 \{\beta_6^0 \zeta_6\} \oplus \mathbb{Z}_2 \{\beta_6^0 \bar{\nu}_6 \nu_{14}\} \oplus \mathbb{Z}_{(2)} \{\beta_{17}^0\}$.
- (8) $\pi_{18}(J_{2,6}^r) = \mathbb{Z}_{2^{m_r^4}} \{\beta_6 \Delta(\sigma_{13})\} \oplus \mathbb{Z}_2 \{\beta_{17} \eta_{17}\} \oplus \mathbb{Z}_2 \{\widehat{\nu}_{12}^2\}$.
- (9) $\pi_{19}(J_{2,6}^r) = \mathbb{Z}_2 \{\beta_6 \nu_6 \sigma_9 \nu_{16}\} \oplus \mathbb{Z}_2 \{\beta_{17} \eta_{17}^2\} \oplus \mathbb{Z}_{2^{m_r^4}} \{\widehat{\delta_r^{16} \sigma_{12}}\}$.

Proof. There are fibration sequence $F_{\bar{p}_{12}} \xrightarrow{\bar{\tau}_{12}} J_{2,6}^r \xrightarrow{\bar{p}_{12}} S^{12}$, with $Sk_{27}(F_{\bar{p}_{12}}) \simeq S^6 \vee S^{17}$. Consider the Lemma 4.1 for $k = 6, 11 \leq m \leq 21$.

For (1), (2), (6) and (7) with $r \geq 1$ of Lemma 6.1, they are easily obtained by the following isomorphism

$$\pi_i(F_{\bar{p}_{12}}) \xrightarrow[\cong]{\bar{\tau}_{12^*}} \pi_i(J_{2,6}^r), i = 11, 12, 16, 17.$$

For $\pi_{17}(J_{2,6}^0)$ of (7), by [39, Lemma 6.2], $\Delta(\iota_{13})\nu_{11} = \pm\Delta(\nu_{13}) = \pm 2\bar{\nu}_6$, it is get by

$$\bar{\delta}_{17^*}^{12,0}(\nu_{12}^2) = \bar{\beta}_{6^*}^0(\Delta(\iota_{13}))_*(\nu_{11}^2) = \bar{\beta}_6^0(\pm\Delta(\nu_{13}))\nu_{14} = 2\bar{\beta}_6^0\bar{\nu}_6\nu_{14}.$$

For (3) of Lemma 6.1, we have the following exact sequence

$$0 \rightarrow \mathbb{Z}_4 \{\bar{\beta}_6 \sigma''\} = \pi_{13}(F_{\bar{p}_{12}}) \xrightarrow{\bar{\tau}_{12^*}} \pi_{13}(J_{2,6}^r) \xrightarrow{\bar{p}_{12^*}} \pi_{13}(S^{12}) = \mathbb{Z}_2 \{\eta_{12}\} \rightarrow 0.$$

There is $-\mathbf{x} \in \{\beta_6, \gamma_2, \eta_{11}\}$ such that $\bar{p}_{12^*}(\mathbf{x}) = \eta_{12}$ and $2\mathbf{x} \in \beta_6\{2^r \Delta(\iota_{13}), \eta_{11}, 2\iota_{12}\}$. By [39, Page 47], $\Delta(\iota_{13})\eta_{11} = \Delta(\eta_{13}) = 0$. By Lemma 3.9

$$\{2^r \Delta(\iota_{13}), \eta_{11}, 2\iota_{12}\} \ni (2^{r-1} \Delta(\iota_{13}))\eta_{11}^2 = 0 \pmod{\{0\}}.$$

Hence $2\mathbf{x} = 0$, which implies $\pi_{13}(J'_{2,6}) \xrightarrow{\bar{p}_{12}^{12*}} \mathbb{Z}_2\{\eta_{12}\}$ is split onto.

For (4) of Lemma 6.1, since $\gamma_{2^*}(v_{11}) = 2^r\Delta(\iota_{13})v_{11} = 2^r\Delta(v_{13}) = 2^{r+1}\bar{v}_6$ by [39, Lemma 6.2], we have

$$0 \rightarrow \mathbb{Z}_{2^{m_{r+1}^3}}\{\bar{\beta}_6\bar{v}_6\} \oplus \mathbb{Z}_2\{\beta_6\varepsilon_6\} = \text{Coker}(\bar{\delta}_{14^*}^{12,r}) \xrightarrow{\bar{\tau}_{12^*}} \pi_{14}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_2\{\eta_{12}^2\} \rightarrow 0.$$

Since the order 2 element $\widehat{\eta}_{12}\eta_{13} \in \pi_{14}(J'_{2,6})$ is the lift of η_{12}^2 by \bar{p}_{12} , where $\widehat{\eta}_{12} \in \pi_{13}(J'_{2,6})$ is the lift of η_{12} by \bar{p}_{12} , we get $\pi_{14}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_2\{\eta_{12}^2\}$ is split onto.

For (5) of Lemma 6.1, we have the following exact sequence

$$0 \rightarrow \mathbb{Z}_2\{\bar{\beta}_6 v_6^3\} \oplus \mathbb{Z}_2\{\bar{\beta}_6 \mu_6\} \oplus \mathbb{Z}_2\{\beta_6 \eta_6 \varepsilon_7\} = \pi_{15}(F_{\bar{p}_{12}}) \xrightarrow{\bar{\tau}_{12^*}} \pi_{15}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_{2^{m_{r+1}^3}}\{\delta_{r+1}v_{12}\} \rightarrow 0.$$

$\exists -\mathbf{x}_r \in \{\beta_6, \gamma_2, \delta_{r+1}v_{11}\}$ such that $\bar{p}_{12}\mathbf{x}_r = \delta_{r+1}v_{12}$ and $2^{m_{r+1}^3}\mathbf{x}_r \in \beta_6\{2^r\Delta(\iota_{13}), \delta_{r+1}v_{11}, 2^{m_{r+1}^3}\iota_{14}\}$, where $\{2^r\Delta(\iota_{13}), \delta_{r+1}v_{11}, 2^{m_{r+1}^3}\iota_{14}\} \subset \{\Delta(\iota_{13}), (2^r\iota_{11})\delta_{r+1}v_{11}, (2^{m_{r+1}^3-1}\iota_{14}), 2\iota_{14}\} = \{\Delta(\iota_{13}), 0, 2\iota_{14}\} \ni 0 \pmod{\{0\}}$. Hence $\{2^r\Delta(\iota_{13}), \delta_{r+1}v_{11}, 2^{m_{r+1}^3}\iota_{14}\} = \{0\}$ implies that $2^{m_{r+1}^3}\mathbf{x}_r = 0$. So $\pi_{15}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_{2^{m_{r+1}^3}}\{\delta_{r+1}v_{12}\}$ is split onto.

For (8) of Lemma 6.1, since $\gamma_{2^*}(\sigma_{11}) = 2^r\Delta(\iota_{13})\sigma_{11} = 2^r l' \Delta(\sigma_{13})$ where l' is the inverse of the odd integer l in \mathbb{Z}_{16} which is given by $\Delta(\sigma_{13}) = l\Delta(\iota_{13})\sigma_{11}$. we have

$$0 \rightarrow \mathbb{Z}_{2^{m_r^4}}\{\bar{\beta}_6\Delta(\sigma_{13})\} \oplus \mathbb{Z}_2\{\bar{\beta}_{17}\eta_{17}\} = \pi_{18}(F_{\bar{p}_{12}}) \xrightarrow{\bar{\tau}_{12^*}} \pi_{18}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_2\{v_{12}^2\} \rightarrow 0.$$

$\exists -\mathbf{x}_r \in \{\beta_6, \gamma_2, v_{11}^2\}$ such that $\bar{p}_{12}\mathbf{x}_r = v_{12}^2$ and $2\mathbf{x}_r \in \beta_6\{2^r\Delta(\iota_{13}), v_{11}^2, 2\iota_{17}\}$, where $\{2^r\Delta(\iota_{13}), v_{11}^2, 2\iota_{17}\} \ni 2^{r-1}\Delta(\iota_{13})v_{11}^2\eta_{17} = 0 \pmod{\langle 2\Delta(\sigma_{13}) \rangle}$. Hence $2\mathbf{x}_r = 2t_r\beta_6\Delta(\sigma_{13}), t_r \in \mathbb{Z}$, which implies $\pi_{18}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_2\{v_{12}^2\}$ is split onto.

For (9) of Lemma 6.1, we have

$$0 \rightarrow \mathbb{Z}_2\{\bar{\beta}_6 v_6 \sigma_9 v_{16}\} \oplus \mathbb{Z}_2\{\bar{\beta}_{17} \eta_{17}^2\} = \pi_{19}(F_{\bar{p}_{12}}) \xrightarrow{\bar{\tau}_{12^*}} \pi_{19}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_{2^{m_r^4}}\{\delta_r^{16} \sigma_{12}\} \rightarrow 0.$$

$\exists -\mathbf{x}_r \in \{\beta_6, \gamma_2, \delta_r^{16}\sigma_{11}\}$ such that $\bar{p}_{12}\mathbf{x}_r = \delta_r^{16}\sigma_{12}$ and $2^{m_r^4}\mathbf{x}_r \in \beta_6\{2^r\Delta(\iota_{13}), \delta_r^{16}\sigma_{11}, 2^{m_r^4}\iota_{18}\}$, where $\{2^r\Delta(\iota_{13}), \delta_r^{16}\sigma_{11}, 2^{m_r^4}\iota_{18}\}$ is a coset of subgroup $(2^r\Delta(\iota_{13})) \circ \pi_{19}(S^{11}) + \pi_{19}(S^6) \circ (2^{m_r^4}\iota_{19}) = \{0\}$. For $r = 1$, $\{2\Delta(\iota_{13})\iota_{11}, 8\sigma_{11}, 2\iota_{18}\} \ni \Delta(\iota_{13})(8\sigma_{11})\eta_{18} = 0$; For $r \geq 2$, $\{2^r\Delta(\iota_{13}), \delta_r^{16}\sigma_{11}, 2^{m_r^4}\iota_{18}\} \subset \{2\Delta(\iota_{13}), 0, 2\iota_{18}\} = \{0\}$. So $2^{m_r^4}\mathbf{x}_r = 0, r \geq 1$, which implies $\pi_{19}(J'_{2,6}) \xrightarrow{\bar{p}_{12^*}} \mathbb{Z}_{2^{m_r^4}}\{\delta_r^{16} \sigma_{12}\}$ is split onto. \square

Lemma 6.2. For $r \geq 0$, $\gamma_3^r = 2^r a_0 \beta_{17} + 2^r b_r \beta_6 \zeta_6 + 2^r c_r \beta_6 \bar{v}_6 v_{14}$, where a_0 is an odd integer, $b_r \in \mathbb{Z}_8, c_r \in \mathbb{Z}_4$.

Proof. By (7) of Lemma 6.1, assume $\gamma_3^r = a_r \beta_{17} + b'_r \beta_6 \zeta_6 + c'_r \beta_6 \bar{v}_6 v_{14} \in \pi_{17}(J_{2,6}^r)$, where $a_r \in \mathbb{Z} \subset \mathbb{Z}_{(2)}$, $b'_r \in \mathbb{Z}_8$, $c'_0 \in \mathbb{Z}_2$, $c'_r \in \mathbb{Z}_4$, $r' \geq 1$. There is the following isomorphism

$$\begin{aligned} \mathbb{Z}_8 \oplus \mathbb{Z}_2 &\cong \pi_{18}(S^7) \xrightarrow[\cong]{\bar{\delta}_{17^*}^{12,0}} \pi_{17}(F_{p_7^0}) \cong \pi_{17}(J_{3,6}^0) = \pi_{17}(J_{2,6}^0) / \langle \gamma_3^0 \rangle \\ &= \frac{\mathbb{Z}_8 \{ \beta_6^0 \zeta_6 \} \oplus \mathbb{Z}_2 \{ \beta_6^0 \bar{v}_6 v_{14} \} \oplus \mathbb{Z}_{(2)} \{ \beta_{17}^0 \}}{\langle a_0 \beta_{17}^0 + b'_0 \beta_6^0 \zeta_6 + c'_0 \beta_6^0 \bar{v}_6 v_{14} \rangle}. \end{aligned}$$

If $a_0 = 2^{t_0} a'_0$, a'_0 is odd, $0 \leq t_0 \in \mathbb{Z}$, then we get the short exact sequence

$$0 \rightarrow \mathbb{Z}_8 \{ \beta_6^0 \zeta_6 \} \oplus \mathbb{Z}_2 \{ \beta_6^0 \bar{v}_6 v_{14} \} \xrightarrow{I} \pi_{17}(J_{2,6}^0) / \langle \gamma_3^0 \rangle \xrightarrow{P} \mathbb{Z}_{2^{t_0}} \{ \beta_{17}^0 \} \rightarrow 0$$

where the definition of inclusion I and quotient P are obvious. So $t_0 = 0$, i.e., a_0 is odd. By the equations (12), (15) and diagram (10) for $k = 6, s = r \geq 1, t = 0$, we get

$$\begin{aligned} \bar{g}_0^r (a_r \beta_{17}^r + b'_r \beta_6^r \zeta_6 + c'_r \beta_6^r \bar{v}_6 v_{14}) &= 2^{2r} (a_0 \beta_{17}^0 + b'_0 \beta_6^0 \zeta_6 + c'_0 \beta_6^0 \bar{v}_6 v_{14}), \text{ with } g_0^r \beta_6^r = \beta_6^0. \\ (g_0^r)|_{J_2} = j_1^6 q_1^6 + 2^r j_2^{17} q_2^{17} + j_1^6 \theta q_2^{17}, \quad \theta &= b_\theta \zeta_6 + c_\theta \bar{v}_6 v_{14} \in \pi_{17}(S^6) = \mathbb{Z}_8 \{ \zeta_6 \} \oplus \mathbb{Z}_4 \{ \bar{v}_6 v_{14} \}. \end{aligned}$$

$$\begin{array}{ccccc} S^{17} \xrightarrow{j_2^{17}} S^6 \vee S^{17} &\simeq & J_{2,6}^{\gamma,r} &\xrightarrow{I_2^{\gamma,r}} & F_{\bar{p}_{12}^r} &\xrightarrow{\bar{\tau}_{12}^r} & J_{2,6}^r, \\ & & \downarrow (g_0^r)|_{J_2} & & \downarrow g_0^r & & \downarrow \bar{g}_0^r \\ S^{17} \xrightarrow{j_2^{17}} S^6 \vee S^{17} &\simeq & J_{2,6}^{\gamma,0} &\xrightarrow{I_2^{\gamma,0}} & F_{\bar{p}_{12}^0} &\xrightarrow{\bar{\tau}_{12}^0} & J_{2,6}^0. \end{array}$$

$$\begin{aligned} \bar{g}_0^r \beta_{17}^r &= \bar{g}_0^r (\bar{\tau}_{12}^r I_2^{\gamma,r} j_2^{17}) = \bar{\tau}_{12}^0 I_2^{\gamma,0} (g_0^r)|_{J_2} j_2^{17} = \bar{\tau}_{12}^0 I_2^{\gamma,0} (2^r j_2^{17} + j_1^6 \theta) = 2^r \beta_{17}^0 + b_\theta \beta_6^0 \zeta_6 + c_\theta \beta_6^0 \bar{v}_6 v_{14}. \\ \Rightarrow 2^r a_r \beta_{17}^0 + (a_r b_\theta + b'_r) \beta_6^0 \zeta_6 + (a_r c_\theta + c'_r) \beta_6^0 \bar{v}_6 v_{14} &= 2^{2r} a_0 \beta_{17}^0 + 2^{2r} b'_0 \beta_6^0 \zeta_6. \end{aligned}$$

Comparing the coefficients of the torsion free element β_{17}^0 , we get $a_r = 2^r a_0$. Then $2^r a_0 b_\theta + b'_r = 2^{2r} b'_0 \in \mathbb{Z}_8$ and $2^r a_0 c_\theta + c'_r = 0 \in \mathbb{Z}_2$ imply $b'_r = 2^r b_r$, $c'_r = 2c_r''$. Hence

$$\gamma_3^r = 2^r a_0 \beta_{17} + 2^r b_r \beta_6 \zeta_6 + 2c_r'' \beta_6 \bar{v}_6 v_{14}.$$

By the equations (12) and (15) for $k = 6, s = r \geq 2, t = 1$, we get

$$\bar{g}_1^r (2^r a_0 \beta_{17}^r + 2^r b_r \beta_6^r \zeta_6 + 2c_r'' \beta_6^r \bar{v}_6 v_{14}) = 2^{2(r-1)} (2a_0 \beta_{17}^1 + 2b_1 \beta_6^1 \zeta_6 + 2c_1'' \beta_6^1 \bar{v}_6 v_{14}).$$

Comparing the coefficients of $\beta_6^r \bar{v}_6 v_{14}$ on both sides of above equation, $2c_r'' = 0$ ($r \geq 2$). \square

Lemma 6.3.

$$\pi_{17}(J_{3,6}^r) = \begin{cases} \mathbb{Z}_2 \{ a_0 \check{\beta}_{17} + b_1 \check{\beta}_6 \zeta_6 + c_1 \check{\beta}_6 \bar{v}_6 v_{14} \} \oplus \mathbb{Z}_8 \{ \check{\beta}_6 \zeta_6 \} \oplus \mathbb{Z}_4 \{ \check{\beta}_6 \bar{v}_6 v_{14} \}, & r = 1; \\ \mathbb{Z}_4 \{ a_0 \check{\beta}_{17} + b_2 \check{\beta}_6 \zeta_6 \} \oplus \mathbb{Z}_8 \{ \check{\beta}_6 \zeta_6 \} \oplus \mathbb{Z}_4 \{ \check{\beta}_6 \bar{v}_6 v_{14} \}, & r = 2; \\ \mathbb{Z}_{2^r} \{ a_0 \check{\beta}_{17} \} \oplus \mathbb{Z}_8 \{ \check{\beta}_6 \zeta_6 \} \oplus \mathbb{Z}_4 \{ \check{\beta}_6 \bar{v}_6 v_{14} \}, & r \geq 3. \end{cases}$$

$$\pi_{18}(J_{3,6}^r) = \mathbb{Z}_{2^{m_r^4}} \{ \check{\beta}_6 \Delta(\sigma_{13}) \} \oplus \mathbb{Z}_2 \{ \check{\beta}_{17} \eta_{17} \} \oplus \mathbb{Z}_2 \{ I_2^r v_{12}^2 \}.$$

$$\pi_{19}(J_{3,6}^r) = \begin{cases} \mathbb{Z}_2 \{ \check{\beta}_6 v_6 \sigma_9 v_{16} \} \oplus \mathbb{Z}_2 \{ I_2^1 8 \widehat{\sigma}_{12} \} \oplus \mathbb{Z}_4 \{ \widehat{\eta}_{18} \}, & 2\widehat{\eta}_{18} = \check{\beta}_{17}^1 \eta_{17}^2 \quad r = 1; \\ \mathbb{Z}_2 \{ \check{\beta}_6 v_6 \sigma_9 v_{16} \} \oplus \mathbb{Z}_2 \{ \check{\beta}_{17} \eta_{17}^2 \} \oplus \mathbb{Z}_{2^{m_r^4}} \{ I_2^r \delta_r^{16} \widehat{\sigma}_{12} \} \oplus \mathbb{Z}_2 \{ \widehat{\eta}_{18} \}, & r \geq 2. \end{cases}$$

Proof. Consider the Lemma 4.3 for $m = 17, 18, 19, k = 6$.

$\pi_m(J_{3,6}^r)$ for $m = 17, 18$ are easily obtained from the isomorphisms

$$\frac{\pi_{17}(J_{2,6}^r)}{\langle \gamma_3^r \rangle} \xrightarrow[\cong]{I_{2*}^r} \pi_{17}(J_{3,6}^r); \quad \pi_{18}(J_{2,6}^r) \xrightarrow[\cong]{I_{2*}^r} \pi_{18}(J_{3,6}^r).$$

For $\pi_{19}(J_{3,6}^r)$, we get $0 \rightarrow \pi_{19}(J_{2,6}^r) \xrightarrow{I_{2*}^r} \pi_{19}(J_{3,6}^r) \xrightarrow{\check{p}_{18*}} \pi_{19}(S^{18}) = \mathbb{Z}_2\{\eta_{18}\} \rightarrow 0$.
 $\exists -\mathbf{x}_r \in \{I_2^r, \gamma_3^r, \eta_{17}\} \subset \pi_{19}(J_{3,6}^r)$, such that $\check{p}_{12*}(\mathbf{x}_r) = \eta_{18}$ and $2\mathbf{x}_r \in I_2^r\{\gamma_3^r, \eta_{17}, 2\iota_{18}\}$.

From Lemma 3.9, Lemma 6.2 and by $\zeta_6\eta_{17} \in \langle 8\Delta(\sigma_{13}) \rangle \subset \pi_{18}(S^6) = \mathbb{Z}_{16}\{\Delta(\sigma_{13})\}$,

$$\{\gamma_3^r, \eta_{17}, 2\iota_{18}\} \ni 2^{r-1}(a_0\beta_{17} + b_r\beta_6\zeta_6 + c_r\beta_6\bar{v}_6v_{14})\eta_{17}^2 = 2^{r-1}\beta_{17}\eta_{17}^2 \pmod{\langle 2\pi_{19}(J_{2,6}^r) \rangle}.$$

Hence $2\mathbf{x}_r = 2^{r-1}\check{\beta}_{17}\eta_{17}^2 + 2I_2^r\theta_r$ for some $\theta_r \in \pi_{19}(J_{2,6}^r)$. Take $\mathbf{x}'_r := \mathbf{x}_r - I_2^r\theta_r$, then $o(\mathbf{x}'_1) = 4$ with $2\mathbf{x}'_1 = \check{\beta}_{17}\eta_{17}^2$ and $o(\mathbf{x}'_r) = 2$ for $r \geq 2$. Thus we get the $\pi_{19}(J_{3,6}^r)$. \square

Remark 6.4. In fact, by our calculation we find that the groups $\pi_i(J_{3,6}^r)$ and $\pi_i(P^7(2^r))$ for $i \leq 22$, under isomorphism, have nothing to do with the undetermined coefficients b_r, c_r in Lemma 6.2.

- $\pi_m(P^7(2^r)), m = 11, 12$.

Consider diagram (6) for $k = 6, m = 11, 12$, we get the isomorphism $\pi_m(J_{2,6}^r) \cong \pi_i(F_7) \xrightarrow[\cong]{\tau_{7*}} \pi_m(P^7(2^r))$. Hence (1) (2) of Lemma 6.1 implies

$$\pi_{11}(P^7(2^r)) \cong \mathbb{Z}_{2^r}, r \geq 1; \quad \pi_{12}(P^7(2^r)) \cong \mathbb{Z}_2, r \geq 1.$$

- $\pi_{13}(P^7(2^r))$.

There is the exact sequence

$$\mathbb{Z}_8\{\sigma'\} = \pi_{14}(S^7) \xrightarrow{\partial_{13,*}^{7,r}} \pi_{13}(F_7) \xrightarrow{\tau_{7*}} \pi_{13}(P^7(2^r)) \xrightarrow{p_{7*}} \mathbb{Z}_2\{v_7^2\} \rightarrow 0.$$

By (3) of Lemma 6.1, assume that $\partial_{13,*}^{7,r}(\sigma') = x_r\beta_6\sigma'' + y_r\widehat{\eta_{12}}$, $x_r \in \mathbb{Z}_4, y_r \in \mathbb{Z}_2$.

Lemma 2.7 and Corollary 3.4 imply the commutative diagrams

$$\begin{array}{ccc} \pi_{14}(S^7) & \xrightarrow{\partial_{13,*}^{7,r}} & \pi_{13}(F_7) \xleftarrow[\cong]{I_{2*}^r} \pi_{13}(J_{2,6}^r) \xrightarrow{\bar{p}_{12*}} \pi_{13}(S^{12}); & \pi_{14}(S^7) & \xrightarrow{\partial_{13,*}^{7,r}} & \pi_{13}(F_7) \\ H_2 \downarrow & & H_2 \downarrow & \Sigma \downarrow & & \Sigma^\infty \downarrow & \downarrow E^\infty \\ \pi_{14}(S^{13}) & \xrightarrow{(2^r\iota_{13})_*} & \pi_{14}(S^{13}) & \xrightarrow{\cong} & \pi_{14}(S^{13}) & \xrightarrow{\mathbb{Z}_{16}\{\sigma\} = \pi_{13}^s(S^6)} & \xrightarrow{(2^r\iota)_*} & \pi_{13}^s(S^6) \end{array}$$

$$0 = H_2'\partial_{13,*}^{7,r}(\sigma') = H_2'(x_r\beta_6\sigma'' + y_r\widehat{\eta_{12}}) = y_r\eta_{13} \Rightarrow y_r = 0.$$

$(2^r \iota)_* \Sigma^\infty(\sigma') = 2^{r+1} \sigma$ [39, Lemma 5.14] and the commutative triangle in Corollary 3.4 implies $E^\infty \partial_{13*}^{7,r}(\sigma') = E^\infty(x_r \beta_6 \sigma'') = k_r x_r \Sigma^\infty(\sigma'') = 4x_r k_r \sigma$ where k_r is odd. So $4x_r k_r = 2^{r+1} \in \mathbb{Z}_{16}$, i.e., $x_r = \pm 2^{r-1} \in \mathbb{Z}_4$. We get $\partial_{13*}^{7,r}(\sigma') = \pm 2^{r-1} \beta_6 \sigma''$. So

$$0 \rightarrow \mathbb{Z}_{2^{m^2_{r-1}}} \{\beta_6 \sigma''\} \oplus \mathbb{Z}_2 \{\widehat{\eta}_{12}\} \xrightarrow{\tau_{7*}} \pi_{13}(P^7(2^r)) \xrightarrow{p_{7*}} \mathbb{Z}_2 \{v_7^2\} \rightarrow 0.$$

There is $-\mathbf{x} \in \{i_6, 2^r \iota_6, v_6^2\}$ such that $p_{7*}(\mathbf{x}) = v_7^2$ and $2\mathbf{x} \in i_6 \{2^r \iota_6, v_6^2, 2\iota_{12}\}$. Note that $(2^r \iota_6) \sigma'' = 2^r \sigma'' \pm \binom{2^r}{2} [\iota_6, \iota_6] H_2(\sigma'') = 2^r \sigma'' \pm \binom{2^r}{2} \Delta(\iota_{13}) \eta_{11}^2 = 2^r \sigma''$. Then

$$\{2^r \iota_6, v_6^2, 2\iota_{12}\} \ni 2^{r-1} v_6^2 \eta_{12} = 0 \pmod{\langle 2\sigma'' \rangle}.$$

So $2\mathbf{x} = 2ti_6 \sigma''$. This implies $\pi_{13}(P^7(2^r)) \xrightarrow{p_{7*}} \mathbb{Z}_2 \{v_7^2\}$ is split onto. Thus

$$\pi_{13}(P^7(2^r)) \cong \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_{2^{m^2_{r-1}}}, r \geq 1.$$

- $\pi_{14}(P^7(2^r))$.

Consider diagram (6) for $k = 6, m = 14$ implies the exact sequence

$$\mathbb{Z}_2 \{\sigma' \eta_{14}\} \oplus \mathbb{Z}_2 \{\bar{v}_7\} \oplus \mathbb{Z}_2 \{\varepsilon_7\} = \pi_{15}(S^7) \xrightarrow{\partial_{14*}^{7,r}} \pi_{14}(F_7) \xrightarrow{\tau_{7*}} \pi_{14}(P^7(2^r)) \xrightarrow{p_{7*}} \mathbb{Z}_{2^{m^3_r}} \{\delta_r \sigma'\} \rightarrow 0.$$

$\partial_{14*}^{7,r}(\varepsilon_7) = 0$ and $\partial_{14*}^{7,r}(\bar{v}_7) = \beta_6(2^r \iota_6) \bar{v}_6 = 2^{2r} \beta_6 \bar{v}_6 = 0$ (Lemma 8.1). By Lemma 3.2,

$$\begin{aligned} \partial_{14*}^{7,r}(\sigma' \eta_{14}) &= \partial_{13*}^{7,r}(\sigma') \eta_{13} = \pm 2^{r-1} \beta_6 \sigma'' \eta_{13} = 2^{r+1} \beta_6 \bar{v}_6 = 0 \quad (\sigma'' \eta_{13} = 4\bar{v}_6 \text{ by [39, (7.4)]}). \\ \Rightarrow \text{Coker}(\partial_{14*}^{7,r}) &= \pi_{14}(F_7) = \mathbb{Z}_{2^{m^3_{r+1}}} \{\beta_6 \bar{v}_6\} \oplus \mathbb{Z}_2 \{\beta_6 \varepsilon_6\} \oplus \mathbb{Z}_2 \{\widehat{\eta}_{12} \eta_{13}\}. \end{aligned}$$

For $r \leq 2$, $\delta_r \sigma' = 2^{2-r} \Sigma \sigma''$. There is $-\mathbf{x}_r \in \{i_6, 2^r \iota_6, 2^{2-r} \sigma''\}$ such that $p_{7*}(\mathbf{x}_r) = \delta_r \sigma'$ and $2^r \mathbf{x}_r \in i_6 \{2^r \iota_6, 2^{2-r} \sigma'', 2^r \iota_{13}\}$. By (vii) of Lemma 8.1, $2^r \mathbf{x}_r = 2^r t_r i_6 \bar{v}_6$ for some integer t_r . Then $\mathbf{x}_r - t_r i_6 \bar{v}_6$ is the order 2^r lift of $\delta_r \sigma'$ by p_7 , i.e., $\pi_{14}(P^7(2^r)) \xrightarrow{p_{7*}} \mathbb{Z}_{2^{m^3_r}} \{\delta_r \sigma'\}$ is split onto for $r \leq 2$.

For $r \geq 3$, the diagram (10) for $k = 6, s = r, t = 2$ induces the following diagram

$$\begin{array}{ccc} \mathbb{Z}_8 \{\beta_6^r \bar{v}_6\} \hookrightarrow \pi_{14}(J_{2,6}^r) \cong \pi_{14}(F_7^r) & \xrightarrow{\tau_{7*}^r} & \pi_{14}(P^7(2^r)) \\ \parallel & \downarrow \chi_{2*}^r & \downarrow \bar{\chi}_{2*}^r \\ \mathbb{Z}_8 \{\beta_6^2 \bar{v}_6\} \hookrightarrow \pi_{14}(J_{2,6}^2) \cong \pi_{14}(F_7^2) & \xrightarrow{\tau_{7*}^2} & \pi_{14}(P^7(4)) \end{array}$$

Since $(\tau_{7*}^2)|_{res} : \mathbb{Z}_8 \{\beta_6^2 \bar{v}_6\} \hookrightarrow \pi_{14}(P^7(4))$ is split into, so is $(\tau_{7*}^r)|_{res} : \mathbb{Z}_8 \{\beta_6^r \bar{v}_6\} \hookrightarrow \pi_{14}(P^7(2^r))$, i.e., there is an epimorphism $\pi_{14}(P^7(2^r)) \xrightarrow{P_r} \mathbb{Z}_8 \{\beta_6^r \bar{v}_6\}$ such that $P_r(\tau_{7*}^r)|_{res} = id$, i.e., $P_r(i_6^r \bar{v}_6) = \beta_6^r \bar{v}_6$. Let $\mathbf{y}_r \in \pi_{14}(P^7(2^r))$ such that $p_{7*}(\mathbf{y}_r) = \sigma'$. Note that the

order 4 element $\bar{\chi}_r^2 \mathbf{x}_2$ satisfies $p_{7*}(\bar{\chi}_r^2 \mathbf{x}_2) = 2\sigma'$, where $\mathbf{x}_2 \in \pi_{14}(P^7(4))$ has been given before which satisfies $p_{7*}(\mathbf{x}_2) = 2\sigma'$. Thus $p_{7*}(2\mathbf{y}_r - \bar{\chi}_r^2 \mathbf{x}_2) = 0$, i.e., $\text{Ker}(p_{7*}) \ni 2\mathbf{y}_r - \bar{\chi}_r^2 \mathbf{x}_2 = \tau_{6*}(a\beta_6 \bar{v}_6 + b\beta_6 \varepsilon_6 + c\widehat{\eta_{12}\eta_{13}}) = ai_6 \bar{v}_6 + bi_6 \varepsilon_6 + c\tau_6 \widehat{\eta_{12}\eta_{13}}$, $a \in \mathbb{Z}_8, b, c \in \mathbb{Z}_2$. Hence $8\mathbf{y}_r = 4ai_6 \bar{v}_6$. So $4a\beta_6^r \bar{v}_6 = P_r(4ai_6 \bar{v}_6) = 8P_r(\mathbf{y}_r) = 0 \in \mathbb{Z}_8\{\beta_6^r \bar{v}_6\}$. We get $4a = 0 \in \mathbb{Z}_8$, hence $8\mathbf{y}_r = 0$, which implies $o(\mathbf{y}_r) = 8$. Thus $\pi_{14}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_{2m^3} \{\delta_r \sigma'\}$ is also split onto for $r \geq 3$. Now we have

$$\pi_{14}(P^7(2^r)) \cong \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_{2m^3} \oplus \mathbb{Z}_{2m^3} (r \geq 1).$$

- $\pi_{15}(P^7(2^r))$.

Consider diagram (6) for $k = 6, m = 15$ implies the exact sequence

$$\begin{aligned} \pi_{16}(S^7) &\xrightarrow{\partial_{15*}^{7,r}} \pi_{15}(F_7) \xrightarrow{\tau_{7*}} \pi_{15}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_2\{\sigma' \eta_{14}\} \oplus \mathbb{Z}_2\{\bar{v}_7\} \oplus \mathbb{Z}_2\{\varepsilon_7\} \rightarrow 0, \\ \text{where } \pi_{16}(S^7) &= \mathbb{Z}_2\{\sigma' \eta_{14}^2\} \oplus \mathbb{Z}_2\{v_7^3\} \oplus \mathbb{Z}_2\{\mu_7\} \oplus \mathbb{Z}_2\{\eta_7 \varepsilon_8\} \text{ and} \\ \partial_{15*}^{7,r}(v_7^3) &= \partial_{15*}^{7,r}(\mu_7) = \partial_{15*}^{7,r}(\eta_7 \varepsilon_8) = 0; \partial_{15*}^{7,r}(\sigma' \eta_{14}^2) = \partial_{14*}^{7,r}(\sigma' \eta_{14}) \eta_{14} = 0. \\ &\Rightarrow 0 \rightarrow \pi_{15}(F_7) \xrightarrow{\tau_{7*}} \pi_{15}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_2\{\sigma' \eta_{14}\} \oplus \mathbb{Z}_2\{\bar{v}_7\} \oplus \mathbb{Z}_2\{\varepsilon_7\} \rightarrow 0. \end{aligned} \quad (41)$$

For $r = 1$, $\pi_{15}(P^7(2)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_4^{\oplus 2} \oplus \mathbb{Z}_8$ [43, Theorem 5.13].

Corollary 3.4 implies the commutative diagram

$$\begin{array}{ccccc} \pi_{15}(J'_{2,6}) \cong \pi_{15}(F_7) & \xrightarrow{\tau_{7*}} & \pi_{15}(P^7(2^r)) & \xrightarrow{P_{7*}} & \pi_{15}(S^7) \\ \downarrow E^\infty & & \downarrow \Sigma^\infty & & \downarrow \Sigma^\infty \\ \mathbb{Z}_2\{v_3\} \oplus \mathbb{Z}_2\{\mu\} \oplus \mathbb{Z}_2\{\eta \varepsilon\} = \pi_{15}^s(S^6) & \xrightarrow{i_*} & \pi_{15}^s(P^7(2^r)) & \xrightarrow{P_*} & \pi_{15}^s(S^7) = \mathbb{Z}_2\{v\} \oplus \mathbb{Z}_2\{\varepsilon\} \end{array} \quad (42)$$

For $r \geq 2$, By Lemma 3.8, it is easy to get the bottom sequence is split. By the right commutative square in Lemma 3.3, we get $\mathbb{Z}_2\{\beta_6 v_6^3\} \oplus \mathbb{Z}_2\{\beta_6 \mu_6\} \oplus \mathbb{Z}_2\{\beta_6 \eta_6 \varepsilon_6\} \xrightarrow{(\tau_{7*})|_{res}} \pi_{15}(P^7(2^r))$ is split into. By the first proposition given in [2], we get $2^{r+1}\pi_{15}(P^7(2^r)) = 0 (r \geq 2)$. Thus for $r = 2$, $8\pi_{15}(P^7(4)) = 0$ implies that $\mathbb{Z}_8\{\widehat{\delta_{r+1} v_{12}}\} \xrightarrow{(\tau_{7*})|_{res}} \pi_{15}(P^7(2^r))$ is split into. So the sequence (41) is split. For $r \geq 3$, the diagram (9) for $s = r, t = 2$ induces the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_{15}(F_7^2) & \xrightarrow{\tau_{7*}^2} & \pi_{15}(P^7(4)) & \longrightarrow & \pi_{15}(S^7) \longrightarrow 0 \\ & & \downarrow \chi_{r*}^2 & & \downarrow \bar{\chi}_r^2 & & \parallel \\ 0 & \longrightarrow & \pi_{15}(F_7^r) & \xrightarrow{\tau_{7*}^r} & \pi_{15}(P^7(2^r)) & \longrightarrow & \pi_{15}(S^7) \longrightarrow 0 \end{array}$$

Since the top sequence in the above diagram is split, so is the bottom one. Now we have

$$\pi_{15}(P^7(2^r)) \cong \mathbb{Z}_2^{\oplus 6} \oplus \mathbb{Z}_{2m^3}, r \geq 2.$$

- $\pi_{16}(P^7(2^r))$.

Consider diagram (6) for $k = 6, m = 16$ implies the exact sequence

$$0 \rightarrow \text{Coker}(\partial_{16*}^{7,r}) \xrightarrow{\tau_{7*}} \pi_{16}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_2\{\sigma'\eta_{14}^2\} \oplus \mathbb{Z}_2\{v_7^3\} \oplus \mathbb{Z}_2\{\mu_7\} \oplus \mathbb{Z}_2\{\eta_7\varepsilon_8\} \rightarrow 0$$

$$\text{Coker}(\partial_{16*}^{7,r}) = \mathbb{Z}_{2m_r^3}\{\beta_6 v_6 \sigma_9\} \oplus \mathbb{Z}_2\{\beta_6 \eta_6 \mu_7\}.$$

Since $\pi_{15}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_2\{\sigma'\eta_{14}\} \oplus \mathbb{Z}_2\{\bar{v}_7\} \oplus \mathbb{Z}_2\{\varepsilon_7\}$ is an epimorphism, let \mathbf{x}, \mathbf{y} and $\mathbf{z} \in \pi_{15}(P^7(2^r))$ be the lifts of $\sigma'\eta_{14}, \bar{v}_7$ and ε_7 by p_7 respectively. By $\bar{v}_7 \eta_{15} = v_7^3$ and $\varepsilon_7 \eta_{15} = \eta_7 \varepsilon_8$ [39, (7.3), (7.5)], $\mathbf{x}\eta_{15}, \mathbf{y}\eta_{15}$ and $\mathbf{z}\eta_{15} \in \pi_{15}(P^7(2^r))$ are the lifts of $\sigma'\eta_{14}^2, v_7^3$ and $\eta_7 \varepsilon_8$ respectively. For μ_7 , there is $-\mathbf{x}_r^\mu \in \{i_6, 2^r \iota_6, \mu_6\}$ such that $p_{7*}(\mathbf{x}_r^\mu) = \mu_7$ and $2\mathbf{x}_r^\mu \in i_6\{2^r \iota_6, \mu_6, 2\iota_{15}\}$, where $\{2^r \iota_6, \mu_6, 2\iota_{15}\} \ni 2^{r-1} \mu_6 \eta_{15} = 2^{r-1} \eta_6 \mu_7 \pmod{(2^r \iota_6) \circ \pi_{16}(S^6) + \pi_{16}(S^6) \circ (2\iota_{15})} = \langle 2v_6 \sigma_9 \rangle$. Hence there are integers t_r such that $2\mathbf{x}_1^\mu = i_6 \eta_6 \mu_7 + 2t_1 i_6 v_6 \sigma_9$ and $2\mathbf{x}_r^\mu = 2t_r i_6 v_6 \sigma_9, r \geq 2$. Then $\mathbf{y}_r^\mu := \mathbf{x}_r^\mu - t_r i_6 v_6 \sigma_9$ is a lift of μ_7 by p_7 with $o(\mathbf{y}_1^\mu) = 4$ and $o(\mathbf{y}_r^\mu) = 2(r \geq 2)$. Hence

$$\pi_{16}(P^7(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 4} \oplus \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2^{\oplus 5} \oplus \mathbb{Z}_{2m_r^3}, & r \geq 2. \end{cases}$$

- $\pi_{17}(P^7(2^r))$.

By diagram (6) for $k = 6, m = 17$ and $\pi_{18}(S^7) = \mathbb{Z}_8\{\zeta_7\} \oplus \mathbb{Z}_2\{\bar{v}_7 v_{15}\}$

$$\partial_{16*}^{7,r}(\zeta_7) = \beta_6(2^r \iota_6)(\zeta_6) = 2^r \beta_6 \zeta_6;$$

$$\partial_{16*}^{7,r}(\bar{v}_7 v_{15}) = \beta_6(2^r \iota_6)(\bar{v}_6 v_{14}) = \beta_6(2^{2r} \bar{v}_6) v_{14} = 0 \text{ by Lemma 8.1.}$$

$$\Rightarrow 0 \rightarrow \text{Coker}(\partial_{17*}^{7,r}) \xrightarrow{\tau_{7*}} \pi_{17}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_{2m_r^3}\{\delta_r v_7 \sigma_{10}\} \oplus \mathbb{Z}_2\{\eta_7 \mu_8\} \rightarrow 0$$

where $\text{Coker}(\partial_{17*}^{7,r}) \cong \mathbb{Z}_{2m_r^3} \oplus \mathbb{Z}_{2m_r^3} \oplus \mathbb{Z}_4$, by Corollary 6.3.

Since $\pi_{16}(P^6(2^r)) \xrightarrow{P_{6*}} \mathbb{Z}_{2m_r^3}\{\delta_r v_6 \sigma_9\} \oplus \mathbb{Z}_2\{\eta_6 \mu_7\}$ is split onto, so is the epimorphism $\pi_{17}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_{2m_r^3}\{\delta_r v_7 \sigma_{10}\} \oplus \mathbb{Z}_2\{\eta_7 \mu_8\}$ by Lemma 3.6. Hence

$$\pi_{17}(P^7(2^r)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_4 \oplus \mathbb{Z}_{2m_r^3}^{\oplus 3}, r \geq 1.$$

- $\pi_{18}(P^7(2^r))$.

By diagram (6) for $k = 6, m = 18$ and $\pi_{19}(S^7) = 0$, we get

$$0 \rightarrow \pi_{18}(J_{3,6}^r) \cong \pi_{18}(F_7) \xrightarrow{\tau_{7*}} \pi_{17}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_{2m_r^3}\{\zeta_7\} \oplus \mathbb{Z}_2\{v_7 v_{15}\} \rightarrow 0$$

Since $\pi_{17}(P^6(2^r)) \xrightarrow{q_{1P_{6*}}} \mathbb{Z}_{2m_r^3}\{\delta_r \zeta_6\}$ is split onto, there is $\widetilde{\delta_r \zeta_6} \in \pi_{17}(P^6(2^r))$ with $o(\widetilde{\delta_r \zeta_6}) = 2m_r^3$ such that $p_{6*}(\widetilde{\delta_r \zeta_6}) = \delta_r \zeta_6$. Then $\Sigma \widetilde{\delta_r \zeta_6}$ is an order $2m_r^3$ element such that $p_{7*}(\Sigma \widetilde{\delta_r \zeta_6}) = \delta_r \zeta_7$.

For $r = 1$, consider commutative diagram (42) for $r = 1$.

In the bottom exact sequence of (42), by the Toda bracket in stable case [39, Page 32], $\exists \tilde{v} \in \langle i, 2\iota, \bar{v} \rangle$ such that $p_*(\tilde{v}) = \bar{v} \in \pi_{15}^s(S^7)$ and $2\tilde{v} \in i \langle 2\iota, \bar{v}, 2\iota \rangle$. $\langle 2\iota, \bar{v}, 2\iota \rangle \ni \bar{v}\eta = i\nu^3 \pmod{2\pi_9^s(S^0)} = \{0\}$. Thus $2\tilde{v} = \nu^3$ and $o(\tilde{v}) = 4$. Since $\pi_{15}(P^7(2)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_4^{\oplus 2} \oplus \mathbb{Z}_8$ and $\sigma'\eta_{14} \in \pi_{15}(S^7)$ has an order 8 lift in $\pi_{15}(P^7(2))$ by p_7 [43, Page 82], there is an order 4 lift $\tilde{v}_7 \in \pi_{15}(P^7(2))$ of \bar{v}_7 by p_7 . Hence $p_*(\Sigma^\infty \tilde{v}_7) = \bar{v}$, which implies that $\Sigma^\infty \tilde{v}_7 - \tilde{v} \in \text{Ker}(p_*) = \text{Im}(i_*) \cong \mathbb{Z}_2^{\oplus 3}$. So $2\Sigma^\infty \tilde{v}_7 = 2\tilde{v} = i\nu^3$. Since $2\tilde{v}_7 \in \text{Ker}(p_{7*}) = \text{Im}(\tau_{7*})$, assume

$$2\tilde{v}_7 = a_1\tau_7\beta_6\nu_6^3 + a_2\tau_7\beta_6\mu_6 + a_3\tau_7\beta_6\eta_6\varepsilon_7 + 2b\tau_7\overline{\delta_{r+1}\nu_{12}}, a_1, a_2, a_3, b \in \mathbb{Z}_2.$$

$$\text{Then } i\nu^3 = \Sigma^\infty(2\tilde{v}_7) = \Sigma^\infty(a_1\tau_7\beta_6\nu_6^3 + a_2\tau_7\beta_6\mu_6 + a_3\tau_7\beta_6\eta_6\varepsilon_7 + 2b\tau_7\overline{\delta_{r+1}\nu_{12}})$$

$$= a_1i\nu^3 + a_2i\mu + a_3i\eta\varepsilon \text{ (by Corollary 3.4) .}$$

$$\Rightarrow a_1 = 1, a_2 = a_3 = 0. \Rightarrow 2\tilde{v}_7 = i_6\nu_6^3 + 2b\tau_7\overline{\delta_{r+1}\nu_{12}}.$$

Let $\mathbf{v} = \tilde{v}_7 - b\tau_7\overline{\delta_{r+1}\nu_{12}}$. Then $2\mathbf{v} = i_6\nu_6^3$ and $p_{7*}(\mathbf{v}) = \bar{v}_7$. Then $p_{7*}(\mathbf{v}\nu_{15}) = \bar{v}_7\nu_{15}$ and $2\mathbf{v}\nu_{15} = (2\mathbf{v})\nu_{15} = i_6\nu_6^4 = i_6\bar{v}_6\eta_{14}\nu_{15} = 0$ [39, Lemma 6.3, Proposition 5.8].

For $r \geq 2$, from the sequence (41), there is an order 2 element $\tilde{v}_7 \in \pi_{15}(P^7(2^r))$ such that $p_{7*}(\tilde{v}_7) = \bar{v}_7$. Then the order 2 element $\tilde{v}_7\nu_{15} \in \pi_{18}(P^7(2^r))$ satisfies $p_{7*}(\tilde{v}_7\nu_{15}) = \bar{v}_7\nu_{15}$.

Hence $\pi_{17}(P^7(2^r)) \xrightarrow{p_{7*}} \mathbb{Z}_{2m_r^3}\{\zeta_7\} \oplus \mathbb{Z}_2\{\nu_7\nu_{15}\}$ is split onto for $r \geq 1$.

$$\pi_{18}(P^7(2^r)) \cong \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_{2m_r^3} \oplus \mathbb{Z}_{2m_r^4} (r \geq 1).$$

- $\pi_{19}(P^7(2^r))$.

By diagram (6) for $k = 6, m = 19$ and $\pi_{20}(S^7) = \mathbb{Z}_2\{\nu_7\sigma_{10}\nu_{17}\}$, we get

$$\pi_{19}(P^7(2^r)) \cong \pi_{19}(F_7) \cong \pi_{19}(J_{3,6}^r) \cong \begin{cases} \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4, & r = 1; \\ \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_{2m_r^4}, & r \geq 2. \end{cases}$$

6.2. Homotopy groups of $P^8(2^r)$

There are cofibration and fibration sequences

$$S^7 \xrightarrow{2^r\iota_7} S^7 \xrightarrow{i_7} P^8(2^r) \xrightarrow{p_8} S^8; \quad \Omega S^8 \xrightarrow{\partial^8} F_8 \xrightarrow{\tau_8} P^8(2^r) \xrightarrow{p_8} S^8$$

From Lemma 4.4, $Sk_{20}(F_8) \simeq J_{2,7}^r \simeq S^7 \vee S^{14}$, hence here $\beta_7 = j_1^7$ and $\beta_{14} = j_2^{14}$.

$$Sk_{27}(F_8) \simeq J_{3,7}^r \simeq (S^7 \vee S^{14}) \cup_{\gamma_3^r = \pm 2^r [j_1^7, j_2^{14}]} CS^{20}.$$

Lemma 6.5.

- (1) $\pi_{20}(J_{3,7}^r) = \mathbb{Z}_2\{\check{\beta}_7\nu_7\sigma_{10}\nu_{17}\} \oplus \mathbb{Z}_2\{\check{\beta}_{14}\nu_{14}^2\} \oplus \mathbb{Z}_{2^r}\{[\check{\beta}_7, \check{\beta}_{14}]\}$.
- (2) $\pi_{21}(J_{3,7}^r) = \mathbb{Z}_8\{\check{\beta}_7\sigma'\sigma_{14}\} \oplus \mathbb{Z}_4\{\check{\beta}_7\kappa_7\} \oplus \mathbb{Z}_{16}\{\check{\beta}_{14}\sigma_{14}\} \oplus \mathbb{Z}_2\{[\check{\beta}_7, \check{\beta}_{14}]\eta_{20}\}$.

Proof. (1) comes from the isomorphism $\pi_{20}(J'_{2,7})/\langle \gamma_3 \rangle \xrightarrow[\cong]{I'_{2*}} \pi_{20}(J'_{3,7})$.

(2) comes from the following isomorphism by Lemma 4.3 for $m = 21, k = 7$

$$\pi_{21}(S^7 \vee S^{14}) \cong \pi_{21}(J'_{2,7}) \xrightarrow[\cong]{I'_{2*}} \pi_{21}(J'_{3,7}).$$

□

- $\pi_{13}(P^8(2^r)) = \mathbb{Z}_2\{i_7 v_7^2\}$, which is easily obtained by the diagram (6) for $k = 7, m = 13$.

- $\pi_{14}(P^8(2^r))$.

Consider the diagram (6) for $k = 7, m = 14$.

$$\begin{aligned} \mathbb{Z}_{(2)}\{\sigma_8\} \oplus \mathbb{Z}_8\{\Sigma\sigma'\} &= \pi_{15}(S^8) \xrightarrow{\partial_{14*}^{8,r}} \pi_{14}(F_8) \xrightarrow{\tau_{8*}} \pi_{14}(P^8(2^r)) \xrightarrow{P_{8*}} \mathbb{Z}_2\{v_8^2\} \rightarrow 0 \\ \pi_{14}(J'_{2,7}) \cong \pi_{14}(F_8) &= \mathbb{Z}_{(2)}\{\beta_{14}\} \oplus \mathbb{Z}_8\{\beta_7\sigma'\}, \partial_{14*}^{8,r}(\Sigma\sigma') = \beta_7(2^r \iota_7)\sigma' = 2^r \beta_7\sigma'. \end{aligned}$$

By Lemma 5.10, $\partial_{14*}^{8,r}(\sigma_8) = 2^r \beta_{14} + y_r \beta_7\sigma'$, where $\mathbb{Z}_8 \ni y_r =$

$r = 1$	$r = 2$	$r = 3$	$r \geq 4$
odd	± 2	4	0

Thus $Coker(\partial_{14*}^{8,r}) = \begin{cases} \mathbb{Z}_{2^{r+1}}\{\beta_{14}\} \oplus \mathbb{Z}_{2^{r-1}}\{2\beta_{14} + \beta_7\sigma'\}, & r \leq 3; \\ \mathbb{Z}_{2^r}\{\beta_{14}\} \oplus \mathbb{Z}_8\{\beta_7\sigma'\}, & r \geq 4. \end{cases}$

Since $\pi_{13}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_2\{v_7^2\}$ is split onto, so is the epimorphism $\pi_{14}(P^8(2^r)) \xrightarrow{P_{8*}} \mathbb{Z}_2\{v_8^2\}$ by Lemma 3.6. Hence

$$\pi_{14}(P^8(2^r)) \cong \begin{cases} \mathbb{Z}_2 \oplus \mathbb{Z}_{2^{r-1}} \oplus \mathbb{Z}_{2^{r+1}}, & r \leq 3; \\ \mathbb{Z}_2 \oplus \mathbb{Z}_8 \oplus \mathbb{Z}_{2^r}, & r \geq 4. \end{cases}$$

- $\pi_{15}(P^8(2^r))$.

The diagram (6) for $k = 7, m = 15$ implies the exact sequence

$$\pi_{16}(S^8) \xrightarrow{\partial_{15*}^{8,r}} \pi_{15}(F_8) \xrightarrow{\tau_{8*}} \pi_{15}(P^8(2^r)) \xrightarrow{P_{8*}} \mathbb{Z}_{2^{m^3}}\{\delta_r \Sigma\sigma'\} \rightarrow 0$$

where $\pi_{16}(S^8) = \mathbb{Z}_2\{\sigma_8 \eta_{15}\} \oplus \mathbb{Z}_2\{\Sigma\sigma' \eta_{15}\} \oplus \mathbb{Z}_2\{\bar{v}_8\} \oplus \mathbb{Z}_2\{\varepsilon_8\}$,

$$\pi_{15}(F_8) = \mathbb{Z}_2\{\beta_7\sigma' \eta_{14}\} \oplus \mathbb{Z}_2\{\beta_7 \bar{v}_7\} \oplus \mathbb{Z}_2\{\beta_7 \varepsilon_7\} \oplus \mathbb{Z}_2\{\beta_{14} \eta_{14}\},$$

$$\partial_{15*}^{8,r}(\Sigma\sigma' \eta_{15}) = \partial_{15*}^{8,r}(\bar{v}_8) = \partial_{15*}^{8,r}(\varepsilon_8) = 0,$$

$$\partial_{15*}^{8,r}(\sigma_8 \eta_{15}) = \partial_{14*}^{8,r}(\sigma_8) \eta_{14} = (2^r \beta_{14} + y_r \beta_7\sigma') \eta_{14} = 2^{r-1} \beta_7\sigma' \eta_{14}.$$

$$\Rightarrow Coker(\partial_{15*}^{8,r}) = \vartheta_r \mathbb{Z}_2\{\beta_7\sigma' \eta_{14}\} \oplus \mathbb{Z}_2\{\beta_7 \bar{v}_7\} \oplus \mathbb{Z}_2\{\beta_7 \varepsilon_7\} \oplus \mathbb{Z}_2\{\beta_{14} \eta_{14}\}.$$

Since $\pi_{14}(P^7(2^r)) \xrightarrow{P_{7*}} \mathbb{Z}_{2^{m^3}}\{\delta_r \sigma'\}$ is split onto, so is the epimorphism $\pi_{15}(P^8(2^r)) \xrightarrow{P_{8*}} \mathbb{Z}_{2^{m^3}}\{\delta_r \Sigma\sigma'\}$ by Lemma 3.6. Hence

$$\pi_{15}(P^8(2^r)) \cong \vartheta_r \mathbb{Z}_2 \oplus \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_{2^{m^3}}, r \geq 1.$$

- $\pi_{16}(P^8(2^r))$.

Consider the diagram (6) for $k = 7, m = 16$.

$$\begin{aligned}
\pi_{17}(S^8) &\xrightarrow{\partial_{16*}^{8,r}} \pi_{16}(F_8) \xrightarrow{\tau_{8*}} \pi_{16}(P^8(2^r)) \xrightarrow{p_{8*}} \vartheta_r \mathbb{Z}_2\{\sigma_8\eta_{15}\} \oplus \mathbb{Z}_2\{\Sigma\sigma'\eta_{15}\} \oplus \mathbb{Z}_2\{\bar{v}_8\} \oplus \mathbb{Z}_2\{\varepsilon_8\} \rightarrow 0 \\
\pi_{17}(S^8) &= \mathbb{Z}_2\{\sigma_8\eta_{15}^2\} \oplus \mathbb{Z}_2\{\Sigma\sigma'\eta_{15}^2\} \oplus \mathbb{Z}_2\{v_8^3\} \oplus \mathbb{Z}_2\{\mu_8\} \oplus \mathbb{Z}_2\{\eta_8\varepsilon_9\}, \\
\pi_{16}(J_{2,7}^r) &\cong \pi_{16}(F_8) = \mathbb{Z}_2\{\beta_7\sigma'\eta_{14}^2\} \oplus \mathbb{Z}_2\{\beta_7v_7^3\} \oplus \mathbb{Z}_2\{\beta_7\mu_7\} \oplus \mathbb{Z}_2\{\beta_7\eta_7\varepsilon_8\} \oplus \mathbb{Z}_2\{\beta_{14}\eta_{14}^2\} \\
\partial_{16*}^{8,r}(\Sigma\sigma'\eta_{15}^2) &= \partial_{16*}^{8,r}(v_8^3) = \partial_{16*}^{8,r}(\mu_8) = \partial_{16*}^{8,r}(\eta_8\varepsilon_9) = 0, \\
\partial_{16*}^{8,r}(\sigma_8\eta_{15}^2) &= \partial_{15*}^{8,r}(\sigma_8\eta_{15})\eta_{15} = 2^{r-1}\beta_7\sigma'\eta_{14}^2. \\
\Rightarrow \text{Coker}(\partial_{16*}^{8,r}) &= \vartheta_r \mathbb{Z}_2\{\beta_7\sigma'\eta_{14}^2\} \oplus \mathbb{Z}_2\{\beta_7v_7^3\} \oplus \mathbb{Z}_2\{\beta_7\mu_7\} \oplus \mathbb{Z}_2\{\beta_7\eta_7\varepsilon_8\} \oplus \mathbb{Z}_2\{\beta_{14}\eta_{14}^2\}.
\end{aligned}$$

For $\alpha = \sigma'\eta_{14}, \bar{v}_7, \varepsilon_7$, there is $-\mathbf{x} \in \{i_7, 2^r\iota_7, \alpha\}$ such that $p_{8*}(\mathbf{x}) = \Sigma\alpha$ and $2\mathbf{x} \in \{2^r\iota_7, \alpha, 2\iota_{15}\}$ which is a coset of subgroup $(2^r\iota_7) \circ \pi_{16}(S^7) + \pi_{16}(S^7) \circ (2^r\iota_{16}) = \{0\}$.

For $\alpha = \sigma'\eta_{14}$, since $i_7\sigma'\eta_{14}^2 = 0$ for $r = 1$, by Lemma 8.2 (viii), $2\mathbf{x} = 0$ for $r \geq 1$.

For $\alpha = \bar{v}_7$, since $\pi_{17}(S^8) \cong \mathbb{Z}_2^{\oplus 5}$, by [39, Proposition 1.3, Lemma 6.3] and Lemma 3.9

$$\Sigma\{2^r\iota_7, \bar{v}_7, 2\iota_{15}\} = -\Sigma\{2^r\iota_7, \bar{v}_7, 2\iota_{15}\} \supset \{2^r\iota_8, \bar{v}_8, 2\iota_{16}\}_1 \ni \begin{cases} \bar{v}_8\eta_{16} = v_8^3 \pmod{\langle \Sigma\sigma'\eta_{15}^2 \rangle}, & r = 1; \\ 0 \pmod{\{0\}}, & r \geq 2. \end{cases}$$

The injection $\Sigma : \pi_{16}(S^7) \rightarrow \pi_{17}(S^8)$ implies that $2\mathbf{x} = i_7v_7^3$ for $r = 1$ and $2\mathbf{x} = 0$ for $r \geq 2$.

For $\alpha = \varepsilon_7$, $\{2^r\iota_7, \varepsilon_7, 2\iota_{15}\} \ni 2^{r-1}\eta_7\varepsilon_8 \pmod{\{0\}}$. Hence $2\mathbf{x} = i_7\eta_7\varepsilon_8$ for $r = 1$ and $2\mathbf{x} = 0$ for $r \geq 2$.

For $\sigma_8\eta_{15} \in \text{Ker}(\partial_{15*}^{8,r})$ with $r \geq 2$, by the same proof as that of Lemma 5.2, we have the following lemma

Lemma 6.6. *There is a $\tilde{\sigma}_8 \in [P^{15}(2^{r-1}), P^8(2^r)]$ such that the following square commutes*

$$\begin{array}{ccc}
P^{15}(2^{r-1}) & \xrightarrow{p_{15}^{r-1}} & S^{15} \\
\downarrow \tilde{\sigma}_8 & & \downarrow \sigma_8 \\
P^8(2^r) & \xrightarrow{p_8^r} & S^8
\end{array}$$

Moreover, $p_{8*}(\tilde{\sigma}_8\widetilde{\eta}_{15}) = \sigma_8\eta_{15}$ where $\widetilde{\eta}_{15} \in \pi_{16}(P^{15}(2^{r-1}))$ is a lift of η_{15} by p_{15}^{r-1} and $o(\tilde{\sigma}_8\widetilde{\eta}_{15}) = 4$ and 2 for $r = 2$ and $r \geq 3$ respectively.

Now we get

$$\pi_{16}(P^8(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_4^{\oplus 2}, & r = 1; \\ \mathbb{Z}_2^{\oplus 7} \oplus \mathbb{Z}_4, & r = 2; \\ \mathbb{Z}_2^{\oplus 9}, & r \geq 3; \end{cases}$$

- $\pi_{17}(P^8(2^r))$.

The diagram (6) for $k = 7, m = 17$ implies the exact sequence

$$\begin{aligned} \pi_{18}(S^8) &\xrightarrow{\partial_{17*}^{8,r}} \pi_{17}(F_8) \xrightarrow{\tau_{8*}} \pi_{17}(P^8(2^r)) \xrightarrow{p_{8*}} \text{Ker}(\partial_{16*}^{8,r}) \rightarrow 0 \text{ where} \\ \text{Ker}(\partial_{16*}^{8,r}) &= \vartheta_r \mathbb{Z}_2\{\sigma_8 \eta_{15}^2\} \oplus \mathbb{Z}_2\{\Sigma \sigma' \eta_{15}^2\} \oplus \mathbb{Z}_2\{v_8^3\} \oplus \mathbb{Z}_2\{\mu_8\} \oplus \mathbb{Z}_2\{\eta_8 \varepsilon_9\}; \\ \pi_{18}(S^8) &= \mathbb{Z}_8\{\sigma_8 v_{15}\} \oplus \mathbb{Z}_8\{v_8 \sigma_{11}\} \oplus \mathbb{Z}_2\{\eta_8 \mu_9\}; \\ \pi_{17}(F_8) &= \mathbb{Z}_8\{\beta_7 v_7 \sigma_{10}\} \oplus \mathbb{Z}_2\{\beta_7 \eta_7 \mu_8\} \oplus \mathbb{Z}_8\{\beta_{14} v_{14}\}. \\ \partial_{17*}^{8,r}(v_8 \sigma_{11}) &= 2^r \beta_7 v_7 \sigma_{10}, \quad \partial_{17*}^{8,r}(\eta_8 \mu_9) = 0. \end{aligned}$$

By [39, (7.19)], $\sigma' v_{14} = x v_7 \sigma_{10}$, x is odd integer.

$$\begin{aligned} \partial_{17*}^{8,r}(\sigma_8 v_{15}) &= \partial_{14*}^{8,r}(\sigma_8) v_{14} = (2^r \beta_{14} + y_r \beta_7 \sigma') v_{14} = 2^r \beta_{14} v_{14} + x y_r \beta_7 v_7 \sigma_{10}. \\ \Rightarrow \text{Coker}(\partial_{17*}^{8,r}) &= \begin{cases} \mathbb{Z}_2\{\beta_7 \eta_7 \mu_8\} \oplus \mathbb{Z}_4\{\beta_{14} v_{14}\}, & r = 1; \\ \mathbb{Z}_2\{\beta_7 v_7 \sigma_{10} + 2\beta_{14} v_{14}\} \oplus \mathbb{Z}_2\{\beta_7 \eta_7 \mu_8\} \oplus \mathbb{Z}_8\{\beta_{14} v_{14}\}, & r = 2; \\ \mathbb{Z}_{2^{m^3-1}}\{\beta_7 v_7 \sigma_{10}\} \oplus \mathbb{Z}_2\{\beta_7 \eta_7 \mu_8\} \oplus \mathbb{Z}_8\{\beta_{14} v_{14}\}, & r \geq 3. \end{cases} \end{aligned}$$

i.e. $\text{Coker}(\partial_{17*}^{8,r}) \cong \mathbb{Z}_{2^{m^3-1}} \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_{2^{m^3+1}}$, $r \geq 1$.

Since $\pi_{16}(P^8(2^r)) \xrightarrow{p_{8*}} \vartheta_r \mathbb{Z}_2\{\sigma_8 \eta_{15}\} \oplus \mathbb{Z}_2\{\Sigma \sigma' \eta_{15}\} \oplus \mathbb{Z}_2\{\bar{v}_8\} \oplus \mathbb{Z}_2\{\varepsilon_8\}$ is epimorphic, let $\tilde{\alpha}$ be the lift of $\alpha = \sigma_8 \eta_{15}$ ($r \geq 2$), $\Sigma \sigma' \eta_{15}$, \bar{v}_8 , ε_8 by p_8 . Then the order 2 element $\tilde{\alpha} \eta_{16}$ is the lift of $\sigma_8 \eta_{15}^2$, $\Sigma \sigma' \eta_{15}^2$, v_8^3 , $\eta_8 \varepsilon_9$ by p_8 .

For μ_7 , there is $-\mathbf{x}_r^\mu \in \{i_7, 2^r \iota_7, \mu_7\}$ such that $p_{8*}(\mathbf{x}_r^\mu) = \mu_8$ and $2\mathbf{x}_r^\mu \in i_7\{2^r \iota_7, \mu_7, 2\iota_{16}\}$, where $\{2^r \iota_7, \mu_7, 2\iota_{17}\} \ni 2^{r-1} \mu_7 \eta_{16} = 2^{r-1} \eta_7 \mu_8 \pmod{(2^r \iota_7) \circ \pi_{17}(S^7) + \pi_{17}(S^7) \circ (2\iota_{16})} = \langle 2v_7 \sigma_{10} \rangle$. Hence there are integers t_r such that $2\mathbf{x}_1^\mu = i_7 \eta_7 \mu_8 + 2t_1 i_7 v_7 \sigma_{10}$ and $2\mathbf{x}_r^\mu = 2t_r i_7 v_7 \sigma_{10}$, $r \geq 2$. Then $\mathbf{y}_r^\mu := \mathbf{x}_r^\mu \mathbf{x}_r^\mu - t_r i_7 v_7 \sigma_{10}$ is a lift of μ_8 by p_8 with $o(\mathbf{y}_1^\mu) = 4$ and $o(\mathbf{y}_r^\mu) = 2$ ($r \geq 2$). Hence

$$\pi_{17}(P^8(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_4^{\oplus 2}, & r = 1; \\ \mathbb{Z}_2^{\oplus 6} \oplus \mathbb{Z}_{2^{m^3-1}} \oplus \mathbb{Z}_8, & r \geq 2. \end{cases}$$

• $\pi_{18}(P^8(2^r))$.

The diagram (6) for $k = 7, m = 18$ implies the exact sequence

$$\begin{aligned} \mathbb{Z}_8\{\zeta_8\} \oplus \mathbb{Z}_2\{\bar{v}_8 v_{16}\} &= \pi_{19}(S^8) \xrightarrow{\partial_{18*}^{8,r}} \pi_{18}(F_8) \xrightarrow{\tau_{8*}} \pi_{18}(P^8(2^r)) \xrightarrow{p_{8*}} \text{Ker}(\partial_{17*}^{8,r}) \rightarrow 0 \text{ where} \\ \text{Ker}(\partial_{17*}^{8,r}) &= \begin{cases} \mathbb{Z}_4\{4\sigma_8 v_{15} + 2v_8 \sigma_{11}\} \oplus \mathbb{Z}_2\{\eta_8 \mu_9\}, & r = 1; \\ \mathbb{Z}_8\{2\sigma_8 v_{15} + v_8 \sigma_{11}\} \oplus \mathbb{Z}_2\{4\sigma_8 v_{15}\} \oplus \mathbb{Z}_2\{\eta_8 \mu_9\}, & r = 2; \\ \mathbb{Z}_8\{v_8 \sigma_{11}\} \oplus \mathbb{Z}_4\{2\sigma_8 v_{15}\} \oplus \mathbb{Z}_2\{\eta_8 \mu_9\}, & r = 3; \\ \mathbb{Z}_8\{v_8 \sigma_{11}\} \oplus \mathbb{Z}_8\{\sigma_8 v_{15}\} \oplus \mathbb{Z}_2\{\eta_8 \mu_9\}, & r \geq 4. \end{cases} \\ \pi_{18}(F_8) &= \mathbb{Z}_8\{\beta_7 \zeta_7\} \oplus \mathbb{Z}_2\{\beta_7 \bar{v}_7 v_{15}\}. \quad \partial_{18*}^{8,r}(\zeta_8) = 2^r \beta_7 \zeta_7, \quad \partial_{18*}^{8,r}(\bar{v}_8 v_{16}) = 0. \\ \Rightarrow 0 \rightarrow \text{Coker}(\partial_{18*}^{8,r}) &= \mathbb{Z}_{2^{m^3}}\{\beta_7 \zeta_7\} \oplus \mathbb{Z}_2\{\beta_7 \bar{v}_7 v_{15}\} \xrightarrow{\tau_{8*}} \pi_{18}(P^8(2^r)) \xrightarrow{p_{8*}} \text{Ker}(\partial_{17*}^{8,r}) \rightarrow 0. \end{aligned}$$

In order to solve the extension problem of the above short exact sequence, we firstly compute the $\pi_{20}(P^{10}(2^r))$.

From (i) of Lemma 4.4, $Sk_{26}(F_{10}) \simeq J_{2,9}^r \simeq S^9 \vee S^{18}$. Then by the diagram (6) for $k = 9, m = 20$, we get the short exact sequence

$$\mathbb{Z}_{2^{m_3}}\{\beta_9\zeta_9\} \oplus \mathbb{Z}_2\{\beta_9\bar{\nu}_9\nu_{17}\} \oplus \mathbb{Z}_2\{\beta_{18}\eta_{18}^2\} \xrightarrow{\tau_{10^*}^{10}} \pi_{20}(P^{10}(2^r)) \xrightarrow{P^{10^*}} \mathbb{Z}_{2^{r-1}}\{\delta_r\sigma_{10}\nu_{11}\} \oplus \mathbb{Z}_2\{\eta_{10}\mu_{11}\}.$$

We have the following commutative squares

$$\begin{array}{ccc} \mathbb{Z}_2\{\beta_9^r\bar{\nu}_9\nu_{17}\} \xrightarrow{(\tau_{10^*}^r)|_{res}} \pi_{20}(P^{10}(2^r)) & ; & \mathbb{Z}_{2^{m_3}}\{\beta_9^r\zeta_9\} \xrightarrow{(\tau_{10^*}^r)|_{res}} \pi_{20}(P^{10}(2^r)) \\ \parallel & \downarrow \bar{\chi}_{1^*} & \downarrow \cong \\ \mathbb{Z}_2\{\beta_9^1\bar{\nu}_9\nu_{17}\} \xrightarrow{(\tau_{10^*}^1)|_{res}} \pi_{20}(P^{10}(2)) & & \mathbb{Z}_{2^{m_3}}\{\zeta\} \xrightarrow{i_*} \pi_{20}^s(P^{10}(2^r)) \xrightarrow{P_*} \mathbb{Z}_2\{\eta\mu\}. \end{array}$$

where the left commutative square is induced by the diagram (9) for $s = r \geq 2, t = 1$ and the right commutative square is from Corollary 3.4.

For $\eta_8\mu_9 \in Ker(\partial_{17^*}^{8,r})$, there is an order 2 element $\bar{\mu}_8\eta_{17} \in \pi_{18}(P^8(2^r))$ such that $p_{8^*}(\bar{\mu}_8\eta_{17}) = \mu_8\eta_{17} = \eta_8\mu_9$, where $\bar{\mu}_8 \in \pi_{17}(P^8(2^r))$ is the lift of μ_8 by p_8 . Then the order 2 elements $\Sigma^2(\bar{\mu}_8\eta_{17})$ and $\Sigma^\infty(\bar{\mu}_8\eta_{17})$ are the lifts of $\eta_{10}\mu_{11}$ and $\eta\mu$ by p_{10} and p (in stable case) respectively. Hence $\pi_{20}(P^{10}(2)) \xrightarrow{P^{10^*}} \mathbb{Z}_2\{\eta_{10}\mu_{11}\}$ and $\pi_{20}^s(P^{10}(2^r)) \xrightarrow{P_*}$

$\mathbb{Z}_2\{\eta\mu\}$ are split onto, which imply the monomorphisms $\mathbb{Z}_2\{\beta_9\bar{\nu}_9\nu_{17}\} \xrightarrow{(\tau_{10^*}^1)|_{res}} \pi_{20}(P^{10}(2))$ and $\mathbb{Z}_{2^{m_3}}\{\zeta\} \xrightarrow{i_*} \pi_{20}^s(P^{10}(2^r))$ are split into. Hence the monomorphisms

$\mathbb{Z}_2\{\beta_9^r\bar{\nu}_9\nu_{17}\} \xrightarrow{(\tau_{10^*}^r)|_{res}} \pi_{20}(P^{10}(2^r))$ and $\mathbb{Z}_{2^{m_3}}\{\beta_9\zeta_9\} \xrightarrow{(\tau_{10^*}^r)|_{res}} \pi_{20}(P^{10}(2^r))$ are also split into. Consider the following commutative diagram where the left map is isomorphic by Lemma 3.3

$$\begin{array}{ccc} \mathbb{Z}_{2^{m_3}}\{\beta_7\zeta_7\} \oplus \mathbb{Z}_2\{\beta_7\bar{\nu}_7\nu_{15}\} & \xrightarrow{\tau_{8^*}} & \pi_{18}(P^8(2^r)) \\ \downarrow E^2 \cong & & \downarrow \Sigma^2 \\ \mathbb{Z}_{2^{m_3}}\{\beta_9\zeta_9\} \oplus \mathbb{Z}_2\{\beta_9\bar{\nu}_9\nu_{17}\} & \xrightarrow{(\tau_{10^*}^r)|_{res}} & \pi_{20}(P^{10}(2^r)) \end{array}$$

Since the bottom monomorphism is split into, so is the top one. Hence

$$\pi_{18}(P^8(2^r)) \cong \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_{2^{m_3}} \oplus \mathbb{Z}_{2^{m_3}} \oplus \mathbb{Z}_{2^{m_3}}.$$

• $\pi_{19}(P^8(2^r)) \cong Ker(\partial_{18^*}^{8,r}) \cong \mathbb{Z}_{2^{m_3}} \oplus \mathbb{Z}_2$, which is easily obtained by $\pi_{19}(F_8) \cong \pi_{19}(S^7 \vee S^{14}) = 0$.

• $\pi_{20}(P^8(2^r))$

The diagram (6) for $k = 7, m = 20$ implies the exact sequence

$$\mathbb{Z}_2\{\sigma_8\nu_{15}^2\} \oplus \mathbb{Z}_2\{\nu_8\sigma_{11}\nu_{18}\} = \pi_{21}(S^8) \xrightarrow{\partial_{20^*}^{8,r}} \pi_{20}(F_8) \xrightarrow{\tau_{8^*}} \pi_{20}(P^8(2^r)) \xrightarrow{P_{8^*}} \pi_{20}(S^8) = 0$$

where $\pi_{20}(F_8) = \mathbb{Z}_2\{\beta_7\nu_7\sigma_{10}\nu_{17}\} \oplus \mathbb{Z}_2\{\beta_{14}\nu_{14}^2\} \oplus \mathbb{Z}_{2^r}\{[\beta_7, \beta_{14}]\}$;

$\partial_{20*}^{\delta,r}(\sigma_8 v_{15}^2) = \partial_{17*}^{\delta,r}(\sigma_8 v_{15})v_{17} = (2^r \beta_{14} v_{14} + x y_r \beta_7 v_7 \sigma_{10})v_{17} = (1 - \vartheta_r) \beta_7 v_7 \sigma_{10} v_{17}$, with integers x and y_r from the previous computation of $\pi_{14}(P^8(2^r))$ and $\pi_{17}(P^8(2^r))$ respectively; $\partial_{20*}^{\delta,r}(v_8 \sigma_{11} v_{18}) = \beta_7 (2^r \iota_7)(v_7 \sigma_{10} v_{17}) = 0$. Hence

$$\pi_{20}(P^8(2^r)) \cong \text{Coker}(\partial_{20*}^{\delta,r}) \cong \vartheta_r \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_{2^r}.$$

- $\pi_{21}(P^8(2^r))$

The diagram (6) for $k = 7, m = 21$ implies the exact sequence

$$\mathbb{Z}_{16}\{\sigma_8^2\} \oplus \mathbb{Z}_8\{\Sigma \sigma' \sigma_{15}\} \oplus \mathbb{Z}_4\{\kappa_8\} = \pi_{22}(S^8) \xrightarrow{\partial_{21*}^{\delta,r}} \pi_{21}(F_8) \xrightarrow{\tau_{8*}} \pi_{21}(P^8(2^r)) \xrightarrow{p_{8*}} \text{Ker}(\partial_{20*}^{\delta,r}) \rightarrow 0$$

where $\text{Ker}(\partial_{20*}^{\delta,r}) = \vartheta_r \mathbb{Z}_2\{\sigma_8 v_{15}^2\} \oplus \mathbb{Z}_2\{v_8 \sigma_{11} v_{18}\}$;

$$\pi_{21}(J_{3,7}^r) \cong \pi_{21}(F_8) = \mathbb{Z}_8\{\beta_7 \sigma' \sigma_{14}\} \oplus \mathbb{Z}_4\{\beta_7 \kappa_7\} \oplus \mathbb{Z}_{16}\{\beta_{14} \sigma_{14}\} \oplus \mathbb{Z}_2\{[\beta_7, \beta_{14}] \eta_{20}\};$$

$$\partial_{21*}^{\delta,r}(\sigma_8^2) = \partial_{14*}^{\delta,r}(\sigma_8) \sigma_{14} = (2^r \beta_{14} + y_r \beta_7 \sigma') \sigma_{14} = 2^r \beta_{14} \sigma_{14} + y_r \beta_7 \sigma' \sigma_{14};$$

$$\partial_{21*}^{\delta,r}(\Sigma \sigma' \sigma_{15}) = \beta_7 (2^r \iota_7)(\sigma' \sigma_{14}) = 2^r \beta_7 \sigma' \sigma_{14}; \quad \partial_{21*}^{\delta,r}(\kappa_8) = \beta_7 (2^r \iota_7)(\kappa_7) = 2^r \beta_7 \kappa_7.$$

Hence $\text{Coker}(\partial_{21*}^{\delta,r}) = A_r \oplus \mathbb{Z}_{2^m} \{\beta_7 \kappa_7\} \oplus \mathbb{Z}_2\{[\beta_7, \beta_{14}] \eta_{20}\}$ with

$$\mathbb{Z}_{2^{m_{r+1}}} \oplus \mathbb{Z}_{2^{m_{r-1}}} \cong A_r = \begin{cases} \mathbb{Z}_4\{\beta_{14} \sigma_{14}\}, & r = 1; \\ \mathbb{Z}_8\{\beta_{14} \sigma_{14}\} \oplus \mathbb{Z}_2\{2\beta_{14} \sigma_{14} + \beta_7 \sigma' \sigma_{14}\}, & r = 2; \\ \mathbb{Z}_{16}\{\beta_{14} \sigma_{14}\} \oplus \mathbb{Z}_4\{2\beta_{14} \sigma_{14} + \beta_7 \sigma' \sigma_{14}\}, & r = 3; \\ \mathbb{Z}_{16}\{\beta_{14} \sigma_{14}\} \oplus \mathbb{Z}_8\{2\beta_{14} \sigma_{14}\}, & r \geq 4. \end{cases}$$

For $v_8 \sigma_{11} v_{18}$, there is $-\mathbf{x} \in \{i_7, 2^r \iota_7, v_7 \sigma_{10} v_{17}\}$ such that $p_{8*}(\mathbf{x}) = v_8 \sigma_{11} v_{18}$ and $2\mathbf{x} \in i_7\{2^r \iota_7, v_7 \sigma_{10} v_{17}, 2\iota_{20}\}$, where $\{2^r \iota_7, v_7 \sigma_{10} v_{17}, 2\iota_{20}\} \ni 2^{r-1} v_7 \sigma_{10} v_{17} \eta_{20} = 0 \pmod{(2^r \iota_7) \circ \pi_{21}(S^7) + \pi_{21}(S^7) \circ (2\iota_{21})} = 2\pi_{21}(S^7)$ by Lemma 3.9. Hence $2\mathbf{x} = 2i_7 \theta$ for some $\theta \in \pi_{21}(S^7)$. Let $\mathbf{x}' = \mathbf{x} - i_7 \theta$. Then \mathbf{x}' is a lift of $v_8 \sigma_{11} v_{18}$ by p_8 with $o(\mathbf{x}') = 2$.

For $r \geq 2$ and $\sigma_8 v_{15}^2 \in \text{Ker}(\partial_{20*}^{\delta,r})$, the commutative square in Lemma 6.6 induces the following commutative square

$$\begin{array}{ccc} \pi_{21}(P^{15}(2^{r-1})) & \xrightarrow{p_{15*}^{r-1}} & \pi_{21}(S^{15}) = \mathbb{Z}_2\{v_{15}^2\} \longrightarrow 0 \\ \downarrow \tilde{\sigma}_{8*} & & \downarrow \sigma_{8*} \\ \pi_{21}(P^8(2^r)) & \xrightarrow{p_{8*}^r} & \pi_{21}(S^8) \end{array} .$$

Since $\pi_{13}(P^7(2^{r-1})) \xrightarrow{p_{7*}^r} \mathbb{Z}_2\{v_7^2\}$ is split onto, so is the epimorphism $\pi_{21}(P^{15}(2^{r-1})) \xrightarrow{p_{15*}^{r-1}} \mathbb{Z}_2\{v_{15}^2\}$ by Lemma 3.6, i.e., \exists order 2 element $\mathbf{y} \in \pi_{21}(P^{15}(2^{r-1}))$ such that $p_{15*}^{r-1}(\mathbf{y}) = v_{15}^2$. Hence $\tilde{\sigma}_8 \mathbf{y} \in \pi_{21}(P^8(2^r))$ is the lift of $\sigma_8 v_{15}^2$ by p_8^r with $o(\tilde{\sigma}_8 \mathbf{y}) = 2$.

Hence

$$\pi_{21}(P^8(2^r)) \cong \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_{2^m} \oplus \mathbb{Z}_{2^{m_{r-1}}} \oplus \mathbb{Z}_{2^{m_{r+1}}}.$$

Because of the limitation of length, we stop the computation of homotopy groups of Moore spaces here. In the next section we show that we are able to compute γ_3 for some other 2-cell complexes.

7. Some remarks

In the vast majority of cases, the generators of the above homotopy groups of Moore spaces are easily obtained according to the calculation process.

For example, the generators of $\pi_{14}(P^6(2^r))$ are given as follows

$$\pi_{14}(P^6(2)) = \mathbb{Z}_2\{i_5v_5^3\} \oplus \mathbb{Z}_2\{i_5\mu_5\} \oplus \mathbb{Z}_{2^r}\{[i_5, \tau_6\beta_{10}]\} \oplus \mathbb{Z}_2\{\widetilde{4v_6}\} \oplus \mathbb{Z}_4\{\widetilde{\varepsilon}_6\}$$

with $2\widetilde{\varepsilon}_6 = i_5\eta_5\varepsilon_6$. For $r \geq 2$

$$\pi_{14}(P^6(2^r)) = \mathbb{Z}_2\{i_5v_5^3\} \oplus \mathbb{Z}_2\{i_5\mu_5\} \oplus \mathbb{Z}_2\{i_5\eta_5\varepsilon_6\} \oplus \mathbb{Z}_{2^r}\{[i_5, \tau_6\beta_{10}]\} \oplus \mathbb{Z}_{2m_r^3}\{\delta_r\widetilde{v_6}\} \oplus \mathbb{Z}_2\{\widetilde{\varepsilon}_6\}.$$

The following theorem generalizes the results of $\pi_i(P^n(2))$ for $n = 9, 10, 11$ in [43] to $\pi_i(P^n(2^r))$ for any $r \geq 1$. The calculation is not difficult since it only needs the homotopy of $J_2(M_{2^r\iota_{n-1}}, S^{n-1})$ and we omit the proof here.

Theorem 7.1.

$$\pi_{15}(P^9(2^r)) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_{2m_{r-1}^3} \oplus \mathbb{Z}_{2^{r+1}}.$$

$$\pi_{16}(P^9(2^r)) \cong \vartheta_r\mathbb{Z}_2 \oplus \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_{2m_r^4}.$$

$$\pi_{17}(P^9(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_4^{\oplus 2}, & r = 1; \\ \mathbb{Z}_2^{\oplus 7} \oplus \mathbb{Z}_4, & r = 2; \\ \mathbb{Z}_2^{\oplus 9} & r \geq 3. \end{cases}$$

$$\pi_{17}(P^{10}(2^r)) \cong \mathbb{Z}_2^{\oplus 3} \oplus \mathbb{Z}_{2m_r^4}.$$

$$\pi_{18}(P^{10}(2^r)) \cong \begin{cases} \mathbb{Z}_2 \oplus \mathbb{Z}_4^{\oplus 2} \oplus \mathbb{Z}_8, & r = 1; \\ \mathbb{Z}_2^{\oplus 6} \oplus \mathbb{Z}_{2^{r+1}} & r \geq 2. \end{cases}$$

$$\pi_{19}(P^{11}(2^r)) \cong \begin{cases} \mathbb{Z}_2^{\oplus 2} \oplus \mathbb{Z}_4^{\oplus 2}, & r = 1; \\ \mathbb{Z}_2^{\oplus 5} \oplus \mathbb{Z}_{2^r} & r \geq 2. \end{cases}$$

At last, I would like to mention that our methods are also useful to calculate higher dimensional homotopy groups of other 2-cell complexes. For the suspended complex projective plane $\Sigma\mathbb{C}P^2$ and quaternionic projective plane $\Sigma\mathbb{H}P^2$, we have the cofibration sequences

$$S^4 \xrightarrow{\eta_3} S^3 \xrightarrow{i_3^\eta} \Sigma\mathbb{C}P^2 \xrightarrow{p_5^\eta} S^5; \quad S^8 \xrightarrow{\nu_5} S^5 \xrightarrow{i_5^\nu} \Sigma\mathbb{H}P^2 \xrightarrow{p_9^\nu} S^9$$

$J_2(M_{\eta_3}, S^4) \simeq S^3 \cup_{\gamma_2^\eta=[\iota_3, \eta_3]} e^7 \simeq S^3 \vee S^7$; $J_2(M_{\nu_5}, S^8) \simeq S^5 \cup_{\gamma_2^\nu=[\iota_5, \nu_5]} e^{13} \simeq S^5 \vee S^{13}$;
By the following lemma we can get the homotopy types of $Sk_{14}(F_{p_5^\eta})$ and $Sk_{28}(F_{p_9^\nu})$

$$i.e., \quad J_3(M_{\eta_3}, S^4) \simeq (S^3 \vee S^7) \cup_{\gamma_3^\eta} e^{11} \quad \text{and} \quad J_3(M_{\nu_5}, S^8) \simeq (S^5 \vee S^{13}) \cup_{\gamma_3^\nu} e^{21}$$

which enables us to calculate the $\pi_i(\Sigma\mathbb{C}P^2)$ and $\pi_i(\Sigma\mathbb{H}P^2)$ for $i \leq 13$ and 27 respectively if the relevant homotopy groups of spheres are known.

Lemma 7.2. $\gamma_3^\eta = [j_1^3, j_2^7]\eta_9$; $\gamma_3^\nu = a[j_1^5, j_2^{13}]\nu_{17}$, where a is an odd integer.

Proof. We only prove the second conclusion of this lemma, while the first is easier to obtain by the similar proof.

Consider the following commutative diagrams of cofibrations and fibrations

$$\begin{array}{ccc} S^8 & \xrightarrow{\iota_8} & S^8 & \longrightarrow & * & \xrightarrow{p_9^0} & S^9 & ; & \Omega S^9 & \xrightarrow{\partial^{9,0}} & F_9^0 & \longrightarrow & * & \xrightarrow{p_9^0} & S^9 \\ \parallel & & \downarrow \nu_5 & & \downarrow \bar{\chi}_\nu^0 & & \parallel & & \parallel & & \downarrow \chi_\nu^0 & & \downarrow \bar{\chi}_\nu^0 & & \parallel \\ S^8 & \xrightarrow{\nu_5} & S^5 & \longrightarrow & \Sigma \mathbb{H}P^2 & \xrightarrow{p_9^\nu} & S^9 & & \Omega S^9 & \xrightarrow{\partial^{9,\nu}} & F_{p_9^\nu} & \longrightarrow & \Sigma \mathbb{H}P^2 & \xrightarrow{p_9^\nu} & S^9 \end{array}$$

where $\bar{\chi}_\nu^0$ is induced by the left commutative square of the first diagram and the χ_ν^0 is induced by the right commutative square of the second diagram.

Let $\bar{g}_\nu^0 := (\chi_\nu^0)|_{J_2}$, $J_{2,\nu} := J_2(M_{\nu_5}, S^8)$, from Lemma 2.5 and Lemma 2.6, we get

$$\begin{array}{ccc} S^8 & \xrightarrow{\beta_8} & J_{2,8}^0 \simeq S^8 \cup_{2\sigma_8 - \Sigma\sigma'} e^{16} & \xrightarrow{\bar{p}_{16}^0} & S^{16} & ; & S^{23} & \xrightarrow{\gamma_3^0} & J_{2,8}^0 \\ \downarrow \nu_5 & & \downarrow \bar{g}_\nu^0 & & \downarrow \nu_{13} & & \downarrow \nu_{20} & & \downarrow \bar{g}_\nu^0 \\ S^5 & \xrightarrow{j_1^5} & J_{2,\nu} \simeq S^5 \vee S^{13} & \xrightarrow{p_9^\nu} & S^9 & & S^{20} & \xrightarrow{\gamma_3^\nu} & J_{2,\nu} \end{array} \quad (43)$$

$$\Rightarrow \bar{g}_\nu^0 \beta_8 \simeq j_1^5 \nu_5 \quad \text{and} \quad \bar{g}_\nu^0 \gamma_3^0 \simeq \gamma_3^\nu \nu_{20}. \quad (44)$$

From Theorem 1.1 and Lemma 2.4, $\gamma_3^\nu \in [j_1^5, j_1^5 \nu_5, j_1^5 \nu_5]$ which is a coset of the subgroup

$$A_3^\nu := [\pi_{16}(S^5 \vee S^{13}), j_1^5] + [\pi_{13}(S^5 \vee S^{13}), j_1^5 \nu_5] + [\pi_{13}(S^5 \vee S^{13}), j_1^5 \nu_5]$$

By $\pi_{16}(S^5 \vee S^{13}) = \mathbb{Z}_2\{j_1^5 \zeta_5\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \bar{\nu}_8\} \oplus \mathbb{Z}_2\{j_1^5 \nu_5 \bar{\epsilon}_8\} \oplus \mathbb{Z}_8\{j_2^{13} \nu_{13}\}$ and $\pi_{13}(S^5 \vee S^{13}) = \mathbb{Z}_2\{j_1^5 \epsilon_5\} \oplus \mathbb{Z}_{(2)}\{j_2^{13}\}$, we have $A_3^\nu = \langle [j_1^5, j_2^{13}]\nu_{17} \rangle$. Since $\Sigma[\iota_5, \nu_5, \nu_5] = 0$ and $\Sigma : \pi_{20}(S^5) \rightarrow \pi_{21}(S^6)$ is an injection [39, (10.14), (10.15)], by Proposition 2.3,

$$\begin{aligned} [j_1^5, j_1^5 \nu_5, j_1^5 \nu_5] \supset j_1^5[\iota_5, \nu_5, \nu_5] &= \{0\} \Rightarrow [j_1^5, j_1^5 \nu_5, j_1^5 \nu_5] = \langle [j_1^5, j_2^{13}]\nu_{17} \rangle. \\ \Rightarrow \gamma_3^\nu &= a[j_1^5, j_2^{13}]\nu_{17}, \text{ for some } a \in \mathbb{Z}_8. \end{aligned} \quad (45)$$

There is the cofibration sequence $S^{15} \xrightarrow{2\sigma_8 - \Sigma\sigma'} S^8 \xrightarrow{\beta_8} J_{2,8}^0 \xrightarrow{\bar{p}_{16}^0} S^{16}$. From (7), we get $Sk_{24}(F_{p_{16}^0}) \simeq J_{2,8}^{\gamma,0} \simeq S^8 \vee S^{23}$. Then Lemma 4.3 for $k = 8, r = 0, m = 23$ gives

$$\begin{aligned} 0 \rightarrow \text{Coker}(\bar{\partial}_{23^*}^{16,0}) &\xrightarrow{\bar{\tau}_{16}^0} \pi_{23}(J_{2,8}^0) \xrightarrow{\bar{p}_{16^*}^0} \text{Ker}(\bar{\partial}_{22^*}^{16,0}) = \mathbb{Z}_2\{8\sigma_{16}\} \rightarrow 0, \\ \text{Coker}(\bar{\partial}_{23^*}^{16,0}) &= \mathbb{Z}_2\{\bar{\beta}_8 \sigma_8 \bar{\nu}_{15}\} \oplus \mathbb{Z}_2\{\bar{\beta}_8 \sigma_8 \epsilon_{15}\} \oplus \mathbb{Z}_8\{\bar{\beta}_8 \Sigma \rho''\} \oplus \mathbb{Z}_2\{\bar{\beta}_8 \bar{\epsilon}_8\} \oplus \mathbb{Z}_{(2)}\{\bar{\beta}_{23}\} \end{aligned}$$

where $\sigma_8 \bar{\nu}_{15}, \sigma_8 \epsilon_{15}, \Sigma \rho''$ and $\bar{\epsilon}_8$ are the generators of $\pi_{23}(S^8)$.

From Lemma 2.8, $\bar{p}_{16}^0 \gamma_3^0 \simeq 0$. Hence $\gamma_3^0 \in \text{Ker}(\bar{p}_{16}^0)$ implies

$$\gamma_3^0 = \bar{\tau}_{16}^0(\bar{\beta}_8 \theta_0 + a_0 \bar{\beta}_{23}) = \beta_8 \theta_0 + a_0 \beta_{23} \text{ for } \theta_0 \in \pi_{23}(S^8) \text{ and some integer } a_0 \in \mathbb{Z}_{(2)}.$$

Moreover, $\pi_{23}(J_{2,8}^0)/\langle \gamma_3^0 \rangle \cong \pi_{23}(J_{3,8}^0) \cong \pi_{23}(F_9^0) \cong \pi_{24}(S^9) \cong \mathbb{Z}_{16} \oplus \mathbb{Z}_2^{\oplus 3}$.

Let $a_0 = 2^{t_0} a'_0$ where integer $t_0 \geq 0$ and integer a'_0 is odd. By comparing the order of the groups of the first and last two items in above isomorphisms, we get $2^{t_0+7} = 2^7$.

Hence $t_0 = 0$, i.e., a_0 is odd.

The first diagram in (43) induces the following diagram

$$\begin{array}{ccccccccc} S^{23} & \xrightarrow{j_2^{23}} & J_{2,8}^{\gamma,0} & \xrightarrow{I_2^{\gamma,0}} & F_{\bar{p}_{16}^0} & \xrightarrow{\bar{\tau}_{16}^0} & J_{2,8}^0 & \xrightarrow{\bar{p}_{16}^0} & S^{16} \\ & & \downarrow (g_\nu^0)|_{J_2} & & \downarrow g_1^s & & \downarrow \bar{g}_\nu^0 & & \downarrow \nu_{13} \\ S^{17} & \xrightarrow{j_2^{17}} & J_2(M_{\gamma_2^s}, S^8) \simeq S^5 \vee S^{17} & \xrightarrow{I_2^{\gamma,\nu}} & F_{q_2^{13}} & \xrightarrow{\bar{\tau}_{13}^\nu} & J_{2,\nu} & \xrightarrow{q_2^{13}} & S^{13} \end{array}$$

$$\text{with } (g_\nu^0)|_{J_2} = j_1^5 \nu_5 q_1^8 + j_2^{17} \nu_{17}^2 q_2^{23} + j_1^5 \theta_1 q_2^{23}, \theta_1 \in \pi_{23}(S^5),$$

where $I_2^{\gamma,\nu}$ and $\bar{\tau}_{13}^\nu$ are canonical inclusions. By (2) and the definition of β_{23} in the Section 4, we have

$$\bar{\tau}_{13}^\nu I_2^{\gamma,\nu} j_1^5 \simeq j_1^5 \quad \text{and} \quad \beta_{23} = \bar{\tau}_{16}^0 I_2^{\gamma,0} j_2^{23}.$$

Since $\pi_{17}(J_{2,\nu}) = \pi_{17}(S^5 \vee S^{13}) = \mathbb{Z}\{[j_1^5, j_2^{13}]\} \oplus j_1^5 \circ \pi_{17}(S^5)$ and on the other hand $\pi_{17}(J_{2,\nu}) = \mathbb{Z}\{I_2^{\gamma,\nu} \bar{\tau}_{13}^\nu j_2^{17}\} \oplus j_1^5 \circ \pi_{17}(S^5)$ which is obtained by the exact sequence of $\pi_{17}(-)$ induced by the fibration sequence $\Omega S^{13} \rightarrow F_{q_2^{13}} \xrightarrow{\bar{\tau}_{13}^\nu} J_{2,\nu} \xrightarrow{q_2^{13}} S^{13}$, we get

$$I_2^{\gamma,\nu} \bar{\tau}_{13}^\nu j_2^{17} = \pm [j_1^5, j_2^{13}] + j_1^5 \varrho, \varrho \in \pi_{17}(S^5).$$

So

$$\begin{aligned} \bar{g}_\nu^0 \beta_{23} &\simeq \bar{g}_\nu^0 \bar{\tau}_{16}^0 I_2^{\gamma,0} j_2^{23} \simeq I_2^{\gamma,\nu} \bar{\tau}_{13}^\nu (g_\nu^0)|_{J_2} j_2^{23} \simeq I_2^{\gamma,\nu} \bar{\tau}_{13}^\nu (j_1^5 \nu_5 q_1^8 + j_2^{17} \nu_{17}^2 q_2^{23} + j_1^5 \theta_1 q_2^{23}) j_2^{23} \\ &\simeq I_2^{\gamma,\nu} \bar{\tau}_{13}^\nu j_2^{17} \nu_{17}^2 + I_2^{\gamma,\nu} \bar{\tau}_{13}^\nu j_1^5 \theta_1 \simeq \pm [j_1^5, j_2^{13}] \nu_{17}^2 + j_1^5 \theta'_1, \text{ where } \theta'_1 = \varrho \nu_{17}^2 + \theta_1. \\ &\Rightarrow \bar{g}_\nu^0 \gamma_3^0 = \bar{g}_\nu^0 (\beta_8 \theta_0 + a_0 \beta_{23}) \simeq j_1^5 \nu_5 \theta_0 + a_0 (\pm [j_1^5, j_2^{13}] \nu_{17}^2 + j_1^5 \theta'_1), \text{ by (44)} \\ &\simeq \pm a_0 [j_1^5, j_2^{13}] \nu_{17}^2 + j_1^5 \theta_2 \in \pi_{23}(J_{2,\nu}), \theta_2 = \theta'_1 + \nu_5 \theta_0 \in \pi_{23}(S^5). \end{aligned}$$

Since a_0 is odd and $\pi_{23}(J_{2,\nu}) = \pi_{23}(S^5 \vee S^{13}) = j_1^5 \circ \pi_{23}(S^5) \oplus j_2^{13} \circ \pi_{23}(S^{13}) \oplus \mathbb{Z}_8 \{[j_1^5, j_2^{13}] \nu_{17}^2\}$, $\simeq \bar{g}_\nu^0 \gamma_3^0 \neq 0$. From (44) and (45),

$$0 \neq \bar{g}_\nu^0 \gamma_3^0 \simeq \gamma_3^\nu \nu_{20} = a [j_1^5, j_2^{13}] \nu_{17} \Rightarrow a \text{ is odd.}$$

We complete the proof of this lemma. \square

Acknowledgement. The author was partially supported by National Natural Science Foundation of China (Grant No. 11701430).

References

- [1] M. Arkowitz, *The generalized Whitehead product*, Pacific J Math. 12(1962) 7-23.
- [2] M.G. Barratt, *On spaces of finite characteristic*, Quart. J. Math. Oxford 11 (1960), 124-136.
- [3] H.J. Baues, *On homotopy classification problems of J. H. C. Whitehead*, Algebraic topology, Göttingen 1984 (1985) 17-55.
- [4] H.J. Baues, M. Hennes, *The homotopy classification of $(n-1)$ -connected $(n+3)$ -dimensional polyhedra, $n \geq 4$* , Topology 30 (1991), 373-408.
- [5] J.M. Boardman, B. Steer, *On Hopf invariants*. Comment. Math. Helv. 42(1967) 180–221.
- [6] W.D. Chen, J. Wu, *Decomposition of loop spaces and periodic problem on π_** , Algebr. Geom. Topol. 13 (2013), 3245–3260.
- [7] F.R. Cohen, *Stable Homotopy*, Lecture Notes in Mathematics 165 Berlin-Heidelberg: Springer, 1970.
- [8] F.R. Cohen, *A course in some aspects of classical homotopy theory*, 1987.DOI:10.1007/BFb0078738.
- [9] F.R. Cohen, L.R. Taylor, *Homology of Function Spaces*, Math. Z. 198(1988) 299-316.
- [10] F.R. Cohen, J. Wu, *A remark on the homotopy groups of $\Sigma^n \mathbb{R}P^2$* , Contem. Math. 181(1995), 65-81.
- [11] T. Cutler, T. So, *The homotopy type of a once-suspended 6-manifold and its applications*, Topology Appl. 318(2022) 108213.
- [12] M. Golasiński, T. de Melo, *On the higher Whitehead product*, J Homotopy and Related Stru. 11(2016) 825-845.
- [13] B. Gray, *On the homotopy groups of mapping cones*, Proc. Lond. Math. Soc. 26(1973) 497-520.
- [14] K.A. Hardie, *Higher Whitehead products*, Quart. J. Math. 12(1961) 241-249.
- [15] K.A. Hardie, *On a construction of E.C.Zeeman*, J. Lond. Math. Soc. 35(1960) 452-464.
- [16] A. Hatcher, *Spectral Sequences in Algebraic Topology*, (Unpublished) <http://www.math.cornell.edu/hatcher>.

- [17] R.Z. Huang, *Homotopy of gauge groups over non-simply connected five dimensional manifolds*, Sci. China Math. 64(2021) 1061-1092.
- [18] R.Z. Huang, *Suspension homotopy of 6-manifolds*, Alg. Geom. Topol. 23(2023) 439-460.
- [19] R.Z. Huang, P.C. Li, *Suspension homotopy of simply connected 7- manifolds*, Sci. China Math. (2023), in press, arXiv:2208.13145.
- [20] I.M. James, *On the homotopy groups of certain pairs and triads*, Quart. J. Math. Oxford 5(1954) 260-270.
- [21] I.M. James, *Reduced product spaces*, Ann. of Math. (2) 62 (1955), 170-179.
- [22] I.M. James, *Note on cup-products*, Proc. Amer. Math. Soc. 8(1957) 374-383.
- [23] T.Jin T, Z.J. Zhu, *The $n+3, n+4$ dimensional homotopy groups of A_n^2 -complexes*, arXiv: 2402.11226 (2024).
- [24] X.G. Liu, *On the Moore space $P^n(8)$ and its homotopy groups*(in Chinese), Chinese Ann. Math. Ser. A 28(2007) 305-318.
- [25] P.C. Li, Z.J. Zhu, *The homotopy decomposition of the suspension of a non-simply-connected five-manifold*, Proc. R. Soc. Edinb. A (2024) 1-29. doi:10.1017/prm.2024.49.
- [26] K. Morisugi, J. Mukai, *Lifting to mod 2 Moore spaces*, J. Math. Soc. Japan 52(2000) 515-533.
- [27] J. Mukai, T. Shinpo, *Some homotopy groups of the mod 4 Moore space*, J. Fac. Sci. Shinshu Univ. 34(1999) 1-14.
- [28] J. Mukai, *A remark on Toda's result about the suspension order of the stunted real projective space*, Memo. of. Facu. of Kyushu Univ. Ser. A 42(1988) 87-94.
- [29] J. Mukai, *Note on existence of the unsatable Adams map*, Kyushu J. Math. 49(1995) 271-279.
- [30] J. Mukai, *Generators of some homotopy groups of the mod2 Moore space of dimension 3 or 5*, Kyushu J. Math. 55(2001) 63-73.
- [31] J. Mukai, *The S^1 -transfer map and homotopy groups of suspended complex projective spaces*, Math. J. Okayama Univ. 24(1982) 179-200.
- [32] J. Neisendorfer, *Algebraic methods in unstable homotopy theory*, Cambridge University Press, 2010.

- [33] N. Oda, *Unstable homotopy groups of spheres*, Bull Inst Adv Res Fukuoka Univ, 44 (1979) 49-151.
- [34] G.J. Porter, *Higher order Whitehead products*, Topology 3(1965) 123-135.
- [35] G.J. Porter, *Higher order Whitehead products and Postnikov systems*, Illinois J. Math. 11(1967) 414-416.
- [36] P. Selick, J. Wu, *The functor A^{min} on p -local spaces*, Math. Z. 253 (3) (2006), 435-451.
- [37] T. So, S. Theriault, *The suspension of a 4-manifold and its applications*, Isr. J. Math. (2022), in press, arXiv: 1909.11129.
- [38] B. Steer, *Transgression in sphere-bundles*, Topology 2(1963), 1-10
- [39] H. Toda, *Composition Methods in Homotopy Groups of Spheres*, Annals of Mathematics Studies, Princeton University Press, Princeton, 1962.
- [40] H. Toda, *Generalized Whitehead Products and Homotopy Groups of Spheres*, J. Inst. Polytech. Osaka City Univ. Ser. A 3(1952) 43-82.
- [41] G.W. Whitehead, *Elements of Homotopy Theory*, Graduate Texts in Mathematics 61, Springer-Verlag, 1978.
- [42] J. Wu, *On combinatorial calculations for the James Hopf maps*, Topology 37 (1998), 1011-1023.
- [43] J. Wu, *Homotopy Theory of the Suspensions of the Projective Plane*, Mem. Amer. Math. Soc. 162(2003), No. 769.
- [44] K. Yamaguchi, *The group of self-homotopy equivalences of S^2 -bundles over S^4 .I.*, Kodai Math. J. 9(1986) 308-326.
- [45] J.X. Yang, J. Mukai, J. Wu, *On the homotopy groups of the suspended quaternionic projective plane and the applications*, Preprint (2023) arXiv:2301.06776.
- [46] Z.J. Zhu, J.Z. Pan, *The decomposability of smash product of \mathbf{A}_n^2 complexes*, Homology Homotopy and Applications 19(2017) 293-318.
- [47] Z.J. Zhu, P.C. Li, J.Z. Pan, *Periodic problem on homotopy groups of Chang complexes $C_r^{n+2,r}$* , Homology Homotopy and Applications 21.2 (2019) 363-375.
- [48] Z.J. Zhu, J.Z. Pan, *The local hyperbolicity of \mathbf{A}_n^2 -complexes*, Homology Homotopy and Applications 23.1 (2021) 367-386.

[49] Z.J. Zhu, J.Z. Pan, *2-local unstable homotopy groups of A_3^2 -Complexes*, Sci. China Math. 67 (2024) 607–626.

[50] Z.J. Zhu, T. Jin, *The relative James construction and its application to homotopy groups*, Topology and its Applications 356 (2024) 109043

8. Appendix

Lemma 8.1. (i) $\Sigma v'\sigma' = 2\Sigma\varepsilon'$. (ii) $\mu_4\eta_{13} = \eta_4\mu_5$.
 (iii) $v_5\eta_8^2\sigma_{10} = v_5^4 + v_5\eta_8\varepsilon_9$. (iv) $(2^r\iota_5)v_5\sigma_8 = 2^r v_5\sigma_8$.
 (v) $(2^r\iota_6)\bar{v}_6 = 2^{2r}\bar{v}_6$.

Proof. (i) By $\Sigma v'\sigma' \in \pi_{14}(S^4) = \mathbb{Z}_8\{v_4\sigma'\} \oplus \mathbb{Z}_4\{\Sigma\varepsilon'\} \oplus \mathbb{Z}_2\{\eta_4\mu_5\}$, assume that

$$\Sigma v'\sigma' = xv_4\sigma' + y\Sigma\varepsilon' + z\eta_4\mu_5, \quad x \in \mathbb{Z}_8, y \in \mathbb{Z}_4, z \in \mathbb{Z}_2. \quad (46)$$

Then $2v_5\Sigma\sigma' = \Sigma(\Sigma v'\sigma') = xv_5\Sigma\sigma' + y\Sigma^2\varepsilon' + z\eta_5\mu_6$. From (7.10), (7.16) of [39], $2v_5\sigma_8 = v_5\Sigma\sigma' = \pm\Sigma^2\varepsilon'$ implies that $4v_5\sigma_8 = (2x \pm 2y)v_5\sigma_8 + z\eta_5\mu_6 \in \pi_{15}(S^5) = \mathbb{Z}_8\{v_5\sigma_8\} \oplus \mathbb{Z}_2\{\eta_5\mu_6\}$. Hence $z = 0$, $2(x \pm y) = 4$. Multiply both sides of equation (46) by 2,

$$2\Sigma v'\sigma' = 2xv_4\sigma' + 2y\Sigma\varepsilon'. \quad (47)$$

$2\Sigma v'\sigma' = \Sigma v'(2\sigma') = \Sigma(v'\sigma'')$ since $2\sigma' = \Sigma\sigma''$ by Lemma 5.14 of [39]. Consider Hopf invariant H_2 on both sides of (47), $0 = H_2(2\Sigma v'\sigma') = H_2(2xv_4\sigma') = xH_2(v_4\Sigma\sigma'') = xH_2(v_4)\Sigma\sigma'' = x\Sigma\sigma'' = 2x\sigma' \in \pi_{14}(S^7) = \mathbb{Z}_8\{\sigma'\}$. So $x = 4t$ ($t = 0, 1$) implies $y = 2$ and (46) becomes $\Sigma v'\sigma' = 4tv_4\sigma' + 2\Sigma\varepsilon' = 2tv_4\Sigma\sigma'' + 2\Sigma\varepsilon'$. By Proposition 2.2, Lemma 3.1 and Lemma 5.14 of [39], $H_2(\Sigma v'\sigma') = (\Sigma v' \wedge v')H_2(\sigma') = \Sigma(-\Sigma^3 v' \Sigma^6 v')\eta_{13} = -4v_7^2\eta_{13} = 0$. On the other hand, $H_2(\Sigma v'\sigma') = 2tH_2(v_4\Sigma\sigma'') = 2tH_2(v_4)\Sigma\sigma'' = 2t\Sigma\sigma'' = 4t\sigma'$. Thus $4t\sigma' = 0$ implies $t = 2t'$ and $x = 8t' = 0 \in \mathbb{Z}_8$. So $\Sigma v'\sigma' = 2\Sigma\varepsilon'$.

(ii) Let $\mu_4\eta_{13} = 4x'v_4\sigma' + 2y'\Sigma\varepsilon' + z'\eta_4\mu_5$, $x', y', z' \in \mathbb{Z}$. By page.66 of [39], $\eta_5\mu_6 = \Sigma\mu_4\eta_{13} = \pm 4y'v_5\sigma_8 + z'\eta_5\mu_6$, implies $2|y', z' = 1$. Hence $\mu_4\eta_{13} = 4x'v_4\sigma' + \eta_4\mu_5$. By $0 = H(\mu_4\eta_{13}) = H_2(4x'v_4\sigma' + \eta_4\mu_5) = 4x'\sigma'$, we get $2|x'$. So $\mu_4\eta_{13} = \eta_4\mu_5$.

(iii) $v_5\eta_8^2\sigma_{10} = v_5\eta_8((\Sigma^2\sigma')\eta_{16} + \bar{v}_9 + \varepsilon_9) = v_5\eta_8(\bar{v}_9 + \varepsilon_9) = v_5^3 + v_5\eta_8\varepsilon_9$. where the first, second and the last equations are from (7.4), Lemma 5.14 and (7.3) of [39] respectively.

(iv) By the same proof as that of Lemma A.1 of [49], $(2^r\iota_8)\sigma_8 = 2^{2r}\sigma_8 - 2^{r-1}(2^r - 1)\Sigma\sigma'$.

$(2^r\iota_5)v_5\sigma_8 = v_5((2^r\iota_8)\sigma_8) = 2^{2r}v_5\sigma_8 - 2^{r-1}(2^r - 1)v_5\Sigma\sigma' = 2^{2r}v_5\sigma_8 - 2^r(2^r - 1)v_5\sigma_8 = 2^r v_5\sigma_8$.

(v) By [33, Proposition 2.10], $H_2(\bar{v}_6) = l\nu_{11}$, l is odd, and $\Delta(v_{13}) = \pm(\iota_{13})v_{11} = \pm 2\bar{v}_6$ [39, Lemma 6.2], we have

$(2^r \iota_6) \bar{v}_6 = 2^r \bar{v}_6 \pm \binom{2^r}{2} [\iota_6, \iota_6] H_2(\bar{v}_6) = 2^r \bar{v}_6 \pm \binom{2^r}{2} l \Delta(\iota_{13}) \nu_{11} = 2^r \bar{v}_6 \pm \binom{2^r}{2} l \Delta(\nu_{13}) = 2^r \bar{v}_6 \pm 2^r (2^r - 1) \bar{v}_6 = 2^{2r} \bar{v}_6$ or $(2^{r+1} - 2^{2r}) \bar{v}_6$. Moreover $H_2((2^r \iota_6) \bar{v}_6) = \Sigma(2^r \iota_5 \wedge 2^r \iota_5) H_2(\bar{v}_6) = 2^{2r} \nu_{11} = H_2(2^{2r} \bar{v}_6)$. We get $(2^r \iota_6) \bar{v}_6 = 2^{2r} \bar{v}_6$. \square

Lemma 8.2.

- (i) $\{2\iota_3, \nu' \eta_6, 2\iota_7\} = \{\nu' \eta_6^2\}$; $\{2^r \iota_3, \nu' \eta_6, 2\iota_7\} = \{0\} (r \geq 2)$.
- (ii) $\{4\iota_4, \nu_4 \eta_7, 2\iota_8\} = \{\Sigma \nu' \eta_7^2\}$; $\{2^r \iota_4, \nu_4 \eta_7, 2\iota_8\} = \{0\} (r \geq 3)$.
- (iii) $\{2\iota_4, \varepsilon_4, 2\iota_{12}\} = \{\eta_4 \varepsilon_5\}$; $\{2^r \iota_4, \varepsilon_4, 2\iota_{12}\} = \{0\} (r \geq 2)$.
- (iv) $\{2\iota_4, \mu_4, 2\iota_{13}\} = \eta_4 \mu_5 + \langle 2\nu_4 \sigma', 2\Sigma \varepsilon' \rangle$; $\{2^r \iota_4, \mu_4, 2\iota_{13}\} = \langle 2\nu_4 \sigma', 2\Sigma \varepsilon' \rangle (r \geq 2)$.
- (v) $\{2\iota_5, \sigma''', 2\iota_{12}\} = \{0\}$.
- (vi) $\{2\iota_5, \varepsilon_5, 2\iota_{13}\} = \{\eta_5 \varepsilon_6\}$; $\{2^r \iota_5, \varepsilon_5, 2\iota_{13}\} = \{0\}, r \geq 2$.
- (vii) $\{2\iota_6, 2\sigma'', 2\iota_{13}\} = \langle 2\bar{v}_6 \rangle$; $\{4\iota_6, \sigma'', 4\iota_{13}\} = \langle 4\bar{v}_6 \rangle$.
- (viii) $\{2\iota_7, \sigma' \eta_{14}, 2\iota_{15}\} = \{\sigma' \eta_{14}^2\}$; $\{2^r \iota_7, \sigma' \eta_{14}, 2\iota_{15}\} = \{0\} (r \geq 2)$;

Proof. (i) For $r \geq 1$, $\{2^r \iota_3, \nu' \eta_6, 2\iota_7\}$ is a coset of subgroup $(2^r \iota_3) \circ \pi_8(S^3) + \pi_8(S^3) \circ (2\iota_8) = \{0\}$, implies that $\{2^r \iota_3, \nu' \eta_6, 2\iota_7\}$ has only one element. Then (i) of this lemma comes from the following arguments

$$\begin{aligned} \{2^r \iota_3, \nu' \eta_6, 2\iota_7\} &\supset \{2^r \nu', \eta_6, 2\iota_7\} \supset (2^{r-1} \nu') \{2\iota_6, \eta_6, 2\iota_7\} \text{ by [39, Prop. 1.2]} \\ &\supset (2^{r-1} \nu') \{2\iota_6, \eta_6, 2\iota_7\}_1 \ni 2^{r-1} \nu' \eta_6^2, \text{ by [39, (1.15) and Corollary 3.7].} \end{aligned}$$

(ii) For $r \geq 2$, $\{2^r \iota_4, \nu_4 \eta_7, 2\iota_8\}$ is a coset of subgroup $\pi_9(S^4) \circ (2\iota_9) + (2^r \iota_4) \circ \pi_9(S^4) = \{0\}$ which implies that $\{2^r \iota_4, \nu_4 \eta_7, 2\iota_8\}$ has only one element.

$$\begin{aligned} \{2^r \iota_4, \nu_4 \eta_7, 2\iota_8\} &\supset \{(2^r \iota_4) \nu_4, \eta_7, 2\iota_8\} = \{2^{2r} \nu_4 - 2^{r-1} (2^r - 1) \Sigma \nu', \eta_7, 2\iota_8\} \\ &\subset \{2^{2r} \nu_4, \eta_7, 2\iota_8\} + \{2^{r-1} (2^r - 1) \Sigma \nu', \eta_7, 2\iota_8\} = \{2^{r-1} \Sigma \nu', \eta_7, 2\iota_8\}. \end{aligned}$$

Thus (ii) of Lemma 8.2 is obtained by (ii) of this lemma.

(iii) By Lemma 3.9 and [39, (7.5)], $\{2^r \iota_4, \varepsilon_4, 2\iota_{12}\} \ni 2^{r-1} \varepsilon_4 \eta_{12} = 2^{r-1} \eta_4 \varepsilon_5 \pmod{\pi_{13}(S^4) \circ (2\iota_{13}) + (2^r \iota_4) \circ \pi_{13}(S^4) = \{0\}}$ which implies the (iii) of Lemma 8.2.

(iv) $\{2^r \iota_4, \mu_4, 2\iota_{13}\}$ is a coset of subgroup $\pi_{14}(S^4) \circ (2\iota_{14}) + (2^r \iota_4) \circ \pi_{14}(S^4) = \langle 2\nu_4 \sigma', 2\Sigma \varepsilon' \rangle \subset \pi_{14}(S^4)$. For $r = 1$, by Corollary 3.7 of [39] and (ii) of Lemma 8.1, $\mu_4 \eta_{13} = \eta_4 \mu_5 \in \{2\iota_4, \mu_4, 2\iota_{13}\}$. Hence $\{2\iota_4, \mu_4, 2\iota_{13}\} = \eta_4 \mu_5 + \langle 2\nu_4 \sigma', 2\Sigma \varepsilon' \rangle$. For $r \geq 2$, $\{2^r \iota_4, \mu_4, 2\iota_{13}\} \supset (2^{r-1} \iota_4) \circ \{2\iota_4, \mu_4, 2\iota_{13}\} \ni (2^{r-1} \iota_4) \eta_4 \mu_5 = 0$ implies $\{2^r \iota_4, \mu_4, 2\iota_{13}\} = \langle 2\nu_4 \sigma', 2\Sigma \varepsilon' \rangle$.

(v) $\{2\iota_5, \sigma''', 2\iota_{12}\}$ is a coset of subgroup $(2\iota_5) \circ \pi_{13}(S^5) + \pi_{13}(S^5) \circ (2\iota_{13}) = \{0\}$. $-\Sigma\{2\iota_5, \sigma''', 2\iota_{12}\} \subset \{2\iota_6, \Sigma \sigma''', 2\iota_{13}\}_1 \ni \Sigma \sigma''' \eta_{13} = (2\sigma'') \eta_{13} = 0 \pmod{\langle 2\bar{v}_6 \rangle}$.

Thus $\Sigma\{2\iota_5, \sigma''', 2\iota_{12}\} \subset \langle 2\bar{v}_6 \rangle$ implies $\{2\iota_5, \sigma''', 2\iota_{12}\} = 0$ by [39, Theorem 7.1].

(vi) $\{2^r \iota_5, \varepsilon_5, 2\iota_{13}\}$ is a coset of $(2^r \iota_5) \circ \pi_{14}(S^5) + \pi_{14}(S^5) \circ (2\iota_{14}) = \{0\}$. Moreover $\{2^r \iota_5, \varepsilon_5, 2\iota_{13}\} \ni (2^{r-1} \iota_5) \varepsilon_5 \eta_{13} = 2^{r-1} \eta_5 \varepsilon_6$ by Lemma 3.9. This implies (vi) of this lemma.

(vii) For $r \leq 2$, $\{2^r \iota_6, 2^{2-r} \sigma'', 2^r \iota_{13}\} \subset \{2\iota_6, (2^{r-1} \iota_6)(2^{2-r} \sigma''), 2^r \iota_{13}\} = \{2\iota_6, 2\sigma'', 2^r \iota_{13}\} = \{2\iota_6, \Sigma\sigma''', 2^r \iota_{13}\} \supset \{2\iota_6, \Sigma\sigma''', 2\iota_{13}\}(2^{r-1} \iota_{14}) \ni 2^{r-1} \Sigma\sigma''' \eta_{13} = 0$. $\{2\iota_6, \Sigma\sigma''', 2^r \iota_{13}\}$ is a coset of subgroup $(2\iota_6) \circ \pi_{14}(S^6) + \pi_{14}(S^6) \circ (2^r \iota_{14}) = \langle 2^r \bar{v}_6 \rangle$ ($r \leq 2$) by (v) of Lemma 8.1 and so is $\{2^r \iota_6, 2^{2-r} \sigma'', 2^r \iota_{13}\}$. Hence $\{2^r \iota_6, 2^{2-r} \sigma'', 2^r \iota_{13}\} = \langle 2^r \bar{v}_6 \rangle$.

(viii) $\{2^r \iota_7, \sigma' \eta_{14}, 2\iota_{15}\}$ is a coset of subgroup $(2^r \iota_7) \circ \pi_{16}(S^7) + \pi_{16}(S^7) \circ (2^r \iota_{16}) = \{0\}$ for $r \geq 1$. Then from [39, Propostion 1.2] and Lemma 3.9, we get $\{2^r \iota_7, \sigma' \eta_{14}, 2\iota_{15}\} \supset \{2^r \sigma', \eta_{14}, 2\iota_{15}\} \ni 2^{r-1} \sigma' \eta_{14}^2$, which implies (viii) of this lemma.

□