

Accordingly, the present paper assumes the popular radix-10, aiming to calculate the exact number of the rightmost (trailing) digits of G that matches the tail-end figures of the endless string $\dots 35796399618993967905496638003222348723967018485186439059104575627262464195387$ of frozen digits originated by the increasing height power tower $\underbrace{3^{3^{\dots^3}}}_{b\text{-times}}$ as b grows over and over [4].

Defintion 1.1. From here on, let us indicate the integer tetration $a \uparrow\uparrow b := \begin{cases} a & \text{if } b = 1 \\ a^{(b-1)a} & \text{if } b \geq 2 \end{cases}$ as ${}^b a$ (where $a, b \in \mathbb{Z}^+$) so that we can compactly write ${}^b 3$ instead of $3 \uparrow\uparrow b$, and consequently note that $\exists! n \in \mathbb{Z}^+ : {}^n 3 = G$.

Thus, the goal of Section 2 is to determine the exact value of the positive integer r such that ${}^{n+1}3 \equiv G \pmod{10^{r-1}}$ and ${}^{n+1}3 \not\equiv G \pmod{10^r}$, which is equivalent to asking to find, for any $c \in \mathbb{Z}^+$, the value of r such that ${}^{n+c}3 \equiv G \pmod{10^{r-1}}$ and ${}^{n+c}3 \not\equiv G \pmod{10^r}$.

In the following sections, we will indicate as $\text{slog}_3(G)$ the base 3 super-logarithm of Graham's number (see Definition 2.1, and see also [1, p. 68] for the *log star* notation of the *iterated logarithm*) to prove that $r = \text{slog}_3(G)$.

Lastly, Section 3 shows that, for each positive integer c , the difference between the $\text{slog}_3(G)$ -th rightmost digit of G and the corresponding digit of ${}^{\text{slog}_3(G)+c}3$ is 6 if the $\text{slog}_3(G)$ -th least significant digit of G belongs to the set $\{6, 7, 8, 9\}$, whereas the mentioned difference is equal to -4 if this does not occur.

In the end, for every $c \in \mathbb{Z}^+$, ${}^{\text{slog}_3(G)+c}3 - G \equiv 4 \cdot 10^{\text{slog}_3(G)-1} \pmod{10^{\text{slog}_3(G)}}$, and so the $\text{slog}_3(G)$ -th rightmost digit of $3^G - G$ is 4.

2 The last $\text{slog}_3(G) - 1$ digits of G

This section is devoted to proving that the rightmost digit of G that is not equal to the corresponding digit of any base 3 tetration whose (integer) hyperexponent is greater than n is the $\text{slog}_3(G)$ -th, where the mentioned super-logarithm is defined as follows.

Defintion 2.1. For positive integers p and q , we define the integer super-logarithm $\text{slog}_p(q)$ by $\text{slog}_p(1) = 0$ and $\text{slog}_p(p^q) = \text{slog}_p(q) + 1$ so that $\text{slog}_p(p^q) = \text{slog}_p(\log_p(q)) + 2$ is certainly true as long as $p > 1$.

Consequently, the base 3 super logarithm of any positive integer n is such that $\text{slog}_3({}^n 3) = \text{slog}_3({}^{n-1}3) + 1 = \dots = \text{slog}_3({}^2 3) + n - 2 = \text{slog}_3({}^1 3) + n - 1 = \text{slog}_3({}^0 3) + n = \text{slog}_3(1) + n$, and thus $\text{slog}_3({}^n 3) = n$ since $\text{slog}_3(1) = 0$ (by Definition 2.1).

Hence, ${}^n 3 = G$ implies $n = \text{slog}_3(G)$.

Although there is not a general agreement on the definition of the super-logarithm for non-integers [9], let us point out that this is not the case for Definition 2.1, where we are considering only strictly positive integer entries.

Now, by assuming radix-10 as usual, we introduce the ‘‘constancy of the congruence speed’’, a peculiar property of every tetration base not a multiple of 10 proved by the author in recent years [6] (see also the related OEIS sequences A317905 and A373387).

Defintion 2.2. Let $m \in \mathbb{N}_0$ and assume that $a \in \mathbb{N} - \{0, 1\}$ is not a multiple of 10. Then, given $b^{-1}a \equiv b^a \pmod{10^m}$ and $b^{-1}a \not\equiv b^a \pmod{10^{m+1}}$, for all $b \in \mathbb{Z}^+$, $V(a, b)$ returns the nonnegative integer such that $b^a \equiv b^{b+1}a \pmod{10^{m+V(a,b)}}$ and $b^a \not\equiv b^{b+1}a \pmod{10^{m+V(a,b)+1}}$, and we define $V(a, b)$ as the *congruence speed* of the base a at the given height of its hyperexponent b . Furthermore, let $\bar{b} := \min_b \{b \in \mathbb{Z}^+ : V(a, b) = V(a, b+k) \text{ for all } k \in \mathbb{Z}^+\}$ so that we define as *constant congruence speed* of a the positive integer $V(a) := V(a, \bar{b})$.

For any $a, b \in \mathbb{N} - \{0, 1\}$, the congruence speed of the tetration b^a should always return the exact number of the rightmost digits of the result that freeze by going from b^a to $b^{b+1}a$ and that were not already stable digits at height $b^{-1}a$, but there is one (disputable) exception to the above-mentioned general correspondence: it is the case $a = 5$ with $b = 2$, as $15 = 5$, $25 = 3125$, and $35 \equiv 25 \pmod{10^5}$ so that we could argue that the stated definition would imply $V(5, 2) = 4 = V(5, 3)$ even if 25 is only a 4-digit number, where the last one has been already frozen at height 1 (i.e., $25 \equiv 5 \pmod{10^1}$); for a proof that this is the only possible collision that can occur, for every a and b greater than 1, see [<https://math.stackexchange.com/questions/4863065/>].

Then, since [7, Equation (16)] let us easily calculate the value of the constant congruence speed of every tetration base not a multiple of 10, we can use it to prove Theorem 2.3. On specific, here we are only interested in line 11 of the above-mentioned Equation (16), stating that the constant congruence speed of each tetration base, let us indicate it as a^* , which is congruent to 3 modulo 20 and such that 5 is not equal to the absolute value of the difference between the rightmost digit of a^* that is different from the corresponding digit of $\dots 99834030970896579486665776138023544317662666830362972182803640476581907922943$ (i.e., $\{5^{2^n}\}_\infty - \{2^{5^n}\}_\infty$) and the mentioned key digit of $\dots 30362972182803640476581907922943$, is given by the 5-adic valuation of $(a^*)^2 + 1$ (and then $a^* := \dots 000003$ implies that the constant congruence speed of a^* is equal to the maximum number of times that 5 divides $3^2 + 1$, so $V(3) = 1$ follows).

Theorem 2.3. Let m and n be positive integers such that $m > n$. In radix-10, $m3 \equiv G \pmod{10^{n-1}}$ and $m3 \not\equiv G \pmod{10^n}$ occurs if and only if $n = slog_3(G)$.

Proof. Theorem 2.3 states that the rightmost digit of Graham's number that is not stable (i.e., a digit that does not match the corresponding figure of the endless string $\dots 27262464195387$ of frozen digits returned by $3^{3^{3^{\dots}}}$) is the figure in position $slog_3(G)$, by counting from right to left of G .

Our strategy to prove this theorem is to show that, in the decimal numeral system, the congruence speed of the base 3 tetration is equal to 0 if and only if its hyperexponent is 1 and that the congruence speed becomes certainly constant starting from height 3 (i.e., the congruence speed of 3^3 is equal to the congruence speed of $3^{3+c}3$ for any given $c \in \mathbb{Z}^+$ since a sufficient condition for the constancy of the congruence speed of every tetration base $a > 1$ not a multiple of 10 is provided by $b \geq \tilde{\nu} + 2$, where b is the hyperexponent of a and $\tilde{\nu}(a)$ is described by [7, Definition 2.1]), so $V(3, b) = V(3) = 1$ for every $b \in \mathbb{N} - \{0, 1, 2, 3\}$. Then, we only need to check the case $b = 2$ to directly conclude that $V(3, b) = V(3) = 1$ if and only if $b \geq 2$.

It is enough to derive that the total number of the rightmost stable digits of $n3$ is $n - 1$.

In detail, from [7, Definition 2.1], it follows that $\tilde{\nu}(3) := \nu_5(a^2 + 1)$, where $\nu_5(\dots)$ indicates the 5-adic valuation of the argument. Page 449 of the mentioned paper gives $b \geq \tilde{\nu}(a) + 2$ as a

sufficient (but not necessary) condition on $b := b(a)$ for the constancy of the congruence speed of any tetration base a whose last digit is 3 or 7. Since $\nu_5(a^2 + 1) + 2 = 3$, we only have to calculate the total number of the rightmost digits of ${}^3 3$ that do not change as we move to ${}^4 3$. Then, it is not difficult to see that ${}^3 3 = 987$ and ${}^4 3 \equiv 387 \pmod{10^3}$ so that $V(3, 1) + V(3, 2) + V(3, 3) = 2$.

Now, [7, Equation (16), line 11] implies that the base 3 tetration is characterized by a unit constant congruence speed since $V(3) = \nu_5(3^2 + 1) = \nu_5(5 \cdot 2) = 1$.

Hence, given the fact that $G = {}^n 3 = g_{64}$ implies $n > 3$ (see Equation (1.1)), the total number of stable digits of G is equal to $2 + 1 \cdot V(3) + (n - 3) = 2 + n - 3 = n - 1$.

Thus, the least significant digit of ${}^n 3$ that is not stable is the one immediately to the left of the $(n - 1)$ -th rightmost digit of ${}^n 3$. Since $n = \text{slog}_3({}^n 3) = \text{slog}_3(G)$, we have finally shown that Graham's number has exactly $\text{slog}_3(G) - 1$ frozen digits and the proof is complete. \square

Lastly, let us note that we can shorten the proof of Theorem 2.3 by invoking [5, Lemma 1]. Still, here we have opted for a more general approach not preliminarily restricted by the selection of the only tetration base taken into account by Lemma 1 of that 2020 paper.

3 The asymptotic phase shift of 3

Since Theorem 2.3 implies that the $\text{slog}_3(G)$ -th rightmost digit of G and the $\text{slog}_3(G)$ -th rightmost digit of 3^G cannot be equal, we can push our investigation even further by asking, for any given positive integer c , which is the $\text{slog}_3(G)$ -th rightmost figure of the difference between ${}^{\text{slog}_3(G)} 3$ (i.e., Graham's number) and ${}^{\text{slog}_3(G)+c} 3$ (e.g., if $c := 719$, then we are interested in the $\text{slog}_3(G)$ -th least significant digit of $G - {}^{\text{slog}_3(G)+729} 3$).

Given the fact that the $\text{slog}_3(G)$ -th rightmost digit of ${}^{\text{slog}_3(G)} 3 - {}^{\text{slog}_3(G)+c} 3$ does not depend on c (since we are assuming $c \in \mathbb{Z}^+$ by hypothesis) and by observing that ${}^{\text{slog}_3(G)} 3 - {}^{\text{slog}_3(G)+c} 3 < 0$, we will solve the problem above by showing that

$$\frac{{}^{\text{slog}_3(G)} 3 - {}^{\text{slog}_3(G)+1} 3}{10^{\text{slog}_3(G)-1}} \equiv 6 \pmod{10}. \quad (3.1)$$

Proposition 3.1. Let n and m be two positive integers such that $m > n$. The congruence class modulo 10 of the difference between the n -th rightmost digit of ${}^n 3$ and the n -th rightmost digit of ${}^m 3$ is 4 or 6. Furthermore, if n is odd, then the n -th least significant digit of ${}^n 3 - {}^m 3$ is 4, otherwise the n -th least significant digit of ${}^n 3 - {}^m 3$ is 6.

Defintion 3.2. Let (a, b) be a pair of positive integers and consider radix-10. For any given tetration base $a \neq 1$ not a multiple of 10, we call *phase shift* of a at height b the congruence class modulo 10 of the difference between the rightmost non-stable digit of ${}^b a$, say the $(\#S(a, b) + 1)$ -th by counting positions from right to left (see [7, Equation (1)]), and the $(\#S(a, b) + 1)$ -th rightmost digit of ${}^{b+1} a$ (e.g., the phase shift of 5 at height 4 is 5 since ${}^4 5 \equiv 68408203125 \pmod{10^{11}}$ and ${}^5 5 \equiv 18408203125 \pmod{10^{11}}$ so that $\frac{68408203125 - 18408203125}{10^{11}-1} \equiv 5 \pmod{10}$). In particular, if \bar{b} (see Definition 2.2) indicates the smallest hyperexponent of a such that its congruence speed is constant (i.e., $V(a, \bar{b}) = V(a)$), we call *asymptotic phase shift* of a (APS_a) the 4-iteration-cycle of the phase shifts generated by $\bar{b}, \bar{b} + 1, \bar{b} + 2$, and $\bar{b} + 3$ (e.g., the asymptotic phase shift of 64 is the cycle $[8, 8, 8, 8]$ that can be halved twice and thus reduced to $[8]$ since $[8, 8, 8, 8] \rightarrow [8, 8] \rightarrow [8]$,

while the asymptotic phase shift of 169 is $[4, 8, 6, 2]$ which cannot be further reduced since $[4, 8]$ is different from $[6, 2]$.

Let $b \in \mathbb{N} - \{0, 1, \dots, \bar{b} - 1\}$. For any given a greater than 1 and not a multiple of 10, a noteworthy property of the phase shift is that it only depends on the congruence modulo 4 of b , which means that the phase shift of a at height b is equal to the phase shift of a at height $b + 4$, the phase shift of a at height $b + 1$ is the same as its phase shift at height $b + 5$, and so forth.

In this regard, we note that if the asymptotic phase shift has four distinct elements, then the first entry plus the third one and the second entry plus the fourth one are both equal to 10, while if the asymptotic phase shift consists of only two entries, their sum is also 10. Moreover, for any given tetration base a as above, all the entries of $\text{APS}(a)$ are constrained to alternatively belong to only one of the following 3 sets, with no exception (see Appendix): $\{2, 4, 6, 8\}$, $\{1, 3, 7, 9\}$, $\{5\}$.

Consequently, although the asymptotic phase shift can have four distinct entries, it is sufficient to calculate only the phase shift at heights b and $b + 1$ in order to fully map the phase shifts of the given tetration base at every height above $\bar{b} - 1$ (e.g., we can safely set $b := \tilde{v}(a) + 2$, where $\tilde{v}(a)$ is stated by [7, Definition 2.1]).

In Chapters 3, 4, 6, and 7 of the 2011 monograph entitled “La strana coda della serie $n^{n^{\cdot^{\cdot^{\cdot}}}}$ ”, the author deeply analyzed the phase shifts of every integer tetration base not a multiple of 10 and the related 4-iteration-cycles (which can be sometimes halved once or twice, depending on a). Here are various examples of the asymptotic phase shift for as many tetration bases coprime to 10:

$$\begin{aligned}
 \text{APS}(9) &= [2]; \\
 \text{APS}(11) &= [4]; \\
 \text{APS}(83) &= [6]; \\
 \text{APS}(53) &= [8]; \\
 \text{APS}(33) &= [2, 8], \text{APS}(39) = [8, 2]; \\
 \text{APS}(43) &= [4, 6], \text{APS}(41) = [6, 4]; \\
 \text{APS}(7) &= [2, 6, 8, 4], \text{APS}(73) = [4, 2, 6, 8], \text{APS}(31) = [8, 4, 2, 6], \text{APS}(29) = [6, 8, 4, 2]; \\
 \text{APS}(23) &= [2, 4, 8, 6], \text{APS}(13) = [6, 2, 4, 8], \text{APS}(77) = [8, 6, 2, 4], \text{APS}(19) = [4, 8, 6, 2]; \\
 \text{APS}(51) &= [5]; \\
 \text{APS}(101) &= [9]; \\
 \text{APS}(901) &= [1, 9]; \\
 \text{APS}(301) &= [3, 9, 7, 1]; \\
 \text{APS}(701) &= [7, 9, 3, 1].
 \end{aligned}$$

In particular, if $a = 3$, the asymptotic phase shift is $[4, 6, 4, 6] \rightarrow [4, 6]$, and then we alternate between the phase shift 4 and the phase shift 6, over and over, according to the parity of b . This modular recurrence is shown in Figure 1, where the difference between each red digit and the corresponding dark digit, right below the red one, is 4 or -6 so that $4 \equiv -6 \pmod{10}$ confirms the stated rule.

Accordingly, we see that the phase shifts of the tetration base 3 at heights 1 to 50 are fully described by the two congruence classes modulo 10 of the following collection of pairs of differences: (3 - 7, 2 - 8; 9 - 3, 9 - 5; 5 - 9, 5 - 1; 0 - 4, 0 - 6; 0 - 4, 6 - 2; 2 - 6, 6 - 2; 3 - 7, 6 - 2; 2 - 6, 9 - 5; 3 - 7, 9 - 5; 0 - 4, 4 - 0; 7 - 1, 3 - 9; 1 - 5, 4 - 0; 5 - 9, 7 - 3; 0 - 4, 0 - 6; 4 - 8, 5 - 1; 1 - 5, 2 - 8; 0 - 4, 2 - 8; 7 - 1, 4 - 0; 3 - 7, 0 - 6; 5 - 9, 7 - 3; 8 - 2, 1 - 7; 4 - 8, 8 - 4; 9 - 3, 6 - 2; 8 - 2, 6 - 2; 9 - 3, 4 - 0; ...).

Hence, we have (-4, -6; 6, 4; -4, 4; -4, -6; -4, 4; -4, 4; -4, 4; -4, 4; -4, 4; -4, 4; 6, -6; -4, 4; -4, 4; -4, -6; -4, 4; -4, -6; -4, -6; 6, 4; -4, -6; -4, 4; 6, -6; -4, 4; 6, 4; 6, 4; 6, 4; ...), which can be split into the odd-hyperexponent difference collection (red entries) (-4, 6, -4, -4, -4, -4, -4, -4, -4, -4, 6, -4, -4, -4, -4, -4, -4, 6, -4, -4, 6, -4, 6, 6, 6, ...) and the even-hyperexponent difference collection (green entries) (-6, 4, 4, -6, 4, 4, 4, 4, 4, 4, -6, 4, 4, -6, 4, -6, -6, 4, -6, 4, -6, 4, 4, 4, 4, ...).

Thus, each red entry belongs to the congruence class 6 modulo 10 while all the green entries are congruent to 4 modulo 10.

For every positive integer b , we easily obtain the b -th rightmost digit of ${}^{b+1}3$ from the b -th rightmost digit of b3 by knowing that the asymptotic phase shift of 3 is [4, 6]. In detail, for each positive odd integer b , 4 plus the b -th least significant digit of b3 is congruent modulo 10 to the b -th least significant digit of ${}^{b+1}3$ (see Figure 1). Conversely, for each positive even integer b , 6 plus the b -th least significant digit of b3 is congruent modulo 10 to the b -th least significant digit of ${}^{b+1}3$.

Let us call B_0 the b -th rightmost digit of b3 and let B_1 indicate the b -th rightmost digit of ${}^{b+1}3$ so that

- if $b \equiv 1 \pmod{2}$ and $B_0 \in \{0, 1, 2, 3, 4, 5\}$, then $B_1 = B_0 + 4$;
- if $b \equiv 1 \pmod{2}$ and $B_0 \in \{6, 7, 8, 9\}$, then $B_1 = ((B_0 + 4) - 10)$ and thus $B_1 = B_0 - 6$;
- if $b \equiv 0 \pmod{2}$ and $B_0 \in \{0, 1, 2, 3\}$, then $B_1 = B_0 + 6$;
- if $b \equiv 0 \pmod{2}$ and $B_0 \in \{4, 5, 6, 7, 8, 9\}$, then $B_1 = ((B_0 + 6) - 10)$ and thus $B_1 = B_0 - 4$.

Thus, given $a := 3$, if $b \equiv 1 \pmod{2}$, then $0 \mapsto 4, 1 \mapsto 5, 2 \mapsto 6, 3 \mapsto 7, 4 \mapsto 8, 5 \mapsto 9, 6 \mapsto 0, 7 \mapsto 1, 8 \mapsto 2, 9 \mapsto 3$, whereas if $b \equiv 0 \pmod{2}$, then $0 \mapsto 6, 1 \mapsto 7, 2 \mapsto 8, 3 \mapsto 9, 4 \mapsto 0, 5 \mapsto 1, 6 \mapsto 2, 7 \mapsto 3, 8 \mapsto 4, 9 \mapsto 5$.

As a consequence, for any given $c \in \mathbb{Z}^+$, the difference between the $\text{slog}_3(G)$ -th rightmost digit of Graham's number and the $\text{slog}_3(G)$ -th rightmost digit of ${}^{\text{slog}_3(G)+c}3$ belongs to the set $\{-4, 6\}$. On the other hand, the difference between the $(\text{slog}_3(G) + 1)$ -th rightmost digit of ${}^{\text{slog}_3(G)+1}3$ and the $(\text{slog}_3(G) + 1)$ -th rightmost digit of ${}^{\text{slog}_3(G)+1+c}3$ is necessarily equal to -6 or 4 .

Now, we observe that $\text{slog}_3(G)$ is congruent to 1 modulo 2 since G is an iterated base 3 integer pentation and 3 is congruent to 1 modulo 2, so we are interested only in the elements of the odd-hyperexponent phase shift set (which we have already verified to belong to the congruence class 6 modulo 10).

Therefore, for any given positive integer c , we can finally conclude that the $\text{slog}_3(G)$ -th least significant digit of $G - {}^{\text{slog}_3(G)+c}3$ is 4 (while the $(\text{slog}_3(G) + 1)$ -th least significant digit of ${}^G3 - {}^{\text{slog}_3(G)+1+c}3$ is 6, indeed).

4 Conclusion

From the trivial observation that Graham's number, G , satisfies (by definition) $G = {}^{slog_3(G)}3$, we have shown that, in the popular decimal system, the last $slog_3(G) - 1$ digits of G are the same as any power tower of the form $3^{3^{\cdot^{\cdot^3}}$ which is at least equal to ${}^{slog_3(G)}3$. Then, as the height of the mentioned power tower goes beyond $slog_3(G)$, the difference between the $slog_3(G)$ -th rightmost digit of G and the $slog_3(G)$ -th rightmost digit of $3^{3^{\cdot^{\cdot^3}}$ becomes congruent to 6 modulo 10, and the stated congruence class does not change anymore as the height of $3^{3^{\cdot^{\cdot^3}}$ continues to grow.

In conclusion, for any positive integer c , we can state that 4 is the $slog_3(G)$ -th least significant digit of ${}^{slog_3(G)+c}3 - G$ while ${}^{slog_3(G)+c}3 - G \equiv 0 \pmod{10^{slog_3(G)-1}}$.

References

- [1] T. H. Cormen, C. E. Leiserson, R. L. Rivest, and C. Stein (2022). *Introduction to Algorithms (Fourth Edition)*, The MIT Press, Cambridge, Massachusetts / London, England.
- [2] R. L. Graham and B. L. Rothschild (1971). Ramsey's Theorem for n -Parameter Sets, *Transactions of the American Mathematical Society*, 159, 257–292.
- [3] N. McWhirter (1979). *Guinness Book of World Records, 1980*, Sterling Publishing Company, New York.
- [4] OEIS Foundation Inc. (2024). *A133613*, The Online Encyclopedia of Integer Sequences. Accessed: Sep. 23 2024, Available online at: <http://oeis.org/A133613>.
- [5] M. Ripà (2020). On the constant congruence speed of tetration, *Notes on Number Theory and Discrete Mathematics*, 26(3), 245–260.
- [6] M. Ripà (2021). The congruence speed formula, *Notes on Number Theory and Discrete Mathematics*, 27(4), 43–61.
- [7] M. Ripà and L. Onnis (2022). Number of stable digits of any integer tetration, *Notes on Number Theory and Discrete Mathematics*, 28(3), 441–457.
- [8] M. Ripà (2024). *Congruence speed of the tetration bases ending with 0*, arXiv:2402.07929v1 [math.NT]. Accessed: Sep. 25 2024, Available online at: <https://arxiv.org/pdf/2402.07929>.
- [9] L. M. Surhone, M. T. Timpledon, and S. F. Marseken (2010). *Super-Logarithm: Mathematics, Tetration, Exponentiation, Nth Root, Logarithm, Abel Function, Logistic Function, Fixed Point, Iterated Logarithm*, Betascript Publishing, Beau Bassin, Mauritius.
- [10] E. W. Weisstein (2024). *Graham's Number*, MathWorld – A Wolfram Web Resource. Accessed: Sep. 23 2024, Available online at: <https://mathworld.wolfram.com/GrahamsNumber.html>.

Appendix

The asymptotic phase shift is well defined only for the tetration bases that are not a multiple of 10 (Definition 3.2) since, for every a ending with 0, there is not any hyperexponent $\bar{b} \in \mathbb{Z}^+$ such that the congruence speed of a is constant, and Equation (2) of [8] describes the very fast growth of the congruence speed of the (nonzero) tetration bases ending with 0.

Nevertheless, let $10^c \mid a$ and $10^{c+1} \nmid a$, where c is a strictly positive integer (as usual). Then, call $h(a)$ the right-hand nonzero digit of such tetration bases multiple of 10 (e.g., $h(200050) = 5$ and $h(81743000) = 3$). We note that, for each hyperexponent $b \in \mathbb{N} - \{0, 1, 2\}$, the phase shift of a at height b is trivially equal to the least significant nonzero digit of $(\frac{a}{c})^{(b-1)a}$, and thus it is 1 if and only if $h(a) \in \{1, 3, 7, 9\}$, 6 if and only if $h(a) \in \{2, 4, 6, 8\}$, and 5 if and only if $h(a) = 5$ (since $b > 2$ implies that $b-1$ is a multiple of 2 and, consequently, is congruent to 0 (mod 4) so that, as long as $10^c \mid a$, the phase shift of a at any height $b \geq 3$ equals the rightmost nonzero digit of $(\frac{a}{c})^4$).

If the last digit of a is 5 (i.e., $5 \mid a$ and $2 \nmid a$), we observe that the asymptotic phase shift is always [5] and, as we also assume that $a \neq 5$, the congruence speed of the mentioned tetration bases is certainly constant from height 3 (see [7, Section 2.1]).

For every $a \in \mathbb{N} - \{0, 1\}$ not a multiple of 10, we conjecture that $\text{APS}(a) \in \{[2], [4], [6], [8], [2, 8], [8, 2], [4, 6], [6, 4], [2, 6, 8, 4], [4, 2, 6, 8], [8, 4, 2, 6], [6, 8, 4, 2], [2, 4, 8, 6], [6, 2, 4, 8], [8, 6, 2, 4], [4, 8, 6, 2], [5], [9], [1, 9], [3, 9, 7, 1], [7, 9, 3, 1]\}$ (and furthermore, if a is also even, only the cases [2], [4], [6], [8], [2, 8], [8, 2], [4, 6], [6, 4], [4, 2, 6, 8], [8, 4, 2, 6], [6, 8, 4, 2], [2, 4, 8, 6], [6, 2, 4, 8], and [4, 8, 6, 2] can occur, given the fact that the congruence speed of every tetration base congruent to 2, 4, 6, 8 modulo 10 becomes constant at most at height 3 [7, p. 448]).