

EXTENSIONS OF THE CONSTANT FAMILY OF HARISH-CHANDRA PAIRS OF $SL_2(\mathbb{R})$

EYAL SUBAG

ABSTRACT. We study and classify algebraic families of Harish-Chandra pairs over the complex affine line and over the complex projective line with generic fiber that is isomorphic to the Harish-Chandra pair of $SL_2(\mathbb{R})$.

CONTENTS

1. Introduction	1
2. Algebraic families of Harish-Chandra pairs	3
2.1. Families of Harish-Chandra pairs over affine base	4
3. Extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{A}_{\mathbb{C}}^1$	5
3.1. Classification of extensions	5
3.2. The universality of $(\mathfrak{g}(1), SO(2, \mathbb{C}))$	9
4. Some properties of extensions	10
4.1. Morphisms between extensions	10
4.2. Morphisms between localizations of extensions	11
4.3. Canonical realization of $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ inside the constant family	12
4.4. The center of $\mathcal{U}(\mathfrak{g}(n))$	13
5. Extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{P}_{\mathbb{C}}^1$	14
5.1. Extensions over $\mathbb{P}_{\mathbb{C}}^1$	14
5.2. Classification of extensions over $\mathbb{P}_{\mathbb{C}}^1$	15
6. Declarations	16
References	16

1. INTRODUCTION

The study of algebraic families of Harish-Chandra pairs and modules was initiated in [6, 7]. The main motivation there was a rigorous mathematical framework for certain limits of Lie groups and their representations that are common in physics and known as *contractions* e.g., see [11, 13]. Many of the algebraic families of Harish-Chandra pairs in the literature are built out of a real reductive Lie group $G(\mathbb{R})$. Such a group has a maximal compact subgroup $K(\mathbb{R})$. The pair (\mathfrak{g}, K) where \mathfrak{g} is the complexification of the Lie algebra of $G(\mathbb{R})$ and K the complexification of $K(\mathbb{R})$, is the Harish-Chandra pair of $G(\mathbb{R})$. The simplest construction of an algebraic family of Harish-Chandra pairs over a smooth complex algebraic variety \mathbf{X} , out of $G(\mathbb{R})$ is a constant family. Explicitly, the pair $(\mathfrak{g}, K)/\mathbf{X} := (\mathcal{O}_{\mathbf{X}} \otimes_{\mathbb{C}} \mathfrak{g}, \mathbf{X} \times_{\text{spec}(\mathbb{C})} K)$ is a constant algebraic family of Harish-Chandra pairs, where $\mathcal{O}_{\mathbf{X}}$ is the sheaf of regular functions on \mathbf{X} .

Families of modules for $(\mathfrak{g}, K)/\mathbf{X}$ naturally appear in representation theory of real reductive groups see e.g., [16, 1, 17].

In [6], two constructions associating with a real reductive Lie group $G(\mathbb{R})$, a non-constant algebraic family of Harish-Chandra pairs over $\mathbb{A}_{\mathbb{C}}^1$ were given. The first family $(\mathfrak{g}_c, \mathbf{K})$ is called *the contraction family* while the second family $(\mathfrak{g}_d, \mathbf{K})$ is called *the deformation family*. Both families share the same constant family of groups $\mathbf{K} = \mathbb{A}_{\mathbb{C}}^1 \times_{\text{spec}(\mathbb{C})} K$ and their restriction to $\mathbb{C}^\times \subset \mathbb{A}_{\mathbb{C}}^1$ is isomorphic to the constant family $(\mathfrak{g}, K)/\mathbb{C}^\times$ associated with $G(\mathbb{R})$. These families have interesting applications.

Families of Harish-Chandra modules for the deformation family were used to give a new algebraic construction for the Mackey-Higson bijection [14]. Full proof for existence of the Mackey-Higson bijection was given in [3]. The bijection has a deep connection with the Baum-Connes conjecture [2, 5, 8, 9, 12].

The contraction family and its modules, in the case of $G(\mathbb{R}) = SO(3, 1)$, were shown to play a central role in the fundamental quantum mechanical system of the hydrogen atom [10, 15].

It is natural to ask whether other similar families of Harish-Chandra pairs can be associated with a real reductive Lie group. The purpose of this paper is to classify and study all families of Harish-Chandra pairs over $\mathbb{A}_{\mathbb{C}}^1$ and over $\mathbb{P}_{\mathbb{C}}^1$ whose restriction to \mathbb{C}^\times is isomorphic to the constant family $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ associated with $SL_2(\mathbb{R})$. Such families of Harish-Chandra pairs are called extensions of the constant family $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$.

We shall now review the structure of the paper and some of our main results. In Section 2 we recall the definitions of all relevant algebraic families. In Section 3 we classify all extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over the complex affine line. We prove that the space of isomorphism classes of such families is in bijection with \mathbb{N}_0 . As explained in Section 2, replacing an extension $(\mathfrak{g}, \mathbf{K})$ by the space of global sections of \mathfrak{g} and the fiber K of \mathbf{K} does not lose any information. In our case $K = SO(2, \mathbb{C})$ and \mathfrak{g} becomes a Lie algebra over the polynomial ring $\mathbb{C}[x]$. Irreducible algebraic representations of $SO(2, \mathbb{C})$ are parameterized by \mathbb{Z} . We let \mathfrak{g}_m denote the K isotopic piece of global sections of \mathfrak{g} corresponding to the irreducible representation with parameter $m \in \mathbb{Z}$.

Theorem 1.1 (Sec. 3, Thm. 3.5). *Let $(\mathfrak{g}, \mathbf{K})$ be an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. Then \mathbf{K} is isomorphic to $\mathbb{A}^1(\mathbb{C}) \times_{\text{spec}(\mathbb{C})} SO(2, \mathbb{C})$ and there exists a basis $\{X, Y, H\}$ of the space of global sections of \mathfrak{g} and $n \in \mathbb{N}_0$, such that $H \in \mathfrak{g}_0$, $X \in \mathfrak{g}_2$, $Y \in \mathfrak{g}_{-2}$ and*

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = x^n H.$$

Moreover, the section H is unique and the sections X and Y are unique up to simultaneous conjugation by $SO(2, \mathbb{C})$.

Denoting the extension corresponding to n by $(\mathfrak{g}(n), SO(2, \mathbb{C}))$. We show that $(\mathfrak{g}(0), SO(2, \mathbb{C}))$ is the constant family, $(\mathfrak{g}(1), SO(2, \mathbb{C}))$ is the contraction family and $(\mathfrak{g}(2), SO(2, \mathbb{C}))$ is the deformation family. We prove that $(\mathfrak{g}(1), SO(2, \mathbb{C}))$ is universal in the sense of the next theorem.

Theorem 1.2 (Sec. 3, Thm. 3.7). *Any extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ is isomorphic to a pullback of $(\mathfrak{g}(1), SO(2, \mathbb{C}))$.*

In Section 4 we show that any extension has a canonical realization inside the constant family. More precisely, for $\mathfrak{g} := \mathfrak{sl}_2(\mathbb{C})$, we let \mathfrak{g}_m , denote the isotypic spaces with respect to $SO(2, \mathbb{C})$ corresponding to the irreducible representation parameterized by $m \in \mathbb{Z}$. A global section F of the constant family $\mathfrak{g}(0)$ is the same thing as an algebraic function from $\mathbb{A}_{\mathbb{C}}^1$ into \mathfrak{g} . Such a function has a well defined n -th derivative $F^{(n)}$ for any $n \in \mathbb{N}_0$. For $n \in \mathbb{N}$, we let $\mathfrak{l}(n)$ be the subspace of the constant family $\mathfrak{g}(0)$ consisting of all sections whose derivatives at 0 up to degree n belong to $\mathfrak{g}_0 \oplus \mathfrak{g}_2$, that is

$$\mathfrak{l}(n) := \{F \in \mathfrak{g}(0) \mid F^{(m)}(0) \in \mathfrak{g}_0 \oplus \mathfrak{g}_2, \forall m \in \{0, 1, \dots, n-1\}\}.$$

Theorem 1.3 (Sec. 4, Thm. 4.6). *For $n \in \mathbb{N}$, the pair $(\mathfrak{l}(n), SO(2, \mathbb{C}))$ is a sub pair of $(\mathfrak{g}(0), SO(2, \mathbb{C}))$ that is isomorphic to $(\mathfrak{g}(n), SO(2, \mathbb{C}))$.*

We also study the space of morphisms between extensions and the structure of the center of the universal enveloping algebra of each $\mathfrak{g}(n)$. In Section 5 we classify extensions over $\mathbb{P}_{\mathbb{C}}^1$.

The author is grateful to Joseph Bernstein for useful remarks. The author thanks Spyridon Afentoulidis-Almpanis for his comments on an earlier version of the manuscript.

2. ALGEBRAIC FAMILIES OF HARISH-CHANDRA PAIRS

In this section we recall the definition of an algebraic family of Harish-Chandra pairs over a complex algebraic variety as given in [6].

Let \mathbf{X} be a smooth complex algebraic variety. We denote its structure sheaf of regular functions by $\mathcal{O}_{\mathbf{X}}$. An algebraic family of complex Lie algebras \mathfrak{g} over \mathbf{X} is a locally free sheaf of $\mathcal{O}_{\mathbf{X}}$ -modules which is also a sheaf of Lie algebras over \mathbf{X} . In particular \mathfrak{g} is equipped with Lie brackets

$$[_, _] : \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathfrak{g},$$

which is $\mathcal{O}_{\mathbf{X}}$ bi-linear, skew-symmetric and satisfying the Jacobi identity. An algebraic family of complex algebraic groups \mathbf{G} over \mathbf{X} is a smooth morphism between smooth complex algebraic varieties

$$\mathbf{G} \longrightarrow \mathbf{X},$$

that carries the structure of a group scheme over \mathbf{X} . For every family of complex algebraic groups there is an associated \mathbf{G} -equivariant family $\text{Lie}(\mathbf{G})$ of complex Lie algebras over \mathbf{X} . An algebraic family of Harish-Chandra pairs over \mathbf{X} is a pair $(\mathfrak{g}, \mathbf{K})$ with \mathbf{K} being an algebraic family of complex algebraic groups over \mathbf{X} , \mathfrak{g} being a \mathbf{K} -equivariant algebraic family of Lie algebras over \mathbf{X} . Along with a \mathbf{K} -equivariant morphism of Lie algebras

$$j : \text{Lie}(\mathbf{K}) \longrightarrow \mathfrak{g},$$

such that the action of $\text{Lie}(\mathbf{K})$ on \mathfrak{g} that is obtained as the differential of the action of \mathbf{K} on \mathfrak{g} coincides with the action obtained by j composed with the adjoint action of \mathbf{G} on \mathfrak{g} .

Example 2.1. Let (\mathfrak{g}, K) be a complex Harish-Chandra pair. Note that this is the same thing as an algebraic family of Harish-Chandra pairs over the trivial one point complex algebraic variety. Then for any smooth complex affine algebraic variety $(\mathbf{X}, \mathcal{O}_{\mathbf{X}})$, the pair $(\mathcal{O}_{\mathbf{X}} \otimes_{\mathbb{C}} \mathfrak{g}, \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K)$ is an algebraic family of Harish-Chandra pairs with all of its fibers canonically isomorphic to (\mathfrak{g}, K) . The equivariant Lie algebra morphism $j : \text{Lie}(\mathbf{X} \times_{\text{Spec}(\mathbb{C})} K) = \mathcal{O}_{\mathbf{X}} \otimes_{\mathbb{C}} \text{Lie}(K) \rightarrow \mathfrak{g}$ is given by $j(f \otimes Y) = f j_0(Y)$ for every $f \in \mathcal{O}_{\mathbf{X}}$ and $Y \in \text{Lie}(K)$, and where $j_0 : \text{Lie}(K) \rightarrow \mathfrak{g}$ is the morphism that is given with (\mathfrak{g}, K) .

A family of Harish-Chandra pairs as in the last example is called a *constant* family and we shall denote it by $(\mathfrak{g}, K)/\mathbf{X}$. Similarly a family of groups of the form $\mathbf{X} \times_{\text{Spec}(\mathbb{C})} K$ is called constant, and a family of Lie algebras of the form $\mathcal{O}_{\mathbf{X}} \otimes_{\mathbb{C}} \mathfrak{g}$ is called constant. Note that a section of $\mathcal{O}_{\mathbf{X}} \otimes_{\mathbb{C}} \mathfrak{g}$ over an open subset $U \subset \mathbf{X}$ is canonically identified with an algebraic function from U into \mathfrak{g} .

2.1. Families of Harish-Chandra pairs over affine base. In this section we describe families of Harish-Chandra pairs with a constant group scheme and over affine base in terms of modules over a commutative ring. This setup appears in many places throughout the paper.

Assume that \mathbf{X} is a smooth complex affine algebraic variety. Denote its ring of global regular functions by $R_{\mathbf{X}}$. Let $(\mathfrak{g}, \mathbf{K})$ be an algebraic family of Harish-Chandra pairs over \mathbf{X} with $\mathbf{K} = \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K$ for some complex algebraic group K . By looking on its space of global sections, such a family is equivalent to following data:

- (1) A locally free $R_{\mathbf{X}}$ -module \mathfrak{g} that is also a Lie algebra over $R_{\mathbf{X}}$. (Note that by abuse of notation we denote the family of Lie algebras and its space of global sections by the same symbol).
- (2) An algebraic action $\alpha : K \rightarrow \text{Aut}_{R_{\mathbf{X}}}(\mathfrak{g})$ of K on \mathfrak{g} by automorphisms of Lie algebras over $R_{\mathbf{X}}$.
- (3) A K -equivariant embedding of complex Lie algebras $j : \text{Lie}(K) \rightarrow \mathfrak{g}$.

Such that for every $Y \in \mathfrak{g}, Z \in \text{Lie}(K)$,

$$[j(Z), Y] = d\alpha(Z)Y,$$

where $d\alpha$ stands for the action of $\text{Lie}(K)$ on \mathfrak{g} via endomorphisms of Lie algebras over $R_{\mathbf{X}}$, $d\alpha : \text{Lie}(K) \rightarrow \text{End}_{R_{\mathbf{X}}}(\mathfrak{g})$, that is the differential of α .

Hence under the above mentioned assumptions we replace the pair $(\mathfrak{g}, \mathbf{K})$ by (\mathfrak{g}, K) where in the latter pair \mathfrak{g} is a Lie algebra over $R_{\mathbf{X}}$ and K a complex algebraic group.

Keeping the above assumptions of \mathbf{X} being smooth affine and \mathbf{K} being constant, if $(\mathfrak{g}_1, \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K_1)$ and $(\mathfrak{g}_2, \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K_2)$ are two algebraic families of Harish-Chandra pairs over \mathbf{X} with K_1, K_2 complex algebraic groups, then a morphism of pairs from $(\mathfrak{g}_1, \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K_1)$ to $(\mathfrak{g}_2, \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K_2)$ is given by a pair of maps $(\psi_{\mathfrak{g}}, \psi_K)$ with $\psi_{\mathfrak{g}}$ being a morphism of Lie algebras over $R_{\mathbf{X}}$ and $\psi_K : K_1 \rightarrow K_2$ being a morphism of complex algebraic groups such that the following two conditions hold.

- (1) *Twisted equivariance:* $\psi_{\mathfrak{g}}(\alpha_1(k)X) = \alpha_2(\psi_K(k))\psi_{\mathfrak{g}}(X), \forall X \in \mathfrak{g}_1, k \in K_1$, and where $\alpha_i : K_i \rightarrow \text{Aut}_{R_{\mathbf{X}}}(\mathfrak{g}_i)$ is the action map for $i \in \{1, 2\}$.

- (2) *Compatibility with embeddings:* $\psi_{\mathfrak{g}}(j_1(Z)) = j_2(d\psi_K(Z))$, $\forall Z \in \text{Lie}(K_1)$ and where $j_i : \text{Lie}(K_i) \rightarrow \mathfrak{g}_i$ is the K_i -equivariant embedding for $i \in \{1, 2\}$, and $d\psi_K : \text{Lie}(K_1) \rightarrow \text{Lie}(K_2)$ the differential of ψ_K .

We denote the space of such morphisms by

$$\text{Hom}_{\text{pairs}}((\mathfrak{g}_1, K_1), (\mathfrak{g}_2, K_2)).$$

Remark 2.2. When $\psi_{\mathfrak{g}}$ is an isomorphism and the center of \mathfrak{g}_1 is trivial then compatibility with embeddings follows from twisted equivariance.

Remark 2.3. For $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}_{\text{pairs}}((\mathfrak{g}_1, K_1), (\mathfrak{g}_2, K_2))$ with a one-to-one differential $d\psi_K : \text{Lie}(K_1) \rightarrow \text{Lie}(K_2)$, we define a pullback pair $\psi_K^*(\mathfrak{g}_2, K_2)$ of (\mathfrak{g}_2, K_2) with respect to ψ_K , in which $\psi_K^*(\mathfrak{g}_2, K_2) = (\mathfrak{g}_2, K_1)$ and the action of K_1 on \mathfrak{g}_2 is given by $\alpha_2 \circ \psi_K : K_1 \rightarrow \text{Aut}_{R_{\mathfrak{X}}}(\mathfrak{g}_2)$, and the K_1 -equivariant Lie algebras embedding is given by $j_2 \circ d\psi_K : \text{Lie}(K_1) \rightarrow \mathfrak{g}_2$. Of course, if $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}_{\text{pairs}}((\mathfrak{g}_1, K_1), (\mathfrak{g}_2, K_2))$ then $(\psi_{\mathfrak{g}}, \mathbb{I}) \in \text{Hom}_{\text{pairs}}((\mathfrak{g}_1, K_1), \psi_K^*(\mathfrak{g}_2, K_2))$.

3. EXTENSIONS OF $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ OVER $\mathbb{A}_{\mathbb{C}}^1$

In this section we classify all extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{A}_{\mathbb{C}}^1$ and show that any extension is a pullback of the contraction family.

3.1. Classification of extensions. Here we classify all extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{A}_{\mathbb{C}}^1$.

Throughout the rest of the paper, we let \mathbf{X} be the complex algebraic variety $\mathbb{A}_{\mathbb{C}}^1$, and let \mathbf{X}_0 the subvariety consisting of \mathbf{X} take away a single point. Fixing a coordinate function on \mathbf{X} we shall identify \mathbf{X} with \mathbb{C} and its ring of global regular functions with $\mathbb{C}[x]$, and \mathbf{X}_0 with \mathbb{C}^\times and its ring of global regular functions with $\mathbb{C}[x, x^{-1}]$.

Definition 3.1. An algebraic family of Harish-Chandra pairs $(\mathfrak{g}, \mathbf{K})$ over $\mathbb{A}_{\mathbb{C}}^1$ is called an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$, if \mathbf{K} is a constant algebraic family of complex algebraic groups and the restriction of $(\mathfrak{g}, \mathbf{K})$ to \mathbb{C}^\times is isomorphic to $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$.

Theorem 3.2. *Let $(\mathfrak{g}, \mathbf{K})$ be an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. Then \mathbf{K} is isomorphic to $\mathbb{A}^1(\mathbb{C}) \times_{\text{Spec}(\mathbb{C})} SO(2, \mathbb{C})$ and there exists a basis $\{X, Y, H\}$ of the space of global sections of \mathfrak{g} and $n \in \mathbb{N}_0$, such that $H \in j(\text{Lie}(\mathbf{K}))$ and*

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = x^n H.$$

Moreover, the section H is unique up to a sign and for a fixed H the sections X and Y are unique up to simultaneous conjugation by $SO(2, \mathbb{C})$.

Proof. Since \mathbf{K} is a constant family there exists a complex algebraic group K such that $\mathbf{K} = \mathbf{X} \times_{\text{Spec}(\mathbb{C})} K$. Let $\iota : \mathbf{X}_0 \rightarrow \mathbf{X}$ be the inclusion morphism. Since $\iota^* \mathbf{K} \simeq \mathbf{X}_0 \times_{\text{Spec}(\mathbb{C})} K \simeq \mathbf{X}_0 \times_{\text{Spec}(\mathbb{C})} SO(2, \mathbb{C})$, without loss of generality we can assume that $K = SO(2, \mathbb{C})$. Since \mathbf{X} and \mathbf{X}_0 are smooth affine algebraic varieties and since all group schemes are constant we are in the setup of Subsection 2.1 and we can consider the corresponding spaces of global sections. In this description of global sections over affine base, the restriction $\iota^* \mathfrak{g} = \mathfrak{g}|_{\mathbf{X}_0}$, of \mathfrak{g} to \mathbf{X}_0 is given by the localization $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}$.

By our assumptions the family $(\mathfrak{g}|_{X_0}, SO(2, \mathbb{C}))$ is isomorphic to $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. We realize $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))$ concretely as

$$\begin{aligned}\mathfrak{sl}_2(\mathbb{C}) &= \left\{ \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \middle| a, b, c \in \mathbb{C} \right\} \\ SO(2, \mathbb{C}) &= \left\{ k(z) := \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} \middle| z \in \mathbb{C}^\times \right\}.\end{aligned}$$

We shall realize $\text{Lie}(SO(2, \mathbb{C}))$ as

$$\mathfrak{so}_2(\mathbb{C}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & -a \end{pmatrix} \middle| a \in \mathbb{C} \right\}.$$

The embedding $\tilde{j}_0 : \text{Lie}(SO(2, \mathbb{C})) \longrightarrow \mathbb{C}[x] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C})$ is given by $\tilde{j}_0(Z) = 1 \otimes j_0(Z)$ for every $Z \in \text{Lie}(SO(2, \mathbb{C}))$, where $j_0 : \text{Lie}(SO(2, \mathbb{C})) \hookrightarrow \mathfrak{sl}_2(\mathbb{C})$ is the inclusion map. Any isomorphism

$$(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}_{\text{pairs}}((\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C}), K), (\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}, K))$$

has exactly two possibilities for the algebraic isomorphisms ψ_K of $SO(2, \mathbb{C})$. We denote them by ψ_K^\pm , explicitly

$$\psi_K^\pm(g) = g^{\pm 1}, \quad \forall g \in K.$$

We first assume that $\psi_K = \psi_K^+ = \mathbb{I}_K$. Since \mathfrak{g} is a K module (as a complex vector space) it has a canonical isotypical decomposition $\mathfrak{g} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{g}_n$ where for any $k(z) \in SO(2, \mathbb{C})$ and $W \in \mathfrak{g}_n$, we have $\alpha(k(z))W = z^n W$. Since the actions of $\mathbb{C}[x]$ and K commute each \mathfrak{g}_n is a $\mathbb{C}[x]$ -module. Since $\mathbb{C}[x]$ is a principal ideal domain and since \mathfrak{g} is a free $\mathbb{C}[x]$ -module then each \mathfrak{g}_n is a free $\mathbb{C}[x]$ -module. Similarly the isotypical decomposition of $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}$ is given by $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g} = \bigoplus_{n \in \mathbb{Z}} (\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}_n)$. The twisted equivariance property and the fact that $\psi_K = \mathbb{I}_K$ imply that $\psi_{\mathfrak{g}}$ is an honest K -equivariant morphism of Lie algebras over $\mathbb{C}[x, x^{-1}]$. The $SO(2, \mathbb{C})$ isotypic decomposition of $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C})$ is given by

$$\left(\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} e_{12} \right) \oplus \left(\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} (e_{11} - e_{22}) \right) \oplus \left(\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} e_{21} \right)$$

where e_{ij} is the two-by-two matrix having a unique non-zero entry at the (i, j) -th entry which is equal to 1, and the action of K on $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C})$ is determined by

$$\begin{aligned}k(z) \cdot 1 \otimes (e_{11} - e_{22}) &= 1 \otimes (e_{11} - e_{22}) \\ k(z) \cdot (1 \otimes e_{12}) &= z^2 \otimes e_{12} \\ k(z) \cdot (1 \otimes e_{21}) &= z^{-2} \otimes e_{21}.\end{aligned}$$

The chain of K -equivariant $\mathbb{C}[x]$ -linear maps

$$\begin{aligned}\mathfrak{g} &\longrightarrow \mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g} \longrightarrow \mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C}) \\ s &\longmapsto 1 \otimes s \quad \longmapsto (\psi_{\mathfrak{g}}^\pm)^{-1}(1 \otimes s)\end{aligned}$$

takes a basis over $\mathbb{C}[x]$ for each isotypic piece to a basis over $\mathbb{C}[x, x^{-1}]$ of the corresponding isotypic piece. Hence \mathfrak{g}_n is a rank-one $\mathbb{C}[x]$ -module for $n \in \{-2, 0, 2\}$ and zero otherwise. For $n \in \{-2, 0, 2\}$, we let X_n be a basis over $\mathbb{C}[x]$ of \mathfrak{g}_n . Since $\psi_{\mathfrak{g}}$ restricts to an isomorphism from any isotypic component of $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C})$ to the

corresponding isotypic component of $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}$, there must be $c_-, c_0, c_+ \in \mathbb{C}^\times$ and $m_-, m_0, m_+ \in \mathbb{Z}$ such that

$$\begin{aligned}\psi_{\mathfrak{g}}(1 \otimes e_{21}) &= c_- x^{m_-} \otimes X_{-2} \\ \psi_{\mathfrak{g}}(1 \otimes (e_{11} - e_{22})) &= c_0 x^{m_0} \otimes X_0 \\ \psi_{\mathfrak{g}}(1 \otimes e_{12}) &= c_+ x^{m_+} \otimes X_2.\end{aligned}$$

For $i, j, k \in \{-2, 0, 2\}$, we let $c_{i,j}^k(t) \in \mathbb{C}[x]$ be the structure constants of the Lie algebra \mathfrak{g} with respect to the ordered basis $\{X_{-2}, X_0, X_2\}$. Note that $c_{i,j}^k(x)$ are also the structure constants of $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}$ with respect to the ordered basis $\{1 \otimes X_{-2}, 1 \otimes X_0, 1 \otimes X_2\}$. Clearly

$$[\psi_{\mathfrak{g}}(1 \otimes (e_{11} - e_{22})), \psi_{\mathfrak{g}}(1 \otimes e_{12})] = \psi_{\mathfrak{g}}([1 \otimes (e_{11} - e_{22}), 1 \otimes e_{12}])$$

which is equivalent to

$$\begin{aligned}[c_0 x^{m_0} \otimes X_0, c_+ x^{m_+} \otimes X_2] &= \psi_{\mathfrak{g}}(2(1 \otimes e_{12})) = 2c_+ x^{m_+} \otimes X_2 \iff \\ \sum_{k \in \{-2, 0, 2\}} c_0 x^{m_0} c_+ x^{m_+} c_{0,2}^k(x) \otimes X_k &= 2c_+ x^{m_+} \otimes X_2.\end{aligned}$$

Hence $c_{0,2}^{-2}(x) \equiv 0$, $c_{0,2}^0(x) \equiv 0$, and $c_0 x^{m_0} c_{0,2}^2 = 2$. As a result $c_{0,2}^2(x) = \frac{2}{c_0} x^{-m_0}$ and $-m_0 \in \mathbb{N}_0$. Similarly, by examining the commutators $[\psi_{\mathfrak{g}}(1 \otimes (e_{11} - e_{22})), \psi_{\mathfrak{g}}(1 \otimes e_{21})]$ and $[\psi_{\mathfrak{g}}(1 \otimes e_{12}), \psi_{\mathfrak{g}}(1 \otimes e_{21})]$ we obtain that $c_{0,-2}^0(x) \equiv 0$, $c_{0,-2}^2(x) \equiv 0$, $c_{0,-2}^{-2}(x) = -\frac{2}{c_0} x^{-m_0}$, $c_{2,-2}^{-2}(x) \equiv 0$, $c_{2,-2}^2(x) \equiv 0$, $c_{2,-2}^0(x) = \frac{c_0}{c_- c_+} x^{m_0 - m_2 - m_-}$ and $m_0 - m_2 - m_- \in \mathbb{N}_0$.

We define $n = m_0 - m_+ - m_-$. and $H = c_0 X_0$, $X = c_+ X_2$, $Y = c_- X_{-2}$, and observe that

$$[H, X] = 2x^{-m_0} X, [H, Y] = -2x^{-m_0} Y, [X, Y] = x^n H.$$

The embedding $1 \otimes j$ of $\text{Lie}(K)$ into $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}$ is given by

$$(1 \otimes j)(Z) = 1 \otimes j(Z), \quad \forall Z \in \text{Lie}(K).$$

By compatibility with embeddings of $(\psi_{\mathfrak{g}}, \psi_K)$,

$$\begin{aligned}(1 \otimes j)(d\psi_K(e_{11} - e_{22})) &= \psi_{\mathfrak{g}}(\tilde{j}_0(e_{11} - e_{22})) \iff \\ 1 \otimes j(e_{11} - e_{22}) &= \psi_{\mathfrak{g}}(1 \otimes (e_{11} - e_{22})).\end{aligned}$$

Hence $1 \otimes j(e_{11} - e_{22}) = c_0 x^{m_0} \otimes X_0 = x^{m_0} \otimes H$. This equality is in $\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}$. It implies that $j(e_{11} - e_{22}) = x^{m_0} H$ in \mathfrak{g} . Recall that $m_0 \in -\mathbb{N}_0$ hence m_0 must be equal to 0. Hence the commutation relations of the bases $\{H, X, Y\}$ are as stated in the theorem. In addition $H = j(e_{11} - e_{22}) \in 1 \otimes \text{Lie}(K) \subset \text{Lie}(\mathbf{K})$.

When

$$(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}_{\text{pairs}}((\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C}), K), (\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}, K))$$

is an isomorphism in which $\psi_K(g) = \psi_K^{-1}(g) = g^{-1}$ for any $g \in K$, then as explained in Remark 2.3,

$$(\psi_{\mathfrak{g}}, \mathbb{I}_K) \in \text{Hom}_{\text{pairs}}((\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C}), K), \psi_K^*(\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}, K)).$$

This allows us to use the previous case with $\psi_K = \mathbb{I}_K$ and get the required basis. In this case the basis $\{X, Y, H\}$ must satisfy $X \in \mathfrak{g}_{-2}$, $Y \in \mathfrak{g}_2$ and $H \in \mathfrak{g}_0$ in contrary to

case with $\psi_K = \mathbb{I}_K$, in which $X \in \mathfrak{g}_2$, $Y \in \mathfrak{g}_{-2}$ and $H \in \mathfrak{g}_0$. This finishes the proof of existence.

We shall now prove uniqueness. Given two bases with the required properties, $B := \{H, X, Y\}$ and $B' := \{H', X', Y'\}$, then the $\mathbb{C}[x]$ -linear map $L : \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying $L(H) = H'$, $L(X) = X'$, $L(Y) = Y'$ is an isomorphism of Lie algebras over $\mathbb{C}[x]$. Both H and H' are non zero complex multiple of X_0 and so there is $\alpha_H \in \mathbb{C}^\times$ such that $L(H) = H' = \alpha_H H$. Since the eigenvalues of ad_H and of $\text{ad}_{H'}$ must be equal to $\{-2, 0, 2\}$ we must have $\alpha_H = \pm 1$. If $\alpha_H = 1$, each of the elements X and X' is a basis for \mathfrak{g}_2 and as a result the two are proportional by an invertible element of $\mathbb{C}[x]$. Similarly each of Y and Y' is a basis for \mathfrak{g}_{-2} and as a result the two are proportional by an invertible element of $\mathbb{C}[x]$. The commutation relations imply that the two proportionality constants are the inverse of each other. Namely, there exist $\lambda \in \mathbb{C}^\times = (\mathbb{C}[x])^\times$ such that

$$X' = \lambda X, Y = \lambda^{-1} Y'.$$

For any square root of λ , the matrix $k(\sqrt{\lambda}) \in SO(2, \mathbb{C})$ satisfies

$$k(\sqrt{\lambda})X = \lambda X = X', k(\sqrt{\lambda})Y = \lambda^{-1} Y = Y'.$$

The case of $\alpha_H = -1$ is handled similarly. \square

Remark 3.3. By replacing $SO(2, \mathbb{C})$ with $O(2, \mathbb{C})$ in the above theorem, we can show that any two bases with the required properties are in the same $O(2, \mathbb{C})$ -orbit.

Corollary 3.4. *There is a bijection between \mathbb{N}_0 and the set of equivalence classes of extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$.*

Proof. We start by showing that if two extensions have bases as in the last theorem with the same $n \in \mathbb{N}_0$, then the extensions are isomorphic. As before we shall only consider the spaces of global sections for the families of Lie algebras and replace the constant group schemes with fiber $K = SO(2, \mathbb{C})$ by their fiber. Let (\mathfrak{g}, K) and (\mathfrak{g}', K) be two extensions having bases $B := \{X, Y, H\}$ and $B' := \{X', Y', H'\}$ of the corresponding spaces of global sections such that $H \in j(\text{Lie}(\mathbf{K}))$, $H' \in j'(\text{Lie}(\mathbf{K}))$ and

$$\begin{aligned} [H, X] &= 2X, [H, Y] = -2Y, [X, Y] = x^n H, \\ [H', X'] &= 2X', [H', Y'] = -2Y', [X', Y'] = x^n H'. \end{aligned}$$

Define $\nu_{\mathfrak{g}}^\pm : \mathfrak{g} \rightarrow \mathfrak{g}'$ via

$$\begin{aligned} \nu_{\mathfrak{g}}^+(H) &= H', \quad \nu_{\mathfrak{g}}^+(X) = X', \quad \nu_{\mathfrak{g}}^+(Y) = Y', \\ \nu_{\mathfrak{g}}^-(H) &= -H', \quad \nu_{\mathfrak{g}}^-(X) = X', \quad \nu_{\mathfrak{g}}^-(Y) = Y'. \end{aligned}$$

Both maps respect the Lie brackets, and exactly one of $(\nu_{\mathfrak{g}}^\pm, \mathbb{I}_K)$, is twisted-equivariance. By Remark, 2.2 it will also be compatible with the embeddings and hence an isomorphism of pairs.

Now given $n \in \mathbb{N}_0$, we define an algebraic family of Lie algebras \mathfrak{g} via a basis $B := \{X, Y, H\}$ for its space of global sections, namely $\mathfrak{g} = \mathbb{C}[x]Y \oplus \mathbb{C}[x]H \oplus \mathbb{C}[x]X$. We define Lie brackets on \mathfrak{g} via

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = x^n H.$$

We define an action of $SO(2, \mathbb{C})$ on \mathfrak{g} by declaring that the isotypic pieces are $\mathfrak{g}_0 = \mathbb{C}[x]H$, $\mathfrak{g}_2 = \mathbb{C}[x]X$, $\mathfrak{g}_{-2} = \mathbb{C}[x]Y$. The embedding $j : \mathfrak{so}(2, \mathbb{C}) \rightarrow \mathfrak{g}$ is the unique \mathbb{C} linear map satisfying $j(e_{11} - e_{22}) = H$. This is indeed an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. \square

From the last proof we see that Theorem 3.2 can be reformulated as follows.

Theorem 3.5. *Let $(\mathfrak{g}, \mathbf{K})$ be an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. Then \mathbf{K} is isomorphic to $\mathbb{A}^1(\mathbb{C}) \times_{\text{Spec}(\mathbb{C})} SO(2, \mathbb{C})$ and there exists a basis $\{X, Y, H\}$ of the space of global sections of \mathfrak{g} and $n \in \mathbb{N}_0$, such that $H \in \mathfrak{g}_0$, $X \in \mathfrak{g}_2$, $Y \in \mathfrak{g}_{-2}$ and*

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = x^n H.$$

Moreover, the section H is unique and the sections X and Y are unique up to simultaneous conjugation by an element of $SO(2, \mathbb{C})$.

For every $n \in \mathbb{N}_0$, we let $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ be a fixed representative for the algebraic family of Harish-Chandra pairs corresponding to n from the last corollary. We fix a basis $\{X_n, Y_n, H_n\}$ for $\mathfrak{g}(n)$ such that $H_n \in j(\text{Lie}(SO(2, \mathbb{C}))) = \mathfrak{g}(n)_0$, $X_n \in \mathfrak{g}(n)_2$, $Y_n \in \mathfrak{g}(n)_{-2}$ and

$$[H_n, X_n] = 2X_n, [H_n, Y_n] = -2Y_n, [X_n, Y_n] = x^n H_n.$$

For these choices we have $j(e_{11} - e_{22}) = H_n$. We shall keep these notations throughout the paper.

Remark 3.6. We note that the family $(\mathfrak{g}(0), SO(2, \mathbb{C}))$ is the constant family $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}$, the family $(\mathfrak{g}(1), SO(2, \mathbb{C}))$ is the *contraction family* of $SL_2(\mathbb{R})$, see section 2.1.3 of [6], and the family $(\mathfrak{g}(2), SO(2, \mathbb{C}))$ is the *deformation family* of $SL_2(\mathbb{R})$, see section 2.1.2 of [6].

3.2. The universality of $(\mathfrak{g}(1), SO(2, \mathbb{C}))$. The next result shows that $(\mathfrak{g}(1), SO(2, \mathbb{C}))$ is universal among extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$.

Theorem 3.7. *Any extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ is isomorphic to a pullback of $(\mathfrak{g}(1), SO(2, \mathbb{C}))$.*

Proof. Let $(\mathfrak{g}, \mathbf{K})$ be an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. Recall that for any $\mu : \mathbb{A}_{\mathbb{C}}^1 \rightarrow \mathbb{A}_{\mathbb{C}}^1$, $\mu^*(\mathfrak{g})$ is the quotient of the $\mathbb{C}[x]$ -module $\mathbb{C}[x] \otimes_{\mathbb{C}} \mathfrak{g}$ by the submodule generated by

$$\{1 \otimes fX - (f \circ \mu) \otimes X \mid f \in \mathbb{C}[x], X \in \mathfrak{g}\}.$$

By abuse of notation the cosets in $\mu^*(\mathfrak{g})$ shall be denoted by their representatives. The formula

$$[f \otimes X, g \otimes Y] = fg \otimes [X, Y], \quad f \otimes X, g \otimes Y \in \mathbb{C}[x] \otimes_{\mathbb{C}} \mathfrak{g}$$

defines Lie brackets on $\mu^*(\mathfrak{g})$. By corollary 3.4, we can assume that there is some $n \in \mathbb{N}_0$, such that $(\mathfrak{g}, \mathbf{K})$ is equal to $(\mathfrak{g}(n), SO(2, \mathbb{C}))$. For $\mu_n(x) = x^n$, the $\mathbb{C}[x]$ -linear map $L : \mathfrak{g}(n) \rightarrow \mu_n^* \mathfrak{g}(1)$ satisfying

$$L(X_n) = 1 \otimes X_1, \quad L(Y_n) = 1 \otimes Y_1, \quad L(H_n) = 1 \otimes H_1$$

together with $\psi_K^\dagger : SO(2, \mathbb{C}) \rightarrow SO(2, \mathbb{C})$ defines an isomorphism of families of Harish-Chandra pairs from $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ onto $(\mu_n^* \mathfrak{g}(1), SO(2, \mathbb{C}))$. \square

The next corollary shows that any extension is integrable, that is, it comes from an algebraic family of complex algebraic groups.

Corollary 3.8. *Let $(\mathfrak{g}, \mathbf{K})$ be an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$. Then there is an algebraic family \mathbf{G} of complex algebraic groups over $\mathbb{A}_{\mathbb{C}}^1$ containing \mathbf{K} and having its Lie algebra $\text{Lie}(\mathbf{G})$ equal to \mathfrak{g} .*

Proof. By the last Theorem we can assume that the extension $(\mathfrak{g}, \mathbf{K})$ is equal to $(\mu_n^* \mathfrak{g}(1), SO(2, \mathbb{C}))$ for some $n \in \mathbb{N}_0$. In [4] it was shown that for the contraction family $(\mathfrak{g}_c, \mathbf{K})$ of a real reductive group $G(\mathbb{R})$ there exists an algebraic family of complex algebra groups \mathbf{G}_c over $\mathbb{A}_{\mathbb{C}}^1$ containing \mathbf{K} and whose Lie algebra is equal to \mathfrak{g}_c . We let \mathbf{G}_1 be the above mentioned family of groups in the case of $G(\mathbb{R}) = SL_2(\mathbb{R})$. In this case $(\mathfrak{g}_c, \mathbf{K}) = (\mathfrak{g}(1), SO(2, \mathbb{C}))$. Since in our case the operation of pulling back with respect to μ_n and the operation of taking a Lie algebra for a family of groups commute, it follows that $\mu_n^* \mathbf{G}_1$ will have the required properties. \square

Remark 3.9. Using Theorem 3.7 we can give the space of equivalence classes of extensions the structure of a monoid in which multiplication is determined by composition of the maps μ_n defining the pullbacks. With this monoid structure the bijection in Corollary 3.4 is an isomorphism of monoids.

4. SOME PROPERTIES OF EXTENSIONS

In this sections we study the space of morphisms between any two extensions. We give a canonical embedding of any extension in the constant family and study the center of the universal enveloping algebra of any extension.

4.1. Morphisms between extensions. In this section we study the space of morphisms between a given pair of extensions. For $m, n \in \mathbb{N}_0$, we introduce the notations

$$\begin{aligned} \text{Hom}(m, n) &:= \text{Hom}_{\text{pairs}}((\mathfrak{g}(m), SO(2, \mathbb{C})), (\mathfrak{g}(n), SO(2, \mathbb{C}))), \\ \text{Hom}(m, n)_{\pm} &:= \{(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}(m, n) \mid \psi_K(k) = k^{\pm 1}, \forall k \in SO(2, \mathbb{C})\}. \end{aligned}$$

Note that by remark 2.3, we have a bijection between $\text{Hom}(m, n)_+$ and $\text{Hom}(m, n)_-$. We denote the space of complex polynomials in one variable of degree at most n by $\mathbb{C}_n[x] = \text{span}_{\mathbb{C}}\{1, x, x^2, \dots, x^n\}$ and its subset of homogeneous polynomials by

$$\mathbb{C}_n^{\text{hom}}[x] = \{f \in \mathbb{C}_n[x] \mid \exists k \in \mathbb{N}_0, f(tx) = t^k f(x), \forall t, x \in \mathbb{C}\}.$$

Theorem 4.1. *The space $\text{Hom}(m, n)$ is nonzero if and only if $m \geq n$. For $m \geq n$,*

$$\text{Hom}(m, n)_+ \simeq \mathbb{C}_{m-n}^{\text{hom}}[x] \simeq \text{Hom}(m, n)_-,$$

as sets.

Proof. For $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}(m, n)_+$, we must have

$$\psi_{\mathfrak{g}}(X_m) = f_X(x)X_n, \quad \psi_{\mathfrak{g}}(Y_m) = f_Y(x)Y_n, \quad \psi_{\mathfrak{g}}(H_m) = f_H(x)H_n$$

for some $f_X, f_Y, f_H \in \mathbb{C}[x]$. Since $\psi_{\mathfrak{g}}$ is a morphism of Lie algebras, the commutation relations imply that

$$f_H(x)f_X(x) = f_X(x), \quad f_H(x)f_Y(x) = f_Y(x), \quad f_X(x)f_Y(x)x^n = x^m f_H(x).$$

If $f_X(x) \equiv 0$, we must have $f_Y(x) \equiv 0$ and $f_H(x) \equiv 0$ and hence $\psi_{\mathfrak{g}}$ is the zero map. Similarly if $f_Y(x) \equiv 0$, $\psi_{\mathfrak{g}}$ is the zero map. Now assume that $f_X(x)f_Y(x)$ is not the zero polynomial then $f_H(x) \equiv 1$ and we must have $f_X(x)f_Y(x)x^n = x^m$. Hence we must have $n \leq m$ and $f_X(x)f_Y(x) = x^{m-n}$. In particular the only possibilities are

$$f_X(x) = cx^k, f_Y(x) = c^{-1}x^{m-n-k}, \quad c \in \mathbb{C}^\times, k \in \{0, 1, 2, \dots, m-n\}.$$

We shall denote the above mentioned morphism by

$$\psi_{\mathfrak{g},m,n}^{c,k,+1} : \mathfrak{g}(m) \longrightarrow \mathfrak{g}(n)$$

The map sending $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}(m, n)_+$ to the unique $f_X(x) \in \mathbb{C}_{m-n}^{\text{hom}}[x]$ such that $\psi_{\mathfrak{g}}(X_m) = f_X(x)X_n$ is a bijection. Similarly for $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}(m, n)_-$, for $\psi_{\mathfrak{g}}$ to be nonzero we must have $n \leq m$ and there exist $c \in \mathbb{C}^\times, k \in \{0, 1, 2, \dots, m-n\}$, such that

$$\psi_{\mathfrak{g}}(X_m) = cx^k Y_n, \quad \psi_{\mathfrak{g}}(Y_m) = c^{-1}x^{m-n-k} X_n, \quad \psi_{\mathfrak{g}}(H_m) = -H_n.$$

We shall denote the above mentioned morphism by

$$\psi_{\mathfrak{g},m,n}^{c,k,-1} : \mathfrak{g}(m) \longrightarrow \mathfrak{g}(n)$$

The map sending $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}(m, n)_-$ to the unique $f_X(x) \in \mathbb{C}_{m-n}^{\text{hom}}[x]$ such that $\psi_{\mathfrak{g}}(X_m) = f_X(x)Y_n$ is a bijection. \square

Corollary 4.2. *Any extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ can be embedded in the constant family*

$$(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{A}_{\mathbb{C}}^1 = (\mathfrak{g}(0), SO(2, \mathbb{C})).$$

The following Lemma shall be needed later on.

Lemma 4.3. *For any $m, n, p \in \mathbb{N}_0$, such that $m \geq n \geq p$, any $k_1 \in \{0, 1, \dots, m-n\}$, $k_2 \in \{0, 1, \dots, n-p\}$, $c_1, c_2 \in \mathbb{C}^\times$, $s_1, s_2 \in \{-1, +1\}$,*

$$\psi_{\mathfrak{g},m,n}^{c_1,k_1,s_1} \circ \psi_{\mathfrak{g},n,p}^{c_2,k_2,s_2} = \psi_{\mathfrak{g},m,p}^{c_1 c_2, k_1+k_2, s_1 s_2}$$

The proof is by direct calculation.

4.2. Morphisms between localizations of extensions. In this section we study the space of morphisms between restrictions to \mathbb{C}^\times of a given pair of extensions. This shall be needed in the last section of the paper where we classify extensions over $\mathbb{P}_{\mathbb{C}}^1$.

Recall that $\iota^* \mathfrak{g}(n) = \mathfrak{g}(n)|_{\mathbb{C}^\times} = \mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}[x]} \mathfrak{g}(n)$. For $m, n \in \mathbb{N}_0$, we define

$$\text{Hom}^0(m, n) := \text{Hom}_{\text{pairs}}((\iota^* \mathfrak{g}(m), SO(2, \mathbb{C})), (\iota^* \mathfrak{g}(n), SO(2, \mathbb{C}))).$$

$$\text{Hom}^0(m, n)_{\pm} := \{(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}^0(m, n) | \psi_K(k) = k^{\pm 1}, \forall k \in SO(2, \mathbb{C})\}.$$

We let $\mathbb{C}((x))$ be the ring of formal Laurent series in one variable x over \mathbb{C} , and let $\mathbb{C}((x))^{\text{hom}}$ be its subset of homogeneous elements. Explicitly,

$$\mathbb{C}((x))^{\text{hom}} := \{cx^n | n \in \mathbb{Z}, c \in \mathbb{C}\}.$$

Theorem 4.4. *For any $m, n \in \mathbb{N}_0$, there are bijections such that*

$$\text{Hom}^0(m, n)_+ \simeq \mathbb{C}((x))^{\text{hom}} \simeq \text{Hom}^0(m, n)_-.$$

We omit the proof that is similar to the proof of Theorem 4.1. For a later use we describe the hom spaces explicitly. For every $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}^0(m, n)_+$ such that $\psi_{\mathfrak{g}} \neq 0$, we must have $\psi_{\mathfrak{g}}(1 \otimes H_m) = 1 \otimes H_n$, and there exist $c \in \mathbb{C}^\times, k \in \mathbb{Z}$ such that

$$\psi_{\mathfrak{g}}(1 \otimes X_m) = cx^k \otimes X_n, \quad \psi_{\mathfrak{g}}(1 \otimes Y_m) = c^{-1}x^{m-n-k} \otimes Y_n.$$

We shall denote the above mentioned morphism by

$$\psi_{\mathfrak{g}^*, m, n}^{c, k, +1} : \iota^* \mathfrak{g}(m) \longrightarrow \iota^* \mathfrak{g}(n).$$

For every $(\psi_{\mathfrak{g}}, \psi_K) \in \text{Hom}^0(m, n)_-$ such that $\psi_{\mathfrak{g}} \neq 0$, we must have $\psi_{\mathfrak{g}}(1 \otimes H_m) = -1 \otimes H_n$, and there exist $c \in \mathbb{C}^\times, k \in \mathbb{Z}$ such that

$$\psi_{\mathfrak{g}}(1 \otimes X_m) = cx^k \otimes Y_n, \quad \psi_{\mathfrak{g}}(1 \otimes Y_m) = c^{-1}x^{m-n-k} \otimes X_n.$$

We shall denote the above mentioned morphism by

$$\psi_{\mathfrak{g}^*, m, n}^{c, k, -1} : \iota^* \mathfrak{g}(m) \longrightarrow \iota^* \mathfrak{g}(n).$$

4.3. Canonical realization of $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ inside the constant family. In a previous section we saw that we can embed $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ in the constant family $(\mathfrak{g}(0), SO(2, \mathbb{C}))$. However, this embedding is not unique and moreover unless $n = 0$ its image inside the constant family is not unique. In this section we show that for even n there is a canonical realization of $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ inside the constant family. With an extra assumption we show that for any n (even or odd) there is a canonical realization of $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ inside the constant family.

Recall that every $\sum_{i=1}^m f_i(x) \otimes Z_i \in \mathfrak{g}(0) = \mathbb{C}[x] \otimes_{\mathbb{C}} \mathfrak{sl}_2(\mathbb{C})$ is canonically identified with the algebraic function from \mathbb{C} into $\mathfrak{sl}_2(\mathbb{C})$ that is defined via

$$t \longmapsto \sum_{i=1}^m f_i(t) Z_i.$$

For such functions we have a well defined derivative of any order via differentiating the coefficients that are polynomial functions. We denote the n -th derivative of a function $F \in \mathfrak{g}(0)$ by $F^{(n)}$. By convention $F^{(0)} = F$.

4.3.1. *The even case.*

Theorem 4.5. *For any $k \in \mathbb{N}$, the subset*

$$\mathfrak{s}(2k) := \{F \in \mathfrak{g}(0) \mid F^{(m)}(0) \in \mathfrak{so}(2, \mathbb{C}), \forall m \in \{0, 1, \dots, k-1\}\}$$

is sub Lie algebra over $\mathbb{C}[x]$ that together with $SO(2, \mathbb{C})$ is a sub pair of $(\mathfrak{g}(0), SO(2, \mathbb{C}))$ that is isomorphic to $(\mathfrak{g}(2k), SO(2, \mathbb{C}))$.

Proof. We let $\{X, Y, H\}$ be an ordered basis of $\mathfrak{sl}_2(\mathbb{C})$ such that $H \in \mathfrak{so}(2, \mathbb{C})$, $X \in (\mathfrak{sl}_2(\mathbb{C}))_2$, $Y \in (\mathfrak{sl}_2(\mathbb{C}))_{-2}$ and

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = H.$$

The set $\{X_0 := 1 \otimes X, Y_0 := 1 \otimes Y, H_0 := 1 \otimes H\}$ is an ordered basis of $\mathfrak{g}(0)$ with the same structure constants as $\{X, Y, H\}$. Note that the notation $\{X_0, Y_0, H_0\}$ is consistent with our fixed basis for $\mathfrak{g}(0)$ from Section 3.1. Clearly $B := \{1 \otimes H, x^k \otimes X, x^k \otimes Y\}$ is linearly independent set over $\mathbb{C}[x]$ that belongs to $\mathfrak{s}(2k)$. Any element F of $\mathfrak{g}(0)$

can be written uniquely in the form $f_X(x) \otimes X + f_Y(x) \otimes Y + f_H(x) \otimes H$, for some $f_X(x), f_Y(x), f_H(x) \in \mathbb{C}[x]$. For any $m \in \mathbb{N}_0$,

$$F^{(m)} = f_X^{(m)}(x) \otimes X + f_Y^{(m)}(x) \otimes Y + f_H^{(m)}(x) \otimes H.$$

Hence $F \in \mathfrak{s}(2k)$ if and only if

$$f_X^{(m)}(0)X + f_Y^{(m)}(0)Y + f_H^{(m)}(0)H \in \mathbb{C}H, \quad \forall m \in \{0, 1, \dots, k-1\}.$$

This shows that B is a basis for $\mathfrak{s}(2k)$. Since

$$\begin{aligned} [1 \otimes H, x^k \otimes X] &= 2x^k \otimes X \\ [1 \otimes H, x^k \otimes Y] &= -2x^k \otimes Y \\ [x^k \otimes X, x^k \otimes Y] &= x^{2k} \otimes H. \end{aligned}$$

it follows that $(\mathfrak{s}(2k), SO(2, \mathbb{C})) \simeq (\mathfrak{g}(2k), SO(2, \mathbb{C}))$. \square

4.3.2. The general case. For $n \in \mathbb{N}$, there are many different isomorphic copies of $\mathfrak{g}(n)$ in $\mathfrak{g}(0)$. For even n , as explained in the last theorem, there is a natural choice for such a copy, given by $\mathfrak{s}(n)$. Here we shall make a different choice that will work equally well for all $n \in \mathbb{N}$. This choice is natural for a given positive system of $\mathfrak{sl}_2(\mathbb{C})$.

As in the proof of the last Theorem, we let $\{X, Y, H\}$ be an ordered basis of $\mathfrak{g} := \mathfrak{sl}_2(\mathbb{C})$ such that $H \in \mathfrak{so}(2, \mathbb{C}) = \mathfrak{g}_0$, $X \in \mathfrak{g}_2$, $Y \in \mathfrak{g}_{-2}$ and

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = H.$$

Theorem 4.6. *For any $n \in \mathbb{N}$, the subset*

$$\mathfrak{l}(n) := \{F \in \mathfrak{g}(0) \mid F^{(m)}(0) \in \mathfrak{g}_0 \oplus \mathfrak{g}_2, \forall m \in \{0, 1, \dots, n-1\}\}$$

is sub Lie algebra over $\mathbb{C}[x]$ that together with $SO(2, \mathbb{C})$ is a sub pair of $(\mathfrak{g}(0), SO(2, \mathbb{C}))$ that is isomorphic to $(\mathfrak{g}(n), SO(2, \mathbb{C}))$.

Proof. As in the proof of the last Theorem, we can show that the set $\{1 \otimes X, x^n \otimes Y, 1 \otimes H\}$ is a basis over $\mathbb{C}[x]$ for $\mathfrak{l}(n)$ satisfying

$$\begin{aligned} [1 \otimes H, 1 \otimes X] &= 2(1 \otimes X) \\ [1 \otimes H, x^n \otimes Y] &= -2(x^n \otimes Y) \\ [1 \otimes X, x^n \otimes Y] &= x^n(1 \otimes H). \end{aligned}$$

This implies that $(\mathfrak{l}(n), SO(2, \mathbb{C})) \simeq (\mathfrak{g}(n), SO(2, \mathbb{C}))$. \square

4.4. The center of $\mathcal{U}(\mathfrak{g}(n))$. In this section we explicitly describe the center of the universal enveloping algebra of $\mathfrak{g}(n)$. This has applications to representation theory of $\mathfrak{g}(n)$ and of $(\mathfrak{g}(n), SO(2, \mathbb{C}))$.

As above we fix a basis $\{X, Y, H\}$ of $\mathfrak{g} := \mathfrak{sl}_2(\mathbb{C})$ such that $H \in \mathfrak{so}(2, \mathbb{C})$ and

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = H.$$

It is well known that $\mathcal{Z}(\mathfrak{g})$, the center of the universal enveloping algebra of \mathfrak{g} is equal to the polynomial algebra $\mathbb{C}[\Omega_{\mathfrak{g}}]$, where $\Omega_{\mathfrak{g}}$ is the canonical Casimir element which is given in terms of the mentioned basis by

$$\Omega_{\mathfrak{g}} = \frac{1}{8}H^2 + \frac{1}{4}(XY + YX).$$

In this section we shall realize the extensions $(\mathfrak{g}(n), SO(2, \mathbb{C}))$ concretely inside the constant family $(\mathfrak{g}(0), SO(2, \mathbb{C}))$ via the isomorphism $(\mathfrak{g}(n), SO(2, \mathbb{C})) \simeq (\mathfrak{h}(n), SO(2, \mathbb{C})) \subset (\mathfrak{g}(0), SO(2, \mathbb{C}))$. We shall not explicitly mention the isomorphism and simply work with $\mathfrak{h}(n)$ and the basis

$$B_n := \{H_n := 1 \otimes H, X_n := 1 \otimes X, Y_n := x^n \otimes Y\} \subset \mathfrak{g}(0)$$

of $\mathfrak{h}(n)$. We define

$$\Omega_n := x^n \otimes \Omega_{\mathfrak{g}} \in \mathbb{C}[x] \otimes_{\mathbb{C}} \mathcal{Z}(\mathfrak{g}) \simeq \mathcal{Z}(\mathfrak{h}(0)).$$

Theorem 4.7. *The center of the universal enveloping algebra of $\mathfrak{h}(n)$ is equal to the commutative polynomial algebra over $\mathbb{C}[x]$ generated by Ω_n inside $\mathbb{C}[x] \otimes_{\mathbb{C}} \mathcal{Z}(\mathfrak{g}) \simeq \mathcal{Z}(\mathfrak{g}(0))$. In particular $\mathcal{Z}(\mathfrak{h}(n)) = \mathbb{C}[x, \Omega_n]$.*

Proof. We first note that

$$\Omega_n = \frac{x^n}{8} H_n^2 + \frac{1}{4} (X_n Y_n + Y_n X_n) \in \mathcal{Z}(\mathfrak{h}(0)).$$

The proof is similar to the proof of Lemma 6.4 in [14]. \square

Remark 4.8. The family of commutative $\mathbb{C}[x]$ -algebras $\mathcal{Z}(\mathfrak{h}(n))$ has a canonical subspace (which is a free rank one $\mathbb{C}[x]$ -module) consisting of all elements of $\mathcal{Z}(\mathfrak{h}(n))$ that are of degree two in the sense of PBW-filtration and act on the trivial $\mathcal{U}(\mathfrak{h}(n))$ -module $\mathbb{C}[x]$ by zero. This subspace, known as the Casimir subfamily of $\mathcal{Z}(\mathfrak{h}(n))$, has Ω_n as its basis. The basis and hence Ω_n is unique up to a nonzero complex multiplicative constant.

5. EXTENSIONS OF $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ OVER $\mathbb{P}_{\mathbb{C}}^1$

In this section we classify extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{P}_{\mathbb{C}}^1$.

5.1. Extensions over $\mathbb{P}_{\mathbb{C}}^1$. Following tradition we realize $\mathbb{P}_{\mathbb{C}}^1$ via homogeneous coordinates as $\{[z_1; z_2] \in \mathbb{C}^2\} / \sim$ where $[z_1; z_2] \sim [cz_1; cz_2]$ for any $c \neq 0, z_1, z_2 \in \mathbb{C}$. We identify \mathbb{C}^\times with $\{[z_1; z_2] \in \mathbb{P}_{\mathbb{C}}^1 | z_1 z_2 \neq 0\}$.

Definition 5.1. An algebraic family of Harish-Chandra pairs $(\mathfrak{g}, \mathbf{K})$ over $\mathbb{P}_{\mathbb{C}}^1$ is called an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$, if \mathbf{K} is a constant algebraic family of complex algebraic groups and the restriction of $(\mathfrak{g}, \mathbf{K})$ to \mathbb{C}^\times is isomorphic to $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$.

For $i \in \{1, 2\}$, we define $U_i := \{[z_1; z_2] \in \mathbb{P}_{\mathbb{C}}^1 | z_i \neq 0\}$ and $\varphi_i : U_i \rightarrow \mathbb{C}$, via $\varphi_1([z_1; z_2]) = z_2 z_1^{-1}, \varphi_2([z_1; z_2]) = z_1 z_2^{-1}$. Then $U_1 \cup U_2$ is an open affine cover of $\mathbb{P}_{\mathbb{C}}^1$ and φ_i are algebraic isomorphisms. Moreover $\varphi_2^{-1} \circ \varphi_1|_{U_1 \cap U_2} : U_1 \cap U_2 \rightarrow U_1 \cap U_2$ is an isomorphism of affine algebraic varieties. It is well known that a sheaf on $\mathbb{P}_{\mathbb{C}}^1$ is the same thing as a pair of sheaves, one on U_1 and the other on U_2 , that agree on $U_1 \cap U_2$. Using the isomorphisms φ_i , a sheaf on $\mathbb{P}_{\mathbb{C}}^1$ is an ordered pair of sheaves $(\mathcal{F}_1, \mathcal{F}_2)$ both on \mathbb{C} , together with an isomorphism $\psi : \mathcal{F}_1|_{\mathbb{C}^\times} \rightarrow \mathcal{F}_2|_{\mathbb{C}^\times}$ between their restrictions to \mathbb{C}^\times . We shall denote the corresponding sheaf on $\mathbb{P}_{\mathbb{C}}^1$ by $(\mathcal{F}_1, \mathcal{F}_2, \psi)$ and its isomorphism class by $[(\mathcal{F}_1, \mathcal{F}_2, \psi)]$.

5.2. Classification of extensions over $\mathbb{P}_{\mathbb{C}}^1$.

Theorem 5.2. *The collection of all the equivalence classes*

$$\left[\left((\mathfrak{g}(m), K), (\mathfrak{g}(n), K), (\psi_{\mathfrak{g}^*, m, n}^{1, k, 1}, \psi_K^1) \right) \right],$$

as m and n vary in \mathbb{N}_0 and k vary in \mathbb{Z} , is an exhaustive list of equivalence classes of extensions of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{P}_{\mathbb{C}}^1$. Moreover, the representatives in the list are pairwise non-isomorphic.

Proof. From the above discussion together with Theorem 3.2 and Theorem 4.4 any extension over $\mathbb{P}_{\mathbb{C}}^1$ is isomorphic to some

$$\left((\mathfrak{g}(m), K), (\mathfrak{g}(n), K), (\psi_{\mathfrak{g}^*, m, n}^{c, k, \pm 1}, \psi_K^\pm) \right).$$

Note that the automorphism

$$(\psi_{\mathfrak{g}, m, m}^{c^{-1}, 0, \pm 1}, \psi_K^{\pm 1}) : (\mathfrak{g}(m), K) \longrightarrow (\mathfrak{g}(m), K)$$

satisfies $\psi_{\mathfrak{g}^*, m, m}^{c^{-1}, 0, \pm 1} = 1 \otimes \psi_{\mathfrak{g}, m, m}^{c^{-1}, 0, \pm 1}$ where

$$(1 \otimes \psi_{\mathfrak{g}, m, m}^{c^{-1}, 0, \pm 1}, \psi_K^{\pm 1}) : (\iota^* \mathfrak{g}(m), K) \longrightarrow (\iota^* \mathfrak{g}(m), K)$$

is the induced map between the corresponding localizations. Since

$$(\psi_{\mathfrak{g}^*, m, n}^{c, k, \pm 1}, \psi_K^\pm) \circ (\psi_{\mathfrak{g}^*, m, m}^{c^{-1}, 0, \pm 1}, \psi_K^{\pm 1}) = (\psi_{\mathfrak{g}^*, m, n}^{1, k, 1}, \psi_K^1),$$

we may assume that $c = 1$ and ψ_K is the identity map. This proves exhaustiveness. We now show that the representatives in the above list are pairwise non-isomorphic. Assume that for $i \in \{1, 2\}$,

$$\left((\mathfrak{g}(m_i), K), (\mathfrak{g}(n_i), K), (\psi_{\mathfrak{g}^*, m_i, n_i}^{1, k_i, 1}, \psi_K^1) \right)$$

is an extension of $(\mathfrak{sl}_2(\mathbb{C}), SO(2, \mathbb{C}))/\mathbb{C}^\times$ over $\mathbb{P}_{\mathbb{C}}^1$. Assume in addition that the two extensions are isomorphic. This means that $(\mathfrak{g}(m_1), K)$ is isomorphic to $(\mathfrak{g}(m_2), K)$ and $(\mathfrak{g}(n_1), K)$ is isomorphic to $(\mathfrak{g}(n_2), K)$ and the two isomorphisms are compatible with $(\psi_{\mathfrak{g}^*, m_1, n_1}^{1, k_1, 1}, \psi_K^1)$ and $(\psi_{\mathfrak{g}^*, m_2, n_2}^{1, k_2, 1}, \psi_K^1)$. Hence we must have $m_1 = m_2$ and $n_1 = n_2$. We denote the corresponding isomorphisms by $(\psi_{\mathfrak{g}, m_1, m_1}^{c_m, 0, s_m}, \psi_K^{s_m})$ and $(\psi_{\mathfrak{g}, n_1, n_1}^{c_n, 0, s_n}, \psi_K^{s_n})$. Compatibility means that the following diagram, is commutative

$$\begin{array}{ccc} (\iota^* \mathfrak{g}(m_1), K) & \xrightarrow{(\psi_{\mathfrak{g}^*, m_1, n_1}^{1, k_1, 1}, \psi_K^1)} & \iota^* \mathfrak{g}(n_1), K \\ \downarrow (1 \otimes \psi_{\mathfrak{g}, m_1, m_1}^{c_m, 0, s_m}, \psi_K^{s_m}) & & \downarrow (1 \otimes \psi_{\mathfrak{g}, n_1, n_1}^{c_n, 0, s_n}, \psi_K^{s_n}) \\ (\iota^* \mathfrak{g}(m_1), K) & \xrightarrow{(\psi_{\mathfrak{g}^*, m_1, n_1}^{1, k_2, 1}, \psi_K^1)} & (\iota^* \mathfrak{g}(n_1), K) \end{array}$$

Noting that $\psi_{\mathfrak{g}^*, m_1, m_1}^{c_m, 0, s_m} = 1 \otimes \psi_{\mathfrak{g}, m_1, m_1}^{c_m, 0, s_m}$ and $\psi_{\mathfrak{g}^*, n_1, n_1}^{c_n, 0, s_n} = 1 \otimes \psi_{\mathfrak{g}, n_1, n_1}^{c_n, 0, s_n}$ and using Lemma 4.3, commutativity of the diagram is equivalent to

$$(\psi_{\mathfrak{g}^*, m_1, n_1}^{c_n, k_1, s_n}, \psi_K^{s_n}) = (\psi_{\mathfrak{g}^*, m_1, n_1}^{c_m, k_2, s_m}, \psi_K^{s_m}).$$

Hence we must have $c_n = c_m$, $k_1 = k_2$ and $s_n = s_m$. Hence the two extensions are equal. \square

By Grothendieck's theorem any algebraic vector bundle on $\mathbb{P}_{\mathbb{C}}^1$ is a direct sum of line bundles. From the last Theorem we can see what is the corresponding decomposition into line bundles of any extension over $\mathbb{P}_{\mathbb{C}}^1$. Here is a precise statement.

Corollary 5.3. *As algebraic K -equivariant vector bundles*

$$\left((\mathfrak{g}(m), SO(2, \mathbb{C})), (\mathfrak{g}(n), SO(2, \mathbb{C})), (\psi_{\mathfrak{g}^*, m, n}^{1, k, 1}, \psi_K^1) \right)$$

is isomorphic to

$$\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(0) \oplus \mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(-k) \oplus \mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(k + n - m).$$

Proof. Since $\psi_{\mathfrak{g}^*, m, n}^{1, k, 1}(H_m) = H_n$, the subsheaf generated by H_m , and H_n is isomorphic to $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(0)$. It is the K isotypic piece corresponding to the trivial representation of K . Since $\psi_{\mathfrak{g}^*, m, n}^{1, k, 1}(X_m) = x^k X_n$, the subsheaf generated by X_m , and X_n is isomorphic to $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(-k)$. It is the K isotypic piece corresponding to the one dimensional representation of K in which $k(z)$ acts via multiplication by z^2 . Similarly, the subsheaf generated by Y_m , and Y_n is isomorphic to $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(k + n - m)$. It is the K isotypic piece corresponding to the one dimensional representation of K in which $k(z)$ acts via multiplication by z^{-2} . \square

6. DECLARATIONS

Ethical Approval. Not applicable.

Funding. This research was supported by the ISRAEL SCIENCE FOUNDATION (grant No. 1040/22).

Availability of data and materials. Not applicable.

REFERENCES

- [1] Jeffrey D. Adams, Marc A. A. van Leeuwen, Peter E. Trapa, and David A. Vogan, Jr. Unitary representations of real reductive groups. *Astérisque*, (417):x + 174, 2020.
- [2] Alexandre Afgoustidis. On the analogy between real reductive groups and Cartan motion groups: a proof of the Connes-Kasparov isomorphism. *J. Funct. Anal.*, 277(7):2237–2258, 2019.
- [3] Alexandre Afgoustidis. On the analogy between real reductive groups and Cartan motion groups: the Mackey-Higson bijection. *Camb. J. Math.*, 9(3):551–575, 2021.
- [4] Dan Barbasch, Nigel Higson, and Eyal Subag. Algebraic families of groups and commuting involutions. *Internat. J. Math.*, 29(4):1850030, 18, 2018.
- [5] Paul Baum, Alain Connes, and Nigel Higson. Classifying space for proper actions and K -theory of group C^* -algebras. In *C^* -algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 240–291. Amer. Math. Soc., Providence, RI, 1994.
- [6] Joseph Bernstein, Nigel Higson, and Eyal Subag. Algebraic families of Harish-Chandra pairs. *Int. Math. Res. Not. IMRN*, (15):4776–4808, 2020.
- [7] Joseph Bernstein, Nigel Higson, and Eyal Subag. Contractions of representations and algebraic families of Harish-Chandra modules. *Int. Math. Res. Not. IMRN*, (11):3494–3520, 2020.
- [8] Nigel Higson. The Baum-Connes conjecture. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, pages 637–646, 1998.
- [9] Nigel Higson. The Mackey analogy and K -theory. In *Group representations, ergodic theory, and mathematical physics: a tribute to George W. Mackey*, volume 449 of *Contemp. Math.*, pages 149–172. Amer. Math. Soc., Providence, RI, 2008.

- [10] Nigel Higson and Eyal Subag. Families of symmetries and the hydrogen atom. *Adv. Math.*, 408:Paper No. 108586, 61, 2022.
- [11] E. Inonu and E. P. Wigner. On the contraction of groups and their representations. *Proc. Nat. Acad. Sci. U.S.A.*, 39:510–524, 1953.
- [12] V. Lafforgue. Banach KK -theory and the Baum-Connes conjecture. In *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*, pages 795–812. Higher Ed. Press, Beijing, 2002.
- [13] E. M. Subag, E. M. Baruch, J. L. Birman, and A. Mann. Strong contraction of the representations of the three-dimensional Lie algebras. *J. Phys. A*, 45(26):265206, 25, 2012.
- [14] Eyal Subag. The algebraic Mackey-Higson bijections. *J. Lie Theory*, 29(2):473–492, 2019.
- [15] Eyal M. Subag. Symmetries of the hydrogen atom and algebraic families. *J. Math. Phys.*, 59(7):071702, 20, 2018.
- [16] Vincent van der Noort. *Analytic parameter dependence of Harish-Chandra modules for real reductive Lie groups—a family affair*. Thesis, Utrecht University, 2009.
- [17] Nolan R. Wallach. The dependence on parameters of the inverse functor to the K -finite functor. *Represent. Theory*, 26:94–121, 2022.