

APPLYING NUMEROSITY TO SURREAL INTEGRATION

ILYA SHNITMAN

ABSTRACT

We introduce the concept of *numerosity*, a measure of a set's size, more precise than cardinality and satisfying the Euclid's principle "the whole is greater than a part" as a surreal number and provide an explicit formula for expressing numerosities of some sequences that have no accumulation points as surreal expressions, with examples. Then we introduce a surreal-valued function that shares many properties with the Dirac Delta distribution and employ it to derive some integration properties of surreal values. Finally, we provide a formula for integrating a surreal-valued function over surreal domain, employing numerosity in a way, similar to Lebesgue measure and derive some novel formulas, connecting surreal numbers via integration. The suggested here method of surreal integration is different from methods suggested by other authors in not satisfying the property of linearity regarding infinite factors. In our opinion, the struggle to keep this rule intact is the reason for many failed attempts to define the surreal integration. For the suggested formulas, code in Wolfram Mathematica language is provided.

1. Mapping numerosities to surreal numbers.

Several authors[1][2][3] have introduced the concept of numerosity as a measure of set's size, striving to satisfy the Euclid's principle "the whole is greater than a part", which can be formally represented by the following equation: $N(A \cup B) = N(A) + N(B) - N(A \cap B)$, where N is the numerosity. The approach described in this paper makes an attempt to do so in the most simple way, yet allowing to explicitly derive values of numerosities in closed form.

Comb-form of the numerosity. We formally represent the numerosity of a nowhere dense set $S \subset \mathbb{R}$ with elements s_n as a (generally, divergent) integral

$N(S) = \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \delta(x - s_n) dx$, where $\delta(x)$ is the Dirac Delta distribution. Indeed, naturally

partial sums of this integral increase by one each time they encounter an element of S . We will call integrals of this form the "comb-form" of the numerosity. We assume that numerosity is invariant regarding changing of the sign of the elements of a set, so

$$\int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \delta(x - s_n) dx = \int_{-1}^{\infty} \sum_{n=0}^{\infty} \delta(x - |s_n|) dx$$

Smooth-form of the numerosity. Let $\phi_n(x)$ to be a bounded real function with finite support $[a_n, b_n]$, area under the curve $\int_{a_n}^{b_n} \phi_n(x) dx = 1$ and center of mass at s_n : $\int_{a_n}^{b_n} x \phi_n(x) dx = s_n$. We postulate that these two integrals are equal: $\int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \delta(x - s_n) dx = \int_0^{\infty} \sum_{n=0}^{\infty} (\phi_n(x) + \phi_n(-x)) dx$. The later one is one of the smooth-forms of $N(S)$.

The surreal numbers can be considered an H-field with natural embedding of the germs of the functions (satisfying the Hard field requirements) at infinity into surreals, such that the germ of identity function $[x]$ is mapped to ω [4]. In such setting, if $w \in \mathbb{J}$, a purely-infinite surreal number, corresponding to the germ at infinity of the function $f(x)$, we postulate equivalence of the following divergent integral to w :

$$f(a) + \int_a^{\infty} f'(t) dt = w \tag{1}$$

where a is an arbitrary real number.

In this way, smooth forms of numerosities can be mapped to surreal numbers.

EXAMPLE. Find a subset of \mathbb{R} , whose numerosity is $\ln \omega$. The number $\ln \omega$ is purely-infinite, because its Conway normal form $\omega^{1/\omega}$ has no finite and infinitesimal terms. Under canonical embedding and the above-mentioned equivalence,

$$\ln \omega = [\ln x] = \ln 1 + \int_1^{\infty} (\ln t)' dt = \int_1^{\infty} \frac{1}{t} dt.$$

Now, we take the last integral and separate it into pieces of area equal to 1, in our case:

$$\int_1^{\infty} \frac{1}{t} dt = \int_1^e \frac{1}{t} dt + \int_e^{e^2} \frac{1}{t} dt + \dots = \sum_{k=0}^{\infty} \int_{e^k}^{e^{k+1}} \frac{1}{t} dt.$$

The centers of mass of each integral are located at $s_k = e^{k+1} - e^k$, so the set $\{e^{k+1} - e^k\} : k \in \mathbb{N}_0$ has the numerosity $\ln \omega$.

EXPLICIT FORMULA. The above-mentioned definitions can be expressed in compact form as follows: $N(\{a_k\}) = (D\Delta^{-1}a_k)^{[-1]}(\omega)$, where $f^{[-1]}$ is the inverse function operator. The formula works not only for purely-infinite numbers but for those containing finite and infinitesimal parts, so we suggest calling the result of applying this formula the full numerosity and after trimming the infinitesimal part, the refined numerosity $\overline{N}(\{a_k\})$. Differences in refined numerosities between different sequences, calculated according to the method suggested in this article coincide with those obtained with the method suggested by James Propp in[5].

MATHEMATICA CODE. Following is the Mathematica code for finding the surreal expression of the numerosity of a non-negative, growing sequence a_k with no accumulation points:

```
a[k] := k^2;
SolveValues[D[Sum[a[k], k], k] == \[Omega], k] /. C[1] -> 0 //
FullSimplify // Expand
```

The opposite process, finding a sequence with given numerosity as a surreal expression S can be done with the following code:

```
S = Log[\[Omega]];
DifferenceDelta[Integrate[Normal[SolveValues[S == k, \[Omega]]], k], k] /. C[1]
-> 0 // FullSimplify // Expand
```

The obtained list of numerosities of subsets of reals can be seen in Appendix 1.

2. Surreal Delta function

Suppose we want to obtain a functional to count the number of roots of a function on an interval. For a differentiable function $f(x)$, the following expression does the thing, where applicable:

$\int_a^b \delta(f(x))|f'(x)|dx$. From now on we will call the operator $\tilde{\delta}$, defined as $\tilde{\delta}(f(x)) = \delta(f(x))|f'(x)|$, where x is the integration variable, “surreal delta function”. We will further generalize it.

First of all, we should notice that $\tilde{\delta}(x)$ behaves just like $\delta(x)$ in all contexts where its argument is the integration variable itself, and not some non-trivial function of it. Particularly, its Laplace and Fourier transforms are the same as of conventional Delta distribution.

Second thing to notice is that it satisfies the following property: $\tilde{\delta}(ax) = \tilde{\delta}(x)$, which coincides with the similar property of the unit impulse function, unlike the property of the conventional Delta distribution $\delta(ax) = \frac{1}{|a|}\delta(x)$.

Let us consider the Fourier transform of the constant function $f(x) \equiv 1$:

$$F(s) = \mathcal{F}[1](s) = \int_{-\infty}^{\infty} e^{-st} dt$$

Formally, the right-hand expression is equal to 0 at any $s \neq 0$ (by Cesaro regularization). But at $s = 0$ we get the divergent integral $\int_{-\infty}^{\infty} 1dt$. This divergent integral corresponds to the surreal number 2ω .

Consequently, we can formally write:

$$F(x) = \mathcal{F}[1](x) = \begin{cases} 2\omega, & \text{if } x = 0 \\ 0, & \text{if } x \neq 0 \end{cases}$$

Applying inverse Fourier transform, $\mathcal{F}^{-1}[F](t) = 1 = 2\pi\mathcal{F}^{-1}[\delta](t) = 2\pi\mathcal{F}^{-1}[\tilde{\delta}](t)$. This allows us to generalize the Surreal Delta function this way:

$$\tilde{\delta}(x) = \begin{cases} \frac{\omega}{\pi}, & \text{if } x = 0 \\ 0, & \text{if } x \neq 0 \end{cases} \quad (2)$$

We can see that the property $\tilde{\delta}(ax) = \tilde{\delta}(x)$ holds with this piece-wise definition. This function $\tilde{\delta}(x)$ is a full-fledged surreal-valued function, and can be evaluated at any point. Particularly, since

$\int_{-1}^1 \tilde{\delta}(x) dx = 1$, we can consider it (in a sense) a derivative of $\text{sign}(x)$ function. This shows us that an infinite value of the derivative at a point indicates a jump the function makes there.

Expanding ω^p as a divergent integral, Surreal Delta function can be raised to any power:

$$\tilde{\delta}(x)^p = \begin{cases} \frac{p}{\pi^p} \int_0^\infty x^{p-1} dx, & \text{if } x = 0 \\ 0, & \text{if } x \neq 0. \end{cases} \quad (3)$$

It should be noted that the Surreal Delta function is not differentiable in terms of surreal-valued functions.

3. Integration of a surreal value over a subset of reals

Generalizing relation (3) to any analytic function f of ω , we have:

$$\int_{-\infty}^{\infty} \left(\begin{cases} f(\tilde{\delta}(0)) & x = 0 \\ 0 & x \neq 0 \end{cases} \right) dx = f'(a) + \frac{1}{\pi} \int_a^\infty f''(x) dx, \quad (4)$$

for arbitrary real a where the expression converges. This is equal to $f'(\tilde{\delta}(0))$, if $f'(\tilde{\delta}(0)) \in \mathbb{J}$ according to the definition (1).

Noticing that our H-field is equipped with operation of derivation $\partial()$, we can re-write it like this:

$$\int_{-\infty}^{\infty} \left(\begin{cases} w & x = 0 \\ 0 & x \neq 0 \end{cases} \right) dx = \pi \partial(w), \quad (5)$$

where $w \in \mathbb{J}$. But if we are integrating over a set where the expression under integral can take surreal values at several points, it is natural to multiply this result by the numerosity of the support:

$$\int_S w dt = \pi N(S) \partial(w) \quad (6)$$

One can easily notice that surreal integration defined this way does not generally satisfy the linearity against an infinite factor α :

$$\int_S (\alpha f) \neq \alpha \int_S f$$

For instance, if $u(x)$ is a unit impulse function $u(x) = \begin{cases} 1 & x = 0 \\ 0 & x \neq 0 \end{cases}$, then

$$\int_{-1}^1 \omega u(x) dx = \pi \neq \omega \int_{-1}^1 u(x) dx = 0$$

In our opinion, the struggle to keep this rule intact is the reason for many failed attempts to define the surreal integration. Still, there are ongoing efforts by other authors to keep this rule [6].

As a consequence,

$$\frac{1}{\pi} \int_S \omega dt = \int_S \tilde{\delta}(0) dt = N(S). \quad (7)$$

4. Numerosity of the continuum

A NOTE ON NOTATION. Since the surreal integration may depend on whether a single point is included in the integration interval, from now on, in the context of real limits $a, b \in \mathbb{R}$, we use notation \int_a^b to denote an integral where the limits of integration, a and b are half-included, in other words, $\int_a^b = \frac{1}{2} \left(\int_{(a,b)} + \int_{[a,b]} \right)$.

FIRST UNCOUNTABLE ORDINAL. The automorphisms of the surreal field give us some freedom in choosing the first uncountable ordinal ω_1 . We have freedom to define it, using formula (7), as the numerosity of the interval $[0, 1)$:

$$\omega_1 = \int_{[0,1)} \tilde{\delta}(0) dt = \frac{1}{\pi} \int_{[0,1)} \omega dt = \frac{1}{\pi} \int_0^1 \omega dt \quad (8)$$

Immediately, one can see from formula (6) that ω_1 can be represented also this way:

$$\omega_1 = \frac{1}{\pi} \int_0^\infty \ln \omega dt \quad (9)$$

It is important that only one of the limits of the integration to be included (or the both half-included) because the finite part of a numerosity corresponds to the Euler's characteristic of a set, and we want the finite part of ω_1 to be zero.

5. Integrating surreal function over surreal domain

EXPLICIT FORMULA. We assume that the Newton-Leibnitz formula is valid for integration over surreals. As such, symbolically obtained antiderivative of an analytic function allows to calculate definite integrals on surreals. Based on this assumption and on formula (6), we propose the following explicit formula. Let $F(x) \in \mathbb{J} + \mathbb{R}$ to be a surreal-valued function with only purely-infinite and finite parts, defined on surreal domain and $a, b \in \text{No}$. Then,

$$\int_a^b F(t)dt = \pi\omega_1 \int_{a(\text{formal})}^b \partial(F(t))dt + \int_a^b \text{fin } F(t)dt, \quad (10)$$

where the first integral should be taken formally, the way as if all the surreal numbers under the integral were real constants, and with help of the Newton-Leibniz formula. In the second integral, the notation `fin` means taking the finite part. For the second integral also, the Newton-Leibniz formula can be used.

In our proposed method the infinitesimal part of the expression under the integral does not affect the result of integration and can be ignored.

MATHEMATICA CODE. Here is the Mathematica code for integration of an arbitrary surreal function over surreal domain using the formula (10). For simplicity, in the input line, `w` and `W` stand for ω and for ω_1 respectively (if you want to use an arbitrary surreal in input, you should specify it as `A[w]` or `B[w]` to distinguish it from reals).

```
f[x_]=Exp[x*w];
a=0;
b=a;
f[x_]=f[x]/. W->W[w];
Unprotect[Power];0^0=1;Protect[Power];
int=Pi*w*Integrate[D[f[t],w],{t,a,b}]+Integrate[Fin[f[t]],{t,a,b}];
int=FullSimplify[Normal[If[a==b,Pi*D[f[a],w],int],int]/. W[w]:>W/. Derivative[1]
[W][w]->d[W]/. FullForm[Derivative[d_][W]]:>(d^d)[W]]];
Print[Inactivate[Integrate[f[x]/. W[w]->w/. Derivative[1][W][w]->d[W]/.
FullForm[Derivative[d_][W][w]]:>(d^d)[W]/. w->w/. W->Subscript[w,1],{x,a/. w-
>w/. W->Subscript[w,1],b/. w->w/. W->Subscript[w,1}],Integrate],"=",int/. w-
>w/. W->Subscript[w,1]]
```

It leaves the integration of the finite part unevaluated because finding the finite part of a surreal number `Fin[f[t]]` is a non-trivial problem (although straightforward if the Conway normal form of the expression is known) but for some numbers in $N_0(\omega_1)$ one can try the regularization via Laplace transform:

```
Fin[e_] := FullSimplify[(PowerExpand[e] /. Log[w] -> 0 /. 1/w -> 0 /.
1/w^(p_) :> 0 /. w -> 0) +
Limit[Evaluate[LaplaceTransform[D[e, w], w, x]] +
Evaluate[LaplaceTransform[D[e, w], w, -x]], x -> 0]/2 /.
Infinity -> 0 /. -Infinity -> 0 /. FullForm[Derivative[d_][W][p_]] :>
0]
```

Alternatively, one can utilize the built-in Mathematica's regularization machinery:

```
Fin[e_] := (PowerExpand[f[t]] /. W[w] -> 0 /. Log[w] -> 0 /. 1/w -> 0 /.
1/w^(p_) :> 0 /. w -> 0 /. 1/W[w] -> 0) +
Limit[FullSimplify[s*Sum[D[e, w] /. w -> s*w, {w, 1, Infinity}],
Regularization -> "Dirichlet"], s -> 0]
```

The list of some surreal indefinite integrals is given in Appendix 2.

APPENDIX 1. Numerosities of real sequences. The following is a list of calculated full numerosities (that is, with infinitesimal parts included) N and refined numerosities (that is without infinitesimal parts), \bar{N} of various subsets of reals ($k \in \mathbb{N}_0$):

- $N(\{\mathbb{N}\}) = N(\{k + 1\}) = \omega - 1/2$

- $N(\mathbb{Z}) = 2\omega$
- $N(\{2k\}) = \frac{\omega}{2} + \frac{1}{2}$
- $N(\{2k + 1\}) = \frac{\omega}{2}$
- $N(\{ak + b\}) = \frac{\omega}{a} + \frac{1}{2} - \frac{b}{a}$
- $N(\{k^2\}) = \frac{1}{6}\sqrt{36\omega + 3} + \frac{1}{2}; \quad \bar{N} = \sqrt{\omega} + \frac{1}{2}$
- $N(\{1/3 + k + k^2\}) = \sqrt{\omega}$
- $N(\{k^4\}) = \frac{1}{30}\sqrt{30\sqrt{900\omega + 30} + 225} + \frac{1}{2}; \quad \bar{N} = \sqrt[4]{\omega} + \frac{1}{2}$
- $N(\{a^k\}) = \frac{\ln\left(\frac{(a-1)\omega}{\ln a}\right)}{\ln a}$

APPENDIX 2. Antiderivatives of some surreal functions ($c \in \mathbb{R}, c > 0$ and $\alpha \in \text{No}, \alpha \neq 0$).

- $\int_0^\alpha \omega^c dt = \pi\alpha c\omega_1\omega^{c-1}$
- $\int_0^\alpha \omega^x dx = \frac{\pi((\alpha \ln \omega - 1)\omega^\alpha + 1)\omega_1}{\omega(\ln \omega)^2}$
- $\int_0^\alpha x^\omega dx = \frac{\pi\alpha^{\omega+1}((\omega + 1) \ln \alpha - 1)\omega_1}{(\omega + 1)^2}$
- $\int_0^\alpha e^\omega dx = e^\omega \pi\alpha\omega_1$
- $\int_0^\alpha c^\omega dx = c^\omega \pi\alpha\omega_1 \ln c$
- $\int_0^\alpha e^{c\alpha\omega} dx = \frac{\pi(e^{c\alpha\omega}(c\alpha\omega - 1) + 1)\omega_1}{c\omega^2}$
- $\int_0^\alpha \ln \omega dx = \frac{\pi\alpha\omega_1}{\omega}$
- $\int_0^\alpha \omega^\omega dx = \alpha\omega^\omega\omega_1\pi(\ln \omega + 1)$
- $\int_0^\alpha \omega^{\omega_1} dx = \alpha\pi\omega_1\omega^{\omega_1-1}(\omega\partial(\omega_1)\ln \omega + \omega_1)$
- $\int_0^\alpha \omega_1^\omega dx = \alpha\pi\omega_1^\omega(\omega\partial(\omega_1) + \ln \omega_1\omega_1)$

APPENDIX 3. Numerosities of shapes in 3D space. The numerosities of closed shapes in Euclidean 3-dimensional space can be obtained using the following formula: $V\omega_1^3 + 1/2S\omega_1^2 + \theta L\omega_1 + \chi$,

where V is volume, S is surface area, L is the length of the edges, θ is the external angle of the edges, and χ is the Euler's characteristic.

- Closed square with side a : $(a\omega_1 + 1)^2 = a^2\omega_1^2 + 2a\omega_1 + 1$
- Closed cube: $(a\omega_1 + 1)^3 = a^3\omega_1^3 + 3a^2\omega_1^2 + 3a\omega_1 + 1$
- Circle: $2\pi\omega_1 R$
- Closed disc: $\pi R^2\omega_1^2 + \pi R\omega_1 + 1$
- Empty sphere: $4\pi\omega_1^2 R^2 + 2$
- Closed ball: $4/3\pi\omega_1^3 R^3 + 2\pi\omega_1^2 R^2 + 1$
- Infinite solid cylinder: $2\pi\omega R^2\omega_1^3 + 2\pi R\omega\omega_1 + 2\omega\omega_1$
- Solid pseudosphere: $2/3\pi R^3\omega_1^3 + 2\pi R^2\omega_1^2 + \pi R\omega_1 + 2\omega\omega_1$
- Two Gabriel's horns glued together along the plane $x = 0$:
 $2\pi\omega_1^3 + 2\pi\omega_1^2 \ln \omega + \pi\omega_1 + 2\omega\omega_1$

References

- 1: Vieri Benci, Mauro Di Nasso, Numerosities of labelled sets: a new way of counting, 2003, [https://doi.org/10.1016/S0001-8708\(02\)00012-9](https://doi.org/10.1016/S0001-8708(02)00012-9)
- 2: Andreas Blass, Mauro Di Nasso, Marco Forti, Quasi-selective ultrafilters and asymptotic numerosities, 2012, <https://arxiv.org/abs/1011.2089>
- 3: Kenny Easwaran, Alan Hájek, Paolo Mancosu, Graham Oppy, Theories of Numerosities, 2023, <https://plato.stanford.edu/entries/infinity/numerosities.html>
- 4: Alessandro Berarducci, Vincenzo Mantova, Surreal numbers, derivations and transseries, 2015, <https://arxiv.org/abs/1503.00315>
- 5: James Propp, A new game with infinity, 2018, <https://mathenchant.wordpress.com/2018/09/16/a-new-game-with-infinity/>
- 6: Ovidiu Costin, Philip Ehrlich, Integration on the Surreals, 2022, <https://doi.org/10.48550/arXiv.2208.14331>