Characterizing extremal dependence on a hyperplane

Phyllis Wan*

Abstract

Quantifying the risks of extreme scenarios requires understanding the tail behaviours of variables of interest. While the tails of individual variables can be characterized parametrically, the extremal dependence across variables can be complex and its modeling remains one of the core problems in extreme value analysis. Notably, existing measures for extremal dependence, such as angular components and spectral random vectors, reside on nonlinear supports, such that statistical models and methods designed for linear vector spaces cannot be readily applied. In this paper, we show that the extremal dependence of d asymptotically dependent variables can be characterized by a class of random vectors residing on a (d-1)-dimensional hyperplane. This translates the analyses of multivariate extremes to that on a linear vector space, opening up the potentials for the application of existing statistical techniques, particularly in statistical learning and dimension reduction. As an example, we show that a lower-dimensional approximation of multivariate extremes can be achieved through principal component analysis on the hyperplane. Additionally, through this framework, the widely used Hüsler-Reiss family for modelling extremes is characterized by the Gaussian family residing on the hyperplane, thereby justifying its status as the Gaussian counterpart for extremes.

Keywords and phrases: multivariate extreme value statistics; extremal dependence structure; dimension reduction AMS 2010 Classification: 62G32 (62H05; 60G70).

1 Introduction

Extreme events, despite their rare occurrences, entail high risks for the society. Quantifying the risks of extreme scenarios plays an important role in preventing and mitigating catastrophic outcomes. The aim of extreme value analysis is to provide mathematically justified tools to model observed rare events and estimate the risks for those not in the observed range.

A general framework for modeling extremes is the *peak-over-threshold* framework, in which one considers the distribution of observations over a high threshold. In the univariate case, this framework is well-studied and widely used. The sample observations exceeding a high threshold converge to the class of generalized Pareto distributions, parametrized by a scale parameter and a shape parameter. This allows for straightforward statistical inference using likelihood techniques. For an overview, see e.g., Coles (2001).

^{*}Erasmus University Rotterdam; Econometric Institute, Burg. Oudlaan 50, 3062 PA Rotterdam, the Netherlands; email: wan@ese.eur.nl

The multivariate case, on the other hand, requires simultaneous considerations of the marginal tails and the extremal dependence. The former can be approached by applying univariate techniques, while the latter can be separated from the former by standardizing the marginals of the data. Even so, modeling extremal dependence remains a core problem in extreme value analysis as its structure may be complex and cannot be summarized by a finite-dimensional model.

There are two common approaches in the literature to geometrically characterize the tail dependence of a random vector $\mathbf{Y} = (Y_1, \ldots, Y_d)$.

• Angular component Θ : Let F_1, \ldots, F_d denote the marginal cdf of Y_1, \ldots, Y_d . Consider the marginal transformation

$$\tilde{\mathbf{X}} = (\tilde{X}_1, \dots, \tilde{X}_d) = \left(\frac{1}{1 - F_1(Y_1)}, \dots, \frac{1}{1 - F_d(Y_d)}\right),$$

such that $\tilde{X}_1, \ldots, \tilde{X}_d$ follow the standard Pareto distribution. Then conditional on the norm of $\tilde{\mathbf{X}}$ being large for a pre-specified norm $\|\cdot\|$, we have

$$\frac{\tilde{\mathbf{X}}}{r} \left\| \|\tilde{\mathbf{X}}\| > r \xrightarrow{d} R \cdot \Theta, \quad \text{as } r \to \infty, \right\|$$
(1.1)

where Θ is a random vector on the positive unit sphere $\{\mathbf{v} \in [0, \infty)^d | \|\mathbf{v}\| = 1\}$ and R is a standard Pareto variable independent of Θ . Here the law of Θ is called the *angular measure* or the *spectral measure*. This characterization is derived from the framework of *multivariate regular variation*. For a detailed overview, see e.g., Chapter 6 of Resnick (2007).

• Spectral random vector S: Consider an alternative marginal transformation

$$\mathbf{X} = (X_1, \dots, X_d) = (-\log(1 - F_1(Y_1)), \dots, -\log(1 - F_d(Y_d))), \quad (1.2)$$

such that X_1, \ldots, X_d follow the standard exponential distribution. Then conditional on the maximum component of **X** being large, we have

$$\mathbf{X} - r \cdot \mathbf{1} | \max(\mathbf{X}) > r \stackrel{d}{\to} \mathbf{Z}, \quad \text{as } r \to \infty,$$
(1.3)

where \mathbf{Z} has the stochastic representation

$$\mathbf{Z} :\stackrel{a}{=} E \cdot \mathbf{1} + \mathbf{S}_{\mathbf{z}}$$

such that **S** is a random vector on the irregular support $\{\mathbf{v} \in \mathbb{R}^d | \max(\mathbf{v}) = 0\}$ and *E* is a standard exponential random variable independent of **S**. Here **S** is called the *spectral random vector*. This characterization results from the framework of *multivariate peak-over-threshold*, see Rootzén and Tajvidi (2006) and Rootzén et al. (2018).

The two characterizations are connected as (1.3) is equivalent to (1.1) using the L_{∞} -norm. Both Θ and **S** can be used to summarize the extremal dependence structure. However, notice that the

supports of Θ and **S** are both nonlinear and induces intrinsic dependence between the dimensions. This poses nontrivial constraints for the construction of statistical models and their inference.

In this paper, we focus on the random vector \mathbf{X} with standard exponential margins and consider a different representation of the extremal dependence. We study the distribution of \mathbf{X} conditional on the component mean $\bar{X} = \frac{1}{d} \sum_{k=1}^{d} X_k$ being large. In the case where the tail of \mathbf{X} has asympotitically dependent components, we show that

$$\mathbf{X} - r \cdot \mathbf{1} \,|\, \bar{X} > r \stackrel{d}{\to} \mathbf{Z}^*, \quad \text{as } r \to \infty,$$

where the limiting distribution \mathbf{Z}^* can be represented as

$$\mathbf{Z}^* :\stackrel{a}{=} E \cdot \mathbf{1} + \mathbf{V} - \boldsymbol{\mu}_{\mathbf{V}},$$

such that

- V belongs to the class of centered random vectors on the hyperplane $\mathbf{1}^{\perp} := {\mathbf{v} | \mathbf{v}^T \mathbf{1} = 0}$ satisfying the moment condition $E[e^{\max(\mathbf{V})}] < \infty;$
- $\mu_{\mathbf{V}}$ is a constant vector determined by the distribution of **V**;
- E is a standard exponential random variable independent of \mathbf{V} .

We term \mathbf{V} the profile random vector.

There are two particular attractive properties in the characterization of profile random vectors. First, the class of profile random vectors \mathbf{V} resides on a linear vector space and is closed under finite addition and scalar multiplication. This allows for straightforward adaptation of existing statistical techniques based on linear operations, which may not be readily applied in the case of the angular component Θ or the spectral random vector \mathbf{S} . As an example, we illustrate the use of principal component analysis to achieve a lower-dimensional approximation of tail dependence structure.

Second, profile random vectors with Gaussian distributions result in the Hüsler-Reiss family (Hüsler and Reiss, 1989). The Hüsler-Reiss family is defined as the class of nontrivial tail dependence of Gaussian triangular arrays. It is one of the most widely used parametric models for extremal dependence. Despite its link to the Gaussian family, the analytical form of Hüsler-Reiss models is not easy to handle mathematically. Using profile random vectors, analyses for Hüsler-Reiss models can be translated to analyses for Gaussian models on the hyperplane $\mathbf{1}^{\perp}$.

The remainder of the paper is structured as follows. Section 2 recalls the multivariate peakover-threshold framework for modeling multivariate extremes. Section 3 introduces the diagonal peak-over-threshold framework and the profile random vectors, presenting their links to the peakover-threshold framework and spectral random vectors. Section 4 studies the case of Gaussian profile random vectors, namely the Hüsler-Reiss models. Section 5 discusses the application of principal component analysis on profile random vectors to achieve lower-dimensional approximation for extremes. The paper concludes with some discussions in Section 6, including what happens in the case where the components might be asymptotically independent. All proofs are postponed to the appendix.

Notation

Throughout the paper, boldface symbols are used to denote vectors, usually of length d. We write $\mathbf{0} = (0, \ldots, 0)$ and $\mathbf{1} = (1, \ldots, 1)$, where the lengths of the vector may depend on the context. For a vector $\mathbf{x} = (x_1, \ldots, x_d)$, denote its maximum component and component mean by $\max(\mathbf{x}) = \max(x_1, \ldots, x_d)$ and $\bar{x} = \frac{1}{d} \sum_{k=1}^d x_k$, respectively. When applied to vectors, mathematical operations, such as addition, multiplication, exponentiation, maximum and minimum are taken to be component-wise. Comparison between vectors are also considered component-wise, except for the notation $\mathbf{x} \notin \mathbf{y}$, which is interpreted as the event where $x_k > y_k$ for at least one k. Last but not the least, $\mathbf{1}^{\perp} := \{\mathbf{v} | \mathbf{v}^T \mathbf{1} = 0\}$ is used to denote hyperplane perpendicular to the vector $\mathbf{1}$.

2 Background on multivariate extremes

2.1 Multivariate generalized Pareto distributions

Let **X** be a random vector in \mathbb{R}^d . To study the tail of **X**, a common assumption is that there exist sequences of normalizing vectors $\{\mathbf{a}_n\}$ and $\{\mathbf{b}_n\}$ such that the component-wise maxima of **X** converges, i.e.,

$$\lim_{n \to \infty} P\left(\frac{\max_{i=1,\dots,n} \mathbf{X}_i - \mathbf{b}_n}{\mathbf{a}_n} \le \mathbf{x}\right) = G(\mathbf{x}),\tag{2.1}$$

where \mathbf{X}_i , i = 1, 2, ... are i.i.d. copies of \mathbf{X} . The limiting distribution G is then called a *generalized* extreme value distribution and we say that \mathbf{X} is in the domain of attraction of G, denoted as $\mathbf{X} \in DA(G)$. Each marginal of G follows a univariate generalized extreme value distribution, which can be parametrized by

$$G_k(x_k) = \exp\left\{-\left(1 + \gamma_k(x_k - \mu_k)/\alpha_k\right)^{-\frac{1}{\gamma_k}}\right\}, \quad 1 + \gamma_k(x_k - \mu_k)/\alpha_k > 0,$$

where $\gamma_k, \mu_k \in \mathbb{R}$ and $\alpha_k > 0$. In the case where $\gamma_k = 0$, $G_k(x_k)$ is interpreted as the limit $G_k(x_k) = \exp\{-\exp(-(x_k - \mu_k)/\alpha_k)\}$. The dependence structure of G cannot be parametrized and may be complex. For background on multivariate generalized extreme value distributions and their domains of attraction, see e.g., de Haan and Ferreira (2006).

The setting of this paper closely follows the *multivariate peak-over-threshold* framework, which is briefly recalled in the following. Assume that \mathbf{X} is in the domain of attraction of G. Then following elementary calculation from (2.1), the distribution of exceedances of \mathbf{X} , conditional on \mathbf{X} 'being extreme', converges to

$$\max\left\{\frac{\mathbf{X} - \mathbf{b}_n}{\mathbf{a}_n}, \boldsymbol{\eta}\right\} \le \mathbf{x} \left| \mathbf{X} \nleq \mathbf{b}_n \stackrel{d}{\to} \mathbf{Z}, \quad n \to \infty.$$
(2.2)

Here $\boldsymbol{\eta}$ is the vector of lower end points of the marginal distribution of G such that $\eta_k = \mu_k - \alpha_k / \gamma_k$ if $\gamma_k > 0$ and $\eta_k = -\infty$ otherwise. The limit distribution \mathbf{Z} is called a *multivariate generalized Pareto distribution* and has distribution function

$$H(\mathbf{z}) := P(\mathbf{Z} \le \mathbf{z}) = \frac{\ln G(\mathbf{z} \land \mathbf{0}) - \ln G(\mathbf{z})}{\ln G(\mathbf{0})}.$$
(2.3)

The conditional event of being extreme $\{\mathbf{X} \nleq \mathbf{b}_n\}$ is interpreted as $\{\exists k \ s.t. \ X_k > b_{nk}\}$, meaning that at least one of the X_k 's exceeds a high threshold. The marginal distribution Z_k may not be absolute continuous as it can have mass on $\{Z_k = 0\}$. Conditional on $\{Z_k > 0\}$, the marginal Z_k follows a univariate generalized Pareto distribution:

$$P(Z_k > z | Z_k > 0) = (1 + \gamma_j z / \sigma_k)_+^{-1/\gamma_k},$$

where $x_{+} = \max(x, 0)$ and $\sigma_{k} := \alpha_{k} - \gamma_{k}\mu_{k}$. A multivariate generalized Pareto distribution can therefore be characterized by $\boldsymbol{\sigma}, \boldsymbol{\gamma}$, the probabilities $P(Z_{k} > 0)$ for $1 \le k \le d$, and the dependence structure. For an overview on multivariate peak-over-threshold and multivariate generalized Pareto distributions, see Rootzén and Tajvidi (2006) and Rootzén et al. (2018).

2.2 Marginal standardization and stochastic representation

To focus exclusively on the extremal dependence structure of a random vector, we assume that the margins of **X** are standardized to the standard exponential distribution following transformation (1.2). Then the convergence of component-wise maxima (2.1) can be reformulated with $\mathbf{a}_n = \mathbf{1}$ and $\mathbf{b}_n = \log(n) \cdot \mathbf{1}$ as

$$\lim_{n \to \infty} P\left(\max_{i=1,\dots,n} \mathbf{X}_i - \log(n) \cdot \mathbf{1} \le \mathbf{x}\right) = G(\mathbf{x}),$$

where the marginal distributions of G follows a Gumbel distribution with $\gamma_k = \mu_k = 0$ and $\alpha_k = 1$ for all $k = 1, \ldots, d$. The convergence of exceedances (2.2) can be re-formulated as

$$\mathbf{X} - r \cdot \mathbf{1} \mid \max(\mathbf{X}) \ge r \stackrel{d}{\to} \mathbf{Z}, \quad r \to \infty,$$
 (2.4)

where **Z** is a multivariate generalized Pareto distribution with $\gamma = 0$, $\sigma = 1$ and $P(Z_1 > 0) = \cdots = P(Z_d > 0)$. Such a multivariate generalized Pareto distribution is said to be a standardized multivariate generalized Pareto distribution.

Rootzén et al. (2018) showed that the class of standardized multivariate generalized Pareto distributions can be represented stochastically by a class of random vectors on the L-shaped support $\{\mathbf{v} | \max(\mathbf{v}) = 0\}$.

Proposition 2.1 (Theorems 6 and 7 of Rootzén et al. (2018)). Let S be the class of random vectors $\mathbf{S} \in (-\infty, 0]^d$ such that $P(\max(\mathbf{S}) = 0) = 1$, $P(S_j > -\infty) > 0$, $1 \leq j \leq d$, and $E[e^{S_1}] = \cdots = E[e^{S_d}]$. Then a standardized multivariate generalized Pareto distribution \mathbf{Z} admits the representation

$$\mathbf{Z} \stackrel{d}{=} E \cdot \mathbf{1} + \mathbf{S},\tag{2.5}$$

where $\mathbf{S} \in \mathcal{S}$ and E is a standard exponential random variable independent of \mathbf{S} . Conversely, any $\mathbf{S} \in \mathcal{S}$ characterizes a standardized multivariate generalized Pareto distribution \mathbf{Z} through (2.5).

Here \mathbf{S} is referred to as the *spectral random vector* associated with \mathbf{Z} . Effectively, the spectral random vector is the limit

$$\mathbf{X} - \max(\mathbf{X}) \cdot \mathbf{1} \mid \max(\mathbf{X}) \ge r \stackrel{d}{\to} \mathbf{S}, \quad r \to \infty,$$

representing the tail of **X** being diagonally projected onto the L-shaped support $\{\mathbf{v} | \max(\mathbf{v}) = 0\}$.

2.3 Asymptotic dependence and extreme directions

Consider the support of a standardized multivariate generalized Pareto distribution $\mathbb{E} = \{\mathbf{v} | \| \mathbf{v} \|_{\infty} \geq 0\}$. Given all subsets $J \subseteq \{1, \ldots, d\}$, \mathbb{E} can be decomposed into the disjoint union of $\bigcup_{J \subseteq \{1, \ldots, d\}} \mathbb{E}_J$, where

$$\mathbb{E}_J = \{ \|\mathbf{v}\|_{\infty} \ge 0 : v_j > -\infty \text{ iff } j \in J \}.$$

If $P(\mathbf{Z} \in \mathbb{E}_J) > 0$, then J is called an extreme direction of \mathbf{X} (Mourahib et al., 2024). Intuitively, this means that there is a positive probability the variables X_j 's for $j \in J$ are large together while the other variables are not. In the case where $\{1, \ldots, d\}$ is the only extreme direction, that is, the multivariate generalized Pareto distribution has support $\{\mathbf{v} | \| \mathbf{v} \|_{\infty} \ge 0, v_j > -\infty, j = 1, \ldots, d\}$, we say that the components of \mathbf{X} are asymptotically dependent. The corresponding spectral random vector satisfies $P(S_j > -\infty) = 1, 1 \le j \le d$.

In this paper, we focus on the scenario where the components of \mathbf{X} are asymptotically dependent. Under this assumption, we show that the extremal dependence structure can be modeled with an alternative, advantageous characterization. On the other hand, a generic tail dependence structure can be constructed via a mixture model with factors of asymptotic dependent components. Specifically, Mourahib et al. (2024) showed that a multivariate generalized Pareto distribution \mathbf{Z} can be represented by a mixture model whose factors consist of

$$\mathbf{Z}_J = \mathbf{Z} \,|\, \{\mathbf{Z} \in \mathbb{E}_J\}$$

for every extreme direction J of \mathbf{X} . Each \mathbf{Z}_J is denegerate on the components in J^c and hence can be modeled by a |J|-dimensional multivariate generalized Pareto distribution with asymptotically dependent components.

3 Diagonal peak-over-threshold and profile random vectors

3.1 Diagonal peak-over-threshold

In this section, we consider a different peak-over-threshold framework. Instead of conditioning on $\{\max(\mathbf{X}) \geq r\}$, consider conditioning on $\{\bar{X} > r\}$, where the component mean of \mathbf{X} exceeds a high threshold. We have the following proposition.

Proposition 3.1. Let $\mathbf{X} \in DA(G)$ be a random vector such that (2.4) holds with a standardized multivariate generalized Pareto distribution \mathbf{Z} with asymptotic dependent components. Then

$$\mathbf{X} - r \cdot \mathbf{1} \mid \bar{X} \ge r \stackrel{d}{\to} \mathbf{Z}^*, \quad r \to \infty,$$
(3.1)

where

$$\mathbf{Z}^* \stackrel{d}{=} \mathbf{Z} \mid \{ \mathbf{Z}^T \mathbf{1} \ge 0 \}.$$
(3.2)

We call the limiting distribution \mathbf{Z}^* a diagonal multivariate generalized Pareto distribution. If a pair of standardized and diagonal multivariate generalized Pareto distributions (\mathbf{Z}, \mathbf{Z}^*) satisfies (3.2), then we say they are associated. **Remark 3.2.** In the case where the components of \mathbf{X} are not asymptotically dependent, the components of \mathbf{Z} have mass on $-\infty$, resulting in the possibility of $\{\mathbf{Z}^T \mathbf{1} > 0\}$ having probability 0. This paper focuses on the scenario where \mathbf{X} has asymptotically dependent components. The scenario for random vectors with asymptotically independent components is considered in an ensuing work and briefly discussed in Section 6.

Remark 3.3. Proposition 3.1 does not explicitly assume that \mathbf{X} has unit exponential margins. Instead, random vectors with marginal distributions that behaves similarly to the unit exponential in the tail can also be considered.

3.2 Profile random vectors

As stated in Proposition 2.1, the class of standardized multivariate generalized Pareto distributions can be characterized by the class of spectral random vector \mathbf{S} on the L-shaped space $\{\mathbf{v} | \max(\mathbf{v}) = 0\}$. In the following proposition, we show that the class of diagonal multivariate generalized Pareto distributions can be characterized by a class of random vectors on the hyperplane $\mathbf{1}^{\perp} := \{\mathbf{v} | \mathbf{v}^T \mathbf{1} = 0\}$.

Proposition 3.4. Let \mathcal{V} be the class of random vectors $\mathbf{V} \in \mathbf{1}^{\perp}$ such that $E[\mathbf{V}] = \mathbf{0}$ and $E[e^{\max(\mathbf{V})}] < \infty$. Then any diagonal multivariate generalized Pareto distribution \mathbf{Z}^* has the stochastic representation

$$\mathbf{Z}^* \stackrel{d}{=} E \cdot \mathbf{1} + \mathbf{V} - \boldsymbol{\nu}_{\mathbf{V}},\tag{3.3}$$

for some $\mathbf{V} \in \mathcal{V}$, where E is a standard exponential variable independent of \mathbf{V} and $\nu_{\mathbf{V}}$ is the constant vector such that

$$\boldsymbol{\nu}_{\mathbf{V}} = \log\left(\frac{E\left[e^{\mathbf{V}}\right]}{\left(\prod_{k=1}^{d} E\left[e^{V_{k}}\right]\right)^{1/d}}\right).$$
(3.4)

We name **V** the profile random vector associated with \mathbf{Z}^* . Conversely, any $\mathbf{V} \in \mathcal{V}$ defines a diagonal multivariate generalized Pareto distribution via (3.3), with $\nu_{\mathbf{V}}$ as defined in (3.4).

As will be shown in the following subsection, the profile random vector \mathbf{V} and the spectral random vector \mathbf{S} have a one-to-one correspondence for asymptotically dependent random vectors and hence can both be used to characterize extremal dependence. Note that the class of profile random vectors $\mathcal{V} = {\mathbf{V} \in \mathbf{1}^{\perp} | E[\mathbf{V}] = \mathbf{0}, E[e^{\max(\mathbf{V})}] < \infty}$ resides on a linear vector space and is closed under finite addition and scalar multiplication. This provides a context to apply statistical analysis based on linear techniques to analyze extremes, as we shall see in Section 5 for the example of principal component analysis.

3.3 Link between spectral and profile random vectors

Given a pair of associated standardized and diagonal multivariate generalized Pareto distributions $(\mathbf{Z}, \mathbf{Z}^*)$, let **S** and **V** be the corresponding spectral and profile random vectors. This subsection

establishes the link between associated \mathbf{S} and \mathbf{V} . To present our results, we consider a pair of transformations of \mathbf{S} and \mathbf{V} .

Define the \mathbf{T} -generator of \mathbf{S} to be

$$\mathbf{T} := \mathbf{S} - \bar{S} \cdot \mathbf{1},\tag{3.5}$$

and the **U**-generator of \mathbf{V} to be

 $\mathbf{U} := \mathbf{V} - \boldsymbol{\nu}_{\mathbf{V}},$

where $\nu_{\mathbf{V}}$ is as defined in (3.4). Then the **T**-generators form the class of random vectors $\mathcal{T} = \{\mathbf{T} \in \mathbf{1}^{\perp} | E[e^{T_1 - \max(\mathbf{T})}] = \cdots = E[e^{T_d - \max(\mathbf{T})}] < \infty\}$ and the **U**-generators form the class of random vectors $\mathcal{U} = \{\mathbf{U} \in \mathbf{1}^{\perp} | E[e^{U_1}] = \cdots = E[e^{U_d}] < \infty\}$. A pair of **T**- and **U**-generators is said to be associated if their corresponding spectral and profile random vectors are associated. The **S** and **V** can be easily retrieved from **T** and **U** by $\mathbf{S} = \mathbf{T} - \max(\mathbf{T})$ and $\mathbf{V} = \mathbf{U} - E[\mathbf{U}]$.

The relationship between \mathbf{T} and \mathbf{U} is given as follows.

Proposition 3.5. Let T and U be associated T- and U-generators. Then

$$\mathbf{U} \mid \{ \max(\mathbf{U}) = s \} \stackrel{d}{=} \mathbf{T} \mid \{ \max(\mathbf{T}) = s \}, \quad \forall s \ge 0.$$
(3.6)

Given the distribution of $\max(\mathbf{T})$, the distribution of $\max(\mathbf{U})$ can be obtained from

$$P(\max(\mathbf{U}) \le s) = \frac{\int_0^s P(\max(\mathbf{T}) \le t)e^{-t}dt + e^{-s}P(\max(\mathbf{T}) \le s)}{E\left[e^{-\max(\mathbf{T})}\right]}, \quad s \ge 0.$$
(3.7)

Conversely, given the distribution of $\max(\mathbf{U})$, the distribution of $\max(\mathbf{T})$ can be obtained from

$$P(\max(\mathbf{T}) \le s) = \frac{e^s P(\max(\mathbf{U}) \le s) - \int_0^s P(\max(\mathbf{U}) \le t) e^t dt}{E\left[e^{\max(\mathbf{U})}\right]}, \quad s \ge 0.$$
(3.8)

The relationship between \mathbf{T} and \mathbf{U} can also be stated through the following stochastic transformations.

Corollary 3.6. Let \mathbf{T} and \mathbf{U} be associated \mathbf{T} - and \mathbf{U} -generators. Then given a unit exponential variable E independent of \mathbf{T} ,

$$\mathbf{T} | \{ \max(\mathbf{T}) < E' \} \stackrel{d}{=} \mathbf{U}.$$

$$(3.9)$$

Given a unit exponential variable E' independent of \mathbf{U} ,

$$\mathbf{U} | \{ \max(\mathbf{U}) \ge r - E \} \stackrel{d}{\to} \mathbf{T}, \quad r \to \infty.$$
(3.10)

In the case where $\max(\mathbf{T})$ and $\max(\mathbf{U})$ are absolutely continuous, the link can be simplified via density functions.

Corollary 3.7. If $\max(\mathbf{T})$ is absolutely continuous and admits density $f_{\max(\mathbf{T})}$, then $\max(\mathbf{U})$ is absolutely continuous with density

$$f_{\max(\mathbf{U})}(s) = \frac{1}{E\left[e^{-\max(\mathbf{T})}\right]} \cdot f_{\max(\mathbf{T})}(s) \cdot e^{-s}.$$

$$f_{\max(\mathbf{T})}(s) = \frac{1}{E\left[e^{\max(\mathbf{U})}\right]} \cdot f_{\max(\mathbf{U})}(s) \cdot e^{s}.$$

Remark 3.8. The names \mathbf{T} - and \mathbf{U} -generators are inherited from Rootzén et al. (2018), who proposed that given a spectral random vector \mathbf{S} , any random vector \mathbf{T} such that

$$\mathbf{S} = \mathbf{T} - \max(\mathbf{T}) \cdot \mathbf{1},$$

is a \mathbf{T} -generator for \mathbf{S} , and any random vector \mathbf{U} such that

$$\frac{E\left[\max\left(\mathbf{y}e^{\mathbf{U}}\right)\right]}{E\left[e^{U_{1}}\right]} = \frac{E\left[\max\left(\mathbf{y}e^{\mathbf{S}}\right)\right]}{E\left[e^{S_{1}}\right]}, \quad \forall \mathbf{y} \in [0,\infty)^{d},$$

is a U-generator for S. It can be shown that our definitions of T and U corresponds to the unique T- and U-generators for S on 1^{\perp} .

3.4 Generating random vector with specific profile random vectors

Finally, it is straightforward to generate random vectors whose extremal dependence is characterized by a given profile random vector \mathbf{V} .

Proposition 3.9. Let **X** be a random vector in \mathbb{R}^d defined by

$$\mathbf{X} \stackrel{d}{=} E \cdot \mathbf{1} + \mathbf{V} - \boldsymbol{\nu}_{\mathbf{V}},$$

where $\mathbf{V} \in \mathbf{1}^{\perp}$ is a centered random vector satisfying $E[e^{\max(\mathbf{V})}] < \infty$, $\nu_{\mathbf{V}}$ is as defined in (3.4), and E is a standard exponential random variable independent of \mathbf{V} . Then \mathbf{X} satisfies (2.4) and (3.1). Its diagonal multivariate generalized Pareto distribution is characterized by profile random vector \mathbf{V} and its standardized multivariate generalized Pareto distribution is characterized by associated spectral random vector \mathbf{S} .

4 Gaussian profile random vectors

Any parametric family on \mathcal{V} induces a parametric family for profile random vectors. For example, let $\tilde{\mathbf{V}} = (\tilde{V}_1, \ldots, \tilde{V}_d)$ be a random vector with independent Gumbel components $\tilde{V}_k \stackrel{iid}{\sim} \tilde{V}$ such that $P(\tilde{V} \leq v) = \exp[-\exp(-\alpha v)]$ for some $\alpha > 0$. Then $\mathbf{V} := \tilde{\mathbf{V}} - \left(\frac{1}{d}\sum_{k=1}^{d} \tilde{V}_k\right) \cdot \mathbf{1}$ is the profile random vector for the well-known *multivariate logistic model*. More parametric examples can be derived from that of **U**-generators in Kiriliouk et al. (2019).

In this section, we focus on the case where the profile random vector follows a Gaussian distribution on the hyperplane $\mathbf{1}^{\perp}$. This results in the family of Hüsler-Reiss models, the class of distributions describing the non-trivial tail limit of Gaussian triangular arrays (Hüsler and Reiss, 1989), which we briefly recall in the following. Consider a Gaussian random vector with unit variance $\mathbf{X} \sim N(\mathbf{0}, \Sigma)$ where $\Sigma_{kk} = 1, k = 1, \ldots, d$. For any $i \neq j$, in the case where $\Sigma_{ij} < 1$, it can be shown that the components X_i and X_j are asymptotically independent in the tail (Sibuya, 1960). In order to construct nontrivial extremal dependence, consider instead a Gaussian triangular array $\mathbf{X}_i^{(n)} \sim N(\mathbf{0}, \Sigma^{(n)}), i = 1, \ldots, n$ where $\Sigma_{kk}^{(n)} = 1, k = 1, \ldots, d$. Assume that the elements of $\Sigma^{(n)}$ converge to 1 such that

$$\log(n) \cdot (\mathbf{1}\mathbf{1}^T - \Sigma^{(n)}) \to \Gamma = (\Gamma_{ij})_{1 \le i,j \le d}.$$

Here Γ satisfies that $\Gamma_{ij} = E(W_i - W_j)^2$ for some centered multivariate Gaussian random vector $\mathbf{W} = (W_1, \ldots, W_d)$ and is called the *variogram* of \mathbf{W} .

A Hüsler-Reiss model parametrized by Γ is characterized by the limiting tail distribution of $\mathbf{X}^{(n)}$, whose generalized extreme value distribution is defined as the limit

$$\lim_{n \to \infty} P\left(\frac{\max_{i=1,\dots,n} \mathbf{X}_i^{(n)} - \mathbf{b}_n}{\mathbf{a}_n} \le \mathbf{x}\right) = G_{\Gamma}(\mathbf{x}),$$

for suitable normalizing sequences $\{\mathbf{a}_n\}$ and $\{\mathbf{b}_n\}$. While not as easy to handle mathematically as the Gaussian distribution, the Hüsler-Reiss models remain the one of the most widely used parametric family for multivariate extremes and is often referred to as the Gaussian counterpart for extremes.

The following proposition shows that the profile random vector of a Hüsler-Reiss model is a Gaussian random vector on the hyperplane $\mathbf{1}^{\perp}$.

Proposition 4.1. The profile random vector of the Hüsler-Reiss model parametrized by Γ is

$$\mathbf{V} \sim N\left(\mathbf{0}, \Sigma\right),$$

where

$$\Sigma := -\frac{1}{2} \left(I - \frac{\mathbf{1}^T \mathbf{1}}{d} \right) \Gamma \left(I - \frac{\mathbf{1}^T \mathbf{1}}{d} \right).$$
(4.1)

In other words, V is the unique centered Gaussian random vector on $\mathbf{1}^{\perp}$ with variogram Γ .

Remark 4.2. Proposition 4.1 was independently derived in an unpublished manuscript by Johan Segers in 2019. In the special case where the variogram matrix Γ is of rank (d-1) and the Hüsler-Reiss multivariate generalized Pareto distribution **Z** admits a density, this result was proven in Corollary 3.7 of Hentschel et al. (2024).

It is also straightforward to construct random vectors with Hüsler-Reiss extremal dependence structure characterized by a given variogram matrix Γ . Let **X** be a random vector defined by

$$\mathbf{X} \stackrel{d}{=} E \cdot \mathbf{1} + \mathbf{V} - \boldsymbol{\nu}_{\mathbf{V}},$$

where $\mathbf{V} \sim N(\mathbf{0}, \Sigma)$ for Σ as defined in (4.1), $\nu_{\mathbf{V}}$ is as defined in (3.4), and E is a standard exponential random variable independent of \mathbf{V} . From Proposition 3.9, the tail of \mathbf{X} follows a Hüsler-Reiss model parametrized by Γ .

In the recent literature on Hüsler-Reiss models, Γ is often assumed to be the variogram of a full-rank Gaussian vector such that the resulting multivariate generalized Pareto distribution \mathbf{Z} admits a density. In this case, the resulting Σ is of rank d-1 and has the eigen decomposition $\Sigma = \sum_{k=1}^{d-1} \lambda_k \mathbf{u}_k \mathbf{u}_k^T$ where $\lambda_k > 0$ for $k = 1, \ldots, d-1$. The last eigenvector $\mathbf{u}_d = \frac{1}{\sqrt{d}} \mathbf{1}$ corresponds to eigenvalue $\lambda_d = 0$. Its pseudo-inverse $\Theta = \sum_{k=1}^{d-1} \frac{1}{\lambda_k} \mathbf{u}_k \mathbf{u}_k^T$ embeds the conditional independence information in the tail and serves as a precision matrix to the extremal graphical model. For extremal graphical models and the precision matrices of Hüsler-Reiss graphical models, see Engelke and Hitz (2020), Hentschel et al. (2024) and Wan and Zhou (2023).

The result in Proposition 4.1 generalizes to Hüsler-Reiss models of all ranks. In fact, Section 5 illustrates that being able to characterize lower-rank models for multivariate extremes allows the possibility of lower-dimensional approximation for tail data.

5 Principal component analysis

In this section, we illustrate the application of principal component analysis to achieve a lowerdimensional approximation to the extremal dependence structure.

Principal component analysis is a classical technique in multivariate analysis for finding lower dimensional representations of a random vector while retaining most of its variability. Given a centered random vector $\mathbf{X} \in \mathbb{R}^d$, principal component analysis identifies the linear subspace $\mathcal{S}_p^* \subset \mathbb{R}^d$ of dimension p < d such that the L_2 -distance between \mathbf{X} and its projection $\prod_{\mathcal{S}_p^*} \mathbf{X}$ onto \mathcal{S}_p^* is minimized:

$$\mathcal{S}_p^* := \arg \min_{\mathcal{P}} E \| \Pi_{\mathcal{P}} \mathbf{X} - \mathbf{X} \|_2^2.$$

This is achieved by considering the orthonormal eigenvectors $\mathbf{v}_1, \ldots, \mathbf{v}_d$ of the covariance matrix $E(\mathbf{X}\mathbf{X}^T)$ with ordered eigenvalues $\lambda_1 \geq \ldots \geq \lambda_d \geq 0$. The projection of $\Pi_{\mathbf{v}_k}\mathbf{X}$ onto the subspace spanned by \mathbf{v}_k is called the k-th principal component of \mathbf{X} . The optimal subspace \mathcal{S}_p^* is the span of $\mathbf{v}_1, \ldots, \mathbf{v}_p$ and the best p-dimensional approximation of \mathbf{X} is the sum of its first p principal components

$$\Pi_{\mathcal{S}_p^*} \mathbf{X} = \Pi_{\mathbf{v}_1} \mathbf{X} + \dots + \Pi_{\mathbf{v}_p} \mathbf{X}.$$

Previous literature applying principal component analysis to extremes has focused on applying the principal component analysis to the angular component Θ , see Cooley and Thibaud (2019) and Drees and Sabourin (2021). However, Θ resides on the unit sphere { $\mathbf{v} \in [0, \infty)^d || \mathbf{v} || = 1$ }, which is not a linear subspace. Hence any lower dimensional approximation of Θ via principal component analysis will no longer result in an angular component.

In this section, let us consider constructing a lower dimensional approximation of a profile random vector \mathbf{V} via principal component analysis. First, given the moment constraint $E[e^{\max(\mathbf{V})}] < \infty$, the covariance matrix $E[\mathbf{V}\mathbf{V}^T]$ always exists. Second, since $\mathbf{V} \in \mathbf{1}^{\perp}$, the last eigenvector \mathbf{v}_d is equal to $\mathbf{1}/\sqrt{d}$ with eigenvalue $\lambda_d = 0$, and hence

$$\mathbf{V} = \Pi_{\mathbf{v}_1} \mathbf{V} + \dots + \Pi_{\mathbf{v}_{d-1}} \mathbf{V}.$$

Each principal component $\Pi_{\mathbf{v}_k} \mathbf{V}$ is a profile random vector on its own and can be interpreted as the extremal dependence along direction \mathbf{v}_k . For any p < d-1, the *p*-dimensional approximation of \mathbf{V} is

$$\Pi_{\mathcal{S}_n^*}\mathbf{V} = \Pi_{\mathbf{v}_1}\mathbf{V} + \dots + \Pi_{\mathbf{v}_p}\mathbf{V},$$

which also defines a profile random vector. This induces a lower-dimensional approximation for the associated diagonal multivariate generalized Pareto distribution \mathbf{Z}^* , standardized multivariate generalized Pareto distribution \mathbf{Z} and spectral random vector \mathbf{S} .

Recall from Proposition 4.1 that a Hüsler-Reiss model has profile random vector $\mathbf{V} \sim N(\mathbf{0}, \Sigma)$, where Σ is any positive semidefinite matrix on $\mathbf{1}^{\perp}$. Let $\mathbf{v}_1, \ldots, \mathbf{v}_d$ be the eigenvectors of Σ corresponding to ordered eigenvalues $\lambda_1 \geq \ldots \geq \lambda_d$. Then $\mathbf{v}_d = \mathbf{1}/\sqrt{d}$ and $\lambda_d = 0$. Each principal component of \mathbf{V} can be written as

$$\Pi_{\mathbf{v}_k} \mathbf{V} \sim N(\mathbf{0}, \mathbf{v}_k \mathbf{v}_k^T).$$

Coversely, for k = 1, ..., d - 1, let \mathbf{V}_k be independent profile random vectors such that $\mathbf{V}_k \sim N(\mathbf{0}, \mathbf{v}_k \mathbf{v}_k^T)$. The **V** can be written as

$$\mathbf{V} \stackrel{d}{=} \mathbf{V}_1 + \dots + \mathbf{V}_{d-1}.$$

In other words, the dependence structure of a Hüsler-Reiss model can be decomposed into that of at most (d-1) Hüsler-Reiss models, each of whose dependence structure is concentrated on one specific direction. The *p*-dimensional approximation of **V** is achieved by

$$\mathbf{V} \stackrel{d}{\approx} \mathbf{V}_1 + \dots + \mathbf{V}_p.$$

In conventional PCA, the discarded principal components describe directions where the variation of the data is minimized. In the PCA for profile random vectors, the discarded principal components describe the directions where the extremal dependence is strong enough to be approximated by complete dependence. Consider the trivial case where **V** can be approximated by the trivial constant **0**, then the diagonal multivariate generalized Pareto distribution $\mathbf{Z}^* \stackrel{d}{=} E \cdot \mathbf{1}$ lies on the vector **1**, meaning that all components are completely dependent in the tail.

6 Discussions

In this paper, we propose to characterize the extremal dependence of a multivariate random vector by a measure the hyperplane $\mathbf{1}^{\perp}$, namely the profile random vectors. The main advantage of the profile random vectors is that they reside on a linear vector space and are closed under finite addition and scalar multiplication. This provides a context to apply statistical analysis based on linear techniques to analyze the extremes. We have illustrated that principal component analysis can be applied naturally to achieve a lower-dimensional representation of the extremal dependence structure. Other possible applications include unsupervised learning, such as clustering, or supervised classification, such as linear discriminant analysis. In addition, the widely used Hüsler-Reiss models are characterized by Gaussian profile random vectors. On one hand, this opens up the possibility for alternative and potentially more efficient inference for the Hüsler-Reiss models. On the other hand, this provides a setting to extend the Hüsler-Reiss models to mixture models, in parallel to Gaussian mixture models.

The scenario which this paper has not discussed is when a random vector has asymptotically independent components. This will be explored in future work but we present below a small illustration of what could happen. Consider the simple example of a two-dimensional vector $\mathbf{X} = (X_1, X_2)$ with standard exponential margins. Denote $\mathbf{Y} = (Y_1, Y_2) = (e^{X_1}, e^{X_2})$ which has standard Pareto margins. Then projecting the tail of \mathbf{X} onto the hyperplane $\{(x_1, x_2)|x_1 + x_2 = 0\}$ is equivalent to projecting the tail of \mathbf{Y} to $\{(y_1, y_2)|(y_1y_2)^{1/2} = 1\}$. In the case where X_1 and X_2 (hence Y_1 and Y_2) are asymptotically independent, the projection reveals the dependence between the two components that is characterized by hidden regular variation, see e.g. Maulik and Resnick (2004) for more details. In the case where the dimension of the vector $d \geq 3$, additional consideration should also be given to the scenario that the extremal dependence is the combination of multiple extremal directions.

Acknowledgement

This research is supported by the Veni grant from the Dutch Research Council (VI.Veni.211E.034). The author would like to thank Anja Janßen, Chen Zhou, and other participants of Oberwolfach Workshop *Mathematics, Statistics, and Geometry of Extreme Events in High Dimensions* (2024) for extensive comments and discussions.

A Proofs

Proof of Proposition 3.1. Consider the conditional distribution $\mathbf{X} - r \cdot \mathbf{1} | \bar{X} \ge r$, we have

$$P(\mathbf{X} - r \cdot \mathbf{1} \le \mathbf{z} | \bar{X} \ge r) = \frac{P(\mathbf{X} - r \cdot \mathbf{1} \le \mathbf{z}, X \ge r | \max(\mathbf{X}) \ge r)}{P(\bar{X} \ge r | \max(\mathbf{X}) \ge r)} \\ = \frac{P(\mathbf{X} - r \cdot \mathbf{1} \le \mathbf{z}, \bar{X} \ge r | \max(\mathbf{X}) \ge r)}{P((\mathbf{X} - r \cdot \mathbf{1})^T \mathbf{1} \ge 0 | \max(\mathbf{X}) \ge r)}.$$

Taking the limit $r \to \infty$, we have

$$\lim_{r \to \infty} P(\mathbf{X} - r \cdot \mathbf{1} \le \mathbf{z} | \bar{X} \ge r) = \frac{\lim_{r \to \infty} P(\mathbf{X} - r \cdot \mathbf{1} \le \mathbf{z}, \bar{X} \ge r | \max(\mathbf{X}) \ge r)}{\lim_{r \to \infty} P((\mathbf{X} - r \cdot \mathbf{1})^T \mathbf{1} \ge 0 | \max(\mathbf{X}) \ge r)} = \frac{P(\mathbf{Z} \le \mathbf{z}, \mathbf{Z}^T \mathbf{1} \ge 0)}{P(\mathbf{Z}^T \mathbf{1} \ge 0)}.$$

To take the last equality, it remains to justify that $P(\mathbf{Z}^T \mathbf{1} \ge 0) > 0$. Since the components of \mathbf{X} and hence \mathbf{Z} are asymptotically dependent, we have $P(S_j > -\infty) = 1$ for $1 \le j \le d$. Hence there exists M > 0 such that $P(\min(\mathbf{S}) > -M) > 0$. We have

$$P(\mathbf{Z}^T \mathbf{1} \ge 0) = P((E \cdot \mathbf{1} + \mathbf{S})^T \mathbf{1} \ge 0)$$

$$\geq P(\min(\mathbf{S}) > -M, E > M)$$

= $P(\min(\mathbf{S}) > -M) \cdot P(E > M) > 0.$

Therefore $\mathbf{Z}^* \stackrel{d}{=} \mathbf{Z} \mid \mathbf{Z}^T \mathbf{1} \ge 0.$

Proofs of Proposition 3.4. To prove this proposition, we make use of the definitions of **T**-generator and **U**-generator introduced in Section 3.3.

A **T**-generator of **S** is defined by $\mathbf{T} := \mathbf{S} - \bar{S} \cdot \mathbf{1}$. From Proposition 3.1,

$$\mathbf{Z}^* \stackrel{d}{=} \mathbf{Z} \,|\, \mathbf{Z}^T \mathbf{1} \ge 0.$$

Since **Z** can be written as $\mathbf{Z} = E \cdot \mathbf{1} + \mathbf{T} - \max(\mathbf{T}) \cdot \mathbf{1}$, the conditional event i

$$\{\mathbf{Z}^T \mathbf{1} \ge 0\} = \{E \cdot d + \mathbf{T}^T \mathbf{1} - \max(\mathbf{T}) \cdot d \ge 0\} = \{E \cdot d - \max(\mathbf{T}) \cdot d \ge 0\} = \{E - \max(\mathbf{T}) \ge 0\},\$$

following the fact that $\mathbf{T} \in \mathbf{1}^{\perp}$ and hence $\mathbf{T}^T \mathbf{1} = 0$. Therefore

$$\mathbf{Z}^* \stackrel{d}{=} \left(\left(E - \max \mathbf{T} \right) \cdot \mathbf{1} + \mathbf{T} \right) \mid E \ge \max(\mathbf{T}).$$

For any $s \ge 0$ and Borel set $B \subseteq \mathbf{1}^{\perp}$,

$$\begin{split} P(E - \max(\mathbf{T}) \ge s, \mathbf{T} \in B | E \ge \max(\mathbf{T})) &= \frac{P(E - \max(\mathbf{T}) \ge s, \mathbf{T} \in B)}{P(E \ge \max(\mathbf{T}))} \\ &= \frac{\int_s^\infty P(\max(\mathbf{T}) \le t - s, \mathbf{T} \in B)e^{-t}dt}{\int_0^\infty P(\max(\mathbf{T}) \le t)e^{-t}dt} \\ u = t - s &= \frac{\int_0^\infty P(\max(\mathbf{T}) \le u, \mathbf{T} \in B)e^{-(u+s)}du}{\int_0^\infty P(\max(\mathbf{T}) \le t)e^{-t}dt} \\ &= e^{-s} \cdot \frac{\int_0^\infty P(\max(\mathbf{T}) \le u, \mathbf{T} \in B)e^{-u}du}{\int_0^\infty P(\max(\mathbf{T}) \le t)e^{-t}dt} \end{split}$$

Take $B = \mathbf{1}^{\perp}$, then

$$P(E - \max(\mathbf{T}) \ge s | E \ge \max(\mathbf{T})) = e^{-s}$$

Take s = 0, then

$$P(\mathbf{T} \in B | E \ge \max(\mathbf{T})) = \frac{\int_0^\infty P(\max(\mathbf{T}) \le u, \mathbf{T} \in B) e^{-u} dt}{\int_0^\infty P(\max(\mathbf{T}) \le t) e^{-t} dt}$$

Therefore the conditional distribution of $E - \max(\mathbf{T}) | E \ge \max(\mathbf{T})$ is again a unit exponential distribution and $E - \max(\mathbf{T})$ and \mathbf{T} are conditionally independent given $E \ge \max(\mathbf{T})$. Define

$$\mathbf{U} \stackrel{a}{:=} \mathbf{T} | \max(\mathbf{T}) \le E, \tag{A.1}$$

then

$$\mathbf{Z}^* \stackrel{d}{=} E' \cdot \mathbf{1} + \mathbf{U}$$

where E' is a unit exponential distribution independent of **U**. Since the **T**-generators form the class of vectors \mathcal{T} , from (A.1), the vectors **U** form the class of random vectors \mathcal{U} .

It remains to show that there is a one-to-one correspondence between $\mathbf{U} \in \mathcal{U}$ and $\mathbf{V} \in \mathcal{V}$ via

$$U = V + \nu_V$$

Given $\mathbf{U} \in \mathcal{U}$, we have

$$E[\mathbf{U}] \le E[\max(\mathbf{U})] \cdot \mathbf{1} \le E[e^{\max(\mathbf{U})} - 1] \cdot \mathbf{1} < \infty$$

Therefore we can construct $\mathbf{V} = \mathbf{U} - E[\mathbf{U}] \in \mathcal{V}$. Given any $\mathbf{V} \in \mathcal{V}$, we seek to find a constant vector $\boldsymbol{\nu} \in \mathbf{1}^{\perp}$ such that

$$\mathbf{U} = \mathbf{V} - \boldsymbol{\nu} \in \mathcal{U}.$$

this holds if and only if

$$E[e^{V_1-\nu_1}] = \dots = E[e^{V_d-\nu_d}] =: M.$$

Since $\nu \in \mathbf{1}^{\perp}$, we have

$$M^{d} = \prod_{k=1}^{d} E[e^{V_{k} - \nu_{k}}] = \left(\prod_{k=1}^{d} E[e^{V_{k}}]\right) \cdot e^{-\sum_{k=1}^{d} \nu_{k}} = \prod_{k=1}^{d} E[e^{V_{k}}].$$

Therefore $\boldsymbol{\nu}$ must take value in $\boldsymbol{\nu} = \boldsymbol{\nu}_{\mathbf{V}}$ where

$$e^{\boldsymbol{\nu}_{\mathbf{V}}} = \frac{E\left[e^{\mathbf{V}}\right]}{M} = \frac{E\left[e^{\mathbf{V}}\right]}{\left(\prod_{k=1}^{d} E\left[e^{V_{k}}\right]\right)^{1/d}}.$$

Proof of Proposition 3.5. From (A.1), for any $s \ge 0$

$$P(\max(\mathbf{U}) \le s) = P(\max(\mathbf{T}) \le s | \max(\mathbf{T}) \le E).$$

Therefore

$$\max(\mathbf{U}) \stackrel{d}{=} \max(\mathbf{T}) \,|\, \max(\mathbf{T}) \,\leq E.$$

Given any Borel set $B \subseteq \mathbf{1}^{\perp}$,

$$P(\mathbf{U} \in B | \max(\mathbf{U}) = s) = P(\mathbf{T} \in B | \max(\mathbf{T}) = s, \max(\mathbf{T}) \le E)$$
$$= P(\mathbf{T} \in B | \max(\mathbf{T}) = s, E \ge s)$$
$$= P(\mathbf{T} \in B | \max(\mathbf{T}) = s).$$

Therefore

$$\mathbf{U} \mid \{\max(\mathbf{U}) = s\} \stackrel{d}{=} \mathbf{T} \mid \{\max(\mathbf{T}) = s\}.$$

For any $s \ge 0$,

$$P(\max(\mathbf{U}) \le s) = P(\max(\mathbf{T}) \le s | \max(\mathbf{T}) \le E)$$

$$= \frac{P(\max(\mathbf{T}) \leq s, \max(\mathbf{T}) \leq E)}{P(\max(\mathbf{T}) \leq E)}$$

$$= \frac{\int_{0}^{s} P(\max(\mathbf{T}) \leq t)e^{-t}dt + \int_{s}^{\infty} P(\max(\mathbf{T}) \leq s)e^{-t}dt}{\int_{0}^{\infty} P(\max(\mathbf{T}) \leq t)e^{-t}dt}$$

$$\stackrel{u=e^{-t}}{=} \frac{\int_{0}^{s} P(\max(\mathbf{T}) \leq t)e^{-t}dt + P(\max(\mathbf{T}) \leq s)\int_{s}^{\infty} e^{-t}dt}{\int_{0}^{1} P(\max(\mathbf{T}) \leq -\log(u))du}$$

$$= \frac{\int_{0}^{s} P(\max(\mathbf{T}) \leq t)e^{-t}dt + e^{-s} \cdot P(\max(\mathbf{T}) \leq s)}{\int_{0}^{1} P(e^{-\max(\mathbf{T})} \geq u)du}$$

$$= \frac{\int_{0}^{s} P(\max(\mathbf{T}) \leq t)e^{-t}dt + e^{-s} \cdot P(\max(\mathbf{T}) \leq s)}{E[e^{-\max(\mathbf{T})}]}$$

Given U, let T be a random vector for whom the joint distribution $(\mathbf{T}|_{\max(\mathbf{T})}, \max(\mathbf{T}))$ is defined via (3.6) and (3.8). It can be seen that $\mathbf{T} \in \mathcal{T}$. Let Z be the standardized multivariate generalized Pareto generated by T and let Z^{*} be its associated diagonal multivariate generalized Pareto distribution. Denote the U-generator for Z^{*} by U_T obtained from T via (3.6) and (3.7). It suffices to show that

$$\mathbf{U_T} \stackrel{d}{=} \mathbf{U}.$$

Since

$$\mathbf{U}_{\mathbf{T}}|_{\max(\mathbf{U}_{\mathbf{T}})} \stackrel{d}{=} \mathbf{T}|_{\max(\mathbf{T})} \stackrel{d}{=} \mathbf{U}|_{\max(\mathbf{U})},$$

it suffices to show that

$$\max(\mathbf{U}_{\mathbf{T}}) \stackrel{d}{=} \max(\mathbf{U}).$$

By definition

$$P(\max(\mathbf{U}_{\mathbf{T}}) \le s) \propto \int_0^s P(\max(\mathbf{T}) \le t) e^{-t} dt + e^{-s} P(\max(\mathbf{T}) \le s)$$

Plug in

$$P(\max(\mathbf{T}) \le s) \propto e^s P(\max(\mathbf{U}) \le s) - \int_0^s P(\max(\mathbf{U}) \le t) e^t dt$$

from (3.8), we have

$$\begin{split} P(\max(\mathbf{U_T}) \leq s) &\propto e^{-s} \left(e^s P(\max(\mathbf{U}) \leq s) - \int_0^s P(\max(\mathbf{U}) \leq t) e^t dt \right) \\ &+ \int_0^s \left(e^t P(\max(\mathbf{U}) \leq t) - \int_0^t P(\max(\mathbf{U}) \leq u) e^u du \right) e^{-t} dt \\ &= \underbrace{P(\max(\mathbf{U}) \leq s)}_{\text{Term I}} - \underbrace{e^{-s} \int_0^s P(\max(\mathbf{U}) \leq t) e^t dt}_{\text{Term II}} + \underbrace{\int_0^s P(\max(\mathbf{U}) \leq t) dt}_{\text{Term III}} \\ &- \underbrace{\int_0^s \int_0^t P(\max(\mathbf{U}) \leq u) e^{u-t} du dt}_{\text{Term IV}}. \end{split}$$

Term IV =
$$\int_0^s \int_0^t P(\max(\mathbf{U}) \le u)e^{u-t} du dt$$

= $\int_0^s \int_u^s P(\max(\mathbf{U}) \le u)e^{u-t} dt du$
= $\int_0^s P(\max(\mathbf{U}) \le u)e^u(e^{-u} - e^{-s}) du$
= $\int_0^s P(\max(\mathbf{U}) \le u) du - e^{-s} \int_0^s P(\max(\mathbf{U}) \le u)e^u du$
= Term III – Term II.

Hence

$$P(\max(\mathbf{U}_{\mathbf{T}}) \le s) \propto \text{Term I} = P(\max(\mathbf{U}) \le s),$$

and

 $\mathbf{U_T} \stackrel{d}{=} \mathbf{U}.$

Proof of Corollary 3.6. Since (3.9) is a direct result during the proof of Proposition 3.5, it suffices to prove (3.10). Furthermore, given (3.6), it suffices to prove

$$\max(\mathbf{U}) \mid \{\max(\mathbf{U}) \ge r - E\} \xrightarrow{d} \max(\mathbf{T}), \quad r \to \infty.$$
(A.2)

For any x > 0 and r > x,

$$P\left(\max(\mathbf{U}) \leq x | \max(\mathbf{U}) \geq r - E\right)$$

$$= \frac{P\left(r - E < \max(\mathbf{U}) \leq x\right)}{P\left(\max(\mathbf{U}) \geq r - E\right)}$$

$$= \frac{\int_{r-x}^{\infty} P\left(r - t < \max(\mathbf{U}) \leq x\right) e^{-t} dt}{\int_{0}^{\infty} P\left(\max(\mathbf{U}) \geq r - t\right) e^{-t} dt}$$

$$= \frac{\int_{r-x}^{\infty} P\left(\max(\mathbf{U}) \leq x\right) e^{-t} dt - \int_{r-x}^{\infty} P\left(\max(\mathbf{U}) \leq r - t\right) e^{-t} dt}{\int_{0}^{\infty} P\left(\max(\mathbf{U}) \geq r - t\right) e^{-t} dt}$$

$$\stackrel{u=r-t}{=} \frac{P\left(\max(\mathbf{U}) \leq x\right) e^{-r+x} - \int_{-\infty}^{x} P\left(\max(\mathbf{U}) \leq u\right) e^{u-r} du}{\int_{-\infty}^{r} P\left(\max(\mathbf{U}) \geq u\right) e^{u} du}$$

$$= \frac{P\left(\max(\mathbf{U}) \leq x\right) e^{x} - \int_{0}^{x} P\left(\max(\mathbf{U}) \leq u\right) e^{u} du}{\int_{-\infty}^{\infty} P\left(\max(\mathbf{U}) \geq u\right) e^{u} du}$$

$$\stackrel{r \to \infty}{\to} \frac{P\left(\max(\mathbf{U}) \leq x\right) e^{x} - \int_{0}^{x} P\left(\max(\mathbf{U}) \leq u\right) e^{u} du}{\int_{-\infty}^{\infty} P\left(\max(\mathbf{U}) \geq u\right) e^{u} du}$$

$$= \frac{P\left(\max(\mathbf{U}) \leq x\right) e^{x} - \int_{0}^{x} P\left(\max(\mathbf{U}) \leq u\right) e^{u} du}{\int_{-\infty}^{\infty} P\left(\max(\mathbf{U}) \geq u\right) e^{u} du}$$

$$= \frac{P\left(\max(\mathbf{U}) \leq x\right) e^{x} - \int_{0}^{x} P\left(\max(\mathbf{U}) \leq u\right) e^{u} du}{I_{-\infty}^{\infty} P\left(\max(\mathbf{U}) \geq e^{u}\right) de^{u}}$$

Therefore (A.2) is proved.

Proof of Corollary 3.7. The result follows by taking the derivatives of both sides of (3.7) or (3.8) with respect to s.

Proof of Proposition 3.9. The convergence to diagonal multivariate generalized Pareto distribution \mathbf{Z}^* is trivial by looking at the conditional distribution of $\mathbf{X} - r \cdot \mathbf{1}$ given $\bar{X} > r$. It remains to show the convergence to the standardized multivariate generalized Pareto distribution \mathbf{Z} .

Denote the **U**-generator $\mathbf{U} := \mathbf{V} - \boldsymbol{\nu}_{\mathbf{V}}$. We have

$$\mathbf{X} - r \cdot \mathbf{1} \mid \max(\mathbf{X}) \ge r$$

= $(E \cdot \mathbf{1} + \mathbf{U} - r \cdot \mathbf{1}) \mid E + \max(\mathbf{U}) \ge r$
= $(E + \max(\mathbf{U}) - r) \cdot \mathbf{1} + (\mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1}) \mid E + \max(\mathbf{U}) \ge r.$

Given s > 0 and any Borel set $B \in {\mathbf{v} | \max(\mathbf{v}) = 0}$, observe that

$$P(E + \max(\mathbf{U}) - r \ge s, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r)$$

$$= \underbrace{P(E + \max(\mathbf{U}) - r \ge s, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r, \max(\mathbf{U}) \le r)}_{I}$$

$$\cdot \underbrace{P(\max(\mathbf{U}) < r | E + \max(\mathbf{U}) \ge r)}_{1 - II}$$

$$+ \underbrace{P(E + \max(\mathbf{U}) - r \ge s, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r, \max(\mathbf{U}) > r)}_{\le 1}$$

$$\cdot \underbrace{P(\max(\mathbf{U}) \ge r | E + \max(\mathbf{U}) \ge r)}_{II}.$$

First consider part II:

$$II = P(\max(\mathbf{U}) \ge r | E + \max(\mathbf{U}) \ge r)$$

$$= \frac{P(\max(\mathbf{U}) \ge r)}{P(E + \max(\mathbf{U}) \ge r)}$$

$$= \frac{P(\max(\mathbf{U}) \ge r)}{\int_0^r P(\max(\mathbf{U}) \ge r - t) e^{-t} dt + \int_r^\infty 1 \cdot e^{-t} dt}$$

$$\stackrel{u=t-s}{=} \frac{P(\max(\mathbf{U}) \ge r)}{\int_0^r P(\max(\mathbf{U}) \ge u) e^{u-r} du + e^{-r}}$$

$$= \frac{e^r P(\max(\mathbf{U}) \ge r)}{\int_0^r P(\max(\mathbf{U}) \ge u) e^u du + 1}.$$

Note that

$$E[e^{\max(\mathbf{U})}] = \int_0^\infty P\left(e^{\max(\mathbf{U})} \ge t\right) dt \stackrel{u=\log(t)}{=} \int_0^\infty P\left(\max(\mathbf{U}) \ge u\right) e^u du.$$

Therefore as $r \to \infty$, the numerator of II

$$P\left(\max(\mathbf{U}) \ge r\right) e^r \to 0,$$

and the denominator

$$\int_0^r P\left(\max(\mathbf{U}) \ge u\right) e^u du + 1 \to E[e^{\max(\mathbf{U})}] + 1 < \infty.$$

Therefore

II =
$$P(\max(\mathbf{U}) \ge r | E + \max(\mathbf{U}) \ge r) \to 0, \quad r \to \infty.$$

Now consider part I:

$$\begin{split} \mathbf{I} &= P\left(E + \max(\mathbf{U}) - r \geq s, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r \right) \\ &= \frac{P\left(E + \max(\mathbf{U}) - r \geq s, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B, \max(\mathbf{U}) \leq r \right)}{P\left(E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r \right)} \\ &= \frac{\int_{s}^{\infty} P\left(s - t \leq \max(\mathbf{U}) - r \leq 0, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B\right) e^{-t} dt}{P\left(E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r \right)} \\ u = t - s \quad \frac{e^{-s} \int_{0}^{\infty} P\left(-u \leq \max(\mathbf{U}) - r \leq 0, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B\right) e^{-u} du}{P\left(E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r \right)} \\ &= e^{-s} \cdot \frac{P\left(E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B\right)}{P\left(E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r\right)} \\ &= e^{-s} \cdot P\left(\mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \geq r, \max(\mathbf{U}) \leq r \right). \end{split}$$

From Corollary 3.6, as $r \to \infty$,

$$\mathbf{U} - \max(\mathbf{U})|_{E+\max(\mathbf{U}) \ge r} \xrightarrow{d} \mathbf{T} - \max(\mathbf{T}) = \mathbf{S}.$$

Hence

$$P (\mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r)$$

$$= P (\mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r, \max(\mathbf{U}) \le r) \cdot \underbrace{P(\max(\mathbf{U}) < r | E + \max(\mathbf{U}) \ge r)}_{1 - \Pi \to 1}$$

$$+ \underbrace{P (\mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) > r, \max(\mathbf{U}) \le r)}_{\leq 1}$$

$$\cdot \underbrace{P(\max(\mathbf{U}) > \geq r | E + \max(\mathbf{U}) \ge r)}_{\Pi \to 0}$$

$$\rightarrow P(\mathbf{S} \in B).$$

Consequently, we have

$$P(\mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r, \max(\mathbf{U}) \le r) \rightarrow P(\mathbf{S} \in B)$$

and

$$P(E + \max(\mathbf{U}) - r \ge s, \mathbf{U} - \max(\mathbf{U}) \cdot \mathbf{1} \in B | E + \max(\mathbf{U}) \ge r) \rightarrow e^{-s} \cdot P(\mathbf{S} \in B).$$

This shows that

$$\mathbf{X} - r \cdot \mathbf{1} | \max(\mathbf{X}) \ge r \xrightarrow{d} \mathbf{Z} \stackrel{d}{=} E \cdot \mathbf{1} + \mathbf{S}$$

where \mathbf{S} is the spectral random vector associated with \mathbf{V} .

Proof of Proposition 4.1. In the case where Σ is of rank (d-1) and the standardized multivariate generalized Pareto distribution **Z** admits a density, the proof directly follows from Corollary 3.7 of Hentschel et al. (2024).

Assume that Γ and Σ are of rank lower than (d-1). Define

$$\Gamma_m := \Gamma + \frac{2}{m} (\mathbf{1}\mathbf{1}^T - I), \quad m = 1, 2, \dots$$

Then each Γ_m is a rank (d-1)-variogram and $\Gamma_m \to \Gamma$ as $m \to \infty$.

Let $\{G_m, \mathbf{Z}_m, \mathbf{Z}_m^*, \mathbf{V}_m\}$ be the generalized extreme value distribution, standardized multivariate generalized Pareto distribution, diagonal multivariate generalized Pareto distribution and profile random vector of the Hüsler-Reiss model parametrized by Γ_m , respectively. Let $\{G, \mathbf{Z}, \mathbf{Z}^*, \mathbf{V}\}$ that of the Hüsler-Reiss model parametrized by Γ . For any $\mathbf{x} \in (0, \infty)^d$,

$$G_m(\mathbf{x}) \to G(\mathbf{x}).$$

 $\mathbf{Z}_m \stackrel{d}{\to} \mathbf{Z},$

 $\mathbf{Z}_m^* \stackrel{d}{
ightarrow} \mathbf{Z}^*$

 $\mathbf{V}_m \stackrel{d}{\rightarrow} \mathbf{V}.$

From (2.3) this implies that

hence

and

We have

$$\mathbf{V}_m = N(\mathbf{0}, \Sigma_m) \stackrel{d}{\to} N(\mathbf{0}, \Sigma)$$

Therefore

$$\mathbf{V} \sim N(\mathbf{0}, \Sigma).$$

References

Coles, S. (2001). An introduction to statistical modeling of extreme values, Volume 208. Springer.

Cooley, D. and E. Thibaud (2019). Decompositions of dependence for high-dimensional extremes. Biometrika 106(3), 587–604.

de Haan, L. and A. Ferreira (2006). Extreme value theory: an introduction. Springer.

- Drees, H. and A. Sabourin (2021). Principal component analysis for multivariate extremes. *Electronic Journal of Statistics* 15, 908–943.
- Engelke, S. and A. S. Hitz (2020). Graphical models for extremes. Journal of the Royal Statistical Society Series B: Statistical Methodology 82(4), 871–932.
- Hentschel, M., S. Engelke, and J. Segers (2024). Statistical inference for hüsler–reiss graphical models through matrix completions. *Journal of the American Statistical Association*, 1–13.
- Hüsler, J. and R.-D. Reiss (1989). Maxima of normal random vectors: between independence and complete dependence. *Statistics & Probability Letters* 7, 283–286.
- Kiriliouk, A., H. Rootzén, J. Segers, and J. L. Wadsworth (2019). Peaks over thresholds modeling with multivariate generalized pareto distributions. *Technometrics* 61(1), 123–135.
- Maulik, K. and S. Resnick (2004). Characterizations and examples of hidden regular variation. Extremes $\gamma(1)$, 31–67.
- Mourahib, A., A. Kiriliouk, and J. Segers (2024). Multivariate generalized pareto distributions along extreme directions. *Extremes*, 1–34.
- Resnick, S. I. (2007). *Heavy-tail phenomena: probabilistic and statistical modeling*, Volume 10. Springer Science & Business Media.
- Rootzén, H., J. Segers, and J. L. Wadsworth (2018). Multivariate generalized pareto distributions: Parametrizations, representations, and properties. *Journal of Multivariate Analysis 165*, 117– 131.
- Rootzén, H. and N. Tajvidi (2006). Multivariate generalized pareto distributions. *Bernoulli* 12(5), 917–930.
- Sibuya, M. (1960). Bivariate extreme statistics. Annals of the Institute of Statistical Mathematics 11(2), 195–210.
- Wan, P. and C. Zhou (2023). Graphical lasso for extremes. arXiv preprint arXiv:2307.15004.