

Extremal spectral radius and g -good r -component connectivity *

Wenxiu Ding, Dan Li[†], Yu Wang

College of Mathematics and System Science, Xinjiang University, Urumqi 830046, China

Abstract: For $F \subseteq V(G)$, if $G - F$ is a disconnected graph with at least r components and each vertex $v \in V(G) \setminus F$ has at least g neighbors, then F is called a g -good r -component cut of G . The g -good r -component connectivity of G , denoted by $c\kappa_{g,r}(G)$, is the minimum cardinality of g -good r -component cuts of G . Let $\mathcal{G}_n^{k,\delta}$ be the set of graphs of order n with minimum degree δ and g -good r -component connectivity $c\kappa_{g,r}(G) = k$. In the paper, we determine the extremal graphs attaining the maximum spectral radii among all graphs in $\mathcal{G}_n^{k,\delta}$. A subset $F \subseteq V(G)$ is called a g -good neighbor cut of G if $G - F$ is disconnected and each vertex $v \in V(G) \setminus F$ has at least g neighbors. The g -good neighbor connectivity $\kappa_g(G)$ of a graph G is the minimum cardinality of g -good neighbor cuts of G . The condition of g -good neighbor connectivity is weaker than that of g -good r -component connectivity, and there is no requirement on the number of components. As a counterpart, we also study similar problem for g -good neighbor connectivity.

Keywords: Spectral radius; g -good r -component connectivity; g -good neighbor connectivity

AMS Classification: 05C50; 05C35

1 Introduction

In the paper, we only consider simple connected graphs. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G)$ of order $n = |V(G)|$ and edge set $E(G)$. The adjacency matrix $A(G)$ of G is an $n \times n$ matrix whose (i, j) entry is 1 if v_i is adjacent to v_j in G , and 0 otherwise. It is obvious that $A(G)$ is a real symmetric matrix. Thus its eigenvalues are real numbers. The largest eigenvalue of $A(G)$ is called the spectral radius of G , denoted by $\rho(G)$. Let $N_G(v)$ be the set of neighbors of v in G and $N_G[v] = N_G(v) \cup \{v\}$. The degree of the vertex v is $d_G(v) = |N_G(v)|$. A graph G is called t -regular if every vertex has the same degree equal to t . Denote by $\delta(G)$ (or for short δ) and $\Delta(G)$ the minimum and maximum degrees of the vertices of G . We know $\delta(G) \leq \rho(G) \leq \Delta(G)$, and the equality holds in either of these inequalities if and only if G is regular.

Connectivity is a fundamental concept in graph theory. For a set $F \subseteq V(G)$ (or $F \subseteq E(G)$), the notation $G - F$ represents the graph obtained by removing the vertices (or edges) in F from G . If $G - F$ is disconnected, then F is called a vertex (or edge) cut. The classic connectivity (connectivity for short) of G , denoted by $\kappa(G)$, is defined as the cardinality $|F|$ of the smallest set $F \subseteq V(G)$ such that $G - F$ is either disconnected or a graph with a single vertex. The edge connectivity of a graph G , denoted by $\lambda(G)$, is the minimal number of edges whose removal produces a disconnected graph. The study of connectivity has been shown to be very important to graphs and has many applications in measuring reliability and fault-tolerance networks, such as [3, 14]. In general, the larger $\kappa(G)$ or $\lambda(G)$ is, the more reliable the network is. It is well

*Supported by NSFC (Nos. 12361071 and 11901498).

[†]Corresponding author. E-mail: ldjedu@163.com.

known that $\kappa(G) \leq \lambda(G) \leq \delta(G)$. A graph G is called maximally edge connected or λ -optimal if $\lambda(G) = \delta(G)$ and maximally vertex connected if $\kappa(G) = \delta(G)$. However, the classic connectivity has certain limitations and underestimates the resilience of the network. To measure the fault tolerance of an interconnection network more accurately, Harary [8] introduced the concept of the conditional connectivity by placing some requirements on the components of $G - F$, where F is a subset of edges or vertices. In 1984, Chartrand, Kapoor, Lesniak and Lick [4] introduced the r -component connectivity, as a generalization of the classic connectivity. The r -component connectivity $c\kappa_r(G)$ of a non-complete graph G is the minimum number of vertices whose deletion results in a disconnected graph with at least r components or a graph with fewer than r vertices. Clearly, $c\kappa_2(G) = \kappa(G)$. In addition, Latifi, Hedge and Naraghi-Pour [9] introduced the concept of g -good neighbor connectivity. A subset $F \subseteq V(G)$ is called a g -good neighbor cut of G if $G - F$ is disconnected and each vertex $v \in V(G) \setminus F$ has at least g neighbors. The g -good neighbor connectivity of G , denoted by $\kappa_g(G)$, is the cardinality of a minimum g -good neighbor cut of G . However, the above mentioned conditional connectivity is only a condition that restricts the components of $G - F$. In some cases, we need to have various restrictions on the components of $G - F$. Recently, Zhu, Zhang, Zou and Ye [21] gave the concept of the g -good r -component connectivity. For $F \subseteq V(G)$, if $G - F$ is disconnected and there are at least r components and each vertex $v \in V(G) - F$ has at least g neighbors, then F is called a g -good r -component cut of G . The g -good r -component connectivity of G , denoted by $c\kappa_{g,r}(G)$, is the minimum cardinality of g -good r -component cuts of G . As a significant tool, g -good r -component connectivity has improved the accuracy of reliability and fault tolerance analysis of networks.

Brualdi and Solheid [2] proposed the following general problem, which became one of the classical problems of spectral graph theory.

Problem 1. [2] Given a set \mathcal{G} of graphs, find $\min\{\rho(G) : G \in \mathcal{G}\}$ and $\max\{\rho(G) : G \in \mathcal{G}\}$, and characterize the corresponding extremal graphs.

A lot of results concerning the Brualdi-Solheid problem were presented, and some of these results were exhibited in a recent monograph on the spectral radius by Stevanović [16]. This paper focuses on the research connectivity from spectral perspectives. The spectral conditions for the connectivity of graphs have been well investigated. Berman and Zhang [1] studied the spectral radius of graphs with n vertices and k cut vertices and described the graph that has the maximal spectral radius in this class. The union $G_1 \cup G_2$ is defined to be $G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$. The join $G_1 \vee G_2$ is obtained from $G_1 \cup G_2$ by adding all the edges joining a vertex of G_1 to a vertex of G_2 . Denote by K_n the complete graph with order n . $K_k \vee (K_1 \cup K_{n-k-1})$ is shown to be the graph with the maximal spectral radius among all graphs of order n with connectivity $\kappa(G) \leq k$ in [10, 19]. Lu and Lin [13] proved that $K_k \vee (K_{\delta-k+1} \cup K_{n-\delta-1})$ is the graph with the maximum spectral radius among all graphs of order n with $\kappa(G) \leq k \leq \delta(G)$. Recently, Fan, Gu and Lin [7] extended some results on classic connectivity. They determined the graphs with maximal spectral radius among all graphs of order n with given minimum degree δ and r -component connectivity. On the other hand, the spectral conditions for the connectivity of digraphs have received a lot of attention of researchers, see [7, 11]. Moreover, the spectral conditions for the edge-connectivity of graphs have also been well investigated by many researchers, see [6, 15, 17].

Since the g -good r -component connectivity is consider as a generalization of the connectivity, based on Problem 1, motivated by the above spectral results, we are interested in studying the spectral condition for g -good r -component connectivity.

Problem 2. Which graphs attain the maximum spectral radii among all connected graphs of order n with fixed minimum degree δ and g -good r -component connectivity $c\kappa_{g,r}(G)$?

For convenience, we use k instead of $c\kappa_{g,r}(G)$. Let $\mathcal{G}_n^{k,\delta}$ be the set of graphs of order n with minimum degree δ and g -good r -component connectivity k . Note that there is no bound between k and δ . Let $G_{n,(g+1)^{r-1}}^{\delta,0}$ be the graph obtained from $K_1 \cup (K_{k-1} \vee (K_{n-(r-1)(g+1)-k} \cup (r-1)K_{g+1}))$ by

adding δ edges between the isolated vertex K_1 and K_{k-1} . Let $G_{n,(g+1)^{r-1}}^{\delta-g,g}$ be the graph obtained from $K_1 \cup (K_k \vee (K_{n-(r-1)(g+1)-k} \cup (r-2)K_{g+1} \cup K_g))$ by adding $\delta - g$ edges between the isolated vertex K_1 and K_k , and adding g edges between K_1 and K_g . Let $G_{n,(g+1)^{r-1}}^{k-1,\delta-k+1}$ be the graph obtained from $K_1 \cup (K_{k-1} \vee (K_{n-(r-1)(g+1)-k} \cup (r-1)K_{g+1}))$ by adding $k-1$ edges between the isolated vertex K_1 and K_{k-1} , and adding $\delta - k + 1$ edges between K_1 and $K_{n-(r-1)(g+1)-k}$. Let $G_{n,(g+1)^{r-1}}^{0,\delta}$ be the graph obtained from $K_1 \cup (K_{n-(r-1)(g+1)-1} \cup (r-1)K_{g+1})$ by adding $\delta - r + 1$ edges between the isolated vertex K_1 and $K_{n-(r-1)(g+1)-1}$, and adding one edge between K_1 and each K_{g+1} , respectively (see Figure 1). In this paper, we mainly give the answer to Problem 2. Then we have the following result.




Figure 1: $G_{n,(g+1)^{r-1}}^{\delta,0}$, $G_{n,(g+1)^{r-1}}^{\delta-g,g}$, $G_{n,(g+1)^{r-1}}^{k-1,\delta-k+1}$ and $G_{n,(g+1)^{r-1}}^{0,\delta}$

Theorem 1. Let $G \in \mathcal{G}_n^{k,\delta}$, where $n \geq k + r(g+1)$. Then we have the following statements.

- (I) If $k > \delta$ and $\delta < g$, then $\rho(G) \leq \rho(G_{n,(g+1)^{r-1}}^{\delta,0})$ with equality if and only if $G \cong G_{n,(g+1)^{r-1}}^{\delta,0}$;
- (II) If $k > \delta \geq g$, then $\rho(G) \leq \rho(G_{n,(g+1)^{r-1}}^{\delta-g,g})$ with equality if and only if $G \cong G_{n,(g+1)^{r-1}}^{\delta-g,g}$;
- (III) If $2 \leq k \leq \delta < g$, then $\rho(G) \leq \rho(G_{n,(g+1)^{r-1}}^{k-1,\delta-k+1})$ with equality if and only if $G \cong G_{n,(g+1)^{r-1}}^{k-1,\delta-k+1}$;
- (IV) If $1 = k \leq \delta < g$, then $\rho(G) \leq \rho(G_{n,(g+1)^{r-1}}^{0,\delta})$ with equality if and only if $G \cong G_{n,(g+1)^{r-1}}^{0,\delta}$;
- (V) If $k \leq \delta$ and $g \leq \delta < g+k$, then $\rho(G) \leq \rho(G_{n,(g+1)^{r-1}}^{\delta-g,g})$ with equality if and only if $G \cong G_{n,(g+1)^{r-1}}^{\delta-g,g}$;
- (VI) If $\delta \geq g+k$, then $\rho(G) \leq \rho(K_k \vee (K_{n-k-(\delta-k+1)(r-1)} \cup (r-1)K_{\delta-k+1}))$ with equality if and only if $G \cong K_k \vee (K_{n-k-(\delta-k+1)(r-1)} \cup (r-1)K_{\delta-k+1})$.

The rest of the paper is structured as follows. In Section 2, we determine the graphs attaining the maximum spectral radii among all connected graphs with given minimum degree and g -good

r -component connectivity. In Section 3, we extend the result of extremal spectral radius and characterize the graphs with maximal spectral radii among all connected graphs with given minimum degree and g -good neighbor connectivity.

2 Proof of Theorem 1

In this section, we present the proof of Theorem 1. First of all, we list several lemmas that will be used in the follows. Note that for a connected graph G , there is a positive unit eigenvector \mathbf{x} of $A(G)$ corresponding to $\rho(G)$. Such a unit eigenvector is also called Perron vector of $A(G)$. The component of \mathbf{x} corresponding to a vertex $v \in V(G)$ is usually written as $x(v)$.

Lemma 1. [15] *Let u, v be two distinct vertices of a connected graph G , and let \mathbf{x} be the Perron vector of $A(G)$.*

- (1) *If $N_G(v) \setminus \{u\} \subset N_G(u) \setminus \{v\}$, then $x(u) > x(v)$;*
- (2) *If $N_G(v) \subseteq N_G[u]$ and $N_G(u) \subseteq N_G[v]$, then $x(u) = x(v)$.*

Lemma 2. [15] *Let G be a connected graph and G' be a proper subgraph of G . Then $\rho(G') < \rho(G)$.*

The following lemma, due to Wu, Xiao and Hong [18], shows an edge transformation which increases the spectral radius.

Lemma 3. [18] *Let G be a connected graph and $\rho(G)$ be the spectral radius of $A(G)$. Let u, v be two vertices of G . Suppose that $v_1, v_2, \dots, v_s \in N_G(v) \setminus N_G(u)$ with $1 \leq s \leq d_G(v)$, and G^* is the graph obtained from G by deleting the edges vv_i and adding the edges uv_i for $1 \leq i \leq s$. Let \mathbf{x} be the Perron vector of $A(G)$. If $x(u) \geq x(v)$, then $\rho(G) < \rho(G^*)$.*

Lemma 4. [5] *Let $n = \sum_{i=1}^t n_i + s$. If $n_1 \geq n_2 \geq \dots \geq n_t \geq p$ and $n_1 < n - s - p(t - 1)$, then*

$$\rho(K_s \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_t})) < \rho(K_s \vee (K_{n-s-p(t-1)} \cup (t-1)K_p)).$$

For any vertex $v \in V(G)$ and any subset $F \subseteq V(G)$, let $N_F(v) = N_G(v) \cap F$ and $d_F(v) = |N_G(v) \cap F|$. Now we shall give a proof of Theorem 1.

Proof of Theorem 1. Suppose that G is a connected graph attaining the maximum spectral radius among all connected graph with minimum degree δ and g -good r -component connectivity $c\kappa_{g,r}(G) = k$. By the definition of g -good r -component connectivity, there exists some nonempty subset $F \subseteq V(G)$ with $|F| = k$ such that $G - F$ is disconnected and there are at least r components and each vertex $v \in V(G) \setminus F$ has at least g neighbors. Let B_1, B_2, \dots, B_q be the components of $G - F$, where $q \geq r$, and let $|V(B_i)| = n_i$ for $1 \leq i \leq q$. It is easy to see that $n_i \geq g + 1$ for $1 \leq i \leq q$. In fact, we can deduce that $q = r$ by Lemma 2. Now, we divide the proof into the following two cases.

Case 1. $k > \delta$. Choose a vertex $u \in V(G)$ such that $d_G(u) = \delta$. Suppose that $d_F(u) = t$. Next, we will discuss the case in two subcases.

Subcase 1.1. $\delta < g$.

By the definition of g -good r -component connectivity, we have $d_G(v) \geq g$ for $v \in V(G) \setminus F$. So $u \in F$. We first assert that $t = \delta$. Otherwise, $t \leq \delta - 1$. By the maximality of $\rho(G)$, we can obtain that $G - u \cong K_{k-1} \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r})$. Notice that $d_F(u) = t$ and $|N_G(u) \cap (V(G) \setminus F)| \geq 1$. Let $N_F(u) = \{w_1, w_2, \dots, w_t\}$ and $F \setminus N_F[u] = \{w_{t+1}, w_{t+2}, \dots, w_{k-1}\}$, and let $P_i = N_G(u) \cap V(K_{n_i})$ and $|P_i| = p_i$ for $1 \leq i \leq r$. Then $\sum_{i=1}^r p_i = \delta - t$. Let \mathbf{x} be the Perron vector of $A(G)$. By symmetry,

we may say that $x(v) = x_i$ for $v \in P_i$ and $x(v) = x'_i$ for $v \in V(K_{n_i}) \setminus P_i$, where $1 \leq i \leq r$. Without loss of generality, assume that $x_1 = \max\{x_i \mid 1 \leq i \leq r\}$. Note that $x(w_1) = x(w_i)$ for $2 \leq i \leq t$ and $x(w_{t+1}) = x(w_i)$ for $t+2 \leq i \leq k-1$. Therefore, by $A(G)\mathbf{x} = \rho(G)\mathbf{x}$, we have

$$\rho(G)x(u) = tx(w_1) + \sum_{i=1}^r p_i x_i, \quad (1)$$

$$\rho(G)x_i = tx(w_1) + (k-t-1)x(w_{t+1}) + (p_i-1)x_i + (n_i-p_i)x'_i + x(u), \quad (2)$$

$$\rho(G)x'_i = tx(w_1) + (k-t-1)x(w_{t+1}) + p_i x_i + (n_i-p_i-1)x'_i, \quad (3)$$

$$\rho(G)x(w_{t+1}) = tx(w_1) + (k-t-2)x(w_{t+1}) + \sum_{i=1}^r p_i x_i + \sum_{i=1}^r (n_i-p_i)x'_i, \quad (4)$$

where $1 \leq i \leq r$. From (1)-(3), we get

$$\begin{aligned} & \rho(G) \left(\sum_{i=2}^r p_i x_i + \sum_{i=2}^r (n_i-p_i)x'_i - x(u) \right) \\ &= \left(\sum_{i=2}^r n_i - 1 \right) tx(w_1) + \sum_{i=2}^r n_i(k-t-1)x(w_{t+1}) + \sum_{i=2}^r (n_i-1)p_i x_i + \sum_{i=2}^r (n_i-1)(n_i-p_i)x'_i \\ & \quad + \sum_{i=2}^r p_i x(u) - \sum_{i=1}^r p_i x_i \\ &> \sum_{i=2}^r n_i(k-t-1)x(w_{t+1}) - \sum_{i=1}^r p_i x_i \quad (\text{since } n_i \geq g+1, g > \delta, t \geq 0 \text{ and } 0 \leq p_i < n_i \text{ for } 2 \leq i \leq r) \\ &\geq (r-1)(g+1)(k-t-1)x(w_{t+1}) - (\delta-t)x_1 \\ & \quad (\text{since } n_i \geq g+1, \sum_{j=1}^r p_j = \delta-t \text{ and } x_1 \geq x_i \text{ for } 2 \leq i \leq r) \\ &> (\delta-t)(x(w_{t+1}) - x_1) \quad (\text{since } r \geq 2, g > \delta \text{ and } k \geq \delta+1). \end{aligned}$$

Combining this with (2) and (4), we have

$$(\rho(G)+1)(x(w_{t+1}) - x_1) = \sum_{i=2}^r p_i x_i + \sum_{i=2}^r (n_i-p_i)x'_i - x(u) > \frac{(\delta-t)(x(w_{t+1}) - x_1)}{\rho(G)},$$

from which we obtain

$$\frac{(\rho^2(G) + \rho(G) - \delta + t)(x(w_{t+1}) - x_1)}{\rho(G)} > 0.$$

Note that G contains K_{k-1+n_i} as proper subgraphs, where $1 \leq i \leq r$. Thus, $\rho(G) > \rho(K_{k-1+n_i}) = k+n_i-2 > \delta$ due to $n_i \geq g+1 > \delta+1$ and $k \geq \delta+1$. Combining this with $t \geq 0$, we can deduce that $\rho^2(G) + \rho(G) - \delta + t > 0$. This demonstrates that $x(w_{t+1}) > x_1$. Let $G' = G - \{uv \mid v \in P_i, 1 \leq i \leq r\} + \{uw_j \mid t+1 \leq j \leq \delta\}$. Obviously, $G' \in \mathcal{G}_n^{k,\delta}$. According to Lemma 3, we have

$$\rho(G') > \rho(G),$$

which contradicts the maximality of $\rho(G)$. This implies that $t = \delta$. In what follows, we shall prove that $G \cong G_{n,(g+1)^{r-1}}^{\delta,0}$. Without loss of generality, suppose that $n_1 \geq n_2 \geq \dots \geq n_r$. By the maximality of $\rho(G)$ and $N_G[u] \subseteq F$, we have $G-u \cong K_{k-1} \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r})$, where $\sum_{i=1}^r n_i = n-k$ and $n_r \geq g+1$. Denote $V(K_{n_i}) = \{v_i^1, v_i^2, \dots, v_i^{n_i}\}$. If $n_i = g+1$ for all $2 \leq i \leq r$, then the result follows. If there exists $n_j \geq g+2$ for some $2 \leq j \leq r$. Let $x(v_1^l) = x_1$ for $1 \leq l \leq n_1$ and $x(v_j^i) = x_j$ for $2 \leq j \leq r$ and $1 \leq i \leq n_j$, then

$$(\rho(G) - n_j + 1)(x_1 - x_j) = (n_1 - n_j)x_1 \geq 0$$

due to $n_1 \geq n_j$. Since G contains K_{k-1+n_j} as a proper subgraph and $k \geq \delta + 1 \geq 2$, it follows that $\rho(G) > \rho(K_{k-1+n_j}) = k + n_j - 2 \geq n_j$, and hence $x_1 \geq x_j$. Let $G'' = G - \{v_j^i v_j^h \mid 1 \leq i \leq g+1, g+2 \leq h \leq n_j\} + \{v_1^l v_j^h \mid 1 \leq l \leq n_1, g+2 \leq h \leq n_j\}$. Then $G'' - F$ contains r components $K_{n-(r-1)(g+1)-k}, K_{g+1}, \dots, K_{g+1}$, $|N_{G''}(v) \cap (V(G'') \setminus F)| \geq g$ for $v \in V(G'') \setminus F$ and $d_{G''}(u) = \delta$. So $G'' \in \mathcal{G}_n^{k, \delta}$. According to Lemma 3, we have $\rho(G'') > \rho(G)$, which also leads to a contradiction. This indicates that $G \cong G_{n, (g+1)^{r-1}}^{\delta, 0}$, as desired.

Subcase 1.2. $\delta \geq g$. Recall that $u \in V(G)$ such that $d_G(u) = \delta$ and $d_F(u) = t$. We will prove the following four claims.

Claim 1. $u \notin F$.

Otherwise, $u \in F$. Using the similar analysis as the proof of subcase 1.1, we can obtain that if $u \in F$, then $t = \delta$, i.e., $N_G(u) \subseteq F$. By the maximality of $\rho(G)$, we can deduce that $G - u \cong K_{k-1} \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r})$, where $\sum_{i=1}^r n_i = n - k$ and $n_i \geq g + 1$. Without loss of generality, we may consider that $n_1 \geq n_2 \geq \dots \geq n_r$. Let $N_F(u) = \{w_1, w_2, \dots, w_\delta\}$, $F \setminus N_F[u] = \{w_{\delta+1}, w_{\delta+2}, \dots, w_{k-1}\}$ and $V(K_{n_i}) = \{v_i^1, v_i^2, \dots, v_i^{n_i}\}$ for $1 \leq i \leq r$. Assume that \mathbf{x} is the Perron vector of $A(G)$. By symmetry, we say $x(v) = x_i$ for $v \in V(K_{n_i})$, where $1 \leq i \leq r$. Observe that $x(w_1) = x(w_i)$ for $2 \leq i \leq \delta$ and $x(w_{\delta+1}) = x(w_i)$ for $\delta + 2 \leq i \leq k - 1$. Since $N_G(v_1^i) \setminus \{w_1\} \subset N_G(w_1) \setminus \{v_1^i\}$ for $1 \leq i \leq n_1$, it follows that $x_1 < x(w_1)$ by Lemma 1. Furthermore, if $n_i \geq n_j$, then we can deduce that $(\rho(G) - n_j + 1)(x_i - x_j) = (n_i - n_j)x_i \geq 0$. Note that G contains K_{k-1+n_j} as a proper subgraph and $k \geq \delta + 1 \geq 2$, then $\rho(G) > \rho(K_{k-1+n_j}) = k + n_j - 2 \geq n_j$, and hence $x_1 \geq x_j$. Moreover,

$$\begin{aligned} \rho(G)x(u) &= \delta x(w_1), \\ \rho(G)x_i &= \delta x(w_1) + (k - \delta - 1)x(w_{\delta+1}) + (n_i - 1)x_i, \\ \rho(G)x(w_1) &= (\delta - 1)x(w_1) + (k - \delta - 1)x(w_{\delta+1}) + x(u) + \sum_{i=1}^r n_i x_i, \end{aligned}$$

where $1 \leq i \leq r$, from which we have

$$\begin{aligned} & (n - k - n_r)x_r + gx(u) - gx(w_1) \\ &= \frac{(\rho(G) + 1)((n - k - n_r)x_r + gx(u) - gx(w_1))}{\rho(G) + 1} \\ &= \frac{(n - k - n_r)\delta x(w_1) + (n - k - n_r - g)(k - \delta - 1)x(w_{\delta+1}) + (n - k - n_r)n_r x_r - g \sum_{i=1}^r n_i x_i}{\rho(G) + 1} \\ &\geq \frac{(n - k - n_r)\delta x(w_1) + (n - k - n_r)n_r x_r - gn_r x_r - g \sum_{i=1}^{r-1} n_i x_i}{\rho(G) + 1} \quad (\text{since } n - k - n_r > g \text{ and } k \geq \delta + 1) \\ &\geq \frac{(n - k - n_r)\delta x(w_1) + (n - k - n_r - g)n_r x_r - g(n - k - n_r)x_1}{\rho(G) + 1} \\ & \quad (\text{since } \sum_{i=1}^{r-1} n_i = n - k - n_r \text{ and } x_1 \geq x_j \text{ for } 2 \leq j \leq r - 1) \\ &= \frac{(n - k - n_r)(\delta x(w_1) - gx_1) + (n - k - n_r - g)n_r x_r}{\rho(G) + 1} \\ &> 0 \quad (\text{since } \delta \geq g, x(w_1) > x_1 \text{ and } n - k - n_r > g) \end{aligned} \tag{5}$$

and

$$x_{r-1} - x(u) = \frac{(k - \delta - 1)x(w_{\delta+1}) + (n_{r-1} - 1)x_{r-1}}{\rho(G)} \geq 0. \tag{6}$$

Suppose that $E_1 = \{v_r^1 v \mid v \in V(G) \setminus (F \cup V(K_{n_r}))\} + \{uv_r^i \mid 2 \leq i \leq g + 1\}$ and $E_2 = \{uw_j \mid \delta - g + 1 \leq j \leq \delta\}$. Let $G_1 = G + E_1 - E_2$ and $F_1 = F - \{u\} + \{v_r^1\}$. Then $|F_1| = k$, $G_1 - F_1$ contains r

components $K_{n_1}, K_{n_2}, \dots, K_{n_{r-1}}, K_{n_r-1} \cup \{u\}$ and $d_{G_1}(u) = \delta$. One can verify that $G_1 \in \mathcal{G}_n^{k, \delta}$. Let \mathbf{y} be the Perron vector of $A(G_1)$. By symmetry, $y(v) = y_i$ for $v \in V(K_{n_i})$, where $1 \leq i \leq r-1$, $y(v_r^i) = y_r$ for $2 \leq i \leq g+1$ and $y(v_r^j) = y_r'$ for $g+2 \leq j \leq n_r$. Notice that $y(w_1) = y(w_i)$ for $2 \leq i \leq \delta - g$ and $y(v_r^1) = y(w_i)$ for $\delta - g + 1 \leq i \leq k-1$. According to Lemma 1, we get $y(u) < y_r$ and $y_r' < y_r$, then

$$\begin{aligned}\rho(G_1)y_i &= (\delta - g)y(w_1) + (k - \delta + g)y(v_r^1) + (n_i - 1)y_i, \\ \rho(G_1)y_r &= (\delta - g)y(w_1) + (k - \delta + g)y(v_r^1) + (g - 1)y_r + (n_r - g - 1)y_r' + y(u) \\ &< (\delta - g)y(w_1) + (k - \delta + g)y(v_r^1) + (n_r - 1)y_r,\end{aligned}$$

where $1 \leq i \leq r-1$, from which we have

$$(\rho(G_1) - n_r + 1)(y_i - y_r) > (n_i - n_r)y_i \geq 0,$$

due to $n_i \geq n_r$. Observe that G_1 contains K_{k+n_r-1} as a proper subgraph, $\rho(G_1) > \rho(K_{k+n_r-1}) = k + n_r - 2 \geq n_r$. Thus $y_i > y_r$ for $1 \leq i \leq r-1$. Similarly, if $n_i \geq n_j$, then $y_i \geq y_j$. Combining these with (5) and (6), we obtain

$$\begin{aligned}& \mathbf{y}^T(\rho(G_1) - \rho(G))\mathbf{x} \\ &= \mathbf{y}^T(A(G_1) - A(G))\mathbf{x} \\ &= \sum_{v_r^1, v \in E_1} (x(v_r^1)y(v) + x(v)y(v_r^1)) + \sum_{uv_r^i \in E_1} (x(u)y(v_r^i) + x(v_r^i)y(u)) - \sum_{uw_j \in E_2} (x(u)y(w_j) + x(w_j)y(u)) \\ &\geq (n - k - n_r)(x_r y_{r-1} + x_{r-1} y(v_r^1)) + g(x(u)y_r + x_r y(u)) - g(x(u)y(v_r^1) + x(w_1)y(u)) \\ &\quad (\text{since } x_i \geq x_{r-1} \text{ and } y_i \geq y_{r-1} \text{ for } 1 \leq i \leq r-2) \\ &> (n - k - n_r)x_r y_{r-1} + gx(u)y_r - gx(w_1)y(u) \quad (\text{since } n - k - n_r > g, x_{r-1} \geq x(u) \text{ and } g \geq 0) \\ &> y(u)((n - k - n_r)x_r + gx(u) - gx(w_1)) \quad (\text{since } y_{r-1} > y_r > y(u)) \\ &> 0 \quad (\text{by (5)}),\end{aligned}$$

and hence $\rho(G_1) > \rho(G)$, which contradicts the maximality of $\rho(G)$. This demonstrates that $u \notin F$, proving Claim 1.

By Claim 1, $u \notin F$, and so $u \in V(B_i)$ for some $1 \leq i \leq r$. Without loss of generality, we may assume that $u \in V(B_r)$. Then $d_{B_r}(u) \geq g$ and $t = d_F(u) = \delta - d_{B_r}(u) \leq \delta - g$. We can get the following claim.

Claim 2. $t = \delta - g$.

If $\delta = g$, then $t = 0$. Clearly, the result holds. Next, we consider $\delta > g$. Suppose on the contrary that $t < \delta - g$, i.e., $d_{B_r}(u) = \delta - t \geq g + 1$. Let $N_F(u) = \{w_1, w_2, \dots, w_t\}$, $F \setminus N_F(u) = \{w_{t+1}, w_{t+2}, \dots, w_k\}$ and $N_{B_r}(u) = \{v_r^1, v_r^2, \dots, v_r^{\delta-t}\}$. Again by the maximality of $\rho(G)$, we can deduce that $G - u \cong K_k \vee (K_{n_1} \cup \dots \cup K_{n_{r-1}} \cup K_{n_r-1})$. Assume that \mathbf{x} is the Perron vector of $A(G)$. By symmetry, we may say $x(v) = x_r$ for $v \in N_{B_r}(u)$, $x(v) = x_r'$ for $v \in V(B_r) \setminus N_{B_r}[u]$ and $x(v) = x_i$ for $v \in V(K_{n_i})$, where $1 \leq i \leq r-1$. Notice that $x(w_1) = x(w_i)$ for $2 \leq i \leq t$ and $x(w_{t+1}) = x(w_i)$ for $t+2 \leq i \leq k$, then

$$\begin{aligned}\rho(G)x(w_{t+1}) &= tx(w_1) + (k - t - 1)x(w_{t+1}) + (\delta - t)x_r + (n_r - 1 - (\delta - t))x_r' + \sum_{i=1}^{r-1} n_i x_i, \\ \rho(G)x_r &= tx(w_1) + (k - t)x(w_{t+1}) + (\delta - t - 1)x_r + (n_r - 1 - (\delta - t))x_r' + x(u),\end{aligned}$$

from which we obtain that

$$(\rho(G) + 1)(x(w_{t+1}) - x_r)$$

$$\begin{aligned}
&= \sum_{i=1}^{r-1} n_i x_i - x(u) \\
&= \frac{1}{\rho(G)} \left(\sum_{i=1}^{r-1} n_i (\rho(G)x_i) - \rho(G)x(u) \right) \\
&= \frac{\sum_{i=1}^{r-1} n_i (tx(w_1) + (k-t)x(w_{t+1}) + (n_i-1)x_i) - (tx(w_1) + (\delta-t)x_r)}{\rho(G)} \\
&= \frac{(\sum_{i=1}^{r-1} n_i - 1)tx(w_1) + \sum_{i=1}^{r-1} n_i(k-t)x(w_{t+1}) + \sum_{i=1}^{r-1} n_i(n_i-1)x_i - (\delta-t)x_r}{\rho(G)} \\
&\geq \frac{(r-1)(g+1)(k-t)x(w_{t+1}) - (\delta-t)x_r}{\rho(G)} \quad (\text{since } n_i \geq g+1 \text{ for } 1 \leq i \leq r-1, t \geq 0 \text{ and } g \geq 0) \\
&> \frac{(\delta-t)(x(w_{t+1}) - x_r)}{\rho(G)} \quad (\text{since } r \geq 2, g \geq 0 \text{ and } k > \delta).
\end{aligned}$$

It follows that $\frac{(\rho^2(G) + \rho(G) - \delta + t)(x(w_{t+1}) - x_r)}{\rho(G)} > 0$. Note that G contains K_k as a proper subgraph. Thus, $\rho(G) > \rho(K_k) = k - 1 \geq \delta$. Combining this with $t \geq 0$, we can deduce that $\rho^2(G) + \rho(G) - \delta + t > 0$. This suggests that $x(w_{t+1}) > x_r$. Let $G_2 = G - \{uv_r^i \mid g+1 \leq i \leq \delta-t\} + \{uw_j \mid t+1 \leq j \leq \delta-g\}$. It is obvious that $G_2 \in \mathcal{G}_n^{k,\delta}$. According to Lemma 3, we have $\rho(G_2) > \rho(G)$, which contradicts the maximality of $\rho(G)$. Therefore, $t = \delta - g$, completing the proof of Claim 2.

Without loss of generality, suppose that $n_1 \geq n_2 \geq \dots \geq n_r$. If $n_1 = n_r$, then $u \in V(B_i)$ for $1 \leq i \leq r$. If $n_1 > n_r$, then we have the following claim.

Claim 3. $u \in V(B_i)$ for $n_i = n_r$.

If $g = 0$, then $t = \delta$. Clearly, $u \in V(B_i)$ for $n_i = n_r = 1$. Therefore, the result holds. Next, we consider $g \geq 1$. Suppose on the contrary that $u \in V(B_j)$ for $n_j > n_r$. Without loss of generality, assume that $u \in V(B_1)$. We obtain $G - u \cong K_k \vee (K_{n_1-1} \cup K_{n_2} \cup \dots \cup K_{n_r})$ by the maximality of $\rho(G)$. Let $N_F(u) = \{w_1, w_2, \dots, w_{\delta-g}\}$ and $F \setminus N_F(u) = \{w_{\delta-g+1}, w_{\delta-g+2}, \dots, w_k\}$. Let \mathbf{x} be the Perron vector of $A(G)$. By symmetry, $x(v) = x_1$ for $v \in N_{B_1}(u)$, $x(v) = x'_1$ for $v \in V(B_1) \setminus N_{B_1}[u]$ and $x(v) = x_i$ for $v \in V(K_{n_i})$, where $2 \leq i \leq r$. Notice that $x(w_1) = x(w_i)$ for $2 \leq i \leq \delta - g$ and $x(w_{\delta-g+1}) = x(w_i)$ for $\delta - g + 2 \leq i \leq k$, then

$$\begin{aligned}
\rho(G)x_r &= (\delta - g)x(w_1) + (k - \delta + g)x(w_{\delta-g+1}) + (n_r - 1)x_r, \\
\rho(G)x'_1 &= (\delta - g)x(w_1) + (k - \delta + g)x(w_{\delta-g+1}) + gx_1 + (n_1 - g - 2)x'_1, \\
\rho(G)x_1 &= (\delta - g)x(w_1) + (k - \delta + g)x(w_{\delta-g+1}) + (g - 1)x_1 + (n_1 - g - 1)x'_1 + x(u),
\end{aligned}$$

from which we have

$$x_1 = x'_1 + \frac{x(u)}{\rho(G) + 1}. \quad (7)$$

Since G contains K_{k+n_1-1} as a proper subgraph and $n_1 > n_r$, it follows that $\rho(G) > \rho(K_{k+n_1-1}) = k + n_1 - 2$ and $\rho(G) - n_r + 1 \geq \rho(G) - n_1 + 2 > 0$. Then we get

$$x'_1 = \frac{\rho(G) - n_r + 1}{\rho(G) - n_1 + 2}x_r + \frac{g(x_1 - x'_1)}{\rho(G) - n_1 + 2} > x_r. \quad (8)$$

Moreover, we have $x_1 > x(u)$ by Lemma 1 and

$$\rho(G)x(u) = (\delta - g)x(w_1) + gx_1,$$

$$\rho(G)x(w_{\delta-g+1}) = (\delta - g)x(w_1) + (k - \delta + g - 1)x(w_{\delta-g+1}) + gx_1 + (n_1 - g - 1)x'_1 + \sum_{i=2}^r n_i x_i,$$

from which we get that

$$\begin{aligned}
x_r - x(u) &= \frac{1}{\rho(G)}(\rho(G)x_r - \rho(G)x(u)) \\
&= \frac{(k - \delta + g)x(w_{\delta-g+1}) + (n_r - 1)x_r - gx_1}{\rho(G)} \\
&> \frac{g(x(w_{\delta-g+1}) + x_r - x_1)}{\rho(G)} \quad (\text{since } k > \delta \text{ and } n_r \geq g + 1) \\
&= \frac{g}{\rho^2(G)}(\rho(G)x(w_{\delta-g+1}) + \rho(G)x_r - \rho(G)x_1) \\
&= \frac{g}{\rho^2(G)}[(\delta - g)x(w_1) + (k - \delta + g - 1)x(w_{\delta-g+1}) + (n_r - 1)x_r + \sum_{i=2}^r n_i x_i + x_1 - x(u)] \\
&> 0 \quad (\text{since } \delta \geq g, k \geq \delta + 1, g \geq 1, n_r \geq g + 1 \text{ and } x_1 > x(u)). \tag{9}
\end{aligned}$$

Denote $V(B_1) = \{v_1^1, v_1^2, \dots, v_1^{n_1-1}, u\}$, $N_{B_1}(u) = \{v_1^1, v_1^2, \dots, v_1^g\}$ and $V(K_{n_i}) = \{v_i^1, v_i^2, \dots, v_i^{n_i}\}$ for $2 \leq i \leq r$. Suppose that $E'_1 = \{uv_r^i \mid 1 \leq i \leq g\} + \{v_r^{n_r} v_1^j \mid 1 \leq j \leq n_1 - 1\}$ and $E'_2 = \{uv_1^i \mid 1 \leq i \leq g\} + \{v_r^{n_r} v_r^j \mid 1 \leq j \leq n_r - 1\}$. Let $G_3 = G + E'_1 - E'_2$, then $V(B'_1) = \{v_1^1, \dots, v_1^{n_1-1}, v_r^{n_r}\}$ and $V(B'_r) = \{v_r^1, \dots, v_r^{n_r-1}, u\}$. So $G_3 - F$ contains r components $K_{n_1}, K_{n_2}, \dots, K_{n_{r-1}}, K_{n_r-1} \cup \{u\}$ and $d_{G_3}(u) = \delta$. Obviously, $G_3 \in \mathcal{G}_n^{k, \delta}$. Combining this with (7)-(9), we have

$$\begin{aligned}
&\rho(G_3) - \rho(G) \\
&\geq \mathbf{x}^T (A(G_3) - A(G)) \mathbf{x} \\
&= 2 \sum_{uv_r^i \in E'_1} x(u)x(v_r^i) + 2 \sum_{v_r^{n_r} v_1^j \in E'_1} x(v_r^{n_r})x(v_1^j) - 2 \sum_{uv_1^i \in E'_2} x(u)x(v_1^i) - 2 \sum_{v_r^{n_r} v_r^j \in E'_2} x(v_r^{n_r})x(v_r^j) \\
&= 2gx(u)x_r + 2gx_r x_1 + 2(n_1 - g - 1)x_r x'_1 - 2gx(u)x_1 - 2(n_r - 1)x_r x_r \\
&\geq 2gx(u)x_r + \frac{2gx_r x(u)}{\rho(G) + 1} + 2n_r x_r x'_1 - 2gx(u)x'_1 - \frac{2gx(u)x(u)}{\rho(G) + 1} - 2(n_r - 1)x_r x_r \quad (\text{since } n_1 \geq n_r + 1) \\
&> 2gx(u)x_r + 2n_r x_r x'_1 - 2gx(u)x'_1 - 2n_r x_r x_r + 2x_r x_r \quad (\text{since } x_r > x(u) \text{ and } g \geq 1) \\
&> 2n_r x_r (x'_1 - x_r) - 2gx(u)(x'_1 - x_r) \\
&= 2(n_r x_r - gx(u))(x'_1 - x_r) \\
&> 0 \quad (\text{since } n_r \geq g + 1, x_r > x(u) \text{ and } x'_1 > x_r).
\end{aligned}$$

Hence $\rho(G_3) > \rho(G)$, which contradicts the maximality of $\rho(G)$. This implies that $u \in V(B_i)$ for $n_i = n_r$, proving Claim 3.

In what follows, we shall prove that $G \cong G_{n, (g+1)^{r-1}}^{\delta-g, g}$. In fact, by the maximality of $\rho(G)$ and Claims 1-3, we get $G - u \cong K_k \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_{r-1}} \cup K_{n_r-1})$ for $n_1 \geq n_2 \geq \dots \geq n_r \geq g + 1$, where $\sum_{i=1}^r n_i = n - k$. Then we have the following claim.

Claim 4. $n_i = g + 1$ for $2 \leq i \leq r$.

Otherwise, there exists $n_j \geq g + 2$ for some $2 \leq j \leq r$. Let $V(K_{n_i}) = \{v_i^1, v_i^2, \dots, v_i^{n_i}\}$ for $1 \leq i \leq r - 1$, $N_{B_r}(u) = \{v_r^1, v_r^2, \dots, v_r^g\}$ and $V(B_r) \setminus N_{B_r}[u] = \{v_r^{g+1}, v_r^{g+2}, \dots, v_r^{n_r-1}\}$. Assume that \mathbf{x} is the Perron vector of $A(G)$. By symmetry, $x(v_i^l) = x_i$ for $1 \leq i \leq r - 1$ and $1 \leq l \leq n_i$, $x(v_r^l) = x_r$ for $1 \leq i \leq g$ and $x(v_r^j) = x'_r$ for $g + 1 \leq j \leq n_r - 1$. Observe that $x(w_1) = x(w_i)$ for $2 \leq i \leq \delta - g$ and $x(w_{\delta-g+1}) = x(w_i)$ for $\delta - g + 2 \leq i \leq k$. According to Lemma 1, we have $x'_r < x_r$ and $x(u) < x_r$. Furthermore,

$$\rho(G)x_1 = (\delta - g)x(w_1) + (k - \delta + g)x(w_{\delta-g+1}) + (n_1 - 1)x_1,$$

$$\begin{aligned}\rho(G)x_r &= (\delta - g)x(w_1) + (k - \delta + g)x(w_{\delta-g+1}) + (g - 1)x_r + (n_r - g - 1)x'_r + x(u) \\ &< (\delta - g)x(w_1) + (k - \delta + g)x(w_{\delta-g+1}) + (n_r - 1)x_r,\end{aligned}$$

from which we obtain that

$$(\rho(G) - n_r + 1)(x_1 - x_r) > (n_1 - n_r)x_1 \geq 0$$

due to $n_1 \geq n_r$. Note that G contains K_{k+n_1} as a proper subgraph, it follows that $\rho(G) > \rho(K_{k+n_1}) = k + n_1 - 1 > n_r$ due to $k \geq \delta + 1 \geq 2$. Thus $x_1 > x_r$. Similarly, we have $x_1 \geq x_j$ for $2 \leq j \leq r - 1$. Let $G_4 = G - \{v_j^i v_j^l \mid 2 \leq j \leq r - 1, 1 \leq i \leq g + 1, g + 2 \leq l \leq n_j\} + \{v_1^p v_j^l \mid 1 \leq p \leq n_1, 2 \leq j \leq r - 1, g + 2 \leq l \leq n_j\} - \{v_r^i v_r^j \mid 1 \leq i \leq g, g + 1 \leq j \leq n_r - 1\} + \{v_1^p v_r^j \mid 1 \leq p \leq n_1, g + 1 \leq j \leq n_r - 1\}$. One can verify that $G_4 \in \mathcal{G}_n^{k, \delta}$. According to Lemma 3, we have $\rho(G_4) > \rho(G)$, which also leads to a contradiction.

Based on the above results, we can conclude that $G \cong G_{n, (g+1)^{r-1}}^{\delta-g, g}$, as desired.

Case 2. $k \leq \delta$. Recall that $u \in V(G)$ such that $d_G(u) = \delta$ and $d_F(u) = t$. Next, we will discuss the case in three subcases.

Subcase 2.1. $\delta < g$.

By the definition of g -good r -component connectivity, we have $d_G(v) \geq g$ for $v \in V(G) \setminus F$. So $u \in F$. Then we consider the following two ways according to the value of k .

In the case of $k \geq 2$. Since $u \in F$ and $|F| = k \leq \delta$, we have $t \leq k - 1$, then we will get the following claim.

Claim 5. $t = k - 1$.

Otherwise, $t \leq k - 2$. Then $|N_G(u) \cap (V(G) \setminus F)| \geq 2$. By the maximality of $\rho(G)$, we can obtain that $G - u \cong K_{k-1} \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r})$. Let $N_F(u) = \{w_1, w_2, \dots, w_t\}$ and $F \setminus N_F[u] = \{w_{t+1}, w_{t+2}, \dots, w_{k-1}\}$, and let $P_i = N_G(u) \cap V(K_{n_i})$ and $|P_i| = p_i$ for $1 \leq i \leq r$. Then $\sum_{i=1}^r p_i = \delta - t$. Let \mathbf{x} be the Perron vector of $A(G)$. By symmetry, we may say $x(v) = x_i$ for $v \in P_i$ and $x(v) = x'_i$ for $v \in V(K_{n_i}) \setminus P_i$, where $1 \leq i \leq r$. Without loss of generality, assume that $x_1 = \max\{x_i \mid 1 \leq i \leq r\}$. Note that $x(w_1) = x(w_i)$ for $2 \leq i \leq t$ and $x(w_{t+1}) = x(w_i)$ for $t + 2 \leq i \leq k - 1$. Using the similar analysis as the proof of subcase 1.1, we get $x(w_{t+1}) > x_1$. Let $P_1 \cup P_2 \cup \dots \cup P_r = \{v_1, \dots, v_{k-t-1}, \dots, v_{\delta-t}\}$ and $G^* = G - \{uv_i \mid 1 \leq i \leq k - t - 1\} + \{uw_j \mid t + 1 \leq j \leq k - 1\}$. Clearly, $G^* \in \mathcal{G}_n^{k, \delta}$. According to Lemma 3, we have

$$\rho(G^*) > \rho(G),$$

which contradicts the maximality of $\rho(G)$. This implies that $t = k - 1$.

By Claim 5, we get $|N_G(u) \cap (V(G) \setminus F)| = \delta - k + 1$, i.e., $\sum_{i=1}^r p_i = \delta - k + 1$. We still consider that $x_1 = \max\{x_i \mid 1 \leq i \leq r\}$. Then we have the following claim.

Claim 6. $p_1 = \delta - k + 1$.

Otherwise, $p_1 \leq \delta - k$. Then $\sum_{i=2}^r p_i \geq 1$. Denote $P_i = \{v_i^1, v_i^2, \dots, v_i^{p_i}\}$ and $V(B_i) \setminus P_i = \{v_i^{p_i+1}, v_i^{p_i+2}, \dots, v_i^{n_i}\}$, where $1 \leq i \leq r$. By the maximality of $\rho(G)$, we have $G - u \cong K_{k-1} \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r})$ and $d_F(u) = k - 1$. Let \mathbf{x} be the Perron vector of $A(G)$. By symmetry, $x(w_1) = x(w_j)$ for $2 \leq j \leq k - 1$, $x(v) = x_i$ for $v \in P_i$ and $x(v) = x'_i$ for $v \in V(K_{n_i}) \setminus P_i$, where $1 \leq i \leq r$. We first assert that $n_1 \geq n_i$ for $2 \leq i \leq r$. If not, there exists $n_j > n_1$ for some $2 \leq j \leq r$. By $A(G)\mathbf{x} = \rho(G)\mathbf{x}$, we obtain

$$\rho(G)x_i = (k - 1)x(w_1) + (p_i - 1)x_i + (n_i - p_i)x'_i + x(u),$$

$$\rho(G)x'_i = (k-1)x(w_1) + p_i x_i + (n_i - p_i - 1)x'_i,$$

where $1 \leq i \leq r$, from which we have

$$x_i = x'_i + \frac{x(u)}{\rho(G) + 1} \quad (10)$$

and

$$(\rho(G) + 1)(x_1 - x_j) = p_1 x_1 + (n_1 - p_1)x'_1 - p_j x_j - (n_j - p_j)x'_j \geq 0 \quad (11)$$

due to $x_1 \geq x_j$. Let $G_1 = G + \{v_j^{n_j} v \mid v \in V(K_{n_1})\} - \{v_j^{n_j} v' \mid v' \in V(K_{n_j}) \setminus \{v_j^{n_j}\}\}$. Clearly, $G_1 \in \mathcal{G}_n^{k, \delta}$. Combining this with (11), we get

$$\begin{aligned} \rho(G_1) - \rho(G) &\geq \mathbf{x}^T (A(G_1) - A(G)) \mathbf{x} \\ &= 2x'_j p_1 x_1 + 2x'_j (n_1 - p_1)x'_1 - 2x'_j p_j x_j - 2x'_j (n_j - p_j - 1)x'_j \\ &= 2x'_j [p_1 x_1 + (n_1 - p_1)x'_1 - p_j x_j - (n_j - p_j - 1)x'_j] \\ &> 0 \quad (\text{by 11}). \end{aligned}$$

Hence $\rho(G_1) > \rho(G)$, which contradicts the maximality of $\rho(G)$. This suggests that $n_1 \geq n_i$ for $2 \leq i \leq r$. Furthermore, let $G_2 = G - \{uv \mid v \in P_i, 2 \leq i \leq r\} + \{uv^j \mid p_1 + 1 \leq j \leq \delta - k + 1\}$. One can easily verify that $G_2 \in \mathcal{G}_n^{k, \delta}$ and $p_1 = \delta - k + 1$. If $\rho(G_2) > \rho(G)$, then the results follows. If not, assume that \mathbf{y} is the Perron vector of $A(G_2)$. By symmetry, we may say $y(v_j^i) = y_j$ for $2 \leq j \leq r$ and $1 \leq i \leq n_j$, $y(v_1^i) = y_1$ for $1 \leq i \leq \delta - k + 1$ and $y(v_1^j) = y'_1$ for $\delta - k + 2 \leq j \leq n_1$. Note that $y(w_1) = y(w_i)$ for $2 \leq i \leq k - 1$ and $y_1 > y'_1$ by Lemma 1, then

$$\begin{aligned} \rho(G_2)y_1 &= (k-1)y(w_1) + y(u) + (\delta - k)y_1 + (n_1 - \delta + k - 1)y'_1, \\ \rho(G_2)y'_1 &= (k-1)y(w_1) + (\delta - k + 1)y_1 + (n_1 - \delta + k - 2)y'_1 > (k-1)y(w_1) + (n_1 - 1)y'_1, \\ \rho(G_2)y_j &= (k-1)y(w_1) + (n_j - 1)y_j, \end{aligned}$$

where $2 \leq j \leq r$, from which we get

$$y_1 = y'_1 + \frac{y(u)}{\rho(G_2) + 1}. \quad (12)$$

Recall that $n_1 \geq n_j$ for $2 \leq j \leq r$, we can deduce that $(\rho(G_2) - n_1 + 1)(y'_1 - y_j) > (n_1 - n_j)y_j \geq 0$. Observe that G_2 contains K_{n_1} as a proper subgraph, thus $\rho(G_2) > \rho(K_{n_1}) = n_1 - 1$. It follows that $y'_1 > y_j$ for $2 \leq j \leq r$. Combining this with (10) and (12), we obtain

$$\begin{aligned} &\mathbf{y}^T (\rho(G_2) - \rho(G)) \mathbf{x} \\ &= \mathbf{y}^T (A(G_2) - A(G)) \mathbf{x} \\ &= (\delta - k + 1 - p_1)(x(u)y(v_1^j) + x(v_1^j)y(u) - x(u)y(v) - x(v)y(u)) \\ &> (\delta - k + 1 - p_1)(x(u)y_1 + x'_1 y(u) - x(u)y'_1 - x_1 y(u)) \quad (\text{since } y'_1 > y_j \text{ and } x_1 \geq x_j \text{ for } 2 \leq j \leq r) \\ &= (\delta - k + 1 - p_1) \left(\frac{x(u)y(u)}{\rho(G_2) + 1} - \frac{x(u)y(u)}{\rho(G) + 1} \right) \quad (\text{by (10) and (12)}) \\ &= (\delta - k + 1 - p_1) \cdot \frac{(\rho(G) - \rho(G_2))x(u)y(u)}{(\rho(G) + 1)(\rho(G_2) + 1)} \\ &\geq 0 \quad (\text{since } \rho(G_2) \leq \rho(G)), \end{aligned}$$

a contradiction. This implies that $p_1 = \delta - k + 1$, proving Claim 6.

Without loss of generality, we may assume that $n_1 \geq n_2 \geq \dots \geq n_r$. By the maximality of $\rho(G)$, we can obtain that $G - u \cong K_{k-1} \vee (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r})$, where $n_r \geq g + 1$ and $\sum_{i=1}^r n_i = n - k$. Then we have the following claim.

Claim 7. $n_i = g + 1$ for $2 \leq i \leq r$.

Otherwise, there exists $n_j \geq g + 2$ for some $2 \leq j \leq r$. Denote $V(K_{n_i}) = \{v_i^1, v_i^2, \dots, v_i^{n_i}\}$ for $1 \leq i \leq r$ and $P_1 = \{v_1^1, v_1^2, \dots, v_1^{\delta-k+1}\}$. Assume that \mathbf{x} is the Perron vector of $A(G)$. By symmetry, we say $x(v_j^i) = x_j$ for $2 \leq j \leq r$ and $1 \leq i \leq n_j$, $x(v_i^1) = x_1$ for $1 \leq i \leq \delta - k + 1$ and $x(v_1^j) = x_1'$ for $\delta - k + 2 \leq j \leq n_1$. Notice that $x(w_1) = x(w_i)$ for $2 \leq i \leq k - 1$ and $x_1 > x_1'$ by Lemma 1. Furthermore,

$$\begin{aligned}\rho(G)x_1' &= (k-1)x(w_1) + (\delta - k + 1)x_1 + (n_1 - \delta + k - 2)x_1' > (k-1)x(w_1) + (n_1 - 1)x_1', \\ \rho(G)x_j &= (k-1)x(w_1) + (n_j - 1)x_j,\end{aligned}$$

where $2 \leq j \leq r$, from which we obtain that

$$(\rho(G) - n_1 + 1)(x_1' - x_j) > (n_1 - n_j)x_j \geq 0$$

due to $n_1 \geq n_j$. Note that G contains K_{n_1} as a proper subgraph, it follows that $\rho(G) > \rho(K_{n_1}) = n_1 - 1$. Thus $x_1' > x_j$ and $x_1 > x_j$. Let $G' = G - \{v_j^i v_j^l \mid 2 \leq j \leq r, 1 \leq i \leq g + 1, g + 2 \leq l \leq n_j\} + \{v_1^h v_j^l \mid 1 \leq h \leq n_1, 2 \leq j \leq r, g + 2 \leq l \leq n_j\}$. It is not difficult to see that $G' \in \mathcal{G}_n^{k, \delta}$. According to Lemma 3, we have $\rho(G') > \rho(G)$, which contradicts the maximality of $\rho(G)$.

Based on the above results, we can conclude that $G \cong G_{n, (g+1)^{r-1}}^{k-1, \delta-k+1}$, as desired.

In the case of $k = 1$. Recall that $u \in F$ and $q = r$. It is worth noting that G is a connected graph, then $\delta \geq r$, $\sum_{i=1}^r p_i = \delta$ and $p_i \geq 1$ for $1 \leq i \leq r$. Without loss of generality, assume that $n_1 \geq n_2 \geq \dots \geq n_r$. Notice that $n_r \geq g + 1$. By the maximality of $\rho(G)$ and using the similar analysis as the proof of Claims 6 and 7, we can obtain that $G \cong G_{n, (g+1)^{r-1}}^{0, \delta}$.

Subcase 2.2. $g \leq \delta < g + k$. Note that $g \neq 0$. Otherwise, $\delta < k$, which contradicts $k \leq \delta$.

Recall that $u \in V(G)$ such that $d_G(u) = \delta$ and $d_F(u) = t$. First of all, we assert $u \notin F$. Otherwise, we consider the following two ways according to the value of k .

In the case of $k \geq 2$. By $u \in F$ and the maximality of $\rho(G)$, using the similar analysis as the proof of Claims 5-7, we can deduce that $G \cong G_{n, (g+1)^{r-1}}^{k-1, \delta-k+1}$ and $n_1 = n - (r-1)(g+1) - k \geq n_2 = \dots = n_r = g + 1$. Let $V(K_{n_i}) = \{v_i^1, v_i^2, \dots, v_i^{n_i}\}$, $P_1 = \{v_1^1, v_1^2, \dots, v_1^{\delta-k+1}\}$ and $N_F(u) = \{w_1, w_2, \dots, w_{k-1}\}$. Assume that \mathbf{x} is the Perron vector of $A(G)$ and $\rho(G) = \rho$. By symmetry, we may say $x(v_i^1) = x_1$ for $1 \leq i \leq \delta - k + 1$, $x(v_1^j) = x_1'$ for $\delta - k + 2 \leq j \leq n_1$ and $x(v) = x_2$ for $v \in V(G) \setminus (F \cup V(K_{n_i}))$. We have $x(w_1) > x_1 > x_1'$ by Lemma 1. Moreover,

$$\begin{aligned}\rho x_2 &= (k-1)x(w_1) + g x_2, \\ \rho x(u) &= (k-1)x(w_1) + (\delta - k + 1)x_1, \\ \rho x_1' &= (k-1)x(w_1) + (\delta - k + 1)x_1 + (n_1 - \delta + k - 2)x_1' > (k-1)x(w_1) + (n_1 - 1)x_1', \\ \rho x(w_1) &= (k-2)x(w_1) + x(u) + (\delta - k + 1)x_1 + (n_1 - \delta + k - 1)x_1' + (r-1)(g+1)x_2,\end{aligned}$$

from which we have $(\rho - n_1 + 1)(x_1' - x_2) > (n_1 - 1 - g)x_2 \geq 0$. Since G contains K_{n_1} as a proper subgraph, $\rho(G) > \rho(K_{n_1}) = n_1 - 1$. It follows that $x_1' > x_2$. Combining this with $x_1 > x_1'$, we get $x_1 > x_2$. Then

$$\begin{aligned}& (n_1 - \delta + k - 1)(x(u) + x_1') + g x(u) + [(r-2)(g+1) + \delta - k + 1]x_2 - (k-1 - \delta + g)x(w_1) \\ &= \frac{(\rho + 1)\{(n_1 - \delta + k - 1)(x(u) + x_1') + g x(u) + [(r-2)(g+1) + \delta - k + 1]x_2 - (k-1 - \delta + g)x(w_1)\}}{\rho + 1} \\ &= \frac{1}{\rho + 1} [(2n_1 + (r-2)(g+1))(k-1)x(w_1) + (2n_1 + k - \delta - 1)(\delta - k + 1)x_1 + n_1 x(u)]\end{aligned}$$

$$\begin{aligned}
& + (n_1 - \delta + k - 1)(n_1 - g)x'_1 + (g + 1)(2(r - 1) + r(\delta - k) - g)x_2] \\
& > \frac{n_1(k - 1)x(w_1) + n_1(\delta - k + 1)x_1 - g(g + 1)x_2}{\rho + 1} \quad (\text{since } r \geq 2, n_1 \geq g + 1 \text{ and } g + k > \delta \geq k) \\
& > \frac{n_1\delta x_2 - g(g + 1)x_2}{\rho + 1} \quad (\text{since } x(w_1) > x_1 > x_2) \\
& \geq 0 \quad (\text{since } n_1 \geq g + 1 \text{ and } \delta \geq g). \tag{13}
\end{aligned}$$

Note that $\delta - k < g$. Let $E_1 = \{uv^i \mid \delta - k + 2 \leq i \leq n_1\} + \{uv^l_j \mid 2 \leq j \leq r - 1, 1 \leq l \leq n_j\} + \{uv^h_r \mid 1 \leq h \leq g\}$, $E_2 = \{v_r^{g+1}w_i \mid \delta - g + 1 \leq i \leq k - 1\}$ and $G_1 = G - E_2 + E_1$. Clearly, $G_1 \in \mathcal{G}_n^{k, \delta}$. Assume that \mathbf{y} is the Perron vector of $A(G_1)$. By symmetry, we may say that $y(v_1^i) = y_1$ for $1 \leq i \leq n_1$, $y(v_j^i) = y_2$ for $2 \leq j \leq r - 1$ and $1 \leq i \leq n_j$, $y(v_r^j) = y_r$ for $1 \leq j \leq g$, $y(w_1) = y(w_i)$ for $2 \leq i \leq \delta - g$ and $y(w_j) = y(u)$ for $\delta - g + 1 \leq j \leq k - 1$. We have $y(u) > y_1$ and $y_r > y(v_r^{g+1})$ by Lemma 1, then

$$\begin{aligned}
\rho(G_1)y_1 &= (\delta - g)y(w_1) + (k - \delta + g)y(u) + (n_1 - 1)y_1, \\
\rho(G_1)y_r &= (\delta - g)y(w_1) + (k - \delta + g)y(u) + (g - 1)y_r + y(v_r^{g+1}) \\
&< (\delta - g)y(w_1) + (k - \delta + g)y(u) + gy_r,
\end{aligned}$$

from which we have

$$(\rho(G_1) - n_1 + 1)(y_1 - y_r) > (n_1 - g - 1)y_r \geq 0.$$

Since G_1 contains K_{n_1} as a proper subgraph, $\rho(G_1) > \rho(K_{n_1}) = n_1 - 1$. It follows that $y_1 > y_r$. Combining this with $y_r > y(v_r^{g+1})$, we have $y_1 > y(v_r^{g+1})$. Furthermore,

$$\begin{aligned}
& \mathbf{y}^T(\rho(G_1) - \rho(G))\mathbf{x} \\
&= \mathbf{y}^T(A(G_1) - A(G))\mathbf{x} \\
&= \sum_{uv^i \in E_1} (x(u)y(v_1^i) + x(v_1^i)y(u)) + \sum_{uv^l_j \in E_1} (x(u)y(v_j^l) + x(v_j^l)y(u)) + \sum_{uv^h_r \in E_1} (x(u)y(v_r^h) + x(v_r^h)y(u)) \\
&\quad - \sum_{v_r^{g+1}w_i \in E_2} (x(v_r^{g+1})y(w_i) + x(w_i)y(v_r^{g+1})) \\
&= (n_1 - \delta + k - 1)(x(u)y_1 + x'_1y(u)) + (r - 2)(g + 1)(x(u)y_2 + x_2y(u)) + g(x(u)y_r + x_2y(u)) \\
&\quad - (k - 1 - \delta + g)(x_2y(u) + x(w_1)y(v_r^{g+1})) \\
&= (n_1 - \delta + k - 1)(x(u)y_1 + x'_1y(u)) + (r - 2)(g + 1)x(u)y_2 + [(r - 2)(g + 1) + \delta - k + 1]x_2y(u) \\
&\quad + gx(u)y_r - (k - 1 - \delta + g)x(w_1)y(v_r^{g+1}) \\
&> y(v_r^{g+1}) \{ (n_1 - \delta + k - 1)(x(u) + x'_1) + gx(u) + [(r - 2)(g + 1) + \delta - k + 1]x_2 - (k - 1 - \delta + g)x(w_1) \} \\
&\quad (\text{since } y(u) > y_1 > y_r > y(v_r^{g+1}) \text{ and } r \geq 2) \\
&> 0 \quad (\text{by 13}),
\end{aligned}$$

and hence $\rho(G_1) > \rho(G)$, which contradicts the maximality of $\rho(G)$. It is easy to see that $d_{G_1}(v_r^{g+1}) = \delta$. This demonstrates that if $k \geq 2$, then $u = v_r^{g+1} \notin F$.

In the case of $k = 1$. By $u \in F$ and the maximality of $\rho(G)$, using the similar analysis as the proof of Claims 6 and 7, we can obtain that $G \cong G_{n, (g+1)^{r-1}}^{0, \delta}$ and $n_1 = n - (r - 1)(g + 1) - 1 \geq n_2 = \dots = n_r = g + 1$. Note that $g \leq \delta < g + 1$, then $\delta = g$. Denote $V(B_i) = \{v_i^1, v_i^2, \dots, v_i^m\}$, $P_1 = \{v_1^1, v_1^2, \dots, v_1^{\delta-r+1}\}$ and $P_j = \{v_j^1\}$ for $2 \leq j \leq r$. Let $G_2 = G + \{uv^i \mid \delta - r + 2 \leq i \leq n_1\} + \{uv^l_j \mid 2 \leq j \leq r - 1, 1 \leq l \leq n_j\} + \{uv^h_r \mid 1 \leq h \leq g\}$. Obviously, $G_2 \in \mathcal{G}_n^{k, \delta}$ and $d_{G_2}(v_r^{g+1}) = \delta$. According to Lemma 2, we get $\rho(G_2) > \rho(G)$, which contradicts the maximality of $\rho(G)$. This means that if $k = 1$, then $u = v_r^{g+1} \notin F$.

Based on the above results, we can conclude that if $g \leq \delta < g + k$, then $u \notin F$. Next, using the similar analysis as the proof of Claims 2-4, we can deduce that $G \cong G_{n,(g+1)^{r-1}}^{\delta-g,g}$, as desired.

Subcase 2.3. $\delta \geq g + k$.

Recall that $|V(B_i)| = n_i$ for $1 \leq i \leq q$ and $q = r$. Therefore,

$$\rho(G) \leq \rho(K_k \vee (K_{n_1} \cup K_{n_2} \cup \cdots \cup K_{n_r})),$$

with equality if and only if $G \cong K_k \vee (K_{n_1} \cup K_{n_2} \cup \cdots \cup K_{n_r})$. Without loss of generality, we may assume that $n_1 \geq n_2 \geq \cdots \geq n_r$. Obviously, $n_r \geq \delta - k + 1$ because the minimum degree of G is δ . Combining this with Lemma 4, we get

$$\rho(K_k \vee (K_{n_1} \cup K_{n_2} \cup \cdots \cup K_{n_r})) \leq \rho(K_k \vee (K_{n-k-(\delta-k+1)(r-1)} \cup (r-1)K_{\delta-k+1})),$$

with equality if and only if $(n_1, n_2, \dots, n_r) = (n-k-(\delta-k+1)(r-1), \delta-k+1, \dots, \delta-k+1)$. By the maximality of $\rho(G)$, we conclude that $G \cong K_k \vee (K_{n-k-(\delta-k+1)(r-1)} \cup (r-1)K_{\delta-k+1})$.

This completes the proof. \square

3 The maximum spectral radius of graphs with g -good neighbor connectivity

Recall that $F \subseteq V(G)$ is called a g -good neighbor cut of G if $G - F$ is disconnected and each vertex $v \in V(G) \setminus F$ has at least g neighbors. The g -good neighbor connectivity of G , denoted by $\kappa_g(G)$ or κ_g , is the minimum cardinality of g -good neighbor cuts of G . Obviously, $\kappa_0(G) = \kappa(G)$. As a conditional connectivity, the g -good neighbor connectivity has improved networks' fault tolerance and has been extensively employed in the analysis of various networks, see [12, 20]. Similar to g -good r -component connectivity, based on the Brualdi-Solheid problem, it is natural for us to consider the following problem.

Problem 3. Which graphs attain the maximum spectral radii among all connected graphs of order n with fixed minimum degree δ and g -good neighbor connectivity $\kappa_g(G)$?

Let $\mathcal{G}_n^{\kappa_g, \delta}$ be the set of graphs of order n with minimum degree δ and g -good neighbor connectivity κ_g . The condition of g -good neighbor connectivity is weaker than that of g -good r -component connectivity, and there is no requirement on the number of components. Therefore, we know that the graphs which achieve the maximum spectral radii among all graphs in $\mathcal{G}_n^{\kappa_g, \delta}$ must be in $\mathcal{G}_n^{k, \delta}$. Furthermore, we have $r = 2$ by Lemma 2. Let $G_{n,g+1}^{\delta,0}$ be the graph obtained from $K_1 \cup (K_{\kappa_g-1} \vee (K_{n-g-1-\kappa_g} \cup K_{g+1}))$ by adding δ edges between the isolated vertex K_1 and K_{κ_g-1} . Let $G_{n,g+1}^{\delta-g,g}$ be the graph obtained from $K_1 \cup (K_{\kappa_g} \vee (K_{n-g-1-\kappa_g} \cup K_g))$ by adding $\delta - g$ edges between the isolated vertex K_1 and K_{κ_g} , and adding g edges between K_1 and K_g . Let $G_{n,g+1}^{\kappa_g-1, \delta-\kappa_g+1}$ be the graph obtained from $K_1 \cup (K_{\kappa_g-1} \vee (K_{n-g-1-\kappa_g} \cup K_{g+1}))$ by adding $\kappa_g - 1$ edges between the isolated vertex K_1 and K_{κ_g-1} , and adding $\delta - \kappa_g + 1$ edges between K_1 and $K_{n-g-1-\kappa_g}$. Let $G_{n,g+1}^{0,\delta}$ be the graph obtained from $K_1 \cup (K_{n-g-2} \cup K_{g+1})$ by adding $\delta - 1$ edges between the isolated vertex K_1 and K_{n-g-2} , and adding one edge between K_1 and K_{g+1} , respectively. Then we have the following result.

Theorem 2. Let $G \in \mathcal{G}_n^{\kappa_g, \delta}$, where $n \geq \kappa_g + 2(g+1)$. Then we have the following statements.

(I) If $\kappa_g > \delta$ and $\delta < g$, then $\rho(G) \leq \rho(G_{n,g+1}^{\delta,0})$ with equality if and only if $G \cong G_{n,g+1}^{\delta,0}$;

(II) If $\kappa_g > \delta \geq g$, then $\rho(G) \leq \rho(G_{n,g+1}^{\delta-g,g})$ with equality if and only if $G \cong G_{n,g+1}^{\delta-g,g}$;

- (III) If $2 \leq \kappa_g \leq \delta < g$, then $\rho(G) \leq \rho(G_{n,g+1}^{\kappa_g-1, \delta-\kappa_g+1})$ with equality if and only if $G \cong G_{n,g+1}^{\kappa_g-1, \delta-\kappa_g+1}$;
- (IV) If $1 = \kappa_g \leq \delta < g$, then $\rho(G) \leq \rho(G_{n,g+1}^{0, \delta})$ with equality if and only if $G \cong G_{n,g+1}^{0, \delta}$;
- (V) If $\kappa_g \leq \delta$ and $g \leq \delta < g + \kappa_g$, then $\rho(G) \leq \rho(G_{n,g+1}^{\delta-g, g})$ with equality if and only if $G \cong G_{n,g+1}^{\delta-g, g}$;
- (VI) If $\delta \geq g + \kappa_g$, then $\rho(G) \leq \rho(K_{\kappa_g} \vee (K_{n-\delta-1} \cup K_{\delta-\kappa_g+1}))$ with equality if and only if $G \cong K_{\kappa_g} \vee (K_{n-\delta-1} \cup K_{\delta-\kappa_g+1})$.

References

- [1] A. Berman, X.D. Zhang, On the spectral radius of graphs with cut vertices, *J. Combin. Theory Ser. B* **83** (2001) 233-240.
- [2] R.A. Brualdi, E.S. Solheid, On the spectral radius of complementary acyclic matrices of zeros and ones, *SIAM J. Algebraic Discrete Methods* **7** (1986) 265-272.
- [3] F. Cao, D.Z. Du, D.F. Hsu, S.H. Teng, Fault tolerance properties of pyramid networks, *IEEE Trans. Comput.* **48** (1999) 88-93.
- [4] G. Chartrand, S.F. Kapoor, L. Lesniak, D.R. Lick, Generalized connectivity in graphs, *Bull. Bombay Math. Colloq.* **2** (1984) 1-6.
- [5] D.D. Fan, S. Goryainov, X.Y. Huang, H.Q. Lin, The spanning k -trees, perfect matchings and spectral radius of graphs, *Linear Multilinear Algebra* **70** (2022) 7264-7275.
- [6] D.D. Fan, X.F. Gu, H.Q. Lin, Spectral radius and edge-disjoint spanning trees, *J. Graph Theory* **104** (2023) 697-711.
- [7] D.D. Fan, X.F. Gu, H.Q. Lin, l -connectivity, l -edge-connectivity and spectral radius of graphs, *arXiv preprint arXiv:2309.05247* (2023).
- [8] F. Harary, Conditional connectivity, *Networks* **13** (1983) 347-357.
- [9] S. Latifi, M. Hedge, M. Naraghi-Pour, Conditional connectivity measures for large multiprocessor systems, *IEEE Trans. Comput.* **43** (1994) 218-222.
- [10] J. Li, W.C. Shiu, W.H. Chan, A. Chang, On the spectral radius of graphs with connectivity at most k , *J. Math. Chem.* **46** (2009) 340-346.
- [11] H.Q. Lin, J.L. Shu, Y.R. Wu, G.L. Yu, Spectral radius of strongly connected digraphs, *Discrete Math.* **312** (2012) 3663-3669.
- [12] H.Q. Liu, X.L. Hu, S. Gao, The g -good-neighbor conditional diagnosability of locally exchanged twisted cubes, *Comput. J.* **63** (2020) 80-90.
- [13] H.L. Lu, Y.Q. Lin, Maximum spectral radius of graphs with given connectivity, minimum degree and independence number, *J. Discrete Algorithms* **31** (2015) 113-119.
- [14] M.J. Ma, The connectivity of exchanged hypercubes, *Discrete Math. Algorithms Appl.* **2** (2010) 213-220.
- [15] W.J. Ning, M. Lu, K. Wang, Maximizing the spectral radius of graphs with fixed minimum degree and edge connectivity, *Linear Algebra Appl.* **540** (2018) 138-148.
- [16] D. Stevanović, Spectral radius of graphs, *Birkhäuser/Springer, Cham*, 2018, 83-130.
- [17] Y. Wang, H.Q. Lin, Y.Z. Tian, Extremal spectral radius and essential edge-connectivity, *Discrete Math.* **347** (2024) 113948.
- [18] B.F. Wu, E.L. Xiao, Y. Hong, The spectral radius of trees on k pendant vertices, *Linear Algebra Appl.* **395** (2005) 343-349.
- [19] M.L. Ye, Y.Z. Fan, H.F. Wang, Maximizing signless Laplacian or adjacency spectral radius of graphs subject to fixed connectivity, *Linear Algebra Appl.* **433** (2010) 1180-1186.

- [20] Q.R. Zhou, H. Liu, B.L. Cheng, Y. Wang, Y.J. Han, J.X. Fan, Fault tolerance of recursive match networks based on g -good-neighbor fault pattern, *Appl. Math. Comput.* **461** (2024) 128318.
- [21] B. Zhu, S.M. Zhang, J.Y. Zou, C.F. Ye, Two kinds of conditional connectivity of hypercubes, *AKCE Int. J. Graphs Comb.* **19** (2022) 255-260.