On Novel Approach for Computing Distance based Indices of Anti-tuberculosis Drugs

D.C. Gunawardhana^{1*}, G.H.J. Lanel², K.K.K.R. Perera³, and A.G.M.J. Gunarathna⁴

¹Department of Mathematical Sciences, Faculty of Applied Sciences, South Eastern University of Sri Lanka

²Department of Mathematics, Faculty of Applied Sciences, University of Sri Jayewardenapura, Sri Lanka

³Department of Mathematics, Faculty of Science, University of Kelaniya, Sri Lanka ⁴Department of Chemistry, Faculty of Science, University of Kelaniya, Sri Lanka

November 6, 2024

Abstract

This work aims to assess the molecular architectures of anti-tuberculosis drugs using both degree-based topological indices and novel distance based indices. We can represent the chemical arrangement as a graph, with atoms serving as the vertices and connections as the edges. Here, the multi bonds were considered as multi edges and included all the hydrogen atoms. Also, we consider three dimensional molecular graph. As a result, the actual bond lengths have been used for computation of new distance based indices. Compared to numerous studies, this is a significant improvement. Furthermore, the investigation of these indices includes a study on the quantitative structure-property relationship (QSPR). The research demonstrates a notable correlation between these indicators and the physical attributes of anti-tuberculosis drugs. Here, Since we reduced the some of existing critical assumptions in the literature, chemists and pharmaceutical professionals might potentially eliminate the need for clinical trials by employing this theoretical model. These models would enable them to predict the characteristics of anti-tuberculosis medications.

Keywords— Degree based indices, distance based indices, Anti-tuberculosis drugs¹

¹* Corresponding author.

Email addresses: rehan@seu.ac.lk (D.C. Gunawardhana), ghjlanel@sjp.ac.lk (Prof. G.H.J. Lanel), kkkrperera@kln.ac.lk (Dr. K.K.K.R. Perera), medhagunaratna@kln.ac.lk(Dr. A.G.M.J. Gunaratna)

1 Introduction

Tuberculosis(TB) is a bacterial infection caused by *Mycobacterium tuberculosis*, leading to primarily lung damage. The primary site of TB infection is the lungs, although it can also impact other organs such as the brain, spine and kidneys. TB continues to be a significant global health issue, well-known for its notorious bacterial infection. TB is transmitted into the air when individuals with lung TB cough, sneeze, or spit upon another. An individual just need inhalation of a small number of germs to become sick. WHO says that, annually, a staggering 10 million individuals become afflicted with tuberculosis (TB). Not with standing its preventability and curability, tuberculosis claims the lives of 1.5 million individuals annually, therefore establishing itself as the leading infectious cause of mortality worldwide [17]. Although most individuals who contract tuberculosis reside in low- and middle-income nations, the disease is prevalent worldwide. A majority of individuals afflicted with tuberculosis are concentrated in eight nations, namely Bangladesh, China, India, Indonesia, Nigeria, Pakistan, Philippines, and South Africa [17].

Despite the significant reduction in TB-related deaths ascribed to the advent of antibiotics, the investigation of the molecular structures of infections and potential treatments has become more vital due to the rise of antibiotic-resistant genotypes.

Hence, the examination of molecular graphs has acquired considerable significance in the field of cheminformatics, including chemistry, mathematics, and information science. An in-depth understanding of the molecular organization of a chemical compound is crucial in the field of chemistry. A molecular graph is a visual representation of the chemical formula of a molecule, with atoms displayed as vertices and chemical bonds as edges. The computation of topological indices that quantify different properties of molecular structures is crucially facilitated by the graphical depiction of these structures.

Numerical values derived from the molecular graph can be used to depict the structural properties of a molecule. The aforementioned techniques have been widely utilized in studies concerning Quantitative Structure-Activity Relationships (QSAR) and Quantitative Structure-Property Relationships (QSPR). Such investigations are essential for predicting the biological activities and physicochemical properties of chemical substances. The Wiener index, developed by Harry Wiener in 1947, was the initial topological statistic to demonstrate a pronounced correlation with the boiling points of alkane compounds [11]. Further studies have shown a relationship between the Wiener index and several properties of molecules, including critical point constants [12], density, surface tension, viscosity [13], and van der Waals surface area [14].

Despite the proven utility of topological indices in predicting certain properties, it is important to recognize that no single indication can accurately correlate with all the physical attributes of a chemical entity. This limitation emphasizes the need of doing additional study on new topological indices that can provide more comprehensive forecasting capabilities in QSAR/QSPR studies.

Object of this work is to assess the advancement of topological indices in enhancing the accuracy of modeling chemical and biological properties of compounds. Specifically, the project will examine their application in addressing antibiotic resistance and enhancing the efficiency of drug discovery. Thus, this work makes a substantial contribution to the field of cheminformatics by greatly enhancing the knowledge of molecular topology. These findings have far-reaching implications in scientific disciplines such as pharmacology, toxicology, and materials research.

The graph G(V, E) with the set of vertices V(G) and the set of edges E(G) is connected if there exists a connection between any pair of vertices in G. The graph theoretical distance between two vertices u and v is denoted as $d(u, v) = d_G(u, v)$ and is defined as the length of shortest path between u and v in graph G. The number of vertices of G adjacent to a given vertex v is the "degree" of this vertex and will be denoted by $d_v(G)$. In all most all the studies, researchers have used two dimensional(2*D*) molecular graph and d(u, v), the graph theoretical distance. For molecular graphs, d(u, v) = 1. Then it is impossible to get meaningful results by considering the graph theoretical lengths. Therefore, we propose to use the actual bond length of three dimensional(3*D*) molecular structure of relevant medicine instead of d(u, v). When it comes to 3*D* molecular graph, let bl(u, v) denotes the geometrical distance between two adjacent atoms.

In [15], M. Adanan et al. have computed the degree-based indices for anti-tuberculosis drugs without considering multi bonds and omitting hydrogen atoms. Similarly, in the almost all studies, researchers have omitted the C-H bond and double/ tipple bonds purposefully to get the required results. Due to the their approaches, the important information which provide from the real structure might be missed. In our work, we consider the full molecule structure to compute the required results.

Some of the degree-based topological indices which we consider in this study are defined as follows.

Definition 1. ABC index [1] is proposed by Estrada et al. defined as,

$$ABC(G) = \sum_{uv \in E(G)} \sqrt{\frac{d(u) + d(v) - 2}{d(u)d(v)}}$$

Definition 2. The Randic index is proposed by Milan Randic [2], as

$$R(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d(u)d(v)}}$$

Definition 3. The sum connectivity index [3] is proposed by Zhou and Trinjstic in, as

$$S(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d(u) + d(v)}}$$

Definition 4. The GA index is proposed by Vukicevic et al. [4] in

$$GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{d(u)d(v)}}{d(u) + d(v)}.$$

Definition 5. The first and second Zagreb indices [5] are proposed by Gutman and Trinajestic, as

$$M_1(G) = \sum_{uv \in E(G)} d(u) + d(v)$$
$$M_2(G) = \sum_{uv \in E(G)} d(u)d(v).$$

Definition 6. The Harmonic index [6] is proposed by Fajtlowiczet al. as,

$$H(G) = \sum_{uv \in E(G)} \frac{2}{d(u) + d(v)}$$

Definition 7. The hyper-Zagreb index [7] is proposed by Shirdelet al. as,

$$HM(G) = \sum_{uv \in E(G)} (d(u) + d(v))^2$$

Definition 8. The third Zagreb index [8] is proposed by Fath-Tabar et al. as

$$ZG_3(G) = \sum_{uv \in E(G)} |d(u) - d(v)|$$

Definition 9. The forgotten index [9] is proposed by Furtula et al. as,

$$F(G) = \sum_{uv \in E(G)} d^2(u) + d^2(v).$$

Definition 10. The symmetric division index [10] is proposed in as,

$$SSD(G) = \sum_{uv \in E(G)} \left(\frac{P}{Q} + \frac{Q}{P}\right)$$

where, $P = \min \{d(u), d(v)\}, Q = \max \{d(u), d(v)\}.$

Definition 11. The Wiener index [16] is denoted by W(G) and defined as,

$$W(G) = \frac{1}{2} \sum_{u,v \in V(G)} d(u,v).$$

Definition 12. *In 1995, M.J. Klein et al. introduced the Hyper Wiener index [18] denoted by WW(G) and defined as,*

$$WW(G) = \frac{1}{2} \left(\sum_{u, v \in V(G)} d(u, v) + \sum_{u, v \in V(G)} d(u, v)^2 \right).$$

Definition 13. *Plavšić et al.*[19] *and Ivanciuc et al.* [20] *independently introduced the Harary index denoted by Ha*(*G*) *and defined as,*

$$Ha(G) = \sum_{u,v \in V(G)} \frac{1}{d(u,v)}.$$

Definition 14. In order to characterize alkanes by an integer, in 1989, Schultz introduced, the Schultz index [21] denoted by Sch(G) and defined as,

$$Sch(G) = \sum_{u,v \in V(G)} (d_G(u) + d_G(v))d(u,v)$$

Definition 15. In 1996, S. Klavžar and I. Gutman introduced Modified Schultz index/ Gutman index [22] is denoted by Gut(G) and is defined as,

$$Gut(G) = \sum_{u,v \in V(G)} (d_G(u) \cdot d_G(v)) d(u,v).$$

Definition 16. In 1982, Alexandru T. Balaban [23, 24] introduced the Balaban index denoted by J(G) and defined as,

$$J(G) = \frac{m}{m-n+2} \sum_{u,v \in V(G)} \frac{1}{d(u,v)+1}.$$

Definition 17. The Klein index is denoted by K(G) and is defined as,

$$K(G) = \sum_{u,v \in V(G)} \frac{d(u,v)}{d(u) + d(v)}.$$

Methods

Inspired by the above distance-based indices, based on the actual bond lengths, here we introduce the following distanced based indices. In this new approach, we suggest that the actual bond lengths of three dimensional chemical structure should capture more information than two dimensional molecular graph structure of a particular drug. Also, Most of applications on topological indices, have used the hydrogen suppression of molecular graphs to get the results with relevant physical-chemical properties of the chemical structures. Also, they have considered the multi-bonds as single bond, in the molecule structure when they calculate the topological indices. But, we suggest that omitting some or all hydrogens, and considering multi-bonds as single bonds , the significant number of information of the molecule is lost. Computing the topological indices without omitting hydrogen and multi-bonds, would give more reliable results than computing indices of hydrogen suppressed structure.

In this study, we compare two computing methods of some topological indices of anti tuberculosis drugs. The methods are as listed below.

- i. The traditional method- In this method, the following assumptions have been made.
 - The hydrogen suppressed molecule structure is studied.
 - The multi-bonds are studied as the single bonds.
 - The two dimensional molecular structure is studied.
 - The graph theoretical distances of two dimensional(2D) chemical graphs are used.
- ii. The proposed method,- In this method
 - The hydrogen atoms aren't omitted.
 - The multi bonds are studied as multi edges.
 - Three dimensional real molecular structure is studied.
 - The geometrical distances of the 3D chemical graphs are considered.

In our work, the proposed method will be used for the comparison of existing degree based indices and novel distance based indices for anti tuberculosis drugs.

Results and discussion

In this section, we compute the degree-based topological indices of the anti-tuberculosis drugs. Also, we calculate the correlation coefficient of those indices and physical properties of relevant drugs.

Now we define new distance based indices based on a property of an edge of the molecular graph. Let bl(u, v) be the weight of the edge uv of the molecular graph G. In the context of molecular graph, we take bl(u, v) as the geometrical bond length of three dimensional molecular graph. We define the new distance based 3D indices, namely Enhanced Wiener index(EW), Enhanced Hyper Wiener index(EHW), Enhanced Second Weiner index(EWW), Enhanced Harary index(EHa), Enhanced Balaban index(EB), Enhanced Shultz index(ES), Enhanced Gutman index(EG) and Modified Klein index(K) as follows.

Definition 18. The Enhanced Wiener index is denoted by EW(G) and is defined as,

$$EW(G) = \sum_{u,v \in V(G)} bl(u,v).$$

Definition 19. The Enhanced second hyper Wiener index is denoted by EHW(G) and is defined as,

$$EHW(G) = \sum_{u,v \in V(G)} bl(u,v)^2.$$

Definition 20. The Enhanced Hyper Wiener index is denoted by EHW(G) and is defined as,

$$EWW(G) = \left(\sum_{u,v \in V(G)} bl(u,v) + \sum_{u,v \in V(G)} bl(u,v)^2\right).$$

Definition 21. The Enhanced Harary index is denoted by EHa(G) and is defined as,

$$EHa(G) = \sum_{u,v \in V(G)} \frac{1}{bl(u,v)}.$$

Definition 22. The Enhanced Schultz index is denoted by ES(G) and is defined as,

$$ES(G) = \sum_{u,v \in V(G)} (d_G(u) + d_G(v))bl(u,v).$$

Definition 23. The Enhanced Gutman index is denoted by EG(G) and is defined as,

$$EG(G) = \sum_{u,v \in V(G)} (d_G(u) \cdot d_G(v)) bl(u,v).$$

Definition 24. The Enhanced Balaban index is denoted by MB(G) and is defined as,

$$EB(G) = \sum_{u,v \in V(G)} \frac{1}{bl(u,v)+1}.$$

Definition 25. The Enhanced Klein index is denoted by K(G) and is defined as,

$$EK(G) = \sum_{u,v \in V(G)} \frac{bl(u,v)}{d(u) + d(v)}.$$

We investigate the relationships between existing degree-based indices, existing distance/degree distance indices, and new distance/degree distance indices with physical properties of antituberculosis drugs, including Boiling point (BP), Melting point (MP), Flash point (FP), Enthalpy of vaporization (EV), Molar refraction (MR), Polarizability (PL), Surface tension(ST) and Molar volume (MV).

Regression models for the novel distance based indices:

Linear regression entails fitting a linear equation to observable data to model the connection between two variables. This theory posits that the first variable serves as an explanatory variable, whereas the second variable functions as a dependent variable. The regression model enables the calculation of linear regression values for each topological index, as demonstrated below:

$P = a + b \left[TI \right]$

The dependent variable is represented by P, while the independent parameter is represented by TI. The parameter P will denote the physical or chemical properties of the pharmaceutical employed in the treatment of tuberculosis, while the parameter TI will signify the estimated values of characteristics or topological indices.

- Boiling Point=70.4 + 6.963 [EW(G)] Flash point =43.5 + 3.738 [EW(G)] Enthalphy of vaporization=28.5+0.908 [EW(G)] Molar refraction=-6.3 + 1.526 [EW(G)] Polarizability=-2.51 + 0.6053 [EW(G)] Molar Volume=-6.4 + 4.133 [EW(G)]
- 2. Boiling Point=90.1 + 5.077 [EHW(G)] Flash point =55.0 + 2.716 [EHW(G)] Enthalphy of vaporization=29.2+0.679 [EHW(G)] Molar refraction=-3.4 + 1.129 [EHW(G)] Polarizability=-1.35 + 0.4476 [EHW(G)] Molar Volume=1.1 + 3.062 [EHW(G)]
- 3. Boiling Point=74.7 + 2.971 [EWW(G)] Flash point =44.6 + 1.603 [EWW(G)] Enthalphy of vaporization=28.9+0.3888 [EWW(G)] Molar refraction=-6.1 + 0.6566 [EWW(G)] Polarizability=-2.46 + 0.2604 [EWW(G)] Molar Volume=-6.3 + 1.780 [EWW(G)]
- 4. Boiling Point=147 + 8.91 [*EHa*(*G*)]
 Flash point =93.3 + 4.59 [*EHa*(*G*)]
 Enthalphy of vaporization=44.0+1.053 [*EHa*(*G*)]
 Molar refraction=36.7 + 1.425 [*EHa*(*G*)]

Polarizability=14.54 + 0.565 [EHa(G)]Molar Volume=102.2 + 4.05 [EHa(G)]

- 5. Boiling Point=64.6 + 20.90 [EB(G)]Flash point =44.9 + 11.02 [EB(G)]Enthalphy of vaporization=28.4 + 2.702 [EB(G)]Molar refraction=-4.1 + 4.424 [EB(G)]Polarizability=-1.63 + 1.754 [EB(G)]Molar Volume=0.5 + 11.94 [EB(G)]
- 6. Boiling Point=351 + 0.483 [ES (G)] Flash point =213.0 + 0.211 [ES (G)] Enthalphy of vaporization=76.7+0.0398 [ES (G)] Molar refraction=61.2 + 0.0875 [ES (G)] Polarizability=24.3 + 0.0347 [ES (G)] Molar Volume=185.9 + 0.212 [ES (G)]
- 7. Boiling Point=135.5 + 0.579 [EG(G)] Enthalphy of vaporization=41.6 + 0.07 [EG(G)] Flash point =77.4 + 0.3125 [EG(G)] Molar refraction=-0.6 + 0.1387 [EG(G)] Polarizability=-0.25 + 0.0550 [EG(G)] Molar Volume=12.2 + 0.3712 [EG(G)]

These distance/ degree-distance indices do not show linear relationships with MP and ST.



(a) Amikacin [26]



(b) Bedaquiline [27]

0

Н

Н



(c) Clofazimine [28]



(f) Ethionamide [31]



(i) Levofloxacin [34]

н.0



(e) Ethambutol [30]



(d) Delamanid [29]

(g) Imipenem [32]



(j) Linezolid [35]

(k) Moxifloxacin [36]

(1) P-Aminosalicyclic acid [37]

н ^{- N} . н

Figure 1: Chemical Structures of Medicines Resource:Chemical structures are taken from PubChem



(a) Amikacin

(b) Bedaquiline



(c) Clofazimine

(d) Delamanid

Figure 2: Chemical Structures of Medicines Resource:Chemical structures are taken from PubChem



(g) Isoniazid

(h) Levofloxacin

Figure 2: Chemical Structures of Medicines (Ctd.) Resource:Chemical structures are taken from PubChem





(i) Linezolid

(j) Moxifloxacin





(k) P-Aminosalicyclic

(l) Pyrazinamide





11

Figure 2: Chemical Structures of Medicines (Ctd.)

Table 1 shows the computed degree based topological indices for the antituberculosis drugs using the proposed method.

Name of Medicine	M1	M2	Η	ABC	R	S	GA	HM	ZG3	F
Amikacin	482	659	31.898	115.568	35.699	36.185	76.446	2974	140	1656
Bedaquiline	563	914	26.495	149.257	29.711	33.112	77.52	3953	105	2125
Clofazizime	493	823	22.057	133.181	24.464	28	67.163	3513	81	1867
Delamanid	491	754	24.56	126.377	27.479	30.386	69.969	3305	111	1797
Ethambutol	210	266	13.893	48.089	16.181	15.902	32.462	1266	74	734
Ethionamide	160	246	8.26	41.181	9.267	10.088	22.977	1076	36	584
Isoniazid	134	210	6.99	35.634	7.677	8.5	19.643	900	26	480
Levofloxacin	352	534	18.071	89.446	20.28	22.28	50.45	2354	84	1286
Linezolid	315	479	15.919	80.122	17.862	19.641	44.984	2109	75	1151
Moxifloxacin	396	601	20.486	100.722	23.069	25.023	56.726	2652	94	1450
p_Aminosalicylic	140	226	7.652	37.734	8.272	9.043	20.779	958	24	506
Pyrazinamide	124	198	6.086	32.898	6.525	7.557	18.078	838	24	442
Rifampin	454	730	23.319	48.116	24.319	28.586	68.849	3028	58	1568
Terizidone	274	430	13.502	71.33	14.981	16.734	38.964	1864	58	1004

Table 1: Values of degree based indices for the proposed method

According to Table 1, the values of degree based indices are in the range 6.086-3953 for the proposed method. The Table 2 shows the physical properties for anti-tuberculosis drugs. Those physical properties are used to compute correlations with topological indices.

Name of Medicine	BP	MP	FP	EV	MR	PL	ST	MV
Amikacin	981.8	203.5	547.6	162.2	134.9	53.5	103.3	363.9
Bedaquiline	702.7	176	378.8	108	156.2	61.9	52.6	420.1
Clofazimine	566.9	210	296.7	85.1	136.2	54	47.1	366.1
Delanamide	653.7	193	349.1	96.3	127.7	50.6	50	368
Ethambutol	345.3	89	113.7	68.3	58.6	23.2	38.1	207
Ethionamide	247.9	163	103.7	46.5	49	19.4	39.8	142
Isoniazid	251.97	172	251	NA	36.9	14.6	57.8	110.2
Levofloxacin	571.5	224	299.4	90.1	91.1	36.1	70.3	244
Linezolid	585.5	177	307.9	87.5	83	32.9	47.7	259
Moxifloxacin	636	270	338.7	98.8	101.8	40.4	60.6	285
P-Aminosalicylic acid	380.8	145	184.1	66.3	39.3	15.6	83.4	102.7
Pyrazinamide	173.3	190	119.1	54.1	31.9	12.6	60.7	87.7
Rifampin	937.4	183	561.3	153.5	213.1	84.5	48	611.7
Terizidone	NA	175	NA	NA	76.1	30.2	62.5	198.9

Table 2: Physical properties of Anti-tuberculosis drugs

Resource: These values are taken from Chemspider.

Also, Table 3 shows the correlation coefficients of degree-based indices and physical properties for the proposed method.

	BP	MP	FP	EV	MR	PL	ST	MV
M1	0.855	0.454	0.782	0.741	0.894	0.894	0.052	0.869
M2	0.808	0.464	0.748	0.677	0.898	0.898	-0.0014	0.868
Η	0.922	0.394	0.839	0.86	0.856	0.856	0.225	0.838
ABC	0.609	0.467	0.512	0.429	0.602	0.602	0.097	0.557
R	0.904	0.387	0.813	0.835	0.829	0.829	0.226	0.811
S	0.912	0.418	0.83	0.834	0.877	0.877	0.171	0.857
GA	0.903	0.439	0.832	0.815	0.909	0.909	0.104	0.887
HM	0.813	0.462	0.745	0.682	0.885	0.885	0.003	0.856
ZG3	0.768	0.365	0.642	0.668	0.615	0.615	0.251	0.603
F	0.815	0.459	0.74	0.683	0.872	0.872	0.018	0.843

Table 3: Correlation Coefficients of Degree based indices and physical properties

Table 4 gives the computed values of the existing distance /degree distance based indices for the selected antituberculosis drugs. In this process, the edge lengths are considered as the graph theoretical distances.

Name of Medicine	W	HW	WW	BB	Sch	Gut
Amikacin	84	84	168	84	482	659
Bedaquiline	84	84	168	84	563	914
Clofazizime	73	73	146	73	493	823
Delamanid	76	76	152	76	491	754
Ethambutol	37	37	74	37	210	266
Ethionamide	25	25	50	25	160	246
Isoniazid	21	21	42	21	134	210
Levofloxacin	55	55	110	55	352	534
Linezolid	49	49	98	49	315	479
Moxifloxacin	62	62	124	62	396	601
p_Aminosalicylic	22	22	44	22	140	226
Pyrazinamide	19	19	38	19	124	198
Rifampin	71	71	142	71	454	730
Terizidone	42	42	84	42	274	430

Table 4: Distance based indices for anti-tuberculosis drugs when distances are graph theoretic distances

As in Table 4, when we consider the graph theoretic distance Weiner index, Hyper Weiner index, second hyper Weiner index, Schultz index and Gutman index, d(u, v) = 1. Then, Weiner index, Hyper Weiner index gives the same values. Also, Shultz index and Gutman Index automatically become the first and second Zagreb indices respectively. In the context of chemical graphs, the Schultz index and Gutman index is meaningless since they coincide with the first and second Zagreb indices when d(u, v) equals graph theoretical distance.

Table 5 presents the computed values of the novel distance/degree-distance based topological indices of the anti-tuberculosis drugs for the proposed drugs.

Name of Medicine	EW	EHW	EWW	EHa	EB	ES	EG
Amikacin	106.826	139.916	246.742	68.096	37.326	640.1	925.689
Bedaquiline	108.992	144.008	253	65.957	36.776	747.862	1253.408
Clofazizime	94.553	126.143	220.696	55.75	31.289	659.764	1129.253
Delamanid	98.8	130.276	229.076	59.35	33.199	650.598	1029.164
Ethambutol	45.285	56.925	102.21	30.999	16.768	267.159	361.86
Ethionamide	32.396	42.974	75.37	19.755	10.971	212.636	338.29
Isoniazid	26.501	26.501	60.439	16.909	9.329	173.582	280.853
Levofloxacin	71.011	93.224	164.235	43.35	24.137	465.607	732.138
Linezolid	62.989	82.35	145.339	82.35	21.557	413.231	648.148
Moxifloxacin	79.752	104.49	184.242	49.132	27.28	523.354	826.327
p_Aminosalicylic	27.849	35.875	63.724	17.724	9.766	183.369	305.664
Pyrazinamide	24.279	31.404	55.683	15.072	8.375	161.539	263.782
Rifampin	99.659	140.702	240.361	50.882	29.598	18.653	1008.919
Terizidone	53.972	70.432	124.403	1.649	18.473	360.422	584.208

Table 5: Novel distance/degree distance based topological indices values of medicines

In the Table 5, the computed novel distance/ degree distance indices were computed using the proposed method. The lengths of edges were taken as the actual bond lengths of three dimensional chemical structure of the relevant anti-tuberculosis drugs.

The correlation values of novel distance/ degree distance indices with physical properties of the given antituberculosis drugs are presented in the Table 6.

Table 6: Correlation c	of new distance	/degree-distance	based indices and	physical	properties
	,	<u> </u>			

	BP	MP	FP	EV	MR	PL	ST	MV
EW	0.905	0.437	0.835	0.818	0.918	0.918	0.099	0.898
EHW	0.912	0.431	0.838	0.830	0.937	0.937	0.083	0.918
EWW	0.910	0.436	0.843	0.825	0.931	0.931	0.090	0.911
EHa	0.785	0.324	0.695	0.645	0.654	0.654	0.011	0.671
EB	0.888	0.430	0.805	0.795	0.869	0.869	0.139	0.847
ES	0.454	0.421	0.341	0.269	0.378	0.378	0.169	0.331
EG	0.816	0.466	0.756	0.690	0.902	0.902	-0.002	0.871

Table 6 presents correlations values of the physical properties of the anti-tuberculosis drugs with the novel distance/ degree-distance based indices.

Figure 3: Correlation of physical properties and novel distance/ degree-distance based topological indices.



Figure 4 illustrates values of the chosen degree based topological indices for each anti-tuberculosis drug for traditional method.



Figure 4: Comparison of degree based topological indices for the traditional method

According to Figure 4, the values of the topological indices vary within 0 - 1600 corresponding to the given drugs. Figure 5 shows values of degree based topological indices for anti-tuberculosis drugs for the proposed method.



Figure 5: Comparison of degree based topological indices for the proposed method

According to Figure 5, the values of degree based indices for the proposed method are within the range 0 - 4000 for relevant drugs.

Figure 6 illustrates the values of the novel distance/ degree-distance topological indices for the proposed method for the given anti-tuberculosis medicines.



Figure 6: Comparison of distance based topological indices for the proposed method

According to Figure 6, we can see the values of novel distance /degree-distance topological indices have been varied within 0 - 1200.

Computation of statistical parameters.

Table 7 - 13 show the statistical parameters for the linear QSPR model for EW(G), EHW(G), EWW(G), EHa(G), EB(G), ES(G) and EG(G). In these tables, N- number of observations, a, b- regression coefficients, r- correlation coefficient, $R^2(adj)$ - coefficient of determination(adjusted).

	N	a	b	r	$R^2(adj)$	F value	p value	Indicator
BP	13	70.4	6.963	0.905	0.8033	50	0	Significant
MP	14	146.5	0.557	0.437	0.1231	2.82	0.119	Not Significant
FP	13	43.5	3.738	0.835	0.6696	25.32	0	Significant
EV	12	28.5	0.908	0.818	0.6362	20.24	0.001	Significant
MR	14	-6.3	1.526	0.918	0.8300	64.48	0	Significant
PL	14	-2.51	0.6053	0.918	0.8300	64.49	0	Significant
ST	14	55.1	0.055	0.099	0	0.1200	0.735	Not Significant
MV	14	-6.4	4.133	0.898	0.7899	49.86	0	Significant

Table 7: Statistical parameters for the linear QSPR model for EW(G)

	N	a	b	r	$R^2(adj)$	<i>F</i> -value	<i>p</i> - value	Indicator
BP	13	90.1	5.077	0.912	0.8157	54.1	0	Significant
MP	14	148.7	0.398	0.431	0.1182	2.74	0.124	Not significant
FP	13	55	2.716	0.838	0.6746	25.88	0	Significant
EV	12	29.2	0.679	0.83	0.6575	22.12	0.001	Significant
MR	14	-3.4	1.129	0.937	0.8684	86.75	0	Significant
PL	14	-1.35	0.4476	0.937	0.8684	86.78	0	Significant
ST	14	55.8	0.033	0.083	0	0.08	0.779	Not significant
MV	14	1.1	3.062	0.918	0.8296	64.28	0	Significant

Table 8: Statistical parameters for the linear QSPR model for EHW(G)

Table 9: Statistical parameters for the linear QSPR model for EWW(G)

	N	a	b	r	$R^2(adj)$	F-value	<i>p</i> -value	Indicator
BP	13	74.7	2.971	0.91	0.8119	52.81	0	Significant
MP	14	147.1	0.236	0.436	12.25	2.81	0.119	Not significant
FP	13	44.6	1.603	0.843	0.6846	27.05	0	Significant
EV	12	28.9	0.389	0.825	0.6492	21.35	0.001	Significant
MR	14	-6.1	0.6566	0.931	0.8548	77.54	0	Significant
PL	14	-2.46	0.2604	0.931	0.8548	77.55	0	Significant
ST	14	55.4	0.0211	0.09	0	0.1	0.759	Not significant
MV	14	-6.3	1.78	0.911	0.8153	58.39	0	Significant

Table 10: Statistical parameters for the linear QSPR model for EHa(G)

	N	a	b	r	$R^2(adj)$	<i>F</i> -value	<i>p</i> -value	Indicator
BP	13	147	8.91	0.785	0.5822	17.72	0.001	Significant
MP	14	161.3	0.542	0.324	0.0304	1.41	0.258	Not significant
FP	13	93.3	4.59	0.695	0.4357	10.27	0.008	Significant
EV	12	44	1.053	0.645	0.3583	7.14	0.023	Significant
MR	14	36.7	1.425	0.654	0.3796	8.95	0.011	Significant
PL	14	14.54	0.565	0.654	0.3794	8.95	0.011	Significant
ST	14	58.38	0.008	0.011	0	0	0.971	Not significant
MV	14	102.2	4.05	0.671	0.4038	9.8	0.009	Significant

Table 11: Statistical parameters for the linear QSPR model for EB(G)

	N	a	b	r	$R^2(adj)$	<i>F</i> -value	<i>p</i> -value	Indicator
BP	13	64.6	20.9	0.888	0.769	40.94	0	Significant
MP	14	145.8	1.68	0.43	0.1173	2.73	0.125	Not significant
FP	13	44.9	11.02	0.805	0.6152	20.18	0.001	Significant
EV	12	28.4	2.702	0.795	0.5947	17.14	0.002	Significant
MR	14	-4.1	4.424	0.869	0.7348	37.03	0	Significant
PL	14	-1.63	1.754	0.869	0.7349	37.03	0	Significant
ST	14	53.4	0.235	0.139	0	0.24	0.635	Not significant
MV	14	0.5	11.94	0.847	0.6936	30.43	0	Significant

	N	a	b	r	$R^2(adj)$	<i>F</i> -value	<i>p</i> -value	Indicator
BP	13	351	0.483	0.454	0.1337	2.85	0.119	Not significant
MP	14	154.3	0.0748	0.421	10.9	2.59	0.133	Not significant
FP	13	213	0.211	0.341	0.0362	1.45	0.254	Not significant
EV	12	76.7	0.0398	0.269	0	0.78	0.399	Not significant
MR	14	61.2	0.0875	0.378	0.0715	2	0.183	Not significant
PL	14	24.3	0.0347	0.378	0.0715	2	0.183	Not significant
ST	14	53.65	0.0129	0.169	0	0.35	0.565	Not significant
MV	14	185.9	0.212	0.331	0.0357	1.48	0.247	Not significant

Table 12: Statistical parameters for the linear QSPR model for ES(G)

Table 13: Statistical parameters for the linear QSPR model for EG(G)

	N	a	b	r	$R^2(adj)$	<i>F</i> -value	<i>p</i> -value	Indicator
BP	13	135.5	0.579	0.816	0.6358	21.95	0.001	Significant
MP	14	145.6	0.0549	0.466	0.1515	0.0332	0.093	Not significant
FP	13	77.4	0.3125	0.756	0.5332	14.71	0.003	Significant
EV	12	41.6	0.07	0.69	0.4231	9.07	0.013	Significant
MR	14	-0.6	0.1387	0.902	0.7982	52.42	0	Significant
PL	14	-0.25	0.05502	0.902	0.7981	52.4	0	Significant
ST	14	58.8	-0.0001	-0.002	0	0	0.995	Not significant
MV	14	12.2	0.3712	0.871	0.7395	37.9	0	Significant

This section has been focused on the calculation of Regression parameters where the sample size, constant or intercept of y, slope, and the correlation coefficient are indicated by N, A, b, and r respectively. r^2 gives the variability percentage of response variable which is explicated by the linear model. Influence of each term tests the null hypothesis which indicates zero coefficients, by using its *p*-value. Conversely, a higher (insignificant) *p*-value indicates that changes in the predictor are not related with changes in the response. Assume we are doing a test where the null hypothesis states that all of the regression coefficients are equal to zero. This test proceeds evaluating the models with no explanatory variables and then measures the enhancement of the model with additive coefficients, by providing an *F* value as the outcome. Somehow, in this scenario, model is unable to provide precise predictions. The Tables 7 - 13 show the statistical parameters of linear QSPR models for a range of topological indices. Each model gives a significant result, where the *p* value is lower or equal to the significance level (0.05).

Tuberculosis (TB) is a bacterial infection caused by the Mycobacterium tuberculosis bacteria. This affects the proper functioning of lungs, while threatening the survival of human beings by diffusing to other body parts, specially brain and spine.

A molecular descriptor is a mathematical formula that can be used on any graph representing a molecule structure. Specifically, a chemical structure is represented by a singular numerical value. Graph theory identifies this numerical value as a topological descriptor, which can be correlated with a molecular element and form a molecular index or specifically a topological index (TI). Topology of the graph is used to describe by a topology index which is a result of the transformation of a molecular graph, where calculating these indices could assist in further examining some sort of physio-chemical molecular aspects.

By calculating these topological indices, it is feasible to assess mathematical values and conduct additional investigation into certain physiochemical aspects of a molecule. Topological indices are created by transforming

a molecular graph into a numerical value that represents the graph's topology. Molecular topological indices have a crucial function in mathematical chemistry, particularly in the study of quantitative structure-property relationship (QSPR) and quantitative structure-activity relationship (QSAR).

Table 2 consists with values of physical properties and Table 1 consists with degree based topological indices for the proposed technique, while Table 3 and Figure 3 indicate their correlation. Table 4 consists of the values of distance based indices for the traditional method while Table 5 illustrates the novel distance/ degree distance based topological indices for the proposed method. Also, Table 6 gives the the correlations for novel distance/degree distance based indices. Upon examining correlation coefficients horizontally for physical properties under consideration, we see that W(G) gives highest correlation coefficient for Molar refraction (r = 0.918) and polarizability (r = 0.918). HW(G) gives highest correlation coefficient for molar refraction and Polarizability (r = 0.937). WW(G) gives highest correlation coefficient for the molar refraction r = 0.931, polarizability (r = 0.931). Ha(G) gives highest correlation coefficient for boiling point (r = 0.785). BB(G) gives highest correlation coefficient for boiling point r = 0.888. Sch(G) gives highest correlation coefficient for boiling point (r = 0.454). Gut(G) gives highest correlation coefficient for molar refraction, polarizability (r = 0.902). When we look vertically, boiling point has also good correlation with HW(G), WW(G), i.e., r = 0.912 and (r = 0.910)respectively. Flash point has good correlation with WW(G), i.e., r = 0.843. Molar refraction has also good correlation with HW(G) and WW(G), i.e., r = 0.937 and r = 0.931. Polarizability has also good correlation with HW(G) and WW(G), i.e., (r = 0.937), r = 0.931. Molar volume has also good correlation with HW(G), WW(G), i.e., r = 0.918, 0.911 respectively.

Table 7 - 13 show different statistical parameters of correlation between values of five distance based and two degree-distance based topological indices and eight physical properties of medicine. The main drawback of our study is the lack of satisfactory correlate findings among degree-based topological index, and the melting point and surface tension of anti-tuberculosis drugs. However, the theoretical analysis of our study could strengthen the prediction of anti-tuberculosis medicinal attributes with zero involvement of experimentation.

Figure 4 shows the comparison of degree based indices for traditional method. Here the values of topological indices vary upto 1600. Also, Figure 5 shows the comparison of the existing degree based topological indices for the proposed method and here the values of topological indices vary within the range 0 - 4000. Next, Figure 6 illustrates the comparison of distance based/degree distance topological indices for proposed method while the value of the indices vary within the range 0 - 1200.

Conclusions

The current study investigates the drugs that are employed to treat tuberculosis. For each drug, the study employs a variety of degree-based numerical descriptors, distance-based topological indices, and newly introduced distance/degree distance-based indices. A total of 14 anti-tuberculosis pharmaceuticals have been investigated using 10 degree-based indices, 6 distance-based indices, and 6 novel distance/degree distance-based indices.

Here we introduced the method to compute the topological indices. When the computing the topological indices, we considered the three dimensional molecular structures of the relevant medications. So, Using parameters of the three dimensional chemical graph to build topological indices, those indices capture more information than the respective two dimensional chemical graphs. Also, in the proposed method, we haven't omitted the hydrogen atoms and considered the structures with all the hydrogen atoms. Also, multi bonds of the structures are considered as multi edges of the three dimensional chemical graphs.

This study investigated the existing 10 degree based topological indices applying the using the aforesaid points.

We suggest that using the proposed method to compute degree based numerical descriptors should give the more reliable results with minimum assumptions when compared to existing computing techniques in cheminformatics.

Also, the novel 6 distance/ degree distance topological indices were introduced using the actual bond length of each edges as a parameter of indices. Using the actual bond length for the topological index, we suggested that the index should give the more information than using the degree of vertices and/or graph theoretical lengths.

These indicators may serve as predictive variables in QSPR studies. The physicochemical properties of antituberculosis medicines are examined to evaluate the predictive efficacy of these parameters. The results indicate a highly significant correlation, demonstrating a robust positive linear association among molecular volume, boiling point, enthalpy, flash point, molar refractivity, vapor pressure, and TIs.

Research suggests that this theoretical analysis could assist chemists and individuals in the pharmaceutical sector in predicting the features of anti-tuberculosis medications without the need for experimentation. It is conceivable that varying combinations of these medications could be employed to treat distinct illnesses, contingent upon the scope of the topological indices calculated in this study. In this study, we have determined the correlation coefficient for several topological indices. This finding will assist chemists in developing novel medications by combining drugs that are highly positively correlated.

Future works

As a future work, studying the connection between the molecular structure and physical characteristics of drugs provides valuable knowledge for predicting the effectiveness and safety of new treatments. Researchers can expedite the pharmaceutical creation process by accurately associating innovative topological indices with numerous physical features, such as solubility, bio-availability, and stability. The objective of this approach is to provide a resilient structure for assessing prospective drug candidates throughout the first stages of their development, with the aim of improving effectiveness and minimizing expenses in pharmaceutical research. Also, numerical methods can be used to obtain best fitted curve (an interpolated curve) of topological index and the relevant physical property.

References

- Estrada, E., Torres, L.A., Rodriguez, L., & Gutman, I. (1998). AN ATOM-BOND CONNECTIVITY IN-DEX: MODELLING THE ENTHALPY OF FORMATION OF ALKANES. *Indian journal of chemistry. Sect. A: Inorganic, physical, theoretical & analytical, 37*, 849-855.
- M. Randi, On Characterization of molecular branching, *Journal of the American Chemical Society*, 97, pp. 6609–6615, 1975.
- [3] B. Zhou and N. Trinajstic, On general sum-connectivity index, J Math Chem, 47, pp. 210–218, 2010.
- [4] D. V.-B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, *J Math Chem*, 46, pp. 1369–1376, 2009.
- [5] I. Gutman, Degree based topological indices, Croatica Chemica Acta, vol. 86, pp. 351–361, 2013.
- [6] S. Fajtlowicz, On conjectures of gra;tti II, Congressus Numerantium, vol. 60, pp. 189–197, 1987.

- [7] G. H. Shirdel, H. RezaPour, and A. M. Sayadi, The Hyper Zagreb Index of Graph Operations, *Iranian J. Math. Chem.*, vol. 4, no. 2, pp. 213–220, 2013.
- [8] A. Astanesh-Asl and G. H Fath-Tabar, "Computing the first and third Zagreb polynomials of certained product of graphs", *Iranian J. Math. Chem.*, vol. 2, no. 2, pp. 73–78, 2011.
- [9] B. Furtula and I. Gutman, "A forgotton topological index", J Math Chem, vol 53, pp. 213–220, 2015.
- [10] V. Alexander, Upper and lower bounds of symmetric division deg index, J Math Chem, vol 5, no. 2, p. 9198, 2014.
- [11] H. Wiener, "Structural determination of paraffin boiling points," *Journal of the American Chemical Society, vol. 69*, pp. 17–20, 1947.
- [12] L. I. Stiel and G. Thodos, "The normal boiling points and critical constants of saturated aliphatic hydrocarbons," *AIChE Journal*, vol. 8, no. 4, Article ID 5276529, 1962.
- [13] D. H. Rouvray and B. C. Crafford, "The dependence of physical-chemical properties on topological factors," *South African Journal of Science, vol.* 72, p. 47, 1976.
- [14] I. Gutman and T. Krtvlyesi, "Wiener indices and molecular surfaces," *Zeitschrift fr Naturforschung, vol.* 50, Article ID6696671, 1995
- [15] Adnan, M., Bokhary, S., Abbas, G., & Iqbal, T. (02 2022). Degree-Based Topological Indices and QSPR Analysis of Antituberculosis Drugs. J. Chem, 2022, 1–17. doi:10.1155/2022/5748626.
- [16] H. Wiener, (1947). Structural determination of the paraffin boiling points, J. Am. Chem. Soc. 69 17-20.
- [17] World Health Organization: WHO. (2019, May 13). Tuberculosis. https://www.who.int/ health-topics/tuberculosis#tab=tab_1.
- [18] D.J. Klein, I. Lukovits, I. Gutman, On the definition of the hyper-Wiener index for cycle-containing structures, J. Chem. Inf. Comput. Sci. 35 (1995) 50–52.
- [19] O. Ivanciuc, T.S. Balaban, A.T. Balaban, Reciprocal distance matrix, related local vertex invariants and topological indices, J. Math. Chem. 12 (1993) 309–318.
- [20] D. Plavšić, S. Nikolić, N. Trinajstić, Z. Mihalić, On the Harary index for the characterization of chemical graphs, J. Math. Chem. 12 (1993) 235–250
- [21] Schultz, H.P. (1989) Topological Organic Chemistry 1. Graph Theory and Topological Indices of Alkanes. Journal Chemical Information and Computational Science, 29, 227-228.
- [22] Klavzar, S. and Gutman, I. (1996) A Comparison of the Schultz Molecular Topological Index with the Wiener Index. Journal Chemical Information and Computational Science, 36, 1001-1003.
- [23] A. T. Balaban, Highly discriminating distance based numerical descriptor, Chem. Phys. Lett. 89 (1982) 399–404.
- [24] A. T. Balaban, Topological indices based on topological distances in molecular graphs, Pure Appl. Chem. 55 (1983) 199–206.

- [25] Parveen S, Farooq FB, Awan NUH, Fanja R, Choudhary MF. Topological indices of drugs used in rheumatoid arthritis treatment and its QSPR modeling. J Maths (2022) 2022:1–11. doi:10.1155/2022/1562125
- [26] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 37768, Amikacin. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Amikacin.
- [27] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 5388906, Bedaquiline. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Bedaquiline.
- [28] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 2794, Clofazimine. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Clofazimine.
- [29] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 6480466, Delamanid. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Delamanid.
- [30] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 14052, Ethambutol. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Ethambutol.
- [31] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 2761171, Ethionamide. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Ethionamide.
- [32] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 104838, Imipenem. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Imipenem.
- [33] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 3767, Isoniazid. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Isoniazid.
- [34] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 149096, Levofloxacin. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Levofloxacin.
- [35] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 441401, Linezolid. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Linezolid.
- [36] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 152946, Moxifloxacin. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/Moxifloxacin.
- [37] National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 4649, 4-Aminosalicylic acid. Retrieved August 3, 2024 from https://pubchem.ncbi.nlm.nih.gov/compound/4-Aminosalicylic-acid.