

Phenotypic heterogeneity in temporally fluctuating environments

Alexander P Browning^{1*} and Sara Hamis²

¹*Mathematical Institute, University of Oxford, Oxford, United Kingdom*

²*Department of Information Technology, Uppsala University, Uppsala, Sweden*

December 17, 2024

Abstract

Many biological systems regulate phenotypic heterogeneity as a fitness-maximising strategy in uncertain and dynamic environments. Analysis of such strategies is typically confined both to a discrete set of environmental conditions, and to a discrete (often binary) set of phenotypes specialised to each condition. In this work, we extend theory on both fronts to encapsulate both a discrete and continuous spectrum of phenotypes arising in response to two broad classes of environmental fluctuations that drive changes in the phenotype-dependent growth rates; specifically, stochastic environments that are temporally uncorrelated (specifically, white-noise processes) and correlated (specifically, Poisson and Ornstein-Uhlenbeck processes). For tractability, we restrict analysis to an exponential growth model, and consider biologically relevant simplifications that pertain to the relative timescale of phenotype switching. These assumptions yield a series of analytical and semi-analytical expressions that reveal environments in which both discrete and continuous phenotypic heterogeneity is evolutionary advantageous.

Keywords: bet-hedging, stochasticity, persists, adaptation, environmental fluctuations

*Corresponding author: browning@maths.ox.ac.uk

1 Introduction

Heterogeneity is ubiquitous to biology [1–4]. Many bacteria strains, for example, employ phenotypic heterogeneity or *persistence* as a bet-hedging strategy against environmental variability [5–7]. Similar strategies are also employed in cancer, a disease in which persistence is now recognised as an important contributor to drug resistance [8, 9]. Such hedging strategies can yield not only an immediate fitness advantage to organisms [10], but are thought to be essential for long-term survival by safeguarding populations against extinction [11, 12].

The degree to which populations bet-hedge is thought to be strongly associated with environment. For instance, so-called high persistence mutants of *Escherichia coli* carry an evolutionary advantage in more frequently stressful environments, but can be outcompeted in less stressful conditions [10, 13]. Complicating the study of bet-hedging is fundamental uncertainty around the environments under which a given organism evolved. Research efforts focus, therefore, on the emergence of evolutionary stable strategies in response to experimentally (or theoretically) prescribed environments [14, 15]. Most typical are environmental models that comprise periodic switching between a finite number of conditions [10, 16]: the presence or absence of an drug, antibiotic, or other stressor, for instance, or the relative prevalence of various nutrient sources [17].

Driven in part by advances in experimental capability, recent years have seen the emergence of a wider spectrum of arguably more realistic experimental and theoretical models of environmental variability [18]. In addition to stochastic analogues of the aforementioned discrete models of environment (typically driven by inhomogeneous Poisson processes) [11, 19–21], recent theoretical work enables interrogation of bet-hedging in environments driven by more general stochastic processes [13, 22–25]. Our recent work [13], among others [19, 22], draws on the similarities between cellular and financial hedging to demonstrate that heterogeneity can be evolutionary advantageous in environments driven by white-noise processes that yield growth rates with constant mean. These works also study what we refer to as *continuously fluctuating environments*: environments that manifest continuous changes in growth rate, such as those described by Ornstein-Uhlenbeck processes or generalisations thereof. There is, however, little general theory available to elucidate bet-hedging in arbitrarily fluctuating environments.

Microorganisms regulate phenotypic variability through gene expression, such that otherwise genetically identical individuals exhibit differences in traits [3]. In bet-hedging bacteria, for example, persisters exhibit dormancy and antibiotic tolerance [6]. Such quiescent phenotypes are typically characterised by a metabolism that differs from regular proliferative bacteria: this manifests as both lower proliferation rates (in growing conditions) and lower death rates (in stressful conditions) [6, 26, 27]. Persisters can be formed both actively (i.e., in response to environmental stressors through stress response hormones including (p)ppGpp [28–30]) and passively (i.e., through environment-independent phenotype switching [6]). Passive switching may be regulated through gene expression, whereby sufficiently high levels of a persistence gene may correspond to the quiescent state. This expression-threshold theory underscores the overwhelmingly common view of heterogeneity as a discrete (and often binary) set of otherwise




Figure 1. Population dynamics model. We assume that cell proliferation is mediated by symmetric division (top), while Phenotypic heterogeneity is regulated through switching where, upon distribution, each cell is assigned a phenotype according to a discrete or continuous target distribution (bottom).

homogeneous subpopulations [31,32]. Subpopulation level variation is typically not considered in the analysis of bet-hedging strategies outside a small number of largely theoretical works [33–36]. While models that capture such continuous heterogeneity are widely studied in the mathematical literature (typically in the context of cancer plasticity [37, 38]), there are few studies that investigate continuous heterogeneity in stochastically fluctuating environments [39].

Theoretical models of bet-hedging range in scale from intracellular studies of the pathways responsible for persister regulation [40], to high-level studies of populations that regulate heterogeneity through phenotype switching [26]. At the population-level, works study the evolutionarily stable strategy (ESS), or fitness-maximising strategy, under a prescribed growth model coupled to an environmental process [41–43]. Growth is typically modelled as exponential, representative of batch- or continuous-culture experiments, where cell concentrations are kept sufficiently low that intercellular competition—which may obfuscate or otherwise alter the evolutionary dynamics—is negligible. Study of the ESS then corresponds to a stochastic optimisation problem in the Malthusian fitness or, in the case that organisms are able to actively react to the environment, a stochastic optimal control problem [13, 44].

In this work, we develop mathematical theory to study the environmental conditions in which bet-hedging is advantageous. Throughout, we consider a Malthusian-like measure of fitness given by the expected per-capita growth rate [27, 41, 42]

$$\varphi = \mathbb{E} \left(\frac{1}{n} \frac{dn}{dt} \right) = \mathbb{E} \left(\frac{d \log n}{dt} \right), \quad (1)$$

where $n(t)$ denotes the overall size (or concentration) of a phenotypically structured population. We make the standard assumption that phenotypic traits are heritable through symmetric exponential division, and that phenotypic heterogeneity is regulated through a phenotype switching strategy (Fig. 1) [6, 10]. Evolutionary stable strategies are, therefore, switching strategies that maximise Eq. (1). In Section 2, we consider a fitness maximising phenotype switching strategy in an environment driven by a pair of temporally uncorrelated white-noise processes. We then, in Section 3, extend the analysis to temporally correlated environments driven by an inhomogeneous Poisson process and by a family of continuously varying environments driven by an Ornstein-Uhlenbeck process and characterised by a specified long-term distribution

of growth rates. In all but the simplest of cases, the calculation of population fitness (Eq. (1)) is both numerically and analytically intractable. Thus, we make progress by studying a series of biologically relevant limiting cases based around the relative time-scales of environmental fluctuations and phenotypic-switching. This approach allows us to determine sufficient conditions for parameter regimes in which discrete and continuous heterogeneity is advantageous.

2 Growth in environments driven by temporally uncorrelated white-noise

We first consider a cell population subject to growth rate fluctuations driven by white-noise. The population is phenotypically structured such that each individual is associated with a value of a *phenotype index*, denoted by $\varepsilon \in [0, 1]$. Two individuals with the same phenotypic index are assumed to behave identically. In practice, the phenotype index may be multidimensional and determined by the expression of a potentially large number of genes; in this work, we restrict our analysis to scalar ε .

In our previous work [13], we describe a two-phenotype model of regular bacteria and persisters driven by two independent white-noise processes. We now consider a more general model through a phenotype-dependent growth rate, $\lambda(\varepsilon)$, of the form

$$\lambda(\varepsilon) = \mu(\varepsilon) + s_1(\varepsilon)\xi_1 + s_2(\varepsilon)\xi_2 \quad (2)$$

where ξ_i are independent Gaussian variables (i.e., growth is driven by two independent white-noise processes), $\mu(\varepsilon)$ gives the time-averaged growth rate, and $s_1(\varepsilon)$ and $s_2(\varepsilon)$ together determine the correlation structure and growth rate variance. One interpretation of a growth rate driven by N independent white-noise processes is of N nutrient sources. While the problem loses analytical tractability for large but finite N , a similar analysis can be conducted in the limit $N \rightarrow \infty$ by considering that the growth rates are given by a Gaussian process with specified autocovariance function $\sigma(\varepsilon_1, \varepsilon_2)$.

Formulation of an appropriate model for the dynamics in the presence of an arbitrary number of phenotypes is nontrivial. Many existing models of bacterial persistence implicitly assume that the phenotypic state is inherited upon proliferation [10], while some studies assume the opposite. Following the majority, we assume that the phenotypic state of individuals is heritable, while phenotypic transitions are distributive: all individuals switch phenotype to that chosen from a genetically-encoded *target distribution*, which we denote by $\hat{p}(\varepsilon)$ (Fig. 1). The switching rate, denoted by ω , may, in general, vary between phenotypes; for convenience we assume it to be phenotype-independent. Altogether, the concentration of cells with phenotype index ε_i , denoted by $r(\varepsilon_i, t)$, is governed by the stochastic partial differential equation (SPDE)

$$\frac{\partial r(\varepsilon_i, t)}{\partial t} = \underbrace{\lambda(\varepsilon_i)r(\varepsilon_i, t)}_{\text{Proliferation}} + \underbrace{\omega \hat{p}(\varepsilon_i)n(t) - \omega r(\varepsilon_i, t)}_{\text{Phenotype switching}}, \quad (3)$$

where $n(t) = \sum_i r(\varepsilon_i, t)$ (or the integral in the case of continuous ε) is the overall cell concen-

tration, governed by the stochastic differential equation (SDE)

$$\frac{dn(t)}{n} = \mathbb{E}_{p(\varepsilon,t)}(\mu(\varepsilon)) dt + \sqrt{\sum_{i=1}^2 \mathbb{E}_{p(\varepsilon,t)}^2(s_i(\varepsilon))} dW(t). \quad (4)$$

Here, we denote by $p(\varepsilon, t) = r(\varepsilon, t)/n(t)$ the current distribution of phenotypes in the population, and by $\mathbb{E}_{f(\varepsilon)}$ an expectation with respect to the probability mass or density function $f(\varepsilon)$. Extensions to the growth model (Eq. (3)) could include a relatively slow diffusion term such that the phenotype of an individual also undergoes a relatively slow random walk throughout the phenotypic space [37].

Analysis of the full SPDE (Eq. (3)) is analytically intractable. Hence, we make initial progress by considering a hypothetical scenario whereby the population is able to *control* a fixed phenotype distribution, denoted $p(\varepsilon)$, and studying the distribution that maximises population fitness, as determined by Eq. (1). This scenario is analogous to assuming that the switching rates ω are sufficiently large that the actual phenotype distribution, $p(\varepsilon, t)$, is in a *quasi-steady-state* (QSS). In the case of general ω we must resort to a numerical approximation, which is only tractable in the case that the target distribution, $\hat{p}(\varepsilon)$, is discrete.

Eq. (4) defines a geometric Brownian motion where the fitness is given by

$$\varphi = \mathbb{E} \left(\frac{d \log n}{dt} \right) = \mathbb{E}_{p(\varepsilon)}(\mu(\varepsilon)) - \frac{1}{2} \sum_{i=1}^2 \mathbb{E}_{p(\varepsilon)}^2(s_i(\varepsilon)). \quad (5)$$

Thus, we wish to study the effect of a controlled population composition, $p(\varepsilon)$, on the fitness of the population. In particular, we wish to establish parameter regimes whereby a heterogeneous strategy carries a fitness advantage over a homogeneous strategy.

2.1 Discrete phenotypic states

While Eq. (4) allows for a relatively arbitrary phenotypic structure, we begin our analysis by building on our previous work [13] and analysing a two phenotype population structure [6, 10]. Growth rates are given by

$$\lambda(\varepsilon) = \begin{cases} \mu_1 + \sigma_1 \xi_1, & \varepsilon = 1, \\ \mu_2 + \sigma_2 \rho \xi_1 + \sigma_2 \sqrt{1 - \rho^2} \xi_2, & \varepsilon = 0, \end{cases} \quad (6)$$

where $\sigma_i^2 = \text{Var}(\lambda_i)$ represents the variance of each growth rate, and $\rho := \text{Corr}(\lambda_1, \lambda_2)$ represents the correlation. An example set of fluctuating growth rates is given in Fig. 2.

In isolated homogeneous populations, the effective growth rate (or fitness) of each phenotype is given by

$$\varphi_i = \mu_i - \frac{\sigma_i^2}{2}. \quad (7)$$

Without loss of generality, we take $\mu_1 = 1$ and assume that the phenotypes are ordered such that $\varphi_2 < \varphi_1$. For bet-hedging systems (i.e., bacterial persistence), we may additionally assume that $\mu_2 \ll 1$, however, this need not be the case. For example, phenotypes may be specialised




Figure 2. Correlated white-noise growth in a discrete two-phenotype population. (a) Correlated growth rates in a highly proliferative phenotype (blue, $\varepsilon = 1$), and a slow-growing dormant phenotype (red, $\varepsilon = 0$). The population is able to increase its effective growth rate from $\varphi(1) = 0.5$ to $\varphi(q^*) \approx 0.56$ using an optimal heterogeneous strategy with $\hat{p}(0) = q^* \approx 0.32$. (b) Stationary distribution of $p(0, t)$ (proportion of dormant cells) arising from a switching strategy with $\omega = 1$ and target distribution $\hat{p}(0) = q^*$ approximated using repeated simulation of the SDE (yellow) and through the Fokker-Planck partial differential equation (blue). (c) Numerical calculations for fitness as a function of the target distribution $\hat{p}(\varepsilon)$ for increasing values of the switching rate ω . For fast switching rates, the optimal target distribution is similar to the optimal controlled distribution. Even for relatively slow switching rates, the optimal target distribution is similar to the optimal controlled distribution. Parameters are set to $\mu_1 = 1, \mu_2 = 0.3, \sigma_1 = 1, \sigma_2 = 0.1, \rho = -0.5$.

to different nutrient sources and thus manifest similar expected growth rates.

To build the model for the heterogeneous population, we follow the QSS assumption and thus assume that the population is able to modulate a constant proportion $q := p(0)$ of the dormant phenotype through phenotype switching. We therefore wish to establish conditions under which a heterogeneous population (i.e., $q > 0$) is evolutionary advantageous over a homogeneous population (i.e., $q = 0$).

Applying Eq. (5) yields a quadratic expression in q for the effective growth rate of the (potentially) heterogeneous population, denoted $\varphi(q)$, and given alongside a full derivation in the supplementary material. Since we can recover homogeneous populations of each phenotype by setting $p(i) = 1$ for $i = 0, 1$, we have $\varphi_1 = \varphi(1)$ and $\varphi_2 = \varphi(0)$. Given that $\varphi_1 > \varphi_2$ by our previous assumption, the optimal proportion, q^* , that maximises the effective growth rate will be on the interior $q^* \in (0, 1)$ (else $q^* = 1$) provided that (1) $\varphi''(q)$ is strictly negative, and that (2) $q^* > 0$. Both conditions hold provided that

$$\mu_2 > \mu_1 + \rho\sigma_1\sigma_2 - \sigma_1^2, \quad (8)$$

in which case the optimal proportion is given by

$$q^* = \frac{\mu_2 - \mu_1 + \sigma_1^2 - \rho\sigma_1\sigma_2}{\sigma_1^2 - 2\rho\sigma_1\sigma_2 + \sigma_2^2}. \quad (9)$$

Given that $\varphi_1 > \varphi_2$, which places an upper bound on μ_2 , we further note that heterogeneity is only advantageous in this model if

$$\frac{\sigma_1^2 + \sigma_2^2}{2} > \rho\sigma_1\sigma_2. \quad (10)$$




Figure 3. Unimodal model of continuous phenotypic variability. (a) We consider correlation functions that are either continuously differentiable at $\Delta_\varepsilon = 0$ (blue; $\rho_1(\Delta_\varepsilon) = \exp(-5\Delta_\varepsilon^2)$) or not (red; $\rho_2(\Delta_\varepsilon) = \exp(-5|\Delta_\varepsilon|)$). (b-c) The fitness relative to $\eta = 0$ is computed as a function of phenotype variance η^2 using numerical quadrature (coloured curves) and the asymptotic expressions (black). The optimal phenotype ε^* is (b) on the interior of the domain at $\varepsilon^* = 0.5$ and (c) at $\varepsilon^* = 1$. In (b-c), the variance model is given by Eq. (12) with $k = 1$ and (b) $\gamma = \sqrt{2}$ and (c) $\gamma = 1$.

Clearly, the range of scenarios in which a two-phenotype strategy is advantageous is reduced as $\rho \rightarrow 1$ and increases as $\rho \rightarrow -1$. For the example presented in Fig. 1, the population may employ a second phenotype to increase the fitness by approximately 12% from 0.5 to 0.56 through heterogeneity with $q^* \approx 0.32$ (parameter values are given in the figure caption).

2.1.1 Finite switching speed

The aforementioned results hold only for sufficiently large and phenotype-independent ω such that the population is, effectively, able to control its composition in the face of growth rate fluctuations. To investigate the effect of finite ω , we consider that the stationary distribution for $p(0, t) := x(t)$ for the actual population composition is given by a boundary value problem on $x(t) \in [0, 1]$ arising out of the one-dimensional Fokker-Planck equation for $x(t)$ (full details are available in the supplementary material). This equation admits, up to a normalisation constant, an analytical solution, as shown in Fig. 2c. The expectation in Eq. (5) can then be taken using numerical quadrature to obtain the fitness for general ω .

Results in Fig. 2a show the fitness as a function of ω and the now target dormant proportion q , in addition to the optimal target composition for each ω . First, it is clear that the optimal target composition converges to the theoretical optimum (Eq. (9)) for sufficiently large ω . Secondly, we find that the optimum remains relatively close to the large ω limit, although the fitness advantage of the homogeneous strategy is reduced for small ω . For white-noise environments, we conjecture that the fitness may be a monotonic function of the switching rate: specifically, a monotonically increasing function in cases where heterogeneity is advantageous, and a monotonically decreasing function in cases where the homogeneous strategy is favoured.

2.2 Continuously distributed phenotypes

We now consider that an otherwise homogeneous population undergoes mutations that yield a continuous distribution of phenotypes throughout the phenotypic space. We assume that the distribution is initially unimodal, and develops over relatively slow evolutionary timescales so

that the initial variance of the phenotype distribution, denoted by η^2 , is sufficiently small. In Section 2.2.2, we extend this analysis to a bimodal model where both populations in the discrete two-phenotype model develop subpopulation heterogeneity simultaneously.

2.2.1 Unimodal strategy

The first model of continuous phenotypic variability is unimodal and local in that that an initially homogeneous and otherwise evolutionarily stable population undergoes small mutations that lead to relatively small differences in the growth rate between individuals. The growth rate is assumed to be given by

$$\lambda(\varepsilon) := \mu(\varepsilon) + \sigma(\varepsilon)\xi \quad (11)$$

where $\xi \sim \mathcal{N}(0, 1)$. We also assume that both $\mu(\varepsilon)$ and $\sigma(\varepsilon)$ are monotonically increasing functions of ε .

To provide a set of analytically tractable results, we focus our analysis on the case that $\sigma^2(\varepsilon)$ is quadratic with form

$$\sigma^2(\varepsilon) = \gamma^2 [k(\varepsilon^2 - \varepsilon) + \varepsilon], \quad (12)$$

such that k represents the curvature and γ^2 represents the variance at $\varepsilon = 1$. For $k = 0$, the variance scales linearly with growth rate, for $k < 0$ the variance is convex such that the variance increases more quickly at smaller values of ε , and for $k > 0$ a faster increase in variance is seen at larger values of ε . We additionally consider that $\mu(\varepsilon) = \varepsilon$ with a corresponding maximum at $\mu(1) = 1$. This latter choice can be made without loss of generality as the phenotype index itself is arbitrary and a temporal rescaling can always be applied to enforce a given maximum growth rate.

We first determine the phenotype in a homogeneous strategy, denoted by ε^* , that maximises the fitness metric Eq. (5). It can easily be seen that phenotypes on the interior $\varepsilon^* \in (0, 1)$ can only be optimal for $k > 0$. More generally, this observation is true for all concave variance functions: a necessary (but not sufficient) condition for the optimal phenotype to lie on the interior of the domain is that $\sigma^2(\varepsilon)$ is *not* linear or strictly concave. For the quadratic model, we state the optimal phenotype for various γ in Table 1.

Next, we suppose that the population develops heterogeneity such that the equilibrium phenotype distribution in the population is approximately normal such that $\varepsilon \sim \mathcal{N}_{(0,1)}(\varepsilon^*, \eta^2)$ for $\eta \ll 1$, where $\mathcal{N}_{(0,1)}$ indicates a truncated Gaussian distribution on $\varepsilon \in [0, 1]$.

Table 1. Optimal single phenotype ε^* for the continuously distributed phenotype model with quadratic variance function given by Eq. (12). The parameter γ^2 determines the growth rate variance at $\varepsilon = 1$, and k determines the concavity (or convexity) of the variance as a function of ε .

	$\gamma^2 < 2$	$\gamma^2 > 2$
Linear	1	0
Convex	$\min \left(1, \max \left[0, \frac{1}{2} + \frac{1}{k} \left(\frac{1}{\gamma^2} - \frac{1}{2} \right) \right] \right)$	
Concave	1	0

To extend the growth model to a continuum of phenotypes, we define a correlation function,

$$\rho(\Delta_\varepsilon) = \text{Corr}(\lambda(\varepsilon^* + \Delta_\varepsilon), \lambda(\varepsilon^*)), \quad (13)$$

that describes the correlation between growth rates, relative to the optimal phenotype, ε^* . We always assume that the correlation function is an even function. This formulation corresponds to the general model (Eq. (2)) with

$$s_1(\varepsilon) = \rho(\varepsilon - \varepsilon^*)\sigma(\varepsilon), \quad (14a)$$

$$s_2(\varepsilon) = \sigma(\varepsilon)\sqrt{1 - \rho^2(\varepsilon - \varepsilon^*)}. \quad (14b)$$

The integrals required to compute the expected growth rate will, generally, be intractable. However, we make progress by considering that $0 < \eta \ll 1$ and studying the behaviour of $s_1(\varepsilon)$ and $s_2(\varepsilon)$ within a small neighbourhood of ε^* . The sign of the first-order correction term thus determines whether continuous heterogeneity is advantageous for small η , and can be interpreted as a sufficient (albeit not necessary) condition for heterogeneity to present some advantage.

We first consider cases where the optimal single phenotype, ε^* , lies on the interior of the domain. Results can be summarised by two subcases, which depend on the behaviour of the correlation function near $\Delta_\varepsilon = 0$ (as shown in Fig. 3a):

1. $\rho(\Delta_\varepsilon)$ is differentiable at $\Delta_\varepsilon = 0$. To leading order, we have that

$$\varphi_\eta = \varphi_0 - \underbrace{\frac{\sigma(\varepsilon^*)\left((\pi - 2)\sigma(\varepsilon^*)\rho''(0) + \pi\sigma''(\varepsilon^*)\right)}{2\pi}}_{\text{Leading order correction}} \eta^2 + \mathcal{O}(\eta^3). \quad (15)$$

2. $\rho(\Delta_\varepsilon)$ is not differentiable at $\Delta_\varepsilon = 0$. To leading order, we have that

$$\varphi_\eta = \varphi_0 + \frac{\sqrt{2}\sigma^2(\varepsilon^*)|\rho'_-(\varepsilon^*)|(\sqrt{\pi} - \Gamma^2(\frac{3}{4}))}{\pi} \eta + \mathcal{O}(\eta^2), \quad (16)$$

where $\rho'_-(\varepsilon^*) := \lim_{\Delta_\varepsilon \rightarrow 0^-} \rho'(\Delta_\varepsilon)$.

In the second case we can readily see that the leading order correction term is positive, and conclude therefore that there exists (at least sufficiently small) choices of η such that $\varphi_\eta \geq \varphi_0$, for all choices of $\sigma(\varepsilon)$. In the first case, heterogeneity will be advantageous provided that

$$|\rho''(0)| > \frac{\pi\sigma''(\varepsilon^*)}{(\pi - 2)\sigma(\varepsilon^*)},$$

a condition that will always hold for concave $\sigma(\varepsilon)$, and will also hold for convex $\sigma(\varepsilon)$ provided that $|\rho''(0)|$ is sufficiently large. We verify these results in Fig. 3b by comparing the asymptotic expansion for the fitness to that calculated numerically through quadrature.

We proceed with a similar analysis for the case where $\varepsilon^* = 1$ on the exterior of the domain. The approach is similar, however asymptotically ε now has a half-Gaussian distribution with scale parameter η that can no longer be interpreted directly as the phenotype variance (the

variance in this case is given by $(1 - 2/\pi)\eta^2$). Similarly, we can characterise the results into two cases that depend on the behaviour of the correlation function at $\Delta_\varepsilon = 0$:

1. $\rho(\Delta_\varepsilon)$ is differentiable at $\Delta_\varepsilon = 0$. To leading order, we have that

$$\varphi_\eta = \varphi_0 + \sqrt{\frac{2}{\pi}} \left(\sigma(1)\sigma'(1) - \frac{1}{\sqrt{\pi}} \right) \eta + \mathcal{O}(\eta^2). \quad (17)$$

2. $\rho(\Delta_\varepsilon)$ is not differentiable at $\Delta_\varepsilon = 0$. To leading order, we have that

$$\varphi_\eta \sim \varphi_0 + \left(\frac{\sqrt{2}\sigma(1)(\sigma(1)\rho'_-(0) + \sigma'(1))}{\sqrt{\pi}} - \frac{\sqrt{2}\sigma^2(1)\rho'_-(0)\Gamma(\frac{3}{4})^2}{\pi} - 1 \right) \eta + \mathcal{O}(\eta^2). \quad (18)$$

In the first case, heterogeneity will be advantageous if $\sigma(1)\sigma'(1) > 1$. For the quadratic model, this will hold, provided $(1+k)\omega^2 > 2$. We again verify these results in Fig. 2c by comparing the asymptotic expansion for the fitness to that calculated numerically through quadrature.

2.2.2 Bimodal strategy

The next model of continuous heterogeneity that we consider is based on the binary model studied in Section 2.1. We assume that a population develops continuous heterogeneity from an otherwise evolutionary stable discrete strategy, such that the population distribution is a mixture

$$p(\varepsilon) = q^* \mathcal{N}_{(0,1)}(0, \eta_1) + (1 - q^*) \mathcal{N}_{(0,1)}(1, \eta_2), \quad (19)$$

where q^* is the optimal discrete proportion of individuals at $\varepsilon = 0$ (Eq. (9)).

For correlation functions that are differentiable at both $\Delta_\varepsilon = 0$ and $\Delta_\varepsilon = 1$, the fitness is given, to leading order, by

$$\varphi_{(\eta_1, \eta_2)} = \varphi_0 + \sqrt{\frac{2}{\pi}} \left(c_1 q^* \eta_1 + c_2 (1 - q^*) \eta_2 \right), \quad (20)$$

where

$$\begin{aligned} c_1 &= \mu'(0) - (1 - q^*)\sigma'(0)\sigma(1)\rho(1) - \left(q^*\sigma'(0) + (1 - q^*)|\rho'(1)|\sigma(1) \right) \sigma(0), \\ c_2 &= -\mu'(1) - q^*\sigma(0)\sigma(1)\sqrt{|\rho''(0)|(1 - \rho(1)^2)} + \left(q^*\sigma(0)\rho(1) + (1 - q^*)\sigma(1) \right) \sigma'(1). \end{aligned}$$

Thus, continuous heterogeneity is advantageous if either c_1 or c_2 are positive.

In Fig. 4, we explore the effect of continuous heterogeneity arising out of the discrete strategy demonstrated in Fig. 2. Here, we take $\sigma(\varepsilon)$ to be of a quadratic form with $k = 0.3$ and $\rho(\Delta_\varepsilon)$ to be differentiable, with all other parameters chosen such that $\sigma(0) = \sigma_2$, $\sigma(1) = \sigma_1$, and $\rho(1) = \rho$ correspond to earlier analysis in Fig. 2. In Fig. 4a, we demonstrate that Eq. (19) tends to a uniform distribution for $\eta_1, \eta_2 \gg 1$. Results Fig. 4b show, both numerically and through




Figure 4. Bivariate continuous model of heterogeneity. We consider that the population in Fig. 2a develops continuous heterogeneity at both $\varepsilon = 0$ (with variance η_1^2) and at $\varepsilon = 1$ (with variance η_2^2), such that the target distribution follows a mixture of truncated Gaussian distributions. (a) Example target distributions for $(\eta_1, \eta_2) = (0.4, 0.4)$ and $(\eta_1, \eta_2) = (0.1, 0.1)$. For either $\eta_1, \eta_2 \gg 0$, the target distribution approaches a uniform distribution (this is seen clearly for $\eta_1 = \eta_2 = 0.4$). (b) Numerical approximation of the fitness, relative to that for a homogeneous strategy with all cells at $\varepsilon = 1$. Shown also is the perturbation expansion for $\eta_1, \eta_2 \ll 0$ (Eq. (20)).

the perturbation expansion, that continuous heterogeneity presents a fitness advantage in the neighbourhood of $\varepsilon = 0$, and a fitness disadvantage near $\varepsilon = 1$.

3 Growth in temporally correlated fluctuating environments

We now turn to randomly fluctuating environments that are temporally correlated. Specifically, we investigate phenotypic heterogeneity under environments driven by homogeneous Poisson processes, and environments in which *changes* are driven by white-noise processes and are, therefore, differentiable everywhere. An example suite of such environments is given in Fig. 5.

The dynamics are now described by

$$\frac{\partial r(\varepsilon_i, t)}{\partial t} = \underbrace{\lambda(\varepsilon_i, z(t))r(\varepsilon_i, t)}_{\text{Proliferation}} + \underbrace{\omega \hat{p}(\varepsilon)n(t) - \omega r(\varepsilon, t)}_{\text{Phenotype switching}}, \quad (21)$$

where we now explicitly denote the growth rate as a function of the environment, denoted by the stochastic process $z(t)$. We restrict our analysis to environmental fluctuations that are




Figure 5. Continuously fluctuating environment models for the growth rate of the fitness phenotype. (a) A homogeneous Poisson process drives transitions between a “growth” and a “stress” state. (b–d) Fluctuating environments with stationary distributions, shown in (e), that are either Gaussian (red) or Gamma (orange) distributed. Shown in (b,c) are Ornstein-Uhlenbeck driven environments that are relatively fast (b) or slow (c). In (d), the growth rate is driven by a second-order Ornstein-Uhlenbeck process such that the derivative of the growth rate is also continuously differentiable.

driven by a single stochastic process representing fluctuations in a single variable (e.g., nutrient or inhibitor concentration or temperature).

It is useful to rewrite Eq. (21) in terms of the total cell concentration and the phenotype distribution, denoted by $n(t)$ and $p(\varepsilon, t) = r(\varepsilon, t)/n(t)$, respectively, giving

$$\frac{d \log n(t)}{dt} = \int_0^1 \lambda(\varepsilon, z(t)) p(\varepsilon, t) d\varepsilon, \quad (22a)$$

$$\frac{\partial p(\varepsilon, t)}{\partial t} = p(\varepsilon, t) \left[\lambda(\varepsilon, z) - \mathbb{E}_{p(\varepsilon, t)}(\lambda(\varepsilon, z)) - \omega \right] + \omega \hat{p}(\varepsilon). \quad (22b)$$

From Eq. (22a), we obtain an expression for the fitness

$$\varphi = \mathbb{E}_W \mathbb{E}_{p(\varepsilon, t)}(\lambda(\varepsilon, z)), \quad (23)$$

where the outer expectation is taken with respect to the measure, denoted by W , driving fluctuations in $z(t)$.

While the phenotype *target distribution*, $\hat{p}(\varepsilon)$, is assumed to be genetically encoded and fixed, we expect $p(\varepsilon, t)$ to be correlated with $z(t)$. For example, when environmental conditions permit growth, we expect the population to be dominated by growing phenotypes, whereas when the net growth rate is low (or negative), we expect dormant phenotypes to dominate. In the limit that environmental fluctuations are slow compared to the timescales relating to phenotype switching, we expect this correlation to become perfect such that $p(\varepsilon, t) = \tilde{p}(\varepsilon, z)$. Conversely, in the limit that environmental fluctuations occur on a much faster timescale than that of phenotype switching, we expect $p(\varepsilon, t)$ to become independent of $z(t)$.

Both expectations in Eq. (23) are, in general, analytically and computationally intractable. Even for a discrete model in which $p(\varepsilon, t)$ is a discrete distribution comprising n phenotypes, numerical approaches to calculate the fitness will involve the solution of a n -dimensional Fokker-Planck equation for the joint stationary density of $z(t)$ and $p(\varepsilon, t)$. We make progress by studying the regime in which environmental fluctuations occur on a significantly slower timescale to the cellular dynamics. Thus, we consider a QSS approximation to Eq. (22b), denoted $\tilde{p}(\varepsilon, z)$, where,

on the cellular dynamics timescale (the fast timescale), the current state of the z is assumed to be fixed. The QSS approximation is given as the solution to

$$0 = \tilde{p}(\varepsilon, z) \left[\lambda(\varepsilon, z) - \mathbb{E}_{\tilde{p}(\varepsilon, z)}(\lambda(\varepsilon, z)) - \omega \right] + \omega \hat{p}(\varepsilon), \quad (24)$$

which is a root finding problem in $\mathbb{E}_{\tilde{p}(\varepsilon, z)}(\lambda(\varepsilon, z))$, the primary quantity of interest in the calculation of the fitness (Eq. (23)). We note that Eq. (24) is only well-posed for $\lambda(\varepsilon, z) - \omega \neq \mathbb{E}_{\tilde{p}(\varepsilon, z)}(\lambda(\varepsilon, z))$ for all ε . If this condition does not hold, a degeneracy will form in the solution to $\tilde{p}(\varepsilon, z)$, corresponding to a domination by a single phenotype.

3.1 Poisson model of discrete transitions

The most common theoretical and experimental model of environmental fluctuations is that of discrete transitions between a finite number of conditions (e.g., presence or absence of antibiotic [10], or transitions between nutrient sources [17]). The Poisson model [11, 19] represents a stochastic analogue of the periodically switching environments studied in [10]. Here, the environment may take one of a finite number of discrete conditions, where switches between states are governed by an inhomogeneous Poisson process. In Fig. 5a, we demonstrate such a model that switches between a “growth state” and a “stress state”, demonstrated using the net growth rate of a phenotype in state $\varepsilon = 1$ (for example, a regular proliferative bacterium). In the Poisson model, transitions between conditions are fully characterised by the expected time spent in each state, denoted by $\tau_0 := \alpha\theta^{-1}$ and $\tau_1 := (1 - \alpha)\theta^{-1}$ for the stress and growth conditions, respectively. Similarly, cell behaviour is fully characterised by functions that map the environmental state onto the growth rate for each phenotype, denoted by $f_0(\varepsilon)$ and $f_1(\varepsilon)$. In the Poisson model, θ denotes the timescale of environmental fluctuations such that $\theta \rightarrow 0$ corresponds to the QSS regime, and $\theta \rightarrow \infty$ to the fast regime. For simplicity, we assume that cells with phenotype $\varepsilon = 0$ are completely dormant, and thus always experience a net growth rate of zero.

In their seminal analysis, Kussell et al. [10] study the growth of a two-phenotype (regular and persister) bet-hedging population in a deterministic analogue of the Poisson environment, for which an exact expression for the strain fitness is available. In Fig. 6, we reproduce their analysis in a stochastically switching environment to demonstrate the fitness advantage gained in wild-type bacteria pursuing a homogeneous strategy as a function of two environmental parameters: α , the proportion of time spent in the stressed state; and θ^{-1} , the period.

Results for general θ are constructed from a numerical solution to the Fokker-Planck equation for the joint distribution of the proportion of dormant cells, $x(t)$, and the environment, $z(t)$ (see the supplementary material). A small artificial diffusion term is added to the governing advection equation for $x(t)$ to regularise the problem (diffusion coefficient $\beta = 10^{-4}$). The QSS and fast environment limits are solved exactly. In the limit that the environment changes quickly (i.e., $\theta \rightarrow \infty$), the heterogeneous strategy is only advantageous for very stressful environments (i.e., for $\alpha \rightarrow 1$), whereas in the limit that the environment changes slowly (i.e., $\theta \rightarrow 0$), the heterogeneous strategy is favoured for a range of α . Importantly, we see that the potential




Figure 6. Discrete heterogeneity in an environment with fluctuations driven by a Poisson process. Phenotype switching occurs according to a target distribution with $q = 0.1$ at rate $\omega = 1$. The environment switches between growth and stress conditions (see Fig. 5a) for a given proportion of time in stress such that the average cycle has duration θ^{-1} . The fitness advantage, relative to that of a homogeneous strategy with $q = 0$, is calculated using a numerical solution to the Fokker-Planck equation for general θ . Note that regions of advantage and disadvantage are shown on a different scale. We also calculate the QSS limit ($\theta \rightarrow 0$) and the fast limit ($\theta \rightarrow \infty$).

fitness advantage gained through a heterogeneous strategy is markedly high in comparison to the magnitude of any advantage that gained through a homogeneous strategy.

3.2 Continuously distributed growth rates

We now extend our framework to a more general model of environmental fluctuation, characterised by the long-term distribution of growth rates for each phenotype, denoted by $\mathcal{D}(\varepsilon)$. We consider that fluctuations in $z(t)$ are now governed by a scalar mean-reverting Ornstein-Uhlenbeck process

$$dz(t) = -\theta z(t)dt + \sqrt{2\theta}dW_t, \quad (25)$$

with stationary distribution $z(t) \sim \mathcal{N}(0, 1)$. The reversion strength parameter θ carries an analogous interpretation to that in the Poisson process model and characterises the relative timescale of environmental fluctuations. To relate the environment to the growth rate of each phenotype we apply the probability inverse transformation such that

$$\lambda(\varepsilon, z) = F_{\mathcal{D}(\varepsilon)}^{-1}(\Phi(z)), \quad (26)$$

where $\Phi(\cdot)$ is the distribution function for the standard Gaussian distribution, and $F_{\mathcal{D}(\varepsilon)}^{-1}(\cdot)$ is the quantile function for $\mathcal{D}(\varepsilon)$. For simplicity, we specify only $\mathcal{D}(1)$, and assume that a general $\mathcal{D}(\varepsilon)$ is chosen such that

$$\lambda(\varepsilon, z) = \varepsilon\lambda(1, z) = \varepsilon F_{\mathcal{D}(1)}^{-1}(\Phi(z)); \quad (27)$$

While other scalings in ε are possible, this choice is in some sense arbitrary (if we assume that the growth rates remain ordered within phenotype space), since the phenotype index may be itself rescaled without loss of generality.

In Fig. 5b,c, we simulate a pair of environments in both the fast ($\theta = 10$) and slow ($\theta = 0.1$) regimes. Distinguishing the environments are their marginal distributions, constructed from both a symmetric Gaussian distribution and a translated Gamma distribution with negative skewness (Fig. 5e). The mean growth rates and the proportions of time spent with a negative growth rate (i.e., in stress) are chosen to match that of the Poisson environment shown in Fig. 5a and studied in the previous section. In Fig. 5d we show realisations of a corresponding pair of environments driven by a second-order Ornstein-Uhlenbeck process, such that the growth rate is not only continuous but smooth.

3.2.1 Two discrete phenotypic states

We begin our analysis by again considering a population comprising two discrete phenotypic states, $\varepsilon \in \{0, 1\}$. For the binary model, phenotype switching can be expressed by considering the switching rates directly, denoted by a and b , such that

$$\begin{aligned}\frac{dr_0(t)}{dt} &= \lambda(0, z)r_0(t) - ar_0(t) + br_1(t), \\ \frac{dr_1(t)}{dt} &= \lambda(1, z)r_1(t) + ar_0(t) - br_1(t),\end{aligned}\tag{28}$$

where $r_\varepsilon(t)$ denotes the density of individuals in phenotypic state ε . The rates a and b are related to the target distribution and switching rate through $a = \omega(1 - q)$ and $b = \omega q$ where $q := \hat{p}(0)$. As earlier, we denote the probability mass $p(0, t) = x(t)$ such that $p(1, t) = 1 - x(t)$. The dynamics of $x(t)$ are thus governed by

$$\frac{dx(t)}{dt} = h(x, z) := b - x(t) \left(a + b + (\lambda(1, z) - \lambda(0, z))(1 - x(t)) \right),\tag{29}$$

which is coupled to the SDE governing the environment (Eq. (25)), and where $\lambda(\varepsilon, z)$ is given by Eq. (27).

For general θ , the fitness can be expressed by

$$\varphi = \int_{-\infty}^{\infty} \int_0^1 (x\lambda(0, z) + (1 - x)\lambda(1, z)) u(x, z) dx dz,\tag{30}$$

where $u(x, z)$ is the joint stationary distribution governed by a two-dimensional stationary Fokker-Planck equation which may be solved numerically; difficulties associated with the lack of a diffusion term in Eq. (29) can be overcome as for the Poisson environment by introducing a small diffusion term with coefficient $\beta = 10^{-5}$ (full details are available in the supplementary material). This inclusion provides good agreement with simulation results for large θ , however produces poor results as $\theta \rightarrow 0$. Within the QSS regime, we consider that the cellular dynamics are at steady state, such that $h(x, z) = 0$, giving, through Eq. (29), a functional relationship between $x(t)$ and $z(t)$ which we denote $x = \tilde{x}(z)$. Further substituting




Figure 7. Discrete heterogeneity in continuously fluctuating environments. (a,b) An approximate solution to the Fokker-Planck equation for the joint distribution of the $\varepsilon = 1$ growth rate and proportion of cells in the dormant state (for a target distribution with $q = 0.1$ with switching speed $\omega = 0.5$). Results shown for the (a) Gaussian and (b) Gamma environments. (c,d) We approximate the fitness as a function of θ through repeated simulation of the SDE (50,000 realisations, shown is the mean and a 95% confidence interval). We approximate the fitness using (red) the viscosity solution to the PDE and (blue, for Gaussian only) a ODE-based moment approximation to the SDE. Both approximations are valid for θ sufficiently large. Also shown is the QSS fitness (dash-dot, $\theta \rightarrow 0$); fast fitness (dot, $\theta \rightarrow \infty$); and the fitness for the homogeneous strategy (black dash).

in Eq. (27), we can obtain a readily interpretable relationship between $x(t)$ and the growth rate of the phenotype at $\varepsilon = 1$. At a QSS, we apply the law of iterated expansion to see that

$$\varphi = \int_{-\infty}^{\infty} (\tilde{x}(z)\lambda(0, z) + (1 - \tilde{x}(z))\lambda(1, z))\Phi'(z) dz, \quad (31)$$

where $\Phi'(z)$ is the density function for the standard Gaussian distribution.

We study in Fig. 7 the relationship between fitness and θ for both the Gaussian and Gamma environments introduced in Fig. 5. In Fig. 7a,b we see through numerical solution to the Fokker-Planck equation the convergence to the QSS regime; as $\theta \rightarrow 0$, the density converges to a perfect correlation on the curve given by $x = \tilde{x}(z)$. In Fig. 7c,d we compare the fitness calculated from the Fokker-Planck equation (which is valid only for θ sufficiently large) to repeated simulation of the coupled SDE system, and an approximation based on a moment closure solution to the SDE system [45]. We observe in all three methodologies a monotonic relationship between fitness and θ that connects the QSS regime as $\theta \rightarrow 0$ to the regime in which $\theta \rightarrow \infty$. Further, it follows directly from Eq. (30) that the fitness as $\theta \rightarrow \infty$, for all environments, is lower than that for a homogeneous strategy.

While we are not able to prove that the relationship between fitness and θ is monotonic, we can see that a sufficient (albeit not necessary) condition for existence of a regime where heterogeneity is advantageous is simply that the QSS fitness is greater than that for the homogeneous strategy. Two further advantages of studying the QSS regime are numerical tractability, as computation of the fitness for the two-phenotype case (Eq. (31)) involves only the numerical calculation of an integral, and tractability for a more general (potentially continuously distributed) population structure.

QSS results in Fig. 8 show the fitness of a heterogeneous strategy with a general target distribution, q , and switching rate, ω , relative to a homogeneous strategy. The most obvious result is that heterogeneity is not evolutionarily stable for large switching rates: an analogous result




Figure 8. Quasi-steady-state fitness for a discrete strategy in a continuously fluctuating Gamma environment. We calculate the slow environment limit of the fitness as a function of the target dormant proportion, q , and the switching rate, ω . Fitness shown is relative to that for the homogeneous strategy.

was observed by [16], who study responsive adaptation of bacteria to switching (Poisson-like) environments. Secondly, we see that fitness is a decreasing function of the switching rate, paradoxically suggesting that a heterogeneous population that does not permit phenotype switching (and therefore, cannot regulate heterogeneity) has the highest fitness. However, implicit in the QSS is the assumption that cellular dynamics rate parameters are sufficiently separated from that of the environment, such that $\omega \gg \theta$. Therefore, we take our results in Fig. 8 to demonstrate the existence of regimes in which a heterogeneous strategy is advantageous, rather than a demonstration of the optimal or evolutionarily stable switching strategy, which is only defined for finite θ .

3.2.2 Bivariate continuously distributed phenotypes

We next consider that an initially discretely structured population develops continuous heterogeneity such that the target distribution is given by the Gaussian mixture in Eq. (19). We again consider a perturbation expansion in the QSS fitness,

$$\varphi = \mathbb{E}_z \mathbb{E}_{\tilde{p}(\varepsilon, z)}(\lambda(\varepsilon, z)) \sim c_0 + c_1 \eta_1 + c_2 \eta_2, \quad (32)$$

such that $c_1 > 0$ indicates a fitness advantage to the population in employing continuous heterogeneity in the neighbourhood of $\varepsilon = 0$, and $c_2 > 0$ an analogous fitness advantage near $\varepsilon = 1$.

Full details of the methodology are provided in the supplementary material. In summary, we construct the perturbation expansion by considering that $\tilde{p}(\varepsilon, z)$ can be, for fixed z , represented as a mixture

$$\tilde{p}(\varepsilon, z) = \beta \tilde{p}_0(\varepsilon, z) + (1 - \beta) \tilde{p}_1(\varepsilon, z) \quad (33)$$

where the variance of each component is of order η_1 and η_2 , respectively. Substituting into




Figure 9. Bivariate continuous heterogeneity in a continuously fluctuating Gamma environment. (a) We calculate the fitness using a QSS approximation to the equilibrium population composition (blue surface) and a perturbation expansion (black dashed). (b) The perturbation coefficients c_1 and c_2 (of η_1 and η_2 , respectively) as a function of q and ω . The dash-dot line indicates the boundary between regions with negative (red) and positive (blue) c_1 . Dashed lines correspond to that in Fig. 8; regions to the left are those in which the discrete heterogeneous strategy is advantageous.

Eq. (24), taking a moment expansion [46], and assuming that higher-order central moments vanish, we obtain a leading-order approximation for the moments of $\tilde{p}_i(\varepsilon, z)$ in terms of η_1 and η_2 . The coefficients in Eq. (32) can then be calculated by numerically integrating over z .

Results in Fig. 9a show the fitness for the Gamma environment studied in Fig. 7d, for general η_1 and η_2 , in addition to the perturbation expansion for $\eta_i \ll 1$. The availability of a numerically tractable perturbation expansion allows us to establish regimes where continuous heterogeneity is advantageous by investigating the coefficients c_1 and c_2 as functions of the strategy (q, ω) . In Fig. 9b we show that c_2 is, for the environment considered, always negative (i.e., it is never evolutionarily stable to present with continuous heterogeneity around the phenotype with the highest average fitness), while c_1 is negative for all but sufficiently large ω and q . Further investigation reveals that $c_1 > 0$ only in regions where the heterogeneous strategy is itself disadvantageous; thus, continuous heterogeneity near $\varepsilon = 0$ gives a fitness advantage only in the case that the existence of a phenotype at $\varepsilon = 0$ is itself disadvantageous.




Figure 10. Continuously fluctuating optimal phenotype. (a) The growth rate of each phenotype is given by a Gaussian centered at the current value of the environment variable $z(t)$, modelled as an Ornstein-Uhlenbeck process with mean $\varepsilon = 0.5$. Time shown on the left axis, with growth rate at $t = 9$ h and $t = 10$ h is shown on the right axis. The phenotype $\varepsilon = 0.5$ is the optimal phenotype if a homogeneous strategy is employed. (b) Fitness (relative to a homogeneous strategy) calculated for a continuous target distribution with variance η^2 , calculated numerically (blue) and using a perturbation expansion (black dashed). Other parameters are given by $\xi = 0.5$, $k = 1$, $\sigma = 0.1$.

3.3 Fluctuations in the fittest phenotype

We have, thusfar, restricted our study to growth rates that are monotonic in the phenotype index. Such environments are typically considered in the context of bet-hedging: the single optimal phenotype is often on the edge of the phenotype space, and the evolutionary stable strategy involves regulating a proportion of cells at the two extrema. We now extend this analysis to a scenario in which the environment induces changes in the optimal phenotype itself to represent, for example, temperature specialisation of bacteria in environments in which the temperature fluctuates continuously [47].

We again assume that environmental fluctuations are driven by an Ornstein-Uhlenbeck process (Eq. (25)), relating the environment to the growth rate more directly through

$$\lambda(\varepsilon, w) = k \exp\left(\frac{-(\varepsilon - w)^2}{\xi^2}\right), \quad (34)$$

where $w = \frac{1}{2} + \sigma z \sim \mathcal{N}(0.5, \sigma)$ describes the (fluctuating) location of the optimal phenotype. In Fig. 10a, we show a realisation of the environment, demonstrating the time-dependent coupling between the environment and the growth rate of every phenotype. While many other choices of $\lambda(\varepsilon, z)$ are reasonable, our specific choice enables us to make analytical progress toward a perturbation solution for the fitness in the case that the population develops continuous heterogeneity around an otherwise optimal singular phenotype. We also note that, as the governing equation for the population composition depends only on a difference in growth rates, we can, without loss of generality, restrict our study to functions such that $\min \lambda(\varepsilon, w) = 0$.

Due to symmetry, the single optimal phenotype (i.e., in a homogeneous strategy) is located at $\varepsilon^* = \frac{1}{2}$. We again consider that the population undergoes mutations that yield a continuously distributed target distribution $\hat{p}(\varepsilon) = \mathcal{N}(\varepsilon^*, \eta)$. Substituting into Eq. (24), assuming that $\eta \ll 1$,

and integrating over z , we find that the fitness is given by

$$\varphi \sim \frac{k\xi}{\sqrt{\xi^2 + 2\sigma^2}} + \left(\frac{4k^2\sigma^2}{\xi\omega(\xi^2 + 4\sigma^2)^{3/2}} - \frac{k\xi}{(\xi^2 + 2\sigma^2)^{3/2}} \right) \eta^2 + \mathcal{O}(\eta^3). \quad (35)$$

For a specific choice of parameters, we demonstrate in Fig. 10b that the expansion conforms with a numerical solution generated for general η . Both the expansion and the numerical results clearly demonstrate the existence of a regime in which continuous heterogeneity is advantageous. While the relationship between fitness and η is clearly not always monotonic, the positivity of the correction term in Eq. (35) gives a sufficient condition for the existence of an evolutionarily stable heterogeneous strategy. Furthermore, we can see that heterogeneity is immediately (i.e., for small η) disadvantageous in cases where either $\omega \gg 0$, if $\sigma \rightarrow 0$, or if $\xi \rightarrow 0$.

4 Discussion

Many biological systems regulate and exploit phenotypic heterogeneity to achieve functional aims [1, 26]. The phenomenon of bet-hedging, in particular, is highly studied, particularly through the lens of a discrete set of highly specialised phenotypes [6, 10, 48]. Ecological theory commonly describes diversification strategies as an evolutionary response to environmental uncertainty [19]. In this context, experimental and mathematical models of environmental fluctuations are predominantly characterised by an effective growth rate that switches either periodically or stochastically between a finite set of conditions: a growth-promoting environment, and a stressful environment, for example. More general and arguably more realistic models of environment are comparatively less studied. Our goal in the present work is to provide mathematical theory to demonstrate the existence of regimes where phenotypic heterogeneity is advantageous as an evolutionary response to a set of generalisable stochastic environments.

4.1 Environments in which discrete heterogeneity is advantageous

We first establish regimes in which the standard model of bet-hedging—that of two specialised phenotypes—is advantageous in both temporally uncorrelated (i.e., white-noise) and correlated (i.e., Poisson and Ornstein-Uhlenbeck) environments. For both environment classes, we make biologically relevant simplifying assumptions to render the calculation of population fitness analytically tractable. In white-noise environments, we consider that a population is able to regulate its phenotype composition. This simplification allows us to present a set of analytical results (Eqs. (8) to (10)) that reveal parameter regimes in which it is evolutionary stable to regulate a proportion of near-dormant cells that present individually a lower mean growth rate. These regimes become more restrictive as the correlation between the inter-phenotype growth rates tends to unity. Conversely, regimes become less restrictive if growth rates are un- or inversely-correlated; representative, for example, of regimes in which phenotypes specialise to different nutrient sources that fluctuate independently or inversely (the latter arising where a fixed *total* amount of nutrient is available).

Environments that are temporally correlated motivate a different set of biologically relevant

simplifications. Specifically, we focus for analytical tractability on the scenario that environmental changes occur on a much slower timescale to that of the cellular dynamics such that the subpopulation structure is at a QSS. Demonstrating the existence of regimes under QSS is thus a sufficient (albeit not necessary) condition for the existence of such regimes in all timescales. The second advantage of this simplification is that we need only characterise the environment with an equilibrium distribution of growth rates. Our initial results reveal a possible balance between the timescales of phenotype switching and environmental changes. In particular, results in Fig. 8 show that the heterogeneous strategy becomes disadvantageous in populations that present fast switching between phenotypes. Advantages are gained, therefore, in regimes where phenotype switching is sufficiently slow, but not as slow as that of the environmental timescales. These somewhat paradoxical results are also seen in [16], and can be explained by considering the antithetical scenario that environmental fluctuations occur very quickly. In such a regime, the environment and population structure will be statistically independent (which is also the case if switching is sufficiently fast), and it is straightforward to see from Eq. (31) that fitness will be lower than in the optimal homogeneous strategy.

4.2 Environments in which bimodal continuous heterogeneity is advantageous

While the theoretical study of bet-hedging strategies that comprise investment in a discrete set of phenotypes is extensive, there is comparatively very little theory to allow study of strategies that account for diversification strategies that are continuous throughout phenotype space. While extraneous factors, such as gene expression noise, mean that within-subpopulation variability is likely omnipresent, it is routine for theoretical models of bet-hedging to assume that cells within each subpopulation (e.g., dormant or proliferative) behave identically. Our most important result is a demonstration of regimes in which such subpopulation heterogeneity is evolutionary stable. In particular, we apply a perturbative approach to study the effect of small within-subpopulation variance on the overall population fitness in two main scenarios: first, that an otherwise-optimal homogeneous population develops heterogeneity (unimodal), and secondly that continuous heterogeneity develops in an initially two-phenotype population (bimodal). For environments driven by white-noise processes, we establish clear parameter regimes in which continuous heterogeneity is advantageous (Eqs. (15) to (18)). As our asymptotic results are dependent only on the local behaviour of the growth rate variance and correlations, we expect our findings to hold for all differentiable choices of these functions. Results for the bimodal scenario are less analytically interpretable, however we demonstrate in Fig. 4 that, for a particular choice of parameters, only continuous heterogeneity in the dormant phenotype is advantageous.

In temporally correlated environments, analogous perturbative correction terms at QSS are analytically intractable. We do, however, derive semi-analytical expressions that allow numerical computation these corrections terms, and therefore allow us to establish regimes in which continuous heterogeneity may be advantageous. For a particular choice of environment, results in Fig. 9 show that such subpopulation variance is always immediately disadvantageous. This does not rule out a heterogeneous strategy that is advantageous for a significantly high

variance, as seen in Fig. 3c for an environment driven by white-noise. These results have further parallels to our findings for temporally uncorrelated environments: semi-analytical expressions for the correction terms (supplementary material) suggest that a necessary condition for a positive correction at the proliferative state may be a negative correction at the dormant state. An aspect of continuous heterogeneity that we have not considered is one that arises in some sense extrinsically out of gene expression noise. That is, a discretely structured population may be evolutionarily favoured *in theory*, however can only be regulated at the cost of some level of subpopulation variance. Further study using our perturbative approach is likely to elucidate the cost-benefit trade-off for bet-hedging in discretely structured populations.

4.3 Unimodal continuous heterogeneity

The classical view of bet-hedging is risk-spreading: the main body of our work has involved environments that present risk to a homogeneous strategy through periods of time that experience negative growth rates. Our final result (Fig. 10), however, demonstrates that continuous unimodal heterogeneity can itself be advantageous in the absence of an explicit stressor if the environment drives fluctuations in the optimal phenotype. Again applying the perturbative approach, we provide for a specific growth rate model an analytically tractable expression for the correction term (Eq. (35)), providing a sufficient condition for an evolutionary stable continuous subpopulation structure. One motivation for this scenario may be that of temperature specialisation in populations of bacteria subject to relatively slow fluctuations in temperature. The optimal homogeneous strategy is to specialise to the mean temperature, while our results show that the population can increase its fitness by introducing small amounts of phenotypic variability in which isogenic cells specialise to different temperatures.

4.4 Competition and high-dimensional phenotype spaces

We restrict our analysis to a simple cellular dynamics model comprising heritable traits and an exponentially growing population; the latter motivated by serial-dilution or continuous-culture experiments where the population density is kept sufficiently low that cell-to-cell competition is negligible. A key theoretical and experimental challenge is that bet-hedging strategies are likely to vary significantly between growth and stationary phase. A body of relatively recent work explores fluctuations in environmental carrying capacity (rather than growth rate), and could, in principle, be extended to allow for continuous variability and more general environments [20,21,24,49]. The formulation of a more biologically realistic model of subpopulation regulation is likely to be driven by future experimentation. In an exponentially growing population within continuous culture, it is clear that active phenotypic switching is required to avoid extinction of subpopulations that are for the majority of the time less fit. In the cancer modelling literature, it is typical for such transitions between phenotypes to be local and modelled as diffusion through a (potentially constrained) phenotype space [37]. Our modelling approach provides a framework in which to experimentally test subpopulation-regulation models, by predicting structures for arbitrarily (albeit slowly) fluctuating environments. In most cases, however, a phenotype is

likely to result from the expression of a large number of genes, and may manifest as a high-dimensional distribution of behaviours [50]. In the QSS regime, we anticipate that our theory can be extended to a two-dimensional phenotype space and to two-dimensional environmental fluctuations.

5 Conclusion

It is fast becoming understood that heterogeneity is not only ubiquitous to biology, but plays a key role in the function of many biological systems. We elucidate this function in populations exposed to a range of stochastically fluctuating environments. Importantly, we demonstrate regimes in which regulation of both discrete and continuous heterogeneity provides a fitness advantage to populations. Overall, our work enables future theoretical and experimental work to investigate and eventually exploit phenotypic heterogeneity in biologically realistic environments.

Data availability

Code used to produce the results is available at <https://github.com/ap-browning/heterogeneity>.

Acknowledgements

APB thanks the Mathematical Institute, University of Oxford, for a Hooke Research Fellowship. SH was funded by Wenner-Gren Stiftelserna/the Wenner-Gren Foundations (WGF2022-0044) and the Kjell och Märta Beijer Foundation.

References

- [1] Elsasser WM. 1984 Outline of a theory of cellular heterogeneity. *Proceedings of the National Academy of Sciences* **81**, 5126–5129.
- [2] Wilkinson DJ. 2009 Stochastic modelling for quantitative description of heterogeneous biological systems. *Nature Reviews Genetics* **10**, 122–133.
- [3] Ackermann M. 2015 A functional perspective on phenotypic heterogeneity in microorganisms. *Nature Reviews Microbiology* **13**, 497–508.
- [4] Gough A, Stern AM, Maier J, Lezon T, Shun TY, Chennubhotla C, Schurdak ME, Haney SA, Taylor DL. 2016 Biologically relevant heterogeneity: metrics and practical insights. *SLAS Discovery* **22**, 213–237.
- [5] Bigger JW. 1944 Treatment of staphylococcal infections with penicillin by intermittent sterilisation. *The Lancet* **244**, 497–500.
- [6] Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. 2004 Bacterial persistence as a phenotypic switch. *Science* **305**, 1622–1625.
- [7] de Jong IG, Haccou P, Kuipers OP. 2011 Bet hedging or not? A guide to proper classification of microbial survival strategies. *BioEssays* **33**, 215–223.
- [8] Turke AB, Zejnullahu K, Wu YL, Song Y, Dias-Santagata D, Lifshits E, Toschi L, Rogers A, Mok T, Sequist L, Lindeman NI, Murphy C, Akhavanfard S, Yeap BY, Xiao Y, Capelletti M, Iafrate AJ, Lee C, Christensen

- JG, Engelman JA, Jänne PA. 2010 Preexistence and clonal selection of MET amplification in EGFR mutant NSCLC. *Cancer Cell* **17**, 77–88.
- [9] Sun X, Wu LF, Altschuler SJ, Hata AN. 2024 Targeting therapy-persistent residual disease. *Nature Cancer* **5**, 1298–1304.
- [10] Kussell E, Kishony R, Balaban NQ, Leibler S. 2005 Bacterial persistence: a model of survival in changing environments. *Genetics* **169**, 1807–1814.
- [11] Wolf DM, Vazirani VV, Arkin AP. 2005 Diversity in times of adversity: probabilistic strategies in microbial survival games. *Journal of Theoretical Biology* **234**, 227–253.
- [12] Thakur B, Meyer-Ortmanns H. 2023 Controlling the mean time to extinction in populations of bacteria. *Entropy* **25**, 755.
- [13] Browning AP, Sharp JA, Mapder T, Baker CM, Burrage K, Simpson MJ. 2021 Persistence as an optimal hedging strategy. *Biophysical Journal* **120**, 133–142.
- [14] Beaumont HJE, Gallie J, Kost C, Ferguson GC, Rainey PB. 2009 Experimental evolution of bet hedging. *Nature* **462**, 90–93.
- [15] Van den Bergh B, Michiels JE, Wenseleers T, Windels EM, Boer PV, Kestemont D, Meester LD, Verstrepen KJ, Verstraeten N, Fauvart M, Michiels J. 2016 Frequency of antibiotic application drives rapid evolutionary adaptation of *Escherichia coli* persistence. *Nature Microbiology* **1**, 16020.
- [16] Thattai M, van Oudenaarden A. 2004 Stochastic gene expression in fluctuating environments. *Genetics* **167**, 523–530.
- [17] Lambert G, Kussell E, Kussel E. 2014 Memory and fitness optimization of bacteria under fluctuating environments. *PLoS Genetics* **10**, e1004556.
- [18] Letten AD, Ludington WB. 2023 Pulsed, continuous or somewhere in between? Resource dynamics matter in the optimisation of microbial communities. *The ISME Journal* **17**, 641–644.
- [19] Müller J, Hense B, Fuchs T, Utz M, Pötzsche C. 2013 Bet-hedging in stochastically switching environments. *Journal of Theoretical Biology* **336**, 144–157.
- [20] Taitelbaum A, West R, Assaf M, Mabilia M. 2020 Population dynamics in a changing environment: random versus periodic switching. *Physical Review Letters* **125**, 048105.
- [21] Hernández-Navarro L, Asker M, Rucklidge AM, Mabilia M. 2023 Coupled environmental and demographic fluctuations shape the evolution of cooperative antimicrobial resistance. *Journal of the Royal Society Interface* **20**, 20230393.
- [22] Hidalgo J, Pigolotti S, Muñoz MA. 2015 Stochasticity enhances the gaining of bet-hedging strategies in contact-process-like dynamics. *Physical Review E* **91**, 032114.
- [23] Liu M, Rubenstein DR, Liu WC, Shen SF. 2019 A continuum of biological adaptations to environmental fluctuation. *Proceedings of the Royal Society B* **286**, 20191623.
- [24] Taitelbaum A, West R, Mabilia M, Assaf M. 2023 Evolutionary dynamics in a varying environment: Continuous versus discrete noise. *Physical Review Research* **5**, L022004.
- [25] Sireci M, Muñoz MA, Grilli J. 2023 Environmental fluctuations explain the universal decay of species-abundance correlations with phylogenetic distance. *Proceedings of the National Academy of Sciences* **120**, e2217144120.
- [26] Kussell E, Leibler S. 2005 Phenotypic diversity, population growth, and information in fluctuating environments. *Science* **309**, 2075–2078.
- [27] Williams PD, Hastings A. 2011 Paradoxical persistence through mixed-system dynamics: towards a unified perspective of reversal behaviours in evolutionary ecology. *Proceedings of the Royal Society B: Biological Sciences* **278**, 1281–1290.
- [28] Day T. 2016 Interpreting phenotypic antibiotic tolerance and persister cells as evolution via epigenetic inheritance. *Molecular Ecology* **25**, 1869–1882.

- [29] Irving SE, Choudhury NR, Corrigan RM. 2021 The stringent response and physiological roles of (pp)pGpp in bacteria. *Nature Reviews Microbiology* **19**, 256–271.
- [30] de Groot DH, Tjalma AJ, Bruggeman FJ, van Nimwegen E. 2023 Effective bet-hedging through growth rate dependent stability. *Proceedings of the National Academy of Sciences* **120**, e2211091120.
- [31] Verstraete L, Van den Bergh B, Verstraeten N, Michiels J. 2021 Ecology and evolution of antibiotic persistence. *Trends in Microbiology* **30**, 466–479.
- [32] Fridman O, Goldberg A, Ronin I, Shores N, Balaban NQ. 2014 Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. *Nature* **513**, 418–421.
- [33] Svardal H, Rueffler C, Hermisson J. 2011 Comparing environmental and genetic variance as adaptive response to fluctuating selection. *Evolution* **65**, 2492–2513.
- [34] Levien E, Kondev J, Amir A. 2020 The interplay of phenotypic variability and fitness in finite microbial populations. *Journal of the Royal Society Interface* **17**, 20190827.
- [35] Mateu JC, Sireci M, Muñoz MA. 2021 Phenotypic-dependent variability and the emergence of tolerance in bacterial populations. *PLoS Computational Biology* **17**, e1009417.
- [36] Kerr R, Jabbari S, Blair JMA, Johnston IG. 2022 Dynamic Boolean modelling reveals the influence of energy supply on bacterial efflux pump expression. *Journal of the Royal Society Interface* **19**, 20210771.
- [37] Chisholm RH, Lorenzi T, Lorz A, Larsen AK, de Almeida LN, Escargueil A, Clairambault J. 2015 Emergence of drug tolerance in cancer cell populations: an evolutionary outcome of selection, nongenetic instability, and stress-induced adaptation. *Cancer Research* **75**, 930–939.
- [38] Browning AP, Crossley RM, Villa C, Maini PK, Jenner AL, Cassidy T, Hamis S. 2024 Identifiability of heterogeneous phenotype adaptation from low-cell-count experiments and a stochastic model. *bioRxiv* p. 2024.08.19.608540.
- [39] Bod’ová K, Szép E, Barton NH. 2021 Dynamic maximum entropy provides accurate approximation of structured population dynamics. *PLoS Computational Biology* **17**, e1009661.
- [40] Garcia-Bernardo J, Dunlop M. 2015 Noise and low-level dynamics can coordinate multicomponent bet hedging mechanisms. *Biophysical Journal* **108**, 184–193.
- [41] Tuljapurkar S. 1990 *Population Dynamics in Variable Environments*. Lecture Notes in Biomathematics. Springer-Verlag.
- [42] Metz JAJ. 2008 Fitness. In Jorgensen S, Fath BD, editors, *Encyclopedia of Ecology* vol. 2 pp. 1599–1612.
- [43] Bukkuri A, Pienta KJ, Hockett I, Austin RH, Hammarlund EU, Amend SR, Brown JS. 2023 Modeling cancer’s ecological and evolutionary dynamics. *Medical Oncology* **40**, 109.
- [44] Mononen T, Kuosmanen T, Cairns J, Mustonen V. 2023 Understanding cellular growth strategies via optimal control. *Journal of the Royal Society Interface* **20**, 20220744.
- [45] Sukys A, Grima R. 2021 MomentClosure.jl: automated moment closure approximations in Julia. *Bioinformatics* **38**, 289–290.
- [46] Browning AP, Drovandi C, Turner IW, Jenner AL, Simpson MJ. 2022 Efficient inference and identifiability analysis for differential equation models with random parameters. *PLoS Computational Biology* **18**, e1010734.
- [47] Lambros M, Pechuan-Jorge X, Biro D, Ye K, Bergman A. 2021 Emerging adaptive strategies under temperature fluctuations in a laboratory evolution experiment of *Escherichia coli*. *Frontiers in Microbiology* **12**, 724982.
- [48] Jolly MK, Kulkarni P, Weninger K, Orban J, Levine H. 2018 Phenotypic plasticity, bet-hedging, and androgen independence in prostate cancer: role of non-genetic heterogeneity. *Frontiers in Oncology* **8**, 50.
- [49] Wienand K, Frey E, Mobilia M. 2018 Eco-evolutionary dynamics of a population with randomly switching carrying capacity. *Journal of The Royal Society Interface* **15**, 20180343.
- [50] Ispolatov MD, Ispolatov I. 2017 Diversity and coevolutionary dynamics in high-dimensional phenotype spaces. *The American Naturalist* **189**, 105–120.

Supplementary material for “Phenotypic heterogeneity in temporally fluctuating environments”

Alexander P Browning¹ and Sara Hamis²

¹*Mathematical Institute, University of Oxford, Oxford, United Kingdom*

²*Department of Information Technology, Uppsala University, Uppsala, Sweden*

December 17, 2024

Contents

S1 White-noise environments	2
S1.1 Binary model	2
S1.2 Binary model (finite switching speed)	2
S1.3 Unimodal continuous heterogeneity	3
S1.4 Bimodal continuous heterogeneity	4
S2 Poisson environment	4
S2.1 Binary model (Fokker-Planck)	4
S2.2 Binary model (QSS and fast regimes)	5
S3 Continuously fluctuating environments	5
S3.1 Binary model (Fokker-Planck)	5
S3.2 Bimodal continuous fluctuations	6
S3.3 Fluctuations in the fittest phenotype	8

¹Corresponding author: browning@maths.ox.ac.uk

S1 White-noise environments

S1.1 Binary model

The growth rates and target distribution are given by

$$\lambda(\varepsilon) = \begin{cases} \mu_1 + \sigma_1 \xi_1, & \varepsilon = 1, \\ \mu_2 + \sigma_2 \rho \xi_1 + \sigma_2 \sqrt{1 - \rho^2} \xi_2, & \varepsilon = 0, \end{cases}$$

$$p(\varepsilon) = \begin{cases} 1 - q & \varepsilon = 1, \\ q & \varepsilon = 0. \end{cases}$$

Thus, the fitness is given by

$$\begin{aligned} \varphi &= \mathbb{E}_{p(\varepsilon)}(\mu(\varepsilon)) - \frac{1}{2} \sum_{i=1}^2 \mathbb{E}_{p(\varepsilon)}^2(s_i(\varepsilon)), \\ &= \mu_1 - \frac{\sigma_1^2}{2} + q(\mu_2 - \mu_1 + \sigma_1^2 - \rho\sigma_1\sigma_2) + q^2 \left(\rho\sigma_1\sigma_2 - \frac{\sigma_1^2 + \sigma_2^2}{2} \right), \end{aligned}$$

which has a critical point at

$$q^* = \frac{\mu_2 - \mu_1 + \sigma_1^2 - \rho\sigma_1\sigma_2}{\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2}.$$

S1.2 Binary model (finite switching speed)

We first apply Itô's lemma to obtain an SDE for the proportion of the population in the dormant phenotype, $p(0, t) := x(t)$, given by

$$\begin{aligned} dx &= f_x(x) dt + g_x(x) dW, \\ f_x(x) &= b - ax - bx - x(1-x)(\mu_1 - \mu_2 - (1-x)\sigma_1^2 + (1-2x)\rho\sigma_1\sigma_2 + x\sigma_2^2), \\ g_x(x) &= x(1-x)\sqrt{\sigma_1^2 - 2\rho\sigma_1\sigma_2 + \sigma_2^2}, \end{aligned}$$

for $a = \omega(1 - q)$ and $b = \omega q$.

The Fokker-Planck equation governing the equilibrium distribution of x is given by

$$0 = -\frac{\partial(f_x(x)u(x))}{\partial x} + \frac{1}{2} \frac{\partial^2(g_x^2(x)u(x))}{\partial x^2},$$

subject to the usual boundary conditions that $u(0) = u(1) = 0$, and that $\int_0^1 u(x) dx = 1$.

Integrating once we obtain

$$0 = -f_x(x)u(x) + \frac{1}{2} \frac{\partial(g_x^2(x)u(x))}{\partial x},$$

which we solve using integrating factor to arrive at

$$u(x) \propto \frac{1}{g^2(x)} \exp\left(2 \int_0^x \frac{f_x(s)}{g_x^2(s)} ds\right). \quad (\text{S2})$$

The integral in the exponent of eq. (S2) can be computed analytically, yielding

$$u(x) = \frac{C}{g^2(x)} e^{\alpha(x)},$$

where

$$\alpha(x) = \frac{2x(\mu_1 - \mu_2 - \sigma_1^2 + \rho\sigma_1\sigma_2) + x^2(\sigma_1^2 - 2\rho\sigma_1\sigma_2 + \sigma_2^2) - 2a \log(1-x) - 2b \log(x)}{\sqrt{\sigma_1^2 - 2\rho\sigma_1\sigma_2 + \sigma_2^2}},$$

and where C is a normalisation constant to be determined using quadrature to ensure that $u(x)$ is a valid probability density function.

S1.3 Unimodal continuous heterogeneity

We here assume that $p(\varepsilon) = \mathcal{N}_{(0,1)}(\varepsilon^*, \eta^2)$ for $\eta \ll 1$ and wish to calculate the fitness

$$\varphi = \mathbb{E}_{p(\varepsilon)}(\mu(\varepsilon)) - \frac{1}{2} \sum_{i=1}^2 \mathbb{E}_{p(\varepsilon)}^2(s_i(\varepsilon)), \quad (\text{S3})$$

in the case that $\mu(\varepsilon) = \varepsilon$ and where

$$s_1(\varepsilon) = \rho(\varepsilon - \varepsilon^*)\sigma(\varepsilon), \quad (\text{S4a})$$

$$s_2(\varepsilon) = \sigma(\varepsilon)\sqrt{1 - \rho^2(\varepsilon - \varepsilon^*)}. \quad (\text{S4b})$$

For η sufficiently small, we can consider firstly that s_1 and s_2 are well approximated in the vicinity of ε^* by their respective Taylor expansions, and secondly that $p(\varepsilon)$ is very well approximated by an untruncated distribution (in the case that $\varepsilon^* = 1$, an untruncated half Gaussian distribution). Together, these assumptions yield analytical expressions for the respective expectations in eq. (S3) (see Mathematica code).

Care must be taken for the case that ε^* is on the interior of the domain and in which $\rho(\cdot)$ is not differentiable: in this case, we note that $\rho(\cdot)$ is symmetric, and thus consider a piecewise formulation of eq. (S4).

$$s_1(\varepsilon) = \begin{cases} \rho_-(\varepsilon - \varepsilon^*)\sigma(\varepsilon) & \varepsilon \leq \varepsilon^*, \\ \rho_-(\varepsilon^* - \varepsilon)\sigma(\varepsilon) & \varepsilon > \varepsilon^*, \end{cases},$$

$$s_2(\varepsilon) = \begin{cases} \sigma(\varepsilon)\sqrt{1 - \rho_-^2(\varepsilon - \varepsilon^*)} & \varepsilon \leq \varepsilon^*, \\ \sigma(\varepsilon)\sqrt{1 - \rho_-^2(\varepsilon^* - \varepsilon)} & \varepsilon > \varepsilon^*, \end{cases},$$

where $\rho_-(\Delta)$ gives the correlation function for $\varepsilon \leq \varepsilon^*$.

S1.4 Bimodal continuous heterogeneity

We proceed in a similar manner to in Section S1.3, however for a slightly more general model without the same restrictions on $\mu(\varepsilon)$ (to enable a direct comparison to the results in Fig. 2 of the main document), although here we only present the expansion for a form of $\rho(\cdot)$ that is differentiable at $\Delta = 0$.

The population composition is given by the mixture

$$p(\varepsilon) = q\mathcal{N}_{(0,1)}(0, \eta_1) + (1 - q)\mathcal{N}_{(0,1)}(1, \eta_2),$$

such that the population tends to a discrete binary structure for $\eta_1, \eta_2 \rightarrow 0$. The required expectations in the fitness (given by eq. (S3)) are calculated for each component of the mixture separately under the assumption that, for η_1 sufficiently small, the density of $\mathcal{N}_{(0,1)}(0, \eta_1)$ is approximately that of a half Gaussian distribution (and similar for the distribution near $\varepsilon = 1$ for $\eta_2 \ll 1$). Full details are given in the Mathematica code.

S2 Poisson environment

S2.1 Binary model (Fokker-Planck)

We again denote the current proportion of cells in the dormant state ($\varepsilon = 0$) by $x(t)$, with dynamics given by

$$\frac{dx}{dt} = f_x(x, \lambda(z)) = b - x(a - b - \lambda(z)(1 - x)) \quad (\text{S6})$$

with $a = \omega(1 - q)$, $b = \omega q$, and where $\lambda(z)$ gives the growth rate of cells at $\varepsilon = 1$ (cells at $\varepsilon = 0$ are assumed to be completely dormant). The environment is given by a homogeneous Poisson process with rates given by $s_1 = \theta/(1 - \alpha)$ and $s_2 = \theta/\alpha$ for the growing and stressful states respectively (in which the growth rates are $\lambda(1) = \lambda_1$ and $\lambda(-1) = \lambda_2$, respectively).

The joint stationary probability density $u(x, \lambda)$ is governed by the coupled system of partial differential equations

$$\begin{aligned} \frac{\partial}{\partial x} \left(f_x(x) u(x, \lambda_1) \right) &= -s_1 u(x, \lambda_1) + s_2 u(x, \lambda_2), \\ \frac{\partial}{\partial x} \left(f_x(x) u(x, \lambda_2) \right) &= s_1 u(x, \lambda_1) - s_2 u(x, \lambda_2), \end{aligned} \quad (\text{S7})$$

subject to the constraint that $\int_0^1 u(x, \lambda_1) + u(x, \lambda_2) dx = 1$.

The lack of a diffusion term in eq. (S7) renders the problem numerically unstable. We thus regularise the problem by introducing a small diffusion term such that the dynamics of $u(x, \lambda(z))$ are given, approximately, by

$$\begin{aligned} \frac{\partial}{\partial x} \left(f_x(x) u(x, \lambda_1) \right) &= -s_1 u(x, \lambda_1) + s_2 u(x, \lambda_2) + \beta \frac{\partial^2 u(x, \lambda_1)}{\partial x^2}, \\ \frac{\partial}{\partial x} \left(f_x(x) u(x, \lambda_2) \right) &= s_1 u(x, \lambda_1) - s_2 u(x, \lambda_2) + \beta \frac{\partial^2 u(x, \lambda_2)}{\partial x^2}, \end{aligned} \quad (\text{S8})$$

where β is small (specifically, we set $\beta = 10^{-4}$). We then solve eq. (S8) numerically using a finite volume discretisation alongside a linear solver¹. The fitness is then given by

$$\varphi = \sum_{i=1}^2 \int_0^1 \lambda_i (1-x) u(x, \lambda_i) dx,$$

which we calculate using quadrature¹.

S2.2 Binary model (QSS and fast regimes)

At QSS, we assume in eq. (S6) that $dx/dt = 0$. This yields an expression for x in terms of $\lambda(z)$, denoted $\tilde{x}(\lambda)$:

$$\tilde{x}(\lambda) = \frac{-a + b + \lambda + \sqrt{4b\lambda + (-a + b + \lambda)^2}}{2\lambda}.$$

The fitness is then given by

$$\varphi = \sum_{i=1}^2 (1 - \tilde{x}(\lambda_i)) \lambda_i u(\lambda_i),$$

where $u(\lambda_i)$, the probability mass function for the environment, is given simply by

$$u(\lambda) = \begin{cases} 1 - \alpha & \lambda = \lambda_1, \\ \alpha & \lambda = \lambda_2. \end{cases}$$

In the fast regime, x and λ are uncorrelated; thus, we take λ to be at its mean value, given by

$$\bar{\lambda} = \sum_{i=1}^2 \lambda_i u(\lambda_i),$$

such that x is given by $\tilde{x}(\bar{\lambda})$. The fitness is given simply by

$$\varphi = (1 - \tilde{x}(\bar{\lambda})) \bar{\lambda}.$$

S3 Continuously fluctuating environments

S3.1 Binary model (Fokker-Planck)

The coupled dynamics of $x(t)$ and $z(t)$ are given by the system of SDEs

$$\begin{aligned} dx(t) &= f_x(x, \lambda(z)) dt + \sqrt{2\beta} dW_1, \\ dz(t) &= -\theta z(t) dt + \sqrt{2\theta} dW_2, \end{aligned} \tag{S9}$$

where we have added an artificial diffusion term of magnitude β to regularise the problem, and where

$$f_x(x, \lambda(z)) = b - x \left(a - b - (\lambda(1, z) - \lambda(0, z))(1 - x) \right).$$

¹library/poisson/solve_binary_poisson.jl

The joint stationary distribution, $u(x, z)$, is given by the solution to the two-dimensional Fokker-Planck equation

$$\frac{\partial}{\partial x}(f_x(x, z)u(x, z)) - \theta \frac{\partial}{\partial z}(zu(x, z)) = \beta \frac{\partial^2 u(x, z)}{\partial x^2} + \theta \frac{\partial^2 u(x, z)}{\partial z^2}, \quad (\text{S10})$$

subject to the condition that $u(x, z)$ must be a valid probability density function.

We solve eq. (S10) numerically using a linear finite volume discretisation, combined with an iterative solve for the resultant fast linear system (GMRES)². The fitness is then given by

$$\varphi = \int_{-\infty}^{\infty} \int_0^1 \left(x\lambda(0, z) + (1-x)\lambda(1, z) \right) u(x, z) dx dz, \quad (\text{S11})$$

which we calculate numerically using quadrature².

S3.2 Bimodal continuous fluctuations

The target distribution is given now by the mixture

$$\hat{p}(\varepsilon) = q\hat{p}_1(\varepsilon) + (1-q)\hat{p}_2(\varepsilon),$$

where $\hat{p}_1(\varepsilon)$ is the density function for the truncated Gaussian $\mathcal{N}_{(0,1)}(0, \eta_1)$, and $\hat{p}_2(\varepsilon)$ that for $\mathcal{N}_{(0,1)}(1, \eta_2)$.

At QSS, the population composition distribution is given by the solution to

$$0 = \tilde{p}(\varepsilon, z) \left[\lambda(\varepsilon, z) - \mathbb{E}_{\tilde{p}(\varepsilon, z)}(\lambda(\varepsilon, z)) - \omega \right] + \omega \hat{p}(\varepsilon), \quad (\text{S12})$$

in which we are primarily interested in (for a fixed z , or equivalently, a fixed λ) the quantity $\mathbb{E}_{\tilde{p}(\varepsilon, z)}(\lambda(\varepsilon, z))$. Given the structure of the target distribution as a mixture, our approach is to assume that $\tilde{p}(\varepsilon, z)$ is also given by the mixture

$$\tilde{p} = \gamma \tilde{p}_1 + (1-\gamma) \tilde{p}_2, \quad (\text{S13})$$

where γ is also to be determined, and where, for brevity, we drop the notation giving function arguments of density functions explicitly. By definition, we assume that the mass of \tilde{p}_1 is concentrated around $\varepsilon = 0$ (specifically, within the support of p_1), and similar for \tilde{p}_2 near $\varepsilon = 1$.

Before proceeding, we define the following quantities

$$\begin{aligned} E_1 &= \mathbb{E}_{\tilde{p}_1}(\varepsilon) \sim \mathcal{O}(\eta_1), & E_2 &= \mathbb{E}_{\tilde{p}_2}(1-\varepsilon) \sim \mathcal{O}(\eta_2), \\ V_1 &= \mathbb{E}_{\tilde{p}_1}(\varepsilon^2) \sim \mathcal{O}(\eta_1^2), & V_2 &= \mathbb{E}_{\tilde{p}_2}((1-\varepsilon)^2) \sim \mathcal{O}(\eta_2^2), \end{aligned}$$

where the order follows from the assumption that \tilde{p}_i will remain within the support of \hat{p} .

We will soon determine the leading order asymptotic behaviour of E_1 and E_2 , from which

²library/ornstein-uhlenbeck/solve_binary_ou.jl

we can calculate the expected growth rate by considering first that (for a fixed z)

$$\mathbb{E}_{\tilde{p}}(\lambda(\varepsilon, z)) = \gamma \mathbb{E}_{\tilde{p}_1}(\lambda(\varepsilon, z)) + (1 - \gamma) \mathbb{E}_{\tilde{p}_2}(\lambda(\varepsilon, z)). \quad (\text{S15})$$

By considering individually the behaviour near $\varepsilon = 0$ and $\varepsilon = 1$, we can see that

$$\begin{aligned} \lambda(\varepsilon, z) \sim \lambda(0, z) + \lambda'(0, z)\varepsilon &\quad \Rightarrow \quad \mathbb{E}_{\tilde{p}_1}(\lambda(\varepsilon, z)) \sim \lambda(0, z) + \lambda'(0, z)E_1, \\ \lambda(\varepsilon, z) \sim \lambda(1, z) - \lambda'(1, z)(1 - \varepsilon) &\quad \Rightarrow \quad \mathbb{E}_{\tilde{p}_2}(\lambda(\varepsilon, z)) \sim \lambda(1, z) - \lambda'(1, z)E_2. \end{aligned}$$

Finally, we can determine γ in terms of E_1 and E_2 by considering that the *net transfer of phenotypes* from near $\varepsilon = 0$ (and similarly in the reverse direction) is still given by $\omega(1 - q)$, while the order of integration and differentiation can be exchanged to see that the net growth of phenotypes near $\varepsilon = 0$ is simply given by $\mathbb{E}_{\tilde{p}_1}(\lambda(\varepsilon))$. Thus, we can use the result from Section S2.2 to see that the QSS proportion of cells near the dormant state is given by

$$\gamma = \frac{-a + b + \Delta\lambda + \sqrt{4b\lambda + (-a + b + \Delta\lambda)^2}}{2\Delta\lambda}, \quad (\text{S16})$$

for $a = \omega(1 - q)$, $b = \omega q$ and where $\Delta\lambda = \mathbb{E}_{\tilde{p}_2}(\lambda(\varepsilon)) - \mathbb{E}_{\tilde{p}_1}(\lambda(\varepsilon))$ (such that we consider that, equivalently, the net growth rate of the dormant population to be zero; results relating to the population distribution are insensitive to such growth rate translation, or equivalently, scaling at the population level).

To determine asymptotic expressions for E_1 and E_2 , we first substitute eq. (S13) into eq. (S12), to obtain

$$0 = (\gamma\tilde{p}_1 + (1 - \gamma)\tilde{p}_2) \left[\lambda - \gamma \mathbb{E}_1(\lambda) - (1 - \gamma) \mathbb{E}_2(\lambda) - \omega \right] + \omega(qp_1 + (1 - q)p_2), \quad (\text{S17})$$

Recalling that eq. (S17) must apply for all ε , we next assume that the density of \tilde{p}_1 and p_1 is negligible near $\varepsilon = 1$, and similar for \tilde{p}_2 and p_2 near $\varepsilon = 0$. Following several substitutions and by considering behaviour up to $\mathcal{O}(\eta_i)$, we obtain

$$\begin{aligned} 0 &= \sqrt{\frac{2}{\pi}} q\omega\eta_1 - E_1\gamma \left[\omega + (1 - \gamma)(\lambda(1, z) - \lambda(0, z)) \right], \\ 0 &= \sqrt{\frac{2}{\pi}} (1 - q)\omega\eta_2 - E_2(1 - \gamma) \left[\omega - \gamma(\lambda(1, z) - \lambda(0, z)) \right]. \end{aligned}$$

It remains to substitute the expression for γ (eq. (S16)), and consider matched order terms to obtain asymptotic expressions for E_1 and E_2 . Substituting the result back into eq. (S15), we obtain

$$\mathbb{E}_{\tilde{p}}(\lambda(\varepsilon, z)) = c_0(z) + c_1(z)\eta_1 + c_2(z)\eta_2, \quad (\text{S19})$$

where

$$\begin{aligned} c_0(z) &= \frac{1}{2} \left(\Delta(z) - \omega + \sqrt{\Omega} \right), \\ c_1(z) &= \frac{\lambda'(0, z) \left([(2q-1)\omega - \Delta(z)] + 1 \right)}{\sqrt{2\pi\Omega(z)}}, \\ c_2(z) &= \frac{\lambda'(1, z) \left([(2q-1)\omega - \Delta(z)] - 1 \right)}{\sqrt{2\pi\Omega(z)}}, \end{aligned}$$

for

$$\begin{aligned} \Delta(z) &:= \lambda(1, z) - \lambda(0, z), \\ \Omega(z) &:= (\omega - \Delta(z))^2 + 4(1-q)\omega\Delta(z), \end{aligned}$$

and where $\lambda'(\varepsilon, z)$ indicates a derivative with respect to ε .

An asymptotic expression for the fitness is, therefore, given by

$$\varphi \sim C_0 + C_1\eta_1 + C_2\eta_2,$$

where

$$C_i = \int_{-\infty}^{\infty} c_i(z) \Phi'(z) dz.$$

S3.3 Fluctuations in the fittest phenotype

We consider that the growth rate is given by (for now, general) $\lambda(\varepsilon, w)$, where the environment, w , fluctuates according to an Ornstein-Uhlenbeck process with equilibrium distribution $w \sim \mathcal{N}(0, \sigma)$. For simplicity, we consider that $\lambda(\varepsilon, w)$ is symmetric about ε , such that the optimal phenotype in a homogeneous strategy is given by $\varepsilon^* = 0$ (in the main document, we translate the phenotype index such that $\varepsilon^* = \frac{1}{2}$ so that the phenotype index remains bounded $\varepsilon \in [0, 1]$).

The target distribution is given by $\hat{p}(\varepsilon) = \mathcal{N}(0, \eta)$, for $\eta \ll 1$. For a fixed value of the environment (i.e., in the QSS regime), in the vicinity of $\varepsilon^* = 0$ we approximate

$$\lambda(\varepsilon, w) \sim \lambda(0, w) + \lambda'(0, w)\varepsilon + \frac{\lambda''(0, w)}{2}\varepsilon^2, \quad (\text{S21})$$

where $\lambda'(\varepsilon, w)$ denotes a derivative with respect to ε .

Next, we substitute eq. (S25) into the governing equation for \tilde{p} (eq. (S12)), multiply by ε^n , and integrate to yield

$$0 = \lambda'(0, w)E_{n+1} + \frac{1}{2}\lambda''(0, w)E_{n+2} - \lambda'(0, w)E_1E_n - \frac{1}{2}\lambda''(0, w)E_2E_n - \omega E_n + \omega M_n, \quad (\text{S22})$$

where we have defined

$$E_n = \mathbb{E}_{\tilde{p}}(\varepsilon^n) \quad \text{and} \quad M_n = \mathbb{E}_{\tilde{p}}(\varepsilon^n). \quad (\text{S23})$$

If we proceed as before and assume that $E_1 \sim \mathcal{O}(\eta)$, we find that the equation can only

be satisfied at $\mathcal{O}(\eta)$ through the trivial solution $E_1 = 0$ that later leads to a contradiction. Thus, we progress to the next order and assume that potentially both $E_1, E_2 \sim \mathcal{O}(\eta^2)$, while $E_n \sim \mathcal{O}(\eta^3)$ for $n \geq 3$. For $n = 1$ and $n = 2$, we obtain up to order $\mathcal{O}(\eta^2)$:

$$\begin{aligned} 0 &= \lambda'(0, w)E_2 - \omega E_1, \\ 0 &= -\omega E_2 + \omega\eta^2, \end{aligned}$$

such that $E_1 = \lambda'(0)\omega^{-1}/\eta^2$ and $E_2 = \eta^2$. Substituting back into eq. (S25), we find that

$$\mathbb{E}_{\bar{p}}(\lambda(\varepsilon, w)) \sim c_0(w) + c_2(w)\eta^2, \quad (\text{S25})$$

where the first order correction vanishes, and

$$\begin{aligned} c_0(w) &= \lambda(0, w), \\ c_2(w) &= \frac{\lambda'(0, w)^2}{\omega} + \frac{\lambda''(0, w)}{2}. \end{aligned}$$

As in the previous section, the fitness is thus given by

$$\varphi \sim C_0 + C_2\eta^2,$$

where

$$C_i = \int_{-\infty}^{\infty} c_i(z)\phi_w(w) dw, \quad (\text{S27})$$

where $\phi_w(w)$ is the density function for the random variable w .

For the specific formulation examined in the main document

$$\lambda(\varepsilon, w) = k \exp\left(\frac{-(\varepsilon - w)^2}{\xi^2}\right),$$

we can compute the integral in eq. (S27) analytically to produce the expression given in the main document.