

Algorithm for motivic Hilbert zeta function of some curve singularities

Wenhao Zhu^a, Yizi Chen^b, Hussein Mourtada^a

^a*Imj-Prg, Paris Cité University, Paris, 75013, France*

^b*ETH Zurich, Zurich, 8093, Switzerland*

Keywords: Algebraic geometry, punctual Hilbert scheme, singularities, semigroup, syzygy.

Abstract

We develop an algorithm for computing the motivic Hilbert zeta function for curve singularities with a monomial valuation group or for singular curves defined by $y^k = x^n$, where $\gcd(k, n) = 1$. It is well known that the Hilbert scheme of points on a smooth curve is isomorphic to the symmetric product of the curve. However, the geometrical structure of Hilbert scheme of points on singular curves remains less understood. The algorithm we propose computes the motivic Hilbert zeta function, $Z_{(C, O)}^{Hilb}(q) \in K_0(Var_{\mathbb{C}})[[q]]$, for such curve singularities. This function is represented as a series with coefficients in the Grothendieck ring of varieties over \mathbb{C} .

The main computational challenge arises from the infinity of Γ . To address this, we approximate Γ by truncating it to a finite subset to allow effective algorithm operation. We also analyze the time complexity and estimate the range of the effective finite length of Γ necessary for reliable results. The Python implementation of our algorithm is available at https://github.com/whaozhu/motivic_hilbert.

1. Introduction

We begin by reviewing the background of the motivic Hilbert zeta function, as presented in [1]. Let k be an algebraically closed field and $K_0(Var_k)$ the Grothendieck ring of varieties over k . Suppose X is a variety over k . M. Kapranov defined the motivic zeta function of X as the power series [2]:

$$Z_X^{mot}(t) := \sum_{n \geq 0} [Sym^n X] t^n \in K_0(Var_k)[[t]] \quad (1)$$

(where $[Sym^0 X] = 1$ by convention.) The motivic zeta function is an invariant of X . When X is defined over a finite field, one can obtain the Hasse-Weil zeta function via the point-counting measure:

$$Z_X(t) = \exp\left(\sum_{n \geq 1} \frac{N_n}{n} t^n\right) = \sum_{n \geq 0} \#(Sym^n X)(\mathbb{F}_q) t^n \in \mathbb{Q}[[t]] \quad (2)$$

Email address: wenhaozhu@imj-prg.fr (Wenhao Zhu)

where $N_n = \#(X(\mathbb{F}_{q^n}))$ be the number of \mathbb{F}_{q^n} -points of X .

However, $Z_X^{mot}(t)$ is not sensitive to the singularities of X . For example when X is a cuspidal cubic, $Z_X^{mot}(t) = Z_{\mathbb{P}^1}^{mot}(t)$.

A more refined invariant that accounts for the singularities of X is the motivic Hilbert zeta function [1]:

$$Z_X^{Hilb}(t) := \sum_{n \geq 0} [Hilb^n(X)] t^n \in K_0(Var_k)[[t]] \quad (3)$$

where $Hilb^n(X)$ is the Hilbert scheme of points parametrizing length n subschemes of X with dimension 0. When X is a smooth curve, $Hilb^n(X)$ coincides with $Sym^n(X)$, so the Hilbert and usual motivic zeta functions coincide. When X is a reduced curve over k , the motivic Hilbert zeta functions of curves are rational [1].

Let C be a reduced plane curve with singular points p_1, \dots, p_l . By Corollary 2.2 in [1], the motivic Hilbert zeta function can be reduced to the local calculation.

$$Z_C^{Hilb}(t) = Z_{C^{sm}}^{Hilb}(t) \prod_{i=1}^l Z_{(C, p_i)}^{Hilb}(t) \quad (4)$$

where C^{sm} denotes the smooth locus of X , $Z_{(C, p_i)}^{Hilb}(t)$ is the motivic zeta function associating with the punctual Hilbert scheme which is the locus parametrizing subschemes supported on the singularity p_i .

The goal of this paper is to develop an algorithm to compute the motivic Hilbert zeta function $Z_{(C, O)}^{Hilb}(q) \in K_0(Var_{\mathbb{C}})[[t]]$ for a germ of singular curve (C, O) with monomial valuation semigroup in the sense of G.Pfister and J.H.M.Steenbrink [3] and for singular curves defined by $y^k = x^n$.

In Section 2, we recall the works on the punctual Hilbert scheme by G. Pfister and J. H. M. Steenbrink [3]. Let (C, O) be the germ of a reduced unbranched curve with a singular point $O \in C$. For a positive integer ℓ , the punctual Hilbert scheme $Hilb^\ell(C, O)$ is the parameter space for colength ℓ ideals in the complete local ring $\widehat{\mathcal{O}_{C, O}}$. We denote it by $C^{[\ell]}$ for convenience.

The Hilbert scheme $C^{[\ell]}$ can be embedded into a Grassmannian variety $Gr(\delta, 2\delta)$ as shown in [3, 4], where δ represents the δ -invariant of (C, O) . Through this embedding, one can derive the equations that define $C^{[\ell]}$ (Theorem 3 in [3]). However, it is difficult to study the geometry of $C^{[\ell]}$ using this equations because we should consider the image of $C^{[\ell]}$ under the Plücker embedding, $Gr(\delta, 2\delta) \hookrightarrow \mathbb{P}(\bigwedge^\delta \mathbb{C}^{2\delta})$. Such calculations are very complicated.

In the unbranched curve case, when $\ell \geq c$, the punctual Hilbert scheme $C^{[\ell]}$ coincides with the compactified Jacobian $\overline{\mathcal{JC}}$ [4], where c is the conductor of Γ . However, progress in obtaining explicit formulas for compactified Jacobians' homology and motivic class has been comparatively slow. G. Lusztig and J. Smelt [5] computed the homology of the curve defined by $y^k = x^n$ with $\gcd(n, k) = 1$. J. Piontkowski extended these results by computing the homology of compactified Jacobians for curves with one Puiseux pair, parameterized as $(x, y) = (t^n, t^m + \dots)$. Additionally, G. Gorsky, M. Mazin, and A. Oblomkov [6] derived the Poincaré polynomials for compactified Jacobians of plane curve singularities with Puiseux

exponents $(nd, md, md + 1)$. A key insight is that the intersection of \overline{JC} with Schubert cells forms an affine space, and these cells, along with their dimensions, have been given combinatorial interpretations, particularly in the context of q, t -Catalan combinatorics [7, 8, 9, 10].

In [11], the authors showed that, for each $\ell \geq 1$, $C^{[\ell]}$ admits a similar stratification (See Equation (5)) and provide defining equations for each stratum (Proposition 12 in [11]). Given a subsemimodule Δ of Γ , the corresponding stratum parametrizes ideals with a valuation set equal to Δ , denoted by $C^{[\Delta]}$. The main task of this paper is to compute the class of the punctual Hilbert scheme $[C^{[\ell]}]$ in the Grothendieck ring using the stratification. The essential part is determining all possible subsemimodules of Γ .

In section 3, we develop a method to determine all possible subsemimodules of Γ corresponding to a tree structure. Inspired by Y.Soma and M.Watari [12], one can obtain a new subsemimodule $d_i(\Delta)$ by deleting i -th minimal generators of a Γ - subsemimodule Δ . The key to ensuring that all possible subsemimodules are obtained lies in the existence of the Frobenius element i.e. the maximal element of $\Gamma \setminus \Delta$, denote it by γ_Δ . The union of Δ and γ_Δ is also a subsemimodule of Γ , denoted by $m(\Delta)$. And γ_Δ is one of the minimal generator of $m(\Delta)$. So we can delete it to obtain Δ . We begin with the semimodule corresponding to the valuation set of the maximal ideal. By viewing subsemimodules as vertices and deletions as edges, we construct a graph. Furthermore, if we restrict the edges to those corresponding to the deletion of Frobenius elements, we obtain a tree G_Γ .

In section 4, we use the main theorem in paper [13] asserts that in the case C is curve with a monomial valuation group or defined by $y^k = x^n$, its valuation group denoted to be Γ , an edge of the tree G_Γ between two vertexes (Δ and $m(\Delta)$) represents a geometric morphism between $C^{[\Delta]}$ and $C^{[m(\Delta)]}$ (Theorem 1). This morphism captures the relationship between the classes of these strata in the Grothendieck ring (Corollary 3).

In section 5, we introduce an Algorithm 1 for computing the motivic Hilbert zeta function of monomial curve singularities, which involves several interdependent Algorithm 2 through Algorithm 4. The algorithms implement truncation set $\Gamma_{\geq n}$ to ensure computational feasibility. Each sub-algorithm contributes a specific component to the overall computation: identifying minimal generators, computing syzygies, and generating semimodule sets from the valuation semigroup. Through these steps, the algorithm efficiently constructs the motivic Hilbert zeta function. The time complexity of the algorithm is analyzed in detail.

In Section 6, we introduce N as the minimal effective bound of Γ in Algorithm 1 and provide an estimate for its range (Theorem 3).

In section 7, we will treat specially the curve singularities of type W_8 (i.e. the curve singularities whose complete local rings are isomorphic to $\mathbb{C}[[t^4, t^5, t^6]]$). Y.Soma [14] and M.Watari [12] studied the geometry of punctual Hilbert of a plane curve singularities of type $A_{2k}(k \geq 1)$, E_6 and E_8 . And recently, M. Watari computed the motivic Hilbert zeta function of such type curve singularities [15]. We also can compute the motivic Hilbert zeta function of such specific type curve singularities differently by using Algorithm 1(See examples in <https://github.com/whaozhu/hilbert>.)

2. Punctual Hilbert scheme and monomial semigroup

In this section, we recall the basic concept in [3]. Let A be a complete local ring of an irreducible curve singularity over an algebraically closed field k such that its normalization \bar{A} is a discrete valuation ring. Let $v : \bar{A} \rightarrow \mathbb{N} \cup \{\infty\}$ be the corresponding discrete valuation. We let $\Gamma := v(A \setminus \{0\})$ denote the semigroup of A , $\bar{I}(n) := \{f \in \bar{A} | v(f) \geq n\}$ the ideal of \bar{A} , $I(n) := \bar{I}(n) \cap A$ for $n \in \mathbb{N}$. Define $c := \min\{n | \bar{I}(n) \subset A\}$ and $\delta := \dim_k(\bar{A}/A) = \#(\mathbb{N} \setminus \Gamma)$ of A . Then $\delta + 1 \leq c \leq 2\delta$, $c = 2\delta$ if and only if A is Gorenstein[16].

Let (C, O) be the germ of a unibranch curve singularity with complete local ring $A := \widehat{\mathcal{O}_{C,O}} \subset \mathbb{C}[[t]]$. For integers $\ell \geq 1$, the punctual Hilbert scheme (C, O) are defined as

$$C^{[\ell]} := \{I | I \text{ is an ideal of } A \text{ with colength } \ell\}$$

Pfister and Steenbrink [3] defined a closed subset \mathcal{M} of the Grassmannian $Gr(\delta, A/I(2\delta))$

$$\mathcal{M} := \{W \in Gr(\delta, \bar{A}/I(2\delta)) | W \text{ is an } A\text{-submodules of } \bar{A}/I(2\delta)\}$$

Proposition 1. [3]. *For $\ell \geq 0$, there exists a closed embedding $\phi_\ell : C^{[\ell]} \rightarrow \mathcal{M}$ and the map are bijective when $\ell \geq c$.*

Since $c \leq 2\delta$, for the computation of motivic Hilbert zeta function, we only need to consider the finite terms.

Next setp, we will introduce a stratification of Hilbert scheme $C^{[\ell]}$. Let Γ be the valuation semigroup of C . Let Δ be a Γ -subsemimodule if Δ is a subset of Γ and $\Delta + \Gamma \subset \Delta$. Given an ideal I , we denote $\Gamma(I) := v(I \setminus \{0\})$ which is a subsemimodule of Γ . We will study the moduli space of ideals with given Γ -subsemimodule:

$$C^{[\Delta]} := \{I | I \text{ is an ideal of } A \text{ with } \Gamma(I) = \Delta\}$$

We introduce Lemma 5.1.24 in [17]:

Lemma 1. [17] *For positive integer ℓ , an ideal I of A belongs to $C^{[\ell]}$ if and only if $\#(\Gamma \setminus \Gamma(I)) = \ell$.*

Let $\mathcal{D}_\ell := \{\Delta \subset \Gamma | \Delta \text{ is } \Gamma\text{-semimodule with } \#(\Gamma \setminus \Delta) = \ell\}$ be the set of semimodule with ℓ number of complement set of Γ . Then following the Lemma 1, we have

$$C^{[\ell]} = \bigsqcup_{\Delta \in \mathcal{D}_\ell} C^{[\Delta]} \tag{5}$$

Remark 1. *As notation above, $C^{[\Delta]}$ is the intersection of $C^{[\ell]}$ with the Schubert cell of the Grassmannian $Gr(\delta, A/I(2\delta))$ (Lemma 5 in [3]). So $C^{[\Delta]}$ is a locally closed set of $C^{[\ell]}$.*

Grothendieck ring $K_0(Var_{\mathbb{C}})$ is the ring generated by the isomorphism class $[X]$ of varieties of X , the sum and product coming from disjoint union and direct product respectively. By the stratification, we express $[C^{[\ell]}] = \sum_{\Delta \in \mathcal{D}_\ell} [C^{[\Delta]}]$ in $K_0(Var_{\mathbb{C}})$.

In the following section, we will introduce a method for determining all possible elements in \mathcal{D}_ℓ .

Definition 1. [3] A monomial curve singularity over k is an irreducible curve singularity with local ring isomorphic to $A = k[[t^{a_1}, \dots, t^{a_m}]]$ with $\gcd(a_1, \dots, a_m) = 1$.

Then we can assume Γ is generated by the minimal set of generators for Γ .

$$\Gamma = \langle a_1, \dots, a_m \rangle$$

Definition 2. [3] A semigroup $\Gamma \subset \mathbb{N}$ is called monomial if $0 \in \Gamma$, $\#(\mathbb{N} \setminus \Gamma) < \infty$ and each reduced and irreducible curve singularity with semigroup Γ is a monomial curve singularity.

Proposition 2. [3] For a semigroup $\Gamma \subset \mathbb{N}$, the following are equivalent:

- (1) Γ is a monomial semigroup.
- (2) Γ is semigroup form:
 - (i) $\Gamma_{m,s,b} := \{im \mid i = 0, 1, \dots, s\} \cup [sm + b, \infty)$ with $1 \leq b < m$, $s \geq 1$,
 - (ii) $\Gamma_{m,r} := \{0\} \cup [m, m + r - 1] \cup [m + r + 1, \infty)$ with $2 \leq r \leq m - 1$,
 - (iii) $\Gamma_m := \{0, m\} \cup [m + 2, 2m] \cup [2m + 2, \infty)$ with $m \geq 3$.
- (3) $0 \in \Gamma$, Suppose that $\Gamma = \#(\mathbb{N} \setminus \Gamma) \leq \infty$ and the following hold:
if $x \in \mathbb{N} \setminus \Gamma$ and $c(x) := \min\{n \in \mathbb{N} \mid [n, \infty) \subset \Gamma \cup (x + \Gamma)\}$, then $\Gamma \cap (x + \Gamma) \subset [c(x), \infty)$.

Lemma 2. [3][13] For $y \in \mathbb{Z}$ and $x \in \mathbb{N} \setminus \Gamma$, then $c(x) + y = \min\{m \in \mathbb{N} \mid [m, \infty) \subset (\Gamma + y) \cup (x + y + \Gamma)\}$.

Corollary 1. [3][13] Given a Γ -module Δ , for $\gamma_1 > \gamma_2$ is two generators of Δ , then for any $\sigma \in \text{Syz}(\langle \gamma_1, \gamma_2 \rangle)$, we have $\sigma \geq c(\Delta)$.

3. Tree structure associated with valuation semigroup

Let (C, O) be the germ of a unibranch plane curve singularity with complete local ring $A := \widehat{\mathcal{O}_{C,O}} \subset \mathbb{C}[[t]]$. Let $\Gamma = v(A \setminus 0)$. Recall that, for integer $\ell \geq 0$, let $\mathcal{D}_\ell = \{\Delta \subset \Gamma \mid \Delta \text{ is } \Gamma\text{-semimodule with } \#(\Gamma \setminus \Delta) = \ell\}$ be the set of semimodule with ℓ number of complement set of Γ . Then, by the Lemma 1, we have

$$C^{[\ell]} = \bigsqcup_{\Delta \in \mathcal{D}_\ell} C^{[\Delta]}$$

For Δ , the subsemimodule of Γ , we denote $\Delta = (\gamma_1, \dots, \gamma_m)$ is minimally generated by $\gamma_1 < \dots < \gamma_m$ (i.e. $\Delta = \sum_{i=1}^m (\gamma_i + \Gamma)$ and $\Delta \not\supseteq \sum_{i=1, i \neq j}^m (\gamma_i + \Gamma)$ for $\forall j \in \{1, \dots, m\}$).

Lemma 3. [12] Let Δ be a Γ -submodule which has minimal generators $\{\gamma_1, \dots, \gamma_m\}$. If $\Delta \in \mathcal{D}_\ell$, then $\Delta \setminus \{\gamma_i\} \in \mathcal{D}_{\ell+1}$, for any $i = 1, \dots, m$.

We denote $\mathcal{D}_{\ell,m} := \{\Delta \in \mathcal{D}_\ell \mid \Delta \text{ admits } m \text{ generators}\}$. Then $\mathcal{D}_\ell = \bigcup_{m \geq 0} \mathcal{D}_{\ell,m}$. Hence, for positive integers $\ell \geq 0$, $i \leq m$ we can define canonical maps:

$$\begin{aligned} d_{\ell,i} : \mathcal{D}_{\ell,m} &\rightarrow \mathcal{D}_{\ell+1} \\ \Delta &\mapsto \Delta \setminus \{\text{the } i\text{-th generator of } \Delta\} \end{aligned}$$

A natural question arises: Can we find all possible elements of $\mathcal{D}_{\ell+1}$ from \mathcal{D}_ℓ using the maps $d_{\ell,i}$? In other words, is $\bigcup_{m \geq 1, i \leq m} d_{\ell,i}(\mathcal{D}_{\ell,m})$ equal to $\mathcal{D}_{\ell+1}$? The answer is affirmative, as demonstrated by Lemma 4 and further explained in Remark 2.

We denote the Frobenius element $\gamma_\Delta := \max(\Gamma \setminus \Delta)$.

Lemma 4. *If $\Delta \in \mathcal{D}_{\ell+1}$, then $\Delta \cup \{\gamma_\Delta\} \in \mathcal{D}_\ell$.*

Remark 2. *We can obtain any $\Delta \in \mathcal{D}_{\ell+1}$ from \mathcal{D}_ℓ by $d_{\ell,i}$ for some i .*

Proof. γ_Δ is one of the generators of $\Delta \cup \{\gamma_\Delta\}$ as Γ -module and $\Delta \cup \{\gamma_\Delta\} \in \mathcal{D}_\ell$. □

So we can define canonical maps:

$$\begin{aligned} m_{\ell+1} : \mathcal{D}_{\ell+1} &\rightarrow \mathcal{D}_\ell \\ \Delta &\mapsto \Delta \cup \{\gamma_\Delta\} \end{aligned}$$

We denote $V_\ell := \mathcal{D}_\ell$, $1 \leq \ell \leq c$. $E_\ell := \{(m_\ell|_\Delta)^{-1} \mid \Delta \in \mathcal{D}_\ell\}$, $2 \leq \ell \leq c$.

$V := \bigcup_{i=1}^c V_i$, $E := \bigcup_{i=2}^c E_i$. We define a finite graph $G_\Gamma := (V, E)$ which is a tree, where E_ℓ are edges between V_ℓ and $m_\ell(V_\ell) \subset V_{\ell-1}$.

Example 1. $A = k[[t^3, t^4]]$, $\Gamma = \langle 3, 4 \rangle = \{0, 3, 4, 6, 7, 8, 9, \dots\}$, $c = 6$, $(3, 4) = \{3, 4, 6, 7, 8, 9, \dots\}$. We have $d_{1,1}((3, 4)) = \{4, 6, 7, 8, 9, \dots\} = (4, 6)$, $d_{1,2}((3, 4)) = \{3, 6, 7, 8, 9, \dots\} = (3, 8)$, $d_{2,1}((4, 6)) = (6, 7, 8)$ etc. Then we can obtain a tree:




Figure 1: Tree for the case of E_6 type singularity

4. Piecewise fibration

In this section, we revisit the key results from the paper [13]. Let (C, O) be the germ of a unibranch plane curve singularity with semigroup Γ , where Γ is either monomial or of the form $\langle \alpha_1, \alpha_2 \rangle$. We denote Γ by $\langle \alpha_1, \dots, \alpha_e \rangle$. Then by section 3, we can construct a tree from Γ . The edges in the tree illustrate the relationships between the classes of cells of Hilbert scheme corresponding to its two vertices in the Grothendieck ring (Corollary 3).

Let $\Delta = (\gamma_1, \dots, \gamma_m)$. The syzygy of Δ is also Γ -subsemimodule defined by $Syz(\Delta) := \{\sigma \in \Delta \mid \sigma = \gamma_{i_1} + b_1 = \gamma_{i_2} + b_2, \exists \gamma_{i_1} \neq \gamma_{i_2}, b_1, b_2 \in \Gamma\}$ which is generated by $\{\sigma_1, \dots, \sigma_n\}$ as Γ -semimodule. $\gamma_\Delta = \max(\Gamma \setminus \Delta)$. $m(\Delta) = \Delta \cup \{\gamma_\Delta\}$. For any element $I \in C^{[\Delta]}$, there exists a set of generators of I of the form $\{f_{\gamma_1}(t), \dots, f_{\gamma_m}(t)\}$, where $f_{\gamma_j} = t^{\gamma_j} + \sum_{k \in \Gamma > \gamma_j \setminus \Delta} \lambda_j^{k-\gamma_j} t^k$ for some $\lambda_j^{k-\gamma_j} \in \mathbb{C}$.

There exists a canonical morphism:

$$C^{[\Delta]} \rightarrow C^{[m(\Delta)]}, \quad \langle f_{\gamma_1}(t), \dots, f_{\gamma_m}(t) \rangle \mapsto \langle f_{\gamma_1}(t), \dots, f_{\gamma_m}(t), t^{\gamma_\Delta} \rangle \quad (6)$$

Theorem 1. [13] For a unibranch plane curve singularity C with valuation semigroup $\Gamma = \langle n, k \rangle$, the canonical morphism $C^{[\Delta]} \rightarrow C^{[m(\Delta)]}$ is a piecewise fibration.

Corollary 2. *According the proof of Theorem 1. for a unibranch plane curve singularity C with monomial valuation semigroup, the canonical morphism $C^{[\Delta]} \rightarrow C^{[m(\Delta)]}$ is a piecewise fibration.*

Corollary 3. *[13] For a unibranch plane curve singularity C with monomial valuation semigroup or $\Gamma = \langle n, k \rangle$, then given a subsemimodule $\Delta = (\gamma_1, \dots, \gamma_m) \subset \Gamma$, $Syz(\Delta) = (\sigma_1, \dots, \sigma_n)$, we have*

$$[C^{[\Delta]}] = [C^{[m(\Delta)]}] \cdot \mathbb{L}^{|\gamma_i|_{\gamma_i < \gamma_\Delta} - |\sigma_i|_{\sigma_i < \gamma_\Delta}}$$

Example 2. $C = \{y^4 = x^7\} \subset \mathbb{C}^2$, $\Gamma = \langle 4, 7 \rangle = \{4, 7, 8, 11, 12, 14, 15, 16, 18, \rightarrow\}$, $\Delta = (8, 11) = \{8, 11, 12, 15, 16, 17, 18, 19, 20, 22, \rightarrow\}$, $\gamma_\Delta = 21$, $m(\Delta) = (8, 11, 21) = \{8, 11, 12, 15, 16, 18, \rightarrow\}$. $Syz(\Delta) = (15, 32)$.

Since $\{\gamma_i \mid \gamma_i < \gamma_\Delta\} = \{8, 11\}$, $\{\sigma_i \mid \sigma_i < \gamma_\Delta\} = \{15\}$, then $[C^{[(8,11)]]} = [C^{[(8,11,21)]]} \cdot \mathbb{L}$.

5. Algorithm

In this section, we introduce Algorithm 1 to compute the motivic Hilbert zeta function of monomial curve singularities. This algorithm relies on a series of sub-algorithms, from Algorithm 2 to Algorithm 4, each of which performs critical steps that contribute to the overall computation process.

We define the notation $[a, b] := \{n \in \mathbb{Z} \mid a \leq n \leq b\}$ for the integers $a \leq b$.

Algorithm 1 takes as input the germ of a unibranch plane curve singularity (C, O) with monomial complete local ring $\widehat{\mathcal{O}}_{C,O} = \mathbb{C}[[t^{\alpha_1}, \dots, t^{\alpha_e}]]$ with a monomial valuation semigroup Γ or $\Gamma = \langle \alpha_1, \alpha_2 \rangle$ and its conductor c . Since Γ is an infinite set, handling it directly on a computer is impractical. To address this, we implement the truncated set $\Gamma_{\geq n} = \Gamma \cap [0, n]$, for some n which allows our Algorithm 1 through Algorithm 4 to operate within practical limits. We will discuss how to determine n . In Section 6, we discuss methods for determining an appropriate value of n . According to Theorem 3, setting $n = (c - 1)(\alpha_1 + 2)$ guarantees that the algorithm functions correctly.

The output of Algorithm 1 is the motivic Hilbert zeta function $Z_{(C,O)}^{Hilb}(q) = \sum_{\Delta} [C^{[\Delta]}] q^{\ell(\Delta)}$. Initializing $\mathcal{D}_1 = \{\Delta_1\}$, where $\Delta_1 = \Gamma \setminus \{0\} = (\alpha_1, \dots, \alpha_e)$. According to Section 3, there exists a tree G_Γ associated Γ , using Algorithm 4, we obtain $\mathcal{D}_2, \dots, \mathcal{D}_c$ (in Algorithm 1 Line 1). Set up a loop over $\ell = 1, \dots, c$ for calculating each coefficient term $[C^\ell] = \sum_{\Delta \in \mathcal{D}_\ell} [C^\Delta]$ in the motivic Hilbert zeta function.

In the main loop: for each $\Delta \in \mathcal{D}_\ell$, we compute γ_Δ the Frobenius element of Δ and use Algorithm 3 to find $Syz(\Delta)$, the syzygy of Δ . Using Algorithm 2, we calculate $T_\Delta = \{\gamma_1, \dots, \gamma_m\}$, the minimal generators of Δ , and $T_{Syz(\Delta)} = \{\sigma_1, \dots, \sigma_n\}$ the minimal generators of $Syz(\Delta)$. For the contribution counts: we compute $C_1 = \#\{\gamma_i \mid \gamma_i < \gamma_\Delta\}$, $C_2 = \#\{\sigma_i \mid \sigma_i < \gamma_\Delta\}$.

Algorithm 1 Motivic Hilbert zeta function of curve singularity

Require: C is a curve singularity with valuation semigroup $\Gamma = \langle \alpha_1, \dots, \alpha_e \rangle$. c is the conductor of Γ .

Ensure: Motivic Hilbert zeta function $Z_{(C,O)}^{Hilb}(q) = \sum_{\Delta} [C^{[\Delta]}] q^{\ell(\Delta)}$.

```
1:  $\mathcal{D}_1 = \{\Delta_1 = (\alpha_1, \dots, \alpha_e)\}$ ,  $\mathcal{D}_2 = \{\Delta_{2,i} = (\Delta_1 \setminus \{\alpha_i\})\}, \dots, \mathcal{D}_c$ ;
2: for  $\ell = 1, \dots, c$  do
3:   for  $\Delta \in \mathcal{D}_\ell$  do
4:      $\gamma_\Delta = \max(\Gamma \setminus \Delta) \leftarrow$  Frobenius element of  $\Delta$ ;
5:      $T_\Delta = \{\gamma_1, \dots, \gamma_m\} \leftarrow$  minimal generators of  $\Delta$ ;
6:     compute  $Syz(\Delta)$ ;
7:      $T_{Syz(\Delta)} = \{\sigma_1, \dots, \sigma_n\} \leftarrow$  minimal generators of  $Syz(\Delta)$ ;
8:      $m(\Delta) = \Delta \cup \{\gamma_\Delta\}$ ;
9:      $C_1 = 0$ ;
10:    for  $\gamma_i \in T_\Delta$  do
11:      if  $\gamma_i < \gamma_\Delta$  then
12:         $C_1 \leftarrow C_1 + 1$ ;
13:      end if
14:    end for
15:     $|\gamma_i|_{\gamma_i < \gamma_\Delta} \leftarrow C_1$ ;
16:     $C_2 = 0$ ;
17:    for  $\sigma_i \in T_{Syz(\Delta)}$  do
18:      if  $\sigma_i < \gamma_\Delta$  then
19:         $C_2 \leftarrow C_2 + 1$ ;
20:      end if
21:    end for
22:     $|\sigma_i|_{\sigma_i < \gamma_\Delta, \sigma_i \prec \gamma_\Delta} \leftarrow C_2$ ;
23:     $[C^{[\Delta]}] \leftarrow [C^{[m(\Delta)}]] \cdot \mathbb{L}^{|\gamma_i|_{\gamma_i < \gamma_\Delta} - |\sigma_i|_{\sigma_i < \gamma_\Delta, \sigma_i \prec \gamma_\Delta}}$ ;
24:    end for
25:     $[C^\ell] = \sum_{\Delta \in \mathcal{D}_\ell} [C^\Delta]$ ;
26:  end for
27:  $Z_{(C,O)}^{Hilb}(q) \leftarrow 1 + \sum_{\ell \geq 1} [C^{[\ell]}] q^\ell$ ;
28: return  $Z_{(C,O)}^{Hilb}(q)$ ;
```

Algorithm 2 Minimal generators of Δ as Γ -module, T_Δ

Require: C is a curve singularity with valuation semigroup $\Gamma = \langle \alpha_1, \dots, \alpha_e \rangle$. c is the conductor of Γ . Δ is subsemimodule of Γ . We assume that $\Delta = \{a_1, \dots, a_s\}$ such that $a_i < a_{i+1}$, where $s = \#(\Delta)$ and $\#(\Gamma \setminus \Delta) = \ell$.

Ensure: Minimal generators of Δ as Γ -module T_Δ .

```
1: let  $T_\Delta = \{a_1\}$ ;  
2: for  $i = 2, \dots, s$  do  
3:   for  $x \in T_\Delta$  do  
4:      $r_i \leftarrow a_i - x$ ;  
5:     if  $r_i \in \Gamma$ : then  
6:       break;  
7:     end if  
8:     if  $r_i \notin \Gamma$  then  
9:        $T_\Delta \leftarrow T_\Delta \cup \{a_i\}$ ;  
10:    end if  
11:  end for  
12: end for  
13: return  $T_\Delta$ ;
```

Algorithm 3 Sygyzy of Δ , $Syz(\Delta)$

Require: C is a curve singularity with valuation semigroup $\Gamma = \langle \alpha_1, \dots, \alpha_e \rangle$. c is the conductor of Γ . Δ is subsemimodule of Γ . We assume that $\Delta = \{a_1, \dots, a_{s-1}\}$ such that $a_i < a_{i+1}$, where $s = \#(\Delta)$. Minimal generators of Δ as Γ -module, $T_\Delta = \{\gamma_1, \dots, \gamma_m\}$.

Ensure: Sygyzy of Δ , $Syz(\Delta)$.

```
1: let  $Syz(\Delta) = \{\}$ ;  
2: for  $i = 1, \dots, s - 1$  do  
3:    $count = 0$ ;  
4:   for  $j = 0, \dots, m - 1$  do  
5:      $r_{i,j} \leftarrow a_i - \gamma_j$ ;  
6:     if  $r_{i,j} \in \Gamma$  then  
7:        $count \leftarrow count + 1$ ;  
8:     end if  
9:     if  $count \geq 2$  then  
10:      break;  
11:    end if  
12:  end for  
13:  if  $count = 2$  then  
14:     $Syz(\Delta) \leftarrow Syz(\Delta) \cup \{a_i\}$ ;  
15:  end if  
16: end for  
17: return  $Syz(\Delta)$ ;
```

Algorithm 4 The tree obtained from Γ, G_Γ

Require: C is a curve singularity with valuation semigroup $\Gamma = \langle \alpha_1, \dots, \alpha_e \rangle$. c is the conductor of Γ .

Ensure: $\ell = 1, \dots, c$, \mathcal{D}_ℓ , the set of Γ -subsemimodule Δ such that $\#(\Gamma \setminus \Delta) = \ell$.

```

1: let  $\mathcal{D}_1 = \{\Delta_1 = (\alpha_1, \dots, \alpha_e)\}$ ,  $(\alpha_1, \dots, \alpha_e) = \Gamma \setminus \{0\}$ ;
2: for  $\ell = 1, \dots, c - 1$  do
3:    $\mathcal{D}_{\ell+1} = \{\}$ ;
4:   for  $\Delta \in \mathcal{D}_\ell$  do
5:     Compute the set of minimal generators of  $\Delta$ ;  $T_\Delta = \{\gamma_1, \dots, \gamma_m\}$ ,  $m \leq \alpha_1$  .
6:     for  $\gamma_i \in T_\Delta$  do
7:        $\Delta_i = \Delta \setminus \{\gamma_i\}$ ;
8:        $\mathcal{D}_{\ell+1} \leftarrow \mathcal{D}_{\ell+1} \cup \{\Delta_i\}$ 
9:     end for
10:  end for
11: end for
12: return  $\mathcal{D}_\ell, \ell = 1, \dots, c$ ;

```

In the final part of this section, we analyze the time complexity of the algorithms. Let $n = (c - 1)(\alpha_1 + 2)$ represent the bound of Γ and assume Γ admits finite length $n^* = n - \delta$ with conductor c .

Lemma 5. *Let Δ is subsemimodule of Γ with $\#(\Gamma \setminus \Delta) = \ell$. The time complexity of Algorithm 2 is $O(n^* \alpha_1 (n^* - \ell))$.*

Proof. For the outer loop i.e. for $i = 2, \dots, s = n^* - \ell$. it runs at most $O(n^* - \ell)$ times. For the inner loop, i.e., for $x \in T_\Delta$, the size of the set of minimal generators of Γ -module is at most α_1 . The inner loop would iterate over the size which is α_1 . Each inner loop iteration contains a membership check in Γ , which takes $O(n^*)$ time. Thus, the overall time complexity is $O(n^* \alpha_1 (n^* - \ell))$. \square

For the same reason, the time complexity of Algorithm 3 is $O(n^* \alpha_1 (n^* - \ell))$.

Let $\#(\mathcal{D}_\ell) = \chi_\ell$. By the corollary in [13], χ_ℓ are equal to Euler number of $C^{[\ell]}$.

Lemma 6. *The time complexity of Algorithm 4 is $\sum_{\ell=1}^{c-1} O(\chi_\ell \cdot n^* \alpha_1 (n^* - \ell))$.*

Proof. You only need to notice that, in line 7, the complexity of deleting an element in Δ is $O(1)$. The reason is that if we treat Δ as a sorted list since the element we delete is one of the generators of Δ as Γ -module, it must be at the relative beginning of the list. Then it takes $O(1)$ times. If we treat Δ as a set, deleting a specified element takes $O(1)$ times. Hence the total complexity is

$$\sum_{\ell=1}^{c-1} [O(\chi_\ell \cdot n^* \alpha_1 (n^* - \ell)) + O(\chi_\ell \cdot \alpha_1)] = \sum_{\ell=1}^{c-1} O(\chi_\ell \cdot n^* \alpha_1 (n^* - \ell))$$

□

Theorem 2. *The time complexity of Algorithm 1 is $\sum_{\ell=1}^c O(c \cdot \chi_\ell \cdot n^* \cdot \alpha_1 \cdot (n^* - \ell))$.*

Proof. By the Lemma 5, the overall time complexity is

$$\sum_{\ell=1}^c O(c \cdot \chi_\ell \cdot n^* \cdot \alpha_1 \cdot (n^* - \ell))$$

□

6. Effective finite length of Γ

Since $\Gamma = \langle \alpha_1, \dots, \alpha_c \rangle$ is a numerical semigroup, we have $\#(\mathbb{N} \setminus \Gamma) < \infty$, meaning that Γ is an infinite set. To implement our algorithm computationally, we face the challenge that a computer cannot directly handle an infinite set. Therefore, we approximate Γ by truncating it to a finite subset, ensuring our algorithm operates effectively. Specifically, we choose a positive integer n such that $\Gamma_{\leq n} = \Gamma \cap [0, n]$ allows the Algorithm 1 to function correctly.

Definition 3. *If N is the smallest integer for which Algorithm 1 operates correctly with the truncated set $\Gamma_{\leq N}$, we refer N to the minimal effective bound of Γ . Additionally, we define $N^* = \#(\Gamma \cap [0, N])$ is defined as the minimal effective length of Γ .*

Since Algorithm 1 relies on Algorithm 2, Algorithm 3, and Algorithm 4, determining N , the minimal effective bound of Γ , requires analyzing each of these sub-algorithms in detail.

Lemma 7. *The minimal effective bound of Algorithm 2 is sufficient to ensure effective operation for all $\Delta \in \mathcal{D}_\ell, 1 \leq \ell \leq c$, then it is less than $\gamma_{\Delta_c} + c - 1$, where $\gamma_{\Delta_c} = \max\{\gamma_\Delta\}_{\Delta \in \mathcal{D}_c}$.*

Proof. There exists $\Delta_c \in \mathcal{D}_c$ such that $\gamma_{\Delta_c} = \max\{\gamma_\Delta\}_{\Delta \in \mathcal{D}_c}$, it is the largest Frobenius element among all $\Delta \in \mathcal{D}_c$, where $\gamma_\Delta = \max(\Gamma \setminus \Delta)$. To prove the lemma, it suffices to show that the Algorithm 2 works operates effectively for Δ_c if we implement $\Gamma_{\leq N'}$ in Algorithm 2, where $\Gamma_{\leq N'} = \Gamma \cap [0, N']$, $N' = \gamma_{\Delta_c} + c - 1$. Assume the generators of Δ_c are $\gamma_1 < \dots < \gamma_m$. We only need to prove $\gamma_m \leq N'$, which implies that $\gamma_1, \dots, \gamma_m \in \Gamma'$.

Observing that the absolute difference between any two generators of Δ must be among the gaps $(\mathbb{N} \setminus \Gamma)$, with $c - 1$ as the maximum number of gaps. Hence, $\gamma_m - \gamma_1 \leq c - 1$. Given that $\gamma_1 < \gamma_\Delta$, it follows that $\gamma_m < \gamma_{\Delta_c} + c - 1 = N'$. □

By a similar argument, the minimal effective bound of Algorithm 4 is also $\gamma_{\Delta_c} + c - 1$. However, the minimal effective bound determined for Algorithm 2 may not suffice for Algorithm 3. Specifically, for a given $\Delta \in \mathcal{D}_c$, the generators of $Syz(\Delta)$ may exceed $\gamma_{\Delta_c} + c - 1$. Nonetheless, the minimal effective bound for Algorithm 2 ensures that Algorithm 3 operates effectively within Algorithm 1, as we only need to identify which minimal generator of $Syz(\Delta)$ is smaller than γ_Δ which is less than $\gamma_{\Delta_c} + c - 1$.

Theorem 3. *Let N be the minimal effective bound of Γ in the Algorithm 1. Then, $N \leq (c-1)(\alpha_1 + 2)$.*

Proof. Let $\Delta_c \in \mathcal{D}_c$ such that $\gamma_{\Delta_c} = \max\{\gamma_{\Delta}\}_{\Delta \in \mathcal{D}_c}$. Then $N \leq \gamma_{\Delta_c} + 1 + c$. Set $\Delta_{c-1} = \Delta_c \cup \{\gamma_{\Delta_c}\}$. If $\gamma_{\Delta_c} - \alpha_1 \in \Gamma$, then $\gamma_{\Delta_c} - \alpha_1 \notin \Delta_{c-1}$. Thus, $\gamma_{\Delta_{c-1}} \geq \gamma_{\Delta_c} - \alpha_1$ implying $\gamma_{\Delta_c} - \gamma_{\Delta_{c-1}} \leq \alpha_1$. Continue this process, we find that there exists a t such that $\gamma_{\Delta_t} - \alpha_1 \notin \Gamma$, which implies $\gamma_{\Delta_t} - \alpha_1 \leq c-1$, i.e., $\gamma_{\Delta_t} \leq \alpha_1 + c - 1$. Since $t \geq 2$, we have $\gamma_{\Delta_c} \leq (c-t+1)\alpha_1 + c - 1 \leq (c-1)\alpha_1 + c - 1$. Thus, $N \leq (c-1)(\alpha_1 + 2)$. \square

7. Experimentation

This section provides an example illustrating the computation of the motivic Hilbert zeta function for certain curve singularities.

Example 3. *Let C be a curve singularity with a local ring $\widehat{\mathcal{O}_{C,O}} = \mathbb{C}[[t^4, t^5, t^6]]$, has monomial semigroup $\Gamma = \Gamma_{4,3}$ in the sense of 2, $\Gamma = \langle 4, 5, 6 \rangle$, $c = 8$. According to the bound provided in Theorem 3, we set $n = (c-1)(\alpha_1 + 2) = 7 \times 6 = 42$ meaning that we implement $\Gamma_{\geq 42}$ in our computations using Algorithm 1, where $\Gamma_{\geq 42} = \{0, 4, 5, 6, 8\} \cup [9, 42]$.*

We initialize with $\mathcal{D}_1 = \{(4, 5, 6)\}$, where $(4, 5, 6) = \{0, 4, 5, 6, 8\} \cup [9, 42]$. Applying Algorithm 4 gives us the following sets:

$$\begin{aligned} \mathcal{D}_2 &= \{(5, 6, 8), (4, 6), (4, 5)\}, \\ \mathcal{D}_3 &= \{(6, 8, 9), (5, 8), (5, 6), (4, 11)\}, \\ \mathcal{D}_4 &= \{(8, 9, 10, 11), (6, 9), (6, 8), (5, 12), (4)\}, \\ \mathcal{D}_5 &= \{(9, 10, 11, 12), (8, 10, 11), (8, 9, 11), (8, 9, 10), (6, 13), (5)\}, \\ \mathcal{D}_6 &= \{(10, 11, 12, 13), (9, 11, 12), (9, 10, 12), (9, 10, 11), (8, 11), (8, 10), (8, 9), (6)\}, \\ \mathcal{D}_7 &= \{(11, 12, 13, 14), (10, 12, 13), (10, 11, 13), (10, 11, 12), (9, 12), (9, 11), (9, 10), (8, 15)\}, \\ \mathcal{D}_8 &= \{(12, 13, 14, 15), (11, 13, 14), (11, 12, 14), (11, 12, 13), (10, 13), (10, 12), (10, 11), (9, 16), (8)\}. \end{aligned}$$

The results of the computation of the motivic class of the punctual Hilbert scheme in the Grothendieck ring are as follows:

$$\begin{aligned} [C^{[1]}] &= 1 \\ [C^{[2]}] &= 1 + \mathbb{L} + \mathbb{L}^2 \\ [C^{[3]}] &= 1 + \mathbb{L} + 2\mathbb{L}^2 \\ [C^{[4]}] &= 1 + \mathbb{L} + 2\mathbb{L}^2 + \mathbb{L}^3 \\ [C^{[5]}] &= 1 + \mathbb{L} + 2\mathbb{L}^2 + 2\mathbb{L}^3 \\ [C^{[6]}] &= 1 + \mathbb{L} + 2\mathbb{L}^2 + 3\mathbb{L}^3 + \mathbb{L}^4 \\ [C^{[7]}] &= 1 + \mathbb{L} + 2\mathbb{L}^2 + 3\mathbb{L}^3 + \mathbb{L}^4 \\ [C^{[8]}] &= 1 + \mathbb{L} + 2\mathbb{L}^2 + 3\mathbb{L}^3 + 2\mathbb{L}^4 \end{aligned}$$

Then, we can compute the motivic Hilbert zeta function

$$Z_{(C,O)}^{Hilb}(q) = 1 + \sum_{\ell=1}^7 [C^{[\ell]}]q^{\ell} + [C^{[8]}](q^8 + q^9 + \dots)$$

Let $\Delta = (8) \in \mathcal{D}_8$ such that $\gamma_{\Delta} = \max\{\gamma_{\Delta}\}_{\Delta \in \mathcal{D}_c} = 15$. We find that $N \leq \gamma_{\Delta} + c - 1 = 22 \leq 42 = (c - 1)(\alpha_1 + 2)$ consistent with the result of Theorem 3.

Acknowledgments. We especially grateful to Mounir Hajli, another supervisor of Wenhao Zhu, deeply for his instruction on mathematics and his sustained encouragement to the first author and long term useful discussions and suggestions in this topic. We thank Ilaria Rossinelli for several very helpful discussions. We thank André Belotto da Silva for useful conversations and suggestions. The first author being supported by the funding from the Guangzhou Elites Sponsorship Council under the Oversea Study Program of Guangzhou Elite Project.

References

- [1] D. Bejleri, D. Ranganathan, R. Vakil, Motivic hilbert zeta functions of curves are rational, *Journal of the Institute of Mathematics of Jussieu* 19 (3) (2020) 947–964. doi:<https://doi.org/10.1017/S1474748018000269>.
- [2] M. Kapranov, The elliptic curve in the s-duality theory and eisenstein series for kac-moody groups, *arXiv preprint math/0001005* (2000). doi:<https://doi.org/10.48550/arXiv.math/0001005>.
- [3] G. Pfister, J. H. M. Steenbrink, Reduced hilbert schemes for irreducible curve singularities, *Journal of pure and applied algebra* 77 (1) (1992) 103–116. doi:[https://doi.org/10.1016/0022-4049\(92\)90033-C](https://doi.org/10.1016/0022-4049(92)90033-C).
- [4] J. Piontowski, Topology of the compactified jacobians of singular curves, *Mathematische Zeitschrift* 255 (2007) 195–226. doi:<https://doi.org/10.1007/s00209-006-0021-3>.
- [5] G. Lusztig, J. Smelt, Fixed point varieties on the space of lattices, *Bulletin of the London Mathematical Society* 23 (3) (1991) 213–218. doi:<https://doi.org/10.1112/blms/23.3.213>.
- [6] E. Gorsky, M. Mazin, A. Oblomkov, Generic curves and non-coprime catalans, *arXiv preprint arXiv:2210.12569* (2022).
- [7] E. Gorsky, M. Mazin, M. Vazirani, Affine permutations and rational slope parking functions, *Transactions of the American Mathematical Society* 368 (12) (2016) 8403–8445. doi:<https://doi.org/10.46298/dmtcs.2450>.

- [8] E. Gorsky, M. Mazin, Compactified jacobians and q, t -catalan numbers, i, *Journal of Combinatorial Theory, Series A* 120 (1) (2013) 49–63. doi:<https://doi.org/10.1016/j.jcta.2012.07.002>.
- [9] E. Gorsky, M. Mazin, Compactified jacobians and q, t -catalan numbers, ii, *Journal of Algebraic Combinatorics* 39 (2014) 153–186. doi:<https://doi.org/10.1007/s10801-013-0443-z>.
- [10] J. Hilburn, J. Kamnitzer, A. Weekes, Bfn springer theory, *Communications in Mathematical Physics* 402 (1) (2023) 765–832. doi:<https://doi.org/10.1007/s00220-023-04735-4>.
- [11] A. Oblomkov, J. Rasmussen, V. Shende, The hilbert scheme of a plane curve singularity and the homfly homology of its link, *Geometry & Topology* 22 (2) (2018) 645–691. doi:[DOI:10.2140/gt.2018.22.645](https://doi.org/10.2140/gt.2018.22.645).
- [12] Y. Soma, M. Watari, The punctual hilbert schemes for the curve singularities of types e_6 and e_8 , *Journal of Singularities* 8 (2014) 135–145. doi:[DOI:10.5427/jsing.2014.8j](https://doi.org/10.5427/jsing.2014.8j).
- [13] H. Mourtada, W. Zhu, On the geometry of punctual hilbert schemes of monomial curves, Preprint (2024).
- [14] Y. Soma, M. Watari, The punctual hilbert schemes for the curve singularities of type a_{2d} , *Journal of Singularities* 11 (2015) 152–163. doi:[DOI:10.5427/jsing.2015.11f](https://doi.org/10.5427/jsing.2015.11f).
- [15] M. Watari, Motivic hilbert zeta functions for curve singularities and related invariants, arXiv preprint [arXiv:2403.12545](https://arxiv.org/abs/2403.12545) (2024). doi:<https://doi.org/10.48550/arXiv.2403.12545>.
- [16] J.-P. Serre, *Groupes algébriques et corps de classes*, Thesis (1959).
- [17] D. Pol, *Singularités libres, formes et résidus logarithmiques*, Ph.D. thesis, Angers (2016).