

# Stabilities of intersecting families revisited\*

Yongjiang Wu<sup>a</sup>, Yongtao Li<sup>a</sup>, Lihua Feng<sup>a</sup>, Jiuqiang Liu<sup>b</sup>, Guihai Yu<sup>c,†</sup>

<sup>a</sup>School of Mathematics and Statistics, HNP-LAMA, Central South University  
Changsha, Hunan, 410083, China

<sup>b</sup>Department of Mathematics, Eastern Michigan University  
Ypsilanti, MI, 48197, USA

<sup>c</sup>College of Big Data Statistics, Guizhou University of Finance and Economics  
Guiyang, Guizhou, 550025, China

November 25, 2024

## Abstract

The well-known Erdős–Ko–Rado theorem states that for  $n > 2k$ , every intersecting family of  $k$ -sets of  $[n] := \{1, \dots, n\}$  has at most  $\binom{n-1}{k-1}$  sets, and the extremal family consists of all  $k$ -sets containing a fixed element (called a full star). The Hilton–Milner theorem provides a stability result by determining the maximum size of a uniform intersecting family that is not a subfamily of a full star. The further stabilities were studied by Han and Kohayakawa (2017) and Huang and Peng (2024). Two families  $\mathcal{F}$  and  $\mathcal{G}$  are called cross-intersecting if for every  $F \in \mathcal{F}$  and  $G \in \mathcal{G}$ , the intersection  $F \cap G$  is non-empty. Let  $k \geq 1, t \geq 0$  and  $n \geq 2k + t$  be integers. Frankl (2016) proved that if  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  are cross-intersecting families, and  $\mathcal{F}$  is non-empty and  $(t+1)$ -intersecting, then  $|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k-t}{k} + 1$ . Recently, Wu (2023) sharpened Frankl’s result by establishing a stability variant. The aim of this paper is two-fold. Inspired by the above results, we first prove a further stability variant that generalizes both Frankl’s result and Wu’s result. Secondly, as an interesting application, we illustrate that the aforementioned results on cross-intersecting families could be used to establish the stability results of the Erdős–Ko–Rado theorem. More precisely, we present new short proofs of the Hilton–Milner theorem, the Han–Kohayakawa theorem and the Huang–Peng theorem. Our arguments are more straightforward, and it may be of independent interest.

**AMS Classification:** 05C65; 05D05

**Key words:** Erdős–Ko–Rado theorem; Cross-intersecting; Stability

## 1 Introduction

For any two integers  $a \leq b$ , let  $[a, b] = \{a, a+1, \dots, b\}$  and simply let  $[n] = [1, n]$ . Let  $2^{[n]}$  denote the power set of  $[n]$ . For any  $0 \leq k \leq n$ , let  $\binom{[n]}{k}$  denote the collection of all its  $k$ -element subsets. A family  $\mathcal{F} \subseteq 2^{[n]}$  is called  $k$ -uniform if  $\mathcal{F} \subseteq \binom{[n]}{k}$ . A family  $\mathcal{F} \subseteq 2^{[n]}$  is called  $t$ -intersecting if  $|F \cap F'| \geq t$  for all  $F, F' \in \mathcal{F}$ . If  $t = 1$ ,  $\mathcal{F}$  is simply called *intersecting*. Two families  $\mathcal{F}, \mathcal{G} \subseteq 2^{[n]}$  are said to be *isomorphic* if there exists a permutation  $\sigma$  on  $[n]$  such that  $\mathcal{G} = \{\{\sigma(x) : x \in F\} : F \in \mathcal{F}\}$ , and we write  $\mathcal{F} \cong \mathcal{G}$ .

\*E-mail addresses: wuyjmath@163.com (Y. Wu), yтли0921@hnu.edu.cn (Y. Li), fenglh@163.com (L. Feng), jiuqiang68@126.com (J. Liu), yuguilai@126.com (G. Yu).

<sup>†</sup>Corresponding author

In 1961, Erdős, Ko and Rado [8] determined the maximum size of  $k$ -uniform intersecting families by proving that if  $k \geq 1$ ,  $n \geq 2k$  and  $\mathcal{F} \subseteq \binom{[n]}{k}$  is an intersecting family, then

$$|\mathcal{F}| \leq \binom{n-1}{k-1}.$$

For  $n > 2k$ , the equality holds if and only if  $\mathcal{F} = \{F \in \binom{[n]}{k} : x \in F\}$  for some  $x \in [n]$ . There are many different proofs and methods for proving the Erdős–Ko–Rado theorem; see, e.g., the Katona cycle method [30], the probabilistic method [1], the algebraic methods [20, 27, 35] and other combinatorial methods [6, 13, 28, 29, 33]. For more results on extremal set theory, we refer the interested readers to the comprehensive surveys [17, 7].

## 1.1 Cross-intersecting families

Two families  $\mathcal{F}, \mathcal{G} \subseteq 2^{[n]}$  are called *cross-intersecting* if  $|F \cap G| \geq 1$  for any  $F \in \mathcal{F}$  and  $G \in \mathcal{G}$ .

In 1967, Hilton and Milner [25] gave the following result.

**Theorem 1.1** (Hilton–Milner [25]). *Let  $k \geq 1$  and  $n \geq 2k$  be positive integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be non-empty cross-intersecting families. Suppose that  $|\mathcal{G}| \geq |\mathcal{F}|$ . Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k}{k} + 1.$$

*For  $n > 2k$ , the equality holds if and only if  $\mathcal{F} = \{F_1\}$  for some  $F_1 \in \binom{[n]}{k}$  and  $\mathcal{G} = \{G \in \binom{[n]}{k} : G \cap F_1 \neq \emptyset\}$ , or one more possibility when  $k = 2, \mathcal{F} = \mathcal{G} \cong \{G \in \binom{[n]}{2} : 1 \in G\}$ .*

This result initiated the study of finding the maximum of the sum of sizes of cross-intersecting families. There are many generalizations of Hilton–Milner’s result in the past years; see, e.g., [9, 11, 15, 16, 18, 22, 38, 39]. In particular, Frankl [9] gave the following inequality, which was applied to establish the stability result of Katona theorem [29].

**Theorem 1.2** (Frankl [9]). *Let  $k \geq 1, t \geq 0$  and  $n \geq 2k + t$  be integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be cross-intersecting families. If  $\mathcal{F}$  is  $(t+1)$ -intersecting and  $|\mathcal{F}| \geq 1$ , then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k-t}{k} + 1.$$

*For  $n > 2k + t$ , the equality holds if and only if  $\mathcal{F} = \{F_1\}$  for some  $F_1 \in \binom{[n]}{k+t}$  and  $\mathcal{G} = \{G \in \binom{[n]}{k} : G \cap F_1 \neq \emptyset\}$ , or one more possibility when  $k = 2, \mathcal{F} = \{[t+1] \cup \{i\} : i \in \{t+2, t+3, \dots, n\}\}$  and  $\mathcal{G} = \{G \in \binom{[n]}{k} : G \cap [t+1] \neq \emptyset\}$  under isomorphism.*

Recently, the second author and Wu [34] sharpened Theorem 1.2 in the case  $t = 1$  during the study on stabilities for non-uniform  $t$ -intersecting families. Subsequently, Wu [40] showed the following more general extension under the constraint  $|\mathcal{F}| \geq 2$ .

**Theorem 1.3** (Wu [40]). *Let  $k \geq 3, t \geq 0$  and  $n \geq 2k + t$  be integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be cross-intersecting families. If  $\mathcal{F}$  is  $(t+1)$ -intersecting and  $|\mathcal{F}| \geq 2$ , then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k-t}{k} - \binom{n-k-t-1}{k-1} + 2.$$

For  $n > 2k + t$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2\}$  for some  $F_1, F_2 \in \binom{[n]}{k+t}$  with  $|F_1 \cap F_2| = k + t - 1$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{k} : G \cap F_1 \neq \emptyset \text{ and } G \cap F_2 \neq \emptyset\right\}$ , or two more possibilities when  $k = 3$ , namely,  $\mathcal{F} = \{[t+2] \cup \{i\} : i \in \{t+3, t+4, \dots, n\}\}$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{3} : G \cap [t+2] \neq \emptyset\right\}$ , or  $\mathcal{F} = \left\{F \in \binom{[n]}{t+3} : [t+1] \subseteq F\right\}$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{3} : G \cap [t+1] \neq \emptyset\right\}$  under isomorphism.

Inspired by Theorems 1.2 and 1.3, we investigate the further stability for family  $\mathcal{F}$  with  $|\mathcal{F}| \geq 3$ . The main result of our paper is listed as below.

**Theorem 1.4** (Main result). *Let  $k \geq 4, t \geq 0$  and  $n \geq 2k + t$  be integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be cross-intersecting families. If  $\mathcal{F}$  is  $(t+1)$ -intersecting and  $|\mathcal{F}| \geq 3$ , then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k-t}{k} - \binom{n-k-t-1}{k-1} - \binom{n-k-t-2}{k-2} + 3.$$

For  $n > 2k+t$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2, F_3\}$  with  $F_i := [k+t-1] \cup \{k+t-1+i\}$  for each  $i \in [3]$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{k} : G \cap F_i \neq \emptyset \text{ for any } i = 1, 2, 3\right\}$ , or one more possibility when  $k = 4, \mathcal{F} = \{[t+3] \cup \{i\} : i \in \{t+4, \dots, n\}\}$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{4} : G \cap [t+3] \neq \emptyset\right\}$  under isomorphism.

**Organization.** The proof of Theorem 1.4 will be provided in Section 2. Before showing the proof, we would like to illustrate the significance of Theorem 1.4 for the stability results of the Erdős–Ko–Rado theorem. More precisely, we show that the above results on cross-intersecting families can be applied to prove the celebrated Hilton–Milner theorem, the Han–Kohayakawa theorem as well as the Huang–Peng theorem; see the forthcoming Theorems 1.5, 1.6 and 1.7, respectively. The second main contribution of this paper is to provide new short methods to prove these stability results. The detailed discussions will be presented in Section 3. To ensure our presentation clearly, some tedious arguments will be postponed to the Appendix.

## 1.2 Applications: Stabilities for Erdős–Ko–Rado’s theorem

In this section, we show that our result is extremely useful for the study of stabilities of  $k$ -uniform intersecting families. Indeed, we will provide a simple unified approach to proving several important stability results for the Erdős–Ko–Rado theorem; see, e.g., [25, 23, 26].

A family  $\mathcal{F}$  that satisfies  $\mathcal{F} = \{F \in \binom{[n]}{k} : x \in F\}$  is called a *full star* centered at  $x$ , and  $\mathcal{F}$  is called *EKR* or a *star* if  $\mathcal{F}$  is contained in a full star. Such family is a “trivial” example of intersecting family. If  $\mathcal{F}$  is not EKR, then  $\cap_{F \in \mathcal{F}} F = \emptyset$  and  $\mathcal{F}$  is called *non-trivial*.

In 1967, Hilton and Milner [25] determined the maximum size of a non-trivial  $k$ -uniform intersecting family. We denote  $\mathcal{HM}(n, k) = \{F \in \binom{[n]}{k} : 1 \in F, F \cap [2, k+1] \neq \emptyset\} \cup \{[2, k+1]\}$ . In the case  $k = 3$ , we denote  $\mathcal{T}(n, 3) = \{F \in \binom{[n]}{3} : |F \cap [3]| \geq 2\}$ .

**Theorem 1.5** (Hilton–Milner [25]). *Let  $k \geq 2$  and  $n \geq 2k + 1$  be integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  be an intersecting family. If  $\mathcal{F}$  is not EKR, then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1.$$

For  $k = 2$  or  $k \geq 4$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{HM}(n, k)$ ; For  $k = 3$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{HM}(n, 3)$  or  $\mathcal{T}(n, 3)$ .

A family  $\mathcal{F}$  is called *HM* if  $\mathcal{F}$  is isomorphic to a subfamily of  $\mathcal{HM}(n, k)$ . In 2017, Han and Kohayakawa [23] took stability further and determined the maximum  $k$ -uniform intersecting family

that is neither EKR nor HM, and not a subfamily of  $\mathcal{T}(n, 3)$  if  $k = 3$ . We shall mention that for  $k \geq 4$ , this was already solved by Hilton and Milner [25] in 1967. Before introducing their result, let us describe some set families.

Let  $k \geq 3$  and  $n \geq 2k + 1$  be integers. Let  $i \in [1, k - 1]$ . For any  $E \in \binom{[n]}{k-1}$ , any  $J \in \binom{[n] \setminus E}{i+1}$ ,  $x_0 \in J$  and  $J_i = J \setminus \{x_0\}$ , we define

$$\mathcal{J}_i(n, k) = \left\{ F \in \binom{[n]}{k} : x_0 \in F, F \cap (E \cup \{j\}) \neq \emptyset \text{ for each } j \in J_i \right\} \cup \{E \cup \{j\} : j \in J_i\}.$$

Furthermore, let  $i \in [2, k]$ . For any  $E \in \binom{[n]}{i}$  and  $x_0 \in [n] \setminus E$ , we define

$$\mathcal{G}_i(n, k) = \left\{ G \in \binom{[n]}{k} : E \subseteq G \right\} \cup \left\{ G \in \binom{[n]}{k} : x_0 \in G, G \cap E \neq \emptyset \right\}.$$

Note that  $\mathcal{G}_2(n, 3)$  is isomorphic to  $\mathcal{T}(n, 3)$ .

**Theorem 1.6** (Han–Kohayakawa [23]). *Let  $k \geq 3$  and  $n \geq 2k + 1$  be integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  be an intersecting family. If  $\mathcal{F}$  is neither EKR nor HM, and  $\mathcal{F} \not\subseteq \mathcal{G}_2(n, 3)$  for  $k = 3$ . Then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 2.$$

*For  $k = 3$  or  $k \geq 5$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{J}_2(n, k)$ ; For  $k = 4$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{J}_2(n, 4)$ ,  $\mathcal{G}_2(n, 4)$  or  $\mathcal{G}_3(n, 4)$ .*

Han and Kohayakawa [23] also asked what is the maximum  $k$ -uniform intersecting family which is neither EKR nor HM, and not a subfamily of  $\mathcal{J}_2(n, k)$ , and not a subfamily of  $\mathcal{G}_2(n, 4)$ ,  $\mathcal{G}_3(n, 4)$  if  $k = 4$ . In 2017, Kostochka and Mubayi [31] answered this question for large enough  $n$ . In 2024, Huang and Peng [26] completely solved this question for all  $n \geq 2k + 1$ .

To state their result, we define the family  $\mathcal{K}_2(n, k)$  as follows. For any  $E_1, E_2 \in \binom{[n]}{k}$  with  $|E_1 \cap E_2| = k - 2$ , and  $x_0 \in [n] \setminus (E_1 \cup E_2)$ , let

$$\mathcal{K}_2(n, k) = \left\{ G \in \binom{[n]}{k} : x_0 \in G, G \cap E_1 \neq \emptyset \text{ and } G \cap E_2 \neq \emptyset \right\} \cup \{E_1, E_2\}.$$

**Theorem 1.7** (Huang–Peng [26]). *Let  $k \geq 4$  and  $\mathcal{F} \subseteq \binom{[n]}{k}$  be an intersecting family which is neither EKR nor HM, and  $\mathcal{F} \not\subseteq \mathcal{J}_2(n, k)$ , in addition,  $\mathcal{F} \not\subseteq \mathcal{G}_2(n, 4)$  and  $\mathcal{F} \not\subseteq \mathcal{G}_3(n, 4)$  if  $k = 4$ .*

*(i) If  $2k + 1 \leq n \leq 3k - 3$ , then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1} - 2 \binom{n-k-1}{k-1} + \binom{n-k-3}{k-1} + 2.$$

*For  $k \geq 5$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{K}_2(n, k)$ ; For  $k = 4$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{K}_2(n, 4)$  or  $\mathcal{J}_3(n, 4)$ .*

*(ii) If  $n \geq 3k - 2$ , then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3.$$

*For  $k = 4$  or  $k \geq 6$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{J}_3(n, k)$ ; For  $k = 5$ , the equality holds if and only if  $\mathcal{F}$  is isomorphic to  $\mathcal{J}_3(n, 5)$  or  $\mathcal{G}_4(n, 5)$ .*

Apart from the above stabilities, there are a large number of results involving the stabilities of  $t$ -intersecting families; see, e.g., [2, 4, 5, 21, 34, 37] and references therein. As mentioned before, various methods involving the celebrated Erdős–Ko–Rado theorem are presented in the literature. Motivated by these methods, there are also many alternative proofs for the Hilton–Milner theorem; see, e.g., [3, 10, 12, 15, 28, 33]. For the Han–Kohayakawa theorem, Kupavskii [32] gave an alternative proof only in the case  $k \geq 5$ . Unfortunately, the cases  $k = 3$  and  $k = 4$  can not be proven by following his method in a straightforward way. The arguments in these cases are relatively more complicated since some exceptional extremal families are appeared. Very recently, Ge, Xu and Zhao [22] provided an alternative proof using a robust linear algebra method, and they also obtained the  $t$ -th level stability result for the Erdős–Ko–Rado theorem. It seems that the original proofs of Han and Kohayakawa [23] and Huang and Peng [26] are quite technical. Therefore, it is interesting to give new short proofs of the Han–Kohayakawa theorem as well as the Huang–Peng theorem.

The second goal of this paper is to show the applications of aforementioned results involving cross-intersecting families to the stability results for the Erdős–Ko–Rado theorem. We will give short proofs for the Hilton–Milner theorem, the Han–Kohayakawa theorem and the Huang–Peng theorem in a unified way. Our approach is based on applying Theorems 1.2, 1.3 and 1.4, respectively. The main purpose of this paper is to present a unified simple approach to proving the stability results for intersecting families, so we will not completely expand on these full arguments in the case  $k = 3$  or 4. For interested readers, we will provide detailed proofs in the Appendix using our unified framework. In addition, it seems feasible to further extend our result on cross-intersecting families to give a short proof of the stability result in [22], we do not realize the details here, since the argument for small integer  $k$  is a little bit more involved.

## 2 Proof of Theorem 1.4

To prove Theorem 1.4, we review some fundamental notations and results about the shifting operation. Let  $\mathcal{F} \subseteq 2^{[n]}$  be a family and  $1 \leq i < j \leq n$ . The *shifting operator*  $s_{i,j}$ , discovered by Erdős, Ko and Rado [8], is defined as  $s_{i,j}(\mathcal{F}) = \{s_{i,j}(F) : F \in \mathcal{F}\}$ , where

$$s_{i,j}(F) = \begin{cases} (F \setminus \{j\}) \cup \{i\} & \text{if } j \in F, i \notin F \text{ and } (F \setminus \{j\}) \cup \{i\} \notin \mathcal{F}, \\ F & \text{otherwise.} \end{cases}$$

Obviously, we have  $|s_{i,j}(F)| = |F|$  and  $|s_{i,j}(\mathcal{F})| = |\mathcal{F}|$ . A frequently used property is that  $s_{i,j}$  maintains the  $t$ -intersecting property of a family. A family  $\mathcal{F} \subseteq 2^{[n]}$  is called *shifted* if for all  $F \in \mathcal{F}$ ,  $i < j$  with  $i \notin F$  and  $j \in F$ , then  $(F \setminus \{j\}) \cup \{i\} \in \mathcal{F}$ . It is well-known that every intersecting family can be transformed to a shifted intersecting family by applying shifting operations repeatedly. There are many nice properties for shifted families. For example, if  $\mathcal{F}$  is a shifted family and  $\{a_1, \dots, a_k\} \in \mathcal{F}$  with  $a_1 < \dots < a_k$ , then for any set  $\{b_1, \dots, b_k\}$  with  $b_1 < \dots < b_k$  and  $b_i \leq a_i$  for each  $i \in [1, k]$ , we have  $\{b_1, \dots, b_k\} \in \mathcal{F}$ . For  $\mathcal{F} \subseteq 2^{[n]}$  and  $i \in [n]$ , we denote  $\mathcal{F}(\bar{i}) = \{F \in \mathcal{F} : i \notin F\}$  and  $\mathcal{F}(i) = \{F \setminus \{i\} : i \in F \in \mathcal{F}\}$ .

**Lemma 2.1.** *Let  $k \geq 1, t \geq 0$  be integers. Let  $\mathcal{F} \subseteq 2^{[n]}$  be a shifted  $(t+1)$ -intersecting family. Then  $\mathcal{F}(\bar{1})$  is  $(t+2)$ -intersecting. Moreover, if  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  is a shifted  $(t+1)$ -intersecting family and  $n \geq 2k+t$ , then  $\mathcal{F}(n)$  is  $(t+1)$ -intersecting.*

**Proof.** We may assume that  $\mathcal{F}(\bar{1}) \neq \emptyset$ . For any  $F_1, F_2 \in \mathcal{F}(\bar{1})$ , let  $j \in F_1 \cap F_2$ . By shiftedness, we have  $(F_1 \setminus \{j\}) \cup \{1\} \in \mathcal{F}$ . It follows from the  $(t+1)$ -intersecting property of  $\mathcal{F}$  that  $|F_1 \cap F_2| = |((F_1 \setminus \{j\}) \cup \{1\}) \cap F_2| + 1 \geq t+2$ . In addition, we may assume that  $\mathcal{F}(n) \neq \emptyset$ . For any  $E_1, E_2 \in \mathcal{F}(n)$ , we have  $E_1 \cup \{n\}, E_2 \cup \{n\} \in \mathcal{F}$ . Observe that  $|E_1 \cup E_2 \cup \{n\}| \leq 2(k+t) - (t+1) \leq n-1$ .

So there exists  $x \notin E_1 \cup E_2 \cup \{n\}$  such that  $E_1 \cup \{x\} \in \mathcal{F}$ . It is immediate that  $|E_1 \cap E_2| = |(E_1 \cup \{x\}) \cap (E_2 \cup \{n\})| \geq t + 1$ , as desired.  $\square$

**Lemma 2.2.** *Let  $k \geq 1, t \geq 0$  and  $n \geq k + t + 3$  be integers. If  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  is shifted and  $|\mathcal{F}| \geq 3$ , then  $|\mathcal{F}(\bar{n})| \geq 3$ .*

**Proof.** If  $\mathcal{F}(n) = \emptyset$ , then  $|\mathcal{F}(\bar{n})| = |\mathcal{F}| \geq 3$ . If  $\mathcal{F}(n) \neq \emptyset$ , then there exists  $F \in \mathcal{F}$  such that  $n \in F$ . Note that  $n \geq k + t + 3$ . So there are at least three different elements  $x, y, z \in [n] \setminus F$ . By shiftedness,  $(F \setminus \{n\}) \cup \{x\}, (F \setminus \{n\}) \cup \{y\}, (F \setminus \{n\}) \cup \{z\} \in \mathcal{F}(\bar{n})$ , and the result follows.  $\square$

**Lemma 2.3** (See [8]). *Let  $\mathcal{F} \subseteq 2^{[n]}$  and  $\mathcal{G} \subseteq 2^{[n]}$  be cross-intersecting, and  $\mathcal{F}$  be  $t$ -intersecting. Then  $s_{i,j}(\mathcal{F})$  and  $s_{i,j}(\mathcal{G})$  are also cross-intersecting, and  $s_{i,j}(\mathcal{F})$  is  $t$ -intersecting.*

**Lemma 2.4.** *Let  $t$  be an non-negative integer. Let  $k \geq 1$  and  $n \geq 2k + t$  be positive integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k+t}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be shifted cross-intersecting families. Then  $\mathcal{F}(n)$  and  $\mathcal{G}(n)$  are cross-intersecting.*

**Proof.** The case for  $\mathcal{F}(n) = \emptyset$  or  $\mathcal{G}(n) = \emptyset$  is trivial. If  $\mathcal{F}(n) \neq \emptyset$  and  $\mathcal{G}(n) \neq \emptyset$ , then there are  $F_1 \in \mathcal{F}(n)$  and  $G_1 \in \mathcal{G}(n)$  such that  $F_1 \cup \{n\} \in \mathcal{F}$  and  $G_1 \cup \{n\} \in \mathcal{G}$ . Observe that  $|F_1 \cup G_1 \cup \{n\}| \leq 2k + t - 1 \leq n - 1$ . So there exists  $x \notin F_1 \cup G_1 \cup \{n\}$  such that  $F_1 \cup \{x\} \in \mathcal{F}$ . Then we have  $|F_1 \cap G_1| = |(F_1 \cup \{x\}) \cap (G_1 \cup \{n\})| \geq 1$ , as desired.  $\square$

Next we define the lexicographic order on the  $k$ -element subsets of  $[n]$ . We say that  $F$  is smaller than  $G$  in the lexicographic order if  $\min(F \setminus G) < \min(G \setminus F)$  holds. For  $0 \leq m \leq \binom{n}{k}$ , let  $\mathcal{L}(n, k, m)$  be the family of the first  $m$   $k$ -sets in the lexicographic order. The following lemma will be used in the next section, for convenience, we list it below; see [14, p.266] for a detailed proof.

**Lemma 2.5** (See [24]). *Let  $k, \ell, n$  be positive integers with  $n > k + \ell$ . If  $\mathcal{F} \subseteq \binom{[n]}{k}$  and  $\mathcal{G} \subseteq \binom{[n]}{\ell}$  are cross-intersecting, then  $\mathcal{L}(n, k, |\mathcal{F}|)$  and  $\mathcal{L}(n, \ell, |\mathcal{G}|)$  are cross-intersecting.*

**Proof of Theorem 1.4.** For fixed  $k$  and  $t$ , we apply induction on  $n \geq 2k + t$ . First let us consider the base case  $n = 2k + t$ . For any  $F \in \binom{[2k+t]}{k+t}$ , the cross-intersecting property of  $\mathcal{F}$  and  $\mathcal{G}$  implies that  $F \notin \mathcal{F}$  or  $[2k + t] \setminus F \notin \mathcal{G}$ . It follows that  $|\mathcal{F}| + |\mathcal{G}| \leq \binom{2k+t}{k}$ , as desired.

Now suppose that  $n \geq 2k + t + 1$  and the result holds for integers less than  $n$  and fixed  $k$  and  $t$ . By Lemma 2.3, we may assume that  $\mathcal{F}$  and  $\mathcal{G}$  are shifted. It follows from Lemma 2.2 that  $|\mathcal{F}(\bar{n})| \geq 3$ . Clearly,  $\mathcal{F}(\bar{n})$  is  $(t + 1)$ -intersecting. Note that  $\mathcal{F}(\bar{n})$  and  $\mathcal{G}(\bar{n})$  are cross-intersecting. By induction hypothesis, we get

$$|\mathcal{F}(\bar{n})| + |\mathcal{G}(\bar{n})| \leq \binom{n-1}{k} - \binom{n-k-t-1}{k} - \binom{n-k-t-2}{k-1} - \binom{n-k-t-3}{k-2} + 3.$$

We proceed the proof by considering the following two cases.

**Case 1.** Suppose that  $|\mathcal{F}(n)| = 0$ .

Since  $|\mathcal{F}(\bar{n})| \geq 3$ , there exist  $F_1, F_2, F_3 \in \mathcal{F}(\bar{n}) = \mathcal{F}$ . For any permutation  $\sigma$  on  $[3]$ , we consider the following families:

$$\begin{aligned} \mathcal{G}_{\sigma(1)} &= \left\{ G \in \binom{[n-1]}{k-1} : G \cap F_{\sigma(1)} = \emptyset \right\}, \\ \mathcal{G}_{\sigma(2)} &= \left\{ G \in \binom{[n-1]}{k-1} : G \cap F_{\sigma(1)} \neq \emptyset, G \cap F_{\sigma(2)} = \emptyset \right\}, \\ \mathcal{G}_{\sigma(3)} &= \left\{ G \in \binom{[n-1]}{k-1} : G \cap F_{\sigma(1)} \neq \emptyset, G \cap F_{\sigma(2)} \neq \emptyset, G \cap F_{\sigma(3)} = \emptyset \right\}. \end{aligned}$$

Notice that  $|\mathcal{G}_{\sigma(1)}| = \binom{n-k-t-1}{k-1}$  and  $|\mathcal{G}_{\sigma(2)}| \geq \binom{n-k-t-2}{k-2}$ . In addition, there exist  $x_1 \in \mathcal{F}_{\sigma(1)}$  and  $x_2 \in \mathcal{F}_{\sigma(2)}$  such that  $x_1, x_2 \notin \mathcal{F}_{\sigma(3)}$ . If  $x_1 \neq x_2$ , then  $|\mathcal{G}_{\sigma(3)}| \geq \binom{n-k-t-3}{k-3}$ ; if  $x_1 = x_2$ , then  $|\mathcal{G}_{\sigma(3)}| \geq \binom{n-k-t-2}{k-2}$ . So  $|\mathcal{G}_{\sigma(3)}| \geq \binom{n-k-t-3}{k-3}$  holds in all cases. Since  $\mathcal{G}_{\sigma(i)} \cap \mathcal{G}_{\sigma(j)} = \emptyset$  for any  $i \neq j$  and  $\mathcal{F} = \mathcal{F}(\bar{n})$  and  $\mathcal{G}(n)$  are cross-intersecting, we have

$$|\mathcal{G}(n)| \leq \binom{n-1}{k-1} - \sum_{j=1}^3 |\mathcal{G}_{\sigma(j)}| \leq \binom{n-1}{k-1} - \binom{n-k-t-1}{k-1} - \binom{n-k-t-2}{k-2} - \binom{n-k-t-3}{k-3}.$$

Next suppose that the equality holds. Then  $\mathcal{F}(\bar{n}) = \{F_1, F_2, F_3\}$  and for any permutation  $\sigma$  on  $[3]$ , we have  $|\mathcal{G}_{\sigma(2)}| = \binom{n-k-t-2}{k-2}$  and  $|\mathcal{G}_{\sigma(3)}| = \binom{n-k-t-3}{k-3}$ . If there exist  $i \neq j \in [3]$  such that  $|F_i \cap F_j| \leq k+t-2$ , then let  $\sigma = (1i)(2j)$ . So there exist  $y_1 \neq y_2 \in \mathcal{F}_{\sigma(1)}$  such that  $y_1, y_2 \notin \mathcal{F}_{\sigma(2)}$ . This implies that  $|\mathcal{G}_{\sigma(2)}| \geq \binom{n-k-t-2}{k-2} + \binom{n-k-t-3}{k-2}$ , a contradiction. Thus  $\mathcal{F}(\bar{n}) = \{F_1, F_2, F_3\}$  with  $|F_i \cap F_j| = k+t-1$  for any  $1 \leq i < j \leq 3$ . We claim that  $|\cap_{i=1}^3 F_i| = k+t-1$ . Otherwise we can write  $F_1 = M \cup \{a, b\}, F_2 = M \cup \{a, c\}, F_3 = M \cup \{b, c\}$ , where  $|M| = k+t-2$  and  $a, b, c$  are pairwise different. But then for any permutation  $\sigma$  on  $[3]$ , we have  $|\mathcal{G}_{\sigma(3)}| \geq \binom{n-k-t-2}{k-2}$ , a contradiction. Therefore, by the shiftedness of  $\mathcal{F}$ , we conclude that the equality holds if and only if  $\mathcal{F}(\bar{n}) = \{F_1, F_2, F_3\}$  with  $F_i := [k+t-1] \cup \{k+t-1+i\}$  for each  $i \in [3]$  and  $\mathcal{G}(n) = \left\{ G \in \binom{[n-1]}{k-1} : G \cap F_i \neq \emptyset \text{ for any } i = 1, 2, 3 \right\}$ . It follows that

$$|\mathcal{F}| + |\mathcal{G}| = |\mathcal{F}(\bar{n})| + |\mathcal{G}(\bar{n})| + |\mathcal{G}(n)| \leq \binom{n}{k} - \binom{n-k-t}{k} - \binom{n-k-t-1}{k-1} - \binom{n-k-t-2}{k-2} + 3.$$

The equality holds if and only if  $\mathcal{F} = \{F_1, F_2, F_3\}$  with  $F_i := [k+t-1] \cup \{k+t-1+i\}$  for each  $i \in [3]$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{k} : G \cap F_i \neq \emptyset \text{ for any } i \in [3] \right\}$ .

**Case 2.** Suppose that  $|\mathcal{F}(n)| \geq 1$ .

By Lemma 2.4,  $\mathcal{F}(n) \subseteq \binom{[n-1]}{k+t-1}$  and  $\mathcal{G}(n) \subseteq \binom{[n-1]}{k-1}$  are cross-intersecting. By Lemma 2.1,  $\mathcal{F}(n)$  is  $(t+1)$ -intersecting. Since  $n \geq 2k+t+1$  and  $k-1 \geq 3$ , applying Theorem 1.2 to  $\mathcal{F}(n)$  and  $\mathcal{G}(n)$  yields

$$|\mathcal{F}(n)| + |\mathcal{G}(n)| \leq \binom{n-1}{k-1} - \binom{n-k-t}{k-1} + 1.$$

By Theorem 1.2 and the shiftedness of  $\mathcal{F}$ , the equality holds if and only if  $\mathcal{F}(n) = \{[k+t-1]\}$  and  $\mathcal{G}(n) = \left\{ G^* \in \binom{[n-1]}{k-1} : G^* \cap [k+t-1] \neq \emptyset \right\}$ . Note that  $\mathcal{F}(\bar{n})$  and  $\mathcal{G}(n)$  are cross-intersecting. By the shiftedness of  $\mathcal{F}$ , if  $\mathcal{F}(n) = \{[k+t-1]\}$  and  $\mathcal{G}(n) = \left\{ G^* \in \binom{[n-1]}{k-1} : G^* \cap [k+t-1] \neq \emptyset \right\}$ , then  $\mathcal{F}(\bar{n}) = \{[k+t-1] \cup \{i\} : i \in \{k+t, \dots, n-1\}\}$ ,  $\mathcal{G}(\bar{n}) = \left\{ G \in \binom{[n-1]}{k} : G \cap [k+t-1] \neq \emptyset \right\}$ .

Therefore, we conclude that

$$\begin{aligned} |\mathcal{F}| + |\mathcal{G}| &= |\mathcal{F}(\bar{n})| + |\mathcal{G}(\bar{n})| + |\mathcal{F}(n)| + |\mathcal{G}(n)| \\ &\leq \binom{n-1}{k} - \binom{n-k-t-1}{k} - \binom{n-k-t-2}{k-1} - \binom{n-k-t-3}{k-2} + 3 \\ &\quad + \binom{n-1}{k-1} - \binom{n-k-t}{k-1} + 1 \\ &= \binom{n}{k} - \binom{n-k-t}{k} - \binom{n-k-t-1}{k-1} - \binom{n-k-t-2}{k-2} + 3 + 1 - \binom{n-k-t-3}{k-4} \\ &\leq \binom{n}{k} - \binom{n-k-t}{k} - \binom{n-k-t-1}{k-1} - \binom{n-k-t-2}{k-2} + 3. \end{aligned}$$

If the equality holds, then  $1 = \binom{n-k-t-3}{k-4}$ , i.e.,  $k = 4$ , and  $\mathcal{F} = \{[t+3] \cup \{i\} : i \in \{t+4, \dots, n\}\}$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{4} : G \cap [t+3] \neq \emptyset \right\}$ .

In the above discussion, we have determined the extremal families  $\mathcal{F}$  and  $\mathcal{G}$  that attain the required upper bound under the shifting assumption by Lemma 2.3. For completeness, we also need to characterize the extremal families in general case. Assume that the extremal families  $\mathcal{F}'$  and  $\mathcal{G}'$  satisfy  $s_{i,j}(\mathcal{F}') = \mathcal{F}$  and  $s_{i,j}(\mathcal{G}') = \mathcal{G}$ . Note that  $i < j$ . When  $\mathcal{F} = \{F_1, F_2, F_3\}$  with  $F_i := [k+t-1] \cup \{k+t-1+i\}$  for each  $i \in [3]$ , if  $i, j \in [k+t-1]$ , then  $\mathcal{F}' = \mathcal{F}$ . If  $i \in [k+t-1]$  and  $j \in [k+t, k+t+2]$ , by symmetry, we assume that  $i = 1$  and  $j = k+t$ , then  $\mathcal{F}' = \{[2, k+t] \cup \{q\} : q \in \{1, k+t+1, k+t+2\}\}$ . Thus  $\mathcal{F}' \cong \mathcal{F}$ . If  $i \in [k+t-1]$  and  $j \notin [1, k+t+2]$ , we may assume that  $i = 1$ , then  $\mathcal{F}' = \{[2, k+t-1] \cup \{q, j\} : q \in [k+t, k+t+2]\}$ . Thus  $\mathcal{F}' \cong \mathcal{F}$ . If  $i, j \in [k+t, k+t+2]$ , then  $\mathcal{F}' = \mathcal{F}$ . If  $i \in [k+t, k+t+2]$  and  $j \notin [1, k+t+2]$ , we may assume that  $i = k+t$  and  $j = k+t+3$ , then  $\mathcal{F}' = \{[1, k+t-1] \cup \{q\} : q \in [k+t+1, k+t+3]\}$ . Thus  $\mathcal{F}' \cong \mathcal{F}$ . If  $i, j \notin [1, k+t+2]$ , then  $\mathcal{F}' = \mathcal{F}$ . Therefore, we conclude that in the case  $\mathcal{F} = \{F_1, F_2, F_3\}$  with  $F_i := [k+t-1] \cup \{k+t-1+i\}$  for each  $i \in [3]$ , we have  $\mathcal{F}' \cong \mathcal{F}$ . The family  $\mathcal{G}'$  is just the maximal subfamily of  $\binom{[n]}{k}$  that is cross-intersecting with  $\mathcal{F}'$ . The same is true when  $\mathcal{F}$  is the other case. Let  $\mathcal{F}_0$  and  $\mathcal{G}_0$  be the original families before applying the shifting operations. In other words,  $\mathcal{F}$  and  $\mathcal{G}$  are obtained from  $\mathcal{F}_0$  and  $\mathcal{G}_0$  by applying a series of shifting operations. Then we have  $\mathcal{F}_0 = \mathcal{F}$  and  $\mathcal{G}_0 = \mathcal{G}$ , under isomorphism.  $\square$

### 3 Revisiting stabilities of intersecting families

As promised, we will present a unified framework to revisit several stability results for the Erdős–Ko–Rado theorem. To begin with, we fix the following notations.

$$\begin{aligned}\mathcal{F}(i, j) &= \{F \setminus \{i, j\} : i, j \in F \in \mathcal{F}\}, \\ \mathcal{F}(i, \bar{j}) &= \mathcal{F}(\bar{j}, i) = \{F \setminus \{i\} : i \in F, j \notin F, F \in \mathcal{F}\}, \\ \mathcal{F}(\bar{i}, \bar{j}) &= \{F : i, j \notin F \in \mathcal{F}\}.\end{aligned}$$

#### 3.1 Short proof of the Hilton–Milner theorem

To more clearly illustrate our approach, we first give a short new proof of Hilton–Milner’s bound in Theorem 1.5. It is worth mentioning that Frankl and Tokushige [15] provided a short proof of the bound in Theorem 1.5 by using the Katona theorem on shadows for intersecting families [29]. Different from their approach, we here present a short proof of Theorem 1.5 by using Theorem 1.2 only, avoiding the use of the Katona intersecting shadow theorem.

**Proof of Theorem 1.5.** Suppose that  $\mathcal{F}$  is not EKR and  $|\mathcal{F}|$  is maximal. As is well-known, the shifting operation preserves the intersecting property. Applying shifting operations to  $\mathcal{F}$  repeatedly, we can arrive at the following two cases:

- (A) After applying all shifting operations, we get a shifted family that is not EKR.
- (B) After some shifting operations, at some point, the resulting family is EKR.

**Case (A).** Suppose that  $\mathcal{H}$  is obtained from  $\mathcal{F}$  by applying all shifts and  $\mathcal{H}$  is not EKR. Then we obtain  $\mathcal{H}(\bar{1}) \neq \emptyset$ . By Lemma 2.1,  $\mathcal{H}(\bar{1})$  is 2-intersecting. Note that  $\mathcal{H}(\bar{1}) \subseteq \binom{[2, n]}{k-1}$  and  $\mathcal{H}(1) \subseteq \binom{[2, n]}{k-1}$  are cross-intersecting. Applying Theorem 1.2 (by setting  $t = 1$ ), we get

$$|\mathcal{F}| = |\mathcal{H}| = |\mathcal{H}(\bar{1})| + |\mathcal{H}(1)| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1.$$

Moreover, the above equality holds if and only if  $\mathcal{H}(\bar{1}) = \{[2, k+1]\}$  and  $\mathcal{H}(1) = \{H \in \binom{[2, n]}{k-1} : H \cap [2, k+1] \neq \emptyset\}$ , or in the case  $k = 3$ , there is one more possibility:  $\mathcal{H}(\bar{1}) = \{\{2, 3\} \cup \{i\} : 4 \leq i \leq n\}$



and  $\mathcal{H}(1) = \{H \in \binom{[2,n]}{2} : H \cap \{2, 3\} \neq \emptyset\}$ . Consequently, we get  $\mathcal{H} = \mathcal{HM}(n, k)$ , or  $\mathcal{H} = \mathcal{T}(n, 3)$ . It is easy to check that if  $\mathcal{H}'$  is an extremal family and  $s_{i,j}(\mathcal{H}') = \mathcal{H}$ , then  $\mathcal{H}' \cong \mathcal{H}$ .

**Case (B).** Let  $\mathcal{G} \subseteq \binom{[n]}{k}$  be obtained from  $\mathcal{F}$  by applying some shifting operations such that  $\mathcal{G}$  is not EKR, but after the shifting operation  $s_{i,j}$ , the resulting family  $s_{i,j}(\mathcal{G})$  is EKR. In this case, all sets of  $s_{i,j}(G)$  contain the element  $i$ , it follows that  $\mathcal{G}(\bar{i}, \bar{j}) = \emptyset$  and  $\{i, j\} \cap G \neq \emptyset$  for every  $G \in \mathcal{G}$ . By the maximality of  $|\mathcal{G}|$ , we have  $\mathcal{G}(i, j) = \binom{[n] \setminus \{i, j\}}{k-2}$ . Note that  $\mathcal{G}(i, \bar{j}) \subseteq \binom{[n] \setminus \{i, j\}}{k-1}$  and  $\mathcal{G}(\bar{i}, j) \subseteq \binom{[n] \setminus \{i, j\}}{k-1}$  are non-empty cross-intersecting. So Theorem 1.1 can be applied. Since  $\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j}) = \emptyset$ , the inequality in Theorem 1.1 holds strictly. Then

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}| = |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ &< \binom{n-2}{k-2} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} + 1 \\ &= \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1, \end{aligned}$$

as desired. This completes the proof of Theorem 1.5.  $\square$

### 3.2 Short proof of the Han–Kohayakawa theorem

To make the proof clearer, we divide it into case  $k \geq 4$  and case  $k = 3$ . Firstly, let us consider the case  $k \geq 4$  and present some tools as follows.

For two integers  $k, \ell$  with  $n \geq k + \ell$  and a family  $\mathcal{F} \subseteq \binom{[n]}{k}$ , we define

$$\mathcal{D}_\ell(\mathcal{F}) = \left\{ D \in \binom{[n]}{\ell} : \exists F \in \mathcal{F}, D \cap F = \emptyset \right\}.$$

Then  $\mathcal{F}$  and  $\mathcal{G} \subseteq \binom{[n]}{\ell}$  are cross-intersecting if and only if  $\mathcal{F} \cap \mathcal{D}_k(\mathcal{G}) = \emptyset$  or equivalently  $\mathcal{G} \cap \mathcal{D}_\ell(\mathcal{F}) = \emptyset$ . Moreover, for given  $\mathcal{F}$ ,  $\mathcal{G} = \binom{[n]}{\ell} \setminus \mathcal{D}_\ell(\mathcal{F})$  is the largest family that is cross-intersecting with  $\mathcal{F}$ , and for given  $\mathcal{G}$ ,  $\mathcal{F} = \binom{[n]}{k} \setminus \mathcal{D}_k(\mathcal{G})$  is the largest family that is cross-intersecting with  $\mathcal{G}$ .

We need the following lemma to characterize the uniqueness of extremal families.

**Lemma 3.1** (See [19, 36]). *Let  $n > k + \ell$ ,  $\mathcal{F} \subseteq \binom{[n]}{k}$  and  $|\mathcal{F}| = \binom{n-r}{k-r}$  for some  $r \leq k$ . Then  $|\mathcal{D}_\ell(\mathcal{F})| \geq \binom{n-r}{\ell}$ , with strict inequality unless  $\mathcal{F} = \{F \in \binom{[n]}{k} : R \subseteq F\}$  for some  $R \in \binom{[n]}{r}$ .*

The following lemma is a direct consequence of Theorem 1.3.

**Lemma 3.2.** *Let  $k \geq 3$  and  $n \geq 2k$  be positive integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be cross-intersecting families. Suppose that  $|\mathcal{G}| \geq |\mathcal{F}| \geq 2$ , then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k}{k} - \binom{n-k-1}{k-1} + 2.$$

*For  $n > 2k$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2\}$  for some  $F_1, F_2 \in \binom{[n]}{k}$  with  $|F_1 \cap F_2| = k-1$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{k} : G \cap F_1 \neq \emptyset \text{ and } G \cap F_2 \neq \emptyset \right\}$ , or two more possibilities when  $k = 3$ , namely,  $\mathcal{F} = \{[2] \cup \{i\} : i \in \{3, 4, \dots, n\}\}$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{3} : G \cap [2] \neq \emptyset \right\}$ , or  $\mathcal{F} = \mathcal{G} = \left\{ G \in \binom{[n]}{3} : 1 \in G \right\}$ , under isomorphism.*

**Proof.** If  $n = 2k$ , then  $\binom{n}{k} - \binom{n-k}{k} - \binom{n-k-1}{k-1} + 2 = \binom{2k}{k}$ . Hence the upper bound for  $n = 2k$  is trivial. Suppose that  $n > 2k$ . We denote  $\mathcal{F}' := \mathcal{L}(n, k, |\mathcal{F}|)$  and  $\mathcal{G}' := \mathcal{L}(n, k, |\mathcal{G}|)$ . By Lemma

2.5, we know that  $\mathcal{F}'$  and  $\mathcal{G}'$  are cross-intersecting. Since  $|\mathcal{G}'| \geq |\mathcal{F}'| \geq r$ , we have  $\mathcal{F}' \subseteq \mathcal{G}'$ , which implies that  $\mathcal{F}'$  is intersecting. Then setting  $t = 0$  in Theorem 1.3 yields

$$|\mathcal{F}| + |\mathcal{G}| = |\mathcal{F}'| + |\mathcal{G}'| \leq \binom{n}{k} - \binom{n-k}{k} - \binom{n-k-1}{k-1} + 2.$$

Moreover, for  $n > 2k$ , the equality holds if and only if  $\mathcal{F}' = \{[k] \cup \{i\} : i \in \{k, k+1\}\}$  and  $\mathcal{G}' = \{G \in \binom{[n]}{k} : G \cap ([k] \cup \{i\}) \neq \emptyset, i \in \{k, k+1\}\}$ , or two more possibilities when  $k = 3$ , namely,  $\mathcal{F}' = \{[2] \cup \{i\} : i \in \{3, 4, \dots, n\}\}$  and  $\mathcal{G}' = \{G \in \binom{[n]}{3} : G \cap [2] \neq \emptyset\}$ , or  $\mathcal{F}' = \mathcal{G}' = \{G \in \binom{[n]}{3} : 1 \in G\}$ . We need to return to the structure of  $\mathcal{F}$  and  $\mathcal{G}$ . In the first extremal case, we have  $|\mathcal{F}| = |\mathcal{F}'| = 2$ . Denote  $\mathcal{F} = \{F_1, F_2\}$ . Then  $|F_1 \cap F_2| = k-1$  otherwise the upper bound would be smaller. Thus  $\mathcal{G} = \{G \in \binom{[n]}{k} : G \cap F_i \neq \emptyset, i \in [2]\}$ . In the second extremal case, we have  $|\mathcal{F}| = |\mathcal{F}'| = n-2 = \binom{n-2}{3-2}$  and  $|\mathcal{G}| = |\mathcal{G}'| = \binom{n}{3} - \binom{n-2}{3}$ . For a fixed family  $\mathcal{F} \subseteq \binom{[n]}{3}$ , the maximality of  $|\mathcal{F}| + |\mathcal{G}|$  yields  $\mathcal{G} = \binom{[n]}{3} \setminus \mathcal{D}_3(\mathcal{F})$ . So we get  $|\mathcal{D}_3(\mathcal{F})| = \binom{n-2}{3}$ , which implies that the equality case of Lemma 3.1 occurs. Thus, it follows from Lemma 3.1 that  $\mathcal{F} = \{[2] \cup \{i\} : i \in \{3, 4, \dots, n\}\}$  and  $\mathcal{G} = \{G \in \binom{[n]}{3} : G \cap [2] \neq \emptyset\}$  under isomorphism. In the third extremal case, we have  $|\mathcal{F}| = |\mathcal{G}| = \binom{n-1}{2}$ . Similarly, we get  $|\mathcal{D}_3(\mathcal{F})| = \binom{n-1}{3}$ . By Lemma 3.1, we obtain  $\mathcal{F} = \{F \in \binom{[n]}{3} : 1 \in F\}$  and  $\mathcal{G} = \{G \in \binom{[n]}{3} : 1 \in G\}$  under isomorphism.  $\square$

**Proof of Theorem 1.6 for  $k \geq 4$ .** Suppose that  $k \geq 4$ ,  $\mathcal{F} \subseteq \binom{[n]}{k}$  is neither EKR nor HM, and  $|\mathcal{F}|$  is maximal. We need to consider the following three cases.

**Case 1.** After applying all possible shifting operations to  $\mathcal{F}$  repeatedly, the resulting family is still neither EKR nor HM.

Note that the resulting family is shifted and has the same size as  $\mathcal{F}$ . In this case, we may assume that  $\mathcal{F}$  is shifted. Since  $\mathcal{F}$  is neither EKR nor HM, we have  $\mathcal{F}(\bar{1}) \neq \emptyset$  and  $|\mathcal{F}(\bar{1})| \geq 2$ . By Lemma 2.1,  $\mathcal{F}(\bar{1})$  is 2-intersecting. Note that  $\mathcal{F}(\bar{1}) \subseteq \binom{[n] \setminus \{1\}}{k}$  and  $\mathcal{F}(1) \subseteq \binom{[n] \setminus \{1\}}{k-1}$  are cross-intersecting. Since  $k-1 \geq 3$ , applying Theorem 1.3 (for  $t = 1$ ) yields

$$|\mathcal{F}| = |\mathcal{F}(\bar{1})| + |\mathcal{F}(1)| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 2.$$

According to the extremal families in Theorem 1.3, we have  $\mathcal{F} \cong \mathcal{J}_2(n, k)$  or two more possibilities when  $k = 4$ ,  $\mathcal{F} \cong \mathcal{G}_3(n, 4)$  or  $\mathcal{F} \cong \mathcal{G}_2(n, 4)$ .

**Case 2.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{k}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G})$  is EKR.

It is clear that  $\cap_{H \in s_{i,j}(\mathcal{G})} H = \{i\}$  and  $\{i, j\} \cap G \neq \emptyset$  for all  $G \in \mathcal{G}$ . Then  $\mathcal{G}(\bar{i}, \bar{j}) = \emptyset$ . It follows that  $|\mathcal{G}(i, \bar{j})| \geq 2$  and  $|\mathcal{G}(\bar{i}, j)| \geq 2$ . In addition, we have  $\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j}) = \emptyset$ . By the maximality of  $|\mathcal{G}|$ , we have  $\mathcal{G}(i, j) = \binom{[n] \setminus \{i, j\}}{k-2}$ . Note that  $\mathcal{G}(i, \bar{j}) \subseteq \binom{[n] \setminus \{i, j\}}{k-1}$  and  $\mathcal{G}(\bar{i}, j) \subseteq \binom{[n] \setminus \{i, j\}}{k-1}$  are cross-intersecting, and  $k-1 \geq 3$ . Note that the inequality in Lemma 3.2 holds strictly. Then we get

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ &< \binom{n-2}{k-2} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 2 \\ &= \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 2. \end{aligned}$$

**Case 3.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{k}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G})$  is HM.

We first have that  $s_{i,j}(\mathcal{G})$  is HM at  $i$ . Then  $|\mathcal{G}(\bar{i}, \bar{j})| \leq 1$ . By Case 2, we may assume that  $s_{i,j}(\mathcal{G})$  is not EKR. If  $|\mathcal{G}(\bar{i}, \bar{j})| = 0$ , then  $|\mathcal{G}(i, \bar{j})| \geq 2$  and  $|\mathcal{G}(\bar{i}, j)| \geq 2$ . In addition, we have  $|\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j})| = 1$ . Observe that the inequality in Lemma 3.2 still holds strictly for  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$ . So the same argument with Case 2 works. Now assume that  $|\mathcal{G}(\bar{i}, \bar{j})| = 1$ . Since  $\mathcal{G}(i, j)$  and  $\mathcal{G}(\bar{i}, \bar{j})$  are cross-intersecting, we have  $|\mathcal{G}(i, j)| \leq \binom{n-2}{k-2} - \binom{n-k-2}{k-2}$ . In addition, by the properties of  $\mathcal{G}$ ,  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting,  $\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j}) = \emptyset$ ,  $|\mathcal{G}(i, \bar{j})| \geq 1$  and  $|\mathcal{G}(\bar{i}, j)| \geq 1$ . The inequality in Theorem 1.1 holds strictly. Then  $|\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \leq \binom{n-2}{k-1} - \binom{n-k-1}{k-1}$ . Consequently,

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| + 1 \\ &\leq \binom{n-2}{k-2} - \binom{n-k-2}{k-2} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} + 1 \\ &< \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 2. \end{aligned}$$

By Case 1-Case 3, we complete the proof of Theorem 1.6 for  $k \geq 4$ .  $\square$

In order to prove Theorem 1.6 for  $k = 3$ , we need to establish the corresponding versions of  $k = 2$  for Theorem 1.3 and Lemma 3.2. For simplicity, we postpone the proof in Appendix B.

### 3.3 Short proof of the Huang–Peng theorem

In Theorem 1.7, suppose that there exists  $x \in [n]$  such that there are only two sets  $E_1$  and  $E_2 \in \mathcal{F}$  missing  $x$ . If  $|E_1 \cap E_2| = k - i$  and  $i \geq 2$ , then

$$\begin{aligned} |\mathcal{F}| &\leq \binom{n-1}{k-1} - 2\binom{n-k-1}{k-1} + \binom{n-k-i-1}{k-1} + 2 \\ &\leq \binom{n-1}{k-1} - 2\binom{n-k-1}{k-1} + \binom{n-k-3}{k-1} + 2. \end{aligned}$$

The equality holds if and only if  $|E_1 \cap E_2| = k - 2$ , i.e.,  $\mathcal{F} = \mathcal{K}_2(n, k)$ . Moreover, it is not hard to see that

$$|\mathcal{J}_3(n, k)| = \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3.$$

Note that for  $2k + 1 \leq n \leq 3k - 3$ ,  $|\mathcal{K}_2(n, k)| \geq |\mathcal{J}_3(n, k)|$ , and the equality holds only for  $k = 4$ ; for  $n \geq 3k - 2$ ,  $|\mathcal{K}_2(n, k)| < |\mathcal{J}_3(n, k)|$ . Therefore, in the following, we may assume that for any  $x \in [n]$ , there are at least three sets in  $\mathcal{F}$  not containing  $x$ . Then it suffices to show that

$$|\mathcal{F}| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3.$$

For  $k = 5$ , the equality holds only for  $\mathcal{F} = \mathcal{J}_3(n, k)$  or  $\mathcal{G}_4(n, k)$ . For every other  $k$ , the equality holds only for  $\mathcal{F} = \mathcal{J}_3(n, k)$ . We divide the proof into case  $k \geq 5$  and case  $k = 4$ . For  $k \geq 5$ , we need the following tools.

Similar to Lemma 3.2, we can translate Theorem 1.4 into the following lemma.

**Lemma 3.3.** *Let  $k \geq 4$  and  $n \geq 2k$  be positive integers. Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be cross-intersecting families. Suppose that  $|\mathcal{G}| \geq |\mathcal{F}| \geq 3$ . Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - \binom{n-k}{k} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 3.$$

For  $n > 2k$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2, F_3\}$  with  $F_i := [k-1] \cup \{k-1+i\}$  for each  $i \in [3]$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{k} : G \cap F_i \neq \emptyset \text{ for any } i = 1, 2, 3 \right\}$ , or one more possibility when  $k = 4$ ,  $\mathcal{F} = \{[3] \cup \{i\} : i \in \{4, \dots, n\}\}$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{4} : G \cap [3] \neq \emptyset \right\}$ , under isomorphism.

**Proof of Theorem 1.7 for  $k \geq 5$ .** Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  be an maximum intersecting family which is neither EKR nor HM, and  $\mathcal{F} \not\subseteq \mathcal{J}_2(n, k)$ . For any  $x \in [n]$ , let  $|\mathcal{F}(\bar{x})| \geq 3$ . Furthermore, for any  $\mathcal{G}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, k)$ , we may assume that  $|\mathcal{G}(\bar{x})| \geq 3$ . We need to deal with the following four cases.

**Case 1.** Applying all shifting operations to  $\mathcal{F}$  repeatedly, the resulting shifted family is still not EKR, not HM and not a subfamily of  $\mathcal{J}_2(n, k)$ .

Without loss of generality, we may assume that  $\mathcal{F}$  is shifted. First note that  $|\mathcal{F}(\bar{1})| \geq 3$  and by Lemma 2.1,  $\mathcal{F}(\bar{1})$  is 2-intersecting. In addition,  $\mathcal{F}(\bar{1}) \subseteq \binom{[n] \setminus \{1\}}{k}$  and  $\mathcal{F}(1) \subseteq \binom{[n] \setminus \{1\}}{k-1}$  are cross-intersecting. Since  $k-1 \geq 4$ , Theorem 1.4 ( $t=1$ ) leads to

$$|\mathcal{F}| = |\mathcal{F}(\bar{1})| + |\mathcal{F}(1)| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3,$$

and for  $n \geq 2k+1$ , the equality holds if and only if  $\mathcal{F} \cong \mathcal{J}_3(n, k)$  or when  $k=5$ ,  $\mathcal{F} \cong \mathcal{G}_4(n, 5)$ .

**Case 2.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{k}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, k)$ ,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G})$  is EKR.

It is clear that  $\cap_{H \in s_{i,j}(\mathcal{G})} H = \{i\}$  and  $\{i, j\} \cap G \neq \emptyset$  for all  $G \in \mathcal{G}$ . Hence,  $\mathcal{G}(\bar{i}, \bar{j}) = \emptyset$ . This implies that  $|\mathcal{G}(i, \bar{j})| = |\mathcal{G}(\bar{j})| \geq 3$  and  $|\mathcal{G}(\bar{i}, j)| = |\mathcal{G}(\bar{i})| \geq 3$ . In addition, we have  $\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j}) = \emptyset$ . By the maximality of  $|\mathcal{G}|$ , we have  $\mathcal{G}(i, j) = \binom{[n] \setminus \{i, j\}}{k-2}$ . Note that  $\mathcal{G}(i, \bar{j}) \subseteq \binom{[n] \setminus \{i, j\}}{k-1}$  and  $\mathcal{G}(\bar{i}, j) \subseteq \binom{[n] \setminus \{i, j\}}{k-1}$  are cross-intersecting. Note that the inequality in Lemma 3.3 holds strictly for  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$ . Then

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ &\leq \binom{n-2}{k-2} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 2 \\ &< \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3. \end{aligned}$$

**Case 3.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{k}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, k)$ ,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G})$  is HM.

First of all,  $s_{i,j}(\mathcal{G})$  is HM at  $i$ . Therefore, we obtain  $|\mathcal{G}(\bar{i}, \bar{j})| \leq 1$ . By Case 2, we may assume that  $s_{i,j}(\mathcal{G})$  is not EKR. If  $|\mathcal{G}(\bar{i}, \bar{j})| = 0$ , then  $|\mathcal{G}(i, \bar{j})| \geq 3$  and  $|\mathcal{G}(\bar{i}, j)| \geq 3$ . In addition, we have  $|\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j})| = 1$ . By Lemma 3.3, the same argument with Case 2 works.

Now assume that  $|\mathcal{G}(\bar{i}, \bar{j})| = 1$ . Since  $\mathcal{G}(i, j)$  and  $\mathcal{G}(\bar{i}, \bar{j})$  are cross-intersecting, we have  $|\mathcal{G}(i, j)| \leq \binom{n-2}{k-2} - \binom{n-k-2}{k-2}$ . Furthermore,  $|\mathcal{G}(i, \bar{j})| \geq 3 - |\mathcal{G}(\bar{i}, \bar{j})| = 2$  and  $|\mathcal{G}(\bar{i}, j)| \geq 2$ . Observe that  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting. Since  $|\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j})| = 0$ , the inequality in Lemma 3.2 holds strictly for  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$ . We get  $|\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \leq \binom{n-2}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 1$ . Consequently,

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| + 1 \\ &\leq \binom{n-2}{k-2} - \binom{n-k-2}{k-2} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} + 2 \\ &< \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3. \end{aligned}$$

**Case 4.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{k}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, k)$ ,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G}) \subseteq \mathcal{J}_2(n, k)$ .

By conditions, we have  $\mathcal{G}(\bar{i}, \bar{j}) \subseteq \{E_1, E_2\}$  for some  $|E_1 \cap E_2| = k - 1$  and so  $|\mathcal{G}(\bar{i}, \bar{j})| \leq 2$ . By Cases 2 and 3, we may assume that  $s_{i,j}(\mathcal{G})$  is neither EKR nor HM. Therefore, if  $|\mathcal{G}(\bar{i}, \bar{j})| = 0$ , then  $|\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j})| = 2$ . In addition,  $|\mathcal{G}(i, \bar{j})| \geq 3$  and  $|\mathcal{G}(\bar{i}, j)| \geq 3$ . By Lemma 3.3, the same argument with Case 2 works. If  $|\mathcal{G}(\bar{i}, \bar{j})| = 1$ , then  $|\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j})| = 1$ . In addition, we have  $|\mathcal{G}(i, \bar{j})| \geq 2$  and  $|\mathcal{G}(\bar{i}, j)| \geq 2$ . By Lemma 3.2, the same argument with Case 3 works.

Now assume that  $|\mathcal{G}(\bar{i}, \bar{j})| = 2$ . Then  $\mathcal{G}(\bar{i}, \bar{j}) = \{E_1, E_2\}$  for some  $|E_1 \cap E_2| = k - 1$ . Considering the following families:

$$\begin{aligned}\mathcal{G}_1 &= \left\{ G \in \binom{[n] \setminus \{i, j\}}{k-2} : G \cap E_1 = \emptyset \right\}, \\ \mathcal{G}_2 &= \left\{ G \in \binom{[n] \setminus \{i, j\}}{k-2} : G \cap E_1 \neq \emptyset, G \cap E_2 = \emptyset \right\}.\end{aligned}$$

Note that  $\mathcal{G}_1 \cap \mathcal{G}_2 = \emptyset$ ,  $\mathcal{G}(i, j)$  and  $\mathcal{G}(\bar{i}, \bar{j})$  are cross-intersecting. Then we have

$$|\mathcal{G}(i, j)| \leq \binom{n-2}{k-2} - |\mathcal{G}_1| - |\mathcal{G}_2| = \binom{n-2}{k-2} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3}.$$

Moreover, we have  $|\mathcal{G}(\bar{i}, j) \cap \mathcal{G}(i, \bar{j})| = 0$ ,  $|\mathcal{G}(i, \bar{j})| \geq 3 - |\mathcal{G}(\bar{i}, \bar{j})| = 1$  and  $|\mathcal{G}(\bar{i}, j)| \geq 1$ . Observe that  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting. The inequality in Theorem 1.1 holds strictly. We get  $|\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \leq \binom{n-2}{k-1} - \binom{n-k-1}{k-1}$ . Therefore, we obtain

$$\begin{aligned}|\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| + 2 \\ &\leq \binom{n-2}{k-2} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} + 2 \\ &< \binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-k-2}{k-2} - \binom{n-k-3}{k-3} + 3,\end{aligned}$$

as desired. This completes the proof.  $\square$

It remains to prove Theorem 1.7 for  $k = 4$ . We shall establish the corresponding versions of Theorem 1.4 and Lemma 3.3 for  $k = 3$ . We defer the detailed discussions to Appendix C.

## Acknowledgement

Yongtao Li was supported by the Postdoctoral Fellowship Program of CPSF (No. GZC20233196). Lihua Feng was supported by the NSFC (Nos. 12271527, 12471022 and 12071484). Guihai Yu was supported by the NSFC (Nos. 12461062 and 11861019) and Natural Science Foundation of Guizhou (Nos. [2020]1Z001 and [2021]5609). Yongtao Li would like to thank Prof. Biao Wu and Dr. Zixiang Xu for the inspiring discussions on extremal set problems over the years.

## References

- [1] N. Alon, J.H. Spencer, The Probabilistic Method, Fourth edition, John Wiley & Sons, Inc., Hoboken, NJ, 2016.
- [2] J. Balogh, W. Linz, Short proofs of three results about intersecting systems, *Combin. Theory* 4 (2024), No. 4.
- [3] D. Bulavka, R. Woodroffe, A short proof of the Hilton–Milner Theorem, (2024), arXiv:2411.02513.

- [4] M. Cao, B. Lv, K. Wang, The structure of large non-trivial  $t$ -intersecting families of finite sets, *European J. Combin.* 97 (2021), No. 103373.
- [5] M. Cao, M. Lu, B. Lv, K. Wang, Nearly extremal non-trivial cross  $t$ -intersecting families and  $r$ -wise  $t$ -intersecting families, *European J. Combin.* 120 (2024), No. 103958.
- [6] D.E. Daykin, Erdős–Ko–Rado from Kruskal–Katona, *J. Combin. Theory Ser. A* 17 (1974) 254–255.
- [7] D. Ellis, Intersection problems in extremal combinatorics: Theorems, techniques and questions old and new, (2021), [arXiv:2107.06371v8](https://arxiv.org/abs/2107.06371v8).
- [8] P. Erdős, C. Ko, R. Rado, Intersection theorems for systems of finite sets, *Q. J. Math. Oxford* 2 (1961) 313–320.
- [9] P. Frankl, New inequalities for cross-intersecting families, *Mosc. J. Comb. Number Theory* 6 (2016) 27–32.
- [10] P. Frankl, A simple proof of the Hilton–Milner theorem, *Mosc. J. Comb. Number Theory* 8 (2019) 97–101.
- [11] P. Frankl, On the maximum of the sum of the sizes of non-trivial cross-intersecting families, *Combinatorica* 44 (2024) 15–35.
- [12] P. Frankl, Z. Füredi, Non-trivial intersecting families, *J. Combin. Theory Ser. A* 41 (1986) 150–153.
- [13] P. Frankl, Z. Füredi, A new short proof of the EKR theorem, *J. Combin. Theory Ser. A* 119 (2012) 1388–1390.
- [14] P. Frankl, A. Kupavskii, A size-sensitive inequality for cross-intersecting families, *European J. Combin.* 62 (2017) 263–271.
- [15] P. Frankl, N. Tokushige, Some best possible inequalities concerning cross-intersecting families, *J. Combin. Theory Ser. A* 61 (1992) 87–97.
- [16] P. Frankl, N. Tokushige, Some inequalities concerning cross-intersecting families, *Combin. Probab. Comput.* 7 (1998) 247–260.
- [17] P. Frankl, N. Tokushige, Invitation to intersection problems for finite sets, *J. Combin. Theory, Ser. A* 144 (2016) 157–211.
- [18] P. Frankl, J. Wang, Improved bounds on the maximum diversity of intersecting families, *European J. Combin.* 118 (2024), No. 103885.
- [19] Z. Füredi, J.R. Griggs, Families of finite sets with minimum shadows, *Combinatorica* 6 (1986) 355–363.
- [20] Z. Füredi, K.-W. Hwang, P.M. Weichsel, A proof and generalizations of the Erdős–Ko–Rado theorem using the method of linearly independent polynomials, in: *Topics in Discrete Mathematics*, in: *Algorithms Combin.*, vol. 26, Springer, Berlin, 2006, pp. 215–224.
- [21] J. Gao, H. Liu, Z. Xu, Stability through non-shadows, *Combinatorica* 43 (2023) 1125–1137.
- [22] G. Ge, Z. Xu, X. Zhao, Algebraic approach to stability results for Erdős–Ko–Rado theorem, 16 pages, (2024), [arXiv:2410.22676](https://arxiv.org/abs/2410.22676).

- [23] J. Han, Y. Kohayakawa, The maximum size of a non-trivial intersecting uniform family that is not a subfamily of the Hilton–Milner family, *Proc. Amer. Math. Soc.* 145 (2017) 73–87.
- [24] A.J.W. Hilton, The Erdős–Ko–Rado theorem with valency conditions, in: Unpublished Manuscript, 1976.
- [25] A.J.W. Hilton, E.C. Milner, Some intersection theorems for systems of finite sets, *Q. J. Math.* 18 (1967) 369–384.
- [26] Y. Huang, Y. Peng, Stability of intersecting families, *European J. Combin.* 115 (2024), No. 103774.
- [27] H. Huang, Y. Zhao, Degree versions of the Erdős–Ko–Rado theorem and Erdős hypergraph matching conjecture, *J. Combin. Theory Ser. A* 150 (2017) 233–247.
- [28] G. Hurlbert, V. Kamat, New injective proofs of the Erdős–Ko–Rado and Hilton–Milner theorems, *Discrete Math.* 341 (2018) 1749–1754.
- [29] G.O.H. Katona, Intersection theorems for systems of finite sets, *Acta Math. Hungar.* 15 (1964) 329–337.
- [30] G.O.H. Katona, A simple proof of the Erdős–Chao Ko–Rado theorem, *J. Combin. Theory Ser. B* 13 (1972) 183–184.
- [31] A. Kostochka, D. Mubayi, The structure of large intersecting families, *Proc. Amer. Math. Soc.* 145 (2017) 2311–2321.
- [32] A. Kupavskii, Structure and properties of large intersecting families, (2018), arXiv:1810.00920.
- [33] A. Kupavskii, D. Zakharov, Regular bipartite graphs and intersecting families, *J. Combin. Theory Ser. A* 155 (2018), 180–189.
- [34] Y. Li, B. Wu, Stabilities for non-uniform  $t$ -intersecting families, *Electron. J. Combin.* 31 (4) (2024), #P4.3.
- [35] L. Lovász, On the Shannon capacity of a graph, *IEEE Trans. Inform. Theory* 25 (1979) 1–7.
- [36] M. Mörs, A generalization of a theorem of Kruskal, *Graphs and Combinatorics* 1 (1985) 167–183.
- [37] J. O’Neill, J. Verstraëte, Non-trivial  $d$ -wise intersecting families, *J. Combin. Theory Ser. A* 178 (2021), No. 105369.
- [38] C. Shi, P. Frankl, J. Qian, On non-empty cross-intersecting families, *Combinatorica* 42 (2022) 1513–1525.
- [39] J. Wang, H. Zhang, Nontrivial independent sets of bipartite graphs and cross-intersecting families, *J. Combin. Theory Ser. A* 120 (2013) 129–141.
- [40] B. Wu, A refined result on cross-intersecting families, *Discrete Appl. Math.* 339 (2023) 149–153.

## A The case $k = 3$ in the Han–Kohayakawa theorem

Define  $\mathcal{A} = \{\{1, 2\} \cup \{i\} : i \in [3, n]\} \cup \left\{G \in \binom{[n]}{2} : G \cap \{1, 2\} \neq \emptyset\right\}$ .

**Lemma A.1.** *Let  $n \geq 6$  be an integer. Let  $\mathcal{F} \subseteq \binom{[n]}{3}$  and  $\mathcal{G} \subseteq \binom{[n]}{2}$  be cross-intersecting families. Suppose that  $|\mathcal{F}| \geq 2$  and  $\mathcal{F}$  is 2-intersecting. If  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\mathcal{A}$ , then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{2} - \binom{n-3}{2} - \binom{n-4}{1} + 2.$$

*The equality holds if and only if  $\mathcal{F} = \{F_1, F_2\}$  for some  $F_1, F_2 \in \binom{[n]}{3}$  with  $|F_1 \cap F_2| = 2$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{2} : G \cap F_1 \neq \emptyset \text{ and } G \cap F_2 \neq \emptyset\right\}$ .*

**Proof.** If  $|\mathcal{F}| = 2$ , then the result holds clearly. Now suppose that  $|\mathcal{F}| \geq 3$ . Let  $x_1, \dots, x_5 \in [n]$  be five different numbers. If there exists  $\{\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, x_2, x_5\}\} \subseteq \mathcal{F}$ , then  $\mathcal{G} \subseteq \{\{x_1, i\}, i \in [n] \setminus \{x_1\}\} \cup \{\{x_2, i\}, i \in [n] \setminus \{x_1, x_2\}\}$ . Since  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\mathcal{A}$ , we have  $\{a, b, c\} \in \mathcal{F}$  for three different elements  $a, b, c \in [n]$  satisfying  $|\{x_1, x_2\} \cap \{a, b, c\}| \leq 1$ . Since  $|\{a, b, c\} \cap \{x_1, x_2, x_3\}| \geq 2$  and  $|\{a, b, c\} \cap \{x_1, x_2, x_4\}| \geq 2$ , we get  $\{a, b, c\} = \{x_1, x_3, x_4\}$  or  $\{a, b, c\} = \{x_2, x_3, x_4\}$ . But  $|\{x_1, x_3, x_4\} \cap \{x_1, x_2, x_5\}| = 1$  and  $|\{x_2, x_3, x_4\} \cap \{x_1, x_2, x_5\}| = 1$ , a contradiction. Therefore, we may assume that  $\{\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, x_3, x_4\}\} \subseteq \mathcal{F}$ . Then  $\mathcal{G} \subseteq \{\{x_1, i\}, i \in [n] \setminus \{x_1\}\} \cup \{\{x_2, x_3\}, \{x_2, x_4\}, \{x_3, x_4\}\}$ . So  $|\mathcal{G}| \leq n + 2$ . Since  $\mathcal{F}$  is 2-intersecting, we must have  $\mathcal{F} \subseteq \{\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, x_3, x_4\}, \{x_2, x_3, x_4\}\}$ . Thus  $|\mathcal{F}| \leq 4 \leq n - 2$ . It follows that  $|\mathcal{F}| + |\mathcal{G}| \leq 2n = \binom{n}{2} - \binom{n-3}{2} - \binom{n-4}{1} + 2$ . It is not hard to see that the inequality holds strictly. This completes the proof.  $\square$

Define  $\mathcal{B} = \{\{1\} \cup \{i\} : i \in [2, n]\}$ . The next lemma deals with the case  $k = 2$  for Lemma 3.2.

**Lemma A.2.** *Let  $n \geq 4$  be an integer. Let  $\mathcal{F} \subseteq \binom{[n]}{2}$  and  $\mathcal{G} \subseteq \binom{[n]}{2}$  be cross-intersecting families. Suppose that  $|\mathcal{G}| \geq |\mathcal{F}| \geq 2$  and  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\mathcal{B}$ . Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{2} - \binom{n-2}{2} - \binom{n-3}{1} + 2.$$

*For  $n \geq 5$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2\}$  for some  $F_1, F_2 \in \binom{[n]}{2}$  with  $|F_1 \cap F_2| = 1$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{2} : G \cap F_1 \neq \emptyset \text{ and } G \cap F_2 \neq \emptyset\right\}$ .*

**Proof.** First note that  $\binom{n}{2} - \binom{n-2}{2} - \binom{n-3}{1} + 2 = n + 2$ . If  $|\mathcal{F}| = 2$ , then the result holds clearly. So we may assume that  $|\mathcal{F}| \geq 3$ . Let  $x_1, \dots, x_6 \in [n]$  be six different numbers (if exist). If there exists  $\{\{x_1, x_2\}, \{x_3, x_4\}, \{x_5, x_6\}\} \subseteq \mathcal{F}$ , then  $\mathcal{G} = \emptyset$ , a contradiction. So we may assume that  $\{\{x_1, x_2\}, \{x_1, x_3\}\} \subseteq \mathcal{F}$ , note that  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\mathcal{B}$ , so there are some  $u \neq v \in [n] \setminus \{x_1\}$  such that  $\{u, v\} \in \mathcal{F}$  or  $\{u, v\} \in \mathcal{G}$ .

If  $\{u, v\} \in \mathcal{F}$ , then  $\mathcal{G} \subseteq \{\{x_1, u\}, \{x_1, v\}, \{x_2, x_3\}\}$ . So  $|\mathcal{G}| \leq 3$ . Then  $|\mathcal{F}| + |\mathcal{G}| \leq 6 \leq n + 2$ .

If  $\{u, v\} \in \mathcal{G}$ , then  $\{u, v\} = \{x_2, x_3\}$ . Observe that  $\mathcal{G} \subseteq \{\{x_1, i\}, i \in [n] \setminus \{x_1\}\} \cup \{\{x_2, x_3\}\}$ . Since  $|\mathcal{G}| \geq |\mathcal{F}| \geq 3$ , we have  $\{\{x_1, a\}, \{x_1, b\}, \{x_2, x_3\}\} \subseteq \mathcal{G}$  for some  $a \neq b \in [n] \setminus \{x_1\}$ . Then the cross-intersecting property of  $\mathcal{F}$  and  $\mathcal{G}$  implies that  $\mathcal{F} \subseteq \{\{x_1, x_2\}, \{x_1, x_3\}, \{a, b\}\}$ . Since  $|\mathcal{F}| \geq 3$ , we have  $\mathcal{F} = \{\{x_1, x_2\}, \{x_1, x_3\}, \{a, b\}\}$  and  $a \in \{x_2, x_3\}$  or  $b \in \{x_2, x_3\}$ . It follows that  $\mathcal{G} = \{\{x_1, a\}, \{x_1, b\}, \{x_2, x_3\}\}$ . Consequently,  $|\mathcal{F}| + |\mathcal{G}| \leq 6 \leq n + 2$ . This completes the proof.  $\square$

**Proposition A.3.** *Let  $n \geq 5$  be an integer. Let  $\mathcal{F} \subseteq \binom{[n]}{2}$  and  $\mathcal{G} \subseteq \binom{[n]}{2}$  be cross-intersecting families and  $|\mathcal{F} \cap \mathcal{G}| \leq 1$ . Suppose that  $|\mathcal{F}| \geq 2$  and  $|\mathcal{G}| \geq 2$ . Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{2} - \binom{n-2}{2} - \binom{n-3}{1} + 1.$$



**Proof.** If  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\{\{1\} \cup \{i\} : i \in [2, n]\}$ , then the result holds by Lemma A.2. If  $\mathcal{F} \cup \mathcal{G}$  is isomorphic to a subfamily of  $\{\{1\} \cup \{i\} : i \in [2, n]\}$ , note that  $|\mathcal{F}| + |\mathcal{G}| = |\mathcal{F} \cup \mathcal{G}| + |\mathcal{F} \cap \mathcal{G}|$ , then  $|\mathcal{F} \cap \mathcal{G}| \leq 1$  implies that  $|\mathcal{F}| + |\mathcal{G}| \leq n < \binom{n}{2} - \binom{n-2}{2} - \binom{n-3}{1} + 1$ , as required.  $\square$

**Proof of Theorem 1.6 for  $k = 3$ .** Suppose that  $\mathcal{F} \subseteq \binom{[n]}{3}$  is neither EKR nor HM, and not a subfamily of  $\mathcal{G}_2(n, 3)$ . In addition,  $|\mathcal{F}|$  is maximal. Since  $\mathcal{F}$  is not a subfamily of  $\mathcal{G}_2(n, 3)$ , we have that  $\mathcal{F}(\bar{1}) \cup \mathcal{F}(1)$  is not isomorphic to a subfamily of  $\mathcal{A}$ . According to Lemma A.1, Proposition A.3 and the proof of Theorem 1.6 for  $k \geq 4$ , we only need to deal with the following case.

**Case 1.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{3}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{G}_2(n, 3)$ ,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G}) \subseteq \mathcal{G}_2(n, 3)$ .

By conditions, there exist three different elements  $x_0, x_1, x_2$  such that

$$s_{i,j}(\mathcal{G}) \subseteq \left\{ G \in \binom{[n]}{3} : \{x_1, x_2\} \subseteq G \right\} \cup \left\{ G \in \binom{[n]}{3} : x_0 \in G, G \cap \{x_1, x_2\} \neq \emptyset \right\}.$$

Hence,  $i \in \{x_0, x_1, x_2\}$  and  $j \notin \{x_0, x_1, x_2\}$ . By symmetry, let us assume that  $x_0 = i$ . This implies that  $\mathcal{G}(i, j) \subseteq \{\{x_1\}, \{x_2\}\}$  and so  $|\mathcal{G}(i, j)| \leq 2$ . Moreover,  $\mathcal{G}(\bar{i}, \bar{j}) \subseteq \{\{x_1, x_2, y\} : y \in [n] \setminus \{x_1, x_2, i, j\}\}$  and so  $|\mathcal{G}(\bar{i}, \bar{j})| \leq n - 4$ . Observe that  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) = \emptyset$  or  $\{x_1, x_2\}$ . By the maximality of  $|\mathcal{G}|$ , we may assume that  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) = \{x_1, x_2\}$ . Since  $\mathcal{G}$  is not a subfamily of  $\mathcal{G}_2(n, 3)$ , we have  $\mathcal{G}(i, \bar{j}) \neq \{x_1, x_2\}$  and  $\mathcal{G}(\bar{i}, j) \neq \{x_1, x_2\}$ . Then  $|\mathcal{G}(i, \bar{j})| \geq 2$  and  $|\mathcal{G}(\bar{i}, j)| \geq 2$ . Note that  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting. By Proposition A.3, we obtain

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(\bar{i}, \bar{j})| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ &\leq 2 + n - 4 + \binom{n-2}{2} - \binom{n-4}{2} - \binom{n-5}{1} + 1 \\ &< \binom{n-1}{2} - \binom{n-4}{2} - \binom{n-5}{1} + 2, \end{aligned}$$

as required.  $\square$

## B The case $k = 4$ in the Huang–Peng theorem

Denote  $\mathcal{C} = \{[3] \cup \{i\} : i \in [4, n]\} \cup \left\{ G \in \binom{[n]}{3} : G \cap [3] \neq \emptyset \right\}$ .

**Lemma B.1.** *Let  $n \geq 7$  be an integer. Let  $\mathcal{F} \subseteq \binom{[n]}{4}$  and  $\mathcal{G} \subseteq \binom{[n]}{3}$  be cross-intersecting families. Suppose that  $|\mathcal{F}| \geq 3$  and  $\mathcal{F}$  is 2-intersecting. In addition,  $\mathcal{F}$  and  $\mathcal{G}$  are both shifted. If  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\mathcal{C}$ , then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - \binom{n-4}{3} - \binom{n-5}{2} - \binom{n-6}{1} + 3.$$

For  $n \geq 8$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2, F_3\}$  with  $F_i := [3] \cup \{3+i\}$  for each  $i \in [3]$  and  $\mathcal{G} = \left\{ G \in \binom{[n]}{3} : G \cap F_i \neq \emptyset \text{ for any } i = 1, 2, 3 \right\}$ , under isomorphism.

**Proof.** We proceed by applying induction on  $n \geq 7$ . The base case  $n = 7$  is trivial. Suppose that  $n \geq 8$  and the result holds for integers less than  $n$ . Since  $\mathcal{F}$  and  $\mathcal{G}$  are shifted, by Lemma 2.2, we have  $|\mathcal{F}(\bar{n})| \geq 3$ . Clearly,  $\mathcal{F}(\bar{n})$  is 2-intersecting. Note that  $\mathcal{F}(\bar{n})$  and  $\mathcal{G}(\bar{n})$  are cross-intersecting. By induction hypothesis, we get

$$|\mathcal{F}(\bar{n})| + |\mathcal{G}(\bar{n})| \leq \binom{n-1}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 3.$$

If  $|\mathcal{F}(n)| = 0$ , then the same argument as Case 1 of Theorem 1.4 works.

If  $|\mathcal{F}(n)| = 1$ , let  $\mathcal{F}' = \{[3] \cup \{i\} : i \in [4, n-1]\}$ . Then  $\mathcal{F}' \subseteq \mathcal{F}(\bar{n})$ . If  $\mathcal{F}' = \mathcal{F}(\bar{n})$ , then  $\mathcal{F} = \{[3] \cup \{i\} : i \in [4, n]\}$ . So  $\mathcal{G} \subseteq \left\{G \in \binom{[n]}{3} : G \cap [3] \neq \emptyset\right\}$ . But  $\mathcal{F} \cup \mathcal{G}$  is isomorphic to a subfamily of  $\mathcal{C}$ , a contradiction. Hence,  $\mathcal{F}' \subsetneq \mathcal{F}(\bar{n})$ . Then we have  $\mathcal{G}(n) \subsetneq \left\{G^* \in \binom{[n-1]}{2} : G^* \cap [3] \neq \emptyset\right\}$ . This implies that  $|\mathcal{G}(n)| < \binom{n-1}{2} - \binom{n-4}{2}$ . Consequently,

$$|\mathcal{F}| + |\mathcal{G}| = |\mathcal{F}(\bar{n})| + |\mathcal{G}(\bar{n})| + |\mathcal{G}(n)| + 1 < \binom{n}{3} - \binom{n-4}{3} - \binom{n-5}{2} - \binom{n-6}{1} + 4.$$

That is,  $|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - \binom{n-4}{3} - \binom{n-5}{2} - \binom{n-6}{1} + 3$ . Note that  $\mathcal{F}(\bar{n})$  and  $\mathcal{G}(n)$  are shifted on  $[n-1]$ . If the equality holds, then  $\mathcal{G}(n) = \left\{G \in \binom{[n-1]}{2} : G \cap [3] \neq \emptyset\right\} \setminus \{3, n-1\}$ . Since  $\mathcal{F}' \subsetneq \mathcal{F}(\bar{n})$  and  $\mathcal{F}(\bar{n})$  is shifted, we have  $\{1, 2, 4, 5\} \in \mathcal{F}(\bar{n})$ . Observe that  $\{3, n-2\} \in \mathcal{G}(n)$  and  $n-2 \geq 6$ . So  $\{1, 2, 4, 5\} \cap \{3, n-2\} = \emptyset$ . But  $\mathcal{F}(\bar{n})$  and  $\mathcal{G}(n)$  are cross-intersecting, a contradiction. Therefore, we have  $|\mathcal{F}| + |\mathcal{G}| < \binom{n}{3} - \binom{n-4}{3} - \binom{n-5}{2} - \binom{n-6}{1} + 3$ .

If  $|\mathcal{F}(n)| \geq 2$ , since  $\mathcal{F} \cup \mathcal{G}$  is not isomorphic to a subfamily of  $\mathcal{C}$ , we have that  $\mathcal{F}(n) \cup \mathcal{G}(n)$  is not isomorphic to a subfamily of  $\mathcal{A}$ . In addition, by Lemmas 2.1 and 2.4,  $\mathcal{F}(n)$  is 2-intersecting,  $\mathcal{F}(n)$  and  $\mathcal{G}(n)$  are cross-intersecting. Then applying Lemma A.1 to  $\mathcal{F}(n)$  and  $\mathcal{G}(n)$  yields  $|\mathcal{F}(n)| + |\mathcal{G}(n)| \leq \binom{n-1}{2} - \binom{n-4}{2} - \binom{n-5}{1} + 2$ . Combining this and the previous inequality about  $\mathcal{F}(\bar{n})$  and  $\mathcal{G}(\bar{n})$ , we obtain

$$\begin{aligned} |\mathcal{F}| + |\mathcal{G}| &= |\mathcal{F}(\bar{n})| + |\mathcal{G}(\bar{n})| + |\mathcal{F}(n)| + |\mathcal{G}(n)| \\ &\leq \binom{n}{3} - \binom{n-4}{3} - \binom{n-5}{2} - \binom{n-6}{1} + 3 - \binom{n-7}{1} \\ &< \binom{n}{3} - \binom{n-4}{3} - \binom{n-5}{2} - \binom{n-6}{1} + 3, \end{aligned}$$

as required.  $\square$

Define  $\mathcal{P} = \{\{1, 2\} \cup \{i\} : i \in [3, n]\} \cup \left\{G \in \binom{[n]}{3} : G \cap \{1, 2\} \neq \emptyset\right\}$  and  $\mathcal{Q} = \left\{Q \in \binom{[n]}{3} : 1 \in Q\right\}$ .

**Proposition B.2.** *Let  $n \geq 7$  be an integer. Let  $\mathcal{F} \subseteq \binom{[n]}{3}$  and  $\mathcal{G} \subseteq \binom{[n]}{3}$  be non-empty cross-intersecting families and  $|\mathcal{F} \cap \mathcal{G}| \leq 2$ . Suppose that  $|\mathcal{F}| \geq 3$  and  $|\mathcal{G}| \geq 3$ . Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - \binom{n-3}{3} - \binom{n-4}{2} - \binom{n-5}{1} + 2.$$

**Proof.** If  $\mathcal{F} \cup \mathcal{G}$  is isomorphic to a subfamily of  $\mathcal{P}$  or  $\mathcal{Q}$ , then  $|\mathcal{F} \cup \mathcal{G}| \leq \binom{n}{3} - \binom{n-3}{3}$  or  $|\mathcal{F} \cup \mathcal{G}| \leq \binom{n-1}{2}$ . Since  $|\mathcal{F} \cap \mathcal{G}| \leq 2$ , we obtain

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - \binom{n-2}{3} + 2 < \binom{n}{3} - \binom{n-3}{3} - \binom{n-4}{2} - \binom{n-5}{1} + 2.$$

Now suppose that  $\mathcal{F} \cup \mathcal{G}$  is neither isomorphic to a subfamily of  $\mathcal{P}$  nor a subfamily of  $\mathcal{Q}$ . If  $|\mathcal{F}| = 3$  or  $|\mathcal{G}| = 3$ , the result holds clearly. So let us assume that  $|\mathcal{F}| \geq 4$  and  $|\mathcal{G}| \geq 4$ .

Let  $x_1, \dots, x_5 \in [n]$  be five different numbers. When either  $\mathcal{F}$  or  $\mathcal{G}$  is a star. By symmetry, we may assume that  $\mathcal{F}$  is a star at  $x_1$ . If any four sets of  $\mathcal{F}$  contain  $x_1, x_2$ , then  $G \cap \{x_1, x_2\} \neq \emptyset$  for any  $G \in \mathcal{G}$ . So  $\mathcal{F} \cup \mathcal{G}$  is isomorphic to a subfamily of  $\mathcal{P}$ , a contradiction. If  $\{\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, x_2, x_5\}, \{x_1, a, b\}\} \subseteq \mathcal{F}$  for some  $a \neq b$  and  $a, b \notin \{x_1, x_2\}$ , then  $\mathcal{G} \subseteq \left\{G \in \binom{[n]}{3} : x_1 \in G\right\} \cup \{\{x_2, a, i\} : i \in [n] \setminus \{x_1, x_2, a\}\} \cup \{\{x_2, b, i\} : i \in [n] \setminus \{x_1, x_2, a, b\}\} \cup \{\{x_3, x_4, x_5\}\}$ . It follows that  $|\mathcal{F}| + |\mathcal{G}| = |\mathcal{F} \cup \mathcal{G}| + |\mathcal{F} \cap \mathcal{G}| \leq \binom{n-1}{2} + 2n - 4$ , which is smaller than

the required upper bound. If  $\{\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, a, b\}, \{x_1, a, d\}\} \subseteq \mathcal{F}$  for three different elements  $a, b, d$  and  $a \notin \{x_1, x_2, x_3, x_4\}$  and  $b, d \notin \{x_1, x_2\}$ , then  $\mathcal{G} \subseteq \left\{G \in \binom{[n]}{3} : x_1 \in G\right\} \cup \{\{x_2, a, i\} : i \in [n] \setminus \{x_1, x_2, a\}\} \cup \{\{x_2, b, d\}, \{x_3, x_4, a\}\}$ . So  $|\mathcal{F}| + |\mathcal{G}| = |\mathcal{F} \cup \mathcal{G}| + |\mathcal{F} \cap \mathcal{G}| \leq \binom{n-1}{2} + n + 1$ . This is also smaller than the required upper bound. If  $\{\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, a, b\}, \{x_1, c, d\}\} \subseteq \mathcal{F}$  for four different elements  $a, b, c, d$  and  $a, b, c, d \notin \{x_1, x_2\}$ , then we similarly have  $|\mathcal{F}| + |\mathcal{G}| \leq \binom{n-1}{2} + 6$ . This is also smaller than the required upper bound. If  $\{\{x_1, x_2, x_3\}, \{x_1, x_4, x_5\}, \{x_1, x_6, x_7\}, \{x_1, x_8, x_9\}\} \subseteq \mathcal{F}$  for nine different elements  $x_1, \dots, x_9 \in [n]$ , then  $G \cap \{x_1\} \neq \emptyset$  for any  $G \in \mathcal{G}$ . So  $\mathcal{F} \cup \mathcal{G}$  is isomorphic to a subfamily of  $\mathcal{Q}$ , a contradiction.

It remains to consider the case that  $\mathcal{F}$  and  $\mathcal{G}$  are both non-trivial. Although we can get the conclusion by a similar discussion as above. For simplicity, we use an old result due to Frankl and Tokushige.

**Lemma B.3** (See [16, 11]). *Let  $n \geq 2k + 1 \geq 5$  be an integer. Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  and  $\mathcal{G} \subseteq \binom{[n]}{k}$  be non-trivial cross-intersecting families. Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{k} - 2 \binom{n-k}{k} + \binom{n-2k}{k} + 2.$$

For  $k \geq 3$ , the equality holds if and only if  $\mathcal{F} = \{F_1, F_2\}$  for some disjoint  $F_1, F_2 \in \binom{[n]}{k}$  and  $\mathcal{G} = \left\{G \in \binom{[n]}{k} : G \cap F_i \neq \emptyset \text{ for any } i = 1, 2\right\}$ .

Note that for  $2k + 1 \leq n \leq 3k - 1$ , the upper bound of Lemma B.3 is reduced to  $\binom{n}{k} - 2 \binom{n-k}{k} + 2$ . Returning to our proof. Note that  $|\mathcal{F}| \geq 4$ . By Lemma B.3, we have  $|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - 2 \binom{n-3}{3} + 1$  for  $7 \leq n \leq 8$ , and  $|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - 2 \binom{n-3}{3} + \binom{n-6}{3} + 1$  for  $n \geq 9$ . By simple calculations, we know that these bounds are smaller or equal to the required upper bound. This completes the proof of Proposition B.2.  $\square$

**Lemma B.4.** *Let  $n \geq 7$  be an integer. Let  $x_1 \neq x_2 \in [n]$  and  $\mathcal{F}, \mathcal{G} \subseteq \left\{A \in \binom{[n]}{3} : A \cap \{x_1, x_2\} \neq \emptyset\right\}$  be cross-intersecting families. Let  $\mathcal{R} = \left\{A \in \binom{[n]}{3} : \{x_1, x_2\} \subseteq A\right\}$ . Suppose that  $\mathcal{R} \subsetneq \mathcal{F}$ ,  $\mathcal{R} \subsetneq \mathcal{G}$  and  $(\mathcal{F} \setminus \mathcal{R}) \cap (\mathcal{G} \setminus \mathcal{R}) = \emptyset$ . Then*

$$|\mathcal{F}| + |\mathcal{G}| \leq \binom{n}{3} - \binom{n-3}{3} - \binom{n-4}{2} - \binom{n-5}{1} + 2.$$

**Proof.** By conditions and symmetry, we may assume that  $\{x_1, a, b\} \in \mathcal{F}$  for some  $a \neq b \in [n] \setminus \{x_1, x_2\}$ . Then

$$\mathcal{G} \subseteq \left\{G \in \binom{[n]}{3} : x_1 \in G\right\} \cup \{\{x_2, a, i\} : i \in [n] \setminus \{x_1, x_2, a\}\} \cup \{\{x_2, b, i\} : i \in [n] \setminus \{x_1, x_2, a, b\}\}.$$

If  $\mathcal{F}$  is a star at  $x_1$ , or  $\mathcal{F}$  is a star at  $x_1$  and there is at most a  $\{x_2, c, d\} \in \mathcal{F}$  for some  $c \neq d \in [n] \setminus \{x_1, x_2\}$ , note that  $(\mathcal{F} \setminus \mathcal{R}) \cap (\mathcal{G} \setminus \mathcal{R}) = \emptyset$ , then we have  $|\mathcal{F}| + |\mathcal{G}| \leq n - 2 + 1 + \binom{n-1}{2} + n - 3 + n - 4 = \binom{n-1}{2} + 3n - 8$ . Note that  $\binom{n}{3} - \binom{n-3}{3} - \binom{n-4}{2} - \binom{n-5}{1} + 2 = \binom{n-1}{2} + \binom{n-2}{2} + 3$ . For  $n \geq 7$ , let  $g(n) = \binom{n-1}{2} + \binom{n-2}{2} + 3 - (\binom{n-1}{2} + 3n - 8) = \binom{n-2}{2} - 3n + 11 = \frac{1}{2}(n^2 - 11n + 28)$ . Then  $h(n) \geq 0$ . So the result follows. By symmetry, the same case holds for  $\mathcal{G}$ .

Next we consider the following three cases.

(i) If there exist  $\{x_2, c, d\}, \{x_2, c, e\} \in \mathcal{F}$  for three different elements  $c, d, e \in [n] \setminus \{x_1, x_2\}$ , and  $\{x_1, a, b\}, \{x_1, a, y\} \in \mathcal{F}$  for three different elements  $a, b, y \in [n] \setminus \{x_1, x_2\}$ , then

$$\mathcal{G} \subseteq \mathcal{R} \cup \{\{x_1, c, i\} : i \in [n] \setminus \{x_1, x_2, c\}\} \cup \{\{x_1, e, d\}, \{x_2, b, y\}\} \cup \{\{x_2, a, i\} : i \in [n] \setminus \{x_1, x_2, a\}\}.$$

So  $|\mathcal{G}| \leq 3n - 6$ .

(ii) If there exist  $\{x_2, c, d\}, \{x_2, e, f\} \in \mathcal{F}$  for four different elements  $c, d, e, f \in [n] \setminus \{x_1, x_2\}$ , and  $\{x_1, a, b\}, \{x_1, a, y\} \in \mathcal{F}$  for three different elements  $a, b, y \in [n] \setminus \{x_1, x_2\}$ , then

$$\mathcal{G} \subseteq \mathcal{R} \cup \{\{x_1, c, i\} : i \in \{e, f\}\} \cup \{\{x_1, d, i\} : i \in \{e, f\}\} \cup \{\{x_2, a, i\} : i \in [n] \setminus \{x_1, x_2, a\}\} \\ \cup \{\{x_2, b, y\}\}.$$

So  $|\mathcal{G}| \leq 2n$ .

(iii) If there exist  $\{x_2, c, d\}, \{x_2, e, f\} \in \mathcal{F}$  for four different elements  $c, d, e, f \in [n] \setminus \{x_1, x_2\}$ , and  $\{x_1, a, b\}, \{x_1, y, z\} \in \mathcal{F}$  for four different elements  $a, b, y, z \in [n] \setminus \{x_1, x_2\}$ , then

$$\mathcal{G} \subseteq \mathcal{R} \cup \{\{x_1, c, i\} : i \in \{e, f\}\} \cup \{\{x_1, d, i\} : i \in \{e, f\}\} \cup \{\{x_2, a, i\} : i \in \{y, z\}\} \\ \cup \{\{x_2, b, i\} : i \in \{y, z\}\}.$$

So  $|\mathcal{G}| \leq n + 6$ .

Returning to case (i), we shall assume that cases (ii) and (iii) are not true. Then  $|\mathcal{F}| \leq n - 2 + 2(n - 3) = 3n - 8$ . Therefore, we have  $|\mathcal{F}| + |\mathcal{G}| \leq 6n - 14 \leq \binom{n-1}{2} + \binom{n-2}{2} + 3$ . For case (ii), we shall assume that case (iii) is not true. In addition, we may assume that at least two elements of  $\{\{x_1, c, i\} : i \in \{e, f\}\} \cup \{\{x_1, d, i\} : i \in \{e, f\}\}$  are in  $\mathcal{G}$ . Then  $|\mathcal{F}| \leq n - 2 + n - 2 + n - 3 = 3n - 7$ . Therefore, we have  $|\mathcal{F}| + |\mathcal{G}| \leq 5n - 7 \leq \binom{n-1}{2} + \binom{n-2}{2} + 3$ . For case (iii), we may assume that at least two elements of  $\{\{x_1, c, i\} : i \in \{e, f\}\} \cup \{\{x_1, d, i\} : i \in \{e, f\}\}$  are in  $\mathcal{G}$ , and at least two elements of  $\{\{x_2, a, i\} : i \in \{y, z\}\} \cup \{\{x_2, b, i\} : i \in \{y, z\}\}$  are in  $\mathcal{G}$ . Then  $|\mathcal{F}| \leq n - 2 + n - 2 + n - 2 = 3n - 6$ . Therefore, we have  $|\mathcal{F}| + |\mathcal{G}| \leq 4n \leq \binom{n-1}{2} + \binom{n-2}{2} + 3$ . This completes the proof.  $\square$

**Proof of Theorem 1.7 for  $k = 4$ .** Let  $\mathcal{F} \subseteq \binom{[n]}{4}$  be an maximum intersecting family which is neither EKR nor HM, and  $\mathcal{F} \not\subseteq \mathcal{J}_2(n, 4), \mathcal{G}_2(n, 4), \mathcal{G}_3(n, 4)$ . For any  $x \in [n]$ , let  $|\mathcal{F}(\bar{x})| \geq 3$ . Furthermore, for any  $\mathcal{G}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, 4), \mathcal{G}_2(n, 4), \mathcal{G}_3(n, 4)$ , we may assume that  $|\mathcal{G}(\bar{x})| \geq 3$ . Since  $\mathcal{F}$  is not a subfamily of  $\mathcal{G}_3(n, 4)$ , we have that  $\mathcal{F}(\bar{1}) \cup \mathcal{F}(1)$  is not isomorphic to a subfamily of  $\mathcal{C}$ . According to Lemma B.1, Proposition B.2 and the proof of Theorem 1.7 for  $k \geq 5$ , we only need to deal with the following two cases.

**Case 1.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{4}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, 4), \mathcal{G}_2(n, 4), \mathcal{G}_3(n, 4)$ ,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G}) \subseteq \mathcal{G}_2(n, 4)$ .

By conditions, there exists  $x_0, x_1, x_2$  such that

$$s_{i,j}(\mathcal{G}) \subseteq \left\{ G \in \binom{[n]}{4} : \{x_1, x_2\} \subseteq G \right\} \cup \left\{ G \in \binom{[n]}{4} : x_0 \in G, G \cap \{x_1, x_2\} \neq \emptyset \right\}.$$

Hence,  $i \in \{x_0, x_1, x_2\}$  and  $j \notin \{x_0, x_1, x_2\}$ . By symmetry, we may assume that  $x_0 = i$ . This implies that  $|\mathcal{G}(i, j)| \leq 2n - 7$ . By the maximality of  $|\mathcal{G}|$ , we have  $|\mathcal{G}(\bar{i}, \bar{j})| = \binom{n-4}{2}$ . Observe that  $\mathcal{G}(i, \bar{j}), \mathcal{G}(\bar{i}, j) \subseteq \left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : A \cap \{x_1, x_2\} \neq \emptyset \right\}$ . Let  $\mathcal{R} = \left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : \{x_1, x_2\} \subseteq A \right\}$ . By the maximality of  $|\mathcal{G}|$  and  $\mathcal{G} \not\subseteq \mathcal{G}_2(n, 4)$ , we may assume that  $\mathcal{R} \subsetneq \mathcal{G}(i, \bar{j})$  and  $\mathcal{R} \subsetneq \mathcal{G}(\bar{i}, j)$ . Note that  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) \subseteq \mathcal{R}$ . So  $(\mathcal{G}(i, \bar{j}) \setminus \mathcal{R}) \cap (\mathcal{G}(\bar{i}, j) \setminus \mathcal{R}) = \emptyset$ . Since  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting and  $n - 2 \geq 7$ , by Lemma B.4, we obtain

$$|\mathcal{F}| = |\mathcal{G}(i, j)| + |\mathcal{G}(\bar{i}, \bar{j})| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ \leq 2n - 7 + \binom{n-4}{2} + \binom{n-2}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 2 \\ < \binom{n-1}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 3.$$

**Case 2.** Suppose that  $\mathcal{F}$  is not shifted, and there exists  $\mathcal{G} \subseteq \binom{[n]}{4}$  obtained from  $\mathcal{F}$  by repeated shifting operations such that  $\mathcal{G}$  is neither EKR nor HM, and  $\mathcal{G} \not\subseteq \mathcal{J}_2(n, 4)$ ,  $\mathcal{G}_2(n, 4)$ ,  $\mathcal{G}_3(n, 4)$ ,  $|\mathcal{G}| = |\mathcal{F}|$ , but  $s_{i,j}(\mathcal{G}) \subseteq \mathcal{G}_3(n, 4)$ .

By conditions, there exist  $x_0, x_1, x_2, x_3$  such that

$$s_{i,j}(\mathcal{G}) \subseteq \left\{ G \in \binom{[n]}{4} : \{x_1, x_2, x_3\} \subseteq G \right\} \cup \left\{ G \in \binom{[n]}{4} : x_0 \in G, G \cap \{x_1, x_2, x_3\} \neq \emptyset \right\}.$$

Hence,  $i = x_0$ , or  $i \in \{x_1, x_2, x_3\}$  and  $j \notin \{x_0, x_1, x_2, x_3\}$ .

If  $i = x_0$  and  $j \in \{x_1, x_2, x_3\}$ , let  $j = x_1$ . Then  $\mathcal{G}(\bar{i}, \bar{j}) = \emptyset$  and  $|\mathcal{G}(i, j)| \leq \binom{n-2}{2}$ . Observe that  $\mathcal{G}(i, \bar{j}), \mathcal{G}(\bar{i}, j) \subseteq \left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : A \cap \{x_2, x_3\} \neq \emptyset \right\}$ . Let  $\mathcal{R} = \left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : \{x_2, x_3\} \subseteq A \right\}$ . By the maximality of  $|\mathcal{G}|$  and  $\mathcal{G} \not\subseteq \mathcal{G}_3(n, 4)$ , we may assume that  $\mathcal{R} \subsetneq \mathcal{G}(i, \bar{j})$  and  $\mathcal{R} \subsetneq \mathcal{G}(\bar{i}, j)$ . Note that  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) \subseteq \mathcal{R}$ . So  $(\mathcal{G}(i, \bar{j}) \setminus \mathcal{R}) \cap (\mathcal{G}(\bar{i}, j) \setminus \mathcal{R}) = \emptyset$ . Since  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting and  $n - 2 \geq 7$ , by Lemma B.4, we obtain

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ &\leq \binom{n-2}{2} + \binom{n-2}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 2 \\ &< \binom{n-1}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 3. \end{aligned}$$

If  $i = x_0$  and  $j \notin \{x_1, x_2, x_3\}$ , then  $|\mathcal{G}(i, j)| \leq 3n - 12$  and  $|\mathcal{G}(\bar{i}, \bar{j})| \leq n - 5$ . Observe that  $\mathcal{G}(i, \bar{j}), \mathcal{G}(\bar{i}, j) \subseteq \left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : A \cap \{x_1, x_2, x_3\} \neq \emptyset \right\}$  and  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) \subseteq \{x_1, x_2, x_3\}$ . Since  $\mathcal{G}$  is not a subfamily of  $\mathcal{G}_3(n, 4)$ , we have that  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are non-empty,  $\mathcal{G}(i, \bar{j}) \neq \{x_1, x_2, x_3\}$  and  $\mathcal{G}(\bar{i}, j) \neq \{x_1, x_2, x_3\}$ . By the maximality of  $|\mathcal{G}|$ , we may assume that  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) = \{x_1, x_2, x_3\}$ . So  $|\mathcal{G}(i, \bar{j})| \geq 2$  and  $|\mathcal{G}(\bar{i}, j)| \geq 2$ . Note that  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  are cross-intersecting. Applying Lemma 3.2 to  $\mathcal{G}(i, \bar{j})$  and  $\mathcal{G}(\bar{i}, j)$  yields

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + |\mathcal{G}(\bar{i}, \bar{j})| + |\mathcal{G}(i, \bar{j})| + |\mathcal{G}(\bar{i}, j)| \\ &\leq 3n - 12 + n - 5 + \binom{n-2}{3} - \binom{n-5}{3} - \binom{n-6}{2} + 1 \\ &= \binom{n-2}{3} - \binom{n-5}{3} - \binom{n-6}{2} + 4n - 16 \\ &< \binom{n-1}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 3, \end{aligned}$$

where the last inequality holds by direct computation.

If  $i \in \{x_1, x_2, x_3\}$  and  $j \notin \{x_0, x_1, x_2, x_3\}$ , we may assume that  $x_1 = i$ . Then  $\mathcal{G}(i, j) \subseteq \{x_2, x_3\} \cup \{\{x_0, a\} : a \in [n] \setminus \{i, j, x_0\}\}$  and  $\mathcal{G}(\bar{i}, \bar{j}) \subseteq \left\{ G \in \binom{[n] \setminus \{i, j\}}{4} : x_0 \in G, G \cap \{x_2, x_3\} \neq \emptyset \right\}$ . So  $|\mathcal{G}(i, j)| \leq n - 2$  and  $|\mathcal{G}(\bar{i}, \bar{j})| \leq \binom{n-3}{3} - \binom{n-5}{3}$ . Let

$$\mathcal{S} = \left\{ S \in \binom{[n] \setminus \{i, j\}}{3} : x_0 \in S, S \cap \{x_2, x_3\} \neq \emptyset \right\}, \quad \mathcal{T} = \left\{ T \in \binom{[n] \setminus \{i, j\}}{3} : x_0 \in T \right\}.$$

Observe that  $\mathcal{G}(i, \bar{j}), \mathcal{G}(\bar{i}, j) \subseteq \mathcal{T} \cup \{\{x_2, x_3, a\} : a \in [n] \setminus \{i, j, x_2, x_3, x_0\}\}$ . Moreover, we have  $\mathcal{G}(i, \bar{j}) \cap \mathcal{G}(\bar{i}, j) \subseteq \mathcal{S}$ . By the maximality of  $|\mathcal{G}|$  and  $\mathcal{G} \not\subseteq \mathcal{G}_3(n, 4)$ , we may assume that  $\mathcal{S} \subsetneq \mathcal{G}(i, \bar{j})$  and  $\mathcal{S} \subsetneq \mathcal{G}(\bar{i}, j)$ . Note that  $(\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}) \cap (\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}) = \emptyset$ , and both  $\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  and  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}$  are subfamilies of

$$\left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : x_0 \in A, A \cap \{x_2, x_3\} = \emptyset \right\} \cup \{\{x_2, x_3, a\} : a \in [n] \setminus \{i, j, x_2, x_3, x_0\}\}.$$

In addition,  $\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  and  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}$  are cross-intersecting.

Next we consider the following four cases, and we claim that in all these cases, the inequality  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq \binom{n-5}{2} + 2$  holds for  $n \geq 9$ . Once we have established this inequality. Then

$$\begin{aligned} |\mathcal{F}| &= |\mathcal{G}(i, j)| + 2|\mathcal{S}| + |\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j)| \\ &\leq n - 2 + 2 \left( \binom{n-3}{2} - \binom{n-5}{2} \right) + \binom{n-5}{2} + 2 + \binom{n-3}{3} - \binom{n-5}{3} \\ &= \binom{n-1}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 2 \\ &< \binom{n-1}{3} - \binom{n-5}{3} - \binom{n-6}{2} - \binom{n-7}{1} + 3, \end{aligned}$$

proving the Case 2. Let us finish the proof by proving the claim.

(i) If there exist  $\{x_2, x_3, u\}, \{x_2, x_3, v\}, \{x_2, x_3, w\} \in \mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  for three different elements  $u, v, w \in [n] \setminus \{i, j, x_2, x_3, x_0\}$ , then  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S} \subseteq \{\{x_2, x_3, a\} : a \in [n] \setminus \{i, j, x_2, x_3, x_0, u, v, w\}\}$ . So  $|\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq n - 8$ .

(ii) If there exist  $\{x_2, x_3, u\}, \{x_2, x_3, v\} \in \mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  for  $u \neq v \in [n] \setminus \{i, j, x_2, x_3, x_0\}$ , then  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S} \subseteq \{\{x_2, x_3, a\} : a \in [n] \setminus \{i, j, x_2, x_3, x_0, u, v\}\} \cup \{x_0, u, v\}$ . So  $|\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq n - 6$ .

(iii) If there exists  $\{x_2, x_3, u\} \in \mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  for  $u \in [n] \setminus \{i, j, x_2, x_3, x_0\}$ , then  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S} \subseteq \{\{x_2, x_3, a\} : a \in [n] \setminus \{i, j, x_2, x_3, x_0, u\}\} \cup \{\{x_0, u, a\} : a \in [n] \setminus \{i, j, x_2, x_3, x_0, u\}\}$ . So  $|\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq 2n - 12$ .

(iv) If there is no such  $\{x_2, x_3, u\} \in \mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  for  $u \in [n] \setminus \{i, j, x_2, x_3, x_0\}$ , then  $\mathcal{G}(i, \bar{j}) \setminus \mathcal{S} \neq \emptyset$  implies that there exists  $\{x_0, y, z\} \in \mathcal{G}(i, \bar{j}) \setminus \mathcal{S}$  for some  $y \neq z \in [n] \setminus \{i, j, x_2, x_3, x_0\}$ . So  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S} \subseteq \{\{x_2, x_3, a\} : a \in \{y, z\}\} \cup \left( \left\{ A \in \binom{[n] \setminus \{i, j\}}{3} : x_0 \in A, A \cap \{x_2, x_3\} = \emptyset \right\} \setminus \{x_0, y, z\} \right)$ . Therefore, we have  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq \binom{n-5}{2} + 2$ .

Returning to case (i), since  $\mathcal{G}(\bar{i}, j) \setminus \mathcal{S} \neq \emptyset$ , by symmetry and case (iii), we have  $|\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq 2n - 12$ . Thus  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq 3n - 20 < \binom{n-5}{2} + 2$ .

For case (ii), we may assume that case (i) is not true. If there exists  $\{x_2, x_3, a\} \in \mathcal{G}(\bar{i}, j) \setminus \mathcal{S}$  for some  $a \in [n] \setminus \{i, j, x_2, x_3, x_0, u, v\}$ , then by symmetry and case (iii), note that case (i) is not true, we have  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| \leq 2 + n - 6 = n - 4$ . Then  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq 2n - 10 \leq \binom{n-5}{2} + 2$ . Otherwise by symmetry and case (iv), we also have  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq \binom{n-5}{2} + 2$ .

We have completed the proofs of cases (i), (ii), (iv). For case (iii), we may assume that cases (i) and (ii) are not true. Furthermore, by symmetry and cases (i), (ii) (iv), we may assume that there is only one  $\{x_2, x_3, a\} \in \mathcal{G}(\bar{i}, j) \setminus \mathcal{S}$  for some  $a \in [n] \setminus \{i, j, x_2, x_3, x_0, u\}$ . Then  $|\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq 1 + n - 6 = n - 5$  and  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| \leq n - 5$ . Therefore, we obtain  $|\mathcal{G}(i, \bar{j}) \setminus \mathcal{S}| + |\mathcal{G}(\bar{i}, j) \setminus \mathcal{S}| \leq 2n - 10 \leq \binom{n-5}{2} + 2$ . This completes the proof.  $\square$