

Regular structures of an intractable enumeration problem: a diagonal recurrence relation of monomer-polymer coverings on two-dimensional rectangular lattices

Yong Kong
 Department of Biostatistics
 School of Public Health
 Yale University
 New Haven, CT 06520, USA
 email: yong.kong@yale.edu

Contents

1	Introduction	2
2	A two-dimensional sequence	5
3	Proof of the main theorem	10
3.1	Outline of the proof	10
3.2	The two sets of recurrence identities: \mathcal{P}_1 and \mathcal{P}_2	12
3.2.1	The set of recurrence identities in \mathcal{P}_1	13
3.2.2	The set of recurrence identities in \mathcal{P}_2	14
3.3	The first quadrant Q_1	16
3.4	The third quadrant Q_3	20
3.4.1	Summary	20
3.4.2	The contributions of $h_{2s-i,s-j}^{(1)}$ cancels the vertical recurrences of \mathcal{P}_1	21
3.4.3	The contributions of $h_{2s-i,s-j}^{(2)}$ cancel the horizontal recurrences of \mathcal{P}_1	21
3.4.4	The contributions of $h_{2s-i,s-j'}^{(3)}$ vanish	23

3.4.4.1	Outline of the proof	23
3.4.4.2	Transforming the inner sum	24
3.4.4.3	Recurrence relation for the new double sum $U(s, i, j)$	28
3.5	The diagonal of the third quadrant Q_3	33
3.6	The fourth quadrant Q_4	34
3.6.1	Summary	34
3.6.2	The contributions from the first term of $h_{i,j}$	36
3.6.3	The contributions from the second term $\beta_{i,j}^{2h}$ cancels those from the third term $\beta_{i,j}^{3u}$	
3.7	The right hand side	40
3.7.1	The right hand side from \mathcal{P}_1	41
3.7.2	The right hand side from \mathcal{P}_2	43
A	Some properties of the two-dimensional sequence	47
B	Gosper and Zeilberger algorithms	50
C	Some binomial coefficient identities	53
D	Some sums used in section 3.7 that are Gosper-summable	54

Abstract

The enumeration of polymer coverings on two-dimensional rectangular lattices is considered as “intractable”. We prove that the number of coverings of s polymer satisfies a simple recurrence relation $\sum_{i=0}^{2s} (-1)^i \binom{2s}{i} a_{n-i, m-i} = 2^s (2s)! / s!$ on a $n \times m$ rectangular lattice with open boundary conditions in both directions.

1 Introduction

The main result of this paper is Theorem 1, which gives a diagonal recurrence relation for the number of coverings of polymers on a $n \times m$ rectangular lattice with open boundary conditions in both directions.

Enumerations of rigid rodlike molecules on lattices has been studied by researchers in diverse fields of physics, mathematics, and theoretical computer science for a long time. In the monomer-polymer model, a linear rigid polymer covers k adjacent lattice sites, with no lattice site occupied by more than one polymer. The polymers are called k -mers, and those unoccupied lattice sites are called monomers.

Physicists have a long history of studying this model for phase transitions in equilibrium statistical mechanics [16, 4, 2, 18]. When $k = 2$, the model becomes the well-known monomer-dimer model, which is closely related to the Ising model [15, 8, 3, 14]. While it has been shown that the special case of close packed dimer problem can be solved analytically for any *planar* lattices using the Pfaffian method [8], no solution has been found for the general monomer-dimer problem where unoccupied lattice sites (monomers) are allowed, despite decades of effort.

This stark dichotomy in the computational complexity of the monomer-dimer problem has captivated researchers in theoretical computer science, particularly those focused on computational complexity theory. It has been shown that the enumeration of monomer-dimer configurations in *planar* lattices is $\#P$ -complete [7], which indicates the problem is computationally “intractable”. For counting problems, the class $\#P$ plays the same role as the more familiar NP class does for decision problems (such as the well-known satisfiability problem). The $\#P$ -complete problems are at least as hard as the NP-complete problems in computational complexity hierarchy. Currently “P versus NP” problem is perhaps the major outstanding problem in theoretical computer science. The prevailing view on the P vs NP problem is that P is not equal to NP - there are fundamental differences between these two complexity classes, and the boundaries between “easy” and “intractable” problems are impenetrable.

Even though the problem of enumeration of monomer-polymer coverings is now considered as “intractable”, regular structures have been found for the problem. For example, a simple recurrence relation was found for the enumeration of polymer coverings on two-dimensional lattice strips [13]. In this paper we will explore more regular structures in this problem, and prove a simple *diagonal* recurrence relation for square lattices (Theorem 1). Aesthetically and practically, the results on square lattices are more appealing than the narrow strips. Various optimization techniques have been used to extend the width of the lattice strips in the past, for example, [11, 9, 10, 12, 4], but still the computational complexity grows exponentially with the lattice width. The diagonal recurrence relation, as proven in this paper, allows us to investigate previously uncharted regions with larger lattice sizes that were inaccessible using earlier methods.

The main result is stated below. Consider a two-dimensional regular lattice with a width of n and a length of m , with open boundary conditions in both directions. Let s denote the number of k -mers in the lattice, and

denote the number of configurations of the s k -mers on the lattice by $a_{n,m}$. Then $a_{n,m}$ follows the following diagonal recurrence relation.

Theorem 1. *The number of configurations of s k -mers on a $n \times m$ lattice with open boundary conditions in both directions satisfies the following recurrence*

$$\sum_{i=0}^{2s} (-1)^i \binom{2s}{i} a_{n-i, m-i} = 2^s \frac{(2s)!}{s!}, \quad (1)$$

for $n \geq (k+1)s$ and $m \geq (k+1)s$.

From this recurrence we can obtain another recurrence directly.

Corollary 1.1.

$$\sum_{i=0}^{2s+1} (-1)^i \binom{2s+1}{i} a_{n-i, m-i} = 0,$$

for $n \geq (k+1)s+1$ and $m \geq (k+1)s+1$.

Proof. Use the identity

$$\binom{2s+1}{i} = \binom{2s}{i} + \binom{2s}{i-1}.$$

□

The proof of Theorem 1 is based on the following recurrence relation on a lattice strip [13].

Lemma 1 (Recurrence on a lattice strip). *For given k , n and s , the following recursive relation holds for a lattice strip with open boundary conditions in both directions.*

$$\sum_{i=0}^s (-1)^i \binom{s}{i} a_{n, m-i, s} = c(n, k)^s, \quad n \geq k \text{ and } m \geq ks, \quad (2)$$

where $c(n, k) = 2n - k + 1$.

Remark 1. *The $c(n, k)$ on rhs of Eq. (2) is a constant that depends on the boundary conditions as well as n and k , but not m or s . For different boundary conditions $c(n, k)$ can take different values [13]. Note also that Eq. (2) holds when $n < k$, but with a different value of $c(n, k)$ [13]. In this paper we only need the conditions stated in Lemma 1.*

Remark 2. *Note that the rhs of Eq. (1) depends only on s , the number of k -mers. It does not depend on k , n , or m . This is in contrast to the recurrence of lattice strips (Lemma 1), where the constant on the rhs depends on s as well as n , the width of the lattice strip.*

The paper is organized as follows. Since the proof of Theorem 1 depends crucially on a two-dimensional sequence, in section 2 this sequence is defined and its properties that are used in the proof will be elucidated. In section 3 Theorem 1 is proved. The proof needs to evaluate many single and double summations, and the major tools for summation evaluation are Gosper-algorithm and Zeilberger-algorithm. These algorithms are outlined in Appendix B.

Throughout the paper we use k for the length of the polymer (the k -mer), n and m for the width and length of the lattice, respectively, and s for the number of the polymers. The exceptions are in Appendix B on Gosper and Zeilberger algorithms, where k is conventional used as the summation index, and n as the summation parameter.

2 A two-dimensional sequence

The proof of the main theorem depends critically on a two-dimensional sequence, which acts as the coefficients for the recurrence identities that are added to specified lattice sites inside the square of size $(2s + 1) \times (2s + 1)$. This sequence will be significant for our studies.

As the discussions below and their uses in the proofs show, this sequence in several aspects looks like the sequence of binomial coefficients, but with an extra parameter s , the number of k -mers covering the lattice. It serves as the counterpart to set \mathcal{P}_2 , just as the binomial coefficient does for set \mathcal{P}_1 (see section 3.2).

Below we first give a recursive definition of the sequence, then derive an explicit formula for the ordinary generating function of the sequence, from which the explicit expression of the sequence is obtained.

The explicit expressions for the sequence have *three terms*. As the proof of the main theorem shows, each term plays a distinct role in building up or canceling out the coefficients of the recurrence identities added to each position in the lattice square.

In Appendix A we show some other interesting properties of this two-dimensional sequence. There we derive the double generating function of the

sequence, and show that each element of the sequence is an integer.

Definition 1. The coefficient sequence $h_{s,i,j}$ is defined recursively as

$$h_{s,i,j} = \begin{cases} \binom{s+j-1}{j} \frac{s-j}{s}, & i = 0, \\ -\frac{s-j+1}{i} h_{s,i-1,j-1} - \frac{j-i}{i} h_{s,i-1,j}, & i > 0, \\ 0, & j \leq i. \end{cases} \quad (3)$$

As examples, the sequence for $s = 4$ is shown in Table 1, and the sequence for $s = 5$ is shown in Table 2.

Table 1: The sequence of $h_{4,i,j}$.

$i \setminus j$	1	2	3	4
0	3	5	5	0
1		-14	-20	-5
2			24	15
3				-13

Table 2: The sequence of $h_{5,i,j}$.

$i \setminus j$	1	2	3	4	5
0	4	9	14	14	0
1		-25	-55	-70	-14
2			65	125	56
3				-85	-79
4					41

Define the ordinary generating function for the sequence as

$$H_{s,i} = \sum_{j=i+1} h_{s,i,j} x^j.$$

We have the following recursive relation for the generating function $H_{s,i}$, where H' is the derivative of H w.r.t. x . Since s is fixed, below we omit s in $H_{s,i}$ and $h_{s,i,j}$.

Lemma 2 (Recursive relation of the generating function). *For $i > 1$,*

$$H_i = - \left(\frac{sx}{i} - 1 \right) H_{i-1} + \frac{x(x-1)}{i} H'_{i-1}. \quad (4)$$

Proof. This can be proved directly from the definition of the sequence (Eq. (3)):

$$\begin{aligned} H_i &= \sum_{j=i+1} h_{i,j} x^j \\ &= - \sum_{j=i+1} \left[\frac{s-j+1}{i} h_{i-1,j-1} + \frac{j-i}{i} h_{i-1,j} \right] x^j \\ &= - \frac{(s+1)x}{i} \sum_{j=i} h_{i-1,j} x^j + \frac{x}{i} \sum_{j=i} (j+1) h_{0,j} x^j - \frac{1}{i} \sum_{j=i} (j-i) h_{i-1,j} x^j \\ &= - \frac{(s+1)x}{i} H_{i-1} + \frac{x}{i} [xH'_{i-1} + H_{i-1}] - \frac{1}{i} [xH'_{i-1} - iH_{i-1}] \\ &= - \left(\frac{sx}{i} - 1 \right) H_{i-1} + \frac{x(x-1)}{i} H'_{i-1}. \end{aligned}$$

□

From this recursive relation of the generating function, we obtain the explicit expression for H_i , containing three terms:

Proposition 1 (Explicit formula for H_i). *The explicit formula for H_i is*

$$\begin{aligned} H_i &= \sum_{j=0}^i (-1)^{j+1} \binom{s}{j} x^j \\ &\quad + (-1)^{i+1} \binom{2s}{i} \frac{x^{i+1}}{(1-x)^{s+1}} \\ &\quad + (-1)^i (2s-i) \binom{2s}{i} \sum_{t=0}^i (-1)^t \binom{i}{t} \frac{1}{2s-t} \frac{1}{(1-x)^{s-t}}. \end{aligned} \quad (5)$$

Proof. The formula can be proved by induction on i using the recurrence Eq. (4). When $i = 0$, the coefficient $h_{0,j}$ can be written as two parts,

$$h_{0,j} = \binom{s+j-1}{j} - \frac{j}{s} \binom{s+j-1}{j}.$$

Since

$$\sum_{j=1}^s \binom{s+j-1}{j} x^j = -1 + \frac{1}{(1-x)^s},$$

the generating function H_0 can be calculated as

$$\begin{aligned} H_0 &= -1 + \frac{1}{(1-x)^s} - \frac{1}{s} x D_x \left[\frac{1}{(1-x)^s} \right] \\ &= -1 - \frac{1}{(1-x)^{s+1}} + \frac{1}{(1-x)^s}, \end{aligned}$$

which agrees with Eq. (5) term by term. Here D_x is the differentiation operator d/dx .

For H_{i+1} , we consider the three terms separately and prove that each of them satisfies the recurrence of Eq. (4).

For the first term of H_{i+1} , the recursive relation of Eq. (4) gives

$$\begin{aligned} & - \left(\frac{sx}{i+1} - 1 \right) H_i^{(1)} + \frac{x(x-1)}{i+1} H_i^{(1)'} \\ &= - \left(\frac{sx}{i+1} - 1 \right) \sum_{j=0}^i (-1)^{j+1} \binom{s}{j} x^j + \frac{x-1}{i+1} \sum_{j=0}^i (-1)^{j+1} j \binom{s}{j} x^j. \quad (6) \end{aligned}$$

The second term in Eq. (6) can be transformed to

$$\begin{aligned} & \frac{x}{i+1} \sum_{j=0}^i (-1)^{j+1} j \binom{s}{j} x^j - \frac{x}{i+1} \sum_{j=0}^i (-1)^{j+1} j \binom{s}{j} x^{j-1} \\ &= \frac{x}{i+1} \sum_{j=0}^i (-1)^{j+1} j \binom{s}{j} x^j + \frac{x}{i+1} \sum_{j=0}^{i-1} (-1)^{j+1} (j+1) \binom{s}{j+1} x^j \\ &= \frac{sx}{i+1} \sum_{j=0}^i (-1)^{j+1} \binom{s}{j} x^j + (-1)^i \binom{s}{i+1} x^{i+1}, \end{aligned}$$

where the identity $j \binom{s}{j} + (j+1) \binom{s}{j+1} = s \binom{s}{j}$ is used in the second step to combine the two sums. The above expression, when combined with the first term of Eq. (6), shows it equals to $H_{i+1}^{(1)}$

$$H_{i+1}^{(1)} = \sum_{j=0}^{i+1} (-1)^{j+1} \binom{s}{j} x^j.$$

For the second term of H_{i+1} , after factoring out common factors,

$$\begin{aligned}
& - \left(\frac{sx}{i+1} - 1 \right) H_i^{(2)} + \frac{x(x-1)}{i+1} H_i^{(2)'} \\
&= (-1)^{i+1} \binom{2s}{i} \frac{x^i}{(1-x)^{s+1}} \left[\left(\frac{sx}{i+1} - 1 \right) x + \frac{x(x-1)}{i+1} \left(i+1 + \frac{(s+1)x}{1-x} \right) \right] \\
&= (-1)^{i+2} \binom{2s}{i+1} \frac{x^{i+2}}{(1-x)^{s+1}},
\end{aligned}$$

which equals to $H_{i+1}^{(2)}$.

For the third term of H_{i+1} , consider the following two equations:

$$- \left(\frac{sx}{i+1} - 1 \right) H_i^{(3)} + \frac{x(x-1)}{i+1} H_i^{(3)'} = \sum_{t=0}^i F_1, \tag{7}$$

$$\begin{aligned}
(-1)^{i+1} (2s-i-1) \binom{2s}{i+1} \sum_{t=0}^{i+1} (-1)^t \binom{i+1}{t} \frac{1}{2s-t} \frac{1}{(1-x)^{s-t}} &= \sum_{t=0}^i F_2 \\
&+ \binom{2s}{i+1} \frac{1}{(1-x)^{s-i-1}},
\end{aligned} \tag{8}$$

where the summands are

$$\begin{aligned}
F_1 &= (-1)^{i+t} (2s-i) \binom{2s}{i} \binom{i}{t} \frac{1}{2s-t} \frac{1}{(1-x)^{s-t}} \left[- \left(\frac{sx}{i+1} - 1 \right) - \frac{x(s-t)}{i+1} \right], \\
F_2 &= (-1)^{i+t+1} (2s-i-1) \binom{2s}{i+1} \binom{i+1}{t} \frac{1}{2s-t} \frac{1}{(1-x)^{s-t}}.
\end{aligned}$$

The sum with summand F_1 in Eq. (7) is the result of substituting $H_i^{(3)}$ into the recurrence Eq. (4). The left side of Eq. (8) is in the form of $H_{i+1}^{(3)}$ according to Eq. (5), and the the right side is its rearrangement. We will demonstrate that the right sides of these two equations are equal, thereby proving the correctness of $H_{i+1}^{(3)}$.

The difference of F_1 and F_2 is Gosper-summable, with the antidifference

$$(-1)^{i+t+1} \binom{2s}{i+1} \binom{i}{t-1} \frac{1}{(1-x)^{s-t}}.$$

Hence the sum of $F_1 - F_2$ is

$$\sum_{t=0}^i (F_1 - F_2) = \binom{2s}{i+1} \frac{1}{(1-x)^{s-i-1}},$$

which is the second term in Eq. (8). This shows that the right sides of Eq. (7) and Eq. (8) are equal. \square

From the explicit expression of the generating function Eq. (5), the explicit expression of the sequence can be obtained by extracting the coefficient of x from $H_i(x)$.

Corollary 1.2 (Explicit formula for $h_{i,j}$). *For $i \geq 0$,*

$$\begin{aligned} h_{i,j} = & (-1)^{j+1} \binom{s}{j} [j \leq i] \\ & + (-1)^{i+1} \binom{2s}{i} \binom{s+j-i-1}{s} \\ & + (-1)^i (2s-i) \binom{2s}{i} \sum_{t=0}^i (-1)^t \frac{1}{2s-t} \binom{i}{t} \binom{s-t-1+j}{j}, \end{aligned} \quad (9)$$

where $[\cdot]$ in the first term is Iverson's convention: $[\ell] = 1$ if ℓ is true, otherwise $[\ell] = 0$ [6]. This term is to make $h_{i,j} = 0$ for $j \leq i$.

3 Proof of the main theorem

3.1 Outline of the proof

The two main ingredients of the proof are the recurrence identities on the *lattice strips*, where the recurrences are on one of the indices of $a_{n,m}$ (Lemma 1) [13], and the sequence $h_{s,i,j}$ discussed in the previous section (Eq. (9) of Corollary 1.2).

Consider a $(2s+1) \times (2s+1)$ square inside the two-dimensional lattice, with the top-right coordinate as (n, m) . To clarify the terminology, we call the diagonal from lower-left corner to the upper-right corner as the *main diagonal*, or simply *diagonal*, instead of anti-diagonal in the context of matrices. Inside this square, by choosing appropriate coefficients as weights, we add the recurrence identities of the lattice strips (Eq. (2)), in both horizontal and

vertical directions symmetrically. After summing up these identities, along the diagonal of this square the coefficients of $a_{n-i,m-i}$ will be twice that given in Theorem 1, while at the off-diagonal positions the coefficients of $a_{n-i,m-j}$ will vanish at each lattice site. Meanwhile, the sum of the right hand sides of these recurrences gives twice that of the right hand of Eq. (1).

Since the lattice have free boundary conditions in both directions, the recurrence of the lattice strips of Lemma 1 applies to both horizontal and vertical directions. For the vertical direction, the recurrence is

$$e_{n,m}^v := \sum_{j=0}^s (-1)^j \binom{s}{j} a_{n,m-j} = c(n, k)^s, \quad m \geq ks, \quad (10)$$

and for the horizontal direction, the recurrence is

$$e_{n,m}^h := \sum_{i=0}^s (-1)^i \binom{s}{i} a_{n-i,m} = c(m, k)^s, \quad n \geq ks. \quad (11)$$

The top-right corner of the square that we are focusing on has the coordinates (n, m) . In general, lattice sites inside the $(2s + 1) \times (2s + 1)$ square have coordinates in the form of $(n - i, m - j)$, for $0 \leq i, j \leq 2s$. Below we abbreviate the coordinates as (i, j) when there is no ambiguity. Hence the top-right corner is labeled as $(0, 0)$, and the bottom-left corner $(2s, 2s)$.

We use the largest coordinates in the recurrence relations Eq. (10) and Eq. (11) to label the identities, and by using abbreviation just mentioned, at the lattice site (i, j) the recurrence relations are

$$e_{i,j}^v := \sum_{j'=0}^s (-1)^{j'} \binom{s}{j'} a_{n-i,m-j-j'} = c(n-i, k)^s,$$

$$e_{i,j}^h := \sum_{i'=0}^s (-1)^{i'} \binom{s}{i'} a_{n-i-i',m-j} = c(m-j, k)^s.$$

To facilitate the discussions, we divide the necessary recurrence identities of the lattice strips used in the proof into two sets, \mathcal{P}_1 and \mathcal{P}_2 , and denote the corresponding weights at lattice site (i, j) by $b_{i,j}$ and $\bar{h}_{i,j}$, respectively. The details of \mathcal{P}_1 and \mathcal{P}_2 are discussed in the following subsections. When the recurrence identities in \mathcal{P}_1 or \mathcal{P}_2 are summed up, they introduce terms such as $a_{n-i,m-j}$. We use $\alpha_{i,j}$ and $\beta_{i,j}$ to denote the coefficient of these terms

at position (i, j) .

$$\alpha_{i,j} = \text{coefficient of } a_{n-i,m-j} \text{ in } \left(\sum_{e_{i',j'}^v \in \mathcal{P}_1} b_{i',j'} e_{i',j'}^v + \sum_{e_{i',j'}^h \in \mathcal{P}_1} b_{i',j'} e_{i',j'}^h \right),$$

$$\beta_{i,j} = \text{coefficient of } a_{n-i,m-j} \text{ in } \left(\sum_{e_{i',j'}^v \in \mathcal{P}_2} \bar{h}_{i',j'} e_{i',j'}^v + \sum_{e_{i',j'}^h \in \mathcal{P}_2} \bar{h}_{i',j'} e_{i',j'}^h \right),$$

where $\bar{h}_{i,j}$ is related to the two-dimensional sequence discussed in section 2 (see below).

Our goals are to prove, for the left hand side of all these added up identities of lattice strips,

$$\alpha_{i,i} + \beta_{i,i} = (-1)^i \binom{2s}{i} \times 2, \quad (12)$$

$$\alpha_{i,j} + \beta_{i,j} = 0, \quad i \neq j,$$

and for the right hand side,

$$\sum_{e_{i',j'}^v \in \mathcal{P}_1} b_{i',j'} c(n-i', k)^s + \sum_{e_{i',j'}^h \in \mathcal{P}_2} \bar{h}_{i',j'} c(n-i', k)^s = 2^s \frac{(2s)!}{s!}. \quad (13)$$

3.2 The two sets of recurrence identities: \mathcal{P}_1 and \mathcal{P}_2

In this section we describe how to add the lattice strip recurrences in the vertical and horizontal directions with appropriate coefficients, so that the terms in the diagonal have the correct values of the main theorem, while the left side of the off-diagonal terms vanish (Eq. (12)).

We introduce two sets of recurrence identities, \mathcal{P}_1 and \mathcal{P}_2 . For the recurrence identities in the set \mathcal{P}_1 , binomial coefficients are used as weights, while for the set \mathcal{P}_2 the two-dimensional sequence introduced above in section 2 is used as weights.

We will prove the main result quadrant by quadrant inside the $(2s+1) \times (2s+1)$ square. The four quadrants are naturally defined by their four corners (counterclockwise): $Q_1 = \{(s, s), (0, s), (0, 0), (s, 0)\}$, $Q_2 = \{(s, s), (s, 0), (2s, 0), (2s, s)\}$, $Q_3 = \{(s, s), (2s, s), (2s, 2s), (s, 2s)\}$, and $Q_4 = \{(s, s), (s, 2s), (0, 2s), (0, s)\}$.

As we can see from Figure 1, Figure 2, Figure 3, and Figure 4, the lattice sites in different quadrants are affected differently by the added vertical and horizontal recurrence identities. We will treat each quadrant separately.

3.2.1 The set of recurrence identities in \mathcal{P}_1

In the set of \mathcal{P}_1 , we put the following vertical recurrence relations at lattice site (i, j) :

$$(-1)^j \binom{s}{j} e_{i,j}^v, \quad (14)$$

for $0 \leq i \leq s$, $0 \leq j \leq i$ in the first quadrant, and $s < i \leq 2s$, $i - s \leq j \leq s$ in the second quadrant. In other word, the weight $b_{i,j}^v$ (the superscript v is for vertical) is

$$b_{i,j}^v = (-1)^j \binom{s}{j}.$$

Similarly, horizontal recurrence relations are put in the set of \mathcal{P}_1 with the same coefficients in a symmetrical way with respect to the main diagonal. Figure 1 schematically shows the recurrence relation identities in \mathcal{P}_1 for $s = 4$, and Figure 2 for $s = 5$. The red arrowhead at the position (i, j) , which point downwards, indicates the vertical recurrence identity $e_{i,j}^v$, while the blue arrowheads, which point to the left, are for the horizontal recurrence identity $e_{i,j}^h$. The numbers at the lattice sites are the coefficients (weights) $b_{i,j}$ for the recurrence identity starting at that lattice site (Eq. (14)).

Note that since the length of the recurrence relation of lattice strips is $s + 1$, by construction the terms introduced by the identities in the set \mathcal{P}_1 are restricted inside the $(2s + 1) \times (2s + 1)$ square.

As shown below, the summation of the recurrence identities in \mathcal{P}_1 gives twice the coefficients of Eq. (1) along the diagonal, but it also introduces nonzero entries of the off-diagonal terms ($a_{n-i,m-j}$, where $i \neq j$). The set of recurrence identities in \mathcal{P}_2 (described below) is used to cancel these off-diagonal terms.

For the identities of vertical recurrences in the set \mathcal{P}_1 , when $0 \leq i \leq s$, the affected sites are in the range of $0 \leq j \leq s + i$; and when $s < i \leq 2s$, the affected sites are in the range of $i - s \leq j \leq 2s$. By simple variable changes we see that the coefficient of $a_{n-i,m-j}$ induced by the vertical recurrences in

the set \mathcal{P}_1 is given by

$$\alpha_{i,j}^v = (-1)^j \sum_{j'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{j'} \binom{s}{j-j'}. \quad (15)$$

By symmetry, the coefficient of $a_{n-i,m-j}$ at (i, j) for the horizontal recurrences to the left direction is

$$\alpha_{i,j}^h = (-1)^i \sum_{i'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{i'} \binom{s}{i-i'}. \quad (16)$$

The coefficient of $a_{n-i,m-j}$ induced by the set \mathcal{P}_1 is thus the sum of these two numbers.

Lemma 3. *The coefficient of $a_{n-i,m-j}$ induced by the set \mathcal{P}_1 is*

$$\alpha_{i,j}^{\mathcal{P}_1} = (-1)^j \sum_{j'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{j'} \binom{s}{j-j'} + (-1)^i \sum_{i'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{i'} \binom{s}{i-i'}. \quad (17)$$

From this expression it can be seen that for the main diagonal where $i = j$, the contribution of the identities of the set \mathcal{P}_1 is twice the value of Theorem 1.

Corollary 1.3. *The coefficient of $a_{n-i,m-i}$ along the diagonal induced by the set \mathcal{P}_1 is*

$$\alpha_{i,i}^{\mathcal{P}_1} = (-1)^i \binom{2s}{i} \times 2, \quad (18)$$

which is twice the coefficient of $a_{n-i,m-j}$ of Theorem 1.

3.2.2 The set of recurrence identities in \mathcal{P}_2

For the recurrence relation identities in \mathcal{P}_2 , the coefficients used are $h_{i,j}$, instead of binomial coefficients. In the first quadrant, we put the following vertical recurrence relations at lattice site (i, j) :

$$h_{i,j} e_{i,j}^v, \quad 0 \leq i < s, i < j \leq s.$$




Figure 1: (Color online) The set of recurrence identities in \mathcal{P}_1 for $s = 4$. The arrows labeled in red are for the vertical recurrences; those in blue for the horizontal recurrences. The numbers are the coefficients b_{ij} (weights) for the recurrences.

In the second quadrant, at the position (i, j) , we put

$$(-1)^s h_{2s-i, s-j} e_{i,j}^v, \quad s < i \leq 2s, 0 \leq j < i - s.$$

Similarly as for the set \mathcal{P}_1 , horizontal recurrence identities are put symmetrically about the main diagonal in the set \mathcal{P}_2 .

Figure 3 schematically shows the recurrence identities in \mathcal{P}_2 for $s = 4$, and Figure 4 for $s = 5$. The red arrowhead at the position (i, j) indicates the vertical recurrence relation identity $e_{i,j}^v$, while the blue arrowhead for the horizontal recurrence identity $e_{i,j}^h$. The numbers are the coefficients $\bar{h}_{i,j}$ (weights) for the recurrence identity starting at the lattice site. As the terms introduced by the recurrence identities in \mathcal{P}_1 , terms introduced by the recur-




Figure 2: (Color online) The set of recurrence identities in \mathcal{P}_1 for $s = 5$.

recurrence identities in \mathcal{P}_2 are restricted inside the $(2s + 1) \times (2s + 1)$ square with corners at $(0, 0)$ and $(2s, 2s)$.

Remark 3. *The restrictions on m and n in Theorem 1 is s greater than those on a lattice strip (Lemma 1). This is due to the fact that the square is of size $2s + 1$. The conditions have to be true for the smallest lattice sites where the recurrence identities are added, which are $n - s$ and $m - s$. It is clear from the Figures.*

3.3 The first quadrant Q_1

From Figure 3 and Figure 4 we see that the recurrence identities in \mathcal{P}_2 do not introduce terms in the main diagonal inside the first quadrant Q_1 . Let consider the strict lower-right triangle of the first quadrant, which excludes




Figure 3: (Color online) The set of recurrence identities in \mathcal{P}_2 for $s = 4$. The arrows labeled in red are for the vertical recurrences; those in blue for the horizontal recurrences. The numbers are the coefficients $\bar{h}_{i,j}$ (weights) for the recurrences.

the main diagonal. In this triangle, both horizontal and vertical recurrence identities in \mathcal{P}_1 introduce terms (Figure 1 and Figure 2), while only vertical recurrences in \mathcal{P}_2 have effects (Figure 3 and Figure 4). These identities are $e_{i,j}^v$, where $0 \leq i < s$ and $i + 1 \leq j \leq s$. Below we prove Eq. (12) for the first quadrant.

Partial proof of Theorem 1 in the first quadrant. For the lattice site (i, j) of the strict lower-right triangle of the first quadrant Q_1 , $0 \leq i < s$ and $i + 1 \leq j \leq s$. The coefficient of $a_{n-i, m-j}$ contributed by the identities from \mathcal{P}_1 is




Figure 4: (Color online) The set of recurrence identities in \mathcal{P}_2 for $s = 5$.

given by Eq. (17) of Lemma 3, which simplifies to

$$\begin{aligned}
\alpha_{i,j} &= (-1)^j \sum_{j'=0}^i \binom{s}{j'} \binom{s}{j-j'} + (-1)^i \sum_{i'=0}^i \binom{s}{i'} \binom{s}{i-i'} \\
&= (-1)^j \sum_{j'=0}^i \binom{s}{j'} \binom{s}{j-j'} + (-1)^i \binom{2s}{i},
\end{aligned} \tag{19}$$

where the binomial identity Eq. (66) is used in the second step.

The contribution from identities of \mathcal{P}_2 is given by

$$\beta_{i,j} = \sum_{j'=i+1}^j (-1)^{j-j'} h_{i,j'} \binom{s}{j-j'} = \sum_{j'=0}^j (-1)^{j-j'} h_{i,j'} \binom{s}{j-j'},$$

where we use the fact that $h_{i,j} = 0$ for $j \leq i$ to extend the range of summation in the second step. If we did not change the range of the summation, the proof would be more involved for the third term of $h_{i,j}$, due to nonstandard boundary conditions.

The contribution from the first term of $h_{i,j}$ is

$$\beta_{i,j}^{(1)} = (-1)^{j+1} \sum_{j'=0}^i \binom{s}{j'} \binom{s}{j-j'}. \quad (20)$$

The contribution from the second term of $h_{i,j}$ is

$$\begin{aligned} \beta_{i,j}^{(2)} &= \sum_{j'=i+1}^j (-1)^{j-j'} h_{i,j'}^{(2)} \binom{s}{j-j'} \\ &= (-1)^{i+1} \binom{2s}{i} \sum_{j'=i+1}^j (-1)^{j-j'} \binom{s+j'-i-1}{s} \binom{s}{j-j'} \\ &= (-1)^{i+1} \binom{2s}{i}. \end{aligned} \quad (21)$$

The summation in the second step is evaluated to 1, which can be obtained by Lemma 25 in Appendix C, by making variable changes $j' - i - 1 \rightarrow j'$ and $j - i - 1 \rightarrow j$. The identity is a special case of $t = 0$ in Eq. (67).

The contribution from the third term of $h_{i,j}$ is,

$$\beta_{i,j}^{(3)} = \sum_{j'=0}^j (-1)^{j-j'} h_{i,j'}^{(3)} \binom{s}{j-j'}.$$

Exchange the order of summation in the above equation, the inner sum becomes

$$\sum_{j'=0}^j (-1)^{j'} \binom{s-t-1+j'}{j'} \binom{s}{j-j'}.$$

Lemma 25 shows that the sum vanishes.

Combining above results of Eq. (19), Eq. (20), and Eq. (21), we obtain

$$\alpha_{i,j} + \beta_{i,j} = 0$$

in the strict lower-right triangle of the first quadrant Q_1 .

By symmetry, we can also prove a similar result for the strict upper-left triangle of the first quadrant Q_1 : the contributions of identities from \mathcal{P}_1 and \mathcal{P}_2 sum to zero. Since the recurrences in \mathcal{P}_2 do not affect the main diagonal in the first quadrant, together with Corollary 1.3, we prove that Eq. (12) holds in the first quadrant Q_1 . \square

3.4 The third quadrant Q_3

3.4.1 Summary

The third quadrant is more complicated than the first quadrant. In this section we focus on the strict upper-left triangle of the third quadrant, which excludes the main diagonal. Once we prove the theorem in this triangle, by symmetry we also prove the theorem in the strict lower-right triangle. The case of diagonal in the third quadrant has its unique characteristics, and is discussed in section 3.5.

On the strict upper-left triangle above the diagonal of the third quadrant where $i > j$, only the vertical recurrences from \mathcal{P}_2 (those labeled in red of Figure 3 and Figure 4) contribute, while both vertical and horizontal recurrences from \mathcal{P}_1 contribute in this triangle (Figure 1 and Figure 2).

We prove that the contributions from the first term of $\bar{h}_{i,j}$ in \mathcal{P}_2 cancel those from the vertical recurrences of \mathcal{P}_1 ,

$$\alpha_{i,j}^v + \beta_{i,j}^{(1)} = 0,$$

the contributions from the second term of $\bar{h}_{i,j}$ in \mathcal{P}_2 cancel those from the horizontal recurrences of \mathcal{P}_1 ,

$$\alpha_{i,j}^h + \beta_{i,j}^{(2)} = 0,$$

while the contributions from the third term of $\bar{h}_{i,j}$ in \mathcal{P}_2 vanish

$$\beta_{i,j}^{(3)} = 0.$$

To prove $\beta_{i,j}^{(3)} = 0$, we transform the double sum involving the third term of $h_{i,j}$ into a new form as a sum of a simple term and a new double sum. The motivation for this transform is that the new double sum satisfies a simple recurrence relation. With this new form, we show that these two terms cancel each other.

In the third quadrant at position (i, j) , the vertical recurrences in \mathcal{P}_2 contribute (see section 3.2.2, Figure 3 and Figure 4)

$$\beta_{i,j}^v = (-1)^s \sum_{j'=0}^s (-1)^{j-j'} h_{2s-i, s-j'} \binom{s}{j-j'}. \quad (22)$$

We deal each of the three terms of $h_{i,j}$ separately below.

3.4.2 The contributions of $h_{2s-i, s-j}^{(1)}$ cancels the vertical recurrences of \mathcal{P}_1

The first term of $h_{i,j}$ is $h_{i,j}^{(1)} = (-1)^{j+1} \binom{s}{j} [j \leq i]$. This term in the sum of $\beta_{i,j}^v$ above (Eq. (22)) is simplified as

$$\begin{aligned} \beta_{i,j}^{(1)} &= (-1)^s \sum_{j'=0}^s (-1)^{j-j'} (-1)^{s-j'+1} \binom{s}{s-j'} \binom{s}{j-j'} [s-j' \leq 2s-i] \\ &= (-1)^{j+1} \sum_{j'=i-s}^s \binom{s}{j'} \binom{s}{j-j'}, \end{aligned} \quad (23)$$

which cancels the contributions from the vertical recurrences of \mathcal{P}_1 (Eq. (15)).

Lemma 4.

$$\alpha_{i,j}^v + \beta_{i,j}^{(1)} = 0.$$

3.4.3 The contributions of $h_{2s-i, s-j}^{(2)}$ cancel the horizontal recurrences of \mathcal{P}_1

In the upper-left triangle of the third quadrant Q_3 , the contributions from the horizontal recurrences of \mathcal{P}_1 are

$$(-1)^i \sum_{i'=j-s}^s \binom{s}{i'} \binom{s}{i-i'}.$$

Since $i \geq j$, the boundary conditions in the sum are standard, and Chu-Vandermonde identity Eq. (66) applies.

Lemma 5. *For $i \geq j$ in upper-left triangle, the contributions from the horizontal recurrences of \mathcal{P}_1*

$$\alpha_{i,j}^h = (-1)^i \sum_{i'=j-s}^s \binom{s}{i'} \binom{s}{i-i'} = (-1)^i \binom{2s}{i}. \quad (24)$$

The sum of the second term of $h_{2s-i, s-j'}$ in the upper-left triangle of the third quadrant Q_3 takes the form of

$$\beta_{i,j}^{(2)} = (-1)^{s+i+j+1} \binom{2s}{i} \sum_{j'=0}^{i-1} (-1)^{j'} \binom{i-j'-1}{s} \binom{s}{j-j'}.$$

Let first prove the following.

Lemma 6.

$$\sum_{j'=0}^{i-1} (-1)^{j'} \binom{i-j'-1}{s} \binom{s}{j-j'} = \begin{cases} (-1)^{s+j}, & s \leq j < i, \\ 0, & \text{otherwise.} \end{cases} \quad (25)$$

Proof. Zeilberger's algorithm shows that the summand satisfies the first order homogeneous recurrence

$$(i-j-1)F(j, j') + (i-j-1)F(j+1, j') = 0, \quad (26)$$

with certificate

$$R = \frac{(i-j')(-s+j-j')}{j+1-j'}.$$

Note that the coefficient polynomial of $F(j+1, j')$ in the recurrence has zero at $j = i-1$, so the recurrence breaks up at the term $S(i)$.

For $j \leq i-1$, let evaluate $S(i-1)$. In this case the summand becomes

$$(-1)^{j'} \binom{i-j'-1}{s} \binom{s}{i-j'-1},$$

which vanishes except when $j' = i-1-s$, hence the sum is evaluated as $S(i-1) = (-1)^{s+i+1}$. From the recurrence Eq. (26) $S(j) = (-1)^{s+j}$ when $j < i$.

When $j \geq i$, the condition for the first binomial in the summand to be nonzero is

$$j' \leq i-s-1,$$

while the condition for the second binomial to be nonzero is

$$j' \geq j-s \geq i-s,$$

which show that the summand is zero for all j' , hence $S(j) = 0$ for $j \geq i$. \square

From this identity we have

Corollary 1.4. *In the upper-left triangle of the third quadrant Q_3 , the contribution of the second term of $h_{i,j}$ from \mathcal{P}_2 is*

$$\beta_{i,j}^{(2)} = (-1)^{i+1} \binom{2s}{i}. \quad (27)$$

By Eq. (24) and Eq. (27) we prove that

Corollary 1.5. *In the upper-left triangle of the third quadrant Q_3 ,*

$$\alpha_{i,j}^h + \beta_{i,j}^{(2)} = 0.$$

3.4.4 The contributions of $h_{2s-i,s-j'}^{(3)}$ vanish

3.4.4.1 Outline of the proof In this section we prove $\beta_{i,j}^{3v} = 0$ in the strict upper-left triangle of the third quadrant. Compared to the proofs of contributions from $h_{2s-i,s-j'}^{(1)}$ and $h_{2s-i,s-j'}^{(2)}$, this proof is more complicated, since a double sum is involved. The third term $h_{2s-i,s-j'}^{(3)}$ itself is a sum, thus by Eq. (22) the contributions from the third term in upper-left triangle where $i > j$ are represented as a double sum (Eq. (28) below). To prove $\beta_{i,j}^{3v} = 0$, we transform the double sum into two parts, the first part is a simple closed form expression, and the second part is a new double sum that satisfies a simple recurrence. We then prove that these two parts cancel each other.

To do this, we transform the inner sum of the double sum

$$\beta_{i,j}^{3v} = (-1)^s \sum_{j'=0}^s (-1)^{j-j'} h_{2s-i,s-j'}^{(3)} \binom{s}{j-j'}$$

to two parts (Proposition 2). The first part is a simple closed form expression (Lemma 7), and the second part is a new sum. This new sum is a correction term that is needed when $j' < i - s$ (Lemma 8). This is due to a zero in the coefficient polynomial of the lowest recurrence term, which interrupts the recurrence in the downwards direction of recurrence on j' . Some of the results of this section will be used in the section for the fourth quadrant Q_4 .

3.4.4.2 Transforming the inner sum The explicit form of full double sum involving $h_{2s-i, s-j'}^{(3)}$ of $\beta_{i,j}^v$ is

$$(-1)^{s+i} i \binom{2s}{i} \sum_{j'=0}^s (-1)^{j-j'} \binom{s}{j-j'} \sum_{t=0}^{2s-i} \frac{(-1)^t}{2s-t} \binom{2s-i}{t} \binom{2s-t-1-j'}{s-j'}. \quad (28)$$

The explicit form of the inner sum with t as the summation index is

$$\sum_{t=0}^{2s-i} \frac{(-1)^t}{2s-t} \binom{2s-i}{t} \binom{2s-t-1-j'}{s-j'}. \quad (29)$$

Zeilberger's algorithm shows the summand obeys a second order recurrence on j'

$$-(j'-s)(i-j'-s-1)F(j', t) + (2ij' - is - 2j'^2 + 2i - 5j' + s - 3)F(j'+1, t) - (j'+2)(i-j'-2)F(j'+2, t) = 0, \quad (30)$$

with the certificate

$$R = \frac{t(j'-s)(2s-t)}{(-2s+t+1+j')}.$$

Note that the lowest coefficient polynomial of the recurrence (the coefficient of $F(j', t)$) has a zero at $j' = i - s - 1$, which is within the range of summation in the outer sum of variable j' .

Below we show that when $j' \geq i - s$, the inner sum has a closed form.

Lemma 7. *When $j' \geq i - s$, the sum of Eq. (29) has the following closed form:*

$$\sum_{t=0}^{2s-i} \frac{(-1)^t}{2s-t} \binom{2s-i}{t} \binom{2s-t-1-j'}{s-j'} = \frac{(-1)^{s+i+j'}}{i} \binom{s}{j'} \binom{2s}{i}^{-1}. \quad (31)$$

Proof. It is easy to check that the closed form on the right side of Eq.(31) obeys the same second order recurrence Eq. (30). We will show that both sides have the same values at $j' = s$ and $j' = s - 1$.

When $j' = s$, the summand becomes

$$F(i, t) = \frac{(-1)^t}{2s-t} \binom{2s-i}{t}.$$

The summand obeys the first order homogeneous recurrence on i

$$iF(i, t) + (-i + 2s)F(i + 1, t) = 0,$$

with the certificate

$$R = \frac{(2s - t)t}{i - 2s}.$$

The sum can be evaluated from the recurrence as

$$\frac{(-1)^i}{i} \binom{2s}{i}^{-1},$$

which equals the right hand side of Eq. (31) when $j' = s$.

For $j' = s - 1$, the summand is

$$F(i, t) = (-1)^t \frac{s - t}{2s - t} \binom{2s - i}{t}.$$

which obeys the first order homogeneous recurrence

$$-(i - 2s + 1)F(i, t) + (i - 2s + 1)(i - 2s)F(i + 1, t) = 0,$$

with the certificate

$$R = \frac{-(2s - t)t(is - 2s^2 - s + 2st - it + i)}{(s - t)(i - 2s)}.$$

The recurrence is solved to yield

$$\frac{(-1)^{i+1}s}{i} \binom{2s}{i}^{-1},$$

which agrees with the right hand side of Eq. (31) when $j' = s - 1$, hence the identity of Eq. (29) is proved. \square

The above Lemma only holds when $j' \geq i - s$. For other values of j' that we need in the sum of Eq. (28), that is, $0 \leq j' \leq i - s - 1$, a correction term is needed. This is due to a zero in the lowest coefficient polynomial of the recurrence Eq. (30). The coefficient polynomial of $F(j', t)$ vanishes at

$$j' = i - s - 1.$$

Because of this zero, the recurrence cannot go on without ambiguity on the downshift direction of j' .

The correction term is given in the following Lemma.

Lemma 8 (The correction term). *When $j' < i - s$, the correction term is*

$$\sum_{t=1}^{s-j'} \frac{(-1)^{t+1}}{t} \binom{i-j'-1}{s+t-1} \binom{s}{t-1} \binom{s-j'}{t}^{-1}. \quad (32)$$

To prove the Lemma, we need first to show that the value of the correction term when $j' = i - s - 1$, plus the value of the right side of Eq. (31) evaluated at $j' = i - s - 1$, is the correct value of the original sum (Eq.(29)). This is done in Lemma 9.

Lemma 9 (Value at $j' = i - s - 1$). *When $j' = i - s - 1$, the original sum of Eq. (29) equals to*

$$\frac{1}{2s-i+1} + \frac{(-1)^{2i-1}}{i} \binom{s}{-s+i-1} \binom{2s}{i}^{-1}. \quad (33)$$

where the first term is the value of the correction term Eq. (32) at $j' = i - s - 1$, and the second term is the value of the right side of Eq. (31) at $j' = i - s - 1$.

Proof. When $j' = i - s - 1$, the summand of the original sum of Eq. (29) obeys a homogeneous second order recurrence

$$(i-2s-1)iF(i, t) - (2i-s+1)(i-2s)F(i+1, t) + (i-s+1)(i-2s+1)F(i+2, t) = 0,$$

with the certificate

$$-\frac{(i-2s-1)s(2s-t)t}{(i-2s)(-3s+t+i)}.$$

It can be show that expression (33) obeys this recurrence by direct calculation.

For initial conditions, when $i = 2s$, the first binomial coefficient of the summand of the original sum vanishes for all values of t except when $t = 0$. In this case, the sum takes a value of $1/2$, which agrees with the expression Eq. (33).

when $i = 2s - 1$, the first binomial coefficient of the summand of the original sum vanishes for all values of t except when $t = 0$ or $t = 1$. The sum of these two terms are

$$\frac{1}{4} \frac{3s-1}{2s-1},$$

which agrees with the expression Eq. (33) with direct substitutions. This proves the Lemma. \square

Next we prove that the correction term Eq.(32) vanishes when $j' \geq i - s$.

Lemma 10 (The correction term vanishes when $j' \geq i - s$). *When $j' \geq i - s$, the correction term Eq.(32) vanishes.*

Proof. The first binomial coefficient in the numerator of the summand of Eq.(32) vanishes when $j' > i - s - t$. Since the index of summation $t \geq 1$, this binomial coefficient vanishes for all the range of t , so is the sum. \square

By the the previous two Lemmas, we proceed to prove the correctness of Lemma 8.

Proof of Lemma 8. When $j' = i - s - 1$, the summand of Eq.(32) is non-vanishing only when $t = 1$, and the above correction term takes a value of

$$\frac{1}{2s - i + 1}.$$

The summand of the correction term

$$F(j', t) = \frac{(-1)^{t+1}}{t} \binom{i - j' - 1}{s + t - 1} \binom{s}{t - 1} \binom{s - j'}{t}^{-1}$$

satisfies the first order inhomogeneous recurrence (with j' as the parameter)

$$(-j' + s)F(j', t) + (j' + 1)F(j' + 1, t) = \frac{s(j' - s)}{(-j' + s)(i - j' - 1)} \binom{i - j' - 1}{s}, \quad (34)$$

with the certificate

$$R = \frac{(s + t - 1)(j' - s)}{i - j' - 1}.$$

It can be checked that the inhomogeneous recurrence of Eq. (34) obeys the recurrence (30) that original sum satisfies.

The only things left are to confirm that the initial conditions also agree. In Lemma 10 we show the correction term vanishes when $j' \geq i - s$, and in Lemma 9 we show the initial condition agrees at $j' = i - s - 1$. By taken these results together, the correctness of Eq.(32) is proved. \square

Putting all above results together, we prove in the upper-left triangle of the third quadrant Q_3 , for $0 \leq j' \leq s$, the following result holds.

Proposition 2 (Transform of the inner sum). *For $0 \leq j' \leq s$, the inner sum of $\beta_{i,j}^{3v}$ is transformed to*

$$\begin{aligned} & \sum_{t=0}^{2s-i} \frac{(-1)^t}{2s-t} \binom{2s-i}{t} \binom{2s-t-1-j'}{s-j'} \\ &= \frac{(-1)^{s+i+j'}}{i} \binom{s}{j'} \binom{2s}{i}^{-1} + \sum_{t=1}^{s-j'} \frac{(-1)^{t+1}}{t} \binom{i-j'-1}{s+t-1} \binom{s}{t-1} \binom{s-j'}{t}^{-1}. \end{aligned} \quad (35)$$

Denote the first term of the rhs of Eq. (35) by $w(s, i, j')$. By putting the outer summation back (Eq. (28)), and by using identity (66), the sum of $w(s, i, j')$ can be evaluated to a closed form.

Lemma 11.

$$(-1)^{s+i+j} i \binom{2s}{i} \sum_{j'=0}^s (-1)^{j'} w(s, i, j') \binom{s}{j-j'} = (-1)^j \binom{2s}{j}.$$

Hence in the strict upper-left triangle of the third quadrant Q_3 we transform the double sum involving the third term of $h_{i,j}$ into another form.

Proposition 3 (Transformed double sum). *In the strict upper-left triangle of the third quadrant Q_3 , the double summation of the third term of $\beta_{i,j}^v$ takes a new form of*

$$\beta_{i,j}^{3v} = (-1)^j \binom{2s}{j} + U(s, i, j), \quad (36)$$

with

$$U(s, i, j) = (-1)^{s+i+j} i \binom{2s}{i} \sum_{j'=0}^s \sum_{t=1}^{s-j'} \frac{(-1)^{j'+t+1}}{t} \binom{i-j'-1}{s+t-1} \binom{s}{t-1} \binom{s}{j-j'} \binom{s-j'}{t}^{-1}. \quad (37)$$

3.4.4.3 Recurrence relation for the new double sum $U(s, i, j)$ With this new transformed form of $\beta_{i,j}^{3v}$, we can obtain a simple recurrence for the double sum using the multivariable version of Zeilberger's algorithm [19, 1].

Proposition 4 (Recurrence of the double sum). *The double sum satisfies the following first order inhomogeneous recurrence*

$$-U(s, i, j) + U(s, i + 1, j) = g(s, i, j), \quad (38)$$

where the inhomogeneous term $g(s, i, j)$ is described in Proposition 6.

Proof. If we leave out the factor $(-1)^{s+j+1}$, the summand

$$F(i; j', t) = (-1)^{i+t+j'} \frac{i}{t} \binom{2s}{i} \binom{i-j'-1}{s+t-1} \binom{s}{t-1} \binom{s}{j-j'} \binom{s-j'}{t}^{-1}$$

satisfies a first order recurrence

$$-F(i; j', t) + F(i + 1; j', t) = \Delta_{j'}(i; j', t) + \Delta_t(i; j', t),$$

where $\Delta_{j'}$ and Δ_t are the forward difference operators on j' and t , respectively (see Appendix B).

The certificates are

$$R_{j'} = 0, \\ R_t = \frac{(s - j' - t + 1)(s + t - 1)}{(i - j' + i + 1 - s - t)}.$$

By summing the variables j' and t on the left side of the recurrence, and by telescoping on the right side, the double sum satisfies a first order inhomogeneous recurrence

$$-U(s, i, j) + U(s, i + 1, j) = g(s, i, j),$$

where $g(s, i, j)$ is a single sum. The inhomogeneous term $g(s, i, j)$ is evaluated in Proposition 6. \square

Since Proposition 6 demonstrates that when $i > j$, the inhomogeneous term vanishes, by the recurrence of Eq. (38) in Proposition 4 $U(s, i, j)$ is independent of i . Below we show explicitly that when $i = 2s$, the double sum $U(s, 2s, j)$ equals to $(-1)^{j+1} \binom{2s}{j}$, hence the double sum $U(s, i, j)$ takes this value when $i > j$.

Lemma 12 (Value of the double sum at $i = 2s$). *When $i = 2s$,*

$$U(s, 2s, j) = (-1)^{j+1} \binom{2s}{j}.$$

Proof. When $i = 2s$, the inner sum of Eq. (37) is Gosper-summable, with the antidifference as

$$\frac{(-1)^{s+j+j'+t}(s+t-1)}{t} \binom{-j'+2s-1}{s+t-1} \binom{s}{t-1} \binom{s}{j-j'} \binom{s-j'}{t}^{-1}.$$

By using this expression of the antidifference, the inner sum of t is evaluated as

$$(-1)^{j+1} \binom{s}{j'} \binom{s}{j-j'} + (-1)^{s+j'+j} \binom{s}{j-j'} \binom{-j'+2s-1}{s-1}.$$

The sum of the first term over j' is evaluated as

$$(-1)^{j+1} \binom{2s}{j}$$

by using identity (66). We show that the sum over j' of the second term in the antidifference vanishes.

It turns out that the second term is also Gosper-summable w.r.t. j' , with the antidifference as

$$(-1)^{s+j'+j} \frac{s}{j-2s} \binom{2s-j'}{s} \binom{s-1}{j-j'}.$$

This antidifference vanishes at both ends at $j' = s+1$ and $j' = 0$ when $j \geq s$, hence the sum is zero. □

From this Lemma and Proposition 4 we prove that the double sum $U(s, i, j)$ takes the value of $(-1)^{j+1} \binom{2s}{j}$ in the strict upper-left triangle of the third quadrant.

Corollary 1.6 (Double sum in the strict upper-left triangle). *The double sum in the strict upper-left triangle, where $i > j$, has the following value*

$$U(s, i, j) = (-1)^{j+1} \binom{2s}{j}.$$

From Proposition 3 and Corollary 1.6 we see that $\beta_{i,j}^{(3)}$ vanishes in the strict upper-left triangle of the third quadrant.

Proposition 5. *In the strict upper-left triangle of the third quadrant,*

$$\beta_{i,j}^{(3)} = 0.$$

Now we deal with the inhomogeneous term $g(s, i, j)$ of the recurrence relation Eq. (38). As mentioned in Appendix B, for multi-variable multisum with nonstandard boundary conditions, the application of Zeilberger's algorithm becomes more complicated. For double sums with nonstandard boundary conditions, two single sums will appear in the inhomogeneous terms (see the Lemmas 22, 23 and 24 in Appendix B). If the running index is the same as the parameter of the recurrence, additional single sums will be needed to cover the elements on the boundary.

In our case here we will use i as the recurrence parameter. Based on the relation of the summation indices t and j' of the double sum $U(s, i, j)$ (Eq. (37)), Lemma 24 will be used in the following. Since one of the certificates $R_{j'} = 0$, we need to only work on the single sum related to R_t (Proposition 4).

After putting back the factor $(-1)^{s+j+1}$, the inhomogeneous term of the recurrence in Proposition 4 is given below. The case of $i = j$ will be used in the next section when we discuss the terms on the diagonal (section 3.5).

Proposition 6 (Inhomogeneous term).

$$g(s, i, j) = \begin{cases} (-1)^{j+1} \binom{2s}{i}, & s \leq i = j, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Denote the summand of the double sum of Eq. (37) by F .

Using the notation of Lemma 24, for the double sum $U(s, i, j)$, we have

$$\begin{aligned} u_{j'} &= 0, & v_{j'} &= s, \\ u_t &= 1. \end{aligned}$$

When $t = 1$,

$$G_t(i, j', 1) = R_t F \Big|_{t=1} = (-1)^{i+j'+1} \binom{2s}{i} \binom{i-j'-1}{s-1} \binom{s}{j-j'}, \quad (39)$$

and when $t = s + 1 - j'$,

$$G_t(i, j', s + 1 - j') = R_t F \Big|_{t=s+1-j'} = 0.$$

Hence the inhomogeneous term is

$$g(s, i, j) = - \sum_{j'=0}^s G_t(i, j', 1).$$

The summand $G_t(i, j', 1)$ is Gosper-summable, with antidifference $W(j')$ as

$$W(j') = (-1)^{i+j'} \binom{2s}{i} \binom{i-j'}{i-j} \binom{i-j-1}{i-s-j'}.$$

We discuss two cases $i > j$ and $i = j$ separately for $W(j')$. The case of $i = j$ will be used in the discussions of the terms along the diagonal.

When $i > j$, for $W(j')$ to be nonzero, the second binomial coefficient needs $j' \geq j$, while the third binomial coefficient needs $-j-1 \geq j'-s$, which is $j \leq j'-s+1$. But we have the range for j' and j as $j' \leq s \leq j$, so $j' = j = s$ is the only choice; this leads to $s \leq 1$; so the only combinations for $W(j')$ to be nonzero is when $(s = 1, i = 2, j = 1, j' = 1)$. In this case $W(j') = -1$, a constant. Hence When $i > j$, $g(s, i, j) = 0$.

For $i = j$, there is a pole in the antidifference $W(j')$, so we use Zeilberger's algorithm to do the sum. The summand $G_t(i, j') = G_t(i, j', 1)$ when evaluated at $i = j$ is

$$G_t(i, j') = (-1)^{i+j'+1} \binom{2s}{i} \binom{i-j'-1}{s-1} \binom{s}{i-j'},$$

which obeys a first order homogeneous recurrence

$$(-2s+i)G_t(i, j') + (i+1)G_t(i+1, j') = 0,$$

with the certificate as

$$R = \frac{(i^2 + 3i - is - 3s + j's - 2ij' + j'^2 - 2j' + 1)(i-j')(i-j'-s)}{(i-j'-s+1)(i-j'+1)}.$$

The sum $g(s, i, i) = g(s, i) = \sum_{j'=0}^s G_t(i, j')$ can be obtained as

$$\begin{aligned} g(s, i) &= \frac{2s-i+1}{i} g(s, i-1) \\ &= \frac{2s-i+1}{i} \frac{2s-i+2}{i-1} \cdots \frac{s}{s+1} g(s, s) \\ &= \frac{s!^2}{i!(2s-i)!} g(s, s). \end{aligned}$$

When $i = s$, $g(s, s)$ can be easily obtained, since only one nonvanishing term remains for the sum when $j' = 0$:

$$g(s, s) = (-1)^s \binom{2s}{s}.$$

Hence we have for $i = j$,

$$g(s, i) = (-1)^s \binom{2s}{i}.$$

□

3.5 The diagonal of the third quadrant Q_3

We show in this section that overall contributions from the recurrence relations in the set \mathcal{P}_2 in the third quadrant diagonal is zero. Since we have already shown that the recurrence identities in the set \mathcal{P}_1 give the correct coefficients along the diagonal (by a factor of 2) (Corollary 1.3), the main theorem is proved for the diagonal in the third quadrant.

First, from Eq. (23) we see that when $i = j$, the boundary condition of the summation of $\beta_{i,j}^{(1)}$ becomes standard, so the identity (66) can be used to evaluate the sum. Hence the contribution from the first term of $h_{i,j}$ of the vertical recurrence identities in the set \mathcal{P}_2 is

$$\beta_{i,i}^{(1)} = (-1)^{i+1} \binom{2s}{i}. \quad (40)$$

The horizontal recurrences are not long enough to touch the diagonal in the third quadrant (Figure 3 and Figure 4).

We have also proved that the contribution from the second term of $h_{i,j}$ of the recurrence identities in the set \mathcal{P}_2 vanishes in the diagonal (Lemma 6).

For the third term of $h_{i,j}$, we first show that the double sum in Proposition 3 (the correction term) vanishes in the diagonal.

Lemma 13 (Double sum vanishes). *When $i = j$,*

$$U(s, i, i) = 0.$$

Proof. The double sum $U(s, i, j)$ follows the recurrence of Eq. (38) on the diagonal,

$$-U(s, i, i) + U(s, i + 1, i) = (-1)^{i+1} \binom{2s}{i},$$

but we have shown that $U(s, i + 1, i) = (-1)^{i+1} \binom{2s}{i}$ in Corollary 1.6, hence $U(s, i, i) = 0$. □

From the above Lemma we see the contribution of the third term of $h_{i,j}$ of the recurrence identities in the set \mathcal{P}_2 is given by the first term of Eq. (36),

$$\beta_{i,i}^{(3)} = (-1)^i \binom{2s}{i}. \quad (41)$$

Putting Eq. (40), Eq. (25), and Eq. (41) together, we have

Proposition 7. *In the diagonal of the third quadrant, the contributions from the recurrence identities in the set \mathcal{P}_2 is zero.*

$$\beta_{i,i} = \beta_{i,i}^{(1)} + \beta_{i,i}^{(2)} + \beta_{i,i}^{(3)} = 0.$$

Partial proof of the main theorem in the third quadrant. By combining Lemma 4, Corollary 1.5, and Proposition 5, we prove that in the strict upper-left triangle of the third quadrant, the terms from the recurrence identities from the sets \mathcal{P}_1 and \mathcal{P}_2 cancel each other out. By symmetry, this also holds for the strict lower-right triangle in the third quadrant. By combining Corollary 1.3 and Proposition 7 we prove that the terms in the diagonal of the third quadrant have the correct values of the main theorem, by a factor of two. Hence the left side of the main theorem Eq. (12) is proved in the third quadrant. \square

3.6 The fourth quadrant Q_4

3.6.1 Summary

In the fourth quadrant Q_4 , $0 \leq i \leq s$ and $s \leq j \leq 2s$. As for the third quadrant, the situation is also complicated, but in a different way from the third quadrant, since at each lattice site, both horizontal and vertical recurrence identities from both \mathcal{P}_1 and \mathcal{P}_2 contribute (Figure 1, Figure 2, Figure 3, and Figure 4).

The site (s, s) is discussed in the Q_3 section, and it is excluded from the following discussions of Q_4 .

At each position (i, j) in the fourth quadrant Q_4 , where $0 \leq i \leq s$ and

$s \leq j \leq 2s$, the contributions from \mathcal{P}_1 are

$$\begin{aligned} \alpha_{i,j}^v + \alpha_{i,j}^h &= (-1)^j \sum_{j'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{j'} \binom{s}{j-j'} + (-1)^i \sum_{i'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{i'} \binom{s}{i-i'} \\ &= (-1)^j \sum_{j'=j-s}^i \binom{s}{j'} \binom{s}{j-j'} + (-1)^i \sum_{i'=j-s}^i \binom{s}{i'} \binom{s}{i-i'}. \end{aligned} \quad (42)$$

The contributions from \mathcal{P}_2 are

$$\begin{aligned} \beta_{i,j}^v &= \sum_{j'=\max(i+1,j-s)}^s (-1)^{j-j'} h_{i,j'} \binom{s}{j-j'} = \sum_{j'=0}^s (-1)^{j-j'} h_{i,j'} \binom{s}{j-j'} \\ \beta_{i,j}^h &= (-1)^s \sum_{i'=0}^i (-1)^{i-i'} h_{2s-j,s-i'} \binom{s}{i-i'} = (-1)^s \sum_{i'=0}^{j-s} (-1)^{i-i'} h_{2s-j,s-i'} \binom{s}{i-i'}. \end{aligned} \quad (43)$$

We will prove the following identities. The contributions from the first term of the vertical (horizontal) recurrence identities in \mathcal{P}_2 cancel those from the vertical (horizontal) identities in \mathcal{P}_1 ,

$$\begin{aligned} \alpha_{i,j}^v + \beta_{i,j}^{1v} &= 0, \\ \alpha_{i,j}^h + \beta_{i,j}^{1h} &= 0. \end{aligned} \quad (44)$$

The contributions from the second term of the vertical (horizontal) recurrence identities in \mathcal{P}_2 cancel those from the third term of the horizontal (vertical) identities in \mathcal{P}_2 ,

$$\begin{aligned} \beta_{i,j}^{2v} + \beta_{i,j}^{3h} &= 0, \\ \beta_{i,j}^{2h} + \beta_{i,j}^{3v} &= 0. \end{aligned} \quad (45)$$

Combining Eq. (44) and Eq. (45) we prove that

$$\alpha_{i,j} + \beta_{i,j} = 0$$

in the fourth quadrant.

3.6.2 The contributions from the first term of $h_{i,j}$

Lemma 14 (The first term of $h_{i,j}$). *The contributions from the first term of the vertical (horizontal) recurrence identities in \mathcal{P}_2 cancel those from the vertical (horizontal) identities in \mathcal{P}_1 (Eq. (44)).*

Proof. The contribution from the vertical identities of \mathcal{P}_1 is

$$\alpha_{i,j}^v = (-1)^j \sum_{j'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{j'} \binom{s}{j-j'} = (-1)^j \sum_{j'=j-s}^i \binom{s}{j'} \binom{s}{j-j'}, \quad (46)$$

while the contribution from horizontal identities of \mathcal{P}_1 is

$$\alpha_{i,j}^h = (-1)^i \sum_{i'=\max(0,i-s,j-s)}^{\min(s,i,j)} \binom{s}{i'} \binom{s}{i-i'} = (-1)^i \sum_{i'=j-s}^i \binom{s}{i'} \binom{s}{i-i'}. \quad (47)$$

The contribution of the first term of $h_{i,j'}$ in the sum of $\beta_{i,j}^v$ is simplified as

$$\beta_{i,j}^{1v} = \sum_{j'=j-s}^i (-1)^{j-j'} (-1)^{j'+1} \binom{s}{j'} \binom{s}{j-j'} = (-1)^{j+1} \sum_{j'=j-s}^i \binom{s}{j'} \binom{s}{j-j'},$$

which cancels the first term in Eq. (46).

The first term of $h_{2s-j,s-i'}$ in the summand of $\beta_{i,j}^h$ is

$$(-1)^{s-i'+1} \binom{s}{s-i'} [s-i' \leq 2s-j],$$

which is simplified in the sum as

$$\begin{aligned} \beta_{i,j}^{1h} &= (-1)^s \sum_{i'=0}^i (-1)^{i-i'} (-1)^{s-i'+1} \binom{s}{s-i'} [s-i' \leq 2s-j] \binom{s}{i-i'} \\ &= (-1)^{i+1} \sum_{i'=j-s}^i \binom{s}{i'} \binom{s}{i-i'}. \end{aligned}$$

This cancels the second term in Eq. (47). □

3.6.3 The contributions from the second term $\beta_{i,j}^{2h}$ cancels those from the third term $\beta_{i,j}^{3v}$

Proposition 8. *In the fourth quadrant Q_4 , the contributions from the second term of the horizontal recurrence identities in \mathcal{P}_2 cancel those from the third term of the vertical identities in \mathcal{P}_2*

$$\beta_{i,j}^{2h} + \beta_{i,j}^{3v} = 0.$$

where

$$\beta_{i,j}^{2h} = (-1)^{s+i+j} \binom{2s}{j} \sum_{i'=0}^s (-1)^{i'} \binom{j-i'-1}{s} \binom{s}{i-i'},$$

$$\beta_{i,j}^{3v} = (-1)^{i+j+1} (2s-i) \binom{2s}{i} \sum_{j'=0}^s \sum_{t=0}^i (-1)^{t+j'} \frac{1}{2s-t} \binom{i}{t} \binom{s-t-1+j'}{j'} \binom{s}{j-j'}.$$

To prove Proposition 8 we would like to express the *inner sum* w.r.t. t in $\beta_{i,j}^{3v}$ in two parts, a simple closed form term and a second correction sum, as we did for the third quadrant Q_3 . Unlike the third quadrant, the correction term is for $j' > i$.

Below we first make this transform for the inner sum of $\beta_{i,j}^{3v}$.

Lemma 15 (Transform of the inner sum). *The inner sum of $\beta_{i,j}^{3v}$ w.r.t. t can be rewritten as*

$$\sum_{t=0}^i (-1)^t \frac{1}{2s-t} \binom{i}{t} \binom{s-t-1+j'}{j'} =$$

$$\frac{(-1)^{i+j'}}{2s-i} \binom{s}{j'} \binom{2s}{i}^{-1} + \sum_{t=1}^{j'} \frac{(-1)^{t+1}}{t} \binom{s}{t-1} \binom{s+j'-i-1}{s+t-1} \binom{j'}{t}^{-1}.$$

Proof. Use the following changes of variables on Eq. (35):

$$\begin{aligned} i &\rightarrow 2s - i, \\ j' &\rightarrow s - j'. \end{aligned} \tag{48}$$

□

Now the sum $\beta_{i,j}^{3v}$ can be written in two parts:

Lemma 16.

$$\beta_{i,j}^{3v} = (-1)^{j+1} \binom{2s}{j} + (-1)^{i+j} (2s-i) \binom{2s}{i} \sum_{j'=0}^s \sum_{t=1}^{j'} \frac{(-1)^{j'+t}}{t} \binom{s}{t-1} \binom{s+j'-i-1}{s+t-1} \binom{s}{j-j'} \binom{j'}{t}^{-1}.$$

Proof. The outer sum w.r.t. j' can be similarly carried out on the first term of Lemma 15 as in Lemma 11. \square

We prove Proposition 8 by showing that $\beta_{i,j}^{2h}$ and $-\beta_{i,j}^{3v}$ satisfy the same recurrence and have the same initial conditions. The following Lemma shows the recurrence satisfied by $\beta_{i,j}^{2h}$.

Lemma 17 (Recurrence of $\beta_{i,j}^{2h}$). *The sum $\beta_{i,j}^{2h}$ satisfies the following first order inhomogeneous recurrence:*

$$\beta_{i+1,j}^{2h} - \beta_{i,j}^{2h} = (-1)^{i+j+s+1} \binom{2s}{i+1} \binom{2s-i-1}{j-i-1} \binom{j-i-2}{j-s-1}. \quad (49)$$

Proof. The summand of $\beta_{i,j}^{2h}$

$$F(i, j') = (-1)^{s+i+j+j'} \binom{2s}{j} \binom{j-i'-1}{s} \binom{s}{i-i'}$$

obeys a first order recurrence shown by the Zeilberger's algorithm

$$(-i-1+j)F(i, j') + (i-j+1)F(i+1, j') = G(i, j'+1) - G(i, j'),$$

with the certificate as

$$R = \frac{(j-j')(-s+i-j')}{i+1-j'}.$$

Sum up with j' from 0 to s by telescoping to obtain Eq. (49), as the value of $G(i, s+1) - G(i, 0) = -G(i, 0)$, since $G(i, s+1) = 0$. \square

The recurrence satisfied by $\beta_{i,j}^{3v}$ is given below.

Lemma 18 (Recurrence of $\beta_{i,j}^{3v}$). *The double sum in $\beta_{i,j}^{3v}$ of Lemma 16 satisfies the following first order inhomogeneous recurrence:*

$$\beta_{i+1,j}^{3v} - \beta_{i,j}^{3v} = (-1)^{i+j+s} \binom{2s}{i+1} \binom{2s-i-1}{j-i-1} \binom{j-i-2}{j-s-1}. \quad (50)$$

Proof. The summand F of the double summation in $\beta_{i,j}^{3v}$

$$F(i; j', t) = (-1)^{i+j+j'+t} \frac{2s-i}{t} \binom{2s}{i} \binom{s}{t-1} \binom{s+j'-i-1}{s+t-1} \binom{s}{j-j'} \binom{j'}{t}^{-1}$$

satisfies a first order inhomogeneous recurrence by the multi-variable multi-sum version of Zeilberger's algorithm

$$-F(i, j', t) + -F(i+1, j', t) = g(s, i, j),$$

with the certificates

$$R_{j'} = 0, \\ R_t = \frac{-(j'-t+1)(s+t-1)}{(-s-j'+i+1)(i+1)}.$$

Sum up both sides on j' and t , the right side telescopes, with two single sums left (Lemma 23 in Appendix B). In this case, since $R_{j'} = 0$ and $R_t F(t = j'+1) = 0$, there is only one single sum left:

$$\begin{aligned} RHS &= - \sum_{j'=1}^s R_t F(t=1) \\ &= (-1)^{i+j} \sum_{j'=1}^s (-1)^{j'} \frac{s(i-2s)}{(-s-j'+i+1)(i+1)} \binom{2s}{i} \binom{s+j'-i-1}{s} \binom{s}{j-j'}. \end{aligned}$$

The summand is Gosper-summable, with the antidifference as

$$(-1)^{i+j+j'+1} \binom{2s}{i+1} \binom{s+j'-i-2}{j-i-1} \binom{j-i-2}{j-j'}.$$

Since the antidifference vanishes at $j' = 1$, set $j' = s+1$ in the antidifference to get the sum, which proves Eq.(50). \square

Proof of Proposition 8. Lemma 17 and Lemma 18 show that $\beta_{i,j}^{2h}$ and $-\beta_{i,j}^{3v}$ satisfy the same first order inhomogeneous recurrence. We only need to check that $\beta_{i,j}^{2h}$ and $-\beta_{i,j}^{3v}$ share the same initial value. At $i = 0$,

$$\beta_{0,j}^{2h} = (-1)^{s+j} \binom{2s}{j} \binom{j-1}{s},$$

while the value of $\beta_{i,j}^{3v}$ when $i = 0$ is

$$\beta_{0,j}^{3v} = (-1)^j \sum_{j'=0}^s (-1)^{j'} \binom{s-1+j'}{j'} \binom{s}{j-j'} = (-1)^{s+j+1} \binom{2s}{j} \binom{j-1}{s} = -\beta_{0,j}^{2h}.$$

This proves Proposition 8. \square

We summarize the results for the fourth quadrant.

Partial proof of the main theorem in the fourth quadrant. Lemma 14 proves the identities of Eq. (44), and Proposition 8 proves the second identity of (45). By symmetry, the first identity of (45) can be proved in a similar way. Put these results together, the second identity of Eq. (12) is proved in the fourth quadrant. \square

By symmetry, a similar conclusion is reached for the second quadrant Q_2 .

3.7 The right hand side

All the recurrence identities from the sets \mathcal{P}_1 and \mathcal{P}_2 introduce terms on the right hand side of the equation. In this section we calculate the sum of these numbers on the right hand side. By symmetry we only need to look at the vertical contributions from \mathcal{P}_1 and \mathcal{P}_2 . In the following the superscript v is omitted.

For two-dimensional lattice strips, the right hand side of the recurrences depend on the boundary conditions of the horizontal (vertical) direction if the recurrence is in the vertical (horizontal) direction. For recurrence in the vertical direction, the right hand side is $(2n + \text{constant})^s$. Below we generalize slightly and assume the right hand side of the vertical recurrence is

$$(\lambda n + \eta)^s.$$

We will show that for the diagonal recurrence in the square, the right hand side depends only on λ and s , and not on n , m , nor η .

We discuss the sets \mathcal{P}_1 and \mathcal{P}_2 separately.

3.7.1 The right hand side from \mathcal{P}_1

For vertical recurrences, the right hand side constant is the same for each *column*. The following Lemma gives the sum of the coefficients $b_{i,j}$ of the recurrence identities in \mathcal{P}_1 at a given column i (Figure 1 and Figure 2, red arrowheads), $x(i)$ (for $0 \leq i \leq s$) and $x'(i)$ (for $s < i \leq 2s$).

Lemma 19.

$$\begin{aligned} x(i) &= \sum_{j'=0}^i (-1)^{j'} \binom{s}{j'} = (-1)^i \binom{s-1}{i}, \quad 0 \leq i \leq s, \\ x'(i) &= \sum_{j'=i-s}^s (-1)^{j'} \binom{s}{j'} = (-1)^{s+i} \binom{s-1}{2s-i}, \quad s < i \leq 2s. \end{aligned} \tag{51}$$

Proof. The summand of the first equation is Gosper-summable, with the antidifference shown in Lemma 27.

The second identity can be obtained from the first by changing the variable $i \rightarrow 2s - i$ and $j' \rightarrow s - j'$. \square

Hence for vertical recurrences from \mathcal{P}_1 , at column $n - i$ when $0 \leq i \leq s$, the contributions from the right side are

$$X(i) = c(n - i)^s x(i) = (-1)^i \binom{s-1}{i} c(n - i)^s.$$

Note that when $i = s$, $X(s) = 0$.

When $s < i \leq 2s$,

$$X'(i) = c(n - i)^s x'(i) = (-1)^{s+i} \binom{s-1}{2s-i} c(n - i)^s.$$

The total contributions of verticals from \mathcal{P}_1 (Figure 1 and Figure 2, red arrowheads) are the sum of $X(i)$ and $X'(i)$ over all the columns in the square:

$$\sum_{i'=0}^{s-1} X(i') + \sum_{i'=s+1}^{2s} X'(i').$$

We are going to calculate this sum.

Proposition 9 (The right hand side of vertical recurrences in \mathcal{P}_1). For $c(n) = \lambda n + \eta$,

$$\sum_{i'=0}^{s-1} X(i') + \sum_{i'=s+1}^{2s} X(i') = \lambda^s (s+1)!. \quad (52)$$

First we prove the following identities:

Lemma 20.

$$\sum_{i=0}^s (-1)^{i+s} i^{s+1-t} \binom{s}{i} = \begin{cases} \frac{1}{2} s (s+1)!, & t = 0, \\ s!, & t = 1, \\ 0, & t > 1. \end{cases} \quad (53)$$

Proof. These identities are special cases of a known identity involving Stirling number of the second kind:

$$\sum_{i=0}^s (-1)^{i+s} i^p \binom{s}{i} = s! \left\{ \begin{matrix} p \\ s \end{matrix} \right\},$$

where $\left\{ \begin{matrix} p \\ s \end{matrix} \right\}$ is the Stirling number of the second kind [6, Eq. (6.19)]. Let $p = s + 1$ in above identity and use the fact that

$$\left\{ \begin{matrix} s+1 \\ s \end{matrix} \right\} = \binom{s+1}{2}$$

to obtain the first identity for $t = 0$.

Let $p = s$ and use

$$\left\{ \begin{matrix} s \\ s \end{matrix} \right\} = 1$$

to obtain the second identity for $t = 1$.

When $t > 1$, we have $s + 1 - t < s$. In this case the Stirling number of the second kind vanishes. □

With Lemma 20 we proceed to prove Proposition 9.

Proof of Proposition 9. By making a change of summation variable in the second sum $i' \rightarrow 2s - i'$, the two sum in Eq.(52) can be combined into a

single sum

$$\sum_{i'=0}^{s-1} X(i') + \sum_{i'=s+1}^{2s} X'(i') = \sum_{i'=0}^{s-1} (-1)^i \binom{s-1}{i} [c(n-i)^s + (-c(n-2s+i))^s]. \quad (54)$$

If we write $c_0 = c(n)$, then

$$\begin{aligned} c(n-i) &= c_0 - \lambda i, \\ -c(n-2s+i) &= 2\lambda s - c_0 - \lambda i. \end{aligned}$$

Expand $c(n-i)^s$ and $(-c(n-2s+i))^s$ in Eq. (54) to obtain a double summation

$$\sum_{i=0}^{s-1} (-1)^i \binom{s-1}{i} \sum_{t=0}^s \binom{s}{t} (c_0^t + (2\lambda s - c_0)^t) (-\lambda i)^{s-t}.$$

Lemma 20 shows that for the outer summation over i , the sum does not vanish only for two values of t of the inner sum, i.e., $t = 0$ and $t = 1$.

When $t = 0$, we have

$$\sum_{i=0}^{s-1} (-1)^{i+s} i^s \binom{s-1}{i} = -\frac{1}{2}(s-1)s!$$

by Lemma 20 after making the variable change from s to $s-1$.

At $t = 1$, the sum over i takes a value of $(s-1)!$. The summation of Eq. (54) is thus the sum of these two values only, multiplied by the factors in the inner sum that are independent of i . After bringing back these factors, the two terms for the sum are

$$\begin{aligned} -\frac{(s-1)s!}{2} \times 2\lambda^s, & \quad t = 0, \\ (s-1)! \times 2\lambda^s s^2, & \quad t = 1. \end{aligned}$$

Hence the summation in Eq. (52) takes the value as the sum of these two terms, which proves Proposition 9. \square

3.7.2 The right hand side from \mathcal{P}_2

For the recurrence identities in the vertical direction from the set \mathcal{P}_2 , at column i when $0 \leq i \leq s$, the contributions to the right hand side are

(Figure 3 and Figure 4, red arrowheads)

$$Y(i) = c(n-i)^s \sum_{j'=i+1}^s h(i, j'),$$

and when $s < i \leq 2s$,

$$Y'(i) = (-1)^s c(n-i)^s \sum_{j'=0}^{i-s-1} h(2s-i, s-j').$$

The total contributions of verticals from \mathcal{P}_2 are

$$\sum_{i'=0}^{s-1} Y(i') + \sum_{i'=s+1}^{2s} Y(i').$$

For the three parts of $h_{i,j}$, the first part does not contribute, since $j' > i$ in the sums of $Y(i)$ and $s - j' > 2s - i$ in the sums of $Y'(i)$. As for the binomial coefficients $b_{i,j}$ in the set \mathcal{P}_1 (Eq. (51)), we will derive analogous sums for the second and third parts of $h_{i,j}$ for each column.

For the second part of $h_{i,j}$, the column sum of the coefficients in the first quadrant is given by

Lemma 21.

$$y^{(2)}(i) = (-1)^{i+1} \binom{2s}{i} \sum_{j'=i+1}^s \binom{s+j'-i-1}{s} = (-1)^{i+1} \binom{2s}{i} \binom{2s-i}{s+1}. \quad (55)$$

Proof. The summand is Gosper-summable with antidifference shown in Lemma 26. \square

The column sum of the third part of $h_{i,j}$ involves a double sum. We have for the third part of $h_{i,j}$ in the first quadrant as

Proposition 10.

$$\begin{aligned} y^{(3)}(i) &= (-1)^i (2s-i) \binom{2s}{i} \sum_{t=0}^i \sum_{j'=i+1}^s \frac{(-1)^t}{2s-t} \binom{i}{t} \\ &= (-1)^i \frac{2s-i}{s} \binom{2s}{i} \binom{2s-i-1}{s} \left[1 - \frac{1}{2} \binom{2s-1}{s}^{-1} \right] \binom{s-t-1+j'}{j'}. \end{aligned} \quad (56)$$

Proof. The inner sum of j' in Eq. (56) is Gosper-summable as shown in Lemma 26, hence Eq. (56) can be written as

$$y^{(3)}(i) = (-1)^i (2s - i) \binom{2s}{i} \sum_{t=0}^i (-1)^t \frac{1}{2s - t} \binom{i}{t} \left[\binom{2s - t}{s - t} - \binom{s - t + i}{s - t} \right].$$

Consider the above equation as a difference between two single sum, and calculate each sum separately. For the first sum, the summand

$$F = (-1)^t \frac{1}{2s - t} \binom{i}{t} \binom{2s - t}{s - t}$$

satisfies a first order homogeneous recurrence

$$(s - i - 1)F(i, t) + (i - 2s + 1)F(i + 1, t) = 0,$$

with the certificate as

$$R = \frac{t(2s - t)}{i - t + 1}.$$

With initial value at $i = 0$ as

$$\frac{1}{2s} \binom{2s}{s},$$

the sum can be evaluated as

$$(-1)^i (2s - i) \binom{2s}{i} \sum_{t=0}^i (-1)^t \frac{1}{2s - t} \binom{i}{t} \binom{2s - t}{s - t} = \frac{1}{s} \binom{2s - i - 1}{s}. \quad (57)$$

For the second sum, the summand

$$F = (-1)^t \frac{1}{2s - t} \binom{i}{t} \binom{s - t + i}{s - t}$$

satisfies a first order homogeneous recurrence

$$-(i + 1)(i - s + 1)F(i, t) + (i + 1)(i - 2s + 1)F(i + 1, t) = 0,$$

with the certificate as

$$R = \frac{t(2s - t)(s - t + i + 1)}{i - t + 1}.$$

With initial value at $i = 0$ as

$$\frac{1}{2s},$$

the sum can be evaluated as

$$(-1)^i (2s-i) \binom{2s}{i} \sum_{t=0}^i (-1)^t \frac{1}{2s-t} \binom{i}{t} \binom{s-t+i}{s-t} = \frac{1}{2s} \binom{2s-i-1}{s} \binom{2s-1}{s}^{-1}. \quad (58)$$

Eq. (56) is the difference between Eq. (57) and Eq. (58) .

□

Putting together the results for $y^{(2)}(i)$ (Eq. (55)) and $y^{(3)}(i)$ (Eq. 56), we obtain the column sum of the vertical recurrence identities from \mathcal{P}_2 in the first quadrant and the second quadrant as

Proposition 11 (Sum of vertical identities in \mathcal{P}_2 at column i).

$$\begin{aligned} y(i) &= \sum_{j'=i+1}^s h(i, j') = (-1)^i \binom{s-1}{i} \left[\frac{1}{s} \binom{2s}{s-1} - 1 \right], \\ y'(i) &= \sum_{j'=0}^{i-s-1} h(2s-i, s-j') = (-1)^i \binom{s-1}{2s-i} \left[\frac{1}{s} \binom{2s}{s-1} - 1 \right], \end{aligned} \quad (59)$$

where the second equation is obtained from the first one by changing $i \rightarrow 2s - i$.

Proposition 12 (Total contributions to the right hand side from vertical identities in \mathcal{P}_2). For $c(n) = \lambda n + \eta$,

$$\sum_{i'=0}^{s-1} Y(i') + \sum_{i'=s+1}^{2s} Y(i') = \left[\frac{1}{s} \binom{2s}{s-1} - 1 \right] \lambda^s (s+1)!. \quad (60)$$

Proof. By comparing Eq. (59) for the set \mathcal{P}_2 with Eq. (51) for the set \mathcal{P}_1 , we see that the sum of coefficients $h_{i,j}$ for the set \mathcal{P}_2 for each column has the same factor

$$\binom{s-1}{i}$$

as that for the set \mathcal{P}_1 . This factor is dependent on i . An extra factor

$$\frac{1}{s} \binom{2s}{s-1} - 1$$

is independent of i . Hence the total contribution from the set \mathcal{P}_2 is that of the set \mathcal{P}_1 (Eq. (52)) multiplied by this factor. \square

The total contribution for the vertical identities from all the recurrence identities from the sets \mathcal{P}_1 and \mathcal{P}_2 is the sum of Eq. (52) and Eq. (60). By symmetry the total contribution from the vertical and horizontal identities is twice this number. Taking into account that LHS is also the sum of vertical and horizontal identities with a factor of 2 (the first identity of Eq. (12)), this factor of 2 cancels. Hence the total contribution for RHS is

Proposition 13 (Overall RHS).

$$RHS = \lambda^s \binom{2s}{s} s!. \quad (61)$$

For two-dimensional monomer-polymer model, $\lambda = 2$, and the main Theorem 1 is proved.

A Some properties of the two-dimensional sequence

In this Appendix we discuss some properties of the two-dimensional sequence.

First, we evaluate the double generating function of the sequence. Define

$$\mathcal{H}(s) = \sum_{i=0} H_i(x) z^i = \sum_{i=0} \sum_{j=i+1} h_{i,j} x^j z^i.$$

Proposition 14 (Double generating function). *The double generating function of the sequence $h_{i,j}$ is*

$$\mathcal{H}(s) = -\frac{(1-xz)^s}{1-z} - \frac{x(1-xz)^{2s}}{(1-x)^{s+1}} + \frac{(1-xz)^{2s}}{(1-x)^s(1-z)}. \quad (62)$$

Proof. The double generating function is obtained from the three parts of $H_i(x)$ (Eq. 5).

The first part can be obtained by exchanging the order of sums and use geometric sum and binomial theorem to obtain

$$\mathcal{H}^{(1)}(s) = -\frac{(1-xz)^s}{1-z}.$$

The second part can also be obtained by binomial theorem:

$$\mathcal{H}^{(2)}(s) = -\frac{x(1-xz)^{2s}}{(1-x)^{s+1}}.$$

The third part involves a double sum. By exchanging the order of sums, the sum over i has the following summand

$$F = (-1)^i (2s-i) \binom{2s}{i} \binom{i}{t} z^i.$$

Zeilberger's method shows that the sum $S(t)$ satisfies the following first order homogeneous recurrence

$$-(2s-t-1)zS(t) + (z-1)(t+1)S(t+1) = 0,$$

with the certificate

$$R = i - t.$$

With initial value at $t = 0$ as

$$S(0) = -2s(1-z)^{2s-1},$$

the sum is evaluated as

$$S(t) = -(2s)z^t(z-1)^{2s-t-1} \binom{2s-1}{t}.$$

Then the sum of over t can be evaluated by Zeilberger's method again with s as the parameter. For the summand

$$F = (2s)z^t(z-1)^{2s-t-1} \binom{2s-1}{t} \frac{1}{2s-t} \frac{1}{(1-x)^{s-t}},$$

the sum obeys the following first order homogeneous recurrence

$$(xz-1)^2 S(s) + (x-1)S(s+1) = 0,$$

with the certificate

$$R = \frac{t(z-1)(2xz+2zs-3+2zxs+z-4s-zxt+t)}{(2s+2-t)(2s-t+1)}.$$

With initial value at $s = 0$ as

$$S(0) = \frac{1}{1-z},$$

the sum is evaluated as

$$S(s) = \mathcal{H}^{(3)}(s) = \frac{(1-xz)^{2s}}{(1-x)^s(1-z)}.$$

□

This two-dimensional sequence has many interesting properties. We point out a few such properties, which will not be used explicitly in the proof of the main theorem.

First we point out that the sequence takes only integer values.

Corollary 1.7 (All items are integer). *All items in the sequence are integers.*

Proof. This is obvious from the double generating function Eq. (62). □

From the explicit expressions (Eq. (62) or Eq. (5) or Eq. (9)), a few simple expressions for the first few items can be obtained:

Corollary 1.8. *The general expression of the first few terms of $h_{s,i,j}$:*

$$\begin{aligned} h_{s,0,1} &= s-1, & h_{s,0,2} &= \frac{1}{2}(s+1)(s-2), & h_{s,0,3} &= \frac{1}{6}(s+1)(s+2)(s-3), \\ h_{s,1,2} &= -\frac{1}{2}s(3s-5), & h_{s,1,3} &= -\frac{1}{6}s(s+1)(5s-14), \\ h_{s,2,3} &= \frac{1}{6}s(7s^2-21s+8). \end{aligned}$$

The sum of the column is given by

Corollary 1.9 (Sum of column). *The column sum of the sequence is given by*

$$\sum_{i=0}^{j-1} h_{s,i,j} = (-1)^{j+1}(s-1) \binom{s-1}{j-1}.$$

Proof. Take limit of the double generating function at $z = 1$ to get

$$(s-1)(1-x)^{s-1}x.$$

□

As a simple corollary of Eq. (9), the sequence always vanishes at the point $(s, 0, s)$.

B Gosper and Zeilberger algorithms

In the proof we need to evaluate many sums, some are single sums, while others are double sums. Two powerful methods to evaluate sums are Gosper's algorithm and Zeilberger's algorithm.

Gosper's algorithm is a procedure to find *indefinite* sum of hypergeometric terms [5, 6, 17]. Given a hypergeometric terms t_n , the algorithm finds, if it exists, a hypergeometric term z_n , such that

$$z_{n+1} - z_n = t_n.$$

The antidifference z_n for the indefinite sum is analogous to the antiderivative for the indefinite integration. Gosper's algorithm gives a definitive answer to the question of whether a given hypergeometric term can be indefinitely summed. If the answer is confirmative, we call the summand Gosper-summable.

Many summands, however, are not Gosper-summable. Zeilberger's algorithm, or the method of *creative telescoping* [20, 6, 17], makes many more summands become summable. The method is an extension of Gosper's algorithm. It shows that for the sum

$$f(n) = \sum_{k=u}^v F(n, k), \quad (63)$$

it is always possible to find a recurrence

$$\sum_{j=0}^J b_j(n) F(n+j, k) = G(n, k+1) - G(n, k), \quad (64)$$

in which the coefficients $\{b_j(n)\}_0^J$ are polynomials in n , and $R(n, k) = G(n, k)/F(n, k)$ is a rational function of n and k . In the proof we list $R(n, k)$ for each sum, which can be used to check the correctness of the recurrence relation of the summand.

Since the coefficients $b_j(n)$ on the left side of the recurrence are independent of the summation index k , Eq. (64) can be summed from u (the lower bound of the summation index k) to v (the upper bound) to obtain the sum $f(n)$:

$$\sum_{j=0}^J b_j(n) f(n+j) = G(n, u+1) - G(n, v),$$

where the right side telescopes. For the *standard boundary conditions* [19], where the sum includes all nonzero values of the summand (and the support lies inside a larger set in which the summand remains well defined and vanishes), the right hand side of the above equation vanishes, so the equation gives a homogeneous recurrence relation for the sum. For nonstandard boundary conditions, however, the right hand side may not vanish, leading to an inhomogeneous recurrence for the sum. For the sums in our case, they most often have nonstandard boundary conditions due to the geometric restrictions imposed on the lower and upper boundaries, hence they often have inhomogeneous recurrence relations.

The algorithm is extended to multi-variable multi-sum summations [19, 1]. For a double sum with summation indices i and j ,

$$f(n) = \sum_{i=u_i}^{v_i} \sum_{j=u_j}^{v_j} F(n, i, j), \quad (65)$$

the recurrence of Eq. (64) is in the form of

$$\sum_{\ell=0}^J b_\ell(n) F(n + \ell, i, j) = \Delta_i G_i(n, i, j) + \Delta_j G_j(n, i, j),$$

where Δ_i and Δ_j are the forward difference operators on i and j , respectively.

Just as the single sums mentioned above, the double sums in our proof often have nonstandard boundary conditions. As the authors commented in [19], under nonstandard boundary conditions, “the boundary terms ‘snowball’ so the application would become rapidly more complicated.”

Whether summation indices are dependent of each other also affects the recurrence relation of the sum. Below we list a few such scenarios for the summation indices of double sums. In each case the inhomogeneous term is a sum of two sums with a single variable. Here we use u_ℓ and v_ℓ for the lower and upper boundaries, respectively, with $\ell = i$ or j for the summation indices. They are assumed to be independent of the summation parameter n . The proofs can be obtained by direct calculations.

Lemma 22. *For the double sum with independent lower and upper boundaries,*

$$\sum_{i=u_i}^{v_i} \sum_{j=u_j}^{v_j} F(n, i, j),$$

the sum satisfies an inhomogeneous recurrence relations with the inhomogeneous term as

$$\begin{aligned} & \sum_{i=u_i}^{v_i} \sum_{j=u_j}^{v_j} (\Delta_i G_i(n, i, j) + \Delta_j G_j(n, i, j)) = \\ & \sum_{i=u_i}^{v_i} (G_j(n, i, v_j + 1) - G_j(n, i, u_j)) + \sum_{j=u_j}^{v_j} (G_i(n, v_i + 1, j) - G_i(n, u_i, j)). \end{aligned}$$

Lemma 23. For the double sum with inner sum's upper boundary as the summation index of the outer sum,

$$\sum_{i=u_i}^{v_i} \sum_{j=u_j}^i F(n, i, j),$$

the sum satisfies an inhomogeneous recurrence relations with the inhomogeneous term as

$$\begin{aligned} & \sum_{i=u_i}^{v_i} \sum_{j=u_j}^i (\Delta_i G_i(n, i, j) + \Delta_j G_j(n, i, j)) = \\ & \sum_{i=u_j}^{v_i} (G_j(n, i, i + 1) - G_j(n, i, v_j)) + \sum_{j=u_j}^{v_i} (G_i(n, v_i + 1, j) - G_i(n, j, j)). \end{aligned}$$

Lemma 24. For the double sum

$$\sum_{i=u_i}^{v_i} \sum_{j=u_j}^{v_i-i} F(n, i, j),$$

the sum satisfies an inhomogeneous recurrence relations with the inhomogeneous term as

$$\begin{aligned} & \sum_{i=u_i}^{v_i} \sum_{j=u_j}^i (\Delta_i G_i(n, i, j) + \Delta_j G_j(n, i, j)) = \\ & \sum_{i=u_i}^{v_i} (G_j(n, i, v_i + 1 - i) - G_j(n, i, u_j)) + \sum_{j=u_j}^{v_i} (G_i(n, v_i + 1 - j, j) - G_i(n, u_i, j)). \end{aligned}$$

Remark 4. *The choice of summation parameter also affects the recurrence obtained by the Zeilberger's method. If one of the running indices of the double sum Eq. (65) is used as a parameter (u_ℓ or v_ℓ), the inhomogeneous term needs to include additional single sums for the boundary terms.*

Remark 5. *Note that in the proof we use F and S to denote the generic summand and sum under discussion.*

C Some binomial coefficient identities

One of the binomial coefficient identities that is used several times in the proof is the well-known Chu-Vandermonde identity.

$$\sum_j \binom{a}{j} \binom{b}{c-j} = \binom{a+b}{c}. \quad (66)$$

Another binomial coefficient identity used in the proof of the result in the first quadrant is listed below.

Lemma 25.

$$\sum_{j'=0}^j (-1)^{j'} \binom{s-t+j'}{j'} \binom{s}{j-j'} = (-1)^j \binom{j-t}{j}. \quad (67)$$

Proof. Rearrange the binomials in the summand,

$$\sum_{j'=0}^j (-1)^{j'} \binom{s-t+j'}{j'} \binom{s}{j-j'} = \frac{s!(j-t)!}{(s-t)!j!} \sum_{j'=0}^j (-1)^{j'} \binom{j}{j'} \binom{s-t+k}{j-t}.$$

By using identity

$$\sum_{k=0}^n \binom{n}{k} \binom{a+k}{m} = (-1)^n \binom{a}{m-n}, \quad (68)$$

Eq. (67) can be obtained. Eq. (68) can be found in many references, such as Prudnikov 4.2.5. Eq. (55) or Knuth 1.2.6. Eq. (23). \square

D Some sums used in section 3.7 that are Gosper-summable

Below we list some summands that appear several times in section 3.7 that are Gosper-summable.

Lemma 26. *For the summand*

$$F = \binom{s + b + j'}{j'},$$

with j' as the index of summation, the antidifference is

$$\binom{s + b + j'}{j' - 1}.$$

Lemma 27. *For the summand*

$$F = (-1)^{j'} \binom{s}{j'},$$

with j' as the index of summation, the antidifference is

$$(-1)^{j'} \binom{s - 1}{j' - 1}.$$

References

- [1] M. Apagodu and D. Zeilberger. Multi-variable zeilberger and almkvist–zeilberger algorithms and the sharpening of wilf–zeilberger theory. *Adv. Appl. Math.*, 37(2):139–152, Aug. 2006.
- [2] D. Dhar and R. Rajesh. Entropy of fully packed hard rigid rods on d-dimensional hypercubic lattices. *Phys. Rev. E*, 103(4), 2021.
- [3] M. E. Fisher. On the dimer solution of planar Ising models. *J. Math. Phys.*, 7(10):1776–1781, 1966.
- [4] A. Ghosh, D. Dhar, and J. L. Jacobsen. Random trimer tilings. *Phys Rev E Stat Nonlin Soft Matter Phys*, 75(1 Pt 1):011115, 2007.

- [5] R. W. Gosper. Decision procedure for indefinite hypergeometric summation. *Proc. National Acad. Sci.*, 75(1):40–42, Jan. 1978.
- [6] R. Graham, D. Knuth, and O. Patashnik. *Concrete Mathematics : A Foundation For Computer Science, Second Edition*. Addison-Wesley Professional, 2nd edition edition, 1994.
- [7] M. Jerrum. Two-dimensional monomer dimer systems are computationally intractable. *J. Stat. Phys.*, 48(1-2):121–134, 1987. Erratum: 59, 1087-1088 (1990).
- [8] P. W. Kasteleyn. Dimer statistics and phase transitions. *J. Math. Phys.*, 4(2):287–293, 1963.
- [9] Y. Kong. Logarithmic corrections in the free energy of monomer-dimer model on plane lattices with free boundaries. *Phys. Rev. E*, 74:011102, Jul 2006.
- [10] Y. Kong. Monomer-dimer model in two-dimensional rectangular lattices with fixed dimer density. *Phys. Rev. E*, 74:061102, Dec 2006.
- [11] Y. Kong. Packing dimers on $(2p + 1) \times (2q + 1)$ lattices. *Phys. Rev. E*, 73:016106, Jan 2006.
- [12] Y. Kong. Asymptotics of the monomer-dimer model on two-dimensional semi-infinite lattices. *Phys. Rev. E*, 75:051123, May 2007.
- [13] Y. Kong. Recurrence solution of monomer-polymer models on two-dimensional rectangular lattices. *Phys. Rev. E*, 2024. accepted.
- [14] B. M. McCoy and T. T. Wu. *The Two-Dimensional Ising Model*. Harvard University Press, Cambridge, Massachusetts, 1973.
- [15] L. Onsager. Crystal statistics. I. A two-dimensional model with an order-disorder transition. *Phys. Rev.*, 65(3/4):117–149, 1944.
- [16] L. Onsager. The effects of shape on the interaction of colloidal particles. *Ann. New York Acad. Sci.*, 51(4):627–659, 1949.
- [17] M. Petkovšek, H. Wilf, and D. Zeilberger. *A=B*. AK Peters, Ltd., Wellesley, MA, USA, 1996.

- [18] L. R. Rodrigues, J. F. Stilck, and W. G. Dantas. Entropy of rigid k-mers on a square lattice. *Phys. Rev. E*, 107(1), 2023.
- [19] H. S. Wilf and D. Zeilberger. An algorithmic proof theory for hypergeometric (ordinary and “ q ”) multisum/integral identities. *Invent. Math.*, 108(3):575–633, June 1992.
- [20] D. Zeilberger. The Method of Creative Telescoping. *J. Symb. Comput.*, 11(3):195–204, Mar. 1991.