

Chance-Constrained Set Multicover Problem

Shunyu Yao, Neng Fan*, Pavlo Krokhmal

Department of Systems and Industrial Engineering, University of Arizona
Tucson, AZ 85721, USA

Abstract

We consider a variant of the set covering problem with uncertain parameters, which we refer to as the chance-constrained set multicover problem (CC-SMCP). In this problem, we assume that there is uncertainty regarding whether a selected set can cover an item, and the objective is to determine a minimum-cost combination of sets that covers each item i at least k_i times with a prescribed probability. To tackle CC-SMCP, we employ techniques of enumerative combinatorics, discrete probability distributions, and combinatorial optimization to derive exact equivalent deterministic reformulations that feature a hierarchy of bounds, and develop the corresponding outer-approximation (OA) algorithm. Additionally, we consider reducing the number of chance constraints via vector dominance relations and reformulate two special cases of CC-SMCP using the “log-transformation” method and binomial distribution properties. Theoretical results on sampling-based methods, i.e., the sample average approximation (SAA) method and the importance sampling (IS) method, are also studied to approximate the true optimal value of CC-SMCP under a finite discrete probability space. Our numerical experiments demonstrate the effectiveness of the proposed OA method, particularly in scenarios with sparse probability matrices, outperforming sampling-based approaches in most cases and validating the practical applicability of our solution approaches.

Keywords: Set Multicover Problem, Chance-Constrained Optimization, Integer Linear Programming

1 Introduction

We consider the following *chance-constrained set multicover problem* (CC-SMCP):

$$\text{[CC-SMCP]} \quad \min_x \sum_{j \in [n]} c_j x_j \quad (1a)$$

$$\text{s.t.} \quad \mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{ij} x_j \geq k_i \right] \geq 1 - \epsilon_i, \quad \forall i \in [m], \quad (1b)$$

$$x \in B \subseteq \{0, 1\}^n \quad (1c)$$

*Corresponding Author: nfan@arizona.edu

where there are m items, indexed by the set $[m] := \{1, \dots, m\}$, and n subsets of these m items, indexed by the set $[n] := \{1, \dots, n\}$. Here $c \in \mathbb{R}^n$ is the vector of weights, or costs of the subsets in $[n]$, $k_i \in \mathbb{Z}_+$, $\tilde{a}_{ij} \in \{0, 1\}$ is a Bernoulli random variable indicating whether item $i \in [m]$ belongs to set $j \in [n]$ or not, and $\epsilon_i \in (0, 1)$ is a prespecified allowed failure probability for item i . The set B represents some other deterministic side constraints on the decision variables x . For instance, in the facility location problem, B can represent constraints requiring that demands at several points must be serviced by the established facilities, or it can be the budget restriction on the number of facilities to be located such as $\{x \in \{0, 1\}^n : \sum_{j \in [n]} x_j \leq U\}$ where at most U facilities can be located.

This problem aims to determine a subset $S \subseteq [n]$ that covers each element $i \in [m]$ at least k_i times with high probability while minimizing the total cost/weight of covering. Note that when $k_i = 1$ for all $i \in [m]$, the problem becomes probabilistic set covering (PSC) problem.

Our problem constitutes a variant of the set covering problem (SCP), a well-known combinatorial optimization problem which serves as a model to a variety of real-world applications (Hall and Hochbaum 1992), including marketing (Özener et al. 2013), security (Lessin et al. 2018), telecommunications (Grötschel et al. 1992), scheduling (Smith and Wren 1988), production planning (Vasko et al. 1989), facility location (Gunawardane 1982, Huang et al. 2010), and vehicle routing (Bramel and Simchi-Levi 1997, 2002), etc. Although many of these applications can be modeled as SCPs, for reliability/backup purposes they are often treated as set multicover problems. As an example of set multicover problems, consider the problem of determining the locations of emergency service facilities, where we aim to establish fire stations across n specific locations to cover m target regions (Hall and Hochbaum 1992). A higher level of service is provided when there are multiple fire stations within a short distance of a building at risk, as opposed to having only one nearby. The objective is to identify optimal locations for fire stations to guarantee that all target regions can receive the necessary level of service, while simultaneously minimizing overall costs. In this context, the connection parameter a_{ij} indicates whether the service provided by candidate location j can cover target i and it may depend on their distances, e.g., $a_{ij} = 1$ if target i is near location j and $a_{ij} = 0$ otherwise.

In practical emergency scenarios (e.g., natural disasters), it is often the case that the connection between target i and open facility j may be randomly disrupted. In that case, it is natural to model a_{ij} as a Bernoulli random variable, denoted by \tilde{a}_{ij} . The probability $P[\tilde{a}_{ij} = 1] := p_{ij}$ can be interpreted as the likelihood of successfully reaching target i from the location j within a limited amount of time in an emergency. Therefore, intuitively, the goal is to identify optimal locations for fire stations to achieve the required level of service of all targets with a high probability. From the perspective of stochastic optimization, this problem can be formulated as a chance-constrained programming model when we incorporate uncertainty as previously described. In addition to the aforementioned application of CC-SMCP, Beraldi and Ruszczyński (2002) considered a game to select arcs on a graph to cover adversaries' path with a high probability. Ahmed and Papageorgiou (2013) presented an example to cover m targets by placing sensors at n potential sensor sites. Shen and Jiang (2022) proposed another emergency facility location example of chance-constrained SCP. These chance-constrained SCP models can be easily generalized to incorporate backup coverage, i.e., define a parameter k_i to reflect the coverage level of target i .

As a powerful paradigm to model optimization problems with uncertainty, chance-constrained programming arises in a wide variety of applications such as finance (Lemus Rodriguez 1999), healthcare (Tanner and Ntaimo 2010), power systems (Van Ackooij et al. 2011), transportation (Dinh et al. 2018), network design (Wang 2007) and wireless communications (Soltani et al. 2013), etc. From the computational perspective, chance-constrained programs are very challenging to solve primarily for two reasons (Küçükyavuz and Jiang 2022). First, given a candidate solution x , it can be computationally demanding to check whether x is feasible or not (Ahmed and Papageorgiou 2013); and, second, the feasible region characterized by chance constraints is nonconvex and even disconnected in general (Ahmed and Papageorgiou 2013, Nemirovski and Shapiro 2007). There are two main approximation techniques to mitigate the aforementioned challenges: convex conservative approximation (Ben-Tal et al. 2009,

Nemirovski and Shapiro 2007) and sample average approximation (SAA) (Calafiore and Campi 2005, Kleywegt et al. 2002, Luedtke and Ahmed 2008), both of which can efficiently identify feasible solutions with a guarantee of performance. The strategy of the former is to formulate a convex optimization problem and produce a feasible solution with a high probability. One drawback of this approach is that it becomes challenging when decision variables are discrete. The latter is to solve an approximation problem based on an independent Monte Carlo sample of random data (Kleywegt et al. 2002). This approach may not require knowledge of the distribution of the random parameters, however, it may need drawing a large number samples from the true distribution to obtain feasible solutions with a guarantee. For other specific relaxations and approximations for chance-constrained problems (such as finite scenario approximation, CVaR approximation, Bonferroni approximation, etc), we refer the reader to (Ahmed and Xie 2018, Lejeune and Prékopa 2018).

In many problems of interest, the decision vector x is binary (as is the case in the present paper) and this special structure can be exploited to derive stronger formulations and specialized algorithms. We refer to such chance-constrained problems with purely binary variables as *chance-constrained combinatorial optimization* problems (Küçükyavuz and Jiang 2022). Luedtke (2014) proposed a branch-and-cut decomposition algorithm for finding exact solutions of chance-constrained problems having discrete distributions with finite support. For chance-constrained bin packing problems, Song et al. (2014) proposed an efficient coefficient strengthening method and lifted probabilistic cover inequalities for chance-constrained bin packing problems. Later, Wang et al. (2021) studied bilinear formulation of chance-constrained multiple bin packing problem and used the lifting techniques to identify cover, clique, and projection inequalities to strengthen the bilinear formulation. Recently, Zhang et al. (2020) proposed Dantzig–Wolfe formulations suited to a branch-and-price (B&P) algorithm to solve two versions of the chance-constrained stochastic bin packing (CCSBP) problem. For chance-constrained assignment problems, Wang et al. (2022) considered a chance-constrained assignment problem to develop valid inequalities and derived lifted cover inequalities based on a bilinear reformulation of this problem. For chance-constrained dominating set problems, Sun and Fan (2019) investigated the reliable connected power dominating set problem that met two requirements including the connectivity of the phasor measurement unit (PMU) subgraph and the reliability of its connectivity. For chance-constrained traveling salesman problems (TSPs), Padberg and Rinaldi (1989) proposed a branch-and-cut approach to a TSP with side constraints, and Campbell and Thomas (2008) presented two recourse models and a chance-constrained model for probabilistic TSP with deadlines. For chance-constrained knapsack problems, Yoda and Prékopa (2016) determined sufficient conditions for the convexity of the formulation under different discrete distributions, and several approximate but more tractable formulations that could provide near-optimal solutions were derived in (De 2017, Han et al. 2016, Klopfenstein and Nace 2008). For more references on chance-constrained combinatorial problems and approaches, we refer the reader to Section 2 of survey (Küçükyavuz and Jiang 2022).

In the context of chance-constrained SCP models, prior studies have examined two types of uncertainty: right-hand-side (RHS) uncertainty and left-hand-side (LHS) uncertainty. For RHS uncertainty, Beraldi and Ruszczyński (2002) developed a specialized branch-and-bound algorithm based on the enumeration of p -efficient points. Later, Saxena et al. (2010) derived the polarity cuts to obtain a stronger formulation and improve the computational performance of this enumeration approach. For LHS uncertainty (as in this paper), Fischetti and Monaci (2012) developed cutting plane approaches to the individual chance-constrained problem where all components of the Bernoulli random vector are independent. Further, Wu and Küçükyavuz (2019) proposed an exact approach to the probabilistic partial SCP where there existed an efficient probability oracle to retrieve the probability of any events under the true distribution. For distributionally robust chance-constrained SCPs, Ahmed and Papageorgiou (2013) studied individual chance constraints under moment-based ambiguity sets, and recently, Shen and Jiang (2022) explored joint chance constraints with Wasserstein ambiguity.

As we can see, while there are several works in the literature related to chance-constrained SCPs, it appears that prior studies have paid less attention to chance-constrained set multicover problems, which

represent a natural extension of SCPs.

In this paper, we focus on individual chance constraints with LHS uncertainty in set multicover problems, where both decision and random variables are purely binary. Our contributions may be summarized as follows:

1. Using techniques of enumerative combinatorics, discrete probability distributions, and combinatorial optimization, we derive exact deterministic reformulations of CC-SCMP that are based on hierarchies of bounds, which are in turn utilized in construction of an outer-approximation (OA) algorithm for CC-SMCP.
2. We investigate reducing the number of chance constraints by considering vector dominance relations defined in an appropriate partially ordered set, which is a part of the presolving method.
3. We consider reformulations for two special cases of CC-SMCP by employing the “log-transformation” method and utilizing properties of the binomial distribution, respectively.
4. We present some theoretical results on SAA method to approximate the true optimal value of CC-SMCP under a finite discrete probability space. We also studied the importance sampling (IS) method and obtained a sufficient condition for selecting the optimal IS estimator.
5. We conduct several numerical experiments to validate the effectiveness of our OA method compared to the sampling-based approaches. The numerical results illustrate that our OA method can solve CC-SMCP effectively with a sparse probability matrix.

The rest of this paper is organized as follows. Section 2 develops an exact deterministic reformulation for CC-SMCP and proposes an outer-approximation algorithm to address CC-SMCP. Some presolving methods such as reducing the number of chance constraints are introduced as well. Reformulations for two special cases of CC-SMCP are investigated in Section 3. Section 4 studies two sampling-based methods to approximate the optimal value of CC-SMCP. Section 5 presents some computational experiments to evaluate the effectiveness of the proposed model and solution approaches. Finally, Section 6 concludes the paper.

2 Deterministic Reformulations and Solution Approaches

2.1 Equivalent reformulation

In this section, we consider a deterministic reformulation of CC-SMCP using some combinatorial methods. For ease of exposition, we call the probability $\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_{ij}x_j \geq k_i\right]$ the *cover probability* of the item i . Let X_i be the set of all x that satisfy the corresponding cover constraint for the i th item, i.e.,

$$X_i = \left\{ x \in \{0, 1\}^n : \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_{ij}x_j \geq k_i\right] \geq 1 - \epsilon_i \right\},$$

and thus the problem (1) can be expressed as

$$\min \left\{ \sum_{j \in [n]} c_j x_j : x \in X_i, i = 1, \dots, m; \quad x \in B \subseteq \{0, 1\}^n \right\}. \quad (2)$$

We call the set X_i the *probabilistic covering set* of the item i . We use i to index the item and j to index the set throughout the paper. In what follows, for the sake of simplicity, we ignore the index i and study

the reformulation of X_i (simplified as X) so that an equivalent deterministic reformulation of the form (2) can be obtained for CC-SMCP.

We assume that \tilde{a}_j 's are independent random variables throughout the paper (note that a_j 's are not necessarily identically distributed). Let S be a subset of $[n]$, and $S^c := [n] \setminus S$, and define the events $A_j(x) := \{\tilde{a}_j x_j = 1\}$ and $A_j^c(x) := \{\tilde{a}_j x_j = 0\}$. We also define $A_S(x) := \bigcap_{j \in S} A_j(x)$ and $A_S^c(x) := [\bigcap_{j \in S} A_j(x)]^c$. A closed-form expression for the cover probability $\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \geq k \right]$ can be developed using the inclusion-exclusion principle, stated as the following lemma:

Lemma 1 *Let \tilde{a}_j be independent Bernoulli random variables with $\mathbb{P}(\tilde{a}_j = 1) = p_j$ for each $j \in [n]$, x_j be binary variables, and $k \in \mathbb{Z}_+$. Then*

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \geq k \right] = \sum_{\ell=k}^n (-1)^{\ell-k} \binom{\ell-1}{\ell-k} h_\ell(x), \quad (3)$$

where $h_\ell(x) = \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \mathbb{P}[A_S(x)] = \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} x_j p_j$, $\ell = k, \dots, n$.

Thus, based on Lemma 1, the probabilistic covering set X can be reformulated into the following deterministic 0-1 nonlinear set

$$X = \left\{ x \in \{0, 1\}^n : \sum_{\ell=k}^n (-1)^{\ell-k} \binom{\ell-1}{\ell-k} \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} x_j p_j \geq 1 - \epsilon \right\}. \quad (4)$$

Remark 1 (Relationship to the Poisson binomial distribution) *If we define $Z := \sum_{j=1}^n \tilde{a}_j x_j$, then Z is a random variable that follows Poisson binomial distribution with a collection of n independent yes/no experiments with different success probabilities $x_1 p_1, x_2 p_2, \dots, x_n p_n$. Actually, from the proof of Lemma 1, we can obtain an explicit expression for its probability mass function.*

Remark 2 (The probability of the complement of the covering event) *From the proof of Lemma 1, we can also obtain the probability*

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \leq k - 1 \right] = \sum_{d=0}^{k-1} \sum_{\ell=d}^n (-1)^{\ell-d} \binom{\ell}{d} h_\ell(x).$$

In Section 2.2, we will explore some lower and upper bounds for this probability and present a outer-approximation algorithm that utilizes these findings.

For more references about the inclusion–exclusion principle and combinatorial identities, we refer the reader to Section A.1 of the appendix and (Prudnikov et al. 1998, Stanley 2011).

Now we consider the linearization of a general polynomial term:

$$y = \prod_{j \in S} x_j, \quad (5)$$

where $S \subseteq [n]$ and $x_j \in \{0, 1\}, j \in S$. Since x_j are binary variables, y is also a binary variable. Typically, there are two commonly used approaches to linearize the cross product (5). The first approach establishes the equivalence of the nonlinear equation (5) to two linear inequalities.

Lemma 2 (Glover and Woolsey (1974)) *Let $s = |S|$. Equality (5) holds if and only if*

$$\sum_{j \in S} x_j - y \leq s - 1, \quad -\sum_{j \in S} x_j + sy \leq 0, \quad x_j \in \{0, 1\}, j \in S, \quad y \in \{0, 1\}. \quad (6)$$

If we allow the use of more constraints in exchange for relaxing the binary variable y in (6), then we obtain the so-called *standard linearization* (Glover and Woolsey 1974):

$$y = \prod_{j \in S} x_j \Leftrightarrow \left\{ \sum_{j \in S} x_j - y \leq s - 1, \quad y \leq x_j \quad \forall j \in S, \quad x_j \in \{0, 1\} \quad \forall j \in S, \quad y \geq 0 \right\}. \quad (7)$$

Indeed, from the experimental results reported below in Section 5, it can be suggested that (7) may be preferable to (6) in spite of the increased number of constraints.

To obtain a linearization of X given by (4), let T_ℓ be a set of ℓ indices (out of n indices), $\binom{[n]}{\ell}$ be a collection of all sets of ℓ indices such that $\left| \binom{[n]}{\ell} \right| = \binom{n}{\ell}$, and let

$$y_{T_\ell} = \prod_{j \in T_\ell} x_j, \quad T_\ell \in \binom{[n]}{\ell}, \quad \ell = k, \dots, n. \quad (8)$$

Taking the first linearization approach as an example (combining Lemma 1 and 2), i.e., substituting (8) into equation (3) and adding constraints as defined in system (6), we obtain the following result:

Theorem 1 *The chance-constrained set*

$$X = \left\{ x \in \{0, 1\}^n : \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \geq k \right] \geq 1 - \epsilon \right\}$$

admits the following linearized deterministic reformulation:

$$\sum_{\ell=k}^n (-1)^{\ell-k} \binom{\ell-1}{\ell-k} \sum_{T_\ell \in \binom{[n]}{\ell}} \left(\prod_{j \in T_\ell} p_j \right) y_{T_\ell} \geq 1 - \epsilon \quad (9a)$$

$$\sum_{j \in T_\ell} x_j - y_{T_\ell} \leq \ell - 1, \quad \forall T_\ell \in \binom{[n]}{\ell}, \quad \ell = k, \dots, n \quad (9b)$$

$$- \sum_{j \in T_\ell} x_j + \ell y_{T_\ell} \leq 0, \quad \forall T_\ell \in \binom{[n]}{\ell}, \quad \ell = k, \dots, n \quad (9c)$$

$$x_j \in \{0, 1\}, \quad \forall j \in [n]; \quad y_{T_\ell} \in \{0, 1\} \quad \forall T_\ell \in \binom{[n]}{\ell}, \quad \ell = k, \dots, n \quad (9d)$$

For small values of n or when k is very close to n , the original CC-SMCP problem (2) can be directly solved using the aforementioned linearization techniques (an equivalent BIP). However, the above reformulation (9) has some significant drawbacks. First, the number of variables and constraints included in this reformulation may have exponential growth as the number of sets increases, making it very time-consuming to compute feasible solutions even for moderately sized problems. Second, the product of probabilities can be very small, and the summation of these products may lead to a substantial *cumulative rounding error* (also known as *accumulation of roundoff error*). Third, an optimal solution often assigns a value of one to numerous decision variables, making the traditional column generation or branch-and-price approach unsuitable for this reformulation. Next, we will discuss several ways to alleviate these computational challenges. We first propose an outer-approximation algorithm in Section 2.2, and discuss how to reduce the number of chance-constraints by using vector domination relations in Section 2.3. Then we investigate two special cases of CC-SMCP in Section 3, which allow for additional simplifications.

2.2 Outer approximations

In this section, we propose an outer-approximation (OA) algorithm for CC-SMCP with small k . We first notice that the probabilistic covering set X is equivalent to

$$X = \left\{ x \in \{0, 1\}^n : \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \leq k - 1 \right] \leq \epsilon \right\},$$

which can be further reformulated into the following 0-1 nonlinear set (as mentioned in Remark 2)

$$X = \left\{ x \in \{0, 1\}^n : \sum_{d=0}^{k-1} \sum_{\ell=d}^n (-1)^{\ell-d} \binom{\ell}{d} \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} x_j p_j \leq \epsilon \right\}.$$

Next we derive some lower and upper bounds for the probability $\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \leq k - 1 \right]$.

Theorem 2 *Let \tilde{a}_j be independent random variables with $\mathbb{P}(\tilde{a}_j = 1) = p_j$ for each $j \in [n]$, x_j 's be binary variables, and $k \in \mathbb{Z}_+$. Then*

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \leq k - 1 \right] = \begin{cases} g_n(x) \leq g_t(x), & \text{if } t \text{ is even} \\ g_n(x) \geq g_t(x), & \text{if } t \text{ is odd} \end{cases} \quad (10)$$

where

$$g_t(x) := \sum_{d=0}^{k-1} \sum_{\ell=d}^{t+d} (-1)^{\ell-d} \binom{\ell}{d} \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} x_j p_j, \quad t = 0, \dots, n.$$

Moreover, if for any fixed $S \subseteq [n]$ with $|S| = d \leq \ell < n$, we have

$$\sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} \mathbb{P}[A_T(x)] \geq \sum_{\substack{S \subseteq T' \subseteq [n] \\ |S|=d, |T'|=\ell+1}} \mathbb{P}[A_{T'}(x)], \quad (11)$$


then the following hierarchy of inequalities holds:

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \leq k - 1 \right] = g_n(x) \leq \dots \leq g_4(x) \leq g_2(x) \leq g_0(x), \quad (12)$$


$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j x_j \leq k - 1 \right] = g_n(x) \geq \dots \geq g_5(x) \geq g_3(x) \geq g_1(x). \quad (13)$$

Figure 1 illustrates how upper and lower bounds converge to the probability $\mathbb{P}[\sum_{j=1}^n \tilde{a}_j \leq k]$ as t increases. In these experiments, we set $n = 20$, $k = 3$ and let $x_j = 1$ for each $j \in [n]$. Specifically, in Figure 1(a), we assume that random variables \tilde{a}_j are i.i.d. with Bernoulli distribution with $\mathbb{P}(\tilde{a}_j = 1) = 0.15$, for each $j \in [n]$. In Figure 1(b), we assume that random variables follow different Bernoulli distributions, such that $\mathbb{P}(\tilde{a}_j = 1) = 0.1, j \in \{1, \dots, 7\}$, $\mathbb{P}(\tilde{a}_j = 1) = 0.2, j \in \{8, \dots, 13\}$, $\mathbb{P}(\tilde{a}_j = 1) = 0.3, j \in \{14, \dots, 17\}$ and $\mathbb{P}(\tilde{a}_j = 1) = 0.5, j \in \{18, \dots, 20\}$. As can be seen in these two figures, the upper and lower bounds converge rapidly to the final probability, within about 4 – 5 iterations.

Note that condition (11) that guarantees monotonicity of bounds for all values of t is rather strong; as seen in Figure 1, the sequence $g_t(x)$ does not exhibit monotonic behavior for smaller values of t . However,



(a) $n = 20, k = 3, \mathbb{P}(\tilde{a}_j^t = 1) = 0.15, \forall j \in [n]$.



(b) $n = 20, k = 3, \mathbb{P}(\tilde{a}_j^t = 1) \in \{.1, .2, .3, .5\}$

Figure 1: Convergence of bounds for the cover probability $\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j \leq k\right]$.

it can be demonstrated that, after a sufficient number of iterations, the sequence $g_t(x)$ will eventually become monotone increasing or decreasing for odd and even t , respectively.

We need additional notations in the following. Let $g_t(x) = \sum_{d=0}^{k-1} g_{t,d}(x)$, and $g_{t,d}(x) = \sum_{\ell=d}^{t+d} (-1)^{\ell-d} \binom{\ell}{d} \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} x_j p_j$.

From the proof of Theorem 2, for any $0 \leq d \leq k-1$, we have

$$\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j = d\right] = \begin{cases} g_{n,d}(x) \leq g_{t,d}(x), & \text{if } t \text{ is even} \\ g_{n,d}(x) \geq g_{t,d}(x), & \text{if } t \text{ is odd} \end{cases} \quad (14)$$

Lemma 3 Given d where $0 \leq d \leq k-1$, if for some odd t , $g_{t,d}(x) \leq g_{t+2,d}(x)$, then $g_{\tau,d}(x) \leq g_{\tau+2,d}(x)$, for all odd τ such that $t \leq \tau \leq n-2$; Similarly, if for some even t , $g_{t,d}(x) \geq g_{t+2,d}(x)$, then $g_{\tau,d}(x) \geq g_{\tau+2,d}(x)$, for all even τ such that $t \leq \tau \leq n-2$.

Theorem 3 Let $t_{\min}^{\text{odd}} := \min\{t \mid t \text{ is odd}, 1 \leq t \leq n-2, g_{t,d}(x) \leq g_{t+2,d}(x), 0 \leq d \leq k-1\}$. Then $g_{\tau}(x) \leq g_{\tau+2}(x)$ for all odd τ such that $t_{\min}^{\text{odd}} \leq \tau \leq n-2$; Similarly, let $t_{\min}^{\text{even}} := \min\{t \mid t \text{ is even}, 0 \leq t \leq n-2, g_{t,d}(x) \geq g_{t+2,d}(x), 0 \leq d \leq k-1\}$. Then $g_{\tau}(x) \geq g_{\tau+2}(x)$, for all even τ such that $t_{\min}^{\text{even}} \leq \tau \leq n-2$.

One of the popular methods of global optimization consists in approximating the given problem by a sequence of easier problems, such that the solutions of the approximating problems converge to an optimal solution of the given problem. If the feasible sets of the approximating problems contain the feasible set of the original problem, this method is known as *outer approximation*. According to the above theorem, we can propose an outer-approximation algorithm by using the lower bounds to solve CC-SMCP to the optimality.

Let $g_{t_i}^{(i)}(x) := \sum_{d=0}^{k-1} \sum_{\ell=d}^{t_i+d} (-1)^{\ell-d} \binom{\ell}{d} \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} x_{ij} p_{ij}$ and initialize $t_i := 1$ for each $i \in [m]$. We start by solving the following relaxed problem:

$$\nu := \min \left\{ \sum_{j \in [n]} c_j x_j : g_{t_i}^{(i)}(x) \leq \epsilon_i, \forall i \in [m], x \in B \subseteq \{0, 1\}^n \right\}. \quad (15)$$

Clearly, ν provides a lower bound for the optimal value of problem (2) by Theorem 2. Let \bar{x} be an optimal solution to problem (15), and check whether $\bar{x} \in X_i$ for every $i \in [m]$ by using Lemma 1. If $\bar{x} \in X_i$ for each $i \in [m]$, then \bar{x} is also an optimal solution to the chance-constrained problem (2). Otherwise, we have $\bar{x} \notin X_i$ for some $i \in [m]$, replace t_i by $t_i + 2$ in $g_{t_i}^{(i)}(x)$ and solve an updated relaxed problem (15),

obtaining an updated solution \bar{x} . The process is repeated until $\bar{x} \in X_i$ for each $i \in [m]$. Algorithm 1 summarizes this outer-approximation method for solving CC-SMCP.

Algorithm 1: Outer Approximation Algorithm

- 1 Initialize $t_i := 1$ for each $i \in [m]$;
 - 2 Solve the relaxed problem (15) to obtain a solution \bar{x} and a lower bound ν for the original problem (2);
 - 3 **if** $\bar{x} \in X_i$ for each $i \in [m]$ **then**
 - 4 | **return** the optimal solution \bar{x} and the optimal value ν
 - 5 **else if** $\bar{x} \notin X_i$ for some $i \in [m]$ **then**
 - 6 | set $t_i := t_i + 2$
 - 7 **end**
 - 8 Go to line 2 and repeat this process until $\bar{x} \in X_i$ for every $i \in [m]$.
-

It is noteworthy that in each iteration, new constraints as well as decision variables are added to the relaxed problem (15). That is, both column and constraint generation procedures are included in our algorithm. Since $g_{t_i}^{(i)}(x)$ approaches the probability $\mathbb{P}\left[\sum_{j \in [n]} \tilde{a}_{ij} x_j \leq k_i - 1\right]$ as the parameter t_i increases, the sequence of solutions of these relaxed problems will eventually converge to an optimal solution of the given problem within a finite number of iterations.

2.3 Reducing the number of chance constraints

In this section, we propose an approach to reducing the number of constraints in the chance-constrained problem (2), using vector dominance relations defined in a partially ordered set (also called poset). We first consider problem (2) under the assumption that for any fixed set $j \in [n]$, $P(\tilde{a}_{ij} = 1) = p_j \forall i \in [m]$, i.e., the set j covers each item $i \in [m]$ with equal probability p_j . Then, the following lemma provides an approach to reduce the number of chance constraints in (2).

Lemma 4 *If for any fixed set $j \in [n]$, $\mathbb{P}(\tilde{a}_{ij} = 1) = p_j$, $\forall i \in [m]$ and $i_1, i_2 \in [m]$ are two rows with $k_{i_1} \geq k_{i_2}$ and $\epsilon_{i_1} \leq \epsilon_{i_2}$, then $X_{i_1} \cap X_{i_2} = X_{i_1}$, where the probabilistic covering sets X_{i_1} and X_{i_2} correspond to the i_1 -th and i_2 -th items, respectively.*

For a more general setting, we define a partially ordered set by considering the relation between the coefficients of the constraints in the chance-constrained problem.

Let $P(\tilde{a}_{ij} = 1) =: p_{ij}$ for each $i \in [m]$ and $j \in [n]$. Given two vectors $v^{i_1} := (p_{i_1 1}, p_{i_1 2}, \dots, p_{i_1 n}, -k_{i_1}, \epsilon_{i_1})$ and $v^{i_2} := (p_{i_2 1}, p_{i_2 2}, \dots, p_{i_2 n}, -k_{i_2}, \epsilon_{i_2})$ of length $n + 2$ corresponding to the i_1^{th} and i_2^{th} row of the chance constraints (1b), the relation $v^{i_1} \leq v^{i_2}$ means that $v_t^{i_1} \leq v_t^{i_2}$ for every $t = 1, \dots, n + 2$, whereas the relation $v^{i_1} \not\leq v^{i_2}$ means that there exists at least one index t such that $v_t^{i_1} > v_t^{i_2}$. The relations “ \geq ” and “ $\not\leq$ ” can also be defined similarly. We say v^{i_1} and v^{i_2} are *comparable* if $v^{i_1} \leq v^{i_2}$ or $v^{i_2} \leq v^{i_1}$. Otherwise they are *incomparable*. Let $\mathcal{C} := \{v^1, v^2, \dots, v^m\}$ be the set of all vectors corresponding to the associated chance-constraints. Clearly, the relation “ \leq ” is a *partial order* on the set \mathcal{C} , as it is reflexive, antisymmetric, and transitive. Then the following result generalizes Lemma 4:

Lemma 5 *If there exist two items $i_1, i_2 \in [m]$ such that $v^{i_1} \leq v^{i_2}$, then $X_{i_1} \cap X_{i_2} = X_{i_1}$, where the probabilistic covering sets X_{i_1} and X_{i_2} correspond to the i_1 -th and i_2 -th items, respectively.*

We consider minimal and maximal elements of the partially ordered set (\mathcal{C}, \leq) . An element $v \in \mathcal{C}$ is a *minimal* element if there is no element $u \in \mathcal{C}$ and $u \neq v$ such that $u \leq v$. An element $v \in \mathcal{C}$ is a *maximal* element if there is no element $u \in \mathcal{C}$ and $u \neq v$ such that $v \leq u$. Based on Lemma 5, the chance-constrained problem (2) can be further simplified by only focusing on the chance-constraints whose rows correspond to the minimal elements of \mathcal{C} .

Theorem 4 Let $\mathcal{C}' \subseteq \mathcal{C}$ be the set of all minimal elements of \mathcal{C} , and $I \subseteq [m]$ be the rows corresponding to the set \mathcal{C}' . Then the chance-constrained problem (2) is equivalent to the following problem

$$\min \left\{ \sum_{j \in [n]} c_j x_j : x \in X_i, \forall i \in I, x \in B \subseteq \{0, 1\}^n \right\}$$

It can be readily seen that the worst-case complexity of finding all minimal elements of the poset (\mathcal{C}, \leq) is $O(m^2n)$, since if no elements are comparable, then we need to compare every element to every other element in order to determine that they are all minimal elements. In this case, the number of comparisons between elements is $O(m^2)$, and comparing every two elements takes $O(n + 2) = O(n)$.

Remark 3 Note that no two distinct elements in \mathcal{C}' are comparable, and for any vector $v \in \mathcal{C} \setminus \mathcal{C}'$, v is comparable to at least one element in \mathcal{C}' . Thus, \mathcal{C}' is a maximal antichain of the poset (\mathcal{C}, \leq) , where an antichain is a subset of a partially ordered set such that any two distinct elements in the subset are incomparable and a maximal antichain is an antichain that is not a proper subset of any other antichain.

2.4 Presolving method

Based on the analyses conducted in the previous and following sections, we can propose a presolving method to simplify chance-constrained problem and improve the efficiency of our algorithm. At the first step, we use vector dominance relations to reduce the problem’s complexity by identifying and eliminating redundant chance-constraints in the problem (2) (see details in Section 2.3). Denote the collection of remaining items by I . If $k_i = 1$ for some item $i \in I$, then use the “log-transformation” technique to reformulate the probabilistic covering set X_i (see details in Section 3.1). And if the item i is covered by each set $j \in [n]$ with equal probability p_i for some $i \in I$, then reformulate the probabilistic covering set X_i using the method mentioned in Theorem 5 (see details in Section 3.2). Finally, we solve the simplified problem using the outer-approximation algorithm (see details in Section 2.2).

To sum up, we first attempt to reduce the number of chance-constraints by only considering the set of all minimal vectors corresponding to the associated chance-constraints. Further, we employ two distinct reformulation techniques (as stated in Section 3) to handle two specific cases. These reformulation techniques will be more efficient in the implementation process, as they involve significantly less variables and constraints compared to those in Theorem 1. Additionally, it is worth mentioning that at the final step, in the process of checking feasibility of our solution, we can utilize the discrete Fourier transform (DFT) method to compute the PMF and CDF of Poisson binomial distribution instead of using Lemma 1 directly. The DFT method provides us with an effective way to mitigate the cumulative rounding error. For more information about the DFT method, please refer to [Hong \(2013\)](#).

3 Reformulation for Two Special Cases of CC-SMCP

3.1 Reformulation for probabilistic set covering problem

Suppose $k_i = 1$ for $i = 1, \dots, m$. As mentioned before, when $k_i = 1$ for each $i \in [m]$, the problem (1) becomes the probabilistic set covering (PSC) problem, which has already been well studied by [Haight et al. \(2000\)](#), [Fischetti and Monaci \(2012\)](#), and [Ahmed and Papageorgiou \(2013\)](#). In the independent case, the probabilistic covering set X can be further simplified using “log-transformation”. We include it here for the sake of completeness.

Proposition 1 (Ahmed and Papageorgiou (2013), Fischetti and Monaci (2012), Haight et al. (2000))

Let \tilde{a}_j be independent random variable with $\mathbb{P}(\tilde{a}_j = 1) = p_j$ for each $j \in [n]$, x_j 's be binary variables, and $k = 1$ in the probabilistic covering set X . Then

$$X = \left\{ x \in \{0, 1\}^n : \sum_{j \in [n]} \log(1 - p_j)x_j \leq \log \epsilon \right\}. \quad (16)$$

It follows that, when $k_i = 1$ for each $i \in [m]$, CC-SMCP is equivalent to the following deterministic binary linear program (BIP):

$$\min \left\{ \sum_{j \in [n]} c_j x_j : \sum_{j \in [n]} \log(1 - p_{ij})x_j \leq \log \epsilon_i, \forall i \in [m], x \in B \subseteq \{0, 1\}^n \right\}.$$

3.2 Case when cover probabilities of each item are equal

Assume that for any fixed item $i \in [m]$, we have $\mathbb{P}(\tilde{a}_{ij} = 1) = p_i, \forall j \in [n]$. That is, the item i is covered by each set $j \in [n]$ with equal probability p_i . We first notice that in this case, the value of the probability $\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_{ij}x_j \geq k_i \right]$ is only related to the number of selected sets in the solution. In this case, given a candidate solution $\bar{x} \in \{0, 1\}^n$, the following lemma provides an efficient way to compute the probability $\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right]$ rather than using Lemma 1 directly:

Lemma 6 Let \tilde{a}_j be independent random variable with $\mathbb{P}(\tilde{a}_j = 1) = p$ for each $j \in [n]$, $\bar{x} \in \{0, 1\}^n$ with $\sum_{j \in [n]} \bar{x}_j = d$, and $k \in \mathbb{Z}_+$. Then

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] = \sum_{\ell=k}^d \binom{d}{\ell} p^\ell (1-p)^{d-\ell}. \quad (17)$$

The above result provides an efficient way to compute the cover probability $\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right]$ without knowing the exact value of \bar{x}_j , and it only depends on the value of the summation $\sum_{j \in [n]} \bar{x}_j$. However, it has a severe drawback: due to the computer's inability to represent some numbers exactly, the calculation may result in the large cumulative rounding error that cannot be ignored, especially when d is a large number. Indeed, the calculation (17) including a series of summation operations of very small numbers can lead the cumulative error to be especially problematic. Even if the rounding error from a single operation (addition, product) is small, the cumulative error from many operations may be significant. Taking the above considerations into account, we develop an alternative equivalent explicit formula to compute the cover probability. This formula can be readily implemented in practice, as numerous software packages offer efficient methods for calculating the values of certain special functions. Specifically, we utilize the beta function $B(a, b)$ and the incomplete beta function $B(q; a, b)$:

$$B(a, b) = \int_0^1 t^{a-1} (1-t)^{b-1} dt \quad \text{and} \quad B(q; a, b) = \int_0^q t^{a-1} (1-t)^{b-1} dt.$$

Then, the next lemma provides another computational method that reduces the cumulative rounding error to obtain the cover probability:

Lemma 7 Let \tilde{a}_j be independent random variables with $\mathbb{P}(\tilde{a}_j = 1) = p$ for each $j \in [n]$, $\bar{x} \in \{0, 1\}^n$ with $\sum_{j \in [n]} \bar{x}_j = d$, and $k \in \mathbb{Z}_+$. Then

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] = k \binom{d}{k} \sum_{\ell=k}^d (-1)^{\ell-k} \binom{d-k}{\ell-k} \frac{p^\ell}{\ell} = k \binom{d}{k} \int_0^p u^{k-1} (1-u)^{d-k} du, \quad (18)$$

or equivalently,

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] = \binom{d}{k} p^k {}_2F_1(k-d, k; k+1; p) = I_p(k, d-k+1), \quad (19)$$

where ${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{a^{(n)} b^{(n)}}{c^{(n)}} \frac{z^n}{n!} = 1 + \frac{ab}{c} \frac{z}{1!} + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{z^2}{2!} + \dots$, $|z| < 1$, is the hypergeometric function and $I_q(a, b) = \frac{B(q, a, b)}{B(a, b)}$ is the regularized incomplete beta function.

Next we derive an equivalent reformulation for the probabilistic covering set X based on the above results. Considering CC-SMCP with a single chance-constraint:

$$\min \left\{ \sum_{j \in [n]} c_j x_j : x \in X, x \in B \subseteq \{0, 1\}^n \right\}, \quad (20)$$

we explore properties of a solution to problem (20) when $P(\tilde{a}_j = 1) = p$, for each $j \in [n]$:

Lemma 8 *Let \tilde{a}_j be independent random variables with $\mathbb{P}(\tilde{a}_j = 1) = p$ for each $j \in [n]$. If \bar{x} is a feasible solution to the problem (20) with $\sum_{j=1}^n \bar{x}_j = d$, then for any $\hat{x} \in B \subseteq \{0, 1\}^n$ with $\sum_{j=1}^n \hat{x}_j = \hat{d} \geq d$, \hat{x} is also a feasible solution to the problem (20).*

Lemmas 7 and 8 imply the following result:

Theorem 5 *Let \tilde{a}_j be independent random variables with $\mathbb{P}(\tilde{a}_j = 1) = p$ for each $j \in [n]$. Then x is a feasible solution to problem (20) if and only if x is a feasible solution to the following deterministic covering problem:*

$$\min \left\{ \sum_{j \in [n]} c_j x_j : \sum_{j \in [n]} x_j \geq \bar{d}, x \in B \subseteq \{0, 1\}^n \right\},$$

where

$$\bar{d} := \min \{d \in \mathbb{Z}_+ : I_p(k, d-k+1) \geq 1-\epsilon, k \leq d \leq n\}. \quad (21)$$

That is, the probabilistic covering set X admits the deterministic equivalent reformulation

$$X = \left\{ x \in \{0, 1\}^n : \sum_{j \in [n]} x_j \geq \bar{d} \right\}.$$

Proof. The necessity follows directly from Lemma 7, and the sufficiency from Lemma 8. ■

For any feasible solution \bar{x} to problem (20), a simple (but potentially weak) lower bound for the sum $\sum_{j \in [n]} \bar{x}_j$ can be obtained using Markov's inequality.

Proposition 2 *Let \tilde{a}_j be independent random variables with $\mathbb{P}(\tilde{a}_j = 1) = p$ for each $j \in [n]$, and \bar{x} be a feasible solution to the chance-constrained problem (20) with $\sum_{j=1}^n \bar{x}_j = d$, then*

$$d := \sum_{j \in [n]} \bar{x}_j \geq \max \left\{ k, \left\lceil \frac{k(1-\epsilon)}{p} \right\rceil \right\}.$$

Actually, from Lemma 7 and 8, we can notice that for any fixed k and p , the function $I_p(k, d-k+1)$ is monotone non-decreasing with respect to d . Thus, the tightest lower bound for $d := \sum_{j \in [n]} \bar{x}_j$ (i.e., the optimal value of problem (21)) can be obtained by the binary search. Based on the above analysis, we can provide the following algorithm to solve CC-SMCP when cover probabilities of each item are equal: we first use binary search algorithm to obtain \bar{d}_i for each item $i \in [m]$ where

$$\bar{d}_i := \min \{d_i \in \mathbb{Z}_+ : I_p(k_i, d_i - k_i + 1) \geq 1 - \epsilon_i, k_i \leq d_i \leq n\},$$

and choose the maximum d_i to satisfy all the chance constraints, i.e., let $\bar{d} := \max\{\bar{d}_i, i \in [m]\}$. Finally, we solve the following deterministic covering problem to the optimality:

$$\min \left\{ \sum_{j \in [n]} c_j x_j : \sum_{j \in [n]} x_j \geq \bar{d}, x \in B \subseteq \{0, 1\}^n \right\}.$$

Remark 4 Note that if $B = \{0, 1\}^n$, then we can obtain an explicit formula for the optimal value of the chance-constrained problem (2) when cover probabilities of each item are equal to each other. Suppose that the sequence c'_1, \dots, c'_n is an order of c_1, \dots, c_n satisfying $c'_1 \leq c'_2 \leq \dots \leq c'_n$. Based on the above algorithm, the optimal value of the chance-constrained problem (2) can be written as $\sum_{j=1}^{\bar{d}} c'_j$.

4 Sampling-Based Approaches

In the following sections, we discuss two sampling-based methods, sample average approximation and importance sampling, to solve CC-SMCP on a finite discrete probability space.

4.1 Sample average approximation

In this section, we consider CC-SMCP on a finite discrete probability space $(\Omega, 2^\Omega, \mathbb{P}_N)$ where $\Omega = \{\omega_1, \dots, \omega_N\}$ and $p(\omega_i) = \mathbb{P}_N(\omega = \omega_i)$. The sample average approximation (SAA) method approximates the true distribution via a finite empirical distribution, \mathbb{P}_N . We assume that the samples are independent and identically distributed (i.i.d.) with $p(\omega) = 1/N, \omega \in \Omega$. The SAA formulation of CC-SMCP is

$$\min \quad \mathbf{c}^T \mathbf{x} \tag{22a}$$

$$\text{s.t.} \quad \tilde{A}_i(\omega) \mathbf{x} + M_i(\omega)(1 - z_i(\omega)) \geq k_i, \quad \forall \omega \in \Omega, \forall i \in [m] \tag{22b}$$

$$\frac{1}{N} \sum_{\omega \in \Omega} z_i(\omega) \geq 1 - \alpha_i, \quad \forall i \in [m] \tag{22c}$$

$$\mathbf{z}_i \in \{0, 1\}^N, \quad \forall i \in [m] \tag{22d}$$

$$\mathbf{x} \in B \subseteq \{0, 1\}^n \tag{22e}$$

where $\tilde{A}_i(\omega)$ is the i -th row of the coefficient matrix \tilde{A} in scenario ω , $M_i(\omega)$ is the big-M coefficient that guarantees feasibility whenever any element in $z_i(\omega)$ is equal to zero, and $\alpha_i \in [0, 1)$ is the risk level that may be different from ϵ_i in CC-SMCP.

Remark 5 Generally, SAA is a sampling-based method commonly employed to approximate the function/distribution $f(\mathbf{x}, \xi)$ that cannot be observed or computed directly. Here we choose to use the term ‘‘SAA’’ since the idea of our method is the same as SAA.

Often, the linear programming (LP) relaxation of the SAA reformulation (22) is very weak because of the big-M coefficients that are introduced to model the chance constraints. Fortunately, note that in CC-SMCP problem, we have $\tilde{A} \in \{0, 1\}^{m \times n}$, $\mathbf{x} \in \{0, 1\}^n$ and $k_i \in \mathbb{Z}_+$. In this case, we can set big-M coefficient $M_i(\omega) := k_i$ for each $\omega \in \Omega$ to obtain a big-M-free SAA reformulation:

$$[\text{SAA}] \quad \min \left\{ \mathbf{c}^T \mathbf{x} : \tilde{A}_i(\omega) \mathbf{x} \geq k_i z_i(\omega), \forall \omega \in \Omega, \forall i \in [m], (22c) - (22e) \right\}. \tag{23}$$

For ease of analysis, we rewrite the SAA reformulation as follows:

$$\nu_\alpha^N := \min \{ \mathbf{c}^T x : x \in X_\alpha^N \}, \tag{24}$$

where

$$X_\alpha^N := \left\{ x \in B : \frac{1}{N} \sum_{\omega \in \Omega} \mathbb{I} \left(\tilde{A}_i(\omega) x \geq k_i \right) \geq 1 - \alpha_i, \quad \forall i \in [m] \right\},$$

where $\mathbb{I}(\cdot)$ is the indicator function. Motivated by theoretical results in [Luedtke and Ahmed \(2008\)](#) regarding SAA for chance-constrained problems with joint probabilistic constraints, that, we study approximations of chance-constrained combinatorial optimization problems with individual probabilistic constraints in this paper. And our goal is to establish statistical relationships between CC-SMCP and its SAA reformulation (24) for $\alpha_i \geq 0, \forall i \in [m]$. We assume that CC-SMCP has an optimal solution x^* and a finite optimal value ν_ϵ^* . And for each $i \in [m]$, we use $q_i(x^*)$ to denote the probability of the event $\{\tilde{A}_i(\omega)x^* \leq k_i - 1\}$. By Lemma 1, we have $q_i(x^*) = \sum_{d=0}^{k_i-1} \sum_{\ell=d}^n (-1)^{\ell-d} \binom{\ell}{d} h_\ell(x^*)$ and $q_i(x^*) \leq \epsilon_i$ since $x^* \in X_i$ is a solution to CC-SMCP. We now establish a bound on the probability that ν_α^N yields a lower bound for ν_ϵ^* .

Theorem 6 *Assume that ν_ϵ^* and ν_α^N are the optimal values to CC-SMCP and the SAA reformulation (24), respectively. Then*

$$\mathbb{P} [\nu_\alpha^N \leq \nu_\epsilon^*] \geq 1 - \sum_{i=1}^m I_{\epsilon_i}(\lfloor \alpha_i N \rfloor + 1, N - \lfloor \alpha_i N \rfloor), \quad (25)$$

where $I_x(a, b)$ is the regularized incomplete beta function. Moreover, if $\epsilon_i \leq \alpha_i$ for each $i \in [m]$, then

$$\mathbb{P} [\nu_\alpha^N \leq \nu_\epsilon^*] \geq 1 - m \exp(-\kappa_1 N), \quad (26)$$

where $\kappa_1 := \min_{i \in [m]} \{(\alpha_i - \epsilon_i)^2 / (\alpha_i + \epsilon_i)\}$.

Theorem 6 states that, when the risk parameter $\alpha_i > \epsilon_i$ for each $i \in [m]$, the SAA method yields a lower bound to the true optimal value of CC-SMCP with probability approaching one exponentially fast as N increases. Further, given a confidence $1 - \delta$, where $0 < \delta < 1$, Theorem 6 ensures that $\nu_\alpha^N \leq \nu_\epsilon^*$ with probability of at least $1 - \delta$, if we choose $\epsilon_i < \alpha_i$ for any $i \in [m]$ and the sample size

$$N \geq \frac{1}{\kappa_1} \ln \frac{m}{\delta}.$$

In fact, with this choice of α_i and N , we have

$$\mathbb{P} [\nu_\alpha^N \leq \nu_\epsilon^*] \geq 1 - m \exp(-\kappa_1 N) \geq 1 - \delta.$$

Next we investigate conditions under which an optimal solution of SAA problem (24) is a feasible solution to CC-SMCP. Let $X_\epsilon := \bigcap_{i \in [m]} X_{\epsilon, i}$ where $X_{\epsilon, i} := \{x \in B : \mathbb{P}[\tilde{A}_i x \geq k_i] \geq 1 - \epsilon_i\}$. We assume that $\alpha_i < \epsilon_i$ for each $i \in [m]$. The idea is that if the risk level α_i is less than ϵ_i for each $i \in [m]$, then, given a sufficiently large sample size N , the feasible region X_α^N will be a subset of X_ϵ . Consequently, any optimal solution to the SAA reformulation (24) must be feasible for CC-SMCP. We have the following theorem:

Theorem 7 *Assume that $\alpha_i < \epsilon_i$ for each $i = 1, \dots, m$, and let $\kappa_2 := 2 \min_{i \in [m]} (\epsilon_i - \alpha_i)^2$. Then*

$$\mathbb{P} [X_\alpha^N \subseteq X_\epsilon] \geq 1 - m |B \setminus X_\epsilon| \exp\{-\kappa_2 N\}.$$

The above theorem also provides us a way to estimate the sample size N such that the feasible solutions of SAA reformulation (24) are feasible to CC-SMCP with a high probability (confidence) $1 - \delta$, if $\alpha_i < \epsilon_i$ for each $i \in [m]$ and we choose the sample size

$$N \geq \kappa_2^{-1} \ln \frac{m |B \setminus X_\epsilon|}{\delta}.$$

Moreover, note that $B \subseteq \{0, 1\}^n$ in CC-SMCP, we can take

$$N \geq \frac{1}{\kappa_2} \ln \frac{m}{\delta} + \frac{n}{\kappa_2} \ln(2)$$

to attain the required confidence.

We are interested in determining the conditions under which the optimal value ν_α^N of the SAA problem will converge to the true optimal value ν_ϵ^* of CC-SMCP with probability one as N approaches infinity for $\alpha = \epsilon$. Notice that Theorem 6 provides a trivial lower bound for the probability $\mathbb{P}[\nu_\epsilon^N \leq \nu_\epsilon^*]$ when $\alpha_i = \epsilon_i$ for any $i \in [m]$. In the following, we will delve further by combining Theorem 6 and Theorem 7 to establish a lower bound for the probability $\mathbb{P}[\nu_\epsilon^N = \nu_\epsilon^*]$. Let X_ϵ^* be the set of optimal solutions to CC-SMCP, and define

$$\underline{\alpha}_i := \max \left\{ \mathbb{P} \left[\tilde{A}_i x \leq k_i - 1 \right] : x \in X_\epsilon^* \right\}, \quad \forall i \in [m],$$

and $\underline{\alpha} := (\underline{\alpha}_1, \dots, \underline{\alpha}_m)$. By definition, we have $\underline{\alpha}_i \leq \epsilon_i$ for any $i \in [m]$ and $\nu_{\underline{\alpha}}^* = \nu_\epsilon^*$. Let $X_\alpha^N = \bigcap_{i \in [m]} X_{\alpha,i}^N$, where $X_{\alpha,i}^N := \left\{ x \in B : \frac{1}{N} \sum_{\omega \in \Omega} \mathbb{I} \left(\tilde{A}_i(\omega) x \geq k_i \right) \geq 1 - \alpha_i \right\}$, as well as

$$\bar{\alpha}_i := \min \left\{ \mathbb{P} \left[\tilde{A}_i x \leq k_i - 1 \right] : x \in B \setminus X_{\epsilon,i} \right\}, \quad \forall i \in [m],$$

and $\bar{\alpha} := (\bar{\alpha}_1, \dots, \bar{\alpha}_m)$. Without loss of generality, we assume that $B \setminus X_{\epsilon,i} \neq \emptyset$. Otherwise, since $B \setminus X_{\epsilon,i} = \emptyset$ implies $B \subseteq X_{\epsilon,i}$, we can remove the i th chance-constraint in CC-SMCP. By definition, we have $\bar{\alpha}_i > \epsilon_i$ for any $i \in [m]$. Then we have the following theorem:

Theorem 8 *Assume that $\underline{\alpha}_i < \epsilon_i$ for each $i \in [m]$, ν_ϵ^* and ν_α^N are the optimal values to CC-SMCP and the SAA reformulation (24), respectively. Then*

$$\mathbb{P}[\nu_\epsilon^N = \nu_\epsilon^*] \geq 1 - m(|B \setminus X_\epsilon| + 1) \exp\{-\kappa_3 N\},$$

where $\kappa_3 := \min \left\{ \min_{i \in [m]} \left\{ (\underline{\alpha}_i - \epsilon_i)^2 / (\underline{\alpha}_i + \epsilon_i) \right\}, 2 \min_{i \in [m]} \left\{ (\bar{\alpha}_i - \epsilon_i)^2 \right\} \right\}$.

Note that the assumption that $\underline{\alpha}_i < \epsilon_i$ for each $i \in [m]$ is mild, because $B \subseteq \{0, 1\}^n$ is finite and there are only a finite number of values of ϵ_i such that $\underline{\alpha}_i = \epsilon_i$. This fact inspires us to add a random perturbation uniformly distributed in $[-\gamma_i, \gamma_i]$ to ϵ_i when $\underline{\alpha}_i = \epsilon_i$, where γ_i can be arbitrarily small, then the assumption will hold with probability one.

Theorem 8 proves that solving a sample approximation with $\alpha = \epsilon$ will yield an exact optimal solution with probability approaching one exponentially fast with N . However, the sample size N required to ensure a reasonably high probability of obtaining the optimal solution will be at least proportional to κ_3^{-1} and therefore may be very large. Hence, Theorem 8, which reflects the qualitative behavior of the sample approximation with $\alpha = \epsilon$, may not be suitable for estimating the required sample size. In Section 5, we will observe that SAA exhibits good performance even with a small sample size N .

4.2 Importance sampling

Importance sampling (IS) is probably the most popular approach to reduce the variance of an estimator. In the case of rare event estimation, this also means increasing the occurrence of rare events. Here we briefly review the basic ideas of IS based on CC-SMCP as an example. For more details about IS methods and applications, we refer the interested reader to (Barrera et al. 2016, Rubino et al. 2009).

Let

$$q_i(x) := \mathbb{P}[\tilde{A}_i x < k] = \mathbb{E}_{\tilde{a}}[\mathbb{I}(\tilde{A}_i x < k_i)], \quad \forall i \in [m], \quad (27)$$

where $\tilde{A}_i := (\tilde{a}_{i1}, \dots, \tilde{a}_{in})$ is a row random vector. In the following we will assume $m = 1$, and drop the row subscript i like what we have done in Section 2. We want to estimate $q(x)$ for all $x \in X$. For a given sample $\tilde{a}(\omega_1), \dots, \tilde{a}(\omega_N)$ of size N from the distribution of \tilde{a} , a natural approximation of $q(x)$ in (27) is the SAA estimator:

$$\hat{q}^{SAA}(x) := \frac{1}{N} \sum_{\ell=1}^N \mathbb{I}(\tilde{a}(\omega_\ell) x < k). \quad (28)$$

Note that $\hat{q}^{SAA}(x)$ is an unbiased estimator of $q(x)$, since

$$\mathbb{E}_{\tilde{a}} [\hat{q}^{SAA}(x)] = \frac{1}{N} \sum_{\ell=1}^N \mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}(\omega_\ell)x < k)] = q(x).$$

Now let us consider \hat{a} a new random vector and let $\hat{a}(\omega_1), \dots, \hat{a}(\omega_N)$ be i.i.d. copies of \hat{a} . Define

$$\hat{q}^{IS}(x) := \frac{1}{N} \sum_{\ell=1}^N \mathbb{I}(\hat{a}(\omega_\ell)x < k) L(\hat{a}(\omega_\ell)), \quad (29)$$

where $L(\cdot)$ is the likelihood ratio $L(\hat{a}) = \prod_{j=1}^n \left(\frac{p_j}{\hat{p}_j}\right)^{\hat{a}_j} \left(\frac{1-p_j}{1-\hat{p}_j}\right)^{1-\hat{a}_j}$ with $p_j = P(\tilde{a}_j = 1)$ and $\hat{p}_j = P(\hat{a}_j = 1)$ for each $j \in [n]$. In this case, L is the ratio between the respective probabilities mass functions, since both \tilde{a} and \hat{a} have discrete support. Notice that for any function $f(\cdot) : \mathbb{R}^n \mapsto \mathbb{R}$, we have

$$\mathbb{E}_{\tilde{a}}[f(\tilde{a})] = \mathbb{E}_{\hat{a}}[f(\hat{a})L(\hat{a})]. \quad (30)$$

Based on the above observation, we obtain $\hat{q}^{IS}(x)$ is also an unbiased estimator of $q(x)$, since

$$\mathbb{E}_{\hat{a}} [\hat{q}^{IS}(x)] = \frac{1}{N} \sum_{\ell=1}^N \mathbb{E}_{\hat{a}} [\mathbb{I}(\hat{a}(\omega_\ell)x < k) L(\hat{a}(\omega_\ell))]$$

and

$$\mathbb{E}_{\hat{a}} [\mathbb{I}(\hat{a}x < k) L(\hat{a})] = \mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}x < k)] = q(x).$$

Further, the variance of the SAA estimator $\hat{q}^{SAA}(x)$ in (28) is

$$\text{Var} [\hat{q}^{SAA}(x)] = \frac{1}{N} (q(x) - q(x)^2) = \frac{1}{N} \mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}x < k)] - \frac{1}{N} q(x)^2$$

whereas the variance of the IS estimator $\hat{q}^{IS}(x)$ in (29) is given by

$$\text{Var} [\hat{q}^{IS}(x)] = \frac{1}{N} \mathbb{E}_{\hat{a}} [\mathbb{I}(\hat{a}x < k)^2 L(\hat{a})^2] - \frac{1}{N} q(x)^2 = \frac{1}{N} \mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}x < k) L(\tilde{a})] - \frac{1}{N} q(x)^2, \quad (31)$$

where the second equality follows from (30). Note that if we choose a “good” IS distribution in such a way that the event $\mathbb{I}(\tilde{a}x < k)$ becomes more likely under that distribution provided that $L(\tilde{a}) \leq 1$, the variance of the IS estimator will be smaller than that of the standard SAA estimator.

To use the IS estimator $\hat{q}^{IS}(x)$ in the formulation of CC-SMCP, we only need to replace the chance-constraint (1b) by

$$\hat{A}_i(\omega)\mathbf{x} \geq k_i z_i(\omega), \forall \omega \in \Omega, \forall i \in [m] \quad (32a)$$

$$\sum_{\omega \in \Omega} L(\hat{A}_i(\omega)) (1 - z_i(\omega)) \leq N\epsilon_i, \forall i \in [m] \quad (32b)$$

$$z_i(\omega) \in \{0, 1\}, \forall \omega \in \Omega \quad (32c)$$

where the definitions of $\hat{A}_i(\omega)$ and $z_i(\omega)$ are similar as those in the SAA reformulation (23). Note that if we choose $L(\hat{A}_i(\omega)) \equiv 1$ for each $i \in [m]$, then the obtained reformulation will exactly match the SAA reformulation as indicated in equation (23).

The remaining problem becomes finding a “good” IS estimator by minimizing the the variance of $\hat{q}^{IS}(x)$. Suppose all components of \tilde{a} and \hat{a} are independent, from the expression (31), we consider minimizing the term

$$\begin{aligned} \mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}x < k) L(\tilde{a})] &= \mathbb{E}_{\tilde{a}} \left[\mathbb{I} \left(\sum_{j=1}^n \tilde{a}_j x_j < k \right) \prod_{j=1}^n \left(\frac{p_j}{\hat{p}_j} \right)^{\tilde{a}_j} \left(\frac{1-p_j}{1-\hat{p}_j} \right)^{1-\tilde{a}_j} \right] \\ &= \mathbb{E}_{\tilde{a}} \left[\mathbb{I} \left(\sum_{j=1}^n \tilde{a}_j x_j < k \right) \prod_{j=1}^n \left(\frac{p_j(1-\hat{p}_j)}{\hat{p}_j(1-p_j)} \right)^{\tilde{a}_j} \prod_{j=1}^n \left(\frac{1-p_j}{1-\hat{p}_j} \right) \right] \end{aligned} \quad (33)$$

For each $j \in [n]$, let

$$\lambda_j := \log \left(\frac{p_j(1-\hat{p}_j)}{\hat{p}_j(1-p_j)} \right) = \log \left(\frac{1/\hat{p}_j-1}{1/p_j-1} \right)$$

Note that $\lambda_j \geq 0$ when $\hat{p}_j \leq p_j$ which is the case that IS distribution works by reducing the cover probability of each set j such that the event $\sum_{j=1}^n \tilde{a}_j x_j < k$ happens more often. By plugging λ_j into equation (33), it follows that

$$\mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}x < k)L(\tilde{a})] = \mathbb{E}_{\tilde{a}} \left[\mathbb{I} \left(\sum_{j=1}^n \tilde{a}_j x_j < k \right) \exp \left(\sum_{j=1}^n \lambda_j \tilde{a}_j \right) \prod_{j=1}^n \left(e^{-\lambda_j} p_j + (1-p_j) \right) \right], \quad (34)$$

where the equation follows from $\frac{1-p_j}{1-\hat{p}_j} = e^{-\lambda_j} p_j + (1-p_j)$. The task of minimizing $\mathbb{E}_{\tilde{a}} [\mathbb{I}(\tilde{a}x < k)L(\tilde{a})]$ requires solving a multidimensional stochastic nonlinear problem, which is quite challenging. Alternatively, our approach focuses on minimizing the largest term within the expectation, that is

$$B_x(\lambda) := \max_{\substack{\tilde{a}: \sum_{j=1}^n \tilde{a}_j x_j < k \\ \tilde{a} \in \{0,1\}^n}} \exp \left(\sum_{j=1}^n \lambda_j \tilde{a}_j \right) \prod_{j=1}^n \left(e^{-\lambda_j} p_j + (1-p_j) \right). \quad (35)$$

Note that $B_x(\lambda) \geq 0$. We consider minimizing $B_x(\lambda)$ over $\lambda \geq 0$. Then we have the following theorem that reduces the minimization problem to a one-dimensional problem, which can be efficiently solved by using some numerical methods.

Theorem 9 *Suppose that $0 < p_j < 1$ for all $j = 1, \dots, n$. Let x be the solution that satisfies $\sum_{j=1}^n x_j = u$ and $n - u + k - 1 < \sum_{j=1}^n p_j x_j$. Then the function $B_x(\lambda)$ is convex and there exists $\lambda_x^* \in \mathbb{R}_+$ such that the vector λ defined as $\lambda_j = \lambda_x^*$ for each $j = 1, \dots, n$ minimizes $B_x(\lambda)$. Moreover, λ_x^* and $\hat{p}_j(\lambda_x^*)$ satisfy*

$$\sum_{j=1}^n \hat{p}_j(\lambda_x^*) = n - u + k - 1 \text{ and } \hat{p}_j(\lambda_x^*) = \frac{e^{-\lambda_x^*} p_j}{e^{-\lambda_x^*} p_j + (1-p_j)}. \quad (36)$$

Remark 6 *Note that if $\sum_{j=1}^n x_j < k$, then all components of vector \tilde{a} in $B_x(\lambda)$ should be one, which implies the optimal solutions to minimizing $B_x(\lambda)$ should be $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$, since $\nabla \log B_x(\lambda) \geq 0$ for all $\lambda \geq 0$. In this case, due to the definition of λ_j , we have $\hat{p}_j = p_j$ for each $j \in [n]$.*

Theorem 9 provides a method for selecting good IS parameters for a given solution $x \in X$. To choose the importance sampling estimator for the whole set X , based on the conditions in Theorem 9, we notice that a reasonable lower bound for the sum $\sum_{j=1}^n x_{ij}$ can be $\lceil (n + k_i - 1)/2 \rceil$. This is because $n - u + k_i - 1 < \sum_{j=1}^n p_{ij} x_{ij} \leq \sum_{j=1}^n x_{ij} = u$. In the following numerical experiments, we will use $\lceil (n + k_i - 1)/2 \rceil$ as an estimator for the sum $\sum_{j=1}^n x_{ij}$, and then calculate the value for \hat{p}_{ij} using equations (36). However, as we do not know the exact value of the sum $\sum_j x_{ij}$, this choice can sometimes be too radical to obtain a feasible solution to the CC-SMCP, as observed in the experimental section.

5 Numerical Experiments

In this section, we study the computational performance of our proposed methods for solving the CC-SMCP. We compare the outer-approximation (OA) algorithm with two different linearization techniques to the sampling-based algorithms, i.e., the SAA and IS approaches, in terms of both time and effectiveness. The detailed experimental settings are listed at the beginning of the following subsections.

All experiments are conducted using Python 3.9 with the optimization solver Gurobi 10.0.2 on a Linux server running CentOS 7 with one AMD EPYC 7642 48-core processor (2.4GHz) and 192 RAM. For all experiments, we use a time limit of 3,600 seconds and report the running time in CPU seconds. All other control parameters remain at their default settings within the branch-and-cut framework of Gurobi. Overall, in our experiments, we compare the performance of the following four solution approaches:

- OA-I: outer approximation approach with the first linearization technique (Lemma 2) and presolving process applied (i.e., presolving + OA + linearization-I)
- OA-II: outer approximation approach with the second linearization technique (standard linearization (7)) and presolving process applied (i.e., presolving + OA + linearization-II)
- SAA: sample average approximation approach (with reformulation (23))
- IS: importance sampling approach (with reformulation (32))

5.1 Computational performance of methods for CC-SMCP

To evaluate the performance of solution methods, we randomly generate *sparse* probability matrix $P = (p_{ij})_{m \times n}$: we assume that the i th row can only be covered by at most 12 columns when $k_i \geq 2$. That is, for each row i where $k_i \geq 2$, we randomly choose $n' \leq 12$ columns and sample the probability of success $p_{ij} = \mathbb{P}[\tilde{a}_{ij} = 1]$ uniformly from the interval $[0.9, 1]$. For the other $n - n'$ columns in the row i , we set $p_{ij} = 0$. Each covering number k_i is randomly selected from $\{1, 2, 3\}$, and costs c_{ij} in the objective function are all set to 1. Further, we set the deterministic set $B = \{0, 1\}^n$, and the risk parameter $\epsilon_i = 0.05$ and 0.1 for each $i \in [m]$ in these experiments, reflecting the natural assumption that we want to cover the item i with high probability. We implement the presolving process before we apply the OA method to solve each instance. For the SAA method, we let the risk parameter $\alpha_i := \epsilon_i$ in problem (23) for each $i \in [m]$, and generate each scenario $\omega \in \Omega$ independently according to the probability matrix P . For the IS approach, we set $\hat{p}_{ij} := 0$ for any column j where $p_{ij} = 0$. For columns $j \in [n]$ where $p_{ij} \neq 0$, we choose $u = \lceil (n' + k_i - 1)/2 \rceil$ and compute the IS estimator \hat{p}_{ij} using equations (36).

Table 1 lists some comparison results between four different methods: OA-I, OA-II, SAA and IS. We use “Val” to denote the best achievable value obtained among four different methods within one hour for the corresponding instance. For OA methods, if an instance has been solved within one hour, then the “Val” of OA methods should be the optimal value of the instance. The symbol ‘-’ in ‘Time’ columns indicates that the method did not obtain an optimal solution within the time limit of 3,600 seconds. Similarly, if ‘-’ appears in the ‘Val’ columns, it denotes that the method was unable to obtain a feasible solution during the last iteration. The gaps (Gap) reported are the percent by which the output value of SAA or IS is below/above the optimal value generated by the OA methods, defined as $\text{Gap} = (\text{Val} - \text{opt})/\text{opt}$, where opt denotes the optimal value of an instance and “Val” represents the objective value obtained from the SAA or IS approach. Specifically, a negative (positive) gap indicates the output value of SAA or IS is below (above) the optimal value. Since the output of SAA or IS depends on sampling results, an “INF” is placed in the column “Gap (%)” if the corresponding SAA or IS problem is infeasible.

As can be seen in Table 1, the OA approaches perform well on these instances with a sparse probability matrix, and the OA-II method outperforms the OA-I method in all 38 cases. Sampling-based methods (SAA and IS) provide good approximations for CC-SMCP on these instances, with output values close to the optimal ones. The solution time for SAA and IS may be less than that of OA methods when n or m increases. Note that most gaps in SAA and all gaps in IS are negative, indicating that sampling-based methods often struggle to find feasible solutions. Hence, developing methods and techniques for faster identification of feasible solutions in sampling-based approaches remains an area of interest.

Note that even the presence of a zero gap in the SAA/IS results of Table 1 does not guarantee that the SAA/IS has generated a feasible solution for CC-SMCP. To explore deeply on the sampling-based methods, we conduct some extra experiments. We test the sampling-based approaches on two instances with $n = 50, m = 30$ and $n = 100, m = 50$ generated in the aforementioned way. For the experiments, we consider $\epsilon_i = \epsilon \in \{0.05, 0.1\}$ and take independent samples of size $N \in \{30, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600\}$, and for each sample size we take 50 different replications. We study how the risk parameter ϵ and the sample size N affect the performance of SAA/IS. To better evaluate the performance of the sampling-based approaches, we define two performance measures:

Table 1: Comparison between OA and sampling-based methods ($p_{ij} \in [0.9, 1], N = 200$)

Parameters			OA-I			OA-II			SAA		IS			
n	m	ϵ	Time	Iterations	Val	Time	Iterations	Val	Time	Val	Gap(%)	Time	Val	Gap(%)
30	10	0.05	3.86	3	7	0.52	2	7	0.15	7	0.00	0.26	7	0.00
30	10	0.1	1.39	2	7	0.58	2	7	0.42	7	0.00	0.21	7	0.00
30	20	0.05	3.55	2	11	1.24	2	11	0.44	11	0.00	0.63	11	0.00
30	20	0.1	7.92	4	9	1.16	2	9	1.33	9	0.00	0.51	8	-11.11
30	30	0.05	5.90	2	13	4.05	2	13	8.37	13	0.00	2.82	13	0.00
30	30	0.1	2.62	2	11	1.13	2	11	12.12	11	0.00	1.16	11	0.00
30	50	0.05	17.79	3	13	2.28	2	13	2.30	13	0.00	2.22	13	0.00
30	50	0.1	13.70	3	13	2.24	2	13	3.67	13	0.00	4.19	13	0.00
30	100	0.05	19.71	2	19	3.29	3	19	4.26	19	0.00	3.86	19	0.00
30	100	0.1	21.63	3	18	5.52	2	18	6.36	18	0.00	3.42	17	-5.56
30	150	0.05	25.69	2	20	5.80	3	20	6.20	20	0.00	7.22	20	0.00
30	150	0.1	16.82	3	19	9.42	3	19	8.09	18	-5.26	9.65	19	0.00
50	30	0.05	21.94	2	14	7.44	2	14	3.21	14	0.00	1.59	13	-7.14
50	30	0.1	32.04	4	12	6.63	3	12	10.37	13	8.33	2.54	11	-8.33
50	50	0.05	50.87	3	17	14.39	3	17	2.78	17	0.00	1.97	16	-5.88
50	50	0.1	195.57	4	14	50.22	4	14	6.42	14	0.00	2.00	12	-14.29
50	100	0.05	83.83	3	24	16.98	2	24	8.13	24	0.00	6.19	24	0.00
50	100	0.1	26.59	2	23	22.74	3	23	12.66	22	-4.35	6.92	22	-4.35
50	150	0.05	26.93	2	33	20.35	3	33	9.76	33	0.00	10.99	33	0.00
50	150	0.1	21.88	3	29	13.02	3	29	11.31	29	0.00	12.08	29	0.00
100	50	0.05	-	3	28	148.56	2	28	15.74	27	-3.57	4.64	27	-3.57
100	50	0.1	2019.72	4	23	42.88	2	23	147.91	25	8.70	9.56	24	4.35
100	100	0.05	592.19	2	40	122.96	2	40	15.64	40	0.00	9.80	39	-2.50
100	100	0.1	1310.52	3	35	108.59	3	35	30.20	35	0.00	49.53	36	2.86
100	150	0.05	439.65	3	46	183.10	4	46	0.19	INF	INF	18.07	46	0.00
100	150	0.1	1660.78	5	44	200.79	3	44	49.03	42	-4.55	65.74	43	-2.27
300	50	0.05	198.26	3	43	71.47	2	43	28.08	43	0.00	29.08	41	-4.65
300	50	0.1	248.80	4	38	93.53	3	38	31.69	35	-7.89	37.32	34	-10.53
300	100	0.05	-	2	61	1437.51	4	61	102.19	61	0.00	72.79	57	-6.56
300	100	0.1	-	2	54	796.72	3	54	89.33	52	-3.70	92.51	52	-3.70
300	150	0.05	-	1	-	-	3	78	192.23	78	0.00	135.22	76	-2.56
300	150	0.1	-	2	-	720.15	2	70	366.51	67	-4.29	145.31	69	-1.43
300	200	0.05	-	2	88	1349.81	3	88	212.94	86	-2.27	180.20	87	-1.14
300	200	0.1	-	2	-	1210.56	3	77	685.08	73	-5.19	276.36	74	-3.90
300	250	0.05	2650.57	3	114	787.82	3	114	283.98	113	-0.88	262.12	112	-1.75
300	250	0.1	-	2	-	1249.90	3	102	843.70	101	-0.98	474.56	100	-1.96
300	300	0.05	3434.26	3	121	1358.17	4	121	453.30	120	-0.83	336.46	120	-0.83
300	300	0.1	-	4	112	1989.73	6	112	717.31	108	-3.57	561.76	110	-1.79




Figure 2: Feasibility-ratio and optimality-ratio curves of SAA and IS as functions of sample size N

feasibility ratio and *optimality ratio*, and consider these two measures as functions of the sample size N . Specifically, for each N , the feasibility ratio consists on the number of replications that output feasible solutions divided by the number of total replications, and the optimality ratio consists on the number of replications that generate optimal solutions divided by the number of total replications. We can use Lemma 1 to verify the feasibility of the solution provided by SAA/IS, and check the optimality by comparing the objective value generated by SAA/IS with the optimal value produced by the OA approaches. Figure 2 illustrates feasibility-ratio and optimality-ratio curves of SAA/IS as functions of N .

Notice that the feasibility ratio and optimality ratio of SAA increase as the sample size N becomes larger. This makes sense as the larger the sample size N , the more likely SAA is to approximate the true distribution. Consequently, it becomes more likely for SAA to generate a feasible or optimal solution for CC-SMCP. This observation aligns with the theoretical findings previously presented in Theorem 7 and 8. However, practical experiments show that a smaller N can also yield a feasible or optimal solution.

For IS curves, the feasibility ratio and optimality ratio also increase with larger sample sizes N , but this effect is less pronounced compared to SAA curves. Moreover, it is important to note that the performance of IS is generally worse than that of SAA in almost all cases. This may be because IS is more sensitive to the choice of the importance sampling estimator \hat{p}_{ij} , which is crucial for the performance of IS.

Furthermore, we compute the average solution time of SAA and IS as well as their 95% confidence intervals, as illustrated in Figure 3. As we observe from Figure 3, the solution time for the IS at beginning iterations is similar to the solution time for SAA, but when the sample size N becomes large, IS becomes significantly faster than SAA. This may be because the IS method is more efficient in generating feasible solutions satisfying constraints (32b) rather than constraints (22c) in the SAA reformulation.

5.2 Additional Experiments

In addition to the above experiments, we also conduct several experiments on investigating the effect of risk parameter ϵ and the performance of methods for checking infeasibility and special cases, please find more details in Section C of the appendix to this paper.




Figure 3: The average solution time of the SAA and IS methods as a function of N , where the 95% confidence intervals are indicated as the shaded area

6 Conclusion

In this paper, we have investigated CC-SMCP with individual chance constraints under LHS uncertainty, where both decision and random variables are purely binary. We derived exact deterministic reformulations and proposed an outer-approximation (OA) algorithm for CC-SMCP using some combinatorial methods. This OA method has significantly alleviated computational challenges and reduced cumulative rounding errors. Additionally, we established statistical relationships between CC-SMCP and its sample average approximation (SAA) reformulation and theoretically demonstrated the effectiveness of the SAA method in generating optimality bounds and feasible solutions for CC-SMCP. We also studied the importance sampling (IS) method and obtained a sufficient condition for selecting the optimal IS estimator. Finally, several computational experiments were conducted to validate the effectiveness of OA method (with two different linearization techniques) and sampling-based approaches (SAA and IS).

References

- Ahmed S, Papageorgiou DJ (2013) Probabilistic set covering with correlations. *Operations Research* 61(2):438–452.
- Ahmed S, Xie W (2018) Relaxations and approximations of chance constraints under finite distributions. *Mathematical Programming* 170:43–65.
- Barrera J, Homem-de Mello T, Moreno E, Pagnoncelli BK, Canessa G (2016) Chance-constrained problems and rare events: an importance sampling approach. *Mathematical Programming* 157:153–189.
- Ben-Tal A, El Ghaoui L, Nemirovski A (2009) *Robust optimization*, volume 28 (Princeton university press).
- Beraldi P, Ruszczyński A (2002) The probabilistic set-covering problem. *Operations Research* 50(6):956–967.
- Bramel J, Simchi-Levi D (1997) On the effectiveness of set covering formulations for the vehicle routing problem with time windows. *Operations Research* 45(2):295–301.
- Bramel J, Simchi-Levi D (2002) Set-covering-based algorithms for the capacitated vrp. *The vehicle routing problem*, 85–108 (SIAM).
- Calafiore G, Campi MC (2005) Uncertain convex programs: randomized solutions and confidence levels. *Mathematical Programming* 102:25–46.
- Campbell AM, Thomas BW (2008) Probabilistic traveling salesman problem with deadlines. *Transportation Science* 42(1):1–21.
- De A (2017) Boolean function analysis meets stochastic optimization: An approximation scheme for stochastic knapsack. *arXiv preprint arXiv:1712.00918* .

- Dinh T, Fukasawa R, Luedtke J (2018) Exact algorithms for the chance-constrained vehicle routing problem. *Mathematical Programming* 172(1-2):105–138.
- Fischetti M, Monaci M (2012) Cutting plane versus compact formulations for uncertain (integer) linear programs. *Mathematical Programming Computation* 4(3):239–273.
- Glover F, Woolsey E (1974) Converting the 0-1 polynomial programming problem to a 0-1 linear program. *Operations research* 22(1):180–182.
- Grötschel M, Monma CL, Stoer M (1992) Computational results with a cutting plane algorithm for designing communication networks with low-connectivity constraints. *Operations Research* 40(2):309–330.
- Gunawardane G (1982) Dynamic versions of set covering type public facility location problems. *European Journal of Operational Research* 10(2):190–195.
- Haight RG, Revelle CS, Snyder SA (2000) An integer optimization approach to a probabilistic reserve site selection problem. *Operations research* 48(5):697–708.
- Hall NG, Hochbaum DS (1992) The multicovering problem. *European Journal of Operational Research* 62(3):323–339.
- Han J, Lee K, Lee C, Choi KS, Park S (2016) Robust optimization approach for a chance-constrained binary knapsack problem. *Mathematical Programming* 157:277–296.
- Hong Y (2013) On computing the distribution function for the poisson binomial distribution. *Computational Statistics & Data Analysis* 59:41–51.
- Huang R, Kim S, Menezes MB (2010) Facility location for large-scale emergencies. *Annals of Operations Research* 181:271–286.
- Kleywegt AJ, Shapiro A, Homem-de Mello T (2002) The sample average approximation method for stochastic discrete optimization. *SIAM Journal on optimization* 12(2):479–502.
- Klopfenstein O, Nace D (2008) A robust approach to the chance-constrained knapsack problem. *Operations Research Letters* 36(5):628–632.
- Küçükyavuz S, Jiang R (2022) Chance-constrained optimization under limited distributional information: A review of reformulations based on sampling and distributional robustness. *EURO Journal on Computational Optimization* 10:100030.
- Lejeune MA, Prékopa A (2018) Relaxations for probabilistically constrained stochastic programming problems: review and extensions. *Annals of Operations Research* 1–22.
- Lemus Rodriguez GJ (1999) *Portfolio optimization with quantile-based risk measures*. Ph.D. thesis, Massachusetts Institute of Technology.
- Lessin AM, Lunday BJ, Hill RR (2018) A bilevel exposure-oriented sensor location problem for border security. *Computers & Operations Research* 98:56–68.
- Luedtke J (2014) A branch-and-cut decomposition algorithm for solving chance-constrained mathematical programs with finite support. *Mathematical Programming* 146(1-2):219–244.
- Luedtke J, Ahmed S (2008) A sample approximation approach for optimization with probabilistic constraints. *SIAM Journal on Optimization* 19(2):674–699.
- Mitzenmacher M, Upfal E (2005) *Probability and Computing: Randomized Algorithms and Probabilistic Analysis* (Cambridge University Press), ISBN 9780521835404, URL <https://books.google.com/books?id=0bAY16d7hvkC>.
- Nemirovski A, Shapiro A (2007) Convex approximations of chance constrained programs. *SIAM Journal on Optimization* 17(4):969–996.
- Özener ÖÖ, Ergun Ö, Savelsbergh M (2013) Allocating cost of service to customers in inventory routing. *Operations Research* 61(1):112–125.

- Padberg M, Rinaldi G (1989) A branch-and-cut approach to a traveling salesman problem with side constraints. *Management Science* 35(11):1393–1412.
- Prudnikov A, Brychkov YA, Integrals OM (1998) Series: Volume 2 special functions.
- Rubino G, Tuffin B, et al. (2009) *Rare event simulation using Monte Carlo methods*, volume 73 (Wiley Online Library).
- Saxena A, Goyal V, Lejeune MA (2010) Mip reformulations of the probabilistic set covering problem. *Mathematical programming* 121(1):1–31.
- Shen H, Jiang R (2022) Chance-constrained set covering with wasserstein ambiguity. *Mathematical Programming* 1–54.
- Smith BM, Wren A (1988) A bus crew scheduling system using a set covering formulation. *Transportation Research Part A: General* 22(2):97–108.
- Soltani NY, Kim SJ, Giannakis GB (2013) Chance-constrained optimization of ofdma cognitive radio uplinks. *IEEE transactions on wireless communications* 12(3):1098–1107.
- Song Y, Luedtke JR, Küçükyavuz S (2014) Chance-constrained binary packing problems. *INFORMS Journal on Computing* 26(4):735–747.
- Stanley RP (2011) *Enumerative Combinatorics: Volume 1* (USA: Cambridge University Press), 2nd edition, ISBN 1107602629.
- Sun O, Fan N (2019) The probabilistic and reliable connected power dominating set problems. *Optimization Letters* 13(5):1189–1206.
- Tanner MW, Ntaimo L (2010) Iis branch-and-cut for joint chance-constrained programs with random technology matrices. *Eur. J. Oper. Res* 207(1):290–296.
- Van Ackooij W, Zorgati R, Henrion R, Möller A (2011) Chance constrained programming and its applications to energy management. *Stochastic Optimization-Seeing the Optimal for the Uncertain* 291–320.
- Vasko FJ, Wolf FE, Stott Jr KL (1989) A set covering approach to metallurgical grade assignment. *European Journal of Operational Research* 38(1):27–34.
- Wang J (2007) The β -reliable median on a network with discrete probabilistic demand weights. *Operations research* 55(5):966–975.
- Wang S, Li J, Mehrotra S (2021) Chance-constrained multiple bin packing problem with an application to operating room planning. *INFORMS Journal on Computing* 33(4):1661–1677.
- Wang S, Li J, Mehrotra S (2022) A solution approach to distributionally robust joint-chance-constrained assignment problems. *INFORMS Journal on Optimization* 4(2):125–147.
- Wu HH, Küçükyavuz S (2019) Probabilistic partial set covering with an oracle for chance constraints. *SIAM Journal on Optimization* 29(1):690–718.
- Yoda K, Prékopa A (2016) Convexity and solutions of stochastic multidimensional 0-1 knapsack problems with probabilistic constraints. *Mathematics of Operations Research* 41(2):715–731.
- Zhang Z, Denton BT, Xie X (2020) Branch and price for chance-constrained bin packing. *INFORMS Journal on Computing* 32(3):547–564.

Appendices

A Preliminaries

A.1 Inclusion–exclusion principle

Lemma A.1.1 (Combinatorial version) For finite sets A_1, \dots, A_n , one has the identity

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{\emptyset \neq S \subseteq [n]} (-1)^{|S|+1} \left| \bigcap_{j \in S} A_j \right|. \quad (1)$$

Lemma A.1.2 (Probabilistic version) For events A_1, \dots, A_n in a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, one has the identity

$$\mathbb{P} \left(\bigcup_{i=1}^n A_i \right) = \sum_{k=1}^n \left((-1)^{k-1} \sum_{\substack{I \subseteq [n] \\ |I|=k}} \mathbb{P} \left(\bigcap_{i \in I} A_i \right) \right) \quad (2)$$

A.2 Bonferroni inequalities

For events A_1, \dots, A_n in a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, we define

$$S_1 := \sum_{i=1}^n \mathbb{P}(A_i), \quad S_2 := \sum_{1 \leq i_1 < i_2 \leq n} \mathbb{P}(A_{i_1} \cap A_{i_2}), \dots, \quad S_k := \sum_{1 \leq i_1 < \dots < i_k \leq n} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k})$$

for all integers k in $\{1, \dots, n\}$. Then we have the following inequalities:

Lemma A.2.1 For odd $k \in \{1, \dots, n\}$,

$$\sum_{j=1}^k (-1)^{j-1} S_j \geq \mathbb{P} \left(\bigcup_{i=1}^n A_i \right) = \sum_{j=1}^n (-1)^{j-1} S_j \quad (3)$$

Lemma A.2.2 For even $k \in \{2, \dots, n\}$,

$$\sum_{j=1}^k (-1)^{j-1} S_j \leq \mathbb{P} \left(\bigcup_{i=1}^n A_i \right) = \sum_{j=1}^n (-1)^{j-1} S_j \quad (4)$$

A.3 Multiplicative Chernoff bound

Lemma A.3.1 (Mitzenmacher and Upfal (2005)) Let $Y \sim \text{Binomial}(n; p)$ and $\mu = \mathbb{E}[Y]$. For any $\delta \geq 0$,

$$\mathbb{P}[Y \geq (1 + \delta)\mu] \leq \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}} \right)^\mu \leq \exp \left(\frac{-\delta^2 \mu}{2 + \delta} \right).$$

A.4 Hoeffding's inequality

Lemma A.4.1 (Hoeffding (1994)) Let Y_1, \dots, Y_N be independent random variables such that $a_i \leq Y_i \leq b_i$ almost surely. Then for all $t > 0$,

$$\mathbb{P} \left[\sum_{i=1}^N (Y_i - \mathbb{E}[Y_i]) \geq t \right] \leq \exp \left\{ - \frac{2t^2}{\sum_{i=1}^N (b_i - a_i)^2} \right\}.$$

B Proofs

B.1 Proof of Lemma 1

Note that $\mathbb{P}[A_j(x)] = \mathbb{P}[\tilde{a}_j x_j = 1] = x_j p_j$ because $x_j \in \{0, 1\}$. We first derive a closed-form expression for the probability $\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j = k\right]$, which can be regarded as the probability of exactly k out of n events $A_1(x), \dots, A_n(x)$ occurring. We have

$$\begin{aligned} \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j = k\right] &= \sum_{\substack{S \subseteq [n] \\ |S|=k}} \mathbb{P}[A_S(x) \cap \bigcap_{j \in S^c} A_j^c(x)] = \sum_{\substack{S \subseteq [n] \\ |S|=k}} \mathbb{P}[A_S(x)] (1 - \mathbb{P}[\bigcup_{j \in S^c} A_j(x)]) \\ &= \sum_{\substack{S \subseteq [n] \\ |S|=k}} \left(\mathbb{P}[A_S(x)] - \mathbb{P}\left[\bigcup_{j \in S^c} A_{S \cup \{j\}}(x)\right] \right), \end{aligned}$$

where the last two equalities follow from independence. Moreover, by the inclusion-exclusion principle, the probability of finite unions of events

$$\begin{aligned} \mathbb{P}\left[\bigcup_{j \in S^c} A_{S \cup \{j\}}(x)\right] &= \sum_{j_1 \in S^c} \mathbb{P}[A_{S \cup \{j_1\}}(x)] - \sum_{\substack{j_1 < j_2 \\ j_1, j_2 \in S^c}} \mathbb{P}[A_{S \cup \{j_1, j_2\}}(x)] + \dots + (-1)^{n-|S|-1} \mathbb{P}[A_{[n]}(x)] \\ &= \sum_{S \subsetneq T \subseteq [n]} (-1)^{|T|-|S|-1} \mathbb{P}[A_T(x)]. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j = k\right] &= \sum_{\substack{S \subseteq [n] \\ |S|=k}} \left(\mathbb{P}[A_S(x)] - \sum_{S \subsetneq T \subseteq [n]} (-1)^{|T|-|S|-1} \mathbb{P}[A_T(x)] \right) = \sum_{\substack{S \subseteq [n] \\ |S|=k}} \sum_{S \subsetneq T \subseteq [n]} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \\ &= \sum_{\ell=k}^n \sum_{\substack{S \subseteq [n] \\ |S|=k}} \sum_{\substack{S \subsetneq T \subseteq [n] \\ |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] = \sum_{\ell=k}^n (-1)^{\ell-k} \binom{\ell}{k} h_\ell(x), \end{aligned} \quad (5)$$

where $h_\ell(x) = \sum_{\substack{T \subseteq [n] \\ |T|=\ell}} \mathbb{P}[A_T(x)] = \sum_{\substack{T \subseteq [n] \\ |T|=\ell}} \prod_{j \in T} x_j p_j$, for all $\ell = k, \dots, n$. And the last equality holds because for each ℓ , $h_\ell(x)$ is counted $\binom{\ell}{k}$ times. Then we derive the closed-form expression for the probability $\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j \geq k\right]$, that is, the probability of at least k out of n events $A_1(x), \dots, A_n(x)$ occurring:

$$\begin{aligned} \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j \geq k\right] &= \sum_{d=k}^n \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j = d\right] = \sum_{d=k}^n \sum_{\ell=d}^n (-1)^{\ell-d} \binom{\ell}{d} h_\ell(x) = \sum_{\ell=k}^n \sum_{d=k}^{\ell} (-1)^{\ell-d} \binom{\ell}{d} h_\ell(x) \\ &= \sum_{\ell=k}^n (-1)^{\ell-k} \sum_{d=k}^{\ell} (-1)^{d-k} \binom{\ell}{d} h_\ell(x) = \sum_{\ell=k}^n (-1)^{\ell-k} \binom{\ell-1}{\ell-k} h_\ell(x), \end{aligned}$$

where the last equality follows from the combinatorial identity $(-1)^{k-1} \binom{\ell-1}{k-1} = \sum_{d=0}^{k-1} (-1)^d \binom{\ell}{d} = -\sum_{d=k}^{\ell} (-1)^d \binom{\ell}{d}$.

B.2 Proof of Lemma 4

$X_{i_1} \subseteq X_{i_2}$ follows from

$$\mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{i_2 j} x_j \geq k_{i_2} \right] = \mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{i_1 j} x_j \geq k_{i_2} \right] \geq \mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{i_1 j} x_j \geq k_{i_1} \right] \geq 1 - \epsilon_{i_1} \geq 1 - \epsilon_{i_2},$$

for any $x \in X_{i_1}$.

B.3 Proof of Lemma 5

Similar to the proof of Lemma 4, we have

$$\mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{i_2 j} x_j \geq k_{i_2} \right] \geq \mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{i_1 j} x_j \geq k_{i_2} \right] \geq \mathbb{P} \left[\sum_{j \in [n]} \tilde{a}_{i_1 j} x_j \geq k_{i_1} \right] \geq 1 - \epsilon_{i_1} \geq 1 - \epsilon_{i_2},$$

for any $x \in X_{i_1}$. Thus, $X_{i_1} \subseteq X_{i_2}$.

B.4 Proof of Theorem 2

From the proof of Lemma 1, for any fixed $S \subseteq [n]$ with $|S| = d$, we have

$$\mathbb{P}[A_S(x) \cap A_j^c(x)] = \mathbb{P}[A_S(x)] - \mathbb{P} \left[\bigcup_{j \in S^c} A_{S \cup \{j\}}(x) \right] = \sum_{S \subseteq T \subseteq [n]} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)].$$

Applying Bonferroni inequalities to the probability $\mathbb{P} \left[\bigcup_{j \in S^c} A_{S \cup \{j\}}(x) \right]$ yields

$$\begin{aligned} \mathbb{P}[A_S(x) \cap A_j^c(x)] &= \sum_{S \subseteq T \subseteq [n]} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] = \sum_{\ell=d}^{d+n} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \\ &\leq \sum_{\ell=d}^{d+t} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)], t \text{ is even} \end{aligned}$$

and

$$\mathbb{P}[A_S(x) \cap A_j^c(x)] = \sum_{\ell=d}^{d+n} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \geq \sum_{\ell=d}^{d+t} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)], t \text{ is odd}$$

Moreover, if the inequality

$$\sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} \mathbb{P}[A_T(x)] \geq \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell+1}} \mathbb{P}[A_T(x)],$$

holds for any fixed $S \subseteq [n]$ with $|S| = d \leq \ell < n$, then we have

$$\mathbb{P}[A_S(x) \cap A_j^c(x)] = \sum_{\ell=d}^{d+n} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \leq \dots \leq \sum_{\ell=d}^{d+2} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \quad (6a)$$

$$\leq \sum_{\ell=d}^d \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \quad (6b)$$

and

$$\begin{aligned} \mathbb{P}[A_S(x) \cap_{j \in S^c} A_j^c(x)] &= \sum_{\ell=d}^{d+n} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \geq \cdots \geq \sum_{\ell=d}^{d+3} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \\ &\geq \sum_{\ell=d}^{d+1} \sum_{\substack{S \subseteq T \subseteq [n] \\ |S|=d, |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \end{aligned}$$

Taking the summations on both sides of inequalities (6), we obtain

$$\begin{aligned} \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j \leq k-1\right] &= \sum_{d=0}^{k-1} \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j = d\right] = \sum_{d=0}^{k-1} \sum_{\substack{S \subseteq [n] \\ |S|=d}} \mathbb{P}\left[A_S(x) \cap_{j \in S^c} A_j^c(x)\right] \\ &= \sum_{d=0}^{k-1} \sum_{\substack{S \subseteq [n] \\ |S|=d}} \sum_{\ell=d}^{d+n} \sum_{\substack{S \subseteq T \subseteq [n] \\ |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \leq \cdots \leq \sum_{d=0}^{k-1} \sum_{\substack{S \subseteq [n] \\ |S|=d}} \sum_{\ell=d}^{d+2} \sum_{\substack{S \subseteq T \subseteq [n] \\ |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \\ &\leq \sum_{d=0}^{k-1} \sum_{\substack{S \subseteq [n] \\ |S|=d}} \sum_{\ell=d}^d \sum_{\substack{S \subseteq T \subseteq [n] \\ |T|=\ell}} (-1)^{|T|-|S|} \mathbb{P}[A_T(x)] \end{aligned}$$

After some algebra (similar as proof in eq. (5) of Lemma 1), we conclude that

$$\begin{aligned} \mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j \leq k-1\right] &= \sum_{d=0}^{k-1} \sum_{\ell=d}^{d+n} (-1)^{\ell-d} \binom{\ell}{d} h_{\ell}(x) \leq \cdots \leq \sum_{d=0}^{k-1} \sum_{\ell=d}^{d+2} (-1)^{\ell-d} \binom{\ell}{d} h_{\ell}(x) \\ &\leq \sum_{d=0}^{k-1} \sum_{\ell=d}^d (-1)^{\ell-d} \binom{\ell}{d} h_{\ell}(x) \end{aligned}$$

which is equivalent to

$$\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j \leq k-1\right] = g_n(x) \leq \cdots \leq g_4(x) \leq g_2(x) \leq g_0(x),$$

and the sequence of inequalities

$$\mathbb{P}\left[\sum_{j=1}^n \tilde{a}_j x_j \leq k-1\right] = g_n(x) \geq \cdots \geq g_5(x) \geq g_3(x) \geq g_1(x)$$

holds similarly.

B.5 Proof of Lemma 3

For any fixed d such that $0 \leq d \leq k-1$, we consider the difference between $g_{t+2,d}(x)$ and $g_{t,d}(x)$. Since $g_{t+2,d}(x) - g_{t,d}(x) \geq 0$ for some odd t , we have

$$\begin{aligned} g_{t+2,d}(x) - g_{t,d}(x) &= \binom{d+t+1}{d} \left(\sum_{\substack{S \subseteq [n] \\ |S|=d+t+1}} \prod_{j \in S} x_j p_j \right) - \frac{d+t+2}{t+2} \binom{d+t+1}{d} \sum_{\substack{S \subseteq [n] \\ |S|=d+t+2}} \prod_{j \in S} x_j p_j \\ &= \binom{d+t+1}{d} \left(\sum_{\substack{S \subseteq [n] \\ |S|=d+t+1}} \prod_{j \in S} x_j p_j \right) \left(1 - \frac{d+t+2}{t+2} \left(\sum_{\substack{q \in [n] \setminus S \\ |S|=d+t+1}} x_q p_q \right) \right) \geq 0. \end{aligned}$$

It follows that $1 - \frac{d+t+2}{t+2} \left(\sum_{\substack{q \in [n] \setminus S \\ |S|=d+t+1}} x_q p_q \right) \geq 0$. Hence, for any odd τ such that $t \leq \tau \leq n-2$, we have

$$\begin{aligned} g_{\tau+2,d}(x) - g_{\tau,d}(x) &= \binom{d+\tau+1}{d} \left(\sum_{\substack{S \subseteq [n] \\ |S|=d+\tau+1}} \prod_{j \in S} x_j p_j \right) \left(1 - \frac{d+\tau+2}{\tau+2} \left(\sum_{\substack{q \in [n] \setminus S \\ |S|=d+\tau+1}} x_q p_q \right) \right) \\ &\geq \binom{d+\tau+1}{d} \left(\sum_{\substack{S \subseteq [n] \\ |S|=d+\tau+1}} \prod_{j \in S} x_j p_j \right) \left(1 - \frac{d+t+2}{t+2} \left(\sum_{\substack{q \in [n] \setminus S \\ |S|=d+t+1}} x_q p_q \right) \right) \geq 0. \end{aligned}$$

For even τ , the statement holds similarly.

B.6 Proof of Theorem 3

From Lemma 3, it follows that such t_{\min}^{odd} and t_{\min}^{even} exist. Therefore, for any odd τ such that $t_{\min}^{\text{odd}} \leq \tau \leq n-2$, we have $g_{\tau+2}(x) - g_{\tau}(x) = \sum_{d=0}^{k-1} (g_{\tau+2,d}(x) - g_{\tau,d}(x)) \geq 0$. For even τ , the statement holds similarly.

B.7 Proof of Lemma 6

We denote the indices of sets in the corresponding candidate solution \bar{x} by $D := \{j : \bar{x}_j = 1, j \in [n]\}$. Then

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] = \mathbb{P} \left[\sum_{j \in D} \tilde{a}_j \geq k \right].$$

Note that the random variable $\sum_{j \in D} \tilde{a}_j \sim \text{Binomial}(d, p)$ since a_j 's are independent Bernoulli random variables with the same success probability p . Thus,

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] = \sum_{\ell=k}^d \mathbb{P} \left[\sum_{j \in D} \tilde{a}_j = \ell \right] = \sum_{\ell=k}^d \binom{d}{\ell} p^{\ell} (1-p)^{d-\ell}.$$

B.8 Proof of Lemma 7

We consider the extended binomial coefficients by allowing α to be an arbitrary number (negative, rational, real), expressed as

$$\binom{\alpha}{k} := \frac{\alpha(\alpha-1)(\alpha-2)\cdots(\alpha-k+1)}{k(k-1)(k-2)\cdots 1} \quad \text{for } k \in \mathbb{N} \text{ and } \alpha \in \mathbb{R}.$$

With this definition one has a generalization of the binomial formula

$$(x+y)^\alpha = \sum_{\ell=0}^{+\infty} \binom{\alpha}{\ell} x^\ell y^{\alpha-\ell}, \quad \text{for any } x, y \in (0, 1) \text{ and } \alpha \in \mathbb{R}.$$

We still use $D := \{j : \bar{x}_j = 1, j \in [n]\}$ to denote the indices of sets in the corresponding candidate solution \bar{x} , and use the rising Pochhammer symbol $q^{(n)} = \binom{q+n-1}{n} n!$ in the proof. By Lemma 1, we have

$$\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] = \sum_{\ell=k}^n (-1)^{\ell-k} \binom{\ell-1}{\ell-k} \sum_{\substack{S \subseteq [n] \\ |S|=\ell}} \prod_{j \in S} \bar{x}_j p_j = \sum_{\ell=k}^d (-1)^{\ell-k} \binom{\ell-1}{\ell-k} \sum_{\substack{S \subseteq D \\ |S|=\ell}} \prod_{j \in S} \bar{x}_j p_j \quad (10)$$

$$= \sum_{\ell=k}^d (-1)^{\ell-k} \binom{\ell-1}{\ell-k} \binom{d}{\ell} p^\ell = \sum_{\ell=k}^d (-1)^{\ell-k} \binom{\ell-1}{k-1} \binom{d}{\ell-1} \frac{d-\ell+1}{\ell} p^\ell \quad (11)$$

$$= \sum_{\ell=k}^d (-1)^{\ell-k} \binom{d}{k-1} \binom{d-k+1}{d-\ell+1} \frac{d-\ell+1}{\ell} p^\ell \quad (12)$$

$$= k \binom{d}{k} \sum_{\ell=k}^d (-1)^{\ell-k} \binom{d-k}{\ell-k} \frac{p^\ell}{\ell} = k \binom{d}{k} \sum_{t=0}^{d+k} (-1)^t \binom{d-k}{t} \frac{p^{k+t}}{k+t} \quad (*)$$

$$= \binom{d}{k} p^k \sum_{t=0}^{+\infty} \binom{t+k-d-1}{t} \frac{kp^t}{k+t} = \binom{d}{k} p^k \sum_{t=0}^{+\infty} (k-d)^{(t)} \cdot \frac{k}{k+t} \cdot \frac{p^t}{t!} \quad (13)$$

$$= \binom{d}{k} p^k \sum_{t=0}^{+\infty} (k-d)^{(t)} \cdot \frac{k^{(t)}}{(k+1)^{(t)}} \cdot \frac{p^t}{t!} = \binom{d}{k} p^k {}_2F_1(k-d, k; k+1; p) \quad (**)$$

$$= I_p(k, d-k+1) \quad (***)$$

where the equation (10) follows from the definition of D , the equation (11) follows from the combinatorial identity $\binom{n}{k} = \frac{n}{k} \binom{n-1}{k-1}$, the equation (12) follows from $\binom{n}{h} \binom{n-h}{k} = \binom{n}{k} \binom{n-k}{h}$, the equation (13) follows from $(-1)^k \binom{n}{k} = \binom{k-n-1}{k}$ and the definition of the rising Pochhammer symbol $q^{(n)}$, the equation (***) follows from the fact $I_x(a, b) = x^a {}_2F_1(a, 1-b; a+1, x)/(aB(a, b))$ and the equations (*), (**) and (***) indicate the conclusion.

Moreover, letting

$$h(u) := k \binom{d}{k} \sum_{\ell=k}^d (-1)^{\ell-k} \binom{d-k}{\ell-k} \frac{u^\ell}{\ell} = k \binom{d}{k} \sum_{t=0}^{+\infty} (-1)^t \binom{d-k}{t} \frac{u^{k+t}}{k+t},$$

we have $h(0) = 0$ and note that

$$\frac{dh}{du} = k \binom{d}{k} u^{k-1} \sum_{t=0}^{+\infty} (-1)^t \binom{d-k}{t} u^t = k \binom{d}{k} u^{k-1} (1-u)^{d-k}.$$

Hence,

$$k \binom{d}{k} \int_0^p u^{k-1} (1-u)^{d-k} du = h(p) - h(0) = h(p),$$

which completes our proof.

B.9 Proof of Lemma 8

We first study the monotonicity of the CDF of a binomial distribution. Let $F(k; n, p)$ be the CDF of a binomial distribution. Then for any fixed k and p , the function $F(k; n, p)$ is monotone nonincreasing with respect to n , where $n \in \mathbb{N}$. In fact, by the law of total probability,

$$\begin{aligned}
F(k; n+1, p) &= \mathbb{P} \left[\sum_{j=1}^{n+1} \tilde{a}_j \leq k \right] \\
&= \mathbb{P} \left[\sum_{j=1}^{n+1} \tilde{a}_j \leq k \mid \sum_{j=1}^n \tilde{a}_j \leq k-1 \right] \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \leq k-1 \right] + \mathbb{P} \left[\sum_{j=1}^{n+1} \tilde{a}_j \leq k \mid \sum_{j=1}^n \tilde{a}_j \geq k \right] \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \geq k \right] \\
&= \mathbb{P} \left[\sum_{j=1}^{n+1} \tilde{a}_j \leq k \mid \sum_{j=1}^n \tilde{a}_j \leq k-1 \right] \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \leq k-1 \right] + \mathbb{P} \left[\sum_{j=1}^{n+1} \tilde{a}_j \leq k \mid \sum_{j=1}^n \tilde{a}_j = k \right] \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j = k \right] \\
&= 1 \cdot \sum_{\ell=0}^{k-1} \binom{n}{\ell} p^\ell (1-p)^{n-\ell} + (1-p) \cdot \binom{n}{k} p^k (1-p)^{n-k} = \sum_{\ell=0}^k \binom{n}{\ell} p^\ell (1-p)^{n-\ell} - \binom{n}{k} p^{k+1} (1-p)^{n-k} \\
&= \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \leq k \right] - \binom{n}{k} p^{k+1} (1-p)^{n-k} \leq F(k; n, p).
\end{aligned}$$

Let $D := \{j : \bar{x}_j = 1, j \in [n]\}$ and $\hat{D} := \{j : \hat{x}_j = 1, j \in [n]\}$. It is equivalent to prove $\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \hat{x}_j \geq k \right] \geq 1 - \epsilon$. Note that the cover probability

$$\begin{aligned}
\mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \hat{x}_j \geq k \right] &= \mathbb{P} \left[\sum_{j \in \hat{D}} \tilde{a}_j \geq k \right] = \mathbb{P} \left[\sum_{j=1}^{\hat{d}} \tilde{a}_j \geq k \right] \quad (\tilde{a}_j \text{'s are i.i.d random variables}) \\
&\geq \mathbb{P} \left[\sum_{j=1}^d \tilde{a}_j \geq k \right] = \mathbb{P} \left[\sum_{j \in D} \tilde{a}_j \geq k \right] \quad (\text{by monotone property of } F(k; n, p)) \\
&= \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] \geq 1 - \epsilon. \quad (\bar{x} \text{ is a feasible solution})
\end{aligned}$$

This completes our proof.

B.10 Proof of Proposition 2

First notice that $d \geq k$ since we cannot use less than k sets to cover the item at least k times with a predefined probability. Furthermore, by the Markov's inequality, we have

$$1 - \epsilon \leq \mathbb{P} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \geq k \right] \leq \frac{\mathbb{E} \left[\sum_{j=1}^n \tilde{a}_j \bar{x}_j \right]}{k} = \frac{\sum_{j=1}^n \bar{x}_j \mathbb{E} [\tilde{a}_j]}{k} = \frac{dp}{k}.$$

Therefore, $d := \sum_{j \in [n]} \bar{x}_j \geq \max \left\{ k, \left\lceil \frac{k(1-\epsilon)}{p} \right\rceil \right\}$.

B.11 Proof of Theorem 6

Let x^* be an optimal solution to CC-SMCP. Then $q_i(x^*) := \mathbb{P}[\tilde{A}_i(\omega)x^* \leq k_i - 1] \leq \epsilon_i$ for each $i \in [m]$. By the definition of X_α^N , $x^* \in X_\alpha^N$ if and only if no more than $\lfloor \alpha_i N \rfloor$ times the event $\{\tilde{A}_i(\omega)x^* \leq k_i - 1\}$

happens in N trials for each $i \in [m]$. That is, for any $i \in [m]$,

$$\sum_{\omega \in \Omega} \mathbb{I}(\tilde{A}_i(\omega)x \geq k_i) \geq (1 - \alpha_i)N \Leftrightarrow \sum_{\omega \in \Omega} \mathbb{I}(\tilde{A}_i(\omega)x \leq k_i - 1) \leq \lfloor \alpha_i N \rfloor.$$

Also, note that if $x^* \in X_\alpha^N$, then $\nu_\alpha^N \leq \nu^*$. Let E_i be the event that $\{\tilde{A}_i(\omega)x^* \leq k_i - 1\}$ happens at most $\lfloor \alpha_i N \rfloor$ times. By Boole's inequality (union bound), we have

$$\mathbb{P}[\nu_\alpha^N \leq \nu_\epsilon^*] \geq \mathbb{P}[x^* \in X_\alpha^N] = \mathbb{P}[\cap_{i=1}^m E_i] \geq 1 - \sum_{i=1}^m (1 - \mathbb{P}[E_i]).$$

Moreover, for any $i \in [m]$,

$$\mathbb{P}[E_i] = \sum_{\ell=0}^{\lfloor \alpha_i N \rfloor} \binom{N}{\ell} [q_i(x^*)]^\ell [1 - q_i(x^*)]^{N-\ell} \geq \sum_{\ell=0}^{\lfloor \alpha_i N \rfloor} \binom{N}{\ell} \epsilon_i^\ell (1 - \epsilon_i)^{N-\ell}$$

where the inequality follows from the fact that the CDF of the binomial distribution, $F(k; n, p)$, is non-increasing with respect to p . Therefore,

$$\mathbb{P}[\nu_\alpha^N \leq \nu_\epsilon^*] \geq 1 - \sum_{i=1}^m \sum_{\ell=\lfloor \alpha_i N \rfloor + 1}^N \binom{N}{\ell} \epsilon_i^\ell (1 - \epsilon_i)^{N-\ell} = 1 - \sum_{i=1}^m I_{\epsilon_i}(\lfloor \alpha_i N \rfloor + 1, N - \lfloor \alpha_i N \rfloor).$$

Moreover, we define Y_i to be a binomial random variable with parameters (N, ϵ_i) . Using the Chernoff Bound in Theorem A.3.1, we obtain if $\epsilon_i \leq \alpha_i$ for some $i \in [m]$, then

$$\mathbb{P}[Y_i \geq \alpha_i N] = \mathbb{P}\left[Y_i \geq \epsilon_i N \left(1 + \left(\frac{\alpha_i}{\epsilon_i} - 1\right)\right)\right] \leq \exp\left(\frac{-(\alpha_i - \epsilon_i)^2 N}{\alpha_i + \epsilon_i}\right) \leq \exp(-\kappa_1 N).$$

where $\kappa_1 := \min_{i \in [m]} \{(\alpha_i - \epsilon_i)^2 / (\alpha_i + \epsilon_i)\}$. Therefore, if $\epsilon_i \leq \alpha_i$ for each $i \in [m]$, then

$$\mathbb{P}[\nu_\alpha^N \leq \nu_\epsilon^*] \geq 1 - \sum_{i=1}^m \mathbb{P}[Y_i \geq \alpha_i N] \geq 1 - \sum_{i=1}^m \exp\left(\frac{-(\alpha_i - \epsilon_i)^2 N}{\alpha_i + \epsilon_i}\right) \geq 1 - m \exp(-\kappa_1 N).$$

B.12 Proof of Theorem 7

We define the indicator random variable $Y_{i,\omega}$ by $Y_{i,\omega} = 1$ if $\tilde{A}_i(\omega)x \geq k_i$ and $Y_{i,\omega} = 0$ otherwise. Recall that $X_\epsilon := \cap_{i \in [m]} X_{\epsilon,i}$ where $X_{\epsilon,i} := \left\{x \in B : \mathbb{P}[\tilde{A}_i x \geq k_i] \geq 1 - \epsilon_i\right\}$. Then for any $x \in B \setminus X_{\epsilon,i}$, we obtain $\mathbb{E}[Y_{i,\omega}] = \mathbb{P}[\tilde{A}_i(\omega)x \geq k_i] < 1 - \epsilon_i$. Let

$$X_{\alpha,i}^N := \left\{x \in B : \frac{1}{N} \sum_{\omega \in \Omega} \mathbb{I}(\tilde{A}_i(\omega)x \geq k_i) \geq 1 - \alpha_i\right\}, \forall i \in [m].$$

Note that $X_\alpha^N = \cap_{i \in [m]} X_{\alpha,i}^N$ and $x \in X_{\alpha,i}^N$ if and only if $(1/N) \sum_{\omega \in \Omega} Y_{i,\omega} \geq 1 - \alpha_i$. Hoeffding's inequality yields, for any $x \in B \setminus X_{\epsilon,i}$,

$$\mathbb{P}[x \in X_{\alpha,i}^N] = \mathbb{P}\left[\frac{1}{N} \sum_{\omega \in \Omega} Y_{i,\omega} \geq 1 - \alpha_i\right] \leq \mathbb{P}\left[\sum_{\omega \in \Omega} (Y_{i,\omega} - \mathbb{E}[Y_{i,\omega}]) \geq N(\epsilon_i - \alpha_i)\right] \leq \exp\{-2N(\epsilon_i - \alpha_i)^2\}.$$

Therefore, we can obtain an upper bound for the probability

$$\mathbb{P}[X_\alpha^N \not\subseteq X_\epsilon] = \mathbb{P}[\exists x \in X_\alpha^N \text{ where } x \in B \setminus X_\epsilon] = \mathbb{P}\left[\bigcup_{x \in B \setminus X_\epsilon} \{x \in X_\alpha^N\}\right]$$

$$\begin{aligned}
&= \mathbb{P} \left[\bigcup_{i \in [m]} \bigcup_{x \in B \setminus X_{\epsilon,i}} \left\{ x \in \bigcap_{i \in [m]} X_{\alpha,i}^N \right\} \right] \leq \sum_{i \in [m]} \sum_{x \in B \setminus X_{\epsilon,i}} \mathbb{P} [x \in X_{\alpha,i}^N] \\
&\leq \sum_{i \in [m]} \sum_{x \in B \setminus X_{\epsilon,i}} \exp \{ -2N(\epsilon_i - \alpha_i)^2 \} \leq m|B \setminus X_{\epsilon}| \exp \{ -\kappa_2 N \}
\end{aligned}$$

where $\kappa_2 := 2 \min_{i \in [m]} (\epsilon_i - \alpha_i)^2$. This completes our proof.

B.13 Proof of Theorem 8

We will use the following lemma in the proof:

Lemma B.13.1 *Let X_{ϵ} be the feasible region of CC-SMCP. Then*

$$X_{\epsilon} = X'_{\bar{\alpha}},$$

where $X'_{\bar{\alpha}} := \bigcap_{i \in [m]} X'_{\bar{\alpha},i}$ and $X'_{\bar{\alpha},i} := \left\{ x \in B : \mathbb{P}[\tilde{A}_i x \leq k_i - 1] < \bar{\alpha}_i \right\}$ for each $i \in [m]$.

Proof (of Lemma B.13.1). We prove that $X_{\epsilon,i} = X'_{\bar{\alpha},i}$ for each $i \in [m]$. Clearly, $X_{\epsilon,i} \subseteq X'_{\bar{\alpha},i}$ for any $i \in [m]$ since $\mathbb{P}[\tilde{A}_i x \leq k_i - 1] \leq \epsilon_i < \bar{\alpha}_i$ for every $x \in X_{\epsilon,i}$. Conversely, if $x \in X'_{\bar{\alpha},i}$ for a fixed i , then by the definition of $\bar{\alpha}_i$, we have $x \in X_{\epsilon,i}$. ■

Proof (of Theorem 8). By the definition of $\underline{\alpha}_i$, we have $\nu_{\underline{\alpha}}^* = \nu_{\epsilon}^*$. Theorem 6 implies

$$\mathbb{P} [\nu_{\epsilon}^N > \nu_{\epsilon}^*] = \mathbb{P} [\nu_{\epsilon}^N > \nu_{\underline{\alpha}}^*] \leq m \exp \left(- \min_{i \in [m]} \left\{ \frac{(\underline{\alpha}_i - \epsilon_i)^2 N}{(\underline{\alpha}_i + \epsilon_i)} \right\} \right).$$

Further, observe that the proof of Theorem 7 can be modified to show the slightly stronger result:

$$\mathbb{P} [X_{\alpha}^N \subseteq X'_{\epsilon}] \geq 1 - m|B \setminus X'_{\epsilon}| \exp \left\{ -2N \min_{i \in [m]} (\epsilon_i - \alpha_i)^2 \right\},$$

where $X'_{\epsilon} := \bigcap_{i \in [m]} X'_{\epsilon,i}$ and $X'_{\epsilon,i} := \left\{ x \in B : \mathbb{P}[\tilde{A}_i x \leq k_i - 1] < \epsilon_i \right\}$ for each $i \in [m]$. (In the proof, we consider each $x \in X \setminus X'_{\epsilon}$ and replace X_{ϵ} with X'_{ϵ} . The remainder of the proof is identical to the original proof.) Applying this result yields

$$\mathbb{P} [X_{\epsilon}^N \subseteq X'_{\bar{\alpha}}] \geq 1 - m|B \setminus X'_{\bar{\alpha}}| \exp \left\{ -2N \min_{i \in [m]} (\bar{\alpha}_i - \epsilon_i)^2 \right\}.$$

Besides, if $X_{\epsilon}^N \subseteq X_{\epsilon}$ then we have $\nu_{\epsilon}^N \geq \nu_{\epsilon}^*$. By Lemma B.13.1, it follows that

$$\mathbb{P} [\nu_{\epsilon}^N < \nu_{\epsilon}^*] \leq \mathbb{P} [X_{\epsilon}^N \not\subseteq X_{\epsilon}] = \mathbb{P} [X_{\epsilon}^N \not\subseteq X'_{\bar{\alpha}}] \leq m|B \setminus X'_{\bar{\alpha}}| \exp \left\{ -2N \min_{i \in [m]} (\bar{\alpha}_i - \epsilon_i)^2 \right\}.$$

Therefore,

$$\begin{aligned}
\mathbb{P} [\nu_{\epsilon}^N \neq \nu_{\epsilon}^*] &\leq \mathbb{P} [\nu_{\epsilon}^N > \nu_{\epsilon}^*] + \mathbb{P} [\nu_{\epsilon}^N < \nu_{\epsilon}^*] \leq m \exp \left(- \min_{i \in [m]} \left\{ \frac{(\underline{\alpha}_i - \epsilon_i)^2 N}{(\underline{\alpha}_i + \epsilon_i)} \right\} \right) + m|B \setminus X_{\epsilon}| \exp \left\{ -2N \min_{i \in [m]} (\bar{\alpha}_i - \epsilon_i)^2 \right\} \\
&\leq m (|B \setminus X_{\epsilon}| + 1) \exp \{ -\kappa_3 N \}.
\end{aligned}$$

B.14 Proof of Theorem 9

To prove the theorem, we need the following lemma:

Lemma B.14.1 *For $n \geq 1$, let $p_j, j = 1, \dots, n$ be numbers such that $1 > p_1 \geq \dots \geq p_n > 0$. Given an integer w with $1 \leq w \leq n$, consider the following problem:*

$$\min_{\lambda \in \mathbb{R}_+^n} \max_{\substack{z_j \in \{0,1\}^n \\ \sum_j z_j = w}} \sum_{j=1}^n z_j \lambda_j + \sum_{j=1}^n \log \left(e^{-\lambda_j} p_j + (1 - p_j) \right). \quad (14)$$

Then, there exists an optimal solution to the above problem that satisfies $\lambda_1 \geq \dots \geq \lambda_n \geq 0$.

Proof (of Lemma B.14.1). Suppose that $\lambda := (\lambda_1, \dots, \lambda_n)$ is an optimal solution to the problem and there exists some $j < n$ such that $\lambda_j < \lambda_{j+1}$. We defined $\bar{\lambda}$ as $\bar{\lambda}_j := \lambda_{j+1}$, $\bar{\lambda}_{j+1} := \lambda_j$ and $\bar{\lambda}_\ell := \lambda_\ell$ for $\ell \neq \{j, j+1\}$. We will show that $\bar{\lambda}$ has no worse function value than λ . First notice that

$$\max_{\substack{z_j \in \{0,1\}^n \\ \sum_j z_j = w}} \sum_{j=1}^n z_j \lambda_j = \max_{\substack{z_j \in \{0,1\}^n \\ \sum_j z_j = w}} \sum_{j=1}^n z_j \bar{\lambda}_j,$$

which is equal to the sum of the largest w components of the vector λ . Therefore, we only need to compare the remaining part of the objective function. Define Δ as the difference in the objection function between λ and $\bar{\lambda}$, i.e.,

$$\begin{aligned} \Delta &= \sum_{j=1}^n z_j \lambda_j + \sum_{j=1}^n \log \left(e^{-\lambda_j} p_j + (1 - p_j) \right) - \left(\sum_{j=1}^n z_j \bar{\lambda}_j + \sum_{j=1}^n \log \left(e^{-\bar{\lambda}_j} p_j + (1 - p_j) \right) \right) \\ &= \sum_{j=1}^n \log \left(e^{-\lambda_j} p_j + (1 - p_j) \right) - \sum_{j=1}^n \log \left(e^{-\bar{\lambda}_j} p_j + (1 - p_j) \right) \\ &= \log \left(e^{-\lambda_j} p_j + (1 - p_j) \right) + \log \left(e^{-\lambda_{j+1}} p_{j+1} + (1 - p_{j+1}) \right) \\ &\quad - \log \left(e^{-\bar{\lambda}_j} p_j + (1 - p_j) \right) - \log \left(e^{-\bar{\lambda}_{j+1}} p_{j+1} + (1 - p_{j+1}) \right) \end{aligned}$$

Since $\bar{\lambda}_j = \lambda_{j+1}$ and $\bar{\lambda}_{j+1} = \lambda_j$, it follows that

$$\Delta = \log \left(\frac{e^{-\lambda_j} p_j + (1 - p_j)}{e^{-\lambda_{j+1}} p_j + (1 - p_j)} \right) - \log \left(\frac{e^{-\lambda_j} p_{j+1} + (1 - p_{j+1})}{e^{-\lambda_{j+1}} p_{j+1} + (1 - p_{j+1})} \right) = \log \left(\frac{e^{-\lambda_j} - e^{-\lambda_{j+1}}}{e^{-\lambda_{j+1}} + \frac{1}{p_j} - 1} + 1 \right) - \log \left(\frac{e^{-\lambda_j} - e^{-\lambda_{j+1}}}{e^{-\lambda_{j+1}} + \frac{1}{p_{j+1}} - 1} + 1 \right).$$

Note that the term inside the log is positive, since $-\lambda_j > -\lambda_{j+1}$. Further, since $p_j \geq p_{j+1}$, it follows that $1/p_j - 1 \leq 1/p_{j+1} - 1$. Therefore, we conclude that $\Delta \geq 0$. \blacksquare

Proof (of Theorem 9). We consider obtaining an optimal solution by minimizing

$$\log(B_x(\lambda)) = \max_{\tilde{a}: \sum_{j=1}^n \tilde{a}_j x_j < k} \sum_{j=1}^n \lambda_j \tilde{a}_j + \sum_{j=1}^n \log \left(e^{-\lambda_j} p_j + (1 - p_j) \right).$$

We consider a vector $\tilde{a} := (z_1, z_2, \dots, z_n) \in \{0, 1\}^n$ such that $z_j = 1$ if $x_j = 0$, and $\sum_{j=1}^n z_j = n - u + k - 1$. Without loss of generality, we assume the index set $\{j : z_j = 1\}$ corresponds to $\{1, \dots, n - u + k - 1\}$. By lemma B.14.1, it follows that minimizing $\log(B_x(\lambda))$ over $\lambda \geq 0$ amounts to solving the following problem:

$$\min_{\lambda \in \mathbb{R}_+^n} \phi(\lambda) := \sum_{j=1}^{n-u+k-1} \lambda_j + \sum_{j=1}^n \log \left(e^{-\lambda_j} p_j + (1 - p_j) \right) \quad (15a)$$

$$\lambda_j \geq \lambda_{j+1} \quad j = 1 \dots n-1 \quad (15b)$$

$$\lambda_n \geq 0 \quad (15c)$$

Note that the objective function $\phi(\lambda)$ is strictly convex in λ . In fact, its second derivatives are

$$\frac{\partial^2 \phi}{\partial \lambda_j^2} = \frac{e^{-\lambda_j} p_j (1-p_j)}{(e^{-\lambda_j} p_j + (1-p_j))^2} > 0, \quad \frac{\partial^2 \phi}{\partial \lambda_{j_1} \partial \lambda_{j_2}} = 0.$$

which implies the above problem (15) has a unique optimal solution that can be found by using Karush-Kuhn-Tucker(KKT) conditions:

$$1_{(1 \leq n-u+k-1)} - \frac{p_j e^{-\lambda_j}}{e^{-\lambda_j} p_j + (1-p_j)} - \mu_1 = 0 \quad (16a)$$

$$1_{(j \leq n-u+k-1)} - \frac{p_j e^{-\lambda_j}}{e^{-\lambda_j} p_j + (1-p_j)} + \mu_{j-1} - \mu_j = 0, \quad j = 2, \dots, n \quad (16b)$$

$$\mu_j (\lambda_{j+1} - \lambda_j) = 0, \quad j = 1, \dots, n-1 \quad (16c)$$

$$\mu_n \lambda_n = 0 \quad (16d)$$

$$\mu_j \geq 0, \quad j = 1, \dots, n \quad (16e)$$

where $(\mu_j)_{j \in [n-1]}$ is the Lagrangian multipliers of constraints (15b) and μ_n is the Lagrangian multiplier of constraint (15c).

We consider a particular choice of vectors μ and λ defined as follows:

$$\mu_n := 0, \quad (17)$$

$$\mu_j := \min\{j, n-u+k-1\} - \sum_{\ell=1}^j \frac{e^{-\lambda^*}}{e^{-\lambda^*} p_\ell + (1-p_\ell)}, \quad j = 1, \dots, n-1 \quad (18)$$

$$\lambda_1 = \dots = \lambda_n := \lambda^* \quad (19)$$

where $\lambda^* \in \mathbb{R}_+$ solves the following equation

$$\eta(\lambda^*) := \sum_{j=1}^n \frac{e^{-\lambda^*} p_j}{e^{-\lambda^*} p_j + (1-p_j)} = n-u+k-1. \quad (20)$$

Note that we can always find such λ^* , since the function $\eta(\lambda)$ is continuous and decreasing, and

$$\eta(0) = \sum_{j=1}^n p_j \geq \sum_{j=1}^n p_j x_j > n-u+k-1,$$

$$\lim_{\lambda \rightarrow +\infty} \eta(\lambda) = 0 < n-u+k-1.$$

Then we claim that μ and λ satisfy the KKT conditions (16a) – (16e). First note that equations (18) imply (16a) and (16b). Equations (16c) follow from (19), Equation (16d) follows from (17). Finally, (20) implies that

$$\sum_{\ell=1}^j \frac{e^{-\lambda^*} p_\ell}{e^{-\lambda^*} p_\ell + (1-p_\ell)} < n-u+k-1, \quad j = 1, \dots, n-1.$$

Additionally, since each term on the left-hand side of the above summation is less than 1, we have

$$\sum_{\ell=1}^j \frac{e^{-\lambda^*} p_\ell}{e^{-\lambda^*} p_\ell + (1-p_\ell)} < j, \quad j = 1, \dots, n.$$

The above two observations yield inequalities (16e).

C Additional Experiments

C.1 The performance of checking infeasibility

In this section, we investigate computational performance of four approaches for checking infeasibility by reducing the value of the success probability p_{ij} . Similar to the settings in Section 5.1, we select up to 12 columns at random for each row i where $k_i \geq 2$. However, this time we uniformly sample the success probability $p_{ij} = P[\tilde{a}_{ij} = 1]$ from the range $[0.2, 0.6]$. For the remaining $n - n'$ columns in row i , we still assign a success probability p_{ij} of 0. All other settings remain consistent with those in Section 5.1. By reducing the success probability p_{ij} , we can generate infeasible instances with a high probability. If an infeasible instance cannot be obtained for fixed n, m , and ϵ , then this process should be repeated until an infeasible instance is found.

Table 2 presents the computational performance of four approaches for detecting infeasibility of instances. Observe that SAA and IS are efficient in detecting infeasibility, while OA methods perform well in all cases except for some instances when $n = 300$. It is worth noting that even if the corresponding SAA or IS problem is infeasible, this does not necessarily mean that the original CC-SMCP problem is infeasible, as reflected in Table 1.

C.2 Experiments on special cases

In this section, we conduct a series of experiments on the special cases outlined in Section 3. For the PSC problem, extensive experiments have already been conducted in the previous literature (Ahmed and Papageorgiou 2013, Fischetti and Monaci 2012, Haight et al. 2000). Therefore, our main focus is on the case when cover probabilities of each item are equal, as discussed in Section 3.2.

In the following experiments, the covering parameter k_i is randomly selected from $\{2,3\}$ and all other experimental settings remain consistent with those in Section 5.1. Given that we are working with a *sparse* probability matrix P in our experimental setup, this implies that for a given item i , it can either be covered with a set with an equal probability p_i or not be covered at all by that set. This assumption extends the case discussed in Section 3.2 and motivates us to avoid computing \bar{d} . Instead, we aim to address the following problem:

$$\min \left\{ \sum_{j \in [n]} c_j x_j : \sum_{j \in J_i} x_j \geq \bar{d}_i, \forall i \in [m], x \in B \subseteq \{0, 1\}^n \right\},$$

where $J_i = \{j \in [n] : p_{ij} \neq 0\}$. This problem takes into account the nonzero coefficients of the probability matrix for each item i and integrates the corresponding subproblems together.

Table 3 presents the computational performance of three methods when applied to special cases. In this table, “SC” refers to the method described in Section 3.2 that is used to address these special cases. “BS”, “Solving” and “Total” indicate the time spent on binary search, the time required to solve the reformulated problem, and the overall time (BS+ Solving) for the SC method, respectively.

From Table 3, we can see that the SC method successfully solves all instances very quickly, indicating that the reformulation technique introduced in Section 3.2 can significantly reduce solution time compared to the general case in Section 5.1. Additionally, the SC method outperforms sampling-based methods in almost all cases. This may imply that sampling-based methods are not very effective for special cases when n or m increases.

C.3 The effect of risk parameter ϵ_i

The results from Table 1 indicate that OA approaches can solve the instances to optimality when $\epsilon_i := \epsilon$ is small, which is commonly the case. However, it remains an intriguing question to investigate how well

Table 2: Computational performance of methods for checking infeasibility ($p_{ij} \in [0.2, 0.6]$, $N = 200$)

Parameters			OA-I		OA-II		SAA	IS
n	m	ϵ	Time	Iterations	Time	Iterations	Time	Time
30	10	0.05	0.26	2	0.04	2	0.00	0.00
30	10	0.1	0.13	2	0.05	2	0.00	0.00
30	20	0.05	1.30	2	0.88	2	0.01	0.01
30	20	0.1	0.81	2	0.93	2	0.01	0.01
30	30	0.05	0.37	2	0.18	2	0.01	0.01
30	30	0.1	0.46	2	0.20	2	0.01	0.01
30	50	0.05	2.90	2	1.62	2	0.02	0.02
30	50	0.1	2.66	2	1.24	2	0.02	0.02
30	100	0.05	0.00	1	0.01	1	0.04	0.03
30	100	0.1	0.00	1	0.00	1	0.04	0.03
30	150	0.05	0.01	1	0.01	1	0.05	0.05
30	150	0.1	0.01	1	0.01	1	0.05	0.05
50	50	0.05	1.18	2	0.35	2	0.02	0.02
50	50	0.1	0.32	2	0.35	2	0.02	0.02
50	100	0.05	0.00	1	0.01	1	0.05	0.07
50	100	0.1	0.00	1	0.01	1	0.05	0.04
50	150	0.05	1.69	2	1.03	2	0.08	0.07
50	150	0.1	1.60	2	1.09	2	0.07	0.07
50	200	0.05	0.01	1	0.01	1	0.11	0.10
50	200	0.1	0.01	1	0.01	1	0.11	0.10
50	250	0.05	0.01	1	0.01	1	0.15	0.13
50	250	0.1	0.01	1	0.01	1	0.15	0.13
50	300	0.05	0.01	1	0.02	1	0.18	0.19
50	300	0.1	0.01	1	0.02	1	0.18	0.17
100	50	0.05	0.00	1	0.00	1	0.04	0.03
100	50	0.1	0.00	1	0.00	1	0.04	0.04
100	100	0.05	0.01	1	0.01	1	0.09	0.07
100	100	0.1	0.01	1	0.01	1	0.08	0.08
100	150	0.05	13.27	2	3.17	2	0.14	0.11
100	150	0.1	9.34	2	3.18	2	0.14	0.11
300	50	0.05	0.00	1	0.00	1	0.08	0.07
300	50	0.1	0.00	1	0.00	1	0.08	0.07
300	100	0.05	740.40	2	955.05	2	0.18	0.15
300	100	0.1	747.88	2	162.76	2	0.18	0.15
300	150	0.05	0.01	1	0.01	1	0.27	0.23
300	150	0.1	487.47	2	1406.14	2	0.25	0.25
300	200	0.05	0.01	1	0.02	1	0.36	0.31
300	200	0.1	0.01	1	0.02	1	0.35	0.31
300	250	0.05	0.02	1	0.02	1	0.44	0.45
300	250	0.1	0.02	1	0.02	1	0.45	0.42
300	300	0.05	0.02	1	0.03	1	0.56	0.81
300	300	0.1	0.02	1	0.03	1	0.54	0.47

Table 3: Computational performance of methods on special cases ($p_{ij} \in [0.9, 1], N = 200$)

Parameters			SC				SAA			IS		
n	m	ϵ	BS	Solving	Total	Val	Time	Val	Gap(%)	Time	Val	Gap(%)
30	10	0.05	0.00	0.00	0.00	9	0.41	9	0.00	1.04	9	0.00
30	10	0.1	0.00	0.00	0.00	8	0.29	7	-12.50	0.10	7	-12.50
30	20	0.05	0.00	0.00	0.00	15	3.28	16	6.67	1.77	15	0.00
30	20	0.1	0.00	0.00	0.00	14	5.41	13	-7.14	3.34	15	7.14
30	30	0.05	0.00	0.00	0.00	17	1.48	18	5.88	0.71	17	0.00
30	30	0.1	0.00	0.00	0.00	17	25.02	15	-11.76	1.77	17	0.00
30	50	0.05	0.00	0.00	0.00	21	2.01	21	0.00	1.84	22	4.76
30	50	0.1	0.00	0.00	0.00	20	7.75	19	-5.00	2.34	19	-5.00
30	100	0.05	0.01	0.00	0.01	24	0.78	25	4.17	1.80	25	4.17
30	100	0.1	0.01	0.00	0.01	24	2.09	24	0.00	2.90	24	0.00
30	150	0.05	0.03	0.00	0.03	25	3.26	26	4.00	1.37	24	-4.00
30	150	0.1	0.01	0.00	0.01	24	2.45	23	-4.17	2.58	24	0.00
50	50	0.05	0.00	0.00	0.00	27	7.71	27	0.00	2.51	27	0.00
50	50	0.1	0.00	0.00	0.00	25	11.00	25	0.00	3.22	26	4.00
50	100	0.05	0.01	0.00	0.01	35	0.03	INF	INF	0.02	INF	INF
50	100	0.1	0.01	0.00	0.01	33	7.11	34	3.03	2.88	33	0.00
50	150	0.05	0.01	0.00	0.01	39	0.03	INF	INF	0.04	INF	INF
50	150	0.1	0.01	0.00	0.01	38	14.95	35	-7.89	3.15	38	0.00
50	200	0.05	0.01	0.00	0.01	39	13.97	37	-5.13	4.10	39	0.00
50	200	0.1	0.01	0.05	0.06	35	152.25	33	-5.71	11.23	35	0.00
50	250	0.05	0.01	0.00	0.01	INF	0.12	INF	0.00	0.07	INF	INF
50	250	0.1	0.01	0.06	0.07	40	30.12	40	0.00	6.87	40	0.00
50	300	0.05	0.02	0.00	0.02	INF	0.20	INF	0.00	0.05	INF	INF
50	300	0.1	0.02	0.00	0.02	42	20.48	41	-2.38	6.90	42	0.00
100	50	0.05	0.00	0.00	0.01	36	49.49	35	-2.78	6.87	35	-2.78
100	50	0.1	0.00	0.09	0.09	33	155.75	31	-6.06	27.11	33	0.00
100	100	0.05	0.01	0.00	0.01	58	10.91	55	-5.17	4.62	56	-3.45
100	100	0.1	0.01	0.00	0.01	51	56.68	50	-1.96	14.32	50	-1.96
100	150	0.05	0.01	0.05	0.06	59	103.79	58	-1.69	32.99	58	-1.69
100	150	0.1	0.01	0.01	0.01	53	1040.21	53	0.00	198.95	54	1.89
300	50	0.05	0.00	0.01	0.01	71	836.67	70	-1.41	7.28	67	-5.63
300	50	0.1	0.00	0.06	0.06	66	948.75	64	-3.03	102.95	64	-3.03
300	100	0.05	0.00	0.01	0.01	105	0.03	INF	INF	190.44	105	0.00
300	100	0.1	0.01	0.05	0.06	100	-	97	-3.00	-	98	-2.00
300	150	0.05	0.01	0.07	0.08	135	0.06	INF	INF	0.02	INF	INF
300	150	0.1	0.01	0.03	0.05	126	0.03	INF	INF	267.93	123	-2.38
300	200	0.05	0.01	0.01	0.02	148	0.08	INF	INF	256.41	147	-0.68
300	200	0.1	0.01	0.01	0.02	138	-	139	0.72	1068.16	139	0.72
300	250	0.05	0.01	0.00	0.01	INF	0.09	INF	0.00	0.05	INF	INF
300	250	0.1	0.01	0.10	0.12	135	-	134	-0.74	-	132	-2.22
300	300	0.05	0.01	0.09	0.10	159	0.09	INF	INF	0.05	INF	INF
300	300	0.1	0.02	0.09	0.11	149	-	146	-2.01	-	147	-1.34

these approaches work for larger values of ϵ . In this section, we calculate the average solution time for four approaches as a function of ϵ , considering cases where $n = 50, m = 30$ and $n = 100, m = 50$. For each approach, we conduct five replications at each ϵ to compute the average time. All other settings remain consistent with those in Section 5.1. Figure 4 illustrates the average solution time of four different approaches as a function of ϵ on two different instances.

From Figure 4, we observe that OA-I exhibits high variability, with significant spikes around $\epsilon = 0.1$ and $\epsilon = 0.5$. SAA displays a sudden large spike at $\epsilon = 0.9$, but maintains a low solution time in other cases. This could be attributed to the fact that the LP relaxation of SAA reformulation (24) is too weak to provide a strong lower bound at $\epsilon = 0.9$. In contrast, the OA-II and IS approaches are more stable, generally maintaining low and consistent solution times across different values of ϵ . OA-II improves upon OA-I, possibly by incorporating stronger reformulations that reduce computational time, resulting in more stable performance across various ϵ values.




Figure 4: The average solution time of four approaches as a function of ϵ on two instances

References

- Hoeffding W (1994) Probability inequalities for sums of bounded random variables. *The collected works of Wassily Hoeffding* 409–426.
- Mitzenmacher M, Upfal E (2005) *Probability and Computing: Randomized Algorithms and Probabilistic Analysis* (Cambridge University Press), ISBN 9780521835404, URL <https://books.google.com/books?id=0bAY16d7hvkC>.