

**Economic evaluation of empowering older patients transitioning from hospital to home:  
Trial-based analysis of the 'Your Care Needs You' intervention**

Alfredo Palacios<sup>1\*</sup>, Simon Walker<sup>1</sup>, Beth Woods<sup>1</sup>, Gerry Richardson<sup>1</sup>

<sup>1</sup>Centre for Health Economics, University of York, York, UK

This version: 24th October 2024

**Abstract**

**Background:** Transitioning from hospital to home presents significant challenges, particularly for patients aged 75 and above. The 'Your Care Needs You' (YCNy) intervention, evaluated through a cluster randomised controlled trial (cRCT), aims to empower these patients by providing essential information and skills to prepare them for being at home. This study evaluates the cost-effectiveness of the YCNy intervention compared to standard care within the UK National Health Service (NHS) over the trial's duration (90 days post-discharge).

**Methods:** The economic evaluation utilised individual patient-level data from the YCNy cRCT. The analysis was conducted from the perspective of the NHS and Personal Social Services. Data from a routine dataset, including hospitalisations, was combined with case report forms (CRFs) detailing patient socio-demographics and EQ-5D health scores. The analysis calculated the adjusted mean differences in costs and Quality-Adjusted Life Years (QALYs) between the YCNy intervention and control groups. In our base case, Multilevel Mixed-Effects Generalised Linear Models (MME-GLM) were employed to account for the hierarchical data structure. In an alternative scenario, Seemingly Unrelated Regression (SUR) models were used to address correlations between costs and QALYs. Additional alternative scenarios and a probabilistic sensitivity analysis were conducted to assess the robustness of the findings.

**Results:** The base case results indicated the YCNy intervention reduced costs (£269) and improved outcomes (QALY gain of 0.0057). The net health benefit (NHB) of the YCNy intervention using a cost-effectiveness threshold of £15,000/QALY was 0.0246. These findings suggest the YCNy intervention dominates usual care, with a high probability (89%) of the YCNy intervention being cost-effective at a £15,000 per QALY threshold. The results remained robust across several alternative scenario analyses.

---

\* Corresponding author. Email: [alfredo.palacios@york.ac.uk](mailto:alfredo.palacios@york.ac.uk)

Discussion: The YCNY intervention could represent a promising and potentially cost-effective approach for enhancing the safety and experience of older adults transitioning from hospital to home in the UK. These findings support the integration of patient-centred care models like YCNY into healthcare policies, potentially fostering more sustainable and effective care strategies within the NHS.

**Keywords:** Economic evaluation in healthcare, Trial-based economic evaluation, Elderly healthcare management, Transitional care evaluation.

## 1. Introduction

The transition from hospital to home is a vulnerable period for elderly patients, marked by a high risk of emergency readmissions that can lead to poor health outcomes (Mor and Besdine, 2011; Naylor et al., 1999). In the United Kingdom National Health Service (NHS), recent statistics indicate that about 19.6% of patients aged 75 and over are readmitted to hospitals in an emergency within 30 days of their initial discharge (NHS, 2022). Notably, a high proportion of these readmissions, approximately 30%, are considered preventable with more effective transitional care and support systems (Auerbach et al., 2016; Blunt et al., 2015; van Walraven et al., 2011). This significant rate of emergency readmission not only imposes a substantial financial burden on the healthcare system but also compromises the quality of life for this patient group, underscoring the urgent need for effective transitional care interventions.

The 'Your Care Needs You' (YCNy) intervention is designed to empower patients by providing them with crucial knowledge and tools essential for managing their care following hospital discharge. The YCNy intervention comprises three structured components directed at patient engagement: (i) a booklet that encourages patient involvement in their healthcare and provides guidance across four key areas—health monitoring, medication management, daily activities, and steps for escalation of care when needed, (ii) an informational film that reinforces the importance of the YCNy intervention and addresses patients' health beliefs and emotional concerns, and (iii) a comprehensive care summary given to patients at discharge, which includes practical information and resources for social support. Additionally, hospital staff are encouraged to facilitate patient engagement with these resources and to provide the necessary social and emotional support to enhance the effectiveness of the YCNy intervention (Baxter et al., 2020).

The YCNy intervention was implemented and assessed through a cluster randomised controlled trial (cRCT) across forty NHS wards in England (the trial was registered as ISRCTN17062524). The study allocated wards to either the YCNy intervention or a care-as-usual arm, stratified by specialty and patient demographics. Primary outcomes focused on 30-day unplanned readmission rates, with secondary metrics including readmissions at 60 and 90 days, time until readmission, and readmission duration. While the initial 30-day outcome did not reveal a significant difference between the two groups (OR 0.93; 95% CI, 0.78 to 1.10;  $p=0.372$ ), the medium-term analysis indicated a favourable trend towards intervention effectiveness. Specifically, there was a statistically significant 13%

reduction in the total number of readmissions over the 90-day period for wards receiving the YCNY intervention (IRR 0.87; 95% CI, 0.76 to 0.99;  $p=0.039$ ). This suggests a potential delayed benefit of the YCNY intervention in improving post-discharge care and reducing the likelihood of readmissions. Although these findings are promising in terms of patient outcomes, the economic implications of adopting such transitional care models remain unexplored.

A systematic review of economic evaluations focusing on interventions aimed at preventing hospital readmissions identified 50 relevant studies (Nuckols et al., 2017). Among these, 14 specifically addressed interventions targeting the elderly population. Seven of these studies concentrated on elderly individuals with a specific health condition, such as heart failure, while the remaining seven encompassed the general elderly population. The findings from these seven economic studies focused on the general elderly population indicated that the interventions were generally cost-saving (Gillespie et al., 2009; Graham et al., 2015; Graves et al., 2009; Jayadevappa et al., 2006; Naylor et al., 1994, 1999; Saleh et al., 2012). However, it is important to note that six out of these seven studies were cost-comparison analyses rather than full economic evaluations, thereby limiting the comprehensiveness of their economic assessments. This highlights a gap in comprehensive economic evaluations within this critical area of healthcare, underscoring the need for more detailed and methodologically rigorous studies to better inform policy and practice.

The aim of this study is to conduct a within-trial economic evaluation of the 'Your Care Needs You' (YCNY) intervention from the perspective of the NHS and Personal Social Services (PSS). In line with the trial's design, this evaluation focuses on the 90-day period post-discharge to provide an analysis of both YCNY intervention and healthcare costs, alongside Quality-Adjusted Life Years (QALYs). This evaluation is essential for the NHS as it seeks to ascertain the cost-effectiveness of the YCNY intervention, thereby determining its 'value for money' and impact on population health. The findings will be crucial in guiding resource allocation to maximise population health and facilitate the integration of patient-centred transitional care models within the healthcare system.

## **2. Methods**

### ***Study design***

The economic analysis utilised individual patient-level data from the YCNY trial to assess the cost-effectiveness of the YCNY intervention versus usual care. These calculations include YCNY intervention costs, healthcare costs and health outcomes. A cost-effectiveness analysis was conducted which calculated the incremental cost-effectiveness ratio (ICER) based on the adjusted mean differences in costs and QALYs between the YCNY intervention and control groups. Additionally, the net health benefit (NHB) and net monetary benefit (NMB) were calculated using a cost-effectiveness threshold of £15,000 per QALY gained (Martin et al., 2023). A health economics analysis plan (HEAP) was developed prior to conducting the economic evaluation to outline the specific methods and procedures. The HEAP is available upon request.

The analysis was conducted from the perspective of the NHS and Personal Social Services (PSS), aligning with UK guidelines for economic evaluation (NICE, 2022). In line with the clinical trial, the within-trial economic evaluation focuses on a 90-day post-randomization period. Given the trial's short duration, costs and benefits were not discounted. Cost-effectiveness thresholds of £15,000 (Martin et al., 2023), and £20,000 and £30,000 per QALY, as recommended by NICE (2022), were used to evaluate the cost-effectiveness of the YCNY intervention. The analysis adhered to established guidelines for reporting cost-effectiveness studies (Gomes et al., 2012a; Husereau et al., 2022; Ramsey et al., 2015) to ensure comprehensive and transparent reporting. CHEERS 2022 Checklist is presented in Table S1 of the Supplementary material.

### ***Participants***

Full details of the inclusion and exclusion criteria to select participants for the YCNY intervention evaluation are available elsewhere (REF). In short, the inclusion criteria required participants to be aged 75 or older, expected to be discharged to their own or a relative's home, staying for at least one night on a participating ward, and able to give informed consent. Exclusion criteria were participants requiring an interpreter, living outside the study area, expecting transfer to another acute hospital or community rehabilitation unit, among others. Notably, the clinical study excluded patients who died within 30 days without readmission. To mitigate potential bias, these patients were incorporated into this economic evaluation, contributing to both cost and QALY calculations. A summary of the sample sizes across the datasets is provided in Table S2 in the Supplementary material.

## **Data**

The YCNY trial utilises two primary data sources: 1) the Routine Dataset, which includes variables such as unplanned hospital readmissions at 30, 60, and 90 days, length of hospital stays, ward characteristics, and other relevant variables; and 2) the Case Report Form (CRF) Dataset, which contains patient characteristics and EQ-5D measurements and some resource use utilisation (some hospital services, and primary care and social care) for a subsample of individuals from the Routine Dataset. These EQ-5D and resource use measurements were taken at baseline (collected in hospital during the initial admission and following patient recruitment), and at 10, 30, and 90 days post-discharge.

In order to consolidate the unplanned hospital readmissions, the EQ-5D and resource use measurements, all essential for the economic evaluation, we combined both datasets (Routine and CRF datasets) creating a Merged Dataset. Therefore, only patients contributing to both datasets (Routine and CRF datasets) were included in the analysis. The analysis of this Merged dataset followed the “intention-to-treat” (ITT) principle.

## ***Intervention and resource use measures and valuation***

Resource use data collection involved the following categories/resources: 1) hospital services: unplanned readmissions at 30, 60, and 90 days and the duration of inpatient stays, outpatient clinic, day case, and accident and emergency (A&E); 2) primary care services: GP at surgery, home and telephone, nurse at surgery, home and telephone, and therapist; and 3) social services: home care and social worker. While unplanned readmissions at 30, 60, and 90 days and the duration of inpatient stays comes from a routine dataset, the information for rest of the healthcare and social services comes from the CRF to patients and caregivers.

Regarding the YCNY intervention cost, we considered the staff profile and time to be trained and to provide the YCNY intervention. This information was captured in a separate administrative documentation based on expert opinion. The staff profile involved mainly nurses/managers, and the main activities for delivering the YCNY intervention were to discuss with patients about care during admission and discharge, assisting with activities of daily living and providing instructions/education for patient and/or caregiver. For further details about the delivery of the YCNY intervention see Table S3 in the Supplementary material.

All resource utilisation was monetized and expressed in 2022 British pounds, using published UK unit cost estimates (Jones et al., 2023; NHS, 2023). The cost of implementing the YCNY intervention was calculated based on staff salary data (Jones et al., 2023).

### ***Outcome measures and valuation***

The primary economic outcome is Quality-Adjusted Life Years (QALYs), derived from health-related quality of life scores obtained through the EQ-5D-5L instrument. These scores were mapped to the EQ-5D-3L value set using the Hernandez-Alava et al. (2022) mapping algorithm, following UK guidance. Outcome measurements were collected at baseline (during the initial hospital admission and post-recruitment) and at 10, 30, and 90 days post-discharge (via postal questionnaires). QALYs were calculated using the area under the curve approach.

### ***Missing data***

The dataset was scrutinised for missing data. The nature and extent of the missing data were assessed to determine the most appropriate imputation method. For missing data at baseline, within-cluster mean imputation was used independently of the YCNY intervention allocation (Faria et al., 2014; Taljaard et al., 2008).

In our base case, we assumed that data was missing at random and multiple imputation methods through chained equations (MICE) by arm were employed (Faria et al., 2014; White et al., 2011). The number of imputed datasets was defined according to the percentage of incomplete cases (White et al., 2011), and the analysis was conducted using the 'mi impute' package in Stata. In the imputation model were considered the same variables as in the main analysis and the hierarchical structure of the data (Díaz-Ordaz et al., 2013; Gomes et al., 2013; Graham, 2009).

For the specific case of EQ-5D variables, the decision about imputing the domains or the EQ-5D index was based on the sample size and the proportion of missing data, as suggested by Simons et al. (2015). Given the potential situation of missing values for the resource use information coming from CRFs, we will conduct multiple imputation for resources with missing values lower 60%, and these resources will be part of the main cost-effectiveness analysis. For the rest of resources, that is, those with a percentage of missing values higher than 60%, we will present a resource use comparison between YCNY intervention and usual care considering a complete case analysis.

### ***Statistical methods***

Mean costs and QALYs for each group were presented alongside their adjusted mean differences, with 95% confidence intervals around the differences in means (Willan et al., 2004). In accordance with the clinical study, the mean differences for costs were controlled by ward specialty, readmission rate at baseline (ward level variable), percentage of patients over 75 years (ward level variable), and sex (patient level variable). The mean differences for QALYs were controlled by EQ-5D at baseline and the same variables as the analysis of mean difference for costs.

For the base case analysis, multilevel mixed-effects generalised linear models (MME-GLM) were estimated for analysing differences in mean costs and outcomes considering the hierarchical structure of the data (wards in hospitals as clusters) (Gomes et al., 2012b). Seemingly unrelated regression (SUR) models with robust standard errors were also considered to account for potential intragroup (cluster) correlations between costs and QALYs. Costs and QALYs were combined to calculate an incremental cost-effectiveness ratio (ICER), as well as the net health benefit (NHB) using the cost-effectiveness threshold of £15,000 per QALY (Martin et al., 2023).

Non-parametric bootstrap was also used to produce the cost-effectiveness plane, representing the uncertainty in incremental cost and effect estimates, and the probability of YCNY intervention being cost-effective at different thresholds. Given that different approaches have been proposed to combine multiple imputation and bootstrap, we followed a possible approach consisting of drawing bootstrap samples from each of the imputed dataset separately and then pooling the estimates (Leurent et al., 2018; Schomaker and Heumann, 2018).

### ***Sensitivity analyses***

Following Leurent et al. (2018), we modify the potentially multiply-imputed data to reflect possible departures from the MAR assumption. In our case, we considered that the potential missing HRQoL data may be MNAR, while the MAR assumption is likely to hold for the potential missing cost data. This approach involves use MI to impute the missing values under an MAR assumption, modify the MAR-imputed data to reflect a range of plausible MNAR scenarios by multiplying the imputed values by +/-10% (assumption), and analyse the resulting dataset as one would a usual multiply-imputed dataset.



### 3. Results

#### *Participants*

Table 1 presents the baseline characteristics of 455 participants in 35 hospital wards included in the Merged dataset, divided into YCNY intervention (patients=216, wards=16) and control (patients=239, wards=19) groups. Both groups are similar in terms of the individual patient level characteristics, such as age and gender. Some differences are observed in the ward level variables: the YCNY intervention group shows a slightly lower baseline readmission rate and percentage of patients over 75 years old compared to the control group. For an analysis comparing the baseline patient and ward characteristics between the Routine dataset and the Merged dataset, please see Table S4 in the Supplementary material. For a description of the missing data by intervention arms, see Table S5 in the Supplementary material.

**Table 1.** Baseline characteristics.

Variable	YCNY Intervention	Control	Total	p-value
	Patients=222 Wards=16	Patients=246 Wards=19	Patients=468 Wards=35	
Age	82.8 (5.4)	83.4 (5.5)	83.1 (5.4)	0.24
Dummy sex, 1=Male, %	48.2 (50.1)	39.4 (49.0)	43.6 (49.6)	0.056
EQ-5D at baseline	0.50 (0.31)	0.48 (0.33)	0.49 (0.32)	0.44
Ward: baseline readmission rate, %	18.4 (6.6)	19.7 (5.6)	19.1 (6.1)	0.028
Ward: patients over 75 yo, %	69.5 (27.6)	74.3 (19.3)	72.0 (23.7)	0.028
Ward: Dummy specialty, 1=Eldery & interm. care, %	53.2 (50.0)	55.7 (49.8)	54.5 (49.9)	0.58

#### *Healthcare resource use and costs*

Table 2 presents the number and percentage of unplanned readmissions within the trial follow-up period (90 days) for both the YCY intervention and control groups. In the YCNY intervention group 73% of patients had zero readmissions, compared to 67% in the control group. The percentage of

participants with one readmission was higher in the Control group (23%) than in the YCNY intervention group (18%). Both groups were similar in the percentage of participants with two readmissions, at approximately 7.2%. The Control group had a higher percentage of participants with three or more compared to the YCNY intervention group.

**Table 2.** Hospital readmissions within 90 days by arm. Data are presented as N (%)

Readmissions	YCNY intervention	Control	Total
0	162 (72.97)	166 (67.48)	328 (70.09)
1	41 (18.47)	56 (22.76)	97 (20.73)
2	16 (7.21)	18 (7.32)	34 (7.26)
3	2 (0.9)	5 (2.03)	7 (1.5)
4	1 (0.45)	0 (0)	1 (0.21)
5	0 (0)	1 (0.41)	1 (0.21)
Mean (SD)	0.3739 (0.6990)	0.4553 (0.7746)	0.4167 (0.7400)

An unadjusted comparative analysis of the YCNY intervention costs, healthcare costs, and total costs between the YCNY intervention and control groups suggest that the mean YCNY intervention cost was £94.32, while the hospitalisation costs averaged £1,542 in the YCNY intervention group and £1,870 in the control group, resulting in reduction of £327. Overall, the total costs for the YCNY intervention group were £1,637, compared to £1,870 for the control group, reflecting a decrease of £233. For a comparative analysis of other hospital services, primary care and social care services between arms based on a complete case analysis, see Table S6 in the Supplementary material. For further details about the intervention cost and the unplanned readmission cost between arms, see Table S7 in the Supplementary material.

### **Health outcomes**

Table 3 presents the unadjusted mean EQ-5D-3L scores across four time points: baseline, 10 days, 30 days, and 90 days, and QALYs comparing the YCNY intervention and control arms based on imputed values. For a summary of the EQ-5D-5L values per arm across time based on complete cases, see Table S8 in the Supplementary material, and for a description of the missing values pattern EQ-5D-3L scores see Table S9, Figure S1 and Table S10 in the Supplementary material.

At baseline, the mean EQ-5D-3L scores were similar between the groups, with the YCNY intervention arm slightly higher. Over time, the YCNY intervention group consistently showed higher mean EQ-5D scores compared to the control group.

**Table 3.** Unadjusted mean for EQ-5D-3L and QALYs variables between arms. Results based on imputed data.

Variable	YCNY Intervention (N=222)	Control (N=246)
EQ-5D at Baseline	0.5010	0.4782
EQ-5D at 10 days	0.4788	0.4498
EQ-5D at 30 days	0.4747	0.4384
EQ-5D at 90 days	0.4945	0.4156
QALYs	0.1191	0.1072

### ***Cost-effectiveness***

Table 4 presents a summary of the main cost-effectiveness results. For our base case (MME-GLM), the adjusted differences in total costs and total QALYs were -£268.78 [95% CI: -889.67 - 352.11] and 0.0057 [95% CI: -0.0108 - 0.0223], respectively. In the alternative specification (SUR model), the adjusted differences in total costs and total QALYs were -£233.75 and 0.0071, respectively (the unadjusted mean total costs and total QALYs by arm are presented in Table S11 in the Supplementary material). Given that the YCNY intervention is associated with lower costs and a slight QALY increment, the YCNY intervention could be considered “Dominant” relative to usual care.

**Table 4.** Cost-effectiveness results. Monetary values expressed in British pounds of 2022. Results based on imputed data.

		Base case: MME-GLM	Alternative case: SUR model
Total costs	Adj. Mean Control*	1,574.29	1,576.43
	Adj. Mean YCNY intervention*	1,305.51	1,342.68

	Adj. Mean Difference*	-268.78	-233.75
	SE Difference	316.78	333.57
	95% CI Difference	-889.67 - 352.11	-887.54 - 420.04
Total QALYs	Adj. Mean Control**	0.0835	0.0841
	Adj. Mean YCNY intervention**	0.0892	0.0912
	Adj. Mean Difference**	0.0057	0.0071
	SE Difference	0.0084	0.0077
	95% CI Difference	-0.0108 - 0.0223	-0.0080 - 0.0222
ICER		YCNY Dominant	YCNY Dominant

Notes: \*Results adjusted by ward specialty, readmission rate at baseline (ward level variable), percentage of patients over 75 years (ward level variable), and sex (patient level variable). \*\*Results adjusted by EQ-5D at baseline ward specialty, readmission rate at baseline (ward level variable), percentage of patients over 75 years (ward level variable), and sex (patient level variable). MME-GLM: Multilevel mixed-effects generalised linear models, SUR: Seemingly unrelated regression, QALY: quality-adjusted life year, ICER: incremental cost-effectiveness ratio.

### **Sensitivity analysis**


Table 5 presents the cost-effectiveness of the YCNY intervention under various assumptions regarding missing not at random (MNAR) quality-of-life data, while Figure 1 shows the corresponding cost-effectiveness planes (the corresponding cost-effectiveness acceptability curves are presented in Figure S2 in Supplementary material). The table compares different scenarios by rescaling the MNAR parameters for both the control and YCNY groups. In the base case scenario (Scenario 1), which assumes missing at random (MAR), the YCNY intervention is dominant with an incremental cost of -£269 and an incremental QALY gain of 0.0057, resulting in an 89% probability of being cost-effective. Other scenarios adjust the rescaling parameters, for example, reducing imputed quality-of-life values by 5% or 10%. Across all scenarios (Scenarios 2-7), the YCNY intervention consistently remains dominant, with incremental QALY gains between 0.0027 and 0.0057. The probability of cost-effectiveness at a threshold of £15,000 per QALY varies from 87% to 90%, underscoring the robustness of the YCNY intervention's cost-effectiveness under different assumptions of missing data. All of these results are based on imputed data.

**Table 5.** Cost-effectiveness of YCNY intervention under different MNAR assumptions for missing quality-of-life data. Results based on imputed data.

Scenario number	MNAR rescaling parameters <sup>a</sup>		Incremental cost <sup>b</sup> (£) [95% CI]	Incremental QALYs [95% CI]	NHB [95% CI]	Prob. cost-effective <sup>c</sup> (%)
	C control	C YCNY				
1 (Basecase, MAR)	1	1	-268.78 [-889.67 - 352.11]	0.0057 [-0.0108 - 0.0223]	0.0246 [-0.0190 - 0.0682]	89
2	1	0.95	-268.78 [-889.67 - 352.11]	0.0027 [-0.0135 - 0.0189]	0.0231 [-0.0205 - 0.0667]	87
3	0.95	1	-268.78 [-889.67 - 352.11]	0.0057 [-0.0103 - 0.0216]	0.0261 [-0.0174 - 0.0670]	90
4	0.95	0.95	-268.78 [-889.67 - 352.11]	0.0043 [-0.0117 - 0.0202]	0.0246 [-0.0189 - 0.0681]	89
5	0.95	0.90	-268.78 [-889.67 - 352.11]	0.0029 [-0.0130 - 0.0187]	0.0231 [-0.0204 - 0.0666]	87
6	0.90	0.95	-268.78 [-889.67 - 352.11]	0.0058 [-0.0098 - 0.0214]	0.0261 [-0.0174 - 0.0656]	90
7	0.90	0.90	-268.78 [-889.67 - 352.11]	0.0044 [-0.0111 - 0.0200]	0.0246 [-0.0189 - 0.0680]	89

Notes: All results are based on imputed data and comparing the YCNY intervention arm to the control arm (n = 468). NHB: net health benefit, MAR: missing at random, MNAR: missing not at random, QALY: quality-adjusted life year, YCNY: Your Care Needs You. CI: confidence interval. <sup>a</sup>How missing quality-of-life data are assumed to differ from the MAR-imputed values, for example, c control = 0.9 means that all imputed quality-of-life values in the control arm have been reduced by 10%. <sup>b</sup>Missing costs assumed to be MAR in all scenarios. <sup>c</sup>At a cost-effectiveness threshold of £15,000/QALY.

**Figure 1.** Cost-effectiveness planes of the YCNY intervention under different MNAR assumptions. Results based on imputed data.



Graphs by scenario

*Notes:* Headings in the top of each plane indicate the scenario number and the MNAR rescaling parameters (c control, c YCNY). For example, (0.9, 0.9): imputed quality-of-life values have been reduced by 10% in both arms. MAR: missing at random, MNAR: missing not at random, QALY: quality-adjusted life year, YCNY: Your Care Needs You.

#### 4. Discussion

The study's findings provide detailed insights into the short-term costs and health outcomes associated with the YCNY intervention. In particular, the YCNY intervention was cost savings alongside slight improvements in health outcomes compared to usual care. Although the observed changes in costs and outcomes were minor, this analysis indicates that the YCNY intervention may offer a viable and efficient means of achieving health benefits in resource-constrained environments.

Recent economic studies conducted after Nuckols' systematic review have evaluated several transitional care programs (TCPs) across different healthcare systems. For example, studies in the US (Heo et al., 2023) and Canada (Murmman et al., 2023) have indicated that TCPs can reduce healthcare consumption and associated costs, with implications in the cost-effectiveness of these programs. However, these studies had limitations, such as not using health-related quality of life (HRQoL) as an outcome measure, being single-centre studies, and employing quasi-experimental designs.

Nevertheless, one UK-based study by Sahara et al. (2017) presents similarities to our research. This study evaluated the cost-effectiveness of a Community In-reach Rehabilitation and Care Transition (CIRACT) service compared to traditional hospital-based rehabilitation (THB-Rehab) through a randomised controlled trial involving 250 participants. The trial measured hospital length of stay, unplanned readmission rates at day 28 and day 91, and HRQoL using the EQ-5D-3L at 91 days post-discharge. Their findings indicated an incremental cost of CIRACT over THB-Rehab of £144 (95% CI: -1645 to 1934) and an incremental QALY gain of 0.0404 (95% CI: -0.0566 to 0.1375), resulting in an incremental cost-effectiveness ratio (ICER) per QALY of £2,022 (95% CI: -76,895 to 121,856). However, the study had some limitations and differences with our study, including its single-centre nature (the Queen's Medical Centre, a large teaching hospital in Nottingham, UK), the relative youngest patient population (adults aged 70 years and older), the distant recruitment period (2013-2014), the small sample size, and the no clear statistical management of missing data.

Our study has several strengths. Firstly, our multicentre design, encompassing 15 hospitals and 35 wards, ensures that our results are more representative than those of single-centre studies. Secondly, although our sample size might be considered modest, it is nearly double that of the Sahara et al. study, providing greater statistical power and confidence in our results. Thirdly, our statistical methods are rigorous. Many economic evaluations using data from cluster-randomised trials overlook the importance of accounting for clustering or correlation (Gomes et al., 2012b), which can lead to underestimated statistical uncertainty and inaccurate point estimates (Grieve et al., 2010). In our study, we employed recommended methodologies to address the hierarchical structure of the data and the potential correlation between costs and effects at the ward level. Additionally, we addressed a common issue in economic evaluations: the handling of missing data. Most studies discard observations with missing data and report only complete case analyses (CCAs) (Noble et al., 2012), which can introduce bias if the missingness is related to unobserved values (missing not at random, or MNAR). For instance, patients in poorer health may be less likely to complete quality-of-life questionnaires, skewing results. To mitigate this, we implemented a multiple imputation exercise using chained equations and conducted an extensive scenario analysis, providing a more comprehensive analysis.

Despite these strengths, our study has limitations that should be acknowledged. First, the partial availability of EQ-5D scores from a smaller subset of the patient cohort (collected via CRF from individuals) introduces uncertainty, compounded by substantial missing data on these measurements. Additionally, most of the healthcare resource consumption information (with the exception of hospitalisation) was collected via CRFs. These resources include certain hospital services, primary care

services, and social services. Unfortunately, data on these non-hospitalisation resources exhibited a high percentage of missing values and were, therefore, excluded from the primary cost-effectiveness analysis. However, an analysis of healthcare consumption by trial arm suggests that the control group utilised more primary care and social services compared to the intervention group. Consequently, we believe our cost-effectiveness results may be considered conservative estimates. Third, in our scenario analysis addressing the missing HRQoL data, we rescaled the MAR-imputed HRQoL using a multiplicative factor. However, alternative pattern-mixture approaches, such as applying an additive factor, may have been considered to 'offset' the data. Moreover, with longitudinal data, the assumption of constant departure from MAR over time, as used in this study, may not hold. For instance, the departure could vary with time since withdrawal, potentially increasing over time (Ratitch et al., 2013). Fourth, one challenge with conducting an MNAR scenario analysis is selecting plausible sensitivity parameter values. An alternative approach could be to formally elicit expert beliefs regarding the missing data distribution (White et al., 2007). These experts may include trial investigators, clinicians, or even patients who can provide valuable insights into the missing data mechanisms. Mason et al. (2017), for example, offer a useful framework for eliciting expert opinion on MNAR mechanisms in cost-effectiveness analysis (CEA). Finally, the 90-day follow-up period may not fully capture the medium-term impacts of the YCNY intervention. This limitation is particularly important, as medium- and long-term effects may significantly influence the cost-effectiveness outcomes. Therefore, a long-term economic evaluation, considering the lifetime horizon of these patients, is necessary to thoroughly assess the YCNY intervention.

In conclusion, the YCNY intervention represents a promising, cost-effective approach for improving the safety and experience of older adults transitioning from hospital to home in the UK. These findings support the integration of patient-centred care models like YCNY into healthcare policies, potentially fostering more sustainable and effective care strategies within the NHS. By adopting such interventions, the healthcare system can strive towards better patient outcomes and efficient resource utilisation. Continued research and investment in innovative transitional care programs are essential to realising these benefits.



**Disclaimer**

The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health and Social Care.

**Funding**

This summary presents independent research funded by the National Institute for Health Research (NIHR) under the NIHR Programme Grant for Applied Research (PGfAR) Award, reference RP-PG-1214-20017. Partners at Care Transitions (PACT): Improving Patient Experience and Safety at Transitions of Care.

**Availability of data and materials**

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

**Competing interests**

The authors declare that they have no competing interests.

## References

- Auerbach, A.D., Kripalani, S., Vasilevskis, E.E., Sehgal, N., Lindenauer, P.K., Metlay, J.P., Fletcher, G., Ruhnke, G.W., Flanders, S.A., Kim, C., Williams, M.V., Thomas, L., Giang, V., Herzig, S.J., Patel, K., Boscardin, W.J., Robinson, E.J., Schnipper, J.L., 2016. Preventability and Causes of Readmissions in a National Cohort of General Medicine Patients. *JAMA Intern. Med.* 176, 484–493. <https://doi.org/10.1001/jamainternmed.2015.7863>
- Baxter, R., Murray, J., O'Hara, J.K., Hewitt, C., Richardson, G., Cockayne, S., Sheard, L., Mills, T., Lawton, R., PACT research team, 2020. Improving patient experience and safety at transitions of care through the Your Care Needs You (YCN) intervention: a study protocol for a cluster randomised controlled feasibility trial. *Pilot Feasibility Stud* 6, 123. <https://doi.org/10.1186/s40814-020-00655-5>
- Blunt, I., Bardsley, M., Grove, A., Clarke, A., 2015. Classifying emergency 30-day readmissions in England using routine hospital data 2004-2010: what is the scope for reduction? *Emerg. Med. J.* 32, 44–50. <https://doi.org/10.1136/emered-2013-202531>
- Díaz-Ordaz, K., Kenward, M.G., Grieve, R., 2013. Handling Missing Values in Cost Effectiveness Analyses that Use Data From Cluster Randomized Trials. *J. R. Stat. Soc. Ser. A Stat. Soc.* 177, 457–474. <https://doi.org/10.1111/rssa.12016>
- Faria, R., Gomes, M., Epstein, D., White, I.R., 2014. A guide to handling missing data in cost-effectiveness analysis conducted within randomised controlled trials. *Pharmacoeconomics* 32, 1157–1170. <https://doi.org/10.1007/s40273-014-0193-3>
- Gillespie, U., Alassaad, A., Henrohn, D., Garmo, H., Hammarlund-Udenaes, M., Toss, H., Kettis-Lindblad, A., Melhus, H., Mörlin, C., 2009. A comprehensive pharmacist intervention to reduce morbidity in patients 80 years or older: a randomized controlled trial. *Arch. Intern. Med.* 169, 894–900. <https://doi.org/10.1001/archinternmed.2009.71>
- Gomes, M., Díaz-Ordaz, K., Grieve, R., Kenward, M.G., 2013. Multiple imputation methods for handling missing data in cost-effectiveness analyses that use data from hierarchical studies: an application to cluster randomized trials. *Med. Decis. Making* 33, 1051–1063. <https://doi.org/10.1177/0272989X13492203>
- Gomes, M., Grieve, R., Nixon, R., Edmunds, W.J., 2012a. Statistical methods for cost-effectiveness analyses that use data from cluster randomized trials: a systematic review and checklist for critical appraisal. *Med. Decis. Making* 32, 209–220. <https://doi.org/10.1177/0272989X11407341>
- Gomes, M., Ng, E.S.-W., Grieve, R., Nixon, R., Carpenter, J., Thompson, S.G., 2012b. Developing appropriate methods for cost-effectiveness analysis of cluster randomized trials. *Med. Decis. Making* 32, 350–361. <https://doi.org/10.1177/0272989X11418372>
- Graham, J.W., 2009. Missing data analysis: making it work in the real world. *Annu. Rev. Psychol.* 60, 549–576. <https://doi.org/10.1146/annurev.psych.58.110405.085530>
- Graham, K.L., Wilker, E.H., Howell, M.D., Davis, R.B., Marcantonio, E.R., 2015. Differences between early and late readmissions among patients: a cohort study. *Ann. Intern. Med.* 162, 741–749. <https://doi.org/10.7326/M14-2159>
- Graves, N., Courtney, M., Edwards, H., Chang, A., Parker, A., Finlayson, K., 2009. Cost-effectiveness of an intervention to reduce emergency re-admissions to hospital among older patients. *PLoS One* 4, e7455. <https://doi.org/10.1371/journal.pone.0007455>
- Grieve, R., Nixon, R., Thompson, S.G., 2010. Bayesian hierarchical models for cost-effectiveness analyses that use data from cluster randomized trials. *Med. Decis. Making* 30, 163–175. <https://doi.org/10.1177/0272989X09341752>
- Heo, M., Taaffe, K., Ghadshi, A., Teague, L.D., Watts, J., Lopes, S.S., Tilkemeier, P., Litwin, A.H., 2023. Effectiveness of Transitional Care Program among High-Risk Discharged Patients: A Quasi-Experimental Study on Saving Costs, Post-Discharge Readmissions and Emergency Department Visits. *Int. J. Environ. Res. Public Health* 20. <https://doi.org/10.3390/ijerph20237136>

- Hernández Alava, M., Pudney, S., Wailoo, A., 2022. Estimating the Relationship Between EQ-5D-5L and EQ-5D-3L: Results from a UK Population Study. *Pharmacoeconomics* 41, 199–207. <https://doi.org/10.1007/s40273-022-01218-7>
- Husereau, D., Drummond, M., Augustovski, F., de Bekker-Grob, E., Briggs, A.H., Carswell, C., Caulley, L., Chaiyakunapruk, N., Greenberg, D., Loder, E., Mauskopf, J., Mullins, C.D., Petrou, S., Pwu, R.-F., Staniszewska, S., 2022. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) 2022 Explanation and Elaboration: A Report of the ISPOR CHEERS II Good Practices Task Force. *Value Health* 25, 10–31. <https://doi.org/10.1016/j.jval.2021.10.008>
- Jayadevappa, R., Chhatre, S., Weiner, M., Raziano, D.B., 2006. Health resource utilization and medical care cost of acute care elderly unit patients. *Value Health* 9, 186–192. <https://doi.org/10.1111/j.1524-4733.2006.00099.x>
- Jones, K.C., Weatherly, H., Birch, S., Castelli, A., Chalkley, M., Dargan, A., Forder, J.E., Gao, M., Hinde, S., Markham, S., Ogunleye, D., Premji, S., Roland, D., 2023. Unit Costs of Health and Social Care 2022 Manual 121. <https://doi.org/10.22024/UniKent/01.02.100519>
- Leurent, B., Gomes, M., Faria, R., Morris, S., Grieve, R., Carpenter, J.R., 2018. Sensitivity Analysis for Not-at-Random Missing Data in Trial-Based Cost-Effectiveness Analysis: A Tutorial. *Pharmacoeconomics* 36, 889–901. <https://doi.org/10.1007/s40273-018-0650-5>
- Martin, S., Claxton, K., Lomas, J., Longo, F., 2023. The impact of different types of NHS expenditure on health: Marginal cost per QALY estimates for England for 2016/17. *Health Policy* 132, 104800. <https://doi.org/10.1016/j.healthpol.2023.104800>
- Mason, A.J., Gomes, M., Grieve, R., Ulug, P., Powell, J.T., Carpenter, J., 2017. Development of a practical approach to expert elicitation for randomised controlled trials with missing health outcomes: Application to the IMPROVE trial. *Clin. Trials* 14, 357–367. <https://doi.org/10.1177/1740774517711442>
- Mor, V., Besdine, R.W., 2011. Policy options to improve discharge planning and reduce rehospitalization. *JAMA* 305, 302–303. <https://doi.org/10.1001/jama.2010.2006>
- Murmann, M., Sinden, D., Hsu, A.T., Thavorn, K., Eddeen, A.B., Sun, A.H., Robert, B., 2023. The cost-effectiveness of a nursing home-based transitional care unit for increasing the potential for independent living in the community among hospitalized older adults. *J. Med. Econ.* 26, 61–69. <https://doi.org/10.1080/13696998.2022.2156152>
- Naylor, M., Brooten, D., Jones, R., Lavizzo-Mourey, R., Mezey, M., Pauly, M., 1994. Comprehensive discharge planning for the hospitalized elderly. A randomized clinical trial. *Ann. Intern. Med.* 120, 999–1006. <https://doi.org/10.7326/0003-4819-120-12-199406150-00005>
- Naylor, M.D., Brooten, D., Campbell, R., Jacobsen, B.S., Mezey, M.D., Pauly, M.V., Schwartz, J.S., 1999. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. *JAMA* 281, 613–620. <https://doi.org/10.1001/jama.281.7.613>
- NHS, 2023. NHS England » 2021/22 National Cost Collection Data Publication [WWW Document]. URL <https://www.england.nhs.uk/publication/2021-22-national-cost-collection-data-publication/>
- NHS, 2022. Emergency readmissions to hospital within 30 days of discharge: indirectly standardised percent trends broken down by age bands and sex.
- NICE, 2022. NICE health technology evaluations: the manual.
- Noble, S.M., Hollingworth, W., Tilling, K., 2012. Missing data in trial-based cost-effectiveness analysis: the current state of play. *Health Econ.* 21, 187–200. <https://doi.org/10.1002/hec.1693>
- Nuckols, T.K., Keeler, E., Morton, S., Anderson, L., Doyle, B.J., Pevnick, J., Booth, M., Shanman, R., Arifkhanova, A., Shekelle, P., 2017. Economic Evaluation of Quality Improvement Interventions Designed to Prevent Hospital Readmission: A Systematic Review and Meta-analysis. *JAMA Intern. Med.* 177, 975–985. <https://doi.org/10.1001/jamainternmed.2017.1136>
- Ramsey, S.D., Willke, R.J., Glick, H., Reed, S.D., Augustovski, F., Jonsson, B., Briggs, A., Sullivan, S.D., 2015. Cost-effectiveness analysis alongside clinical trials II—An ISPOR Good Research Practices Task Force report. *Value Health* 18, 161–172. <https://doi.org/10.1016/j.jval.2015.02.001>

- Ratitch, B., O'Kelly, M., Tosiello, R., 2013. Missing data in clinical trials: from clinical assumptions to statistical analysis using pattern mixture models. *Pharm. Stat.* 12, 337–347.  
<https://doi.org/10.1002/pst.1549>
- Sahota, O., Pulikottil-Jacob, R., Marshall, F., Montgomery, A., Tan, W., Sach, T., Logan, P., Kendrick, D., Watson, A., Walker, M., Waring, J., 2017. The Community In-reach Rehabilitation and Care Transition (CIRACT) clinical and cost-effectiveness randomisation controlled trial in older people admitted to hospital as an acute medical emergency. *Age Ageing* 46, 26–32.  
<https://doi.org/10.1093/ageing/afw149>
- Saleh, S.S., Freire, C., Morris-Dickinson, G., Shannon, T., 2012. An effectiveness and cost-benefit analysis of a hospital-based discharge transition program for elderly Medicare recipients. *J. Am. Geriatr. Soc.* 60, 1051–1056. <https://doi.org/10.1111/j.1532-5415.2012.03992.x>
- Schomaker, M., Heumann, C., 2018. Bootstrap inference when using multiple imputation. *Stat. Med.* 37, 2252–2266. <https://doi.org/10.1002/sim.7654>
- Simons, C.L., Rivero-Arias, O., Yu, L.-M., Simon, J., 2015. Multiple imputation to deal with missing EQ-5D-3L data: Should we impute individual domains or the actual index? *Qual. Life Res.* 24, 805–815. <https://doi.org/10.1007/s11136-014-0837-y>
- Taljaard, M., Donner, A., Klar, N., 2008. Imputation strategies for missing continuous outcomes in cluster randomized trials. *Biom. J.* 50, 329–345. <https://doi.org/10.1002/bimj.200710423>
- van Walraven, C., Bennett, C., Jennings, A., Austin, P.C., Forster, A.J., 2011. Proportion of hospital readmissions deemed avoidable: a systematic review. *CMAJ* 183, E391–402.  
<https://doi.org/10.1503/cmaj.101860>
- White, I.R., Carpenter, J., Evans, S., Schroter, S., 2007. Eliciting and using expert opinions about dropout bias in randomized controlled trials. *Clin. Trials* 4, 125–139.  
<https://doi.org/10.1177/1740774507077849>
- White, I.R., Royston, P., Wood, A.M., 2011. Multiple imputation using chained equations: Issues and guidance for practice. *Stat. Med.* 30, 377–399. <https://doi.org/10.1002/sim.4067>
- Willan, A.R., Briggs, A.H., Hoch, J.S., 2004. Regression methods for covariate adjustment and subgroup analysis for non-censored cost-effectiveness data. *Health Econ.* 13, 461–475.  
<https://doi.org/10.1002/hec.843>

## Supplementary material

**Table S1.** CHEERS 2022 Checklist.

Topic	No.	Item	Location where item is reported
<b>Title</b>			
Title	1	Identify the study as an economic evaluation and specify the interventions being compared.	Title
<b>Abstract</b>			
Abstract	2	Provide a structured summary that highlights context, key methods, results, and alternative analyses.	Abstract
<b>Introduction</b>			
Background and objectives	3	Give the context for the study, the study question, and its practical relevance for decision making in policy or practice.	Introduction
<b>Methods</b>			
Health economic analysis plan	4	Indicate whether a health economic analysis plan was developed and where available.	Methods, Line 5
Study population	5	Describe characteristics of the study population (such as age range, demographics, socioeconomic, or clinical characteristics).	Methods, Line 10
Setting and location	6	Provide relevant contextual information that may influence findings.	Methods, Third Paragraph
Comparators	7	Describe the interventions or strategies being compared and why chosen.	Methods, Fourth Paragraph
Perspective	8	State the perspective(s) adopted by the study and why chosen.	Methods, Fifth Paragraph
Time horizon	9	State the time horizon for the study and why appropriate.	Methods, Fifth Paragraph
Discount rate	10	Report the discount rate(s) and reason chosen.	Methods, Fifth Paragraph

Selection of outcomes	11	Describe what outcomes were used as the measure(s) of benefit(s) and harm(s).	Methods, Sixth Paragraph
Measurement of outcomes	12	Describe how outcomes used to capture benefit(s) and harm(s) were measured.	Methods, Sixth Paragraph
Valuation of outcomes	13	Describe the population and methods used to measure and value outcomes.	Methods, Sixth Paragraph
Measurement and valuation of resources and costs	14	Describe how costs were valued.	Not reported
Currency, price date, and conversion	15	Report the dates of the estimated resource quantities and unit costs, plus the currency and year of conversion.	Methods, Seventh Paragraph
Rationale and description of model	16	If modelling is used, describe in detail and why used. Report if the model is publicly available and where it can be accessed.	Methods, Eighth Paragraph
Analytics and assumptions	17	Describe any methods for analysing or statistically transforming data, any extrapolation methods, and approaches for validating any model used.	Methods, Last paragraph and Appendix
Characterising heterogeneity	18	Describe any methods used for estimating how the results of the study vary for subgroups.	Methods, Last paragraph
Characterising distributional effects	19	Describe how impacts are distributed across different individuals or adjustments made to reflect priority populations.	Methods, Last paragraph
Characterising uncertainty	20	Describe methods to characterise any sources of uncertainty in the analysis.	Not applicable
Approach to engagement with patients and others affected by the study	21	Describe any approaches to engage patients or service recipients, the general public, communities, or stakeholders (such as clinicians or payers) in the design of the study.	Appendix
<b>Results</b>			
Study parameters	22	Report all analytic inputs (such as values, ranges, references) including uncertainty or distributional assumptions.	Results, first paragraph and Table
Summary of main results	23	Report the mean values for the main categories of costs and outcomes of interest and summarise them in the most appropriate overall measure.	Results, second paragraph

Effect of uncertainty	24	Describe how uncertainty about analytic judgments, inputs, or projections affect findings. Report the effect of choice of discount rate and time horizon, if applicable.	Results, third paragraph
Effect of engagement with patients and others affected by the study	25	Report on any difference patient/service recipient, general public, community, or stakeholder involvement made to the approach or findings of the study	Not reported
<b>Discussion</b>			
Study findings, limitations, generalisability, and current knowledge	26	Report key findings, limitations, ethical or equity considerations not captured, and how these could affect patients, policy, or practice.	Discussion
<b>Other relevant information</b>			
Source of funding	27	Describe how the study was funded and any role of the funder in the identification, design, conduct, and reporting of the analysis	End of manuscript
Conflicts of interest	28	Report authors conflicts of interest according to journal or International Committee of Medical Journal Editors requirements.	End of manuscript

From: Husereau D, Drummond M, Augustovski F, et al. Consolidated Health Economic Evaluation Reporting Standards 2022 (CHEERS 2022) Explanation and Elaboration: A Report of the ISPOR CHEERS II Good Practices Task Force. Value Health 2022;25. doi:10.1016/j.jval.2021.10.008

**Table S2.** Sample sizes across datasets.

Dataset	Initial sample	Primary analysis cohort sample	Primary analysis cohort sample + patients who have died within 30 days and no readmission recorded
Routine	5,450	4,947	5,147
CRF	622	455	468
EE	615	455	468

Notes: CRF: Complete case form; EE: Economic evaluation.

**Table S3.** Intervention delivery per patient.

Activity	Duration (minutes)			Undertaken by
	Usual care wards	Intervention wards	Incremental	
Discussion with patient about care - on admission	5	5	0	Nursing/Manager
Discussion with patient about care - during admission	15	30	15	Nursing/Manager
Discussion with patient about care - during admission	15	25	10	Medical staff
Discussion with patient about care - discharge	15	20	5	Nursing/Manager
Assisting with activities of daily living	70	105	35	Nursing/Manager
Instructions/education for patient and/or caregiver	5	25	20	Nursing/Manager

**Table S4.** Comparison between the EE dataset and the Routine dataset at baseline.

Variable	Routine dataset	EE dataset
		Patients=5,147 Wards=39
Age	84.5 (5.8)	83.1 (5.4)
Dummy sex, 1=Male, %	41.4 (49.3)	43.6 (49.6)
Ward: baseline readmission, %	18.0 (5.3)	19.1 (6.2)
Ward: patients over 75 yo, %	74.7 (22.5)	72.0 (23.7)
Ward: Dummy specialty, 1=Eldery & interm. care, %	59.0 (49.2)	54.5 (49.9)



**Table S5.** Missing values.

Variable	Missing (n)	Total	Missing (%)
<i>Baseline data</i>			
Intervention	0	468	0
Age	0	468	0
Sex	0	468	0
EQ-5D at baseline	9	468	1.92
Ward specialty	0	468	0
Baseline readmission rate	12	468	2.56
Patients over 75 years old	0	468	0
<i>Quality of life</i>			
EQ-5D at 10 days	149	468	31.84
EQ-5D at 30 days	159	468	33.97
EQ-5D at 90 days	213	468	45.51
<i>Resource use - Hospital care</i>			
Hospitalisation	2	468	0.43
Outpatient clinic	284	468	60.68
Day case	367	468	78.42
A&E	359	468	76.71
<i>Resource use - Primary care</i>			
GP at surgery	289	468	61.75
GP at home	313	468	66.88
GP at telephone	295	468	63.03

Nurse at surgery	298	468	63.68
Nurse at home	302	468	64.53
Nurse at telephone	318	468	67.95
Therapist	300	468	64.10
<i>Resource use - Social care</i>			
Home care	304	468	64.96
Social worker	308	468	65.81

**Table S6.** Other healthcare resource use within 90 days by arm and total, complete case analysis.

Healthcare resource use	Intervention		Control		Total	
	N	%	N	%	N	%
<i>Hospital care</i>						
Outpatient clinic	84	66.7	100	68	184	67.4
Day case	42	14.3	59	23.7	101	19.8
A&E	45	20	64	29.7	109	25.7
<i>Primary care</i>						
GP at surgery	82	51.2	97	54.6	179	53.1
GP at home	72	22.2	83	31.3	155	27.1
GP at telephone	83	62.6	90	70	173	66.5
Nurse at surgery	79	43	91	49.4	170	46.5
Nurse at home	80	52.5	86	54.6	166	53.6
Nurse at telephone	74	33.8	76	27.6	150	30.7
Therapist	81	48.1	87	44.8	168	46.4

<i>Social care</i>						
Home care	79	40.5	85	51.8	164	46.3
Social worker	81	17.3	79	24	160	20.6

**Table S7.** Unadjusted mean intervention, healthcare and total costs between arms. Monetary values expressed in British pounds of 2022. Results based on imputed values.

Variable	Intervention	Control	Difference
Intervention costs	94.32	0.00	94.32
Hospitalisation costs	1,502.44	1,812.44	-310.00
Total costs	1,596.76	1,812.44	-215.68

**Table S8.** Unadjusted mean for EQ-5D-5L between arms. Results based on complete cases.


Variable	YCNV Intervention	Control
EQ-5D at Baseline	0.5999	0.5806
EQ-5D at 10 days	0.6007	0.5705
EQ-5D at 30 days	0.6271	0.5814
EQ-5D at 90 days	0.6590	0.5638

**Table S9.** Missing data pattern for the EQ-5D-3L and QALYs variables by intervention and control arms.

Arm: Intervention			
Variable	Missing	Total	% missing
EQ-5D at Baseline	0	222	0
EQ-5D at 10 days	59	222	26.58
EQ-5D at 30 days	65	222	29.28
EQ-5D at 90 days	82	222	36.94

Arm: Control			
Variable	Missing	Total	% missing
EQ-5D at Baseline	0	246	0
EQ-5D at 10 days	59	246	23.98
EQ-5D at 30 days	73	246	29.67
EQ-5D at 90 days	94	246	38.21

**Figure S1.** Pattern of missing data for the EQ-5D-3L and QALYs variables.



**Table S10.** Multilevel mixed-effect logistic estimations for the missingness in EQ-5D-3L measurements.

Variable	Dummy missing EQ-5D at 10 days, 1=Yes	Dummy missing EQ-5D at 30 days, 1=Yes	Dummy missing EQ-5D at 90 days, 1=Yes
Dummy intervention arm	0.161 (0.74)	0.0544 (0.27)	-0.136 (-0.76)
EQ-5D at baseline	-1.018** (-3.20)	-0.986*** (-3.44)	-0.0126 (-0.04)
Dummy ward speciality, 1=Eldery and intermediate care	0.471 (1.12)	0.278 (0.66)	0.134 (0.33)
Ward: baseline readmission, %	0.00225 (0.15)	-0.00128 (-0.11)	-0.00875 (-0.60)
Ward: patients over 75 yo, %	-0.00686 (-1.17)	0.000618 (0.10)	0.000955 (0.12)
Dummy sex, 1=Male	-0.129 (-0.58)	-0.259 (-1.52)	0.102 (0.54)
Constant	-0.418 (-0.89)	-0.471 (-1.31)	-0.437 (-1.02)
N	468	468	468

Notes: t statistics in parentheses. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001.

**Table S11.** Unadjusted mean total costs and total QALYs by arm.

		Intervention	Control
Total cost	Mean	1,636.77	1,869.58
	95% CI	1,210.40 - 2,063.13	1,444.54 - 2,294.62
Total QALYs	Mean	0.1459	0.1354
	95% CI	0.1369 - 0.1548	0.1269 - 0.1439

Figure S2. Cost-effectiveness acceptability curves.

