Packing sets under finite groups via algebraic incidence structures

Norbert Hegyvári^{*} Le Quang Hung[†] Alex Iosevich[‡] Thang Pham[§]

February 25, 2025

Abstract

Let E be a set in \mathbb{F}_p^n and S be a set of maps from \mathbb{F}_p^n to $\mathbb{F}_p^n.$ We define

$$S(E) := \bigcup_{f \in S} f(E) = \{f(x) \colon x \in E, f \in S\}$$

In this paper, we establish sharp lower bounds on the size of S(E) when S consists of matrices from either the special linear group $SL_2(\mathbb{F}_p)$ or the first Heisenberg group $\mathbb{H}_1(\mathbb{F}_p)$. Our proofs are based on novel results on algebraic incidence-type structures associated with these groups. We also discuss higher-dimensional generalizations.

Contents

1	Introduction						
	1.1 The special linear group $SL_2(\mathbb{F}_p)$	3					
	1.2 The first Heisenberg group $\mathbb{H}_1(\mathbb{F}_p)$	7					
2	Incidence structures spanned by $SL_2(\mathbb{F}_p)$	8					
	2.1 Incidence bounds for large sets via Fourier analysis (Theorem 2.1)	10					
	2.2 Incidence bounds for large sets via energies (Theorem 2.2)	14					
	2.3 Two alternative approaches yield weaker bounds	17					
	2.4 Incidence bounds for small sets (Theorem 2.3)	22					
	2.5 Alternative approach with relaxed conditions	26					
3	3 Proof of Propositions 1.3, 1.5 and Theorems 1.4, 1.6						
4	Incidence structures spanned by $\mathbb{H}_1(\mathbb{F}_p)$	28					
5	Proof of Theorem 1.7	32					
6	Acknowledgements	33					
	*Eötvös University and associated member of Alfréd Rényi Institute, Hungary. Email:hegyvari@renyi.hu						

[†]University of Science, Vietnam National University, Hanoi. Email: lequanghung_t65@hus.edu.vn

[‡]Department of Mathematics, University of Rochester. Email: iosevich@math.rochester.edu

[§]University of Science, Vietnam National University, Hanoi. Email: thangpham.math@vnu.edu.vn

1 Introduction

Let E be a Borel set in \mathbb{R}^n and let S be a set of maps from \mathbb{R}^n to \mathbb{R}^n . We define

$$S(E) := \bigcup_{f \in S} f(E) = \{f(x) : x \in E, f \in S\} \subset \mathbb{R}^n.$$

The packing problem asks if it is possible for a set of zero *n*-dimensional Lebesgue measure to contain the image f(E) for all $f \in S$. The study of this problem has a reputed history in the literature, for example, there are sets in the plane of zero Lebesgue measure containing a line segment of unit length in every direction (see [2] and [3]), a circle of radius r for all r > 0 (see [4] and [15]), or a circle centered at x for all x on a given straight line (see [27]). In another direction, the question of finding conditions on S and E such that S(E) has positive Lebesgue measure has been also received a lot of attention. Bourgain [5] and independently Marstrand [18] proved that given a set of circles in the plane, if the centers form a set of positive Lebesgue measure, then the union of circles also has positive Lebesgue measure. Wolff [29] strengthened this result by showing that if the set of centers has Hausdorff dimension s, $0 < s \leq 1$, then the dimension of the set of union of circles is at least 1 + s. For the most recent progress, we refer the reader to [14] and references therein.

This paper is devoted to exploring this topic in the finite field setting. Let \mathbb{F}_p be a finite field of order p, where p is a prime. Let E be a set in \mathbb{F}_p^n and S be a set of maps from \mathbb{F}_p^n to \mathbb{F}_p^n . As over the reals, we define

$$S(E) := \bigcup_{f \in S} f(E) = \left\{ f(x) \colon x \in E, f \in S \right\}.$$

The main question is to bound the size of S(E) from below. We first recall the following result due to R. Orberlin in [21].

Theorem 1.1. Suppose $1 \le d \le n-1$ is an integer, $0 \le \beta \le 1$, and that L is a collection of lines in \mathbb{F}_p^n with $|L| \ge p^{2(d-1)+\beta}$. Then

$$\left|\bigcup_{\ell \in L} \ell\right| \gg p^{d+\beta}.$$
 (1)

Here $\left|\bigcup_{\ell\in L}\ell\right|$ counts the number of points in the union of lines in L.

This theorem is the finite field analog of a conjecture due to D. Oberlin, which states that for an integer $d \ge 1$ and $0 \le \beta \le 1$, if L is set of lines in \mathbb{R}^n with Hausdorff dimension at least $2(d-1)+\beta$, then the Hausdorff dimension of the union of lines in L is at least $d+\beta$. This conjecture has been solved recently by Zahl in [31].

Theorem 1.1 is optimal. To see this, let L be the set of lines contained in p^{β} parallel d-planes. Then, we have $|L| = \frac{p^d(p^d-1)}{p(p-1)}p^{\beta}$ and $\left|\bigcup_{\ell \in L} \ell\right| = p^{d+\beta}$.

We pause here to discuss some connections to the Kakeya set problem. Dvir [9] proved that if a set $P \subset \mathbb{F}_p^n$ contains a line in every direction, then its size is at least $\gg p^n$. We know that in \mathbb{F}_p^n , there are about p^{n-1} distinct directions. If n is even, by choosing d = (n-2)/2 and $\beta = 1$, Theorem 1.1 implies that $|P| \gg p^{(n+2)/2}$. If n is odd, by choosing d = (n-1)/2 + 1 and $\beta = 0$, then Theorem 1.1 implies that $|P| \gg p^{(n+1)/2}$. These lower bounds are of course very weak compared to Dvir's

result, since the distinctness of the directions is not required in the statement of Theorem 1.1.

Note that the conclusion of Theorem 1.1 can be written as $|S(E)| \gg p^{d+\beta}$ where E is a given line and S is a set of rigid-motions with the size can be as large as |S| = |L||O(n-1)|p. When E is a general set in \mathbb{F}_p^d , the third listed author [23] established the following theorem.

Theorem 1.2. Let $E \subset \mathbb{F}_p^n$ and $S \subset O(n) \times \mathbb{F}_p^n$, $n \geq 3$.

1. If $|E| < p^{\frac{n-1}{2}}$, then we have

$$|S(E)| \gg \min\left\{p^n, \ \frac{|E||S|}{p^{n-1}|O(n-1)|}\right\}$$

2. If $p^{\frac{n-1}{2}} \le |E| \le p^{\frac{n+1}{2}}$, then we have

$$|S(E)| \gg \min\left\{p^n, \ \frac{|S|}{p^{\frac{n-1}{2}}|O(n-1)|}\right\}$$

3. If $|E| > p^{\frac{n+1}{2}}$, then we have

$$|S(E)| \gg \min\left\{p^n, \ \frac{|E||S|}{p^n|O(n-1)|}\right\}$$

A direct computation shows that Theorem 1.2(1) implies Theorem 1.1 with d = n - 1.

In this paper, we focus on the case when S consists of matrices from either the special linear group $SL_2(\mathbb{F}_p)$ or the first Heisenberg group $\mathbb{H}_1(\mathbb{F}_p)$. This is motivated by the underlying algebraic structures in these groups that highlight deep connections between Group theory, Geometry, and Combinatorics. Specifically, we aim to determine conditions on S and E such that either the set S(E) covers a positive proportion of all elements in the plane/space or $|S(E)| \gg |E|^{1+\epsilon}$ for some $\epsilon > 0$.

Notations: Throughout the paper, we will write $X \ll_{\alpha} Y$ if $X \leq CY$, where C > 0 is a constant depending on α . If it is clear from the context what C should depend on, we may write only $X \ll Y$. If $X \ll Y$ and $Y \ll X$, we write $X \sim Y$. Furthermore, $X \leq Y$ if $X \ll \log_2 p \cdot Y$.

1.1 The special linear group $SL_2(\mathbb{F}_p)$

We first start with some observations.

Observation 1: It is trivial that $|S(E)| \ge |E|$, and one might hope that $|S(E)| \ge |S|$. So, it implies $|S(E)| \ge |S|^{\frac{1}{2}} |E|^{\frac{1}{2}}$. However, the estimate $|S(E)| \ge |S|$ might not be true. For example, take $E = \{(0,1)\}$ and S being a set of matrices θ in $SL_2(\mathbb{F}_p)$ such that $\theta(0,1) = (1,0)$, then, by Lemma 2.6, S can be as large as $\sim p$, and in this case, one has |S(E)| = 1.

Moreover, there are sets S and E such that $|S(E)| = |S|^{\frac{1}{2}} |E|^{\frac{1}{2}}$. Indeed, let $A \subset B$ be two subgroups of \mathbb{F}_p^* . For each $[x] \in B/A$, fix $x' \in [x]$. Let $S_{[x]}$ be a subset of $\{\theta \in \mathrm{SL}_2(\mathbb{F}_p) : \theta(0,1) = (0,x')\}$

such that $|S_{[x]}| = |B|$. Let $E = \{(0, a) \colon a \in A\}$. Then, we have

$$S_{[x]}(E) = \{0\} \times [x].$$

Therefore, if we choose $S = \bigcup_{[x] \in B/A} S_{[x]}$, we have |S| = |B| |B/A| and

$$S(E) = \bigcup_{[x] \in B/A} \{0\} \times [x] = \{0\} \times B.$$

Then $|S(E)| = |B| = (|B| |B/A|)^{\frac{1}{2}} |A|^{\frac{1}{2}} = |S|^{\frac{1}{2}} |E|^{\frac{1}{2}}.$

Observation 2: There are sets S and E such that $|S(E)| \sim \frac{|S||E|}{p^2} > |E|$. Indeed, fix $0 < \epsilon < 1$, let E be the set of points on the line $\{y = 0\}$. Let $S \subset SL_2(\mathbb{F}_p)$ be a set of all matrices θ such that $\theta(1,0) \in E'$, where E' is a set of $p^{1+\epsilon}$ points on p^{ϵ} lines passing through the origin. Then, by Lemma 2.6, we have $|S| \sim p^{2+\epsilon}$. Therefore, we have

$$|S(E)| = p^{1+\epsilon} \sim \frac{p^{2+\epsilon} \cdot p}{p^2} \sim \frac{|S| |E|}{p^2}.$$

In the first result, we prove that the lower bound $|S||E|/p^2$ actually holds for all sets S and E. **Proposition 1.3.** Let $E \subset \mathbb{F}_p^2$ and $S \subset SL_2(\mathbb{F}_p)$. We have

$$|S(E)| \gg \min\left\{p^2, \frac{|S||E|}{p^2}\right\}.$$

It follows from this theorem that if $|S||E| \gg p^4$ then $|S(E)| \gg p^2$. This condition is optimal in the sense that for all $\epsilon > 0$ there exist sets S and E with $|S||E| \gg p^{4-\epsilon}$ such that $|S(E)| = o(p^2)$. Indeed, for all $\epsilon > 0$, by choosing p large enough, we can find a cyclic subgroup A of \mathbb{F}_p^* with $|A| \sim p^{1-\epsilon}$. Let S be the set of matrices θ in $SL_2(\mathbb{F}_p)$ such that $\theta(0,1) \in \{(-x,0): x \in A\}$. Each such matrix is of the form

$$\begin{bmatrix} * & -x \\ x^{-1} & 0 \end{bmatrix},$$

and $|S| = p^{2-\epsilon}$. We now let E to be the set of points of the form (y, *) with $y \in A$ and $* \in \mathbb{F}_p$. Then $|E| = p^{2-\epsilon}$. Moreover, S(E) is covered by the lines of the from $y = \lambda$ with $\lambda \in A$. So $|S(E)| \leq p^{2-\epsilon}$.

To prove this result, we introduce a new incidence structure between the group $SL_2(\mathbb{F}_p)$ and pairs of points in $\mathbb{F}_p^2 \times \mathbb{F}_p^2$. In particular, for $x, y \in \mathbb{F}_p^2$ and $\theta \in SL_2(\mathbb{F}_p)$, we say (x, y) is incident to θ if $\theta y = x$. Given sets $A, B \subset \mathbb{F}_p^2$ and a set $S \subset SL_2(\mathbb{F}_p)$, the main strategy is to bound the number of incidences between $P = A \times B$ and S. The equation $\theta y = x$ means that the two vectors θy and x are the same in \mathbb{F}_p^2 , thus, it is very natural to make use of tools from discrete abelian Fourier analysis to study this case. The incidence bounds obtained here are of independent interest and are expected to have further applications. A standard argument implies the following optimal incidence estimate (Theorem 2.1 below):

$$\left| I(P,S) - \frac{|P||S|}{p^2} \right| \ll p\sqrt{|S||P|} + |S|.$$
(2)

To deduce Proposition 1.3 from this incidence structure, we set A = S(E) and B = E, then the above upper bound of I(P,S) and the trivial lower bound $I(P,S) \ge |S||E|$ imply our desired estimates on the size of S(E).

In this paper, we are interested in improvements of Proposition 1.3 under structural conditions of E or S.

Our first main theorem is stated as follows.

Theorem 1.4. For any $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds. Let $E \subset \mathbb{F}_p^2$ and $S \subset SL_2(\mathbb{F}_p)$. Assume p is a sufficiently large prime, S is a symmetric subset of $SL_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and $|S \cap gH| < p^{-\frac{\gamma}{2}}|S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$, and any line through the origin contains at most k points from E, then

$$|S(E)| \gg \min\left\{p^2, \max\left\{\frac{|S||E|}{pk}, \frac{|S|^{\frac{1}{2}}|E|}{p^{\frac{1-\epsilon}{2}}k^{\frac{1}{2}}}\right\}\right\}.$$

Sharpness: We observe that conditions on S in the above theorem are natural for further improvements. Indeed, let ℓ_1 and ℓ_2 be two lines passing through the origin and S be the set of all $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta(\ell_1) = \ell_2$. Then, let $E = \ell_1$, we have $S(E) = \ell_2$, and by Lemma 2.6, $|S| \sim p^2$, so

$$|S(E)| = p \sim \frac{|S|^{\frac{1}{2}}|E|}{p}$$

Moreover, we have S = gH, where H is the group of matrices $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta(\ell_1) = \ell_1$ and $g \in SL_2(\mathbb{F}_p)$ is a matrix such that $g(\ell_1) = \ell_2$.

Compared to the proof of Proposition 1.3, that of Theorem 1.4 uses a stronger upper bound, namely,

$$I(S(E) \times E, S) \ll \frac{|S(E)|^{\frac{1}{2}}|E||S|}{p} + k^{\frac{1}{4}}p^{\frac{1-\epsilon}{4}}|S(E)|^{\frac{1}{2}}|E|^{\frac{1}{2}}|S|^{\frac{3}{4}},$$

whose proof is more subtle, relying on a refined discrete Fourier analysis argument in which the following two energies play a crucial role

$$\texttt{Energy}_1 := |\{(x, y, u, v) \in E^4 \colon x \cdot y^{\perp} = u \cdot v^{\perp}\}| \text{ and } \texttt{Energy}_2 := |\{(a, b, c, d) \in S^4 \colon ab = cd\}|.$$

The first energy has been studied intensively in the literature, for example, see [13, 22]. It was showed that

$$\operatorname{Energy}_1 \ll \frac{|E|^4}{p} + pk|E|^2,$$

where k is the maximal number of points from E on a line passing through the origin.

Regarding the second energy, it is not hard to construct examples of S such that the $\text{Energy}_2 \sim |S|^3$, see Section 2.2. However, under the conditions on S as stated in Theorem 1.4, an improved upper

bound of $|S|^3 p^{-\epsilon}$, for some $\epsilon = \epsilon(|S|) > 0$ was proved by Bourgain and Gamburd in [7]. This might be the only place where tools from non-abelian group settings have been used. In this paper, we use their result as a black box.

We now turn our attention to the case of small sets. If S and E are arbitrary sets of small size, then based on Observation 1 and Observation 2, one can guess of the existence of sets such that the size of S(E) is the same as the trivial lower bound |E|. For example, let A be a subgroup of \mathbb{F}_p^* , and let $E = \{\lambda(1,0) : \lambda \in A\}$, and S be a set of matrices θ such that θ maps at least one point in E to (0,1). Depending on the form of p, one can choose E of arbitrary small size, and the size of S can also be chosen arbitrary large in the range (0, p|E|) such that |S(E)| = |E|.

In the setting of small sets, the Fourier discrete analysis argument is not effective, and we use incidence bounds (point-line, point-plane) instead. As a consequence, we derive the following optimal result.

Proposition 1.5. Let p be an odd prime, $S \subseteq SL_2(\mathbb{F}_p)$, and $E \subseteq \mathbb{F}_p^2 \setminus \{(0,0)\}$ be such that $|E| \leq p$, any line passing through the origin contains at most k_1 points from E, and E determines at most k_2 distinct directions through the origin. Then,

$$|S(E)| \gtrsim \min\left\{\frac{|E| |S|^{\frac{1}{2}}}{k_1^{\frac{1}{2}} k_2^{\frac{1}{2}}}, \frac{|E|^{\frac{1}{2}} |S|^{\frac{1}{2}}}{k_1^{1/4}}, \frac{|E| |S|^{\frac{1}{2}}}{k_1}, \frac{|E|^2}{k_1}\right\}$$

Here, by "directions" we mean the number of lines passing through the origin that are required to cover the whole set E.

In order to have |S(E)| > |E|, one would need the conditions that

$$|E| \gg k_1, |S| \ge \max\left\{k_1^2, k_1^{1/2}|E|, k_1k_2\right\}.$$

The lower bound of Proposition 1.5 is attainable from the following example.

For a constant 0 < c < 1, let $\{\ell_i\}_{i=1}^{\frac{p}{c}}$ be a family of $\frac{p}{c}$ lines passing through the origin. For each $1 \le i \le \frac{p}{c}$, choose $a_i \in \ell_i \setminus \{(0,0)\}$, and set $E = \{a_i \colon 1 \le i \le \frac{p}{c}\}$, $S = \mathrm{SL}_2(\mathbb{F}_p)$. Then we have $k_2 = |E| = \frac{p}{c}$, $k_1 = 1$, and $|S(E)| = p^2 - 1$. Thus,

$$|S(E)| \sim \min\left\{\frac{|E||S|^{\frac{1}{2}}}{k_1^{\frac{1}{2}}k_2^{\frac{1}{2}}}, \frac{|E|^{\frac{1}{2}}|S|^{\frac{1}{2}}}{k_1^{1/4}}, \frac{|E||S|^{\frac{1}{2}}}{k_1}, \frac{|E|^2}{k_1}\right\}$$

In the spirit of Theorem 1.4, we provide an " ϵ -improvement" under structural conditions on S.

Theorem 1.6. For any $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds. Let p be a sufficiently large prime, and $E \subseteq \mathbb{F}_p^2 \setminus \{(0,0)\}$ such that $|E| \leq p$, any line passing through the origin contains at most k_1 points from E and E determines at most k_2 distinct directions through the origin. Let $S \subseteq SL_2(\mathbb{F}_p)$ be a symmetric subset such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and

 $|S \cap gH| < p^{-\frac{\gamma}{2}}|S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$. Then,

$$|S(E)| \gtrsim \min\left\{\frac{|E| |S|^{\frac{1}{2}} p^{\epsilon/2}}{k_1^{\frac{1}{2}} k_2^{\frac{1}{2}}}, \frac{|E|^{\frac{1}{2}} |S|^{\frac{1}{2}} p^{\epsilon/2}}{k_1^{1/4}}, \frac{|E| |S|^{\frac{1}{2}} p^{\epsilon/2}}{k_1}, \frac{|E|^2}{k_1}\right\}.$$

One might ask about the higher dimensional case. Let's assume n = 3 for simplicity. It is not hard to check that if we have two triples (x, y, z) and (x', y', z') such that each forms an independent system, then there exists unique $\theta \in SL_3(\mathbb{F}_p)$ such that $\theta x = x'$, $\theta y = y'$, and $\theta z = z'$ if and only if $\det(x, y, z) = \det(x', y', z')$. Here $\det(x, y, z)$ is the determinant of the matrix with columns x, y, and z. If one wishes to apply the method in the two dimensions, then an estimate on the number of tuples $(x, y, z, x', y', z') \in E^6$, $E \subset \mathbb{F}_p^3$, such that $\det(x, y, z) = \det(x', y', z')$ is needed in the first step. This is the L^2 -norm version of earlier results studied in [8, 28]. Notice that this framework only solves the case of large sets, and for the case of small sets in higher dimensions, it appears to be a hard problem due to the limited understanding of incidence bounds. We plan to address this in a subsequent paper.

1.2 The first Heisenberg group $\mathbb{H}_1(\mathbb{F}_p)$

We now move to the case of the Heisenberg group. Let \mathbb{F}_p be a prime field, we denote by $\mathbb{H}_1(\mathbb{F}_p)$ the first Heisenberg group over \mathbb{F}_p , i.e. the group of matrices of the form

$$[x, y, t] := \begin{pmatrix} 1 & x & t \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}, \ x, y, t \in \mathbb{F}_p.$$

In the group $\mathbb{H}_1(\mathbb{F}_p)$, we have

$$[x, y, t] \cdot [x', y', t'] = \left(x + x', y + y', t + t' + \frac{xy' - yx'}{2}\right).$$

This section uses the notation X(E) in order to distinguish with the case of $SL_2(\mathbb{F}_p)$.

We first look at the following examples.

Example 1: Let $X = \mathbb{H}_1(\mathbb{F}_p)$ and let E be the set of all points with the third coordinate belonging to a set of size αp in \mathbb{F}_p . Then it is clear that $|X(E)| \leq \alpha p^3$. This example tells us that in order to cover the whole space or a positive proportion of all elements in \mathbb{F}_p^3 , the condition $|E| \gg p^3$ is needed.

Example 2: There are sets X and E of arbitrary large such that $|X(E)| \ll |X|$. Let $A \subset \mathbb{F}_p$ be an arbitrary set, let X be the set of matrices $[a, b, c] \in \mathbb{H}_1(\mathbb{F}_p)$ with $a, c \in \mathbb{F}_p$ and $b \in A$, let E be the set of points (x, y, z) with $x \in \mathbb{F}_p$, $y \in A$, and $z \in A$. Then we have $|X| = p^2|A|$ and $|E| = p|A|^2$. A direct computation shows that $|X(E)| \ll p^2|A| = |X|$. Thus, for any $0 < \epsilon < 2$, there exist sets $E \subset \mathbb{F}_p^3$ and $X \subset \mathbb{H}_1(\mathbb{F}_p)$ with $|E|, |X| \ge p^{3-2\epsilon}$ such that $|X(E)| \ll |X|$.

Let $\pi_{23}: \mathbb{F}_p^3 \to \mathbb{F}_p^2$ defined by $\pi_{23}(x, y, z) = (y, z)$. The following example shows that if the preimage set $\pi_{23}^{-1}(y, z) \cap E$ is of large size for all $(y, z) \in \pi_{23}(E)$, then the trivial bound |E| might be best possible.

Example 3: There are sets X and E such that $|X(E)| \ll |E|$. Let $A \subset \mathbb{F}_p$ be an arithmetic progression. Let E be a set in \mathbb{F}_p^3 such that each point in E has the last two coordinates belonging to A. Assume that for each $(y, z) \in \pi_{23}(E)$, we have $|\pi_{23}^{-1}(y, z) \cap E| \sim p$. Then, for all sets X containing of matrices of the form $[*, 1, *] \in \mathbb{H}_1(\mathbb{F}_p)$, we have $|X(E)| \ll |E|$.

The main theorem in this section reads as follows.

Theorem 1.7. Let $\epsilon \geq 0$, X be a subset of $\mathbb{H}_1(\mathbb{F}_p)$, and E be a set in $\mathbb{F}_p^3 \setminus (\mathbb{F}_p^2 \times \{0\})$. Assume that for each $(y, z) \in \mathbb{F}_p^2$, we have $|\pi_{23}^{-1}(y, z) \cap E| \leq p^{1-\epsilon}$, then we have

$$|X(E)| \gg \min\left\{p^3, \frac{|X||E|}{p^{3-\frac{\epsilon}{2}}}\right\}.$$

Note that when $\epsilon = 0$, our proof implies directly that

$$|X(E)| \gg \min\left\{p^3, \ \frac{|X||E|}{p^3}\right\}.$$

Similar to the case of the group $SL_2(\mathbb{F}_p)$, to prove Theorem 1.7, our main tool will be an incidence estimate associated to the Heisenberg group $\mathbb{H}_1(\mathbb{F}_p)$. One might also ask about a version of Theorem 1.7 for small sets by using incidence bounds. However, we find it too complicated to pursue in this direction. For simplicity of this paper, we pose it as an open question.

2 Incidence structures spanned by $SL_2(\mathbb{F}_p)$

Let $x, y \in \mathbb{F}_p^2$ and $\theta \in \mathrm{SL}_2(\mathbb{F}_p)$, we say (x, y) is incident to θ if $\theta y = x$. Let $P = A \times B \subseteq \mathbb{F}_p^2 \times \mathbb{F}_p^2$ and $S \subseteq \mathrm{SL}_2(\mathbb{F}_p)$, we denote the number of incidences between P and S by I(P, S).

In this section, we prove the following incidence bound.

Theorem 2.1. Let $P = A \times B \subseteq \mathbb{F}_p^2 \times \mathbb{F}_p^2$ and $S \subseteq SL_2(\mathbb{F}_p)$. Then, we have

$$\left| I(P,S) - \frac{|P||S|}{p^2} \right| \ll p\sqrt{|S||P|} + |S|.$$

Moreover, let k_A and k_B be the maximal number of points from A and B on a line passing through the origin, respectively. Assume that $\min\{k_A, k_B\} = k$, then

$$\left| I(P,S) - \frac{|P| \, |S|}{p^2} \right| \ll p^{\frac{1}{2}} k^{\frac{1}{2}} \sqrt{|S| \, |P|} + |S|.$$

We have some comments on this theorem.

1. This theorem is sharp, i.e. there are sets P and S such that the upper bound is attained.

Let $P = A \times B$ where A, B are the sets of points on two lines through the origin. Let S be the set of all matrices θ in SL₂ (\mathbb{F}_p) such that $\theta(B) = A$. Then, we have $|S| = p(p-1), |P| = p^2$,

and

$$I(P,S) = p(p-1) + p(p-1)^2.$$

Hence,

$$\left| I(P,S) - \frac{|P| |S|}{p^2} \right| = p(p-1)^2 \sim p\sqrt{|P| |S|} + |S|.$$

2. The same result holds for general sets $P \subset \mathbb{F}_p^4$ instead of sets of Cartesian product structures.

If we assume some structural conditions on S, the upper bound of the above theorem can be improved further.

Theorem 2.2. For $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds. Let p be a sufficiently large prime. Let $P = A \times B \subseteq (\mathbb{F}_p^2 \times \mathbb{F}_p^2) \setminus \{(0,0,0,0)\}$, and let S be a symmetric subset of $SL_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and $|S \cap gH| < p^{\frac{-\gamma}{2}}|S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$. Then, we have

$$I(P,S) \ll \frac{|A|^{\frac{1}{2}}|B||S|}{p} + p^{\frac{2-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}}.$$

Moreover, if any line passing through the origin contains at most k points from B, we have

$$I(P,S) \ll \frac{|A|^{\frac{1}{2}}|B||S|}{p} + k^{\frac{1}{4}}p^{\frac{1-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}}.$$

The following example shows that conditions on S are necessary to obtain non-trivial improvements.

Let ℓ_1 and ℓ_2 be two lines passing through the origin. Let A, B be the set of points on ℓ_1, ℓ_2 , respectively. Let S be the set of all $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta(\ell_2) = \ell_1$, and H be the group of matrices $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta(\ell_2) = \ell_2$. Then

$$I(P,S) \sim p|A||B| \sim p^3 \sim \frac{|A|^{\frac{1}{2}}|B||S|}{p} + p^{\frac{2}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}}.$$

Moreover, it is not hard to check that S = gH for some $g \in SL_2(\mathbb{F}_p)$.

For small sets, we first look at an example, which says that I(P, S) could be |P||S|. Let S be a subset of matrices θ in $SL_2(\mathbb{F}_p)$ such that $\theta(0, 1) = (1, 0)$. So the size of S can be arbitrary smaller than p. Let $P = \{\lambda(e_1, e_2) : \lambda \in \mathbb{F}_p^*\}$, where $e_1 = (1, 0), e_2 = (0, 1)$. Then the size of P can be arbitrary smaller than p by choosing λ . With these sets P and S, we have I(P, S) = |P||S|.

When we know better about structures of B and S, then the following theorem is attained.

Theorem 2.3. For $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds.

Let p be a sufficiently large prime. Let $P = A \times B \subseteq (\mathbb{F}_p^2 \times \mathbb{F}_p^2) \setminus \{(0,0,0,0)\}$, and let S be a symmetric subset of $SL_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and $|S \cap gH| < p^{\frac{-\gamma}{2}} |S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$.

1. Assume $|B| \leq p$, any line passing through the origin contains at most k_1 points from B, and

B determines at most k_2 distinct directions through the origin, then

$$I(P,S) \lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{8}} |B|^{\frac{3}{4}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{4}} k_2^{\frac{1}{4}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}}$$

2. Assume $|B| \le p^{8/15}$, and any line passing through the origin contains at most k_1 points from B, then

$$I(P,S) \lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{15}} |B|^{\frac{187}{225}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}}$$

We note that the bound of I(P, S) in this theorem is smaller than |P||S| provided that $|A| \gg 1$.

Remark 2.4. If we remove the factor $p^{\epsilon/4}$ in the incidence bounds of Theorem 2.3, then the conditions on the set S are not required.

2.1 Incidence bounds for large sets via Fourier analysis (Theorem 2.1)

Theorem 2.1 will be proved by using tools from discrete Fourier analysis. We first recall some basic notations.

For $n \in \mathbb{N}$, let $f : \mathbb{F}_p^n \to \mathbb{C}$ be a complex valued function. The Fourier transform of f, denoted by \widehat{f} , is defined by

$$\widehat{f}(m) := p^{-n} \sum_{x \in \mathbb{F}_p^n} \chi(-m \cdot x) f(x),$$

where χ is a nontrivial additive character of \mathbb{F}_p . We have the following basic properties of \widehat{f} .

• The orthogonality property:

$$\sum_{\alpha \in \mathbb{F}_p^n} \chi(\beta \cdot \alpha) = \begin{cases} 0, & \text{if } \beta \neq (0, \dots, 0), \\ p^n, & \text{if } \beta = (0, \dots, 0). \end{cases}$$

• The Fourier inversion formula:

$$f(x) = \sum_{m \in \mathbb{F}_p^n} \chi(m \cdot x) \widehat{f}(m).$$

• The Plancherel formula:

$$\sum_{m \in \mathbb{F}_p^n} \left| \widehat{f}(m) \right|^2 = p^{-n} \sum_{x \in \mathbb{F}_p^n} |f(x)|^2.$$

For $A \subseteq \mathbb{F}_p^d$, by abuse of notation, we also denote its characteristic function by A(x), i.e. A(x) = 1 if $x \in A$ and A(x) = 0 if $x \notin A$.

To proceed further, we need three lemmas. For each $x = (x_1, x_2) \in \mathbb{F}_p^2$, we define $x^{\perp} = (-x_2, x_1)$. Note that $x \cdot y^{\perp}$ measures the area of the triangle with three vertices x, y, and the origin. The first lemma presents a fact that the group $SL_2(\mathbb{F}_p)$ preserves areas of triangles with one vertex pinned at the origin. **Lemma 2.5.** Let x, y, u, v be points in $\mathbb{F}_p^2 \setminus \{(0,0)\}$. If there exists $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta x = u$ and $\theta y = v$, then $x \cdot y^{\perp} = u \cdot v^{\perp}$. In the inverse direction, if x and y are not on the same line passing through the origin and $x \cdot y^{\perp} = u \cdot v^{\perp}$, then there exists unique $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta x = u$ and $\theta y = v$.

Proof. Assume there exists $\theta \in SL_2(\mathbb{F}_p)$ such that $\theta x = u$ and $\theta y = v$. Writing θ in the form

$$\theta = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

where ad - bc = 1, $x = (x_1, x_2)$, and $y = (y_1, y_2)$. We have $u = (ax_1 + bx_2, cx_1 + dx_2), v = (ay_1 + by_2, cy_1 + dy_2)$. Then,

$$u \cdot v^{\perp} = -(ax_1 + bx_2)(cy_1 + dy_2) + (cx_1 + dx_2)(ay_1 + by_2)$$

= -(ad - bc)x_1y_2 + (ad - bc)x_2y_1 = x \cdot y^{\perp}.

In the inverse direction, since $x \neq ky$ for all $k \in \mathbb{F}_p$, we have $x \cdot y^{\perp} \neq 0$. Indeed, writing x as $x = (x_1, x_2) \neq (0, 0)$. Since $x \neq (0, 0)$, we can assume that $x_1 \neq 0$. Therefore, if $y = (y_1, y_2)$ satisfies $x \cdot y^{\perp} = 0$, then x and y belong to a line passing through the origin, a contradiction. Similarly, we obtain $u \cdot v^{\perp} \neq 0$. Let θ be the matrix that maps the basis $\{x, y\}$ to the basis $\{u, v\}$. We show that $\theta \in \mathrm{SL}_2(\mathbb{F}_p)$. Indeed, we write $\theta = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $x = (x_1, x_2)$, and $y = (y_1, y_2)$. This implies $u = (ax_1 + bx_2, cx_1 + dx_2)$ and $v = (ay_1 + by_2, cy_1 + dy_2)$. Therefore, $x \cdot y^{\perp} = u \cdot v^{\perp} = (ad - bc)x \cdot y^{\perp}$, so ad - bc = 1. In other words, $\theta \in SL_2(\mathbb{F}_p)$.

The next lemma tells us that the action of $SL_2(\mathbb{F}_p)$ on the plane \mathbb{F}_p^2 is transitive with multiplicity of $\sim p$. We give a general proof for the case $SL_n(\mathbb{F}_p)$.

Lemma 2.6. For any $m, m' \in \mathbb{F}_p^2 \setminus \{(0,0)\}$, define

$$\mathcal{M}_{2,p}\left(m,m'\right) := \left\{ T \in SL_2\left(\mathbb{F}_p\right) : Tm = m' \right\}.$$

Then $|\mathcal{M}_{2,p}(m,m')| = p.$

Proof. It follows from [20, Theorem 13.3.3] that $SL_2(\mathbb{F}_p)$ is a group of size

$$\left|\operatorname{SL}_{2}\left(\mathbb{F}_{p}\right)\right| = p\left(p^{2} - 1\right).$$

Considering the group action of $\operatorname{SL}_2(\mathbb{F}_p)$ on \mathbb{F}_p^2 , $A: x \mapsto Ax, \forall A \in \operatorname{SL}_2(\mathbb{F}_p), x \in \mathbb{F}_p^2$. For $m \neq (0,0)$, let $\operatorname{Orb}(m)$ be the orbit of m, i.e. $\operatorname{Orb}(m) = \{Tm: T \in \operatorname{SL}_2(\mathbb{F}_p)\}$. Since $m \neq (0,0)$, there exists $m'_1 \in \mathbb{F}_p^2 \setminus \{(0,0)\}$ such that $\{m, m'_1\}$ forms a linear independent system. So, let

$$T'_m = \begin{bmatrix} m & m'_1 \end{bmatrix},$$

we have det $T'_m = \lambda_m \neq 0$. Let $m_1 = \lambda_m^{-1} m'_1$, and

$$T_m = \begin{bmatrix} m & m_1 \end{bmatrix}.$$

Then, det $T_m = \lambda_m \cdot \lambda_m^{-1} = 1$, so $T_m \in \mathrm{SL}_2(\mathbb{F}_p)$. Now, for all $m' \in \mathbb{F}_p^2 \setminus \{(0,0)\}$, we observe that $T_{m'} \circ (T_m)^{-1} \in \mathrm{SL}_2(\mathbb{F}_p)$, and $T_{m'} \circ (T_m)^{-1}(m) = m'$ so $\mathcal{M}_{2,p}(m,m') \neq \emptyset, \forall m,m' \neq (0,0)$.

For $m, m' \in \mathbb{F}_p^2 \setminus \{(0,0)\}$, let T be an element in $\mathcal{M}_{2,p}(m,m')$. Then for all $T' \in \mathcal{M}_{2,p}(m,m')$, there exists $A \in \operatorname{Stab}(m)$ such that TA = T'. This implies that $|\mathcal{M}_{2,p}(m,m')| = |\operatorname{Stab}(m)|$ for all $m, m' \in \mathbb{F}_p^2 \setminus \{(0,0)\}$.

Moreover, for any $m \neq (0,0)$, there is no $T \in SL_2(\mathbb{F}_p)$ such that Tm = (0,0). Hence, we have $Orb(m) = \mathbb{F}_p^2 \setminus \{(0,0)\}$. Thus, by the Orbit-Stabilizer theorem,

$$|\operatorname{Stab}(m)| = \frac{|\operatorname{SL}_2(\mathbb{F}_p)|}{|\operatorname{Orb}(m)|} = \frac{p(p^2 - 1)}{p^2 - 1} = p.$$

This completes the proof.

The next lemma is an L^2 bound for the dot-product function.

Lemma 2.7 ([22]). Let A and B be subsets of \mathbb{F}_p^2 . The number of tuples $(x_1, x_2, y_1, y_2) \in A \times A \times B \times B$ such that $x_1 \cdot x_2^{\perp} = y_1 \cdot y_2^{\perp}$ is at most

$$\frac{|A|^2|B|^2}{p} + Cp^2|A||B|,\tag{3}$$

for some positive constant C.

When $P \subset \mathbb{F}_p^4$ is a general set, the above upper bound can be replaced by $\frac{|P|^2}{p} + Cp^2|P|$. Let k_A and k_B be the maximal number of points from A and B on a line passing through the origin, respectively. Assume that $\min\{k_A, k_B\} = k$, then, with the same argument, the bound (3) can be replaced by

$$\frac{|A|^2|B|^2}{p} + Cpk|A||B|.$$

With these three lemmas in hand, we are ready to prove Theorem 2.1.

Proof of Theorem 2.1. Without loss of generality, we assume that $(0, 0, 0, 0) \notin P$, since this element only contributes |S| incidences to the incidence bound.

By using the Fourier transformation and the Fourier inversion formula, we have

$$I(P,S) = \sum_{\substack{p=(x,y)\in P\\\theta\in S}} 1_{\theta y=x} = \frac{1}{p^2} \sum_{\substack{m\in\mathbb{F}_p^2\\\theta\in S}} \sum_{\substack{(x,y)\in P,\\\theta\in S}} \chi(m \cdot (x - \theta y))$$
$$= \frac{|P||S|}{p^2} + \frac{1}{p^2} \sum_{\substack{m\in\mathbb{F}_p^2\setminus\{(0,0)\}\\\theta\in S}} \sum_{\substack{(x,y)\in P,\\\theta\in S}} \chi(m \cdot (x - \theta y))$$
$$= \frac{|P||S|}{p^2} + p^2 \sum_{\substack{m\neq(0,0)\\\theta\in S}} \sum_{\substack{\theta\in S}} \widehat{P}(-m, \theta^t m).$$

By using the Cauchy-Schwarz inequality, one has

$$\sum_{\theta \in S} \sum_{m \neq (0,0)} \widehat{P}(-m, \theta^T m) \le |S|^{\frac{1}{2}} \left(\sum_{\theta \in \mathrm{SL}_2(\mathbb{F}_p)} \sum_{m_1, m_2 \neq (0,0)} \widehat{P}(-m_1, \theta^t m_1) \overline{\widehat{P}(-m_2, \theta^t m_2)} \right)^{\frac{1}{2}}.$$

We now observe

$$\sum_{\theta \in SL_2(\mathbb{F}_p)} \sum_{m_1, m_2 \neq (0,0)} \widehat{P}(-m_1, \theta^t m_1) \overline{\widehat{P}(-m_2, \theta^t m_2)}$$

= $\frac{1}{p^8} \sum_{\theta} \sum_{m_1, m_2 \neq (0,0)} \sum_{(x_1, y_1), (x_2, y_2)} P(x_1, y_1) P(x_2, y_2) \chi(-m_1 x_1 + \theta^t m_1 y_1) \chi(m_2 x_2 - \theta^t m_2 y_2)$
= $\sum_{m_1, m_2} - \sum_{m_1 = (0,0), m_2 \neq (0,0)} - \sum_{m_1 \neq (0,0), m_2 = (0,0)} - \sum_{m_1 = m_2 = (0,0)}$
=: $I - II - III - IV$.

We now estimate each term separately.

$$I = \frac{1}{p^8} \sum_{\theta} \sum_{m_1, m_2 \in \mathbb{F}_p^2} \sum_{(x_1, y_1), (x_2, y_2)} P(x_1, y_1) P(x_2, y_2) \chi(m_1(\theta y_1 - x_1)) \chi(m_2(\theta y_2 - x_2))$$
$$= \frac{1}{p^4} \sum_{\theta} \sum_{(x_1, y_1), (x_2, y_2)} P(x_1, y_1) P(x_2, y_2) \mathbf{1}_{\theta y_1 = x_1} \mathbf{1}_{\theta y_2 = x_2}.$$

By using Lemma 2.5, one has

$$I = \frac{1}{p^4} \sum_{\theta} \sum_{\lambda \in \mathbb{F}_p} \sum_{\substack{x_1, y_1, x_2, y_2 \\ y_1 = \lambda y_2, x_1 = \lambda x_2}} A(x_1) A(x_2) B(y_1) B(y_2) \mathbf{1}_{\theta y_1 = x_1} \mathbf{1}_{\theta y_2 = x_2}$$
$$+ \frac{1}{p^4} \sum_{x_1, y_1, x_2, y_2} A(x_1) A(x_2) B(y_1) B(y_2) \mathbf{1}_{y_1 \cdot y_2^{\perp} = x_1 \cdot x_2^{\perp}}.$$

Notice that Lemma 2.6 and Lemma 2.5 tell us that the first sum can be bound by at most $p^2|A||B|/p^4$.

By Lemma 2.7, the second sum can be at most

$$\frac{|A|^2|B|^2}{p^5} + C\frac{|A||B|}{p^2}$$

In other words, we have

$$I \le \frac{|A|^2|B|^2}{p^5} + (C+1)\frac{|A||B|}{p^2}.$$

Moreover,

$$IV = \frac{(p^3 - p)|A|^2|B|^2}{p^8} = \frac{|A|^2|B|^2}{p^5} - \frac{|A|^2|B|^2}{p^7}.$$

Thus, $I - IV \ll \frac{|A||B|}{p^2}$. Regarding II,

$$II = \frac{1}{p^8} \sum_{\theta} \sum_{(x_1, y_1)} P(x_1, y_1) \left(\sum_{(x_2, y_2)} P(x_2, y_2) \left(\sum_{m_2 \neq 0} \chi \left(m_2 \left(x_2 - \theta y_2 \right) \right) \right) \right)$$

$$= \frac{1}{p^8} \sum_{\theta} \sum_{(x_1, y_1)} P(x_1, y_1) \left(\sum_{\substack{(x_2, y_2), \\ x_2 = \theta y_2}} P(x_2, y_2) \left(p^2 - 1 \right) - \sum_{\substack{(x_2, y_2), \\ x_2 \neq \theta y_2}} P(x_2, y_2) \right)$$

$$= \frac{1}{p^8} |P| \left(\left(p^2 - 1 \right) \sum_{\substack{(x_2, y_2) \in P, \\ \theta \in \text{SL}_2(\mathbb{F}_p), \\ x_2 = \theta y_2}} 1 - \sum_{\substack{(x_2, y_2) \in P, \\ \theta \in \text{SL}_2(\mathbb{F}_p), \\ x_2 \neq \theta y_2}} 1 \right)$$

$$= \frac{1}{p^8} |P| \left(|A| |B| p \left(p^2 - 1 \right) - |A| |B| \left(p^3 - 2p \right) \right) = \frac{|P|^2}{p^7}.$$

Similarly, we obtain $III = \frac{|P|^2}{p^7}$. Putting all estimates together, we conclude that

$$\left| I(P,S) - \frac{|P||S|}{p^2} \right| \ll p\sqrt{|P||S|} + |S|.$$

This completes the proof.

2.2 Incidence bounds for large sets via energies (Theorem 2.2)

For a set $S \subset SL_2(\mathbb{F}_p)$, we define the energy $\mathbf{E}(S,S)$ by

$$\mathbf{E}(S,S) := \#\{(a,b,c,d) \in S^4 : ab = cd\}.$$

The trivial bound of $\mathbf{E}(S, S)$ is $|S|^3$. When S is a large set, Babai, Nikolov, and Pyber proved in [1] that

$$\mathbf{E}(S,S) \ll p^2 |S|^2 + \frac{|S|^4}{p^3}.$$
(4)

This bound is sharp. To see its sharpness, we provide an example here.

Let S be the set of matrices of the form

$$\begin{bmatrix} * & -x \\ x^{-1} & 0 \end{bmatrix},$$

where $x, * \in \mathbb{F}_p \setminus \{0\}$. Then, S is a subset of $SL_2(\mathbb{F}_p)$ and $|S| = (p-1)^2$. We consider following equation

$$\begin{bmatrix} *_1 & -x \\ x^{-1} & 0 \end{bmatrix} \begin{bmatrix} *_2 & -y \\ y^{-1} & 0 \end{bmatrix} = \begin{bmatrix} *'_1 & -x' \\ (x')^{-1} & 0 \end{bmatrix} \begin{bmatrix} *'_2 & -y' \\ (y')^{-1} & 0 \end{bmatrix},$$
(5)

where all matrices are in S. The equation is equivalent to

$$\begin{cases} *_1 *_2 - xy^{-1} = *'_1 *'_2 - x'(y')^{-1} \\ -y *_1 = -y' *'_1, \\ *_2 x^{-1} = *'_2 (x')^{-1}, \\ -y x^{-1} = -y' x^{-1}. \end{cases}$$

This implies $\frac{y}{y'} = \frac{x}{x'} = \frac{*_2}{*_2} = \frac{*'_1}{*_1}$. Therefore, for fixed $*_1, *'_1, x, x'$, there exist $(p-1)^2$ tuples $(y, y', *_2, *'_2) \in (\mathbb{F}_p \setminus \{0\})^4$ such that the equation (5) holds. In other words, for each pair of matrices $(A, C) \in S^2$, there exist $(p-1)^2$ pairs of matrices $(B, D) \in S^2$ such that AB = CD. Hence,

$$\mathbf{E}(S,S) \gg |S|^2 (p-1)^2 = (p-1)^6 \sim p^2 (p-1)^4 + \frac{(p-1)^8}{p^3} = p^2 |S|^2 + \frac{|S|^4}{p^3}.$$

When the set S is of small size, one would hope to have an upper bound of $\mathbf{E}(S, S)$ that does not depend on p. In this paper, we make use of the following result due to Bourgain and Gamburd in [7] to derive such a bound over prime fields.

Theorem 2.8 (Proposition 2, [7]). Let p be a sufficiently large prime, and let η be a symmetric probability measure on $SL_2(\mathbb{F}_p)$ and $0 < \gamma < \frac{3}{4}$, such that

- (1) $\|\eta\|_{\infty} < p^{-\gamma};$
- (2) $\eta(gH) < p^{\frac{-\gamma}{2}}$ for any proper subgroup $H \subset SL_2(\mathbb{F}_p), g \in SL_2(\mathbb{F}_p);$

(3)
$$\|\eta\|_2 > p^{\frac{-3}{2}+\gamma}$$
.

Then there exists $\epsilon = \epsilon(\gamma) > 0$ such that

$$\|\eta * \eta\|_2 < p^{-\epsilon} \|\eta\|_2.$$

Moreover, following the proof of Theorem 2.8, we have $\epsilon < \gamma$.

For $0 < \gamma < \frac{3}{4}$, let S be a symmetric subset of $\operatorname{SL}_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and for any proper subgroup $H \subsetneq \operatorname{SL}_2(\mathbb{F}_p)$, $g \in \operatorname{SL}_2(\mathbb{F}_p)$ we have $|S \cap gH| < p^{\frac{-\gamma}{2}}|S|$. Let $\mu_S : \operatorname{SL}_2(\mathbb{F}_p) \to \mathbb{R}$ be the function defined by $\mu_S(g) = \frac{1}{|S|}$ if $g \in S$ and $\mu(g) = 0$ if $g \notin S$. Then, μ_S satisfies all conditions of Theorem 2.8. Indeed, we have

- (1) $\|\mu_S\|_{\infty} = \max_g \mu_S(g) = \frac{1}{|S|} < p^{-\gamma},$
- (2) $\mu_S(gH) = \frac{|S \cap gH|}{|S|} < p^{\frac{-\gamma}{2}}$, for any proper subgroup $H \subset SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$,

(3)
$$\|\mu_S\|_2 = \left(\sum_{g \in \mathrm{SL}_2(\mathbb{F}_p)} \mu_S(g)^2\right)^{\frac{1}{2}} = \frac{1}{|S|^{\frac{1}{2}}} > p^{\frac{-3}{2} + \gamma}.$$

Therefore,

$$\|\mu_S * \mu_S\|_2 < p^{-\epsilon} \|\mu_S\|_2 = p^{-\epsilon} \frac{1}{|S|^{\frac{1}{2}}}.$$

One the other hand, $\mu_S * \mu_S(g) = \frac{|\{(a,b) \in S \times S : ab=g\}|}{|S|^2}$. Then,

$$\|\mu_S * \mu_S\|_2^2 = \frac{\mathbf{E}(S,S)}{|S|^4}.$$

In other words, we have proved the following corollary.

Corollary 2.9. For any $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds. Let p be a sufficiently large prime, let S be a symmetric subset of $SL_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and $|S \cap gH| < p^{\frac{-\gamma}{2}}|S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$. Then, we have

$$\boldsymbol{E}(S,S) < \frac{|S|^3}{p^{\epsilon}}.$$
(6)

Proof. For any $\gamma \in (0, \frac{3}{4})$, it follows directly from the previous computation. For any $\gamma \in [3/4, 1)$, we observe that

$$(p^{\gamma}, p^{3-2\gamma}) \subset (p^{\gamma'}, p^{3-2\gamma'}), \ p^{-\frac{\gamma}{2}}|S| \le p^{-\frac{\gamma'}{2}}|S|,$$

for any $\gamma' \in (0, 3/4)$. Then, by choosing $\epsilon = \epsilon(\gamma')$ for any $\gamma' \in (0, 3/4)$, the corollary follows. \Box

Let $A \times B \subseteq \mathbb{F}_p^2 \times \mathbb{F}_p^2$ and $S \subseteq \mathrm{SL}_2(\mathbb{F}_p)$. Then,

$$I(A \times B, S) \le N^{\frac{1}{2}} |A|^{\frac{1}{2}},$$

where N is the number of $(b, b', \theta, \theta') \in B \times B \times S \times S$ such that $\theta b = \theta' b'$.

Indeed, for each $(a,b) \in A \times B$, denote $s_{(a,b)}$ as the number of $\theta \in S$ such that $\theta b = a$. Therefore,

$$I(P,S) = \sum_{a \in A} \left(\sum_{b \in B} s_{(a,b)} \right) \le |A|^{\frac{1}{2}} \left(\sum_{a \in A} \left(\sum_{b \in B} s_{(a,b)} \right)^2 \right)^{\frac{1}{2}}$$
$$= |A|^{\frac{1}{2}} \left(\sum_{a \in A} \left| \left\{ \left(b, b', \theta, \theta' \right) \in B \times B \times S \times S : \theta b = \theta' b' = a \right\} \right| \right)^{\frac{1}{2}}$$
$$\le |A|^{\frac{1}{2}} \cdot N^{\frac{1}{2}}.$$

To bound N, we observe that the equation $\theta b = \theta' b'$ gives $(\theta')^{-1} \theta b = b'$. So, N can be viewed as the number of incidences between $B \times B$ and the multi-set $S^{-1}S$. If $(0,0) \notin B$, by following the proof of Theorem 2.1 identically, we obtain

$$N \ll \frac{|B|^2 |S|^2}{p^2} + p |B| (\mathbf{E}(S,S))^{\frac{1}{2}},$$

and if any line passing through the origin contains at most k points from B, then

$$N \ll \frac{|B|^2 |S|^2}{p^2} + p^{\frac{1}{2}} k^{\frac{1}{2}} |B| (\mathbf{E}(S,S))^{\frac{1}{2}}.$$

As mentioned in (4),

$$\mathbf{E}(S,S) \ll p^2 |S|^2 + \frac{|S|^4}{p^3}.$$

Substituting this bound into $I(A \times B, S)$ implies

$$I(A \times B, S) \ll \frac{|A|^{\frac{1}{2}}|B||S|}{p} + \frac{|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|}{p^{\frac{1}{4}}} + p|A|^{\frac{1}{2}}|S|^{\frac{1}{2}}|B|^{\frac{1}{2}}.$$

Compared to the bound of Theorem 2.1, this result is weaker.

However, if we use Corollary 2.9 instead, then

$$I(P,S) \ll \frac{|A|^{\frac{1}{2}}|B||S|}{p} + p^{\frac{2-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}}.$$

Moreover, if any line passing through the origin contains at most k points from B, then

$$I(P,S) \ll \frac{|A|^{\frac{1}{2}}|B||S|}{p} + k^{\frac{1}{4}}p^{\frac{1-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}}.$$

This completes the proof of Theorem 2.2.

2.3 Two alternative approaches yield weaker bounds

This section presents two different approaches without techniques from Fourier analysis. Although the resulting bounds are weaker compared to those of Theorem 2.1 and Theorem 2.2, the methods will be useful for us when studying the case of small sets.

Theorem 2.10. For any $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds. Let p be a sufficiently large prime. Let $P = A \times B \subseteq (\mathbb{F}_p^2 \times \mathbb{F}_p^2) \setminus \{(0,0,0,0)\}$, and let S be a symmetric subset of $SL_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and $|S \cap gH| < p^{\frac{-\gamma}{2}}|S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and $g \in SL_2(\mathbb{F}_p)$. Assume any line passing through the origin contains at most k points from B. Then,

(1) if $|B| < k^{\frac{1}{2}}p$, we have

$$I(P,S) \lesssim k^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + k^{\frac{1}{4}} p^{\frac{1-\epsilon}{4}} |A|^{\frac{1}{2}} |B|^{\frac{1}{2}} |S|^{\frac{3}{4}},$$

(2) if $|B| \ge k^{\frac{1}{2}}p$, we have

$$I(P,S) \lesssim \frac{|A|^{\frac{1}{2}}|B||S|^{\frac{3}{4}}}{p^{\frac{1+\epsilon}{4}}}.$$

Compared to the bound of Theorem 2.2, which is

$$I(P,S) \ll \frac{|A|^{\frac{1}{2}}|B||S|}{p} + k^{\frac{1}{4}}p^{\frac{1-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}},$$

we can see that

• if $|B| < k^{\frac{1}{2}}p$, then

$$\frac{|A|^{\frac{1}{2}}|B||S|}{p} < k^{\frac{1}{2}}|A|^{\frac{1}{2}}|S|,$$

• if $|B| \ge k^{\frac{1}{2}}p$, then

$$\frac{|A|^{\frac{1}{2}}|B||S|}{p}, k^{\frac{1}{4}}p^{\frac{1-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}} \le \frac{|A|^{\frac{1}{2}}|B||S|^{\frac{3}{4}}}{p^{\frac{1+\epsilon}{4}}},$$

since $|S| < p^{3-2\gamma} < p^{3-\epsilon}$. In other words, Theorem 2.2 is better than Theorem 2.10.

Proof of Theorem 2.10. Since the identity matrix I contributes at most

$$\min\{|A|, |B|\} \ll k^{\frac{1}{4}} p^{\frac{1-\epsilon}{4}} |A|^{\frac{1}{2}} |B|^{\frac{1}{2}} |S|^{\frac{3}{4}}$$

incidences to I(P, S), we may assume without loss generality that $I \notin S$.

For a set $D \subset \mathbb{F}_p^2 \times \mathbb{F}_p^2$ and $\theta \in SL_2(\mathbb{F}_p)$, by $i_D(\theta)$, we mean the number of incidences between θ and the set D.

We have

$$I(P,S) = \sum_{\theta \in S} i_P(\theta) \le |A|^{\frac{1}{2}} \left(I(B \times B, S') \right)^{\frac{1}{2}},$$

where S' be the multi-set of elements of the form $a^{-1}b$, where $a, b \in S$. Set $E = B \times B$. For $\theta \in S'$, let $m(\theta)$ be the multiplicity of θ in S', and for $(x, y) \in E$, let n(x, y) be the number of elements $(u, v) \in E$ such that $(u, v) = \lambda(x, y)$ for some $\lambda \in \mathbb{F}_p^*$.

We now bound the number of incidences between S' and $B \times B$. By $\overline{S'}$, we mean the set of distinct elements in S. We have

$$I(B \times B, S') = \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E} \theta(x,y),$$

where $\theta(x, y) = 1$ if $\theta y = x$, and 0 otherwise.

We now write

$$\begin{split} I(B \times B, S') &= \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E} \theta(x,y) \\ &= \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) = 1} \theta(x,y) + \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) > 1} \theta(x,y) = I + II. \end{split}$$

Let E' be the set of $(x, y) \in E$ such that n(x, y) = 1. To bound I,

$$I = \sum_{\theta \in \overline{S'}, \ i_{E'}(\theta) > 1} m(\theta) \sum_{(x,y) \in E, n(x,y) = 1} \theta(x,y) + \sum_{\theta \in \overline{S'}, \ i_{E'}(\theta) = 1} m(\theta) \sum_{(x,y) \in E, n(x,y) = 1} \theta(x,y) = I_1 + I_2.$$

It is clear that $I_2 \leq |S'| = |S|^2$. By the Cauchy-Schwarz inequality and Lemma 2.7, we have

$$\begin{split} I_{1} &= \sum_{\theta \in \overline{S'}, i_{E'}(\theta) > 1} m(\theta) i_{E'}(\theta) \leq \left(\sum_{\theta \in \overline{S'}, i_{E'}(\theta) > 1} m(\theta)^{2} \right)^{\frac{1}{2}} \left(\sum_{\theta \in \overline{S'}, i_{E'}(\theta) > 1} i_{E'}(\theta)^{2} \right)^{\frac{1}{2}} \\ &\ll (\mathbf{E}(S, S))^{\frac{1}{2}} \left(\sum_{\theta \in \overline{S'}, i_{E'} > 1} \binom{i_{E'}(\theta)}{2} \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S, S))^{\frac{1}{2}} \left(\left| \left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in E : (x_{1}, y_{1}) \neq (x_{2}, y_{2}), \exists \theta \in \overline{S'}, \theta \text{ is incident to } (x_{1}, y_{1}) \text{ and } (x_{2}, y_{2}) \right\} \right| \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S, S))^{\frac{1}{2}} \left(\left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in B^{4} : \exists \theta \in \overline{S'}, \theta x_{1} = y_{1}, \theta x_{2} = y_{2} \right\} \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S, S))^{\frac{1}{2}} \left(\left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in B^{4} : x_{1} \cdot x_{2}^{\perp} = y_{1} \cdot y_{2}^{\perp} \right\} \right)^{\frac{1}{2}} \\ &\ll (\mathbf{E}(S, S))^{\frac{1}{2}} \cdot \left(\frac{|B|^{2}}{p^{\frac{1}{2}}} + k^{\frac{1}{2}} p^{\frac{1}{2}} |B| \right). \end{split}$$

1

1

Thus, using Corollary 2.9, we obtain

$$I \le |S|^2 + \left(\frac{|B|^2}{p^{\frac{1}{2}}} + k^{\frac{1}{2}}p^{\frac{1}{2}}|B|\right)|S|^{\frac{3}{2}}p^{\frac{-\epsilon}{2}}.$$

We now consider II.

$$II = \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) > 1} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, 2^i \le n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 1} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 1} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 1} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 1} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) \in E, n(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta) \sum_{(x,y) < 2^{i+1}} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} m(\theta)$$

By using pigeonhole principle, there exists $\ell = 2^{i_0}$ such that

$$II \lesssim \sum_{\theta \in \overline{S'}} m(\theta) \sum_{(x,y) \in E, \ell \leq n(x,y) < 2\ell} \theta(x,y).$$

We say two pairs (x, y) and (x', y') are in the same congruence class if there exists $\lambda \in \mathbb{F}_p^*$ such that $(x, y) = \lambda(x', y')$. Define E'' to be the set of congruence classes [(x, y)] in E such that $\ell \leq n(x, y) < 2\ell$.

Then we have

$$II \lesssim \ell \sum_{\theta \in \overline{S'}} m(\theta) i_{E''}(\theta) = II_1 + II_2.$$

Here

$$II_1 = \ell \sum_{\theta \in \overline{S'}, \ i_{E''}(\theta) = 1} m(\theta) i_{E''}(\theta), \quad II_2 = \ell \sum_{\theta \in \overline{S'}, \ i_{E''}(\theta) \geq 2} m(\theta) i_{E''}(\theta).$$

As above, we have

 $II_1 \le \ell |S|^2,$

and

$$\begin{split} II_{2} &\leq ((\mathbf{E}(S,S)))^{\frac{1}{2}} \, \ell \left(\sum_{\theta \in \overline{S'}, \ i_{E''}(\theta) \geq 2} i_{E''}(\theta)^{2} \right)^{\frac{1}{2}} \ll (\mathbf{E}(S,S))^{\frac{1}{2}} \left(\sum_{\theta \in \overline{S'}, i_{E''}(\theta) > 1} \ell^{2} \binom{i_{E''}(\theta)}{2} \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S,S))^{\frac{1}{2}} \left(\ell^{2} \left| \left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in E'' \times E'' : \exists \theta \in \overline{S'}, \theta \text{ is incident to } (x_{1}, y_{1}) \text{ and } (x_{2}, y_{2}) \right\} \right| \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S,S))^{\frac{1}{2}} \left(\left| \left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in B \times B \times B \times B : \exists \theta \in \overline{S'}, \theta y_{1} = x_{1}, \theta y_{2} = x_{2} \right\} \right| \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S,S))^{\frac{1}{2}} \left(\left| \left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in B \times B \times B \times B : x_{1} \cdot x_{2}^{\perp} = y_{1} \cdot y_{2}^{\perp} \right\} \right| \right)^{\frac{1}{2}} \\ &\leq (\mathbf{E}(S,S))^{\frac{1}{2}} \cdot \left(\frac{|B|^{2}}{p^{\frac{1}{2}}} + k^{\frac{1}{2}} p^{\frac{1}{2}} |B| \right). \end{split}$$

Putting these bounds together implies

$$II \lesssim \ell |S|^2 + (\mathbf{E}(S,S))^{\frac{1}{2}} \left(\frac{|B|^2}{p^{\frac{1}{2}}} + k^{\frac{1}{2}} p^{\frac{1}{2}} |B| \right).$$

Notice that $\ell \leq k$. So,

$$I(B \times B, S') \lesssim k|S|^2 + \left(\frac{|B|^2}{p^{\frac{1}{2}}} + k^{\frac{1}{2}}p^{\frac{1}{2}}|B|\right)|S|^{\frac{3}{2}}p^{\frac{-\epsilon}{2}},$$

and then

$$I(P,S) \lesssim k^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + k^{\frac{1}{4}} p^{\frac{1-\epsilon}{4}} |A|^{\frac{1}{2}} |B|^{\frac{1}{2}} |S|^{\frac{3}{4}} + p^{\frac{-1-\epsilon}{4}} |A|^{\frac{1}{2}} |B||S|^{\frac{3}{4}}.$$

A direct computation implies that

- 1. if $|B| < k^{\frac{1}{2}}p$, then $I(P,S) \lesssim k^{\frac{1}{2}}|A|^{\frac{1}{2}}|S| + k^{\frac{1}{4}}p^{\frac{1-\epsilon}{4}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{3}{4}};$
- 2. if $|B| \ge k^{\frac{1}{2}}p$, then

 $I(P,S) \lesssim p^{\frac{-1-\epsilon}{4}} |A|^{\frac{1}{2}} |B| |S|^{\frac{3}{4}}.$

This completes the proof.

Theorem 2.11. Let p be a prime and $P = A \times B \subseteq (\mathbb{F}_p^2 \times \mathbb{F}_p^2) \setminus \{(0,0,0,0)\}$. Assume any line passing through the origin contains at most k points from B. Then, we have

$$I(P,S) \lesssim \frac{|A||B||S|^{\frac{1}{2}}}{p^{\frac{1}{2}}} + k^{\frac{1}{2}}p^{\frac{1}{2}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{1}{2}} + k|S|.$$

Compared to the bound of Theorem 2.1, which is

$$I(P,S) \ll \frac{|A||B||S|}{p^2} + k^{\frac{1}{2}}p^{\frac{1}{2}}|A|^{\frac{1}{2}}|B|^{\frac{1}{2}}|S|^{\frac{1}{2}},$$

we can see that

$$\frac{|A||B||S|}{p^2} \le \frac{|A||B||S|^{\frac{1}{2}}}{p^{\frac{1}{2}}},$$

under $|S| \le p^3 - p$. Therefore, Theorem 2.1 is better than Theorem 2.11.

Proof of Theorem 2.11. Since the identity matrix I contributes at most $\min\{|A|, |B|\}$ incidences to I(P, S), we may assume without loss generality that $I \notin S$. For $\theta \in S$ and set $D \subset \mathbb{F}_p^2 \times \mathbb{F}_p^2$, by $i_D(\theta)$ we mean the number of pairs $(x, y) \in D$ such that $\theta y = x$. For $(x, y) \in P$, let n(x, y) be the number of elements $(u, v) \in P$ such that $(u, v) = \lambda(x, y)$ for some $\lambda \in \mathbb{F}_p^*$. Then

$$I(P,S) \le \sum_{\theta \in S} i_P(\theta) = \sum_{\theta \in S} \sum_{(x,y) \in P, n(x,y)=1} \theta(x,y) + \sum_{\theta \in S} \sum_{(x,y) \in P, n(x,y)>1} \theta(x,y)$$
$$=: I_1 + I_2$$

where $\theta(x, y) = 1$ if $\theta y = x$, and 0 ortherwise. Let P' be the set of $(x, y) \in P$ such that n(x, y) = 1. By the Cauchy-Schwarz inequality and Lemma 2.7

$$\begin{split} I_1 &= \sum_{\theta \in S} i_{P'}(\theta) \le |S|^{\frac{1}{2}} \left(\sum_{\theta \in S} i_{P'}(\theta)^2 \right)^{\frac{1}{2}} \ll |S|^{\frac{1}{2}} \left(\sum_{\theta \in S} \binom{i_{P'}(\theta)}{2} + |S| \right)^{\frac{1}{2}} \\ &\le |S|^{\frac{1}{2}} \left(|\{(x_1, y_1, x_2, y_2) \in A \times B \times A \times B \colon \exists \theta \in S, \theta y_1 = x_1, \theta y_2 = x_2\}| + |S|)^{\frac{1}{2}} \\ &\le |S|^{\frac{1}{2}} \left(\left| \left\{ (x_1, y_1, x_2, y_2) \in A \times B \times A \times B \colon x_1 \cdot x_2^{\perp} = y_1 \cdot y_2^{\perp} \right\} \right| + |S|)^{\frac{1}{2}} \\ &\ll |S|^{\frac{1}{2}} \left(\frac{|A||B|}{p^{\frac{1}{2}}} + (pk|A||B|)^{\frac{1}{2}} + |S|^{\frac{1}{2}} \right). \end{split}$$

We now consider I_2 ,

$$I_2 = \sum_{\theta \in S} \sum_{(x,y) \in P, n(x,y) > 1} \theta(x,y) = \sum_{i=1}^{\lfloor \log_2(k) \rfloor} \sum_{\theta \in S} \sum_{(x,y) \in P, 2^i \le n(x,y) < 2^{i+1}} \theta(x,y).$$

By using pigeonhole principle, there exists $\ell = 2^{i_0}$ such that

$$I_2 \lesssim \sum_{\theta \in S} \sum_{(x,y) \in P, \ell \le n(x,y) < 2\ell} \theta(x,y).$$

Define P'' to be the set of representatives (x, y) in P such that $\ell \leq n(x, y) < 2\ell$. Then, by the

Cauchy-Schwarz inequality, Lemma 2.7 and note that $k \ge \ell$ we have

$$\begin{split} I_{2} &\lesssim \ell \sum_{\theta \in S} i_{P''}(\theta) \leq \ell |S|^{\frac{1}{2}} \left(\sum_{\theta \in S} i_{P''}(\theta)^{2} \right)^{\frac{1}{2}} \ll |S|^{\frac{1}{2}} \left(\sum_{\theta \in S} \ell^{2} \binom{i_{P''}(\theta)}{2} + \ell^{2} |S| \right)^{\frac{1}{2}} \\ &\leq |S|^{\frac{1}{2}} \left(\ell^{2} \left| \left\{ (x, y) \in (P'')^{2} : \exists \theta \in S, \theta \text{ is incident to } x \text{ and } y \right\} \right| + \ell^{2} |S| \right)^{\frac{1}{2}} \\ &\leq |S|^{\frac{1}{2}} \left(|\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in A \times B \times A \times B : \exists \theta \in S, \theta y_{1} = x_{1}, \theta y_{2} = x_{2} \} | + \ell^{2} |S| \right)^{\frac{1}{2}} \\ &\leq |S|^{\frac{1}{2}} \left(\left| \left\{ (x_{1}, y_{1}, x_{2}, y_{2}) \in A \times B \times A \times B : x_{1} \cdot x_{2}^{\perp} = y_{1} \cdot y_{2}^{\perp} \right\} \right| + \ell^{2} |S| \right)^{\frac{1}{2}} \\ &\ll |S|^{\frac{1}{2}} \left(\frac{|A||B|}{p^{\frac{1}{2}}} + (pk|A||B|)^{\frac{1}{2}} + k |S|^{\frac{1}{2}} \right), \end{split}$$

Putting these bounds together implies

$$I(P,S) \lesssim |S|^{\frac{1}{2}} \left(\frac{|A||B|}{p^{\frac{1}{2}}} + (pk|A||B|)^{\frac{1}{2}} + k|S|^{\frac{1}{2}} \right).$$

as desired.

2.4 Incidence bounds for small sets (Theorem 2.3)

A skew dot-product energy estimate

Given $B \subset \mathbb{F}_p^2$, this section is devoted to study the magnitude of the set

$$\{(x, y, u, v) \in B^4 \colon x \cdot y^\perp = u \cdot v^\perp\}$$

$$\tag{7}$$

when the size of B is small.

When the set B is of large size, Lemma 2.7 can be used to show that the number of such quadruples is almost the expected value $|B|^4/p$. However, when the size of B is small, say, |B| < p, we have to deal with degenerate structures.

More precisely, let ℓ be a line passing through the origin and B be a subset of $\ell \setminus \{(0,0)\}$, then the number of tuples $(x, y, u, v) \in B^4$ such that $x \cdot y^{\perp} = u \cdot v^{\perp}$ is $|B|^4$. Note that if we remove the ' \perp ' sign, then it will be at most $|B|^3$.

This example shows that we need to have some conditions on the structures of B so that a non-trivial estimate can be obtained.

If B has a few distinct directions through the origin, then, by following Rudnev's argument in [25, Section 3] identically, one obtains

Lemma 2.12. Let B be a subset in \mathbb{F}_p^2 with $|B| \leq p$. Assume any line passing through the origin contains at most k_1 points from B and B determines at most k_2 distinct directions through the origin. The number of quadruples $(x, y, u, v) \in B^4$ such that $x \cdot y^{\perp} = u \cdot v^{\perp}$ is at most $k_1^{\frac{1}{2}}|B|^3 + k_1k_2|B|^2 + k_1^2|B|^2$.

If we only assume an assumption on the number of points on a line passing through the origin, then we have the following lemma.

Lemma 2.13. Let B be a subset in \mathbb{F}_p^2 with $|B| \leq p^{\frac{8}{15}}$. Assume any line passing through the origin contains at most k points from B. The number of quadruples $(x, y, u, v) \in B^4$ such that $x \cdot y^{\perp} = u \cdot v^{\perp}$ is at most $k^{\frac{4}{15}} |B|^{\frac{748}{225}} + k^2 |B|^2$.

Proof of Lemma 2.13

The key ingredient in the proof of Lemma 2.13 is the multi-set version of a point-line incidence bound due to Stevens and De Zeeuw in [26].

Theorem 2.14 (Stevens-de Zeeuw, [26]). Let P be a point set and L a set of lines in \mathbb{F}_p^2 . If $|P| \ll p^{\frac{8}{5}}$, then the number of incidences between P and L, denoted by I(P,L), satisfies

$$I(P,L) \ll |P|^{\frac{11}{15}} |L|^{\frac{11}{15}} + |P| + |L|.$$

Theorem 2.15. Let P be a multi-set of points and L be a multi-set of lines in \mathbb{F}_p^2 . We denote the set of distinct points in P by \overline{P} and the set of distinct lines in L by \overline{L} . For $p \in \overline{P}$ and $\ell \in \overline{L}$, let m(p) and $m(\ell)$ be the multiplicity of p and ℓ , respectively. If $|P| = \sum_{p \in \overline{P}} m(p) \ll p^{\frac{8}{5}}$, then

$$I(P,L) \lesssim |P|^{\frac{7}{15}} |L|^{\frac{7}{15}} \left(\sum_{p \in \overline{P}} m(p)^2 \right)^{\frac{4}{15}} \left(\sum_{\ell \in \overline{L}} m(\ell)^2 \right)^{\frac{4}{15}} + |P| + |L|.$$
(8)

Proof. Our argument to prove this theorem is similar to proof of [17, Lemma 2.12].

Let L_k be the set of lines in \overline{L} of multiplicity $\sim 2^k$, and P_k be the set of points in \overline{P} of multiplicity $\sim 2^k$. Set

$$Q_1 := \sum_{p \in \overline{P}} m(p)^2$$
 and $Q_2 := \sum_{\ell \in \overline{L}} m(\ell)^2$.

Then, it is clear that

$$\sum_{k} 2^{k} |P_{k}| = |P|, \qquad \sum_{k} 2^{2k} |P_{k}| = Q_{1},$$

and

$$\sum_{k} 2^{k} |L_{k}| = |L|, \qquad \sum_{k} 2^{2k} |L_{k}| = Q_{2}.$$

Thus, we have

$$|P_k| \le \min\left\{\frac{|P|}{2^k}, \frac{Q_1}{2^{2k}}\right\}$$
 and $|L_k| \le \min\left\{\frac{|L|}{2^k}, \frac{Q_2}{2^{2k}}\right\}$

We now observe

$$\begin{split} I(P,L) &= \sum_{i,j} I(P_i,L_j) = \sum_{\substack{2^j < Q_2/|L|}} 2^j I(P,L_j) + \sum_{\substack{2^j \ge Q_2/|L|}} I(P,L_j) \\ &= \sum_{\substack{2^i < Q_1/|P|\\ 2^j < Q_2/|L|}} 2^i 2^j I(P_i,L_j) + \sum_{\substack{2^i \ge Q_1/|P|\\ 2^j < Q_2/|L|}} 2^i 2^j I(P_i,L_j) \\ &+ \sum_{\substack{2^i < Q_1/|P|\\ 2^j \ge Q_2/|L|}} 2^i 2^j I(P_i,L_j) + \sum_{\substack{2^i \ge Q_1/|P|\\ 2^j \ge Q_2/|L|}} 2^i 2^j I(P_i,L_j) \\ &=: I + II + III + IV. \end{split}$$

Bounding I: Since $|P_i| \leq |P|/2^i$ and $|L_j| \leq |L|/2^j$, by Theorem 2.14, we obtain

$$I \ll \sum_{\substack{2^i < Q_1/|P|\\2^j < Q_2/|L|\\ \lesssim |P|^{\frac{7}{15}}|L|^{\frac{7}{15}}(Q_1Q_2)^{\frac{4}{15}} + |P| + |L| = \sum_{\substack{2^i < Q_1/|P|\\2^j < Q_2/|L|}} 2^{\frac{4(i+j)}{15}}(|P||L|)^{\frac{11}{15}} + |P| + |L|$$

Bounding II: Using $|P_i| \leq Q_{\frac{1}{2}}^{2i}$ and $|L_j| \leq |L|/2^j$, Theorem 2.14 implies

$$\begin{split} II &:= \sum_{\substack{2^i \ge Q_1/|P|\\2^j < Q_2/|L|}} 2^i 2^j I(P_i, L_j) \le \sum_{\substack{2^i \ge Q_1/|P|\\2^j < Q_2/|L|}} 2^i 2^j \left(\frac{Q_1}{2^{2i}}\right)^{\frac{11}{15}} \left(\frac{|L|}{2^j}\right)^{\frac{11}{15}} + |P| + |L| \\ &\lesssim |P|^{\frac{7}{15}} |L|^{\frac{7}{15}} (Q_1 Q_2)^{\frac{4}{15}} + |P| + |L|. \end{split}$$

Bounding III, IV: Similarly, we also have

$$III, IV \lesssim |P|^{\frac{7}{15}} |L|^{\frac{7}{15}} (Q_1 Q_2)^{\frac{4}{15}} + |P| + |L|.$$

This concludes the proof of the theorem.

With this incidence bound, we are ready to prove Theorem 2.13.

Proof of Theorem 2.13. We have

$$\begin{split} \left| \{ (x, y, u, v) \in B^4 \colon x \cdot y^{\perp} = u \cdot v^{\perp} \} \right| &= \left| \{ (x, y, u, v) \in B^4 \colon x \cdot y^{\perp} = u \cdot v^{\perp} \neq 0 \} \right| \\ &+ \left| \{ (x, y, u, v) \in B^4 \colon x \cdot y^{\perp} = u \cdot v^{\perp} = 0 \} \right| \\ &=: I + II \end{split}$$

Regarding *I*. For $a, u, v \in B$, $u \cdot v^{\perp} \neq 0$ we define $L_{a,u,v}$ to be the multi-set of lines of the form $x \cdot a^{\perp} = u \cdot v^{\perp}$. Let $L = \bigcup_{a,u,v \in B} L_{a,u,v}$. It is clear that the number of such quadruples is at most I(B, L). We note that $|L| \leq |B|^3$. By applying Theorem 2.14, we observe that for each line in L,

its multiplicity is at most $\ll k|B|^{\frac{22}{15}}$. For each $l \in L$, let m(l) be the multiplicity l, we have

$$\sum_{\ell} m(\ell) = |L| \le |B|^3,$$

and

$$\sum_{\ell} m(\ell)^2 \le \max_{\ell} m(\ell) \cdot |B|^3 \ll k|B|^{\frac{22}{15}} \cdot |B|^3.$$

It follows from Theorem 2.15 for B and L that

$$I(B,L) \lesssim |B|^{\frac{7}{15}} |B|^{\frac{21}{15}} |B|^{\frac{4}{15}} \left(k |B|^{\frac{67}{15}}\right)^{\frac{4}{15}} + |B| + |L| \ll k^{\frac{4}{15}} |B|^{\frac{748}{225}}$$

Regarding II. For each $x, u \in B$, there are at most k points y and k points v such that $x \cdot y^{\perp} = u \cdot v^{\perp} = 0$. So

$$II \le k^2 \left| B \right|^2.$$

This completes the proof.

Proof of Theorem 2.3

We follow the proof of Theorem 2.10 with Lemma 2.12 and Lemma 2.13 in place of Lemma 2.7, one has two following bounds

$$I(P,S) \lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{8}} |B|^{\frac{3}{4}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{4}} k_2^{\frac{1}{4}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}},$$

and

$$I(P,S) \lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{15}} |B|^{\frac{187}{225}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}}.$$

This completes the proof. \Box

For applications, a comparison of these two incidence bounds, with $|B| \leq p$, is provided as follows.

1. If $|B| \ge k_2 k_1^{\frac{1}{2}}$, then $I(P,S) \lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{n^{\frac{6}{4}}} + \frac{k_1^{\frac{1}{8}} |B|^{\frac{3}{4}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{n^{\frac{6}{4}}}.$

$$\begin{aligned} 2. \ \text{If} \ k_1^{\frac{165}{298}} k_2^{\frac{225}{298}} &\leq |B| < k_2 k_1^{\frac{1}{2}}, \text{ then} \\ I(P,S) &\lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{4}} k_2^{\frac{1}{4}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} \end{aligned}$$

3. If $|B| < k_1^{\frac{165}{298}} k_2^{\frac{225}{298}}$, then $I(P,S) \lesssim k_1^{\frac{1}{2}} |A|^{\frac{1}{2}} |S| + \frac{k_1^{\frac{1}{2}} |B|^{\frac{1}{2}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}} + \frac{k_1^{\frac{1}{15}} |B|^{\frac{187}{225}} |A|^{\frac{1}{2}} |S|^{\frac{3}{4}}}{p^{\frac{\epsilon}{4}}}.$

2.5 Alternative approach with relaxed conditions

This section is devoted to prove the following analog, which offers a bound which is meaningful when the size of S is large compared to the sizes of A and B.

Theorem 2.16. Let p be a prime and $P = A \times B \subseteq (\mathbb{F}_p^2 \times \mathbb{F}_p^2) \setminus \{0\}$ with $|B| \leq |A| \leq p^{\frac{8}{15}}$ and any lines passing through the origin contains at most k points from B. Then, we have

$$I(P,S) \lesssim k^{\frac{2}{15}} \left|A\right|^{\frac{11}{15}} \left|B\right|^{\frac{209}{225}} \left|S\right|^{\frac{1}{2}} + k|A|^{\frac{1}{2}} |B|^{\frac{1}{2}} |S|^{\frac{1}{2}} + k \left|S\right|.$$

Theorem 2.16 is proved by following the proof of Theorem 2.11 and the next lemma, whose proof is identical to that of Lemma 2.13.

Lemma 2.17. Let $A, B \subset \mathbb{F}_p^2$ with $|B| \leq |A| \leq p^{\frac{8}{15}}$. Assume any line passing through the origin contains at most k points from B and at most k points from A. The number of quadruples $(x, y, u, v) \in A \times A \times B \times B$ such that $x \cdot y^{\perp} = u \cdot v^{\perp}$ is at most $k^{\frac{4}{15}} |A|^{\frac{22}{15}} |B|^{\frac{418}{225}} + k^2 |A| |B|$.

3 Proof of Propositions 1.3, 1.5 and Theorems 1.4, 1.6

Proof of Proposition 1.3. On the one hand, we apply Theorem 2.1 for S and $P = S(E) \times E$ to obtain

$$I(S(E) \times E, S) \ll \frac{|S(E)| |E| |S|}{p^2} + p\sqrt{|S(E)| |E| |S|} + |S|.$$
(9)

On the other hand, for each $x \in E$ and $\theta \in S$, there exists unique $y \in S(E)$ such that (y, x) is incident to θ . So, we obtain

$$|E||S| = I(S(E) \times E, S) \ll \frac{|S(E)||E||S|}{p^2} + p\sqrt{|S(E)||E||S|} + |S|.$$

Solving this inequality, the theorem follows.

Theorem 1.4 will follow from the two following results.

Theorem 3.1. Let $E \subset \mathbb{F}_p^2$ and $S \subset SL_2(\mathbb{F}_p)$.

- a. If $|E| \ge 4p$ and $|S| \gg p^2$, then there exists $x \in E$ such that $|S(E-x)| \gg p^2$.
- b. If any line through the origin contains at most k points from E, then

$$|S(E)| \gg \min\left\{p^2, \frac{|S||E|}{pk}\right\}.$$

Theorem 3.1 (a) is not a part of Theorem 1.4, but we find it to be of independent interest.

Theorem 3.2. For any $\gamma \in (0,1)$, there exists $\epsilon = \epsilon(\gamma) > 0$ such that the following holds.

Let p be a sufficiently large prime. Let $E \subseteq \mathbb{F}_p^2 \setminus \{(0,0)\}$, and let S be a symmetric subset of $SL_2(\mathbb{F}_p)$ such that $p^{\gamma} < |S| < p^{3-2\gamma}$ and $|S \cap gH| < p^{\frac{-\gamma}{2}}|S|$ for any subgroup $H \subsetneq SL_2(\mathbb{F}_p)$ and

	-	-	-

 $g \in SL_2(\mathbb{F}_p)$. Then, we have

$$|S(E)| \gg \min\left\{p^2, \frac{|S|^{\frac{1}{2}}|E|}{p^{1-\frac{\epsilon}{2}}}\right\}.$$

Moreover, if any line passing through the origin contains at most k points from E, we have

$$|S(E)| \gg \min\left\{p^2, \frac{|S|^{\frac{1}{2}}|E|}{p^{\frac{1-\epsilon}{2}}k^{\frac{1}{2}}}\right\}$$

To prove Theorem 3.1(a), we recall the following result from [16].

Lemma 3.3 (Corollary 10, [16]). Let $E \subset \mathbb{F}_p^2$ with $|E| \ge 4p$. Then there exists a point $x \in E$ such that there are at least p/2 lines passing through x and each line contains at least one other point from E.

Proof of Theorem 3.1.

Part a. By Lemma 3.3, there exists a point $x \in E$ such that there are at least p/2 lines passing through x and each line contains at least one other point from E. From each of these lines, we pick one point which is different from x and let E' be the set of those points. Then we have $|E'| \ge p/2$. We observe that

$$|S(E-x)| \ge |S(E'-x)|.$$

Note that E' - x is a translation of E' by x. So, any line passing through the origin contains at most one point from E' - x. Set E'' = E' - x. We now estimate the size of S(E'') from below.

Set $P = E'' \times S(E'')$ and $P' = \{\lambda \cdot u : u \in P, \lambda \neq 0\}$. We have |P'| = (p-1)|P|. Note that $I(P,S) = |E''||S| = \frac{1}{p-1}I(P',S)$. By Theorem 2.1, we have

$$I(P',S) \ll \frac{|P'||S|}{p^2} + p\sqrt{|P'||S|} + |S|.$$

Putting the upper and lower bounds together, the theorem follows.

Part b. The proof is almost the same, we just need to partition the set E into at most k subsets E_i such that each has the same structure as the set E'' in the Part a. So we omit the details. \Box

Theorem 3.2, Proposition 1.5, and Theorem 1.6 are proved by following the proof of Theorem 1.3, but we use Theorems 2.2 and 2.3 (1) in place of Theorem 2.1.

Remark 3.4. Theorem 2.3 (2) implies the following bound

$$|S(E)| \gtrsim \min\left\{\frac{|E|^{0.337} |S|^{\frac{1}{2}} p^{\epsilon/2}}{k_1^{\frac{2}{15}}}, \frac{|E| |S|^{\frac{1}{2}} p^{\epsilon/2}}{k_1}, \frac{|E|^2}{k_1}\right\},\$$

which improves Theorem 1.6 in the range $|E| \le \min\left\{p^{\frac{8}{15}}, k_1^{0.553}k_2^{0.755}\right\}$.

4 Incidence structures spanned by $\mathbb{H}_1(\mathbb{F}_p)$

As in the case of the special linear group, we define an incidence structure as follows. We say the point $(x, y) \in \mathbb{F}_p^3 \times \mathbb{F}_p^3$ is incident to the matrix $\theta \in \mathbb{H}_1(\mathbb{F}_p)$ if $\theta y = x$.

Theorem 4.1. Let $\epsilon \geq 0$. Let $X \subset \mathbb{H}_1(\mathbb{F}_p)$ and $P = A \times B \subset \mathbb{F}_p^6$. Assume that all points in A and B have the third coordinate non-zero, and for each $(y, z) \in \mathbb{F}_p^2$, we have $|\pi_{23}^{-1}(y, z) \cap B| \leq p^{1-\epsilon}$, then the number of incidences between X and P satisfies

$$\left| I(P,X) - \frac{|P||X|}{p^3} \right| \le p^{\frac{3-\epsilon}{2}} |P|^{\frac{1}{2}} |X|^{\frac{1}{2}}.$$

We now turn our attention to proving Theorem 4.1.

Proof of Theorem 4.1. By repeating the argument as in the case of $SL_2(\mathbb{F}_p)$, we have

$$\begin{split} I(P,X) &= \sum_{\substack{(x,y) \in P \\ \theta \in X}} 1_{\theta x = y} = \frac{1}{p^3} \sum_{m \in \mathbb{F}_p^3} \sum_{\substack{(x,y) \in P, \\ \theta \in X}} \chi \left(m \cdot (\theta y - x) \right) \\ &= \frac{|P| |X|}{p^3} + \frac{1}{p^3} \sum_{m \in \mathbb{F}_p^3 \setminus \{(0,0,0)\}} \sum_{\substack{(x,y) \in P, \\ \theta \in X}} \chi \left(m \cdot (\theta y - x) \right) \\ &= \frac{|P| |X|}{p^3} + p^3 \sum_{m \neq (0,0,0)} \sum_{\theta \in X} \widehat{P} \left(-m, \theta^t m \right). \end{split}$$

By using the Cauchy-Schwarz inequality, one has

$$\sum_{\theta \in X} \sum_{m \neq (0,0,0)} \widehat{P}(-m, \theta^t m) \le |X|^{\frac{1}{2}} \left(\sum_{\theta \in \mathbb{H}_1(\mathbb{F}_p)} \sum_{m_1, m_2 \neq (0,0,0)} \widehat{P}(-m_1, \theta^t m_1) \overline{\widehat{P}(-m_2, \theta^t m_2)} \right)^{\frac{1}{2}}.$$

We now observe

$$\sum_{\theta \in \mathbb{H}_{1}(\mathbb{F}_{p})} \sum_{m_{1}, m_{2} \neq (0, 0, 0)} \widehat{P}(-m_{1}, \theta^{t}m_{1}) \overline{\widehat{P}(-m_{2}, \theta^{t}m_{2})}$$

$$= \frac{1}{p^{12}} \sum_{\theta} \sum_{m_{1}, m_{2} \neq (0, 0, 0)} \sum_{(x_{1}, y_{1}), (x_{2}, y_{2})} P(x_{1}, y_{1}) P(x_{2}, y_{2}) \chi(-m_{1}x_{1} + \theta^{t}m_{1}y_{1}) \chi(m_{2}x_{2} - \theta^{t}m_{2}y_{2})$$

$$= \sum_{m_{1}, m_{2}} - \sum_{m_{1} = (0, 0, 0), m_{2} \neq (0, 0, 0)} - \sum_{m_{1} \neq (0, 0, 0), m_{2} = (0, 0, 0)} - \sum_{m_{1} = m_{2} = (0, 0, 0)}$$

$$= I - II - III - IV.$$

We now estimate each term separately.

Regarding IV, it is clear that it is equal to $|P|^2/p^9$. The terms II and III are the same, and we can see that

$$II = \frac{|P|}{p^{12}} \sum_{\theta} \sum_{m_2 \neq (0,0,0)} \sum_{(x_2,y_2)} P(x_2,y_2) \chi(m_2(x_2 - \theta y_2)),$$

which can be written as

$$\frac{|P|(p^2-1)}{p^{12}} \sum_{\theta} \sum_{(x_2,y_2)\in P, x_2=\theta y_2} 1 - \frac{|P|}{p^{12}} \sum_{\theta} \sum_{(x_2,y_2)\in P, x_2\neq \theta y_2} 1.$$

So, $-II \leq \frac{|P|^2}{p^9}$. Regarding I,

$$I = \frac{1}{p^{12}} \sum_{\theta} \sum_{m_1, m_2 \in \mathbb{F}_p^3} \sum_{(x_1, y_1), (x_2, y_2)} P(x_1, y_1) P(x_2, y_2) \chi(m_1(\theta y_1 - x_1)) \chi(m_2(\theta y_2 - x_2))$$
$$= \frac{1}{p^6} \sum_{\theta} \sum_{(x_1, y_1), (x_2, y_2)} P(x_1, y_1) P(x_2, y_2) \mathbf{1}_{\theta y_1 = x_1} \mathbf{1}_{\theta y_2 = x_2}.$$

To proceed further, we need to count the number of quadruples $(x, y, z) \in B$, $(x', y', z') \in A$, $(u, v, w) \in B$, $(u', v', w') \in A$ such that there exists $\theta = [a, b, c] \in \mathbb{H}_1(\mathbb{F}_p)$ and $\theta(x, y, z) = (x', y', z')$, $\theta(u, v, w) = (u', v', w')$.

For such quadruples, one has

$$yw' + zv' = y'w + vz', \ z(u' - u) + w(x - x') = a(vz - wy), \ z = z', w = w'.$$

From here, as in the case of the special linear group, we need to make use of a number of preliminary lemmas, which will be proved later.

Proposition 4.2. Given $A, B \subset \mathbb{F}_p^3$, let N be the number of quadruples $(x, y, z) \in A, (x', y', z') \in B, (u, v, w) \in A, (u', v', w') \in B$ such that

$$yw' + zv' = y'w + vz', \ z = z', w = w'.$$

Suppose in addition that for each $(y,z) \in \mathbb{F}_p^2$, we have $\pi_{23}^{-1}(y,z) \cap A$ and $\pi_{23}^{-1}(y,z) \cap B$ are of sizes at most $p^{1-\epsilon}$, then we have

$$N \le \frac{|A|^2 |B|^2}{p} + p^{3-2\epsilon} |A||B|$$

Proposition 4.3. Given $A, B \subset \mathbb{F}_p^3$, let N' be the number of quadruples $(x, y, z) \in B, (x', y', z') \in A, (u, v, w) \in B, (u', v', w') \in A$ such that

$$yw' + zv' = y'w + vz', vz - wy = 0, zu' + xw' - z'u - x'w = 0, w = w', z = z'$$

Then $N' \leq p|A||B|$.

Lemma 4.4. Given (x, y, z), (x', y', z'), (u, v, w), (u', v', w') in \mathbb{F}_p^3 with $w \neq 0, z \neq 0$, and yw' + zv' = y'w + z'v. Let N'' be the number of matrices $\theta = [a, b, c] \in \mathbb{H}_1(\mathbb{F}_p)$ such that

$$\theta(x, y, z) = (x', y', z')$$

and

$$\theta(u, v, w) = (u', v', w').$$

- If $vz wy \neq 0$, then N'' = 1.
- If vz wy = 0 and z(u' u) = w(x x'), then N'' = p.
- If vz wy = 0 and $z(u' u) \neq w(x x')$, then N'' = 0.

So, combining Propositions 4.2, 4.3, and Lemma 4.4 gives

$$I \le \frac{|P|^2}{p^3} + p^{3-\epsilon}|P|$$

In other words,

$$I - II - III - IV \ll \frac{1}{p^6} \cdot p^{3-\epsilon} |P|,$$

and

$$\left| I(P,X) - \frac{|P||X|}{p^3} \right| \le p^{\frac{3-\epsilon}{2}} |P|^{\frac{1}{2}} |X|^{\frac{1}{2}}.$$

This completes the proof.

The rest of this section is devoted to prove Propositions 4.2, 4.3, and Lemma 4.4.

We first start with Propositions 4.3 and 4.4, since they are much elementary.

Proof of Proposition 4.3. By fixing $(x, y, z) \in A$ and $(u', v', w') \in B$, then y' and v are determined uniquely by

$$yw' + zv' = y'w + vz', vz - wy = 0, zu' + xw' - z'u - x'w = 0$$

With each $u \in \mathbb{F}_p$, x' is determined uniquely. Thus, $N' \leq p|A||B|$.

Proof of Lemma 4.4. A direct computation shows that

$$b = \frac{v' - v}{w} = \frac{y' - y}{z}.$$

Note that z(u'-u) + w(x-x') = a(vz - wy). If $vz - wy \neq 0$, then

$$a = \frac{z(u'-u) + w(x-x')}{vz - wy}.$$

For such a, c is determined uniquely.

If vz - wy = 0 and z(u' - u) = w(x - x'), there are p such matrices θ .

If vz - wy = 0 and $z(u' - u) \neq w(x - x')$, there is no such matrix θ .

We now move to the proof of Proposition 4.2.

By repeating the proof of [10, Theorem 2.1], the following weighted version can be obtained.

Lemma 4.5. Let U, V be two sets in \mathbb{F}_p^4 , and $F: U \to \mathbb{N}$, $G: V \to \mathbb{N}$. Define

$$M = \sum_{\substack{u \in U, v \in V \\ u \cdot v = 0}} F(u)G(v).$$

Then we have

$$\left| M - \frac{(\sum_{u \in U} F(u))(\sum_{v \in V} G(v))}{p} \right| \le p^2 \left(\sum_{u \in U} |F(u)|^2 \right)^{\frac{1}{2}} \cdot \left(\sum_{v \in V} |G(v)|^2 \right)^{\frac{1}{2}}.$$

In our setting, we partition the set A into A_{λ} and the set B into B_{λ} , for $\lambda \in \mathbb{F}_p$, such that any two points in A_{λ} have the same last coordinate which is equal to λ , and the same applies to B_{λ} .

Recall $\pi_{23} \colon \mathbb{F}_p^3 \to \mathbb{F}_p^2$ be the projection onto the two last coordinates. For $A_{\lambda} \subset \mathbb{F}_p^3$, we define $\overline{A_{\lambda}} = \pi_{23}(A_{\lambda})$. For each $x \in \overline{A_{\lambda}} \subset \mathbb{F}_p^2$, define $f(x) = \pi_{23}^{-1}(x) \cap A_{\lambda}$. For $B_{\lambda} \subset \mathbb{F}_p^3$, we define $\overline{B_{\lambda}} = \pi_{23}(B_{\lambda})$. For each $x \in \overline{B_{\lambda}}$, define $g(x) = \pi_{23}^{-1}(x) \cap B_{\lambda}$.

Let $N_{\lambda,\beta}$ be the number of quadruples $(a, a', b, b') \in \overline{A_{\lambda}} \times \overline{B_{\lambda}} \times \overline{B_{\beta}} \times \overline{B_{\beta}}$ such that $a \cdot b' = a' \cdot b$, counted with multiplicity, i.e.

$$N_{\lambda,\beta} = \sum_{a,b,a',b':\ a\cdot b' = a'\cdot b} f(a)g(a')f(b)g(b')$$

We want to bound $N_{\lambda,\beta}$ from above. The next result will be proved by using the above lemma. Lemma 4.6. For fixed λ, β , we have

$$N_{\lambda,\beta} \leq \frac{1}{p} \left(\sum_{x \in \overline{A_{\lambda}}} f(x) \right) \left(\sum_{x \in \overline{A_{\beta}}} f(x) \right) \left(\sum_{y \in \overline{B_{\lambda}}} g(y) \right) \left(\sum_{y \in \overline{B_{\beta}}} g(y) \right) + p \left(\sum_{x \in \overline{A_{\lambda}}} |f(x)|^2 \right)^{\frac{1}{2}} \left(\sum_{x \in \overline{A_{\beta}}} |f(x)|^2 \right)^{\frac{1}{2}} \left(\sum_{y \in \overline{B_{\lambda}}} |g(y)|^2 \right)^{\frac{1}{2}} \left(\sum_{y \in \overline{B_{\beta}}} |g(y)|^2 \right)^{\frac{1}{2}}.$$

Proof. To prove this lemma, we first enlarge the set $A_{\lambda}, A_{\beta}, B_{\lambda}$, and B_{β} . More precisely, define

$$U = \{t(a, a') \colon a \in \overline{A_{\lambda}}, \ a' \in \overline{B_{\lambda}}, \ t \neq 0\},\$$

and

$$V = \{t(b,b') \colon b \in \overline{A_{\beta}}, \ b' \in \overline{B_{\beta}}, \ t \neq 0\}.$$

Also define

$$F(t(a, a')) = f(a)g(a'), \quad G(t(b, b')) = f(b)g(b')$$

for all $t(a, a') \in U$ and $t(b, b') \in V$. If $(a, a', b, b') \in \overline{A_{\lambda}} \times \overline{B_{\lambda}} \times \overline{A_{\beta}} \times \overline{B_{\beta}}$ such that $a \cdot b' = a' \cdot b$, then

$$(a, a') \cdot (-b', b) = 0,$$

which implies

$$t(a,a') \cdot t'(-b',b) = 0$$

for all $t, t' \neq 0$.

Therefore,

$$N_{\lambda,\beta} = \frac{1}{(p-1)^2}M,$$

where

$$M = \sum_{u \in U, v \in V, u \cdot v = 0} F(u)G(v).$$

Applying Lemma 4.5 gives us

$$\begin{split} N_{\lambda,\beta} &\leq \frac{1}{p} \left(\sum_{x \in \overline{A_{\lambda}}} f(x) \right) \left(\sum_{x \in \overline{A_{\beta}}} f(x) \right) \left(\sum_{y \in \overline{B_{\lambda}}} g(y) \right) \left(\sum_{y \in \overline{B_{\beta}}} g(y) \right) \\ &+ p \left(\sum_{x \in \overline{A_{\lambda}}} |f(x)|^2 \right)^{\frac{1}{2}} \left(\sum_{x \in \overline{A_{\beta}}} |f(x)|^2 \right)^{\frac{1}{2}} \left(\sum_{y \in \overline{B_{\lambda}}} |g(y)|^2 \right)^{\frac{1}{2}} \left(\sum_{y \in \overline{B_{\beta}}} |g(y)|^2 \right)^{\frac{1}{2}}. \end{split}$$

This completes the proof.

Using this lemma, we take the sum over all pairs (λ, β) , and obtain

$$N \leq \frac{|A|^2 |B|^2}{p} + p\left(\sum_{x \in \mathbb{F}_p^2} f(x)^2\right) \cdot \left(\sum_{y \in \mathbb{F}_p^2} g(y)^2\right),$$

by using the Cauchy-Schwarz inequality.

The trivial upper bound that $f(x), g(y) \leq p$ gives

$$N \le \frac{|A|^2|B|^2}{p} + p^3|A||B|.$$

If we assume in addition that either $\max_x f(x) \leq p^{1-\epsilon}$ or $\max_x g(x) \leq p^{1-\epsilon}$, then

$$N \le \frac{|A|^2 |B|^2}{p} + p^{3-\epsilon} |A||B|$$

This completes the proof. \Box

5 Proof of Theorem 1.7

With Theorem 4.1 in hand, one can prove Theorem 1.7 easily. To see this, we set B = E and A = X(E). We now estimate the number of incidences between $A \times B$ and X. It is clear that $I(A \times B, X) = |X||B|$. Applying Theorem 4.1 on the number of incidences between $A \times B$ and X, we obtain

$$I(A \times B, X) \le \frac{|A||B||X|}{p^3} + p^{\frac{3-\epsilon}{2}} |X|^{\frac{1}{2}} |A|^{\frac{1}{2}} |B|^{\frac{1}{2}}.$$

_	_	

Putting the lower and upper bounds together and solving the inequality give us the desired bound.

6 Acknowledgements

Thang Pham would like to thank the Vietnam Institute for Advanced Study in Mathematics (VIASM) for the hospitality and for the excellent working condition.

Norbert Hegyvári was supported by the National Research, Development and Innovation Office NKFIH Grant No K-146387. Alex Iosevich was partially supported by the National Science Foundation, NSF DMS 2154232. Thang Pham was partially supported by ERC Advanced Grant no. 882971, "GeoScape", and by the Erdős Center.

References

- L. Babai, N. Nikolov, and L. Pyber, Product growth and mixing in finite groups, Proceedings of the Nineteenth Annual ACM-SIAM Symposium on Discrete Algorithms, (2008), 248–257.
- [2] A. S. Besicovitch, Sur deux questions d'intégrabilité des fonctions, Journal de la Société de Physique et de Mathématique de l'Université de Perm, 2 (1919), 105–123.
- [3] A. S. Besicovitch, On Kakeya's problem and a similar one, Mathematische Zeitschrift, 27(1) (1927), 312–320.
- [4] A. S. Besicovitch and R. Rado, A plane set of measure zero containing circumferences of every radius, Journal of the London Mathematical Society, 43 (1968), 717–719.
- [5] J. Bourgain, Estimations de certaines fonctions maximales, Comptes rendus de l'Académie des sciences, Série 1, Mathématique, **312**(8) (1985), 499–512.
- [6] J. Bourgain, N. Katz, and T. Tao, A sum-product estimate in finite fields, and applications, Geometric & Functional Analysis, 14 (2004), 27–57.
- [7] J. Bourgain and A. Gamburd, Uniform expansion bounds for Cayley graphs of $SL_2(\mathbb{F}_p)$, Annals of Mathematics, **167** (2008) 625–642.
- [8] D. Covert, D. Hart, A. Iosevich, D. Koh, and M. Rudnev, Generalized incidence theorems, homogeneous forms and sum-product estimates in finite fields, European Journal of Combinatorics, 31(1) (2010), 306—319.
- [9] Z. Dvir, On the size of Kakeya sets in finite fields, Journal of the American Mathematical Society, 22(4) (2009), 1093–1097.
- [10] D. Hart, A. Iosevich, D. Koh, and M. Rudnev. Averages over hyperplanes, sum-product theory in vector spaces over finite fields and the Erdős-Falconer distance conjecture, Transactions of the American Mathematical Society, 363(6) (2011), 3255–3275.
- [11] N. Hegyvári and F. Hennecart, A structure result for bricks in Heisenberg groups, Journal of Number Theory, 133(9) (2013), 2999–3006.
- [12] N. Hegyvári and F. Hennecart, Expansion for cubes in the Heisenberg group, Forum Mathematicum, 30(1) (2018), 227–236.

- [13] A. Iosevich, M. Rudnev, and Y. Zhai, Areas of triangles and Beck's theorem in planes over finite fields, Combinatorica, 35(3) (2015), 295–308.
- [14] A. Iosevich, P. Mattila, E. Palsson, M. Q. Pham, T. Pham, S. Senger, and C. Y. Shen, *Packing sets in euclidean space by affine transformations*, arXiv:2405.03087 (2024).
- [15] J. R. Kinney, A thin set of circles, American Mathematical Monthly, 75 (1968), 1077–1081.
- [16] B. Lund, T. Pham, and L. A. Vinh, Distinct spreads in vector spaces over finite fields, Discrete Applied Mathematics, 239 (2018), 154–158.
- [17] B. Lund and G. Petridis, *Bisectors and Pinned Distances*, Discrete & Computational Geometry, 64 (2020), 995–1012.
- [18] J. M. Marstrand, *Packing circles in the plane*, Proceedings of the London Mathematical Society. Third Series, 55 (1987), 37–58.
- [19] G. Mockenhaupt and T. Tao. Restriction and Kakeya phenomena for finite fields, Duke Mathematical Journal, 121(1) (2004), 35–74.
- [20] G. L. Mullen and D. Panario, Handbook of Finite Fields, Chapman and Hall/CRC, (2013).
- [21] R. Oberlin, Unions of lines in F^n , Mathematika, **62** (2016), 738–752.
- [22] T. Pham and L. A. Vinh, Some combinatorial number theory problems over finite valuation rings, Illinois Journal of Mathematics, 61(1-2) (2017), 243–257.
- [23] T. Pham, Triangles with one fixed side-length, a Furstenberg-type problem, and incidences in finite vector spaces, Forum Mathematicum, in press, (2024).
- [24] T. Pham and S. Yoo, Intersection patterns and incidence theorems, arXiv:2304.08004 (2023).
- [25] M. Rudnev, On the number of incidences between points and planes in three dimensions, Combinatorica, 38 (2018), 219–254.
- [26] S. Stevens and F. Zeeuw, An improved point-line incidence bound over arbitrary fields, Bulletin of the London Mathematical Society, 49(5) (2017), 842–858.
- [27] M. Talagrand, Sur la mesure de la projection d'un compact set certaines families de cercles, Bulletin des Sciences Mathématiques, 104 (1980), 225–231.
- [28] L. A. Vinh. On the volume set of point sets in vector spaces over finite fields, Proceedings of the American Mathematical Society, 141(90) (2013), 3067–3071.
- [29] T. Wolff, A Kakeya-type problem for circles, American Journal of Mathematics, 119(5) (1997), 985–1026.
- [30] T. Wolff, Recent work connected with the Kakeya problem, American Mathematical Society, (1999), 129–162.
- [31] J. Zahl, Union of lines in \mathbb{R}^n , Mathematika, **69**(2) (2023), 473–481.