

ON A FAMILY OF DIVISIBLE DESIGN DIGRAPHS

MIKHAIL MUZYCHUK AND GRIGORY RYABOV

ABSTRACT. For every odd prime power q , a family of pairwise nonisomorphic normal arc-transitive divisible design Cayley digraphs with isomorphic neighborhood designs over a Heisenberg group of order q^3 is constructed. It is proved that these digraphs are not distinguished by the Weisfeiler-Leman algorithm and have the Weisfeiler-Leman dimension 3.

Keywords: divisible design digraphs, Cayley digraphs, Weisfeiler-Leman dimension.

MSC: 05B05, 05C60, 05E30.

1. INTRODUCTION

Let $\Gamma = (\Omega, D)$ be a digraph (possibly, with loops). We say that a vertex $\alpha \in \Omega$ *dominates* a vertex $\beta \in \Omega$ if (α, β) is an arc of Γ . The digraph Γ is said to be *regular* of degree k if each vertex of Γ dominates exactly k vertices and is dominated by exactly k vertices. The digraph Γ is said to be *asymmetric* if $(\beta, \alpha) \notin D$ for all $\alpha, \beta \in \Omega$ such that $\alpha \neq \beta$ and $(\alpha, \beta) \in D$.

A k -regular asymmetric digraph on v vertices is called a *divisible design digraph* (DDD for short) with parameters $(v, k, \lambda_1, \lambda_2, m, n)$ if its vertex set can be partitioned into m classes of size n such that for any two distinct vertices α and β , the number of vertices that dominates or being dominated by both α and β is equal to λ_1 if α and β belong to the same class and λ_2 otherwise. The notion of a divisible design graph was introduced in [14] and extended to digraphs in [7]. Several constructions of DDDs can be found in [6, 7, 14, 17, 21].

DDDs are closely related to symmetric (group) divisible designs (see [2, 24] for the definition). More precisely, if Γ is a DDD, then an incidence structure whose points are vertices and blocks are out-neighborhoods of vertices of Γ is a symmetric divisible design called the *neighborhood design* of Γ and denoted by $\mathcal{D} = \mathcal{D}(\Gamma)$. An adjacency matrix of Γ coincides with an incidence matrix of \mathcal{D} . Observe that even if DDDs Γ_1 and Γ_2 are not isomorphic, the neighborhood designs $\mathcal{D}(\Gamma_1)$ and $\mathcal{D}(\Gamma_2)$ can be isomorphic. For example, if Γ_2 is obtained from Γ_1 by a dual Seidel switching, then $\mathcal{D}(\Gamma_1)$ and $\mathcal{D}(\Gamma_2)$ are isomorphic, whereas Γ_1 and Γ_2 can be nonisomorphic (see [12]). An example of nonisomorphic DDDs with isomorphic neighborhood designs can be found in [14]. It seems interesting to ask how many pairwise nonisomorphic DDDs may have the same (up to isomorphism) neighborhood design.

In fact, the latter question is a main motivation of this paper. We construct a family of pairwise nonisomorphic Cayley DDDs Γ_i , $i \in I$, over a Heisenberg group with the same neighborhood design that have such additional properties like normality and arc-transitivity.

The second author was supported by the grant of The Natural Science Foundation of Hebei Province (project No. A2023205045) and the grant of The Israel Science Foundation (project No. 87792731).

By a *Cayley digraph* $\Gamma = \text{Cay}(G, X)$ over a finite group G with a non-empty connection set $X \subseteq G$, we mean the digraph with vertex set G and arc set $D = \{(g, xg) : x \in X, g \in G\}$. The automorphism group $\text{Aut}(\Gamma)$ contains a regular subgroup G_{right} consisting of all right translations of G and Γ is said to be *normal* if G_{right} is normal in $\text{Aut}(\Gamma)$.

The Cayley digraph Γ is a DDD with parameters $(v, k, \lambda_1, \lambda_2, m, n)$ if and only if the connection set X of Γ is a symmetric *divisible difference set* (DDS for short, see [1] for the definition) with parameters $(m, n, k, \lambda_1, \lambda_2)$. In the latter case, the neighborhood design $\mathcal{D}(\Gamma)$ is called a *development* of X and denoted by $\text{dev}(X)$. The point and block sets of $\text{dev}(X)$ are G and $\{Xg : g \in G\}$, respectively. More details on DDSs and Cayley DDDs can be found in [1, 2] and in [8, 9, 18], respectively. Our construction of Cayley DDDs is based on a recent construction of DDSs [20].

To check that Γ_i 's are pairwise nonisomorphic, we compute isomorphisms of a special S -ring (Section 3) whose automorphism group coincides with an automorphism group of each Γ_i . This also allows us to establish normality and arc-transitivity of Γ_i 's. Besides, a knowledge of the above isomorphisms implies some more interesting properties of Γ_i 's which are discussed further. Clearly, Γ_i 's have the same parameters because they have the same neighborhood design. However, they possess a stronger similarity from a combinatorial point of view. Namely, the tensors constructed from Γ_i 's by the Weisfeiler-Leman algorithm [27] are the same, or, in other words, all Γ_i 's are pairwise *WL-equivalent*. Nevertheless, each Γ_i can be distinguished from any other graph by the tensor constructed from Γ_i by the 3-dimensional Weisfeiler-Leman algorithm, i.e. the *Weisfeiler-Leman dimension* (WL-dimension for short) $\dim_{\text{WL}}(\Gamma_i)$ of Γ_i is at most 3. In particular, the isomorphism between Γ_i and any other graph can be verified in polynomial time. For a background and details on the Weisfeiler-Leman algorithm and dimension, we refer the readers to [5, 11, 13, 26].

Given a power q of a prime p , put $\log(q) = \log_p(q)$. A Heisenberg group (of dimension 3) over a finite field \mathbb{F}_q of order q is denoted by $H_3(q)$. The Euler function of a positive integer n is denoted by $\phi(n)$. To sum up the whole previous discussion, we formulate the theorem below which is a main result of the paper.

Theorem 1.1. *Let q be an odd prime power. There exist $I \subseteq \mathbb{F}_q$ and a family Γ_i , $i \in I$, of pairwise nonisomorphic normal arc-transitive divisible design Cayley digraphs with parameters $(q^3, q^2, 0, q, q^2, q)$ over $H_3(q)$ such that:*

- (1) $|I| \geq \left\lfloor \frac{\phi(q+1)}{2 \log(q)} \right\rfloor$,
- (2) the neighborhood designs $\mathcal{D}(\Gamma_i)$ and $\mathcal{D}(\Gamma_j)$ are isomorphic,
- (3) Γ_i and Γ_j are WL-equivalent,
- (4) $\dim_{\text{WL}}(\Gamma_i) \leq 3$

for all $i, j \in I$.

Remark 1. By the estimates for the Euler function (see, e.g., [22, Theorem 15]), we have

$$\frac{\phi(q+1)}{2 \log(q)} \geq \frac{q+1}{2c \log(q) \log(\log(q+1))} \xrightarrow{q \rightarrow \infty} \infty$$

for some constant $c > 0$.

Remark 2. The WL-dimension of the digraphs from Theorem 1.1 is equal to 3 if $q > 9$ and equal to 2 if $q = 3$ (see Proposition 4.4). It would be interesting to determine exact value of the WL-dimension for these graphs in cases $q = 5$ and $q = 9$.

We finish the introduction with a brief outline of the paper. Section 2 contains a necessary background of coherent configurations, S -rings (Schur rings), and WL-dimension. In Section 3, we study an S -ring over a Heisenberg group and its isomorphisms that will be used for the proof of Theorem 1.1. Finally, we prove Theorem 1.1 in Section 3. We use the standard group-theoretical notations in the text. Namely, if G is a group and $H \leq G$, then the centralizer and normalizer of H in G are denoted by $C_G(H)$ and $N_G(H)$, respectively, whereas the center of G is denoted by $Z(G)$. The identity element of G and the set of all nontrivial elements of G are denoted by e and $G^\#$, respectively. The symmetric group of the set Ω is denoted by $\text{Sym}(\Omega)$. If $\Delta \subseteq \Omega$ is invariant under $f \in \text{Sym}(\Omega)$, then the bijection induced by f on Δ is denoted by f^Δ . The holomorph of a group G is denoted by $\text{Hol}(G)$.

2. PRELIMINARIES

Throughout the text, we freely use basic definitions and facts from the theories of coherent configurations and S -rings (Schur rings) and recall in this section some of them which will be crucial for further explanation. For a background of coherent configurations and S -rings, we refer the readers to [5] and [23], respectively, where the terminology used in the paper and all facts not explained in detail are contained.

2.1. Coherent configurations. Let $\mathcal{X} = (\Omega, S)$ be a coherent configuration on a finite set Ω and S the set of all basis relations of \mathcal{X} . If the diagonal of $\Omega \times \Omega$ is a basis relation of \mathcal{X} , then \mathcal{X} is an (*association*) *scheme*. A binary relation on Ω is defined to be a *relation* of \mathcal{X} if it is a union of some basis relations. Given a coherent configuration \mathcal{X}' on Ω , put $\mathcal{X}' \geq \mathcal{X}$ if every basis relation of \mathcal{X} is a relation \mathcal{X}' .

If $s \in S$ and $\alpha \in \Omega$, then n_s denotes the valency of S and $\alpha s = \{\beta \in \Omega : (\alpha, \beta) \in s\}$. A scheme \mathcal{X} such that $n_s = 1$ for all $s \in S$ is said to be *regular*. The set of all fibers of \mathcal{X} is denoted by $F(\mathcal{X})$. If $\Delta \in F(\mathcal{X})$, then the pair

$$\mathcal{X}_\Delta = (\Delta, S_\Delta),$$

where $S_\Delta = \{s \in S : s \subseteq \Delta \times \Delta\}$, is a coherent configuration on Δ . Given $r, s, t \in S$, the corresponding intersection number of \mathcal{X} is denoted by c_{rs}^t .

Let $\mathcal{X} = (\Omega, S)$ and $\mathcal{X}' = (\Omega', S')$ be coherent configurations. An *algebraic isomorphism* from \mathcal{X} to \mathcal{X}' is defined to be a bijection $\varphi : S \rightarrow S'$ such that

$$c_{rs}^t = c_{\varphi(r), \varphi(s)}^{\varphi(t)}$$

for all $r, s, t \in S$. Every algebraic isomorphism can be extended in a natural way to a bijection between the relations of \mathcal{X} and \mathcal{X}' . The set of all algebraic isomorphisms from \mathcal{X} to itself (algebraic automorphisms) forms the subgroup of $\text{Sym}(S)$ denoted by $\text{Aut}_{\text{Alg}}(\mathcal{X})$.

A (*combinatorial*) *isomorphism* from \mathcal{X} to \mathcal{X}' is defined to be a bijection $f : \Omega \rightarrow \Omega'$ such that $S' = f(S)$, where $f(S) = \{f(s) : s \in S\}$ and $f(s) = \{(f(\alpha), f(\beta)) : (\alpha, \beta) \in s\}$. The group $\text{Iso}(\mathcal{X})$ of all isomorphisms from \mathcal{X} onto itself has a normal subgroup

$$\text{Aut}(\mathcal{X}) = \{f \in \text{Iso}(\mathcal{X}) : f(s) = s \text{ for every } s \in S\}$$

called the *automorphism group* of \mathcal{X} . A coherent configuration is said to be *schurian* if S is equal to the set of all orbits of $\text{Aut}(\mathcal{X})$ acting on Ω^2 componentwise. A schurian coherent configuration is said to be *2-minimal* if $\text{Aut}(\mathcal{X})$ does not have a proper subgroup with the same set S of orbits in its componentwise action on Ω^2 .

Every isomorphism of coherent configurations induces in a natural way the algebraic isomorphism of them. However, not every algebraic isomorphism is induced by a combinatorial one. A coherent configuration is said to be *separable* if every algebraic isomorphism from it to any coherent configuration is induced by an isomorphism. The set of all algebraic automorphisms of \mathcal{X} induced by isomorphisms forms the subgroup of $\text{Aut}_{\text{Alg}}(\mathcal{X})$ denoted by $\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{X})$.

Lemma 2.1. [5, Theorem 2.3.33] *Every regular scheme is separable.*

Lemma 2.2. [10, Lemma 9.2] *Let $\mathcal{X} = (\Omega, S)$ be a coherent configuration and $\Delta \in F(\mathcal{X})$ such that for every $\Delta \neq \Delta' \in F(\mathcal{X})$ there is a basis relation $r \subseteq \Delta \times \Delta'$ of valency 1. Then*

- (1) $\text{Aut}(\mathcal{X}_\Delta) \cong \text{Aut}(\mathcal{X})$;
- (2) \mathcal{X} is separable (schurian) whenever \mathcal{X}_Δ so is.

Given $\alpha \in \Omega$, the one-point extension of \mathcal{X} with respect to α , i.e. the smallest coherent configuration \mathcal{X}' on Ω such that $\mathcal{X}' \geq \mathcal{X}$ and $\{\alpha\} \in F(\mathcal{X}')$, is denoted by \mathcal{X}_α . By [5, Proposition 3.3.3(1)],

$$(1) \quad \text{Aut}(\mathcal{X}_\alpha) = \text{Aut}(\mathcal{X})_\alpha.$$

The first statement of the lemma below is [5, Lemma 3.3.5(2)] whereas the second one is [5, Theorem 3.3.7(1)].

Lemma 2.3. *Let $\mathcal{X} = (\Omega, S)$ be a coherent configuration and $\alpha \in \Omega$. Then the following statements hold.*

- (1) *For all $r, s, t \in S$, the binary relation $r \cap (\alpha s \times \alpha t)$ is a relation of \mathcal{X}_α .*
- (2) *If \mathcal{X} is schurian, then $F(\mathcal{X}_\alpha) = \{\alpha s : s \in S\}$.*

2.2. S -rings. Let \mathcal{A} be an S -ring over a group G . The set of all basic sets of \mathcal{A} is denoted by $\mathcal{S} = \mathcal{S}(\mathcal{A})$. Given $X, Y, Z \in \mathcal{S}(\mathcal{A})$, the corresponding structure constant of \mathcal{A} is denoted by c_{XY}^Z . Due to [5, Eq. 2.1.14],

$$(2) \quad |Z|c_{XY}^{Z^{(-1)}} = |X|c_{YZ}^{X^{(-1)}} = |Y|c_{ZX}^{Y^{(-1)}}$$

for all $X, Y, Z \in \mathcal{S}(\mathcal{A})$. If $X \subseteq G$ and $\underline{X} \in \mathcal{A}$, then X is defined to be an \mathcal{A} -set. A subgroup of G which is an \mathcal{A} -set is called an \mathcal{A} -subgroup.

Lemma 2.4. [28, Proposition 22.1] *Let \mathcal{A} be an S -ring over G , $\xi = \sum_{g \in G} c_g g \in \mathcal{A}$, where $c_g \in \mathbb{Z}$, and $c \in \mathbb{Z}$. Then $\{g \in G : c_g = c\}$ is an \mathcal{A} -set.*

If \mathcal{A} is an S -ring over G , then $\mathcal{X} = \mathcal{X}(\mathcal{A}) = (G, S)$, where $S = \mathcal{S}(\mathcal{A}) = \{r(X) : X \in \mathcal{S}(\mathcal{A})\}$ and $r(X) = \{(g, xg) : x \in X, g \in G\}$, is a Cayley scheme over G , i.e. a scheme on the set G such that $\text{Aut}(\mathcal{X}) \geq G_{\text{right}}$ (see [5, Section 2.4]). Due to [5, Theorem 2.4.16], the mapping

$$\mathcal{A} \mapsto \mathcal{X}(\mathcal{A})$$

is a partial order isomorphism between the S -rings and Cayley schemes over G . One can see that $\mathcal{A}_1 \leq \mathcal{A}_2$ if and only if $\mathcal{X}(\mathcal{A}_1) \leq \mathcal{X}(\mathcal{A}_2)$.

Put

$$\text{Iso}(\mathcal{A}) = \text{Iso}(\mathcal{X}(\mathcal{A})) \text{ and } \text{Aut}(\mathcal{A}) = \text{Aut}(\mathcal{X}(\mathcal{A})).$$

Clearly, $\text{Aut}(\mathcal{A}) \geq G_{\text{right}}$. By the definitions, if $f \in \text{Aut}(\mathcal{A})$, $X \in \mathcal{S}(\mathcal{A})$, and $g \in G$, then

$$(3) \quad f(Xg) = Xf(g).$$

The S -ring \mathcal{A} is said to be *normal* if G_{right} is normal in $\text{Aut}(\mathcal{A})$ or, equivalently, $\text{Aut}(\mathcal{A})$ is a subgroup of $\text{Hol}(G) = G_{\text{right}} \rtimes \text{Aut}(G)$. If H is an \mathcal{A} -subgroup of G , then the set of all right and the set of all left H -cosets form imprimitivity systems of $\text{Aut}(\mathcal{A})$.

The S -ring \mathcal{A} is said to be *separable* (*schurian*, *2-minimal*, respectively) if the Cayley scheme $\mathcal{X}(\mathcal{A})$ so is. If \mathcal{A} is schurian, then $\mathcal{S}(\mathcal{A})$ is equal to the set of all orbits of the stabilizer $\text{Aut}(\mathcal{A})_e$ on G and

$$(4) \quad \text{Iso}(\mathcal{A}) = N_{\text{Sym}(G)}(\text{Aut}(\mathcal{A})).$$

For every $K \leq \text{Aut}(G)$, the partition of G into the orbits of K defines the S -ring \mathcal{A} over G . In this case, \mathcal{A} is called *cyclotomic* and denoted by $\text{Cyc}(K, G)$. Clearly, $K \leq \text{Aut}(\mathcal{A})_e$. If \mathcal{A} is cyclotomic, then \mathcal{A} is schurian.

Let \mathcal{A}_1 and \mathcal{A}_2 be S -rings, $\mathcal{X}_1 = \mathcal{X}(\mathcal{A}_1)$, and $\mathcal{X}_2 = \mathcal{X}(\mathcal{A}_2)$. Every algebraic isomorphism from \mathcal{X}_1 to \mathcal{X}_2 induces a bijection φ from $\mathcal{S}(\mathcal{A}_1)$ to $\mathcal{S}(\mathcal{A}_2)$ such that

$$c_{\varphi(X)\varphi(Y)}^{\varphi(Z)} = c_{XY}^Z$$

for all $X, Y, Z \in \mathcal{S}(\mathcal{A})$. For the convenience, further we will refer to the above bijection as an *algebraic isomorphism* from \mathcal{A}_1 to \mathcal{A}_2 . Every such algebraic isomorphism can be extended in a natural way to a bijection between \mathcal{A}_1 - and \mathcal{A}_2 -sets. The group of all algebraic isomorphisms from \mathcal{A} to itself (algebraic automorphisms) and its subgroup consisting of all algebraic automorphisms inducing by the elements of $\text{Iso}(\mathcal{A})$ are denoted by $\text{Aut}_{\text{Alg}}(\mathcal{A})$ and $\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})$, respectively. The groups $\text{Aut}_{\text{Alg}}(\mathcal{A})$ and $\text{Aut}_{\text{Alg}}(\mathcal{X}(\mathcal{A}))$ ($\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})$ and $\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{X}(\mathcal{A}))$, respectively) are equivalent as permutation groups.

It is easy to see that $f \in \text{Iso}(\mathcal{A})$ induces the trivial algebraic automorphism of \mathcal{A} which fixes every basic set if and only if $f \in \text{Aut}(\mathcal{A})$ and hence $f_1, f_2 \in \text{Iso}(\mathcal{A})$ induce the same algebraic isomorphism if and only if $f_2 f_1^{-1} \in \text{Aut}(\mathcal{A})$. Therefore

$$(5) \quad |\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})| = \frac{|\text{Iso}(\mathcal{A})|}{|\text{Aut}(\mathcal{A})|}.$$

2.3. WL-closure and WL-dimension. The material of this subsection is taken from [3] (see also [5, Section 2.6]). The *WL-closure* $\text{WL}(\Gamma)$ of a digraph $\Gamma = (\Omega, D)$ is defined to be the smallest coherent configuration $\mathcal{X} = (\Omega, S)$ such that D is a relation of \mathcal{X} . The tensor of the intersection numbers of $\text{WL}(\Gamma)$ is an output of the Weisfeiler-Leman algorithm whose input graph is Γ . One can check that $\text{Aut}(\text{WL}(\Gamma)) = \text{Aut}(\Gamma)$. So if $\Gamma = \text{Cay}(G, X)$ is a Cayley digraph, then $\text{Aut}(\text{WL}(\Gamma)) \geq G_{\text{right}}$ and hence $\text{WL}(\Gamma)$ is a Cayley scheme over G . For the convenience, further we will refer to the S -ring \mathcal{A} such that $\mathcal{X}(\mathcal{A}) = \text{WL}(\Gamma)$ as the *WL-closure* of the Cayley digraph Γ . Due to the definitions, \mathcal{A} is the smallest (with respect to inclusion) S -ring \mathcal{A} over G such that X is an \mathcal{A} -set.

Lemma 2.5. [5, Theorem 2.6.4] *Let Γ_1 and Γ_2 be digraphs and f an isomorphism from Γ_1 to Γ_2 . Then f is an isomorphism from $\text{WL}(\Gamma_1)$ to $\text{WL}(\Gamma_2)$.*

The next lemma can be deduced straightforwardly by the induction on the number of iterations of the Weisfeiler-Leman algorithm.

Lemma 2.6. *Let $\Gamma_1 = (\Omega_1, D_1)$ and $\Gamma_2 = (\Omega_2, D_2)$ be digraphs and φ an algebraic isomorphism from $\text{WL}(\Gamma_1)$ to $\text{WL}(\Gamma_2)$ such that $\varphi(D_1) = D_2$. Then Γ_1 and Γ_2 are WL-equivalent.*

The lemma below is a corollary of [11, Theorem 2.5] and can be found, e.g., in [3, Lemma 3.1].

Lemma 2.7. *Let Γ be a digraph with vertex set Ω and $\mathcal{X} = \text{WL}(\Gamma)$. Then the following statements hold.*

- (1) $\dim_{\text{WL}}(\Gamma) \leq 2$ if and only if \mathcal{X} is separable.
- (2) If \mathcal{X}_α is separable for some $\alpha \in \Omega$, then $\dim_{\text{WL}}(\Gamma) \leq 3$.

3. AN S -RING OVER A HEISENBERG GROUP

In this section, we recall a construction of a cyclotomic S -ring from [20], where all the details can be found. Let q be an odd prime power and \mathbb{F}_q a finite field of order q . Further till the end of the paper,

$$G = H_3(q) = \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} : x, y, z \in \mathbb{F}_q \right\}$$

is a Heisenberg group of dimension 3 over \mathbb{F}_q . Put

$$Z = Z(G) = \left\{ \begin{pmatrix} 1 & 0 & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} : z \in \mathbb{F}_q \right\}.$$

Clearly, $|G| = q^3$ and $|Z| = q$.

Let $\varepsilon \in \mathbb{F}_q$ be a nonsquare and

$$\mathcal{M} = \mathcal{M}(\varepsilon) = \left\{ \begin{pmatrix} \alpha & \beta \\ \varepsilon\beta & \alpha \end{pmatrix} : \alpha, \beta \in \mathbb{F}_q, (\alpha, \beta) \neq (0, 0) \right\}.$$

Due to [19, Chapter 2.5], the group \mathcal{M} (with the standard matrix multiplication) is a subgroup of $\text{GL}_2(q)$ isomorphic to the multiplicative group $\mathbb{F}_{q^2}^*$. In particular, $|\mathcal{M}| = q^2 - 1$ and \mathcal{M} is cyclic. Given $M = \begin{pmatrix} \alpha & \beta \\ \varepsilon\beta & \alpha \end{pmatrix} \in \mathcal{M}$, let us define $\rho = \rho(M) : G \rightarrow G$ as follows:

$$\rho \left(\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \right) = \begin{pmatrix} 1 & \alpha x + \varepsilon\beta y & F_{\alpha, \beta}(x, y, z) \\ 0 & 1 & \beta x + \alpha y \\ 0 & 0 & 1 \end{pmatrix},$$

where $F_{\alpha, \beta}(x, y, z) = \alpha\beta \left(\frac{x^2}{2} + \varepsilon \frac{y^2}{2} \right) + \varepsilon\beta^2 xy + (\alpha^2 - \varepsilon\beta^2)z$. The mapping $M \mapsto \rho(M)$ is a monomorphism from \mathcal{M} to $\text{Aut}(G)$ (see [20]). Let $K = \{\rho(M) : M \in \mathcal{M}\}$. The group K is a cyclic group of order $q^2 - 1$. Put $\mathcal{A} = \text{Cyc}(K, G)$. The basic sets of \mathcal{A} are the following:

$$\{e\}, Z^\#, Y_i, i \in \mathbb{F}_q,$$

where

$$Y_i = \left\{ \begin{pmatrix} 1 & \alpha & \gamma_i(\alpha, \beta) \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix}, \alpha, \beta \in \mathbb{F}_q, (\alpha, \beta) \neq (0, 0) \right\}$$

for $\gamma_i(\alpha, \beta) = \frac{\alpha\beta}{2} + (\alpha^2 - \varepsilon\beta^2)i$. One can see that $|Y_i| = q^2 - 1$ and $Y_i^{(-1)} = Y_{-i}$ for every $i \in \mathbb{F}_q$.

Given $i \in \mathbb{F}_q$, put $X_i = Y_i \cup \{e\}$. The next lemma immediately follows from [20, Corollary 6.4].

Lemma 3.1. *For every $i \in \mathbb{F}_q$,*

$$\underline{X}_i \cdot \underline{X}_i^{(-1)} = q^2 e + q(\underline{G} - \underline{Z}),$$

i.e. X_i is a DDS with parameters $(q^2, q, q^2, 0, q)$ which is a transversal for Z in G .

Given $i, j \in \mathbb{F}_q^\infty = \mathbb{F}_q \cup \{\infty\}$, put

$$\chi(i) = \begin{cases} -i, & i \neq \infty, \\ \infty, & i = \infty, \end{cases}$$

and

$$\psi(i, j) = \begin{cases} \frac{ij+\delta}{i+j}, & i+j \neq 0, i, j \neq \infty, \\ \infty, & j = -i, \\ i, & i \neq \infty, j = \infty, \\ j, & i = \infty, j \neq \infty, \end{cases}$$

where $\delta = \frac{\varepsilon}{16}$.

Lemma 3.2. [20, Lemma 6.7] *The set \mathbb{F}_q^∞ equipped with the binary operation ψ forms a cyclic group of order $q+1$, where ∞ is the identity element and $\chi(i)$ is the element inverse to $i \in \mathbb{F}_q^\infty$.*

The lemma below collects an information on the structure constants of \mathcal{A} from [20, Eqs. (9),(10),(14),(16),(17)] which we will need further.

Lemma 3.3. *Given $i, j, k \in \mathbb{F}_q$,*

$$c_{Y_i Y_j}^{Z^\#} = \begin{cases} 0, & j = -i, \\ q+1, & j \neq -i, \end{cases}$$

$$c_{Y_i Y_{-i}}^{Y_k} = \begin{cases} q, & k \notin \{i, -i\}, \\ q-1, & k \in \{i, -i\} \text{ and } i \neq 0, \\ q-2, & k = i = -i = 0, \end{cases}$$

and if $i+j \neq 0$, then

$$c_{Y_i Y_j}^{Y_k} = \begin{cases} q+1, & k \notin \{i, j, \psi(i, j)\}, \\ 1, & k = \psi(i, j), \\ q, & i \neq j, \text{ and } k \in \{i, j\}, \\ q-1, & k = i = j. \end{cases}$$

Proposition 3.4. *In the above notations, $\text{Aut}(\mathcal{A}) = G_{\text{right}} \rtimes K$.*

Proof. The S -ring \mathcal{A} is cyclotomic and hence schurian. So each basic set of \mathcal{A} is an orbit of the stabilizer $\text{Aut}(\mathcal{A})_e$. Obviously, $G_{\text{right}} \rtimes K \leq \text{Aut}(\mathcal{A})$. One can see that

$$|\text{Aut}(\mathcal{A})| = |G| |\text{Aut}(\mathcal{A})_e| = |G| |Y_0| |\text{Aut}(\mathcal{A})_{e, y_0}| = q^3 (q^2 - 1) |\text{Aut}(\mathcal{A})_{e, y_0}| = |G| |K| |\text{Aut}(\mathcal{A})_{e, y_0}|,$$

where $y_0 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in Y_0$. So to prove that $\text{Aut}(\mathcal{A}) = G_{\text{right}} \rtimes K$, it suffices to check that the stabilizer $\text{Aut}(\mathcal{A})_{e, y_0}$ is trivial. Let us prove that every $f \in \text{Aut}(\mathcal{A})_{e, y_0}$ is trivial.

Lemma 3.5. *In the above notations, $f^Z = \text{id}_Z$.*

Proof. Let $z \in Z$ and Y a basic set containing zy_0^{-1} . Note that $\{z\} = Yy_0 \cap Z^\#$ because $Y \cup \{e\}$ is a transversal for Z (Lemma 3.1). Therefore

$$\{f(z)\} = f(Yy_0 \cap Z^\#) = Yy_0 \cap Z^\# = \{z\},$$

where the second equality holds by $f(y_0) = y_0$ and Eq. (3). Thus, $f(z) = z$ for every $z \in Z$ and hence $f^Z = \text{id}_Z$. \square

Lemma 3.6. *In the above notations, if $f(y) = y$ for some $y \in G \setminus Z$, then $f(Zy) = Zy$ and $f^{Zy} = \text{id}_{Zy}$. In particular, $f^{Zy_0} = \text{id}_{Zy_0}$.*

Proof. Since Z is an \mathcal{A} -subgroup, the set Zy is a block of $\text{Aut}(\mathcal{A})$. So due to $f(y) = y$, we have $f(Zy) = Zy$. Each X_i is a transversal for Z by Lemma 3.1. Therefore all elements from Zy belong to pairwise distinct basic sets. Together with $f(Zy) = Zy$, this implies that $f^{Zy} = \text{id}_{Zy}$. \square

In view of Lemma 3.1, Lemma 3.5, and Lemma 3.6, it remains to verify that f fixes every $y \in Y_1 \setminus Zy_0$. Since X_0 and X_{-1} are transversals for Z and $y \notin Zy_0$, there exist $i, j, k, l \in \{0, \dots, q-1\}$ such that

$$y \in T = Y_0 z^i \cap Y_0 y_0 z^j \cap Y_{-1} z^k \cap Y_{-1} y_0 z^l \cap Y_1.$$

If $|T| = 1$ or, equivalently, $T = \{y\}$, then we are done. Indeed,

$$\{f(y)\} = f(T) = T = \{y\},$$

where the second equality holds by Eq. (3), Lemma 3.5, and Lemma 3.6.

Further, we establish that $|T| = 1$. Let $t \in T$. Then $t = \begin{pmatrix} 1 & \alpha & \gamma_1(\alpha, \beta) \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix} \in Y_1$ for some $\alpha, \beta \in \mathbb{F}_q$. Let

$$u_1 = \begin{pmatrix} 1 & \alpha_1 & \gamma_0(\alpha_1, \beta_1) \\ 0 & 1 & \beta_1 \\ 0 & 0 & 1 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 1 & \alpha_2 & \gamma_0(\alpha_2, \beta_2) \\ 0 & 1 & \beta_2 \\ 0 & 0 & 1 \end{pmatrix} \in Y_0$$

and

$$u_3 = \begin{pmatrix} 1 & \alpha_3 & \gamma_{-1}(\alpha_3, \beta_3) \\ 0 & 1 & \beta_3 \\ 0 & 0 & 1 \end{pmatrix}, \quad u_4 = \begin{pmatrix} 1 & \alpha_4 & \gamma_{-1}(\alpha_4, \beta_4) \\ 0 & 1 & \beta_4 \\ 0 & 0 & 1 \end{pmatrix} \in Y_{-1}$$

such that

$$t = u_1 z^i = u_2 y_0 z^j = u_3 z^k = u_4 y_0 z^l.$$

The latter equalities are equivalent to the system

$$(6) \quad \begin{cases} \alpha = \alpha_1 = \alpha_2 + 1 = \alpha_3 = \alpha_4 + 1, \\ \beta = \beta_1 = \beta_2 = \beta_3 = \beta_4, \\ \gamma_1(\alpha, \beta) = \gamma_0(\alpha_1, \beta_1) + i = \gamma_0(\alpha_2, \beta_2) + j = \gamma_{-1}(\alpha_3, \beta_3) + k = \gamma_{-1}(\alpha_4, \beta_4) + l. \end{cases}$$

One can see that α_i, β_i can be expressed via α, β using the equations from the first and second lines of System (6) and these expressions can be substituted to the equations from the third line of System (6). As a result, we obtain

$$\alpha^2 - \varepsilon\beta^2 = i = -\frac{\beta}{2} + j = -(\alpha^2 - \varepsilon\beta^2) + k = -\frac{\beta}{2} - ((\alpha - 1)^2 - \varepsilon\beta^2) = l.$$

In particular, $\beta = 2(j - i)$ from the second equality and $\alpha = \frac{k+1+\beta}{2}$ from the fourth one. Therefore System (6) has at most one solution and hence $|T| \leq 1$. On the other hand, $y \in T$. Thus, $T = \{y\}$ as required. \square

Corollary 3.7. *The S -ring \mathcal{A} is normal and 2-minimal.*

Proof. The normality immediately follows from Proposition 3.4. Observe that the basis relations $r(Y_i)$, $i \in \mathbb{F}_q$, of the Cayley scheme $\mathcal{X}(\mathcal{A})$ have size

$$|G|n_{r(Y_i)} = |G||Y_i| = q^3(q^2 - 1) = |\text{Aut}(\mathcal{A})|.$$

Therefore there is no a proper subgroup of $\text{Aut}(\mathcal{A})$ with the same orbits on $G \times G$ as $\text{Aut}(\mathcal{A})$ and hence \mathcal{A} is 2-minimal. \square

Proposition 3.8. *In the above notations, $|\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})| \leq 2 \log(q)$.*

Proof. Note that $|\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})| = |\text{Iso}(\mathcal{A})|/|\text{Aut}(\mathcal{A})|$ by Eq. (5). From Eq. (4) it follows that $\text{Iso}(\mathcal{A}) = N_{\text{Sym}(G)}(\text{Aut}(\mathcal{A}))$. By Proposition 3.4, we have $\text{Aut}(\mathcal{A}) = G_{\text{right}} \rtimes K$, where $|K| = q^2 - 1$. So the group G_{right} is a normal Sylow subgroup of $\text{Aut}(\mathcal{A})$. In particular, G_{right} is characteristic in $\text{Aut}(\mathcal{A})$. Since $\text{Aut}(\mathcal{A})$ is normal in $\text{Iso}(\mathcal{A})$, we conclude that G_{right} is normal in $\text{Iso}(\mathcal{A})$ and hence $\text{Iso}(\mathcal{A}) \leq N_{\text{Sym}(G)}(G_{\text{right}}) = \text{Hol}(G)$. Thus,

$$\text{Iso}(\mathcal{A}) \leq N_{\text{Sym}(G)}(\text{Aut}(\mathcal{A})) \cap \text{Hol}(G) = N_{\text{Hol}(G)}(G_{\text{right}} \rtimes K).$$

The group $N_{\text{Hol}(G)}(G_{\text{right}} \rtimes K)$ coincides with $G_{\text{right}} \rtimes N_{\text{Aut}(G)}(K)$. Indeed, $g_{\text{right}}\tau \in \text{Hol}(G)$, where $g \in G$ and $\tau \in \text{Aut}(G)$, normalizes the group $G_{\text{right}} \rtimes K$ if and only if $G_{\text{right}} \rtimes K^\tau = G_{\text{right}} \rtimes K$. Clearly, if $\tau \in N_{\text{Aut}(G)}(K)$, then the latter equality holds. Suppose that $G_{\text{right}} \rtimes K^\tau = G_{\text{right}} \rtimes K$. Since $\text{GCD}(|G|, |K|) = 1$ and G is solvable, the Schur-Zassenhaus theorem implies that K and K^τ are conjugate in $G_{\text{right}} \rtimes K$. So $K^{h_{\text{right}}} = K^\tau$ for some $h \in G$. Then for every $\sigma \in K$, we have $h_{\text{right}}\sigma h_{\text{right}}^{-1}(e) = \sigma(h^{-1})h = e$ because $h_{\text{right}}\sigma h_{\text{right}}^{-1} \in K^\tau \leq \text{Aut}(G)$. This yields that h is fixed by every element of K and hence the basic set of \mathcal{A} containing h is of size 1. The unique basic set of \mathcal{A} of size 1 is $\{e\}$. So $h = e$ and, consequently, $K^\tau = K$. Therefore $G_{\text{right}} \rtimes K^\tau = G_{\text{right}} \rtimes K$ if and only if $\tau \in N_{\text{Aut}(G)}(K)$ and hence $N_{\text{Hol}(G)}(G_{\text{right}} \rtimes K) = G_{\text{right}} \rtimes N_{\text{Aut}(G)}(K)$. Thus,

$$\text{Iso}(\mathcal{A}) \leq G_{\text{right}} \rtimes N_{\text{Aut}(G)}(K).$$

Due to the latter equality and $\text{Aut}(\mathcal{A}) = G_{\text{right}} \rtimes K$, the statement of the proposition is a consequence of inequality

$$(7) \quad |N_{\text{Aut}(G)}(K)| \leq 2 \log(q)|K|$$

which we are going to prove. Let π be the homomorphism from $\text{Aut}(G)$ to $\text{Aut}(G/Z)$ which maps every $\sigma \in \text{Aut}(G)$ to the induced automorphism of G/Z acting by the rule $\sigma(Zg) = Z\sigma(g)$. The group G/Z is isomorphic to the additive group of $(2l)$ -dimensional vector space over \mathbb{F}_p , where $q = p^l$ for an odd prime p and $l = \log(q)$, and hence

$$\text{Aut}(G/Z) \cong \text{GL}_{2l}(p).$$

The group $\pi(K)$ is a cyclic subgroup of $\text{Aut}(G/Z)$ of order $q^2 - 1$. Moreover, $\pi(K)$ acts transitively on the set of nonidentity elements of G/Z because each orbit of K outside Z together with the identity element is a transversal for Z in G (Lemma 3.1). Therefore $\pi(K)$ is a Singer subgroup of $\text{Aut}(G/Z) \cong \text{GL}_{2l}(p)$. From [16, p. 187] it follows that

$$(8) \quad |N_{\text{Aut}(G/Z)}(\pi(K))| = 2l|\pi(K)|.$$

Let $N_0 = N_{\text{Aut}(G)}(K) \cap \ker(\pi)$ and $\theta \in N_0$. Then $\theta\sigma\theta^{-1} \in K$ for every $\sigma \in K$. Let $g \in G \setminus Z$. The elements $\theta\sigma\theta^{-1}(g)$ and $\sigma(g)$ lie in the same orbit of K because $\theta \in N_{\text{Aut}(G)}(K)$ and in the same Z -coset, namely in $\sigma(Zg)$, because $\theta \in \ker(\pi)$. So $\theta\sigma\theta^{-1}(g) = \sigma(g)$ by Lemma 3.1. Since the above equality holds for all $g \in G \setminus Z$, we conclude that $\theta\sigma\theta^{-1} = \sigma$ for every $\sigma \in K$. Therefore $\theta \in C_{\text{Aut}(G)}(K)$ and hence $N_0 \leq C_{\text{Aut}(G)}(K)$. The group K is regular on each of its orbits outside Z . This yields that $C_{\text{Aut}(G)}(K)$ is semiregular on each orbit of K outside Z . So $|C_{\text{Aut}(G)}(K)| \leq |K|$. One can see that K is abelian and consequently $K \leq C_{\text{Aut}(G)}(K)$. Therefore $C_{\text{Aut}(G)}(K) = K$. The latter implies that $N_0 \leq K_0$, where $K_0 = K \cap \ker(\pi)$. However, each orbit of K outside Z together with the identity element is transversal for Z in G which implies that K_0 is trivial. Thus,

$$(9) \quad |N_0| = |K_0| = 1.$$

Now one can estimate $|N_{\text{Aut}(G)}(K)|$ as follows:

$$|N_{\text{Aut}(G)}(K)| = |N_{\text{Aut}(G)}(K)/N_0| = |\pi(N_{\text{Aut}(G)}(K))| \leq |N_{\text{Aut}(G/Z)}(\pi(K))| = 2l|\pi(K)| = 2l|K|,$$

where the first and fifth equalities hold by Eq. (9), whereas the fourth equality holds by Eq. (8). Thus, Eq. (7) holds and we are done. \square

Given $\tau \in \text{Aut}(\mathbb{F}_q^\infty)$, let us define a bijection $\hat{\tau}$ on the set $\mathcal{S}(\mathcal{A})$ as follows:

$$\hat{\tau}(\{e\}) = \{e\}, \quad \hat{\tau}(Y_k) = Y_{\tau(k)}, \quad k \in \mathbb{F}_q^\infty,$$

where $Y_\infty = Z^\#$.

Proposition 3.9. *The following statements hold.*

- (1) *The mapping $\tau \mapsto \hat{\tau}$ is a monomorphism from $\text{Aut}(\mathbb{F}_q^\infty)$ to $\text{Aut}_{\text{Alg}}(\mathcal{A})$.*
- (2) $|\text{Aut}_{\text{Alg}}(\mathcal{A})| \geq \phi(q+1)$.
- (3) *Given generators i, j of $\mathbb{F}_q^\infty \cong C_{q+1}$, there is an algebraic automorphism of \mathcal{A} which maps Y_i to Y_j .*

Proof. Statement (1) immediately follows from the the definition of $\hat{\tau}$ and inclusion $\hat{\tau} \in \text{Aut}_{\text{Alg}}(\mathcal{A})$ which can be verified straightforwardly using Eq. (2) and Lemma 3.3. For example,

$$c_{\hat{\tau}(Y_i)\hat{\tau}(Y_j)}^{\hat{\tau}(Y_{\psi(i,j)})} = c_{Y_{\tau(i)}Y_{\tau(j)}}^{Y_{\tau(\psi(i,j))}} = c_{Y_{\tau(i)}Y_{\tau(j)}}^{Y_{\psi(\tau(i),\tau(j))}},$$

where the latter equality holds because $\tau \in \text{Aut}(\mathbb{F}_q^\infty)$. Statement (2) holds by Statement (1) and Lemma 3.2. Since i and j are generators of \mathbb{F}_q^∞ , there is $\tau \in \text{Aut}(\mathbb{F}_q^\infty)$ such that $\tau(i) = j$. So $\hat{\tau}(Y_i) = Y_{\tau(i)} = Y_j$ and Statement (3) holds by Statement (1). \square

Corollary 3.10. *The S -ring \mathcal{A} is separable only if $q \in \{3, 5, 9\}$.*

Proof. Due to Proposition 3.8 and Proposition 3.9(2), the S -ring \mathcal{A} is separable only if $\phi(q+1) \leq 2 \log(q)$ which is true if and only if $q \in \{3, 5, 9\}$. \square

4. PROOF OF THEOREM 1.1

Throughout this section, we use the notations from the previous one. Given $i \in \mathbb{F}_q$, put $\Gamma_i = \text{Cay}(G, X_i)$. By Lemma 3.1, Γ_i is a divisible design digraph with parameters $(q^3, q^2, 0, q, q^2, q)$ whose classes are Z -cosets and in which each vertex dominates exactly one vertex from each class. Denote the set of all generators of $\mathbb{F}_q^\infty \cong C_{q+1}$ by I . Clearly, $|I| = \phi(q+1)$.

Lemma 4.1. *In the above notations, $\text{WL}(\Gamma_i) = \mathcal{A}$ for every $i \in I$.*

Proof. Put $\mathcal{B} = \text{WL}(\Gamma_i)$. The S -ring \mathcal{B} is the smallest S -ring over G for which Y_i is a \mathcal{B} -set. So $\mathcal{B} \leq \mathcal{A}$. Let us check the reverse inclusion. Put

$$a_1 = i \text{ and } a_k = \psi(a_{k-1}, i) \text{ for } k \geq 2.$$

Since i is a generator of \mathbb{F}_q^∞ , we conclude that $a_{q+1} = \infty$ and

$$(10) \quad \{a_1, \dots, a_{q+1}\} = \mathbb{F}_q^\infty.$$

Further we are going to prove that Y_{a_k} is a \mathcal{B} -set for every $k \in \{1, \dots, \frac{q+1}{2}\}$. We proceed by induction on k . The base of induction for $k = 1$ follows from the definition of \mathcal{B} . Observe

that $a_k \neq -i$ for $k \in \{1, \dots, \frac{q+1}{2}\}$ because $-i = \chi(i)$ and hence $-i = a_q$. Together with Lemma 3.3 this implies that

$$\underline{Y}_{a_k} \cdot \underline{Y}_i = \underline{Y}_{a_{k+1}} + q(\underline{Y}_{a_k} + \underline{Y}_i) + (q+1)(G^\# - \underline{Y}_{a_{k+1}} - \underline{Y}_{a_k} - \underline{Y}_i).$$

By the induction hypothesis, the left-hand side of the above equality belongs to \mathcal{B} . So $\underline{Y}_{a_{k+1}} \in \mathcal{B}$ by Lemma 2.4 and hence $Y_{a_{k+1}}$ is a \mathcal{B} -set. For every $k \in \{\frac{q+1}{2}, \dots, q\}$, there is $l \in \{1, \dots, \frac{q+1}{2}\}$ such that $Y_{a_k} = Y_{\chi(a_l)} = Y_{-a_l} = Y_{a_l}^{(-1)}$. Therefore Y_{a_k} is a \mathcal{B} -set for every $k \in \{\frac{q+1}{2}, \dots, q\}$. The above discussion together with Eq. (10) yields that Y_i is a \mathcal{B} -set for every $i \in \mathbb{F}_q$. Finally,

$$Z^\# = G^\# \setminus \bigcup_{i \in \mathbb{F}_q} Y_i$$

is a \mathcal{B} -set. Thus, every basic set of \mathcal{A} is a \mathcal{B} -set. This implies that $\mathcal{B} \geq \mathcal{A}$ and hence $\mathcal{B} = \mathcal{A}$ as desired. \square

Corollary 4.2. *For all $i, j \in I$, the digraphs Γ_i and Γ_j are normal, arc-transitive, and WL-equivalent.*

Proof. Due Lemma 4.1, we have $\text{WL}(\Gamma_i) = \text{WL}(\Gamma_j) = \mathcal{A}$ and hence $\text{Aut}(\Gamma_i) = \text{Aut}(\Gamma_j) = \text{Aut}(\mathcal{A})$. So $\text{Aut}(\Gamma_i) = \text{Aut}(\Gamma_j) = G_{\text{right}} \rtimes K$ by Proposition 3.4. This implies that Γ_i and Γ_j are normal. The group $K = \text{Aut}(\Gamma_i)_e = \text{Aut}(\Gamma_j)_e$ acts transitively on Y_i and Y_j which are neighborhoods of e in Γ_i and Γ_j , respectively. Since Γ_i and Γ_j are vertex-transitive, this yields that Γ_i and Γ_j are arc-transitive. Finally, Γ_i and Γ_j are WL-equivalent by Lemma 2.6, Proposition 3.9(3), and Proposition 4.1. \square

Lemma 4.3. *There are at least $\left\lfloor \frac{\phi(q+1)}{2 \log(q)} \right\rfloor$ pairwise nonisomorphic digraphs among Γ_i , $i \in I$.*

Proof. The relation “to be isomorphic” defines an equivalence E on the set $\{\Gamma_i : i \in I\}$. It suffices to verify that there are at least $\left\lfloor \frac{\phi(q+1)}{2 \log(q)} \right\rfloor$ equivalence classes of E . Suppose that Γ_i and Γ_j are isomorphic for some $i, j \in I$ and f is an isomorphism from Γ_i to Γ_j . Then $f \in \text{Iso}(\mathcal{A})$ by Lemma 2.5 and Lemma 4.1. Clearly, f induces an algebraic automorphism $\varphi \in \text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})$ such that $\varphi(Y_i) = Y_j$. Proposition 3.8 implies that $|\text{Aut}_{\text{Alg}}^{\text{ind}}(\mathcal{A})| \leq 2 \log(q)$. Thus, each class of E has size at most $2 \log(q)$ and hence the number of classes is at least $\left\lfloor \frac{\phi(q+1)}{2 \log(q)} \right\rfloor$ as required. \square

Proposition 4.4. *Let $i \in I$. Then $\dim_{\text{WL}}(\Gamma_i) \leq 3$. Moreover, $\dim_{\text{WL}}(\Gamma_i) = 3$ if $q > 9$ and $\dim_{\text{WL}}(\Gamma_i) = 2$ if $q = 3$.*

Proof. At first, let us prove that $\dim_{\text{WL}}(\Gamma_i) \leq 3$. Due to Lemma 2.7(2) and Lemma 4.1, it suffices to verify that the one-point extension \mathcal{X}_e of the Cayley scheme $\mathcal{X} = \mathcal{X}(\mathcal{A})$ is separable. Since \mathcal{A} and hence \mathcal{X} are schurian, $F(\mathcal{X}_e) = \mathcal{S}(\mathcal{A})$ by Lemma 2.3(2). Given $j \in \mathbb{F}_q^*$, put $k = k(j) = \psi(j, 0)$. Lemma 2.3(1) implies that

$$r_j = r(Y_{k(j)}) \cap (Y_0 \times Y_j)$$

is a relation of \mathcal{X}_e . Let $y \in Y_0$. Clearly, $yr_j = Y_k y \cap Y_j$. One can see that $|Y_k y \cap Y_j| = c_{Y_{-k} Y_j}^{Y_0}$. In view of Eq. (2) and Lemma 3.3, we have $c_{Y_{-k} Y_j}^{Y_0} = c_{Y_j Y_0}^{Y_k} = 1$. Therefore r_j is exactly a basis relation of \mathcal{X}_e and $n_{r_j} = 1$. Since each X_j is a transversal for Z in G (Lemma 3.1), $|Y_j y \cap Z^\#| = 1$. So all basis relations of \mathcal{X}_e inside $Y_0 \times Z^\#$ are of valency 1. The above

discussion yields that for every $\Delta \in F(\mathcal{X}_e)$, there is a basis relation of valency 1 inside $Y_0 \times \Delta$. Thus, \mathcal{X}_e is separable whenever $(\mathcal{X}_e)_{Y_0}$ so is by Lemma 2.2(2).

Let us verify that $(\mathcal{X}_e)_{Y_0}$ is separable. From Lemma 2.2(1) it follows that $\text{Aut}((\mathcal{X}_e)_{Y_0}) \cong \text{Aut}(\mathcal{X}_e)$. Eq. (1) implies that $\text{Aut}(\mathcal{X}_e) = \text{Aut}(\mathcal{X})_e$. By Proposition 3.4, we have $\text{Aut}(\mathcal{X}) = \text{Aut}(\mathcal{A}) = G_{\text{right}} \rtimes K$ and hence $\text{Aut}(\mathcal{X})_e = K \cong C_{q^2-1}$. Thus,

$$\text{Aut}((\mathcal{X}_e)_{Y_0}) \cong C_{q^2-1}.$$

Since Y_0 is an orbit of K and $|Y_0| = q^2 - 1$, the group $\text{Aut}((\mathcal{X}_e)_{Y_0})$ is regular. Therefore $(\mathcal{X}_e)_{Y_0}$ is isomorphic to a Cayley scheme over a cyclic group with a regular automorphism group. From [5, Corollary 4.4.4] it follows that $(\mathcal{X}_e)_{Y_0}$ is schurian and hence regular. Thus, $(\mathcal{X}_e)_{Y_0}$ is separable by Lemma 2.1 as required.

If $q > 9$, then Corollary 4.2 and Lemma 4.3 imply that there is at least one digraph non-isomorphic to Γ_i which is WL-equivalent to Γ_i . So $\dim_{\text{WL}}(\Gamma_i) > 2$ and hence $\dim_{\text{WL}}(\Gamma_i) = 3$. If $q = 3$, then it can be verified using the list of small schemes [15] that \mathcal{X} is separable and hence $\dim_{\text{WL}}(\Gamma_i) = 2$ by Lemma 2.7(1). \square

Proposition 4.5. *For all $i, j \in \mathbb{F}_q$, the divisible designs $\mathcal{D}(\Gamma_i) = \text{dev}(X_i)$ and $\mathcal{D}(\Gamma_j) = \text{dev}(X_j)$ are isomorphic.*

Proof. Recall that given $i \in \mathbb{F}_q$, $\text{dev}(X_i) = (G, \mathcal{B}(X_i))$, where $\mathcal{B}(X_i) = \{X_i g : g \in G\}$. We are going to check that $\text{dev}(X_i)$ is isomorphic to $\text{dev}(X_0)$ for every $i \in \mathbb{F}_q^*$ and by that to prove the lemma. To do this, it suffices to find two bijections $f, h : G \rightarrow G$ such that

$$(11) \quad g \in X_0 g_0 \Leftrightarrow f(g) \in X_i h(g_0)$$

for all $g = \begin{pmatrix} 1 & \alpha & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix}$, $g_0 = \begin{pmatrix} 1 & \alpha_0 & \gamma_0 \\ 0 & 1 & \beta_0 \\ 0 & 0 & 1 \end{pmatrix} \in G$. One can see that $g \in X_0 g_0$ if and only if $g g_0^{-1} \in X_0$. Due to the definition of X_0 , the latter is equivalent to

$$(12) \quad \gamma - \gamma_0 = \frac{(\alpha - \alpha_0)(\beta + \beta_0)}{2}.$$

Let $f : \begin{pmatrix} 1 & \alpha & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & \alpha' & \gamma' \\ 0 & 1 & \beta' \\ 0 & 0 & 1 \end{pmatrix}$ and $h : \begin{pmatrix} 1 & \alpha_0 & \gamma_0 \\ 0 & 1 & \beta_0 \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & \alpha_0'' & \gamma_0'' \\ 0 & 1 & \beta_0'' \\ 0 & 0 & 1 \end{pmatrix}$ be mappings from G to itself such that

$$(13) \quad \begin{cases} \alpha = \alpha', \\ \beta = \beta', \\ \gamma = \gamma' - ((\alpha')^2 - \varepsilon(\beta')^2)i, \end{cases}$$

$$(14) \quad \begin{cases} \alpha_0 = \alpha_0'' - \delta, \\ \beta_0 = \beta_0'' - \sigma, \\ \gamma_0 = \gamma_0'' + \frac{(\alpha_0'' - \delta)(\beta_0'' - \sigma)}{2} - \frac{\alpha_0'' \beta_0''}{2} + ((\alpha_0'')^2 - \varepsilon(\beta_0'')^2)i, \end{cases}$$

where

$$(15) \quad \delta = 4i\varepsilon\beta_0'' \text{ and } \sigma = 4i\alpha_0''.$$

At first, let us verify that f and h are bijections. Since the mapping $\gamma \mapsto \gamma'$ is a translation by a function of α and β , we conclude that f is a bijection. Due to Eqs. (14) and (15), we

obtain

$$\begin{pmatrix} \alpha_0 \\ \beta_0 \end{pmatrix} = A \begin{pmatrix} \alpha''_0 \\ \beta''_0 \end{pmatrix},$$

where

$$A = \begin{pmatrix} 1 & -4\varepsilon i \\ -4i & 1 \end{pmatrix}.$$

One can see that $\det(A) = 1 - 16\varepsilon i^2 \neq 0$ because ε is nonsquare. Therefore the mapping

$$\begin{pmatrix} \alpha_0 \\ \beta_0 \end{pmatrix} \mapsto \begin{pmatrix} \alpha''_0 \\ \beta''_0 \end{pmatrix}$$

is a bijection. The mapping $\gamma_0 \mapsto \gamma''_0$ is a translation by a function of α_0 and β_0 which yields that h is a bijection.

Substituting expressions for α , β , γ via α' , β' , γ' from Eq. (13) and for α_0 , β_0 , γ_0 via α''_0 , β''_0 , γ''_0 from Eq. (14) to Eq. (12) and making computations, we obtain

$$\gamma' - \gamma''_0 = \frac{(\alpha' - \alpha''_0)(\beta' + \beta''_0)}{2} + ((\alpha' - \alpha''_0)^2 - \varepsilon(\beta' - \beta''_0)^2)i.$$

The latter is equivalent to $f(g)h(g_0)^{-1} = \begin{pmatrix} 1 & \alpha' & \gamma' \\ 0 & 1 & \beta' \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \alpha''_0 & \gamma''_0 \\ 0 & 1 & \beta''_0 \\ 0 & 0 & 1 \end{pmatrix}^{-1} \in X_i$. Thus, Eq. (11) holds and hence $\text{dev}(X_0)$ and $\text{dev}(X_i)$ are isomorphic. \square

Theorem 1.1 follows from Corollary 4.2, Lemma 4.3, Proposition 4.4, and Proposition 4.5.

REFERENCES

1. *K. T. Arasu, D. Jungnickel, A. Pott*, Divisible difference sets with multiplier -1 , *J. Algebra*, **133** (1990), 35–62.
2. *T. Beth, D. Jungnickel, H. Lenz*, *Design Theory*, 2nd edition, Cambridge University Press, Cambridge (1999).
3. *R. Bildanov, V. Panshin, G. Ryabov*, On WL-rank and WL-dimension of some Deza circulant graphs, *Graphs Combin.*, **37**, No. 6 (2021), 2397–2421.
4. *A. Brouwer, A. Cohen, A. Neumaier*, *Distance-regular graphs*, Springer, Heidelberg (1989).
5. *G. Chen, I. Ponomarenko*, *Coherent configurations*, Central China Normal University Press, Wuhan (2019).
6. *D. Crnković, W. H. Haemers*, Walk-regular divisible design graphs, *Des. Codes Cryptogr.*, **72** (2014) 165–175.
7. *D. Crnković, H. Kharaghani*, Divisible design digraphs, in: *Algebraic Design Theory and Hadamard Matrices*, (C. J. Colbourn, Ed.), Springer Proc. Math. Stat., Springer, New York **133** (2015), 43–60.
8. *D. Crnković, H. Kharaghani, A. Švob*, Divisible design Cayley digraphs, *Discrete Math.*, **343**, No. 4 (2020), Article ID 111784.
9. *D. Crnković, A. Švob*, New constructions of divisible design Cayley graphs, *Graphs Comb.*, **38** (2022) Article number 17.
10. *S. Evdokimov, I. Ponomarenko*, Characterization of cyclotomic schemes and normal Schur rings over a cyclic group, *St. Petersburg Math. J.*, **14**, No. 2 (2003), 189–221.
11. *F. Fuhlbrück, J. Köbler, O. Verbitsky*, Identifiability of graphs with small color classes by the Weisfeiler-Leman algorithm, in: *Proc. 37th International Symposium on Theoretical Aspects of Computer Science*, Dagstuhl Publishing, Germany (2020), 43:1–43:18.

12. *S. Goryainov, W.H. Haemers, V. Kabanov, L. Shalaginov*, Deza graphs with parameters $(n, k, k-1, a)$ and $\beta = 1$, *J. Comb. Des.*, **27** (2019) 188–202.
13. *M. Gröhe*, Descriptive complexity, canonisation, and definable graph structure theory, Cambridge University Press, Cambridge (2017).
14. *W. H. Haemers, H. Kharaghani, M. A. Meulenberg*, Divisible design graphs, *J. Combin. Theory, Ser. A*, **118** (2011) 978–992.
15. *A. Hanaki, I. Miyamoto*, Classification of association schemes with small number of vertices. <http://math.shinshu-u.ac.jp/~hanaki/as/> (2016).
16. *B. Huppert*, Endliche Gruppen, Springer Berlin-Heidelberg-New York, **1** (1967).
17. *V. Kabanov*, New versions of the Wallis-Fon-Der-Flaass construction to create divisible design graphs, *Discrete Math.*, **345**, No. 11 (2022) Article ID 113054.
18. *V. Kabanov, L. Shalaginov*, On divisible design Cayley graphs, *Art Discrete Appl. Math.*, **4**, No. 2 (2021), 1–9.
19. *R. Lidl, H. Niederreiter*, Finite fields, Cambridge University Press, Second edition (1997).
20. *M. Muzychuk, G. Ryabov*, Constructing linked systems of relative difference sets via Schur rings, *Des. Codes Cryptogr.*, **92** (2024), 2615–2637.
21. *D. Panasenko, L. Shalaginov*, Classification of divisible design graphs with at most 39 vertices, *J. Comb. Des.*, **30**, No. 4 (2022), 205–219.
22. *J. B. Rosser, L. Schoenfeld*, Approximate formulas for some functions of prime numbers, *Illinois J. Math.*, **6**, No. 1 (1962), 64–94.
23. *G. Ryabov*, On separability of Schur rings over abelian p -groups, *Algebra Log.*, **57**, No. 1 (2018), 49–68.
24. *R.-H. Schulz, A. G. Spera*, Divisible designs and groups, *Geom. Dedicata*, **44** (1992), 147–157.
25. *I. Schur*, Zur theorie der einfach transitiven Permutationgruppen, *S.-B. Preus Akad. Wiss. Phys.-Math. Kl.*, **18**, No. 20 (1933), 598–623.
26. *B. Weisfeiler (ed.)*, On construction and identification of graphs, Springer-Verlag, Berlin-New York (1976).
27. *B. Weisfeiler, A. Leman*, Reduction of a graph to a canonical form and an algebra which appears in the process, *NTI*, **2**, No. 9 (1968), 12–16.
28. *H. Wielandt*, Finite permutation groups, Academic Press, New York - London (1964).

BEN GURION UNIVERSITY OF THE NEGEV, BEER SHEVA, ISRAEL
Email address: muzychuk@bgu.ac.il

SCHOOL OF MATHEMATICAL SCIENCES, HEBEI KEY LABORATORY OF COMPUTATIONAL MATHEMATICS AND APPLICATIONS, HEBEI NORMAL UNIVERSITY, SHIJIAZHUANG 050024, P. R. CHINA

BEN GURION UNIVERSITY OF THE NEGEV, BEER SHEVA, ISRAEL
Email address: gric2ryabov@gmail.com