Comparison of global networks of third-generation gravitational-wave detectors

Michele Maggiore (D, 1, 2) Francesco Iacovelli (D, 1, 2) Enis Belgacem (D, 1, 2) Michele Mancarella (D, 3) Niccolò Muttoni (D, 1, 2)

¹Département de Physique Théorique, Université de Genève, 24 quai Ernest Ansermet, 1211 Genève 4, Switzerland

²Gravitational Wave Science Center (GWSC), Université de Genève, CH-1211 Geneva, Switzerland

³Aix-Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France

E-mail: Michele.Maggiore@unige.ch, Francesco.Iacovelli@unige.ch, Enis.Belgacem@unige.ch, mancarella@cpt.univ-mrs.fr, Niccolo.Muttoni@unige.ch

Abstract. We study the performances of a world-wide network made by a European thirdgeneration gravitational-wave (GW) detector, together with a 40-km Cosmic Explorer detector in the US, considering three scenarios for the European detector: (1) Einstein Telescope (ET) in its 10-km triangle configuration; (2) ET in its configuration featuring two 15-km L-shaped detectors in different sites, still taken to have all other ET characteristics (underground, and with each detector made of a high-frequency interferometer and a cryogenic low-frequency interferometer); (3) A single L-shaped underground interferometer with the ET amplitude spectral density, either with 15 km or with 20 km arm length. Overall, we find that, if a 2L configuration should be retained for ET, the network made by a single-L European underground detector together with CE-40km could already provide a very interesting intermediate step toward the construction of a full 2L+CE network, and is in any case superior to a 10-km triangle not inserted in an international network.

Contents

T	Intro	oduction	1
2	Resu	lts	3
	2.1	Horizons	3
	2.2	Parameter reconstruction for BBHs	4
	2.3	Parameter reconstruction for BNSs	5
	2.4	Pre-merger alerts	10
	2.5	Stochastic backgrounds	10
3	Conc	lusions	13

1 Introduction

The detection of gravitational waves (GWs) in the last few years has opened a new way of observing the Universe [1-9]. A new generation of ground-based GW detectors, referred as third-generation (3G) or next-generation GW detectors, is under study, to overcome the limitations imposed by current detector infrastructures. The European project is Einstein Telescope (ET) [10–12], while the US community effort is represented by the Cosmic Explorer (CE) project [13–15].

Several studies have been conducted in the last few years, to compare the scientific potential of various 3G detector networks [16, 17] (see also refs. [18–21] for follow-up studies elaborating on various aspects of ref. [17], and refs. [22, 23] for studies of the capabilities of further 3G network configurations). In particular, in ref. [17] two different geometries for ET were compared, a single-site triangular geometry made of 3 nested detectors (with each detector made by a high-frequency interferometer and a cryogenic low-frequency interferometer), and a network made of two identical L-shaped detectors ('2L' in the following), located in two different sites within Europe, but still underground, and both made by a high-frequency interferometer. These different configurations for ET were considered in a ET-only scenario, as well as in a broader world-wide network including also a single 40-km Cosmic Explorer (CE) detector, or two CE detectors with arm-lengths of 20 and 40 km, respectively.

Recently, a report of the NSF MPS AC Subcommittee on Next-Generation Gravitational-Wave Detector Concepts [15] included a network of a single CE-40km together with ET (which was considered in the triangle configuration) in a list of recommended world-wide next-generation networks.¹ The performance of such a network, as well as of a network made by a single CE-40km together with ET in its 2L configuration, was already discussed in detail in [17]. In this paper, we extend the analysis to the situation in which the European detector is made of just a single L-shaped underground detector, with the amplitude spectral density (ASD) of ET, and 15 km or 20 km arm-length. Such a network could also be seen as an intermediate step toward the realization of a larger world-wide detector network, featuring a

¹Note, however, that the baseline configuration for the CE project is a pair of detectors, one with 20 km arms and the other with 40 km arms; see [22] for an exhaustive study of the different CE configurations, with and without ET.



Figure 1. Amplitude spectral densities (ASDs) used in this work. Note that the ASDs are always defined as if we had a single L-shaped detector of the given arm-length; for the triangle, they are then combined as discussed in [17].

2L ET configuration together with a 40-km CE detector. Therefore, we consider the following configurations:

- ET in the 10-km triangle configuration (denoted ET-Δ), together with a single 40-km CE located in the US.
- ET in the configuration of two underground L-shaped detectors with 15 km arms, together with a single 40-km CE located in the US.
- A single L-shaped underground detector with the ASD of ET, with 15 km arms, together with a single 40-km CE located in the US.
- A single L-shaped underground detector with the ASD of ET, with 20 km arms, together with a single 40-km CE located in the US.

We will denote these networks, respectively, as $(\text{ET-}\Delta + \text{CE-}40\text{km})$, (ET-2L + CE-40km), $(1\text{L}_{\text{EU}}^{15\,\text{km}} + \text{CE-}40\text{km})$ and $(1\text{L}_{\text{EU}}^{20\,\text{km}} + \text{CE-}40\text{km})$.² The results for $(\text{ET-}\Delta + \text{CE-}40\text{km})$ and (ET-2L + CE-40km) were already shown in [17], and here we will plot them together with the results for $(1\text{L}_{\text{EU}}^{15\,\text{km}} + \text{CE-}40\text{km})$ and $(1\text{L}_{\text{EU}}^{20\,\text{km}} + \text{CE-}40\text{km})$, which are new. Note that, in ref. [17], the ET-2L case was studied both when the two L-shaped detectors are misaligned, with a relative angle close to 45°, and when they are parallel. Here, in order not to overburden the plots, we will only show the results for (ET-2L + CE-40km) when ET-2L are in the

²We purposely refrain from using a label "ET" for the latter two configurations, reserving it to the currently officially recognized ET configurations, i.e. triangle or 2L. Concerning the locations, we place for definiteness the triangle in the Sardinia candidate site, but essentially no significant difference would be obtained putting it in the Meuse-Rhine (EMR) candidate site. In the ET-2L configuration we put the two L-shaped detectors in the Sardinia and EMR candidate sites; basically the same results would be obtained using the Sardinia candidate site and the recently proposed candidate site in Saxony, since the cord distances are very similar, see footnote 52 in [17]. In the 1L case, we put the single L in the Sardinia candidate site, but again, at the level of our analysis, no significant difference would be obtained putting it in the Meuse-Rhine (EMR) candidate site. For the 40-km CE in the US, we use for definiteness the location and orientation in Table III of [24].

misaligned configuration, which provides the best results for compact binary coalescences; when we wish to emphasize this, we will actually denote this configuration as (ET-2L-mis + CE-40km), otherwise, in the interest of readability, we will just use the notation (ET-2L + CE-40km). We will instead use a generic orientation with respect to the CE detector in the US (we use the same orientations as in [17]). In the (1L_{EU} + CE-40km) configurations, we will set the two detector at 45° with respect to the great circle joining them.

Of course, these three networks have different levels of complexity. The ET- Δ configuration involves 3 nested detectors of 10 km arms, ET-2L involves two detectors with 15 km arms, and the single-L just one detector, of 15-km or 20 km arms (where, in all cases, a detector is actually made by two interferometers, a low-frequency and a high-frequency interferometer, so ET- Δ is made of 6 interferometers, ET-2L of 4, and the single L of 2).³ We already know from the study in [17] that, for coalescing binaries, the ET-2L-15km configuration performs better than the 10-km triangle when they are both considered in isolation, and that the differences become less evident when they are put in a network with a 40-km CE.

Obviously, the configuration with a single L is bound to be less performant than the 2L configuration or the triangle, but our aim is to understand how much we lose in term of science, with the single-L detector, for example in case of delay in the implementation of the full European ET network or in case of prolonged absence of one of the two ET nodes, in a scenario that also features a 40-km CE.

We will also compare with the results that can be obtained with the ET- Δ configuration not inserted in a network with CE, as well as with the results that could be obtained by the most advanced 2G detector network, namely LIGO Hanford, LIGO Livingston, Virgo, KAGRA and LIGO India, using the publicly available best sensitivities that are planned to be achieved by the end of the O5 run [25]. We denote this network as LVKI O5.

Our methodology (astrophysical populations, parameter estimation codes, waveforms, etc.) is identical to the one already followed in Section 3 of ref. [17], whom we refer the reader for more details. In particular, for compact binary coalescences, we perform parameter estimation in the Fisher matrix approximation, using the GWFAST code [26, 27].⁴

2 Results

In this section we present the results of the analysis. The structure of the plots and the definition of the various observables are the same as in [17, 23, 26].

2.1 Horizons

Figure 2 shows the detection horizons for equal-mass non-spinning coalescing binaries, as a function of the total mass of the binary, for the 3G networks considered and, for comparison,

³We stick to the recommended terminology from the 2020 ET Design Report Update, https://apps. et-gw.eu/tds/?r=18715: the high-frequency (HF) and low-frequency (LF) interferometers that make the so-called "xylophone" configuration are indeed referred to as "interferometers". The combination of a HF interferometer and a LF interferometer (whether in a L-shaped geometry, or with arms at 60° as in the triangle configuration) is called a "detector". The whole ensemble of detectors is called an "observatory".

⁴Recently, the effect of correlated noise on parameter estimation has been studied in [28]. Although the results in [28] apply to 2 colocated L-shaped detectors, and not to the triangle configuration, when using the realistic coherence values found in [29], they would point toward the fact that, for parameter estimation of coalescing binaries, correlated noise can be neglected. Correlated noise are, however, significant for stochastic backgrounds in the triangle configuration, as discussed in Section 5.4 of ref. [17], and as we will discuss again below.



Figure 2. Detector horizons for equal mass, non-spinning binaries for the various detector geometries considered. In each panel we also add, for comparison, LVKI O5.

Detector configuration	$z_{\rm hor}(2.7~{ m M}_{\odot})$
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	8.5 9.0 8.1 8.3 3.3
LVKI O5	0.3

Table 1. Horizon redshifts for equal mass non-spinning binaries evaluated at a source-frame total mass $M_{\rm tot} = 2.7 \ {\rm M}_{\odot}$ for the various configurations considered.

also for ET- Δ and LVKI O5. We see that, in terms of detection horizons, all the 3G configurations are quite similar (except for ET- Δ in isolation, which is less performant at low masses), and allow us to make a large jump with respect to the best possible 2G detector network. In particular, for $M_{\rm tot} = 2.7 \, {\rm M}_{\odot}$, a typical value for the total mass of a BNS, the values of the horizons are given in Table 1. We see that, in all cases, the horizons are well beyond the peak of the star formation rate, at $z_p \sim 2-3$, and basically cover the whole range of redhsifts relevant for BNS.

2.2 Parameter reconstruction for BBHs

The results from the accuracy of parameter reconstruction for BBHs is shown in Fig. 3. Table 2 gives, more quantitatively, the number of detections with some cuts on angular localization or on luminosity distance. Here and in the following, all the results are given for one year of data, with the duty cycle computed as in [17].

As expected, among the world-wide networks, the hierarchy is that the best results come from (ET-2L-mis + CE-40km), followed by (ET- Δ + CE-40km), and then by (1L_{EU} +

BBH						
Detector	$\Delta \Omega_{g}$	Detection $_{00\%} \leq$	ons with $\Delta d_L/d_L \leq$			
	1 deg^2	10 deg^2	5×10^{-3}	10^{-2}		
$ET-\Delta + 1CE$	2 447	29924	395	2 901		
ET-2L-mis + 1CE	3 743	36457	575	4301		
$1L_{EU}^{15 \text{ km}} + 1CE\text{-mis}$	1557	19433	199	1603		
$1L_{EU}^{20km} + 1CE$ -mis	2077	23185	283	2116		
$\mathrm{ET} ext{-}\Delta$	35	914	2	28		
LVKI O5	38	599	1	1		

Table 2. Number of detected BBH sources at the considered networks, with different cuts on the sky localization and relative error on the luminosity distance. The results are given for one year of data, with the duty cycle computed as in [17].

CE-40km) [with $(1L_{EU}^{20 \text{ km}} + \text{CE-40km})$ only marginally better than $(1L_{EU}^{15 \text{ km}} + \text{CE-40km})$], but we see that the results are quite similar among these 3G global networks: the differences among the configurations are in general within about a factor of 2, and all these configurations provide a large jump compared to 2G detectors. As far as BBHs are concerned, the $(1L_{EU} + \text{CE-40km})$ option therefore appears to maintain a relevant scientific output, also in the version with 15 km arms. A similar conclusion emerges from Figure 4, which gives the redshift distribution of BBH golden events, i.e. events detected with especially high SNR, or especially good accuracy on luminosity distance, or especially good angular localization. We also observe that each of these international network is better than ET- Δ in isolation, particularly for angular localization and luminosity distance.

2.3 Parameter reconstruction for BNSs

The corresponding results for BNSs are shown in Figure 5. From this we see that, while the results for the SNR distribution and for the accuracy of reconstruction of the chirp mass \mathcal{M}_c , of the symmetric mass ratio η , and of the tidal deformability $\tilde{\Lambda}$ are similar among the 3G world-wide networks considered, a large difference appears in angular localization, luminosity distance, orbit inclination θ_{JN} and polarization angle ψ , where the (ET-2L-mis + CE-40km) and (ET- Δ + CE-40km) configurations are better than (1L_{EU} + CE-40km) by one order of magnitude. This is also shown, for some specific cuts on angular localization or on luminosity distance, in Table 3. The corresponding results for "golden events" are shown in Figure 6.

It is also interesting to observe that the $(1L_{EU} + CE-40km)$ configurations are, in turn, significantly better than the triangle in isolation, in particular for angular localization, luminosity distance and polarization angle.



Figure 3. Cumulative distributions of the number of detections per year, for the SNRs and for the error on the parameters, for BBH signals, for the various detector configurations considered.



Figure 4. Redshift distribution of BBHs detected with SNR ≥ 100 (left column), or relative error on the luminosity distance $\Delta d_L/d_L \leq 0.05$ (central column), or sky location $\Delta \Omega_{90\%} \leq 10 \text{ deg}^2$ (right column) for the various detector configurations considered.

BNS						
Detector configuration	$\Delta \Omega_{0}$	Detection $_{00\%} \leq$	as with $\Delta d_L/d_L \leq$			
	$10 \ \mathrm{deg}^2$	$ 100 \deg^2$	5×10^{-2}	$ 10^{-1} $		
$ET-\Delta + 1CE$	2427	54994	535	4 100		
ET-2L-mis + 1CE	3838	75828	1 0 4 0	7949		
$1L_{EU}^{15 \text{ km}} + 1CE\text{-mis}$	252	5388	59	347		
$1L_{EU}^{20km} + 1CE$ -mis	391	7852	90	506		
$\text{ET-}\Delta$	8	184	8	52		
LVKI O5	31	51	0	1		

Table 3. Number of detected BNS sources at the considered networks, with different cuts on the sky localization and relative error on the luminosity distance. The results are given for one year of data, with the duty cycle computed as in [17].



Figure 5. Cumulative distributions of the number of detections per year, for the SNRs and for the error on the parameters, for BNS signals, for the various detector configurations considered.



Figure 6. Redshift distribution of BNSs detected with SNR ≥ 30 (left column), relative error on the luminosity distance $\Delta d_L/d_L \leq 0.2$ (central column), or sky location $\Delta \Omega_{90\%} \leq 100 \text{ deg}^2$ (right column) for the various detector configurations considered.



Figure 7. Accumulation of the SNR in the various 3G network configurations, for a source with the properties of the BNS GW170817, as the signal sweeps up in frequency. The upper horizontal scale gives the corresponding time to merger. Observe that, for this system, the merger takes place at about 2 kHz, see e.g. Figure 2 of [26].

2.4 Pre-merger alerts

We next consider the ability of the various networks to provide an alert to electromagnetic telescopes before the merger.

Figure 7 shows how the SNR would accumulate in the various 3G networks considered, for an event with the characteristics of GW170817, as the signal sweeps up in frequency. For this event, which was very close in distance, all world-wide 3G network configurations considered have similar performances and rapidly reach a very large SNR. A SNR larger than a detection threshold, say set at SNR = 12, is reached in all case about 10 hours prior to merger.

For multi-messenger studies, however, we also need to get a good angular resolution before merger, in order to give to electromagnetic observatories at least some localization information, and sufficiently early. Table 4 shows the number of BNSs detected with SNR \geq 12 and different cuts on the sky localization, at 30 min, 10 min and 1 min prior to merger, for the various network configurations considered. We show separately the BNS detections where the orbit has a generic inclination, and those close to face-on, defined as those such that $\Theta \equiv \min{\{\iota, 180^\circ - \iota\}}$ is smaller than 15°, which can result more likely in coincident γ -ray burst (GRB) detections.

The results reflect the trends observed in Figure 5 for angular localization, with the (ET-2L-mis + CE-40km) and (ET- Δ + CE-40km) configurations better than (1L_{EU} + CE-40km) by one order of magnitude, and the latter significantly better than ET- Δ in isolation.

2.5 Stochastic backgrounds

Finally, we consider the sensitivity to stochastic GW backgrounds (SGWBs) of these configurations.

BNS							
Detector	Time before	Orientation	Detections with $\Delta \Omega_{90\%} \leq$				
	merger		10 deg^2	$100 \ \mathrm{deg}^2$	$1000 \mathrm{deg}^2$	all sky	
	30 min	All Θ	1	25	229	418	
		$\Theta \le 15^\circ$	0	0	5	37	
$ $ ET- $\Lambda \perp 1$ CE	10 min	All Θ	11	234	1 888	2 4 9 3	
$ E^{1-\Delta} + 10E$		$\Theta \leq 15^\circ$	0	9	64	233	
	1 min	All Θ	77	2 1 4 0	22906	33 0 4 2	
		$\Theta \le 15^{\circ}$	3	76	790	3 1 3 6	
	20 main	All Θ	0	41	307	875	
	30 min	$\Theta \leq 15^\circ$	0	0	9	82	
\mathbf{FT} 2L min $+ 1$ CF	10 min	All Θ	10	363	2521	4 5 4 2	
E_1 -2L-IIIIS + ICE		$\Theta \le 15^\circ$	0	10	79	417	
	1 min	All Θ	101	2824	27880	42804	
	1 min	$\Theta \le 15^\circ$	7	82	909	4057	
	30 min	All Θ	0	10	59	399	
		$\Theta \le 15^\circ$	0	1	2	35	
$1I^{15}km + 1CF$ mia	10 min	All Θ	0	41	373	2 374	
$1L_{EU} + 1CE-mis$		$\Theta \leq 15^\circ$	0	1	14	229	
	1 min	All Θ	8	229	2 343	30344	
		$\Theta \le 15^{\circ}$	0	9	98	2861	
	30 min	All Θ	0	12	96	757	
		$\Theta \leq 15^\circ$	0	0	0	69	
$1L^{20 \text{ km}} \perp 1CE$ mis	10 min	All Θ	2	59	535	3 6 3 7	
$1L_{EU} + 10L-1015$		$\Theta \le 15^\circ$	0	2	25	342	
	1 min	All Θ	15	346	3268	35 086	
		$\Theta \le 15^\circ$	0	14	103	3 3 1 8	
	30 min	All Θ	0	8	39	345	
		$\Theta \le 15^{\circ}$	0	0	2	31	
	10 min 1 min	All Θ	0	28	125	1 5 4 4	
		$\Theta \le 15^\circ$	0	2	4	153	
		All Θ	4	79	372	$7\overline{599}$	
		$\Theta \le 15^{\circ}$	0	3	9	767	

Table 4. Number of BNS detected with SNR ≥ 12 and different cuts on the sky localization at 30 min, 10 min and 1 min prior to merger for the various detector configurations studied in this work. We further report the numbers both for all the sources in the catalog and the sub-sample of sources having an angle $\Theta \leq 15^{\circ}$, which can result more likely in coincident GRB detections.



Figure 8. Power-law integrated sensitivity curves for $h^2 \Omega_{gw}(f)$ for the various configurations studied. These are computed assuming an observational time $T_{obs} = 1$ yr and an SNR threshold of 1. We further report some SGWB examples, namely a flat-in-frequency background with different amplitudes and the AGWB generated by our 1 yr catalogs of BBHs and BNSs. The PLS are computed assuming uncorrelated noise among the detectors of a given network which, particularly for the triangle, might lead to a significant overestimate of the actual sensitivity to stochastic backgrounds, especially at low frequencies.

Detector	SNR for				
configuration	flat SGWB, $h^2 \Omega_{\rm gw} =$			AGWB	
	10^{-10}	10^{-11}	10^{-12}	all BBH	all BNS
$\text{ET-}\Delta + 1\text{CE}$	358.8	35.9	3.6	828.7	234.9
ET-2L-mis + 1CE	285.3	28.5	2.9	698.7	199.7
$1L_{EU}^{15 \text{ km}} + 1CE\text{-mis}$	49.9	5.0	0.5	145.0	42.7
$1L_{EU}^{20 \text{ km}} + 1CE\text{-mis}$	60.1	6.0	0.6	175.2	51.6
$ ext{ET-}\Delta$	285.2	28.5	2.9	621.6	174.5

Table 5. SNRs at the various detector configurations for different stochastic background sources in an observational time $T_{\rm obs} = 1$ yr. The columns 2, 3 and 4 show the results for a flat-in-frequency SGWB with different amplitudes, the fifth and sixth column for the superposition of all the BBH and BNS signals present in our 1 yr catalogs, respectively. All results assume uncorrelated noise among the detectors of a given network.

The sensitivity to stochastic background can be expressed in terms of the power-law integrated sensitivity (PLS) introduced in [30] (see also App. A of [17] for conventions and definitions). Figure 8 shows the PLS of the various network configurations studied, expressed in terms of $h^2 \Omega_{gw}(f)$ where, as usual, $\Omega_{gw}(f) = (1/\rho_c) d\rho_{gw}/d \log f$, $\rho_c = 3H_0^2/(8\pi G)$ is the critical density for closing the Universe, and h is defined from $H_0 = h \times (100 \text{ km s}^{-1} \text{Mpc}^{-1})$ [31].

An important caveat, when interpreting the PLS shown in Figure 8, is that these are obtained assuming that the noise in the different detectors are uncorrelated. More precisely, the output in the *i*-th detector can be written as $s_i(t) = n_i(t) + h_i(t)$, where $n_i(t)$ is the noise of the *i*-th detector and $h_i(t)$ the projection of the GW signal onto the *i*-th detector. Let us recall that, if the noise between two different detectors is uncorrelated, the noise-noise

correlator only grows with the observation time T as \sqrt{T} (as in a random walk) while the GW signal, which is correlated between the two detectors, grows as T (see e.g. Section 7.8.3) of [32] for more accurate discussion). In this way, correlating the output of two detectors and integrating for a long time, we can extract a small signal from a much larger noise. As an order-of-magnitude estimate, correlating for a time T the output of two detectors with a useful bandwidth Δf , we gain a factor of order $\sqrt{2T\Delta f}$ in sensitivity (see eq. 7.246 of [32]). Setting for instance Δf of order of a few hundred Hz, and T = 1 yr, we then gain a factor $\mathcal{O}(10^5)$ in sensitivity. The curves in Figure 8 have indeed been computed assuming uncorrelated noise, and integrating for 1 yr. If, however, a noise is correlated, also the corresponding noise-noise correlator will grow as T rather than \sqrt{T} . In this case, if in some frequency range a correlated noise is above a GW stochastic signal, we can integrate for as long as we want, but it will remain above the signal. This problem is especially important for the triangle, with its three colocated interferometers. In particular, correlated noise are expected at low frequencies, because of Newtonian, seismic and magnetic noise, see [29, 33] and Section 5.4.1 of [17]. This means that, at low frequencies, the actual sensitivity of the triangle to stochastic GW backgrounds could actually be up to five orders of magnitude worse than that shown in Figure 8. The actual sensitivity is in fact difficult to compute, because it depends strongly on the level of correlated noise between the interferometers of the triangle configuration, which is difficult to predict before constructing and commissioning the detector. However, ref. [29], reports correlations of several hundreds meters to a few kilometers in underground seismic measurements from the ET candidate sites, and also elsewhere, in the frequency range from 0.01 to 40 Hz. From this, they project that the effect of correlated Newtonian noise from body waves on the sensitivity of ET in the triangle configurations will be severely degraded at least up to 20 Hz. Even if, to some extent, some correlated noise will be present even between two well-separated L-shaped detectors, in this case the problem will be much less severe, and in particular the correlation of seismic noise will be negligible in the ET frequency band.

In Figure 8 we also show, for comparison, the current best estimate of the astrophysical GW background (AGWB) obtained from the superposition of all BBH signals, and that obtained from all BNS signals (computed as in Section 5.3 of [17], see also [34, 35]). To guide the eye, we further draw horizontal lines corresponding to a stochastic background where $h^2 \Omega_{gw}$ is flat in frequency, and chosen to have the values 10^{-10} , 10^{-11} , or 10^{-12} . Signals for which $h^2 \Omega_{gw}$ is approximately constant in f, at least over the range of frequencies explored by ground-based GW detectors, emerge rather naturally in cosmology, where typical spectra extend over many more decades in frequency, and the range explored by ground-based detectors is comparatively small; an explicit example is provided by some models for cosmic strings, see e.g. Figure 63 of [17]. The SNR for a given stochastic background signal is then computed as in Section 7.8.3 of [32]. The results are shown in Table 5 where, again, we have assumed uncorrelated noise, so the results for the triangle might actually be a gross overestimate of the actual sensitivity.

3 Conclusions

In this paper we have studied the performances of a global international network made by a single 40 km CE detector in the US, together with a European third-generation GW detector. For the latter, we have examined three different options: the two reference configurations for ET, namely a 10 km triangle and a 15-km 2L, and also a 1L configuration, in which in

Europe there is only a single underground L-shaped detector with the ASD of ET. We have denoted these configuration, respectively, as $(\text{ET-}\Delta + \text{CE-}40\text{km})$, (ET-2L + CE-40km) and $(1\text{L}_{\rm EU} + \text{CE-}40\text{km})$. In the latter case, we have considered both 15 km or 20 km arms for the $1\text{L}_{\rm EU}$ detector.

From the point of view of the science case, the main conclusions that emerge from the comparison of these configurations are as follows.

- For the detection horizons, the differences between these configurations are marginal, see Figure 2. All these networks will detect compact binaries up to similar distances, and over the same range of binary masses, from subsolar masses (a potential smoking-gun signature of primordial BHs) to BHs with masses of several thousands M_{\odot} , which provide the bridge toward supermassive BHs.
- For BBHs, the results for parameter estimations are comparable among these networks, within factors of order 2, with (ET-2L + CE-40km) always providing the best results, followed by (ET- Δ + CE-40km) and then by (1L_{EU} + CE-40km); see Figures 3 and 4. This, combined with the results for detector horizons, implies that all aspects of the science case concerning BH astrophysics (demography, origin) are not significantly affected in the (1L_{EU} + CE-40km) configuration, nor is the possibility of detecting and characterizing subsolar-mass primordial BHs or intermediate-mass BH. Similarly, the perspectives for extracting cosmological information by treating BHs as dark sirens (using either the correlation with galaxy catalogs or features in the mass function) are not significantly degraded, as well as the potential for studying deviations from General Relativity or the existence of compact objects from ringdown tests.
- For BNSs the situation is different. We see from Figure 5 that the SNR distribution, and the reconstruction of masses and tidal deformabilities are not very sensitive to the configuration. Therefore, all aspects of the science cased involving NS demography, as well as those related to the extraction of nuclear physics information from measurement of the tidal deformability, would be broadly preserved in the ($1L_{EU} + CE-40$ km) configuration.

In contrast, the $(1L_{EU} + CE-40 \text{km})$ configuration is one order of magnitude less performant than $(ET-\Delta + CE-40km)$ and (ET-2L + CE-40km) on the number of events with given angular localization and with given resolution on the luminosity distance. This could impact multi-messenger astronomy and the possibility of providing pre-merger alerts. For instance, we see from Table 4 that the number of BNSs detected 10 min before merger with a localization better than $100 \, \text{deg}^2$ (in our sample realization and in 1 yr, for our choice of duty cycle) is 363 for (ET-2L + CE-40km), 234 for (ET- Δ + CE-40km), and only 41 for (1L_{EU} + CE-40km). From this point of view it should however be observed that, most likely, electromagnetic telescopes will in any case be able to perform a follow-up of only a limited number of events per year. With $(1L_{\rm EII})$ + CE-40km) we will also have a reduction in the number of the events with particularly good localization, as we see looking at the tails of the distribution in $\Delta\Omega_{90\%}$ in Figure 5. For instance, we can ask what is the cut $\Delta\Omega_{\rm cut}$ that we can impose on angular resolution such that we remain with, say, 10 events with $\Delta\Omega_{90\%} < \Delta\Omega_{\rm cut}$ (we consider a set of 10 events because looking only at the very best event will be too sensitive to statistical fluctuations associated to our sample realization); then, we find

that $\Delta\Omega_{\rm cut} \simeq 1 \, {\rm deg}^2$ for $(1 L_{\rm EU} + {\rm CE}\text{-}40 {\rm km})$, while $\Delta\Omega_{\rm cut} \simeq 0.2 \, {\rm deg}^2$ for $({\rm ET}\text{-}\Delta + {\rm CE}\text{-}40 {\rm km})$ and $\Delta\Omega_{\rm cut} \simeq 0.1 \, {\rm deg}^2$ for $({\rm ET}\text{-}2 L + {\rm CE}\text{-}40 {\rm km})$.

- The sensitivity of the various configurations to stochastic backgrounds is shown in the left panel of Figure 8. The (1L_{EU} + CE-40km) configuration improves significantly with respect to 2G detectors, but is clearly less performant than (ET-Δ + CE-40km) or (ET-2L + CE-40km). As discussed, however, the sensitivity of the configuration involving ET-Δ could be significantly degraded by correlated noise among the colocated detectors, especially at low frequencies.
- Finally, we have found that, for coalescing binaries, a $(1L_{EU} + CE-40km)$ international network provides results that are significantly better than ET- Δ in isolation. On the one hand, this could have been expected because of the large inter-continental baseline of $(1L_{EU} + CE-40km)$; on the other hand, this is non-trivial, considering that ET- Δ is made of six interferometers, while $(1L_{EU} + CE-40km)$ only of three (two making up $1L_{EU}$, and one in CE, that only has a single interferometer per detector).

To sum up, these results indicate that, in an international context in which the US community builds a single 40-km CE in the US, to be set in a network with ET, from the purely scientific point of view the European network should be based on the (ET- Δ + CE-40km) or (ET-2L + CE-40km) configurations, that provide the best results. When correlated with a 40-km CE, the ET- Δ and ET-2L options become quite comparable, with a general overall preference for ET-2L, as discussed in [17] (apart possibly for stochastic backgrounds, where however the effect of correlated noise could disfavor again the ET- Δ configuration).

However, 3G interferometers are enterprises of great technical complexity, requiring significant financial investment, manpower, etc. Should some show-stopper emerge, the result of our analysis indicate that, in the context of a network with a 40-km CE in the US (and only in the context of such a network), a European project based on just a single L-shaped underground detector with the ASD of ET would be a viable (although temporary) back-up solution, that would preserve a relevant part of the science case expected by the next generation of GW detectors. The price to pay in terms of scientific co-leadership, scientific independence and self-consistency is not evaluated in this paper. In particular, in order to guarantee a strategic and leading role for the ET project within the context of the global world-wide GW research, if the "L" geometry is chosen for ET, it is of paramount importance to implement ET in Europe through a network of detectors, each covering the whole ET frequency detection range, in order to minimise the impact of local disturbances, understand and control the noise sources, and benefit from the coherent analysis of the whole network at all frequencies.

Acknowledgements

We thank Eugenio Coccia and Mario Martinez for very useful discussions that stimulated this study, Harald Lück, Michele Punturo and B. Sathyaprakash for useful comments on the manuscript, and Jerome Degallaix and Mikhail Korobko for their useful internal ET review. The work of F.I., M.Mag. and N.M. is supported by the Swiss National Science Foundation (SNSF) grants 200020_191957 and by the SwissMap National Center for Competence in Research. E.B. and M.Mag. are supported by the SNSF grant CRSII5_213497. M.Mag. thanks for the hospitality the IFAE in Barcelona, where part of this work was done. The work of M.Manc. received support from the French government under the France 2030 investment plan, as part of the Initiative d'Excellence d'Aix-Marseille Université – A*MIDEX AMX-22-CEI-02. This work made use of the clusters Yggdrasil and Baobab at the University of Geneva.

References

- LIGO Scientific, Virgo Collaboration, B. P. Abbott *et al.*, "Observation of Gravitational Waves from a Binary Black Hole Merger," *Phys. Rev. Lett.* **116** no. 6, (2016) 061102, arXiv:1602.03837 [gr-qc].
- [2] LIGO Scientific, Virgo Collaboration, B. P. Abbott et al., "GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral," *Phys. Rev. Lett.* 119 no. 16, (2017) 161101, arXiv:1710.05832 [gr-qc].
- [3] LIGO Scientific, Virgo, Fermi-GBM, INTEGRAL Collaboration, B. P. Abbott *et al.*, "Gravitational Waves and Gamma-rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A," *Astrophys. J. Lett.* 848 no. 2, (2017) L13, arXiv:1710.05834 [astro-ph.HE].
- [4] LIGO Scientific, Virgo, Fermi GBM, INTEGRAL, IceCube, AstroSat Cadmium Zinc Telluride Imager Team, IPN, Insight-Hxmt, ANTARES, Swift, AGILE Team, 1M2H Team, Dark Energy Camera GW-EM, DES, DLT40, GRAWITA, Fermi-LAT, ATCA, ASKAP, Las Cumbres Observatory Group, OzGrav, DWF (Deeper Wider Faster Program), AST3, CAASTRO, VINROUGE, MASTER, J-GEM, GROWTH, JAGWAR, CaltechNRAO, TTU-NRAO, NuSTAR, Pan-STARRS, MAXI Team, TZAC Consortium, KU, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS, BOOTES, MWA, CALET, IKI-GW Follow-up, H.E.S.S., LOFAR, LWA, HAWC, Pierre Auger, ALMA, Euro VLBI Team, Pi of Sky, Chandra Team at McGill University, DFN, ATLAS Telescopes, High Time Resolution Universe Survey, RIMAS, RATIR, SKA South Africa/MeerKAT Collaboration, B. P. Abbott *et al.*, "Multi-messenger Observations of a Binary Neutron Star Merger," Astrophys. J. Lett. 848 no. 2, (2017) L12, arXiv:1710.05833 [astro-ph.HE].
- [5] LIGO Scientific, Virgo Collaboration, R. Abbott *et al.*, "GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run," *Phys. Rev. X* 11 (2021) 021053, arXiv:2010.14527 [gr-qc].
- [6] LIGO Scientific, VIRGO, KAGRA Collaboration, R. Abbott *et al.*, "GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run," arXiv:2111.03606 [gr-qc].
- [7] LIGO Scientific, VIRGO, KAGRA Collaboration, R. Abbott *et al.*, "The population of merging compact binaries inferred using gravitational waves through GWTC-3," arXiv:2111.03634 [astro-ph.HE].
- [8] LIGO Scientific, VIRGO, KAGRA Collaboration, R. Abbott *et al.*, "Tests of General Relativity with GWTC-3," arXiv:2112.06861 [gr-qc].
- [9] LIGO Scientific, VIRGO, KAGRA Collaboration, R. Abbott *et al.*, "Constraints on the cosmic expansion history from GWTC-3," arXiv:2111.03604 [astro-ph.CO].
- [10] M. Punturo et al., "The Einstein Telescope: A third-generation gravitational wave observatory," Class. Quant. Grav. 27 (2010) 194002.
- [11] S. Hild et al., "Sensitivity Studies for Third-Generation Gravitational Wave Observatories," Class. Quant. Grav. 28 (2011) 094013, arXiv:1012.0908 [gr-qc].

- [12] M. Maggiore et al., "Science Case for the Einstein Telescope," JCAP 2003 (2020) 050, arXiv:1912.02622 [astro-ph.CO].
- [13] D. Reitze et al., "Cosmic Explorer: The U.S. Contribution to Gravitational-Wave Astronomy beyond LIGO," Bull. Am. Astron. Soc. 51 no. 7, (2019) 035, arXiv:1907.04833 [astro-ph.IM].
- [14] M. Evans et al., "A Horizon Study for Cosmic Explorer: Science, Observatories, and Community," arXiv:2109.09882 [astro-ph.IM].
- [15] M. Evans et al., "Cosmic Explorer: A Submission to the NSF MPSAC ngGW Subcommittee," (6, 2023), arXiv:2306.13745 [astro-ph.IM].
- [16] S. Borhanian and B. S. Sathyaprakash, "Listening to the Universe with Next Generation Ground-Based Gravitational-Wave Detectors," arXiv:2202.11048 [gr-qc].
- [17] M. Branchesi *et al.*, "Science with the Einstein Telescope: a comparison of different designs," *JCAP* 07 (2023) 068, arXiv:2303.15923 [gr-qc].
- [18] A. Puecher, A. Samajdar, and T. Dietrich, "Measuring tidal effects with the Einstein Telescope: A design study," *Phys. Rev. D* 108 no. 2, (2023) 023018, arXiv:2304.05349 [astro-ph.IM].
- [19] S. Bhagwat, C. Pacilio, P. Pani, and M. Mapelli, "Landscape of stellar-mass black-hole spectroscopy with third-generation gravitational-wave detectors," *Phys. Rev. D* 108 no. 4, (2023) 043019, arXiv:2304.02283 [gr-qc].
- [20] G. Franciolini, F. Iacovelli, M. Mancarella, M. Maggiore, P. Pani, and A. Riotto, "Searching for primordial black holes with the Einstein Telescope: Impact of design and systematics," *Phys. Rev. D* 108 no. 4, (2023) 043506, arXiv:2304.03160 [gr-qc].
- [21] F. Iacovelli, M. Mancarella, C. Mondal, A. Puecher, T. Dietrich, F. Gulminelli, M. Maggiore, and M. Oertel, "Nuclear physics constraints from binary neutron star mergers in the Einstein Telescope era," *Phys. Rev. D* 108 no. 12, (2023) 122006, arXiv:2308.12378 [gr-qc].
- [22] I. Gupta *et al.*, "Characterizing Gravitational Wave Detector Networks: From A[#] to Cosmic Explorer," (7, 2023), arXiv:2307.10421 [gr-qc].
- [23] F. Iacovelli, E. Belgacem, M. Maggiore, M. Mancarella, and N. Muttoni, "Combining underground and on-surface third-generation gravitational-wave interferometers," *JCAP* 10 (2024) 085, arXiv:2408.14946 [gr-qc].
- [24] S. Borhanian, "GWBENCH: a novel Fisher information package for gravitational-wave benchmarking," Class. Quant. Grav. 38 (2021) 175014, arXiv:2010.15202 [gr-qc].
- [25] KAGRA, LIGO Scientific, Virgo, VIRGO Collaboration, B. P. Abbott *et al.*, "Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, Advanced Virgo and KAGRA," *Living Reviews in Relativity* 23 no. 1, (Sept., 2020) 3, arXiv:1304.0670 [gr-qc].
- [26] F. Iacovelli, M. Mancarella, S. Foffa, and M. Maggiore, "Forecasting the detection capabilities of third-generation gravitational-wave detectors using GWFAST," *Astrophys. J.* 941 no. 2, (2022) 208, arXiv:2207.02771 [gr-qc].
- [27] F. Iacovelli, M. Mancarella, S. Foffa, and M. Maggiore, "GWFAST: a Fisher information matrix Python code for third-generation gravitational-wave detectors," Astrophys. J. Suppl. 263 (2022) 2, arXiv:2207.06910 [astro-ph.IM].
- [28] I. C. F. Wong, P. T. H. Pang, M. Wils, F. Cireddu, W. Del Pozzo, and T. G. F. Li, "The Potential Impact of Noise Correlation in Next-generation Gravitational Wave Detectors," arXiv:2407.08728 [gr-qc].
- [29] K. Janssens *et al.*, "Correlated 0.01Hz-40Hz seismic and Newtonian noise and its impact on future gravitational-wave detectors," arXiv:2402.17320 [gr-qc].

- [30] E. Thrane and J. D. Romano, "Sensitivity curves for searches for gravitational-wave backgrounds," *Phys. Rev. D* 88 no. 12, (2013) 124032, arXiv:1310.5300 [astro-ph.IM].
- [31] M. Maggiore, "Gravitational wave experiments and early universe cosmology," *Phys. Rept.* 331 (2000) 283-367, arXiv:gr-qc/9909001 [gr-qc].
- [32] M. Maggiore, *Gravitational Waves. Vol. 1: Theory and Experiments*. Oxford University Press, 2007.
- [33] K. Janssens, K. Martinovic, N. Christensen, P. M. Meyers, and M. Sakellariadou, "Impact of Schumann resonances on the Einstein Telescope and projections for the magnetic coupling function," *Phys. Rev. D* 104 no. 12, (2021) 122006, arXiv:2110.14730 [gr-qc].
- [34] E. Belgacem, F. Iacovelli, M. Maggiore, M. Mancarella, and N. Muttoni, "The spectral density of astrophysical stochastic backgrounds," arXiv:2411.04028 [gr-qc].
- [35] E. Belgacem, F. Iacovelli, M. Maggiore, M. Mancarella, and N. Muttoni, "Confusion noise from astrophysical backgrounds at third-generation gravitational-wave detector networks," arXiv:2411.04029 [gr-qc].