

On Banyaga's conjecture on the group of symplectic homeomorphisms

Carole Madengko ^{*1}, Stephane Tchuiaga ^{†2}, and Franck Houenou ^{‡3}

^{1,3}Institut de Mathématiques et de Sciences Physiques, University of Abomey Calavi, Abomey Calavi, Benin

²Department of Mathematics, University of Buea, South West Region, Cameroon

Abstract

This paper addresses Banyaga's conjecture asserting that : the group of strong symplectic homeomorphisms is a proper normal subgroup of the symplectic homeomorphism group of a closed symplectic manifold.

1 Introduction

On a closed symplectic manifold (M, ω) , denote by $G_\omega(M)$ the identity component in the group $Symp(M, \omega)$ consisted of symplectic diffeomorphisms. The group $G_\omega(M)$ carries a Hofer-like topology introduced in [2] and further examined in [11]. Exploring the C^0 and Hofer-like topologies on $G_\omega(M)$, led the authors in [3, 14] to define the group $SG_\omega(M)$ of strong symplectic homeomorphisms as a C^0 -counterpart of $G_\omega(M)$. In [4] Banyaga showed that $SG_\omega(M)$ is a subgroup of $Sympeo_0(M, \omega)$ the identity component in the group of symplectic homeomorphisms, and conjectured the the following

Conjecture 1. The inclusion $SG_\omega(M) \subset Sympeo_0(M, \omega)$ is proper.

When M is simply connected, $SG_\omega(M)$ coincides with $Hameo(M, \omega)$, the hameomorphism group defined by Oh-Müller [10].

As will be detailed in Section 2 any symplectic isotopy $\Phi = \{\phi^t\}$ is generated by a pair (U, \mathcal{H}) , where $U = \{U^t\}$ is a smooth time dependent function on $M \times [0, 1]$ and $\mathcal{H} = \{\mathcal{H}^t\}$ is a smooth family of S -forms or harmonic forms. Therefore, the Hofer-like length of $\Phi = \{\phi^t\}$ is given by

$$l_{\kappa, S}^{(1, \infty)}(\Phi) = \int_0^1 (osc(U^t) + \kappa \|\mathcal{H}^t\|_{L^2}) dt, \quad (1.1)$$

where $osc(\cdot) = \max(\cdot) - \min(\cdot)$, κ is a positive real number and $\|\cdot\|_{L^2}$ is the L^2 -norm on $H^1(M, \mathbb{R})$. The length $l_{\kappa, S}^{(1, \infty)}$ induces metrics $D_{\kappa, S}^1$ and d_{HL} , on $Iso(M, \omega)$ the group of symplectic isotopies and $G_\omega(M)$ respectively.

Definition 1.1. A homeomorphism ϕ is called a finite symplectic energy homeomorphism if there exists a sequence $(\Phi_i)_i = (\{\phi_i^t\})_i$ of symplectic isotopies generated by $(U_i, \mathcal{H}_i)_i$ such that $\phi_i^1 \xrightarrow{C^0} \phi$ and $l_{\kappa, S}^{(1, \infty)}(\Phi_i)$ is bounded for all i .

Denote by $FSHomeo(M)$ the finite energy symplectic

*carolemadengko@gmail.com

†tchuiagas@gmail.com

‡rdjeam@gmail.com

Definition 1.2. [14] A continuous isotopy Φ of homeomorphisms of M is called an \mathcal{S} -topological isotopy if there exists a pair (U, \mathcal{H}) (with U a continuous time dependent function on $M \times [0, 1]$ and \mathcal{H} a continuous family of \mathcal{S} -forms or harmonic forms), and a sequence of symplectic isotopies $\{\Phi_i\}_i$ generated by $((U_i, \mathcal{H}_i))_i$ such that $\lim_{i \rightarrow \infty} D_{\kappa, \mathcal{S}}^1((U_i, \mathcal{H}_i), (U, \mathcal{H})) = 0$ and $\Phi_i \xrightarrow{C^0} \Phi$.

The group $SG_\omega(M)$ of strong symplectic homeomorphisms of M consists of time-one maps of \mathcal{S} -topological isotopies.

If M is simply connected, $FSHomeo(M)$ coincides with the finite energy hamiltonian homeomorphism group $FHomeo(M)$ defined in [6]. It has been shown in [5, 6] that on any closed simply connected surface (Σ, ω) , the inclusions

$$SG_\omega(\Sigma) \subseteq FSHomeo(\Sigma) \subseteq Sympeo_0(\Sigma, \omega), \quad (1.2)$$

are strict. To our knowledge it is not known whether the inclusions (1.2) are still strict on closed surfaces with genus $g > 0$ and higher dimensional closed symplectic manifolds.

In [12], Tchiuiga showed that function

$$\begin{aligned} \Delta(\cdot, \alpha) : G_\omega(M) &\longrightarrow \mathbb{R} \\ \phi^1 &\longmapsto \Delta(\phi^1, \alpha) := \frac{1}{\|\alpha\|_{L^2}} \int_M \left(\int_\gamma (\phi^1)^* \alpha - \alpha \right) \frac{\omega^n}{n!} \end{aligned}$$

is continuous w.r.t the C^0 -topology, hence admits a C^0 -extension say $\chi(\cdot, \alpha) : \overline{G_\omega(M)}^{C^0} \rightarrow \mathbb{R}$. Since $\overline{G_\omega(M)} = Sympeo_0(M, \omega)$ on surfaces, we use the latter extension to proof the following result.

Theorem A. On a closed symplectic surface (Σ_g, ω) of genus $g \geq 1$, $FSHomeo(\Sigma_g)$ is a proper normal subgroup of $Sympeo_0(\Sigma_g, \omega)$.

Theorem A confirms Conjecture 1 for surfaces of genus $g > 0$, and gives an affirmative answer to the following question posed by Banyaga in [4]

Question 1. Is $SG_\omega(M)$ a normal subgroup of $Sympeo_0(M, \omega)$.

The following result generalises Theorem 1.4-[5].

Theorem B. For every $E > 0$, there exists a continuous path $\{\psi^t\}$ of homeomorphism of M such that

1. the path $\{\psi^t\}$ is a uniform limit of a sequence of smooth symplectic isotopies $(\Phi_i)_i = (\{\phi_i^t\})_i$ generated by $(U_i, \mathcal{H}_i)_i$ and $l_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi_i) \leq E$
2. for every sequence $(\varphi_i)_i \subset G_\omega(M)$, satisfying $\psi^1 = \lim_{C^0} \varphi_i$, and for every $\varphi \in G_\omega(M)$, we have $\liminf_{i \rightarrow \infty} d_{HL}(\varphi_i, \varphi) \geq E$

Item 1 of Theorem B tells us that $\psi^1 \in FSHomeo(M)$. Any such ψ^1 can not be in $SG_\omega(M)$. Indeed, suppose $\psi \in SG_\omega(M)$, there exists a sequence $(U_i, \mathcal{H}_i)_i \subseteq \mathfrak{T}(M, \omega, \mathcal{S})$ generating a sequence $(\{\phi_i^t\})_i \subseteq Iso(M, \omega)$ with $(U_i, \mathcal{H}_i)_i \xrightarrow{D_{\lambda, \mathcal{S}}^1} (U', \mathcal{H}')$ and $\phi_i^1 \xrightarrow{C^0} \psi$. Fix l sufficiently large, consider the pair (U_l, \mathcal{H}_l) which generates $\phi_l^1 \in G_\omega(M)$. Then for any $\epsilon > 0$, we have $\liminf_{i \rightarrow \infty} d_{HL}(\phi_i^1, \phi_l^1) \leq \epsilon$. This contradicts item 2 of Theorem B hence $\psi^1 \notin SG_\omega(M)$.

Corollary B. $SG_\omega(M) \neq FSHomeo(M)$

Theorem B completes the proof of Cojecture 1 on any closed symplectic manifold and gives an affirmative answer to question the following question posed by Humilière and Seyfaddini:

Question 2. Are the groups $Hameo(M, \omega)$ and $FHomeo(M)$ distinct?

This paper is organised as follows: In Section 2, we review the construction of norms on $G_\omega(M)$ namely, the Tchiuiga norm $\|\cdot\|^\infty$, and the Hofer-like norm. With the help of the C^0 and Hofer-like topologies, we construct a Hofer-like norm on $FSHomeo(M)$. Section 3 is devoted to prove our results.

2 Preliminaries

Let (M, ω) , be a closed symplectic surface with genus $g \geq 1$, and $\mathcal{Z}^1(M)$, the space of closed 1-forms on M . The group $G_\omega(M)$ coincides with the time one map of the group of symplectic isotopies $Iso(M, \omega)$. For all $\alpha \in \mathcal{Z}^1(M) \setminus \{0\}$ and for any $\Phi = \{\phi^t\} \in Iso(M, \omega)$, we have

$$(\phi^t)^* \alpha - \alpha = d\mathcal{F}_\Phi^\alpha(t), \quad \forall t, \quad (2.1)$$

where $\mathcal{F}_\Phi^\alpha(t) = \int_0^t (\iota_{\dot{\phi}^s} \alpha) \circ \phi^s ds$. Fix $x \in M$, for any $y \in M$, take a curve γ from x to y . Integrating (2.1) over γ yields

$$\int_\gamma (\phi^t)^* \alpha - \alpha = \mathcal{F}_\Phi^\alpha(t)(y) - \mathcal{F}_\Phi^\alpha(t)(x), \quad \forall t$$

Recently, it has been shown in [12] that

$$\Delta(\phi^1, \alpha)_x := \frac{1}{\|\alpha\|_{L^2}} \left\langle [\alpha \wedge \omega^{n-1}], \tilde{S}_\omega(\Phi) \right\rangle - \frac{Vol(M)}{\|\alpha\|_{L^2}} \mathcal{F}_\Phi^\alpha(1)(x), \quad (2.2)$$

where $\tilde{S}_\omega : \widetilde{G_\omega(M)} \rightarrow H^1(\Sigma, \mathbb{R})$ is the flux homomorphism see [1] for more details. From Proposition 3.8–[12], if M has boundary, equation (2.2) has an additional term. In [12], Tchiuaga defined a norm on $G_\omega(M)$ as follows:

$$\|\phi\|^\infty = \sup_{\alpha \in \mathcal{B}(1)} \left(\sup_{x \in M} |\tilde{\Delta}(\phi, \alpha)_x| \right),$$

for any $\phi \in G_\omega(M)$, where $\tilde{\Delta}(\phi, \alpha)_x = \|\alpha\|_{L^2} \Delta(\phi, \alpha)_x$ and $\mathcal{B}(1) = \{\alpha \in \mathcal{Z}^1(\Sigma_g) : \|\alpha\|_{L^2} = 1\}$.

The norm $\|\cdot\|^\infty$ extends to the C^0 - setting: For any $\phi \in Sympeo(\Sigma_g, \omega)$,

$$\|\phi\|^\infty = \sup_{\alpha \in \mathcal{B}(1)} \left(\sup_{x \in M} |\tilde{\chi}(\phi, \alpha)_x| \right), \quad (2.3)$$

where $\tilde{\chi}(\phi, \alpha)_x = \|\alpha\|_{L^2} \chi(\phi, \alpha)_x$ and $\mathcal{B}(1) = \{\alpha \in \mathcal{Z}^1(\Sigma_g) : \|\alpha\|_{L^2} = 1\}$.

2.1 Splitting of closed 1-forms

Let $B^1(M)$, be the space of exact 1-forms and $H^1(M, \mathbb{R})$ the first de Rham cohomology group. Consider a linear section $\mathcal{S} : H^1(M, \mathbb{R}) \rightarrow \mathcal{Z}^1(M)$ of the natural projection $\pi : \mathcal{Z}^1(M) \rightarrow H^1(M, \mathbb{R})$. Any $\alpha \in \mathcal{Z}^1(M)$ splits as:

$$\alpha = \mathcal{S}(\pi(\alpha)) + (\alpha - \mathcal{S}(\pi(\alpha))) \quad (2.4)$$

where $\mathcal{S}(\pi(\alpha))$ is an \mathcal{S} -form and $\alpha - \mathcal{S}(\pi(\alpha))$ is an exact form. Let $\mathbb{H}_\mathcal{S}(M)$ denote the space of all \mathcal{S} -forms. Note that if the linear section \mathcal{S} is trivial then $\mathcal{S}(\pi(\alpha))$ is a harmonic 1-form and equation 2.4 is the Hodge decomposition of closed 1-forms. For more details see [3, 11]. To any symplectic isotopy $\Phi = \{\phi^t\}$ corresponds a unique pair (U, \mathcal{H}) through the following isomorphisms:

$$\begin{array}{ccccccc} Iso(M, \omega) & \longrightarrow & \mathcal{Z}^1(M) & \xrightarrow{\text{Splitting}} & \mathcal{Z}^1(M) & \longrightarrow & \mathfrak{T}(M, \omega, \mathcal{S}) \\ \Phi = \{\phi^t\}_t & \longmapsto & \iota_{\dot{\phi}^t} \omega & \longmapsto & dU^t + \mathcal{H}^t & \longmapsto & (U, \mathcal{H}), \end{array}$$

where $U = \{U^t\}_t \in C_0^\infty(M \times [0, 1], \mathbb{R})$ (the set of zero mean smooth functions on $M \times [0, 1]$), $\mathcal{H} = \{\mathcal{H}^t\}_t$ is a smooth path in $\mathbb{H}_\mathcal{S}(M)$ and $\mathfrak{T}(M, \omega, \mathcal{S})$ is the set of all generators (U, \mathcal{H}) of symplectic isotopies. In the sequel, we write $\phi_{(U, \mathcal{H})}$ to mean the isotopy Φ is generated by (U, \mathcal{H}) . The set $\mathfrak{T}(M, \omega, \mathcal{S})$ forms a group w.r.t the following rule:

$$(U, \mathcal{H}) \rtimes_{\mathcal{S}} (V, \mathcal{K}) = \left(U + V \circ \phi_{(U, \mathcal{H})}^{-1} + \tilde{\mathcal{F}}_{\phi_{(U, \mathcal{H})}^{-1}}^{\mathcal{K}}(t), \quad \mathcal{H} + \mathcal{K} \right), \quad (2.5)$$

the inverse of (U, \mathcal{H}) , denoted $\overline{(U, \mathcal{H})}$, is given by

$$\overline{(U, \mathcal{H})} = \left(-U \circ \phi_{(U, \mathcal{H})} - \tilde{\mathcal{F}}_{\phi_{(U, \mathcal{H})}}^{\mathcal{H}}(t), \quad -\mathcal{H} \right) \quad (2.6)$$

where $\tilde{\mathcal{F}}_{\phi_{(U, \mathcal{H})}}^{\mathcal{H}}(t) = \int_0^t \mathcal{H}^t(\psi_{(U, \mathcal{H})}^s) \circ \psi_{(U, \mathcal{H})}^s ds - \frac{\int_M \left(\int_0^t \mathcal{H}^t(\psi_{(U, \mathcal{H})}^s) \circ \psi_{(U, \mathcal{H})}^s ds \right) \omega}{\int_M \omega}$ for all $t \in [0, 1]$.

Given a symplectic isotopy $\Phi = \{\phi^t\}$ generated by (U, \mathcal{H}) , the $L^{(1, \infty)}$ -version of the Hofer-like length of Φ is given by

$$l_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi) = \int_0^1 (\nu^B(dU^t) + \kappa \|\mathcal{H}^t\|_{L^2}) dt, \quad (2.7)$$

where ν^B is any norm on $B^1(M)$, the coefficient κ is a positive real number. Here, we choose ν^B to be the oscillation norm for more details see [11]. The above Hofer-like length gives rise to the Hofer-like energy e_{HL} , which assigns to any $\psi \in G_\omega(M)$:

$$e_{HL}(\psi) = \inf l_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi)$$

where the infimum is taken over all symplectic isotopies whose time one map is ψ . The $L^{(1, \infty)}$ -Hofer-like norm of ψ is given by

$$\|\psi\|_{HL} = \frac{e_{HL}(\psi) + e_{HL}(\psi^{-1})}{2},$$

the latter norm induces a right invariant distance d_{HL} on $G_\omega(M)$ given by

$$d_{HL}(\phi, \psi) = \|\phi \circ \psi^{-1}\|_{HL} \quad \forall \phi, \psi \in G_\omega(M).$$

The length $l_{\kappa, \mathcal{S}}^{(1, \infty)}$ also induces a metric $D_{\kappa, \mathcal{S}}^1$ on the $Iso(M, \omega)$ given by:

$$D_{\kappa, \mathcal{S}}^1((U, \mathcal{H}), (V, \mathcal{K})) := \frac{D_0^{\kappa, \mathcal{S}}((U, \mathcal{H}), (V, \mathcal{K})) + D_0^{\kappa, \mathcal{S}}(\overline{(U, \mathcal{H})}, \overline{(V, \mathcal{K})})}{2}, \quad \forall \kappa > 0,$$

where

$$D_0^{\kappa, \mathcal{S}}((U, \mathcal{H}), (V, \mathcal{K})) := \int_0^1 (\text{osc}(U_t - V_t) + \kappa \|\mathcal{H}_t - \mathcal{K}_t\|_{L^2}) dt,$$

See [11] for more details on the Hofer-like metrics. In this paper, we will assume $\kappa = 1$.

2.2 From smooth to C^0

C^0 -symplectic topology began with Eliashberg-Gromov rigidity theorem [8, 9] which states that $Symp(M, \omega)$ is C^0 -closed in $Diff(M)$ the group of diffeomorphisms of M . This inspired Oh-Müller [10] to define $Sympeo(M, \omega)$, the symplectic homeomorphism group as

$$\overline{Symp(M, \omega)}^{C^0} \subseteq Homeo(M),$$

where $Homeo(M)$ is the group of homeomorphisms of M . The C^0 -topology on $Homeo(M)$ is the topology induced by the distance:

$$d_0(f, h) = \max(d_{C^0}(f, h), d_{C^0}(f^{-1}, h^{-1})), \quad \forall f, h \in Homeo(M)$$

with $d_{C^0}(f, h) = \sup_{x \in M} d_g(h(x), f(x))$.

On the space of all continuous paths $\lambda : [0, 1] \rightarrow Homeo(M)$ such that $\lambda(0) = id_M$, the C^0 -topology is the topology induced by the distance:

$$\bar{d}(\lambda, \mu) = \max_{t \in [0, 1]} d_0(\lambda(t), \mu(t)).$$

The distance \bar{d} is neither left nor right invariant however, it satisfies the following:

Proposition 2.1. [13] *Let Θ be a smooth isotopy of diffeomorphisms of M . Then there exists a constant $C^\Theta > 0$ depending on Θ such that for all paths Φ, Ψ in $\text{Homeo}(M)$, we have*

$$\bar{d}(\Phi \circ \Theta, \Psi \circ \Theta) \leq C^\Theta \bar{d}(\Phi, \Psi).$$

Given $\phi \in \text{FSHomeo}(M)$, define

$$\|\phi\|_{\widetilde{HL}} := \liminf_{i \rightarrow \infty} \|\phi_i\|_{HL}, \quad (2.8)$$

where the infimum is taken over all $(\phi_i)_i \subset G_\omega(M)$ such that $\phi_i \xrightarrow{C^0} \phi$. The rule $\|\cdot\|_{\widetilde{HL}}$ defines a Hofer-like norm on $\text{FSHomeo}(M)$ indeed:

1. Positivity: clearly $\|\phi\|_{\widetilde{HL}} \geq 0$, for any $\phi \in \text{FSHomeo}(M)$
2. Non-degeneracy: Let $\phi \in \text{FSHomeo}(M)$ and $(\phi_i)_i \subset G_\omega(M)$ such that $\phi_i \xrightarrow{C^0} \phi$. Assume

$$0 = \|\phi\|_{\widetilde{HL}} = \liminf_{i \rightarrow \infty} \|\phi_i\|_{HL}.$$

By definition of \liminf , we have for any $\epsilon > 0$, there exists $\delta > 0$, such that for all $(\phi_i)_i \subseteq G_\omega(M)$, satisfying $d_0(\phi_i, \phi) < \delta$, we have $\|\phi_i\|_{HL} - \epsilon < \|\phi\|_{\widetilde{HL}}$. By assumption, $\|\phi\|_{\widetilde{HL}} = 0$, hence $\|\phi_i\|_{HL} - \epsilon < 0$, i.e, $0 \leq \|\phi_i\|_{HL} < \epsilon$. Letting ϵ tend to zero, we get that $\|\phi_i\|_{HL} \rightarrow 0$, as $i \rightarrow \infty$. Therefore, $\phi_i \xrightarrow{C^0} \phi$, and $\|\phi_i\|_{HL} \rightarrow 0$, as $i \rightarrow \infty$ hence by Theorem 3.3–[11], we have that $\phi = id$.

3. Triangle inequality: let $\phi, \psi \in \text{FSHomeo}(M)$. By definition, $\phi = \lim_{C^0} \phi_i$, and $\psi = \lim_{C^0} \psi_i$, where $(\phi_i)_i$ and $(\psi_i)_i$ are sequences in $G_\omega(M)$ whose Hofer-like norms are bounded. By triangle inequality of $\|\cdot\|_{HL}$, one has

$$\|\phi_i \circ \psi_i\|_{HL} \leq \|\phi_i\|_{HL} + \|\psi_i\|_{HL}. \quad (2.9)$$

Taking the infimum over all $\phi_i \circ \psi_i$ which C^0 converges to $\phi \circ \psi$, on both sides of (2.9) we have

$$\liminf_{i \rightarrow \infty} \|\phi_i \circ \psi_i\|_{HL} \leq \liminf_{i \rightarrow \infty} (\|\phi_i\|_{HL} + \|\psi_i\|_{HL}) \leq \liminf_{i \rightarrow \infty} \|\phi_i\|_{HL} + \liminf_{i \rightarrow \infty} \|\psi_i\|_{HL}$$

The norm $\|\cdot\|_{\widetilde{HL}}$ induces a right invariant distance on $\text{FSHomeo}(M)$ defined as:

$$\tilde{d}_{HL}(\psi, \phi) = \liminf_{i \rightarrow \infty} \|\phi \circ \psi^{-1}\|_{HL}$$

3 Proof of Results

3.1 Proof of Theorem A

The proof Theorem A consists of constructing a homeomorphism $\phi \in \overline{G_\omega(\Sigma_g)}$, such that $\|\phi\|^\infty = \infty$. Then any such ϕ can not be in $\text{FSHomeo}(\Sigma_g)$.

Lemma 3.1. *For any $\phi \in \text{FSHomeo}(\Sigma_g)$, we have that $\|\phi\|^\infty$ is bounded.*

Proof. The proof of Lemma 3.1 follows from the proof of Theorem 6.5–[12] □

Lemma 3.2. *There exists a sequence $(\phi_i)_i \subset G_\omega(\Sigma_g)$, such that $\phi_i \xrightarrow{C^0} \phi$ and $\|\phi_i\|^\infty \rightarrow \infty$, as $i \rightarrow \infty$*

Proof. We proceed by induction on g

Step 1: For $g = 1$, consider the 2 torus $\mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1$ with coordinate system (θ_1, θ_2) and consider the standard symplectic form $\omega = d\theta_1 \wedge d\theta_2$ where $\theta_1, \theta_2 \in [0, 2\pi]$. Define a flow

$$\phi^t(\theta_1, \theta_2) := (\theta_1 + t h(\theta_2), \theta_2), \quad \forall t,$$

with

$$\begin{aligned} h : (0, 2\pi] &\longrightarrow \mathbb{R} \\ \theta_2 &\longmapsto h(\theta_2) := \frac{1}{\theta_2} \end{aligned}$$

The isotopy $\{\phi^t\}_t$ is a non-hamiltonian symplectic isotopy of $\mathbb{T}^2 \setminus \{(\theta_1, 0)\}$. Indeed, equip \mathbb{T}^2 with the flat riemannian metric g_0 , then the 1-forms $d\theta_1$ and $d\theta_2$ are harmonic. By Hodge decomposition, $\iota_{\phi^t} \omega = h d\theta_2 = d(\ln(\theta_2) - \theta_2) + d\theta_2$. Therefore the flow $\{\phi^t\}_t$ is generated by $(U, \mathcal{H}) = (\ln(\theta_2) - \theta_2, d\theta_2)$ and $\tilde{S}_\omega(\{\phi^t\}_t) = [d\theta_2] \neq 0$.

Set $P = (\theta_1, 0)$, the isotopy $\{\phi^t\}_t$ extends to a non-hamiltonian symplectic flow $\{\psi^t\}_t$ on \mathbb{T}^2 by fixing P . The flow $\{\psi^t\}_t$ is differentiable everywhere except at P .

For all $i \in \mathbb{N}^*$, consider the smooth function $h_i : [0, 2\pi] \rightarrow \mathbb{R}$ such that $h_i = h$, on $[\frac{1}{i}, 2\pi]$. Define the map $\phi_i(\theta_1, \theta_2) := (\theta_1 + h_i(\theta_2), \theta_2)$, then $\phi_i \circ (\psi^1)^{-1} \xrightarrow{C^0} id$, as $i \rightarrow \infty$, hence $\psi^1 \in \text{Sympeo}_0(\mathbb{T}^2, \omega)$. Let $\alpha = d\theta_1$, then $(\phi_i)^* \alpha - \alpha = dh_i$. Pick $x \in \mathbb{T}^2$ (fixed), and take a C^1 -curve γ from x to any point $y \in \mathbb{T}^2$.

$$\begin{aligned} \int_{\mathbb{T}^2} \left(\int_{\gamma} (\phi_i)^* \alpha - \alpha \right) \omega &= \int_{\mathbb{T}^2} \left(\int_{\gamma} dh_i \right) \omega \\ &= \int_{\mathbb{T}^2} (h_i(y) - h_i(x)) \omega \end{aligned} \quad (3.1)$$

Set A , to be the area of \mathbb{T}^2 . Since x is fixed, (3.1) becomes

$$\begin{aligned} \int_{\mathbb{T}^2} \left(\int_{\gamma} (\phi_i)^* \alpha - \alpha \right) \omega &= \int_{\mathbb{T}^2} (h_i(y) \omega - A h_i(x)) \\ &\geq \int_{[0, 2\pi] \times [1/i, 2\pi]} h_i(y) \omega - A h_i(x). \end{aligned} \quad (3.2)$$

Taking the limit as i tend to ∞ in equation (3.2), we have $\Delta(\phi_i, \alpha) \rightarrow \infty$ as $i \rightarrow \infty$ hence $\|\phi_i\|^\infty \rightarrow \infty$ as $i \rightarrow \infty$, i.e., $\|\psi^1\|^\infty = \infty$.

Step 2: For $g = 2$, consider 2, tori $\mathbb{T}_1^2, \mathbb{T}_2^2$ and flows $\{\phi_1^t\}_t$ and $\{\phi_2^t\}_t$ defined on each torus as in Step 1. Remove discs \mathbb{D}_1 and \mathbb{D}_2 of radius $R < 1$ from \mathbb{T}_1^2 and \mathbb{T}_2^2 respectively. A 2- genus surface Σ_2 is obtained by gluing $\mathbb{T}_1^2 \setminus \mathbb{D}_1$, and $\mathbb{T}_2^2 \setminus \mathbb{D}_2$ along the boundaries of \mathbb{D}_1 and \mathbb{D}_2 . Define $\varphi \in \text{Sympeo}_0(\Sigma_2, \omega)$ as follows:

$$\varphi = \begin{cases} \phi_1^1 & \text{on } \mathbb{T}_1^2 \setminus \mathbb{D}_1 \\ \phi_2^1 & \text{on } \mathbb{T}_2^2 \setminus \mathbb{D}_2 \end{cases}, \quad (3.3)$$

such that $\phi_j^1 = id$ on $\partial \mathbb{D}_j$, $j = 1, 2$. Consider a smooth symplectic map φ_i defined as follows

$$\varphi_i = \begin{cases} \phi_{1,i}^1 & \text{on } \mathbb{T}_1^2 \setminus \mathbb{D}_1 \\ \phi_{2,i}^1 & \text{on } \mathbb{T}_2^2 \setminus \mathbb{D}_2 \end{cases}, \quad (3.4)$$

where $\phi_{j,i}^1$, $j = 1, 2$ coincide with the map smooth ϕ_i constructed in Step 1 and $\phi_{j,i}^1 = id$ on $\partial \mathbb{D}_j$. Then $\varphi_i \xrightarrow{C^0} \varphi$, as $i \rightarrow \infty$.

For each $j \in \{1, 2\}$ set $\alpha_j = d\theta_j^1$. Consider the projection, $Pr_j : \Sigma_2 \rightarrow \mathbb{T}_j^2 \setminus \mathbb{D}_j$. Let $\alpha \in Z^1(\Sigma_2)$ be given by $\alpha = Pr_1^*(\alpha_1) + Pr_2^*(\alpha_2)$. Let γ a C^1 -curve from x (fixed) to y , in Σ_2 , then

$$\tilde{\Delta}(\varphi_i, \alpha) = \begin{cases} \tilde{\Delta}(\phi_{1,i}^1, \alpha) & \text{on } \mathbb{T}_1^2 \setminus \mathbb{D}_1 \\ \tilde{\Delta}(\phi_{2,i}^1, \alpha) & \text{on } \mathbb{T}_2^2 \setminus \mathbb{D}_2 \end{cases}, \quad (3.5)$$

where

$$\tilde{\Delta}(\phi_{j,i}^1, \alpha) = \langle \tilde{S}_\omega(\phi_{j,i}^t), [\alpha] \rangle + \int_{\partial \mathbb{D}_j} \left(\int_0^1 \mathcal{F}_{\phi_{j,i}^t}^\alpha(t) (\phi_{j,i}^t)^*(\iota_{\phi_{j,i}^t} \omega) dt \right) - Area(\mathbb{T}_j^2 \setminus \mathbb{D}_j) \mathcal{F}_{\phi_{j,i}^t}^\alpha(1)(x). \quad (3.6)$$

See Proposition 3.8–[?] for details. Since for each j , $\phi_{j,i} = id$ on $\partial \mathbb{D}_j$, then

$$\int_{\partial \mathbb{D}_j} \left(\int_0^1 \mathcal{F}_{\phi_{j,i}^t}^\alpha(t) (\phi_{j,i}^t)^*(\iota_{\phi_{j,i}^t} \omega) dt \right) = 0$$

Using (3.1) and (3.2), one can show that the right hand side of equation (3.6) goes to ∞ , as $i \rightarrow \infty$. Therefore, taking the limit as i tends to ∞ in (3.5) yields $\tilde{\Delta}(\varphi_i, \alpha) \rightarrow \infty$ as $i \rightarrow \infty$ and $\|\varphi\|^\infty = \infty$

Step 3 Assume that on a surface Σ_{g-1} of genus $g-1$, there exists a sequence $(\psi_i)_i \subset G_\omega(\Sigma_{g-1})$, such that $\psi_i \xrightarrow{C^0} \psi$ and $\|\psi_i\|^\infty \rightarrow \infty$, as $i \rightarrow \infty$. Remove a disc \mathbb{D}_{g-1} of radius $R < 1$ from Σ_{g-1} and consider the punctured torus $\mathbb{T}^2 \setminus \mathbb{D}$ with \mathbb{D} a disc of radius $R < 1$. A g -genus surface is formed by gluing $\Sigma_{g-1} \setminus \mathbb{D}_{g-1}$ and $\mathbb{T}^2 \setminus \mathbb{D}$ along the boundaries of \mathbb{D}_{g-1} and \mathbb{D} . Define $\vartheta \in \text{Sympeo}_0(\Sigma_g, \omega)$ as follows:

$$\vartheta = \begin{cases} \phi & \text{on } \mathbb{T}^2 \setminus \mathbb{D}_1 \\ \psi & \text{on } \Sigma_{g-1} \setminus \mathbb{D}_{g-1} \end{cases}, \quad (3.7)$$

where ϕ is constructed as in Step 2 and $\psi = id$ on $\partial \Sigma_{g-1} \setminus \mathbb{D}_{g-1}$. Following similar arguments from Step 2, one constructs a sequence $(\vartheta_i)_i$ of symplectic diffeomorphisms and a closed 1-form α such that $\vartheta_i \xrightarrow{C^0} \vartheta$ and $\tilde{\Delta}(\vartheta_i, \alpha)$ tends to infinity as $i \rightarrow \infty$ that is, $\|\vartheta_i\|^\infty \rightarrow \infty$, as $i \rightarrow \infty$. Therefore, by mathematical induction, on any closed symplectic surface Σ_g , of genus $g \geq 1$, there is a sequence $(\phi_i)_i \subset G_\omega(\Sigma_g)$, such that $\phi_i \xrightarrow{C^0} \phi$ and $\|\phi_i\|^\infty \rightarrow \infty$, as $i \rightarrow \infty$. \square

In order to complete the proof Theorem A, we now show that $FSH\text{Homeo}(M)$ is a normal subgroup of $\text{Sympeo}_0(M)$, for all closed symplectic manifolds (M, ω) .

Lemma 3.3. *For any closed symplectic surface (M, ω) , $FSH\text{Homeo}(M)$ is a normal subgroup of $\text{Sympeo}_0(M)$.*

Proof. Let $\varphi \in FSH\text{Homeo}(M)$ and $\psi \in \text{Sympeo}_0(M)$. By definition, $\psi = \lim_{C^0} \psi_i$ with $(\psi_i)_i \subseteq G_\omega(M)$. Consider a smooth sequence $(\psi_{(V_i, \mathcal{K}_i)})_i \subseteq \text{Iso}(M, \omega)$ whose sequence of time one maps $(\psi_i)_i$ converge to φ in the C^0 -topology and the sequence $(V_i, \mathcal{K}_i)_i$ is bounded w.r.t $D_{\kappa, \mathcal{S}}^1$. The sequence $(V_i \circ \psi_i + \tilde{\mathcal{F}}_{\psi_{(V_i, \mathcal{K}_i)}}^{\mathcal{K}_i}(t))$, $\mathcal{K}_i \subset \mathfrak{T}(\Sigma_g, \omega, \mathcal{S})$ generates the sequence of isotopies $(\psi_i^{-1} \circ \psi_{(V_i, \mathcal{K}_i)} \circ \psi_i)_i$ whose time one maps C^0 -converges to $\psi^{-1} \circ \varphi \circ \psi$ and

$$\int_0^1 \left(\text{osc} \left[V_i^t \circ \psi_i + \tilde{\mathcal{F}}_{\psi_{(V_i, \mathcal{K}_i)}}^{\mathcal{K}_i}(t) \right] + \|\mathcal{K}_i^t\|_{L^2} \right) dt \leq \int_0^1 \text{osc}(V_i^t \circ \psi_i) dt + \int_0^1 \text{osc} \left(\tilde{\mathcal{F}}_{\psi_{(V_i, \mathcal{K}_i)}}^{\mathcal{K}_i}(t) \right) dt + \int_0^1 \|\mathcal{K}_i^t\|_{L^2} dt$$

Combining equations (2.10) and (2.25) in [11], we get that

$$\int_0^1 \left(\text{osc} \left[V_i^t \circ \psi_i + \tilde{\mathcal{F}}_{\Psi_i}^{\mathcal{K}_i}(t) \right] + \|\mathcal{K}_i^t\|_{L^2} \right) dt \leq \int_0^1 \left(\text{osc}(V_i^t) dt + BK(g_0) \|\mathcal{K}_i^t\|_{L^2} \right) dt + \int_0^1 \|\mathcal{K}_i^t\|_{L^2} dt$$

for some positive constants B and $K(g_0)$ which depend on a riemannian metric g_0 on M . Therefore,

$$\begin{aligned} \int_0^1 \left(\text{osc} \left[V_i^t \circ \psi_i + \tilde{\mathcal{F}}_{\psi(V_i, \mathcal{K}_i)}^{\mathcal{K}_i}(t) \right] + \|\mathcal{K}_i^t\|_{L^2} \right) dt &\leq \int_0^1 \text{osc}(V_i^t) dt + (BK(g_0) + 1) \int_0^1 \|\mathcal{K}_i^t\|_{L^2} dt \\ &\leq 2(BK(g_0) + 1) \int_0^1 (\text{osc}(V_i^t) + \|\mathcal{K}_i^t\|_{L^2}) dt. \end{aligned} \quad (3.8)$$

The right hand side of inequality (3.8) is bounded since (V_i, \mathcal{K}_i) is bounded w.r.t $D_{\kappa, \mathcal{S}}^1$. \square

3.2 Proof of Theorem B

We need the following Lemma in order to prove Theorem B

Lemma 3.4. *Let $\Phi = \{\phi^t\}$ be a symplectic isotopy. For every $E > 0$, such that $l_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi) < E$, there exist a symplectic isotopy $\Phi' = \{(\phi')^t\}$ such that*

1. $\bar{d}(\Phi, \Phi')$ can be made arbitrarily small
2. $l_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi') < E + \epsilon$, for some arbitrarily positive ϵ
3. for any $\alpha \in \mathcal{Z}^1(M) \setminus \{0\}$, and any symplectic isotopy $\Psi = \{\psi^t\}$ there exists a constant $C > 0$ such that $\Delta((\phi')^1 \circ (\psi^1)^{-1}, \alpha) > CE - \epsilon$.

Proof. Assume $\text{Vol}(M) = 1$, let $\alpha \in \mathcal{Z}^1(M) \setminus \{0\}$ and $\Phi = \{\phi^t\}$ a symplectic isotopy generated by (U, \mathcal{H}) and $l_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi) < E$. By Theorem 6.5–[12], we have $\|\phi\|^\infty \leq Cl_{\kappa, \mathcal{S}}^{(1, \infty)}(\Phi)$ with $C = 3 \max\{C(\omega), \eta, 2\}$. The Hofer norm of the function $\mathcal{F}_\Phi^\alpha(1)$ is given by

$$\begin{aligned} \text{osc}(\mathcal{F}_\Phi^\alpha(1)) &\leq 2 \sup_{x \in \Sigma_g} \left| \int_0^1 (\iota_{\phi^t} \alpha)(x) dt \right| \\ &\leq 2 \sup_x \left| \int_0^1 (\iota_{\phi^t} \omega)(X_\alpha)(x) dt \right| \end{aligned} \quad (3.9)$$

where X_α is the symplectic vector field defined by $\iota_{X_\alpha} \omega = \alpha$. From the splitting in (2.4), inequality (3.9) becomes

$$\begin{aligned} \text{osc}(\mathcal{F}_\Phi^\alpha(1)) &\leq 2 \sup_x \left| \int_0^1 (dU^s + \mathcal{H}^t)(X_\alpha)(x) dt \right| \\ &\leq 2 \left(\sup_x \left| \int_0^1 (dU^t)(X_\alpha)(x) dt \right| + \sup_x \left| \int_0^1 (\mathcal{H}^t)(X_\alpha)(x) dt \right| \right) \end{aligned} \quad (3.10)$$

From the lines of proof of Proposition 6.1–[12], we have

$$\sup_x \left| \int_0^1 (dU^t)(X_\alpha)(x) dt \right| \leq 2 \max_t (\text{osc}(U^t)) \quad (3.11)$$

By Proposition 6.3–[12], for all $\alpha \in \mathcal{Z}^1(M)$, and a harmonic vector field X , one can find a positive constant $C(\alpha)$ such that

$$\sup_{x \in M} |\alpha(X)(x)| \leq C(\alpha) \|\iota_X \omega\|_{L^2}. \quad (3.12)$$

Therefore,

$$\begin{aligned}
\sup_x \left| \int_0^1 (\mathcal{H}^t)(X_\alpha)(x) dt \right| &= \sup_x \left| \int_0^1 \iota_{X_{\mathcal{H}^t} \omega}(X_\alpha)(x) dt \right| \\
&= \sup_x \left| \int_0^1 (\iota_{X_\alpha} \omega)(X_{\mathcal{H}^t})(x) dt \right| \\
&\leq \sup_{x,t} |(\iota_{X_\alpha} \omega)(X_{\mathcal{H}^t})(x)| \\
&= C(\alpha) \sup_t \|\mathcal{H}^t\|_{L^2}
\end{aligned} \tag{3.13}$$

Putting (3.11) and (3.13) in (3.10), we have that

$$\text{osc}(\mathcal{F}_\Phi^\alpha(1)) \leq 2 \max\{2, C(\alpha)\} l_S^{(1, \infty)}(\Phi) < 2 \max\{2, \eta\} E \leq CE,$$

where η is as in Lemma 6.4 item 2. Let $\epsilon > 0$, be arbitrary and D_ϵ a disc of radius ϵ in M , set $a(t) = \sup_{x \in D_\epsilon} \mathcal{F}_\Phi^\alpha(t)(x)$ and $b(t) = \inf_{x \in D_\epsilon} \mathcal{F}_\Phi^\alpha(t)(x)$ for each $t \in [0, 1]$. Consider a function $h : M \rightarrow \mathbb{R}$ compactly supported in D_ϵ such that $h(x) \leq 1$ for all $x \in D_\epsilon$ and $\text{osc}(h) \leq \frac{\epsilon}{CE + \epsilon}$. Let $\phi_{(K_1, 0)}$ and $\phi_{(K_2, 0)}$ be hamiltonian isotopies generated by the functions $K_1(x, t) = (a(t) + \frac{\epsilon - 3CE}{2}) h(x)$ and $K_2(x, t) = (b(t) - \frac{\epsilon - 3CE}{2}) h(x)$ respectively. The symplectic isotopies $\{\phi_1^t\} = \Phi_1 =: \phi_{(K_1, 0)} \circ \Phi$ and $\{\phi_2^t\} = \Phi_2 =: \phi_{(K_2, 0)} \circ \Phi$ are good candidates for Φ' . Indeed, for any symplectic isotopy $\Psi = \{\psi^t\}_t$ and $\alpha \in \mathcal{Z}^1(M)$, we have

$$\Delta(\phi_1^1 \circ (\psi^1)^{-1}, \alpha)_y = \Delta(\phi_2^1 \circ (\psi^1)^{-1}, \alpha)_y + \Delta(\phi_1^1 \circ (\phi_2^1)^{-1}, \alpha)_{\phi_2^1 \circ (\psi^1)^{-1}(y)}, \tag{3.14}$$

for some $y \in M$. But

$$\begin{aligned}
\Delta(\phi_1^1 \circ (\phi_2^1)^{-1}, \alpha)_{\phi_2^1 \circ (\psi^1)^{-1}(y)} &= \frac{1}{\|\alpha\|_{L^2}} \left\langle [\alpha], \tilde{S}_\omega(\Phi_1 \circ \Phi_2^{-1}) \right\rangle - \frac{\text{Vol}(M)}{\|\alpha\|_{L^2}} \mathcal{F}_{\Phi_1 \circ \Phi_2^{-1}}^\alpha(1)(\phi_2^1 \circ (\psi^1)^{-1}(y)) \\
&= -\frac{1}{\|\alpha\|_{L^2}} \mathcal{F}_{\phi_{K_1} \circ \phi_{K_2}^{-1}}^\alpha(1)(\phi_2^1 \circ (\psi^1)^{-1}(y)) \\
&= -\frac{1}{\|\alpha\|_{L^2}} \int_0^1 \alpha(\overbrace{\phi_{K_1}^t \circ (\phi_{K_2}^t)^{-1}}) \circ \phi_{K_1}^t \circ (\phi_{K_2}^t)^{-1}(y) dt \\
&= \frac{1}{\|\alpha\|_{L^2}} \int_0^1 \left(\iota_{\overbrace{\phi_{K_1}^t \circ (\phi_{K_2}^t)^{-1}} \omega} \right) (X_\alpha) \circ \phi_{K_1}^t \circ (\phi_{K_2}^t)^{-1}(y) dt \\
&= \frac{1}{\|\alpha\|_{L^2}} \int_0^1 d(K_1 \# \bar{K}_2)^t(X_\alpha(z)) dt \\
&= \frac{1}{\|\alpha\|_{L^2}} \int_0^1 \frac{d}{ds} ((K_1 \# \bar{K}_2)^t(\phi_\alpha^s(z))) dt
\end{aligned} \tag{3.15}$$

Normalizing α such that $\|\alpha\|_{L^2} = 1$ and integrating (3.15) with respect to s , we have

$$\begin{aligned}
\Delta(\phi_1^1 \circ (\phi_2^1)^{-1}, \alpha)_{\phi_2^1 \circ (\psi^1)^{-1}(y)} &= \int_0^1 \int_0^1 \frac{d}{ds} (K_1 \# \bar{K}_2)^t(\phi_\alpha^s(z)) ds dt \\
&= \int_0^1 [(K_1 \# \bar{K}_2)^t(\phi_\alpha^1(z)) - (K_1 \# \bar{K}_2)^t(z)] dt
\end{aligned} \tag{3.16}$$

Since the function $K_1 \# \bar{K}_2$ is compactly supported in D_ϵ , we have

$$(K_1 \# \bar{K}_2)^t(\phi_\alpha^1(z)) - (K_1 \# \bar{K}_2)^t(z) = \begin{cases} (K_1 \# \bar{K}_2)^t(p), & \text{if } z \notin D_\epsilon \text{ and } \phi_\alpha^1(z) = p \in D_\epsilon, \\ 0, & \text{if } z \notin D_\epsilon \text{ and } \phi_\alpha^1(z) = p \notin D_\epsilon, \\ -(K_1 \# \bar{K}_2)^t(z), & \text{if } z \in D_\epsilon \text{ and } \phi_\alpha^1(z) = p \notin D_\epsilon, \\ (K_1 \# \bar{K}_2)^t(p) - (K_1 \# \bar{K}_2)^t(z), & \text{if } z \in D_\epsilon \text{ and } \phi_\alpha^1(z) = p \in D_\epsilon. \end{cases}$$

For each $z \in D_\epsilon$, we have

$$\begin{aligned}
\int_0^1 (K_1 \# \bar{K}_2)_t(z) dt &= \int_0^1 [K_1^t - K_2^t \circ \phi_{K_2}^t \circ (\phi_{K_1}^t)^{-1}](z) dt \\
&= \int_0^1 \left(a(t) + \frac{\epsilon - 3CE}{2} \right) h(z) dt \\
&\quad - \int_0^1 \left(b(t) - \frac{\epsilon - 3CE}{2} \right) h(\phi_{K_2}^t \circ (\phi_{K_1}^t)^{-1}(z)) dt \\
&\leq \int_0^1 (a(t) - b(t) + \epsilon - 3CE) [h(p) + h(\phi_{K_2}^t \circ (\phi_{K_1}^t)^{-1}(z))] dt \\
&\leq \int_0^1 2(a(t) - b(t) + \epsilon - 3CE) dt \quad \text{since } h(x) \leq 1 \ \forall x \in D_\epsilon \\
&\leq \int_0^1 2(CE + \epsilon - 3CE) dt \quad \text{by definition of } a(t) \text{ and } b(t) \\
&= 2\epsilon - 4CE.
\end{aligned} \tag{3.17}$$

Therefore, (3.16) becomes

$$\Delta(\phi_1^1 \circ (\phi_2^1)^{-1}, \alpha)_{\phi_2^1 \circ (\psi^1)^{-1}(y)} \leq 2\epsilon - 4CE. \tag{3.18}$$

Putting (3.18) in (3.14) we have

$$\Delta(\phi_1^1 \circ (\psi^1)^{-1}, \alpha)_y \leq \Delta(\phi_2^1 \circ (\psi^1)^{-1}, \alpha)_y + 2\epsilon - 4CE \tag{3.19}$$

Therefore, either $\Delta(\phi_2^1 \circ (\psi^1)^{-1}, \alpha)_y \geq 2CE - \epsilon$ or $\Delta(\phi_1^1 \circ (\psi^1)^{-1}, \alpha)_y \leq \epsilon - 2CE$. That is either ϕ_2^1 or ϕ_1^1 satisfies item (3). From Proposition 2.1, we have

$$\bar{d}(\Phi_1, \Phi) = \bar{d}(\phi_{K_2} \circ \Phi, \Phi) \leq C^\Phi \bar{d}(\phi_{K_2}, id),$$

since K_2 is compactly supported in D_ϵ , for an ϵ small enough, $\bar{d}(\Phi_1, \Phi)$ can be made as small as we want. For item 2., we have that

$$\begin{aligned}
l_S^{(1,\infty)}(\Phi_1) &\leq l_S^{(1,\infty)}(\phi_{K_1}) + l_S^{(1,\infty)}(\Phi) \leq \int_0^1 \left(a(t) + \frac{\epsilon - 3CE}{2} \right) \text{osc}(h) dt + E \\
&\leq \left(\text{osc}(\mathcal{F}_\Phi^\alpha(t)(\cdot)) + \frac{\epsilon - 3CE}{2} \right) \text{osc}(h) + E \\
&\leq \left(CE + \frac{\epsilon - 3CE}{2} \right) \text{osc}(h) + E \\
&\leq \left(\frac{\epsilon - CE}{2} \right) \frac{\epsilon}{CE + \epsilon} + E \leq E + \epsilon.
\end{aligned}$$

□

Remark 3.5.

1. The first de Rham cohomology group $H^1(M, \mathbb{R})$, is a finite dimensional vector space over the separable field \mathbb{R} . Hence, $H^1(M, \mathbb{R})$ is separable w.r.t the L^2 - norm ($\|\cdot\|_{L^2}$)
2. $\mathbb{H}_S(M)$ is separable since it is isomorphic to the separable space $H^1(M, \mathbb{R})$. Hence the space of paths $\mathcal{P}\mathbb{H}_S(M)$ from $[0, 1]$ to $\mathbb{H}_S(M)$ is separable w.r.t the norm $\int_0^1 \|\cdot\|_{L^2} dt$
3. The space of normalised functions $C_0^\infty(M \times [0, 1], \mathbb{R})$ is separable when endowed with the $L^{(1,\infty)}$ -norm. Therefore, $\mathfrak{T}(M, \omega, S) = C_0^\infty(M \times [0, 1], \mathbb{R}) \times \mathcal{P}\mathbb{H}_S(M)$ is separable w.r.t $l_S^{(1,\infty)}$

Proof of Theorem B. Let $E > 0$ and $(U, \mathcal{H}) \in \mathfrak{T}(M, \omega, \mathcal{S})$ such that $D_{\lambda, \mathcal{S}}^1((U, \mathcal{H}), (0, 0)) < E$. Consider a dense sequence $(V_j, \mathcal{K}_j)_j \subseteq \mathfrak{T}(M, \omega, \mathcal{S})$, then the corresponding sequence of time-one maps $(\phi_{(V_j, \mathcal{K}_j)}^1)_j \subseteq G_\omega(M)$ is dense w.r.t the Hofer-like topology. Construct inductively a sequence $(U_0, \mathcal{H}_0), (U_1, \mathcal{H}_1), \dots$ in $\mathfrak{T}(M, \omega, \mathcal{S})$: by setting $(U_0, \mathcal{H}_0) = (U, \mathcal{H})$, for each $k \geq 1$ Lemma 3.4 provides us with $(U_k, \mathcal{H}_k) \in \mathfrak{T}(M, \omega, \mathcal{S})$ such that

- (a) $l_S^{(1, \infty)}(\phi_{(U_k, \mathcal{H}_k)}) < E$ for each k ,
- (b) $\bar{d}(\phi_{(U_{k-1}, \mathcal{H}_{k-1})}, \phi_{(U_k, \mathcal{H}_k)}) < \frac{\epsilon}{k}$.
- (c) for any $\alpha \in \mathcal{Z}^1(M) \setminus \{0\}$, $\Delta\left(\phi_{(U_k, \mathcal{H}_k)}^1 \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) > CE - \frac{1}{j}$, for $1 \leq k \leq j$.

Property (b) above implies that $(\phi_{(U_k, \mathcal{H}_k)})_k$ converges uniformly to a continuous path $\{\psi^t\}$ of homeomorphisms of M . Property (a) and (b) together implies item 1 of Theorem B. For item 2, let $(\psi_k) \subseteq G_\omega(M)$, such that $\psi_k \xrightarrow{C^0} \psi^1$. Given $\delta > 0$, there exists infinitely many j for which $d_{HL}(\psi^1, \phi_{(V_j, \mathcal{K}_j)}^1) < \delta$. For such j and for every k , there exists a constant C from Theorem ?? such that

$$\begin{aligned} Cd_{HL}(\psi_k, \psi^1) &\geq Cd_{HL}(\psi_k, \phi_{(V_j, \mathcal{K}_j)}^1) - C\delta \geq \left\| \psi_k \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1} \right\|^\infty - C\delta \\ &\geq \left| \Delta\left(\psi_k \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) \right| - C\delta. \end{aligned} \quad (3.20)$$

Since $\phi_{(U_k, \mathcal{H}_k)}^1 \xrightarrow{C^0} \psi^1 \xleftarrow{C^0} \psi_k$, and $\Delta(\cdot, \alpha)$ is continuous w.r.t the C^0 topology, we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \left| \Delta\left(\phi_{(U_k, \mathcal{H}_k)}^1 \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) \right| &= \left| \Delta\left(\psi^1 \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) \right| \\ &= \lim_{k \rightarrow \infty} \left| \Delta\left(\psi_k \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) \right|. \end{aligned} \quad (3.21)$$

Combining (3.20) (3.21) with property (c) one gets

$$\begin{aligned} C \liminf_{k \rightarrow \infty} d_{HL}(\psi_k, \psi^1) &\geq \lim_{k \rightarrow \infty} \left| \Delta\left(\psi_k \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) \right| - C\delta \\ &= \lim_{k \rightarrow \infty} \left| \Delta\left(\phi_{(U_k, \mathcal{H}_k)}^1 \circ \left(\phi_{(V_j, \mathcal{K}_j)}^1\right)^{-1}, \alpha\right) \right| - C\delta \\ &\geq \lim_{k \rightarrow \infty} \left(CE - \frac{1}{j} - C\delta \right). \end{aligned}$$

Therefore, $\liminf_{k \rightarrow \infty} d_{HL}(\psi_k, \psi^1) \geq E - \delta$, for every $\delta > 0$. □

Acknowledgments :

The first author acknowledges support of the CEA-SMIA (IDA N° 6509-BJ et D5320), the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université), and LabEx CARMIN (ANR-10-LABX-59-01).

References

- [1] A. Banyaga, *The Structure of classical diffeomorphisms groups*, Mathematics and its applications **400**. Kluwer Academic Publisher's Group, Dordrecht, The Netherlands (1997).
- [2] A. Banyaga, A Hofer-like metric on the group of symplectic diffeomorphisms, *Contempt. Math. Amer. Math. Soc. RI.* **512** (2010), 1 – 24.

- [3] A. Banyaga, *On the group of strong symplectic homeomorphisms*, C. R. Math. Acad. Sci. Paris, **346**(15-16)(2008), 867 – 872.
- [4] A. Banyaga. *On The Group of Strong Symplectic Homeomorphisms*, CUBO A Mathematical Journal, **12**(03) (2010) , 49 – 69.
- [5] L. Buhovsky, *On two remarkable groups of area-preserving homeomorphisms*, Journal of Mathematical Physics, Analysis, Geometry,**19**(2) (2023), 339 – 373.
- [6] D. Cristofaro-Gardiner, V. Humilière, and S. Seyfaddini. PFH spectral invariants on the two-sphere and the large scale geometry of Hofer’s metric. Journal of the European Mathematical Society,(2021): n. pag.
- [7] D. Cristofaro-Gardiner, V. Humilière, C. Y. Mak, S. Seyfaddini, and I. Smith. Quantitative Heegaard Floer cohomology and the Calabi invariant. Forum of Mathematics, Pi (2022), 10, E27 1 – 59. doi:10.1017/fmp.2022.18
- [8] Y. Eliashberg, *A theorem on the structure of wave fronts and applications in symplectic topology*, Funct. Anal. Appl. 21 (1987), 227 – 232.
- [9] M. Gromov, *Pseudo-holomorphic curves in symplectic manifolds*, Invent. Math. 82 (1985), 307 – 347.
- [10] Y-G. Oh and S. Müller, *The group of Hamiltonian homeomorphisms and C^0 -symplectic topology*, J. Symp. Geometry **5**(2007), 167 – 225.
- [11] S. Tchuiaga, *On symplectic dynamics*, Differ. Geom. Appl. **61**(2018), 170 – 196.
- [12] S. Tchuiaga, *Hofer-like geometry and flux theory* , J. Dyn. Syst. Geom. Theories 19 (2) (2021) 227 – 270, <https://doi.org/10.1080/172603X.2021.2011110>.
- [13] S. Tchuiaga, C. Madengko *On the simplicity of the group of strong hamiltonian homeomorphisms”* to appear
- [14] S. Tchuiaga, F. Houenou, C. Madengko , A. Nguedakumana *C^0 –Transport of flux geometry* Journal of Topology and its Applications 322 (2022) 108301