

Temperature dependence of charge transport in the half-filled 1D Hubbard model

J. M. P. Carmelo

*Center of Physics of University of Minho and University of Porto, LaPMET, P-4169-007 Oporto, Portugal and
CeFEMA, Instituto Superior Técnico, Universidade de Lisboa,
LaPMET, Av. Rovisco Pais, P-1049-001 Lisboa, Portugal*

P. D. Sacramento

*CeFEMA, Instituto Superior Técnico, Universidade de Lisboa,
LaPMET, Av. Rovisco Pais, P-1049-001 Lisboa, Portugal*

The use of hydrodynamic transport theory seems to indicate that the charge diffusion constant D of the one-dimensional (1D) half-filled Hubbard model, whose Drude weight vanishes, diverges for temperature $T > 0$, which would imply anomalous superdiffusive charge transport. Here the leading term of that constant is derived for low finite temperatures $k_B T / \Delta_\eta \ll 1$ where Δ_η is the Mott-Hubbard gap. It only diverges in the $T \rightarrow 0$ limit, being finite and decreasing upon increasing T within the $k_B T / \Delta_\eta \ll 1$ regime. Our exact results both provide valuable physical information on a complex quantum problem and bring about the interesting unsolved issue of how charge transport evolves from normal diffusive for $k_B T / \Delta_\eta \ll 1$ to anomalous superdiffusive in the $k_B T \rightarrow \infty$ limit.

The repulsive one-dimensional (1D) Hubbard model [1–3] is the paradigmatic quantum system for low-dimensional strongly correlated electron systems and materials. The real part of the charge conductivity of that model reads,

$$\sigma(\omega, T) = 2\pi D^z(T)\delta(\omega) + \sigma_{\text{reg}}(\omega, T). \quad (1)$$

When the charge stiffness $D^z(T)$ in its singular part is finite, the dominant charge transport is ballistic. It is well established that in the thermodynamic limit the charge stiffness exactly vanishes at half filling for all temperatures $T > 0$ [4, 5].

The use of hydrodynamic theory and Kardar-Parisi-Zhang (KPZ) scaling to study dynamical scaling properties of charge transport associated with the regular part of the conductivity $\sigma_{\text{reg}}(\omega, T)$ seems to indicate that the diffusion constant D diverges for $T > 0$, which would imply anomalous superdiffusive charge transport [6–8]. However, the charge transport associated with the regular part of dc charge conductivity that characterizes the sub-ballistic timescales is a very complex quantum problem.

In this Letter we derive the exact expression of the leading term of the charge diffusion constant D of the half-filled 1D Hubbard model for magnetic fields $h \in [0, h_c]$ in the $k_B T / \Delta_\eta \ll 1$ regime. Here Δ_η is the Mott-Hubbard gap and h_c is the critical magnetic field for $T = 0$ fully polarized ferromagnetism. That constant is found to diverge only in the $T \rightarrow 0$ limit, being finite and decreasing upon increasing T within the $k_B T / \Delta_\eta \ll 1$ regime. This reveals normal diffusive charge transport in that low-temperature regime.

Our exact results thus contradict the expectation of Refs. 6–8 that the anomalous superdiffusive charge transport found for high temperatures $T \rightarrow \infty$ applies to all finite temperatures $T > 0$. The regular part of dc charge conductivity is thus a quantum problem more involved

than expected that deserves further investigations. In addition to provide valuable physical information on that complex quantum problem, our results bring about the interesting unsolved issue of how charge transport evolves from normal diffusive for $k_B T / \Delta_\eta \ll 1$ to anomalous superdiffusive for $k_B T \rightarrow \infty$.

The 1D Hubbard model describes N electrons in a lattice of length L with N_a sites. We consider the thermodynamic limit, its Hamiltonian at chemical potential $\mu = 0$ in a magnetic field magnetic field $h \in [0, h_c]$ and under periodic boundary conditions reading,

$$\hat{H} = -t \sum_{\sigma, j} \left[c_{j, \sigma}^\dagger c_{j+1, \sigma} + \text{h.c.} \right] + U \sum_j \hat{\rho}_{j, \uparrow} \hat{\rho}_{j, \downarrow} + \mu_0 h \hat{S}_s^z. \quad (2)$$

Here $c_{j, \sigma}^\dagger$ creates one electron of spin projection σ at site j , $\hat{\rho}_{j, \sigma} = (\hat{n}_{j, \sigma} - 1/2)$, $\hat{n}_{j, \sigma} = c_{j, \sigma}^\dagger c_{j, \sigma}$, $\hat{S}_s^z = \frac{1}{2} \sum_{j=1}^L (\hat{n}_{j, \uparrow} - \hat{n}_{j, \downarrow})$, $\mu_0 = 2\mu_B$, and $h_c = (\sqrt{(4t)^2 + U^2} - U) / \mu_0$.

At fixed field $h < h_c$ and $T = 0$, fully polarized ferromagnetism is also achieved upon increasing U above $U_c = \frac{8t^2}{\mu_0 h} - \frac{1}{2}\mu_0 h$. Upon increasing h , U_c continuously decreases from $U_c = \infty$ at $h = 0$ to $U_c = 0$ at $h = h_c$. The densities read $m_\eta^z = (N_a - N) / L$ and $m_s^z = (N_\uparrow - N_\downarrow) / L$. We use natural units in which the Planck constant, electronic charge, and lattice spacing are equal to one, so that $N_a = L$.

In the last three to four decades, representations in terms of spinons and holons have been widely used to successfully describe the static and dynamical properties of both integrable 1D models and the physics of the materials they describe [9]. Hence such representations became the paradigm of the 1D physics. Although the quantum problem under study here is found to be defined in a small subspace for $k_B T / \Delta_\eta \ll 1$, the suitable identification of the charge carriers requires the use of a general representation that refers to the whole Hilbert space beyond the holon-spinon paradigm [5].

For $u = U/4t > 0$ the global symmetry of the 1D Hubbard model at chemical potential $\mu = 0$ and magnetic field $h = 0$ is larger than $SO(4) = [SU(2) \otimes SU(2)]/Z_2$, as usually assumed, and given by $[SO(4) \otimes U(1)]/Z_2$, which can be written as $[SU(2) \otimes SU(2) \otimes U(1)]/Z_2^2$ and $SO(3) \otimes SO(3) \otimes U(1)$ [10]. This is consistent with the form of the two monodromy matrices of the Bethe-ansatz solution inverse-scattering method [3, 11, 12]. The corresponding η -spin S_η , spin S_s , $S_\eta^z = \frac{1}{2}(L - N)$, and $S_s^z = \frac{1}{2}(N_\uparrow - N_\downarrow)$ are good quantum numbers. Singlet and multiplet refer in the following to η -spin or spin configurations.

Within the representation used here [5], the Bethe-ansatz solution performs a squeezed-space construction [13]. It leads to (i) a set of $n = 1, \dots, \infty$ ηn -squeezed effective lattices, (ii) a set of $n = 1, \dots, \infty$ sn -squeezed effective lattices, and (iii) a τ -squeezed effective lattice whose occupancy configurations generate irreducible representations of the (i) η -spin $SU(2)$ symmetry, (ii) spin $SU(2)$ symmetry, and (iii) τ -translational $U(1)$ symmetry, respectively, in $[SU(2) \otimes SU(2) \otimes U(1)]/Z_2^2$. Each such squeezed effective lattice refers to a corresponding ηn -band, sn -band, and τ -band, respectively, whose discrete momentum values q_j such that $q_{j+1} - q_j = 2\pi/L$ are Bethe-ansatz quantum numbers multiplied by $2\pi/L$ [5].

The squeezed-space construction is such that the (a) N_τ τ -particles and (b) $N_\tau^h = L - N_\tau$ τ -holes of the τ -squeezed effective lattice refer for $u > 0$ to (a) singly occupied sites and (b) unoccupied and doubly occupied sites, respectively, of the electrons for $u \rightarrow \infty$. The corresponding (a) spin and (b) η -spin degrees of freedom are described by the occupancies of (a) $L_s = N_\tau$ physical spins 1/2 and (b) $L_\eta = N_\tau^h$ physical η -spins 1/2 in the (a) sn -squeezed effective lattices and (b) ηn -squeezed effective lattices, respectively. (A $u > 0$ physical η -spin of projection +1/2 and -1/2 refers to a $u \rightarrow \infty$ unoccupied and a doubly occupied site by electrons, respectively.)

The two $SU(2)$ symmetries in $[SU(2) \otimes SU(2) \otimes U(1)]/Z_2^2$ impose that for *all* energy eigenstates the number L_α can be written as $L_\alpha = N_\alpha + \mathcal{N}_\alpha$. Here $N_\alpha = N_{\alpha,+1/2} + N_{\alpha,-1/2} = 2S_\alpha$ where $N_{\alpha,\pm 1/2} = S_\alpha \pm S_\alpha^z$ is the number of *unpaired physical spins* ($\alpha = s$) and *unpaired physical η -spins* ($\alpha = \eta$) that participate in a multiplet configuration and $\mathcal{N}_\alpha = \mathcal{N}_{\alpha,+1/2} + \mathcal{N}_{\alpha,-1/2}$ where $\mathcal{N}_{\alpha,\pm 1/2} = L_\alpha/2 - S_\alpha$ is that of *paired physical spins* ($\alpha = s$) and *paired physical η -spins* ($\alpha = \eta$) that participate in singlet configurations. All the latter are contained in αn -pairs, each involving a number $n = 1, \dots, \infty$ of unbound ($n = 1$) and bound ($n > 1$) singlet pairs. The number $N_{\alpha n}$ of such pairs thus obeys the sum-rule $\sum_{n=1}^{\infty} 2n N_{\alpha n} = \mathcal{N}_\alpha$. The $N_{\alpha n}$ αn -pairs occupy a number $N_{\alpha n}$ of sites of the corresponding αn -squeezed effective lattice. The corresponding $N_{\alpha n}^h = N_\alpha + \sum_{n'=n+1}^{\infty} 2(n' - n) N_{\alpha n'}$ holes of that lattice include the N_α unpaired physical spins ($\alpha = s$) or unpaired physical η -spins ($\alpha = \eta$) and a subset of $\sum_{n'=n+1}^{\infty} 2(n' - n) N_{\alpha n'} < \mathcal{N}_\alpha$

paired physical spins ($\alpha = s$) or unpaired physical η -spins ($\alpha = \eta$) from $n' > n$ $\alpha n'$ -branches.

The Hamiltonian, Eq. (2), in the presence of a uniform vector potential (twisted boundary conditions) remains solvable by the Bethe ansatz [5]. The energy eigenstates described by the Bethe ansatz are η -spin highest-weight states (HWSs) [11] whose $N_\eta = 2S_\eta$ unpaired physical η -spins have the same projection +1/2. For them, the momentum eigenvalues, $P(\Phi)$, have the general form,

$$P(\Phi/L) = P(0) - (L_\eta - \sum_n 2n N_{\eta n}) \frac{\Phi}{L} = P(0) - N_\eta \frac{\Phi}{L}, \quad (3)$$

where $P(0)$ is their value for $\Phi = 0$ [5]. The term $-L_\eta \frac{\Phi}{L}$ in $-(L_\eta - \sum_n 2n N_{\eta n}) \frac{\Phi}{L}$ refers to *all* L_η physical η -spins coupling to the vector potential in the absence of physical η -spins singlet pairing. Indeed, the coupling counter terms $\sum_n 2n N_{\eta n} \frac{\Phi}{L}$ refer to the number $2n$ of paired physical η -spins in each ηn -pair. These counter terms *exactly cancel* the coupling of the corresponding $2n$ paired physical η -spins in each such an ηn -pair. As a result of such counter terms, only the $N_\eta = L_\eta - \sum_n 2n N_{\eta n}$ unpaired physical η -spins couple to the vector potential and thus carry charge current. For non-HWSs, the result is $P(\Phi/L) = P(0) - (N_{\eta,+1/2} - N_{\eta,-1/2}) \frac{\Phi}{L}$.

Hence for *all* energy eigenstates the *charge carriers* are the unpaired physical η -spins. One can generate from each η -spin HWS for which $N_\eta = N_{\eta,+1/2}$ a set of $n_\eta^z = 1, \dots, 2S_\eta$ non-HWSs populated by $N_{\eta,+1/2} = 2S_\eta - n_\eta^z$ and $N_{\eta,-1/2} = n_\eta^z$ unpaired physical η -spins with opposite projection and thus opposite coupling to the vector potential [11]. The elementary charge currents $j_{\pm 1/2}$ carried by each unpaired physical η -spin that populates a HWS and its non-HWSs is then found to read [5],

$$j_{\pm 1/2} = \pm \frac{\langle \hat{J}_{HWS}^z(S_\eta) \rangle}{2S_\eta} = \pm \frac{\langle \hat{J}_{HWS}^z(S_\eta) \rangle}{N_\eta}. \quad (4)$$

Here $\langle \hat{J}_{HWS}^z(S_\eta) \rangle$ denotes the charge current expectation value of the HWS. That of its non-HWSs reads $\langle \hat{J}^z(S_\eta^z) \rangle = \sum_{\sigma=\pm 1/2} j_\sigma N_{\eta,\sigma}$, which can be written as $\langle \hat{J}^z(S_\eta^z) \rangle = \frac{S_\eta^z}{S_\eta} \langle \hat{J}_{HWS}^z(S_\eta) \rangle$. This applies to *all* finite- S_η energy eigenstates.

The $\mu = 0$ ground states and all other $S_\eta = 0$ energy eigenstates have no charge carriers, so that their charge current expectation value vanishes. For general finite- S_η energy eigenstates, a number $N_\eta = 2S_\eta$ of holes out of both the (i) $N_\tau^h = N_\eta + \mathcal{N}_\eta$ τ -band holes and (ii) $N_{\eta n}^h = N_\eta + \sum_{n'=n+1}^{\infty} 2(n' - n) N_{\eta n'}$ holes of ηn -bands with finite occupancy, $N_{\eta n} > 0$, describe the translational degrees of freedom of the N_η unpaired physical η -spins that carry the elementary charge currents $j_{\pm 1/2}$, Eq. (4).

However, $\mathcal{N}_\eta = 0$ and thus $N_{\eta n} = 0$ for $n = 1, \dots, \infty$ for the states that contribute to the quantum problem studied in the following. The translational degrees of freedom of their unpaired physical η -spins are thus described only

by the $N_\tau^h = N_\eta$ τ -holes. In addition, $N_{sn} = 0$ for $n > 1$, their $s1$ -band being full for $s1$ -band momentum values $q \in [-k_{F\downarrow}, k_{F\downarrow}]$ and empty for $|q| \in [k_{F\downarrow}, k_{F\uparrow}]$ where $k_{F\uparrow} = \frac{\pi}{2}(1 + m_s^z)$ and $k_{F\downarrow} = \frac{\pi}{2}(1 - m_s^z)$ [11].

The charge diffusion constant D can be derived by accounting for the Kubo linear response formula for the real part of dc charge conductivity being related to it through the Einstein relation. At very low temperatures, the charge currents contributing to it are generated by processes where upon moving in the τ -squeezed effective lattice the $N_\tau^h = 2S_\eta$ τ -holes that describe the translational degrees of freedom of the $N_\eta = 2S_\eta$ unpaired physical η -spins interchange position with the $N_\tau = L - 2S_\eta$ τ -particles. In the following we identify the motion of the $N_\tau^h = 2S_\eta$ holes with that of such $N_\eta = 2S_\eta$ η -spin-1/2 charge carriers they translationally describe.

The physics associated with the diffusive charge transport is for $U > 0$ and $\mu = 0$ such that in the limit of very low temperatures nearly ballistic transport occurs, but with zero charge stiffness. Nearly ballistic transport with zero spin stiffness also occurs in the gapped spin-1/2 XXZ chain [14]. In the present case, the corresponding charge carriers are actually η -spin triplet pairs that in the τ -squeezed effective lattice refer to two adjacent unpaired physical η -spins with the same projection $+1/2$ or $-1/2$. We call them *η -spin triplet charge carriers* to distinguish from their two unpaired physical η -spins.

Relative to the $\mu = 0$ ground states, the excitation energy and momentum of one such a $S_\eta^z = \pm 1$ η -spin triplet pair read,

$$\delta E = -\varepsilon_\tau(q) - \varepsilon_\tau(q') + 2\mu_B h \quad \text{and} \quad k \equiv \delta P = \pm\pi - q - q', \quad (5)$$

respectively, where the τ -band energy dispersion $\varepsilon_\tau(q)$ is defined in Ref. 11 for $h \in [0, h_c]$. Here $-\varepsilon_\tau(q) > 0$ and $-\varepsilon_\tau(q') > 0$ are the energies for creation of two corresponding τ -holes of τ -band momentum $-q$ and $-q'$, respectively, and the τ -band momentum values shift $\pm\pi/L$ that gives the k 's term $\pm\pi$ and the energy $2\mu_B h$ are due to a $s1$ -band occupancy deviation $\delta N_{s1} = -1$ [11].

At very low temperature the excitation energy of such η -spin triplet pairs, Eq. (5), must be very near its minimum allowed value, the Mott-Hubbard gap,

$$\Delta_\eta = \Delta_\tau + 2\mu_B h, \quad (6)$$

where $\Delta_\tau = -2\varepsilon_\tau(\pi)$. It is plotted in Fig. 1 (a) of Ref. 11, its general expression for $h \in [0, h_c]$ being given in that reference. It involves the excitation energy of the two τ -holes, Eq. (5), at $-q = \pm\pi$ and $-q' = \pm\pi \mp 2\pi/L$.

The translational degrees of freedom of the η -spin triplet charge carriers that play the role of low-temperature diffusing charges are thus described by two τ -holes whose τ -band momentum values $-q$ and $-q'$ are very near $-\pi$ or π , $-q \approx \mp\pi$ and $-q' = -q \pm 2\pi/L$. This gives an excitation momentum $k = \pm\pi - 2q \pm 2\pi/L$ that reads $k = \pm\pi - 2q \approx \mp\pi$ in the thermodynamic limit,

the corresponding two unpaired physical η -spins with the same projection $+1/2$ or $-1/2$ being adjacent in the τ -squeezed effective lattice upon moving with nearly the same τ -band momentum.

$S_\eta^z = 0$ η -spin triplet pairs have the same minimal excitation energy Δ_η yet involve two η -spin carriers with *opposite* projection. They do not contribute to charge transport because their elementary charge currents, Eq. (4), cancel each other. The same applies to *singlet* $\eta1$ -pairs, which are the only ηn -pairs whose minimal excitation energy is also Δ_η , Eq. (6).

The lowest excited states have a single τ -squeezed effective lattice domain wall that refers to a single η -spin triplet pair. The domain wall or diffusing charge associated with the η -spin triplet pair moves ballistically for such lowest excited states. Also for excited states with a finite number and thus vanishing concentration of η -spin triplet pairs, such pairs move ballistically over large τ -squeezed effective lattice distances before interacting with other η -spin triplet pairs. They carry an elementary charge current $j_{\pm 1}(k) = 2j_{\pm 1/2}(q)|_{q=(\pm\pi-k)/2}$ where $j_{\pm 1/2}(q)$ is that carried by each of the two corresponding adjacent unpaired physical η -spins, Eq. (4). It reads [11],

$$j_{\pm 1}(k) = \mp \frac{(k \pm \pi)}{M_t} \quad \text{for} \quad k = \pm\pi - 2q \approx \mp\pi. \quad (7)$$

Here M_t is the charge transport mass of the η -spin triplet pairs such that $1/M_t = |\partial j_{\pm 1}(k)/\partial dk|_{k=\mp\pi}$ whose general expression for $h \in [0, h_c]$ is given in Ref. 11. (See Fig. 1.)

Consistently with nearly ballistic charge transport, but with zero charge stiffness, the leading term of the charge diffusion constant D in the $k_B T/\Delta_\eta \ll 1$ regime can be expressed in terms of the first derivative with respect to m_η^z of the $T = 0$ charge stiffness D^z for $m_\eta^z \ll 1$ [11],

$$D = \frac{e^{\Delta_\eta \beta}}{2} \frac{dD^z}{dm_\eta^z} \Big|_{m_\eta^z=0} = \frac{e^{\Delta_\eta \beta}}{4M_t} \quad \text{where} \quad D^z = \frac{m_\eta^z}{2M_t}. \quad (8)$$

That $D \propto e^{\Delta_\eta \beta}$ for low temperature where $\beta = 1/k_B T$ applies to gapped 1D models in the quantum Sine-Gordon universality class [15]. In the opposite limit of high temperature, there is for some models a relation between D and the stiffness that rather involves a second derivative [6, 16].

To arrive to the expression, Eq. (8), we treated the quantum problem as a dilute gas of excited η -spin triplet pairs, each with energy $\epsilon_\eta(k) = \Delta_\eta + \frac{(k \pm \pi)^2}{2M_\eta}$ and velocity $v_\eta(k) = \frac{k \pm \pi}{M_\eta}$ where $M_\eta = 2M_t$ such that $1/M_\eta = \frac{d^2 \epsilon_\eta(k)}{dk^2}$ is their static mass [11]. Their motion and collisions dominate the charge transport properties, their spacing $x_\eta = (\beta/M_\eta)^{1/2} e^{\Delta_\eta \beta}$ being for $k_B T/\Delta_\eta \ll 1$ much larger than their thermal de Broglie wavelength $\lambda_\eta = (\beta/M_\eta)^{1/2}$. The quantum problem to be solved then refers to the collisions of the η -spin triplet pairs and




FIG. 1. The charge transport mass $M_t = M_\eta/2$ of the η -spin triplet pairs appearing in Eqs. (7) and (8) (a) and its inverse $1/M_t$ (b) plotted for several values of U in units of transfer integral t as a function of spin density for the interval $m_s^z = 2S_s^z/L \in [0, 1]$ that refers to magnetic fields $h \in [0, h_c]$. The charge transport mass $M_t = M_\eta/2 \in [0, 1/2t]$ vanishes in the $U \rightarrow 0$ limit, consistently with charge ballistic transport at $U = 0$, and is enhanced upon increasing $u = U/4t$ and/or the magnetic field h .

corresponding S -matrix. However, the root mean square thermal velocity of such pairs, $v_{\eta,\beta} = 1/\sqrt{M_\eta\beta}$, tends to zero as $k_B T/\Delta_\eta \rightarrow 0$, so that the S -matrix that describes the present quantum problem refers to vanishing exchange incoming or outgoing momenta.

This much simplifies the problem, the leading behaviors of the uniform charge susceptibility and real part of the dc charge conductivity being for the $k_B T/\Delta_\eta \ll 1$ regime and in our units expressed solely in terms of the two above length scales as $\chi = (2/\pi)^{1/2} \frac{\beta}{x_\eta}$ and $\sigma = (2/\pi)^{1/2} \frac{\lambda_\eta}{2}$, respectively [11, 17]. The use of the Einstein relation, $\sigma = \chi D$, then gives the leading behavior $D = \sigma/\chi = \frac{x_\eta \lambda_\eta}{2\beta}$, so that $D = e^{\Delta_\eta \beta}/(4M_t)$ for $k_B T/\Delta_\eta \ll 1$, as given in Eq. (8).

The Mott-Hubbard gap Δ_η , Eq. (6), in the expression, Eq. (8), vanishes at $U = 0$ and continuously increases upon increasing $u = U/4t$. It is plotted as a function of u in Fig. 1 (a) of Refs. [11] and [18] for $m_s^z = 0$ ($h = 0$) and $m_s^z = 1$ ($h = h_c$). Interestingly, in spite of the apparently different expressions $\Delta_\eta = U - 4t + 8t \int_0^\infty d\omega \frac{J_1(\omega)}{\omega(1+e^{2\omega u})}$ at $h = 0$ where $J_1(\omega)$ is a Bessel function and $\Delta_\eta = \sqrt{(4t)^2 + U^2} - 4t$ at $h = h_c$, the two corresponding curves in the figure are very similar, which shows that charge-spin coupling effects are very small at $\mu = 0$. (The inset of that figure shows the amplified deviations between the two lines, the amplified general deviation $\Delta_\eta(m_s^z) - \Delta_\eta(0)$ being shown in Fig. 1 (b) of Ref. 11 for $m_s^z \in [0, 1]$.)

For fields $h \in [0, h_c]$, the inverse transport mass $1/M_t$ in Eqs. (7) and (8) decreases upon increasing U from ∞ for $U \rightarrow 0$, consistently with the $U = 0$ ballistic charge transport, to $2t$ at $U = U_c$. M_t (a) and $1/M_t$ (b) are plotted in Fig. 1 for several values of U in units of t as a function of spin density for the interval $m_s^z = 2S_s^z/L \in [0, 1]$ that refers to magnetic fields $h \in [0, h_c]$.

Concerning the physics behind our results, the present interesting case of nearly ballistic charge transport, but with zero charge stiffness, is consistent with D being proportional to the inverse transport mass of the η -spin triplet charge carriers. Their concentration goes to zero for $k_B T/\Delta_\eta \rightarrow 0$ as $S_\eta/L = 1/x_\eta = (M_\eta/\beta)^{1/2} e^{-\Delta_\eta \beta}$. At the same time, the τ -squeezed effective lattice average distance $x_\eta = (\beta/M_\eta)^{1/2} e^{\Delta_\eta \beta}$ traveled by one η -spin triplet pair before it interacts with other such pairs and ceases to be ballistic diverges exponentially. The interplay between the exponential factors in the concentration of charge carriers and the distance between them is behind D diverging exponentially, Eq. (8). It becomes infinite in the $T \rightarrow 0$ limit, having large finite values at low finite temperatures.

In contrast to the results of Refs. 6–8, which predict that the charge anomalous superdiffusive transport found for high temperatures applies to all finite temperatures $T > 0$, we thus find that for $k_B T/\Delta_\eta \ll 1$ the charge diffusion constant leading term is finite and given by Eq. (8), which implies normal diffusive charge transport. At very low temperatures within the $k_B T/\Delta_\eta \ll 1$ regime, the charge diffusion constant, Eq. (8), actually decreases upon increasing the temperature.

The anomalous superdiffusive spin transport at any finite temperature predicted by hydrodynamic theory (KPZ scaling) for the spin-1/2 XXX chain [11, 19, 20] and half-filled 1D Hubbard model is compatible with the gapless nature of spin excitations. Its prediction of superdiffusive charge transport is then based on the assumption that at half filling η -spin and spin $SU(2)$ symmetries are related by duality, which would imply that spin and charge transport have identical properties. However, the physical reason why concerning charge

transport that prediction fails for $k_B T/\Delta_\eta \ll 1$ is that due to quantum effects the above duality relation is broken by the Mott-Hubbard gap [11].

By both providing the exact expression of the charge diffusion constant D in the $k_B T/\Delta_\eta \ll 1$ regime, Eq. (8), and showing that finite-temperature charge transport is a much more complex quantum problem than predicted by hydrodynamic theory, as at low temperature KPZ scaling is erased by *quantum* effects associated with the Mott-Hubbard gap [11], our results open a new avenue for the further understanding of the charge transport properties of the paradigmatic quantum system for low-dimensional strongly correlated electron systems, the 1D Hubbard model.

However, the results obtained by hydrodynamic theory and KPZ scaling in Refs. 6–8 concerning charge transport are valid at high temperatures $T \rightarrow \infty$ for which the above duality relation emerges, so that it is anomalous superdiffusive in that regime. Combining such a behavior with that found here for the $k_B T/\Delta_\eta \ll 1$ regime and accounting for the lack of phase transitions imposed by the Mermin-Wagner Theorem, there must be a crossover such that D decreases upon enhancing T until reaching a minimum at some intermediate U -dependent temperature and then increases up to $D = \infty$ for $T \rightarrow \infty$. The full understanding of that crossover requires studies on charge transport in the half-filled Hubbard model for temperatures $T \approx \Delta_\eta/k_B$, which is a very complex quantum problem beyond the goals of this Letter.

Our exact results apply to the (i) charge and (ii) spin degrees of freedom of the 1D Hubbard model for (i) $\mu = 0$ and $U > 0$ and (ii) $h = 0$ and $U < 0$, respectively. They provide valuable physical information on a very complex quantum problem [11, 21] and show that finite-temperature charge (spin) transport in the $\mu = 0$ repulsive ($h = 0$ attractive) 1D Hubbard model deserves further studies to clarify the mechanisms under which it evolves from normal diffusive for $k_B T/\Delta_\alpha \ll 1$ to anomalous superdiffusive for $k_B T \rightarrow \infty$ where $\alpha = \eta$ ($\alpha = s$.)

We thank Tomaž Prosen and Subir Sachdev for illuminating discussions. J. M. P. C. and P. D. S. acknowledge support from Fundação para a Ciência e Tecnologia through the Grant UIDB/04650/2020 and UID/CTM/04540/2019, respectively.

-
- [1] E. H. Lieb, F. Y. Wu, *Absence of Mott transition in an exact solution of the short-range one-band model in one dimension*, Phys. Rev. Lett. **20**, 1445 (1968); **21**, 192 (1968).
 [2] M. Takahashi, *One-dimensional Hubbard model at finite temperature*, Progr. Theor. Phys **47**, 69 (1972).
 [3] M. J. Martins and P. B. Ramos, *The quantum inverse*

- scattering method for Hubbard-like models*, Nucl. Phys. B **522**, 413 (1998).
 [4] E. Ilievski and J. De Nardis, *Ballistic transport in the one-dimensional Hubbard model: The hydrodynamic approach*, Phys. Rev. B **96**, 081118(R) (2017).
 [5] J. M. P. Carmelo, S. Nemati, and T. Prosen, *Absence of ballistic charge transport in the half-filled 1D Hubbard model*, Nucl. Phys. B **930**, 418 (2018).
 [6] E. Ilievski, J. De Nardis, M. Medenjak, and T. Prosen, *Superdiffusion in one-dimensional quantum lattice models*, Phys. Rev. Lett. **121**, 230602 (2018).
 [7] M. Fava, B. Ware, S. Gopalakrishnan, R. Vasseur, and S. A. Parameswaran, *Spin crossovers and superdiffusion in the one-dimensional Hubbard model*, Phys. Rev. B **102**, 115121 (2020).
 [8] C. P. Moca, M. A. Werne, A. Valli, T. Prosen, and G. Zaránd, *Kardar-Parisi-Zhang scaling in the Hubbard model*, Phys. Rev. B **108**, 235139 (2023).
 [9] R. G. Pereira, K. Penc, S. R. White, P. D. Sacramento, and J. M. P. Carmelo, *Charge dynamics in half-filled Hubbard chains with finite on-site interaction*, Phys. Rev. B **85**, 165132 (2012).
 [10] J. M. P. Carmelo, S. Östlund, and M. J. Sampaio, *Global $SO(3) \times SO(3) \times U(1)$ symmetry of the Hubbard model on bipartite lattices*, Ann. Phys. **325**, 1550 (2010).
 [11] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.110.L201108> for the model representation and basic quantities used in the studies of this Letter.
 [12] L. D. Faddeev and L. A. Takhtajan, *What is the spin of a spin wave?*, Phys. Lett. A **85**, 375 (1981).
 [13] H. V. Kruis, I. P. McCulloch, Z. Nussinov, and J. Zaanen, *Geometry and the hidden order of Luttinger liquids: The universality of squeezed space*, Phys. Rev. B **70**, 075109 (2004).
 [14] C. Karrasch, J. E. Moore, and F. Heidrich-Meisner, *Real-time and real-space spin and energy dynamics in one-dimensional spin- $\frac{1}{2}$ systems induced by local quantum quenches at finite temperatures*, Phys. Rev. B **89**, 075139 (2014).
 [15] K. Damle and S. Sachdev, *Universal relaxational dynamics of gapped one-dimensional models in the quantum Sine-Gordon universality class*, Phys. Rev. Lett. **95**, 187201 (2005).
 [16] M. Medenjak, C. Karrasch, and T. Prosen, *Lower bounding diffusion constant by the curvature of Drude weight*, Phys. Rev. Lett. **119**, 080602 (2017).
 [17] J. M. P. Carmelo, L. M. Martelo, and P. D. Sacramento, *One- and two-electron spectral function expressions in the vicinity of the upper-Hubbard bands lower limit*, J. Phys.: Condens. Matter **16**, 1375 (2004).
 [18] J. M. P. Carmelo, Čadež, and P. D. Sacramento, *One-particle spectral functions of the one-dimensional Fermionic Hubbard model with one fermion per site at zero and finite magnetic fields*, Phys. Rev. B **103**, 195129 (2021).
 [19] M. Ljubotina, M. Žnidarič, and T. Prosen, *Kardar-Parisi-Zhang physics in the quantum Heisenberg magnet*, Phys. Rev. Lett. **122**, 210602 (2019).
 [20] M. Ljubotina, M. Žnidarič, and T. Prosen, *Spin diffusion from an inhomogeneous quench in an integrable system*, Nat. Commun. **8**, 16117 (2017).
 [21] A. M. Tsvelik, *1 + 1-dimensional sigma model at finite*

temperature, Sov. Phys. JETP **66**, 221 (1987).