On Reductions and Representations of Learning Problems in Euclidean Spaces

Bogdan Chornomaz

Shay Moran[†]

Tom Waknine *

November 19, 2024

Abstract

Many practical prediction algorithms represent inputs in Euclidean space and replace the discrete 0/1 classification loss with a real-valued surrogate loss, effectively reducing classification tasks to stochastic optimization. In this paper, we investigate the expressivity of such reductions in terms of key resources, including dimension and the role of randomness.

We establish bounds on the minimum Euclidean dimension D needed to reduce a concept class with VC dimension d to a Stochastic Convex Optimization (SCO) problem in \mathbb{R}^D , formally addressing the intuitive interpretation of the VC dimension as the number of parameters needed to learn the class. To achieve this, we develop a generalization of the Borsuk-Ulam Theorem that combines the classical topological approach with convexity considerations. Perhaps surprisingly, we show that, in some cases, the number of parameters D must be exponentially larger than the VC dimension d, even if the reduction is only slightly non-trivial. We also present natural classification tasks that can be represented in much smaller dimensions by leveraging randomness, as seen in techniques like random initialization. This result resolves an open question posed by Kamath, Montasser, and Srebro (COLT 2020).

Our findings introduce new variants of *dimension complexity* (also known as *sign-rank*), a wellstudied parameter in learning and complexity theory. Specifically, we define an approximate version of sign-rank and another variant that captures the minimum dimension required for a reduction to SCO. We also propose several open questions and directions for future research.

1 Introduction

Reduction is a fundamental concept in computer science, serving as a basic primitive in both computability and complexity theory. It plays a crucial role in defining complexity classes, such as P and NP, by enabling structured transformations of problems into one another. Through reductions, we can compare the difficulty of different problems, determine their relative complexity, and classify them accordingly. This framework has been instrumental in understanding computational limits and identifying problems that are efficiently solvable or intractable, shaping the study of algorithms and theoretical computer science.

Reductions are also a common and powerful tool in machine learning. Implicitly, reductions are used whenever one solves a problem by translating it into an already solved one. For instance, any linear representation of a classification task, such as those used in kernel machines, can be viewed as a reduction from the original learning task to linear classification (half-spaces). Another prominent example is the use of convex surrogate losses in place of the discrete 0/1 classification loss, which reduces the classification task to (stochastic) convex optimization. Similarly, representation learning,

^{*}Departments of Mathematics, Technion.

[†]Departments of Mathematics, Computer Science, and Data and Decision Sciences, Technion and Google Research. Robert J. Shillman Fellow; supported by ISF grant 1225/20, by BSF grant 2018385, by Israel PBC-VATAT, by the Technion Center for Machine Learning and Intelligent Systems (MLIS), and by the European Union (ERC, GENER-ALIZATION, 101039692). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

which focuses on learning features, and meta-learning, which focuses on learning which learning algorithm to use, can both be interpreted as forms of learning reductions. This perspective is evident, for example, in the formulation of meta-learning by Aliakbarpour, Bairaktari, Brown, Smith, Srebro, and Ullman [2024].

Historically, reductions in Valiant's PAC learning model have been studied since the 1980s, primarily in the form of representations—mapping a concept class we wish to learn into a concept class we know how to learn [Pitt and Warmuth, 1990]. In particular, geometric representations as halfspaces have been thoroughly explored in both learning theory and complexity theory, a partial list includes [Ben-David, Eiron, and Simon, 2002, Linial and Shraibman, 2009, Forster, Krause, Lokam, Mubarakzjanov, Schmitt, and Simon, 2001, Forster, Schmitt, Simon, and Suttorp, 2003, Forster, 2002, Alon, Moran, and Yehudayoff, 2017, Kamath, Montasser, and Srebro, 2020, Hatami, Hosseini, and Meng, 2022].

In this paper, we extend this line of research by studying reductions to arbitrary stochastic convex optimization (SCO) problems [Shalev-Shwartz, Shamir, Srebro, and Sridharan, 2009]. We relax the notion of dimension complexity by examining the minimum dimension *d* for which a given class can be reduced to a *d*-dimensional SCO problem. Additionally, we explore reductions that exploit randomness and agnostic learners, demonstrating their advantages over naive representation-based reductions.

Organization. We begin in Section 2 by presenting our main results. In Section 3, we provide an overview of the central tools and ideas that underpin our proofs. Next, Section 4 presents natural examples of reductions that illustrate the tightness of our results. In Section 5, we cover preliminaries and basic definitions. The remaining sections are devoted to the detailed proofs.

2 Main results

In this section, we assume familiarity with basic concepts in learning theory such as concept classes, loss functions, and VC dimension. These definitions are provided in detail in Section 5.

We focus on reducing realizable-case classification problems to well-studied geometric learning tasks: (i) stochastic convex optimization (SCO) and (ii) linear classification. Our impossibility results extend even to non-uniform (distribution-dependent) learning, where sample complexity may vary depending on the input distribution. We examine how the dimension of the reduced problem relates to the VC dimension of the original classification task, and explore whether introducing randomness can help reduce this dimension.

Both the VC dimension and the Euclidean dimension are key parameters in classification and stochastic convex optimization (SCO), respectively: the VC dimension is the fundamental parameter characterizing PAC learnability, as highlighted by the fundamental theorem of PAC learning [Shalev-Shwartz and Ben-David, 2014]. In SCO, the Euclidean dimension is closely tied to computational and space complexity; for example, the complexity of arithmetic operations—essential for computing gradients (at training time) and predictions (at test time)—scales with this dimension. It also influences model interpretability, as models with fewer parameters are generally easier for humans to understand. Studying the relationship between these dimensions also addresses the intuitive interpretation of the VC dimension as the number of parameters needed to describe a concept class; we discuss this in further detail after Theorem 1.

A natural approach to relate the VC dimension and the Euclidean dimension would be through a sample complexity analysis, such as showing that if a binary concept class C is reducible to an SCO problem in \mathbb{R}^d , then its sample complexity is at most O(d). However, this approach cannot work because, in SCO, the Euclidean dimension does not determine sample complexity; instead, it depends on factors like Lipschitzness and the diameter of the parameter space. To overcome this, we employ a topological argument, specifically a variant of the Borsuk-Ulam Theorem.

Section Organization. In Section 2.1, we formally define reductions between learning tasks. In Section 2.2, we present our main result on reductions from classification to stochastic convex optimiza-

tion, and in Section 2.3, we present our results on representing classification tasks using half-spaces, a well-studied and useful special case of reductions.

2.1 Reductions



Figure 1: A reduction from problem A, which we wish to solve, to problem B, which we can solve. The reduction maps instances of A to instances of B, and solutions of B back to solutions of A. The reduction is successful if, when combined with an algorithm for B, it solves problem A.

To define reductions, we first introduce a simple abstraction of learning problems that captures both binary classification and stochastic convex optimization, as well as other scenarios. We start by establishing the appropriate terminology, and after that, we will illustrate it in examples.

Definition 1 (Learning task). A learning task \mathcal{T} is a tuple $\mathcal{T} = (H, C, Z, P)$, where H is a hypothesis class, $C \subseteq H$ is a concept class, Z is the space of labeled examples, and P is a family of distributions over Z. Additionally, each example $z \in Z$ has an associated loss function $\ell_z : H \to \mathbb{R}_{>0}$. For a distribution $D \in P$, the loss of an hypothesis $h \in H$ is given by $L_D(h) = \mathbb{E}_{z\sim D}[\ell_z(h)]$.

For a learning task \mathcal{T} , our typical goal is to design a *learning rule* A that maps a sequence of examples $S \in Z^n$ to a hypothesis $A(S) \in H$. The goal is to design a learning rule that competes with the best concept in C, such that for every distribution $D \in P$:

$$\mathop{\mathbb{E}}_{S \sim D^n} [L_D(A(S))] \le \operatorname{OPT}_C(D) + o(1),$$

where $OPT_C(D) = \inf_{c \in C} L_D(c)$, and the o(1) term converges to 0 as $n \to \infty$.

A distribution D over Z is called *realizable* if there exists $c \in C$ such that $L_D(c) = 0$; in particular, $OPT_C(D) = 0$. We also say that D is α -realizable, for $\alpha \ge 0$, if $OPT_C(D) \le \alpha$. We say that the task \mathcal{T} is *realizable* if P is the family of all realizable distributions, and is *agnostic* if it is the family of all distributions over Z. Unless specified otherwise, we assume the agnostic setting. As a remark, let us note that the complexity of designing the learning algorithm A is decreasing in H (more options for an output), increasing in C (competing against a bigger class), and increasing in P (more potential inputs).

Example 2 (PAC-learning). In PAC learning model, $C \subseteq \{\pm 1\}^X$ is a class of binary classifiers over a domain X that we want to learn. The labeled examples are $Z = X \times \{\pm 1\}$, and $\ell_{(x,y)}(h) = 1[h(x) \neq y]$ is the 0/1 classification loss. P is the set of realizable distributions in the *realizable* case and the set of all distributions in the *agnostic* case. Finally, H is usually taken to be all functions $H = \{\pm 1\}^X$ restricting it to H = C corresponds to *proper* PAC learning.

Example 3 (PAC-learning for partial concept classes). This model generalizes PAC-learning by letting C be a class of partial binary functions, that is, $C \subseteq \{-1, +1, *\}^X$, where * is treated as "undefined". For $c \in C$, the support $\sup(c)$ is the subset of X on which c is defined. Note, however, that the labeled examples are still $Z = X \times \{\pm 1\}$, that is, we do not allow examples to be undefined. The loss function is defined similarly, that is, for a partial function h on X, $\ell_{(x,y)}(h) = 1[h(x) \neq y]$. In particular, for an example (x, y), the case h(x) = * is treated as a mistake.

The main difference with the classical PAC learning comes from the class of distributions P. Note that now a realizable distribution D should be supported by some partial concept $c \in C$ in the sense that $\mathbb{P}_{x,y\sim D}[x \in \sup(c)] = 1$ (assuming X is finite; in the infinite case, there might be additional subtleties). For example, if every concept in C is supported on only half of the domain, then the same is true for any realizable distribution. This contrasts with canonical PAC, where "typical" distributions are supported on the whole X. Just as before, H is typically the set of all functions on X, and the agnostic case is defined similarly.

One particularly useful (and classical) example of PAC learning with partial concept classes is the class of linear classifiers with margins, which we consider in Theorems 5 and 6.

Example 4 (Stochastic convex optimization). In this setup, H = C is a convex set, which we will call W; Z is an abstract set such that every $z \in Z$ is equipped with a convex function $\ell_z : W \to \mathbb{R}_{\geq 0}$, and P is the set of all distributions over Z (i.e., agnostic setting). All in all, Z might be viewed simply as an index set of a set of convex loss functions, and the goal of the learning algorithm is, having a sample of such functions, to find a point in W that would minimize the expected loss of the given distribution over them.

We will be interested in a *dimension* of an SCO task. For this, we will assume that $W \subseteq \mathbb{R}^d$ and refer to the learning task as to *stochastic convex optimization in* \mathbb{R}^d .

Example 5 (General setting of learning). The general setting of learning, introduced by Vapnik [1998, 1999], can be modeled as a learning task. In this setup, H = C and Z are arbitrary sets, with P being the set of all distributions over Z. In general, there are no assumptions on the loss functions $\ell_z : H \to \mathbb{R}_{\geq 0}$.

In many cases within this framework, we assume the existence of a set of examples X and a set of labels Y, such that $Z = X \times Y$, and the hypothesis space H to be a subset of functions from X to Y (i.e., $H \subseteq Y^X$). In this case, the loss functions take the form $\ell_{(x,y)}(h) = L(h(x), y)$, where $L: Y^2 \to \mathbb{R}_{\geq 0}$ is a fixed loss function that compares the predicted label h(x) with the true label y.

For a learning task $\mathcal{T} = (H, C, Z, P)$ and $\alpha \geq 0$, we say that $h \in H$ is α -optimal for $D \in P$ if $L_D(h) \leq \mathsf{OPT}_C(D) + \alpha$. Note that for a PAC-learning task, 0-optimality that is witnessed by $c \in C$ is equivalent to realizability.

Definition 6 (Reductions). Let $\mathcal{T}_1 = (H_1, C_1, Z_1, P_1)$ and $\mathcal{T}_2 = (H_2, C_2, Z_2, P_2)$ be two learning tasks, and let $\alpha > 0$ and $\beta \ge 0$. An (α, β) -reduction r from \mathcal{T}_1 to \mathcal{T}_2 consists of two maps $r_{in}: Z_1 \to Z_2$ and $r_{out}: H_2 \to H_1$ such that the following holds.

- 1. For every distribution D_1 in P_1 , the distribution $r_{in}(D_1)$ is in P_2 . Here $r_{in}(D_1)$ is the push-forward measure induced by sampling $z \sim D_1$ and mapping it to $r_{in}(z)$.
- 2. For every $D_1 \in P_1$ and $h_2 \in H_2$, if h_2 is α -optimal for $r_{in}(D_1)$ then $r_{out}(h_2)$ is β -optimal for D_1 .

A reduction r is called <u>exact</u> if for every distribution D_1 , realizable by \mathcal{T}_1 , the push-forward distribution $r_{in}(D_1)$ is realizable by \mathcal{T}_2 .

In the above definition, we do not allow $\alpha = 0$ as, potentially, because $OPT_C(\cdot)$ is defined using infimum, a learning task might not contain 0-optimal solutions, in which case a notion of $(0, \beta)$ -reduction would trivialize; however, allowing $\beta = 0$ does not lead to such situations. Note that an

 (α, β) -reduction is an (α', β') reduction for any $\alpha' \leq \alpha$ and $\beta' \geq \beta$. Figure 2 below illustrates how (α, β) -reduction aligns to the approach outlined in Figure 1.



Figure 2: An (α, β) -reduction.

2.2 Reductions to stochastic convex optimization

Reductions from classification problems to stochastic convex optimization (SCO) are a common algorithmic tool, exemplified by the use of surrogate losses (e.g., hinge loss), regularization techniques, kernel methods, and other methods that rely on representing data in Euclidean spaces. In this section, we study the minimum dimension in which a concept class H can be reduced to an SCO task. Our main theorem provides a lower bound on the minimum Euclidean dimension required for such reductions, expressed in terms of the VC and dual VC dimension of the problem.

Theorem 1 (Binary classification vs. stochastic convex optimization). Let $C \subseteq \{\pm 1\}^X$ be a binary concept class. If for some $\beta < 1/2$ and $\alpha > 0$ there exists an (α, β) -reduction from the task of learning C in the realizable case to a stochastic convex optimization task in \mathbb{R}^d , with loss functions $\{\ell_z\}_{z\in Z}$ satisfying $\ell_z(w) < \infty$ for all $z \in Z$, $w \in W$. Then

$$d \ge \max\{ \mathsf{VC}(C), \mathsf{VC}^{\star}(C) - 1 \}.$$

Perhaps surprisingly, Theorem 1 implies that certain VC classes require an exponential increase in dimension for learning by reduction to SCO. See Example 15 for a simple class that exhibits this property. We discuss more classes that witness boundary cases of Theorem 1 in Section 4.1.

It is also worth noting that the conclusion in Theorem 1 extends to certain cases where the loss functions are ∞ -valued. Below, we provide a natural example of such an ∞ -valued SCO task (Example 8) and present an adaptation of the theorem for these reductions (Theorem 2).

From a technical perspective, relating the VC dimension to the Euclidean dimension in SCO is subtle, as they capture fundamentally different aspects of learning complexity: in PAC learning, the VC dimension directly governs sample complexity, whereas in SCO, sample complexity is decoupled from the Euclidean dimension and instead depends on factors such as the Lipschitz constant and the diameter of the parameter space.¹ Instead, the Euclidean dimension in SCO is closely related to other resources, such as space complexity, the complexity of arithmetic operations, and test-time complexity.

VC dimension vs. number of parameters. There is a natural, intuitive interpretation of the VC dimension as the number of parameters needed to encode concepts in a class, and this intuition is supported by natural classes such as half-spaces and axis-aligned boxes. However, it is known that if one allows general parametrization schemes, this interpretation does not hold. A standard example illustrating this is the class $\{x \mapsto \operatorname{sign}(\sin(tx)) : t \in \mathbb{R}\}$, which has an unbounded VC dimension despite being parameterized by a single parameter t. (See also Example 18.) In contrast, Theorem 1 shows that when the parametrization is restricted to be convex, the number of parameters required must be at least as large as the VC dimension, and in some cases, it must be exponentially larger, see Example 15 in Section 4.1 for details. This demonstrates that the VC dimension imposes a meaningful lower bound when parametrizations are constrained to a convex setting.

¹For example, there are one-dimensional SCO problems with an unbounded diameter or Lipschitz constant that are not learnable and infinite-dimensional SCO problems with a bounded diameter and Lipschitz constant that are learnable. (See, e.g. Cutkosky [2024]; Theorems 3.2 and 16.7)

Moreover, the open question posed in the previous paragraph—whether the minimal dimension *d* for which an SCO-reduction exists can be *upper bounded* in terms of the VC dimension—is also of interest here. If such an upper bound exists, it would provide a formal manifestation of the intuition that the VC dimension corresponds to the number of parameters needed to encode and learn the class.

Half-spaces. The following examples (Examples 7 and 8) concern classical reductions from *d*-dimensional linear classification (i.e. learning half-spaces) to stochastic convex optimization in *d*-dimensions.

A half-space is a concept $c_{w,b} : \mathbb{R}^d \to \{\pm 1\}$, parametrized by a pair $w \in \mathbb{R}^d$ and $b \in \mathbb{R}$ and defined as $c_{w,b}(x) = \text{sign}(\langle w, x \rangle + b)$. A half-space is called *homogenous* if b = 0 and *affine* in the general case; if not specified otherwise, we consider half-spaces to be homogenous. Learning half-spaces is a fundamental task in learning theory, and Support Vector Machines (SVMs) are a classical family of algorithms for this problem. SVMs aim to find a consistent half-space whose supporting hyperplane maximizes the margin from the data points. Formally, for each $x \in \mathbb{R}^d$ and $y \in \{\pm 1\}$, the *regularized hinge loss* is defined by

$$\ell_{x,y}(w) = \lambda \cdot ||w||^2 + \max(0, 1 - y\langle w, x \rangle),$$

where $\lambda \ge 0$ is a regularization parameter. In some cases, the regularized hinge loss is alternatively defined as

$$\ell_{x,y}(w) = ||w||^2 + c \cdot \max(0, 1 - y \langle w, x \rangle),$$

where $c \ge 0$ is a regularization parameter. Both definitions are equivalent up to a multiplicative constant, by setting $\lambda = 1/c$. Thus, SVMs can be viewed as a reduction from learning half-spaces to stochastic convex optimization, where the loss function is the regularized hinge loss.

In practice, SVMs are used with regularization parameters in $(0, \infty)$. A key advantage that makes SVMs practical is that the regularized hinge loss can be efficiently minimized even on non-separable data (i.e., data from non-realizable distributions). Theoretically, SVMs provably learn separable distributions with a margin between positive and negative regions. The next examples show that, in the limit as $\lambda \to \infty$ and $c \to 0$, SVMs reduce *d*-dimensional half-spaces to SCO, even without margin assumptions. These limiting cases are often called *hard SVM*. Interestingly, taking λ to infinity and *c* to zero results in different reductions.

Example 7 (SVM with unregularized hinge loss). Let $W = \mathbb{R}^{d+1}$, $Z = \mathbb{R}^d \times \{\pm 1\}$. Define the unregularized hinge loss by

$$\ell_{x,y}(w,a) = \max\left(0, 1 - y(\langle w, x \rangle + a)\right).$$

This defines the unregularized hinge loss SCO task, which is an SCO problem with continuous loss functions but over a non-compact domain W. It corresponds to taking $\lambda \to 0$ in the first formula for the regularized hinge loss. Define the reduction r from the task of learning homogenous halfspaces in \mathbb{R}^d to the above SCO by

$$r_{in}(x,y) = (x,y),$$

$$r_{out}(w,a) = c_{w,a},$$

where $c_{w,a}(x) = \operatorname{sign}(\langle w, x \rangle + a)$. In Section 4.2 we show that for all $\alpha > 0$, r is an (α, α) -reduction.

Example 8 (Hard SVM). Let $W = \mathbb{R}^d$, $Z = \mathbb{R}^d \times \{\pm 1\}$ and define the linear programming loss function by

$$\ell_{x,y}(w) = \begin{cases} ||w||^2 & \text{sign}(\langle x, w \rangle) = y \\ \infty & \text{otherwise.} \end{cases}$$

It corresponds to taking $c \to \infty$ in the second formula for the regularized hinge loss. Note that while in general $\ell_{x,y}$ is neither continuous nor finite, it is convex. In fact, it can be considered a limit case of the regularized hinge loss function

$$\ell_{x,y}(w) = ||w||^2 + c \cdot \max(0, 1 - y\langle w, x \rangle)$$

where $c = \infty$. This example illustrates how ∞ values are useful when one wishes to express hard constraints on the parameter space within the objective function. This allows to express linear programming as an SCO task. The reduction r from the learning homogenous halfspaces to this SCO is then defined in the same way as in Example 7. In Section 4.2 we show that, for all $\alpha > 0$, this is an $(\alpha, 0)$ -reduction.

Theorem 1 does not apply to the last example because the loss functions in the SCO problem are ∞ -valued. Since, as illustrated by the example above, ∞ -valued SCO tasks are natural, we prove the next theorem that applies to *exact* reductions to such tasks.

Theorem 2 (A variant of Theorem 1 for ∞ -valued SCO). Let $C \subseteq \{\pm 1\}^X$ be a binary concept class. Assume that there exists an <u>exact</u> reduction from the task of learning C in the realizable case with randomized hypotheses to a stochastic convex optimization task in \mathbb{R}^d with ∞ -valued loss functions $\ell_z : W \to \mathbb{R}_{\geq 0} \cup \{\infty\}$. Then,

 $d \geq \max\{\operatorname{VC}(C) - 1, \operatorname{VC}^*(C) - 1\}.$

2.3 Geometric representations

In this section, we focus on a special type of reductions called *representations*. This notion is quite natural and, within the learning theory, variants of this definition were studied, for example, in Pitt and Warmuth [1990], Kamath, Montasser, and Srebro [2020], Aliakbarpour, Bairaktari, Brown, Smith, Srebro, and Ullman [2024].

Definition 9 (Representations). Let $C_1 \subseteq \{\pm 1\}^{X_1}$ and $C_2 \subseteq \{\pm 1\}^{X_2}$ be two concept classes and let $\alpha \geq 0$. For $\alpha \geq 0$, an α -representation of C_1 by C_2 is a map $r : X_1 \to X_2$ such that for every distribution D_1 , realizable by C_1 , $r(D_1)$ is α -realizable by C_2 .² The representation r is called exact if it is a 0-representation, that is, it maps realizable distributions into realizable.

In particular, representations are indeed reductions.

Proposition 10 (Representations are reductions). Let $r: X_1 \to X_2$ be an α -representation of C_1 by C_2 . Then for $r_{in}(x, y) = (r(x), y)$ and $r_{out}(h)(x) = h(r(x))$, the pair (r_{in}, r_{out}) is a $(\gamma, \gamma + \alpha)$ -reduction of the task of PAC learning C_1 in a realizable case to PAC-learning C_2 , for any $\gamma \ge 0$.

The representations in the above sense were defined in Pitt and Warmuth [1990]. One of the most popular reductions, especially in the context of studying the expressivity of kernel methods, is the reduction to the class of half-spaces. The dimensionality of this reduction is also known as a *sign-rank*. We follow Kamath et al. [2020] in its formal definition below (see their Definition 1). Note that there the authors consider this definition with different loss functions and use terms *dimension complexity* for the generalized situation and *sign-rank* specifically for the case of 0/1 loss.

Definition 11 (Sign-rank). For a class $C \subseteq \{\pm 1\}^X$, the sign-rank of C is the smallest d for which there exists an exact representation of C by homogenous half-spaces in \mathbb{R}^d . That is, such that there are maps $\varphi \colon X \to \mathbb{R}^d$ and $w \colon C \to \mathbb{R}^d$ such that for all $c \in C$ and $x \in X$, it holds $\operatorname{sign}(w(c), \varphi(x)) = c(x)$.

Sign-rank and the reductions to SCO are closely related. Examples 7 and 8 above show that a class of sign-rank at most d can be reduced to SCO in dimension d. The following result shows that, for finite classes and exact reductions, this almost (with a factor of +1) goes in the other direction. We note that +1 can be removed if we allow representations by affine halfspaces, rather than with only homogenous, and so in the affine case this connection becomes tight.

²Here, with an abuse of notation, we extend r from X_1 to $X_1 \times \{\pm 1\}$ by letting r(x, y) = (r(x), y), and further extend it to distributions D_1 over $X_1 \times \{\pm 1\}$ by pushing-forward: sampling from $r(D_1)$ amounts to sampling $(x, y) \sim D_1$ and outputting r(x, y).

Theorem 3 (Half-spaces are complete for exact reductions to SCO). Let $C \subseteq \{\pm 1\}^X$ be a finite concept class. If for some $\beta < 1/2$ and α there exists an exact (α, β) -reduction from the task of learning C in the realizable case to a stochastic convex optimization task in \mathbb{R}^d , then C has an exact representation by homogenous half-spaces in \mathbb{R}^{d+1} . In other words, the sign-rank of C is at most d + 1.

Note that our setup naturally enables us to consider approximate reductions and representations, where we understand "approximate" as "not exact". Note that for representations it simply means letting $\alpha > 0$, while in the context of reductions, it means something else, which does not boil down to the parameters α and β . Formally:

Definition 12 (Approximate sign-rank and SCO dimension). For a class C and $\alpha < 1/2$ we define an α -sign-rank of C as the smallest d for which C has an α -representation by homogenous half-spaces in \mathbb{R}^d . Similarly, we define an SCO-dimension of C as the smallest d for which there is an (α, β) reduction, for some $\beta < 1/2$, from the realizable learning of C to an SCO task in \mathbb{R}^d .

While Theorem 3 says that, in the exact case, the SCO dimension and the sign-rank are essentially equivalent, we know much less about the approximate case. With respect to (approximate) SCO versus the (exact) sign-rank, from the above, we know that

 $\max{\{VC, VC^{\star} - 1\}} \leq SCO \leq sign-rank.$

It is also known that there are classes with constant VC dimension and unbounded sign-rank [Ben-David et al., 2002, Alon et al., 2017]. However, we do not know if SCO can be substantially smaller than the sign rank, and whether it can be upper bounded by a function of the VC dimension or lower bounded by a function of the sign-rank.

At the same time, the following theorem establishes a separation between the exact and approximate sign-ranks.

Theorem 4 (Exact vs approximate sign-rank). For every integer $d \ge 0$, there exists a finite concept class C_d whose (1/3)-sign-rank is at most d, and the sign-rank is at least $d^{\Omega(\log d)}$.

Note that the class in Theorem 4 is rather simple: its domain is \mathbb{R}^d and the functions are majority votes of the signs of three homogenous halfspaces in \mathbb{R}^d .

A natural way to generalize representations is to allow them to be probabilistic.

Definition 13 (Randomized representations). A δ -confident randomized α -representation of C_1 by C_2 , or (δ, α) -representation for short, is a distribution R over maps $r: X_1 \to X_2$ such that for every distribution D_1 , realizable by C_1 , the distribution $r(D_1)$ is α -realizable by C_2 with probability at least $1-\delta$ over $r \sim R$.

As before, the most canonical representation is the one by half-spaces in \mathbb{R}^d , which gives rise to the definition of a randomized sign-rank^{*}

Definition 14 (Probabilistic sign-rank^{*}). For a class C, $\delta > 0$, and $\alpha < 1/2$, we define a probabilistic δ -confident α -sign-rank^{*} of C, or (δ, α) -sign-rank^{*} for short, as the smallest d for which C has a (δ, α) -representation by homogenous half-spaces in \mathbb{R}^d .

Similar notions of *probabilistic dimension complexity* and *probabilistic sign-rank* were studied in the same paper [Kamath et al., 2020], see their Definitions 2 and 23. The latter is also a well-known term in communication complexity, see [Alman and Williams, 2017]. However, due to an additional swerve space from randomization, neither of them is identical to ours; in particular, we write sign-rank^{*} to separate our notion from the established term in an adjacent area.

Theorems 5 and 6 below establish a separation between the probabilistic and the deterministic sign-ranks for a partial class of linear classifiers over \mathbb{S}^n with margin. Note that the bounds on the probabilistic sign-rank in Theorem 5 are applicable (and stated) not only for our sign-rank^{*}, but also for the two of its abovementioned relatives, see the discussion and the definitions in Section 8.

Theorem 5 (Probabilistic sign-rank of halfspaces with margin). Let C_n be the partial class of linear classifiers with constant margin $\gamma = 1/3$ on the n-dimensional sphere \mathbb{S}^n . Then for $\alpha \in (0, 1/2)$ and $\delta \in (0, 1)$, the (δ, α) -sign-rank^{*} d of C_n is at most

$$d = O\left(\log\frac{1}{\alpha\delta}\right).$$

The randomized (δ, α) -representation witnessing it is linear, that is, the respective distribution is over linear maps $\mathbb{S}^n \to \mathbb{R}^d$, where we treat \mathbb{S}^n as a unit sphere in \mathbb{R}^{n+1} .

Moreover, probabilistic δ -dimension complexity and probabilistic δ -sign-rank are at most $O(\log(1/\delta))$.

Theorem 6 (Deterministic sign-rank of halfspaces with margin). Let C_n be the partial class of linear classifiers with constant margin $\gamma = 1/3$ on the n-dimensional sphere \mathbb{S}^n . Then for $\alpha \in (0, 1/2)$ the (deterministic) α -sign-rank d of C_n is at least

$$d \ge \min \Big\{ \frac{1-\alpha}{\alpha}, n+1 \Big\}.$$

Moreover, if the respective α -representation is continuous, then $d \ge n+1$.

In Kamath et al. [2020], the authors ask whether there is an "infinite" separation between the probabilistic dimension complexity and the sign-rank. Modulo the fact that our class C_n is partial, the above separation gives a positive answer to this question. Indeed, by Theorem 5, the δ -dimension complexity of the family of classes C_n above is uniformly bounded, independently of n. At the same time, by Theorem 6, the (exact deterministic) sign-rank of C_n is unbounded (although the case of exact sign-rank is also known from Theorem 1.5 in Hatami et al. [2022]). Moreover, let $C = \bigcup_{n=1}^{\infty} C_n$; here we abuse the fact that our classes are partial and that C is defined on a disjoint union of domains of the classes C_n . This way, a formal union of (δ, α) -representations of C_n 's gives a (δ, α) -representation of C. Hence, C is a partial class of bounded δ -dimension complexity and infinite sign-rank.

At the same time, many natural questions about the sign-rank and its relaxations can further be asked. We know that, for fixed $\alpha \in (0, 1/2)$ and $\delta \in (0, 1)$

 (δ, α) -sign-rank^{*} $\leq \alpha$ -sign-rank \leq sign-rank.

As argued, for a fixed δ and α , there is an infinite separation between the first and the last one. Moreover, for a fixed δ , there is an exponential in $1/\alpha$ separation between the first and the second. It is, however, left open whether any two of the "adjacent" ranks above are finite together. In particular, we conjecture that, for a fixed α , the constant lower bound in Theorem 6 is too weak, and the α -signrank of C_n goes to infinity with n. This, in particular, would imply an infinite separation between the probabilistic and the approximate sign-ranks.

2.4 A variant of Borsuk-Ulam for closed convex relations

Our proof of Theorem 1 uses a version of the Borsuk-Ulam theorem that combines the classical topological approach with convexity considerations. Recall that the original Borsuk-Ulam theorem states that a sphere \mathbb{S}^d cannot be continuously mapped to \mathbb{R}^d without collapsing a pair of antipodal points. Borsuk-Ulam theorem is a well-established tool in combinatorics, brought to light by Lovász's

proof of Kneser conjecture [Lovász, 1978], but which since then developed into a central tool in topological combinatorics, see [Matoušek, Björner, Ziegler, et al., 2003]. More recently, it has found applications in TCS and learning theory, see Hatami et al. [2022], Chase, Chornomaz, Hanneke, Moran, and Yehudayoff [2024a], Chase, Chornomaz, Moran, and Yehudayoff [2024b].

Theorem 7 (Borsuk-Ulam for closed convex relations). Let W be a compact convex set in \mathbb{R}^k and let G be a closed set inside $\mathbb{S}^d \times W$, where \mathbb{S}^d is a d-dimensional unit sphere. Additionally, suppose that:

- For any $x \in \mathbb{S}^d$, the set $G_x = \{w \in W \mid (x, w) \in G\}$ is nonempty and convex;
- For any $x \in \mathbb{S}^d$, the sets G_x and G_{-x} are disjoint.

Then $k \ge d+1$.

The original Borsuk-Ulam theorem can be seen as a special case of Theorem 7 by setting $G = (x, f(x)) : x \in \mathbb{S}^d$ as the graph of f and W as the convex hull of the image of f.

While formally for the proof of Theorem 1 we use not Theorem 7 itself, but rather its close relative Theorem 9, the latter is technical and its statement is less self-sufficient.

3 Proof overview

Most of the proofs use, in one or another way, the standard toolset of the learning theory, such as the VC dimension, the sign-rank, loss estimates, etc., see Section 5 below for a brief overview. In particular, the central definitions that we introduce and explore, namely, learning task (Definition 1), reduction (Definition 6), and representation (Definition 9), are quite typical, although to our knowledge, our formalization of them is new. Below, we outline the use of techniques that are either uncommon for this area, or that are in some sense specialized.

The main technical ingredient in the proof of reduction from classification to SCO (Theorem 1) is a version of Borsuk-Ulam for closed convex relations (Theorem 7). The proof of the latter relies on the classical Borsuk-Ulam together with some tools from convex geometry (Carathéodory's theorem) and real analysis (partitions of unity), although the proof can be carried out within a more topological toolset by using dense triangulations instead of the last two. The proof of Theorem 1 itself then uses the topological approach to PAC-learning problems similar to the one in Chase et al. [2024b], see the proof of Theorem D there. Namely, the set of realizable distributions of a class is topologized by equipping it with a total variation metric and the antipodality that comes from flipping the signs of the labels. The application of a Borsuk-Ulam-like theorem then hinges on two crucial observations: i) that no hypothesis can simultaneously achieve loss < 1/2 on two antipodal distributions, and ii) that both VC and dual VC dimensions enforce the existence of spheres of comparable dimension in the space of realizable distributions.

We would like to remark that deriving Theorem 1 does not seem to follow from standard samplecomplexity-based considerations. While this result lower-bounds the dimension of the SCO task in terms of the VC dimension of the class, reduced to it, it would be interesting to determine whether, conversely, the minimal dimension d for which a VC class H is reducible to SCO in \mathbb{R}^d can be *upper bounded* in terms of the VC dimension. Additionally, it would be interesting to explore whether a more elementary proof of Theorem 1 can be found.

The completeness of halfspaces for exact reductions to SCO (Theorem 3) is proven using LP duality for a certain game played on the space W, associated with the SCO task. The result about exact reductions to ∞ -valued SCO (Theorem 2) is an easy consequence of it and of the fact that both VC and dual VC lower-bound the sign-rank Alon et al. [2017].

The separation between exact and approximate sign-ranks(Theorem 4) is witnessed by a specific class, whose approximate sign-rank is easily upper-bounded, and sign-rank was shown to be high by Bun, Mande, and Thaler [2021]. For the other separation, between the probabilistic and approximate sign-ranks (Theorems 5 and 6), the probabilistic sign rank of the class in question is upper-bounded using a relative of the Johnson-Lindenstrauss theorem, and its approximate sign-rank is lower-bounded by combining uniform convergence (see Theorem 8 below) with the result in Hatami et al. [2022], establishing the sign-rank for this class; interestingly enough, the latter is also proved by a topological argument that uses Borsuk-Ulam.

4 Examples

4.1 Examples demonstrating the tightness of Theorem 1

The following is an example of a simple class for which the dual VC dimension, and hence the SCO dimension, is exponential in the VC dimension.

Example 15 (Projection functions). Let d be the dimension, and consider the domain $X = \{\pm 1\}^d$ and the class $U_d = \{h_i : i \leq d\}$ of all projection functions defined by $h_i(x_1, \ldots, x_d) = x_i$ on $(x_1, \ldots, x_d) \in X$. This class satisfies: (i) $VC(U_d) \leq \log d$, (ii) $VC^*(U_d) = d$, and (iii) U_d is reducible to stochastic convex optimization in \mathbb{R}^d . This is because U_d can be represented by half-spaces in \mathbb{R}^d : we embed Xinto \mathbb{R}^d trivially (since $X \subseteq \mathbb{R}^d$) and represent each concept h_i as the half-space $(x_1, \ldots, x_d) \mapsto \operatorname{sign}(x_i)$.

From the PAC learning perspective, β -optimality for $\beta < 1/2$ corresponds to weak learnability. That is, a hypothesis h which is (< 1/2)-optimal for a distribution D provides some information about D. On the other hand, 1/2-optimality can always be achieved by a random guess, and hence carries no information about the distribution. In the world of reductions, this is paralleled by the next two examples that show that a learning task can always be reduced to a 1-dimensional SCO for $\beta = 1/2$, as long we allow our hypotheses to be randomized.

Example 16 (PAC-learning with randomized hypotheses). In the setup of Example 2, it is often useful to allow a learner to use randomized hypotheses. We model this by defining $H = [-1,1]^X$ where a randomized hypothesis $h: X \to [-1,1]$ on x outputs 1 with probability $\frac{1+h(x)}{2}$ and -1 with probability $\frac{1-h(x)}{2}$. The loss function is then defined by $\ell_z(h) = \frac{1}{2}|h(x) - y|$, which corresponds to the expected 0/1 loss of h.

Note that $\ell_z(h)$ on a "deterministic" hypothesis h with values in $\{\pm 1\}$ is just a usual 0/1 loss, and hence this model generalizes the classical PAC setting. While this approach is less used in the PAC-learning paradigm, this generalization is common, for example, in the online learning setting, see Section 1.2.2 in Shalev-Shwartz [2012]. For us, it is useful in the context of reductions from PAC-learning to stochastic convex optimization, see Example 17 below.

Example 17 (A trivial reduction). This example demonstrates that any concept class $C \subseteq \{\pm 1\}^X$ can be reduced to a one-dimensional stochastic convex optimization problem with a classification error of $\beta = 1/2$. This highlights the tightness of the assumption $\beta < 1/2$ in Theorem 1.

Consider the task of learning a concept class $C \subseteq \{\pm 1\}^X$ in the realizable setting using a randomized learning rule. So, $H = [-1, 1]^X$, where each $h \in H$ represents a random hypothesis that outputs 1 with probability $\frac{1+h(x)}{2}$ and -1 with probability $\frac{1-h(x)}{2}$. The loss is the expected zero-one loss, $\ell_{(x,y)}(h) = \frac{1}{2}|h(x) - y|$.

We reduce to a stochastic convex optimization task with W = [-1, 1], $Z = \{\pm 1\}$, and $\ell_z(w) = \frac{1}{2}|z - w|$. The reduction is defined by $r_{in}(x, y) = y$ and $r_{out}(w) = h_w$, where $h_w(x) = w$ for all $x \in X$. We claim that for $\alpha \in [0, 1/2]$, the above is an $(\alpha, \frac{1+\alpha}{2})$ -reduction. In particular, for $\alpha = 0$ it is a $(\alpha = 0, \beta = \frac{1}{2})$ -reduction.

Proof. Let D_1 be a distribution realizable by C and $D_2 = r_{in}(D_1)$. Notice that $L_{D_1}(h_w) = L_{D_2}(w)$ for all $w \in [-1, +1]$. Let $w^* \in W$ be α -optimal with respect to D_2 . We need to show that $r_{out}(w^*) = h_{w^*}$ is $\frac{1+\alpha}{2}$ -optimal with respect to D_1 . Define $p_+ = \mathbb{P}_{(x,y)\sim D_1}[y=+1]$ and $p_- = \mathbb{P}_{(x,y)\sim D_1}[y=-1]$. For all $w \in W$:

$$L_{D_2}(w) = \frac{p_+|1-w| + p_-|1+w|}{2} = \frac{1-(p_+-p_-)w}{2}$$

Optimizing for the above yields that minimum of L_{D_2} on W is $\min\{p_+, p_-\}$. Hence, since w^* is α optimal for D_2 , then $L_{D_1}(h_{w^*}) = L_{D_2}(w^*) \leq \alpha + \min\{p_+, p_-\}$. And since the maximum of L_{D_2} is $\max\{p_+, p_-\}$, this bound can be improved to

$$L_{D_1}(h_{w^*}) \le \min\{\alpha + \min\{p_+, p_-\}, \max\{p_+, p_-\}\} \le \frac{\alpha + \min\{p_+, p_-\} + \max\{p_+, p_-\}}{2} = \frac{1+\alpha}{2}.$$

The following example demonstrates that if we remove the requirement for convex loss functions, any finite concept class can be reduced to a one-dimensional learning task with continuous loss functions.

Example 18 (A non-convex reduction to one dimension). Consider the task of learning a finite concept class $C \subseteq \{\pm 1\}^X$ in the realizable setting using a randomized learning rule. Define the following learning task: Let $Z = X \times \{\pm 1\}, W = [0, 1]$, and for each $c \in C$ pick some unique $w_c \in W$. For each example $(x, y) \in Z$ define the set $V_{x,y} = \{w_c \in W : c(x) = y\}$. Now define the loss function $\ell_{x,y}$ by

$$\ell_{x,y}(w) = \frac{d(w, V_{x,y})}{d(w, V_{x,y}) + d(w, V_{x,-y})}$$

Note that $\ell_{x,y}$ is continuous (but not convex). Define the following reduction from C to the above learning task

$$r_{in}(x,y) = (x,y),$$

$$r_{out}(w)(x) = \operatorname{sign}(\ell_{x,-1}(w) - \ell_{x,1}(w)).$$

We claim that for all $\alpha > 0$ the above is an exact $(\alpha, 2\alpha)$ -reduction.

Proof. The above reduction is exact because $L_{r_{in}(D)}(w_c) = 0$ for any distribution D which is realizable by $c \in C$. To show it is $(\alpha, 2\alpha)$ -reduction it is enough to prove that

$$L_D^{01}(r_{\text{out}}(w)) \le 2L_{r_{\text{in}}(D)}(w).$$

Note that $\ell_{x,-1}(w) + \ell_{x,1}(w) = 1$. Thus, $r_{out}(w)(x) \neq y$ implies $l_{x,y}(w) \geq 1/2$. But then

$$\begin{split} L_D^{01}\big(r_{\texttt{out}}(w)\big) &= \mathop{\mathbb{E}}_{(x,y)\sim D} \left[r_{\texttt{out}}(w)(x) \neq y\right] \\ &\leq \mathop{\mathbb{E}}_{(x,y)\sim D} 2l_{x,y}(w) = 2 \mathop{\mathbb{E}}_{(x,y)\sim r_{\texttt{in}}(D)} l_{x,y}(w) = 2L_{r_{\texttt{in}}(D)}(w), \end{split}$$

as needed.

4.2 Reductions to SVM

In this subsection, we prove statements about the reductions to SVM from Example 7 and Example 8. We restate the above examples in an abridged form, to make apparent the claims that we prove.

Example 7 (SVM with unregularized hinge loss). This SVM is an SCO task with $W = \mathbb{R}^{d+1}$, $Z = \mathbb{R}^d \times \{\pm 1\}$, and, for $(w, a) \in \mathbb{R}^d \times \mathbb{R} = W$,

$$\ell_{x,y}(w,a) = \max\left(0, 1 - y(\langle w, x \rangle + a)\right).$$

Define the reduction r from the task of learning affine halfspaces in \mathbb{R}^d to this task by

$$r_{in}(x, y) = (x, y),$$

$$r_{out}(w, a) = c_{w,a},$$

where $c_{w,a}(x) = \operatorname{sign}(\langle w, x \rangle + a)$. Then for any $\alpha > 0$, r is an (α, α) -reduction.

Proof. Recall that $\ell_{x,y}^{01}(w,a) = [\operatorname{sign}(\langle w,x\rangle+a) \neq y]$. In particular, $\ell_{x,y}^{01}(w,a) = 1$ means that $y(\langle w,x\rangle+a) \geq 0$ and hence $\ell_{x,y}(w,a) \geq 1$. So, for any realizable D and $w, a \in W$, $L_D^{01}(c_{w,a}) \leq L_D(w,a)$. Thus, to prove the validity of the reduction, it suffices to show that $\operatorname{OPT}_W(D) = 0$ for all D realizable by d-dimensional half-spaces.

Let D be a distribution realizable by $c_{w,a}$ and let $\varepsilon > 0$, we claim that for n large enough we have

$$L_D(nw, na+1) < \varepsilon.$$

Indeed,

$$\begin{split} L_D(nw, na+1) &= \mathop{\mathbb{E}}_{x, y \sim D} \max\left(0, 1 - y(\langle nw, x \rangle + na + 1)\right) \\ &\leq \mathop{\mathbb{E}}_{x, y \sim D} \max\left(0, 1 - y(\langle nw, x \rangle + na + 1)\right) \\ &= \mathop{\mathbb{E}}_{x, y \sim D} \max\left(0, 1 - y(\langle nw, x \rangle + na + 1)\right) \\ &= \left[y(\langle w, x \rangle + a) < 1 - y\right] \\ &= \left[y(\langle w, x \rangle + a) \geq 0 \text{ for } x, y \sim D\right] \\ &= \mathop{\mathbb{E}}_{x, y \sim D; \ y = -1} \max\left(0, 1 + 1(\langle nw, x \rangle + na + 1)\right) \\ &= \mathop{\mathbb{E}}_{x, y \sim D; \ y = -1} \max\left(0, 2 + n(\langle w, x \rangle + a)\right) \leq \mathop{\mathbb{E}}_{x, y \sim D; \ y = -1} 2 \\ &= \mathop{\mathbb{E}}_{x, y \sim D; \ y = -1} \max\left(0, 2 + n(\langle w, x \rangle + a)\right) \leq \mathop{\mathbb{E}}_{x, y \sim D; \ y = -1} 2 \\ &= 2 \mathop{\mathbb{E}}_{x, y \sim D} \left[-2/n < \langle w, x \rangle + a \leq 0 \text{ and } y = -1 \leq 0\right] \\ &\xrightarrow{n \to \infty} 2 \mathop{\mathbb{E}}_{x, y \sim D} \left[\langle w, x \rangle + a = 0 \text{ and } y = -1 \leq 0\right] = 0. \end{split}$$

Example 8 (Hard SVM). This SVM is an SCO task with $W = \mathbb{R}^d$, $Z = \mathbb{R}^d \times \{\pm 1\}$, and

$$\ell_{x,y}(w) = \begin{cases} ||w||^2 & \text{sign}(\langle x, w \rangle) = y, \\ \infty & otherwise. \end{cases}$$

Define the reduction r from the task of learning homogenous halfspaces in \mathbb{R}^d to this task by

$$\begin{split} r_{\texttt{in}}(x,y) &= (x,y), \\ r_{\texttt{out}}(w) &= c_w, \end{split}$$

where $c_w(x) = c_{w,0}(x) = \text{sign}(\langle w, x \rangle)$. Then for any $\alpha > 0$, r is an $(\alpha, 0)$ -reduction.

Proof. Note that for every distribution D which is realizable by half-spaces and every $w \in W$ we have that $L_{r_{in}(D)}(w)$ is $||w||^2$ if $\mathbb{P}_{(x,y)\sim D}[\text{sign}(\langle x,w\rangle) = y] = 1$ and infinite otherwise. Hence, for any finite α , every α -optimal solution for L_D achieves a loss of zero on the zero-one loss function L_D^{01} . Thus, r is indeed an $(\alpha, 0)$ -reduction for all $\alpha > 0$. And, by rescaling w we get that $\mathsf{OPT}_W(D) = 0$ so the reduction is exact.

5 Preliminaries

We use standard notation from learning theory, see e.g. Shalev-Shwartz and Ben-David [2014]. In general, X will denote the domain and C a concept class of functions from X to $\{\pm 1\}$. For a given collection of loss functions $\{\ell_z\}_{z\in Z}$ for each distribution D over Z and finite sample $S \subseteq Z$ we define the induced loss functions

$$L_D(w) = \mathop{\mathbb{E}}_{z \sim D} \ell_z(w),$$

$$L_S(w) = \frac{1}{|S|} \sum_{z \in Z} \ell_z(w).$$

Whenever there can be some confusion with other loss functions, we will use $\ell_{x,y}^{01}(c)$ (and similarly L_D^{01}) to denote the zero-one loss function which is 0 if c(x) = y and 1 otherwise.

Definition 19 (VC dimension). We say that a set $\{x_i\}_{i=1}^n$ is shattered by a concept class C if for any labeled sample $S = \{(x_i, y_i)\}_{i=1}^n$ over this set there is some $c \in C$ such that $L_S(c) = 0$. The VC dimension VC(C) of C is the largest number n such that C shatters a set of size n, or infinity if C shatters sets of arbitrary size.

Definition 20 (Dual VC dimension). For a fixed concept class C over a domain X, each $x \in X$ defines a function $h_x : C \to \{\pm 1\}$ by $h_x(c) = c(x)$. The class $C^* = \{h_x : x \in X\}$ is called a dual class of C. The dual VC dimension of C is the VC dimension of C^* , $VC^*(C) = VC(C^*)$.

We say that a concept class C satisfies uniform convergence if there exists a vanishing sequence $\epsilon_n \xrightarrow{n \to \infty} 0$ such that for all distributions D over $X \times \{\pm 1\}$,

$$\mathop{\mathbb{E}}_{S \sim D^n} \left(\sup_{c \in C} |\mathsf{L}_D(c) - \mathsf{L}_S(c)| \right) \le \epsilon_n.$$

The seminal work by Vapnik and Chervonenkis [1971] shows that any concept class C with finite VC dimension satisfies the uniform convergence. The following asymptotically optimal quantitative bound was achieved in the seminal work by Talagrand [1994] using a technique called chaining [Dudley, 1978].

Theorem 8 (Uniform convergence for VC classes). Let C be a concept class with finite VC dimension VC(C) = d. Then for any n > 0 and a distribution D over $X \times \{\pm 1\}$ we have

$$\mathop{\mathbb{E}}_{S \sim D^n} \left(\sup_{c \in C} |L_D(c) - L_S(c)| \right) = O\left(\sqrt{d/n}\right)$$

6 Proofs for reductions to SCO

Our aim for this section is to prove Theorem 1 and Theorem 3. However, a central technical ingredient here is Theorem 9 below, which is a close relative of the Borsuk-Ulam for closed convex relations (Theorem 7), but with a more technical statement. For accessibility, in this and the following sections we restate the theorems that we are going to prove.

Theorem 9 (Borsuk-Ulam for certain relations). Let W be a compact convex set in \mathbb{R}^k and let $G \subseteq \mathbb{S}^d \times W$. Additionally, suppose that:

- For any $x \in \mathbb{S}^d$, the set $G_x = \{w \in W \mid (x, w) \in G\}$ is nonempty;
- For any $x \in \mathbb{S}^d$, the convex hulls of G_x and G_{-x} are disjoint;
- For any $w \in W$, the set $G^w = \{x \in \mathbb{S}^d : (x, w) \in G\}$ is open.

Then
$$k \ge d+1$$
.

Proof. Note that $x \in G^w$ if and only if $y \in G_x$, and, as G_x is nonempty for all x, the family $\{G^w\}_{w \in W}$ is an open cover of \mathbb{S}^d . By compactness, there is a finite $T \subset W$ such that $\{G^t\}_{t \in T}$ is a finite open cover of \mathbb{S}^d . Let $\{\rho_t\}_{t \in T}$ be a partition of unity subordinate to this cover, that is, a family of continuous functions parameterized by $t \in T$ such that each ρ_t is 0 outside of G^t and for all $x \in \mathbb{S}^d$, $\sum_{t \in T} \rho_t(x) = 1$. Partitions of unity are a well-known tool in real analysis, see Theorem 2.13 in Rudin [1987]. We can also explicitly define

$$\rho_t(x) = \frac{d(x, \mathbb{S} \setminus G^t)}{\sum_{t' \in T} d(x, \mathbb{S} \setminus G^{t'})}.$$

Define $\Phi : \mathbb{S}^d \to W$ by

$$\Phi(x) = \sum_{t \in T} \rho_t(x) t.$$

Note that $\rho_t(x) > 0$ implies that $t \in G_x$, so $\Phi(x)$ is a convex combination of elements in G_x . Then, as $\Phi(x)$ and $\Phi(-x)$ are in the convex hulls of G_x and G_{-x} respectively and the latter are disjoint, $\Phi(x) \neq \Phi(-x)$. By the Borsuk-Ulam theorem, this implies $k \ge d+1$, as needed.

Lemma 21. Let $I \subset \mathbb{R}^d$ be an nonempty collection of pointwise positive vectors in \mathbb{R}^d , that is, such that $\alpha_i > 0$ for all $\alpha \in I$ and i = 1, ..., d. Let $X = \{x \in [0, 1]^d : \sum_{i=1}^d x_i = 1\}$ and define $F : X \to \mathbb{R}$ by

$$F(x) = \inf_{\alpha \in I} \langle x, \alpha \rangle.$$

Then F is continuous.

Proof. For $x \in X$, let us take an arbitrary $\varepsilon > 0$ and let $\delta > 0$ to be specified later. Define U_x to be an open neighborhood of x containing the points y such that $|x_i - y_i| < \delta x_i$ whenever $x_i \neq 0$ and for $y_i < \delta$ whenever $x_i = 0$. Let $y \in U_x$, we will show that for δ small enough $|F(x) - F(y)| < 2\varepsilon$. Fix some $\alpha \in I$ such that

$$\langle x, \alpha \rangle < F(x) + \varepsilon.$$

First, we bound F(y) - F(x). Note that

$$\langle y, \alpha \rangle - \langle x, \alpha \rangle \le \sum_{i=1}^{d} |x_i - y_i| \alpha_i \le \sum_{x_i \ne 0} \delta x_i \alpha_i + \sum_{x_i = 0} \delta \alpha_i \le \delta (F(x) + \varepsilon + ||\alpha||_1).$$

So,

$$F(y) \le \langle y, \alpha \rangle \le \langle x, \alpha \rangle + \delta \big(F(x) + \varepsilon + ||\alpha||_1 \big) \le F(x) + \varepsilon + \delta \big(F(x) + \varepsilon + ||\alpha||_1 \big).$$

Now we bound F(x) - F(y). Fix some $\beta \in I$ such that

$$\langle y, \beta \rangle \le F(y) + \varepsilon.$$

Note that

$$\langle x,\beta\rangle - \langle y,\beta\rangle \le \sum_{x_i\neq 0} |x_i - y_i|\beta_i - \sum_{x_i=0} y_i\beta_i \le \sum_{x_i\neq 0} \delta x_i\beta_i \le \frac{\delta}{1-\delta} \sum_{i=1}^a y_i\beta_i \le \frac{\delta}{1-\delta} F(y).$$

Hence,

$$F(x) \le \langle x, \beta \rangle \le \langle y, \beta \rangle + \frac{\delta}{1 - \delta} F(y) \le F(y) + \varepsilon + \frac{\delta}{1 - \delta} \Big(F(x) + \delta \big(F(x) + \varepsilon + ||\alpha||_1 \big) \Big).$$

So taking $\delta > 0$ such that $\frac{\delta}{1-\delta} \left(F(x) + \delta \left(F(x) + \varepsilon + ||\alpha||_1 \right) \right) < \varepsilon$ will imply that

$$|F(x) - F(y)| \le 2\varepsilon.$$

Theorem 1 (Binary classification vs. stochastic convex optimization). Let $C \subseteq \{\pm 1\}^X$ be a binary concept class. If for some $\beta < 1/2$ and $\alpha > 0$ there exists an (α, β) -reduction from the task of learning C in the realizable case to a stochastic convex optimization task in \mathbb{R}^d , with loss functions $\{\ell_z\}_{z \in Z}$ satisfying $\ell_z(w) < \infty$ for all $z \in Z$, $w \in W$. Then

$$d \ge \max\{ \mathsf{VC}(C), \mathsf{VC}^{\star}(C) - 1 \}.$$

We are going to prove Theorem 1 in a slightly more general setting, namely, we allow the learning task to have *randomized hypotheses*, see Example 16 for the definition. This is not a huge generalization, but it provides conformity with subsequent Example 17, which illustrates that the above statement trivializes if we allow $\beta \geq 1/2$.

Proof. Let $r = (r_{in}, r_{out})$ be an (α, β) -reduction from C to a stochastic convex optimization task with a convex loss function $L: W \times Z \to \mathbb{R} \ge 0$, where $W \subseteq \mathbb{R}^d$ is a convex set. To lower bound d in terms of VC(C) and $VC^*(C)$, we can assume, without loss of generality, that C is a finite class over a finite domain X. We achieve this by replacing C with a finite subclass that has the same VC and dual VC dimensions. Consequently, we also assume Z is finite by focusing on $r_{in}(X \times \{\pm 1\})$. Let P be the collection of C-realizable distributions over $X \times \{\pm 1\}$, equipped with the total variation metric

$$\mathsf{TV}(D,D') = \sup_A |D(A) - D'(A)|.$$

where the supremum is over all measurable events A. We define the involution $D \rightarrow -D$ on the set of distributions over Z as

$$-D(A) = D(\{(x, -y) : (x, y) \in A\}).$$

Note that P is not, in general, closed under this involution. That is, -D is not necessarily in P even if D is. We say that $S \subseteq P$ is an n-sphere if there is a homeomorphism $\varphi : \mathbb{S}^n \to S$ such that $\varphi(-u) = -\varphi(u)$ for all $u \in \mathbb{S}^n$. We are now going to show that the existence of an n-sphere implies that $d \ge n + 1$.

Let $S \subseteq P$ with a homeomorphism φ be such *n*-sphere and define $G \subseteq \mathbb{S}^n \times W$ as

$$G = \{(u,w) \ : L_{r_{\mathrm{in}}\left(\varphi(u)\right)}(w) < \mathrm{OPT}_W\big(r_{\mathrm{in}}(\varphi(u))\big) + \alpha/2\}.$$

That is, for $u \in \mathbb{S}^n$, the set $G_u = \{w \in W : (u, w) \in G\} \subseteq W$ is the set of all $w \in W$ witnessing the $\alpha/2$ -optimal loss with respect to $L_{r_{in}(\varphi(u))}$. In order to use Theorem 7 we now need to show that $G_u = \{w \in W : (u, w) \in G\} \subseteq W$ is nonempty, that its convex hull is disjoint from the convex hull of G_{-u} , and that $G^w = \{u \in \mathbb{S}^n : (u, w) \in G\}$ is open.

As $L_{r_{in}(\varphi(u))}$ is a convex function, it trivially follows that G_u is nonempty and convex. We now want to check that for all $u \in \mathbb{S}^n$ the sets G_u and G_{-u} are disjoint.

Indeed, as r is an (α, β) -reduction, $r_{out}(w)$ is β -optimal for $\varphi(x)$ whenever w is α -optimal for $r_{in}(\varphi(u))$; in particular, $L^{01}_{\varphi(u)}(r_{out}(w)) \leq \beta$ for $w \in G_u$. Also, for any $D \in P$, and a randomized hypothesis $h \in [-1, 1]^X$ we have:

$$\begin{split} L_D^{01}(h) + L_{-D}^{01}(h) &= \mathop{\mathbb{E}}_{x,y\sim D} \frac{1}{2} \left| h(x) - y \right| + \mathop{\mathbb{E}}_{x,y\sim -D} \frac{1}{2} \left| h(x) - y \right| \\ &= \mathop{\mathbb{E}}_{x,y\sim D} \frac{1}{2} \left(\left| h(x) - y \right| + \left| h(x) + y \right| \right) = \mathop{\mathbb{E}}_{x,y\sim D} \frac{1}{2} \cdot 2 = 1. \end{split}$$

So for any $w \in G_u$ it holds

$$L^{01}_{\varphi(-u)}(r_{\text{out}}(w)) = 1 - L^{01}_{\varphi(u)}(r_{\text{out}}(w)) \ge 1 - \beta > 1/2 > \beta,$$

and $w \notin G_{-u}$.

We now need to show that $G^w = \{u \in \mathbb{S}^n : (w, u) \in G\}$ is open. Define $p : \mathbb{S}^n \to \mathbb{R}^{|Z|}$ by $p(u) = (p_z(u))_{z \in Z}$ where

$$p_z(u) = \mathop{\mathbb{P}}_{(x,y) \sim \varphi(u)} (z = r_{\text{in}}(x,y)).$$

Note that p is continuous by properties of the total variation metric. Thus if we set $\alpha_w = (\ell_z(w))_{z \in Z} \in \mathbb{R}^{|Z|}$ we get

$$\begin{split} L_{\operatorname{rin}\left(\varphi(u)\right)}(w) &= \sum_{z \in Z} p_z(u)\ell_z(w) = \langle p(u), \alpha_w \rangle, \\ \operatorname{OPT}_W\Big(\operatorname{rin}\left(\varphi(u)\right) \Big) &= \inf_{w \in W} \langle p(u), \alpha_w \rangle. \end{split}$$

Now, by Lemma 21, for all $w \in W$ the function $u \to L_{r_{in}(u)}(w) - OPT_W(r_{in}(u))$ is continuous. Thus, G^w is indeed open.

It can now be seen that G satisfies the condition of Theorem 7, and so $d \ge n+1$. Finally, we refer the reader to the proof of Theorem D in Chase et al. [2024b] for the following statement regarding the existence of spheres: **Proposition 22.** For a class $C \subseteq \{\pm 1\}^X$, there is an n-sphere in the space of realizable distributions of C whenever $n \leq \max{\{VC(C) - 1, VC^*(C) - 2\}}$.

Combining Proposition 22 with $d \ge n+1$, we get $g \ge \max{\{VC(C) - 1, VC^*(C) - 2\}} + 1 = \max{\{VC(C), VC^*(C) - 1\}}$, as needed.

Theorem 2. Let $C \subseteq \{\pm 1\}^X$ be a binary concept class. Assume that there exists an <u>exact</u> reduction from the task of learning C in the realizable case with randomized hypotheses to a stochastic convex optimization task in \mathbb{R}^d with ∞ -valued loss functions $\ell_z : W \to \mathbb{R}_{\geq 0} \cup \{\infty\}$. Then,

$$d \ge \max\{\mathsf{VC}(C) - 1, \mathsf{VC}^*(C) - 1\}.$$

Proof. By Theorem 3 (see the proof in Section 7) the sign-rank of C is at most d + 1. By Alon et al. [2017], both VC(C) and $VC^*(C)$ lower-bound the sign-rank from which the statement of the theorem follows.

6.1 Borsuk-Ulam for closed convex relations

Lemma 23. Let G be as in Theorem 7 and let us define $A_{x,\delta}$, for $x \in \mathbb{S}^d$ and $\delta > 0$, as a convex hull of the set $\{(x', w) \in G : d(x, x') < \delta\}$. Then for any $\varepsilon > 0$ there exist $\delta > 0$ such that for any $x \in \mathbb{S}^d$ and $x' \in A_{x,\delta}$, x' is ε -close to G.

Proof. By Carathéodory's theorem, it is enough to show that any convex combination of d + 1 points from $\{(x', w) \in G : d(x, x') < \delta\}$ is at most ε away from G. Let $\Lambda = \{\lambda \in [0, 1]^{d+1} : \sum_{i=1}^{d} \lambda_i = 1\}$ and define the function $F : G^d \times \Lambda \to \mathbb{R}$ by

$$F(g,\lambda) = d\left(G, \sum_{i=1}^{d+1} \lambda_i g_i\right).$$

By construction, F is continuous and, as $G^d \times \Lambda$ is compact, it is uniformly continuous. Now, for $g = (g_1, g_2, \dots, g_{d+1}) \in G^{d+1}$, where $g_i = (x_i, w_i)$, note that if for some $x' \in \mathbb{S}^d$, all $x_i = x'$, then $w_i \in G_{x'}$ for all i and, by convexity of $G_{x'}$, $\sum_{i=1}^d \lambda_i w_i \in G_{x^*}$ for all $\lambda \in \Lambda$. Hence in this case $F(g, \lambda) = 0$ for all λ . Thus, by the uniform continuity of F, for any $\varepsilon > 0$ there is some $\delta > 0$ such that if $x_1, x_2 \dots, x_{d+1}$ are in a δ -ball around $x' \in \mathbb{S}^d$ then $F(x, \lambda) < \varepsilon$ for all $\lambda \in \Lambda$, which concludes the proof.

Theorem 7 (Borsuk-Ulam for closed convex relations). Let W be a compact convex set in \mathbb{R}^k and let G be a closed set inside $\mathbb{S}^d \times W$, where \mathbb{S}^d is a d-dimensional unit sphere. Additionally, suppose that:

- For any $x \in \mathbb{S}^d$, the set $G_x = \{w \in W \mid (x, w) \in G\}$ is nonempty and convex;
- For any $x \in \mathbb{S}^d$, the sets G_x and G_{-x} are disjoint.

Then $k \geq d+1$.

Proof. Assume towards contradiction that we have such G with $d \leq k$. Let $-G = \{(-x, w) : (x, w) \in G\}$. Note that as $G_x \cap G_{-x} = \emptyset$ for all x, we have $-G \cap G = \emptyset$ and, since both are closed, d(G, -G) > 0. That is, there is some $\varepsilon > 0$ such that $d(g, g') > \varepsilon$ for all $g \in G, g' \in -G$.

Let $\delta > 0$ be a small constant to be chosen later. For each $x \in \mathbb{S}^d$ let $B_{\delta}(x)$ be the ball with radius δ centered at x. By compactness we have some finite $T \subset \mathbb{S}^d$ such that $\{B_{\delta}(t)\}_{t \in T}$ is an open cover of \mathbb{S}^d . Similarly to the proof of Theorem 9, let $\{\rho_t\}_{t \in T}$ be a partition of unity subordinate to this cover.

Now for each $t \in T$ choose some $w_t \in G_t$ in an arbitrary way, and define $\chi : \mathbb{S}^d \to \mathbb{R}^{d+1}$, $\phi : \mathbb{S}^d \to W$, and $\Phi = (\chi, \phi) : \mathbb{S}^d \to \mathbb{R}^{d+1} \times W$ as

$$\chi(x) = \sum_{t \in T} \rho_t(x)t,$$

$$\phi(x) = \sum_{t \in T} \rho_t(x)w_t.$$

Note that in order to define $\chi(x)$ we assume that the unit sphere \mathbb{S}^d is canonically embedded into \mathbb{R}^{d+1} .

As $\rho_t(x) = 0$ whenever $d(x,t) > \delta$, $\chi(x)$ is a convex combination of points inside $B_{\delta}(x)$. As the latter is convex, $d(\Phi(x), (x, \phi(x))) = d(\chi(x), x) \leq \delta$ for all $x \in \mathbb{S}^d$. Also, $\Phi(x)$ is in the convex hull of $\{(x', w) \in G : d(x, x') \leq \delta\}$, so, by Lemma 23, for $\delta > 0$ small enough, $\Phi(x)$ is $(\varepsilon/4)$ -close to G. Additionally, assuming $\delta \leq \varepsilon/4$, for any $x \in \mathbb{S}^d$ we get

$$d(G,(x,\phi(x))) \le d(G,\Phi(x)) + d(\Phi(x),(x,\phi(x))) \le \frac{\varepsilon}{2}.$$

By the definition of -G, $d(G, (-x, \phi(x))) \leq \varepsilon/2$. As x is arbitrary from \mathbb{S}^d , by changing x to -x we get $d(G, (x, \phi(-x))) \leq \varepsilon/2$.

Now, since ϕ is continuous, $d \leq k$ implies, by the Borsuk-Ulam theorem, that there is some $x \in \mathbb{S}^d$ such that $\phi(x) = \phi(-x)$. For such x we have

$$d(G, -G) \le d(G, (x, \phi(x))) + d(-G, (x, \phi(x)))$$

= $d(G, (x, \phi(x))) + d(-G, (x, \phi(-x))) \le \varepsilon$,

which contradicts the fact that $d(G, -G) > \varepsilon$.

7 Proofs for geometric representations

The statement of Proposition 10 below is made more formal than in the main part where it was originally formulated.

Proposition 10 (Representations are reductions). Let $\mathcal{T}_1 = (H_1, C_1, Z_1, P_1)$ be a realizable PAClearning task over domain X_1 , that is, $H_1 = \{\pm 1\}^{X_1}$, $C_1 \subseteq \{\pm 1\}^{X_2}$, $Z_1 = X_1 \times \{\pm 1\}$, and P_1 be the family of all distributions on Z_1 , realizable by C_1 .

Let $r: X_1 \to X_2$ be an α -representation of C_1 by a class $C_1 \subseteq \{\pm 1\}^{X_2}$. Let $\mathcal{T}_2 = (H_2, C_2, Z_2, P_2)$ be a PAC-learning task over X_2 , where $P_2 \supseteq r(P_1)$, that is, P_2 contains the images of all distributions in P_1 under r.

Let us define $r_{in}(x, y) = (r(x), y)$, for $x \in X_1$ and $y \in \{\pm 1\}$, and $r_{out}(h)(x) = h(r(x))$, for $x \in X_1$ and $h \in H_2$. Then the pair of maps r_{in} , r_{out} is a $(\gamma, \gamma + \alpha)$ -reduction for any $\gamma \ge 0$.

Proof. Let $D_1 \in P_1$ and $D_2 = r_{in}(D_1)$. By the condition on P_2 , $D_2 \in P_2$. Suppose $h_2 \in H_2$ is γ -optimal for D_2 , that is, $L_{D_2}(h_2) \leq \mathsf{OPT}_{C_2}(D_2) + \gamma$.

As \mathcal{T}_1 is realizable, so is D_1 . Hence, by the definition of α -representation, $OPT_{C_2}(D_2) \leq \alpha$, and so $L_{D_2}(h_2) \leq \alpha + \gamma$. Let $h_1 = r_{out}(h_2)$. Then

$$L_{D_1}(h_1) = \mathbb{P}_{x,y \sim D_1}[h_1(x) \neq y] = \mathbb{P}_{x,y \sim D_1}[h_2(r(x)) \neq y] = \mathbb{P}_{x,y \sim D_2}[h_2(x) \neq y] = L_{D_2}(h_2).$$

Thus, $L_{D_1}(h_1) = L_{D_2}(h_2) \leq \alpha + \gamma = \alpha + \gamma$. As D_1 is realizable, $OPT_{C_1}(D_1) = 0$, and so h_1 is $(\alpha + \gamma)$ -optimal, as needed.

Theorem 3 (Half-spaces are complete for exact reductions to SCO). Let $C \subseteq \{\pm 1\}^X$ be a finite concept class. If for some $\beta < 1/2$ and α there exists an exact (α, β) -reduction from the task of learning H in the realizable case to a stochastic convex optimization task in \mathbb{R}^d , then C has an exact representation by homogenous half-spaces in \mathbb{R}^{d+1} . In other words, the sign-rank of C is at most d+1.

Proof. Without losing generality, we can assume that for every $x, y \in X \times \{\pm 1\}$ there is some $c \in C$ such that c(x) = y; otherwise, we can restrict the domain to those x's for which this condition is satisfied, and the restricted class will be trivially equivalent to the original one. Recall that in the setup of an SCO task in \mathbb{R}^d we assume that the loss functions are defined on a convex set $W \subseteq \mathbb{R}^d$. We will further assume that this \mathbb{R}^d is embedded into \mathbb{R}^{d+1} as $\mathbb{R}^d = \{\overline{x} \mid x_{d+1} = 1\}$. The reason for this is that now every affine hyperplane in \mathbb{R}^d can be uniquely extended to a homogenous hyperplane in \mathbb{R}_{d+1} .

Fix some $c \in C$ and look at the following game: the player chooses some $w \in W$ and the adversary chooses some $x \in X$, then the player suffers a loss of $L_z(w)$ for $z = r_{in}(x, c(x))$. Strategies for the adversary are distributions over X, which are equivalent to distributions over $X \times \{\pm 1\}$, realizable by c. Since r is exact, for each c-realizable distribution D there is some $w \in W$ such that $L_{r_{in}(D)}(w) = 0$. Hence, by the minimax theorem, there is some $w_c \in W$ such that $L_{r_{in}(D)}(w_c) = 0$ for all such D. Note that we are using the fact that the class is finite for the applicability of the minimax.

Now for each $x, y \in X \times \{\pm 1\}$ define $V_{x,y}$ as a convex hull of the set $\{w_c : c \in C, c(x) = y\}$. By construction, $V_{x,y}$ is convex and compact, also, as we assumed that there is $c \in C$ with c(x) = y, it is nonempty. Moreover, as $L_{r_{in}(x,y)}$ is convex, nonnegative, and 0 on all w_c for $c \in C$ such that c(x) = y, it follows that $L_{r_{in}(x,y)}(w) = 0$ for $w \in V_{x,y}$.

As r is an (α, β) -reduction, for any $w \in V_{x,y}$ the 0/1 loss of $r_{out}(w)$ on the distribution which is supported only on (x, y) is at most $\beta < \frac{1}{2}$. Thus $r_{out}(w)(x) = y$, which implies that $V_{x,1} \cap V_{x,-1} = \emptyset$. As both are convex and compact, by the Hahn-Banach separation theorem, there is an affine hyperplane in \mathbb{R}^d strictly separating them. Recall that, by the choice of an embedding of \mathbb{R}^d into \mathbb{R}^{d+1} , we can extend this hyperplane to a homogenous hyperplane in \mathbb{R}^{d+1} . That is, for every $x \in X$, there is $\varphi_x \in \mathbb{R}^{d+1}$ such that sign $(\langle \varphi_x, w \rangle)$ is y for $w \in V_{x,y}$.

With this, the maps $\varphi : X \to \mathbb{R}^{d+1}$ and $w \colon C \to \mathbb{R}^{d+1}$ defined as $\varphi(x) = \varphi_x$ and $w(c) = w_c$ satisfy $\operatorname{sign}(\langle w(c), \varphi(x) \rangle) = c(x)$ for all $c \in C$ and $x \in X$, witnessing that the sign-rank of C is at most d+1.

Theorem 4 (Exact vs approximate sign-rank). For every integer $d \ge 0$, there exists a finite concept class C_d whose (1/3)-sign-rank is at most d, and the sign-rank is at least $d^{\Omega(\log d)}$.

Proof. Let H_d be the class of half-spaces in \mathbb{R}^d and let $C_d = \{ \mathsf{maj}(h_1, h_2, h_3) : h_1, h_2, h_3 \in H_d \}$ be the class of majority vote of three such half-spaces. It is easy to see that an identity map on \mathbb{R}^d is an (1/3)-representation of C_d by H_d . Indeed, let D be a distribution realizable by C_d and let $c = \mathsf{maj}(h_1, h_2, h_3)$ be an element in C_d with $L_D(c) = 0$. For $i \in \{1, 2, 3\}$, let A_i be the event that $h_i(x) = y$ for a random pair $(x, y) \sim D$. By definition of c, for any x at least two of the events A_1, A_2, A_3 will occur, hence

$$\mathbb{E}\left(1_{A_1} + 1_{A_2} + 1_{A_3}\right) \ge 2.$$

This trivially implies that $\mathbb{E}(1_{A_i}) \geq \frac{2}{3}$ for some $i \in \{1, 2, 3\}$. Equivalently, $L_D(h_i) \leq 1/3$, so the identity map is indeed a (1/3)-representation.

For the second part, in Corollary 1.2 in Bun et al. [2021] the authors prove that a class of 2intersections of certain signs of weighted majorities on a $4m^2$ -dimensional Boolean hypercube has sign-rank $m^{\Omega(\log m)}$. We note that that some details are not stated in Corollary 1.2 explicitly, but can be easily extracted from the proof of their Theorem 1.1. By embedding the hypercube into \mathbb{R}^{4m^2} in a standard way, and noting that a sign of a weighted majority on a hypercube can be expressed by a sign of a homogenous hyperplane, we get that for any d, the sign-rank of the class I_d of intersections of two half-spaces in \mathbb{R}^d is at least $d^{\Omega(\log d)}$. Note that Ω hides coefficients arising from going from $4m^2$ to an arbitrary d.

Finally, it is easy to see that the class I_d can be embedded into C_{d+1} , yielding the desired lower bound on the sign-rank of C_d

8 Several probabilistic sign-ranks and proofs of Theorems 5 and 6

Before proving Theorem 5, let us elaborate on several notions of probabilistic sign-ranks related to our work. We start with *probabilistic dimension complexity* see Definition 2 in Kamath et al. [2020]. Note that the authors consider it for different families of loss functions, but the definition below is specifically for 0/1-loss. We also give all definitions for *partial* concept classes and slightly align the notation in line with ours.

Definition 24 (Probabilistic dimension complexity, Kamath et al. [2020]). For a (partial) class C over domain X and $\delta > 0$, a probabilistic δ -dimension complexity $dc_C(\delta)$ of C is the smallest d for

which there is a distribution R over maps $r: X \to \mathbb{R}^d$ such that for all distributions D over $X \times \{\pm 1\}$, realizable by C, it holds

$$\mathop{\mathbb{E}}_{r \sim R} \left[\inf_{w \in \mathbb{R}^d} \mathop{\mathbb{P}}_{x, y \sim D} \bigl[\mathsf{sign} \langle w, r(x) \rangle \neq y \bigr] \right] \leq \delta.$$

By expanding the intermediate definitions, one can see that this is very close to how we defined the sign-rank^{*}:

Definition 14 (Probabilistic sign-rank^{*}, restated). For a (partial) class C over domain X, $\delta > 0$, and $\alpha < 1/2$, a (δ, α) -sign-rank^{*} $\operatorname{sr}^*_C(\delta, \alpha)$ of C is the smallest d for which there is a distribution R over maps $r: X \to \mathbb{R}^d$ such that for all distributions D over $X \times \{\pm 1\}$, realizable by C, it holds

$$\mathbb{P}_{r\sim R}\left[\inf_{w\in\mathbb{R}^d}\mathbb{P}_{x,y\sim D}\left[\mathsf{sign}\langle w,r(x)\rangle\neq y\right]>\alpha\right]\leq\delta.$$

The following mutual bounds between the two are rather trivial:

Proposition 25. For a (partial) class C, $\delta > 0$, and $\alpha < 1/2$, it holds:

$$\begin{aligned} & \operatorname{sr}^*(\delta/\alpha,\alpha) \leq \operatorname{dc}(\delta), \\ & \operatorname{dc}(\delta+\alpha(1-\delta)) \leq \operatorname{sr}^*(\delta,\alpha), \end{aligned}$$

where, for compliance with the definitions, in the first bound we additionally assume that $\delta < \alpha/2$.

Note that here and below, we drop the underscripts identifying the class in sr, dc, etc., whenever the class is clear from the context.

Proof. Note that the condition on the distribution R witnessing the respective d in Definitions 24 and 14 is stated as for all D, $\mathbb{E}_{r\sim R}[F_D(r)] \leq \delta$ for $dc(\delta)$ and $\mathbb{P}_{r\sim R}[F_D(r) > \alpha] \leq \delta$ for $sr^*(\delta, \alpha)$ respectively. Here, informally, $F_D(r)$ is a fit of the distribution r(D) to the class of half-spaces in \mathbb{R}^d , however, we only need that $F_D(r) \in [0, 1]$. By Markov's inequality, $\mathbb{P}[F_D(r) > \alpha] \leq \delta/\alpha$ whenever $\mathbb{E}[F_D(r) > \alpha] \leq \delta$, and so the same R witnessing $dc(\delta) \leq d$ also witnesses $sr^*(\delta/\alpha, \alpha) \leq d$, yielding the first inequality.

Note that Markov's inequality comes from the fact that the function F_D maximizing $\mathbb{P}[F_D(r) > \alpha]$ provided $\mathbb{E}[F_D(r)] \leq \delta$ is $F_D = \alpha + 0$ w.p. $\delta/\alpha - 0$ and $F_D = 0$ otherwise. In the same spirit, the function that maximizes $\mathbb{E}[F_D(r)]$ provided $\mathbb{P}[F_D(r) > \alpha] \leq \delta$ is $F_D = 1$ w.p. δ and $F_D = \alpha$ otherwise. From this, $\mathbb{E}[F_D(r)] \leq \delta + \alpha(1-\delta)$ whenever $\mathbb{P}[F_D(r) > \alpha] \leq \delta$, yielding, in a similar way, the second inequality.

It is natural to compare the above two definitions to a canonical notion of *probabilistic sign-rank* from communication complexity. Its formulation below is from Definition 23 in Kamath et al. [2020], where it is called *point-wise probabilistic dimension complexity*.

Definition 26 (Probabilistic sign-rank). For a (partial) class C over domain X and $\delta > 0$, a probabilistic δ -sign-rank $\operatorname{sr}_C(\delta)$ of C is the smallest d for which there is a distribution R over pairs of maps $(r: X \to \mathbb{R}^d, \omega: C \to \mathbb{R}^d)$ such that for all $h \in C$ and $x \in \sup(h)$, it holds

$$\mathbb{E}_{r,\omega \sim R} \Big[\mathsf{sign} \langle \omega(h), r(x) \rangle \neq h(x) \Big] \leq \delta.$$

Finally, let us define the following two relaxations of sign-rank, that we will use in the proof of Theorem 5. We will not give them descriptive names, as they are purely technical and used only to give a uniform proof for all of the three versions of sign-rank above.

Definition 27. For a (partial) class C over domain X, $\delta > 0$, and $\alpha < 1/2$, let $\operatorname{sr}_{C}^{\dagger}(\delta)$ and $\operatorname{sr}_{C}^{\dagger}(\delta, \alpha)$ be the smallest d for which there is a distribution R over pairs of maps $(r: X \to \mathbb{R}^{d}, \omega: C \to \mathbb{R}^{d})$ such that for all $h \in C$ and all distributions D, realizable by h, it holds

$$\mathbb{E}_{r,\omega \sim R} \left[\mathbb{P}_{x,y \sim D} \left[\mathsf{sign} \langle w(h), r(x) \rangle \neq y \right] \right] \leq \delta$$

for $\operatorname{sr}_{C}^{\dagger}(\delta)$ and

$$\mathbb{P}_{r,\omega \sim R}\left[\mathbb{P}_{x,y \sim D}\left[\mathsf{sign}\langle w(h), r(x) \rangle \neq y\right] \geq \alpha\right] \leq \delta$$

for $\operatorname{sr}_{C}^{\dagger}(\delta, \alpha)$.

Proposition 28. For a (partial) class C over domain X, $\delta > 0$, and $\alpha < 1/2$, it holds

$$\begin{split} \mathsf{sr}^{\dagger}(\delta/\alpha,\alpha) &\leq \mathsf{sr}^{\dagger}(\delta),\\ \mathsf{sr}^{\dagger}(\delta+\alpha(1-\delta)) &\leq \mathsf{sr}^{\dagger}(\delta,\alpha),\\ \mathsf{dc}(\delta) &\leq \mathsf{sr}(\delta) \leq \mathsf{sr}^{\dagger}(\delta), and\\ \mathsf{sr}^{*}(\delta,\alpha) &\leq \mathsf{sr}^{\dagger}(\delta,\alpha). \end{split}$$

where in the second bound we assume $\delta < \alpha/2$.

Proof. Note that the relation between $\operatorname{sr}^{\dagger}(\delta)$ and $\operatorname{sr}^{\dagger}(\delta, \alpha)$ is the same as between $\operatorname{dc}(\delta)$ and $\operatorname{sr}^{*}(\delta, \alpha)$, so the proof of first two inequalities is similar to Proposition 25. The fact that $\operatorname{dc}(\delta) \leq \operatorname{sr}(\delta)$ is by Proposition 24 in Kamath et al. [2020]. For $\operatorname{sr}(\delta) \leq \operatorname{sr}^{\dagger}(\delta)$, note that for R witnessing $\operatorname{sr}^{\dagger}(\delta) \leq d$, it holds

$$\begin{split} \sup_{h \in C} \sup_{x \in \sup(h)} & \mathbb{E}_{r, \omega \sim R} \Big[\mathsf{sign} \langle \omega(h), r(x) \rangle \neq h(x) \Big] \\ &= \sup_{h \in C} \sup_{\chi_{x,y} \ll h} \mathop{\mathbb{E}}_{r, \omega \sim R} \left[\mathop{\mathbb{P}}_{x, y \sim D} \Big[\mathsf{sign} \langle w(h), r(x) \rangle \neq y \Big] \right] \\ &\leq \sup_{h \in C} \sup_{D \ll h} \mathop{\mathbb{E}}_{r, \omega \sim R} \left[\mathop{\mathbb{P}}_{x, y \sim D} \Big[\mathsf{sign} \langle w(h), r(x) \rangle \neq y \Big] \right] \leq \delta, \end{split}$$

where $\chi_{x,y}$ is a one-point distribution of the example (x, y), and $D \ll h$ denotes that D is realizable by h. Thus, R also witnesses $\operatorname{sr}(\delta) \leq d$ and hence $\operatorname{sr}(\delta) \leq \operatorname{sr}^{\dagger}(\delta)$.

Finally, for R witnessing $sr^{\dagger}(\delta) \leq d$, the same R restricted to the first coordinate trivially witnesses $sr^{*}(\delta) \leq d$, and so $sr^{*}(\delta, \alpha) \leq sr^{\dagger}(\delta, \alpha)$.

We are now ready to prove Theorem 5.

Theorem 5 (Probabilistic sign-rank of halfspaces with margin). Let C_n be the partial class of linear classifiers with constant margin $\gamma = 1/3$ on the n-dimensional sphere \mathbb{S}^n . Then for $\alpha \in (0, 1/2)$ and $\delta \in (0, 1)$, the (δ, α) -sign-rank^{*} $d = \operatorname{sr}^*(\delta, \alpha)$ of C_n is at most

$$d = O\left(\log\frac{1}{\alpha\delta}\right).$$

The randomized (δ, α) -representation witnessing it is linear, that is, the respective distribution is over linear maps $\mathbb{S}^n \to \mathbb{R}^d$, where we treat \mathbb{S}^n as a unit sphere in \mathbb{R}^{n+1} .

Moreover, probabilistic δ -dimension complexity $dc(\delta)$ and probabilistic δ -sign-rank $sr(\delta)$ are at most $O(\log(1/\delta))$.

Proof. We will prove that for given α and δ , $\mathsf{sr}^{\dagger}(\delta \cdot \alpha) \leq O(\log 1/\alpha \delta)$. All the claimed bound then follow from the $\mathsf{sr}^*(\delta) \leq \mathsf{sr}^{\dagger}(\delta)$ and $\mathsf{dc}(\delta) \leq \mathsf{sr}^{\dagger}(\delta)$ bounds from Proposition 28.

Let d > 0, to be chosen later. Note that in this case the domain and the class are both equal to \mathbb{S}^n , and we need to construct a distribution R over pairs of maps $(r: \mathbb{S}^n \to \mathbb{R}^d, \omega: \mathbb{S}^n \to \mathbb{R}^d)$. In fact, we put both maps to be equal to the same random linear map $\mathbb{R}^{n+1} \to \mathbb{R}^d$ whose entries $R_{i,j}$ are independent normal distributions $\mathcal{N}(0, 1)$. We refer to Corollary 20 in Ben-David et al. [2002] in the case of a single half-space for the following statement: For any half-space w in \mathbb{R}^{n+1} with margin γ , and $x \in \mathbb{S}^n$ in its support, so $|\langle w, x \rangle| > \gamma$, it holds

$$\mathbb{P}_{r\sim R}[\operatorname{sign}(\langle r(w), r(x) \rangle) \neq \operatorname{sign}(\langle w, x \rangle)] \leq 4e^{\frac{-d\gamma^2}{8}}.$$

We note that the above statement is a close relative of the Johnson-Lindenstrauss lemma, where this particular construction of the random projection is from Arriaga and Vempala [2006]. In particular, as is usual for this lemma, this estimate does not depend on n. With this we get that for any distribution D realizable by the half-space with margin w we have

$$\mathbb{E}_{r \sim R} \left[\mathbb{P}_{(x,y) \sim D} \left[\mathsf{sign} \left(\langle r(w), r(x) \rangle \right) \neq y \right] \right] = \mathbb{P}_{(x,y) \sim D} \left[\mathbb{P}_{r \sim R} \left[\mathsf{sign} \left(\langle r(w), r(x) \rangle \right) \neq \mathsf{sign} \left(\langle w, x \rangle \right) \right] \right] \le 4e^{\frac{-d\gamma^2}{8}}.$$

But, by recalling that $\gamma = 1/3$ is a fixed constant, the above will be less than $\delta \cdot \alpha$ for $d = \frac{10}{\gamma^2} \log \frac{1}{\alpha \delta} = O(\log(1/\alpha \delta))$, as needed.

We will now go for the proof of Theorem 6, starting with the following lemma.

Lemma 29. For $d \in \mathbb{N}$ and $0 < \alpha < 1/(d+1)$, an α -representation of a (partial) concept class C by half-spaces in \mathbb{R}^d is an exact representation. In particular, if the α -sign-rank of C is at most $d = (1 - \alpha)/\alpha$, then the sign-rank of C is at most d.

Proof. Let $r : X \to \mathbb{R}^d$ be such α -representation, and for any D distribution over $X \times \{\pm 1\}$ let $L_D : \mathbb{R}^d \to [0, 1]$ be the induced loss function, i.e.

$$L_D(w) = \mathop{\mathbb{P}}_{(x,y)\sim D} \left(\mathsf{sign}(\langle w, r(x) \rangle) \neq y \right)$$

. For any $S \subseteq X \times \{\pm 1\}$, let U_S denote the uniform measure on S, and define the set $V_S \subset \mathbb{R}^d$ by

$$V_S = \bigcap_{(x,y)\in S} \{ w \in \mathbb{R}^d : \operatorname{sign}(\langle w, r(x) \rangle) = y \}.$$

Note that V_s is a convex set as an intersection of half-spaces.

Now, let $S \subseteq X \times \{\pm 1\}$ be a finite sample realizable by C. As r is an α -representation, for any $T \subseteq S$ of size at most d + 1 there is $w_T \in \mathbb{R}^d$ such that

$$\frac{1}{|T|} \big| \{ (x,y) \in T : \operatorname{sign} \langle w_T, r(x) \rangle \neq y \} \big| = L_{U_T}(w_T) < \alpha < \frac{1}{|T|}.$$

Hence, $\operatorname{sign}\langle w_T, r(x) \rangle = y$ for all $(x, y) \in T$, and so $w_T \in V_T$. Thus, any intersection of at most (d+1) sets $V_{\{(x,y)\}}$ for $(x,y) \in S$ is non-empty and, by Helly's theorem, the overall intersection V_S is non-empty. Therefore, for each finite S there exists $w_S \in \mathbb{R}^d$ such that $\operatorname{sign}\langle w_S, r(x) \rangle = y$ for all $(x,y) \in S$.

Now the class of half-spaces in \mathbb{R}^d is a learnable class with VC-dimension of d, hence it satisfies the uniform convergence principle and by Theorem 8 for any $\varepsilon > 0$ and distribution D realizable by C, we have that for all n > 0

$$\mathop{\mathbb{E}}_{S \sim D^n} L_D(w_S) = \mathop{\mathbb{E}}_{S \sim D^n} \left(|L_D(w_S) - L_S(w_S)| \right) \le \mathop{\mathbb{E}}_{S \sim D^n} \left(\sup_{w \in \mathbb{R}^d} |L_D(w) - L_S(w)| \right) = O\left(\sqrt{d/n}\right).$$

Hence, for any n > 0 there is some $w \in \mathbb{R}^d$ such that $L_D(w) = O(\sqrt{d/n})$, from which we deduce that OPT(D) = 0 and r is an exact representation.

Theorem 6 (Deterministic sign-rank of halfspaces with margin). Let C_n be the partial class of linear classifiers with constant margin $\gamma = 1/3$ on the n-dimensional sphere \mathbb{S}^n . Then for $\alpha \in (0, 1/2)$ the (deterministic) α -sign-rank d of C_n is at least

$$d \ge \min\left\{\frac{1-\alpha}{\alpha}, n+1\right\}.$$

Moreover, if the respective α -representation is continuous, then $d \ge n+1$.

Proof. Let $r : \mathbb{S}^n \to \mathbb{R}^d$ be an α -representation of C_n with $d < \frac{1-\alpha}{\alpha}$. By Lemma 29, r is an exact representation, so d is at least the sign-rank of C_n . However, by Theorem 1.5 in Hatami et al. [2022], the sign-rank of C_n is exactly n + 1.

Let us now assume that $\alpha \in (0, 1/2)$ and that the α -representation $r : \mathbb{S}^n \to \mathbb{R}^d$ is continuous. Towards contradiction, suppose $d \leq n$. Then, by the Borsuk-Ulam theorem, there is some $x \in \mathbb{S}^n$ such that r(x) = r(-x). Let D be the uniform distribution on (x, 1) and (-x, -1), which is clearly realizable by C. At the same time, trivially, for any $w \in \mathbb{R}^d$, $L_{r(D)}(w) = 1/2$ and so r(D) is not α -realizable for $\alpha < 1/2$.

References

- Maryam Aliakbarpour, Konstantina Bairaktari, Gavin Brown, Adam Smith, Nathan Srebro, and Jonathan R. Ullman. Metalearning with very few samples per task. In Shipra Agrawal and Aaron Roth, editors, The Thirty Seventh Annual Conference on Learning Theory, June 30 - July 3, 2023, Edmonton, Canada, volume 247 of Proceedings of Machine Learning Research, pages 46–93. PMLR, 2024. URL https://proceedings.mlr.press/v247/aliakbarpour24a.html.
- Josh Alman and Ryan Williams. Probabilistic rank and matrix rigidity. In *Proceedings of the 49th* Annual ACM SIGACT Symposium on Theory of Computing, pages 641–652, 2017.
- Noga Alon, Shay Moran, and Amir Yehudayoff. Sign rank versus Vapnik-Chervonenkis dimension. *Sbornik: Mathematics*, 208(12):1724, dec 2017. doi: 10.1070/SM8780. URL https://dx.doi.org/10.1070/SM8780.
- Rosa I. Arriaga and Santosh Vempala. An algorithmic theory of learning: Robust concepts and random projection. *Machine learning*, 63:161–182, 2006.
- Shai Ben-David, Nadav Eiron, and Hans-Ulrich Simon. Limitations of learning via embeddings in Euclidean half spaces. *Journal of Machine Learning Research*, 3:441-461, 2002. URL http://www.jmlr.org/papers/v3/bendavid02a.html.
- Mark Bun, Nikhil S. Mande, and Justin Thaler. Sign-rank can increase under intersection. ACM Transactions on Computation Theory (TOCT), 13(4):1–17, 2021.
- Zachary Chase, Bogdan Chornomaz, Steve Hanneke, Shay Moran, and Amir Yehudayoff. Dual VC dimension obstructs sample compression by embeddings. In COLT 2024 The 37th Annual Conference on Learning Theory, Edmonton, Canada, June 30-July 3, 2024, 2024a.
- Zachary Chase, Bogdan Chornomaz, Shay Moran, and Amir Yehudayoff. Local Borsuk-Ulam, stability, and replicability. In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing*, pages 1769–1780, 2024b.
- Ashok Cutkosky. Lecture notes: Optimization for machine learning, 2024. https://optmlclass.github.io/notes/optforml_notes.pdf.
- Richard M. Dudley. Central limit theorems for empirical measures. *The Annals of Probability*, 6(6): 899–929, 1978. doi: 10.1214/aop/1176995384.
- Jürgen Forster. A linear lower bound on the unbounded error probabilistic communication complexity. J. Comput. Syst. Sci., 65(4):612–625, 2002. doi: 10.1016/S0022-0000(02)00019-3. URL http://dx.doi.org/10.1016/S0022-0000(02)00019-3.
- Jürgen Forster, Matthias Krause, Satyanarayana V. Lokam, Rustam Mubarakzjanov, Niels Schmitt, and Hans-Ulrich Simon. Relations between communication complexity, linear arrangements, and computational complexity. In FST TCS 2001: Foundations of Software Technology and Theoretical Computer Science, 21st Conference, Bangalore, India, December 13-15, 2001, Proceedings, pages 171–182, 2001. doi: 10.1007/3-540-45294-X_15. URL http://dx.doi.org/10.1007/3-540-45294-X_15.

- Jürgen Forster, Niels Schmitt, Hans-Ulrich Simon, and Thorsten Suttorp. Estimating the optimal margins of embeddings in Euclidean half spaces. *Machine Learning*, 51(3):263–281, 2003. doi: 10.1023/A:1022905618164. URL http://dx.doi.org/10.1023/A:1022905618164.
- Hamed Hatami, Kaave Hosseini, and Xiang Meng. A Borsuk-Ulam lower bound for signrank and its application. *Electron. Colloquium Comput. Complex.*, TR22-130, 2022. URL https://eccc.weizmann.ac.il/report/2022/130.
- Pritish Kamath, Omar Montasser, and Nathan Srebro. Approximate is good enough: Probabilistic variants of dimensional and margin complexity. In Jacob D. Abernethy and Shivani Agarwal, editors, *Conference on Learning Theory, COLT 2020, 9-12 July 2020, Virtual Event [Graz, Austria]*, volume 125 of *Proceedings of Machine Learning Research*, pages 2236–2262. PMLR, 2020. URL http://proceedings.mlr.press/v125/kamath20b.html.
- Nathan Linial and Adi Shraibman. Learning complexity vs communication complexity. *Combinatorics*, *Probability & amp; Computing*, 18(1-2):227-245, 2009. doi: 10.1017/S0963548308009656. URL http://dx.doi.org/10.1017/S0963548308009656.
- László Lovász. Kneser's conjecture, chromatic number, and homotopy. Journal of Combinatorial Theory, Series A, 25(3):319–324, 1978.
- Jiří Matoušek, Anders Björner, Günter M Ziegler, et al. Using the Borsuk-Ulam theorem: lectures on topological methods in combinatorics and geometry, volume 2003. Springer, 2003.
- Leonard Pitt and Manfred K. Warmuth. Prediction-preserving reducibility. J. Comput. Syst. Sci., 41(3):430-467, 1990. doi: 10.1016/0022-0000(90)90028-J. URL https://doi.org/10.1016/0022-0000(90)90028-J.
- Walter Rudin. Real and complex analysis (3rd ed.). New York: McGraw-Hill, 1987. ISBN 978-0-07-054234-1.
- Shai Shalev-Shwartz. Online learning and online convex optimization. Foundations and Trends® in Machine Learning, 4(2):107–194, 2012. ISSN 1935-8237. doi: 10.1561/2200000018. URL http://dx.doi.org/10.1561/2200000018.
- Shai Shalev-Shwartz and Shai Ben-David. Understanding machine learning: From theory to algorithms. Cambridge university press, 2014.
- Shai Shalev-Shwartz, Ohad Shamir, Nathan Srebro, and Karthik Sridharan. Stochastic convex optimization. In COLT 2009 - The 22nd Conference on Learning Theory, Montreal, Quebec, Canada, June 18-21, 2009, 2009. URL http://www.cs.mcgill.ca/%7Ecolt2009/papers/018.pdf#page=1.
- Michel Talagrand. Sharper bounds for Gaussian and empirical processes. *The Annals of Probability*, 22(1):28–76, 1994. doi: 10.1214/aop/1176988847.
- Vladimir N. Vapnik. Statistical learning theory. Wiley-Interscience, 1998.
- Vladimir N. Vapnik. An overview of statistical learning theory. *IEEE transactions on neural networks*, 10(5):988–999, 1999.
- V.N. Vapnik and A.Ya. Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory Probab. Appl.*, 16:264–280, 1971. ISSN 0040-585X; 1095-7219/e. doi: 10.1137/1116025.