

ON THE MODEL THEORY OF OPEN INCIDENCE STRUCTURES: THE RANK 2 CASE

GIANLUCA PAOLINI AND DAVIDE EMILIO QUADRELLARO

ABSTRACT. Taking inspiration from [1, 21, 24], we develop a general framework to deal with the model theory of open incidence structures. In this first paper we focus on the study of systems of points and lines (rank 2). This has a number of applications, in particular we show that for any of the following \mathcal{C} :

- (1) (k, n) -Steiner systems (for $2 \leq k < n$);
- (2) generalised n -gons (for $n \geq 3$);
- (3) k -nets (for $k \geq 3$);
- (4) affine planes;
- (5) projective Möbius, Laguerre and Minkowski planes;

all the non-degenerate free structures in \mathcal{C} are elementarily equivalent and their common theory is decidable, strictly stable and with no prime model.

CONTENTS

| | |
|---|----|
| 1. Introduction | 2 |
| 2. Preliminaries and notations | 5 |
| 3. The general framework | 8 |
| 3.1. Finitary conditions | 8 |
| 3.2. HF-orderings | 10 |
| 3.3. The theory T^+ | 14 |
| 3.4. Completeness and stability | 15 |
| 3.5. Forking independence in T^+ | 20 |
| 3.6. Failure of superstability | 27 |
| 3.7. Non-existence of prime models | 29 |
| 4. Applications | 31 |
| 4.1. n -open graphs | 32 |
| 4.2. (k, n) -Steiner systems | 37 |
| 4.3. Generalised n -gons | 46 |
| 4.4. k -nets | 52 |
| 4.5. Affine planes | 56 |
| 4.6. Projective Möbius, Laguerre and Minkowski planes | 62 |
| References | 67 |

Date: December 3, 2024.

2020 *Mathematics Subject Classification.* 03C45, 03C65, 03C98, 05B30, 51B05, 51E05.

The authors were supported by project PRIN 2022 “Models, sets and classifications”, prot. 2022TECZJA. The first author was also supported by INdAM Project 2024 (Consolidator grant) “Groups, Crystals and Classifications”. The second author was also supported by an INdAM post-doc grant. We wholeheartedly thank John Baldwin for useful discussions related to this paper. We are also thankful to Domenico Zambella for comments on a prior draft of this paper.

1. INTRODUCTION

As witnessed by the 1420 pages book “Handbook of Incidence Geometry” [14], the subject of incidence geometry is a vast one, with interactions with various other parts of mathematics, such as algebra, combinatorics, geometry and topology. In this article we develop an abstract framework to deal with the model theory of open incidence structures, and we use it to give model theoretic applications to cases of incidence geometries of interest, focusing here on systems of points and lines.

Originally, the notion of open projective plane was introduced by Hall in [22] as a tool to study free projective planes, which he described as the analogue of free groups in the context of projective geometry. He defined *the free projective plane of rank k* as the inductive closure of the finite configuration π_0^k consisting of a line ℓ , $k - 2$ points on ℓ and two points off of ℓ , under the following operations:

- (a) if two points are not joined by a line, then add a new line that joins them;
- (b) if two lines are parallel, then add a new point intersecting them.

To study the free projective planes, Hall defined the notion of *open plane*: a plane P is open if every finite configuration $C \subseteq P$ contains at least one line ℓ incident with at most two points in C , or it contains at least one point p incident with at most two lines in C . Hall then proved that free projective planes are always open and, conversely, that finitely generated open projective planes are free (the assumption of finite generation is necessary, cf. [26]). In [24] Hyttinen and Paolini used Hall’s notion of open projective plane (which is a first-order notion) to axiomatise the theory of non-degenerate (i.e., infinite) free projective planes, proving various model theoretic results, in particular stability and non-superstability of the theory.

The idea of free constructions does not apply only to projective planes, but to a vast number of cases of incidence geometries. The first generalization of Hall’s construction is probably due to Tits [36], who introduced a notion of free completion in the context of *generalised n -gons*, combinatorial structures which Tits introduced to give a geometric interpretation of groups of Lie type in rank 2 (in fact, generalised n -gons are simply the rank 2 case of spherical buildings). As in the case of projective planes (i.e., generalised 3-gons), one can introduce a notion of *open generalised n -gon*. Relying on this, in [1] Ammer and Tent use the theory of non-degenerate open generalised n -gons to axiomatise the theory of free generalised n -gons, generalizing all the result of Hyttinen and Paolini [24] from $n = 3$ to any $3 \leq n < \omega$. Apart from the increased level of generality, the methods from [1] were decidedly more elegant than those from [24], and made clearer the crucial role played by *Hrushovski constructions* in this setting. Our paper rests on the fundamental insights of [1].

Given [24] and [1], it is thus natural to ask to what extent these results indicate a pattern, and whether, whenever in a given geometric context a notion of open incidence structure is well-defined, then it can be used to axiomatise the free incidence structures in that class. The starting point of this investigation is of course the analysis of systems of points and lines (as in the case of projective planes and generalised n -gons), which we will refer to as *the rank 2 case*. To our knowledge, the most detailed and general study of rank 2 free constructions in incidence geometry is contained in Chapter 13 “Free Constructions” by Funk and Strambach in the already mentioned “Handbook of Incidence Geometry” [14]. There it is shown that, in all naturally occurring theories of incidence structures T , there is a notion of free construction and a corresponding notion of free object of rank k (in the

sense of number of generators). In particular, in [14, Chapter 13] a general theory of free extensions is proposed for T an arbitrary $\forall\exists$ -theory of incidence structures, and this theory is tested on the following cases:

- (1) (k, n) -Steiner systems (for $2 \leq k < n$);
- (2) generalised n -gons (for $n \geq 3$);
- (3) k -nets (for $k \geq 3$);
- (4) affine planes;
- (5) projective Möbius, Laguerre and Minkowski planes.

In particular, Funk and Strambach provide a uniform definition of free completions in incidence geometry, and they also show that one can always define a notion of open configuration (generalizing the definition of open projective plane considered above). Commenting on their results, Funk and Strambach [21, p. 741] remark:

Besides a survey on significant results obtained thus far, the main purpose of this article is the development of a unifying treatment including all classes of incidence geometries which can be characterised by a set Σ of axioms formulated in a first-order language \mathcal{L} (e.g., projective planes, affine planes, generalised n -gons, Benz planes, etc.). By using rather simple model-theoretic tools, we can define the notions of (*hyper-*)*free*, *open*, *confined*, *closed*, (*hyper-*)*free extensions*, and *degenerate geometries* without knowing Σ and \mathcal{L} explicitly. To a very vast extent, we succeed in reformulating and proving the main results concerning free extensions within this general frame. The application of these results reduces to an easy verification of some model-theoretic conditions on the axioms. Thus it becomes clear that the real nature of free extensions in fact lies beyond geometry. We hope that this insight will contribute to stop splitting research on that subject.

Our present contribution is fully in this spirit. In this paper we develop an abstract model-theoretic framework that we believe will cover all concrete instances of free incidence structures and, using that, we extend the results from [1, 24] to *all examples of incidence geometries of rank 2* known to us. In particular, this covers all concrete cases considered in [21] and listed above. We take as a special sign of strength of our approach the fact that we were able to apply our methods also to cases of incidence geometries with notions of parallelism, namely affine planes and Möbius, Laguerre and Minkowski planes (cf. Sections 4.5 and 4.6), at the best of our knowledge this has not been considered elsewhere (apart from [21]). Our main result can be summarised in the following theorem, which vastly generalises [1, 24]:

Main Theorem. *The theory T^+ of open non-degenerate models of T is complete, decidable and strictly stable for any of the following theories T :*

- (1) (k, n) -Steiner systems (for $2 \leq k < n$);
- (2) generalised n -gons (for $n \geq 3$);
- (3) k -nets (for $k \geq 3$);
- (4) affine planes;
- (5) projective Möbius, Laguerre and Minkowski planes.

Thus, for any of the above classes, all the non-degenerate free incidence structures in that class are elementarily equivalent. Further, the theory T^+ does not have a prime model, it is not model complete, and it does not admit quantifier elimination.

Finally, for any $A, B, C \subseteq \mathfrak{M}$ we have that $B \downarrow_A C$ if and only if $\text{acl}(ABC) = F(\text{acl}(AB) \otimes_{\text{acl}(A)} \text{acl}(AC))$ (where \mathfrak{M} is the monster model of T^+ and the right-hand part of the equivalence is defined in Section 3).

Our paper has two parts, an abstract one (Section 3) and one that deals with concrete applications (Section 4). The previous theorem puts together our general results from Section 3 with the applications from Section 4. We stress here however that our results in Section 3 are stated in a purely axiomatic fashion, and we expect to find applications of our methods beyond the scope of applications detailed here, which, as mentioned, are all cases of geometries of rank 2, as in [21], which was one of the main inspirations of the present study. In a work in preparation we intend to explore the general rank k case, with particular focus on connections to diagram geometries and buildings, and thus naturally to the fundamental references [12, 34]. But as the present paper is already of considerable length and covers many case studies we decided to split the treatment of the subject into two separate papers.

In addition to the case studies listed above, we decided to include here a further one (cf. Section 4.1) which has a very different motivation and history than the other ones, i.e., the case of *n-open graphs*. These structures were very recently introduced by Baldwin, Freitag and Mutchnik in [3] to give the first example, for every $n < \omega$, of a stable theory T_n admitting a type p such that $F_{Mb}(p) = n$, and so that $F_{Mb}(p) \leq n$ for all types p . This case fits perfectly in our setting and shows that, apart from the intrinsic geometric interest, our approach could be exploited by model theorists to give counterexample to conjectures in classification theory. From our point of view, an additional key element of interest is that these graphs make for an example where the notion of openness applies, but where strictly speaking there is no notion of free extension (we elaborate this further in Section 3, where we distinguish between arbitrary completions and free algebraic completions). Additionally, *n-open graphs* are one-sorted structures and, with the exception of the case $n = 1$, in which case the corresponding theory is the theory of the free pseudoplane, they do not have a natural interpretation in terms of geometries of rank 2. We show that our framework covers also these structures and we prove a version of the theorem above (cf. Corollary 4.1). We stress that our results in this case are not novel, but provide nonetheless an interesting application of our techniques beyond the context of free incidence geometries of rank 2.

We now come to the technical side of the story, which is intended mainly for the model theorists, but also explains to the general reader the techniques used to establish our results. The fundamental technical tool behind this study is the use of *Hrushovski constructions*. The relevance of this was already realised by the first author in [24, Section 8], but the idea was fully developed only in [1]. In this study we owe a great deal to the elegant treatment of the subject presented in [1] in the case of generalised *n-gons*. At the same time we also make a crucial use of the notion of HF-order; this notion was first introduced in [24], generalizing the existing notion from the incidence geometry literature (cf. e.g. [32]). These HF-orders are simply a technical tool (essentially an application of compactness) and they are not necessary for the development of the subject, but we believe that they make for a very handy instrument for dealing with these kind of Hrushovski constructions.

What we feel to be the most interesting contribution of this study from the technical point of view is the isolation of some axiomatic conditions which make the whole machinery work. The axioms are actually split into different sets, namely

3.2, 3.28, 3.43, 3.60 and 3.65. The set of axioms from 3.2 isolates some conditions on the Hrushovski class of finite structures (\mathcal{K}, \leq_o) under consideration, expanding the axiomatic framework of Baldwin and Shi in [3] to the case of open incidence structures. Crucially, some of the axioms from 3.2 (most notably 3.2(H) and 3.2(I)) are *false* in most Hrushovski constructions arising from a predimension. We believe that this not a defect of our approach but rather one of its main strengths, as our axioms are exactly what makes things work in the more specific context of open incidence structures (a context which is specific but general enough to cover all the known cases of free incidence structures in rank 2). The second set of axioms, namely 3.28, isolates some conditions on infinite models of the intended axiomatisation of the generic for (\mathcal{K}, \leq_o) , namely, the theory T^+ defined in 3.22, which in all concrete cases will correspond to the theory of open incidence structures of the given type (generalising [1, 24]). The assumptions from 3.2 and 3.28 are enough by themselves to prove that the theory T^+ is complete and stable, and thus they already suffice to show that in all intended applications the free incidence structures are all elementarily equivalent. Assumptions 3.43, 3.60 and 3.65 are needed for our additional results: we use them to characterise the relation of forking independence in T^+ , to prove that T^+ is not superstable, and to establish that T^+ does not have a prime model, which are all generalisation of the results from [1, 24]. We crucially notice that, while the assumptions from 3.2, 3.28 and 3.43 hold in all the examples of geometries that we consider in Section 4, the assumptions from 3.60 and 3.65 are not always valid, and they thus provide us with a main *dividing line* for our applications. On the one hand, we have the incidence geometries from our Main Theorem, which are all not superstable and with no prime model; on the other hand, there are structures such as the free pseudoplane which, as well-known, is ω -stable and prime. Our abstract framework covers both cases.

The structure of the paper reflects our axiomatic treatment of the subject. After recalling some preliminary definitions and notational conventions in Section 2, in Section 3 we present our general framework and we prove all our main results about T^+ from the corresponding set of assumptions. We provide a general proof of completeness and stability of T^+ , we characterise forking independence, and we show that T^+ is not superstable and does not have a prime model. In the subsequent Section 4 we first consider the simpler case of n -open graphs (introduced in [3]), and then we deal separately with each case of incidence structures from (1)-(5) of our Main Theorem above, verifying each of the axiomatic conditions isolated in Section 3. We believe that our abstract approach confirms Funk and Strambach's fundamental verdict that "the real nature of free extensions lies beyond geometry".

2. PRELIMINARIES AND NOTATIONS

We introduce in this section some general notions that we use to define the general framework at the heart of this article. We also fix some notation and terminology that we will follow throughout the paper. Firstly, we introduce the general context of the paper and some basic notational convention.

Context 2.1. In this article we shall always work in a finite multi-sorted relational language L . We denote the substructure relation between L -structures as $A \leq B$. We let T to be an L -theory and T_{\forall} its universal fragment. We let (\mathcal{K}, \leq_o) to be a hereditary class of finite models of T_{\forall} and \leq_o to be a strong substructure relation (this means that $A \leq_o B$ implies $A \leq B$ and that $A \leq_o B \in \mathcal{K}$ implies that $A \in \mathcal{K}$).

Notation 2.2. Let κ be a cardinal. We write $A \subseteq_\kappa B$ if $A \subseteq B$ and, additionally, $|A| < \kappa$. Thus in particular $A \subseteq_\omega B$ means that A is a finite subset of B . We write A^κ for the set of all sequences of elements from A of length κ , and we let $A^{<\kappa} = \bigcup_{\alpha < \kappa} A^\alpha$. Let M be an L -structure, κ a cardinal, $\bar{a} \in M^{<\kappa}$ and $A \subseteq M$, we write $\text{tp}_M(\bar{a}/A)$ for the *type* of \bar{a} in M with parameters from A and $\text{tp}_M^{\text{qf}}(\bar{a}/A)$ for the *quantifier-free type* of \bar{a} in M with parameters from A . Given a model M and a subset $A \subseteq M$ we write $\text{acl}_M(A)$ for the algebraic closure of A in M and $\text{acl}_M^{\text{qf}}(A)$ for the algebraic closure of A in M computed in the restricted language with only quantifier-free formulas. We often omit the subscript M when it is clear from the context.

As we spelled out in the introduction, our approach in this paper is mainly graph-theoretical. We thus recall some standard notions from graph theory, which we will often employ in this work. In this paper by a graph we mean a structure of the form (G, E) with E a binary, irreflexive and symmetric relation on G .

Definition 2.3. Let (G, E) be a graph, then we define the following:

- (1) the *valency* of an element $a \in G$ is the size of the set $\{b \in G : aEb\}$ of neighbours of a ;
- (2) the *distance* $d(a, b) = d(a, b/G)$ between two elements $a, b \in G$ is the length n of the shortest path $a = c_0Ec_1E \dots Ec_{n-1}Ec_n = b$ in G , and it is ∞ if there is no such path;
- (3) the *girth* of a graph is the length of its shortest cycle;
- (4) the *diameter* of a graph is the maximal distance between any two elements;
- (5) the graph G is *bipartite* if there are $A, B \subseteq G$ such that $A \cap B = \emptyset$, $A \cup B = G$, and every edge has one vertex in A and one in B .

We next introduce some less standard notions, mostly from [21], which are motivated by the geometric nature of the theories which we consider in this work. The following distinction between incidence and parallelism symbols is essentially from [21, pp. 743, 749].

Definition 2.4.

- (1) Let $R \in L \cup \{=\}$ be a relation symbol of arity $n \geq 2$, we say that R is a *local equivalence relation in the theory T* if for any L -structure $A \models T$ and tuple $a_1, \dots, a_{n-2} \in A$ we have that $R(x, y, a_1, \dots, a_{n-2})$ induces an equivalence relation on A (notice that in the case $n = 2$ the quantifier “for every tuple $a_1, \dots, a_{n-2} \in A$ ” is vacuous). If $R \in L \cup \{=\}$ is not a local equivalence relation, then we say that it is an *incidence relation in the theory T* . We omit the reference to T when it is clear from the context.
- (2) We denote by L_p the subset of L consisting of all local equivalence relations in T , and by L_i the subset of L consisting of all incidence relations in T .
- (3) Let M be an L -structure, $\bar{a} \in M^{<\kappa}$ and $B \subseteq M$. The *parallelism type* $\text{tp}_M^p(\bar{a}/B)$ of \bar{a} in M with parameters in B is the quantifier-free type of $\bar{a} \in M^{<\kappa}$ with parameters in B computed in the restricted signature L_p . The *incidence type* $\text{tp}_M^i(\bar{a}/B)$ of \bar{a} in M with parameters in B is the quantifier-free type of $\bar{a} \in M^{<\kappa}$ with parameters in B computed in the restricted signature L_i . We often omit the subscript M when it is clear from the context.

Remark 2.5. Firstly, notice that to simplify the notation we always assume that in a local equivalence relation R the first two variables represent the arguments of

the equivalence relation, and the rest are the parameters. Secondly, notice that the equality $=$ is a (local) equivalence relation in all theories T . Clearly we have that $\text{tp}_M^i(\bar{a}/B) \cup \text{tp}_M^p(\bar{a}/B) = \text{tp}_M^{\text{qf}}(\bar{a}/B)$, and that quantifier-free formulas containing equality symbols are contained in $\text{tp}_M^p(\bar{a}/B)$. Moreover, we stress that the sentence stating that a certain relation $R \in L$ is a local equivalence relation is universal. It follows that T and T_\forall agree on the distinction between L_i and L_p . We will often use this fact implicitly when working with models of T_\forall .

We stress that the previous notions of parallelism type $\text{tp}^p(\bar{a}/B)$ and incidence type $\text{tp}^i(\bar{a}/B)$ are not standard in the literature, but they are convenient in the present context as we shall deal with theories (e.g., the theory of affine planes) containing both symbols for incidences and symbols for parallelism, which require a different treatment. We need these notions especially in the various definitions of closure, e.g., the following definition of Gaifman closure 2.8 and in the definition of HF-closure from 3.19. We first recall the notion of Gaifman graph (cf. [19, p.26]).

Definition 2.6. Let $M \models T_\forall$ and $L' \subseteq L$.

- (1) The *Gaifman graph* $G_{L'}(M)$ of a structure A is the graph whose set of vertices is M and which has an edge $E(a, b)$ between $a, b \in M$ whenever $A \models R(\sigma(a, b, \bar{c}))$ for some relation $R \in L'$, for some tuple $\bar{c} \in M^{<\omega}$ of the suitable arity and for some permutation σ .
- (2) We write $G(M)$ for the *full Gaifman graph* $G_L(M)$. We write $G_i(M)$ for the *incidence Gaifman graph* $G_{L_i}(M)$, i.e., the Gaifman graph computed in the restricted language of incidence relation symbols in T . We write $G_p(M)$ for the *parallelism Gaifman graph* $G_{L_p}(M)$, i.e., the Gaifman graph computed in the language of local equivalence relations in T . We omit reference to T in the notation as it will always be clear from the context what T is.
- (3) The *distance* $d_{L'}(a, b/M)$ between two elements $a, b \in M$ is their distance in the Gaifman graph $G_{L'}(M)$. We write $d(a, b/M)$ for their distance in $G(M)$, $d_i(a, b/M)$ for their distance in $G_i(M)$ and $d_p(a, b/M)$ for their distance in $G_p(M)$. We write $d(a, b)$, $d_i(a, b)$ and $d_p(a, b)$ when the structure M is clear from the context.
- (4) We extend the notion of distance to arbitrary finite subsets $A, B \subseteq M$ by letting

$$d(A, B) := \min_{a \in A} \left(\min_{b \in B} (d(a, b)) \right),$$

and we let the distance $d(\bar{a}, \bar{b})$ between two finite tuples $\bar{a}, \bar{b} \in M^{<\omega}$ be the same as their distance as subsets. We define $d_i(A, B)$ and $d_p(A, B)$ analogously.

Remark 2.7. We notice that, since relations in L_p are (locally) transitive, the distance $d_p(a, b/M)$ is always either 0 or 1. In particular, $d_p(a, b/M) = 1$ means that there is a local equivalence relation $P \in L_p$ and a tuple \bar{c} such that $M \models P(a, b, \bar{c})$.

We use the previous definitions to define the notion of Gaifman closure. This was introduced in [21, Def. 2] under the name of (restricted) interior construction.

Definition 2.8 (Gaifman Closure). Let $A \subseteq M \models T_\forall$, the *Gaifman closure* $\text{gcl}_M(A)$ of A in M is defined as $\text{gcl}_M(A) = \text{gcl}_M^i(A) \cup \text{gcl}_M^p(A)$, where:

- (1) $\text{gcl}_M^i(A) = \{b \in M : d_i(A, b) \leq 1\}$;
- (2) $\text{gcl}_M^p(A)$ is any subset of M satisfying the two following constraints:

(a) for any $b \in M$ with $d_p(A, b) \leq 1$, there is $b' \in \text{gcl}_M^p(A)$ such that

$$M \models R(b, a, \bar{c}) \iff M \models R(b', a, \bar{c})$$

for all $R \in L_p$, $a \in A$ and $\bar{c} \in A^{<\omega}$;

(b) for all $b, b' \in \text{gcl}_M^p(A)$ there are some $R \in L_p$, $a \in A$, and $\bar{c} \in A^{<\omega}$ s.t.

$$M \models R(b, a, \bar{c}) \text{ and } M \models \neg R(b', a, \bar{c}).$$

In other words, $\text{gcl}_M^p(A)$ is any subset of $\{b \in M : d_p(A, b) \leq 1\}$ containing exactly one element for every distinct parallelism type with parameters in A .

Remark 2.9. Clearly, we have that $A \subseteq \text{gcl}_M^i(A)$. Moreover, since the equality symbol “=” belongs to L_p , we also have that $A \subseteq \text{gcl}_M^p(A)$, since every $a \in A$ is the unique element satisfying the formula $x = a$. We stress that by the definition of $\text{gcl}_M^p(A)$ the Gaifman closure of a set is *not unique*. However, we always work in a context (\mathcal{K}, \leq_o) where the choice of one Gaifman closure over another does not really make a difference. This is captured by Condition 3.2(K) below.

3. THE GENERAL FRAMEWORK

In this paper we investigate the first-order theory of open incidence structures of rank 2, such as open projective planes, open Steiner systems, and similar classes of geometric objects. To cover all these examples we proceed in an axiomatic fashion, and we start in Section 3.1 by introducing a class of finitary conditions (Context 3.2) that carves out a special subclass of Hrushovski constructions that covers all the geometric theories which we are interested in. In Section 3.2 we introduce the main technical device of our article, namely that of HF-orders. In Section 3.3 we introduce the theory T^+ , which is the main object of investigation of the article, and we state the crucial assumptions 3.28 which we later verify for all our concrete examples. In Section 3.4 we prove the main result of the paper, i.e., that under the context from 3.2 and the assumptions from 3.28 the theory T^+ is both complete and stable. In Section 3.5 we list an additional set of assumptions (3.43) and we use them to characterise the relation of forking independence in models of T^+ . Finally, in Sections 3.6 and 3.7 we identify some further conditions which entail that T^+ is respectively not superstable, and that it does not have a prime model, it is not model complete and does not have quantifier elimination.

3.1. Finitary conditions

As we laid out in the previous section, we always work in a finite multi-sorted relational language L . We also fix a class \mathcal{K} of finite structures and a relation \leq_o between elements of \mathcal{K} which makes (\mathcal{K}, \leq_o) hereditary, i.e., if $A \leq_o B \in \mathcal{K}$ then $A \in \mathcal{K}$. We refer to extensions $A \leq_o B$ as *strong extensions* and we provide the following additional definitions.

Definition 3.1. For $A \subseteq B \in \mathcal{K}$, we define the following notions:

- (1) we say that (A, B) is a *one-element extension* if $|B \setminus A| = 1$;
- (2) we say that $A \leq_o B$ is a *minimal (strong) extension* if there is no $A \subseteq C \subsetneq B$ such that $A \leq_o C \leq_o B$;
- (3) we say that (A, B) is an *algebraic extension* if there is some $n < \omega$ such that for every $A \subseteq C \in \mathcal{K}$ we have that C contains at most n copies of B over A ;
- (4) we say that (A, Ab) is a *trivial extension* if $d_{Ab}(A, b) = \infty$, i.e., b has no edge with elements from A in the full Gaifman graph $G(Ab)$.

Additionally, we assume that (\mathcal{K}, \leq_o) satisfies the following conditions.

Context 3.2. We let (\mathcal{K}, \leq_o) be a hereditary class \mathcal{K} of finite models of T_\forall such that \leq_o is a relation with the following properties:

- (A) if $A \in \mathcal{K}$, then $A \leq_o A$;
- (B) if $A \leq_o B$, then A is a substructure of B ;
- (C) if $A, B, C \in \mathcal{K}$ and $A \leq_o B \leq_o C$, then $A \leq_o C$;
- (D) $\emptyset \in \mathcal{K}$ and $\emptyset \leq_o A$, for all $A \in \mathcal{K}$;
- (E) if $A, B, C \in \mathcal{K}$, $A \leq_o C$, and B is a substructure of C , then $A \cap B \leq_o B$;
- (F) if $A \subseteq B, B' \in \mathcal{K}$, $A \leq_o B$ and $f : B \cong_A B'$, then $A \leq_o B'$;
- (G) (\mathcal{K}, \leq_o) satisfies the amalgamation and the joint embedding property;
- (H) if $A \subseteq B \in \mathcal{K}$ and $A \not\leq_o B$, then (A, B) is an algebraic extension;
- (I) there is a finite set $X_{\mathcal{K}} \subseteq \omega$ such that, if $A \leq_o B \in \mathcal{K}$, then there are $n \in X_{\mathcal{K}}$ and $\bar{b} = (b_1, \dots, b_n) \in (B \setminus A)^n$ such that $B \setminus \{b_1, \dots, b_n\} \leq_o B$;
- (J) there is $n_{\mathcal{K}} < \omega$ such that, if $A \leq_o Ab \in \mathcal{K}$ is a one-element extension, then $|\text{gcl}_{Ab}(b) \setminus \{b\}| \leq n_{\mathcal{K}}$ (recall the definition of gcl from 2.8);
- (K) for all $A, B \in \mathcal{K}$, if $A \cap \text{gcl}_B(B \setminus A) \leq_o \text{gcl}_B(B \setminus A)$ then $A \leq_o B$;
- (L) for all $A \in \mathcal{K}$, if (A, Ab) is trivial (in the sense of 3.1(4)), then $A \leq_o Ab \in \mathcal{K}$.

Remark 3.3. We stress that the conditions from 3.2 can be split in two groups. Conditions 3.2(A)-(F) are standard and hold also in Hrushovski constructions arising from a *predimension* δ (cf. [3]). Similarly, Condition 3.2(G) is usually also required in arbitrary Hrushovski constructions. On the contrary, Conditions 3.2(H)-(K) are quite specific of our current combinatorial setting. Condition (H) simply says that configurations that are not strong are always algebraic. As we show later in 3.33, this has the important consequence that every extension of an algebraically closed set (in a model of T^+) is strong. Condition (I) means that every strong extension $A \leq_o B$ in \mathcal{K} can be build *via* a series of strong extensions which are bounded in size. Conditions (J)-(K) capture the essential properties of the Gaifman closure in the class of geometries we are investigating. Condition (J) essentially says that the Gaifman closure of any strong extension is bounded: if $A \leq_o B$, then clearly $\text{gcl}_B(B \setminus A) = \bigcup_{b \in B \setminus A} \text{gcl}_B(b)$, thus by (J) we have that $|\text{gcl}_B(B \setminus A)| \leq (n_{\mathcal{K}} \cdot |B \setminus A|) + |B \setminus A|$. Together, Conditions (I) and (J) have the important consequence that the notion of HF-order becomes first-order expressible (cf. 3.10 and 3.13). Condition (K) essentially says that, in order to determine if an extension (A, B) is strong, it is enough to look at the subextension induced by the Gaifman closure of $B \setminus A$. Together with Condition (J), this has the important consequence that whether an extension $A \leq_o B$ is strong depends on a subset of B of size $\leq (n_{\mathcal{K}} \cdot |B \setminus A|) + |B \setminus A|$. Finally, Condition (L) makes sure that trivial extensions (in the sense of 3.1(4)) of any sort are strong.

Example 3.4. To provide some intuition about the abstract setting from 3.2, we explain how for example the case of open projective planes from [24] fits into this framework. Recall that a *partial projective plane* is a set of lines and points such that any two points are incident with at most one line, and any two lines are incident with at most one point. A partial projective plane P is *open* if every finite subset $A \subseteq P$ contains either a point incident with at most two lines, or it contains a line incident with at most two points. We let \mathcal{K} be the class of finite, partial, open projective planes, and we write $A \leq_o B$ if every nonempty $C \subseteq B \setminus A$ contains at least one element incident to at most two elements from AC . Then, Condition (H)

boils down to the fact that extensions $B \supseteq A$ where each element $b \in B \setminus A$ has at least three incidences are algebraic over A , which can be easily verified. Condition **(I)** holds immediately for $X_{\mathcal{K}} = \{1\}$, since every finite open partial projective plane can be obtained via a sequence of one-element extensions, by adding points incident with at most two lines and lines incident to at most two points (this was shown already by Siebenmann in [32, Lemma 1]). Similarly, Condition **(J)** is true by letting $n_{\mathcal{K}} = 2$, since whenever $A \leq_o Ab$ we have that b is incident to at most two elements from A , i.e., $|\text{gcl}_{Ab}(b) \setminus \{b\}| \leq 2$. Condition **(K)** holds because whether $A \leq_o B$ depends exclusively on the number of incidences of elements in $B \setminus A$, which is clearly preserved when restricting to $\text{gcl}_B(B \setminus A)$. Finally, Condition **(L)** holds because we can always extend a plane $A \in \mathcal{K}$ with either a point or a line which bears no incidence with elements in A .

Notation 3.5. We write T_{\forall} for the universal theory of the class (\mathcal{K}, \leq_o) , i.e., for the theory $\text{Th}_{\forall}(\mathcal{K})$.

We then extend the relation \leq_o from finite structures in \mathcal{K} to arbitrary models of T_{\forall} as follows.

Definition 3.6. Let $A \subseteq B \models T_{\forall}$, then we let $A \leq_o B$ if and only if $A \cap B_0 \leq_o B_0$ for all finite $B_0 \subseteq B$.

Notice that all definitions from 3.1 extend naturally also to the setting with arbitrary structures $A \subseteq B \models T_{\forall}$. Also, if B is finite, then Condition **(E)** makes sure that $A \leq_o B$ holds if and only if $A \leq_o B_0$ for all $B_0 \subseteq B$, thus the previous definition agrees with the original relation \leq_o over the finite models of T_{\forall} (cf. also [3, Def. 2.17]). Finally, we remark that, exactly as observed in [3, Lemma 2.8] the property of being a strong extension is expressible in first-order logic.

Remark 3.7. We notice that our definition of $A \leq_o B$ from 3.6 is equivalent to the extension of the definition of \leq_o from \mathcal{K} to T_{\forall} from [3, 2.17]. Moreover, if A is finite and $A \subseteq B \models T_{\forall}$, then it follows exactly as in [3, 2.8] that $A \leq_o B$ is equivalent to saying that $B \models p(\bar{a})$, where $p(\bar{x})$ is the following type:

$$p(\bar{x}) := \{ \forall \bar{y} (\neg \bigwedge \delta_{(A,C)}(\bar{x}, \bar{y})) : A \not\leq_o C, C \in \mathcal{K} \},$$

where $\delta_{(A,C)}(\bar{x}, \bar{y}) = \bigwedge \text{tp}_C^{\text{qf}}(\bar{a}, \bar{c})$ for \bar{a}, \bar{c} enumerations of A and $C \setminus A$, respectively. Notice that such formulas always exists because L is finite and, moreover, if B is also finite then $p(\bar{x})$ can be chosen to be finite, by considering only those extensions $A \not\leq_o C$ with C of size $\leq |B|$.

By Definition 3.6 the same can be done also for infinite strong extensions $A \leq_o B$. Let $\bar{a} = (a_i)_{i < \kappa}$ be an enumeration of A , then $A \leq_o B$ is equivalent to saying that $B \models p(\bar{a})$, where $p(\bar{x})$ is the following type:

$$p(\bar{x}) := \{ \forall \bar{y} (\neg \bigwedge \delta_{(A_0, C_0)}(\bar{x}, \bar{y})) : A_0 \subseteq_{\omega} A, A_0 \not\leq_o C_0, C_0 \in \mathcal{K} \},$$

where $\bar{x} = (x_i)_{i < \kappa}$, $\delta_{(A_0, C_0)}(\sigma(\bar{x}), \bar{y}) = \bigwedge \text{tp}_{C_0}^{\text{qf}}(\sigma(\bar{a}), \bar{c})$, $\sigma(\bar{a})$ is the subenumeration of A_0 induced by \bar{a} , $\sigma(\bar{x})$ the corresponding subenumeration of $\bar{x} = (x_i)_{i < \kappa}$, and \bar{c} is an enumeration of $C_0 \setminus A_0$.

3.2. HF-orderings

HF-orderings are an important tool from the literature on incidence geometries which allows one to study any extension $A \leq_o B$ by decomposing it in smaller

“atomic” extensions. More precisely, the notion of HF-order was introduced by Siebenmann in [32] in the context of projective planes and later developed further in various references such as [17, 18]. As a way of example, one can associate a linear order to any finite open projective plane, by requiring that each element has at most two incidences with its set of predecessors, and the same can be done to for any free projective plane. In this context, however, HF-orders were always assumed to be wellfounded, thus limiting their scope of application in the context of first-order logic. In fact, as observed in [24, 3.7], if $A = (A, <)$ is an infinite structure with an HF-order (or, for that matter, with any linear order), then its ultrapower A^{ul} will have infinite descending chains. Therefore, in order to apply the technique of HF-orders in the setting of model theory, one also has to make sense of non-wellfounded ones. This approach was taken by Hyttinen and Paolini in [24, 3.4]. Here we introduce a further generalisation of their definition, as we define HF-orderings arbitrarily for any class \mathcal{K} satisfying the conditions from 3.2. Most importantly, we stress the crucial role of conditions 3.2(I)–(J), which are essential to make the notion of HF-order first-order expressible (cf. Remark 3.13 below). We start by fixing some notation and then proceed to defining HF-orders.

Notation 3.8. For any partial order $(X, <)$ and $Y \subseteq X$ we write Y^\downarrow for the downset generated by Y , i.e., $Y^\downarrow = \{b \in X : \exists a \in Y \text{ s.t. } b \leq a\}$. Also, whenever $(L, <)$ is a linear order and $X, Y \subseteq L$, we write $X < Y$ to say that $x < y$ for every $x \in X$ and $y \in Y$.

Definition 3.9. If $(X, <)$ is a partial order, then we say that a set $Y \subseteq X$ is *convex* if $a, c \in Y$ and $a < b < c$ entails $b \in Y$.

Definition 3.10 (HF-order). Let $A \subseteq B \models T_\forall$. An HF-order of B over A is a pair $(<, P_<)$, where:

- (H1) $<$ is a linear order of $B \setminus A$, and $P_<$ is a partition of $B \setminus A$ in convex sets of size $\leq \max(X_{\mathcal{K}})$ (where $X_{\mathcal{K}}$ is given by Condition 3.2(I));
- (H2) for every $b \in B \setminus A$ and for every finite $C_0 \subseteq b^\downarrow \setminus P_<(b)$ we have that

$$AC_0 \leq_o AC_0 P_<(b),$$

where $P_<(b)$ is the unique set in $P_<$ such that $b \in P_<(b)$.

For ease of notation, we generally simply say that $<$ is a HF-order of B over A , and we write P instead of $P_<$ when the ordering $<$ is clear from the context. If there is such an order we also write $A \leq_{\text{HF}} B$. When $A = \emptyset$ we say that $<$ is an HF-order of B instead of saying that $<$ is an HF-order of B over \emptyset .

Notation 3.11. We notice that, if $<$ is an HF-order of B over A , then every piece $X \in P$ is essentially a finite ordered subchain of $B \setminus A$. For this reason, we usually think of these pieces as finite tuples with the order induced by $<$. We write \hat{c} for the tuple (c_1, \dots, c_n) such that $P(c) = \{c_1, \dots, c_n\}$ and $c_1 < \dots < c_n$. Differently, for a set $X \subseteq B \setminus A$ we simply let $\hat{X} = \bigcup_{x \in X} P(x)$. Thus for an element $c \in B \setminus A$ we have that \hat{c} is a tuple, while for a set $C \subseteq B \setminus A$ we have that \hat{C} is a set.

Example 3.12. We spell out what the previous definition means in the case of open projective planes (recall 3.4). Since in this case $X_{\mathcal{K}} = \{1\}$, Condition 3.10(H1) says that $<$ is a linear ordering over $B \setminus A$. Then, Condition 3.10(H2) simply says that each element $b \in B \setminus A$ is incident with at most two elements $c_0, c_1 \in A \cup b^\downarrow$

of the opposite sort. This is exactly the notion of HF-order for projective planes defined in [24, Def. 3.4].

Remark 3.13. We observe that the property of being an HF-order is first-order expressible, by a (possibly infinite) set of formulas, in the language $L \cup \{<, P\}$ where $<$ is a binary relation and P is a relation of arity $\max(X_{\mathcal{K}})$. Notice that the existence of $\max(X_{\mathcal{K}})$ is always guaranteed by Condition 3.2(I). The fact that $<$ is a linear order is clearly expressible by a sentence, and similarly we can express that P is a partition of $B \setminus A$ in convex sets of size $\leq \max(X_{\mathcal{K}})$ (notice that we also need to use some technical device such as repetition of the last occurring element to encode pieces of the partition which have length shorter than $\max(X_{\mathcal{K}})$). Therefore we can encode Condition 3.10(H1) with a first-order formula. Similarly, since by Remark 3.7 we can capture the strong submodel relation \leq_o in first-order logic, we can also express Condition 3.10(H2) by a set of formulas in $L \cup \{<, P\}$.

The following Lemma is essentially a generalisation of [32, Lemma 1] to the abstract setting of a class (\mathcal{K}, \leq_o) satisfying the conditions from 3.2.

Lemma 3.14. Let $A \leq_o C \in \mathcal{K}$, then there is an HF-order of C over A .

Proof. Since $A \leq_o C$, by Condition 3.2(I) there is some tuple $\bar{c}_1 = (c_1^1, \dots, c_1^{k_1}) \in (C \setminus A)^{k_1}$ with $k_1 \in X_{\mathcal{K}}$ and such that

$$C \setminus \{c_1^1, \dots, c_1^{k_1}\} \leq_o C.$$

By Condition 3.2(E) it follows that $A \leq_o C \setminus \{c_1^1, \dots, c_1^{k_1}\}$, thus there is some $\bar{c}_2 = (c_2^1, \dots, c_2^{k_2}) \in (C \setminus A c_1^1, \dots, c_1^{k_1})^{k_2}$ with $k_2 \in X_{\mathcal{K}}$ and such that

$$C \setminus \{c_1^1, \dots, c_1^{k_1} c_2^1, \dots, c_2^{k_2}\} \leq_o C \setminus \{c_1^1, \dots, c_1^{k_1}\}.$$

Since C is finite, after finitely-many steps, we find some $n < \omega$ and tuples $\bar{c}_1, \dots, \bar{c}_n$ such that $C = A \cup \{c_j^i : 1 \leq i \leq k_j, 1 \leq j \leq n\}$. We then let $P(c_j^i) = \{c_j^i : 1 \leq i \leq k_j\}$ and we define

$$c_j^i < c_m^\ell \iff (m < j) \text{ or } (j = m \text{ and } i < \ell).$$

Then $(<, P)$ is a HF-order of C over A . ■

The next proposition is an abstract version of [24, Prop. 3.14]. The proof is essentially the same.

Proposition 3.15. Let $A \leq_o B \models T_{\forall}$, then there is an HF-order $<$ of B over A .

Proof. The proof follows as in [24]. Let X be the set of all finite substructures of B , by 3.14 every $C \in X$ has an HF-order $<_C$ over \emptyset . Let \mathfrak{U} be an ultrafilter on X such that for all $C \in X$ we have that $X_C = \{D \in X : C \subseteq D\} \in \mathfrak{U}$. For every $C \in X$ we let $<_C^1, \dots, <_C^{n(C)}$ be an injective enumeration of all the HF-orderings of C over \emptyset . Then, since \mathfrak{U} is an ultrafilter, it follows that for every $C \in X$ there is a unique HF-ordering $(<_C^*, P_C^*)$ such that:

$$Y_C := \{D \in X : C \subseteq D, <_D \upharpoonright C = <_C^* \text{ and } P_{<_D} \upharpoonright C = P_C^*\} \in \mathfrak{U},$$

where $P_{<_D} \upharpoonright C = \{X \cap C : X \in P_{<_D}\}$. Then for every $c \in B$, there is a structure $C \in X$ such that $P_{<_C^*}(c) = P_{<_D}(c)$ for all $C \subseteq D \in X$. We let $<_* = \bigcup_{C \in X} <_C^*$ and we define a partition P_* by letting, for every $c \in B$, $P_*(c) = P_{<_*^*}(c)$, where C is chosen to satisfy $P_{<_*^*}(c) = P_{<_D}(c)$ for all $C \subseteq D \in X$. It is then easy to verify that $(<_*, P_*)$ is a HF-order of B . ■

Corollary 3.16. Let $A \subseteq B \models T_{\forall}$. The following are equivalent:

- (1) $A \leq_o B$;
- (2) $A \leq_{\text{HF}} B$;

Proof. The direction from (1) to (2) is Proposition 3.15. The direction from (2) to (1) follows from Def. 3.6 and induction on the size of $C \subseteq_{\omega} B \setminus A$. ■

If $<$ is an HF-order of a structure B and $C \subseteq B$, it is often useful to restrict the HF-order of B to the subset C (we did this for example in the proof of 3.15). To this end, we introduce the following definition.

Definition 3.17. Let $A \subseteq B$ and suppose that $<$ is a HF-order of B over A . Let $C \subseteq B$, then the *restriction* of $<$ to C is the pair $(<', P_{<'})$ defined by letting $<' := < \upharpoonright C$ and $P_{<'}(c) = P_{<}(c) \cap C$ for all $c \in C$. With a slight abuse of notation we often write simply $< \upharpoonright C$ for the restriction of the HF-order $(<, P_{<})$ to C .

Remark 3.18. We notice that, if $<$ is a HF-order $<$ of B over A and $C \subseteq B$, then the restriction $< \upharpoonright C$ is always a HF-order of C over $A \cap C$. This follows immediately from Definition 3.10 and Condition 3.2(E).

In stark contrast to the previous remark, if $<$ is a HF-order of B and C an arbitrary subset of B , it is not necessarily the case that $< \upharpoonright B \setminus C$ induces an HF-order of B over C . In fact, as C is an arbitrary subset of B , we could actually even have that $C \not\leq_o B$. We generalise [24, Def. 3.9] and we introduce the notion of HF-closure of a set C relative to a specific HF-order $<$, which is an extension $\text{cl}_{<}(C) \supseteq C$ of size $\leq |C| + \aleph_0$ such that $\text{cl}_{<}(C) \leq_o B$ always holds. To obtain the set $\text{cl}_{<}(C)$ we extend C inductively so that it eventually contains the Gaifman closure (cf. 2.8) of any given element $c \in \text{cl}_{<}(C)$ in its set of predecessor c^{\downarrow} . Crucially, the HF-closure depends also on the starting HF-order $<$ and not exclusively on C .

Definition 3.19 (HF-closure). Let $A \subseteq B \models T_{\forall}$ and suppose $<$ is an HF-order of B over A . Let $C \subseteq B \setminus A$, recall the notion of distance in Gaifman graphs from 2.6, the Gaifman closure from 2.8 and the notation from 3.11. We define:

$$\begin{aligned} \text{cl}_{<}^0(C) &:= \widehat{C} \\ \text{cl}_{<}^{n+1}(C) &:= \text{gcl}_{\text{Acl}_{<}^n(C)^{\downarrow}}(\text{cl}_{<}^n(C)) \\ \text{cl}_{<}(C) &:= \bigcup_{n < \omega} \text{cl}_{<}^n(C). \end{aligned}$$

We refer to $\text{cl}_{<}(C)$ as the HF-closure of C with respect to $<$, as it depends not only on the ambient model B but also on the specific HF-order $<$.

Example 3.20. In the case of an open projective plane P with an HF-order $<$, the HF-closure of a set $C \subseteq P$ is obtained recursively by adding, for every element $c \in \text{cl}_{<}^n(C)$ all the elements $b < c$ which are incident to it. This is essentially [24, Def. 3.9]. In particular, local equivalence relations do not play any role in the case of projective planes. The simplest case where one needs to consider also local equivalence relations is the case of affine planes (cf. Sec. 4.5).

Proposition 3.21. If $A \leq_o B \models T_{\forall}$ and $<$ is an HF-order of B over A , then, for every $C \subseteq B \setminus A$, $|\text{cl}_{<}(C)| \leq |C| + \aleph_0$ and $\text{Acl}_{<}(C) \leq_o B$.

Proof. We first prove that $|\text{cl}_<(C)| \leq |C| + \aleph_0$. By Condition 3.2(J), there is a number $n_{\mathcal{K}}$ such that, for every $n < \omega$ and $c \in \text{cl}_<^n(C)$, $|\text{gcl}_{\text{Acl}_<^{n+1}(C)}(c)| \leq n_{\mathcal{K}} + 1$. By the definition of the Gaifman closure we have that

$$\text{gcl}_{\text{Acl}_<^{n+1}(C)}(\text{cl}_<^n(C)) = \bigcup_{c \in \text{cl}_<^n(C)} \text{gcl}_{\text{Acl}_<^{n+1}(C)}(c),$$

and thus $|\text{gcl}_{\text{Acl}_<^{n+1}(C)}(\text{cl}_<^n(C))| \leq (n_{\mathcal{K}} \cdot |\text{cl}_<^n(C)|) + |\text{cl}_<^n(C)|$.

Therefore, at each step $n \leq \omega$ we add only finitely many new elements to $\text{cl}_<^{n+1}(C)$. It follows that $|\text{cl}_<(C)| \leq |C| + \aleph_0$.

We next show $\text{Acl}_<(C) \leq_o B$. Notice that by 3.16 to show that $\text{Acl}_<(C) \leq_o B$ it suffices to prove that there is an HF-order of B over $\text{Acl}_<(C)$. Since $<$ is a HF-order, it has an associated partition $P_<$ of $B \setminus A$. Let $<' = < \upharpoonright (B \setminus \text{Acl}_<(C))$ and $P' = \{X \setminus \text{cl}_<(C) : X \in P_<\}$. Clearly $<'$ is a linear order, and P' is a partition of $B \setminus \text{Acl}_<(C)$ (notice that by the definition of $\text{cl}_<(C)$ every set $X \in P_<$ is either entirely in P' or in $P \setminus P'$). We thus have that $(<', P')$ verifies Condition 3.10(H1).

It remains to verify Condition 3.10(H2). Let $b \in B \setminus \text{Acl}_<(C)$ and suppose towards contradiction that $\text{Acl}_<(C)D \not\leq_o \text{Acl}_<(C)D\widehat{b}$ for some finite $D \subseteq b^\downarrow \setminus P'(b)$ — and recall that by 3.11 \widehat{b} is an enumeration of $P'(b)$. By Definition 3.6, there is a finite subset $C_0 \subseteq \text{cl}_<(C)$ such that $AC_0D \not\leq_o AC_0D\widehat{b}$. Additionally, we can also assume that C_0 is minimal with this property, i.e., for all $C_1 \subsetneq C_0$ it holds that $AC_1D \leq_o AC_1D\widehat{b}$. We distinguish the following cases.

Case 1. $c < b$ for all $c \in C_0$.

Then $AC_0D \not\leq_o AC_0D\widehat{b}$ contradicts the fact that $<$ is an HF-order of B over A .

Case 2. $b < c$ for some $c \in C_0$.

Case 2.1. $d(c, \widehat{b}/\text{Acl}_<(C)P'(b)^\downarrow) > 1$.

Let $C_1 = C_0 \setminus \{c\}$. Since there is no relation between c and \widehat{b} , it follows that \widehat{b} has the same Gaifman closure in AC_0D and AC_1D , thus by Condition 3.2(K) it follows that $AC_1D \not\leq_o AC_1D\widehat{b}$, contradicting the minimality of C_0 .

Case 2.2. $d_r(c, \widehat{b}/\text{Acl}_<(C)P'(b)^\downarrow) = 1$.

Then there is some $d \in P'(b)$ such that $d \in \text{gcl}_{\text{Acl}_<(C)}(c)$ and so by the definition of HF-closure, it follows that $P'(b) \subseteq \text{cl}_<(C)$, contradicting our choice of b .

Case 2.3. $d_p(c, \widehat{b}/\text{Acl}_<(C)P'(b)^\downarrow) = 1$.

By definition of the Gaifman closure (recall 2.8), there is one way to pick elements in it so that $d \in \text{gcl}_{\text{Acl}_<(C)}(c)$ for some $d \in P'(b)$. It follows as in the previous subcase that $P'(b) \subseteq \text{cl}_<(C)$. This again contradicts our choice of b , and concludes the proof. \blacksquare

3.3. The theory T^+

We come to the main object of investigation of the present article, i.e., the theory T^+ . Given a class (\mathcal{K}, \leq_o) satisfying the conditions from 3.2, we define T^+ as follows.

Definition 3.22. Let T^+ be such that $M \models T^+$ if the following happens:

- (1) $M \models \text{Th}_V(\mathcal{K})$;
- (2) let $A \subseteq_\omega M$ and $A \leq_o A\bar{b}$ be a minimal algebraic extension in \mathcal{K} , if every $A \subseteq D \subseteq M$ with $|D \setminus A| \leq \max(X_{\mathcal{K}})$ amalgamates with $A\bar{b}$ in \mathcal{K} , then M contains a copy of $A\bar{b}$ over A ;

- (3) let $A \subseteq_{\omega} M$ and let $A \leq_{\circ} A\bar{b}$ be a minimal non-algebraic extension in \mathcal{K} , then M contains infinitely many copies of $A\bar{b}$ over A .

Remark 3.23. We notice that the definition of T^+ is indeed first-order. In particular, since $M \models T^+$ entails that $M \models \text{Th}_{\forall}(\mathcal{K})$, it follows that the definition of \leq_{\circ} for arbitrary structures from 3.6 applies also to models of T^+ and their subsets. Also, it follows from Remark 3.7 and the fact that L is finite that, if $A, B \subseteq M$ are finite, then it is possible to express that $A \leq_{\circ} B$ by means of a single formula. Thus, clauses (1) and (3) are expressible by a set of first-order sentences in L . Consider clause (2) and suppose $f_0 : A\bar{b} \rightarrow C$ and $f_1 : D \rightarrow C$ witness that $A\bar{b}$ and D amalgamate in some structure $C \in \mathcal{K}$. Let $C' = f_0(A\bar{b}) \cup f_1(D)$, then we have that $C' \in \mathcal{K}$. Moreover, since $f_0(A\bar{b}) \leq_{\circ} C$ and $f_1(D) \leq_{\circ} C$, it also follows from 3.2(E) that $f_0(A\bar{b}) \leq_{\circ} C'$ and $f_1(D) \leq_{\circ} C'$. Thus $A\bar{b}$ and D amalgamate also in C' , which has size $\leq |A\bar{b}| + |D|$. It thus follows that Clause (2) can also be expressed by a set of first-order sentences.

Remark 3.24. We remark that T^+ is always an inductive theory, i.e., its axiomatisation consists exclusively of $\forall\exists$ -formulas. In particular, this is the case independently of how the structures \mathcal{K} are defined: even if the models in \mathcal{K} are identified as the finite structures of a non-inductive theory T , it is still the case that the associated theory T^+ is inductive (T^+ extends $\text{Th}_{\forall}(\mathcal{K})$ but it does not need to extend T , a priori).

Example 3.25. Let $(\mathcal{K}, \leq_{\circ})$ be the class of finite open projective planes with the relation \leq_{\circ} defined in 3.4. Then we have $M \models T^+$ if the following hold:

- (1) M is open, i.e., every finite subset of M contains either a point incident with at most two lines, or it contains a line incident with at most two points;
- (2) every two points in M are incident to a unique common line, and every two lines in M are incident to a unique common point;
- (3) every point is incident to infinitely many lines, and every line is incident to infinitely many points.

Strictly speaking, the three clauses above may appear different from those in 3.22. However, clause 3.22(1) easily follows from the axiomatisation above, while the amalgamation constraint from 3.22(2) is always verified in the context of partial projective planes, thus making the above axiomatisation simpler. As a matter of fact, the amalgamation constraint plays a (crucial) role only in the case of generalised n -gons. We expand on this issue later in Section 4.3.

3.4. Completeness and stability

The main goal of this section is to show that, under the conditions from 3.2 and the following assumptions in 3.28, the theory T^+ is complete and stable. In the cases of the incidence geometries considered in [21] (and that we address later in Sections 4.2-4.6), this in particular entails that, whenever A and B are two structures in \mathcal{K} with infinite free completions $F(A)$ and $F(B)$ (defined in 3.39), then $F(A)$ and $F(B)$ are elementarily equivalent. We first consider the question of completeness of T^+ . We prove this in Theorem 3.34 by introducing a back-and-forth system between suitable models of T^+ . The key technical ingredient in the proof of 3.34 is the notion of countable $(\mathcal{K}, \leq_{\circ})$ -saturation. We thus start by introducing this concept and recalling the existence of a generic structure in the

context of Hrushovski constructions. We stress that, while the concept of (\mathcal{K}, \leq_o) -saturation is standard in the literature, the notion of countable (\mathcal{K}, \leq_o) -saturation is quite specific of our setting. Implicitly, it also occurs in the setting of Hrushovski constructions with irrational α , as in [31] and [27] (cf. Remark 3.27).

Definition 3.26. A structure M is (\mathcal{K}, \leq_o) -saturated if, for every $A \leq_o B \in \mathcal{K}$, whenever there is $f : A \rightarrow M$ such that $f(A) \leq_o M$ then there is $\hat{f} : B \rightarrow M$ extending f such that $\hat{f}(B) \leq_o M$. A structure M is *countably* (\mathcal{K}, \leq_o) -saturated if, for every countable $A \leq_o B \models T_\forall$, whenever there is $f : A \rightarrow M$ such that $f(A) \leq_o M$ then there is $\hat{f} : B \rightarrow M$ extending f such that $\hat{f}(B) \leq_o M$.

Remark 3.27. As defined in [3, 2.10], one says that a model $M \models T^+$ has the *finite closure property* if every finite subset $A \subseteq M$ is included in a set $A' \supseteq A$ such that $A' \leq_o M$. It follows from [3, Thm. 2.12] that, if (\mathcal{K}, \leq_o) satisfies the conditions from 3.2, then there is always a countable structure M_* (the *generic* of (\mathcal{K}, \leq_o)) which has finite closure, is (\mathcal{K}, \leq_o) -saturated, and is unique up to isomorphism. In the context of Hrushovski constructions this is usually the key element to prove completeness: if one can show that the \aleph_0 -saturated models of a certain theory T are also (\mathcal{K}, \leq_o) -saturated, then it follows by back-and-forth between strong finite subsets of two \aleph_0 -saturated models that T is exactly the theory of the generic M_* of (\mathcal{K}, \leq_o) . However, this strategy crucially requires that we work in a setting where every model satisfies the finite closure property. Crucially, this is *not* the case of the geometries from [21]. In fact, the examples that we provide later to show that open Steiner systems, n -gons, k -nets, affine planes and Benz planes are not superstable also witness that there are models $M \models T^+$ and finite configurations $C \subseteq M$ such that for every finite $C \subseteq D \subseteq_\omega M$ we have $D \not\leq_o M$. For this reason we need to consider the stronger version of saturation that we call countable (\mathcal{K}, \leq_o) -saturation. Our completeness proof will then reproduce the back-and-forth argument that one can find in many Hrushovski constructions, but crucially it requires models of T^+ that are countably (\mathcal{K}, \leq_o) -saturated, and not simply (\mathcal{K}, \leq_o) -saturated. Interestingly, our case resembles the one from [31], [27] and [3, Section 6], i.e., the famous non-superstable case of Hrushovski predimension constructions with “irrational α ”. In particular, although Laskowski’s completeness proof [27, Cor. 5.7] is syntactical, he also mentions that “more model-theoretic proofs of these results are possible by passing to sufficiently saturated elementary extensions” [27, p. 168]. Essentially all that our proofs are doing is to follow this alternative model-theoretic route.

We next identify the following assumptions. We stress that 3.28(C1) is the essential ingredient in the proof of completeness, namely Theorem 3.34. On the other hand, Assumption 3.28(C2) is fundamental for the proof of stability, i.e., Theorem 3.37.

Assumption 3.28. Let (\mathcal{K}, \leq_o) and T^+ be respectively as in 3.2 and 3.22, and recall the definition of \leq_o for infinite structures from 3.6. We assume that T^+ satisfies the following conditions:

- (C1) for every $M \models T^+$ there is a model $N \models T^+$ such that $M \preceq N$ and N is countably (\mathcal{K}, \leq_o) -saturated (recall 3.26);
- (C2) there is $m_{\mathcal{K}} < \omega$ s.t., if $A \models T_\forall$ and A contains at least $m_{\mathcal{K}}$ -many elements of each sort, then there is a structure $C(A) \models T^+$ s.t. $A \leq_o C(A)$ and:
 - (a) $C(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : C(A) \cong C(B)$;

- (b) $C(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_o M$ there is an embedding $\hat{h} : C(A) \rightarrow M$ such that $\hat{h} \upharpoonright A = h$ and $\hat{h}(C(A)) \leq_o M$.

Notation 3.29. Whenever $A \models T_V$, we will say that A is non-degenerate if there is a structure $C(A)$ satisfying the conditions from 3.28(C2) above. In particular, it follows from 3.28(C2) that if A contains at least $m_{\mathcal{K}}$ elements of every sort then it is non-degenerate.

Now, it is important to notice that in all the concrete cases from Section 4 we do not directly verify Assumption 3.28(C1), but rather we check some ancillary technical conditions. We identify these as the Assumptions (D1) and (D2) of the following Lemma 3.30, which says that (D1) and (D2) are sufficient to establish 3.28(C1).

Lemma 3.30. Suppose (\mathcal{K}, \leq_o) satisfies the conditions from 3.2 and, additionally, suppose T^+ satisfies the following assumptions (recall 3.10 and 3.19):

- (D1) if $M \models T^+$, $A \subseteq M$ is finite, $<$ is an HF-order of M and (A, Ab) is a trivial one-element extension, then for all $n < \omega$ there is some $b' \in M$ such that:
- (a) $Ab \cong_A Ab'$;
 - (b) $Ab' \leq_o \text{Acl}_{<}^n(b')$;
- (D2) if $M \models T^+$ is \aleph_1 -saturated, $A \leq_o M$ is countable, and for every trivial one-element extension $A \leq_o Ab$ there is $b' \in M$ such that $Ab \cong_A Ab' \leq_o M$, then for every minimal extension $A \leq_o A\bar{c}$ there is $\bar{d} \in M^{<\omega}$ such that $A\bar{c} \cong_A A\bar{d} \leq_o M$.

Then T^+ also satisfies 3.28(C1), i.e., every model $M \models T^+$ has a countably (\mathcal{K}, \leq_o) -saturated elementary extension M' .

Proof. Let $N \models T^+$ and consider a HF-order $(<', P_{<'})$ of N . We can find an elementary extension M of N which is \aleph_1 -saturated in the language $L \cup \{<, P_{<}\}$, i.e., in the expansion of the language with symbols for the HF-order. Crucially, this is possible by Remark 3.13. We prove that $(M, <, P_{<})$ is countably (\mathcal{K}, \leq_o) -saturated.

Firstly, we prove the claim for extensions of the form $A \leq_o B$ where $|A| \leq \omega$ and $|B \setminus A| < \omega$. By induction, it is clearly sufficient to consider only the minimal strong extensions $A \leq_o A\bar{b}$. Moreover, by Condition (D2), it suffices to consider only the trivial one-element extensions (cf. 3.1(4)).

Let $A \leq_o M$, $|A| \leq \aleph_0$ and suppose that $A \leq_o Ab$ is a trivial extension. Let $A' := \text{cl}_{<}(A)$, then by Proposition 3.21 we have that $A' \leq_o M$ and $|A'| \leq \aleph_0$. Consider the type $p(x/A')$ that says, for every $n < \omega$ and for every finite subset $D \subseteq A'$, that $Dx \leq_o \text{Dcl}_{<}^n(x)$ and that Dx is a trivial extension of D . Notice that this is first-order expressible by reasoning as in Remark 3.7. Then, by Condition (D1), it follows that $p(x/A')$ is satisfiable. We let $b_* \in M$ be a realisation of this type. Clearly $Ab_* \cong_A Ab$. We show that also $Ab_* \leq_o M$ holds.

First of all, notice that since $Ab_* \leq_o M$, it follows by Proposition 3.21 that $\text{cl}_{<}(Ab_*) \leq_o M$, and thus $A'\text{cl}_{<}(b_*) \leq_o M$. Moreover, since b_* satisfies $Db_* \leq_o \text{Dcl}_{<}^n(b_*)$ for every $D \subseteq_\omega A'$, it follows that $A'b_* \leq_o A'\text{cl}_{<}(b_*)$. By transitivity we conclude that $A'b_* \leq_o M$. Now, since b_* is trivial over A' , notice that $\text{gcl}_{A'b_*}(A'b_* \setminus Ab_*) = \text{gcl}_{A'}(A' \setminus A)$. Since we also have that $A \leq_o A'$, it follows by Condition 3.2(K) that $Ab_* \leq_o A'b_*$. By transitivity we conclude that $Ab_* \leq_o M$.

We are now ready to prove the full statement of the Lemma. Consider an arbitrary extension $A \leq_o B$ with $A \leq_o M$ and both A and $B \setminus A$ countable. Fix an enumeration $(b_i)_{i < \omega}$ of $B \setminus A$ and consider the infinitary type $p(\bar{x}, A)$, where $\bar{x} = (x_i)_{i < \omega}$, which says the following:

- (i) for all $n < \omega$, $p \upharpoonright \{x_0, \dots, x_{n-1}\} = \text{tp}^{\text{qf}}(b_0 \dots b_{n-1}/A)$;
- (ii) for all finite $A_0 \subseteq A$ and for all $n < \omega$, $A_0 x_0 \dots x_{n-1} \leq_o M$.

Now, condition (i) can be expressed simply by adding formulas to $p(\bar{x}/A)$ that say that the isomorphic type of x_1, \dots, x_n over A is exactly $\text{tp}^{\text{qf}}(b_0 \dots b_{n-1}/A)$. Condition (ii) is more subtle, but one can see that it is indeed a first-order condition in the light of Remark 3.7. In particular, Clause (ii) is equivalent to let:

$$\forall \bar{y} (\neg \delta_{(A_0 b_0 \dots b_{n-1}, C)}(\bar{a}, x_0 \dots x_{n-1}, \bar{y})) \in p(\bar{x}, A),$$

for all $n < \omega$, $A_0 \subseteq_\omega A$, $A_0 b_0 \dots b_{n-1} \not\leq_o C$ with $C \in \mathcal{K}$, and where \bar{a} is an enumeration of A_0 and the formula $\delta_{(A_0 b_0 \dots b_{n-1}, C)}(\bar{a} x_0 \dots x_{n-1}, \bar{y})$ is defined as in Remark 3.7.

Let $n < \omega$, then since $A \leq_o B$ it follows in particular that $A \leq_o A b_0 \dots b_{n-1}$ (recall 3.6). Since we have already proven the Lemma for the case of finite extensions, it follows that there are $c_0, \dots, c_{n-1} \in M$ such that $\text{tp}^{\text{qf}}(b_0 \dots b_{n-1}/A) = \text{tp}^{\text{qf}}(c_0 \dots c_{n-1}/A)$ and $A c_0 \dots c_{n-1} \leq_o M$. Thus, the type $p(\bar{x}, A)$ is consistent. Since M is \aleph_1 -saturated it then follows that $p(\bar{x}, A)$ is realised in M . Thus we can find a sequence $(d_i)_{i < \omega}$ in M such that $\text{tp}^{\text{qf}}((d_i)_{i < \omega}/A) = \text{tp}^{\text{qf}}((b_i)_{i < \omega}/A)$ and, additionally, $A d_0 \dots d_{n-1} \leq_o M$ for all $n < \omega$. By 3.6 this immediately entails that $A \cup \{d_i : i < \omega\} \leq_o M$. This completes the proof of the lemma. \blacksquare

Remark 3.31. We point out that the previous proof can be easily adapted to show that, if $M \models T^+$ is κ^+ saturated (in the language expanded by symbols $<$ and $P_<$ encoding an HF-order), $A \subseteq M$ and $A \leq_o B$ is a strong extension with $|B| \leq \kappa$, then there is an embedding $f : B \rightarrow M$ with $f \upharpoonright A = \text{id}$ and $f(B) \leq_o M$. This is analogous to the situation from [4, 4.22]. We notice that this is not an accident, as in particular the theories of open graphs defined in [4, 4.20] are a specific instance of the present setting (cf. Section 4.1 below).

Remark 3.32. By Lemma 3.30, it is sufficient to verify Assumptions 3.30(D1) and 3.30(D2) in order to show that 3.28(C1) holds. This is how we will proceed in Section 4 later, where we show that 3.30(D1) and 3.30(D2) are satisfied in all our intended applications. However, we also decided to keep 3.28(C1) and 3.30(D1)–3.30(D2) separate, since to our eyes they have a very different role. While 3.28(C1) captures the essential ingredients of the completeness proof, and also relates to the failure of finite closure, 3.30(D1) and 3.30(D2) are simply technical conditions that we used to prove that 3.28(C1) is verified. We also notice that the assumption that M is \aleph_1 -saturated in 3.30(D2) is used only in the case of n -open graphs from Section 4.1, and it is not needed to verify 3.30(D2) in the other concrete examples.

Recall from 2.2 that we denote by $\text{acl}_B^{\text{qf}}(A)$ the algebraic closure of A in some model B obtained by considering only quantifier-free formulas. We prove the following lemma, which follows essentially from Condition 3.2(H).

Lemma 3.33. Let $B \models T^+$ and $A \subseteq B$ such that $A = \text{acl}_B^{\text{qf}}(A)$ then $A \leq_o B$.

Proof. Suppose there is a finite $B_0 \subseteq B$ such that $A \cap B_0 \not\leq_o B_0$ but $B_0 \setminus A \not\subseteq \text{acl}_B^{\text{qf}}(A)$. Since $B \models T^+$, it follows that for every $n < \omega$ we can find some $C_n \subseteq B$

such that $C_n \in \mathcal{K}$ and C_n contains at least n -many copies of B_0 over $A \cap B_0$. This contradicts 3.2(H). ■

Theorem 3.34. *The theory T^+ is complete.*

Proof. Let $M, N \models T^+$, then by 3.28(C1) we can also assume w.l.o.g. that they are countably (\mathcal{K}, \leq_o) -saturated. We claim that the following set \mathcal{I} is a back-and-forth system between M and N :

$$\mathcal{I} = \{f : A \cong B : A \leq_o M, B \leq_o N \text{ and } |A|, |B| \leq \aleph_0\}.$$

Clearly this suffices, as then by [35, Exercise 1.3.5] we have that M and N are elementary equivalent. Let $f : A \cong B$, where $A \leq_o M$, $B \leq_o N$, $|A| = |B| \leq \aleph_0$. Let $a \in M \setminus A$ and notice that by Lemma 3.33 we immediately have that $\text{acl}_M(Aa) \leq_o M$. Moreover, we clearly have that $A \leq_o \text{acl}_M(Aa)$ and that $\text{acl}_M(Aa)$ is countable. Since N is countably (\mathcal{K}, \leq_o) -saturated it then follows that $f : A \rightarrow B$ lifts to an embedding $\hat{f} : \text{acl}_M(Aa) \rightarrow N$ satisfying $\hat{f}(\text{acl}_M(Aa)) \leq_o N$. This means that $\hat{f} \in \mathcal{I}$ and completes our proof. ■

Theorem 3.35. *Let $A \subseteq B$ be such that $A, B \models T^+$. The following are equivalent:*

- (1) $A = \text{acl}_B(A)$;
- (2) $A \leq_o B$;
- (3) $A \preceq B$.

Proof. This proof generalises [1, Thm. 2.28] to our setting. Direction (1) \Rightarrow (2) follows from Lemma 3.33, and direction (3) \Rightarrow (1) is obvious. We consider the remaining direction (2) \Rightarrow (3). In addition to the assumption that $A \leq_o B$, by 3.28(C1) (and Remark 3.7) we can also assume w.l.o.g. that A and B are countably (\mathcal{K}, \leq_o) -saturated. Now, notice that if $M \leq_o A$ then by transitivity $M \leq_o B$ and, if $N \leq_o B$ and $N \subseteq A$ then it follows from 3.2(E) that $N \leq_o A$. Using this and arguing as in the proof of 3.34 one shows that $A \preceq B$, as in fact for every finite set $D \subseteq A$ of parameters we can consider the following set of isomorphisms \mathcal{I} :

$$\{f : M \cong N : D \subseteq M \leq_o A, D \subseteq N \leq_o B, |M|, |N| \leq \aleph_0, f \upharpoonright D = \text{id}_D\}.$$

By reasoning as in 3.34 one can show that \mathcal{I} is a back-and-forth system, thus completing the proof. ■

We conclude this section by finally showing that, under the assumptions from 3.28, the theory T^+ is stable. We notice that Assumption 3.28(C2) is the fundamental ingredient of the stability proof. This replicates the proof of stability for open projective planes from [24] in the present abstract setting.

Notation 3.36. We denote by \mathfrak{M} the monster model of the theory T^+ in the language expanded with an HF-order. Thus in particular \mathfrak{M} satisfies the saturation property from 3.31. When dealing with the monster model \mathfrak{M} we usually omit the indices and write simply $\text{tp}(a/B)$ instead of $\text{tp}_{\mathfrak{M}}(a/B)$ to refer to the type of an element, and we write simply $\text{acl}(B)$ instead of $\text{acl}_{\mathfrak{M}}(B)$ to refer to the algebraic closure of a set. We do the same for quantifier-free types and quantifier-free algebraic closures.

Theorem 3.37. *The theory T^+ is stable.*

Proof. Let κ be such that $\kappa^{\aleph_0} = \kappa$, $A \models T^+$ with $|A| = \kappa$ and $A \preccurlyeq \mathfrak{M}$. By Proposition 3.35 it follows that $A \leq_o \mathfrak{M}$ and thus by 3.15 we can find an HF-ordering $<$ of \mathfrak{M} over A . Additionally, it also follows from 3.15 that we can extend $<$ to a full HF-order $<'$ of \mathfrak{M} with A as an initial segment.

Let $a, b \in \mathfrak{M} \setminus A$, let $A_0 = \text{cl}_{<}(a)$ and $B_0 = \text{cl}_{<}(b)$. By Proposition 3.21 we have that $|A_0| \leq \aleph_0$, $|B_0| \leq \aleph_0$, and also that $AA_0 \leq_o \mathfrak{M}$, $AB_0 \leq_o \mathfrak{M}$. Since $A \models T^+$, it follows in particular that A already contains infinitely many elements of each sort, thus there are two models $C(AA_0)$ and $C(AB_0)$ that satisfy the conditions from 3.28(C2). With slight abuse of notation, we identify $C(AA_0)$ and $C(AB_0)$ with their image in \mathfrak{M} given by 3.28(C2). It then follows from Condition 3.28(C2) that $C(AA_0) \models T^+$, $C(AB_0) \models T^+$ and $A \leq_o C(AA_0) \leq_o \mathfrak{M}$, $A \leq_o C(AB_0) \leq_o \mathfrak{M}$.

Now, suppose w.l.o.g. that $A_0 \cap A = B_0 \cap A$ and that there is an isomorphism $f : A_0 \cong B_0$ such that $f|_{A_0 \cap A} = \text{id}_{A_0 \cap A}$ and $f(a) = b$. Let $h : AA_0 \rightarrow AB_0$ be such that $f \subseteq h$ and $h|_A = \text{id}_A$, then h is an isomorphism. By 3.28(C2) it follows that h extends to an isomorphism $\hat{h} : C(AA_0) \cong_A C(AB_0)$. By the homogeneity of the monster model \mathfrak{M} it follows that \hat{h} extends to an A -automorphism of \mathfrak{M} . We conclude that, modulo κ -many elements, the type of an element $a \in \mathfrak{M}$ over a set A of size κ is determined by its type over $\text{cl}_{<}(a)$. Since $|\text{cl}_{<}(a)| \leq \aleph_0$ and $\kappa^{\aleph_0} = \kappa$, this completes our proof. \blacksquare

3.5. Forking independence in T^+

Up to this point, we have proved that the theory T^+ is both complete and stable, provided that the conditions from 3.2 and the assumptions from 3.28 are satisfied. Since T^+ is stable, it follows that the relation \perp of forking independence is well-behaved, in the sense that it satisfies the key properties from [35, Thm. 8.55]. In this section we study the relation of forking independence \perp in models of T^+ and we show that, under the (quite minimal) additional assumptions from 3.43 (which we introduce below), we have that \perp is equivalent to a relation \perp^\otimes , defined in purely combinatorial terms. We first recall some preliminary notions about forking, and we introduce some preliminary definitions.

Remark 3.38. In the following, we always use the symbol \perp to denote the relation of *forking independence*, and we refer the reader to [35, §7.1] for its definition. Let $\bar{a} \in \mathfrak{M}^{<\omega}$ and $B, C \subseteq \mathfrak{M}$, we write $\bar{a} \perp_B C$ if the type $\text{tp}(\bar{a}/BC)$ *does not fork* over B , and we write $\bar{a} \not\perp_B C$ if the type $\text{tp}(\bar{a}/BC)$ *does fork* over B . Since by 3.37 the theory T^+ is stable, the relation of forking independence is well-behaved and satisfies the properties from [35, Thm. 8.5.5]. We point out that a consequence of these properties is that, in stable theories, $B \perp_A C$ is equivalent to $\text{acl}(AB) \perp_{\text{acl}(A)} \text{acl}(AC)$.

The key ingredient in the characterisation of forking independence is the notion of free algebraic completion $F(A)$ (we define this later in 3.41), which generalises Hall's free completion procedure from the setting of projective planes to the setting of an arbitrary Hrushovski class (\mathcal{K}, \leq_o) satisfying the assumptions from 3.2. To this end, we start by first introducing the notion of free amalgam in (\mathcal{K}, \leq_o) , and the notion of degree of an algebraic extension.

Definition 3.39 (Free amalgam). Let $A, B, C \models T_\forall$ with $A \subseteq B$, $A \subseteq C$ and $A = B \cap C$, the *free amalgam* of B and C over A is the structure $B \otimes_A C$ such that:

- (1) the domain of $B \otimes_A C$ is $B \cup C$;
- (2) every incidence symbols $R \in L_i$ is interpreted by letting $R^{B \otimes_A C} = R^B \cup R^C$;
- (3) every parallelism symbols $P \in L_p$ is interpreted by letting $B \otimes_A C \models P(b, c, \bar{d})$, for $b, c \in B \cup C$ and $\bar{d} \in (B \cup C)^{<\omega}$ if one of the following conditions hold:
 - (i) $B \models P(b, c, \bar{d})$ (so in particular $b, c \in B$ and $\bar{d} \in B^{<\omega}$);
 - (ii) $C \models P(b, c, \bar{d})$ (so in particular $b, c \in C$ and $\bar{d} \in C^{<\omega}$);
 - (iii) there is some $a \in A$ such that $B \models P(b, a, \bar{d})$ and $C \models P(a, c, \bar{d})$ (so in particular $\bar{d} \in A^{<\omega}$).

Definition 3.40. Let $A \subseteq B \in \mathcal{K}$ and suppose that the extension (A, B) is algebraic in \mathcal{K} (cf. 3.1 above). The *degree* of the extension (A, B) is the greatest number $n < \omega$ such that there is some $C \in \mathcal{K}$ containing n -many isomorphic copies of B over A .

Using the previous definitions, we can define the notion of *free algebraic completion* of some $A \models T_\forall$. This was essentially already introduced by Funk and Strambach in [21, §1.4], where they provide a uniform model-theoretic definition of all instances of free completions from the incidence geometry literature. However, we notice that we take minimal algebraic strong extensions as the building blocks of free completions, while for Funk and Strambach this role is played by the so-called bricks (cf. [21, Def. 3]).

Definition 3.41 (Free completion). Let $A \models T_\forall$, we define its free completion $F(A)$ recursively as follows. Let $A_0 = A$ and suppose A_n is given. We let $(A_n, A_n B_i)_{i < m}$ enumerate all minimal strong algebraic extensions of A_n , up to isomorphism. For every $i < m$, we let n_i be the degree of $(A_n, A_n B_i)$, and for every $1 \leq j \leq n_i$, we let $A_n B_i^j$ be isomorphic copies of $A_n B_i$ such that $A_n B_i^j \cap A_n B_i^k = A$ for $j \neq k$. Then we define:

$$\begin{aligned} A_n^0 &:= A_n B_0^1 \otimes_{A_n} A_n B_0^2 \otimes_{A_n} \cdots \otimes_{A_n} A_n B_0^{n_0} \\ A_n^{i+1} &:= A_n^i B_{i+1}^1 \otimes_{A_n^i} A_n^i B_{i+1}^2 \otimes_{A_n^i} \cdots \otimes_{A_n^i} A_n^i B_{i+1}^{n_{i+1}} \\ A_{n+1} &:= \bigcup_{i < m} A_n^i, \end{aligned}$$

and, finally, we let $F(A) = \bigcup_{n < \omega} A_n$ be the *free (algebraic) completion* of A .

Remark 3.42. By the previous definitions 3.39 and 3.41 the amalgam $B \otimes_A C$ and the free completion $F(A)$ are well-defined L -structures, but notice that they are not necessarily models of T_\forall . To characterise forking independence in T^+ we need additionally that $F(A)$ is a model of T_\forall , and to this end we add the following Assumption 3.43(C4). Notice that in many cases (the non-degenerate cases of the geometries from Sections 4.2-4.6) it will actually be the case that $F(A) \models T^+$, *but this is not necessary to characterise forking*. In fact, our characterisation of the forking independence relation \perp also applies to cases, e.g., the pseudoplane, where $F(A)$ is not a model of T^+ (cf. Sec. 4.1 below).

We introduce the following Assumption 3.43. As we just remarked, Assumption 3.43(C4) guarantees that $F(A) \models T_\forall$. On the other hand, Assumption 3.43(C5) makes sure that, if $A \leq_o F(A) \subseteq B$, then it follows additionally that $F(A) \leq_o B$. This captures a key property of algebraic extensions in all the geometries that we consider. In terms of HF-orderings, one can thus think of algebraic strong extensions as the “first” extensions occurring in the order.

Assumption 3.43. Let (\mathcal{K}, \leq_o) and T^+ be respectively as in 3.2 and 3.22, and assume that T^+ satisfies all conditions from 3.28. In this section we also assume the following conditions.

- (C4) Let $A, B, C \models T_\forall$ with $A \leq_o B$ minimal, $A \leq_o C$ and $A = B \cap C$. If $A \leq_o B$ is non-algebraic, then $B, C \leq_o B \otimes_A C$ and $B \otimes_A C \models T_\forall$. If $A \leq_o B$ is algebraic of degree k (recall 3.40), and C contains $< k$ isomorphic copies of B over A , then $B, C \leq_o B \otimes_A C$ and $B \otimes_A C \models T_\forall$.
- (C5) Let $A \leq_o C \models T_\forall$ and let $B \subseteq C \setminus A$ be such that $A \leq_o AB$ is a minimal algebraic extension, then $AB \leq_o C$.

The following Proposition captures the key properties of the free completion $F(A)$ and it is essentially a corollary of the previous assumptions.

Proposition 3.44. Let $A \models T_\forall$ and suppose Assumption 3.43 holds, then $F(A) \models T_\forall$, $A \leq_o F(A)$, $|F(A)| = |A| + \aleph_0$, and:

- (a) every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : F(A) \cong F(B)$;
(b) for every $M \models T^+$ and $h : A \rightarrow M$ with $h(A) \leq_o M$ there is $\hat{h} : F(A) \rightarrow M$ s.t. $\hat{h} \upharpoonright A = h$ and $\hat{h}(F(A)) \leq_o M$.

Proof. The fact that $|F(A)| = |A| + \aleph_0$ follows immediately by construction. Similarly $F(A) \models T_\forall$ since for every $n < \omega$, $A_n \models T_\forall$, by Assumption 3.43(C4). Now, for every $n < \omega$ in the construction of $F(A)$, it follows from 3.43(C4) that $A_n \leq_o A_n^0 \leq_o \dots \leq_o A_{n+1}$. Inductively, we obtain that $A \leq_o F(A)$.

Consider (a). By the construction of $F(A)$, it is immediate to extend an isomorphism $h : A \cong B$ to an isomorphism $\hat{h} : F(A) \cong F(B)$.

Consider (b). Let $h : A \rightarrow M \models T^+$ with $h(A) \leq_o M$. Suppose inductively that we have found an embedding $h_n \supseteq h$ such that $h_n : A_n \rightarrow M$ and $h_n(A_n) \leq_o M$. Now, since $h_n(A_n) \leq_o M$, every strong algebraic extension of $h_n(A_n)$ amalgamates with every finite subset $D \subseteq M$ (as we have $h_n(A) \leq_o D$). It follows from the definition of T^+ (cf. 3.22) that every minimal strong extension of $h_n(A_n)$ is realised in M . Thus we can extend h_n to an embedding $h_{n+1} : A_{n+1} \rightarrow M$. Moreover, it follows from 3.43(C5) that $h_{n+1}(A_{n+1}) \leq_o M$. Finally, let $\hat{h} = \bigcup_{n < \omega} h_n$, then clearly $\hat{h} : F(A) \rightarrow M$ is an embedding and $\hat{h}(F(A)) \leq_o M$. ■

Remark 3.45. The previous proposition shows that, under the conditions from 3.43, the free algebraic completion $F(A)$ *almost* satisfies the conditions from 3.28(C2). We stress the “almost”: the key complication is that nothing ensures us that the structure $F(A)$ is a model of T^+ , and in particular it could even be that $F(A) = A$. This is what actually happens in those geometries where the class (\mathcal{K}, \leq_o) does not contain any algebraic extension $A \leq_o B$. Most notably, this is what happens in the pseudoplane and in the n -open graphs from Section 4.1. We stress again that the characterisation of forking independence which we give in this section will also apply to these cases, which somehow represent the degenerate case of the general geometric setting which we are studying.

Notation 3.46. Suppose $A \leq_o M \models T^+$, from now on, with some slight abuse of notation, we identify $F(A)$ with its isomorphic image in M (which exists by Proposition 3.44) and we simply write $F(A) \leq_o M$. Notice that, in particular, this justifies the use of the equality symbol (rather than the congruence one) in the Definition 3.55 below.

Remark 3.47. We also notice that, if $A \leq_o B \otimes_A C$, then (by Condition 3.2(K) and Assumption 3.43(C4)) $A \leq_o B$ is algebraic if and only if $AC \leq_o B \otimes_A C$ is algebraic. By Def. 3.41, this in particular means that, if $A \leq_o C$, then $F(C)$ always contains an isomorphic copy of $F(A)$, and additionally $F(A) \leq_o F(C)$. Notice that this is not necessarily the case for arbitrary $A \subseteq C \in \mathcal{K}$, since given an algebraic extensions $A \leq_o B$ it might be the case that $B \otimes_A C \notin \mathcal{K}$.

Using the previous notion of free completion, we can provide a characterisation of the algebraic closure in models of T^+ . The next lemma first shows that, if $A \leq_o B \models T^+$, then every type over A is determined by its quantifier-free part.

Lemma 3.48. Let $A \leq_o B \models T^+$ and $\bar{b}, \bar{c} \in B^n$ for some $n < \omega$. If $\text{tp}^{\text{qf}}(\bar{b}/A) = \text{tp}^{\text{qf}}(\bar{c}/A)$ then $\text{tp}(\bar{b}/A) = \text{tp}(\bar{c}/A)$.

Proof. Let $\bar{b} = (b_1, \dots, b_n)$ and $\bar{c} = (c_1, \dots, c_n)$, let $f_0 : A \cup \{b_1, \dots, b_n\} \rightarrow A \cup \{c_1, \dots, c_n\}$ be a partial isomorphism witnessing that $\text{tp}^{\text{qf}}(\bar{b}/A) = \text{tp}^{\text{qf}}(\bar{c}/A)$, in particular $f_0 \upharpoonright A = \text{id}_A$ and $f_0(b_i) = c_i$ for all $1 \leq i \leq n$. By Proposition 3.44 this extends to an isomorphism $f_1 : F(A\bar{b}) \rightarrow F(A\bar{c})$. Since w.l.o.g. we can assume that A contains at least $m_{\mathcal{K}}$ -many elements of every sort, it follows from Assumption 3.28(C2) that f_1 can be extended to an isomorphism $f_2 : C(F(A\bar{b})) \rightarrow C(F(A\bar{c}))$ between two models $C(F(A\bar{b}))$ and $C(F(A\bar{c}))$ of T^+ . Since again by 3.28(C2) we have that $C(F(A\bar{b}))$ and $C(F(A\bar{c}))$ are essentially two elementary submodels of B containing A , we conclude that f_2 is a partial elementary map witnessing that $\text{tp}(\bar{b}/A) = \text{tp}(\bar{c}/A)$. ■

Remark 3.49. In most of our applications, we have by Corollary 3.70 that the theory T^+ does not have a prime model and is not model complete. It follows in particular that T^+ does not eliminate quantifier and therefore the assumption that $A \leq_o B$ in the previous lemma is necessary. This is the case for example in projective planes and generalised n -gons (cf. also [24] and [1]).

A fundamental ingredient in the characterisation of algebraic closure is the notion of intrinsic closure. We recall the following definition from [3, 2.22].

Definition 3.50. Let $A \subseteq B \models T_{\forall}$, the *intrinsic closure* $\text{icl}_B(A)$ is the set containing all minimal closed extensions of A , i.e., we have:

$$\text{icl}_B(A) := \bigcup \{C : C \subseteq B, A \not\leq_o C \text{ and } A \leq_o C' \text{ for all } C' \subsetneq C\}.$$

Proposition 3.51. Let $A \subseteq B \models T^+$, then $\text{acl}_B(A) = F(\text{icl}_B(A))$.

Proof. By 3.2(H) we immediately have that $\text{icl}_B(A) \subseteq \text{acl}_B(A)$. Moreover, by Definition 3.41 of the free completion $F(\text{icl}_B(A))$ it follows that $F(\text{icl}_B(A)) \subseteq \text{acl}_B(A)$. Consider the converse direction. By Definition 3.50 it follows that $\text{icl}_B(A) \leq_o B$, so then by Proposition 3.44 we obtain $F(\text{icl}_B(A)) \leq_o B$. Thus, by Lemma 3.48, it is sufficient to show that $\text{acl}_B^{\text{qf}}(F(\text{icl}_B(A))) = F(\text{icl}_B(A))$. Consider an element $c \in \text{acl}_B^{\text{qf}}(A) = \text{acl}_B^{\text{qf}}(F(\text{icl}_B(A)))$ and suppose towards contradiction that $c \notin F(\text{icl}_B(A))$. Then, there is a step A_n from the construction of the free completion $F(\text{icl}_B(A))$ (cf. 3.39) such that c , together possibly with some other elements, is algebraic over A_n . Let C be a minimal extension of A_n which contains c and is algebraic over A . Then since $F(\text{icl}_B(A)) \leq_o B$ we also have that $A_n \leq_o C$. It then follows from Definition 3.39 that $C \subseteq A_{n+1} \subseteq F(\text{icl}_B(A))$, and so $c \in F(\text{icl}_B(A))$, contrary to our assumption. We conclude that $\text{acl}_B(A) = F(\text{icl}_B(A))$. ■

We next introduce the notion of canonical amalgam and the independence relation \downarrow^\otimes . We then prove, generalising [24], [1] and [4], that \downarrow^\otimes is equivalent to the relation of forking independence \downarrow in models of T^+ .

Definition 3.52. Let $A \leq_o B \models T_\forall$, we say that A is \leq_o -algebraically closed in B if there is no $A \subsetneq C \subseteq B$ such that $A \leq_o C$ is an algebraic extension.

Definition 3.53 (Canonical amalgam). Let $A, B, C \models T_\forall$ be such that $A \leq_o B$, $A \leq_o C$, $B \cap C = A$ and suppose that A is \leq_o -algebraically closed both in B and C . The *canonical amalgam* $B \oplus_A C$ of B and C over A is the structure

$$B \oplus_A C := F(B \otimes_A C),$$

i.e., it is the free completion of the free amalgam $B \otimes_A C$.

Proposition 3.54. Let $A, B, C \models T_\forall$ be such that $A \leq_o B$, $A \leq_o C$, $B \cap C = A$ and suppose that A is \leq_o -algebraically closed both in B and C , then:

- (a) $B \otimes_A C \models T_\forall$ and $B \oplus_A C \models T_\forall$,
- (b) $A \leq_o B \leq_o B \otimes_A C \leq_o B \oplus_A C$ and $A \leq_o C \leq_o B \otimes_A C \leq_o B \oplus_A C$,
- (c) $F(B \otimes_A C) = F(F(B) \otimes_{F(A)} F(C))$,
- (d) given two isomorphisms $f_0 : B \cong_A B'$ and $f_1 : C \cong_A C'$ there is an isomorphism $f : B \oplus_A C \cong_A B' \oplus_{f(A)} C'$ such that $f|_B = f_0$ and $f|_C = f_1$.

Proof. Clauses (a) and (b) follow immediately from 3.43(C4) and Proposition 3.44. Clause (c) holds by Def. 3.41 and Remark 3.47. Clause (d) follows from Definition 3.39 and Proposition 3.44. \blacksquare

Definition 3.55. Let $A, B, C \subseteq \mathfrak{M}$ (and recall from 3.36 that \mathfrak{M} is the monster model of T^+), we define the relation \downarrow^\otimes by letting

$$B \downarrow_A^\otimes C \iff \text{acl}(ABC) = \text{acl}(AB) \oplus_{\text{acl}(A)} \text{acl}(AC).$$

We recall that we always assume the monster model \mathfrak{M} is sufficiently saturated and homogeneous in the language L expanded by symbols $<, P$ which encode a HF-order. In particular, it then follows by Remark 3.31 that if $A \leq_o B$ are small sets and $A \leq_o \mathfrak{M}$, then there is an embedding $f : B \rightarrow \mathfrak{M}$ such that $f|_A = \text{id}_A$ and $f(B) \leq_o \mathfrak{M}$. The strategy that we employ in the following arguments to characterise the relation of forking independence is exactly the same as in [1], i.e., we use the fact that in stable theories forking independence is the only independence relation satisfying the key properties from Fact 3.56 below (cf. [35, Thm. 8.5.10]).

Fact 3.56. Let T be a stable theory, then the relation of forking independence \downarrow is the only ternary relation on subsets of \mathfrak{M} which satisfies the properties below:

- (a) *Invariance:* let $A, B, B', C \subseteq \mathfrak{M}$, if $B \downarrow_A C$ and $\text{tp}(B/C) = \text{tp}(B'/C)$, then $B' \downarrow_A C$;
- (b) *Local Character:* there is a cardinal κ such that for all finite $B \subseteq \mathfrak{M}$ and $C \subseteq \mathfrak{M}$, there is a subset $A \subseteq \mathfrak{M}$ of size $\leq \kappa$ such that $B \downarrow_A C$;
- (c) *Weak Boundedness:* there is a cardinal μ such that, for all $B, B' \subseteq_\omega \mathfrak{M}$ and $A \subseteq C \subseteq \mathfrak{M}$, there are at most μ -many types $\text{tp}(B'/C)$ such that $B' \downarrow_A C$ and $\text{tp}(B'/A) = \text{tp}(B/A)$;
- (d) *Existence:* for all finite $B \subseteq \mathfrak{M}$ and $A \subseteq C \subseteq \mathfrak{M}$, there is some $B' \subseteq \mathfrak{M}$ such that $B' \downarrow_A C$ and $\text{tp}(B'/A) = \text{tp}(B/A)$;

- (e) *Transitivity*: let $B \subseteq_{\omega} \mathfrak{M}$ and $A \subseteq C \subseteq D \subseteq \mathfrak{M}$, if $B \downarrow_A C$ and $B \downarrow_C D$, then $B \downarrow_A D$;
- (f) *Weak monotonicity*: let $A, B, C, D \subseteq \mathfrak{M}$, if $B \downarrow_A C$ and $D \subseteq C$, then $B \downarrow_A D$.

By the fact above, to show that $\downarrow = \downarrow^{\otimes}$ it suffices to verify that the relation \downarrow^{\otimes} satisfies all the properties of forking independence.

Theorem 3.57. *The relation \downarrow^{\otimes} satisfies all the properties of forking independence from Fact 3.56. Thus, for any $A, B, C \subseteq \mathfrak{M}$, we have that $B \downarrow_A C$ is equivalent to $B \downarrow_A^{\otimes} C$.*

Proof. We verify that \downarrow^{\otimes} satisfies all the properties from Fact 3.56. By [35, Thm. 8.5.10] this establishes that $\downarrow = \downarrow^{\otimes}$.

Invariance: let $A, B, B', C \subseteq \mathfrak{M}$, if $B \downarrow_A^{\otimes} C$ and $\text{tp}(B/C) = \text{tp}(B'/C)$, then $B' \downarrow_A^{\otimes} C$.

Let π be a C -automorphism of \mathfrak{M} witnessing that $\text{tp}(B/C) = \text{tp}(B'/C)$ then $\text{acl}(AB'C) = \text{acl}(A\pi(B)C) = \pi(B) \oplus_A C = B' \oplus_A C$. This shows that $B \downarrow_A^{\otimes} C$ if and only if $B' \downarrow_A^{\otimes} C$.

Weak Monotonicity: let $A, B, C, D \subseteq \mathfrak{M}$, if $B \downarrow_A^{\otimes} C$ and $D \subseteq C$, then $B \downarrow_A^{\otimes} D$. Suppose w.l.o.g. that A is algebraically closed and assume that $\text{acl}(ABC) = \text{acl}(AB) \oplus_A \text{acl}(AC)$. If $D \subseteq C$, then we clearly have that $\text{acl}(AB) \cap \text{acl}(AD) = A$, so $\text{acl}(AB) \otimes_A \text{acl}(AD)$ is well-defined. It then follows (from Condition 3.2(K) and the definition of free amalgam) that

$$\text{acl}(AB) \otimes_A \text{acl}(AD) \leq_o \text{acl}(AB) \otimes_A \text{acl}(AC) \leq_o \text{acl}(ABC)$$

so by Lemma 3.51 it follows that $\text{acl}(AB) \oplus_A \text{acl}(AD)$ is algebraically closed, and thus $\text{acl}(ABD) = \text{acl}(AB) \oplus_A \text{acl}(AD)$. We conclude that $B \downarrow_A^{\otimes} D$.

Weak Boundedness: there is a cardinal μ such that, for all $B, B' \subseteq_{\omega} \mathfrak{M}$ and $A \subseteq C \subseteq \mathfrak{M}$, there are at most μ -many types $\text{tp}(B'/C)$ such that $B' \downarrow_A^{\otimes} C$ and $\text{tp}(B'/A) = \text{tp}(B/A)$.

We claim that $\mu = 1$ suffices. Suppose $B_0 \downarrow_A^{\otimes} C$, $B_1 \downarrow_A^{\otimes} C$ and $\text{tp}(B_0/A) = \text{tp}(B_1/A)$. Then there is an isomorphism $f : B_0 \rightarrow B_1$ such that $f \upharpoonright A = \text{id}_A$. Together with the identity map id_C , we obtain by Proposition 3.54 an isomorphism $\hat{f} : B_0 \oplus_A C \rightarrow B_1 \oplus_A C$ such that $\hat{f} \upharpoonright B_0 = f$ and $\hat{f} \upharpoonright C = \text{id}_C$. It follows that $\text{tp}^{\text{af}}(B_0/C) = \text{tp}^{\text{af}}(B_1/C)$. Finally, notice that by assumption we have

$$C \leq_o B_0 \oplus_A C = \text{acl}(AB_0C) \leq_o \mathfrak{M} \text{ and}$$

$$C \leq_o B_1 \oplus_A C = \text{acl}(AB_1C) \leq_o \mathfrak{M}.$$

So it follows from Lemma 3.48 that $\text{tp}(B_0/C) = \text{tp}(B_1/C)$.

Existence: for all finite $B \subseteq \mathfrak{M}$ and $A \subseteq C \subseteq \mathfrak{M}$, there is some $B' \subseteq \mathfrak{M}$ such that $B' \downarrow_A^{\otimes} C$ and $\text{tp}(B/A) = \text{tp}(B'/A)$.

By the weak monotonicity property (already proved above), we can assume without loss of generality that C is non-degenerate. Now, recall that $B' \downarrow_A^{\otimes} C$ is equivalent to $\text{acl}(B') \downarrow_{\text{acl}(A)}^{\otimes} \text{acl}(C)$, thus we assume without loss of generality that $A = \text{acl}(A)$, $B = \text{acl}(AB)$ and $C = \text{acl}(AC)$. Also, we assume that $B \cap C = A$, as if this is not the case we can find an isomorphic copy of B over A satisfying this constraint. It follows that the structure $B \oplus_A C$ is well-defined, and

$A \leq_o C \leq_o B \oplus_A C$. Moreover, by Assumption 3.28(C2) and the fact that C is non-degenerate, it follows that $C \leq_o C(B \oplus_A C)$. Then, by Remark 3.31, there is an embedding $f : C(B \oplus_A C) \rightarrow \mathfrak{M}$ such that $f(C(B \oplus_A C)) \leq_o \mathfrak{M}$ and $f \upharpoonright C = \text{id}_C$. Let $B' = f(B)$, then since C is non-degenerate, it follows from 3.28(C2) that $f(C(B \oplus_A C)) \preceq \mathfrak{M}$. Thus we obtain that $\text{tp}(B/A) = \text{tp}(B'/A)$. Moreover, since $B' \oplus_A C \leq_o f(C(B \oplus_A C)) \leq_o \mathfrak{M}$, it follows from Lemma 3.51 that $\text{acl}(AB'C) = B' \oplus_A C$, and thus $B \downarrow_A^\otimes C$.

Transitivity: let $B \subseteq_\omega \mathfrak{M}$ and $A \subseteq C \subseteq D \subseteq \mathfrak{M}$, if $B \downarrow_A^\otimes C$ and $B \downarrow_C^\otimes D$, then $B \downarrow_A^\otimes D$.

Suppose $B \downarrow_A^\otimes C$ and $B \downarrow_C^\otimes D$, then $\text{acl}(ABC) = \text{acl}(AB) \oplus_{\text{acl}(A)} \text{acl}(AC)$ and also $\text{acl}(BCD) = \text{acl}(BC) \oplus_{\text{acl}(C)} \text{acl}(CD)$. Then:

$$\begin{aligned} \text{acl}(ABD) &= \text{acl}(BCD) \\ &= \text{acl}(BC) \oplus_{\text{acl}(C)} \text{acl}(CD) \\ &= (\text{acl}(AB) \oplus_{\text{acl}(A)} \text{acl}(AC)) \oplus_{\text{acl}(C)} \text{acl}(CD) \\ &= \text{acl}(AB) \oplus_{\text{acl}(A)} \text{acl}(AD) \end{aligned}$$

which entails that $B \downarrow_A^\otimes D$.

Local Character: there is a cardinal κ such that for all finite $B \subseteq \mathfrak{M}$ and $C \subseteq \mathfrak{M}$, there is a subset $A \subseteq \mathfrak{M}$ of size $\leq \kappa$ such that $B \downarrow_A^\otimes C$.

We show that local character holds with respect to $\kappa = \aleph_0$. Let $B \subseteq \mathfrak{M}$ be finite and consider an arbitrary subset $C \subseteq \mathfrak{M}$. Since $\text{acl}(C) \leq_o \mathfrak{M}$ it follows from Corollary 3.16 that there is a HF-order $<$ of \mathfrak{M} over $\text{acl}(C)$. Thus, again by 3.16, we can extend $<$ to an HF-order $<$ of \mathfrak{M} with $\text{acl}(C)$ as an initial segment.

Consider now the set $B' = \text{cl}_<(B)$ and notice that by Proposition 3.21 we have that $B' \leq_o \mathfrak{M}$ and $|B'| \leq \aleph_0$. From Proposition 3.51 we obtain that $\text{acl}(B') = F(B')$. Let $A = B' \cap \text{acl}(C)$, we claim that $B' \downarrow_A^\otimes C$.

First, suppose that $c \in F(B') \cap \text{acl}(C)$. Since $c \in F(B')$, the distance between c and B' in the Gaifman graph of $F(B')$ is nonzero, i.e., $d(c, B'/F(B')) > 0$ (recall 2.3 and 2.6). Then, since $\text{acl}(C)$ is an initial segment of $<$ and $c \in \text{acl}(C)$, it follows that $c \in \text{cl}_<(B) = B'$. We thus conclude that $c \in A$ and so $\text{acl}(B') \cap \text{acl}(C) = F(B') \cap \text{acl}(C) \subseteq A$. Since $A \subseteq B'$ and $A \subseteq \text{acl}(C)$ we also obtain that $\text{acl}(B') \cap \text{acl}(C) = \text{acl}(A)$ (so notice also $A = \text{acl}(A)$).

Secondly, notice that by definition of the closure $\text{cl}_<(B)$ (cf. 3.19) and the fact that $\text{acl}(C)$ is an initial segment of $<$, we have for any element $c \in C$ incident with some $b \in B' = \text{cl}_<(B)$ that $c \in B'$ as well, which yields $c \in A$. Similarly, if for some $b \in B'$ and $c \in C$ we have that $\mathfrak{M} \models P(b, c, \vec{d})$, for some local equivalence relation $P \in L$ and some tuple $\vec{d} \in A^{<\omega}$, it follows by the same reasoning that there is some $c' \in A$ satisfying $\mathfrak{M} \models P(b, c', \vec{d})$ and $\mathfrak{M} \models P(c, c', \vec{d})$. By Definition 3.39 it follows that $B' \cup \text{acl}(C) = B' \otimes_A \text{acl}(C)$.

Now, since $\text{acl}(C)$ is an initial segment of $<$ we have that $\text{cl}_<(\text{acl}(C) \cup B') = \text{acl}(C) \cup B' \leq_o \mathfrak{M}$. In particular, the identity embedding $\text{acl}(C) \cup B' \leq_o \mathfrak{M}$ extends by Proposition 3.44 to an embedding $F(\text{acl}(C) \cup B') \leq_o \mathfrak{M}$. Notice that, since $B' \cup \text{acl}(C) = B' \otimes_A \text{acl}(C)$, we have $F(\text{acl}(C) \cup B') = F(\text{acl}(C) \otimes_A B')$. From Proposition 3.54(c) and the fact that $A = \text{acl}(A)$, it then follows that $F(B') \cup \text{acl}(C) = F(B') \otimes_A \text{acl}(C)$.

We conclude that:

$$\text{acl}(B') \cup \text{acl}(C) = \text{acl}(B') \otimes_A \text{acl}(C) \leq_o \mathfrak{M}$$

and so by 3.51 we have $\text{acl}(AB'C) = \text{acl}(B') \oplus_A \text{acl}(C)$. This shows that $B' \downarrow_A^\otimes C$. Since the relation \downarrow^\otimes is clearly symmetric by definition, it then follows from the weak monotonicity of \downarrow^\otimes that $B \downarrow_A^\otimes C$. This completes our proof. \blacksquare

We conclude this section by pointing out an equivalent characterisation of forking independence. Recall the definition of intrinsic closure from Definition 3.50, we then have the following corollary.

Corollary 3.58. Let $A, B, C \subseteq \mathfrak{M}$ and suppose that $\text{icl}(A)$ is \leq_o -algebraically closed both in $\text{icl}(AB)$ and $\text{icl}(AC)$. Then the following are equivalent:

- (1) $B \downarrow_A C$,
- (2) $B \downarrow_A^\otimes C$,
- (3) $\text{acl}(ABC) = F(\text{acl}(AB) \cup \text{acl}(AC)) = \text{acl}(AB) \oplus_{\text{acl}(A)} \text{acl}(AC)$,
- (4) $\text{icl}(ABC) = \text{icl}(AB) \cup \text{icl}(AC) = \text{icl}(AB) \otimes_{\text{icl}(A)} \text{icl}(AC)$.

Proof. Immediate from Theorem 3.57 and Proposition 3.51. \blacksquare

In particular we notice that, when there are no algebraic extensions in (\mathcal{K}, \leq_o) , and thus the free algebraic completion $F(A)$ trivialises, the previous corollary gives a more insightful characterisation of forking than Theorem 3.57. In particular, one can easily see that in the case of n -open graphs (cf. Sec. 4.1) the previous corollary provides the same characterisation of forking independence as the one in [4].

3.6. Failure of superstability

It was shown in [24] that the theory of infinite open projective planes is stable but not superstable. Later, this result was generalised in [1] to the case of open generalised n -gons (for $3 \leq n < \omega$). We observe in this section that *most, but not all* of the examples of theories T^+ that satisfy the conditions from 3.2, 3.28 and 3.43 are not superstable. More precisely, we identify in this section a combinatorial condition (3.60) and we prove that, when satisfied, it entails the non-superstability of T^+ . Interestingly, the following condition (3.60) behaves as a first dividing line among our class of intended applications. In fact, while all the geometries considered in Section 4 satisfy the assumptions from 3.2, 3.28 and 3.43, not all of them satisfy Condition 3.60 as well. The crucial counterexamples are the theories T_0^+ and T_1^+ from Section 4.1. While T_0^+ is not very interestingly, as it is simply the theory of an infinite set, T_1^+ is exactly the theory of the free pseudoplane, which is well-known to be ω -stable. This shows that our framework applies to both superstable and not superstable classes of models. Before introducing the combinatorial assumption 3.60, we introduce the technical notion of k -iterate of a finite configuration.

Definition 3.59. Let $C = \{c_0, \dots, c_n\} \in \mathcal{K}$ and $1 \leq k < \omega$, we define its k -iterate $I_k(C)$ as the structure D defined as follows:

- (1) the domain of D is $\{d_j^i : i < k, j \leq n\}$, where d_j^i and d_j^ℓ have the same sort for all $i < \ell < k$ and $j \leq n$, and with the additional constraint that $d_n^i = d_0^{i+1}$ for all $i < k$;
- (2) for every incidence symbol $R \in L_i$ we let $D \models R(d_{j_0}^i, d_{j_1}^i)$ if and only if $D \models R(c_{j_0}, c_{j_1})$, for all $j_0, j_1 \leq n$ and $i < k$;

- (3) for every (local) parallelism symbol $P \in L_p$ we let $D \models P(d_{j_0}^i, d_{j_1}^i, d_{j_2}^i, \dots, d_{j_\ell}^i)$ if $D \models P(c_{j_0}, c_{j_1}, c_{j_2}, \dots, c_{j_\ell})$, for all $j_0, j_1, \dots, j_\ell \leq n$ and $i < k$, and we then transitively close the relation $P(x, y, \bar{d})$ for all $\bar{d} \in D^{<\omega}$.

Assumption 3.60. In addition to the assumptions from 3.28 and 3.43, in this section we also assume that T^+ satisfies the following condition:

- (C6) there is a configuration $C = \{c_0, c_1, \dots, c_n\} \in \mathcal{K}$ such that:
- (a) $c_0 c_1 \leq_o C$ and there is no relation between c_0 and c_1 ;
 - (b) $c_n \not\leq_o C$ and c_n has the same sort of c_0 ;
 - (c) for every $k < \omega$, the k -iterate of C from Definition 3.59 is in \mathcal{K} .

Remark 3.61. We briefly mention that one cannot indeed find such a configuration in the theory T_1^+ of the pseudoplane. This is because, in any configuration $C \in \mathcal{K}$, where \mathcal{K} is the class of finite partial models of T_1^+ , we have for every element $c \in C$ that $c \leq_o C$, as it is easy to verify from Definition 4.2.

As the next proof involves some subtle constructions, we spend some words to provide the reader with some background intuition. The key point of the following proof is to construct an element d_n^0 whose intrinsic closure is infinite. In fact, if we can find an infinite subset $A \subseteq \text{icl}(d_n^0)$ which is not contained in the algebraic closure of any of its finite subsets A' , then we are done, since the characterisation of forking from Theorem 3.57 will then make sure that $d_n^0 \not\downarrow_{A'} A$ for all finite $A' \subseteq_\omega A$ (so as to have a witness for the failure of superstability). In the following proof we do this essentially by constructing d_n^0 so that there is a set $A \leq_o \mathfrak{M}$ such that $A \subseteq \text{icl}(d_n^0)$ and, additionally, there is no relation between the elements of A . This is similar to the configuration from [24, 9.1-9.2], which in turns provides a concrete example of our abstract construction in the setting of free projective planes.

Remark 3.62. We recall that a stable theory T is superstable if, for all finite $B \subseteq \mathfrak{M}$ and $C \subseteq \mathfrak{M}$, there is a finite subset $A \subseteq_\omega C$ such that $B \downarrow_A C$ (cf. [35, Def. 8.6.3]). Thus, superstable theories are exactly those stable theories where the property of local character from Fact 3.56 holds already with respect to finite subsets.

Theorem 3.63. *Suppose T^+ satisfies the assumptions from 3.28, 3.43 and also the assumption from 3.60, then T^+ is not superstable.*

Proof. Let $C = \{c_0, c_1, \dots, c_n\} \in \mathcal{K}$ be as in 3.60. Let $A = \{a_i : i < \omega\} \leq_o \mathfrak{M}$ be a countable set containing infinitely many elements of the same sort as c_1 , with no relation between them. By the saturation of \mathfrak{M} we can find a copy C_0 of C such that $A \leq_o AC_0 \leq_o \mathfrak{M}$ and a_0 plays the role of c_0 in C . Thus letting $C_0 = \{c_0^0, c_1^0, \dots, c_n^0\}$ we have $a_0 = c_0^0$. Now, suppose inductively that we have found configurations C_0, \dots, C_k such that they are all isomorphic to the configuration C from 3.60 and

$$A \leq_o AC_0 \leq_o \dots \leq_o AC_0 \dots C_{k-1} \leq_o AC_0 \dots C_{k-1} C_k \leq_o \mathfrak{M}.$$

Then for every $\ell \leq k$ let $C_\ell = \{c_0^\ell, c_1^\ell, \dots, c_n^\ell\}$ be the enumeration inherited from the isomorphism with C . Additionally, we assume inductively that $c_0^{\ell+1} = c_n^\ell$ for all $\ell < k$ and $c_1^\ell = a_\ell$ for all $\ell \leq k$. Then, by 3.60(c), we can iterate this procedure and obtain (by 3.60(a) and 3.2(K)) that $AC_0 \dots C_{k-1} C_k \leq_o AC_0 \dots C_k C_{k+1}$, where C_{k+1} is a configuration isomorphic to C from 3.60, but where $c_0^{k+1} = c_n^k$ and

$c_1^{k+1} = a_{k+1}$. By the saturation of \mathfrak{M} we then have (modulo isomorphism) that $AC_0 \dots C_k C_{k+1} \leq_{\circ} \mathfrak{M}$. Thus we can continue this construction for all $k < \omega$.

Now, using the saturation of \mathfrak{M} , we can find a sequence of configurations $(D_i)_{i < \omega}$ in \mathfrak{M} such that, for every $i < \omega$, $D_i = \{d_0^i, d_1^i, \dots, d_n^i\}$ is isomorphic to C , $d_n^i = d_0^{i-1}$ and $d_1^i \in A$. Moreover, for all $i < \omega$:

$$A \leq_{\circ} A \cup \bigcup_{i < j} D_j \leq_{\circ} A \cup \bigcup_{i \leq j} D_j \leq_{\circ} \mathfrak{M}.$$

Intuitively, while the chain of configurations $(C_i)_{i < \omega}$ is *ascending*, the chain of configurations $(D_i)_{i < \omega}$ is *descending*. We now want to show that $d_n^0 \not\downarrow_{A'} A$ for all finite $A' \subseteq_{\omega} A$. Since A is well-ordered we can assume without loss of generality that A' is an initial segment of A . Also, we assume without loss of generality that every element of A occurs in some D_i for some $i < \omega$.

First, we claim $A \not\subseteq \text{acl}(A')$. Let $a_n \in A \setminus A'$, this clearly exists because A' is finite. Then since A consists of elements with no relation between them, we have (by Condition 3.2(L)) that $A' \leq_{\circ} A' a_n \leq_{\circ} \mathfrak{M}$. Also, since a_n is a trivial extension of A' , we can find infinitely many extensions $A' \leq_{\circ} A' b_i$ such that $A' b_i \cong_{A'} A' a_n$ for all $i < \omega$. Then, all the elements b_i have the same quantifier-free type over A , thus since $A \leq_{\circ} \mathfrak{M}$ it follows from Lemma 3.48 that they have the same full type over A . Finally, this means that a_n is not algebraic over A' .

Secondly, we claim that every configuration D_i is contained in $\text{acl}(d_n^0)$ and so that $A \subseteq \text{acl}(d_n^0)$. By construction $d_n^0 \not\leq_{\circ} D_0$, thus it follows from Condition 3.2(H) that $D_0 \subseteq \text{acl}(d_n^0)$. Suppose inductively that we have showed $D_i \subseteq \text{acl}(d_n^0)$ for all $i \leq k$. Then since $d_n^{k+1} = d_0^k$ and $d_n^{k+1} \not\leq_{\circ} D_{k+1}$, it follows that $D_{k+1} \subseteq \text{acl}(d_0^k) \subseteq \text{acl}(d_n^0)$. Finally, since we assumed that $A = \{d_1^i : i < \omega\}$, it follows that $A \subseteq \text{acl}(d_n^0)$.

Now, obviously we have that $A \subseteq \text{acl}(A)$ and, by the previous claims, we have $A \subseteq \text{acl}(d_n^0)$ and $A \not\subseteq \text{acl}(A')$. It follows that

$$\text{acl}(d_n^0) \cup \text{acl}(A) \neq \text{acl}(d_n^0) \otimes_{\text{acl}(A')} \text{acl}(A),$$

since by Definition 3.39 in $P \otimes_X Q$ one must have that $P \cap Q = X$. Therefore, we obtain from Theorem 3.57 that $d_n^0 \not\downarrow_{A'} A$ for all finite subsets $A' \subseteq A$. It follows from Remark 3.62 that T^+ is not superstable. \blacksquare

3.7. Non-existence of prime models

We conclude the general part of the article with some negative results about T^+ . In particular, we show that in the specific case where the algebraic free completion $F(A)$ is already a model of T^+ , then the theory T^+ does not have a prime model, it is not model complete, and thus does not admit quantifier elimination. This is the case in all the incidence geometries considered in [21], but crucially it is false in the case of the pseudoplane, i.e., the theory T_1^+ from Section 4.1. We first define an abstract notion of predimension for the classes $(\mathcal{K}, \leq_{\circ})$.

Definition 3.64. Let $\delta : \mathcal{K} \rightarrow \mathbb{N}$ and let, for $A \subseteq B \models T_{\forall}$, $A \leq_{\delta} B$ if and only if $\delta(A \cap B_0) \leq \delta(B_0)$ for every finite $B_0 \subseteq B$. We say that δ is a *predimension function* for $(\mathcal{K}, \leq_{\circ})$ if it satisfies the following properties:

- (a) $A \cong B$ implies $\delta(A) = \delta(B)$;
- (b) $A \leq_{\circ} B$ implies $A \leq_{\delta} B$;
- (c) if $A \leq_{\circ} B$ is minimal, then $\delta(A) = \delta(B)$ if and only if B is algebraic over A ;

(d) if $A \leqslant_\delta B$ and $C \subseteq B$, then $A \cap C \leqslant_\delta C$.

In [21, Def. 11] Funk and Strambach give an explicit definition of a predimension function of all the geometries that they consider, with the exception of the generalised n -gons (for which a predimension function is defined in [1]) and (k, n) -Steiner systems, that we address in Section 4.2. We next identify the additional Assumptions 3.65 needed in this section. Theorem 3.70 establishes that, when these conditions hold, then T^+ does not have a prime model, it is not model complete, and does not have quantifier elimination. Our proof follows both [24] and [1], but in our general abstract setting.

Assumption 3.65. In addition to the assumptions from 3.28 and 3.43, in this section we also assume the following:

- (C7) if A contains at least $m_{\mathcal{K}}$ -many elements of each sort, then $F(A) \models T^+$;
- (C8) if $F(A) \models T^+$ and $B \in \mathcal{K}$, then $F(B)$ embeds into $F(A)$;
- (C9) there exists a predimension function $\delta : \mathcal{K} \rightarrow \mathbb{N}$ which satisfies the properties from Definition 3.64.

Remark 3.66. Notice that, by Proposition 3.44, what Assumption 3.65(C7) essentially says is that Assumption 3.28(C2) holds for exactly $C(A) = F(A)$. It follows in particular that the non-degenerate structures from 3.29 are essentially those structures whose free completion is a model of T^+ . Finally, we also point out that an immediate consequence of Assumption 3.65(C8) is that $F(A)$ is not Hopfian, i.e. there is some $B \subseteq F(A)$ such that $B \neq F(A)$ and $B \cong F(A)$.

Example 3.67. In the case of projective planes, the predimension function $\delta(A)$ is defined as follows. Let I_A be the number of incidences in A , i.e., the number of different unordered pairs $\{a, b\} \subseteq A$ such that $a \mid b$. Then we let $\delta(A) = |A| - 2 \cdot I_A$. The conditions from Assumption 3.64 can be verified by a routine argument. One can see that this is exactly the notion of predimension from [21] and also [35, §10.4]. This notion of predimension and its key properties were already studied by Hall in his seminal paper, see for instance [22, Th. 4.10]

Lemma 3.68. Let $A = F(A_0)$, for $A_0 \in \mathcal{K}$, then every elementary embedding of A into A is surjective.

Proof. Let $\alpha : A \rightarrow A$ be an elementary embedding and let $A = \bigcup_{i < \omega} A_i$ be a witness that $A = F(A_0)$. Obviously we have that $A_0 \leqslant_o A$. Let $A'_0 = \alpha(A_0)$. Now, as α is elementary and $A_0 \leqslant_o A$, we have that $A'_0 \leqslant_o A$ (by 3.7). Furthermore, there is $j < \omega$ such that for every $i \geqslant j$ we have that $A'_0 \subseteq A_i$. Now, fix one such $j < \omega$ and let $i \geqslant j$, then, as $A'_0 \leqslant_o A$, we have that $A'_0 \leqslant_o A_i$. In particular, we have that there is a HF-order $<$ of A_i over A'_0 . Let $a_0 < \dots < a_{m-1}$ be the elements in $A_i \setminus A'_0$ with respect to the order $<$. Suppose that one of the minimal extensions $A'_0 a_0, \dots, a_{\ell-t} \leqslant_o A'_0 a_1, \dots, a_\ell$ is not algebraic, and let $\ell \in [0, m-1]$ be minimal with respect to this property, then (by 3.65(C9)) $\delta(A'_0) = \delta(A'_0 a_1, \dots, a_{\ell-t})$ but $\delta(A'_0 a_1, \dots, a_\ell) > \delta(A'_0)$, on the other hand $\delta(A_i) = \delta(A_0) = \delta(A'_0)$, but this contradicts the fact that $<$ is an HF-order $<$ of A_i over A'_0 . It follows that for every $i \geqslant j$ we have that $A_i \subseteq \text{acl}_A(A'_0) \subseteq \alpha(F(A))$. ■

Lemma 3.69. Assume that $A_0 \in \mathcal{K}$ and $B \models T^+$. If B embeds elementarily in $F(A_0)$, then there is a finite $B_0 \subseteq B$ such that $B \cong F(B_0)$ and $\delta(B_0) \leqslant \delta(A_0)$.

Proof. Let $<$ be the natural HF-order of $F(A_0)$ witnessing that $A := F(A_0) = \bigcup_{i < \omega} A_i$ and such that $\delta(A_i) = \delta(A_0)$ for all $i < \omega$. In particular $<$ is a well-order. Suppose that B embeds elementarily in A and assume w.l.o.g. that this elementary embedding is simply the inclusion $B \subseteq A$. For each $i < \omega$ we let $B_i = A_i \cap B$. Since $B \preceq A$, it follows from 3.35 that $B \leq_o A$. From this we deduce that $B_i \leq_o A_i$ for all $i < \omega$ and so by 3.65(C9) we have $B_i \leq_\delta A_i$ for all $i < \omega$. Since $\delta(A_0)$ is finite and $\delta(A_0) = \delta(A_i)$ for all $i < \omega$, there is an index $k < \omega$ such that $\delta(B_k) \geq \delta(B_j)$ for all $j > k$. Moreover, since we have that $A_k \leq_o A_j$ (for $k < j$), we obtain also $B_k \leq_o B_j$. Therefore, we conclude by 3.64(d) that $\delta(B_k) = \delta(B_j)$ for all $j > k$. Again by 3.64(c), it follows that for all $j > k$ the elements in B_j are algebraic over B_{j-1} . It follows that $B = \text{acl}_B(B_k)$ and so, since $B_k \leq_o B$, we conclude by 3.51 that $F(B_k) \cong \text{acl}_B(B_k)$. Since $\delta(B_k) \leq \delta(A_k) = \delta(A_0)$, this completes our proof. ■

Theorem 3.70. *Suppose T^+ satisfies 3.28, 3.43 and also 3.65, then:*

- (1) T^+ does not have a prime model;
- (2) T^+ is not model complete;
- (3) T^+ does not eliminate quantifiers.

Proof. We first prove that T^+ does not have a prime model. Suppose towards contradiction that B is a prime model of T^+ . Then, by Lemma 3.69, it follows that $B = F(B_0)$ with $B_0 \in \mathcal{K}$ and such that $\delta(B_0)$ is minimal among the non-degenerate configurations in \mathcal{K} . It then follows from 3.65(C8) (and Remark 3.66) that we can choose copies $(A_i : i < \omega)$ of B such that $A_i \subsetneq A_{i+1}$ and let $A = \bigcup_{i < \omega} A_i$. Since the theory T^+ is a $\forall\exists$ -theory (cf. 3.22), it follows that $A \models T^+$. Now, since by assumption B is a prime model of T^+ , there is an elementary embedding $f : B \rightarrow A$. Let $C := f(B)$ and $C_0 = f(B_0)$, then since B_0 is finite there is an index $i < \omega$ such that $B_0 \subseteq A_i$. It then follows, since $A_i \models T^+$ and C_0 generates C , that $C \subseteq A_i$. Then, by 3.35 and the fact that $A_{i+1} \models T^+$, we have:

$$\begin{aligned} C \preceq A &\implies C \leq_o A \\ &\implies C \leq_o A_{i+1} \\ &\implies C \preceq A_{i+1}. \end{aligned}$$

But then $f : B \rightarrow A_{i+1}$ is non-surjective elementary embedding of B into (an isomorphic copy of) itself, which contradicts Lemma 3.68.

We next prove that T^+ is not model complete and does not eliminate quantifiers. Let $A \leq_o B \in \mathcal{K}$ be such that both A and B are non-degenerate, and suppose $\delta(A) < \delta(B)$ (these exists because trivial extensions are always strong and increase the value of δ). Then, by 3.65(C8) there is an embedding $h : F(B) \rightarrow F(A)$. Notice that, if $F(B) \cong F(D)$, then by 3.64(d) we must have that $\delta(B) = \delta(D)$. Therefore, by Lemma 3.69, it follows that h is not elementary, for otherwise we would have $\delta(B) \leq \delta(A)$. Thus h is a non-elementary embedding from $F(B)$ into $F(A)$. This shows that T^+ is not model complete and, as a direct consequence, it does not eliminate quantifiers. ■

4. APPLICATIONS

In the previous Section 3 we have introduced the class (\mathcal{K}, \leq_o) and studied the properties of the theory T^+ under the assumption from 3.28, 3.43, 3.60 and 3.65. In this section we turn to concrete examples of this setting and we apply this abstract

machinery to several distinct concrete cases of open incidence structures. We point out that such examples of incidence geometries can be divided into two families.

On the one hand, we study in Section 4.1 the n -open graphs introduced in [4], of which the free pseudoplane is a major example. We verify in these cases Conditions 3.2, 3.28 and 3.43, and we derive the completeness and stability of the associated theory T^+ in this context. We stress that, crucially, in this case the structure $F(A)$ trivialises and so the completion $C(A)$ from 3.28(C2) is not the free algebraic completion $F(A)$. Interestingly, for $n \geq 2$ the model-theoretic and the combinatorial notions of complexity diverge in this class of examples. While $F(A) \neq C(A)$ seems to indicate that the class in question is rather tame, it is still possible to exhibit configurations satisfying the properties from Assumption 3.60 and verify that T_n^+ is not stable, where T_n^+ is the theory determined as in 3.22 by the class of n -open graphs. All these results concerning n -open graphs were already shown in [4], but with different techniques.

On the other hand, in Sections 4.2–4.6 we study all the examples of open incidence geometries from [21]. From the model-theoretic point of view these cases behave quite similarly: they all verify the assumptions from 3.2, 3.28, 3.43, 3.60 and 3.65. In particular, in these cases we have that the free algebraic completion $F(A)$ is a model of T^+ , provided that A is non-degenerate. It thus follows from our results that the free completions of non-degenerate partial models in \mathcal{K} are all elementary equivalent. On the other hand, this family of examples also showcases several differences from the combinatorial point of view. Two issues are worth noting. Firstly, generalised n -gons make for the only case where minimal strong extensions are not necessarily one-element extensions – we shall expand later on this issue in Section 4.3. Secondly, affine planes and projective Möbius planes exhibit cases of incidence geometries with an additional relation of parallelism (or tangency) between elements of the same sort. As far as we know, the present work is the first model-theoretic study of incidence geometries which also involves such additional notions. However, here as in many other places, our treatment of the subject is largely indebted to the work of Funk and Strambach in [21], as they already identified the technical notion of Gaifman closure (cf. 2.8) that we use to deal with these difficulties. Finally, we point out again that the case of generalised n -gons was already considered in the literature (by [24] for the case $n = 3$ and by [1] for the case $n \geq 3$). The completeness result for T^+ in the case of (k, n) -Steiner systems for $k = 2$ and $n = 3$ was recently showed in the thesis [2]. Independently, this case was considered by Barbina and Casanovas in [9], where they prove both the completeness and the stability of the theory of open non-degenerate $(2, 3)$ -Steiner systems. All the other results from Sections 4.2–4.6 are novel.

4.1. n -open graphs

We consider the case of n -open graphs introduced by Baldwin, Freitag and Mutchnik in [4]. In this section we let $L = \{\downarrow\}$ be the language of graphs. For every $n < \omega$ we introduce a class of graphs (\mathcal{K}_n, \leq_n) and we show it satisfies the conditions from 3.2, 3.28 and 3.43. As a consequence, it follows from our results in Section 3 that the theory T_n^+ defined from (\mathcal{K}_n, \leq_n) as in 3.22 is complete, stable and with $\downarrow = \downarrow^\otimes$. Moreover, for $n \geq 2$, we also show that T_n^+ satisfies Assumption 3.60, and thus that T_n^+ is not superstable. We thus establish the following Theorem 4.1. We stress that all these facts are not novel and they are already known

from the literature, but we believe they provide an interesting application of our general setting. In particular, our proof of the non-superstability of T_n^+ for $n \geq 2$ is significantly different from the one from [4], and it highlights the combinatorial nature of the theories T_n^+ .

Theorem 4.1. *For every $n < \omega$, the theory T_n^+ is complete, stable, and with $\perp = \perp^\otimes$. Moreover, if $n \geq 2$, then T_n^+ is not superstable.*

We start by recalling the definition of n -open graphs from [4, 4.20]. If A is a graph then we denote by $\vartheta_A(a)$ the valency (cf. 2.3) of an element $a \in A$.

Definition 4.2. Let $A \subseteq B$ be graphs, then for every $n < \omega$ we define the following notions.

- (1) We say that B is n -open over A and write $A \leq_n B$ if every non-empty $C \subseteq B \setminus A$ contains an element $c \in C$ with $\vartheta_{AC}(c) \leq n$. We say that B is n -open if $\emptyset \leq_n B$, i.e., if every non-empty $C \subseteq B$ contains some element of valency $\leq n$.
- (2) We say that B is closed over A if $A \not\leq_n B$, i.e., if there is a non-empty $C \subseteq B \setminus A$ where every element $c \in C$ has valency $\vartheta_{AC}(c) \geq n+1$. We say that B is closed if $\emptyset \not\leq_n B$, i.e., if there is a non-empty configuration $C \subseteq B$ where every element $c \in C$ has valency $\vartheta_C(c) \geq n+1$.
- (3) For every $n < \omega$ we let \mathcal{K}_n be the class of all finite n -open graphs, and we let $T_{\forall}^n = \text{Th}_{\forall}(\mathcal{K}_n)$.

The definitions above extend to infinite models of T_{\forall}^n as in 3.28.

Remark 4.3. Each class (\mathcal{K}_n, \leq_n) gives rise to a theory T_n^+ defined as in 3.22. It is straightforward to verify that our theories T_n^+ coincide with the theories T_n defined in [4, 4.21]. In particular, T_0^+ is the theory of an infinite set, while T_1^+ is the theory of the free pseudoplane (cf. also [34]).

Conditions 3.2(A)–(F) follow immediately from the definition of the relation \leq_n . Condition 3.2(I) follows by letting $X_{\mathcal{K}_n} = \{1\}$ and Condition 3.2(J) by letting $n_{\mathcal{K}_n} = n$. Also Condition 3.2(K) and 3.2(L) follow immediately from Definition 4.2. It remains to verify 3.2(G) and 3.2(H).

Proposition 4.4 (3.2(G)). The class (\mathcal{K}_n, \leq_n) has both the amalgamation and joint embedding property.

Proof. Since $\emptyset \in \mathcal{K}_n$, it suffices to verify the amalgamation property. Consider $A, B, C \in \mathcal{K}$ with $A \leq_n B, A \leq_n C$ and $B \cap C = A$. By the definition of the free amalgam $D := B \otimes_A C$ (recall 3.39), we have that $\vartheta_D(b) = \vartheta_B(b)$ for every $b \in B$ and $\vartheta_D(c) = \vartheta_C(c)$ for every $c \in C$. It follows that $B \otimes_A C \in \mathcal{K}_n$ and, moreover, $B \leq_n B \otimes_A C, C \leq_n B \otimes_A C$. This completes our proof. ■

Recall that $\text{gcl}_M(A)$ is the Gaifman closure from 2.8. Then in the setting of graphs we have $\text{gcl}_M(A) = \text{gcl}_M^1(A)$, as there is no parallelism relation in L . We next verify Condition 3.2(H).

Proposition 4.5 (3.2(H)). If $A \subseteq B \in \mathcal{K}$ and $A \not\leq_n B$, then every $C \in \mathcal{K}_n$ contains at most n -many copies of B over A .

Proof. Suppose $A \not\leq_n B$, then by Definition 4.59 there is a subset $B' \subseteq B \setminus A$ such that every element in B' has valency at least $n+1$. Without loss of generality we let $B' = B \setminus A$. Suppose $C \in \mathcal{K}$ contains at least $n+1$ copies $(B_i)_{0 \leq i \leq n}$ of B isomorphic

over A , then we consider the set $D := \bigcup_{0 \leq i \leq n} \text{gcl}_C(B_i \setminus A)$. Now, if $b \in B_i \setminus A$ then by assumption it is incident to at least $n + 1$ -many elements from $\text{gcl}_C(B_i \setminus A)$. Moreover, by construction and the fact that $\text{gcl}_C(B_i \setminus A) = \text{gcl}_C^1(B_i \setminus A)$, we also have that every element $a \in \text{gcl}_C(B_i \setminus A) \cap A$ is incident to at least $n + 1$ -many elements from D , as it is incident to at least one element for each B_i . It follows that every element $d \in D$ has valency $\vartheta_D(d) \geq n + 1$, contradicting $C \in \mathcal{K}_n$. \blacksquare

It follows from Proposition 4.4, Proposition 4.5, and the previous observations, that (\mathcal{K}_n, \leq_n) satisfies the conditions from 3.2. We next prove that it also verifies the assumptions from 3.28. By Lemma 3.30, to show 3.28(C1) it is sufficient to verify Assumptions 3.30(D1) and 3.30(D2).

Proposition 4.6 (3.30(D1)). *If $M \models T_n^+$, $A \subseteq M$ is finite, $<$ is an HF-order of M and $A \leq_n Ac \in \mathcal{K}_n$ is a trivial extension, then for all $\ell < \omega$ there exists $c'_\ell \in M$ s.t.:*

- (a) $Ac \cong_A Ac'$;
- (b) $Ac' \leq_n \text{Acl}_<^\ell(c')$.

Proof. By induction on $\ell < \omega$ we prove that for every $A \leq_n M$ and $A \leq_n Ac \in \mathcal{K}$ we can find c'_ℓ as in (a)-(b). The case for $n = 0$ is easy: it is enough to pick an element $c'_0 > A$ which has no incidence with elements from A . Clearly, we can find such an element in any model $M \models T_n^+$ with an HF-order $<$.

Thus suppose the inductive hypothesis holds with respect to ℓ , we define $c'_{\ell+1}$ so that (a)-(b) are satisfied for it. By assumption, and proceeding recursively, we can find n -many elements d_1, \dots, d_n such that, for each $0 \leq i < n$, we have that the element d_{i+1} has no incidence with elements from $\text{Acl}_<^\ell(\{d_1, \dots, d_i\})$ and

$$\text{Acl}_<^\ell(\{d_1, \dots, d_i\})d_{i+1} \leq_n \text{Acl}_<^\ell(\{d_1, \dots, d_i, d_{i+1}\}).$$

Now, notice that the extension $d_1 \dots d_n \leq_n d_1 \dots d_n b$ with b incident to d_1, \dots, d_n is not algebraic. Since $M \models T_n^+$ there are infinitely many elements in M isomorphic to b over d_1, \dots, d_n . In particular, since $<$ is a HF-order, there is some $c'_{\ell+1} \in M$ such that $d_i < c'_{\ell+1}$ for all $1 \leq i \leq n$. It follows that $c'_{\ell+1}$ is incident only to d_1, \dots, d_n from $(c'_{\ell+1})^\downarrow$. Thus clearly we have that $Ac \cong_A Ac'_{\ell+1}$.

We next claim that $Ac'_{\ell+1} \leq_n \text{Acl}_<^{\ell+1}(c'_{\ell+1})$. To see this, notice that we can define an HF-order $<'$ by first letting $Ac'_{\ell+1} <' d_1 <' \dots <' d_n$. By construction, $c'_{\ell+1}$ is incident only to d_1, \dots, d_n in $\text{cl}_<^{\ell+1}(c'_{\ell+1})$, and, for $1 \leq i < j \leq n$, no element in $\text{cl}_<^\ell(d_i)$ is incident to elements from $\text{cl}_<^\ell(d_j)$. It follows that we can extend the order $<'$ and obtain $Ac'_{\ell+1} \leq_{\text{HF}} Ac'_{\ell+1} \text{cl}_<^\ell(\{d_1, \dots, d_i, d_{i+1}\})$. Since by construction we have that $\text{cl}_<^{\ell+1}(c'_{\ell+1}) = \text{cl}_<^\ell(\{d_1, \dots, d_n\}) \cup \{c'_{\ell+1}\}$ we obtain that $Ac'_{\ell+1} \leq_n \text{Acl}_<^{\ell+1}(c'_{\ell+1})$. This concludes our proof. \blacksquare

Lemma 4.7. *Let $A \leq_n C \models T_\forall^n$ and $c \in C$ be such that $\vartheta_{Ac}(c) = n$, then $Ac \leq_n C$.*

Proof. Consider any finite subset $D \subseteq C$ such that $Ac \subsetneq D$, we need to show that $D \setminus Ac$ contains some element with valency $\leq n$. Now, since $A \leq_n C$ and $D \subseteq C$, there is an element $d \in D \setminus A$ with $\vartheta_{AD}(d) \leq n$. If $c \neq d$ then we have $d \in D \setminus Ac$ and $\vartheta_{AD}(d) \leq n$, and we are done. Thus suppose that c is the unique element in $D \setminus A$ with valency $\leq n$. Since $\vartheta_{Ac}(c) = n$, it follows that $\vartheta_{AD}(c) = n$, and in particular it follows that c is not incident to any element from $D \setminus A$. Then let $D' = D \setminus \{c\}$ and consider an element $d' \in D' \setminus A$ with $\vartheta_{AD'}(d') \leq n - 1$ - this exists since $A \leq_n C$ and $Ac \subsetneq D$. Then c and d' are not incident, whence $\vartheta_{AD}(d') \leq n$.

This shows that every finite subset $D \subseteq C$ such that $Ac \subsetneq D$ contains an element with valency $\leq n$, and thus proves that $Ac \leq_n C$. \blacksquare

Proposition 4.8 (3.30(D2)). Suppose $M \models T_n^+$ is \aleph_1 -saturated and $A \leq_n M$ is countable. If for every trivial one-element extension $A \leq_n Ab$ there is $b' \in M$ such that $Ab \cong_A Ab' \leq_n M$, then for every minimal extension $A \leq_n Ac \models T_\forall$ there is an element $c' \in M$ such that $Ac \cong_A Ac' \leq_n M$.

Proof. Let $M \models T_n^+$ be \aleph_1 -saturated and $A \leq_n M$ be countable. Suppose $A \leq_n Ac \in \mathcal{K}_n$ is a minimal extension with c incident to elements a_1, \dots, a_ℓ from A , for $1 \leq \ell \leq n$. By assumption, we can find elements $b_{\ell+1}, \dots, b_n$ in $M \setminus A$ such that each $b_{\ell+k+1}$ is trivial over $Ab_{\ell+1} \dots b_{\ell+k}$ and $Ab_{\ell+1} \dots b_n \leq_n M$. Then, since $M \models T_n^+$, it follows that there are infinitely many elements which are incident to $a_1, \dots, a_\ell, b_{\ell+1}, \dots, b_n$. In particular, it follows that the type $p(x, Ab_{\ell+1} \dots b_n)$ which says that x is incident to $a_1, \dots, a_\ell, b_{\ell+1}, \dots, b_n$ and $x \neq a$ for all $a \in A$ is consistent. It follows from the fact that M is \aleph_1 -saturated that there is some $c' \in M \setminus A$ incident to $a_1, \dots, a_\ell, b_{\ell+1}, \dots, b_n$. Then, since $Ab_{\ell+1} \dots b_n \leq_o M$, we obtain that c' is incident only to these elements from $Ab_{\ell+1} \dots b_n$. It follows in particular that $Ac \cong_A Ac'$. Moreover, since $\vartheta_{Ab_{\ell+1}, \dots, b_n}(c') = n$, we obtain from Lemma 4.7 that $Ab_{\ell+1} \dots b_n c' \leq_n M$. Since $b_{\ell+1}, \dots, b_n$ are trivial extensions of A , we conclude from 3.2(K) that $Ac' \leq_n Ab_{\ell+1} \dots b_n c' \leq_n M$. \blacksquare

Finally, to verify Assumption 3.28(C2), we introduce a version of the free completion process for n -open graphs, which was already considered in [4, p. 26]. As noticed by the authors, this free construction can be considered as an adaptation of Hall's construction of a free projective plane in the context of n -open graphs. However, we stress that, while in Hall's case the free completion is algebraic over the starting configuration, this is not the case in the setting of n -open graphs. As we pointed out already in the previous section, we believe that this indicates an important difference between the case of n -open graphs and the geometries considered in the later sections. We reflect this in our notation and we use $C(A)$ for the abstract notion of completion needed in 3.28(C2), while we reserve $F(A)$ for the notion of algebraic completion defined in 3.41.

Definition 4.9. Let $A \models T_\forall^n$. We let $A_0 := A$ and, for every $i < \omega$, we let A_{i+1} be obtained from A_i by adding, for every distinct $c_1, \dots, c_n \in A_i$, infinitely many elements which are incident to c_1, \dots, c_n and to no other element from A_i . We then define $C(A) := \bigcup_{i < \omega} A_i$ and we say that $C(A)$ is the *completion* of A .

Proposition 4.10 (3.28(C2)). Let $A \models T_\forall^n$ and suppose that A contains at least n -many elements. Then $C(A) \models T_n^+$, $A \leq_n C(A)$, $|C(A)| = |A| + \aleph_0$ and:

- (a) $C(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : C(A) \cong C(B)$;
- (b) $C(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_n M$ there is an embedding $\hat{h} : C(A) \rightarrow M$ such that $\hat{h}|_A = h$ and $\hat{h}(C(A)) \leq_n M$.

Proof. Now, since it is immediate to verify that the construction of $C(A)$ induces a HF-order of $C(A)$ over A , it follows that $A \leq_n C(A)$. Similarly, it follows immediately by the construction that $|C(A)| = |A| + \aleph_0$. The fact that any isomorphism

$h : A \cong B$ extends to an isomorphism $\hat{h} : C(A) \cong C(B)$ also follows immediately by Definition 4.9. It thus remains to verify that $C(A) \models T_n^+$ and that $C(A)$ is prime.

We verify that, if $A \models T_{\forall}^n$ contains at least n -many elements, then $C(A) \models T_n^+$. Notice that, if $|A| \geq n$ then $C(A)$ is infinite. We verify that $C(A)$ satisfies the axioms from 3.22. Since in \mathcal{K}_n there are no algebraic extensions, it suffices to verify 3.22(1) and 3.22(3). Let $D \subseteq_{\omega} C(A)$, then $D \subseteq A_i$ for some $i < \omega$ and $D \not\subseteq A_j$ for all $j < i$. We assume w.l.o.g. $i > 0$, then any $d \in D \setminus A_{i-1}$ must satisfy $\vartheta_D(d) \leq n$. This shows that every finite substructure of $C(A)$ is a model of T_{\forall}^n , and thus $C(A) \models T_{\forall}^n$. Next, consider any $D \subseteq_{\omega} C(A)$ and some minimal extension $D \leq_n Dc \in \mathcal{K}$. Now, c is incident exactly to elements $d_1, \dots, d_{\ell} \in D$ for some $\ell \leq n$. Pick some new elements $d_{\ell+1}, \dots, d_n \in C(A) \setminus D$ and let A_i be the stage with the smallest index satisfying $\{d_1, \dots, d_{\ell}, d_{\ell+1}, \dots, d_n\} \subseteq A_i$. By construction A_{i+1} contains infinitely-many elements $(c_i)_{i < \omega}$ incident only to $d_1, \dots, d_{\ell}, d_{\ell+1}, \dots, d_n$ in $D \cup \{d_{\ell+1}, \dots, d_n\}$. It follows that for every $i < \omega$ we have $Dc \cong_D Dc_i \subseteq C(A)$. This shows that $C(A) \models T_n^+$.

Now, suppose that $M \models T^+$ and that $h : A \rightarrow M$ is an embedding with $h(A) \leq_n M$. We suppose inductively that there is $h_i \supseteq h$ with $h_i : A_i \rightarrow M$ and $h_i(A_i) \leq_n M$. Let $c \in A_{i+1} \setminus A_i$, then c is incident to exactly n -many elements $d_1, \dots, d_n \in A_i$. Since $M \models T_n^+$ there is an element $c' \in M \setminus h_i(A_i)$ which is incident to $h_i(d_1), \dots, h_i(d_n)$. Since $h_i(A_i) \leq_n M$ it follows that c' is not incident to any other element from $h_i(A_i)$. Let $h_{i+1} := h_i \cup \{(c, c') : c \in A_{i+1} \setminus A_i\}$ where the element c' is chosen as above. It follows immediately from the choice of such elements that $h_{i+1} : A_{i+1} \rightarrow M$ is an embedding. Moreover, since every element in $h_{i+1}(A_{i+1}) \setminus h_{i+1}(A_i)$ has valency n in $h_{i+1}(A_{i+1})$, it follows from Lemma 4.7 that $h_{i+1}(A_{i+1}) \leq_n M$. Finally, letting $\hat{h} = \bigcup_{i < \omega} h_i$, we obtain an embedding $\hat{h} : C(A) \rightarrow M$ satisfying $\hat{h} \upharpoonright A = h$ and $\hat{h}(C(A)) \leq_n M$. \blacksquare

Now, the fact that (\mathcal{K}_n, \leq_n) also verifies the assumptions from 3.43 is an immediate consequence of Proposition 4.5 and the fact that every strong extension $A \leq_n B$ is not algebraic. This latter observation together with the fact that $F(A) \neq C(A)$, means that the conditions from 3.65 do not apply to this class of examples. It remains to consider Assumption 3.60. Since T_0^+ and T_1^+ are respectively the theory of the infinite set and of the pseudoplane, they are both ω -stable and thus do not satisfy 3.60. However, we show that if $n \geq 2$ then T_n^+ satisfies 3.60, thereby also completing the proof that T_n^+ (with $n \geq 2$) is not superstable.

Proposition 4.11. Let $n \geq 2$, then there is a configuration $C \in \mathcal{K}_n$ satisfying the conditions from 3.60.

Proof. Fix some $n \geq 2$, then we construct a configuration $C \in \mathcal{K}_n$ which satisfies the conditions from 3.60. Consider the configuration $C = \{c_0, c_1\} \cup \{c_i : 2 \leq i \leq n+2\}$ defined as follows: (1) for all $2 \leq i \leq n$, c_i is incident to c_j for all $j < i$; (2) c_{n+1} is incident to c_i for $i < n$; (3) c_{n+2} is incident to c_0, c_1, c_n and c_{n+1} .

Clearly C is a graph. Moreover, it is straightforward to verify that the order obtained by letting $c_i <' c_j$ if $i < j$ is an HF-order, thus we have that $C \in \mathcal{K}_n$. We verify conditions (a)-(c) from 3.60.

Firstly, we clearly have from the fact that $<'$ is a HF-order that $c_0, c_1 \leq_n C$. By definition we have that there is no edge between c_0 and c_1 . Thus 3.60(a) holds.

Now, notice that the elements c_0, c_1 are incident to $c_2, c_3, \dots, c_{n+1}, c_{n+2}$, and for every $2 \leq i \leq n+1$, the element c_i is incident to all elements c_j with $0 \leq j \leq n+1$ and $j \neq i$. Hence every element in $C \setminus \{c_{n+2}\}$ has valency $n+1$ in $C \setminus \{c_{n+2}\}$, and so $c_{n+2} \not\leq_n C$. Also L has only one sort, thus all elements in C have the same sort. This shows that 3.60(b) also holds.

Finally, one can continue this construction by taking $c_{n+2+1}, \dots, c_{2n+4}$ such that the configuration $C' = \{c_{n+2}, c_{n+2+1}, \dots, c_{2n+4}\}$ is isomorphic to C . Then the resulting configuration $C \cup C'$ is still a model in \mathcal{K}_n . Since this construction can be continued to obtain a sequence $(c_i)_{i < \omega}$, it follows that also 3.60(c) is verified. ■

4.2. (k, n) -Steiner systems

In this section we start to study the incidence geometries considered in [21], namely those geometries where the free completion of a (non-degenerate) model in \mathcal{K} satisfies T^+ . We first consider (k, n) -Steiner systems, which are an example of incidence geometry consisting of points and blocks. We notice that, recently, there has been a significant interest in the model theory of Steiner systems, as witnessed by the references [5, 6, 7, 8, 23]. Given this interest, and also because this is the first example of incidence geometry from [21] that we consider, we provide details for most proofs, while in subsequent section we will omit the details of those proofs that follow exactly as in the case of Steiner systems. We thus proceed as follows. Firstly, we introduce in 4.2.1 the class \mathcal{K} of finite, open, (k, n) -Steiner systems and we define a relation $A \leq_o B$ in this context. Then, in 4.2.2 we verify all the conditions from 3.2, showing that (\mathcal{K}, \leq_o) satisfies all the properties of the usual Hrushovski constructions (plus the additional 3.2(H)–3.2(L)). Incidentally, these results will also verify the conditions from 3.43, as we shall remark later. In 4.2.3, we verify the assumptions from 3.28, and in particular we show that the free completion $F(A)$ is a model of T^+ , provided that A is a non-degenerate structure in \mathcal{K} . Finally, in 4.2.4 we show that 3.60 and 3.65 are also verified. The following theorem then follows from the results in Section 3.

Theorem 4.12. *Let (\mathcal{K}, \leq_o) be the class of finite, open, partial (k, n) -Steiner systems and let T^+ be defined as in 3.22. Then T^+ is complete, stable, and with $\downarrow = \downarrow^\otimes$. Moreover, T^+ is not superstable, it does not have a prime model, it is not model complete and does not eliminate quantifiers.*

4.2.1. Definition of (\mathcal{K}, \leq_o)

We recall in this section the definition of (k, n) -Steiner systems from [21, p. 748] and we introduce the class \mathcal{K} of partial, finite, open, (k, n) -Steiner systems.

Definition 4.13. Let L be the two-sorted language where the symbols p_0, p_1, \dots denote *points*, the symbols b_0, b_1, \dots denote *blocks* and the symbol $p | b$ means that the point p is incident with the block b . For any $2 \leq k < n$ the theory of (k, n) -Steiner systems consists of the following axioms:

- (1) given k -many distinct points p_0, \dots, p_{k-1} there is a unique block b such that $p_i | b$ for all $i < k$;
- (2) for every block b there are exactly n -many points p_0, \dots, p_{n-1} such that $p_i | b$ for all $i < n$.

The universal fragment of the theory of (k, n) -Steiner systems is axiomatised exactly by the formula saying that, whenever both $\bigwedge_{i < k} p_i | b$ and $\bigwedge_{i < k} p_i | b'$ hold, then $b = b'$. We refer to models of this universal theory as *partial (k, n) -Steiner systems*.

Definition 4.14. For finite partial (k, n) -Steiner systems $A \subseteq B$, we define the following.

- (1) An element $a \in A$ is *hyperfree* in B if it satisfies one of the following conditions:
 - (i) a is a point incident with at most one block from B ;
 - (ii) a is a block incident with at most k -many points from B .
- (2) We say that B is *open over* A and write $A \leq_o B$ if every non-empty $C \subseteq B \setminus A$ contains an element which is hyperfree in AC . We say that B is *open* if $\emptyset \leq_o B$, i.e., if every non-empty $C \subseteq B$ contains some element hyperfree in C .
- (3) We say that B is *closed over* A if $A \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B \setminus A$ which does not contain any element hyperfree in AC . We say that B is *closed* if $\emptyset \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B$ which does not contain any element hyperfree in C .

We let \mathcal{K} be the class of finite, open, partial (k, n) -Steiner systems and we let T_{\forall} be the universal theory of \mathcal{K} . The previous definitions immediately extend to infinite models of T_{\forall} as in 3.6. The notion of hyperfree element in a partial (k, n) -Steiner system is exactly as in [21, p. 753].

We remark what exactly are the algebraic strong extensions in the context of (k, n) -Steiner systems and we show that Condition 3.43(C5) holds. Notice that, although this conditions belongs to the set of assumptions that we introduced in order to characterise the relation of forking independence, in the case of Steiner system it is convenient to prove it first, as it is useful to prove the other conditions.

Remark 4.15. In the case of (k, n) -Steiner systems, a minimal strong extensions $A \leq_o Ac$ is algebraic if c is a point incident with some block from A , or if c is a block incident with exactly k -many points from A .

Lemma 4.16 (3.43(C5)). Let $A \leq_o C \models T_{\forall}$ and let $c \in C$ be such that $A \leq_o Ac$ is an algebraic strong extension. Then $Ac \leq_o C$.

Proof. Consider any finite subset $D \subseteq C$ such that $Ac \subsetneq D$, we need to show that $D \setminus Ac$ contains some element hyperfree in D . Now, since $A \leq_o C$ and $D \subseteq C$, there is an element $d \in D \setminus A$ hyperfree in D . If $c \neq d$ then d also witnesses that $D \setminus Ac$ contains an element hyperfree in D , and we are done. Thus suppose that c is the unique element in $D \setminus A$ hyperfree in D .

Notice that, since the strong extension $A \leq_o Ac$ is algebraic, c is either a point incident to one block in A or a block incident to k -many points in A . In both cases, c is not incident to any element from $D \setminus A$. Then let $D' = D \setminus \{c\}$ and consider an element $d' \in D' \setminus A$ hyperfree in D' – this exists since $A \leq_o C$ and $Ac \subsetneq D$. Since c is incident only to elements from A , it follows that c and d' are not incident, and therefore d' is hyperfree also in D . This shows that every finite subset $D \subseteq C$ such that $Ac \subsetneq D$ contains an element hyperfree in D and thus proves that $Ac \leq_o C$. ■

4.2.2. Conditions 3.2

We now start considering the conditions from 3.2. Conditions (A), (B), (D), (E), (F), (K) and (L) follow immediately from the definition of the relation \leq_o for finite, open, partial (k, n) -Steiner systems. By Definition 4.14 it is clear that Condition 3.2(I) holds with $X_{\mathcal{K}} = \{1\}$, i.e., the minimal extensions are exactly the one-element extensions. It remains to verify conditions (C), (G), (H) and 3.2(J), which we consider separately in the following propositions.

Proposition 4.17 (3.2(C)). If $A, B, C \in \mathcal{K}$ and $A \leq_o B \leq_o C$, then $A \leq_o C$.

Proof. We have to show that for every non-empty $D \subseteq C \setminus A$ there is $d \in D$ which is hyperfree in AD .

Case 1. $D \cap (C \setminus B) \neq \emptyset$.

Let $D' := D \cap (C \setminus B)$. As $\emptyset \neq D' \subseteq C \setminus B$ and $B \leq_o C$ there is $d \in D'$ which is hyperfree in BD' , but then $d \in D \setminus A$ and d is hyperfree in AD' (as it is hyperfree in BD' and $AD' \subseteq BD'$).

Case 2. $D \subseteq B \setminus A$.

This case is clear as $A \leq_o B$. ■

The following proposition shows that the class (\mathcal{K}, \leq_o) verifies 3.2(G). Together with the next remark this also verifies Assumption 3.43(C4).

Remark 4.18 (3.43(C4)). Let $A \subseteq B, C$ be open partial (k, n) -Steiner systems with $B \cap C = A$ and recall the definition of free amalgam from 3.39. We stress that the free amalgam $B \otimes_A C$ is always an L -structure, but it is not necessarily an open partial (k, n) -Steiner system. However, it follows from the proof of Proposition 4.19 that when the conditions from 3.43(C4) are satisfied then $B \otimes_A C \in \mathcal{K}$.

The following proof adapts [1, Prop. 2.21] and [24, Lem. 8.12] to the context of partial (k, n) -Steiner systems.

Proposition 4.19 (3.2(G)). The class (\mathcal{K}, \leq_o) has both the amalgamation and joint embedding property.

Proof. Firstly, notice that the empty model is obviously prime in \mathcal{K} . Since the existence of a prime model together with the fact that \mathcal{K} has the amalgamation property entail that \mathcal{K} also has the joint embedding property, it suffices to check that amalgamation holds in \mathcal{K} .

Consider $A, B, C \in \mathcal{K}$ with $A \leq_o B, A \leq_o C$ and $B \cap C = A$. Notice that, if $B \otimes_A C \in \mathcal{K}$, then it follows immediately from the definitions of free amalgam 3.39 and the definition of hyperfree elements in (k, n) -Steiner systems that $B \leq_o B \otimes_A C$ and $C \leq_o B \otimes_A C$. Thus, it suffices to verify that either there is an embedding $f : B \rightarrow C$ such that $f(B) \leq_o C$ or $B \otimes_A C \in \mathcal{K}$, i.e., $B \otimes_A C$ is a well-defined partial, open, (k, n) -Steiner system. Reasoning by induction, we can assume that $A \leq_o B$ is a minimal strong extension, i.e., that $B \setminus A$ consists of a single element. We distinguish the following cases.

Case 1. $B = A \cup \{b\}$ where b is a block.

Since $A \leq_o Ab$ then b is incident with at most k -many points from A .

Case 1.1. b is incident with exactly k -many points p_0, \dots, p_{k-1} from A and C contains a block b' incident with p_0, \dots, p_{k-1} .

Since $A \leq_o C$ there is no other point in A incident to b' and so the map $f : B \rightarrow C$ such that $f|_A = \text{id}_A$ and $f(b) = b'$ is an embedding. Moreover, since $f(B) = Ab'$ and b' is algebraic over A , it follows by 4.16 and $A \leq_o C$ that $f(B) \leq_o C$.

Case 1.2. b is incident with exactly k points p_0, \dots, p_{k-1} and C does not contain a block b' incident with p_0, \dots, p_{k-1} .

In this case the free amalgam $B \otimes_A C$ is a partial (k, n) -Steiner system and so $B \otimes_A C \in \mathcal{K}$.

Case 1.3. b is incident with strictly less than k points from A .

As in the previous subcase, it follows immediately that $B \otimes_A C \in \mathcal{K}$.

Case 2. $B = A \cup \{p\}$ where p is a point.

Since $A \leq_o Ap$ then p is incident with at most a single block in A .

Case 2.1. p is not incident with any block.

It follows immediately that $B \otimes_A C \in \mathcal{K}$.

Case 2.2. p is incident with a block $b \in A$ and $C \setminus A$ contains less than n points incident to b .

It follows immediately that $B \otimes_A C \in \mathcal{K}$.

Case 2.3. p is incident to b and $C \setminus A$ contains n points p'_0, \dots, p'_{n-1} incident to b .

Since $A \leq_o A \cup \{p'_0, \dots, p'_{n-1}\}$, then each point p'_i is incident only with b . Then, for any $i < n$, the map $f : B \rightarrow C$ defined by letting $f|_A = \text{id}_A$ and $f(p) = p'_i$ is an embedding. Moreover, since $A \leq_o Ap'_i$ is algebraic, it follows from 4.16 that $f(B) \leq_o C$. ■

The next proposition shows that closed extensions of finite, open partial partial (k, n) -Steiner systems are algebraic, i.e., we verify 3.2(H). This condition is specific of our framework and it does not always hold in Hrushovski constructions (cf. [3]).

Proposition 4.20 (3.2(H)). If $A \subseteq B \in \mathcal{K}$ and $A \not\leq_o B$, then there is $m < \omega$ such that every $C \in \mathcal{K}$ contains at most m copies of B over A .

Proof. Suppose $A \not\leq_o B$, then by Definition 4.14 there is a subset $B' \subseteq B \setminus A$ such that every element in B' is not hyperfree in AB' , meaning that every point $p \in B'$ is incident with at least two blocks from AB' and every block $b \in B'$ is incident with at least $k + 1$ points from AB' . Without loss of generality we assume that $B' = B \setminus A$. We distinguish the following cases.

Case 1. $B \setminus A$ contains a point p_0 which is incident with some block $b \in A$.

Then, if $C \in \mathcal{K}$ contains at least $n + 1$ copies of B over A (where n is as in the definition of (k, n) -Steiner systems), in particular it contains $n + 1$ points p_0, \dots, p_n which are all incident with $b \in A$. But this is impossible since b can be incident with at most n points (as C is a partial (k, n) -Steiner system).

Case 2. $B \setminus A$ contains no point incident to blocks in A .

Then, if also no block in $B \setminus A$ is incident to points in A , it follows that every point in $B \setminus A$ is incident with at least two blocks from $B \setminus A$ and every block in $B \setminus A$ is incident with at least $k + 1$ points from $B \setminus A$. But then $B \setminus A$ contains no element hyperfree in $B \setminus A$, contradicting $B \in \mathcal{K}$. Thus $B \setminus A$ contains a block b incident to some point $p \in A$.

Then, let p_0, \dots, p_{m-1} be all the points in A which are incident to some block in $B \setminus A$. We let $B_0 := B$ and let B_1 be a disjoint copy of B_0 over A . We claim that $D = (B_0 \setminus A) \cup (B_1 \setminus A) \cup \{p_0, \dots, p_{m-1}\}$ does not contain any element hyperfree in D . By Definition 4.14 this shows that any C containing at least two copies of B over A is not open, proving our claim. We need to consider the following elements.

Case (a). $p \in B_i \setminus A$, for $i \in \{0, 1\}$.

Since $A \not\leq_o B_i$ we have that p is incident to at least two blocks from B_i . By assumption we have that B_i does not contain points incident to blocks in A , thus p is incident to at least two blocks from $(B_i \setminus A) \subseteq D$.

Case (b). $p = p_i$, for $i < m$.

By choice of the elements p_0, \dots, p_{m-1} we have that $p_i | b_0^i$ for some $b_0^i \in B_0 \setminus A$. Since B_1 is isomorphic to B_0 over A , we can also find some block $b_1^i \in B_1 \setminus A$ such that $p_i | b_1^i$. So p_i is incident to at least two blocks from D .

Case (c). $b \in B_i$, for $i \in \{0, 1\}$.

Since $A \not\leq_o B_i$ we have that every block $b \in B_i$ is incident with at least $k+1$ points from B_i . By the choice of the points p_0, \dots, p_{m-1} , it follows it is incident to at least $k+1$ points from $(B_i \setminus A) \cup \{p_0, \dots, p_{m-1}\} \subseteq D$. This concludes the proof. ■

Finally, the following condition is straightforward to verify, but we provide an argument to show the role of the Gaifman closure, as defined in 2.8.

Proposition 4.21 (3.2(J)). There is $n_{\mathcal{K}} < \omega$ such that, if $A \leq_o Ac \in \mathcal{K}$ is a one-element extension, then $|\text{gcl}_{Ac}(c) \setminus \{c\}| \leq n_{\mathcal{K}}$.

Proof. Let $A \leq_o Ac$, then by Definition 4.14 we have that c is either a point incident to one block from A , or it is a block incident to k -many points from A . In other words, we have from the definition of Gaifman closure from 2.8 that $\text{gcl}_{Ac}(c) \setminus \{c\} = \{b\}$, for $b \in A$ a block, or $\text{gcl}_{Ac}(c) \setminus \{c\} = \{p_1, \dots, p_k\}$, for p_1, \dots, p_k points. Letting $n_{\mathcal{K}} = k$, the claim is verified immediately. ■

4.2.3. Assumptions 3.28

We showed in the previous section that (\mathcal{K}, \leq_o) fits the specific subclass of Hrushovski constructions that we introduced in 3.2. It follows that (\mathcal{K}, \leq_o) determines a theory T^+ , as defined in 3.22. We verify in this section that T^+ satisfies the conditions from 3.28. By Theorem 3.34 this shows that the theory T^+ of open (k, n) -Steiner systems is complete and stable.

We start by proving that Conditions 3.30(D1) and 3.30(D2) hold. Then, by Lemma 3.30, it follows that T^+ also satisfies Assumption 3.28(C1). Notice that we state the condition slightly differently than in 3.30(D1), as we add a third condition to simplify our inductive argument. It is immediate to see that the following proposition entails 3.30(D1). Also, recall that by 3.2(L) in the context of (k, n) -Steiner systems trivial extensions are always strong.

Proposition 4.22 (3.30(D1)). If $M \models T^+$, $A \subseteq M$ is finite, $<$ is an HF-order of M and $A \leq_o Ac$ is a trivial extension, then for all $\ell < \omega$ there is $c' = c'_\ell \in M$ s.t.:

- (a) $Ac \cong_A Ac'$;
- (b) $Ac' \leq_o \text{Acl}_<^\ell(c')$;
- (c) $A < \text{cl}_<^{\ell+1}(c')$ (recall 3.8).

Proof. By induction on $\ell < \omega$ we prove that for every finite $A \subseteq M$ and $A \leq_o Ac \in \mathcal{K}$ we can find c'_ℓ as in (a)-(c). If $\ell = 0$, then it is easy to see that we can find c'_0 as wanted. So suppose the inductive hypothesis and let c be such that $A \leq_o Ac \in \mathcal{K}$ is a trivial extension, we want to define $c'_{\ell+1}$ so that (a)-(c) are satisfied for it.

Case 1. c is a block.

By inductive hypothesis we can find a point d_ℓ^0 such that (a)-(c) is true for A . Additionally, by applying the induction hypothesis and since $<$ is an HF-order, we can find inductively, for every $0 < i < k$, a block b_ℓ^i and a point d_ℓ^i such that $b_\ell^i < d_\ell^i$, $b_\ell^i \mid d_\ell^i$, b_ℓ^i is a trivial extension of $A \cup \bigcup \{\text{cl}_<^\ell(d_\ell^j) : 0 \leq j < i\}$ and (a)-(c) is true with respect to $A \cup \bigcup \{\text{cl}_<^\ell(d_\ell^j) : 0 \leq j < i\}$. Let $c'_{\ell+1}$ be the (unique) block incident to $(d_\ell^i)_{i < k}$, then for every $0 < i < k$, we have that $c'_{\ell+1} \neq b_\ell^i$ since b_ℓ^i is not incident to d_ℓ^0 but $c'_{\ell+1}$ is. Since $<$ is an HF-order, it follows that for every $0 < i < k$ we have

$c'_{\ell+1} > d_\ell^i > b_\ell^i$. Further, notice that by construction we clearly have that:

$$\begin{aligned} \text{cl}_{<}^{\ell+1}(c'_{\ell+1}) &= \{c'_{\ell+1}\} \cup \bigcup \{\text{cl}_{<}^\ell(d_\ell^i) : i < k\}; \\ \text{cl}_{<}^{\ell+2}(c'_{\ell+1}) &= \{c'_{\ell+1}\} \cup \bigcup \{\text{cl}_{<}^{\ell+1}(d_\ell^i) : i < k\}. \end{aligned}$$

Thus we obtain by construction and the induction hypothesis that $Ac \cong_A Ac'_{\ell+1}$ and $A < \text{cl}_{<}^{\ell+2}(c'_{\ell+1})$ (whence also $\text{cl}_{<}^{\ell+2}(c'_{\ell+1})$). Finally, we claim that $Ac'_{\ell+1} \leq_o \text{Acl}_{<}^{\ell+1}(c'_{\ell+1})$. We show how to HF-construct $\text{Acl}_{<}^{\ell+1}(c'_{\ell+1})$ from $Ac'_{\ell+1}$. We start with $Ac'_{\ell+1}$ and define an HF-order $<'$ by adding $d_\ell^{k-1} <' d_\ell^{k-2} <' \dots <' d_\ell^0$ (by the choice of our elements this is an HF-order). Notice now that $c'_{\ell+1}$ is only incident to $(d_\ell^i)_{i < k}$ in $\text{cl}_{<}^{\ell+1}(c'_{\ell+1})$. Also, for every $0 < i < k$ we have by condition (c) that $A \cup \bigcup \{\text{cl}_{<}^\ell(d_\ell^j) : 0 \leq j < i\} < \text{cl}_{<}^{\ell+1}(b_\ell^i)$. Therefore, since $b_\ell^i < d_\ell^i$, it follows that no element in $\text{cl}_{<}^\ell(d_\ell^i)$ is incident to any element in $\text{cl}_{<}^\ell(d_\ell^j)$ for $i \neq j < k$. So using the condition (b) from the inductive hypothesis it is easy to continue to construct the HF-order $<'$.

Case 2. c is a point.

By inductive hypothesis we can find a block b'_ℓ such that (a)-(c) is true for A . In particular $Ab'_\ell \leq_o \text{Acl}_{<}^\ell(b'_\ell)$. Then there is a point $c'_{\ell+1} > b'_\ell$ such that b'_ℓ is incident to $c'_{\ell+1}$. Clearly we have that $\text{cl}_{<}^{\ell+1}(c'_{\ell+1}) = \text{cl}_{<}^\ell(b'_\ell) \cup \{c'_{\ell+1}\}$. From this fact and $Ab'_\ell \leq_o \text{Acl}_{<}^\ell(b'_\ell)$, it is easy to reason as in Case 1 and construct an HF-order witnessing that $Ac'_{\ell+1} \leq_o \text{Acl}_{<}^{\ell+1}(c'_{\ell+1})$. By construction we immediately obtain that $Ac \cong_A Ac'_{\ell+1}$ and, since $\text{cl}_{<}^{\ell+2}(c'_{\ell+1}) = \text{cl}_{<}^{\ell+1}(b'_\ell) \cup \{c'_{\ell+1}\}$, we also have that $A < \text{cl}_{<}^{\ell+2}(c'_{\ell+1})$. \blacksquare

We next consider the other assumption needed to prove that T^+ is complete, which is essentially a corollary of Lemma 4.16.

Proposition 4.23 (3.30(D2)). If $M \models T^+$ is \aleph_1 -saturated, $A \leq_o M$ is countable, and for every trivial extension $A \leq_o Ab$ there is $b' \in M$ such that $Ab \cong_A Ab' \leq_o M$, then for every minimal extension $A \leq_o Ac$ there is $c' \in M$ such that $Ac \cong_A Ac' \leq_o M$.

Proof. Let $M \models T^+$ and $A \leq_o M$. Suppose $A \leq_o Ac$ is a minimal strong extension. We distinguish the following cases.

Case 1. c is a point.

If $A \leq_o Ac$ is a trivial extension then we are immediately done by the assumption. Otherwise, $A \leq_o Ac$ is algebraic and c is incident to exactly one block $b \in A$. Let p_0, \dots, p_ℓ be the points incident to b in A and consider the set $B = \{p_0, \dots, p_\ell, b\}$. Since $M \models T^+$, and since the extension $B \leq_o Bc$ clearly amalgamates (by 4.19) with every $D \subseteq_\omega M$, there is some $c' \in M \setminus A$ incident to b . By the choice of the set B , it also follows that $c' \notin A$ and, since $A \leq_o M$ it follows that c' is incident to no other element in A , yielding $Ac \cong_A Ac'$. By Lemma 4.16, we obtain that $Ac' \leq_o M$.

Case 2. c is a block.

If c is a block and $A \leq_o Ac$ is not trivial, then c is incident to $0 < \ell \leq k$ points $p_0, \dots, p_{\ell-1}$ from A . By assumption, we can find $k - \ell$ new points p_ℓ, \dots, p_{k-1} in $M \setminus A$ which have no incidence with A and such that $A \cup \{p_\ell, \dots, p_{k-1}\} \leq_o M$. Let $c' \in M \setminus A$ be the unique block which is incident to p_0, \dots, p_{k-1} (this exists

because $M \models T^+$). Clearly we have that $Ac \cong_A Ac'$ and, by Lemma 4.16, we obtain $A \cup \{p_\ell, \dots, p_{k-1}\} \cup \{c'\} \leq_o M$. Since p_ℓ, \dots, p_{k-1} are incident with no element in A , we also obtain that $Ac' \leq_o Ac' \cup \{p_\ell, \dots, p_{k-1}\}$. By transitivity we conclude that $Ac' \leq_o M$. \blacksquare

It already follows from the previous propositions and Theorem 3.34 that the theory T^+ of open (k, n) -Steiner systems is complete. We next consider the last condition from 3.28, which is needed to show that T^+ is stable. We start by recalling the free extension process for partial (k, n) -Steiner systems. This makes explicit how the abstract construction of a free completion from 3.41 behaves in this specific setting.

Definition 4.24. Let A be a partial (k, n) -Steiner system. We let $A_0 = A$ and for every $2i < \omega$ we define A_{2i+1} and A_{2i+2} as follows:

- (a) for every sequence of k points p_1, \dots, p_k from A_{2i} which do not have a common incident block in A_{2i} , we add a new block $b \in A_{2i+1}$ which is incident only with p_1, \dots, p_k ;
- (b) for every block $b \in A_{2i+1}$ which is incident only to $\ell < n$ points from A_{2i+1} , we add $n - \ell$ points $p_\ell, \dots, p_{n-1} \in A_{2i+2}$ incident only with b .

Then the structure $F(A) := \bigcup_{i < \omega} A_i$ is called the *free completion* of A and we say that $F(A)$ is freely generated over A . We say that A is *non-degenerate* if $F(A)$ is infinite, and *degenerate* otherwise.

Remark 4.25. Notice that a partial (k, n) -Steiner system A is non-degenerate if and only if it contains at least two blocks or at least $k + 1$ -many points not incident to the same block (cf. [21, p. 756]). By letting $m_{\mathcal{K}} = k + 1$ we thus have that any $A \in \mathcal{K}$ containing at least $m_{\mathcal{K}}$ -many elements of each sort is non-degenerate.

The next proposition provides the key properties of the free completion process for Steiner systems, and it simultaneously verifies Condition 3.28(C2) with $C(A) = F(A)$, thus establishing also Assumption 3.65(C8). It follows that the theory T^+ of open (k, n) -Steiner systems is stable.

Proposition 4.26 (3.28(C2), 3.65(C8)). Suppose that $A \models T_{\forall}$ and that A contains at least $m_{\mathcal{K}} = k + 1$ elements of every sort, then $F(A) \models T^+$, $A \leq_o F(A)$, $|F(A)| = |A| + \aleph_0$ and:

- (a) $F(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : F(A) \cong F(B)$;
- (b) $F(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_o M$ there is an embedding $\hat{h} : F(A) \rightarrow M$ such that $\hat{h} \upharpoonright A = h$ and $\hat{h}(F(A)) \leq_o M$.

Proof. By 3.44 it suffices to show that, if A contains at least $m_{\mathcal{K}} = k + 1$ elements of every sort, then $F(A) \models T^+$. Suppose thus that A is non-degenerate.

The algebraic family of axioms from 3.22(2) is verified immediately by the construction of $F(A)$. In fact, by Remark 4.15 the algebraic extensions of partial (k, n) -Steiner system are exactly those considered in the inductive construction of $F(A)$ and, moreover, for all $i < \omega$ and any algebraic extension $A_i \leq_o A_i c$ and $A_i \leq_o D \subseteq_{\omega} F(A)$, we have that $A_i c \otimes_A D$ is a well-defined (k, n) -Steiner system. We next consider the non-algebraic family of axioms from 3.22(3). Consider a minimal non-algebraic extension $B \leq_o Bc$ with $B \subseteq A_i$, where A_i is the i -th step of the construction from 4.24. We distinguish the following cases.

Case 1. c is a point.

Then $B \leq_o Bc$ is a trivial extension (cf. 3.1). Since A is non-degenerate, for every even $j > i$ there are a block $b_j \in A_{j+1} \setminus A_j$ and a point $c_j \in A_{j+2} \setminus A_{j+1}$ incident to b_j . Then c_j is incident *only* to b_j and so $Bc \cong_B Bc_j \subseteq F(A)$ for all even $j < \omega$.

Case 2. c is a block incident to $\ell < k$ points $p_0, \dots, p_{\ell-1}$ in B .

Since A is non-degenerate, we can find an infinite sequence of indices $i = j_0 < j_1 < j_2 < \dots$ such that for every $0 < m < \omega$ we can find $(k - \ell)$ -many points $p_\ell^{j_m}, \dots, p_{k-1}^{j_m} \in A_{j_m} \setminus A_{j_m-1}$ which are not incident to any common block in A_{j_m} . Consider then, for each $0 < m < \omega$, the block c_{j_m} added in stage $j_m + 1$ of the completion process (or $j_m + 2$, depending on the parity of j_m) which is incident exactly to $p_0, \dots, p_{\ell-1}$ and $p_\ell^{j_m}, \dots, p_{k-1}^{j_m}$. Then $Bc \cong_B Bc_{j_m} \subseteq F(A)$ for all $0 < m < \omega$, proving our claim.

We thus obtained that $F(A)$ satisfies the families of axioms 3.22(2) and 3.22(3). Now for all $a \in F(A)$ let $i(a)$ be the least index such that $a \in A_{i(a)}$. Let $<$ be the partial order given by letting $a < b$ if and only if $i(a) < i(b)$. Then it is immediate to verify that any extension of $<$ into a linear order is an HF-order of $F(A)$, whence we also have that $\emptyset \leq_o F(A)$. It thus follows that $F(A) \models \text{Th}_\forall(\mathcal{K})$, since any counterexample to some $\phi \in \text{Th}_\forall(\mathcal{K})$ would be contained in some $A_i \in \mathcal{K}$. We conclude that $F(A) \models T^+$. \blacksquare

4.2.4. Assumptions 3.60 and 3.65

We consider in this section Conditions 3.60 and 3.65. In order to show that T^+ is not superstable, we first exhibit a configuration in \mathcal{K} witnessing that Assumption 3.60 is satisfied.

Proposition 4.27 (3.60). There is a configuration $C \in \mathcal{K}$ satisfying the conditions from 3.60.

Proof. Recall that k and n are fixed by the definition of (k, n) -Steiner systems. Then we define a configuration C as follows. For simplicity, we let $k = 2$ and $n = 3$, but it is then straightforward to adapt our example to the arbitrary case of $2 \leq k < n$. Let $C = \{c_i : i \leq 12\}$, where $c_0, c_1, c_3, c_5, c_6, c_9, c_{10}, c_{12}$ are points, $c_2, c_4, c_7, c_8, c_{11}$ are blocks, and the incidences between them are specified by the following table.

| | c_2 | c_4 | c_7 | c_8 | c_{11} |
|----------|-------|-------|-------|-------|----------|
| c_0 | × | | × | | |
| c_1 | × | | | × | |
| c_3 | × | × | | | |
| c_5 | | × | × | | |
| c_6 | | × | | × | |
| c_9 | | | × | | × |
| c_{10} | | | | × | × |
| c_{12} | | | | | × |

Then C is a partial $(2, 3)$ -Steiner system and the order $c_0 < c_1 < \dots < c_{12}$ is an HF-order, which means that $C \in \mathcal{K}$. As the other requirements from 3.60 are easily verified, it follows that C is as desired. \blacksquare

Finally, we consider the conditions from 3.65. First, notice that Assumption 3.65(C7) follows immediately from 4.26. Thus it remains to consider 3.65(C8) and 3.65(C9).

Proposition 4.28. If $F(A) \models T^+$ and $B \in \mathcal{K}$, then $F(B)$ embeds into $F(A)$.

Proof. Let $A = \bigcup_{i < \omega} A_i$ as in 4.24. Suppose for simplicity that B contains exclusively points. Then, for $\ell < \omega$ large enough, we can find a stage A_ℓ in the free completion $F(A)$ of A , such that A_ℓ contains an isomorphic copy C of B such that $A_\ell \setminus C$ does not contain any block incident to at least k -many points in C . Then, it is clear from the definition of free completion from 4.24 that the free completion $F(A)$ will contain an isomorphic copy $F(C)$ of $F(B)$. Finally, we notice that if B also contains blocks, then this argument can be adapted by making sure that every two different blocks $b, b' \in B$ are mapped to blocks $c, c' \in C \subseteq A_\ell$ in such a way that all the points incident to c and c' in $A_\ell \setminus C$ have no common incidence in A_ℓ . ■

Finally, we introduce a suitable predimension function for the setting of open (k, n) -Steiner systems. To our knowledge, this had not been considered previously in the literature.

Definition 4.29. Let $A \in \mathcal{K}$ be a finite, open, partial (k, n) -Steiner system, we define a *predimension function* $\delta : \mathcal{K} \rightarrow \omega$ as follows. Let P_A , L_A and I_A denote points, blocks and incidences of A , respectively, then we define

$$\delta(A) = |P_A| + k|L_A| - |I_A|.$$

Proposition 4.30 (3.65(C9)). The predimension function $\delta : \mathcal{K} \rightarrow \omega$ satisfies the properties from Definition 3.64, i.e., if $A, B \in \mathcal{K}$, then:

- (a) $A \cong B$ implies $\delta(A) = \delta(B)$;
- (b) $A \leq_o B$ implies $A \leq_\delta B$;
- (c) if $A \leq_o B$ is minimal, then $\delta(A) = \delta(B)$ if and only if B is algebraic over A ;
- (d) if $A \leq_\delta B$ and $C \subseteq B$, then $A \cap C \leq_\delta C$.

Proof. Clause (a) follows immediately by the definition of the predimension δ . Then, by definition, it is clear that if $A \leq_o Ab$ is an algebraic strong extension, then $\delta(A) = \delta(Ab)$. Differently, if $A \leq_o Ab$ is a non-algebraic strong extension we have that $\delta(A) < \delta(Ab)$. Proceeding inductively, it follows that whenever $A \leq_o C \in \mathcal{K}$ then $\delta(A) \leq \delta(C)$.

It thus remain to verify condition (c). Now, if $X, Y \subseteq A$ are disjoint, we denote by $I(X, Y)$ the set of incidences between X and Y in A . Suppose $A \leq_\delta B$ and let $C \subseteq B$. By 4.29 we have that:

$$\begin{aligned} \delta(A \cap C) &= |P_{A \cap C}| + k|L_{A \cap C}| - |I_{A \cap C}| \\ &= |P_C| - |P_{C \setminus A}| + k|L_C| - k|L_{C \setminus A}| - |I_C| + |I_{C \setminus A}| + I(C, C \setminus A) \\ &= \delta(C) - \delta(C \setminus A) + I(C, C \setminus A). \end{aligned}$$

Since $A \leq_\delta B$ we have that $\delta(A) \leq \delta(AC)$, thus $\delta(A) \leq \delta(A) + \delta(C \setminus A) - I(C, C \setminus A)$ and in particular $\delta(C \setminus A) - I(C, C \setminus A) \geq 0$. It follows that

$$\delta(A \cap C) = \delta(C) - \delta(C \setminus A) + I(C, C \setminus A) \leq \delta(C)$$

which proves $A \cap C \leq_\delta C$. ■

Finally, this concludes the verification of all assumptions from 3.2, 3.28, 3.43, 3.60 and 3.43. Therefore, in light of the results from Section 3, this completes the proof of Theorem 4.12.

4.3. Generalised n -gons

In this section we apply our framework to the case of generalised n -gons. We stress again that this has already been considered in the literature. Most specifically, [24] considered the case of projective planes, i.e., 3-gons, while more recently [1] considered arbitrary generalised n -gons. Possibly, this is most interesting application of our framework, since generalised n -gons are the rank 2 case of spherical buildings and thus have a natural connection to higher-rank geometries, which, as mentioned in the introduction, we intend to explore in a work in preparation. Additionally, generalised n -gons have a vast number of connections to the general mathematical literature; for an overview of such connections we refer the interested reader to [37].

Since our techniques differ from those employed in [1] (although they are very much inspired by them), we provide most of the details of the proofs, except for those proofs which proceed exactly as in the case of Steiner systems, or that are similar to [1] and [24]. In particular, we remark that the case of generalised n -gon is the only concrete examples where a minimal extension $A \leq_o B$ is not necessarily a one-element extension, i.e., where $B \setminus A$ is not necessarily a singleton (see also Remarks 4.35 and 4.39 later). We recall that it is for this reason that in Condition 3.2(I) we introduced an arbitrary finite set $X_{\mathcal{K}}$, which encodes the possible length of minimal strong extensions in \mathcal{K} . To show that our framework accounts also for this more complicated case, we provide several details in the proofs of the conditions from 3.2, 3.28 and 3.43. We will however omit the details of 3.60 and 3.65, as these were essentially already considered in the literature (cf. Remark 4.45). All in all, the following theorem (which was proved already in [34]) holds in the setting of generalised n -gons.

Theorem 4.31. *Let (\mathcal{K}, \leq_o) be the class of finite, open, partial generalised n -gons and let T^+ be defined as in 3.22. Then T^+ is complete, stable, and with $\perp = \perp^{\otimes}$. Moreover, T^+ is not superstable, it does not have a prime model, it is not model complete and does not eliminate quantifiers.*

We start by recalling some graph-theoretic notions, and the graph-theoretic definition of the generalised n -gons (cf. [1, Def. 2.1]). A different and more geometrical definition of n -gons can be found, e.g., in [21, p. 745].

Definition 4.32. Let L be the two-sorted language where the symbols p_0, p_1, \dots denote *points* and the symbols ℓ_0, ℓ_1, \dots denote *lines*, the symbol $p|\ell$ (or $\ell|p$) means that the point p is *incident* with the line ℓ . For every $n \geq 3$, we define a *generalised n -gon* as a bipartite graph (with points and lines) with girth $2n$ and diameter n . A *partial n -gon* is a model of the universal fragment of the theory of generalised n -gons, i.e., it is a bipartite graph with girth at least $2n$.

Remark 4.33. We notice that, by the fact that partial n -gons are bipartite, they do not have any odd-length cycle. Moreover, differently from [1], we do not assume that partial n -gons are always connected.

Definition 4.34. For finite partial n -gons $A \subseteq B$, we define the following.

- (1) A tuple $\bar{a} \in A^{<\omega}$ is *hyperfree in B* if it satisfies one of the following conditions:
 - (i) $\bar{a} = a$ is a *loose end*, i.e., it is a single element incident with at most one element in B ;
 - (ii) \bar{a} is a *clean arc*, i.e., \bar{a} is a chain $a_1 | a_2 | \dots | a_{n-3} | a_{n-2}$ where each a_i for $1 \leq i \leq n-2$ is incident exactly to two elements in B .
- (2) We say that B is *open over A* and write $A \leq_o B$ if every non-empty $C \subseteq B \setminus A$ contains a tuple which is hyperfree in AC . We say that B is *open* if $\emptyset \leq_o B$, i.e., if every non-empty $C \subseteq B$ contains a tuple hyperfree in C .
- (3) We say that B is *closed over A* if $A \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B \setminus A$ which does not contain any tuple hyperfree in AC . We say that B is *closed* if $\emptyset \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B$ which does not contain any tuple hyperfree in C .

We let \mathcal{K} be the set of finite, open, partial n -gons and we let T_{\forall} be the universal theory of \mathcal{K} . The previous definitions extend to infinite models of T_{\forall} as in 3.6.

Remark 4.35. We stress that our notion of hyperfree tuples follows [1, Def. 2.10] and it differs from both [20, p. 351] and [21, p. 753]. In fact, the issue with these latter definitions is that they are introduced at the level of single elements, which creates problems when considering the amalgamation property (we expand on this issue later in Remark 4.39). We thus stress that generalised n -gons are the only concrete examples where we use the full generality of our setting, i.e., where minimal hyperfree extensions are not necessarily one-element extensions but can also consist of longer tuples. In particular, we have that an extension $A \leq_o B \in \mathcal{K}$ is minimal if $B \setminus A = \{c\}$ is a single element incident to at most one element from A , or if $B \setminus A$ is a clean arc.

To see that clean arcs form minimal extensions, notice that if $A \leq_o B$ and $B \setminus A$ is a clean arc \bar{c} , then given $A \subsetneq C \subseteq B$ we have that every element in $B \setminus C$ is incident to two elements from B , but the clean arc \bar{c} is not fully included in $B \setminus C$. This shows that $C \not\leq_o B$ and proves that clean arcs form minimal extensions.

Secondly, we claim that extensions $A \leq_o A\bar{c}$ where \bar{c} is a clean arc are exactly the only minimal algebraic extensions. In fact, if a and b are the endpoints of the clean arc \bar{c} then $d_{A\bar{c}}(a, b) = n-1$. Hence, if $C \in \mathcal{K}$ contains an isomorphic copy of $A\bar{c}$ over A , it follows that a, b belong to two paths of length $n-1$, which together form a $(n-1)$ -gon (i.e., a cycle of length $2n-2$). This contradicts $C \in \mathcal{K}$ and shows that $A \leq_o A\bar{c}$ is algebraic. On the other hand, it is immediate to see that extensions by loose ends are not algebraic.

The next Lemma is essentially Lemma 4.16 but for partial n -gons. We omit the proof as it is fundamentally the same. Notice also that this is exactly Assumption 3.43(C5) for the case of n -gons. As in the setting of Steiner systems, it will often be useful in the subsequent proofs.

Lemma 4.36 (3.43(C5)). Let $A \leq_o C \models T_{\forall}$ and let $\bar{c} \in C^{<\omega}$ be such that $A \leq_o A\bar{c}$ is a minimal algebraic extension. Then $A\bar{c} \leq_o C$.

Proof. The proof follows analogously to the proof of Lemma 4.16. ■

We start by verifying that the class (\mathcal{K}, \leq_o) satisfies the conditions from 3.2. As in the case of Steiner systems, conditions (A), (B), (D), (E), (F), (K) and (L) from 3.2 follow immediately from Definition 4.34. Then, Condition 3.2(C) follows exactly as in the case of Steiner systems and Condition 3.2(H) follows exactly as

in [1, Lemma 2.24]. By Remark 4.35 it is easy to see that Condition 3.2(I) holds by letting $X_{\mathcal{K}} = \{1, n-2\}$ – namely, minimal strong extensions are obtained by adjoining either a loose end or a clean arc of length $n-2$. Condition 3.2(J) holds immediately with $n_{\mathcal{K}} = 1$. Finally, although Condition 3.2(G) is exactly [1, Prop. 2.21], we provide details of this proof to make it clear that it holds also when considering non-connected partial n -gons. As we remark later in 4.39, amalgamation is the main reason why we need to consider minimal extensions which are not one-element extensions. Notice that, by the next remark, it follows as in the case of Steiner systems that the next proof also establishes 3.43(C4).

Remark 4.37 (3.43(C4)). Let $A \subseteq B, C$ be open partial n -gons and recall the definition of free amalgam from 3.39. We notice that, in general, the free amalgam $B \otimes_A C$ is not necessarily an open partial n -gon, as the amalgam may introduce cycles of length $< 2n$. However, it follows from the proof of Proposition 4.38 that when the conditions from 3.43(C4) are satisfied then $B \otimes_A C \in \mathcal{K}$.

The proof of the next proposition is exactly as in [1, Prop. 2.21], but we make all details explicit for the benefit of the reader and to show that it works also when (\mathcal{K}, \leq_o) contains also non-connected open, partial n -gons.

Proposition 4.38 (3.2(G)). The class (\mathcal{K}, \leq_o) has both the amalgamation and joint embedding property.

Proof. Let $A, B, C \in \mathcal{K}$ with $A \leq_o B, A \leq_o C$ and $B \cap C = A$. As in the case of (k, n) -Steiner systems, if we have that $B \otimes_A C \in \mathcal{K}$ then it follows that $B \leq_o B \otimes_A C$ and $C \leq_o B \otimes_A C$. Thus it suffices to verify that either there is an embedding $f : B \rightarrow C$ such that $f(B) \leq_o C$ or $B \otimes_A C \in \mathcal{K}$, i.e., $B \otimes_A C$ is a well-defined open partial n -gon. Reasoning by induction, we assume that $A \leq_o B$ is a minimal strong extension, i.e., we distinguish the following cases.

Case 1. $B \setminus A = \{c\}$ and c is incident to at most one element in A .

Then the free amalgam $B \otimes_A C$ is an open partial n -gon, whence $B \otimes_A C \in \mathcal{K}$.

Case 2. $B \setminus A = \{c_1, \dots, c_{n-2}\}$ and $\bar{c} = (c_1, \dots, c_{n-2})$ is a clean arc with endpoints $a, b \in A$.

Case 2.1. There is a clean arc $\bar{d} = (d_1, \dots, d_{n-2}) \in C^n$ with endpoints $a, b \in A$.

Then define $f : B \rightarrow C$ such that $f \upharpoonright A = \text{id}_A$ and $f(c_i) = d_i$ for all $1 \leq i \leq n-2$. This is obviously an embedding. Moreover, since $f(B) = A\bar{d}$ and \bar{d} is algebraic over A , it follows by 4.16 and $A \leq_o C$ that $f(B) \leq_o C$.

Case 2.2. There is no clean arc $\bar{d} = (d_1, \dots, d_{n-2}) \in C^{n-2}$ with endpoints $a, b \in A$. Now, since $A\bar{c}$ has girth $\geq 2n$, it follows in particular that $d(a, b/A) \geq n+1$, for otherwise we would have a short cycle in $A\bar{c}$. We thus obtain that the shortest cycle in $B \otimes_A C$ to which a and b belong contains at least $2n$ -many elements. Thus, the free amalgam $B \otimes_A C$ is an open partial n -gon in \mathcal{K} . This completes our proof. ■

Remark 4.39. It is crucial that in the previous Proposition 4.38 we regard the extension $A \leq_o A\bar{c}$, with \bar{c} a clean arc, as a minimal extension. In fact, suppose otherwise that one defines hyperfree extensions as in [20, p. 351] and replaces clause (1) from Def. 4.34 with one saying that an element $a \in A$ is hyperfree if and only if it is incident with at most two elements from B . Then we have the following counterexample to the amalgamation property. Let $n > 3$ and consider the partial n -gon $A = \{a_1, a_2, b_1, b_2\}$ with $a_1 \mid a_2, b_1 \mid b_2$, and where a_1, b_1 are points, a_2, b_2

lines. Consider then two extensions Ac and Ad obtained by adding a line c incident to a_1, b_1 , and a point d incident to a_2, b_2 . Clearly, according to the alternative definition above, we have $A \leq_o Ac$ and $A \leq_o Ad$. However, if any $C \in \mathcal{K}$ amalgamates both Ac and Ad , then C contains the short cycle $a_1 | a_2 | d | b_2 | b_1 | c | a_1$, contradicting $C \in \mathcal{K}$. It is easy to adapt this counterexample to other definitions of hyperfree extensions based on single elements, e.g., to [21, p. 753] (or also to the setting where partial n -gons are taken to be connected).

We next consider the assumptions from 3.28. Recall that to show 3.28(C1) it suffices to establish 3.30(D1) and 3.30(D2). We first prove condition 3.30(D1). As in the setting of Steiner systems, we add a further condition (c) which is useful in the induction. Recall also the notation from 3.11: if (\langle, P) is a HF-order of M and $c \in M$, then we write \widehat{c} for the tuple (c_0, \dots, c_k) such that $\{c_0, \dots, c_k\} \in P$ and $c_0 < \dots < c_n$.

Proposition 4.40 (3.30(D1)). If $M \models T^+$, $A \subseteq M$ is finite, \langle is an HF-order of M and $A \leq_o Ac$ is a trivial extension, then for all $\ell < \omega$ there is $c' = c'_\ell \in M$ s.t.:

- (a) $Ac \cong_A Ac'$;
- (b) $Ac' \leq_o \text{Acl}_\langle^\ell(c')$;
- (c) $d(A, c'/\text{Acl}_\langle^{\ell+1}(c')) > \ell + n$.

Proof. We assume w.l.o.g. that n is odd, for the case with n even is analogous. In particular this means that clean arcs have as endpoints elements of the same sort. We proceed similarly to the proof of 4.22. By induction on $\ell < \omega$ we prove that for every $A \subseteq M$ and $A \leq_o Ac \in \mathcal{K}$ we can find c'_ℓ satisfying conditions (a)-(c).

Base case $\ell = 0$.

We find an element c'_ℓ as wanted. Since $M \models T^+$ and \langle is a HF-order, there are cofinitely-many clean arcs in the ordering \langle (for otherwise we would have points at distance $> n$). It follows in particular that we can find a clean arc $\bar{x} = (x_1, \dots, x_{n-2})$ with endpoints $b, c \in M$ such that \widehat{b} and \widehat{c} do not contain any element from A . Similarly, let $\bar{y} = (y_1, \dots, y_{n-2}) > \bar{x}$ be another such clean arc with endpoints $d, e \in M$ such that \widehat{d} and \widehat{e} do not contain any element from A . Additionally, we assume w.l.o.g. that b, c, d, e all have the same sort and that $x_1 < \dots < x_{n-2} < y_1 < \dots < y_{n-2}$. Let $k = \frac{n-1}{2}$. Then in y_{n-2}^\downarrow the elements x_k and y_k must have distance $\geq n+1$, since the elements b, c, d, e are pairwise different and have the same sort. Consider then the clean arc $\bar{z} = (z_1, \dots, z_{n-2})$ between x_k and y_k and notice that we must have $z_i > y_{n-2}$ for all $1 \leq i \leq n-2$. Then we let $c'_0 = z_k$ and from 3.19 it follows that we have:

$$\begin{aligned} \text{cl}_\langle^1(c'_0) &= \{x_1 \dots, x_{n-2}, y_1 \dots, y_{n-2}, z_1 \dots, z_{n-2}\} \\ \text{cl}_\langle^0(c'_0) &= \{z_1 \dots, z_{n-2}\}. \end{aligned}$$

Then, we obtain that $d(A, c'_0/\text{Acl}_\langle^1(c'_0)) > n$, as no element from the tuples $\bar{x}, \bar{y}, \bar{z}$ is incident to elements from A . Similarly, we also have that $Ac'_0 \leq_o \text{Acl}_\langle^0(c'_0)$, as we can define an HF-order \langle' by letting $Ac'_0 \langle' z_{k-1} \langle' \dots \langle' z_1 \langle' z_{k+1} \langle' \dots \langle' z_{n-2}$. Finally, we clearly have that $Ac \cong_A Ac'_0$.

Inductive step.

Now, suppose the inductive hypothesis holds for $\ell \geq n$ and let c be such that $A \leq_o Ac$ is a trivial extension, we want to define $c'_{\ell+1}$ so that (a)-(c) are satisfied for it. By inductive hypothesis we can find a point d'_ℓ satisfying (a)-(c) with respect

to A . Additionally we can find a second point d_ℓ^1 such that d_ℓ^1 is a trivial extension of $\text{Acl}_<^\ell(d_\ell^0)$ and (a)-(c) are true with respect to it. Now, by the choice of d_ℓ^1 and condition (c) it follows that there is no path of length $\leq n$ between d_ℓ^0 and d_ℓ^1 in $\text{cl}_<^\ell(d_\ell^1)$. But then since $d_\ell^0 < d_\ell^1$ and $<$ is a HF-order, it follows that the shortest path between d_ℓ^0 and d_ℓ^1 is a clean arc $\bar{z} = (z_1, \dots, z_{n-2})$ s.t. $d_\ell^1 < z_1 < \dots < z_{n-2}$. Let $c'_{\ell+1} = z_1$, then we have:

$$\begin{aligned} \text{cl}_<^{\ell+2}(c'_{\ell+1}) &= \{z_1, \dots, z_{n-2}\} \cup \text{cl}_<^{\ell+1}(d_\ell^0) \cup \text{cl}_<^{\ell+1}(d_\ell^1), \\ \text{cl}_<^{\ell+1}(c'_{\ell+1}) &= \{z_1, \dots, z_{n-2}\} \cup \text{cl}_<^\ell(d_\ell^0) \cup \text{cl}_<^\ell(d_\ell^1). \end{aligned}$$

Whence, from the induction hypothesis and the display above we immediately obtain that $d(A, c'_{\ell+1}/\text{Acl}_<^{\ell+2}(c'_{\ell+1})) > (\ell+1) + n$. Similarly, it is clear from the choice of $c'_{\ell+1}$ that we also have $Ac \cong_A Ac'_{\ell+1}$. Finally, to see that $Ac'_{\ell+1} \leq_o \text{Acl}_<^{\ell+1}(c'_{\ell+1})$ we show that $Ac'_{\ell+1} \leq_{\text{HF}} \text{Acl}_<^{\ell+1}(c'_{\ell+1})$. We start with $Ac'_{\ell+1}$ and define an HF-order $<'$ by adding $d_\ell^0 <' z_2 <' z_3 \dots <' z_{n-2} <' d_\ell^1$. Crucially, each new element that we add has only one incidence with the previous ones, thus $<'$ is an HF-order. Finally, notice that $c'_{\ell+1}$ is only incident to z_2 and d_ℓ^0 in $\text{cl}_<^{\ell+2}(c'_{\ell+1})$, and also that no element in $\text{cl}_<^\ell(d_\ell^0)$ is incident to any element in $\text{cl}_<^\ell(d_\ell^1)$ by the choice of d_ℓ^1 . So using the inductive hypothesis it is easy to continue to construct the HF-order $<'$. This finally shows that also $Ac'_{\ell+1} \leq_o \text{Acl}_<^{\ell+1}(c'_{\ell+1})$. \blacksquare

The proof of the other assumption needed to prove that T^+ is similar to the case of Steiner systems. As in the setting of Steiner systems, the key ingredient of the next proof is Lemma 4.36.

Proposition 4.41 (3.30(D2)). If $M \models T^+$ is \aleph_1 -saturated, $A \leq_o M$ is countable, and for every trivial extension $A \leq_o Ab$ there is $b' \in M$ s.t. $Ab \cong_A Ab' \leq_o M$, then for every minimal extension $A \leq_o A\bar{c}$ there is $\bar{c}' \in M$ s.t. $A\bar{c} \cong_A A\bar{c}' \leq_o M$.

Proof. Let $M \models T^+$ and $A \leq_o M$. Suppose $A \leq_o A\bar{c}$ is a minimal strong extension. If \bar{c} is a single element with no incidence with elements in A then we are immediately done by the inductive assumption, thus we distinguish the following two cases.

Case 1. $\bar{c} = c$ is an element incident with one element in A .

Suppose w.l.o.g. that c is a point incident with a line $d \in A$. By assumption, we can find an element $d' \in M$ such that $Ad' \leq_o M$ and d' has no incidence with A . Additionally, we can choose d' so that it has an appropriate sort, i.e., it is a point if n is even and it is a line if n is odd. Then, we have that $d(d, d'/M) \geq n+1$, for otherwise $M \setminus Ad'$ contains a short chain contradicting $Ad' \leq_o M$. It follows that $\{d, d', b_1, \dots, b_{n-2}\}$, where $\bar{b} = (b_1, \dots, b_{n-2})$ is a clean arc between d and d' of the appropriate length, amalgamates with every finite subset from M . Since $M \models T^+$ there must be a clean arc $\bar{z} \in M^{n-2}$ with endpoints d and d' . Let z_1 be the first element from \bar{z} , clearly $Ac \cong_A Az_1$. By 4.36 we have that $Ad'\bar{z} \leq_o M$. Moreover, we have that $Az_1 \leq_o Ad'\bar{z}$, since we can obtain $Ad'\bar{z}$ by a sequence of one-element extensions from Az_1 . It follows by transitivity that $Az_1 \leq_o M$.

Case 2. \bar{c} is a clean arc with endpoints in A .

Let $a, b \in A$ be the endpoints of \bar{c} . Since $A \leq_o A\bar{c}$ we must have $d_A(a, b) \geq n+1$. Crucially, $A \leq_o M$, thus $d_M(a, b)$ cannot be strictly less than n . Therefore $A\bar{c}$ amalgamates over A with every subset $D \subseteq M$, and since $M \models T^+$ it follows that there is a clean arc $\bar{d} \in (M \setminus A)^{<\omega}$ with endpoints a and b . Clearly $A\bar{c} \cong_A A\bar{d}$ and, by 4.36, it follows that $A\bar{d} \leq_o M$. This completes our proof. \blacksquare

From the previous propositions and 3.34 it already follows that the theory T^+ is complete. We next exhibit how the free completion process from 3.41 behaves in the setting of generalised n -gons. We notice that what Definition 3.41 delivers is slightly different from the usual notion of free completion for generalised n -gons. In fact, we recall that the free completion process for partial n -gons was originally introduced by Tits [36] and can be found also in [1, Def. 2.4] and [21, p. 755]. In these definitions free completions are defined only respectively to *connected* partial n -gons. Our construction from 3.41 applies also to the non-connected case, and it essentially delivers a minor variant of Tits' free completion process. We stress that, when restricted to connected partial n -gons, our construction is equivalent to the standard one from [36], and thus it is a conservative extensions of Tits' definition.

Definition 4.42. Let A be a partial n -gon. We let $A_0 = A$ and, for every $i < \omega$, we let A_{i+1} be obtained by adding a clean arc $a | z_1 | z_2 | \dots | z_{n-2} | b$ between any two elements $a, b \in A_i$ such that

- (a) either $d(a, b/A_i) = n + 1$,
- (b) or $d(a, b/A_i) = \infty$ and a, b are of the appropriate sort (i.e., a, b have the same sort if n is odd and have different sorts if n is even).

The structure $F(A) := \bigcup_{i < \omega} A_i$ is called the *free completion* of A and we say that $F(A)$ is freely generated over A . We say that A is *non-degenerate* if $F(A)$ is infinite, and *degenerate* otherwise.

Remark 4.43. The standard free completion process for generalised n -gons differs from the one above in that it does not include our clause (b). Without (b), one has that that A generates an infinite model if it is both connected and, either it contains a cycle of length $2n + 2$, or has diameter $\geq n + 2$ (cf. [21, p. 755]). Given our slightly different definition, we can replace the requirement of connectedness with one saying that $|A| \geq 3$ and that A contains element of both sorts. In fact, if this is the case, then clause (b) from 4.42 makes sure that in A_1 we introduce at least one clean arc, and thus A_2 is in turn a connected partial n -gon. In particular, it follows that if A contains at least $m_{\mathcal{K}} = 2n + 2$ elements of both sorts, then $F(A)$ is infinite.

The next proposition verifies simultaneously Assumption 3.28(C2) and Assumption 3.65(C8), since the structure $C(A)$ is exactly $F(A)$.

Proposition 4.44 (3.28(C2), 3.65(C8)). Suppose $A \models T_{\forall}$, and A is non-degenerate, then $F(A) \models T^+$, $A \leq_o F(A)$, $|F(A)| = |A| + \aleph_0$ and:

- (a) $F(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : F(A) \cong F(B)$;
- (b) $F(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_o M$ there is an embedding $\hat{h} : F(A) \rightarrow M$ such that $\hat{h} \upharpoonright A = h$ and $\hat{h}(F(A)) \leq_o M$.

Proof. As in 4.26, it suffices to verify that $F(A) \models T^+$, provided that A is non-degenerate. Let $A' \subseteq_{\omega} F(A)$ and suppose $A' \leq_o A'\bar{c}$ is a minimal algebraic extension which can be amalgamated with every $D \subseteq F(A)$ with $|D \setminus A'| \leq n - 2$. By Remark 4.35 \bar{c} is a clean arc between two elements $a, b \in A'$ and thus in particular $d(a, b/A') \geq n + 1$. Since A is non-degenerate, there is a minimal stage $i < \omega$ such that $\{a, b\} \subseteq A_i$ and $d(a, b/A_i) \leq n + 1$. Then, since $A'\bar{c}$ can be

amalgamated over A' with any subset $D \subseteq F(A)$ with $|D \setminus A'| \leq n - 2$, it follows that $d(a, b/A_i) = n + 1$. By the definition of free completion, there is a clean arc $\bar{d} \in A_{i+1}^{<\omega}$ such that $A'\bar{c} \cong_A A'\bar{d}$. This shows that $F(A)$ satisfies the second family of axioms from 3.22. Then, by reasoning exactly as in the case of Steiner systems, one can also verify that $F(A)$ satisfies axioms 3.22(1) and 3.22(3), thus showing $F(A) \models T^+$. ■

Remark 4.45. In this section we have verified all the conditions from 3.2, the assumptions from 3.28 and those from 3.43. This entails that T^+ is complete, stable and with $\downarrow = \downarrow^\otimes$. We do not provide full details for the fact that T^+ is not superstable and it does not have a prime model. We notice that a suitable notion of predimension for the generalised n -gons can be found in [21] and it was used in [1] to prove that T^+ does not have a prime model. Similarly, the construction of a concrete configuration satisfying Assumption 3.60 can be essentially found in [24, 9.1] for the case of projective planes. This can be adapted to all $n \geq 3$ by replacing with a clean arc of length $n - 2$ those points and lines which have two incidences with the previous elements in the HF-order of the construction.

4.4. k -nets

We consider in this section k -nets, which are a third example of incidence structures of rank 2 and which admit a free algebraic completion. As we make explicit later, we axiomatise k -nets as structures without any parallelism relation, although in other presentations one has a notion of parallelism between lines. This indicates that, in a certain sense, k -nets are a hybrid form of incidence structures that lies between those with parallelism relations and those without. We refer the reader to [29, p. 20] and [21] for more information on k -nets. We proceed as in the previous cases and we verify all the conditions and assumptions from 3.2, 3.28, 3.43, 3.60 and 3.65. One then obtains the following theorem from the results in Section 3.

Theorem 4.46. *Let (\mathcal{K}, \leq_o) be the class of finite, open, partial k -nets and let T^+ be defined as in 3.22. Then T^+ is complete, stable, and with $\downarrow = \downarrow^\otimes$. Moreover, T^+ is not superstable, it does not have a prime model, it is not model complete and does not eliminate quantifiers.*

We recall the axiomatisation of k -nets, which we take from [21, pp. 745-746] with only small modifications.

Definition 4.47. For any $k \geq 3$ we fix the two-sorted language L with one sort for points, one sort for lines, one incidence symbol $|$ between points and lines, and k -many predicates $(P_i)_{i < k}$ for lines. We denote points by p_0, p_1, \dots and lines by ℓ_0, ℓ_1, \dots . The theory of k -nets is axiomatised by the following axioms:

- (1) every line ℓ belongs to some P_i , for $i < k$;
- (2) for all points p and predicates P_i , with $i < k$, there is a unique line ℓ such that $p|\ell$ and $\ell \in P_i$;
- (3) for any $i < j < k$ and lines $\ell \in P_i$ and $\ell' \in P_j$, there is a unique point p such that $p|\ell$ and $p|\ell'$.

The universal fragment of the theory of k -nets is axiomatised by axiom (1) together with the uniqueness requirement from axioms (2) and (3). We call a model of the universal theory of k -nets a *partial k -net*.

We often refer to the predicates $(P_i)_{i < k}$ as the *parallelism type* of lines. We notice that this is a slight abuse of notation, as this notion of parallelism crucially differs from the relation of parallelism from e.g., affine planes or Laguerre planes, as it does not play a role in the definition of hyperfree elements and we do not treat it as a local equivalence relation. As a matter of fact, in other treatments of nets in the literature such predicates are internalised as added sorts, e.g., in [21]. We decided to distinguish these predicates from the two sorts of points and lines to emphasise the fact that nets are also an instance of rank-2 geometries. We refer the interested reader also to [29, 1.2] for a similar presentation.

Definition 4.48. For finite partial k -nets $A \subseteq B$, we define the following.

- (1) An element $a \in A$ is *hyperfree in B* if it satisfies one of the following conditions:
 - (i) a is a point incident with at most two lines from B ;
 - (ii) a is a line incident with at most one point from B .
- (2) We say that B is *open over A* and write $A \leq_o B$ if every non-empty $C \subseteq B \setminus A$ contains an element which is hyperfree in AC . We say that B is *open* if $\emptyset \leq_o B$, i.e., if every non-empty $C \subseteq B$ contains some element hyperfree in C .
- (3) We say that B is *closed over A* if $A \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B \setminus A$ which does not contain any element hyperfree in AC . We say that B is *closed* if $\emptyset \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B$ which does not contain any element hyperfree in C .

We let \mathcal{K} be the set of open partial finite k -nets and we let T_{\forall} be the universal theory of \mathcal{K} . The previous definitions extend to infinite models of T_{\forall} as in 3.6. The notion of hyperfree element in a k -net is exactly as in [21, p. 753].

The following remark makes it explicit what are the algebraic extensions in the setting of nets. The subsequent Lemma is then proved exactly as in the previous cases, and also establishes 3.43(C5).

Remark 4.49. In the case of k -nets, a minimal extensions $A \leq_o Ac$ is *algebraic* if c is a point incident to two lines in A or it is a line incident to some point from A .

Lemma 4.50. Let $A \leq_o C \models T_{\forall}$ and let $c \in C$ be such that $A \leq_o Ac$ is an algebraic strong extension. Then $Ac \leq_o C$.

Proof. This is proved by reasoning exactly as in 4.16 and 4.36. ■

Consider firstly the conditions from 3.2. As in the previous cases, conditions (A), (B), (D), (E), (F), (K) and (L) follow immediately from our definition of the relation \leq_o for finite, open, partial k -nets. Condition 3.2(C) follows exactly as in the case of Steiner systems. Condition (I) follow immediately by letting $X_{\mathcal{K}} = \{1\}$, as it is easy to verify that the the minimal extensions for k -nets are exactly the one-element extensions. Condition (J) follows immediately by letting $n_{\mathcal{K}} = 2$. Conditions (G) and (H) are non-trivial, but we omit their proof as it is essentially identical to the case of Steiner systems. Similarly to that case, one can also verify simultaneously that 3.43(C4) holds. We consider in details the assumptions from 3.28. As in the previous cases we first prove 3.30(D1) and 3.30(D2). We add a third condition (c) to 3.30(D1) to ease our inductive argument.

Proposition 4.51 (3.30(D1)). If $M \models T^+$, $A \subseteq M$ is finite, $<$ is an HF-order of M and $A \leq_o Ac$ is a trivial extension, then for all $n < \omega$ there is $c' = c'_n \in M$ s.t.:

- (a) $Ac \cong_A Ac'$;

- (b) $Ac' \leq_o \text{Acl}_{<}^n(c')$;
(c) $A < \text{cl}_{<}^{n+1}(c')$.

Proof. By induction on $n < \omega$ we prove that for every finite $A \subseteq M$ and $A \leq_o Ac$ we can find c'_n satisfying the conditions (a)-(c). If $n = 0$, then it is easy to see that we can find c'_1 as wanted. So suppose the inductive hypothesis and let c be such that $A \leq_o Ac$ is a trivial extension, we want to define c'_{n+1} so that (a)-(c) are satisfied for it.

Case 1. c is a line (of parallelism type $i < k$).

Firstly, since $k \geq 3$, we notice there are two indices i_0, i_1 both different from i . By inductive hypothesis we can find a line $d_n^0 \in P_{i_0}$ such that $A < \text{cl}_{<}^{n+1}(d_n^0)$ and $Ad_n^0 \leq_o \text{Acl}_{<}^n(d_n^0)$. Also, we can find a further line $d_n^1 \in P_{i_1}$ such that $\text{Acl}_{<}^{n+1}(d_n^0) < \text{cl}_{<}^{n+1}(d_n^1)$ and $\text{Acl}_{<}^{n+1}(d_n^0)d_n^1 \leq_o \text{Acl}_{<}^{n+1}(d_n^0)\text{cl}_{<}^n(d_n^1)$. Consider the unique point p incident to both d_n^0 and d_n^1 , then since $\text{Acl}_{<}^{n+1}(d_n^0) < \text{cl}_{<}^{n+1}(d_n^1)$ it follows that $p > d_n^1 > d_n^0$.

Then, since $<$ is an HF-order and $M \models T^+$, there is a line $c'_{n+1} \in P_i$ such that $c'_{n+1} > p$ and $c'_{n+1} \mid p$. Then we have that:

$$\begin{aligned} \text{cl}_{<}^{n+1}(c'_{n+1}) &= \text{cl}_{<}^n(d_n^0) \cup \text{cl}_{<}^n(d_n^1) \cup \{p, c'_{n+1}\}; \\ \text{cl}_{<}^{n+2}(c'_{n+1}) &= \text{cl}_{<}^{n+1}(d_n^0) \cup \text{cl}_{<}^{n+1}(d_n^1) \cup \{p, c'_{n+1}\}. \end{aligned}$$

It follows by construction that $A < \text{cl}_{<}^{n+2}(c'_{n+1})$ and $Ac \cong_A Ac'_{n+1}$. We next define a HF-order $<'$ witnessing $Ac'_{n+1} \leq_o \text{Acl}_{<}^{n+1}(c'_{n+1})$. It suffices to start from Ac'_{n+1} and let $c'_{n+1} <' p <' d_n^1 <' d_n^0$ (by the choice of our elements this is an HF-order). Notice then that c'_{n+1} is incident only to p in $\text{cl}_{<}^{n+1}(c'_{n+1})$, p is incident only to d_n^0, d_n^1 in $\text{cl}_{<}^{n+1}(c'_{n+1})$, and also by our choice of d_n^1 no element in $\text{cl}_{<}^n(d_n^1)$ is incident to elements in $\text{cl}_{<}^n(d_n^0)$. Thus we can use the inductive hypothesis to continue the construction of an HF-order and obtain that $Ac'_{n+1} \leq_o \text{Acl}_{<}^{n+1}(c'_{n+1})$.

Case 2. c is a point.

We proceed analogously to the previous case. By inductive hypothesis we can find a line $d_n^0 \in P_0$ such that $A < \text{cl}_{<}^{n+1}(d_n^0)$ and $Ad_n^0 \leq_o \text{Acl}_{<}^n(d_n^0)$. Also, we can find a further line $d_n^1 \in P_1$ such that $\text{Acl}_{<}^{n+1}(d_n^0) < \text{cl}_{<}^{n+1}(d_n^1)$ and $\text{Acl}_{<}^{n+1}(d_n^0)d_n^1 \leq_o \text{Acl}_{<}^{n+1}(d_n^0)\text{cl}_{<}^n(d_n^1)$. Let c'_{n+1} be the unique point incident to both d_n^0 and d_n^1 , then since $\text{Acl}_{<}^{n+1}(d_n^0) < \text{cl}_{<}^{n+1}(d_n^1)$ it follows that $c'_{n+1} > d_n^1 > d_n^0$. Clearly we have that $\text{cl}_{<}^{n+1}(c'_{n+1}) = \text{cl}_{<}^n(d_n^0) \cup \text{cl}_{<}^n(d_n^1) \cup \{c'_{n+1}\}$. From this fact and the induction hypothesis, we can reason as in Case 1 and see that $Ac'_{n+1} \leq_o \text{Acl}_{<}^n(c'_{n+1})$. The facts that $A < \text{cl}_{<}^{n+2}(c'_{n+1})$ and $Ac \cong_A Ac'_{n+1}$ are immediately true by construction and the inductive hypothesis. \blacksquare

Proposition 4.52 (3.30(D2)). If $M \models T^+$ is \aleph_1 -saturated, $A \leq_o M$ is countable, and for every trivial extension $A \leq_o Ab$ there is $b' \in M$ s.t. $Ab \cong_A Ab' \leq_o M$, then for every minimal extension $A \leq_o Ac$ there is $c' \in M$ s.t. $Ac \cong_A Ac' \leq_o M$.

Proof. This follows by reasoning exactly as in the case of Steiner systems 4.23. \blacksquare

As in the previous cases, it already follows from the previous propositions that the theory T^+ of open k -nets is complete. To show that also 3.28(C2) holds, and thus establish the stability of T^+ , we consider the free completion process for the setting of k -nets. We thus also verify that 3.65(C8) is verified. It is straightforward to verify that the construction given by 3.39 is equivalent to the following one.

Definition 4.53. Let A be a partial k -net. We let $A_0 = A$ and for every $n < \omega$ we define A_{2n+1}, A_{2n+2} as follows:

- (a) for every pair of lines $\ell_0 \in P_i$ and $\ell_1 \in P_j$ in A_n with $i < j < k$ and such that they are not incident to any common point, we add a new point p in A_{2n+1} incident only to ℓ_0, ℓ_1 ;
- (b) for every point $p \in A_{2n+1}$ and for every $i < k$ such that p is not incident to any line of parallelism type i in A_{2n+1} , we add a new line $\ell \in A_{2n+2}$ of parallelism type i which is incident only to p .

The structure $F(A) := \bigcup_{i < \omega} A_i$ is called the *free completion* of A and we say that $F(A)$ is freely generated over A . We say that A is *non-degenerate* if $F(A)$ is infinite, and *degenerate* otherwise.

Remark 4.54. We notice that, as remarked in [10, p. 13], the only degenerate partial k -nets are the empty set \emptyset , the configuration of a point incident with k -many lines, and any set of lines of the same parallelism type. Then letting $m_{\mathcal{K}} = 2$, it follows that any structure $A \models T_{\mathcal{V}}$ containing at least $m_{\mathcal{K}}$ -many elements of each sort is non-degenerate.

Proposition 4.55 (3.28(C2), 3.65(C8)). Suppose $A \models T_{\mathcal{V}}$, and A is non-degenerate, then $F(A) \models T^+$, $A \leq_o F(A)$, $|F(A)| = |A| + \aleph_0$ and:

- (a) $F(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : F(A) \cong F(B)$;
- (b) $F(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_o M$ there is an embedding $\hat{h} : F(A) \rightarrow M$ such that $\hat{h} \upharpoonright A = h$ and $\hat{h}(F(A)) \leq_o M$.

Proof. The proof proceeds analogously to the case of Steiner systems. ■

We have verified all the conditions from 3.2, 3.28 and 3.43. It remains to consider the assumptions from 3.60 and 3.65. We point out that Assumption 3.65(C7) follows immediately from 4.55, 3.65(C8) can be proved by a similar argument as in the case of Steiner systems, and 3.65(C9) can be verified by using the definition of $\delta(A)$ from [21, Def. 11]. We conclude this section by exhibiting a configuration that witnesses that Assumption 3.60 holds.

Proposition 4.56 (3.60). There is a configuration $C \in \mathcal{K}$ satisfying the conditions from 3.60.

Proof. We define a configuration C as follow. Let $C = \{c_i : 0 \leq i \leq 18\}$, where $c_0, c_1, c_8, c_9, c_{10}, c_{14}, c_{16}, c_{18}$ are points, c_2, c_5, c_{12} are lines of parallelism type P_0 , c_3, c_6, c_{13}, c_{15} are lines of parallelism type P_1 , and c_4, c_7, c_{11}, c_{17} are lines of parallelism type P_2 . Notice that then, since $k \geq 3$, this construction thus applies to all k -nets. Finally, the incidence relations between elements of C are specified by the following table.

| | c_2 | c_3 | c_4 | c_5 | c_6 | c_7 | c_{11} | c_{12} | c_{13} | c_{15} | c_{17} |
|----------|-------|-------|-------|-------|-------|-------|----------|----------|----------|----------|----------|
| c_0 | × | × | × | | | | | | | | |
| c_1 | | | | × | × | × | | | | | |
| c_8 | × | | | | × | | × | | | | |
| c_9 | | × | | | | × | | × | | | |
| c_{10} | | | × | × | | | | | × | | |
| c_{14} | | | | | | | × | × | | × | |
| c_{16} | × | | | | | | | | | × | × |
| c_{18} | | | | | | | | | × | | × |

Then C is a partial k -net and the order $c_0 < c_1 < \dots < c_{18}$ is an HF-order, which means that $C \in \mathcal{K}$. As the other requirements from 3.60 are easily verified, it follows that C is as desired. \blacksquare

4.5. Affine planes

We apply in this section our general framework to the theory of affine planes, which essentially axiomatise the incidence properties of points and lines in the Euclidean plane (see e.g., [15] for a reference to affine planes). Although this case is very similar to the setting of projective planes, affine planes are particularly interesting as they also involve a notion of parallelism between lines, in addition to the relation of incidence between lines (or blocks) and points. Essentially, this is the first case that we are considering where we need the full generality of Definition 2.8, i.e., where we may have that $\text{gcl}_M(A) \neq \text{gcl}_M^1(A)$. We provide details of all proofs where the presence of the relation of parallelism makes a difference. Analogously to the previous cases, the results from this section then establish the following theorem.

Theorem 4.57. *Let (\mathcal{K}, \leq_o) be the class of finite, open, partial affine planes and let T^+ be defined as in 3.22. Then T^+ is complete, stable, and with $\downarrow = \downarrow^\otimes$. Moreover, T^+ is not superstable, it does not have a prime model, it is not model complete and does not eliminate quantifiers.*

We start by recalling the axiomatisation of affine planes from [21, p. 744] and by introducing the class (\mathcal{K}, \leq_o) in this setting.

Definition 4.58. Let L be the two-sorted language where the symbols p_0, p_1, \dots denote *points* and the symbols ℓ_0, ℓ_1, \dots denote *lines*, the symbol $p | \ell$ means that the point p is *incident* with the line ℓ , and the symbol $\ell_0 \parallel \ell_1$ means the two lines ℓ_0 and ℓ_1 are *parallel*. The theory of *affine planes* consists of the following axioms:

- (1) the parallelism relation \parallel is an equivalence relation;
- (2) every two distinct points p_0, p_1 have a unique common incident line $p_0 \vee p_1$;
- (3) every two non-parallel lines ℓ_0, ℓ_1 have a unique common incident point $\ell_0 \wedge \ell_1$;
- (4) given a point p and a line ℓ , there is a unique line ℓ' s.t. $p | \ell'$ and $\ell' \parallel \ell$.

The universal fragment of the theory of affine planes is axiomatised by the axioms (1) saying that parallelism is an equivalence relation and the uniqueness requirements from axioms (2), (3) and (4), i.e., the formulas saying that the line incident to two given points, the point incident to two non-parallel lines, and the line incident to a point and parallel to a line are all unique. We refer to models of this universal theory as *partial affine planes*.

Definition 4.59. For finite partial affine planes $A \subseteq B$, we define the following.

- (1) An element $a \in A$ is hyperfree in B if it satisfies one of the following conditions:
 - (i) a is a point incident with at most two lines from B ;
 - (ii) a is a line incident with at most one point from B ;
 - (iii) a is a line incident with exactly two points from B and not parallel to any other line in B .
- (2) We say that B is *open over* A and write $A \leq_o B$ if every non-empty $C \subseteq B \setminus A$ contains an element which is hyperfree in AC . We say that B is *open* if $\emptyset \leq_o B$, i.e., if every non-empty $C \subseteq B$ contains some element hyperfree in C .
- (3) We say that B is *closed over* A if $A \not\leq_o B$, i.e., if there is a non-empty $C \subseteq B \setminus A$ which does not contain any element hyperfree in AC . We say that B is *closed* if $\emptyset \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B$ which does not contain any element hyperfree in C .

We let \mathcal{K} be the set of open partial finite affine planes and we let T_{\forall} be the universal theory of \mathcal{K} . The previous definitions extend to infinite models of T_{\forall} as in 3.6. The notion of hyperfree element in partial affine planes is from [21, p. 752].

Remark 4.60. As the examples from the previous sections only involved a relation of incidence, it was always clear that the property of being hyperfree was preserved by taking subsets. We point out that this holds also in the setting of affine planes. Suppose that c is hyperfree in Xc and $Y \subseteq X$. If c is of type 4.59(2)(i) or of type 4.59(2)(ii), then this is clear (as we have the ‘‘at most’’). So suppose that c is of type 4.59(2)(iii). If Y contains the two lines which witness that c is of type 4.59(2)(iii), then c is of type 4.59(2)(iii) also in Yc . If Y does not contain at least one of the two lines, then it is of type 4.59(2)(ii) in Yc .

As in the previous cases, we start by recalling what are algebraic strong extensions in the setting of affine planes, and by proving the crucial Lemma 4.62. We provide details of its proof as this is the first case where parallelism relations play a role. Moreover, we also notice that this establishes Assumption 3.43(C5).

Remark 4.61. In partial affine planes an extension $A \leq_o Ac$ is algebraic if:

- (a) c is a point incident with two lines from A ;
- (b) c is a line incident with two points from A and parallel to no other line in A ;
- (c) c is a line incident with a point in A and parallel to some line in A .

Lemma 4.62 (3.43(C5)). Let $A \leq_o C \models T_{\forall}$ and let $c \in C$ be such that $A \leq_o Ac$ is an algebraic strong extension. Then $Ac \leq_o C$.

Proof. Consider a finite subset $D \subseteq C$ such that $Ac \subsetneq D$, we need to show that $D \setminus Ac$ contains some hyperfree element. Now, since $A \leq_o C$ and $D \subseteq C$, there is an element $d \in D \setminus A$ hyperfree in D . If $c \neq d$ then d also witnesses that $D \setminus Ac$ contains a hyperfree element and we are done. Thus suppose that c is the only element in $D \setminus A$ to be hyperfree in D . Since c is algebraic over A , it forms one of the configuration from Remark 4.61.

Case 1. c is a point incident with two lines from A .

Since c is hyperfree in D , it is not incident to lines in $D \setminus A$. It follows that an element $d' \in D \setminus Ac$ is hyperfree in D if and only if it is hyperfree in $D \setminus \{c\}$. Since by assumption c is the unique element in $D \setminus A$ to be hyperfree in D , it follows that $D \setminus Ac$ contains no element hyperfree in D , contradicting $A \leq_o C$.

Case 2. c is a line incident with two points from A .

Then c is neither incident to points in $D \setminus A$, nor parallel to any other line from D . Our claim then follows exactly as in Case 1.

Case 3. c is a line incident to some point in A and parallel to some line $\ell \in A$.

Since c is hyperfree in D , it is not incident to any point from $D \setminus A$. Moreover, if $d' \in D \setminus A$ is such that $d' \parallel c$ then $d' \parallel \ell$. As in the previous cases, it follows that an element $d' \in D \setminus Ac$ is hyperfree in D if and only if it is hyperfree in $D \setminus \{c\}$. Since by assumption c is the unique element in $D \setminus A$ to be hyperfree in D , it follows that $D \setminus Ac$ contains no element hyperfree in D , contradicting $A \leq_o C$. ■

We next consider the conditions from 3.2. As in the previous cases, conditions (A), (B), (D), (E) (F) and (L) follow immediately from the definition of the relation \leq_o for finite, open, partial affine planes. Condition 3.2(C) follows exactly as in the case of Steiner systems. Condition (I) follows immediately for $X_{\mathcal{K}} = \{1\}$ and Condition (J) follows by letting $n_{\mathcal{K}} = 2$. Condition (K) follows by the definition of hyperfree elements in affine planes together with the definition of Gaifman closure 2.8. It remains to verify Conditions (G) and (H). We first consider the amalgamation property and we notice that, together with the next remark, Proposition 4.65 also verifies Assumption 3.43(C4).

Remark 4.63 (3.43(C4)). Let $A \subseteq B, C$ be open partial affine planes with $B \cap C = A$ and recall the definition of free amalgam from 3.39. As in the previous cases, the free amalgam $B \otimes_A C$ is not necessarily an open partial affine planes. However, it follows from the proof of Proposition 4.65 that if the conditions from 3.43(C4) are satisfied then the free amalgam $B \otimes_A C$ is always in \mathcal{K} .

Definition 4.64. Let A be a partial affine plane and $\ell \in A$ be a line. We say that ℓ forms a *trivial parallelism class* in A if $\ell \parallel \ell', \ell' \in A$ entail that $\ell = \ell'$. Also, we denote by $P(A)$ any subset of A containing exactly one line for every parallelism class, i.e., for every $\ell \in A$ there is exactly one $\ell' \in P(A)$ such that $\ell \parallel \ell'$.

Proposition 4.65 (3.2(G)). The class (\mathcal{K}, \leq_o) has both the amalgamation and joint embedding property.

Proof. We proceed as in the previous cases and simply verify the amalgamation property. Consider $A, B, C \in \mathcal{K}$ with $A \leq_o B, A \leq_o C, B \cap C = A$. Notice that if $B \otimes_A C \in \mathcal{K}$, then it follows immediately from the definition of hyperfree elements in affine planes that $B \leq_o B \otimes_A C$ and $C \leq_o B \otimes_A C$. We assume by induction that $B \setminus A$ consists of a single element and we show that either there is an embedding $f : B \rightarrow C$ such that $f(B) \leq_o C$ or $B \otimes_A C \in \mathcal{K}$.

Case 1. $B = A \cup \{\ell\}$ where ℓ is a line incident with at most one point from A .

If ℓ is not incident with any point from A , or if it forms a trivial parallelism class in B , then $B \otimes_A C \in \mathcal{K}$. So suppose that $p \in \ell$ and $\ell \parallel \ell_0$ for some $p, \ell_0 \in A$.

Case 1.1. C does not contain a line parallel with ℓ_0 and incident with p .

Then we immediately have $B \otimes_A C \in \mathcal{K}$.

Case 1.2. C contains a line ℓ' parallel with ℓ_0 and incident to p .

Then since $A \leq_o C$, it follows in particular that ℓ' is incident only with p and so the map f defined by letting $f|_A = \text{id}_A$ and $f(\ell) = \ell'$ is an embedding. Moreover, since $A \leq_o A\ell'$ is algebraic, it follows from 4.62 that $f(B) \leq_o C$.

Case 2. $B = A \cup \{\ell\}$ where ℓ is a line incident with exactly two points p_0, p_1 from A and parallel to no line in A .

Case 2.1. C does not contain a line incident with p_0 and p_1 .

In this case we immediately have $B \otimes_A C \in \mathcal{K}$.

Case 2.2. C contains a line ℓ' incident with p_0 and p_1 .

Since $A \leq_o C$ it follows that ℓ' is incident only with p_0, p_1 and parallel with no line in A . It follows that the map f such that $f|_A = \text{id}_A$ and $f(\ell) = \ell'$ is an embedding. Moreover, since ℓ' is algebraic over A , it follows by 4.62 that $f(B) \leq_o C$.

Case 3. $B = A \cup \{p\}$ where p is a point incident with at most two lines from A .

If p is incident with at most one line, then $B \otimes_A C \in \mathcal{K}$, so we suppose p is incident to exactly two lines $\ell_0, \ell_1 \in A$.

Case 3.1. C does not contain a point incident with ℓ_0 and ℓ_1 .

In this case we immediately have $B \otimes_A C \in \mathcal{K}$.

Case 3.2. C contains a point p' incident with ℓ_0 and ℓ_1 .

Since $A \leq_o C$ it follows that p' is incident only to such lines in A . Then the map f such that $f|_A = \text{id}_A$ and $f(p) = p'$ is an embedding of B into C . Moreover, since $A \leq_o Ap'$ is algebraic, it follows from 4.62 that $f(B) \leq_o C$. ■

Next, we give details of the proof of 3.2(H), since this requires to take parallelism relations into account.

Proposition 4.66 (3.2(H)). If $A \subseteq B \in \mathcal{K}$ and $A \not\leq_o B$, then there is $n < \omega$ such that every $C \in \mathcal{K}$ contains at most n copies of B over A .

Proof. Suppose $A \not\leq_o B$, then by Definition 4.59 there is a subset $B' \subseteq B \setminus A$ such that every element in B' is not hyperfree in AB' . Without loss of generality we let $B' = B \setminus A$. Then every point $p \in B \setminus A$ is incident with at least three lines from B and every line $\ell \in B \setminus A$ is incident with at least three points from B , or it is incident with two points from B while being parallel to some other line from B . For every line $\ell \in B$ we let P_ℓ be the set of points in A that are incident with ℓ , and for every point $p \in B$ we let L_p be the set of lines in A that are incident with p . Suppose $C \in \mathcal{K}$ contains at least three copies $(B_i)_{i < 3}$ of B isomorphic over A , then we claim that the set

$$D = \bigcup_{p \in B_0} L_p \cup \bigcup_{\ell \in B_0} P_\ell \cup \bigcup_{i < 3} (B_i \setminus A)$$

witnesses that C is not open. We need to consider the following elements.

Case 1. $\ell \in L_{p_0}$ for some $p_0 \in B_0$.

By definition $p_0 | \ell$ where $p_0 \in B_0$. Since B_0, B_1, B_2 are all isomorphic over A , there are $p_1 \in B_1$ and $p_2 \in B_2$ such that $p_1 | \ell$ and $p_2 | \ell$. Thus ℓ has three incidences in D and it is not hyperfree.

Case 2. $p \in P_{\ell_0}$ for some $\ell_0 \in B_0$.

By reasoning exactly as in Case 1 it follows that p is incident to three lines in D and thus it is not hyperfree.

Case 3. $p \in B_i \setminus A$ is a point, for $i < 3$.

By assumption p is incident to at least three lines in B_i , and thus in $L_p \cup (B_i \setminus A) \subseteq D$. Thus p is not hyperfree in D .

Case 4. $\ell \in B_i \setminus A$ is a line, for $i < 3$.

Since ℓ is not hyperfree in B_i we have three possible subcases.

Case 4.1. ℓ is incident to three points in B_i .

Then, by definition of the set P_ℓ , it follows that ℓ is incident to three elements in $P_\ell \cup (B_i \setminus A) \subseteq D$. It follows that ℓ is not hyperfree in D .

Case 4.2. ℓ is incident to two points in B_i , and parallel to another line $\ell' \in B_i \setminus A$. Reasoning as in case 4.1, it follows that ℓ is incident to two points also in $P_\ell \cup (B_i \setminus A) \subseteq D$. Then, since $\ell' \in (B_i \setminus A) \subseteq D$, we obtain that ℓ is incident to two points in D while being parallel to some line $\ell' \in D$, showing that ℓ is not hyperfree in D .

Case 4.3. ℓ is incident to two points in B_i while being parallel to some line $\ell' \in A$. Assume w.l.o.g. that $i = 0$. Reasoning as in case 4.1, it follows that ℓ is incident to two points also in $P_\ell \cup (B_0 \setminus A) \subseteq D$. Now, since B_0 and B_1 are isomorphic over A , it follows that there is a line ℓ'' such that $\ell \parallel \ell'$ and $\ell'' \parallel \ell'$. By transitivity of the parallelism relation it follows that $\ell \parallel \ell'' \in (B_1 \setminus A) \subseteq D$. It follows that ℓ is not hyperfree in D . This completes our proof. \blacksquare

We next consider the assumptions from 3.28. As in the previous cases 3.28(C1) follows from 3.30(D1) and 3.30(D2). We add a condition to the statement of 3.30(D1) to ease our inductive argument.

Proposition 4.67 (3.30(D1)). If $M \models T^+$, $A \subseteq M$ is finite, $<$ is an HF-order of M and $A \leq_o Ac \in \mathcal{K}$ is a trivial extension, then for all $n < \omega$ there is $c'_n \in M$ s.t.:

- (a) $Ac \cong_A Ac'$;
- (b) $Ac' \leq_o \text{Acl}_<^n(c')$;
- (c) $A < \text{cl}_<^{n+1}(c')$.

Proof. By induction on $n < \omega$ we prove that for every $A \leq_o M$ and $A \leq_o Ac \in \mathcal{K}$ we can find c'_n as in (a)-(c). If $n = 0$, then it is easy to see that we can find c'_0 as wanted. So suppose the inductive hypothesis and let c be such that $A \leq_o Ac \in \mathcal{K}$ is a trivial extension, we want to define c'_{n+1} so that (a)-(c) hold. We assume without loss of generality that c is a line, since the case for points is analogous.

By inductive hypothesis we can a point d_n^0 which satisfies (a)-(c) with respect to A . Additionally, we can apply the induction hypothesis again to the set $\text{Acl}_<^n(d_n^0)$ and obtain a second point d_n^1 satisfying (a)-(c) with respect to the set $\text{Acl}_<^n(d_n^0)$, whence in particular $\text{Acl}_<^n(d_n^0) < \text{cl}_<^{n+1}(d_n^1)$ and $\text{Acl}_<^n(d_n^0)d_n^1 \leq_o \text{Acl}_<^n(d_n^0)\text{cl}_<^n(d_n^1)$. We let $c'_{n+1} := d_n^0 \vee d_n^1$ and we claim it satisfies conditions (a)-(c) for the value $n + 1$.

Firstly, notice that since $\text{Acl}_<^n(d_n^0) < \text{cl}_<^{n+1}(d_n^1)$ it follows immediately that $d_n^0 < d_n^1 < c'_{n+1}$. Then, it follows from the fact that $<$ is a HF-order of M that the only elements incident to c'_{n+1} occurring before it in the ordering $<$ are d_n^0 and d_n^1 . By construction it follows immediately that $Ac \cong_A Ac'_{n+1}$. Additionally, we have:

$$\begin{aligned} \text{cl}_<^{n+2}(c'_{n+1}) &= \{c'_{n+1}\} \cup \text{cl}_<^{n+1}(d_n^0) \cup \text{cl}_<^{n+1}(d_n^1), \\ \text{cl}_<^{n+1}(c'_{n+1}) &= \{c'_{n+1}\} \cup \text{cl}_<^n(d_n^0) \cup \text{cl}_<^n(d_n^1). \end{aligned}$$

Then, it follows immediately from the induction hypothesis and the choice of d_n^0, d_n^1 that $A < \text{cl}_<^{n+2}(c'_{n+1})$. Finally, we define a HF-order $<'$ witnessing $Ac'_{n+1} \leq_o \text{Acl}_<^n(c'_{n+1})$. It suffices to start from Ac'_{n+1} and let $c'_{n+1} <' d_n^1 <' d_n^0$ (by the choice of our elements this is an HF-order). As remarked above, c'_{n+1} is incident only to d_n^0 and d_n^1 in $\text{cl}_<^{n+1}(c'_{n+1})$, and moreover by our choice of d_n^1 no element in $\text{cl}_<^n(d_n^1)$ is incident to elements in $\text{cl}_<^n(d_n^0)$. Thus, we can use the inductive hypothesis that $Ad_n^0 \leq_o \text{Acl}_<^n(d_n^0)$ and $\text{Acl}_<^n(d_n^0)d_n^1 \leq_o \text{cl}_<^n(d_n^0)\text{cl}_<^n(d_n^1)$ to continue the construction of an HF-order and obtain that $Ac'_{n+1} \leq_o \text{Acl}_<^{n+1}(c'_{n+1})$. This completes our proof. \blacksquare

Proposition 4.68 (3.30(D2)). If $M \models T^+$ is \aleph_1 -saturated, $A \leq_o M$ is countable, and for every trivial extension $A \leq_o Ab$ there is an element $b' \in M$ such that $Ab \cong_A Ab' \leq_o M$, then for every minimal extension $A \leq_o Ac$ there is an element $c' \in M$ such that $Ac \cong_A Ac' \leq_o M$.

Proof. The proof is analogous to the case of Steiner systems (Prop. 4.23). ■

We next need to verify Assumption 3.28(C2), which we do as in the previous cases by simultaneously showing 3.65(C8). We recall the free completion process for affine planes from [21, p. 754], and we point out that this is essentially equivalent to the one given by Definition 3.39. Recall that, after 4.64, given a partial affine plane A we denote by $P(A)$ any set containing exactly one representative for every parallelism class, i.e., for every $\ell \in A$ there is one $\ell' \in P(A)$ such that $\ell \parallel \ell'$.

Definition 4.69 (Free completion). Let A be a partial affine plane. We let $A_0 = A$ and for every $n < \omega$ we define A_{3n+1} , A_{3n+2} and A_{3n+3} as follows.

- (a) First, for every pair of distinct points $p_0, p_1 \in A_{3n}$ not incident to a common line in A_{3n} , we add a new line ℓ incident only to p_0, p_1 and not parallel to any other line from A_{3n} . We denote the resulting partial plane by A_{3n+1} .
- (b) Secondly, for every pair of lines $\ell_0, \ell_1 \in A_{3n+1}$ which are not incident to any common point in A_{3n+1} and which are not parallel, we add a new point p incident only to ℓ_0 and ℓ_1 . We denote the resulting partial plane by A_{3n+2} .
- (c) Thirdly, for every line $\ell \in P(A_{3n+2})$ and every point $p \in A_{3n+2}$ such that A_{3n+2} contains no line parallel to ℓ and incident to p , we add a new line satisfying these two conditions. We denote the resulting partial plane by A_{3n+3} .

The structure $F(A) := \bigcup_{n < \omega} A_n$ is called the *free completion* of A and we say that $F(A)$ is freely generated over A . We say that A is *non-degenerate* if $F(A)$ is infinite, and *degenerate* otherwise.

Remark 4.70. We notice that, as in the projective case, it is enough that A contains four non-collinear points in order for it to be non-degenerate. Let $m_\kappa = 5$, then in particular any partial affine planes A which contains at least m_κ -many elements of both sorts is non-degenerate. Thus the following proposition establishes 3.28(C2). We refer the reader to [30, p. 29] for a precise characterisation of non-degenerate partial affine planes.

Proposition 4.71 (3.28(C2), 3.65(C8)). Suppose $A \models T_\forall$, and A is non-degenerate, then $F(A) \models T^+$, $A \leq_o F(A)$, $|F(A)| = |A| + \aleph_0$ and:

- (a) $F(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : F(A) \cong F(B)$;
- (b) $F(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_o M$ there is an embedding $\hat{h} : F(A) \rightarrow M$ such that $\hat{h} \upharpoonright A = h$ and $\hat{h}(F(A)) \leq_o M$.

Proof. The proof proceeds exactly as in the case of Steiner systems. ■

Finally, we have verified all the conditions from 3.2, 3.28 and 3.43. For what concerns the assumptions from 3.60 and 3.65 we simply notice the following. The configuration from [24, 9.1] does work (with minor modifications) also in the present setting of affine planes, thus we do not provide the full details of the construction.

Similarly, Assumption 3.65(C7) follows from 4.71, 3.65(C8) can be proved by reasoning as in the case of Steiner systems, and 3.65(C9) can be verified by using the definition of $\delta(A)$ from [21, Def. 11].

4.6. Projective Möbius, Laguerre and Minkowski planes

We conclude this article by providing a last application of our framework to the setting of Benz planes. Differently from the previous examples, Benz planes are not a single instance of incidence geometry but rather a family of theories closely related to each other: Möbius planes, Laguerre planes and Minkowski planes. These were originally introduced by Walter Benz in [13] and they axiomatise the structure induced by a 3-dimensional projective space on one of its quadratic sets (cf. [16, §5] and [29, p. 21]). As in the previous cases, we will study Benz planes from the synthetic point of view, essentially as a multigraph with points and blocks.

From our current perspective, the main interest of this family of theories is that, additionally to the relation of incidence (and parallelism) they also involve a notion of tangency between blocks, which is a local equivalence relation parametrised by points. Crucially, it is important to keep in mind that there are in the literature different versions of Benz planes, and it is essential in our context that we consider *projective* Benz planes. We elaborate this further in Remark 4.84. Additionally, we stress that for brevity and simplicity we restrict our treatment only to the case of Möbius planes. The key difference with Laguerre and Minkowski planes is simply that they add respectively one and two parallelism relations between points, and it is then straightforward to adapt our arguments to these settings. We refer the reader to [13] and [21] for more details on Laguerre and Minkowski planes. Finally, as in the previous cases, we obtain the following theorem.

Theorem 4.72. *Let (\mathcal{K}, \leq_o) be the class of finite, open, partial Möbius planes (resp. of Laguerre and Minkowski planes), and let T^+ be defined as in 3.22. Then T^+ is complete, stable, and with $\downarrow = \downarrow^\otimes$. Moreover, T^+ is not superstable, it does not have a prime model, it is not model complete and does not eliminate quantifiers.*

We recall the axiomatisation of projective Möbius planes from [21, p. 748].

Definition 4.73. Let L be the two-sorted language where the symbols p_0, p_1, \dots denote *points* and the symbols b_0, b_1, \dots denote *blocks*, the symbol $p | b$ means that the point p is incident to the block b , and the ternary tangency symbols $\rho(b_0, b_1, p)$ means that the two blocks b_0, b_1 are tangent in p . The theory T of *projective Möbius planes* consists of the following axioms:

- (1) every three different points are incident to a unique block;
- (2) any two blocks tangent to the same point are incident to it, i.e.,

$$\forall p \forall b_0 \forall b_1 (\rho(b_0, b_1, p) \rightarrow (p_0 | b_0 \wedge p_0 | b_1));$$

- (3) given two points p_1, p_2 and a block b_1 such that p_1 is incident to b_1 but p_2 is not incident to b_1 , there is a unique block b_2 incident to p_2 and such that $\rho(b_1, b_2, p_1)$ (we say that this is the unique block incident to p_2 that “touches” b_1 in p_1);
- (4) for every point c , the relation induced by $\rho(x, y, c)$ is an equivalence relation;
- (5) every two different blocks are either tangent in one common point, or they are incident to two common points.

The universal fragment of the theory of projective Möbius planes is axiomatised exactly by the axioms (2) and (4), by the axiom stating that two different blocks are incident with at most two common points, and by the uniqueness requirements from axiom (1) and (3) stating that the block incident to three given points and the touching block are unique. We refer to models of this universal theory as *partial projective Möbius planes*.

Before defining what open Möbius planes are, we introduce the following notion of valency in Möbius planes. This should not be confused with the predimension function δ , although as one can then see from [21, Def. 11], they are closely related.

Definition 4.74. Let $B \in \mathcal{K}$. Given a block $b \in B$, we let $T(b/B)$ be a maximal sets of blocks $b' \in B$ such that $B \models \rho(b, b', p)$ for some $p \in B$ and every $b', b'' \in B$ are not tangent to each other in B . Given an element $c \in B$ we write $I(c/B)$ for the set of elements incident to c in B . We then define the *valency* of an element $c \in B$ as the number $\vartheta(c/B)$ given by

$$\vartheta(c/B) := |I(c/B)| + |T(c/B)|.$$

Notice that, by Def. 2.8, we immediately have that $\vartheta(c/B) = |\text{gcl}_B(c)|$.

Definition 4.75. For finite partial projective Möbius planes $A \subseteq B$, we define the following.

- (1) An element $a \in A$ is hyperfree in B if
 - (i) a is a point with valency $\vartheta(a/B) \leq 2$;
 - (ii) a is a block with valency $\vartheta(a/B) \leq 3$.
- (2) We say that B is *open over* A and write $A \leq_o B$ if every non-empty $C \subseteq B \setminus A$ contains an element which is hyperfree in AC . We say that B is *open* if $\emptyset \leq_o B$, i.e., if every non-empty $C \subseteq B$ contains some element hyperfree in C .
- (3) We say that B is *closed over* A if $A \not\leq_o B$, i.e., if there is a non-empty $C \subseteq B \setminus A$ which does not contain any element hyperfree in AC . We say that B is *closed* if $\emptyset \not\leq_o B$, i.e., if there is a non-empty configuration $C \subseteq B$ which does not contain any element hyperfree in C .

We let \mathcal{K} be the set of open partial projective Möbius planes and we let T_{\forall} be the universal theory of \mathcal{K} . The previous definitions extend to infinite models of T_{\forall} as in 3.6. The notion of hyperfree elements in partial projective Möbius planes is from [21, p. 753].

Remark 4.76. We notice that, in particular, an extension $A \leq_o Ac$ is algebraic if c is a point with valency 2, or if c is a block with valency 3, i.e., if

- (a) c is a point incident with two blocks from A ;
- (b) c is a block incident to three points from A ;
- (c) c is a block incident to two points from A and tangent to a block from A .

We proceed as in the previous cases and we first prove the following key lemma concerning strong algebraic extensions. We also notice that this establishes Assumption 3.43(C5).

Lemma 4.77 (3.43(C5)). Let $A \leq_o C \models T_{\forall}$ and let $c \in C$ be such that $A \leq_o Ac$ is an algebraic strong extension. Then $Ac \leq_o C$.

Proof. The proof follows analogously to the case of affine planes. ■

We now consider the conditions from 3.2. As before, conditions (A), (B), (D), (E), (F), (K) and (L) follow immediately from our definition of the relation \leq_o for finite, open, partial Möbius planes. Condition 3.2(C) follows exactly as in the case of Steiner systems. Condition (I) follow immediately by letting $X_{\mathcal{K}} = \{1\}$ and Condition (J) follows by letting $n_{\mathcal{K}} = 3$. Then, Condition 3.2(G) can be proved similarly to the case of affine planes, and it also verifies 3.43(C4). To showcase how to work with the relation of tangency, and also make it explicit that it is essentially analogous to the parallelism relation of projective planes, we include the proof of Condition 3.2(H).

Proposition 4.78 (3.2(H)). If $A \subseteq B \in \mathcal{K}$ and $A \not\leq_o B$, then there is $n < \omega$ such that every $C \in \mathcal{K}$ contains at most n copies of B over A .

Proof. Suppose $A \not\leq_o B$, then w.l.o.g. we have that every element in $B \setminus A$ is not hyperfree. In particular, for every point $p \in B$ we have $\vartheta(p/B) \geq 3$ and for every block in $b \in B$ we have that $\vartheta(b/B) \geq 4$. For every element $c \in B \setminus A$ we write $I(c/A)$ for the set of elements in A that are incident with c . Suppose $C \in \mathcal{K}$ contains at least four copies $(B_i)_{i < 4}$ of B over A , then we claim that the set

$$D = \bigcup_{c \in B_0 \setminus A} I(c/A) \cup \bigcup_{i < 4} (B_i \setminus A)$$

witnesses that C is not open. We consider the following elements.

Case 1. $d \in I(c/A)$ for some $c \in B \setminus A$.

Then, from the fact that c is incident to d and that B_0, B_1, B_2, B_3 are all isomorphic over A , it follows that $\vartheta(d/D) \geq 4$, showing that d is not hyperfree in D .

Case 2. $c \in B_i \setminus A$ is a point, for $i \leq 3$.

By assumption $\vartheta(c/B_i) \geq 3$. Then, by Definition 4.74 and the definition of D , it follows immediately that $\vartheta(c/D) \geq 3$, whence c is not hyperfree in D .

Case 3. $c \in B_i \setminus A$ is a block, for $i \leq 3$.

We let w.l.o.g. $i = 0$, then by assumption $\vartheta(c/B_0) \geq 4$. We claim that $\vartheta(c/D) \geq 4$. Firstly, notice that every point in B_0 incident to c belongs to $I(c/B_0) \cup (B_0 \setminus A) \subseteq D$, thus $|I(c/B_0)| = |I(c/D)|$. Moreover, consider a block $d \in B_0$ such that $B_0 \models \rho(c, d, p)$ for some point $p \in B_0$. If $p \in B_0 \setminus A$ then we have that $d \in D$, since clearly p is incident with d . If $p \in A$, then there is a block $c' \in B_1 \setminus A$ such that $B_1 \models \rho(c, c', p)$ and $B_1 \models \rho(c', d, p)$. Therefore, D contains a representative of all tangency classes parametrised by points in B_0 . By Definition 4.74 it follows that $\vartheta(c/D) \geq 4$ and so c is not hyperfree in D . This completes our proof. ■

We next consider the assumptions from 3.28. As in the previous cases we establish 3.28(C1) by proving 3.30(D1) and 3.30(D2). We add an extra condition to the statement of the next proposition to ease our inductive argument.

Proposition 4.79 (3.30(D1)). If $M \models T^+$, $A \subseteq M$ is finite, $<$ is an HF-order of M and $A \leq_o Ac \in \mathcal{K}$ is a trivial extension, then for $n < \omega$ there is $c'_n \in M$ s.t.:

- (a) $Ac \cong_A Ac'$;
- (b) $Ac' \leq_o \text{Acl}_{<}^n(c')$;
- (c) $A < \text{cl}_{<}^{n+1}(c')$.

Proof. By induction on $n < \omega$ we prove that for every $A \leq_o M$ and trivial extension $A \leq_o Ac$ we can find c'_n as in (a)-(c). If $n = 0$, then it is easy to see that we can

find c'_0 as wanted. So suppose the inductive hypothesis holds and let c be such that $A \leq_o Ac$ is a trivial extension, we find c'_{n+1} so that (a)-(c) hold.

Case 1. c is a block.

By inductive hypothesis we can find three points p_n^0, p_n^1, p_n^2 such that (a)-(c) hold respectively for A , $\text{Acl}_{<}^n(p_n^0)$ and $\text{Acl}_{<}^n(p_n^0)\text{cl}_{<}^n(p_n^1)$. Let c'_{n+1} be the unique block incident to p_n^0, p_n^1, p_n^2 . By the assumption (c) and the choice of the elements it follows in particular that $c'_{n+1} > p_n^2 > p_n^1 > p_n^0$. We have:

$$\begin{aligned} \text{cl}_{<}^{n+2}(c'_{n+1}) &= \{c'_{n+1}\} \cup \text{cl}_{<}^{n+1}(p_n^0) \cup \text{cl}_{<}^{n+1}(p_n^1) \cup \text{cl}_{<}^{n+1}(p_n^2), \\ \text{cl}_{<}^{n+1}(c'_{n+1}) &= \{c'_{n+1}\} \cup \text{cl}_{<}^n(p_n^0) \cup \text{cl}_{<}^n(p_n^1) \cup \text{cl}_{<}^n(p_n^2). \end{aligned}$$

Hence we immediately have that $A < \text{cl}_{<}^{n+2}(c'_{n+1})$ and also $Ac \cong_A Ac'_{n+1}$. We define a HF-order $<'$ to show that $Ac'_{n+1} \leq_o \text{Acl}_{<}^{n+1}(c'_{n+1})$. We start by letting $c'_{n+1} <' p_n^2 <' p_n^1 <' p_n^0$. By the condition (b) and the choice of p_n^0, p_n^1, p_n^2 we have that $Ap_n^0 \leq_o \text{Acl}_{<}^n(p_n^0)$, $\text{Acl}_{<}^n(p_n^0)p_n^1 \leq_o \text{Acl}_{<}^n(p_n^0)\text{cl}_{<}^n(p_n^1)$ and $\text{Acl}_{<}^n(p_n^0)\text{cl}_{<}^n(p_n^1)p_n^2 \leq_o \text{Acl}_{<}^n(p_n^0)\text{cl}_{<}^n(p_n^1)\text{cl}_{<}^n(p_n^2)$. By condition (c) it follows that there is no incidence between elements of $\text{cl}_{<}^n(p_n^0)$, $\text{cl}_{<}^n(p_n^1)$ and $\text{cl}_{<}^n(p_n^2)$. Also, since $<$ is a HF-order, we have that p_n^0, p_n^1, p_n^2 are the only elements in $\text{cl}_{<}^{n+1}(c'_{n+1})$ to which c'_{n+1} is incident. It follows that we can extend the ordering $<'$ using the induction hypothesis and show that $Ac'_{n+1} \leq_o \text{Acl}_{<}^n(c'_{n+1})$.

Case 2. c is a point.

Similarly to Case 1, we use the inductive hypothesis to find two blocks b_n^0 and b_n^1 such that they satisfy conditions (a)-(c) with respect to A and $\text{Acl}_{<}^n(b_n^0)$. Then we let c'_{n+1} be the unique block incident to b_n^0 and b_n^1 . Then by reasoning as in Case 1 one can verify that conditions (a)-(c) are satisfied. ■

Proposition 4.80 (3.30(D2)). If $M \models T^+$ is \aleph_1 -saturated, $A \leq_o M$ is countable, and for every trivial extension $A \leq_o Ab$ there is $b' \in M$ s.t. $Ab \cong_A Ab' \leq_o M$, then for every minimal extension $A \leq_o Ac$ there is $c' \in M$ s.t. $Ac \cong_A Ac' \leq_o M$.

Proof. This is proved by Lemma 4.77 reasoning as in the previous cases. ■

We next verify Assumption 3.28(C2) for $C(A) = F(A)$, thus establishing also 3.65(C8). We recall the free completion process for projective Möbius planes from [21, p. 755-756] and we leave to the reader to verify that this is equivalent to the construction given by 3.41.

Definition 4.81. Let A be a partial Möbius planes. We let $A_0 = A$ and for every $n < \omega$ we define $A_{4n+1}, A_{4n+2}, A_{4n+3}, A_{4n+4}$ as follows.

- (a) First, for every triple of distinct points $p_0, p_1, p_2 \in A_{4n}$ which are not incident to any common block, we add a new block b incident only to p_0, p_1, p_2 and not tangent any other block from A_{4n} . We denote the resulting partial plane by A_{4n+1} .
- (b) Secondly, for every block $b \in A_{4n+1}$, every point p incident to b and every point q not incident to it, we add a new block b' incident only to p, q in A_{4n+1} , and tangent to b exactly in p . We denote the resulting partial plane by A_{4n+2} .
- (c) Thirdly, for every two non-tangent blocks $b_0, b_1 \in A_{4n+2}$ incident to only one common point $p_0 \in A_{4n+2}$, we add a new point p_1 incident to both b_0, b_1 . We denote the resulting partial plane by A_{4n+3} .

- (d) Finally, for every two non-tangent blocks $b_0, b_1 \in A_{4n+2}$ incident to no common point in A_{4n+3} , we add two new points p_0, p_1 incident to both b_0, b_1 . We denote the resulting partial plane by A_{4n+4} .

The structure $F(A) := \bigcup_{n < \omega} A_n$ is called the free completion of A and we say that $F(A)$ is freely generated over A . We say that A is *non-degenerate* if $F(A)$ is infinite, and *degenerate* otherwise.

Remark 4.82. A partial projective Möbius plane is non-degenerate if it contains at least one block, two points incident to it, and one point not incident to it [21, p. 756]. It follows in particular that any set A containing at least $m_{\mathcal{K}} = 4$ elements of every sort is non-degenerate. Thus the following proposition also establishes 3.28(C2).

Proposition 4.83 (3.28(C2), 3.65(C8)). Suppose $A \models T_{\forall}$, and A is non-degenerate, then $F(A) \models T^+$, $A \leq_o F(A)$, $|F(A)| = |A| + \aleph_0$ and:

- (a) $F(A)$ is unique up to A -isomorphisms, i.e., every isomorphism $h : A \cong B$ extends to an isomorphism $\hat{h} : F(A) \cong F(B)$;
- (b) $F(A)$ is prime over A , i.e., for every $M \models T^+$ and every embedding $h : A \rightarrow M$ with $h(A) \leq_o M$ there is an embedding $\hat{h} : F(A) \rightarrow M$ such that $\hat{h}|_A = h$ and $\hat{h}(F(A)) \leq_o M$.

Proof. The proof proceeds exactly as in the case of Steiner systems. ■

Remark 4.84. The definition of free completion process in Möbius plane makes it clear why the projectivity requirement is essential in the context of Benz planes. In fact, if we drop Clause (d) from Definition 4.81, what we obtain is the free completion process for arbitrary Möbius planes (cf. [21, p. 755-756]). Now, let A be a partial Möbius plane containing two blocks b_0, b_1 and three points p_0, p_1, p_2 such that p_0, p_1 are incident only to b_0 , and p_2 is not incident with neither b_0 nor b_1 . Let $F^a(A)$ be the free affine completion of A , i.e., the model obtained by closing A under all the clauses from Definition 4.81 with the exception of Clause (d). Then $F^a(A)$ satisfies all the axioms of the theory of Möbius with the exception of the projectivity requirement, since the blocks b_0 and b_1 have no common incidence point in $F^a(A)$. At the same time, however, it is straightforward to verify that $F^a(A)$ is open. It follows that $F^a(A)$ and $F(A)$ are two non elementary equivalent models of the theory of open Möbius planes without the projectivity axiom. This shows that the requirement of projectivity is essential to obtain a complete theory T^+ . Notice that this is reflected in the proof of Assumption 3.30(D1), since Case 2 of the proof crucially requires that M is projective.

We thus have verified all the conditions from 3.2, 3.28 and 3.43. It remains to consider 3.60 and 3.65. We notice that 3.65(C7) follows from 4.83, 3.65(C8) is verified by essentially the same argument as in the case of Steiner systems, and 3.65(C9) can be verified by using the definition of $\delta(A)$ from [21, Def. 11]. The next proposition exhibits a configuration which satisfies the conditions from Assumption 3.60. Together with the previous results, this establishes Theorem 4.72.

Proposition 4.85 (3.60). There is a configuration $C \in \mathcal{K}$ satisfying the conditions from 3.60.

Proof. We define a configuration C as follow. Let $C = \{c_i : 0 \leq i \leq 16\}$, where $c_0, c_1, c_2, c_3, c_8, c_9, c_{10}, c_{12}, c_{14}, c_{16}$ are points, $c_4, c_5, c_6, c_7, c_{11}, c_{13}, c_{15}$ are blocks, and the incidence relations between elements of C is specified by the following table.

| | c_4 | c_5 | c_6 | c_7 | c_{11} | c_{13} | c_{15} |
|----------|-------|-------|-------|-------|----------|----------|----------|
| c_0 | × | | × | × | | | |
| c_1 | × | × | | × | | | |
| c_2 | × | × | × | | | | |
| c_3 | | × | × | × | | | |
| c_8 | × | | | | × | | × |
| c_9 | | × | | | × | | × |
| c_{10} | | | × | | × | × | |
| c_{12} | | | | × | × | × | |
| c_{14} | × | | | | | × | × |
| c_{16} | | | | | | × | × |

Crucially, recall that we are working in the projective setting, thus we can always find *two* incidence points between two non-tangent blocks. Then, it follows that C is a partial Möbius plane and the order $c_0 < c_1 < \dots < c_{16}$ is an HF-order, which means that $C \in \mathcal{K}$. As the other requirements from 3.60 are easily verified, it follows that C witnesses that 3.60 holds. ■

REFERENCES

- [1] A. M. Ammer and K. Tent. *On the model theory of open generalized polygons*. Model Th., **3** (2024), no. 2, 433-448.
- [2] C. Alaimo. *Completeness of the theory of open Steiner triple systems*. Master Thesis, University of Turin (2022).
- [3] J. Baldwin and N. Shi. *Stable generic structures*. Ann. Pure Appl. Logic **79** (1996), no. 1, 1-35.
- [4] J. Baldwin, J. Freitag and S. Mutchnik. *Simple homogeneous structures and indiscernible sequence invariants*. arXiv preprint arXiv:2405.08211 (2024).
- [5] J. Baldwin, G. Paolini. *Strongly minimal Steiner systems I: existence*. J. Symb. Log. **86** (2021), no. 4, 1486-1507.
- [6] J. Baldwin. *Strongly minimal Steiner systems II: coordinatization and quasigroups*. Algebra Universalis **84** (2023), no. 2.
- [7] J. Baldwin. *Strongly minimal Steiner Systems III: Path graphs and sparse configurations*. arXiv preprint arXiv:2201.11566 (2024).
- [8] S. Barbina and E. Casanovas. *Model theory of Steiner triple systems*. J. Math. Log. **20** (2020), no. 02.
- [9] S. Barbina and E. Casanovas. *The elementary theory of free Steiner triple systems*. arXiv preprint arXiv:2411.13723 (2024).
- [10] A. Barlotti and K. Strambach. *The geometry of binary systems*. Adv. Math. **49** (1983), no. 1, 1-105.
- [11] A. Baudisch and A. Pillay. *A free pseudospace*. J. Symb. Log. **65** (2000), no. 01, 443-460.
- [12] A. Baudisch, A. Martin-Pizarro and M. Ziegler. *A model theoretic study of right-angled buildings*. J. Eur. Math. Soc. **19** (2017), no. 10.
- [13] W. Benz. *Vorlesungen über Geometrie der Algebren*. Springer, Heidelberg, 1973.
- [14] F. Buekenhout (ed.). *Handbook of incidence geometry*. North-Holland, Amsterdam, 1995.
- [15] F. Buekenhout and P. Cameron. *Projective and affine geometry over division rings*. Handbook of Incidence Geometry, 27-62, North-Holland, Amsterdam, 1995.
- [16] A. Delandtsheer. *Dimensional linear spaces*. Handbook of Incidence Geometry, 193-294, North-Holland, Amsterdam, 1995.
- [17] P. Dembowski. *Freie und offene projektive Ebenen*. Math. Z. **72** (1959/1960), 410-438.

- [18] S. Ditor. *A new kind of free extension for projective planes*. *Canad. Math. Bull.* **5** (1962) 167-170.
- [19] H. D. Ebbinghaus and J. Flum. *Finite model theory*. Springer, Heidelberg, 1995.
- [20] M. Funk and K. Strambach. *On free constructions*. *Manuscripta Math.* **72** (1991), no. 4, 335-374.
- [21] M. Funk and K. Strambach. *Free constructions*. *Handbook of Incidence Geometry*, 739-780, North-Holland, Amsterdam, 1995.
- [22] M. Hall. *Projective planes*. *Trans. Amer. Math. Soc.* **54** (1943), 229-277.
- [23] D. Horsley and B.S. Webb. *Countable homogeneous Steiner triple systems avoiding specified subsystems*. *J. Combin. Theory Ser. A* **180** (2021).
- [24] T. Hyttinen and G. Paolini. *First-order model theory of free projective planes*. *Ann. Pure Appl. Logic* **172** (2021), no. 2.
- [25] M. L. Johnson. *Homomorphisms of free planes*. *Math. Z.* **125** (1972), no. 03, 255-263.
- [26] L. Kopeikina. *Decompositions of projective planes*. *Bull. Acad. Sci. URSS. Sér. Math.* **9** (1945) 495-526.
- [27] M. C. Laskowski. *A simpler axiomatization of the Shelah-Spencer almost sure theories*. *Israel J. Math.* **161** (2007), no. 1, 157-186.
- [28] A. A. Nikitin. *Homomorphisms of freely generated projective planes*. *Algebra Logic* **20** (1981), no. 4, 277-282.
- [29] A. Pasini. *Diagram geometries*. Clarendon Press, Oxford, 1994.
- [30] A. Schleiermacher and K. Strambach. *Freie Erweiterungen in der affinen Geometrie und in der Geometrie der Kreise I*. *Abh. Math. Sem. Univ. Hamburg* **34** (1970) 22-37.
- [31] S. Shelah and J. Spencer. *Zero-one laws for sparse random graphs*. *J. Amer. Math. Soc.* **1** (1988), no. 1, 97-115.
- [32] L. C. Siebenmann. *A characterization of free projective planes*. *Pacific J. Math.* **15** (1965), 293-298.
- [33] K. Tent. *Free polygons, twin trees, and CAT(1)-spaces*. *Pure Appl. Math. Q.* **7** (2011), no. 3, Special Issue: In honor of Jacques Tits, 1037-1052.
- [34] K. Tent. *The free pseudospace is n -ample, but not $(n + 1)$ -ample*. *J. Symb. Log.* **79** (2014), no. 2, 410-428.
- [35] K. Tent and M. Ziegler. *A course in model theory*. Cambridge University Press, Cambridge, 2012.
- [36] J. Tits. *Endliche Spiegelungsgruppen, die als Weylgruppen auftreten*. *Invent. Math.* **43** (1977), no. 3, 283-295.
- [37] H. Van Maldeghem. *Generalized polygons*. Birkhäuser, Basel, 1998.

DEPARTMENT OF MATHEMATICS “GIUSEPPE PEANO”, UNIVERSITY OF TORINO, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY.

Email address: gianluca.paolini@unito.it

Email address: davideemilio.quadrellaro@unito.it

ISTITUTO NAZIONALE DI ALTA MATEMATICA “FRANCESCO SEVERI”, PIAZZALE ALDO MORO 5 00185 ROMA, ITALY.

Email address: quadrellaro@altamatematica.it