WEIGHTED ENUMERATION OF NUMBER FIELDS USING PSEUDO AND SUDO MAXIMAL ORDERS

GAURAV DIGAMBAR PATIL

ABSTRACT. We establish a fundamental theorem of orders (FTO) which allows us to express all orders uniquely as an intersection of 'irreducible orders' along which the index and the conductor distributes multiplicatively. We define a subclass of Irreducible orders named Pseudo maximal orders. We then consider orders (called Sudo maximal orders) whose decomposition under FTO contains only Pseudo maximal orders. These rings can be seen as being "close" to being maximal (ring of integers) and thus there is a limited number of them with bounded index (by X). We give an upper bound for this quantity. We then show that all polynomials which can be sieved using only the Ekedahl sieve correspond to Sudo Maximal Orders. We use this understanding to get a weighted count for the number of number-fields with fixed degree and bounded discriminant using the concept of weakly divisible rings.

1. INTRODUCTION

We define the quantities $S(X:n)$ and $N(X:n)$ as follows:

Definition 1.

$$
S(X:n) := \{K : [K : \mathbb{Q}] = n, Gal(\overline{K}/\mathbb{Q}) \simeq S_n, \text{disc}(K) < X\}
$$
\n
$$
N(X:n) := #S(X:n)
$$

Malle's Conjecture (for *Sn*) in [\[8\]](#page-39-0) tells us to expect

$$
N(X:n)\simeq c_nX.
$$

where c_n is a constant dependent on only *n*. This conjecture is known for small values of *n*, namely $n \leq 5$. However, some lower bounds on the quantity $N(X : n)$ which are known are as follows.

• In [\[7\]](#page-39-1) (2002), Malle showed that

$$
N(X:n)\gg X^{\frac{1}{n}}.
$$

• In [\[5\]](#page-39-2)(2006) , Akshay Venkatesh and Jordan Ellenberg establish

$$
N(X:n)\gg X^{\frac{1}{2}+\frac{1}{n^2}}.
$$

• In [\[2\]](#page-39-3), Bhargava-Shankar-Wang show

$$
N(X:n) \gg_n X^{\frac{1}{2}+\frac{1}{n}}
$$

• In [\[3\]](#page-39-4), the same authors show

$$
N(X:n) \gg_n X^{\frac{1}{2}+\frac{1}{n-1}}.
$$

• In our previous paper [\[10\]](#page-39-5), we showed that

Date: Tuesday 19th November, 2024.

Theorem 1 (Number-field count)**.**

$$
N(X:n) \gg_n X^{\frac{1}{2} + \frac{1}{n-1} + \frac{(n-3)r_n}{(n-1)(n-2)}}
$$

where $r_n = \frac{\eta_n}{n^2 - 4n + 3 - 2\eta_n(n + \frac{2}{n-2})}$ and where η_n is $\frac{1}{5n}$ if *n* is odd and is $\frac{1}{88n^6}$ when *n* is even.

In that paper([\[10\]](#page-39-5)), we also show that the number of rings with rank *n* over \mathbb{Z} with bounded discriminant is

$$
\gg_n X^{\frac{1}{2}+\frac{1}{n-4/3}}.
$$

Since, for every prime *p* the ratio of non-maximal rings is *O*(1/*p* 2), one might expect that *N*(*X* : *n*) is also of the same order. In this paper, we show that that is true up to weighted enumeration and an ϵ in the order. Thus, showing that the liminf of $N(X:n)$ is bounded below by $X^{\frac{1}{2}+\frac{1}{n-4/3}}.$

Each of the above bounds uses different classifications of rings (see Introduction of our previous paper titled "Weakly Divisible Rings" [\[10\]](#page-39-5).).

There are two main obstructions (there are three but we will only look at two) on to getting better and better lower bounds for *N*(*X* : *n*). One side of the problem requires more general parametrizations of rings(as discussed in [\[10\]](#page-39-5)). The first three results given above use monic polynomials parametrizing monogenic rings. Heuristically, the third bound best possible using monogenic rings. This optimality is a result of solving a sieve theoretic problem. The fourth bound is based on binary rings and their parametrization via binary forms. Heuristically, this is the best possible result using just binary rings and a result of solving the second problem which is sieve theoretic in nature. In [\[10\]](#page-39-5) we introduce a different parametrization, namely weakly divisible rings, to get the fifth result. Our other computations heuristically suggests that the lower bound can be improved to ≫ $X^{\frac{1}{2}+\frac{1}{n-4/3}}$, provided we can optimize the seive that sieves for maximal (weakly divisible) rings among all (weakly divisible) rings.

To get lower bounds, one has to restrict the given parametrization to certain local conditions that force the ring under consideration to be the whole ring of integers.

Ekedahl sieve allows us to sieve out points in a parameter space that are lifts of points in a codimension two (or more) Variety over **F***p*. We thus refer to such a condition as Ekedahlian. In general local conditions are a combination of mod p , mod p^2 , etc conditions. A specific type of local condition that is not easily handled by the Ekedahl sieve (in its base form) is often the biggest deterrent to our ability to sieve for maximal rings of integers. In this paper, we show that in some such cases, using the Ekedahl sieve only we can instead get a weighted lower bound of the appropriate order. We do this by carefully looking at what it means for a polynomial to be weakly divisible in terms of the ring associated to it. The concept of weakly divisible polynomials is defined in [\[3\]](#page-39-4) and [\[2\]](#page-39-3).

We define

Definition 2.

$$
N^*(X:n) := \sum_{\mathbb{K} \in S(X,n)} \frac{1}{\sqrt{\mathfrak{disc}(\mathbb{K})}}.
$$

Malle's conjecture for $N(X : n)$ is clearly equivalent (via discrete integration by parts computation) to

$$
N^*(X:n)\simeq 2c_nX^{\frac{1}{2}}.
$$

In our previous paper we showed that, the number of "weakly divisible" rings of rank *n* over **Z** with discriminant less than or equal to *X* is

$$
\gg_n X^{\frac{\frac{1}{2}+\frac{1}{n-\frac{4}{3}}}}.
$$

one would expect that the number of number fields one gets sieving these rings would also be

$$
\gg_n X^{\frac{1}{2}+\frac{1}{n-\frac{4}{3}}}.
$$

This is because the product of the local probabilities of a ring under consideration having a index co-prime to *p* for any given prime *p* is (in all the currently known parametrizations) $\prod (1 O(p^{-2})$), which is bounded below by an absolute constant dependent on only *n*.

In this paper, we will show (using only the Ekedahl Sieve in its base form)

Theorem 2.

$$
N^*(X:n) \gg X^{\frac{1}{n-\frac{4}{3}}-\epsilon}.
$$

As a corollary, this implies

$$
\limsup_{X \to \infty} \frac{N(X:n)}{X^{\frac{1}{2} + \frac{1}{n - \frac{4}{3}} - \epsilon}} = \infty.
$$

If we get some good upper-bound on the size of the space of parametrization used (weakly divisible polynomials) or some sort of equi-distribution, it would follow that

$$
N(X:n) \gg_n X^{\frac{1}{2} + \frac{1}{n - \frac{4}{3}} - \epsilon}
$$

improving the result of our previous paper.

In section [2](#page-3-0) and section [3](#page-4-0) we discuss a Fundamental Theorem of Orders, which describes Orders being described as an intersection of *irreducible orders* in a unique way. For this decomposition of an order, the conductor of the given order distributes multiplicatively along the conductors of the the irreducible orders in its decomposition. The index of an order in its ring of integers also distributes (multiplicatively) along the decomposition of the order into irreducible orders. The above theorem and understanding give us a slightly more nuanced way to understand conductor ideals and give a nice proof of Furtwangler's condition for a conductor ideal of order in a given number field.

We will apply the context of FTO to binary rings and as a consequence show a Dedekind-Kummer Type Theorem for Binary Rings and factorization of associated binary form. This was previously shown by [\[4\]](#page-39-6), which we found in the references of [\[11\]](#page-39-7), after writing this paper.

We discuss small prime splitting restrictions for Binary Rings, a restriction similar to the one the small prime splitting restrictions for Monogenic Rings.

We then show a structure theorem of all 'large' irreducible orders. We call these orders as Pseudo Maximal Orders. We will give a more general structure theorem for irreducible orders in a future paper.

We then define *Sudo Maximal* Orders as orders whose decomposition in the Fundamental Theorem of Orders consists of only'large' irreducible orders (i.e. *Pseudo Maximal orders*). We have a structure theorem for such orders. We bound the number of such orders in a given number field of bounded index. This quantity can be bound in terms of only the bound on the index and the degree of the number field. Notably, this bound is independent of the discriminant of the number field. This

bound can be seen as an upper bound for all orders in a number field with squarefree conductors. This is evidence that indicates that the theorem shown in [\[6\]](#page-39-8) about orders in number fields with degree less than or equal to five, is generalizable to higher degrees. In an upcoming paper, we will do this.

We use the Ekedahl sieve on the space of polynomials and get polynomials which we refer to as Ultra-Weakly divisible polynomials. As a consequence of the Dedekind-Kummer Theorem for Binary Rings and the Ultra-Weakly Divisible condition on Binary Rings associated to Ultra-Weakly Divisible polynomials it follows that the Binary Rings (and Weakly Divisible Rings) associated to Ultra-Weakly Divisible Polynomials are all Sudo Maximal Rings.

Combining with our bound on Sudo Maximal Orders in a number field we show the result in theorem [2.](#page-2-0)

2. DEDEKIND DOMAINS IN NUMBER-FIELDS

Notation 3. Let K denote a number-field, \mathcal{O}_K its ring of integers. Let $M(K)$ denote the set of non-*Archimedean places of* **K** *i.e. the set of non-zero prime ideals in* \mathcal{O}_K *(or maximal ideals). Let* $\mathbb{P}_\mathbb{Z}$ *denote the set of prime numbers in integers.*

For $R \subseteq \mathbb{K}$ *be a ring (not necessarily of finite rank over* \mathbb{Z}) such that $Frac(R) = \mathbb{K}$, let \overline{R} denote its integral *closure.*

Lemma 1. $\forall p \in \mathbb{P}_{\mathbb{Z}} : |R/pR| \leq p^{[\mathbb{K}:\mathbb{Q}]}$.

Proof. We see that R/pR as a $\mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p$ -module.

For *x* ∈ *R* let *x* ∗ denote its image under the canonical map from *R* to *R*/*pR*. We follow our previous observation with another,

if $x_1^*, x_2^*, \dots, x_k^*$ are linearly independent over \mathbb{F}_p , then x_1, x_2, \dots, x_n are linearly independent over **Z**(or equivalently, over **Q**).

This is is easy to see since if x_i 's satisfy a $\mathbb Z$ linear relationship we may re write the relationship with coefficients such that the gcd of all the coefficients is 1 i.e. we may find a_i such that $\sum_{i=1}^{k} a_i x_i = 0$ and $(a_1, a_2, \dots a_k) = 1$ which immediately allows us to see $\sum_{i=1}^k a_i^* x_i^* = 0$ where not all $a_i^* = 0$ as $p \nmid (a_1, a_2, \cdots a_k) = 1.$

Lemma 2. $\forall p \in \mathbb{P}_{\mathbb{Z}} : |R/p^k R| = |R/p R|^k$.

Proof. Multiplication by p^m from $R \longrightarrow p^mR$ and $pR \longrightarrow p^{m+1}R$ are bijections and thus the canonical induced map, $R/pR \longrightarrow p^mR/p^{m+1}R$ is also a bijection, giving us $|p^mR/p^{m+1}R| = |\frac{R}{pR}|$. The lemma follows. \Box

It follows that *R*/*nR* is a finite module for all *n*.

If $x \in R$ then one can see that the numerator of $\mathfrak{Mm}(x)$ (say t) can be expressed as an integral polynomial of *x* and thus $R/xR \subseteq R/tR$ which is a finite set. Implying that every ideal in *R* is finitely generated and thus no-etherian. We also note that for every non zero prime ideal in *R*, *R*/*ρ* is a finite integral domain and hence a field i.e. ρ is a maximal ideal. Thus, \bar{R} is Noetherian, integrally closed domain with Krull dimension equal to one. It follows that *R* is a Dedekind Domain.

We naturally have $\mathcal{O}_K \subset \overline{R}$ for every such ring. (Since, integral closure of $\mathbb Z$ in $\mathbb K$ (or \mathcal{O}_K) will sit in the integral closure of *R* in **K**.)

Every non-zero prime (ρ) in \bar{R} thus naturally corresponds to a prime ($\rho \cap \mathcal{O}_K$) in \mathcal{O}_K . Furthermore, since *R*¯ is a Dedekind domain *R*¯ *^ρ* (*R*¯ localized at *ρ*) is a DVR and hence, a prime *ρ* in *R*¯ will naturally induce a non-Archimedean metric on **K**. Ostrowski's theorem tells us that *M*(**K**) simultaneously denotes the set of all non zero primes in \mathcal{O}_K and all possible non-Archimedean valuations on K. If $v = \rho \cap (\mathcal{O}_\mathbb{K}) \in M(\mathbb{K})$, it follows that $\bar{R}_\rho = (\tilde{\mathcal{O}}_\mathbb{K})_v$. This allows us to see non-zero prime ideals in \bar{R} as a subset of $M(K)$. On the other hand, given any nonempty subset (*S*) of *M*(**K**) one can obtain a Dedekind domain from this set by defining

Notation 4.

$$
R_S := \bigcap_{v \in S} (\mathcal{O}_\mathbb{K})_v.
$$

We thus get the following correspondence,

Theorem 3.

$$
\left\{\text{Dedekind Domains in } \mathbb{K}\right\} \leftrightarrow \left\{\text{ Non-Empty subsets of } M(\mathbb{K})\right\}
$$

Notation 5. *When R is a Dedekind Domain in* **K***, we use the notation*

$$
M(R) := \{ v \in M(K) : R \subseteq (\mathcal{O}_K)_v \}.
$$

We can write the correspondence as

$$
R \leftrightarrow M(R)
$$

$$
R_S \leftrightarrow S.
$$

Remark 4*.* We can let *g* denote the class number of **K** and observe that *v g* can be seen as a principal ideal (t_v) . For $S \subseteq M(K)$ we may define

$$
A(S) := \{ \prod_{v \in S}^* t_v^{n_v} : n_v \in \mathbb{Z}_{\geq 0} \}
$$

where \prod^* denotes finitely supported product. Then, the above correspondence from set to domain can be written as

$$
S\longrightarrow A(S)^{-1}\cdot \mathcal{O}_{K}.
$$

In particular, we notice that for any ring *R*,

$$
\bar{R}=R\mathcal{O}_{K}.
$$

3. ORDERS AND SEPARATION LEMMATA

Notation 6. *For* $R ⊆ K$ *a ring with Frac*(R) = K *, we define* $M(R)$ *to be the set of non-zero prime ideals (same as maximal ideals in this context) of R*.

When R is integrally closed (and thus a Dedekind Domain) this matches up with the understanding that M(*R*) *is the set of valuations corresponding to non-zero prime ideals of R from above.*

We note that for every ring *R* satisfying $Frac(R) = K$, $M(\overline{R})$ can be seen as a cover of $M(R)$.

To see this, we note that for $\rho, \rho' \in M(R)$ we clearly have $\rho + \rho' = R$ since they are distinct maximal ideals. Thus, valuations over *ρ* and *ρ* ′ have to be disjoint.

Notation 7. *For* $\rho \in R$ *, we define*

$$
\bar{\rho} := \{ v \in M(\bar{R}) : R_{\rho} \subseteq (\mathcal{O}_{\mathbb{K}})_v \}.
$$

In other words, $\{\bar{\rho} : \rho \in M(R)\}$ is a partition of $M(\bar{R})$.

We also note that

$$
R=\bigcap_{\rho\in R}R_{\rho}.
$$

We also note that there will only exist finitely many $\rho \in M(R)$ such that $R_\rho \subsetneq (\mathcal{O}_K)_v$. See, remark [5.](#page-5-0)

Notation 8. *We set*

$$
S(R) := \{ \rho \in M(R) : R_{\rho} \subsetneq (\mathcal{O}_{\mathbb{K}})_v \}.
$$

For all $\rho \notin S(R)$, we have $|\bar{\rho}| = 1$.

Definition 9. We say an ring R with Frac(R) = \mathbb{K} is irreducible if $|S(R)| = 1$.

In other words, R is irreducible if and only if $R \neq \bar{R}$ *and* $\exists !\rho$ *, a maximal ideal in R such that* $\forall \rho' \neq \rho$ *in M*(*R*), *we have*

$$
R_{\rho'} = (\mathcal{O}_\mathbb{K})_v
$$

for some $v \in M(\overline{R})$.

We call R an irreducible order*, when R is an order that is irreducible.*

We thus have every ring *R* with $Frac(R) = K$, we have

$$
R = \bigcap_{\rho \in S(R)} (R_{\rho} \cap \bar{R}).
$$
\n(1)

Notation 10. For an order \mathcal{O} , we let $c_{\mathcal{O}}$ denote the conductor of the order \mathcal{O} . In other words,

 $c_{\mathcal{O}} := (\mathcal{O} : \mathcal{O}_{\mathbb{K}})_{\mathbb{K}} = \{x \in \mathbb{K} : x\mathcal{O}_{\mathbb{K}} \subseteq \mathcal{O}\}\$

Remark 5*.* We recall some properties of the conductor ideal of an order.

- $c_{\mathcal{O}}$ is simultaneously an ideal of \mathcal{O} and \mathcal{O}_K .
- $c_{\mathcal{O}}$ is the largest ideal of \mathcal{O}_K sitting inside $\mathcal O$ as a subset.
- \mathcal{O}_ρ is not a DVR (i.e. is not equal to $(\mathcal{O}_K)_v$ for some valuation $v \in M(K)$) if and only if $c_{\mathcal{O}} \subseteq \rho$.

Thus, the radical of $c_{\mathcal{O}}$ in K will be the exactly equal to the product of valuations in $\sqcup_{\rho \in S(\mathcal{O})} \bar{\rho}$.

Lemma 3 (Separation Lemma 1). Let \mathcal{O} *is an order in* \mathcal{O}_K *, and*

$$
\mathcal{O} = \bigcap_{\rho \in S(\mathcal{O})} (\mathcal{O}_{\rho} \cap \mathcal{O}_{\mathbb{K}})
$$

from eq. [\(1\)](#page-5-1)*.*

Let c_{ρ} *denote the conductor of* $\mathcal{O}_{\rho} \cap \mathcal{O}_{K}$ *. We have*

•

$$
[{\mathcal O}_{{\mathbb K}}:{\mathcal O}]=\prod_{\rho\in S({\mathcal O})}[{\mathcal O}_{{\mathbb K}}:({\mathcal O}_\rho\cap{\mathcal O}_{{\mathbb K}})].
$$

• *For* $\rho, \rho' \in S(\mathcal{O})$

and

$$
c_{\mathcal{O}}=\prod_{\rho\in S(\mathcal{O})}c_\rho
$$

 $c_{\rho} + c_{\rho'} = \mathcal{O}_{\mathbb{K}}$

Proof. Since, for distinct primes $\rho, \rho' \in M(\mathcal{O})$ we have $\rho + \rho' = \mathcal{O}$. This implies that, for any primary ideals ∂ , ∂' satisfying $\sqrt{\partial} = \rho$ and $\sqrt{\partial'} = \rho'$, we have $\partial + \partial' = \mathcal{O}$. In other words,

$$
\mathcal{O}/\partial \partial' \simeq \mathcal{O}/\partial \times \mathcal{O}/\partial'.
$$

It follows that

$$
\mathcal{O}\otimes\mathbb{Z}_p=\prod_{\rho\in S(\mathcal{O})}(\mathcal{O}_\rho\cap\mathcal{O}_\mathbb{K})\otimes\mathbb{Z}_p=\prod_{\rho\in S(\mathcal{O})}\mathcal{O}_\rho\otimes\mathbb{Z}_p
$$

 $\mathrm{where}~\mathcal{O}_{\rho}\otimes\mathbb{Z}_p\subseteq \prod_{v\in\bar{\rho}}(\mathcal{O}_\mathbb{K})_v.$

We can look at all of these as \mathbb{Z}_p modules and compare the \mathbb{Z}_p indexes. We get

$$
v_p([\mathcal{O}_{\mathbb{K}}:\mathcal{O}])=\sum_{\rho\in S(\mathcal{O})}v_p([\mathcal{O}_{\mathbb{K}}:(\mathcal{O}_{\rho}\cap\mathcal{O}_{\mathbb{K}})]).
$$

The result follows.

The other part follows naturally from Remark remark [5](#page-5-0) as

$$
\sqrt{c_\rho}=\prod_{v\in\bar{\rho}}v
$$

and the fact that for any distinct primes $\rho, \rho' \in M(R)$,

$$
\bar{\rho}\cap \bar{\rho}'=\phi.
$$

 \Box

Lemma 4 (Separation Lemma 2). *If* $O = O_1 \cap O_2$ *such that* $c_{O_1} + c_{O_2} = O_K$, *then* $S(O_1)$ *and* $S(O_2)$ *can be naturally identified with disjoint subsets of S*(O) *such that*

- $S(\mathcal{O}_1) \sqcup S(\mathcal{O}_2) = S(\mathcal{O})$
- $\bullet \rho \in S(\mathcal{O}_1) \subseteq S(\mathcal{O}) \iff \mathcal{O}_\rho = (\mathcal{O}_1)_\rho$
- $\bullet \rho \in S(\mathcal{O}_2) \subseteq S(\mathcal{O}) \iff \mathcal{O}_{\rho} = (\mathcal{O}_2)_{\rho}.$

Proof. Since, $c_{\mathcal{O}_1} + c_{\mathcal{O}_2} = \mathcal{O}_\mathbb{K}$ let $r_1 + r_2 = 1$ with $r_1 \in c_{\mathcal{O}_1}$ and $r_2 \in c_{\mathcal{O}_2}$.

Fix a prime ideal $\rho \in S(\mathcal{O}_1)$ (that is, $c_{\mathcal{O}_1} \subseteq \rho$). Let $\tau = \rho \cap \mathcal{O}$.

Clearly, τ is a prime ideal in $\mathcal O$ and $\mathcal O_{\tau} \subseteq (\mathcal O_1)_{\rho}$.

If $s \in (O_1)_\rho$ then $s = q/t$ for some $q \in O_1$ and $t \in O_1 \setminus \rho$. We simply note that since $tr_2 \in c_{O_2} \subseteq O_2$ and $tr_2 = t - tr_1 \in \mathcal{O}_1$ we get $tr_2 \in \mathcal{O}$. We further note that $tr_2 \notin \tau$ as $tr_2 \in \tau \subseteq \rho$ would imply *r*₂ ∈ *ρ* and since *r*₁ ∈ *c*_{*O*₁} we get 1 = *r*₁ + *r*₂ ∈ *ρ*.) Similarly, *qr*₂ ∈ *c*_{*O*₂} ⊆ *O*₂ and *qr*₂ = *q* − *qr*₁ ∈ *O*₁ we get $qr_2 \in \mathcal{O}$.

We get $s = q/t = (qr_2)/(tr_2) \in \mathcal{O}_\tau$ implying

$$
\rho \in S(\mathcal{O}_1) \Rightarrow (\mathcal{O}_1)_{\rho} = \mathcal{O}_{\rho \cap \mathcal{O}}
$$

If $\rho, \rho' \in S(\mathcal{O}_1)$ such that $\rho \cap \mathcal{O} = \rho' \cap \mathcal{O}$, then $(\mathcal{O}_1)_{\rho} = (\mathcal{O}_2)_{\rho'}$ and thus $\rho = \rho'$.

Thus, the canonical map $S(\mathcal{O}_1) \longrightarrow S(\mathcal{O})$ given by $\rho \longrightarrow \rho \cap \mathcal{O}$ is injective. Similarly we can identify $S(\mathcal{O}_2)$ with a subset of $S(\mathcal{O})$. To show that $S(\mathcal{O}_1)$ and $S(\mathcal{O}_2)$ are disjoint, we only need to note that if some $\rho \in S(\mathcal{O})$ lies in both $S(\mathcal{O}_1)$ and $S(\mathcal{O}_2)$, we would immediately have $r_1 \in c_{\mathcal{O}_1} \subseteq \rho$
and $r_2 \in c_{\mathcal{O}_2} \subseteq \rho$. Forcing $1 = r_1 + r_2 \in \rho$. Contradicting ρ is a prime ideal. and $r_2 \in c_{\mathcal{O}_2} \subseteq \rho$. Forcing $1 = r_1 + r_2 \in \rho$. Contradicting ρ is a prime ideal.

Remark 6*.* Note that the above lemma tells us that irreducible orders mimic Euclid's property defining prime numbers, that is

If O is an irreducible order, then \mathcal{O}_A , \mathcal{O}_B are orders with $c_{\mathcal{O}_A} + c_{\mathcal{O}_B} = \mathcal{O}_K$, then $\mathcal{O}_A \cap \mathcal{O}_B \subseteq \mathcal{O} \Rightarrow \mathcal{O}_A \subseteq \mathcal{O}$ or $\mathcal{O}_B \subseteq \mathcal{O}$.

Theorem 7 (Fundamental Theorem of Orders)**.** *Every order* O *can be written as an intersection of irreducible orders in a unique way such that the conductors of the irreducible orders are pairwise co-prime. Furthermore, we get that the index (and conductor) of* \mathcal{O} *in* \mathcal{O}_K *will be the product of the indices (and conductors) of the irreducible orders in* \mathcal{O}_K *in the given decomposition.*

Proof. Clearly follows from lemma [3](#page-5-2) and lemma [4.](#page-6-0) □

Notation 11. For any valuation $v \in M(K)$, we denote the ramification index and the inertial degree of v *by e_v* and f_v , respectively. That is, if $v \cap \mathbb{Z} = (p)$ for some prime in \mathbb{Z} , then e_v and f_v are defined by

 $p\mathcal{O}_\mathbb{K} = \prod$

and

$$
\mathfrak{Nm}_{\mathcal{O}_{\mathbb{K}}}(v)=p^{f_v}.
$$

p∈*v*

 v^{e_v}

Definition 12. *For any order* O *and any non-zero prime ideal of* $\rho \in M(O)$ *, we define*

$$
ef(\rho) := \sum_{v \in \bar{\rho}} e_v \cdot f_v.
$$

Definition 13. *If* \mathcal{O} *is an irreducible order with* $S(\mathcal{O}) = \{\rho\}$ *, we define*

$$
ef(\mathcal{O}) := ef(\rho).
$$

4. DEDEKIND-KUMMER TYPE THEOREMS AND THE FUNDAMENTAL THEOREM OF ORDERS.

We recall the traditional versions of Dedekind-Kummer and Dedekind Criterion theorems surrounding polynomials.

4.1. **Dedekind-Kummer Theorem.**

Theorem 8 (Dedekind-Kummer Theorem). *Suppose* $\alpha \in \overline{\mathbb{Z}}$ *satisfies* $\mathbb{Q}[\alpha] = \mathbb{K}$ *, and suppose that* $f(x)$ *is the minimal polynomial of α (will be monic) over* **K***.*

If $p \nmid [\mathcal{O}_K : \mathbb{Z}[\alpha]]$ *and*

$$
f(x) \equiv \prod_i (f_i)^{e_i} \bmod p,
$$

then

$$
p\mathcal{O}_{\mathbb{K}}=\prod_i(\beta_i)^{e_i}
$$

where β_i *are prime ideals in* \mathcal{O}_K *sitting over p, with the understanding that*

$$
\beta_i = p\mathcal{O}_{\mathbb{K}} + f_i(\alpha)\mathcal{O}_{\mathbb{K}}.
$$

Notation 14. *Let* \overline{g} *denote the image of* $g \in \mathbb{Z}[x]$ *in* $\mathbb{F}_p[x]$ *.*

Dedekind's Criterion (see Lemma 3.1 in [\[1\]](#page-39-9)) for index adds

Theorem 9 (Dedekind's Criterion)**.** *We have*

$$
p\big| \big[\mathcal{O}_{\mathbb{K}} : \mathbb{Z}[\alpha] \big] \iff \exists i : 1 \leq i \leq k, \overline{f_i} \big| \frac{f - \prod_i (f_i)^{e_i}}{p}.
$$

Equivalently, we can say

$$
p\big| \big[\mathcal{O}_{\mathbb{K}}:\mathbb{Z}[\alpha]\big] \iff \exists g \in \mathbb{Z}[x]: f \in (p^2, pg, g^2) \subseteq \mathbb{Z}[x]
$$

We will unpack the standard proof of Dedekind-Kummer theorem with the Dedekind's Criterion by talking about the ring $\mathbb{Z}[\alpha]$ rather than \mathcal{O}_K as follows.

Theorem 10. *If* $\mathcal{O} = \mathbb{Z}[\alpha]$ *with* $\alpha \in \mathbb{Z}$ *and if* $f(x)$ *the minimal polynomial of* f *shows the following decomposition modulo p*,

$$
f(x) \equiv \prod_i (f_i)^{e_i} \bmod p,
$$

then, we get

$$
(\mathbb{Z}\backslash p\mathbb{Z})^{-1}\mathcal{O}=\bigcap_i\mathcal{O}_{\rho_i}
$$

and

$$
p\mathcal{O}=\prod_i\beta_i
$$

where βⁱ are primary ideals in O *sitting over p such that*

$$
\sqrt{\beta_i}=\rho_i
$$

 $\rho_i = p\mathcal{O} + f_i(\alpha)\mathcal{O}.$

with the understanding that

are all distinct prime ideals and all prime ideals in O *sitting over p and*

$$
\sum_i ef(\rho_i)=[\mathbb{K}:\mathbb{Q}].
$$

Remark 11*.* The theorem [10](#page-8-0) says that, "The factorization of a monic polynomial modulo *p* informs us about the local ring decomposition of **Z**[*α*] over *p*" and the theorem [9](#page-8-1) is about when a particular factor (corresponding to a particular prime ideal) does or does not correspond to a DVR.

4.2. **Dedekind-Kummer-Dedekind System for Binary Rings and Primitive Binary Forms.**

Theorem 12. *If* $\mathcal{O} = \mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}]$ *with* $\delta \in \overline{\mathbb{Q}}$ *and if* $f(x, y)$ *the minimal primitive binary form such that* $f(\delta, 1) = 0$ *and shows the following decomposition/factorization modulo p,*

$$
f(x,y) \equiv \prod_i (f_i)^{e_i} \bmod p,
$$

then, we get

$$
(\mathbb{Z}\backslash p\mathbb{Z})^{-1}\mathcal{O}=\bigcap_{i}\mathcal{O}_{\rho_{i}}
$$

and

$$
p\mathcal{O}=\prod_i\beta_i
$$

where βⁱ are primary ideals in O *sitting over p such that*

$$
\sqrt{\beta_i}=\rho_i
$$

with the understanding that

• *if* $f_i \neq Y$ *, then*

$$
\rho_i = (p\mathbb{Z}[\delta] + f_i(\delta, 1)\mathbb{Z}[\delta]) \cap \mathbb{Z}[\delta^{-1}]
$$

• *and if* $f_i \neq X$ *, then*

$$
\rho_i = (p\mathbb{Z}[\delta^{-1}] + f_i(1,\delta^{-1})\mathbb{Z}[\delta^{-1}]) \cap \mathbb{Z}[\delta]
$$

such that ρⁱ are all distinct prime ideals and all prime ideals in O *sitting over p and*

$$
\sum_i ef(\rho_i)=[K:Q].
$$

Theorem 13 (Dedekind's Criterion Binary Rings)**.** *We have*

$$
p\vert [\mathcal{O}_{\mathbb{K}}:(\mathbb{Z}[\delta]\cap \mathbb{Z}[\delta^{-1}])] \iff \exists i:1\leq i\leq k,\ \overline{f_i}\vert \frac{\overline{f-\prod_i(f_i)^{e_i}}}{p}.
$$

Equivalently, we can say

$$
p\big| \big[\mathcal{O}_{\mathbb{K}}:(\mathbb{Z}[\delta]\cap\mathbb{Z}[\alpha^{-1}])\big] \iff \exists g\in\mathbb{Z}[x,y]_m : f\in (p^2,pg,g^2)\subseteq\mathbb{Z}[x,y]_n
$$

Before jumping into the proof we will review some properties of binary rings associated to primitive forms.

4.3. **Localizations of Binary Rings associated to primitive forms.**

Let's review Binary rings: All of the following results can be found in [\[11\]](#page-39-7).

Let

$$
f(X,Y) := a_n X^n + a_{n-1} X^{n-1} Y + \dots + a_0 Y^n
$$

denote a binary form of degree *n*. Let $a_n \neq 0$ and suppose that δ denotes the image of *X* in the algebra $\mathbb{Q}[X]/f(X,1)$.

Definition 15. *When* $a_n \neq 0$,

$$
R_f := \mathbb{Z}\langle 1, a_n \delta, a_n \delta^2 + a_{n-1} \delta, \cdots, \sum_{i=0}^{k-1} a_{n-i} \delta^{k-i}, \cdots, \sum_{i=0}^{n-2} a_{n-i} \delta^{n-1-i} \rangle.
$$
 (2)

We set

$$
\langle B_0, B_1, \cdots, B_{n-1} \rangle := \langle 1, a_n \delta, a_n \delta^2 + a_{n-1} \delta, \cdots, \sum_{i=0}^{k-1} a_{n-i} \delta^{k-i}, \cdots, \sum_{i=0}^{n-2} a_{n-i} \delta^{n-1-i} \rangle
$$
 (3)
that is,

$$
B_0 := 1 \tag{4}
$$

$$
B_1 := a_n \delta \tag{5}
$$

$$
B_k := a_n \delta^k + a_{n-1} \delta^{k-1} + \dots + a_{n-k+1} \delta
$$
\n
$$
(6)
$$

$$
B_{n-1} := a_n \delta^{n-1} + a_{n-1} \delta^{n-2} + \dots + a_2 \delta
$$
 (7)

Definition 16. *I_f* denotes the (fractional) ideal class generated by $(1, \delta)$ in R_f

· · · · · ·

· · ·

Theorem 14. *R^f is a ring of rank n*.

Theorem 15. *Properties of R^f :*

(1)

$$
\operatorname{disc}_{\mathbb{Z}}(R_f) = \operatorname{disc}(f) \tag{8}
$$

(2) If δ is invertible, and f is primitive, then

$$
R_f = \mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}]. \tag{9}
$$

- (3) If f is primitive, I_f is invertible in R_f .
- *(4) R^f and I^f invariant a under the natural GL*2(**Z**) *action on binary forms of degree n*.
- *(5) We also have*

$$
R_f + R_f \delta = \mathbb{Z} \langle 1, \delta, B_2, \cdots, B_{n-1} \rangle \tag{10}
$$

and

$$
R_f \cap R_f \delta^{-1} = \mathbb{Z} \langle a_n, B_1, B_2, \cdots, B_{n-1} \rangle \tag{11}
$$

and

$$
(R_f + R_f \delta) \cdot (R_f \cap R_f \delta^{-1}) = \mathbb{Z} \langle \gcd(a_n, a_{n-1}, \cdots, a_0), B_1 - a_1, B_2 - a_2, \cdots, B_{n-1} - a_{n-1} \rangle. \tag{12}
$$

Remark 16*.* We make note of two facts. One is that the product (of modules) in eq. [\(12\)](#page-11-0) is in fact equal to R_f when $gcd(a_n, a_{n-1}, \dots, a_0) = 1$ (that is, *f* is primitive). The second is

$$
R_f/R_f \cap R_f \delta^{-1} \simeq \mathbb{Z}/a_n \mathbb{Z}.
$$
\n(13)

This implies that the prime ideals in R_f containing $R_f \cap R_f \delta^{-1}$ are of the form $pR_f + (R_f \cap R_f \delta^{-1})$ where $p \mid a_n$. We can conclude that for each prime number dividing a_n there exists exactly one prime in R_f containing p and $R_f \cap R_f \delta^{-1}$, and for $p \nmid a_n$ there is no prime ideal in R_f containing p . Furthermore, when $p \mid a_n$ the prime ideal in R_f containing p and $R_f \cap R_f \delta^{-1}$ clearly has norm p .

Notation 17. *Let* $O = \mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}]$. *Let* $f(x, y)$ *denote the minimal primitive binary form satisfying* $f(\delta, 1) = 0.$

Thus, $R_f \simeq \mathcal{O}$.

Lemma 5.

$$
(\mathcal{O}\cap\mathcal{O}\delta^{-1})\mathbb{Z}[\delta]=\mathbb{Z}[\delta]
$$

Proof. We know that from item [5,](#page-10-0)

$$
\mathcal{O}\cap\mathcal{O}\delta^{-1}=\mathbb{Z}\langle a_n,B_1,B_2,\cdots,B_{n-1}\rangle.
$$

Thus, $(\mathcal{O} \cap \mathcal{O} \delta^{-1})\mathbb{Z}[\delta]$ is spanned by the set organized in the table,

We note that $B_{i+1} - B_i \delta = a_{n-i}$ is in the \mathbb{Z} -span of the above set for all $n-1 \ge i \ge 1$. We can also see that $a_0 = -B_{n-1}\delta$ is also in the span. Thus,

$$
1 = \gcd(a_0, a_1, a_2, \cdots, a_n) \in (\mathcal{O} \cap \mathcal{O} \delta^{-1})\mathbb{Z}[\delta].
$$

Result follows. \Box

We note that if $\rho \in M(\mathcal{O})$ (prime ideal in \mathcal{O}) such that

$$
(\mathcal{O}\cap\mathcal{O}\cdot\delta^{-1})\nsubseteq\rho
$$

i.e. the denominator (proper) ideal of δ in $\mathcal O$ is not contained in ρ , then $\exists t \in \mathcal O \cap \mathcal O \cdot \delta^{-1}$ such that $t^{-1} \in \mathcal{O}_\rho$. Since, $t\delta \in \mathcal{O}$ it follows that $\mathbb{Z}[\delta] \subseteq \mathcal{O}_\rho$. For such primes ρ (primes not containing $\mathcal{O} \cap \mathcal{O} \cdot \delta^{-1}$), we note that $\mathbb{Z}[\delta]/\rho \mathbb{Z}[\delta]$ contains \mathcal{O}/ρ and is contained in $\mathcal{O}_\rho/\rho \mathcal{O}_\rho \simeq \mathcal{O}/\rho$ and hence must be equal to ^O/*^ρ* forcing *ρ***Z**[*δ*] to be a prime ideal in **Z**[*δ*]. This identifies *ρ* ∈ *M*(O) with a prime in $\mathbb{Z}[\delta]$ in a localization preserving way. That is to say,

$$
\mathbb{Z}[\delta]_{\rho\mathbb{Z}[\rho]}=\mathcal{O}_{\rho}.
$$

Combining with lemma [5,](#page-11-1) we argue that the only prime ideals in O which will not be mapped to any prime ideal in $\mathbb{Z}[\delta]$ will be exactly those prime ideals containing $\mathcal{O}\cap\mathcal{O}\cdot\delta^{-1}$

So, if we look at the map $M(\mathbb{Z}[\delta]) \longrightarrow M(\mathcal{O})$ given by $\rho \longrightarrow \rho \cap \mathcal{O}$, then this map is clearly the inverse of the localization preserving map above.

4.4. **Proof of Dedekind Kummer in Binary/One-fine Rings or theorem [12.](#page-9-0)**

This is easily observed using the previous understanding of prime ideals in $\mathcal{O} = \mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}]$ and their relationship to prime ideals in **Z**[*δ*] and the identification

$$
\mathbb{Z}[\delta] \simeq \mathbb{Z}[x]/(f(x,1)).
$$

Suppose, $f(x, y)$ has the following decomposition/factorization modulo p .

$$
f(x,y) \equiv y^k \prod_i (f_i)^{e_i} \bmod p,
$$

where f_i are irreducible binary forms modp such that $f_i(x, 1)$ has the same degree in *X* as the binary form f_i in (x,y) , (note that this will not happen if and only if $f_i = y$). We set

$$
\rho_i = (p\mathbb{Z}[\delta] + f_i(\delta, 1)^{e_i}\mathbb{Z}[\delta]).
$$

It follows that

$$
\mathbb{Z}[\delta]/\rho_i \simeq \mathbb{Z}[X]/(p, f_i(x,1)^{e_i}, f(x,1) \simeq \mathbb{Z}[X]/(p, f_i(x,1)) \simeq \mathbb{F}_p[x]/(f_i(x)^{e_i}).
$$

Thus, ρ_i are primary ideals in $\mathbb{Z}[\delta]$ with

$$
\sqrt{\rho_i}=(p, f_i(\delta)).
$$

On the other hand, it is easy to see that

$$
(p, f(x, 1)) = \prod_i (p, f_i(x, 1)^{e_i})
$$

where $i \neq j \implies (p, f_i(x, 1)^{e_i}) + (p, f_j(x, 1)^{e_j}) = \mathbb{Z}[x],$

since $f_i(x, 1)$ mod p and $f_i(x, 1)$ mod p are irreducible distinct polynomials in $\mathbb{F}_p[x]$. Thus,

$$
\mathbb{Z}[\delta]/(p) \simeq \prod_i \mathbb{F}_p[x]/(f_i(x,1)^{e_i}).
$$

We can conclude that $\sqrt{\rho_i}$ are all distinct prime ideals and all prime ideals in $\mathbb{Z}[\delta]$ containing p. This concludes the proof of the Dedekind Kummer theorem for Binary rings associated to primitive forms.

4.5. **Proof of Dedekind's Criterion for Primitive binary forms or theorem [13.](#page-9-1)**

The proof is very much the same as we can look at $\mathbb{Z}[\delta]$ and $\mathbb{Z}[\delta^{-1}]$ separately.

Suppose that $f(x, y)$ is the minimal primitive binary form satisfying $f(\delta, 1) = 0$.

Notation 18. *Given a polynomial* $h(x) \in \mathbb{Z}[x]$ *we let its image in* $\mathbb{F}_p[x]$ *be denoted by* $\bar{h}(x)$ *.*

Thus, if *f*(*x*, *y*) shows the following decomposition/factorization modulo *p*,

$$
f(x,y) \equiv y^k \prod_i (f_i)^{e_i} \bmod p,
$$

where f_i are irreducible binary forms modp such that $f_i(x, 1)$ has the same degree in x as the binary form f_i in (x,y) , (we note that this will not happen if and only if $f_i = y$) then we may set

$$
\beta_i = (p\mathbb{Z}[\delta] + f_i(\delta, 1)\mathbb{Z}[\delta])
$$

as the prime ideal associated to *fⁱ* .

It follows that

$$
\beta_i^2 = (p^2 \mathbb{Z}[\delta] + pf_i(\delta, 1)\mathbb{Z}[\delta] + f_i(\delta, 1)^2 \mathbb{Z}[\delta])
$$

Based on the structure of the prime ideal, we conclude a random element

$$
t = m(\delta) \in \beta_i \text{ if and only if } \bar{f}_i(x,1) | \bar{m}(x). \tag{14}
$$

We will consider two cases:

 $f_i(\delta, 1) \in \beta_i^2 \mathbb{Z}[\delta]_{\beta_i}$ and $f_i(\delta, 1) \notin \beta_i^2 \mathbb{Z}[\delta]_{\beta_i}$. • Case: $f_i(\delta, 1) \in \beta_i^2 \mathbb{Z}[\delta]_{\beta_i}$.

$$
f_i(\delta, 1) \in \beta_i^2 \mathbb{Z}[\delta]_{\beta_i} \implies f_i(\delta, 1) = p^2 A + p f_i(\delta, 1) B + f_i(\delta, 1)^2 C
$$

$$
\iff f_i(\delta, 1) (1 - p B - f_i(\delta, 1) C) = p^2 A
$$

Since, $1 - pB - f_i(\delta, 1)C \notin \beta_i$, it is a unit this means that $p^2R(\delta) = f_i(\delta, 1)S(\delta)$ where $S(\delta)$ is a unit in $\mathbb{Z}[\delta]_{\beta_i}.$

In this case, $\mathbb{Z}[\delta]_{\beta_i}$ is a DVR where the uniformizer can be taken to be p as it generates the maximal ideal in $\mathbb{Z}[\delta]_{\beta_i}$. Furthermore, since the uniformizer of $\mathbb{Z}[\delta]_{\beta_i} \otimes \mathbb{Z}_p$ is the same as the uniformizer of \mathbb{Z}_p , $\mathbb{Z}[\delta]_{\beta_i} \otimes \mathbb{Z}_p$ is an un-ramified extension of \mathbb{Z}_p . This forces $e_i = 1$, which implies $f(x, y) \notin (p^2, p f_i, (f_i)^2)$.

• Case: $f_i(\delta, 1) \notin \beta_i^2 \mathbb{Z}[\delta]_{\beta_i}$.

In this case, $f_i(\delta,1) \in \beta_i \mathbb{Z}[\delta]_{\beta_i} \backslash \beta_i^2 \mathbb{Z}[\delta]_{\beta_i}$. Thus, if $\mathbb{Z}[\delta]_{\beta_i}$ is a DVR then $f_i(\delta,1)$ may be chosen as a uniformizer.

We see $\mathbb{Z}[\delta_i] \otimes \mathbb{Z}_p$ as the ring of integers of a totally ramified extension of a totally unramified extension of \mathbb{Z}_p . Since, $\mathbb{Z}[\delta]/\beta_i$ is a $deg(f_i)$ extension of \mathbb{F}_p . Thus, the ramification degree of $\mathbb{Z}[\delta_i] \otimes \mathbb{Z}_p$ is $deg(f_i)$ and the ramification index is e_i . This will imply that

$$
f_i(\delta,1)^{e_i}=pu
$$

for some unit *u*.

Now substituting $(\delta, 1)$ in $f(x, y) = y^k \prod_j f_j(x, y)^{e_i} + ph(x, y)$ we get

$$
\prod_j f_j(\delta,1)^{e_j} = -ph(\delta,1)
$$

Since for $j \neq i$, $f_j(\delta, 1)$ are all units in $\mathbb{Z}[\delta]_{\beta_i}$, as they, by definition, are distinct irreducible polynomials over **F***p*.

This means that $h(\delta, 1)$ has to be a unit in $\mathbb{Z}[\delta]_{\beta_i}$ i.e. $\bar{f}_i(x, 1) \nmid \bar{h}(x, 1)$ (using eq. [\(14\)](#page-13-0)) which is equivalent to $h(x, y) \notin (p, f_i(x, y)) \iff f(x, y) \notin (p^2, pf_i, f_i^2)$.

To show the other direction, we note that, if $e_i = 1$ then clearly $f(x, y) \notin (p^2, pf_i, f_i^2)$. At the same time,

$$
\prod_j f_j(\delta,1)^{e_j} = -ph(\delta,1)
$$

and for $j \neq i$, $f_i(\delta,1)$ are all units in $\mathbb{Z}[\delta]_{\beta_i}$ which means that $\beta_i \mathbb{Z}[\delta]_{\beta_i} = p\mathbb{Z}[\delta]_{\beta_i}$ making it a principal ideal and thus **Z**[*δ*]*βⁱ* a DVR.

If $e_i > 1$ and $f \notin (p^2, pf_i, f_i^2)$ then $h(x, y) \notin (p, f_i(x, y))$. Subbing in δ we see that this means that $h(\delta, 1)$ is a unit in $\mathbb{Z}[\delta]_{\beta_i}$. At the same time,

$$
\prod_j f_j(\delta,1)^{e_j} = -ph(\delta,1)
$$

and for $j \neq i$, $f_j(\delta,1)$ are all units in $\mathbb{Z}[\delta]_{\beta_i}$ which means that $\beta_i \mathbb{Z}[\delta]_{\beta_i} = f_i(\delta,1) \mathbb{Z}[\delta]_{\beta_i}$ making it a principal ideal and thus **Z**[*δ*]*βⁱ* a DVR. This completes the proof.

5. SMALL PRIME SPLITTING RESTRICTIONS.

Just as a small prime splitting poses an obstruction for a ring to be monogenic, a small prime splitting poses a restriction for a ring to be binary. In this section we will quatify this restriction.

Let K denote a finite extension of Q. Let M_K denote the set of non-archimedean places of K . We fix a prime *p* in **Z** for this section. Let *O* denote an order in \mathcal{O}_K Let $p\mathcal{O} = \delta_1 \cdot \delta_2 \cdots \delta_r$ denote the unique m-primary ideal factorization of the ideal pO in O where $\delta_i + \delta_j = O$. Let τ_i denote the radical of the ideal δ_i in *R*. These will be all the maximal ideals of $\mathcal O$ containing p . For any non zero proper ideal I of O we define

$$
Norm_{\mathcal{O}}(I)=|\mathcal{O}/I|.
$$

And let

$$
T_{\mathcal{O}}(p, f) := \{ \tau_i : Norm(\tau_i) = p^f \} \qquad H(p, f) := \frac{1}{f} \sum_{d \mid f} p^d \mu(\frac{f}{d})
$$

Remark 17. *H*(p , f) is the number of prime ideals in $\mathbb{F}_p[x]$ with the norm p^f .

Theorem 18. *If* O *is binary ring, then*

$$
H(p,1) + 1 \ge |T_{\mathcal{O}}(p,f)|
$$

$$
f \ge 2 \implies H(p,f) \ge |T_{\mathcal{O}}(p,f)|.
$$

Proof. Let $\mathcal{O} = \mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}]$. From Dedekind Kummer Theorem for Binary forms (theorem [12\)](#page-9-0), we have a neat (norm preserving) correspondence from all prime ideals of **Z**[*δ*] containing *p* to either all or all but one prime ideal(that of norm *p*) of O . We identify $\mathbb{Z}[\delta]$ with $\mathbb{Z}[X]/g(x)$, where $g(x) = f(x, 1)$. Let $A = |T_{\mathcal{O}}(p, f)|$.

Consider the following diagram of canonical quotient maps,

$$
\mathbb{Z}[x] \longrightarrow \mathbb{Z}[\delta] \simeq \frac{\mathbb{Z}[x]}{\binom{g(x)}{g(x)}}
$$
\n
$$
\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow
$$
\n
$$
\mathbb{F}_p[x] \longrightarrow \mathbb{F}_p[\delta] \simeq \frac{\mathbb{Z}[x]}{\binom{p}{\mathcal{B}(x) \mathcal{B}(x) \mathcal{B}(x) \mathcal{B}(x)}} \longrightarrow \frac{\mathbb{Z}[\delta]}{\prod_{\tau \in A} \tau} \simeq (\mathbb{F}_{p^f})^{|A|}
$$

Clearly each arrow is surjective. Consider the lower row of maps culminating in the following surjective ring homo-morphism

$$
\mathbb{F}_p[x] \longrightarrow (\mathbb{F}_{p^f})^{|A|}
$$

In particular, there must exist at least |A| distinct prime ideals in $\mathbb{F}_p[x]$ with norm p^f . This corresponds to surjective maps from $\mathbb{F}_p[x]$ to \mathbb{F}_{p^f} up-to $Gal(\mathbb{F}_{p^f}/\mathbb{F}_p)$ action. In other words, the number of degree f elements in \mathbb{F}_{p^f} up-to $Gal(\mathbb{F}_{p^f}/\mathbb{F}_p)$ action. This is $H(p,f)$. Thus, $H(p,f) \ge |A|$. Combining with Dedekind Kummer Theorem for Binary rings, we see that,

$$
f \ge 2 \implies H(p, f) \ge |T_{\mathcal{O}}(p, f)|
$$

$$
H(p, 1) + 1 \ge |T_{\mathcal{O}}(p, f)|.
$$

 \Box

Remark 19*.* If

$$
\mathbb{P}(p, f) := #\{ \text{ prime ideals in } \mathbb{P}(\overline{\mathbb{F}}_p) \text{ with norm } p^f \},
$$

then,

and

$$
\mathbb{P}(p,1) = H(p,1) + 1
$$

$$
f \ge 2 \implies \mathbb{P}(p, f) = H(p, f).
$$

Remark 20*.* Binary rings are best viewed as a subset of orders which are locally monogenic, more specifically there are locally monogenic rings which are almost globally monogenic. Monogenic rings are globally monogenic (simultaneously across all primes). In case of *n* = 2 the concept of freely locally monogenic is equivalent to the version of almost globally monogenic used here. We compare this to, "Every ideal class in a rank two ring occurs in binary quadratic forms." When $n = 3$, this concept of locally monogenic is the same as almost globally monogenic used here, but not freely locally monogenic. We compare this to, "Every ring of integers occurs as a binary ring not every ideal class in ring of integers occurs as a binary ring and associated invertible ideal." We generalize the concept in our thesis and in upcoming papers and relate it to other classification of ring theorems.

Remark 21*.* We compare this condition with the small prime splitting condition in monogenic rings, which is

 $f \geq 1 \implies H(p, f) \geq |T_{\mathcal{O}}(p, f)|.$

In-spite of a very minimal weakening on this condition we note that the rings captured as monogenic rings are expected to be proportion 0 in binary rings (almost globally monogenic rings).

6. PSEUDO MAXIMAL ORDERS AND SUDO MAXIMAL ORDERS.

Remark 22. In general, if we have an irreducible order O with $S(\mathcal{O}) = \{\rho\}$ such that $\overline{\rho} = \{v_1, v_2, \cdots, v_k\}$ and if we let the conductor of \mathcal{O} to be $c_{\mathcal{O}}$, an ideal in \mathcal{O}_K , then we know that

$$
c_{\mathcal{O}}=(v_1)^{a_{v_1}}(v_2)^{a_{v_2}}\cdots(v_k)^{a_{v_k}}
$$

where $a_i \geq 1$.

Furthermore, the since $c_{\mathcal{O}} \subseteq \mathcal{O} \subseteq \mathcal{O}_K$ we see that $\mathcal O$ is completely determined by $\mathcal O/c_{\mathcal{O}}$ a sub-ring of $\mathcal{O}_{K}/c_{\mathcal{O}}$. Hence, we can search for sub-rings of the appropriate complete local finite extension of *p*-adic integers.

From here it is easy to get the Furtwrangler's condition for a conductor ideals by just looking at the ideal $(p^m, c_{\mathcal{O}})$ in a reduced order where $\mathcal O$ where $p^{m+1} \in c_{\mathcal{O}}$ and $p^m \notin c_{\mathcal{O}}$ and observing when it will be an ideal \mathcal{O}_K . This ideal can be an ideal of \mathcal{O}_K if and only if $c_{\mathcal{O}}$ cannot be a conductor ideal. In fact, combining with Fundamental Theorem of Orders or theorem [7](#page-7-0) it gives a more nuanced answer about orders and associated conductors and what can exist. For example, it tells us that

there cannot exist an order O with conductor vw with $f_v = e_v = f_w = e_w = 1$ and $O/v_w \simeq \mathbb{Z}_p^2$. This one is obvious, by things we already know.

6.1. **Classification of Irreducible orders** $\mathcal O$ with $ef(\mathcal O) = 1$ and $ef(\mathcal O) = 2$.

6.1.1. $ef(\mathcal{O}) = 1$.

There exist no irreducible order $\mathcal O$ in $\mathcal O_K$ with $ef(\mathcal O) = 1$ as this would be akin to finding proper unital sub-rings of the p-adic integers \mathbb{Z}_p .

6.1.2. $ef(\mathcal{O}) = 2$.

Lemma 6. If $A = \mathbb{Z}_p + \mathbb{Z}_p \cdot t$ is a algebra of rank 2 over \mathbb{Z}_p then all sub-rings of A which have rank 2 over \Z_p and containing \Z_p are given by $\Z_p+p^k(A)$ for some $k\in\Z^{\geq 0}.$ The index of $\Z_p+p^k(A)$ in A is *p k* .

Proof. If *M* is a sub-ring of *A* which is rank 2 over \mathbb{Z}_p may be seen as $\mathbb{Z}_p + \mathbb{Z}_p r$ where $r = a + bt$ with $a, b \in \mathbb{Z}_p$. We further write $b = p^k u$ where $u \in \mathbb{Z}_p$ is a unit, i.e. $u\mathbb{Z}_p = \mathbb{Z}_p$ Clearly,

$$
\mathbb{Z}_p + \mathbb{Z}_p r = \mathbb{Z}_p + \mathbb{Z}_p bt = \mathbb{Z}_p + \mathbb{Z}_p + \mathbb{Z}_p up^k t = \mathbb{Z}_p + \mathbb{Z}_p (p^k t) = \mathbb{Z}_p + p^k A.
$$

Remark 23. There is no irreducible(non \mathcal{O}_K) order $\mathcal O$ with $ef(\mathcal O) = 1$. Consequence of the Furtwangler condition for an ideal to be a conductor ideal. We give a structure theorem for irreducible orders with $ef(\mathcal{O}) = 2$. Similar Structure theorems can be given easily using corresponding ring classification theories for irreducible orders with $ef(\mathcal{O}) \leq 4$. With some difficulty, the authors also expect such theorems possible for $ef(\mathcal{O}) = 5$ by extending Bhargava's work for quintic rings to **Z***p*.

Definition 19. *A* Pseudo Maximal Order *in* K *is an irreducible order* $\mathcal O$ *satisfying ef*($\mathcal O$) = 2. *See definition [13](#page-7-1) for the definition of e f* .

If O is an irreducible order with $S(O) = \{\rho\}$, then its conductor is made up of prime ideals (in K) in *ρ*. Thus, when

Lemma 7. O *is a Pseudo Maximal Order, then one of the following must be true:*

$$
\overline{\rho} = \{v, w\} \text{ with } f_v = f_w = e_v = e_w = 1
$$

$$
= \{v\} \text{ with } f_v = 2 = 2e_v
$$

$$
= \{w\} \text{ with } 2f_v = 2 = e_v.
$$

Theorem 24 (A). If $\{v, w\} \subseteq M(K)$ with $f_v = f_w = e_v = e_w = 1$, then the irreducible orders with *conductor vaw b are given by*

- *if* $a = b = r$ then there is a unique irreducible order O with conductor $(vw)^r$ given by $\mathbb{Z} + (vw)^r$. *This order will have index in* \mathcal{O}_K *given by* $[\mathcal{O}_K : \mathbb{Z} + (vw)^r] = p^r$ *.*
- *if* $a \neq b$ then there is no irreducible (or otherwise) order with conductor $v^a w^b$.

Proof. Clearly, we are looking for irreducible orders in $\mathbb{Z}_p \oplus \mathbb{Z}_p$ which is treated an extension of \mathbb{Z}_p via diagonal embedding. So, we may write it as $\mathbb{Z}_p + \mathbb{Z}_p t$ where $t = (0, 1)$. Thus every order here must be of the form $\mathbb{Z}_p + p^k(\mathbb{Z}_p \oplus \mathbb{Z}_p)$. Coordinate wise $p = (p, p)$ due to diagonal embedding. Thus, the sub-ring is given by $\mathbb{Z}_p + ((p, 1)^k \cdot (1, p)^k)(\mathbb{Z}_p \oplus \mathbb{Z}_p)$. Intersecting with $\mathcal{O}_\mathbb{K}$ we get the order is given by $\mathbb{Z}+(vw)^k$ whose conductor is easily seen as $(vw)^k$.

Theorem 25 (B). If $\{v\} \subseteq M(K)$ with $f_v = 2 = 2e_v$, then there is a unique irreducible order with *conductor* v^a *which is given by* $\mathbb{Z} + v^a$ *with* $[\mathcal{O}_\mathbb{K} : (\mathbb{Z} + v^a)] = p^a$

Proof. Clearly, we are looking for irreducible orders in the ring of integers of the unique totally unramified extension of degree 2 over \mathbb{Z}_p . Thus, p may be seen as the uniformizer for this completion. So, we may write it as $O_v := \mathbb{Z}_p + \mathbb{Z}_p t$. Thus every order here must be of the form $\mathbb{Z}_p + p^k(O_v)$. Thus, the sub-ring is given by $\mathbb{Z}_p + p^k(O_v)$. Intersecting with $\mathcal{O}_\mathbb{K}$ we get the order is given by $\mathbb{Z} + v^k$ whose conductor is easily seen as v^k .

Theorem 26 (C). *If* $\{v\} \subseteq M(\mathbb{K})$ *with* $2f_v = 2 = e_v$, *then*

- *if a* \geq 2 *and a is even then there is a unique irreducible order with conductor* v^a *which is given by* $\mathbb{Z} + v^a$ with $[\mathcal{O}_K : (\mathbb{Z} + v^a)] = p^{a/2}.$
- *if* $a \geq 1$ and a is odd then there is no irreducible (or otherwise) order with conductor v^a .

Proof. Clearly, we are looking for irreducible orders in the ring of integers of some totally ramified extension of degree 2 over \mathbb{Z}_p . Thus, p may be seen as the square of the uniformizer for this completion. So, we may write it as $O_v := \mathbb{Z}_p + \mathbb{Z}_p t$ where *t* is the uniformizer(since it is totally ramified. Thus, every order here must be of the form $\mathbb{Z}_p + p^k(\mathcal{O}_v)$. Thus, the sub-ring is given by $\mathbb{Z}_p + p^k(O_v)$. Intersecting with $\mathcal{O}_\mathbb{K}$ we get the order is given by $\mathbb{Z} + v^{2k}$ (as $(p) = v^2$, as it is a totally ramified extension) whose conductor is easily seen as v^{2k} . В последните поставите на примерение в поставите на примерение в поставите на примерение в собствение и при
В поставите на примерение в поставите на примерение в поставите на примерение в поставите на примерение в пост

A corollary we will use for counting purposes is,

Corollary 1. Given a set of valuations $S \subseteq M(\mathbb{K})$ such that $\sum_{v \in S} e_v f_v = 2$ then given $r \ge 1$ there exists *a* unique irreducible order O with $S(O) = \{\rho\}$ such that $\overline{\rho} = S$ and $[O_K : O] = p^r$.

Definition 20. *A* Sudo Maximal Order *in* **K** *is an order* O *such that*

$$
\forall \rho \in S(\mathcal{O}): ef(\rho)=2.
$$

We say that this is Restricted Sudo Maximal Order if for every prime p ∈ **Z** *we have at most one prime ideal in S*(O) *which contains p*.

Theorem 27. *The number of Restricted Sudo Maximal Orders in* **K** *with index* ≤ *X is*

$$
\leq A_n X \log(X)^{{n \choose 2} - 1}
$$

where Aⁿ is a constant only dependent on n.

Proof. We note that if *C***K**,*^p ^k* denote the number of Restricted Sudo Maximal Order in **K** with index *p k* , we may look at

$$
S_{\mathbb{K},p}(s):=\sum_{k=0}^\infty C_{\mathbb{K},p^k}p^{-ks}
$$

and look at the L-function

$$
L_{\mathbb{K}}(s):=\prod_{p\in\mathbb{P}}S_{\mathbb{K},p}
$$

Since we know from the above section that

$$
C_{\mathbb{K},p^k} = |\{(v,w): v, w \in M(\mathbb{K}), f_v = f_w = 1, v \neq w, v(p) = w(p) = 1\}|
$$

+
$$
|\{v \in M(\mathbb{K}): f_v = 2v(p) = 1\}| + |\{v \in M(\mathbb{K}): f_v = 1v(p) = 2\}|.
$$

Thus, $C_{\mathbb{K},p^k} \leq \binom{n}{2}$ $\binom{n}{2}$.

It follows that the Dirichlet coefficients of $L_K(s)$ are all between 0 and the Dirichlet coefficients of $\zeta(s)^{n \choose 2}$.

Sum of Dirichlet coefficients up to *X* of $\zeta(s)^{(\frac{n}{2})}$ grows asymptotically like

$$
\frac{X\log(X)^{{n\choose 2}-1}}{{n\choose 2}-1}.
$$

Thus, the number of orders in \mathbb{K} with index $\leq X$ is

$$
\leq A_n \frac{X \log(X)^{{n \choose 2} - 1}}{({n \choose 2} - 1)!}
$$

for some constant A_n only dependant on n .

Remark 28*.* We note that this gives a strong indication that the number of orders in **K** with index bounded by *X* should be of the order of *X*.

6.2. **Strongly divisible-ness and Ultra-weakly divisible polynomials.**

Remark 29*.* Recall that Bhargava-Shankar-Wang define *f* is strongly divisible by *p* (in [\[3\]](#page-39-4) and [\[2\]](#page-39-3)) if p^2 |disc(*f* + *p* · *g*) for any choice of *g* such that $deg(f) \geq deg(g)$.

Recall the theorem.

Theorem 30. p^2 |disc(f) *if and only if one of the following holds*:

- *f is strongly divisible by p if and only if one of the following is true.*
	- *(1) f* mod *p* has a triple root in $\mathbb{P}(\mathbb{F}_p)$
	- *(2) f* mod *p* has two double roots in $\mathbb{P}(\overline{\mathbb{F}}_p)$.
- *If f is not strongly divisible by p, then one of the following holds*
	- *(1) f is weakly divisible by p*. *This is the case where the linear double root in* **P**(**F***p*) *is in* **F***^p (seen as a base affine component) and not at the point at infinity.*
	- *(2) The lead coefficient is divisible by p*² *and the second lead is divisible by p. In other words, the palindromic reverse of f is weakly divisible by p at* 0.*This is the case where the linear double root in* **P**(**F***p*) *is at the point at infinity. (We will try and ignore this case by making sure that the lead coefficient of our polynomial is squarefree or making sure that the leading two coefficients are co-prime)*

Proof.

• We start by showing that if

 $-$ *f* mod *p* has a triple root in $\mathbb{P}(\mathbb{F}_p)$ or

 $-$ *f* mod *p* has two double roots in $\mathbb{P}(\overline{\mathbb{F}}_p)$

then p^2 |disc(*f*). Note that this automatically implies that *f* is strongly divisible by *p*.

Since, we just proved the Dedekind-Kummer Theorem for Binary Rings (theorem [12\)](#page-9-0), let us use it here.

If *f*(*x*, *y*) shows the following decomposition/factorization modulo *p*,

$$
f(x,y) \equiv \prod_i (f_i)^{e_i} \bmod p,
$$

then we know $(f_i)^{e_i}$ corresponds to either a DVR or it corresponds to a proper subring in associated DVR.

Now if some $(f_i)^{e_i}$ corresponds to a proper subring of a DVR, then p will divide the index of corresponding local ring in DVR. This will imply p^2 |disc(*f*).

On the other hand, if it does not correspond to an irreducible order, then *R^f* localized at that prime is a DVR, which corresponds to a degree e_i totally ramified extension of the totally un-ramified extension of degree f_i of the local field \mathbb{Q}_p . This means that p^{e_i-1} divided the discriminant of the corresponding local field extension and thus *p* ∑*i ^ei*−¹ divides the discriminant of **K***^f* . In this case, however, the power of *p* dividing the discriminant of **K***^f* and *f* is the same.

Thus, $p^2|\text{disc}(f + pg)$ for any choice of *g* provided that for some *i*, $deg(f_i) \geq 2$ & $e_i \geq 2$.

Similarly, p^2 |disc($f + pg$) for any choice of *g* provided that for some *i*, $deg(f_i) \geq 1$ & $e_i \geq 3$.

• If p^2 |disc(*f*), then *p*|disc*f*. Thus, *f* has a double root in $\mathbb{P}(\overline{\mathbb{F}}_p)$. If this root is non linear then we fall in the case discussed above which implies *f* is strongly divisible by *p*. Similarly, if the root is linear and the multiplicity is greater than 3 we again fall in the above case.

Thus, since *f* is not strongly divisible by *p*, we may assume that *f* has only one double root in $\mathbb{P}^1(\bar{\mathbb{F}}_p).$ Now if this linear double root is in \mathbb{F}_p , we will translate appropriately so that the double root is at zero. If the linear root is a point at infinity we take the palindromic inverse or the reciprocal polynomial to shift its double root to 0. Then, showing initial polynomial is weakly divisible by *p* as in definition [32](#page-29-0) at the root will be same as showing new *f* is weakly divisible by *p* at 0.

We note that, using lemma [15,](#page-37-0) the discriminant of *f* can be written as

$$
\operatorname{Disc}(f) = 2f(0)\left(\frac{f^{(2)}(0)}{2!}\right)^3 \cdot \Delta_1 + f'(0)^2 \cdot \Delta_2 + f(0)f'(0) \cdot \Delta_3 + f(0)^2 \Delta_4
$$

where

$$
\Delta_1 = \mathfrak{disc}(\frac{f - f(0)Y^n - f'(0)XY^{n-1}}{XY}).
$$

Since, f is not strongly divisible by p , we see that $p\nmid \Delta_1$ (no two double roots) and $p\nmid \frac{f^{(2)}(0)}{2!}$ 2! (no triple linear root).

Since,
$$
p|f(0)
$$
 and $p|f'(0)$ and $p^2|\text{Disc } f$ and $p \nmid \Delta_1$ and $p \nmid \frac{f^{(2)}(0)}{2!}$ it follows that $p^2|f(0)$.

See appendix.

Definition 21. *We say a f ultra weakly divisible if*

 p^2 |disc $(f) \Rightarrow f$ *is weakly divisible by p*.

If we treat $\mathfrak{disc}(f)$ as a function of its coefficients (say a_i by abuse of notation), then this condition $\frac{\partial}{\partial a_i}$ corresponds to $p^2|\text{Disc}(f)|$ and $p|\frac{\partial(\text{disc})}{\partial a_i}|$ *∂aⁱ* (*f*) for all 0 ≤ *i* ≤ *n* (*f* mod *p* is in the singular locus of $\mathfrak{disc}(f) = 0 \text{ mod } p$.

Remark 31*.* While it may seem that the singular locus has a very large co-dimension, it has a component of co-dimension 2.

By definition, if *f* is ultra-weakly divisible, then *f* does not have non linear double root or a linear triple root modulo any prime *p*.

Theorem 32. *If f is ultra-weakly divisible, then there exist a maximal m^f such that f is weakly divisible* by m_f (at some l_f) and $R'_{(f,m_f,l_f)}$ is maximal i.e. $R'_{(f,m_f,l_f)}$ is THE ring of integers for $\mathbb{K}_f\simeq \mathbb{Q}[x]/(f)$. In *fact, if* $\mathfrak{disc}(f) = st^2$ *where s is squarefree, then* $m_f = t$.

Proof. Let *p* denote a prime such that p^2 |disc(*f*). And let $0 \le l_0 < p$ denote the linear double root in \mathbb{F}_p of *f*. Then, we note that if

$$
f(x+l) = f(l) + f'(l) \cdot x + \frac{f^{(2)}(l)}{2!} \cdot x^2 + \dots + a_0 x^n
$$

then using lemma [15](#page-37-0) we see that,

$$
\operatorname{Disc}(f) = f(l) \cdot \Delta_1 + f'(l)^2 \cdot \Delta_2 + f(l) f'(l) \cdot \Delta_3.
$$

Furthermore, if $l \equiv l_0 \mod p$ then we can realize Δ_1 as

$$
\Delta_1 \equiv 4(\frac{f^{(2)}(l)}{2!})^3 \text{disc}(f^*) \bmod p
$$

where $f^* = \frac{f(x)}{(x-1)^2}$ $\frac{f(x)}{(x-l_0)^2}$ ∈ **F**_{*p*}[*x*].

Since *f* is ultra-weakly divisible by *p*, we have *f* is not strongly divisible by *p*. This means that if *l* \equiv *l*₀ mod *p* then $p \nmid \frac{f^2(l)}{2}$ $\frac{1}{2}$ as this will force *f* to not have triple linear root at *l*₀ in **F**_{*p*}. We can further say that that $p \nmid \Delta_1$. If $p \mid \Delta_1$ then it is easy to see that f will have at least two double roots in $\bar{\mathbb{F}}_p$.

Now, by Hensels lemma for any *k* we can find l_k such that $p^k|f'(l_k)$.

Let $D = v_p(\mathfrak{disc}(f))$ and $l \equiv l_k \text{ mod } p^k$ with $0 \le l < p^k$ where $k = \lceil \frac{D}{2} \rceil$.

Then, we note that $p^r | f(l)$ and $r < 2[\frac{D}{2}]$ $\frac{D}{2}$. It follows that $\mathfrak{disc}(f) \equiv p^r \cdot \Delta_1 \bmod p^{r+[D/2]}$ which contradicts the fact that $p^D ||$ disc (f) .

Thus, $p^{2 \cdot [D/2]} |f(l)$ and $p^{[D/2]} |f'(l)$ or *f* is weakly divisible by $p^{[D/2]}$. Thus, *f* is weakly divisible by m_f where $\mathfrak{disc}(f) = s(m_f)^2$ and s is squarefree. This tells us that $R'_{(f, m_f, l_f)}$ is $p^{D-2 \cdot [D/2]}$ which is at most one. This makes $R'_{(f,m_f,l_f)}$ the ring of integers of \mathbb{K}_f as its discriminant is squarefree.

7. POLYNOMIALS THAT CORRESPOND TO DISTINCT RINGS

Definition 22.

$$
W(s:t) := \Big\{ \sum_{i=0}^{n} a_i x^{n-i} \in \mathbb{Z}[x] : 1 \leq a_1 < s/2 < a_0 \leq s, (a_0, a_1) = 1, |a_i| \leq s t^i \text{ for all } 2 \leq i \leq n \Big\}
$$

Remark 33*.* The above set is a subset of a fundamental domain for action of integers by translation on polynomials. That is, for any polynomial *f* there exists at most one integer *l* such that $f(x+l) \in$ $W(s : t)$.

Definition 23.

$$
W(s:t:m) := \{(f,l) : f \in W(s:t), 0 \le l < m, f \text{ is weakly divisible by } m \text{ at } l\}
$$

with the understanding that each element $(f, l) \in W(s : t : m)$ where f is Reduced-m-Polynomial will give distinct rings, see Theorem theorem [50.](#page-35-0)

Lemma 8. *Given* a_0, a_1, \dots, a_{n-2} *, <i>l, m there is a unique choice of* a_{n-1} *and* a_n *satisfying*

\n- $$
B + 1 \le a_{n-1} \le B + m
$$
\n- $$
C + 1 \le a_{n-1} \le C + m^2
$$
\n- $$
a_0 x^n + a_1 x^{n-1} + \cdots + a_n
$$
 is weakly divisible by *m* at *l*.
\n

Proof. Follows from the definition directly. □

Discussion 24. Let B_{ϵ} denote a compact set in

$$
\{f\in\mathbb{R}[x]:H(f)\leq 1, \mathfrak{disc}_\mathbb{Z}(f)\neq 0\}
$$

such that

$$
Vol(B_{\epsilon}) \ge (1 - \epsilon) Vol(\{f : H(f) \le 1\}).
$$

Then for this region we can construct a ρ_B *such that all polynomials f with height* $\geq \rho_B$ *satisfying*

$$
H(f)^n f(\frac{X}{H(f)}) \in B_{\epsilon}
$$

*are Reduced-*1*-Polynomials.*

From theorem [50,](#page-35-0) we can say that polynomials of this type with height

$$
\geq m^{1/(n-2)} \rho_B
$$

are Reduced-m-Polynomials.

Furthermore, we may consider B_ϵ *to be a finite union of disjoint boxes* B_i *(depending on epsilon).*

Thus, correspondingly, if $d_{i,r}$ *are the dimensions of the box* B_i *, we define* $W(s:t:m)_{B_i}$ *to be those polynomials that satisfy*

$$
\frac{H(f)^n}{s}f(\frac{X}{H(f)}) \in B_i
$$

and

$$
\frac{f^{(2)}(0)}{2!} \ge s\rho_B^{n-2}m.
$$

The latter condition a_{n−2} \ge $sp_B^{(n-2)}$ *m ensures that polynomials we count are of 'large enough' (see lemma [14\)](#page-36-0) height to be a Reduced-m-Polynomial.*

Definition 25.

$$
W(s:t:m)^{Red} := \{(f,l) \in W(s,t,m) : f \text{ is a Reduced-m-Polynomial.}\}
$$

7.1. **Ekedahlian Sieve- Quantitative version.**

Let V_n denote the variety in \mathbb{A}^{n+1} which describes the Singular locus of the curve given by $G_n = 0$ where

$$
G_n := \text{diag}(X_0 X^n + X_1 X^{n-1} + \cdots X_n).
$$
 (15)

We describe the structure of the discriminant polynomial as a polynomial of *Xn*.

Lemma 9.

$$
\begin{aligned} \mathfrak{disc}_X(X_0X^n + X_1X^{n-1} + \dots + X_n) &= \\ n^n (X_0X_n)^{n-1} \\ &+ \sum_{i=1}^{n-2} X_n^i \Delta_i \\ &+ X_{n-1}^2 \mathfrak{disc}_X(X_0X^{n-1} + X_1X^{n-2} + \dots + X_{n-1}) \end{aligned}
$$

Proof. Obvious from the the understanding that X_0 α **isc**_{*X*}($X_0X^n + X_1X^{n-1} + \cdots + X_n$) is the resultant of $X_0X^n + X_1X^{n-1} + \cdots + X_n$ and $\frac{\partial}{\partial X}(X_0X^n + X_1X^{n-1} + \cdots + X_n)$ can the consideration of this resultant as the determinant of the Sylvester matrix. See eq. (28) and lemma [15.](#page-37-0)

Let

$$
F_n := \text{Disc}_{X_n}(\text{Disc}_X(X_0X^n + X_1X^{n-1} + \cdots + X_n))
$$
\n(16)

Thus, for any prime *p*, and integer tuple (a_0, a_1, \dots, a_n) ,

$$
(a_0, a_1, \cdots, a_n) \in V_n(\mathbb{F}_p) \Rightarrow p | (G_n(a_0, a_1, \cdots, a_n), F_n(a_0, a_1, \cdots, a_{n-1})).
$$
 (17)

This follows from the fact that G_n , $F_n \in I(V_n)$.

We note that

Remark 34*.* A polynomial *f* is UWD or ultra-weakly divisible if and only if

$$
\forall p \in \mathbb{P} : f \bmod p \notin V_n(\mathbb{F}_p).
$$

We will now look for UWD polynomials in *W*(*s* : *t* : *m*). We will then look for Reduced-*m*polynomials in this set which will allow us to count UWD Rings. We note that UWD polynomials are easily sieve-able via the Ekedahl sieve.

Discussion 26. *Given* $f = \sum_{i=0}^{n} a_i x^{n-i}$ *such that* $f(x)$ *is weakly divisible by m at* 0 *and* $(a_0, a_1) = 1$ *,*

$$
\frac{\text{Disc}(f)}{m^2} \equiv 2(a_{n-2})^3 \text{Disc}(f^*) \bmod m
$$

 ω here $f^* = \sum_{i=0}^{n-2} a_i x^{n-i-2}$. This follows from the structure of the discriminant polynomial in lemma [15.](#page-37-0) *Thus, the only way f can be strongly divisible by p for some* $p|m$ *if*

- *either p*|*an*−2*(this will force a triple root at* 0*)*
- or $p | \text{disc}(f^*)$ (*this will force some other double root in* \mathbb{F}_p *).*

Furthermore, applying the structure of the discriminant (lemma [15\)](#page-37-0) *to f*^{*} *we can say that for every* $a_0, a_1, \cdots, a_{n-3}$ *and p*|*m there are at most n* + 1 *solutions* mod *p* for a_{n-2} *such that*

$$
2(a_{n-2})^3 \text{disc}(\sum_{i=0}^{n-2} a_i x^{n-i-2}) \equiv 0 \bmod p.
$$

We will use the traditional form of the Ekedahl sieve. This will serve us better. Our parameter of *m* which interferes with the sieving in the main term. And since we wish to maximize *m* to be used, we wish to use the fact that *an*−² freely moves in a much larger range than other coefficients and based on the structure of G_n can be used to give a usable lower bound for $f \notin V_n(\mathbb{F}_p)$ for $p|m$ for much larger values of *m*.

Theorem 35. If
$$
s \gg \frac{st^n}{m^2}
$$
, then
\n
$$
|\{(f, l) \in W(s : t : m) : \exists p \in \mathbb{P}_{\mathbb{Z}}, p > M, p \nmid m, f \in V_n(\mathbb{F}_p)\}|
$$
\n
$$
= O(\frac{s^n t^{\frac{n(n+1)}{2}-1}}{m^2}) \cdot (\frac{1}{M \log M} + \frac{m^2 \log(st^n)}{st^n})
$$

Proof. As usual we put this set into the union of 3 sets:

•

$$
S_1 := \{(f, l) \in W(s : t : m) : F_n(f) = 0\}
$$

For any choice of $(a_0, a_1, \dots, a_{n-2}, l)$ such that F_n is non-degenerate as a polynomial in a_{n-1} , then we can have at most $n(n-1)$ possible choices for a_{n-1} .

Inductively, the number of choices for $(a_0, a_1, \dots, a_{n-2})$ for which the polynomial F_n is degenerate as a polynomial in *an*−¹ would be space cut out by all coefficient polynomials in \mathbb{A}^{n-1} . Thus, the number of elements in S_1 is

$$
O_n(\frac{s^n t^{\frac{n(n+1)}{2}-1}}{m^2}) \cdot (\frac{1}{s} + \frac{m}{st^{n-1}}).
$$

•

$$
S_2 := \{ (f, l) \in W(s : t : m) : \exists p \in \mathbb{P}_{\mathbb{Z}}, \frac{s t^n}{m^2} > p > M, p \nmid m, f \in V_n(\mathbb{F}_p) \}
$$

For every such prime *p* the number of solutions (mod*p*) to

$$
G_n(f) \equiv F_n(f) \equiv 0 \bmod p
$$

is $O(p^{n-1})$. Thus, the total number of solutions possible $f \in W(s : t : m)$ is

$$
O(p^{n-1}) \cdot m \cdot (\frac{s}{p} + O(1))^2 \cdot (\frac{2st^2}{p} + O(1)) \cdots (\frac{2st^k}{p} + O(1)) \cdots (\frac{2st^{n-2}}{p} + O(1))
$$

$$
\cdot (\frac{2st^{n-1}}{pm} + O(1)) \cdot (\frac{2st^n}{pm^2} + O(1))
$$

$$
= O(\frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2})(\frac{1}{p^2}).
$$

Summing this value over the given range for primes, we get that |*S*2| is

•

$$
O(\frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2M\log M})
$$

$$
S_3 := \{ (f, l) \in W(s : t : m) : F_n(f) \neq 0, \exists p \in \mathbb{P}_{\mathbb{Z}}, \frac{st^n}{m^2} < p, p \nmid m, f \in V_n(\mathbb{F}_p) \}
$$

In this case, we first pick any a_0 , a_1 , \dots , a_{n-1} . F_n is not zero. We then pick a prime p such that $p|F_n(a_0, a_1, \dots, a_{n-1})$ and $p > \frac{st^n}{m^2}$. The size of F_n is less than $(st^n)^{k_n}$ for some k_n . Thus, the number of distinct prime divisors of F_n is

$$
\ll k_n \log(st^n).
$$

Since, if $p \nmid a_0$, then we can deduce from lemma [9](#page-22-0) that G_n is a polynomial in a_n of degree *n* − 1. If $p|a_0$, then $p \nmid a_1$ (as $(a_0, a_1) = 1$) and thus

$$
G_n(a_0,\cdots,a_n)\equiv a_1^2G_{n-1}(a_1,a_2,\cdots,a_n) \bmod p.
$$

This follows directly from applying lemma [9](#page-22-0) to the palindromic inverse of *f* . We basically do not have to worry about *Gⁿ* mod *p* being degenerate as a polynomial in *an*. Thus, there will be at most $n - 1$ possibilities for a_n as the range it moves through is of size less than p . Therefore, |*S*3| will be

$$
O(\frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2})\cdot \frac{m^2}{st^n}\cdot \log(st^n).
$$

 \Box

Definition 27. Let W_n denote the variety which is union of V_n with the variety given by $X_0 = X_1 = 0$. Let

$$
c_p:=1-\frac{|W_n(\mathbb{F}_p)|}{p^{n+1}}.
$$

Since, for every p we have polynomials with discriminant co-prime to p , c_p is never zero. Furthermore, since V_n is a variety contained in $V(F_n, G_n)$ of co-dimension 2 and $X_0 = X_1 = 0$ is also co-dimension 2 we can conclude that $c_p \geq (1 - \frac{\dot{r}_n}{p^2})$ $\binom{r_n}{p^2}$ for some constant $r_n.$ For $p \nmid m$, c_p will serve as the measure of how many polynomials mod p are not strongly divisible. The addition of this second variety will save us some trouble as the condition $(a_0, a_1) = 1$ will be taken care of internally.

7.2. **Actual sieve.**

Definition 28.

$$
W^*(s:t:m) = \{(f,l) \in W(s,t,m) : \forall p|m, f \notin V_n(\mathbb{F}_p)\}\
$$

Before moving forward, we refer to discussion [24.](#page-21-0) Let B_i denote boxes as there, and let

Definition 29.

$$
W^*(s:t:m)_{B_i} := W(s:t:m)_{B_i} \cap W^*(s:t:m)
$$

and

$$
W^*(s:t:m)_{B_i}^{UWD} := \{(f,l) \in W^*(s:t:m)_{B_i} : \forall p \in \mathbb{P}, f \notin V_n(\mathbb{F}_p)\}\
$$

and

$$
W^*(s:t:m:M)_{B_i} := \{(f,l) \in W(s:t:m)_{B_i}^*: \forall p < M, f \notin V_n(\mathbb{F}_p)\}.
$$

We begin with defining $M' = \prod_{p < M} p$. We will search for solutions to $\forall p < M, f \notin V_n(\mathbb{F}_p)$ by *p*∤*m* looking at the system modulo *M*′ . So we let

$$
C_{M'} = \prod_{\substack{p
$$

We will restrict ourselves to squarefree *m*. For *p*|*m*, we will look at *an*−² only and note that for any choice of a_0, a_1, \dots, a_{n-3} there are at most $n^{\omega(m)}$ possible solutions modulo *m* for a_{n-2} for *f* to be strongly divisible by *m*. See discussion [26.](#page-22-1)

Thus, making boxes of size *M*′ , we see that

$$
|W^*(s:t:m:M)_{B_i}|
$$

\n
$$
= C_M \cdot m \cdot (\frac{s}{2M'} + O(1))^2 \cdot (\frac{2st^2}{M'} + O(1)) \cdot \dots \cdot (\frac{2st^k}{M'} + O(1)) \cdot \dots \cdot (\frac{2st^{n-2}}{M'} + O(n^{\omega(m)}) + O(s\rho_B m))
$$

\n
$$
\cdot (\frac{2st^{n-1}}{M'm} + O(1)) \cdot (\frac{2st^n}{M'm^2} + O(1)) \cdot (M')^{n+1}.
$$

\nIf we have $M' \le \min\{s, \frac{st^n}{m^2}, \frac{t^{n-2}}{m}\}$, we get,
\n
$$
= 2^{n-3}Vol(B_i)C_M \cdot \frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2} (1 + O(\frac{m^2M'}{st^n}) + O(\frac{M'm}{t^{n-2}}) + O(\frac{M'}{s})).
$$

This will not be a problem as we will be taking *M*′ to be small.

We quickly write $|W^*(s:t:m)^{UWD}_{B_i}|$ as

 $|W^*(s:t:m:M)_{B_i}|+O(|\{(f,l)\in W(s:t:m):\exists p\in \mathbb{P}_{\mathbb{Z}},p>M,p\nmid m,f\in V_n(\mathbb{F}_p)\}|)$ Let $C = \lim_{M \to \infty} C_M$. Using theorem [35,](#page-23-0) we get

Theorem 36. If $s \gg \frac{st^n}{m^2}$, then

$$
\begin{aligned} |W^*(s:t:m)^{UWD}_{B_i}| \gg &2^{n-3}Vol(B_i)C \cdot \frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2}(1+O(\frac{m^2e^M}{st^n})+O(\frac{me^M}{t^{n-2}})) \\ &+ O(\frac{s^nt^{\frac{n(n+1)}{2}-1}}{m^2}) \cdot (\frac{1}{M\log M}+\frac{m^2\log(st^n)}{st^n}) \end{aligned}
$$

Setting $e^M = (st^n)^{\epsilon/2}$, we get the following corollary.

Corollary 2. *If* $(st^n)^{\epsilon} \ll s$ *then*

$$
|W^*(s:t:m)^{UWD}_{B_i}| \gg 2^{n-3}Vol(B_i)C \cdot \frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2}(1+O_{\epsilon}(\frac{1}{\log(st^n)})+O(\frac{m^2(st^n)^{\epsilon}}{st^n})+O(\frac{m(st^n)^{\epsilon}}{t^{n-2}})).
$$

Summing over all boxes B_i we get the following corollary.

Corollary 3. *If* $(st^n)^{\epsilon} \ll s$ then

$$
|W^*(s:t:m)| \gg (1-\epsilon)2^{n-3}C\frac{s^{n+1}t^{\frac{n(n+1)}{2}-1}}{m^2}(1+O_{\epsilon}(\frac{1}{\log(st^n)})+O(\frac{m^2(st^n)^{\epsilon}}{st^n})+O(\frac{m(st^n)^{\epsilon}}{t^{n-2}})).
$$

We set $X = \frac{s^{2n-2}t^{n(n-1)}}{m^2}$ $\frac{1}{m^2}$.

Let *S*(*X*, *s*, *m*) := $W^*(s : t : m)$.

Note that this set counts distinct rings.

It follows that

Theorem 37. *For* $n \geq 5$ *,*

$$
S(X,s,m) \gg_n \frac{X^{\frac{1}{2} + \frac{1}{n}} \cdot s^{\frac{2}{n}}}{m^{1 - \frac{2}{n}}}
$$
\n(18)

provided that

- *(1)* $s \gg_{\epsilon} (st^n)^{\epsilon}$
- *(2)* $X^{1-\epsilon} \gg_{\epsilon} s^{2n-2} \cdot m^{n-4-\frac{1}{n-2}}$ *(Condition corresponding to* $t^{n-2} \gg_{\epsilon} m^{1+\epsilon}$ required for injectivity).
- *(3)* $X^{1-\epsilon} \geq e^{s^{n-1}} \cdot m^{2n-4}$ *(Condition corresponding to* $(st^n)^{1-\epsilon} \geq e^{m^2}$ -required so the final coeffi*cient has enough space to very to have polynomials which are weakly divisible by m)*

To count ultra-weakly divisible rings which are weakly divisible by *m* rings, we wish to choose *s* appropriately. Note that within the constraints for a fixed *X*, the rings counted will all be distinct for any choice of *m* and then any choice of *s*.

We can chose *s* such that $st^n \simeq t^{2n-4}$. This will make the condition on *m* a singular condition. May not be optimal.

$$
st^n \simeq t^{2n-4} \iff s \simeq (m^2X)^{\frac{n-4}{(n-1)(3n-8)}}
$$

Both conditions combine to

$$
X^{1-\epsilon} \ge (m^2 X)^{\frac{n-4}{3n-8}} m^{2n-4} \iff X^{\frac{n-2}{3n^2-13n+12}-\epsilon} \ge m
$$

Thus, if we let *S*(*X* : *m*) denote the set of all UWD Rings, weakly divisible by *m*, then we have

$$
S(X:m) \gg \frac{X^{\frac{1}{2} + \frac{1}{n} + \frac{2(n-4)}{n(n-1)(3n-8)}} m^{\frac{2}{n} + \frac{4(n-4)}{n(n-1)(3n-8)}}
$$

$$
T(X) := \sum_{\substack{n-2 \ n \le X^{3n^2-13n+12}} -\epsilon} S(X,m) \gg \sum_{\substack{n-2 \ n \le X^{3n^2-13n+12}} -\epsilon} \frac{X^{\frac{1}{2} + \frac{1}{n} + \frac{2(n-4)}{n(n-1)(3n-8)}} m^{\frac{2}{n} + \frac{4(n-4)}{n(n-1)(3n-8)}}}{m}
$$

$$
\gg_n X^{\frac{1}{2} + \frac{1}{n} + \frac{2(n-4)}{n(n-1)(3n-8)}} (X^{\frac{n-2}{3n^2-13n+12}} - \epsilon)^{\frac{2}{n} + \frac{4(n-4)}{n(n-1)(3n-8)}}
$$

$$
\sim_{\epsilon} X^{\frac{1}{2} + \frac{1}{n - \frac{4}{3}} - \epsilon}
$$

Remark 38*.* We are summing over squarefree *m*.

7.3. **Number-field Counting Strategy.**

The strategy now comes into play by noting that

Theorem 39. *A UWD ring is a Restricted Sudo maximal Ring.*

Proof. Using Dedekind Kummer for Binary Rings i.e.theorem [12,](#page-9-0) we see that factorization of the corresponding UWD polynomial modulo *p* captures the irreducible decomposition of the associated binary ring. We see that if *f* is Ultra-weakly divisible then at every prime *p* it can never have 2 double roots or a triple root. This means that *f* can at most have one double root modulo *p*. The corresponding factor will correspond to either a maximal ring at *p* or an irreducible order with $ef(\mathcal{O}) = 2$. Thus, R_f is restricted Sudo maximal Ring. Recall that Sudo maximal rings were Rings whose theorem [7](#page-7-0) decomposition only consist of irreducible orders \mathcal{O} , satisfying $e f(\mathcal{O}) = 2$. Our description of Restricted Sudo Maximal Rings and Pseudo maximal rings immediately tells us that any intermediate Ring that sits in between a Restricted Sudo maximal Ring and its integral Closure(the ring of integers it sits in) is also a Restricted Sudo maximal Ring. \Box

Let $A(X)$ denote the set of all Restricted Sudo maximal Rings with discriminant $\leq X$. Let $a(X) :=$ $|A(X)|$. Thus, as a corollary we have

Corollary 4. $a(X) \geq T(X)$.

Let

$$
B(X) := \{ frac(R) : R \in A(X) \}.
$$

and $b(X) := |B(X)|$.

On the other hand, we may bound $a(X)$ above by using theorem [27](#page-17-0) as follows.

$$
a(X) \leq \sum_{\mathbb{K}\in B(X)} S_{\mathbb{K}}(X) \leq \sum_{\mathbb{K}\in B(X)} \sqrt{\frac{X}{D_{\mathbb{K}}}} \log(\frac{X}{D_{\mathbb{K}}})^{c_n}
$$

$$
\leq \sqrt{X} \sum_{\mathbb{K}\in B(X)} \frac{1}{\sqrt{D_{\mathbb{K}}}} \log(X)^{c_n}.
$$

Combine with the fact that

$$
a(X) \gg T(X) \gg X^{\frac{1}{2} + \frac{1}{n - \frac{4}{3}} - \epsilon}
$$

We get

$$
N^*(X:n) \ge \sum_{\mathbb{K}\in B(X)} \sqrt{\frac{1}{D_{\mathbb{K}}}} \gg X^{\frac{1}{n-\frac{4}{3}}-\epsilon}
$$

Corollary 5.

$$
\limsup_{X \to \infty} \frac{N(X:n)}{X^{\frac{1}{2} + \frac{1}{n - \frac{4}{3}} - \epsilon}} = \infty.
$$

Remark 40*.* We note that if the number of UWD polynomial tuples (f, m) with discriminant $\lt X$ is of the expected order then we can easily convert this to be an appropriate lower bound for $N(X:n)$.

APPENDIX A. **Defining Weakly Divisible Rings**

Most of this section is from our previous paper on Weakly Divisible Rings.

We recall the following theorem from [\[11\]](#page-39-7) defining Binary Rings.

Let

$$
f(X,Y) := a_n X^n + a_{n-1} X^{n-1} Y + \dots + a_0 Y^n
$$

denote a binary form of degree *n*. Let $a_n \neq 0$ and suppose that δ denotes the image of *X* in the algebra $\mathbb{Q}[X]/(f(X,1)).$

We recall eq. (3) .

Remark 41*.* We refer to the basis given for *R^f* in eq. [\(3\)](#page-10-1) as the canonical basis attached to *f* .

Theorem 42. *When f is integral(i.e. f is a binary form of degree n with integer coefficients), R^f is a ring of rank n over* **Z**.

Definition 30. *We define I^f as the (fractional) ideal class generated by* (1, *δ*) *over R^f* , *when f is integral.*

Definition 31. *When f is integral, R^f is known as the binary ring associated to the binary form f* .

These are a few properties of binary rings.

Proposition 1. *Properties of R^f (when f is integral):*

(1)

$$
\operatorname{disc}_{\mathbb{Z}}(R_f) = \operatorname{disc}(f). \tag{19}
$$

(2) If δ is invertible, and f is primitive, then

$$
R_f := \mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}]. \tag{20}
$$

- (3) If f is primitive, I_f is invertible in R_f .
- *(4) Both R_f and I_f are invariant under the natural* $GL_2(\mathbb{Z})$ *action on binary forms of degree n. In* particular, for $\delta\in\overline{\mathbb Q}\backslash\mathbb Q$ this means that if $\lambda=\frac{a\delta+b}{c\delta+d}$ with a, b, c, $d\in\mathbb Z$ and ad $-bc=\pm1$ then

$$
\mathbb{Z}[\delta] \cap \mathbb{Z}[\delta^{-1}] = \mathbb{Z}[\lambda] \cap \mathbb{Z}[\lambda^{-1}].
$$

You can find the proof of all of these statements in [\[11\]](#page-39-7).

Remark 43*.* Note that one can write down the multiplication table for the defining basis of *R^f* in terms of coefficients of *f* explicitly. Using this table to define binary rings, one can give a definition for R_f without the need for the condition " $a_n \neq 0$ ".

We will note the following part of the multiplication table of the canonical basis of *R^f* (eq. [\(3\)](#page-10-1)), which is easily verified.

Lemma 10. Let $f(X, Y) := a_n X^n + a_{n-1} X^{n-1} Y + \cdots + a_0 Y^n$ denote a binary form of degree n, and let $\langle B_0, B_1, \cdots, B_{n-1} \rangle$ denote the canonical basis for R_f associated to f as in eq. [\(3\)](#page-10-1). Then, we have

$$
B_{n-1} \cdot B_{n-i} = -a_0 \cdot B_{n-i-1} + a_i \cdot B_{n-1}
$$
 (21)

Proof. We make note of the fact, $B_{n-1} = -a_0 \delta^{-1}$. Thus,

$$
B_{n-1} \cdot B_{n-i} = -a_0 \delta^{-1} \cdot \left(\sum_{j=0}^{n-i} a_{n-j} \delta^j \right)
$$

= $-a_0 \cdot \sum_{j=0}^{n-i} a_{n-j} \delta^{j-1}$
= $-a_0 \cdot (B_{n-i-1} + \frac{a_i}{\delta})$
= $-a_0 \cdot B_{n-i-1} + a_i \cdot \left(\frac{-a_0}{\delta} \right)$
= $-a_0 \cdot B_{n-i-1} + a_i \cdot B_{n-1}$.

 \Box

Definition 32. *We say a binary form* $f(x, y) \in \mathbb{Z}[x, y]$ *of degree n is weakly divisible by* $m \in \mathbb{Z}_{>0}$ *if there exists an* $\ell \in \mathbb{Z}$ *such that*

$$
f(l, 1) \equiv 0 \mod m^2
$$

$$
\frac{\partial f}{\partial x}(l, 1) \equiv 0 \mod m
$$

When appropriate we say f is weakly divisible by m at l.

Remark 44*.* We use "weakly divisible" as these polynomials are defined in [\[3\]](#page-39-4) and [\[2\]](#page-39-3) and the main parts of these papers is about establishing an upper bound for the number of polynomials which are weakly divisible by *m* and bounded height, for all large *m*.

Definition 33. *Given a binary form f we set* $f_l = f_l(x, y) := f(x + ly, y)$ *.*

The condition that "f is weakly divisible by *m* at *l*" is equivalent to having integers a_0, a_1, \cdots, a_n such that

$$
f_1 = a_n x^n + a_{n-1} x^{n-1} y + \cdots + a_2 x^2 y^{n-2} + m a_1 x y^{n-1} + m^2 a_0 y^n.
$$

Definition 34. *We define,*

$$
R'_{(f,m)} := \mathbb{Z} \langle B_0, B_1, \cdots, B_{n-2}, \frac{B_{n-1}}{m} \rangle.
$$

Theorem 45. If f is weakly divisible by m at l, then $R'_{(f_l,m)}$ is a ring.

Proof. Let $\langle B_0, B_1, \cdots, B_{n-1} \rangle$ denote the canonical (old) basis of R_{f_l} associated to f_l . Since, f is weakly divisible by *m* at *l*, we can write

$$
f_1 = a_n x^n + a_{n-1} x^{n-1} y + \cdots + a_2 x^2 y^{n-2} + m a_1 x y^{n-1} + m^2 a_0 y^n,
$$

where a_i , $m \in \mathbb{Z}$.

We will show that $R'_{(f_l,m)}$ is a ring, by showing that product of any two elements in the (new) basis given by $\langle B_0, B_1, \cdots, B_{n-2}, \frac{B_{n-1}}{m}\rangle$ (basis for $R'_{(f_l,m)})$ is in the \Z -span of itself. We will achieve this by comparing the product of every two elements in the old basis with the product of the corresponding two elements in the new basis.)

We note that $B_i \cdot B_j$ for $i, j \neq n-1$ can be written as a \mathbb{Z} -linear combination of $\langle B_i \rangle$. This follows directly from the fact that the Z-span of $\langle B_i \rangle$ forms a ring (the prodigal binary ring) (we can also refer to the multiplication tables given in [\[9\]](#page-39-10) or section 2.1 in [\[11\]](#page-39-7)).

It immediately follows that $B_i \cdot B_j$ for $i, j \neq n-1$ can also be written as a \mathbb{Z} -linear combination of our new basis $\langle B_0, B_1, \cdots, B_{n-2}, \frac{B_{n-1}}{m_n} \rangle$ (which only differs from the original basis at the index *n* − 1) by simply replacing B_{n-1} with $m \cdot \frac{B_{n-1}}{m}$ in the multiplication table of the old basis.

On the other hand, lemma [10](#page-29-1) immediately tells us that, when $i \neq 1$

$$
\frac{B_{n-1}}{m} \cdot B_{n-i} = -ma_0 \cdot B_{n-i-1} + a_i \cdot \frac{B_{n-1}}{m}
$$

and

$$
\frac{B_{n-1}}{m} \cdot \frac{B_{n-1}}{m} = -a_0 \cdot B_{n-2} + a_1 \cdot \frac{B_{n-1}}{m}.
$$

Thus, the products of elements in this new basis are **Z**-linear combinations of the same new basis. It follows that $R'_{(f_l,m)}$ is in-fact a ring.

Definition 35. *We say R*′ (*fl* ,*m*) *is the weakly divisible ring (at l with respect to m) associated to f , when f is weakly divisible by m at l. When appropriate we will also represent this ring as R*′ (*f* ,*m*,*l*) .

Remark 46*.* Every binary ring is a weakly divisible ring at every value with respect to 1.

Remark 47*.* We note that weakly divisible rings may also defined by multiplication tables to avoid dependence on a condition like " $a_n \neq 0$ or $a_0 \neq 0$ ".

APPENDIX B. **Effective injectivity of the map:**

$$
(f,m,l)\longrightarrow R'_{(f_l,m)}
$$

B.1. **Matrix of transformation.** For convenience sake, we define $\binom{n}{a}$ $a_n^{\{n\}} := 0$ if $n \geq 0$ and $a < 0$.

We mention a useful matrix of transformation for base change from the canonical basis of *R^f* associated to $f(X, Y)$ to the canonical basis of R_{f_l} associated to f_l .

Theorem 48. We let $f(X,Y) = A_0X^n + A_1X^{n-1}Y + \cdots + A_nY^n$ and $\langle B_0, B_1, \cdots, B_{n-1} \rangle$ be the canonical basis of R_f associated to $f(X,Y)$ and $\langle C_0, C_1, \cdots, C_{n-1}\rangle$ be the canonical basis of R_{f_l} associated to f_l ,

then we have

$$
\langle C_0, C_1, \cdots, C_{n-1} \rangle = \langle B_0, B_1, \cdots, B_{n-1} \rangle \begin{bmatrix} 1 & \binom{n-1}{1}lA_0 & \binom{n-1}{2}l^2A_0 & \cdots & \binom{n-1}{k-1}l^{k-1}A_0 & \cdots & \binom{n-1}{n-1}l^{n-1}A_0 \\ 0 & 1 & \binom{n-2}{1}l & \cdots & \binom{n-2}{k-2}l^2 & \cdots & \binom{n-2}{n-2}l^{n-2} \\ 0 & 0 & 1 & \cdots & \binom{n-3}{k-3}l^{k-3} & \cdots & \binom{n-3}{n-3}l^{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \binom{n-k'}{k-k'}l^{k-k'} & \cdots & \binom{n-k'}{n-k'}l^{n-k'} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \binom{n}{k-n+1}l^{k-n+1} & \cdots & l \\ 0 & 0 & 0 & \cdots & \binom{n}{k-n}l^{k-n} & \cdots & 1 \end{bmatrix}
$$

B.2. **Proof.**

Let $F(x, y) =$ *n* ∑ *i*=0 $a_ix^{n-i}y^i$ denote an irreducible integral binary form. Let δ denote the root of $f(x) := F(x, 1)$. Let $F_l(x, y) := f(x + ly, y) =$ *n* ∑ *i*=0 $b_i x^{n-i} y^i$. Note that $\delta - l$ is the root of $f_l(x) := F_l(x, 1)$. Let $s_i(x) :=$ *i*−1 ∑ *j*=0 $a_j x^{i-j} = a_0 x^i + a_1 x^{i-1} + \dots + a_{i-1} x.$ Let $m_i(x) :=$ *i*−1 ∑ *j*=0 $b_j x^{i-j} = b_0 x^i + b_1 x^{i-1} + \ldots + b_{i-1} x.$ (22)

Thus the basis for R_{F_l} will be given by

 $\mathbb{Z}\langle 1, m_1(\delta-l)+b_1, m_2(\delta-l)+b_2, \cdots, m_k(\delta-l)+b_k, \cdots, m_{n-1}(\delta-l)+b_{n-1}\rangle$ and that of R_f will be given by

$$
\mathbb{Z}\langle 1, s_1(\delta)+a_1, s_2(\delta)+a_2, \cdots, s_k(\delta)+a_k, \cdots, s_{n-1}(\delta)+a_{n-1}\rangle
$$

Lemma 11.

$$
b_k = \sum_{i=0}^k {n-i \choose n-k} a_i
$$
 (23)

Proof.

$$
f(x+l) = \sum_{i=0}^{n} a_i (x+l)^{n-i} = \sum_{i=0}^{n} a_i \sum_{j=0}^{n-i} {n-i \choose j} x^j l^{n-i-j}
$$

=
$$
\sum_{i=0}^{n} \sum_{j=0}^{n-i} a_i {n-i \choose j} x^j l^{n-i-j} = \sum_{j=0}^{n} x^j \sum_{i=0}^{n-j} a_i {n-i \choose j} l^{n-i-j}
$$

Thus,

$$
b_k = \sum_{i=0}^k a_i {n-i \choose n-k} l^{k-i}
$$

 \Box

Lemma 12.

$$
m_k(x) + b_k = x(m_{k-1}(x) + b_{k-1}) + b_k
$$

$$
s_k(x) + a_k = x(s_{k-1}(x) + a_{k-1}) + a_k
$$

Proof. By definition. □

Proposition 2.

$$
m_k(x-l) + b_k = \sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j})l^j + {n-1 \choose k} l^k a_0
$$
 (24)

Proof. We proceed by induction on *k*,

Base Case. $m_1(x-l) = b_0(x-l) + b_1 = (a_0)(x-l) + a_1 + na_0l$ $= (a_0x + a_1) + \binom{n-1}{1}a_0l$ $= {\binom{n-2}{0}}(s_1(x) + a_1) + {\binom{n-1}{1}}a_0$

Induction Hypothesis:

$$
m_k(x-l) + b_k = \sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j})l^j + {n-1 \choose k} l^k a_0
$$

We have from lemma [12,](#page-31-0)

$$
m_{k+1}(x-l) = (x-l)(m_k(x-l) + b_k).
$$

Substituting in our induction hypothesis, we get

$$
m_{k+1}(x-l)
$$
\n
$$
= (x-l)(\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j})l^j + {n-1 \choose k} l^k a_0)
$$
\n
$$
= \sum_{j=0}^{k-1} {n-k+j-1 \choose j} ((s_{k-j}(x) + a_{k-j})l^j x - (s_{k-j}(x) + a_{k-j})l^{j+1}) + {n-1 \choose k} a_0 l^k x - {n-1 \choose k} l^{k+1} a_0
$$
\n
$$
= (\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) x + a_{k-j} x)l^j + {n-1 \choose k} a_0 x l^k)
$$
\n
$$
- (\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j})l^{j+1} + {n-1 \choose k} a_0 l^{k+1})
$$
\n
$$
= (\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j+1}(x))l^j + {n-1 \choose k} s_1 (x) l^k)
$$
\n
$$
- (\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j})l^{j+1} + {n-1 \choose k} a_0 l^{k+1})
$$
\n
$$
= (\sum_{j=0}^{k} {n-k+j-1 \choose j} (s_{k-j+1}(x))l^j - (\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j})l^{j+1} + {n-1 \choose k} a_0 l^{k+1})
$$

$$
= (\sum_{j=0}^{k} {n-k+j-1 \choose j} (s_{k-j+1}(x) + a_{k-j+1}) U - (\sum_{j=0}^{k-1} {n-k+j-1 \choose j} (s_{k-j}(x) + a_{k-j}) U^{j+1})
$$

\n
$$
- {n-1 \choose k} a_0 I^{k+1} - \sum_{j=0}^{k-1} {n-k+j-1 \choose j} a_{k-j+1} U^{j+1}
$$

\n
$$
= \sum_{j=0}^{k} (n-k+j-1) - {n-k+j-2 \choose j-1} (s_{k+1-j} + a_{k+1-j}) U^{j})
$$

\n
$$
- {n-1 \choose k} a_0 I^{k+1} - \sum_{j=0}^{k-1} {n-k+j-1 \choose j} a_{k-j+1} U^{j+1}
$$

\n
$$
= (\sum_{j=0}^{k} {n-k-2+j \choose j} (s_{k+1-j} + a_{k+1-j}) U^{j}) - {n-1 \choose k} a_0 I^{k+1} - \sum_{j=0}^{k} {n-k+j-1 \choose j} a_{k-j+1} U^{j+1}
$$

\n
$$
= (\sum_{j=0}^{k} {n-k-2+j \choose j} (s_{k+1-j} + a_{k+1-j}) U^{j} + {n-1 \choose k+1} a_0 I^{k+1})
$$

\n
$$
= (\sum_{j=0}^{k} {n-k-2+j \choose j} (s_{k+1-j} + a_{k+1-j}) U^{j} + {n-1 \choose k+1} a_0 I^{k+1})
$$

\n
$$
= (\sum_{j=0}^{k} {n-k+1 \choose j} (s_{k+1-j} + a_{k+1-j}) U^{j} + {n-1 \choose k+1} a_0 I^{k+1})
$$

\n
$$
= (\sum_{j=0}^{k} {n-k+2+j \choose j} (s_{k+1-j} + a_{k+1-j}) U^{j} + {n-1 \choose k+1} a_0 I^{k+1})
$$

\n
$$
= (\sum_{j=0}^{k} {n-k+2+j \choose j} (s_{k+1-j} + a_{k+1-j}) U^{j} + {n-1 \choose k+1} a_0 I^{k+1})
$$

\n
$$
= (\sum
$$

 \Box

Substituting *x* = δ in the above Proposition we get the result.

B.3. Given *f* a binary form, we let $\langle B_0, B_1, \cdots, B_{n-1} \rangle$ be the canonical basis of R_f associated to *f* as in eq. [\(3\)](#page-10-1). Let $\langle v_1, v_2, \cdots, v_n \rangle$ denote a real basis for \mathbb{R}^n . We perform a Gram-Schmidt reduction

on the basis using some canonical distance form. We write

$$
\langle v_1, v_2, \cdots, v_n \rangle = \langle B \rangle M[(t_1, t_2, ..., t_n)]
$$

where *B* is an orthonormal ordered basis of vectors, all of which are of the same size. *M* is an upper triangular uni-potent matrix. And $[(t_1, t_2, ..., t_n)]$ denotes a diagonal matrix with t_i placed in the (i, i) th place.

Let v'_i denote the projection of v_i to the space orthogonal to the space spanned by $\{v_1, v_2, \cdots, v_{i-1}\}.$ Then, Gram-Schmidt process forces

$$
|t_i| = ||v_i'||. \t\t(25)
$$

We note that for each $1 \le i \le n$, and any choice of coefficients a_1, \dots, a_i we have

$$
||a_i v_i + \sum_{j=1}^{i-1} a_j v_j|| \geq |a_i||t_i|.
$$

This follows by looking at the component of the given vector (on RHS) along v_i' . Moreover, for each 1 ≤ *i* ≤ *n*, there exist b_1 , \dots , b_{i-1} ∈ **Z** so that

$$
||v_i + \sum_{j=1}^{i-1} b_j v_j|| \leq |t_i| + |t_{i-1}| + \cdots + |t_1|.
$$

One can simply choose b_j such that the $\sum_{j=1}^{i-1} b_j v_j$ approximates the vector $v_i - v'_i$ which clearly lies in the space spanned by $\langle v_1, v_2, \cdots, v_{i-1}\rangle$.

Now if $t_{i+1}/t_i > 2$ for all *i*, one can see that inductively the i^{th} vector in the Minkowski reduced basis of the lattice spanned by $\{v_1,v_2\cdots,v_n\}$ will have the form $\pm v_i+\sum_{j=1}^{i-1}b_jv_j.$

In fact, having

$$
t_i/t_{i-1} > \sqrt{1 + \frac{1}{i}} \text{ for all } 1 \le i \le n
$$

is sufficient to conclude this.

Furthermore, if we just know that,

$$
\frac{t_i}{t_2} \ge 2 \text{ for all } 3 \le i \le n \text{ and } \frac{t_2}{t_1} \ge 2 \tag{26}
$$

then, the first two vectors (which will be unique up to sign) in the Minkowski reduced basis for the lattice $\langle v_1, v_2, \cdots, v_n \rangle$ will have the above form. That is

$$
\{\pm v_1, \pm (v_2 + a \cdot v_1)\}\
$$

will be the first two elements in the Minkowski reduced basis for this lattice spanned by $\langle v_1, v_2, \cdots, v_n \rangle$.

Definition 36. If $\langle v_1, v_2, \cdots, v_n \rangle$ is a basis for \mathbb{R}^n and v'_i denotes the projection of v_i to the space orthogonal *to the space spanned by* $\{v_1, v_2, \cdots, v_{i-1}\}$ *and* $t_i = ||v'_i||$ *, then we say* $\langle v_1, v_2, \cdots, v_n \rangle$ *is* Normally Minkowski Reduced *if tⁱ satisfy*

$$
\frac{t_i}{t_2} \ge 2 \text{ for all } 3 \le i \le n \text{ and } \frac{t_2}{t_1} \ge 2 \tag{27}
$$

Our discussion above gives us the following lemma:

Lemma 13. *If* $\langle B_1, B_2, \cdots, B_n \rangle$ *is* Normally Minkowski Reduced, then

- *(1)* B_1 *is the unique smallest vector in L up to sign.*
- *(2)* $∃a ∈ Z$ *such that,* $B_2 + a ⋅ B_1$ *is the smallest vector in L which is not in* $Z ⋅ B_1$ *up to sign.*

B.4. **Reduced-m-Polynomials.**

Definition 37. *We say a tuple* (*f* , *m*, *l*) *is a Reduced-m-Polynomial (at l) if*

 \bullet The canonical basis for $R'_{(f_l,m)}$ is Normally Minkowski Reduced when seen in $R'_{(f_l,m)}\otimes R\simeq R^r\oplus R^r$ **C** *^s under the canonical norm.*

Remark 49. We will ignore the *l* part as (f, m, l) is a Reduced-*m*-polynomial (at *l*) \iff (f, m, r) is a Reduced-*m*-Polynomial (at *r*) for any real value *r*. This is easy to see from the fact that the matrix in theorem [48](#page-30-0) is upper triangular and hence will leave relative sizes of normal components unchanged.

Theorem 50. *Given f, g polynomials of degree* $n \geq 4$ *such that f is a Reduced-m-polynomial and f is weakly divisible by m at e and g is a Reduced-m*′ *-polynomial and g is weakly divisible by m*′ *at d (m*, *m*′ ≥ 1*)* w ith $R'_{(f_e,m)} = R'_{(g_d,m)}$ then

$$
m = m' \text{ and } \exists r \in \mathbb{Z} : g(x) = f(x + mr - d + e).
$$

Proof. Let β and α denote roots of f and g respectively ($\in \mathbb{R}^r \oplus \mathbb{C}^s$). Let a_0 and b_0 denote the positive leading coefficients of *f* and *g* and *a*¹ and *b*¹ denote the second leading coefficient of *f* and *g* respectively.

We know that the smallest two elements of any Normally Minkowski Reduced basis are unique. Since $R'_{(f_e,m)}=R'_{(g_d,m')}$, comparing the smallest two elements elements in these, we get

$$
\langle 1, a_0 \cdot \beta \rangle \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} = \langle 1, b_0 \cdot \alpha \rangle
$$

where $a = \pm 1$ and $c = \pm 1$. Clearly, $a = 1$ and thus it follows that

$$
\alpha = \frac{\pm a_0}{b_0} \cdot \beta + \frac{b}{b_0}.
$$

$$
\frac{f(\frac{\pm a_0}{b_0} \cdot x + \frac{b}{b_0})}{a_0} = (\pm \frac{a_0}{b_0})^n \frac{g(x)}{b_0}.
$$

This immediately tells us that the matrix of transfer from $\langle 1, a_0\beta, a_0\beta^2 + a_1\beta \rangle$ to $\langle 1, b_0\alpha, b_0\alpha^2 + b_1\alpha \rangle$ is

$$
\begin{bmatrix} 1 & b & * \\ 0 & c & * \\ 0 & 0 & \frac{a_0}{b_0} \end{bmatrix}.
$$

Since $n \geq 4$ the above matrix must also be integral and invertible. It follows that $a_0 = b_0$ and thus $\mathfrak{disc}(f) = \mathfrak{disc}(g)$.

 $\text{Since } R'_{(f_e,m)} = R'_{(g_d,m')}$ we also have

$$
\frac{\operatorname{Disc}(f)}{m^2} = \frac{\operatorname{Disc}(g)}{m'^2},
$$

and we immediately get $m = m'$.

Now, without loss of generality, we may assume $c = 1$, for if $c = -1$ we may change $g(x)$ to $(-1)^n g(-x)$. We let *l* = $\frac{b}{b_0}$ ∈ **Q**. Thus, $f(x+l) = g(x)$.

We make note of the fact that *f* and *g* are weakly divisible by the same value *m*.

Observing the matrix of transformation from R_{f_e} to R_{g_d} given using ??, we notice that the entry in the $(n-1)$ th row and n th column of the matrix of transfer for the canonical bases of R_{f_e} to R_{g_d} is $d - e + l$.

Thus, the entry in the $(n-1)$ th row and n^{th} column of the matrix of transfer for the canonical bases of $R'_{(f_e,m)}$ to $R'_{(g_d,m)}$ is $\frac{d-e+1}{m}$. Thus $r = \frac{d-e+1}{m}$ must be an integer. It follows that $\exists r \in \mathbb{Z} : g(x) =$ $f(x + mr - d + e)$.

Lemma 14. *If f is a real monic polynomial, then there exists a* $\rho_f \in \mathbb{R}$ *(continuously varying with f) such that* $\lambda \rho^n f(\frac{x}{\rho})$ $\frac{x}{\rho}$) *is a Reduced-*1-polynomial for all $\rho \ge \rho_f$ and $\lambda \ge 1$.

 $\int\int_{0}^{x} \rho \geq m^{1/(n-2)} \rho_{f}$ and $\lambda \geq 1$ the polynomial $\lambda \rho^{n} f(\frac{x}{\rho})$ *ρ*) *is a Reduced-m-polynomial.*

Proof. We follow the argument of Bhargava-Shankar-Wang proving in Lemma 5.2 in[\[2\]](#page-39-3). Set

$$
f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_kx^{n-k} + \dots + a_0
$$

and set $a_n = 1$.

We perform a Gram-Schmidt reduction of the basis for *R^f* using the canonical distance form on $R_f \otimes \mathbb{R} \simeq \mathbb{R}^r \oplus \mathbb{C}^s (r + 2s = n)$. We write

$$
\langle 1, \delta, \delta^2 + a_{n-1}\delta, \cdots, \sum_{i=0}^{k-1} a_{n-i}\delta^{k-i}, \cdots, \sum_{i=0}^{n-2} a_{n-i}\delta^{k-i} \rangle = \langle B \rangle M[(t_1, t_2, ..., t_n)]
$$

where *B* is an orthonormal ordered basis of vectors, *M* is an upper triangular unipotent matrix, and $[(t_1, t_2, ..., t_n)]$ denotes a diagonal matrix with those entries along the diagonal. Now we note that if

$$
\langle 1, \delta, \delta^2 + a_{n-1}\delta, \cdots, \sum_{i=0}^{k-1} a_{n-i}\delta^{k-i}, \cdots, \sum_{i=0}^{n-2} a_{n-i}\delta^{n-1-i} \rangle
$$

= $\langle B \rangle M[(t_1, t_2, \cdots, t_k, \cdots, t_n)]$

then, using eq. [\(25\)](#page-34-0), we see that

$$
\langle 1, \lambda(\rho\delta), \lambda((\rho\delta)^2 + a_{n-1}\rho\delta), \cdots, \lambda(\sum_{i=0}^{k-1} a_{n-i}(\rho\delta)^{k-i}), \cdots, \lambda(\sum_{i=0}^{n-2} a_{n-i}(\rho\delta)^{n-1-i}) \rangle
$$

= $\langle B \rangle M_\rho [(t_1, \lambda \rho t_2, \cdots, \lambda \rho^{k-1} t_k, \dots, \lambda \rho^{n-1} t_n)].$

Thus, for each polynomial *f* one may find ρ_f which is a continuous function of *f* such that $\lambda \rho^n f(\frac{x}{\rho})$ *ρ*) is Minkowski reduced for all $\lambda \geq 1$ and $\rho \geq \rho_f.$ One can simply take

$$
\rho_f := \max\{\max_{i\geq 3}\{(\frac{2t_2}{t_i})^{1/(i-2)}\}, \frac{2t_1}{t_2}\}.
$$

Furthermore, we note that translating the polynomial changes the canonical basis by an upper triangular matrix.

Thus, if $g(x) = \lambda \rho^n f(\frac{X}{\rho})$ $\frac{\lambda}{\rho}$), then the canonical basis of $R'_{(g_l,m)}$ after Gram-Schmidt process will look like *λρn*−¹

$$
\langle B \rangle M_{\rho,m,l}[(t_1,\lambda \rho t_2,\cdots,\lambda \rho^{k-1} t_k,\cdots,\lambda \rho^{n-2} t_{n-1},\frac{\lambda \rho^{n-1}}{m} t_n)]
$$

where *Mρ*,*m*,*^l* is a uni-potent upper triangular matrix.

 I t follows that for $\rho \geq m^{1/(n-2)}\rho_f$ and $\lambda \geq 1$, $\lambda \rho^n f(\frac{X}{\rho})$ $\frac{X}{\rho}$) is Reduced-*m*-polynomial.

APPENDIX C. **The structure of the discriminant polynomial**

Let

$$
f(X,Y) := a_0 X^n + a_1 X^{n-1} Y + \cdots + a_n Y^n
$$

denote a binary form of degree *n*. Let $a_n \neq 0$. Let

$$
\mathfrak{disc}(a_0,a_1,\cdots,a_n):=\mathfrak{disc}(f)
$$

denote the discriminant as a polynomial in the coefficients of *f* . Then, we know that

$$
a_0\text{disc}(f) = \text{Res}(f, \frac{\partial}{\partial X}f).
$$

Since, the resultant can be seen as the determinant of the Sylvester matrix. The Sylvester matrix for these polynomials is

The following lemmata are easy to observe from the following:

Lemma 15.

$$
\mathfrak{disc}(f) = a_n \Delta_1 + a_n a_{n-1} \Delta_2 + a_{n-1}^2 \Delta_3 + a_n^2 \Delta_4.
$$

where

and

$$
\Delta_1 = 4a_{n-2}^3 \cdot \mathfrak{disc}(a_0 X^{n-2} + a_1 X^{n-3} Y + \cdots + a_{n-2} Y^{n-2})
$$

$$
\Delta_3 = a_{n-1}^2 \text{disc}(a_0 X^{n-1} + a_1 X^{n-2} Y + \cdots + a_{n-2} XY^{n-2} + a_{n-1} Y^{n-1}).
$$

Proof. We carefully expand the determinant of the Sylvester Matrix along the last two columns. We can clearly see we can write the determinant as

$$
2a_n a_{n-2} \Delta_1 + a_{n-1}^2 \Delta_2 + a_n a_{n-1} \Delta_3 + a_n^2 \Delta_4.
$$
 (29)

We will pick $\Delta_1 \in \mathbb{Z}[a_0, a_1, \cdots, a_{n-2}]$ and as the components of Δ_1 containing a_{n-1} and a_n can be pushed inside Δ_2 , Δ_3 , Δ_4 . One can then choose $\Delta_3 \in \mathbb{Z}[a_0, a_1, \cdots, a_{n-1}]$ as components of Δ_2 containing a_n can be pushed inside Δ_2 and Δ_4 .

Then, Δ_1 can be seen as the determinant of (i.e. the minor corresponding to $a_n \cdot (2a_{n-2})$ where a_n and a_{n-1} are set to zero looks as follows)

Now doing operations $R_n - R_1$, $R_{n+1} - R_2$, $R_{n+1} - R_3$, \cdots , $R_{2n-2} - R_{n-2}$ we get the following matrix with the same determinant.

We note that the above matrix is very close to Sylvester matrix for computing the discriminant of

$$
\frac{f(x,y)-a_nY^n-a_{n-1}XY^{n-1}}{Y}=X(a_0X^{n-2}+a_1X^{n-3}Y+\cdots+a_{n-2}Y^{n-2}).
$$

More particularly, the determinant of the above matrix is twice of the discriminant of the above form. This can be easily observed by expanding the determinant of the above matrix and that of the Sylvester Matrix along the final column.

Clearly, since a_{n-2} is the Resultant of *X* and $a_0X^{n-2} + a_1X^{n-3}Y + \cdots + a_{n-2}Y^{n-2}$, we get

$$
\mathfrak{disc}(\frac{f(x,y)-a_nY^n-a_{n-1}XY^{n-1}}{Y})=a_{n-2}^2\cdot\mathfrak{disc}(a_0X^{n-2}+a_1X^{n-3}Y+\cdots+a_{n-2}Y^{n-2}).
$$

Substituting back into eq. [\(29\)](#page-37-2) we get the appropriate structure for Δ_1 .

.

.

To get the structure of Δ_3 we see that

$$
a_{n-1}^2 \Delta_3 = \mathfrak{disc}(f(X,Y) - a_n Y^n)
$$

by definition.

REFERENCES

- [1] T. Z. Avner Ash, Jos Brakenhoff. Equality of polynomial and field discriminants. *Experimental Mathematics*, 16(3):367–374, 2007.
- [2] M. Bhargava, A. Shankar, and X. Wang. Squarefree values of polynomial discriminants I. *Invent. Math.*, 228(3):1037– 1073, 2022.
- [3] M. Bhargava, A. Shankar, and X. Wang. Squarefree values of polynomial discriminants II, 2022.
- [4] I. D. Corso, R. Dvornicich, and D. Simon. Decomposition of primes in non-maximal orders. *Acta Arithmetica*, 120:231–244, 2005.
- [5] J. S. Ellenberg and A. Venkatesh. The number of extensions of a number field with fixed degree and bounded discriminant. *Annals of Mathematics*, 163:723–741, 2003.
- [6] N. Kaplan, J. Marcinek, and R. Takloo-Bighash. Distribution of orders in number fields. *Research in the Mathematical Sciences*, 2:1–57, 2014.
- [7] G. Malle. On the distribution of galois groups. *Journal of Number Theory*, 92(2):315–329, 2002.
- [8] G. Malle. On the Distribution of Galois groups, II. *Experimental Mathematics*, 13(2):129 – 136, 2004.
- [9] G. D. Patil. *Rings of finite rank over integers,PhD Thesis-University of Toronto*. 2023.
- [10] G. D. Patil. Weakly divisible rings, 2024.
- [11] M. M. Wood. Rings and ideals parameterized by binary *n*-ic forms. *Journal of the London Mathematical Society*, 83(1):208–231, 2011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, BAHEN CENTRE, 40 ST. GEORGE STREET, ROOM 6290, TORONTO, ONTARIO, CANADA, M5S 2E4

Email address: g.patil@mail.utoronto.ca