

LARGE-DEGREE ASYMPTOTIC EXPANSIONS FOR THE JACOBI AND ALLIED FUNCTIONS

GERGŐ NEMES

Dedicated to Roderick S. C. Wong on the occasion of his 80th birthday

ABSTRACT. Simple asymptotic expansions for the Jacobi functions $P_\nu^{(\alpha,\beta)}(z)$ and $Q_\nu^{(\alpha,\beta)}(z)$ for large degree ν , with fixed parameters α and β , are surprisingly rare in the literature, with only a few special cases covered. This paper addresses this notable gap by deriving simple (inverse) factorial expansions for these functions, complemented by explicit and computable error bounds. Additionally, we provide analogous results for the associated functions $Q_\nu^{(\alpha,\beta)}(x)$ and $Q_\nu^{(\alpha,\beta)}(x)$.

1. INTRODUCTION AND MAIN RESULT

In this paper, we derive asymptotic expansions for the Jacobi functions $P_\nu^{(\alpha,\beta)}(z)$ and $Q_\nu^{(\alpha,\beta)}(z)$, as well as for the associated functions $Q_\nu^{(\alpha,\beta)}(x)$ and $Q_\nu^{(\alpha,\beta)}(x)$, in the regime where the degree ν becomes large while α and β are held fixed. These functions are solutions to the Jacobi differential equation

$$(1.1) \quad (1 - z^2) \frac{d^2 w(z)}{dz^2} + (\beta - \alpha - z(\alpha + \beta + 2)) \frac{dw(z)}{dz} + \nu(\nu + \alpha + \beta + 1)w(z) = 0,$$

which is characterised by regular singularities at $z = \pm 1$ and $z = \infty$.

Over the past several decades, significant effort has been dedicated to deriving asymptotic expansions for these functions. Generally, there are two types of expansions: those in terms of elementary functions and uniform expansions involving Bessel functions (a brief summary is provided at the end of this section). A common feature of these results is the complexity of their coefficients. In contrast, the expansions we present in this paper are simple (inverse) factorial expansions with elementary coefficients. Additionally, in Section 2, we provide explicit and computable error bounds for these new expansions under certain conditions on the degree ν and the parameters α and β . It is important to note that, unlike their uniform counterparts, our expansions break down near the transition points $z = \pm 1$ of (1.1). Surprisingly, these simple expansions have not been extensively documented in the literature, except for a few special cases. This omission may be because previous researchers primarily focused on deriving uniform asymptotic results, which naturally possess a more complicated structure. We note that (inverse) factorial series were recently obtained for the associated Legendre and Ferrers functions in [22].

We start by defining the functions under consideration. The Jacobi functions of the first and second kind, with complex degree ν and parameters α and β , are defined in terms of the (regularised) hypergeometric function as follows (cf. [5], [12, Eq. (10.8.18)]):

$$(1.2) \quad P_\nu^{(\alpha,\beta)}(z) = \frac{\Gamma(\nu + \alpha + 1)}{\Gamma(\nu + 1)} \mathbf{F} \left(\begin{matrix} -\nu, \nu + \alpha + \beta + 1 \\ \alpha + 1 \end{matrix}; \frac{1 - z}{2} \right)$$

2020 *Mathematics Subject Classification.* 41A60, 33C05, 33C45.

Key words and phrases. asymptotic expansions, error bounds, Jacobi functions, hypergeometric function.

for $z \in \mathbb{C} \setminus (-\infty, -1]$ and $\nu + \alpha \notin \mathbb{Z}^-$, and

$$(1.3) \quad Q_\nu^{(\alpha, \beta)}(z) = \frac{2^{\nu+\alpha+\beta} \Gamma(\nu + \alpha + 1) \Gamma(\nu + \beta + 1)}{(z-1)^{\nu+\alpha+1} (z+1)^\beta} \mathbf{F} \left(\nu + 1, \nu + \alpha + 1; \frac{2}{1-z} \right)$$

for $z \in \mathbb{C} \setminus (-\infty, 1]$ and $\nu + \alpha, \nu + \beta \notin \mathbb{Z}^-$. For analytic continuation to other Riemann sheets, see [8]. In the case where $\nu = n$ is a non-negative integer, $P_n^{(\alpha, \beta)}(z)$ corresponds to the classical Jacobi polynomials. We will later use the following relation between the two types of Jacobi functions:

$$(1.4) \quad P_\nu^{(\alpha, \beta)}(x) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{\pi} \left(e^{\pi\alpha i} Q_\nu^{(\alpha, \beta)}(x + i\varepsilon) - e^{-\pi\alpha i} Q_\nu^{(\alpha, \beta)}(x - i\varepsilon) \right),$$

which holds for $-1 < x < 1$ and $\nu + \alpha, \nu + \beta \notin \mathbb{Z}^-$ [9, Eq. (2.3)]. The literature offers (at least) two distinct definitions for extending $Q_\nu^{(\alpha, \beta)}(x)$ into the interval $-1 < x < 1$. Szegő defines the associated function $\mathbf{Q}_\nu^{(\alpha, \beta)}(x)$, with complex degree ν and parameters α and β , as follows (cf. [12, Eq. (10.8.22)] and [26, Eq. (4.62.9)]):

$$(1.5) \quad \mathbf{Q}_\nu^{(\alpha, \beta)}(x) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \left(Q_\nu^{(\alpha, \beta)}(x + i\varepsilon) + Q_\nu^{(\alpha, \beta)}(x - i\varepsilon) \right),$$

for $-1 < x < 1$ and $\nu + \alpha, \nu + \beta \notin \mathbb{Z}^-$. Alternatively, Durand [9, Eq. (2.4)] defines the associated function $Q_\nu^{(\alpha, \beta)}(x)$ as

$$(1.6) \quad Q_\nu^{(\alpha, \beta)}(x) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \left(e^{\pi\alpha i} Q_\nu^{(\alpha, \beta)}(x + i\varepsilon) + e^{-\pi\alpha i} Q_\nu^{(\alpha, \beta)}(x - i\varepsilon) \right),$$

for $-1 < x < 1$ and $\nu + \alpha, \nu + \beta \notin \mathbb{Z}^-$. Both functions are real when ν, α and β are real.

The associated Legendre functions $P_\nu^\mu(z)$ and $Q_\nu^\mu(z)$ are related to the Jacobi functions by the expressions

$$P_\nu^\mu(z) = \frac{\Gamma(\nu + 1)}{\Gamma(\nu - \mu + 1)} \left(\frac{z+1}{z-1} \right)^{\mu/2} P_\nu^{(-\mu, \mu)}(z),$$

using (1.2) and [23, Eq. (14.3.6)], and

$$(1.7) \quad Q_\nu^\mu(z) = e^{\pi i \mu} \frac{\Gamma(\nu + 1)}{\Gamma(\nu - \mu + 1)} \left(\frac{z-1}{z+1} \right)^{\mu/2} Q_\nu^{(\mu, -\mu)}(z),$$

using (1.3), [23, Eq. (14.3.10)], and [23, Eq. (14.3.19)]. The Ferrers functions $P_\nu^\mu(x)$ and $Q_\nu^\mu(x)$ are similarly related to the Jacobi function of the first kind and the associated function $\mathbf{Q}_\nu^{(\alpha, \beta)}(x)$ as follows:

$$P_\nu^\mu(x) = \frac{\Gamma(\nu + 1)}{\Gamma(\nu - \mu + 1)} \left(\frac{1+x}{1-x} \right)^{\mu/2} P_\nu^{(-\mu, \mu)}(x),$$

using (1.2) and [23, Eq. (14.3.1)], and

$$Q_\nu^\mu(x) = \frac{\Gamma(\nu + 1)}{\Gamma(\nu - \mu + 1)} \left(\frac{1-x}{1+x} \right)^{\mu/2} Q_\nu^{(\mu, -\mu)}(x),$$

by combining (1.5), (1.7), and [22, Eq. (8.7)]. Thus, the results of this paper can also be directly applied to these functions. However, the resulting expansions differ from those given in [22].

Before presenting our main results, we first establish some notation. For any non-negative integer n and $\mu \in \mathbb{C}$, we define

$$(1.8) \quad a_n(\mu) = \frac{(4\mu^2 - 1^2)(4\mu^2 - 3^2) \cdots (4\mu^2 - (2n-1)^2)}{8^n n!} = \frac{(\frac{1}{2} + \mu)_n (\frac{1}{2} - \mu)_n}{(-2)^n n!},$$

where $(w)_n = \Gamma(w+n)/\Gamma(w)$ denotes the Pochhammer symbol. We will use $B(w_1, w_2)$ to denote the beta function and \mathcal{D} to represent the domain

$$(1.9) \quad \mathcal{D} = \left\{ \xi : \operatorname{Re}(\xi) > 0, |\operatorname{Im}(\xi)| < \frac{\pi}{2} \right\}.$$

Note that $\cosh(2\xi)$ is a biholomorphic bijection between \mathcal{D} and $\mathbb{C} \setminus (-\infty, 1]$. We further define, for $\xi \in \mathcal{D}$ and $\alpha \in \mathbb{C}$,

$$(1.10) \quad C(\xi, \alpha) = \begin{cases} \sin(\pi\alpha) & \text{if } \xi > 0, \\ \mp i e^{\pm\pi i\alpha} & \text{if } 0 < \pm \operatorname{Im}(\xi) < \frac{\pi}{2}. \end{cases}$$

The following two theorems offer inverse factorial expansions for the Jacobi and associated functions.

Theorem 1.1. *Assume that $\xi \in \mathcal{D}$, that $|\sinh \xi|, |\cosh \xi| \geq \varepsilon (> 0)$, and that $\alpha, \beta \in \mathbb{C}$ are bounded. Under these conditions, the Jacobi functions admit the following inverse factorial expansions:*

$$(1.11) \quad 2^{2\nu+\alpha+\beta+1} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha, \beta)}(\cosh(2\xi)) \sim \\ \left(\frac{e^\xi}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^\xi}{\cosh \xi} \right)^{\beta+1/2} e^{2\nu\xi} \sum_{n=0}^{\infty} g_n(-\xi, \alpha, \beta) \frac{\Gamma(2\nu+\alpha+\beta-n+1)}{\Gamma(2\nu+\alpha+\beta+2)} \\ - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} \sum_{m=0}^{\infty} g_m(\xi, \alpha, \beta) \frac{\Gamma(2\nu+\alpha+\beta-m+1)}{\Gamma(2\nu+\alpha+\beta+2)},$$

as $\nu \rightarrow \infty$ within the sector $|\arg \nu| \leq \frac{\pi}{2} - \delta (< \frac{\pi}{2})$, and

$$(1.12) \quad \frac{2^{2\nu+\alpha+\beta+1}}{\pi} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha, \beta)}(\cosh(2\xi)) \sim \\ \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} \sum_{n=0}^{\infty} g_n(\xi, \alpha, \beta) \frac{\Gamma(2\nu+\alpha+\beta-n+1)}{\Gamma(2\nu+\alpha+\beta+2)},$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \pi - \delta (< \pi)$. The coefficients $g_n(\pm\xi, \alpha, \beta)$ are given by

$$g_n(\pm\xi, \alpha, \beta) = \sum_{\ell=0}^n (\pm 1)^\ell a_\ell(\alpha) a_{n-\ell}(\beta) \left(\frac{e^{\pm\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{\pm\xi}}{\cosh \xi} \right)^{n-\ell}.$$

The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere.

Theorem 1.2. *Suppose that $(0 <) \varepsilon \leq \zeta \leq \frac{\pi}{2} - \varepsilon (< \frac{\pi}{2})$, and let $\alpha, \beta \in \mathbb{C}$ be bounded. Define*

$$(1.13) \quad \zeta_{\nu, n, \ell}^{(1)} = (2\nu + \alpha + \beta - n + 1)\zeta - \left(\alpha - \ell + \frac{1}{2}\right) \frac{\pi}{2}, \\ \zeta_{\nu, n, \ell}^{(2)} = (2\nu + \alpha + \beta - n + 1)\zeta + \left(\alpha + \ell + \frac{1}{2}\right) \frac{\pi}{2},$$

where n and ℓ are non-negative integers. Then, the Jacobi function and the associated functions admit the following inverse factorial expansions:

$$(1.14) \quad 2^{2\nu+\alpha+\beta} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha, \beta)}(\cos(2\zeta)) \sim \\ \frac{1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \sum_{n=0}^{\infty} \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \cos\left(\zeta_{\nu, n, \ell}^{(1)}\right) \right) \frac{\Gamma(2\nu+\alpha+\beta-n+1)}{\Gamma(2\nu+\alpha+\beta+2)},$$

$$(1.15) \quad \frac{2^{2\nu+\alpha+\beta+1}}{\pi} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha,\beta)}(\cos(2\zeta)) \sim \frac{1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \sum_{n=0}^{\infty} \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \cos \left(\zeta_{\nu,n,\ell}^{(2)} \right) \right) \frac{\Gamma(2\nu+\alpha+\beta-n+1)}{\Gamma(2\nu+\alpha+\beta+2)},$$

$$(1.16) \quad \frac{2^{2\nu+\alpha+\beta+1}}{\pi} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha,\beta)}(\cos(2\zeta)) \sim \frac{-1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \sum_{n=0}^{\infty} \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \sin \left(\zeta_{\nu,n,\ell}^{(1)} \right) \right) \frac{\Gamma(2\nu+\alpha+\beta-n+1)}{\Gamma(2\nu+\alpha+\beta+2)},$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \pi - \delta$ ($< \pi$).

To our knowledge, inverse factorial-type expansions for these functions have not previously been established in the literature. For additional information on inverse factorial expansions, see, for example, [24].

We note that if 2α and 2β are odd integers, these inverse factorial expansions terminate, providing an exact representation of the corresponding function (see Sections 4 and 5).

The next two theorems provide factorial expansions for the Jacobi functions and their associated functions.

Theorem 1.3. *Suppose that $\xi \in \mathcal{D}$, that $|\sinh \xi|, |\cosh \xi| \geq \varepsilon$ (> 0), and that $\alpha, \beta \in \mathbb{C}$ are bounded. Under these conditions, the Jacobi functions admit factorial expansions given by*

$$(1.17) \quad \frac{\pi P_\nu^{(\alpha,\beta)}(\cosh(2\xi))}{2^{2\nu+\alpha+\beta} B(\nu+\alpha+1, \nu+\beta+1)} \sim \left(\frac{e^\xi}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^\xi}{\cosh \xi} \right)^{\beta+1/2} e^{2\nu\xi} \sum_{n=0}^{\infty} (-1)^n g_n(\xi, \alpha, \beta) \frac{\Gamma(2\nu+\alpha+\beta+2)}{\Gamma(2\nu+\alpha+\beta+n+2)} - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} \sum_{m=0}^{\infty} (-1)^m g_m(-\xi, \alpha, \beta) \frac{\Gamma(2\nu+\alpha+\beta+2)}{\Gamma(2\nu+\alpha+\beta+m+2)}$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \frac{\pi}{2} - \delta$ ($< \frac{\pi}{2}$), and

$$(1.18) \quad \frac{Q_\nu^{(\alpha,\beta)}(\cosh(2\xi))}{2^{2\nu+\alpha+\beta} B(\nu+\alpha+1, \nu+\beta+1)} \sim \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} \sum_{n=0}^{\infty} (-1)^n g_n(-\xi, \alpha, \beta) \frac{\Gamma(2\nu+\alpha+\beta+2)}{\Gamma(2\nu+\alpha+\beta+n+2)}$$

as $\nu \rightarrow \infty$ within the sector $|\arg \nu| \leq \pi - \delta$ ($< \pi$). The coefficients $g_n(\pm\xi, \alpha, \beta)$ are the same as those appearing in the corresponding inverse factorial series. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere.

In the special case where $\operatorname{Re}(\xi) > 0$, $-\frac{\pi}{2} < \operatorname{Im}(\xi) < 0$ (or equivalently, $\operatorname{Im}(z) < 0$), with real parameters α and β and ν approaching infinity along positive real values, the factorial expansion (1.17) was derived by Hahn [17]. Note that some typographical errors appear in the expressions for the error term and coefficients in [17].

We note that when $\operatorname{Re}(2\nu+\alpha+\beta+1) > \operatorname{Re}(\alpha'+\beta')$ and $|\operatorname{Re}(e^{\pm 2\xi})| < \frac{1}{2}$, the infinite series in (1.17) converges to its left-hand side, and similarly, when $\operatorname{Re}(2\nu+\alpha+\beta+1) > \operatorname{Re}(\alpha'+\beta')$ and

$|\operatorname{Re}(e^{-2\xi})| < \frac{1}{2}$, the series in (1.18) converges to the left-hand side. Here, α' and β' are defined in (2.16). For a short proof, refer to the end of Section 6.

Theorem 1.4. *Suppose that $(0 <) \varepsilon \leq \zeta \leq \frac{\pi}{2} - \varepsilon$ ($< \frac{\pi}{2}$), and let $\alpha, \beta \in \mathbb{C}$ be bounded. Define*

$$(1.19) \quad \begin{aligned} \zeta_{\nu, n, \ell}^{(3)} &= (2\nu + \alpha + \beta + n + 1)\zeta - (\alpha + \ell + \frac{1}{2})\frac{\pi}{2}, \\ \zeta_{\nu, n, \ell}^{(4)} &= (2\nu + \alpha + \beta + n + 1)\zeta + (\alpha - \ell + \frac{1}{2})\frac{\pi}{2}, \end{aligned}$$

where n and ℓ are non-negative integers. Then, the Jacobi function and the associated functions admit the following factorial expansions:

$$(1.20) \quad \begin{aligned} &\frac{\pi P_{\nu}^{(\alpha, \beta)}(\cos(2\zeta))}{2^{2\nu + \alpha + \beta + 1} B(\nu + \alpha + 1, \nu + \beta + 1)} \sim \\ &\frac{1}{\sin^{\alpha + 1/2} \zeta \cos^{\beta + 1/2} \zeta} \sum_{n=0}^{\infty} (-1)^n \left(\sum_{\ell=0}^n \frac{a_{\ell}(\alpha) a_{n-\ell}(\beta)}{\sin^{\ell} \zeta \cos^{n-\ell} \zeta} \cos\left(\zeta_{\nu, n, \ell}^{(3)}\right) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)}, \end{aligned}$$

$$(1.21) \quad \begin{aligned} &\frac{Q_{\nu}^{(\alpha, \beta)}(\cos(2\zeta))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} \sim \\ &\frac{1}{\sin^{\alpha + 1/2} \zeta \cos^{\beta + 1/2} \zeta} \sum_{n=0}^{\infty} (-1)^n \left(\sum_{\ell=0}^n \frac{a_{\ell}(\alpha) a_{n-\ell}(\beta)}{\sin^{\ell} \zeta \cos^{n-\ell} \zeta} \cos\left(\zeta_{\nu, n, \ell}^{(4)}\right) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)}, \end{aligned}$$

$$(1.22) \quad \begin{aligned} &\frac{Q_{\nu}^{(\alpha, \beta)}(\cos(2\zeta))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} \sim \\ &\frac{-1}{\sin^{\alpha + 1/2} \zeta \cos^{\beta + 1/2} \zeta} \sum_{n=0}^{\infty} (-1)^n \left(\sum_{\ell=0}^n \frac{a_{\ell}(\alpha) a_{n-\ell}(\beta)}{\sin^{\ell} \zeta \cos^{n-\ell} \zeta} \sin\left(\zeta_{\nu, n, \ell}^{(3)}\right) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)}, \end{aligned}$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \pi - \delta$ ($< \pi$).

In the special case where α and β are real and ν approaches infinity along positive real values, the factorial expansion (1.20) was derived by Hahn [17]. For more information on factorial series, see, for example, [7].

In the case that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$ and $\frac{\pi}{6} < \zeta < \frac{\pi}{3}$, the infinite series on the right-hand sides of (1.20), (1.21) and (1.22) converge to the corresponding functions on the left-hand sides. For a short proof, refer to the end of Section 8.

We mention that if 2α and 2β are odd integers, these factorial expansions terminate, providing an exact representation of the corresponding function (see Sections 6 and 8).

The proofs of Theorems 1.1–1.4 rely on the connection between these functions and hypergeometric functions, Watson's large-parameter asymptotic expansions for hypergeometric functions, and our error-bound theorems presented in Section 2. The proofs of our error bounds are based on convenient representations of the Jacobi functions, which we briefly discuss here for the Jacobi function of the second kind $Q_{\nu}^{(\alpha, \beta)}(z)$. When ν is large, this function is the recessive solution of differential equation (1.1).

(i) The Jacobi function of the second kind can be expressed in terms of the modified Bessel function of the second kind as follows:

$$(1.23) \quad \begin{aligned} 2^{2\nu+\alpha+\beta} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha,\beta)}(\cosh(2\xi)) &= \frac{1}{\sinh^\alpha \xi \cosh^\beta \xi} \\ &\times \frac{1}{\Gamma(2\nu+\alpha+\beta+2)} \int_0^{+\infty} t^{2\nu+\alpha+\beta+1} K_\alpha(t \sinh \xi) K_\beta(t \cosh \xi) dt, \end{aligned}$$

provided that $\xi \in \mathcal{D}$ and $\operatorname{Re}(2\nu+\alpha+\beta+2) > |\operatorname{Re}(\alpha)| + |\operatorname{Re}(\beta)|$ ¹. An immediate consequence of the factor $t^{2\nu+\alpha+\beta+1}$ in the integrand is that when $\operatorname{Re}(\nu)$ is large and positive, the main contribution to this integral comes from large values of t . Thus, we can obtain a large- ν asymptotic expansion by using the well-known large-argument asymptotic expansion of the Bessel function. This approach allows for the straightforward derivation of explicit and computable error bounds.

(ii) Alternatively, the Jacobi function of the second kind can also be represented in terms of the hypergeometric function as follows:

$$(1.24) \quad Q_\nu^{(\alpha,\beta)}(\cosh(2\xi)) = \frac{2\Gamma(\nu+\alpha+1)\Gamma(\nu+\beta+1)}{(\sinh \xi)^{2\alpha} (\cosh \xi)^{2\nu+2\beta+2}} \mathbf{F} \left(\begin{matrix} \nu+1, \nu+\beta+1 \\ 2\nu+\alpha+\beta+2 \end{matrix}; \frac{1}{\cosh^2 \xi} \right),$$

where $\xi \in \mathcal{D}$ and $\nu+\alpha, \nu+\beta \notin \mathbb{Z}^-$ (see [5, Eq. (22)]). Building on ideas by Hahn [17], we will reformulate the hypergeometric function as a double integral, making it suitable for obtaining factorial series with explicit remainder terms.

We conclude this section with a brief overview of existing asymptotic results in the literature. Darboux [6] established the leading-order asymptotic behaviour of the Jacobi polynomials $P_n^{(\alpha,\beta)}(z)$ for $-1 < z < 1$ and $z \in \mathbb{C} \setminus [-1, 1]$, away from the points $z = \pm 1$, in the case of large n with fixed α and β . Watson [28] provided complete asymptotic expansions for the associated hypergeometric functions, with the coefficients in these expansions being notably complicated. In the case of Jacobi polynomials, Watson's results were presented by Szegő [26, Theorem 8.21.9], who also derived a complicated uniform asymptotic expansion in terms of Bessel functions for $P_n^{(\alpha,\beta)}(x)$ for large n over the interval $-1 < x \leq 1$ [25]. Hahn [17] derived factorial series expansions for the Jacobi function $P_n^{(\alpha,\beta)}(z)$ for $-1 < z < 1$ and $\operatorname{Im}(z) < 0$, away from the points $z = \pm 1$, in the case of large positive n with fixed α and β . These expansions were complemented with error bounds for $-\frac{1}{2} < \alpha, \beta < \frac{1}{2}$. McCabe and Connor [20] provided the leading-order asymptotic behaviour of $P_n^{(\alpha,\beta)}(x)$, $Q_n^{(\alpha,\beta)}(x)$, and $Q_n^{(\alpha,\beta)}(x)$ for $0 < x < 1$ in the case of large n with fixed α and β . Several authors have derived uniform asymptotic approximations or expansions of the Jacobi functions in terms of Bessel functions, which remain valid at the endpoint $z = 1$ of the interval $-1 \leq z \leq 1$, with fixed parameters α and β in specific real or complex domains [2, 3, 10, 11, 14, 15, 30, 31, 32, 33, 34]. In many of these works, the expansions are accompanied by error bounds. Jones [18] obtained large- ν uniform asymptotic expansions for the hypergeometric functions associated with $P_\nu^{(\alpha,\beta)}(z)$ for z outside the interval $-1 < z < 1$. Finally, several studies have focused on obtaining large- n asymptotic results for the Jacobi polynomials $P_n^{(\alpha,\beta)}(z)$ that are uniform in α and β within certain real or complex domains [4, 19, 1, 16, 27].

The remainder of the paper is organised as follows. In Section 2, we provide error bounds for the inverse factorial and factorial expansions presented in Theorems 1.1–1.4. Section 3 contains the proofs of Theorems 1.1–1.4. The proofs of the error bounds for the inverse factorial series are given in Sections 4 and 5, while those for the factorial series are provided in Sections 6–8.

¹Note that this condition ensures that none of $\nu+1$, $\nu+\alpha$, $\nu+\beta$, or $\nu+\alpha+\beta$ are negative integers, so the left-hand side of (1.23) is well-defined.

2. ERROR BOUNDS

In the following two theorems, we provide computable error bounds for the inverse factorial expansions (1.11) and (1.12) of the Jacobi functions $P_\nu^{(\alpha, \beta)}(z)$ and $Q_\nu^{(\alpha, \beta)}(z)$, respectively. In Theorem 2.1, we introduce the notation

$$(2.1) \quad \chi(p) = \sqrt{\pi} \frac{\Gamma\left(\frac{p}{2} + 1\right)}{\Gamma\left(\frac{p}{2} + \frac{1}{2}\right)}, \quad p > 0.$$

We remark that $\chi(p) \sim \sqrt{\frac{1}{2}\pi p}$ as $p \rightarrow +\infty$ [23, Eq. (9.7.4)]. For the definition of the domain \mathcal{D} , see (1.9).

Theorem 2.1. *Let $\xi \in \mathcal{D}$ and let N, M both be arbitrary non-negative integers. Let ν, α , and β be complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \max(N, M)$. Then we have*

$$(2.2) \quad 2^{2\nu + \alpha + \beta + 1} B(\nu + 1, \nu + \alpha + \beta + 1) P_\nu^{(\alpha, \beta)}(\cosh(2\xi)) = \\ \left(\frac{e^\xi}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^\xi}{\cosh \xi} \right)^{\beta + 1/2} e^{2\nu\xi} \left(\sum_{n=0}^{N-1} g_n(-\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta - n + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} + R_N^{(P1)}(\nu, \xi, \alpha, \beta) \right) \\ - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta + 1/2} e^{-2\nu\xi} \left(\sum_{m=0}^{M-1} g_m(\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta - m + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} + R_M^{(P2)}(\nu, \xi, \alpha, \beta) \right),$$

where $C(\xi, \alpha)$ is defined in (1.10), and the remainder terms satisfy the following estimates:

$$(2.3) \quad \left| R_N^{(P1)}(\nu, \xi, \alpha, \beta) \right| \leq \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\alpha)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^N \right| \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|} \\ \times \begin{cases} 1 & \text{if } \operatorname{Re}(e^{2\xi}) \leq 1, \\ \min(|1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, 1 + \chi(N + \frac{1}{2})) & \text{if } \operatorname{Re}(e^{2\xi}) > 1 \end{cases} \\ + \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$(2.4) \quad \left| R_N^{(P1)}(\nu, \xi, \alpha, \beta) \right| \leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\cosh \xi} \right)^N \right| \right. \\ \left. + \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\beta) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|} \\ \times \begin{cases} 1 & \text{if } \operatorname{Re}(e^{2\xi}) \leq 1, \\ \min(|1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, 1 + \chi(N + \frac{1}{2})) & \text{if } \operatorname{Re}(e^{2\xi}) > 1, \end{cases}$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$, and

$$(2.5) \quad \left| R_N^{(P1)}(\nu, \xi, \alpha, \beta) \right| \leq \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|} \\ \times \begin{cases} 1 & \text{if } \operatorname{Re}(e^{2\xi}) \leq 1, \\ \min(|1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, 1 + \chi(N + \frac{1}{2})) & \text{if } \operatorname{Re}(e^{2\xi}) > 1, \end{cases}$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. Similarly,

$$(2.6) \quad \left| R_M^{(P2)}(\nu, \xi, \alpha, \beta) \right| \leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_M(\operatorname{Re}(\alpha)) \left(\frac{e^\xi}{\sinh \xi} \right)^M \right| + \sum_{\ell=0}^{M-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} a_\ell(\alpha) a_{M-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{M-\ell} \right| \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - M + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)| < M + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$(2.7) \quad \left| R_M^{(P2)}(\nu, \xi, \alpha, \beta) \right| \leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_M(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\cosh \xi} \right)^M \right| + \sum_{\ell=1}^M \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} a_\ell(\operatorname{Re}(\alpha)) a_{M-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{M-\ell} \right| \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - M + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < M + \frac{1}{2}$, and

$$(2.8) \quad \left| R_M^{(P2)}(\nu, \xi, \alpha, \beta) \right| \leq \sum_{\ell=0}^M \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{M-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{M-\ell} \right| \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - M + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. If $2 \operatorname{Re}(\alpha)$ or $2 \operatorname{Re}(\beta)$ is an odd integer, then the limiting values must to be taken in these bounds. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere. Furthermore, the remainder term $R_M^{(P2)}(\nu, \xi, \alpha, \beta)$ is bounded in absolute value by the corresponding first neglected term and retains the same sign, provided that ξ is positive, ν, α , and β are real, $|\alpha|, |\beta| < \frac{1}{2}$, and $2\nu + \alpha + \beta + 1 > M$.

Theorem 2.2. Let $\xi \in \mathcal{D}$ and let N be an arbitrary non-negative integer. Let ν, α , and β be complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > N$. Then we have

$$(2.9) \quad \frac{2^{2\nu + \alpha + \beta + 1}}{\pi} B(\nu + 1, \nu + \alpha + \beta + 1) Q_\nu^{(\alpha, \beta)}(\cosh(2\xi)) = \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} \left(\sum_{n=0}^{N-1} g_n(\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta - n + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} + R_N^{(Q)}(\nu, \xi, \alpha, \beta) \right),$$

where the remainder term satisfies the following estimates:

$$\begin{aligned} \left| R_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| &\leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\alpha)) \left(\frac{e^\xi}{\sinh \xi} \right)^N \right| \right. \\ &+ \left. \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|}, \end{aligned}$$

provided $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$\begin{aligned} \left| R_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| &\leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\cosh \xi} \right)^N \right| \right. \\ &+ \left. \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|}, \end{aligned}$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$, and

$$\begin{aligned} \left| R_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| &\leq \\ &\sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|}, \end{aligned}$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. If $2 \operatorname{Re}(\alpha)$ or $2 \operatorname{Re}(\beta)$ is an odd integer, then the limiting values must to be taken in these bounds. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere. Furthermore, the remainder term $R_N^{(Q)}(\nu, \xi, \alpha, \beta)$ is bounded in absolute value by the corresponding first neglected term and retains the same sign, provided that ξ is positive, ν, α , and β are real, $|\alpha|, |\beta| < \frac{1}{2}$, and $2\nu + \alpha + \beta + 1 > N$.

The following theorem provides computable error bounds for the inverse factorial expansions (1.14), (1.15), and (1.16) of the Jacobi function $P_\nu^{(\alpha, \beta)}(z)$ and the associated functions $Q_\nu^{(\alpha, \beta)}(z)$ and $Q_\nu^{(\alpha, \beta)}(z)$.

Theorem 2.3. Let $0 < \zeta < \frac{\pi}{2}$ and let N be an arbitrary non-negative integer. Let ν, α , and β be complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > N$. Then we have

$$\begin{aligned} (2.10) \quad &2^{2\nu+\alpha+\beta} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha, \beta)}(\cos(2\zeta)) = \frac{1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \\ &\times \left(\sum_{n=0}^{N-1} \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \cos \left(\zeta_{\nu, n, \ell}^{(1)} \right) \right) \frac{\Gamma(2\nu + \alpha + \beta - n + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} + R_N^{(P)}(\nu, \zeta, \alpha, \beta) \right), \end{aligned}$$

$$\begin{aligned} (2.11) \quad &\frac{2^{2\nu+\alpha+\beta+1}}{\pi} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha, \beta)}(\cos(2\zeta)) = \frac{1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \\ &\times \left(\sum_{n=0}^{N-1} \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \cos \left(\zeta_{\nu, n, \ell}^{(2)} \right) \right) \frac{\Gamma(2\nu + \alpha + \beta - n + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} + R_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right), \end{aligned}$$

$$\begin{aligned} (2.12) \quad &\frac{2^{2\nu+\alpha+\beta+1}}{\pi} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha, \beta)}(\cos(2\zeta)) = \frac{-1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \\ &\times \left(\sum_{n=0}^{N-1} \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \sin \left(\zeta_{\nu, n, \ell}^{(1)} \right) \right) \frac{\Gamma(2\nu + \alpha + \beta - n + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} + R_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right), \end{aligned}$$

where $\zeta_{\nu,n,\ell}^{(1)}$ and $\zeta_{\nu,n,\ell}^{(2)}$ are defined in (1.13), and the remainder terms satisfy the following estimates:

$$(2.13) \quad \left| R_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| R_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_N(\operatorname{Re}(\alpha))}{\sin^N \zeta} \right| \cosh \left(\operatorname{Im} \left(\zeta_{\nu,N,N}^{(1)} \right) \right) \right. \\ \left. + \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} \frac{a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh \left(\operatorname{Im} \left(\zeta_{\nu,N,\ell}^{(1)} \right) \right) \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$(2.14) \quad \left| R_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| R_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_N(\operatorname{Re}(\beta))}{\cos^N \zeta} \right| \cosh \left(\operatorname{Im} \left(\zeta_{\nu,N,0}^{(1)} \right) \right) \right. \\ \left. + \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\beta)}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh \left(\operatorname{Im} \left(\zeta_{\nu,N,\ell}^{(1)} \right) \right) \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$, and

$$(2.15) \quad \left| R_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| R_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq \\ \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh \left(\operatorname{Im} \left(\zeta_{\nu,N,\ell}^{(1)} \right) \right) \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) - N + 1)}{|\Gamma(2\nu + \alpha + \beta + 2)|},$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$, with analogous bounds for $R_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ in which each instance of $\zeta_{\nu,N,\ell}^{(1)}$ ($0 \leq \ell \leq N$) is replaced by $\zeta_{\nu,N,\ell}^{(2)}$. If $2 \operatorname{Re}(\alpha)$ or $2 \operatorname{Re}(\beta)$ is an odd integer, then the limiting values must be taken in these bounds.

In the two theorems below, we present computable error bounds for the factorial expansions (1.17) and (1.18) of the Jacobi functions $P_\nu^{(\alpha,\beta)}(z)$ and $Q_\nu^{(\alpha,\beta)}(z)$, respectively. In these theorems, we use the following notation:

$$(2.16) \quad \alpha' = \begin{cases} \alpha, & \text{if } \operatorname{Re}(\alpha) \geq \frac{1}{2}, \\ \pm\alpha, & \text{if } -\frac{1}{2} < \operatorname{Re}(\alpha) < \frac{1}{2}, \\ -\alpha, & \text{if } \operatorname{Re}(\alpha) \leq -\frac{1}{2}, \end{cases} \quad \beta' = \begin{cases} \beta, & \text{if } \operatorname{Re}(\beta) \geq \frac{1}{2}, \\ \pm\beta, & \text{if } -\frac{1}{2} < \operatorname{Re}(\beta) < \frac{1}{2}, \\ -\beta, & \text{if } \operatorname{Re}(\beta) \leq -\frac{1}{2}. \end{cases}$$

The symbol $\pm\alpha$ indicates that either α or $-\alpha$ can be chosen arbitrarily, with the same meaning applying to $\pm\beta$.

Theorem 2.4. *Let $\xi \in \mathcal{D} \setminus \mathbb{R}^+$ and let N, M both be arbitrary non-negative integers. Let ν, α , and β be complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. Then we have*

$$(2.17) \quad \frac{\pi P_\nu^{(\alpha, \beta)}(\cosh(2\xi))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \left(\frac{e^\xi}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^\xi}{\cosh \xi} \right)^{\beta + 1/2} e^{2\nu\xi} \left(\sum_{n=0}^{N-1} (-1)^n g_n(\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)} + \widehat{R}_N^{(P1)}(\nu, \xi, \alpha, \beta) \right) - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta + 1/2} e^{-2\nu\xi} \times \left(\sum_{m=0}^{M-1} (-1)^m g_m(-\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + m + 2)} + \widehat{R}_M^{(P2)}(\nu, \xi, \alpha, \beta) \right),$$

where $C(\xi, \alpha)$ is defined in (1.10), and the remainder terms satisfy the following estimates:

$$(2.18) \quad \begin{aligned} \left| \widehat{R}_N^{(P1)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ &\times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\alpha)) \left(\frac{e^\xi}{\sinh \xi} \right)^N \right| \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ &\times \left| 1 - e^{4\xi} \right| |\operatorname{csc}(2 \operatorname{Im}(\xi))| + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\beta' + \frac{1}{2})|} \\ &\times \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} \frac{\Gamma(\operatorname{Re}(\alpha') + \ell + \frac{1}{2})}{\Gamma(\alpha' + \ell + \frac{1}{2})} a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \begin{cases} |1 + e^{2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{-2\xi}) \leq 0, \\ |1 + e^{2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{-2\xi}), \end{cases} \end{aligned}$$

provided $-\frac{1}{2} < \operatorname{Re}(\alpha) < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$(2.19) \quad \begin{aligned} \left| \widehat{R}_N^{(P1)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ &\times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\cosh \xi} \right)^N \right| \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ &\times \left| 1 - e^{4\xi} \right| |\operatorname{csc}(2 \operatorname{Im}(\xi))| + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\alpha') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\alpha' + \frac{1}{2})|} \\ &\times \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{\Gamma(\operatorname{Re}(\beta') + N - \ell + \frac{1}{2})}{\Gamma(\beta' + N - \ell + \frac{1}{2})} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \begin{cases} |1 - e^{2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{-2\xi}) < 1, \\ |1 - e^{2\xi}|, & \text{if } \operatorname{Re}(e^{-2\xi}) < 0, \end{cases} \end{aligned}$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $-\frac{1}{2} < \operatorname{Re}(\beta) < N + \frac{1}{2}$, and

$$(2.20) \quad \left| \widehat{R}_N^{(P_1)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ \times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \\ \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times |1 - e^{4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|,$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. Similarly,

$$(2.21) \quad \left| \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ \times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_M(\operatorname{Re}(\alpha)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^M \right| \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + M + 2)} \\ \times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1 \end{cases} + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\beta' + \frac{1}{2})|} \\ \times \sum_{\ell=0}^{M-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} \frac{\Gamma(\operatorname{Re}(\alpha') + \ell + \frac{1}{2})}{\Gamma(\alpha' + \ell + \frac{1}{2})} a_\ell(\alpha) a_{M-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{M-\ell} \right| \\ \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + M + 2)} \times \begin{cases} 1, & \text{if } \operatorname{Re}(e^{2\xi}) \leq -1, \\ |1 + e^{-2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{2\xi}) \leq 0, \\ |1 + e^{-2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{2\xi}), \end{cases}$$

provided $-\frac{1}{2} < \operatorname{Re}(\alpha) < M + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$(2.22) \quad \left| \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ \times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_M(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\cosh \xi} \right)^M \right| \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + M + 2)} \\ \times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1 \end{cases} + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\alpha') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\alpha' + \frac{1}{2})|} \\ \times \sum_{\ell=1}^M \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{\Gamma(\operatorname{Re}(\beta') + M - \ell + \frac{1}{2})}{\Gamma(\beta' + M - \ell + \frac{1}{2})} a_\ell(\operatorname{Re}(\alpha)) a_{M-\ell}(\beta) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{M-\ell} \right| \\ \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + M + 2)} \times \begin{cases} 1, & \text{if } 1 \leq \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{2\xi}) < 1, \\ |1 - e^{-2\xi}|, & \text{if } \operatorname{Re}(e^{2\xi}) < 0, \end{cases}$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $-\frac{1}{2} < \operatorname{Re}(\beta) < M + \frac{1}{2}$, and

$$(2.23) \quad \left| \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ \times \sum_{\ell=0}^M \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{M-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{M-\ell} \right| \\ \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + M + 2)} \times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1, \end{cases}$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. If $2 \operatorname{Re}(\alpha)$ or $2 \operatorname{Re}(\beta)$ is an odd integer, then the limiting values must to be taken in these bounds. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere.

Theorem 2.5. Let $\xi \in \mathcal{D}$ and let N be an arbitrary non-negative integer. Let ν, α , and β be complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. Then we have

$$(2.24) \quad \frac{Q_\nu^{(\alpha, \beta)}(\cosh(2\xi))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \\ \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta + 1/2} e^{-2\nu\xi} \left(\sum_{n=0}^{N-1} (-1)^n g_n(-\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)} + \widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) \right),$$

where the remainder term satisfies the following estimates:

$$\left| \widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ \times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\alpha)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^N \right| \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ \times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1 \end{cases} + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\beta' + \frac{1}{2})|} \\ \times \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} \frac{\Gamma(\operatorname{Re}(\alpha') + \ell + \frac{1}{2})}{\Gamma(\alpha' + \ell + \frac{1}{2})} a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \\ \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \begin{cases} 1, & \text{if } \operatorname{Re}(e^{2\xi}) \leq -1, \\ |1 + e^{-2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{2\xi}) \leq 0, \\ |1 + e^{-2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{2\xi}), \end{cases}$$

provided $-\frac{1}{2} < \operatorname{Re}(\alpha) < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

$$\begin{aligned} \left| \widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ &\times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\cosh \xi} \right)^N \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \right. \\ &\times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1 \end{cases} + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1) \Gamma(\operatorname{Re}(\alpha') + \frac{1}{2})}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)| |\Gamma(\alpha' + \frac{1}{2})|} \\ &\times \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{\Gamma(\operatorname{Re}(\beta') + N - \ell + \frac{1}{2})}{\Gamma(\beta' + N - \ell + \frac{1}{2})} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \begin{cases} 1, & \text{if } 1 \leq \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{2\xi}) < 1, \\ |1 - e^{-2\xi}|, & \text{if } \operatorname{Re}(e^{2\xi}) < 0, \end{cases} \end{aligned}$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $-\frac{1}{2} < \operatorname{Re}(\beta) < N + \frac{1}{2}$, and

$$\begin{aligned} \left| \widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ &\times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1, \end{cases} \end{aligned}$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. If $2 \operatorname{Re}(\alpha)$ or $2 \operatorname{Re}(\beta)$ is an odd integer, then the limiting values must to be taken in these bounds. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere. Furthermore, the remainder term $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta)$ is bounded in absolute value by twice the corresponding first neglected term and retains the same sign, provided that ξ is positive, ν, α , and β are real, $|\alpha|, |\beta| < \frac{1}{2}$, and $2\nu + \alpha + \beta + 1 > \alpha' + \beta'$.

In the following two theorems, we present alternative, less explicit error bounds for the factorial expansions (1.17) and (1.18). In these theorems, the function $\mathbf{g}(w)$ is defined as follows:

$$(2.25) \quad \mathbf{g}(w) = \begin{cases} 1 & \text{if } \operatorname{Re}(w) \leq 0, \\ \left| \frac{w}{\operatorname{Im}(w)} \right| & \text{if } 0 < \operatorname{Re}(w) \leq |w|^2, \\ \frac{1}{|1-w|} & \text{if } \operatorname{Re}(w) > |w|^2, \end{cases}$$

for $w \in \mathbb{C} \setminus [1, +\infty)$ (cf. [17, Eq. (5.5)]). In the special case where ν, α , and β are real and $|\alpha|, |\beta| < \frac{1}{2}$, Theorem 2.6 was also proved by Hahn [17].

Theorem 2.6. *Let $\xi \in \mathcal{D} \setminus \mathbb{R}^+$ and let N, M both be arbitrary non-negative integers. Suppose that ν, α , and β are complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$, and that either $0 < \operatorname{Re}(\alpha' + \beta')$ or $\alpha = \beta = 0$. Then the remainder terms in (2.17) satisfy the following*

estimates:

(2.26)

$$\begin{aligned} \left| \widehat{R}_N^{(P_1)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{|\Gamma(-\alpha' - \beta' + N + 1)|}{\Gamma(\operatorname{Re}(-\alpha' - \beta') + N + 1)} \\ &\times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \max_{0 \leq u \leq \frac{\pi}{2}} \mathfrak{g} \left(\frac{\sin^2 u}{1 + \sin(2u)} \frac{e^\xi}{2 \sinh \xi} + \frac{\cos^2 u}{1 + \sin(2u)} \frac{e^\xi}{2 \cosh \xi} \right), \end{aligned}$$

and

(2.27)

$$\begin{aligned} \left| \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{|\Gamma(-\alpha' - \beta' + M + 1)|}{\Gamma(\operatorname{Re}(-\alpha' - \beta') + M + 1)} \\ &\times \sum_{\ell=0}^M \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{M-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{M-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + M + 2)} \times \max_{0 \leq u \leq \frac{\pi}{2}} \mathfrak{g} \left(-\frac{\sin^2 u}{1 + \sin(2u)} \frac{e^{-\xi}}{2 \sinh \xi} + \frac{\cos^2 u}{1 + \sin(2u)} \frac{e^{-\xi}}{2 \cosh \xi} \right), \end{aligned}$$

provided that $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere.

Theorem 2.7. Let $\xi \in \mathcal{D}$ and let N be an arbitrary non-negative integer. Suppose that ν, α , and β are complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$, and that either $0 < \operatorname{Re}(\alpha' + \beta')$ or $\alpha = \beta = 0$. Then the remainder term in (2.24) satisfies the following estimate:

(2.28)

$$\begin{aligned} \left| \widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) \right| &\leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{|\Gamma(-\alpha' - \beta' + N + 1)|}{\Gamma(\operatorname{Re}(-\alpha' - \beta') + N + 1)} \\ &\times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \\ &\times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \max_{0 \leq u \leq \frac{\pi}{2}} \mathfrak{g} \left(-\frac{\sin^2 u}{1 + \sin(2u)} \frac{e^{-\xi}}{2 \sinh \xi} + \frac{\cos^2 u}{1 + \sin(2u)} \frac{e^{-\xi}}{2 \cosh \xi} \right), \end{aligned}$$

provided that $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. The fractional powers are defined to be positive for positive real ξ and are defined by continuity elsewhere.

The following theorem provides computable error bounds for the factorial expansions (1.20), (1.21), and (1.22) of the Jacobi function $P_\nu^{(\alpha, \beta)}(z)$ and the associated functions $Q_\nu^{(\alpha, \beta)}(z)$ and $Q_\nu^{(\alpha, \beta)}(z)$. In the special case where ν, α , and β are real and $|\alpha|, |\beta| < \frac{1}{2}$, the error bound (2.35) for $\widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta)$ was also proved by Hahn [17].

Theorem 2.8. *Let $\xi \in \mathcal{D}$ and let N be an arbitrary non-negative integer. Let ν , α , and β be complex numbers satisfying $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. Then we have*

(2.29)

$$\begin{aligned} & \frac{\pi P_\nu^{(\alpha, \beta)}(\cos(2\zeta))}{2^{2\nu + \alpha + \beta + 1} B(\nu + \alpha + 1, \nu + \beta + 1)} = \frac{1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \\ & \times \left(\sum_{n=0}^{N-1} (-1)^n \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \cos\left(\zeta_{\nu, n, \ell}^{(3)}\right) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)} + \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) \right), \end{aligned}$$

(2.30)

$$\begin{aligned} & \frac{Q_\nu^{(\alpha, \beta)}(\cos(2\zeta))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \frac{1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \\ & \times \left(\sum_{n=0}^{N-1} (-1)^n \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \cos\left(\zeta_{\nu, n, \ell}^{(4)}\right) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)} + \widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right), \end{aligned}$$

(2.31)

$$\begin{aligned} & \frac{Q_\nu^{(\alpha, \beta)}(\cos(2\zeta))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \frac{-1}{\sin^{\alpha+1/2} \zeta \cos^{\beta+1/2} \zeta} \\ & \times \left(\sum_{n=0}^{N-1} (-1)^n \left(\sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sin^\ell \zeta \cos^{n-\ell} \zeta} \sin\left(\zeta_{\nu, n, \ell}^{(3)}\right) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)} + \widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right), \end{aligned}$$

where $\zeta_{\nu, n, \ell}^{(3)}$ and $\zeta_{\nu, n, \ell}^{(4)}$ are defined in (1.19), and the remainder terms satisfy the following estimates:

(2.32)

$$\begin{aligned} & \left| \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| \widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq 2 \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ & \times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_N(\operatorname{Re}(\alpha))}{\sin^N \zeta} \right| \cosh\left(\operatorname{Im}\left(\zeta_{\nu, N, N}^{(3)}\right)\right) \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ & + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\beta' + \frac{1}{2})|} \\ & \times \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} \frac{\Gamma(\operatorname{Re}(\alpha') + \ell + \frac{1}{2})}{\Gamma(\alpha' + \ell + \frac{1}{2})} \frac{a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh\left(\operatorname{Im}\left(\zeta_{\nu, N, \ell}^{(3)}\right)\right) \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ & \times \begin{cases} 2 \cos \zeta, & \text{if } 0 < \zeta < \frac{\pi}{4}, \\ \csc \zeta, & \text{if } \frac{\pi}{4} \leq \zeta < \frac{\pi}{2}, \end{cases} \end{aligned}$$

provided $-\frac{1}{2} < \operatorname{Re}(\alpha) < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$,

(2.33)

$$\begin{aligned} & \left| \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| \widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq 2 \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ & \times \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_N(\operatorname{Re}(\beta))}{\cos^N \zeta} \right| \cosh\left(\operatorname{Im}\left(\zeta_{\nu, N, 0}^{(3)}\right)\right) \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ & + \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2})|} \\ & \times \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{\Gamma(\operatorname{Re}(\beta') + \ell + \frac{1}{2})}{\Gamma(\beta' + \ell + \frac{1}{2})} \frac{a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh\left(\operatorname{Im}\left(\zeta_{\nu, N, \ell}^{(3)}\right)\right) \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \\ & \times \begin{cases} \sec \zeta, & \text{if } 0 < \zeta < \frac{\pi}{4}, \\ 2 \sin \zeta, & \text{if } \frac{\pi}{4} \leq \zeta < \frac{\pi}{2}, \end{cases} \end{aligned}$$

provided $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $-\frac{1}{2} < \operatorname{Re}(\beta) < N + \frac{1}{2}$,

(2.34)

$$\begin{aligned} & \left| \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| \widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq 2 \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2}) \Gamma(\operatorname{Re}(\beta') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})|} \\ & \times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh\left(\operatorname{Im}\left(\zeta_{\nu, N, \ell}^{(3)}\right)\right) \\ & \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)}. \end{aligned}$$

provided $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$, and

(2.35)

$$\begin{aligned} & \left| \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) \right|, \left| \widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta) \right| \leq 2 \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{|\Gamma(-\alpha' - \beta' + N + 1)|}{\Gamma(\operatorname{Re}(-\alpha' - \beta') + N + 1)} \\ & \times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \cosh\left(\operatorname{Im}\left(\zeta_{\nu, N, \ell}^{(3)}\right)\right) \\ & \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)}, \end{aligned}$$

provided that $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$, and that either $0 < \operatorname{Re}(\alpha' + \beta')$ or $\alpha = \beta = 0$. Analogous bounds hold for $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ in which each instance of $\zeta_{\nu, N, \ell}^{(3)}$ ($0 \leq \ell \leq N$) is replaced by $\zeta_{\nu, N, \ell}^{(4)}$. If $2\operatorname{Re}(\alpha)$ or $2\operatorname{Re}(\beta)$ is an odd integer, then the limiting values must to be taken in these bounds.

3. PROOFS OF THEOREMS 1.1–1.4

The validity of the inverse factorial expansions (1.11) and (1.12) in the special case where $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$, $\operatorname{Re}(\nu) \rightarrow +\infty$, and $\operatorname{Im}(\nu)$ is fixed, follows from Theorems 2.1 and 2.2. In what follows, we will extend the region where these expansions are valid. Define

$$(3.1) \quad \overline{\mathcal{D}} = \{\xi : \operatorname{Re}(\xi) \geq 0, \xi \neq 0, |\operatorname{Im}(\xi)| < \frac{\pi}{2}\}.$$

Note that $\cosh(2\xi)$ maps $\overline{\mathcal{D}}$ bijectively onto the region $\mathbb{C} \setminus (-\infty, 1]$, along with both sides of the open segment $(0, 1)$ on the cut along $(-\infty, 1]$. Using the results of Watson [28, §3–9] applied to

the hypergeometric functions in (1.2) and (1.3), we find that for $\xi \in \overline{\mathcal{D}}$ and $\alpha, \beta \in \mathbb{C}$, the Jacobi functions can be represented as

$$(3.2) \quad 2^{2\nu+\alpha+\beta+1} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha, \beta)}(\cosh(2\xi)) = \left(\frac{e^\xi}{\sinh \xi}\right)^{\alpha+1/2} \left(\frac{e^\xi}{\cosh \xi}\right)^{\beta+1/2} e^{2\nu\xi} I_+ \pm i e^{\pm\pi i \alpha} \left(\frac{e^{-\xi}}{\sinh \xi}\right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi}\right)^{\beta+1/2} e^{-2\nu\xi} I_-,$$

and

$$\frac{2^{2\nu+\alpha+\beta+1}}{\pi} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha, \beta)}(\cosh(2\xi)) = \left(\frac{e^{-\xi}}{\sinh \xi}\right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi}\right)^{\beta+1/2} e^{-2\nu\xi} I_-,$$

where, for fixed α and β , the functions I_\pm have asymptotic expansions of the form

$$(3.3) \quad I_\pm \sim \sum_{n=0}^{\infty} \frac{c_n(e^{\pm\xi})}{\Gamma(2\nu+\alpha+\beta+2)} \frac{2^{2\nu+\alpha+\beta}}{\pi} \Gamma(n+\frac{1}{2}) \frac{\Gamma(\nu+1)\Gamma(\nu+\alpha+\beta+1)}{\nu^{n+1/2}}$$

as $\nu \rightarrow \infty$, within the sector $|\arg \nu| \leq \frac{\pi}{2} - \delta$ ($< \frac{\pi}{2}$) (for I_+) and within $|\arg \nu| \leq \pi - \delta$ ($< \pi$) (for I_-). In (3.3), we assume that ξ is bounded away from the points $0, \pm\frac{\pi}{2}i$. The coefficients $c_n(w)$ are rational functions of w and polynomial functions of α and β . In the expansion (3.2), the upper or lower sign is taken according as $\text{Im}(\xi) \gtrless 0$. Watson does not specify the choice of sign when $\text{Im}(\xi) = 0$, noting that the term $e^{-2\nu\xi} I_-$ is exponentially small compared to $e^{2\nu\xi} I_+$ and thus can be neglected. To derive an expression for $P_\nu^{(\alpha, \beta)}(\cosh(2\xi))$ when $\text{Im}(\xi) = 0$, we use

$$P_\nu^{(\alpha, \beta)}(\cosh(2\xi)) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \left(P_\nu^{(\alpha, \beta)}(\cosh(2\xi + i\varepsilon)) + P_\nu^{(\alpha, \beta)}(\cosh(2\xi - i\varepsilon)) \right)$$

and obtain

$$2^{2\nu+\alpha+\beta+1} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha, \beta)}(\cosh(2\xi)) = \left(\frac{e^\xi}{\sinh \xi}\right)^{\alpha+1/2} \left(\frac{e^\xi}{\cosh \xi}\right)^{\beta+1/2} e^{2\nu\xi} I_+ - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi}\right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi}\right)^{\beta+1/2} e^{-2\nu\xi} I_-,$$

where $C(\xi, \alpha)$ is defined in (1.10). This is consistent with both (2.2) and the fact that $P_\nu^{(\alpha, \beta)}(\cosh(2\xi))$ is real when ξ, ν, α , and β are real. For any fixed complex h , the gamma function has an asymptotic expansion given by

$$(3.4) \quad \Gamma(\nu+h) \sim \sqrt{2\pi} \nu^{\nu+h-1/2} e^{-\nu} \left(1 + \frac{6h^2 - 6h + 1}{12\nu} + \frac{36h^4 - 120h^3 + 120h^2 - 36h + 1}{288\nu^2} + \dots \right),$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \pi - \delta$ ($< \pi$). This result follows from exponentiating the well-known asymptotic expansion of $\log \Gamma(\nu+h)$ [23, Eq. (5.11.8)]. Using this expansion, we can verify the inverse factorial expansion

$$\frac{2^{2\nu+\alpha+\beta}}{\pi} \Gamma(n+\frac{1}{2}) \frac{\Gamma(\nu+1)\Gamma(\nu+\alpha+\beta+1)}{\nu^{n+1/2}} \sim \sum_{k=0}^{\infty} r_{n,k} \Gamma(2\nu+\alpha+\beta-n-k+1),$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \pi - \delta$ ($< \pi$), where the coefficients $r_{n,k}$ are polynomials in α and β . The first two are explicitly given by

$$r_{n,0} = \frac{(2n)!}{2^n n!}, \quad r_{n,1} = \frac{(2n)!}{2^{n+2} n!} (2(\alpha+\beta)^2 + (2n+1)(1+2\alpha+2\beta) - 2n^2).$$

Substituting into (3.3) gives

$$I_{\pm} \sim \sum_{n=0}^{\infty} \left(\sum_{k=0}^n r_{n-k,k} c_{n-k} (e^{\pm\xi}) \right) \frac{\Gamma(2\nu + \alpha + \beta - n + 1)}{\Gamma(2\nu + \alpha + \beta + 2)},$$

as $\nu \rightarrow \infty$, within the sector $|\arg \nu| \leq \frac{\pi}{2} - \delta$ ($< \frac{\pi}{2}$) (for I_+) and within $|\arg \nu| \leq \pi - \delta$ ($< \pi$) (for I_-). Theorem 2.1 and the uniqueness of the coefficients of an asymptotic expansion imply that

$$(3.5) \quad g_n(\mp\xi, \alpha, \beta) = \sum_{k=0}^n r_{n-k,k} c_{n-k} (e^{\pm\xi})$$

when $\xi \in \mathcal{D}$ and $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. Since both sides are rational functions of $e^{\pm\xi}$ and polynomial functions of α and β , an analytic continuation and continuity argument shows that (3.5) holds for $\xi \in \overline{\mathcal{D}}$ and arbitrary $\alpha, \beta \in \mathbb{C}$. This concludes the proof of Theorem 1.1.

We now continue with the proof of Theorem 1.2. Suppose that $0 < \zeta < \frac{\pi}{2}$. Then, by (1.4), we have

$$(3.6) \quad P_{\nu}^{(\alpha,\beta)}(\cos(2\zeta)) = \lim_{\varepsilon \rightarrow 0^+} \frac{i}{\pi} \left(e^{\pi\alpha i} Q_{\nu}^{(\alpha,\beta)}(\cosh(2\varepsilon + 2i\zeta)) - e^{-\pi\alpha i} Q_{\nu}^{(\alpha,\beta)}(\cosh(2\varepsilon - 2i\zeta)) \right),$$

provided that $\nu + \alpha, \nu + \beta \notin \mathbb{Z}^-$. The associated functions $Q_{\nu}^{(\alpha,\beta)}(x)$ and $Q_{\nu}^{(\alpha,\beta)}(x)$ are defined in terms of the Jacobi function of the second kind, through the limits (1.5) and (1.6), respectively. Specifically,

$$(3.7) \quad Q_{\nu}^{(\alpha,\beta)}(\cos(2\zeta)) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \left(Q_{\nu}^{(\alpha,\beta)}(\cosh(2\varepsilon + 2i\zeta)) + Q_{\nu}^{(\alpha,\beta)}(\cosh(2\varepsilon - 2i\zeta)) \right),$$

and

$$(3.8) \quad Q_{\nu}^{(\alpha,\beta)}(\cos(2\zeta)) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2} \left(e^{\pi\alpha i} Q_{\nu}^{(\alpha,\beta)}(\cosh(2\varepsilon + 2i\zeta)) + e^{-\pi\alpha i} Q_{\nu}^{(\alpha,\beta)}(\cosh(2\varepsilon - 2i\zeta)) \right),$$

provided that $\nu + \alpha, \nu + \beta \notin \mathbb{Z}^-$. Since the validity of the inverse factorial series for $Q_{\nu}^{(\alpha,\beta)}(\cosh(2\xi))$ has been established in $\overline{\mathcal{D}}$, we can apply it to the right-hand sides of (3.6)–(3.8). After some straightforward algebra, this yields the desired inverse factorial expansions presented in Theorem 1.2.

The validity of the factorial expansions in Theorem 1.3 for the special case where $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$, $\operatorname{Re}(\nu) \rightarrow +\infty$, and $\operatorname{Im}(\nu)$ is fixed, is established by Theorems 2.4 and 2.5. In the following, we will extend the region of validity for these expansions. Following a similar approach to that used for the inverse factorial series, we find that for $\xi \in \overline{\mathcal{D}}$ and $\alpha, \beta \in \mathbb{C}$, the Jacobi functions can be represented as

$$\frac{\pi P_{\nu}^{(\alpha,\beta)}(\cosh(2\xi))}{2^{2\nu+\alpha+\beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \left(\frac{e^{\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{2\nu\xi} J_+ - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} J_-,$$

and

$$\frac{Q_{\nu}^{(\alpha,\beta)}(\cosh(2\xi))}{2^{2\nu+\alpha+\beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha+1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta+1/2} e^{-2\nu\xi} J_-,$$

where, for fixed α and β , the functions J_{\pm} have asymptotic expansions of the form

$$(3.9) \quad J_{\pm} \sim \sum_{n=0}^{\infty} c_n (e^{\pm\xi}) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{2^{2\nu+\alpha+\beta+1}} \Gamma\left(n + \frac{1}{2}\right) \frac{1}{\nu^{n+\frac{1}{2}} \Gamma(\nu + \alpha + 1) \Gamma(\nu + \beta + 1)}$$

as $\nu \rightarrow \infty$, within the sector $|\arg \nu| \leq \frac{\pi}{2} - \delta$ ($< \frac{\pi}{2}$) (for J_+) and within $|\arg \nu| \leq \pi - \delta$ ($< \pi$) (for J_-). In (3.9), we assume that ξ is bounded away from the points $0, \pm \frac{\pi}{2}i$. Using the expansion (3.4), we can verify the factorial expansion

$$\frac{1}{2^{2\nu+\alpha+\beta+1}} \Gamma\left(n + \frac{1}{2}\right) \frac{1}{\nu^{n+\frac{1}{2}} \Gamma(\nu + \alpha + 1) \Gamma(\nu + \beta + 1)} \sim \sum_{k=0}^{\infty} \frac{q_{n,k}}{\Gamma(2\nu + \alpha + \beta + n + k + 2)},$$

as $\nu \rightarrow \infty$ in the sector $|\arg \nu| \leq \pi - \delta$ ($< \pi$), where the coefficients $q_{n,k}$ are polynomials in α and β . The first two are explicitly given by

$$q_{n,0} = \frac{(2n)!}{2^n n!}, \quad q_{n,1} = -\frac{(2n)!}{2^{n+2} n!} (2(\alpha - \beta)^2 - (2n + 1)(2\alpha + 2\beta + 3) - 2n^2).$$

Substituting into (3.9), we obtain

$$J_{\pm} \sim \sum_{n=0}^{\infty} \left(\sum_{k=0}^n q_{n-k,k} c_{n-k} (e^{\pm \xi}) \right) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + n + 2)},$$

as $\nu \rightarrow \infty$, within the sector $|\arg \nu| \leq \frac{\pi}{2} - \delta$ ($< \frac{\pi}{2}$) (for J_+) and within $|\arg \nu| \leq \pi - \delta$ ($< \pi$) (for J_-). Theorem 2.4 and the uniqueness of the coefficients in an asymptotic expansion imply that

$$(3.10) \quad g_n(\pm \xi, \alpha, \beta) = (-1)^n \sum_{k=0}^n q_{n-k,k} c_{n-k} (e^{\pm \xi})$$

when $\xi \in \mathcal{D} \setminus \mathbb{R}^+$ and $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. Since both sides of this equation are rational functions of $e^{\pm \xi}$ and polynomial functions of α and β , an analytic continuation and continuity argument show that (3.10) holds for $\xi \in \overline{\mathcal{D}}$ and arbitrary $\alpha, \beta \in \mathbb{C}$. This completes the proof of Theorem 1.3.

Since the factorial series for $Q_{\nu}^{(\alpha, \beta)}(\cosh(2\xi))$ has been established in $\overline{\mathcal{D}}$, we can now apply it to the right-hand sides of (3.6)–(3.8). A straightforward algebraic manipulation then leads to the desired factorial expansions, as given in Theorem 1.4.

4. PROOFS OF THEOREMS 2.1 AND 2.2

In this section, we establish the error bounds presented in Theorems 2.1 and 2.2. To this end, we need an estimate for the remainder term in the well-known asymptotic expansion of the modified Bessel function of the second kind, as provided in the following lemma.

Lemma 4.1. *Let N be a non-negative integer, and let μ be an arbitrary complex number. Define the remainder $r_N^{(K)}(w, \mu)$ by*

$$(4.1) \quad K_{\mu}(w) = \sqrt{\frac{\pi}{2w}} e^{-w} \left(\sum_{n=0}^{N-1} \frac{a_n(\mu)}{w^n} + r_N^{(K)}(w, \mu) \right).$$

Then we have the bound

$$\left| r_N^{(K)}(w, \mu) \right| \leq \left| \frac{\cos(\pi \mu)}{\cos(\pi \operatorname{Re}(\mu))} \right| \frac{|a_N(\operatorname{Re}(\mu))|}{|w|^N} \times \begin{cases} 1 & \text{if } |\arg w| \leq \frac{\pi}{2}, \\ \min(|\operatorname{csc}(\arg w)|, 1 + \chi(N + \frac{1}{2})) & \text{if } \frac{\pi}{2} < |\arg w| \leq \pi, \end{cases}$$

provided $|\operatorname{Re}(\mu)| < N + \frac{1}{2}$. Here, $\chi(p)$ is defined in (2.1). If $2\operatorname{Re}(\mu)$ is an odd integer, the limiting value must be taken in this bound. Moreover, the remainder term $r_N^{(K)}(w, \mu)$ does not exceed the first neglected term in absolute value and has the same sign, provided w is positive, μ is real, and $|\mu| < N + \frac{1}{2}$.

Proof. The statement for complex variables follows from Theorem 1.8 and Propositions B.1 and B.3 in [21]. For the real case, see, e.g., [29, pp. 206–207]. \square

We proceed with the proof of Theorem 2.1. The Jacobi function of the first kind can be expressed in terms of modified Bessel functions as

$$2^{\nu-1} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha,\beta)}(z) = \frac{1}{(z-1)^{\alpha/2} (z+1)^{\beta/2}} \\ \times \frac{1}{\Gamma(2\nu+\alpha+\beta+2)} \int_0^{+\infty} t^{2\nu+\alpha+\beta+1} I_\alpha(t\sqrt{z-1}) K_\beta(t\sqrt{z+1}) dt,$$

provided that $\operatorname{Re}(\sqrt{z+1}) > |\operatorname{Re}(\sqrt{z-1})|$ and $\operatorname{Re}(2\nu+\alpha+\beta+2) > -\operatorname{Re}(\alpha) + |\operatorname{Re}(\beta)|$ (combine [5, Eq. (18)] and [13, Eq. (6.8.44)]²). By applying a suitable change of variables, this can be re-written as

$$(4.2) \quad 2^{2\nu+\alpha+\beta} B(\nu+1, \nu+\alpha+\beta+1) P_\nu^{(\alpha,\beta)}(\cosh(2\xi)) = \frac{1}{\sinh^\alpha \xi \cosh^\beta \xi} \\ \times \frac{1}{\Gamma(2\nu+\alpha+\beta+2)} \int_0^{+\infty} t^{2\nu+\alpha+\beta+1} I_\alpha(t \sinh \xi) K_\beta(t \cosh \xi) dt$$

provided that $\xi \in \mathcal{D}$ and $\operatorname{Re}(2\nu+\alpha+\beta+2) > -\operatorname{Re}(\alpha) + |\operatorname{Re}(\beta)|$.

Now, suppose that $|\arg w| \leq \pi$. Under this assumption, the modified Bessel functions satisfy the following relations:

$$I_\mu(w) = \mp \frac{i}{\pi} K_\mu(we^{\mp\pi i}) \pm \frac{i}{\pi} e^{\pm\pi i \mu} K_\mu(w),$$

when $0 < \pm \arg w \leq \pi$, and

$$I_\mu(w) = \frac{i}{2\pi} (K_\mu(we^{\pi i}) - K_\mu(we^{-\pi i})) - \frac{\sin(\pi\mu)}{\pi} K_\mu(w),$$

when $\arg w = 0$ (cf. [23, Eq. (10.34.3)]). By combining these functional equations with the expression (4.1), we obtain

$$(4.3) \quad I_\mu(w) = \frac{e^w}{\sqrt{2\pi w}} \left(\sum_{n=0}^{N-1} (-1)^n \frac{a_n(\mu)}{w^n} + \tilde{r}_N^{(K)}(w, \mu) \right) \\ - \tilde{C}(\arg w, \mu) \frac{e^{-w}}{\sqrt{2\pi w}} \left(\sum_{m=0}^{M-1} \frac{a_m(\mu)}{w^m} + r_M^{(K)}(w, \mu) \right),$$

with

$$(4.4) \quad \tilde{C}(\arg w, \mu) = \begin{cases} \sin(\pi\mu) & \text{if } \arg w = 0, \\ \mp i e^{\pm\pi i \mu} & \text{if } 0 < \pm \arg w \leq \pi, \end{cases}$$

and

$$(4.5) \quad \tilde{r}_N^{(K)}(w, \mu) = \begin{cases} \frac{1}{2} \left(r_N^{(K)}(we^{\pi i}, \mu) + r_N^{(K)}(we^{-\pi i}, \mu) \right) & \text{if } \arg w = 0, \\ r_N^{(K)}(we^{\mp\pi i}, \mu) & \text{if } 0 < \pm \arg w \leq \pi. \end{cases}$$

²Note that the condition $\operatorname{Re}(s + \mu + \nu) > 0$ is missing in reference [13] for formula (6.8.44) to be valid.

Here, N and M are arbitrary non-negative integers. Multiplying the expressions (4.1) and (4.3) together gives

$$(4.6) \quad I_\alpha(t \sinh \xi) K_\beta(t \cosh \xi) = \frac{\exp(-te^{-\xi})}{2\sqrt{\sinh \xi \cosh \xi} t} \left(\sum_{n=0}^{N-1} \sum_{\ell=0}^n (-1)^\ell \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sinh^\ell \xi \cosh^{n-\ell} \xi} \frac{1}{t^n} + R_{N,1}(t, \xi, \alpha, \beta) \right) \\ - C(\xi, \alpha) \frac{\exp(-te^\xi)}{2\sqrt{\sinh \xi \cosh \xi} t} \left(\sum_{m=0}^{M-1} \sum_{\ell=0}^m \frac{a_\ell(\alpha) a_{m-\ell}(\beta)}{\sinh^\ell \xi \cosh^{m-\ell} \xi} \frac{1}{t^m} + R_{M,2}(t, \xi, \alpha, \beta) \right),$$

where

$$(4.7) \quad R_{N,1}(t, \xi, \alpha, \beta) = \tilde{r}_N^{(K)}(t \sinh \xi, \alpha) r_0^{(K)}(t \cosh \xi, \beta) + \sum_{\ell=0}^{N-1} (-1)^\ell \frac{a_\ell(\alpha)}{\sinh^\ell \xi} \frac{1}{t^\ell} r_{N-\ell}^{(K)}(t \cosh \xi, \beta),$$

or

$$(4.8) \quad R_{N,1}(t, \xi, \alpha, \beta) = \tilde{r}_0^{(K)}(t \sinh \xi, \alpha) r_N^{(K)}(t \cosh \xi, \beta) + \sum_{\ell=1}^N \frac{a_{N-\ell}(\beta)}{\cosh^{N-\ell} \xi} \frac{1}{t^{N-\ell}} \tilde{r}_\ell^{(K)}(t \sinh \xi, \alpha),$$

and

$$(4.9) \quad R_{M,2}(t, \xi, \alpha, \beta) = r_M^{(K)}(t \sinh \xi, \alpha) r_0^{(K)}(t \cosh \xi, \beta) + \sum_{\ell=0}^{M-1} \frac{a_\ell(\alpha)}{\sinh^\ell \xi} \frac{1}{t^\ell} r_{M-\ell}^{(K)}(t \cosh \xi, \beta),$$

or

$$(4.10) \quad R_{M,2}(t, \xi, \alpha, \beta) = r_0^{(K)}(t \sinh \xi, \alpha) r_M^{(K)}(t \cosh \xi, \beta) + \sum_{\ell=1}^M \frac{a_{M-\ell}(\beta)}{\cosh^{M-\ell} \xi} \frac{1}{t^{M-\ell}} r_\ell^{(K)}(t \sinh \xi, \alpha).$$

Furthermore, we have defined $C(\xi, \alpha) = \tilde{C}(\arg(\sinh \xi), \alpha)$. The formula (1.10) for $C(\xi, \alpha)$ follows directly from (4.4) and the properties of the hyperbolic sine function. Assume that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \max(N, M)$ and $\operatorname{Re}(2\nu + \alpha + \beta + 2) > -\operatorname{Re}(\alpha) + |\operatorname{Re}(\beta)|$. Substituting equation (4.6) into (4.2), we obtain (2.2) with

$$R_N^{(P1)}(\nu, \xi, \alpha, \beta) = \frac{e^{-(2\nu+\alpha+\beta+1)\xi}}{\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} t^{2\nu+\alpha+\beta} \exp(-te^{-\xi}) R_{N,1}(t, \xi, \alpha, \beta) dt,$$

$$R_M^{(P2)}(\nu, \xi, \alpha, \beta) = \frac{e^{(2\nu+\alpha+\beta+1)\xi}}{\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} t^{2\nu+\alpha+\beta} \exp(-te^\xi) R_{M,2}(t, \xi, \alpha, \beta) dt.$$

We change the integration variable from t to s , setting $t = se^\xi$ for the first integral and $t = se^{-\xi}$ for the second. By applying Cauchy's theorem, the contours of integration can be deformed back onto the positive real axis, leading to the following representations:

$$(4.11) \quad R_N^{(P1)}(\nu, \xi, \alpha, \beta) = \frac{1}{\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} s^{2\nu+\alpha+\beta} e^{-s} R_{N,1}(se^\xi, \xi, \alpha, \beta) ds,$$

$$(4.12) \quad R_M^{(P2)}(\nu, \xi, \alpha, \beta) = \frac{1}{\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} s^{2\nu+\alpha+\beta} e^{-s} R_{M,2}(se^{-\xi}, \xi, \alpha, \beta) ds.$$

From Lemma 4.1, equations (4.5) and (4.7), and the properties of the function $e^\xi \sinh \xi$, we deduce the following bound:

(4.13)

$$\begin{aligned} |R_{N,1}(se^\xi, \xi, \alpha, \beta)| &\leq \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\alpha)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^N \right| \frac{1}{s^N} \\ &\quad \times \begin{cases} 1 & \text{if } \frac{\pi}{2} \leq |\vartheta| \leq \pi, \\ \min(|\csc \vartheta|, 1 + \chi(N + \frac{1}{2})) & \text{if } |\vartheta| < \frac{\pi}{2} \end{cases} \\ &\quad + \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \frac{1}{s^N}, \end{aligned}$$

where $\vartheta = \arg(e^\xi \sinh \xi) \in [-\pi, \pi]$, assuming $\xi \in \overline{\mathcal{D}}$, $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$, and $|\operatorname{Re}(\beta)| < \frac{1}{2}$. For the definition of $\overline{\mathcal{D}}$, refer to (3.1). Similarly, using the alternative expression (4.8), we obtain the bound

(4.14)

$$\begin{aligned} |R_{N,1}(se^\xi, \xi, \alpha, \beta)| &\leq \left(\left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_N(\operatorname{Re}(\beta)) \left(\frac{e^{-\xi}}{\cosh \xi} \right)^N \right| \frac{1}{s^N} \right. \\ &\quad \left. + \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\beta) \left(\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \right| \frac{1}{s^N} \right) \\ &\quad \times \begin{cases} 1 & \text{if } \frac{\pi}{2} \leq |\vartheta| \leq \pi, \\ \min(|\csc \vartheta|, 1 + \chi(N + \frac{1}{2})) & \text{if } |\vartheta| < \frac{\pi}{2}, \end{cases} \end{aligned}$$

provided that $\xi \in \overline{\mathcal{D}}$, $|\operatorname{Re}(\alpha)| < \frac{1}{2}$, and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$. Note that if either $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$, or $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$, then the condition $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \max(N, M)$ automatically implies $\operatorname{Re}(2\nu + \alpha + \beta + 2) > -\operatorname{Re}(\alpha) + |\operatorname{Re}(\beta)|$. Consequently, this latter condition can be omitted. In Section 7 of the paper [22], it is shown that the conditions involving ϑ can be equivalently expressed in terms of ξ as follows:

$$(4.15) \quad \begin{cases} 1 & \text{if } \operatorname{Re}(e^{2\xi}) \leq 1, \\ \min(|1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, 1 + \chi(N + \frac{1}{2})) & \text{if } \operatorname{Re}(e^{2\xi}) > 1. \end{cases}$$

From (4.11), we can infer that

$$\left| R_N^{(P1)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{1}{|\Gamma(2\nu + \alpha + \beta + 2)|} \int_0^{+\infty} s^{\operatorname{Re}(2\nu + \alpha + \beta)} e^{-s} |R_{N,1}(se^\xi, \xi, \alpha, \beta)| ds.$$

Restricting ξ to \mathcal{D} , substituting the estimates from (4.13) and (4.14), and applying (4.15) yields the bounds (2.3) and (2.4), respectively. In a similar manner, from Lemma 4.1, formulae (4.9) and (4.10), and the properties of the function $e^{-\xi} \sinh \xi$, we can deduce the following bounds:

(4.16)

$$\begin{aligned} |R_{M,2}(se^{-\xi}, \xi, \alpha, \beta)| &\leq \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_M(\operatorname{Re}(\alpha)) \left(\frac{e^\xi}{\sinh \xi} \right)^M \right| \frac{1}{s^M} \\ &\quad + \sum_{\ell=0}^{M-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} a_\ell(\alpha) a_{M-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{M-\ell} \right| \frac{1}{s^M}, \end{aligned}$$

provided that $\xi \in \overline{\mathcal{D}}$, $|\operatorname{Re}(\alpha)| < M + \frac{1}{2}$, and $|\operatorname{Re}(\beta)| < \frac{1}{2}$, and

(4.17)

$$\begin{aligned} |R_{M,2}(se^{-\xi}, \xi, \alpha, \beta)| &\leq \left| \frac{\cos(\pi\alpha)\cos(\pi\beta)}{\cos(\pi\operatorname{Re}(\alpha))\cos(\pi\operatorname{Re}(\beta))} a_M(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\cosh\xi}\right)^M \right| \frac{1}{s^M} \\ &\quad + \sum_{\ell=1}^M \left| \frac{\cos(\pi\alpha)}{\cos(\pi\operatorname{Re}(\alpha))} a_\ell(\operatorname{Re}(\alpha)) a_{M-\ell}(\beta) \left(\frac{e^\xi}{\sinh\xi}\right)^\ell \left(\frac{e^\xi}{\cosh\xi}\right)^{M-\ell} \right| \frac{1}{s^M} \end{aligned}$$

provided that $\xi \in \overline{\mathcal{D}}$, $|\operatorname{Re}(\alpha)| < \frac{1}{2}$, and $|\operatorname{Re}(\beta)| < M + \frac{1}{2}$. The bounds (2.6) and (2.7) now follow similarly to those of (2.3) and (2.4). The estimates (2.5) and (2.8) can be derived from (2.4) and (2.7) by applying the inequality

$$(4.18) \quad |a_\ell(\beta)| \leq \left| \frac{\cos(\pi\beta)}{\cos(\pi\operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\beta)) \right|, \quad |\operatorname{Re}(\beta)| < \ell + \frac{1}{2},$$

which follows from [21, Eq. (69)].

Now, assume that $\xi > 0$, and that ν , α , and β are real, with $|\alpha|, |\beta| < \frac{1}{2}$, and $2\nu + \alpha + \beta + 1 > M$. From Lemma 4.1, it follows that

$$r_M^{(K)}(w, \mu) = \theta_M(w, \mu) \frac{a_M(\mu)}{w^M},$$

where $w > 0$, μ is real, $|\mu| < M + \frac{1}{2}$, and $0 < \theta_M(w, \mu) < 1$. Applying this result to (4.9) gives

$$\begin{aligned} R_{M,2}(se^{-\xi}, \xi, \alpha, \beta) &= \theta_M(se^{-\xi} \sinh \xi, \alpha) \theta_0(se^{-\xi} \cosh \xi, \beta) a_M(\alpha) \left(\frac{e^\xi}{\sinh \xi}\right)^M \frac{1}{s^M} \\ &\quad + \sum_{\ell=0}^{M-1} \theta_{M-\ell}(se^{-\xi} \cosh \xi, \beta) a_\ell(\alpha) a_{M-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi}\right)^\ell \left(\frac{e^\xi}{\cosh \xi}\right)^{M-\ell} \frac{1}{s^M} \\ &= \tilde{\theta}_M(s, \xi, \alpha, \beta) \sum_{\ell=0}^{M-1} a_\ell(\alpha) a_{M-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi}\right)^\ell \left(\frac{e^\xi}{\cosh \xi}\right)^{M-\ell} \frac{1}{s^M}, \end{aligned}$$

with a suitable $0 < \tilde{\theta}_M(s, \xi, \alpha, \beta) < 1$, since $a_\ell(\alpha) a_{M-\ell}(\beta) = (-1)^M |a_\ell(\alpha) a_{M-\ell}(\beta)|$ when $|\alpha|, |\beta| < \frac{1}{2}$. Hence, by applying the mean value theorem for improper integrals, we deduce that

$$\begin{aligned} R_M^{(P2)}(\nu, \xi, \alpha, \beta) &= \sum_{\ell=0}^{M-1} a_\ell(\alpha) a_{M-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi}\right)^\ell \left(\frac{e^\xi}{\cosh \xi}\right)^{M-\ell} \\ &\quad \times \frac{1}{\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} \tilde{\theta}_M(s, \xi, \alpha, \beta) s^{2\nu + \alpha + \beta - M} e^{-s} ds \\ &= \Theta_M(\nu, \xi, \alpha, \beta) \sum_{\ell=0}^{M-1} a_\ell(\alpha) a_{M-\ell}(\beta) \left(\frac{e^\xi}{\sinh \xi}\right)^\ell \left(\frac{e^\xi}{\cosh \xi}\right)^{M-\ell} \frac{\Gamma(2\nu + \alpha + \beta - M + 1)}{\Gamma(2\nu + \alpha + \beta + 2)} \end{aligned}$$

where $0 < \Theta_M(\nu, \xi, \alpha, \beta) < 1$. This concludes the proof of Theorem 2.1.

We continue with the proof of Theorem 2.2. The Jacobi function of the second kind can be expressed in terms of the modified Bessel function of the second kind as follows:

$$\begin{aligned} 2^{\nu-1} B(\nu+1, \nu+\alpha+\beta+1) Q_\nu^{(\alpha, \beta)}(z) &= \frac{1}{(z-1)^{\alpha/2} (z+1)^{\beta/2}} \\ &\quad \times \frac{1}{\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} t^{2\nu + \alpha + \beta + 1} K_\alpha(t\sqrt{z-1}) K_\beta(t\sqrt{z+1}) dt, \end{aligned}$$

provided that $\operatorname{Re}(\sqrt{z-1} + \sqrt{z+1}) > 0$ and $\operatorname{Re}(2\nu + \alpha + \beta + 2) > |\operatorname{Re}(\alpha)| + |\operatorname{Re}(\beta)|$ (combine [5, Eq. (21)] and [13, Eq. (6.8.47)]). By applying an appropriate change of variables, this can be rewritten as (1.23). Let N be an arbitrary non-negative integer. Multiplying two copies of the expression in (4.1) yields

$$(4.19) \quad K_\alpha(t \sinh \xi) K_\beta(t \cosh \xi) = \frac{\pi}{2} \frac{\exp(-te^\xi)}{\sqrt{\sinh \xi \cosh \xi} t} \left(\sum_{n=0}^{N-1} \sum_{\ell=0}^n \frac{a_\ell(\alpha) a_{n-\ell}(\beta)}{\sinh^\ell \xi \cosh^{n-\ell} \xi} \frac{1}{t^n} + R_{N,2}(t, \xi, \alpha, \beta) \right),$$

where $R_{N,2}(t, \xi, \alpha, \beta)$ is given by either (4.9) or (4.10), with N in place of M . Assume that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > N$ and $\operatorname{Re}(2\nu + \alpha + \beta + 2) > |\operatorname{Re}(\alpha)| + |\operatorname{Re}(\beta)|$. By substituting equation (4.19) into (1.23), we obtain (2.9) with $R_N^{(Q)}(\nu, \xi, \alpha, \beta) = R_N^{(P2)}(\nu, \xi, \alpha, \beta)$. Therefore, the estimates for $R_N^{(Q)}(\nu, \xi, \alpha, \beta)$ follow directly from those for $R_N^{(P2)}(\nu, \xi, \alpha, \beta)$. Note that if either $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$, or $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$, then the condition $\operatorname{Re}(2\nu + \alpha + \beta + 1) > N$ automatically ensures that $\operatorname{Re}(2\nu + \alpha + \beta + 2) > |\operatorname{Re}(\alpha)| + |\operatorname{Re}(\beta)|$. Therefore, the latter condition can be omitted.

We conclude this section by showing that the inverse factorial expansions (1.11) and (1.12) terminate, providing an exact representation of the corresponding functions when both 2α and 2β are odd integers. Let L denote the positive integer defined by $\max(2|\alpha|, 2|\beta|) = 2L - 1$. Then, by definition (1.8) and Lemma 4.1, we have

$$a_\ell(L - \frac{1}{2}) = 0, \quad r_\ell^{(K)}(w, L - \frac{1}{2}) = 0,$$

for all $\ell \geq L$. Therefore, from (4.5), (4.7), and (4.9), it follows that $R_{2L,1}(t, \xi, \alpha, \beta) = 0$ and $R_{2L,2}(t, \xi, \alpha, \beta) = 0$. Hence, by (4.11) and (4.12), we find that $R_{2L}^{(P1)}(\nu, \xi, \alpha, \beta) = 0$ and $R_{2L}^{(P2)}(\nu, \xi, \alpha, \beta) = 0$, provided $\operatorname{Re}(2\nu + \alpha + \beta + 1) > 2L$. Thus, by Theorems 2.1 and 2.2, the expansions (1.11) and (1.12) terminate, giving exact representations of the corresponding functions. The conditions on the parameters may be relaxed by appealing to analytic continuation.

5. PROOF OF THEOREM 2.3

In this section, we establish the error bounds presented in Theorem 2.3. Suppose that $0 < \zeta < \frac{\pi}{2}$. Assuming that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > N$ and $\operatorname{Re}(2\nu + \alpha + \beta + 2) > |\operatorname{Re}(\alpha)| + |\operatorname{Re}(\beta)|$, and noting that $R_N^{(Q)}(\nu, \xi, \alpha, \beta) = R_N^{(P2)}(\nu, \xi, \alpha, \beta)$, we substitute (2.9) into the right-hand side of (3.6) and apply the representation (4.12). This results in (2.10) with

$$\begin{aligned} R_N^{(P)}(\nu, \zeta, \alpha, \beta) &= \frac{1}{2} e^{-((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i} \lim_{\varepsilon \rightarrow 0^+} R_N^{(Q)}(\nu, \varepsilon + i\zeta, \alpha, \beta) \\ &\quad + \frac{1}{2} e^{((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i} \lim_{\varepsilon \rightarrow 0^+} R_N^{(Q)}(\nu, \varepsilon - i\zeta, \alpha, \beta) \\ &= \frac{e^{-((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i}}{2\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} s^{2\nu + \alpha + \beta} e^{-s} R_{N,2}(se^{-i\zeta}, i\zeta, \alpha, \beta) ds \\ &\quad + \frac{e^{((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i}}{2\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} s^{2\nu + \alpha + \beta} e^{-s} R_{N,2}(se^{i\zeta}, -i\zeta, \alpha, \beta) ds. \end{aligned}$$

A straightforward estimation yields the bound

$$(5.1) \quad \begin{aligned} |R_N^{(P)}(\nu, \zeta, \alpha, \beta)| &\leq \frac{e^{\operatorname{Im}(\zeta_{\nu, N, 0}^{(1)})}}{2\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} s^{\operatorname{Re}(2\nu + \alpha + \beta)} e^{-s} |R_{N,2}(se^{-i\zeta}, i\zeta, \alpha, \beta)| \, ds \\ &\quad + \frac{e^{-\operatorname{Im}(\zeta_{\nu, N, 0}^{(1)})}}{2\Gamma(2\nu + \alpha + \beta + 2)} \int_0^{+\infty} s^{\operatorname{Re}(2\nu + \alpha + \beta)} e^{-s} |R_{N,2}(se^{i\zeta}, -i\zeta, \alpha, \beta)| \, ds. \end{aligned}$$

Using the estimates in (4.16) and (4.17) with $\pm i\zeta$ replacing ξ , we obtain

$$\begin{aligned} |R_{N,2}(se^{\mp i\zeta}, \pm i\zeta, \alpha, \beta)| &\leq \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_N(\operatorname{Re}(\alpha))}{\sin^N \zeta} \right| \frac{1}{s^N} \\ &\quad + \sum_{\ell=0}^{N-1} \left| \frac{\cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\beta))} \frac{a_\ell(\alpha) a_{N-\ell}(\operatorname{Re}(\beta))}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \frac{1}{s^N}, \end{aligned}$$

provided that $|\operatorname{Re}(\alpha)| < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$, and

$$\begin{aligned} |R_{N,2}(se^{\mp i\zeta}, \pm i\zeta, \alpha, \beta)| &\leq \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} \frac{a_N(\operatorname{Re}(\beta))}{\cos^N \zeta} \right| \frac{1}{s^N} \\ &\quad + \sum_{\ell=1}^N \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\beta)}{\sin^\ell \zeta \cos^{N-\ell} \zeta} \right| \frac{1}{s^N}, \end{aligned}$$

assuming $|\operatorname{Re}(\alpha)| < \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < N + \frac{1}{2}$. By substituting these estimates into (5.1) and observing that $\operatorname{Im}(\zeta_{\nu, N, 0}^{(1)}) = \operatorname{Im}(\zeta_{\nu, N, \ell}^{(1)})$ for all $0 \leq \ell \leq N$, we obtain the bounds (2.13) and (2.14), respectively. The estimate (2.15) can be derived from (2.14) by applying the inequality (4.18). The condition $\operatorname{Re}(2\nu + \alpha + \beta + 2) > |\operatorname{Re}(\alpha)| + |\operatorname{Re}(\beta)|$ can be omitted for the same reason explained in the previous section.

The bounds for the remainder terms $R_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ and $R_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ in the truncated inverse factorial series (2.11) and (2.12) can be derived in a similar fashion by combining (2.9) with (3.7) and (3.8), respectively. The details are left for the reader to verify.

The integral representation of $R_N^{(P)}(\nu, \zeta, \alpha, \beta)$, along with the analogous representations for $R_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ and $R_N^{(Q)}(\nu, \zeta, \alpha, \beta)$, as well as the discussion at the end of the previous section, demonstrate that the inverse factorial expansions (1.14), (1.15), and (1.16) terminate, providing an exact representation of the corresponding functions when both 2α and 2β are odd integers.

6. PROOF OF THEOREMS 2.4 AND 2.5

In this section, we establish the error bounds given in Theorems 2.4 and 2.5. To do so, we first need appropriate integral representations of the Jacobi functions. Hahn [17, Eqs. (9.6) and (14.4)] showed that if $\operatorname{Re}(\xi) > 0$, $-\frac{\pi}{2} < \operatorname{Im}(\xi) < 0$ and $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$, then

$$\begin{aligned} \frac{\pi P_\nu^{(\alpha, \beta)}(\cosh(2\xi))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} &= \\ \left(\frac{e^\xi}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^\xi}{\cosh \xi} \right)^{\beta + 1/2} e^{2\nu\xi} Y_+(\xi) &- ie^{-\pi i\alpha} \left(\frac{e^{-\xi}}{\sinh \xi} \right)^{\alpha + 1/2} \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{\beta + 1/2} e^{-2\nu\xi} Y_-(\xi), \end{aligned}$$

where

$$(6.1) \quad Y_{\pm}(\xi) = \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} \\ \times \int_0^1 \int_0^1 t^{\alpha' + \beta'} (1-t)^{2\nu + \alpha + \beta - \alpha' - \beta'} s^{\alpha' - 1/2} (1-s)^{\beta' - 1/2} \\ \times \left(1 \mp ts \frac{e^{\pm\xi}}{2 \sinh \xi}\right)^{\alpha' - 1/2} \left(1 - t(1-s) \frac{e^{\pm\xi}}{2 \cosh \xi}\right)^{\beta' - 1/2} ds dt.$$

(In Hahn's notation, $Y_{\pm}(\xi) = y\left(\frac{e^{\pm\xi}}{2 \cosh \xi}\right)$.) Here, α' and β' are defined in (2.16). Note that, in Hahn's original formulation, ν , α , and β are assumed to be real, but the derivation remains valid for complex parameters as well. By the Schwarz reflection principle, we have $P_{\nu}^{(\alpha, \beta)}(\cosh(2\xi)) = \overline{P_{\nu}^{(\alpha, \beta)}(\cosh(2\bar{\xi}))}$, assuming that ν , α , and β are real. Using this observation along with analytic continuation, we obtain

$$(6.2) \quad \frac{\pi P_{\nu}^{(\alpha, \beta)}(\cosh(2\xi))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \\ \left(\frac{e^{\xi}}{\sinh \xi}\right)^{\alpha + 1/2} \left(\frac{e^{\xi}}{\cosh \xi}\right)^{\beta + 1/2} e^{2\nu\xi} Y_{+}(\xi) - C(\xi, \alpha) \left(\frac{e^{-\xi}}{\sinh \xi}\right)^{\alpha + 1/2} \left(\frac{e^{-\xi}}{\cosh \xi}\right)^{\beta + 1/2} e^{-2\nu\xi} Y_{-}(\xi),$$

where $C(\xi, \alpha)$ is defined in (1.10), $\xi \in \mathcal{D} \setminus \mathbb{R}^{+}$ and $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$.

From the hypergeometric representation (1.24) and Hahn's results [17, Eqs. (8.2) and (8.21)], we can assert that

$$(6.3) \quad \frac{Q_{\nu}^{(\alpha, \beta)}(\cosh(2\xi))}{2^{2\nu + \alpha + \beta} B(\nu + \alpha + 1, \nu + \beta + 1)} = \left(\frac{e^{-\xi}}{\sinh \xi}\right)^{\alpha + 1/2} \left(\frac{e^{-\xi}}{\cosh \xi}\right)^{\beta + 1/2} e^{-2\nu\xi} Y_{-}(\xi),$$

provided that $\xi \in \mathcal{D}$ and $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$.

To continue, we first need to establish the following lemma. Here, $\overline{\mathcal{D}}$ is defined by (3.1).

Lemma 6.1. *Let N be a non-negative integer, and let A and B be complex numbers. Let u , v_1 , and v_2 be real numbers such that $0 < u, v_1, v_2 < 1$. For $\xi \in \overline{\mathcal{D}} \setminus \mathbb{R}^{+}$, define the remainder $\widehat{R}_N^{+}(u, v_1, v_2, \xi, A, B)$, and for $\xi \in \overline{\mathcal{D}}$, define the remainder $\widehat{R}_N^{-}(u, v_1, v_2, \xi, A, B)$ by*

$$(6.4) \quad \left(1 \mp uv_1 \frac{e^{\pm\xi}}{2 \sinh \xi}\right)^{-A} \left(1 - uv_2 \frac{e^{\pm\xi}}{2 \cosh \xi}\right)^{-B} = \\ \sum_{n=0}^{N-1} \left(\sum_{\ell=0}^n \frac{(A)_{\ell} (B)_{n-\ell}}{\ell! (n-\ell)!} \left(\pm v_1 \frac{e^{\pm\xi}}{2 \sinh \xi}\right)^{\ell} \left(v_2 \frac{e^{\pm\xi}}{2 \cosh \xi}\right)^{n-\ell}\right) u^n + \widehat{R}_N^{\pm}(u, v_1, v_2, \xi, A, B).$$

Then, we have the following bounds:

(6.5)

$$\begin{aligned} \left| \widehat{R}_N^+(u, v_1, v_2, \xi, A, B) \right| &\leq \left| \frac{\sin(\pi A) \sin(\pi B)}{\sin(\pi \operatorname{Re}(A)) \sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(A))_N}{N!} \left(v_1 \frac{e^\xi}{2 \sinh \xi} \right)^N \right| u^N \\ &\quad \times |1 - e^{4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))| \\ &\quad + \sum_{\ell=0}^{N-1} \left| \frac{\sin(\pi B)}{\sin(\pi \operatorname{Re}(B))} \frac{(A)_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left(v_1 \frac{e^\xi}{2 \sinh \xi} \right)^\ell \left(v_2 \frac{e^\xi}{2 \cosh \xi} \right)^{N-\ell} \right| u^N \\ &\quad \times \begin{cases} |1 + e^{2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{-2\xi}) \leq 0, \\ |1 + e^{2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{-2\xi}), \end{cases} \end{aligned}$$

provided that $-N < \operatorname{Re}(A) < 1$ and $0 < \operatorname{Re}(B) < 1$,

(6.6)

$$\begin{aligned} \left| \widehat{R}_N^+(u, v_1, v_2, \xi, A, B) \right| &\leq \left| \frac{\sin(\pi A) \sin(\pi B)}{\sin(\pi \operatorname{Re}(A)) \sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(B))_N}{N!} \left(v_2 \frac{e^\xi}{2 \cosh \xi} \right)^N \right| u^N \\ &\quad \times |1 - e^{4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))| \\ &\quad + \sum_{\ell=1}^N \left| \frac{\sin(\pi A)}{\sin(\pi \operatorname{Re}(A))} \frac{(\operatorname{Re}(A))_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left(v_1 \frac{e^\xi}{2 \sinh \xi} \right)^\ell \left(v_2 \frac{e^\xi}{2 \cosh \xi} \right)^{N-\ell} \right| u^N \\ &\quad \times \begin{cases} |1 - e^{2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{-2\xi}) < 1, \\ |1 - e^{2\xi}|, & \text{if } \operatorname{Re}(e^{-2\xi}) < 0, \end{cases} \end{aligned}$$

provided that $0 < \operatorname{Re}(A) < 1$ and $-N < \operatorname{Re}(B) < 1$,

(6.7)

$$\begin{aligned} \left| \widehat{R}_N^-(u, v_1, v_2, \xi, A, B) \right| &\leq \left| \frac{\sin(\pi A) \sin(\pi B)}{\sin(\pi \operatorname{Re}(A)) \sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(A))_N}{N!} \left(v_1 \frac{e^{-\xi}}{2 \sinh \xi} \right)^N \right| u^N \\ &\quad \times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1 \end{cases} \\ &\quad + \sum_{\ell=0}^{N-1} \left| \frac{\sin(\pi B)}{\sin(\pi \operatorname{Re}(B))} \frac{(A)_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left(v_1 \frac{e^{-\xi}}{2 \sinh \xi} \right)^\ell \left(v_2 \frac{e^{-\xi}}{2 \cosh \xi} \right)^{N-\ell} \right| u^N \\ &\quad \times \begin{cases} 1, & \text{if } \operatorname{Re}(e^{2\xi}) \leq -1, \\ |1 + e^{-2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{2\xi}) \leq 0, \\ |1 + e^{-2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{2\xi}), \end{cases} \end{aligned}$$

provided that $-N < \operatorname{Re}(A) < 1$ and $0 < \operatorname{Re}(B) < 1$, and

(6.8)

$$\begin{aligned} \left| \widehat{R}_N^-(u, v_1, v_2, \xi, A, B) \right| &\leq \left| \frac{\sin(\pi A) \sin(\pi B)}{\sin(\pi \operatorname{Re}(A)) \sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(B))_N}{N!} \left(v_2 \frac{e^{-\xi}}{2 \cosh \xi} \right)^N \right| u^N \\ &\times \begin{cases} |1 \pm e^{-2\xi}|, & \text{if } 1 \leq \pm \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-4\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } |\operatorname{Re}(e^{2\xi})| < 1 \end{cases} \\ &+ \sum_{\ell=1}^N \left| \frac{\sin(\pi A)}{\sin(\pi \operatorname{Re}(A))} \frac{(\operatorname{Re}(A))_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left(v_1 \frac{e^{-\xi}}{2 \sinh \xi} \right)^\ell \left(v_2 \frac{e^{-\xi}}{2 \cosh \xi} \right)^{N-\ell} \right| u^N \\ &\times \begin{cases} 1, & \text{if } 1 \leq \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{2\xi}) < 1, \\ |1 - e^{-2\xi}|, & \text{if } \operatorname{Re}(e^{2\xi}) < 0, \end{cases} \end{aligned}$$

provided that $0 < \operatorname{Re}(A) < 1$ and $-N < \operatorname{Re}(B) < 1$. If A or B is an integer, then the limiting values must be taken in these bounds. Moreover, when ξ , A , and B are real with $\xi > 0$ and $0 < A, B < 1$, we have

$$(6.9) \quad 0 < \widehat{R}_N^-(u, v_1, v_2, \xi, A, B) < 2 \sum_{\ell=0}^N \left(\frac{(A)_\ell (B)_{N-\ell}}{\ell! (N-\ell)!} \left(-v_1 \frac{e^{-\xi}}{2 \sinh \xi} \right)^\ell \left(v_2 \frac{e^{-\xi}}{2 \cosh \xi} \right)^{N-\ell} \right) u^N.$$

Proof. Let N be a non-negative integer, let A and B be complex numbers, and let u be a real number such that $0 < u < 1$. For $\xi \in \overline{\mathcal{D}} \setminus \mathbb{R}^+$, define the remainder $\widehat{r}_{N,1}^\pm(u, \xi, A)$, and for $\xi \in \overline{\mathcal{D}}$, define the remainders $\widehat{r}_{N,1}^\pm(u, \xi, A)$ and $\widehat{r}_{N,2}^\pm(u, \xi, B)$ by

$$\left(1 \mp u \frac{e^{\pm \xi}}{2 \sinh \xi} \right)^{-A} = \sum_{n=0}^{N-1} \frac{(A)_n}{n!} \left(\pm \frac{e^{\pm \xi}}{2 \sinh \xi} \right)^n u^n + \widehat{r}_{N,1}^\pm(u, \xi, A)$$

and

$$\left(1 - u \frac{e^{\pm \xi}}{2 \cosh \xi} \right)^{-B} = \sum_{n=0}^{N-1} \frac{(B)_n}{n!} \left(\frac{e^{\pm \xi}}{2 \cosh \xi} \right)^n u^n + \widehat{r}_{N,2}^\pm(u, \xi, B),$$

respectively. Using this notation, the remainders $\widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B)$ can be expressed as

$$(6.10) \quad \begin{aligned} \widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B) &= \widehat{r}_{N,1}^\pm(uv_1, \xi, A) \widehat{r}_{0,2}^\pm(uv_2, \xi, B) \\ &+ \sum_{\ell=0}^{N-1} \frac{(A)_\ell}{\ell!} \left(\pm v_1 \frac{e^{\pm \xi}}{2 \sinh \xi} \right)^\ell u^\ell \widehat{r}_{N-\ell,2}^\pm(uv_2, \xi, B), \end{aligned}$$

or

$$(6.11) \quad \begin{aligned} \widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B) &= \widehat{r}_{0,1}^\pm(uv_1, \xi, A) \widehat{r}_{N,2}^\pm(uv_2, \xi, B) \\ &+ \sum_{\ell=1}^N \frac{(B)_{N-\ell}}{(N-\ell)!} \left(v_2 \frac{e^{\pm \xi}}{2 \cosh \xi} \right)^{N-\ell} u^{N-\ell} \widehat{r}_{\ell,1}^\pm(uv_1, \xi, A). \end{aligned}$$

To obtain the required bounds, we will estimate the remainders $\widehat{r}_{N,1}^\pm(u, \xi, A)$ and $\widehat{r}_{N,2}^\pm(u, \xi, B)$. Using the standard integral representation of the beta function [23, Eq. (5.12.1)], we have

$$\frac{(A)_n}{n!} = \frac{\sin(\pi A)}{\pi} \int_0^1 t^{A+n-1} (1-t)^{-A} dt,$$

valid for $-n < \operatorname{Re}(A) < 1$. Summation and analytic continuation yields

$$(1 - uw)^{-A} = \sum_{n=0}^{N-1} \frac{(A)_n}{n!} u^n w^n + M_N(u, w, A),$$

where

$$M_N(u, w, A) = u^N w^N \frac{\sin(\pi A)}{\pi} \int_0^1 \frac{t^{A+N-1} (1-t)^{-A}}{1-tuw} dt,$$

provided that $0 < u < 1$, $w \in \mathbb{C} \setminus [1, +\infty)$, and $-N < \operatorname{Re}(A) < 1$. For $0 < t, u < 1$, we note that

$$|1 - tuw|^2 = |w|^2 t^2 u^2 - 2 \operatorname{Re}(w) tu + 1 \geq \begin{cases} 1 & \text{if } \operatorname{Re}(w) \leq 0, \\ \frac{(\operatorname{Im}(w))^2}{|w|^2} & \text{if } 0 < \operatorname{Re}(w) \leq |w|^2, \\ |1 - w|^2 & \text{if } \operatorname{Re}(w) > |w|^2. \end{cases}$$

Consequently, we obtain

$$(6.12) \quad |M_N(u, w, A)| \leq u^N |w|^N \frac{|\sin(\pi A)|}{\pi} \int_0^1 \frac{t^{\operatorname{Re}(A)+N-1} (1-t)^{-\operatorname{Re}(A)}}{|1-tuw|} dt \\ \leq \left| \frac{\sin(\pi A)}{\sin(\pi \operatorname{Re}(A))} \frac{(\operatorname{Re}(A))_N}{N!} w^N \right| u^N \times \begin{cases} 1 & \text{if } \operatorname{Re}(w) \leq 0, \\ \left| \frac{w}{\operatorname{Im}(w)} \right| & \text{if } 0 < \operatorname{Re}(w) \leq |w|^2, \\ \frac{1}{|1-w|} & \text{if } \operatorname{Re}(w) > |w|^2. \end{cases}$$

If A is an integer, then the limiting value must to be taken in this bound. Additionally, when $0 < u < 1$, $0 < w \leq \frac{1}{2}$, and $0 < A < 1$, we have

$$(6.13) \quad 0 < M_N(u, w, A) < 2u^N w^N \frac{\sin(\pi A)}{\pi} \int_0^1 t^{A+N-1} (1-t)^{-A} dt = 2 \frac{(A)_N}{N!} w^N u^N.$$

Similarly, when $0 < u < 1$, $w < 0$, and $0 < A < 1$, we obtain

$$(6.14) \quad 0 < M_N(u, w, A) < u^N w^N \frac{\sin(\pi A)}{\pi} \int_0^1 t^{A+N-1} (1-t)^{-A} dt = \frac{(A)_N}{N!} w^N u^N.$$

Observing that

$$(6.15) \quad \widehat{r}_{N,1}^{\pm}(u, \xi, A) = M_N \left(u, \pm \frac{e^{\pm \xi}}{2 \sinh \xi}, A \right), \quad \widehat{r}_{N,2}^{\pm}(u, \xi, B) = M_N \left(u, \frac{e^{\pm \xi}}{2 \cosh \xi}, B \right)$$

and applying (6.12) to the right-hand sides, we obtain the following bounds:

$$(6.16) \quad \left| \widehat{r}_{N,1}^+(u, \xi, A) \right| \leq \left| \frac{\sin(\pi A)}{\sin(\pi \operatorname{Re}(A))} \frac{(\operatorname{Re}(A))_N}{N!} \left(\frac{e^{\xi}}{2 \sinh \xi} \right)^N \right| u^N \\ \times \begin{cases} |1 - e^{2\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{-2\xi}) < 1, \\ |1 - e^{2\xi}|, & \text{if } \operatorname{Re}(e^{-2\xi}) < 0, \end{cases}$$

$$(6.17) \quad \left| \widehat{r}_{N,1}^-(u, \xi, A) \right| \leq \left| \frac{\sin(\pi A)}{\sin(\pi \operatorname{Re}(A))} \frac{(\operatorname{Re}(A))_N}{N!} \left(\frac{e^{-\xi}}{2 \sinh \xi} \right)^N \right| u^N \\ \times \begin{cases} 1, & \text{if } 1 \leq \operatorname{Re}(e^{2\xi}), \\ |1 - e^{-2\xi}| |\csc(2 \operatorname{Im}(\xi))|, & \text{if } 0 \leq \operatorname{Re}(e^{2\xi}) < 1, \\ |1 - e^{-2\xi}|, & \text{if } \operatorname{Re}(e^{2\xi}) < 0, \end{cases}$$

provided $-N < \operatorname{Re}(A) < 1$, and

$$(6.18) \quad \left| \widehat{r}_{N,2}^+(u, \xi, B) \right| \leq \left| \frac{\sin(\pi B)}{\sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(B))_N}{N!} \left(\frac{e^\xi}{2 \cosh \xi} \right)^N \right| u^N \\ \times \begin{cases} |1 + e^{2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{-2\xi}) \leq 0, \\ |1 + e^{2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{-2\xi}), \end{cases}$$

$$(6.19) \quad \left| \widehat{r}_{N,2}^-(u, \xi, B) \right| \leq \left| \frac{\sin(\pi B)}{\sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(B))_N}{N!} \left(\frac{e^{-\xi}}{2 \cosh \xi} \right)^N \right| u^N \\ \times \begin{cases} 1, & \text{if } \operatorname{Re}(e^{2\xi}) \leq -1, \\ |1 + e^{-2\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|, & \text{if } -1 < \operatorname{Re}(e^{2\xi}) \leq 0, \\ |1 + e^{-2\xi}|, & \text{if } 0 < \operatorname{Re}(e^{2\xi}), \end{cases}$$

provided $-N < \operatorname{Re}(B) < 1$.

Straightforward estimations in (6.10) and (6.11) provide the bounds

$$(6.20) \quad \left| \widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B) \right| \leq \left| \widehat{r}_{N,1}^\pm(uv_1, \xi, A) \widehat{r}_{0,2}^\pm(uv_2, \xi, B) \right| \\ + \sum_{\ell=0}^{N-1} \left| \frac{(A)_\ell}{\ell!} \left(\pm v_1 \frac{e^{\pm\xi}}{2 \sinh \xi} \right)^\ell \right| u^\ell \left| \widehat{r}_{N-\ell,2}^\pm(uv_2, \xi, B) \right|,$$

and

$$(6.21) \quad \left| \widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B) \right| \leq \left| \widehat{r}_{0,1}^\pm(uv_1, \xi, A) \widehat{r}_{N,2}^\pm(uv_2, \xi, B) \right| \\ + \sum_{\ell=1}^N \left| \frac{(B)_{N-\ell}}{(N-\ell)!} \left(v_2 \frac{e^{\pm\xi}}{2 \cosh \xi} \right)^{N-\ell} \right| u^{N-\ell} \left| \widehat{r}_{\ell,1}^\pm(uv_1, \xi, A) \right|,$$

respectively. Applying the inequalities in (6.16)–(6.19) to the right-hand side of (6.20) yields the bounds (6.5) and (6.7). Similarly, applying the same inequalities to the right-hand side of (6.21) yields the bounds (6.6) and (6.8).

Using (6.13) and (6.14) on the right-hand sides of (6.15), we obtain the inequalities

$$0 < \widehat{r}_{N,1}^-(u, \xi, A) < \frac{(A)_N}{N!} \left(-\frac{e^{-\xi}}{2 \sinh \xi} \right)^N u^N, \quad 0 < \widehat{r}_{N,2}^-(u, \xi, B) < 2 \frac{(B)_N}{N!} \left(\frac{e^{-\xi}}{2 \cosh \xi} \right)^N u^N,$$

for $\xi > 0$ and $0 < A, B < 1$. Applying these inequalities to the right-hand side of (6.10) yields (6.9). \square

We now proceed with the proof of Theorem 2.4. Assume that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. Substituting into (6.1) and using (6.4) with the values $u = t$, $v_1 = s$, $v_2 = 1 - s$, $A = \frac{1}{2} - \alpha'$, and $B = \frac{1}{2} - \beta'$, while taking into account (6.2), we derive (2.17) with

$$(6.22) \quad \widehat{R}_N^{(P_1)}(\nu, \xi, \alpha, \beta) = \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1) \Gamma(\alpha' + \frac{1}{2}) \Gamma(\beta' + \frac{1}{2})} \\ \times \int_0^1 \int_0^1 t^{\alpha' + \beta'} (1-t)^{2\nu + \alpha + \beta - \alpha' - \beta'} s^{\alpha' - 1/2} (1-s)^{\beta' - 1/2} \widehat{R}_N^+(t, s, 1-s, \xi, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') ds dt,$$

(6.23)

$$\begin{aligned} \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) &= \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} \\ &\times \int_0^1 \int_0^1 t^{\alpha' + \beta'} (1-t)^{2\nu + \alpha + \beta - \alpha' - \beta'} s^{\alpha' - 1/2} (1-s)^{\beta' - 1/2} \widehat{R}_M^-(t, s, 1-s, \xi, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') \, ds \, dt. \end{aligned}$$

In deriving the form (2.17), we used the fact that $a_\ell(\alpha') = a_\ell(\alpha)$ and $a_\ell(\beta') = a_\ell(\beta)$ for all ℓ . Straightforward estimation gives

(6.24)

$$\begin{aligned} \left| \widehat{R}_N^{(P_1)}(\nu, \xi, \alpha, \beta) \right| &\leq \left| \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} \right| \\ &\times \int_0^1 \int_0^1 t^{\operatorname{Re}(\alpha' + \beta')} (1-t)^{\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta')} s^{\operatorname{Re}(\alpha') - 1/2} (1-s)^{\operatorname{Re}(\beta') - 1/2} \\ &\times \left| \widehat{R}_N^+(t, s, 1-s, \xi, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') \right| \, ds \, dt, \end{aligned}$$

(6.25)

$$\begin{aligned} \left| \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) \right| &\leq \left| \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} \right| \\ &\times \int_0^1 \int_0^1 t^{\operatorname{Re}(\alpha' + \beta')} (1-t)^{\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta')} s^{\operatorname{Re}(\alpha') - 1/2} (1-s)^{\operatorname{Re}(\beta') - 1/2} \\ &\times \left| \widehat{R}_M^-(t, s, 1-s, \xi, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') \right| \, ds \, dt. \end{aligned}$$

Restricting ξ to $\mathcal{D} \setminus \mathbb{R}^+$, we substitute the estimates from (6.5), (6.6), (6.7), and (6.8). Noting that $a_\ell(\alpha') = a_\ell(\alpha)$, $a_\ell(\operatorname{Re}(\alpha')) = a_\ell(\operatorname{Re}(\alpha))$, $a_\ell(\beta') = a_\ell(\beta)$ and $a_\ell(\operatorname{Re}(\beta')) = a_\ell(\operatorname{Re}(\beta))$ for all ℓ , this yields the bounds (2.18), (2.19), (2.21), and (2.22), respectively. Note that, for example, in deriving (2.18), the conditions $-\frac{1}{2} < \operatorname{Re}(\alpha') < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta')| < \frac{1}{2}$ apply. However, these are equivalent to $-\frac{1}{2} < \operatorname{Re}(\alpha) < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$. A similar remark applies to the other bounds.

The estimates (2.20) and (2.23) can be derived from (2.18) and (2.21) by applying the following inequality:

(6.26)

$$\begin{aligned} \left| \frac{\Gamma(\operatorname{Re}(\alpha') + \ell + \frac{1}{2})}{\Gamma(\alpha' + \ell + \frac{1}{2})} a_\ell(\alpha) \right| &= \left| \frac{\Gamma(\operatorname{Re}(\alpha') + \ell + \frac{1}{2})}{\Gamma(\alpha' + \ell + \frac{1}{2})} a_\ell(\alpha') \right| \\ &= \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{\Gamma(-\alpha' + \ell + \frac{1}{2})}{\Gamma(-\operatorname{Re}(\alpha') + \ell + \frac{1}{2})} a_\ell(\operatorname{Re}(\alpha')) \right| = \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} \frac{\Gamma(-\alpha' + \ell + \frac{1}{2})}{\Gamma(-\operatorname{Re}(\alpha') + \ell + \frac{1}{2})} a_\ell(\operatorname{Re}(\alpha)) \right| \\ &\leq \frac{\Gamma(\operatorname{Re}(\alpha') + \frac{1}{2})}{|\Gamma(\alpha' + \frac{1}{2})|} \left| \frac{\cos(\pi\alpha)}{\cos(\pi \operatorname{Re}(\alpha))} a_\ell(\operatorname{Re}(\alpha)) \right| \end{aligned}$$

which holds when $\ell \geq 0$ and $|\operatorname{Re}(\alpha')| < \frac{1}{2}$ (or, equivalently, $|\operatorname{Re}(\alpha)| < \frac{1}{2}$). We then use the following observations:

- (i) If $-1 < \operatorname{Re}(e^{-2\xi}) \leq 0$, then $|1 + e^{2\xi}| \leq |1 - e^{4\xi}|$.
- (ii) If $\operatorname{Re}(e^{-2\xi}) > 0$, then $|1 + e^{2\xi}| \leq |1 - e^{4\xi}| |\csc(2 \operatorname{Im}(\xi))|$.
- (iii) If $1 \leq \operatorname{Re}(e^{2\xi})$, then $1 \leq |1 + e^{-2\xi}|$.
- (iv) If $0 \leq \operatorname{Re}(e^{2\xi}) < 1$, then $|1 - e^{-2\xi}| \leq |1 - e^{-4\xi}|$.

(v) If $-1 < \operatorname{Re}(e^{2\xi}) < 0$, then $|1 - e^{-2\xi}| \leq |1 - e^{-4\xi}| |\operatorname{csc}(2 \operatorname{Im}(\xi))|$.

This completes the proof of Theorem 2.4.

To prove Theorem 2.5, assume that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. Substituting into (6.1) and using (6.4) with the values $u = t$, $v_1 = s$, $v_2 = 1 - s$, $A = \frac{1}{2} - \alpha'$, and $B = \frac{1}{2} - \beta'$, while taking into account (6.3), we obtain (2.24), with $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) = \widehat{R}_N^{(P_2)}(\nu, \xi, \alpha, \beta)$ (cf. (6.23)). Thus, the estimates for $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta)$ directly follow from those for $\widehat{R}_N^{(P_2)}(\nu, \xi, \alpha, \beta)$, the latter of which are also valid for $\xi > 0$.

Now, assume that $\xi > 0$, and that ν , α , and β are real, with $|\alpha|, |\beta| < \frac{1}{2}$ and $2\nu + \alpha + \beta + 1 > \alpha' + \beta'$. Since $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) = \widehat{R}_N^{(P_2)}(\nu, \xi, \alpha, \beta)$, substituting (6.9) into (6.23), and observing that $a_\ell(\alpha') = a_\ell(\alpha)$ and $a_\ell(\beta') = a_\ell(\beta)$ for all ℓ , we obtain

$$\begin{aligned} 0 &< \widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) \\ &< 2 \times (-1)^N \sum_{\ell=0}^N a_\ell(\alpha) a_{N-\ell}(\beta) \left(-\frac{e^{-\xi}}{\sinh \xi} \right)^\ell \left(\frac{e^{-\xi}}{\cosh \xi} \right)^{N-\ell} \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + N + 2)} \\ &= 2 \times (-1)^N g_N(-\xi, \alpha, \beta) \frac{\Gamma(2\nu + \alpha + \beta + 2)}{\Gamma(2\nu + \alpha + \beta + N + 2)}. \end{aligned}$$

This concludes the proof of Theorem 2.5.

We proceed by showing that the factorial expansions (1.17) and (1.18) terminate, providing an exact representation of the corresponding functions when both 2α and 2β are odd integers. Let L denote the positive integer defined by $\max(2|\alpha|, 2|\beta|) = 2L - 1$. Then, by definition (6.4), we have

$$\widehat{R}_{2L}^\pm(t, s, 1-s, \xi, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') = 0.$$

Thus, from (6.22) and (6.23), it follows that $\widehat{R}_{2L}^{(P_1)}(\nu, \xi, \alpha, \beta) = 0$ and $\widehat{R}_{2L}^{(P_2)}(\nu, \xi, \alpha, \beta) = 0$. Consequently, by Theorems 2.4 and 2.5, the expansions (1.17) and (1.18) terminate, yielding exact representations of the corresponding functions. These parameter conditions may be relaxed by invoking analytic continuation.

Finally, observe that when

$$\left| \frac{e^{\pm\xi}}{2 \sinh \xi} \right|, \quad \left| \frac{e^{\pm\xi}}{2 \cosh \xi} \right| < 1 \quad \iff \quad |\operatorname{Re}(e^{\pm 2\xi})| < \frac{1}{2},$$

the convergence of the binomial expansion implies $\lim_{N \rightarrow +\infty} \widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B) = 0$. Therefore, by (6.22) and (6.23), it follows that $\lim_{N \rightarrow +\infty} \widehat{R}_N^{(P_1)}(\nu, \xi, \alpha, \beta) = 0$ and $\lim_{M \rightarrow +\infty} \widehat{R}_M^{(P_2)}(\nu, \xi, \alpha, \beta) = 0$, provided that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. As a result, under these conditions, the infinite series in (1.17) converges to its left-hand side. Since $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) = \widehat{R}_N^{(P_2)}(\nu, \xi, \alpha, \beta)$, the series in (1.18) converges to its left-hand side under the same assumptions.

7. PROOF OF THEOREMS 2.6 AND 2.7

In this section, we establish the error bounds stated in Theorems 2.6 and 2.7. To this end, we require alternative estimates for the remainder terms $\widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B)$ defined in Lemma 6.1.

Lemma 7.1. *Let N be a non-negative integer, and let A and B be complex numbers. Let u , v_1 , and v_2 be real numbers such that $0 < u, v_1, v_2 < 1$. For $\xi \in \overline{\mathcal{D}} \setminus \mathbb{R}^+$, define the remainder $\widehat{R}_N^+(u, v_1, v_2, \xi, A, B)$, and for $\xi \in \overline{\mathcal{D}}$, define the remainder $\widehat{R}_N^-(u, v_1, v_2, \xi, A, B)$ by (6.4). Then*

the following bounds hold:

$$(7.1) \quad \left| \widehat{R}_N^\pm(u, v_1, v_2, \xi, A, B) \right| \leq \frac{\Gamma(\operatorname{Re}(A))\Gamma(\operatorname{Re}(B))}{|\Gamma(A)\Gamma(B)|} \frac{|\Gamma(A+B+N)|}{\Gamma(\operatorname{Re}(A+B)+N)} \\ \times \sum_{\ell=0}^N \frac{(\operatorname{Re}(A))_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell!(N-\ell)!} \left| \left(v_1 \frac{e^{\pm\xi}}{2 \sinh \xi} \right)^\ell \left(v_2 \frac{e^{\pm\xi}}{2 \cosh \xi} \right)^{N-\ell} \right| u^N \\ \times \max_{0 \leq q \leq 1} \mathfrak{g} \left(\pm q u v_1 \frac{e^{\pm\xi}}{2 \sinh \xi} + (1-q) u v_2 \frac{e^{\pm\xi}}{2 \cosh \xi} \right),$$

under the conditions $\operatorname{Re}(A), \operatorname{Re}(B) > 0$ and $\operatorname{Re}(A+B) < 1$. Here, \mathfrak{g} is defined by (2.25).

Proof. The proof proceeds similarly to the argument given in [17, Lemma 3] for the special case where A and B are real. We omit the details. \square

We now proceed with the proof of Theorem 2.6. Assume that $\operatorname{Re}(\alpha'), \operatorname{Re}(\beta') < \frac{1}{2}$ and that $0 < \operatorname{Re}(\alpha' + \beta') < \operatorname{Re}(2\nu + \alpha + \beta + 1)$. By restricting ξ to $\mathcal{D} \setminus \mathbb{R}^+$, we substitute the bound (7.1) into the right-hand side of (6.24). Noting that $a_\ell(\operatorname{Re}(\alpha')) = a_\ell(\operatorname{Re}(\alpha))$ and $a_\ell(\operatorname{Re}(\beta')) = a_\ell(\operatorname{Re}(\beta))$ for all ℓ , we obtain the estimate

$$\left| \widehat{R}_N^{(P_1)}(\nu, \xi, \alpha, \beta) \right| \leq \frac{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta') + 1)}{|\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)|} \frac{|\Gamma(-\alpha' - \beta' + N + 1)|}{\Gamma(\operatorname{Re}(-\alpha' - \beta') + N + 1)} \\ \times \sum_{\ell=0}^N \left| \frac{\cos(\pi\alpha) \cos(\pi\beta)}{\cos(\pi \operatorname{Re}(\alpha)) \cos(\pi \operatorname{Re}(\beta))} a_\ell(\operatorname{Re}(\alpha)) a_{N-\ell}(\operatorname{Re}(\beta)) \left(\frac{e^\xi}{\sinh \xi} \right)^\ell \left(\frac{e^\xi}{\cosh \xi} \right)^{N-\ell} \right| \\ \times \frac{|\Gamma(2\nu + \alpha + \beta + 2)|}{\Gamma(\operatorname{Re}(2\nu + \alpha + \beta) + N + 2)} \times \max_{0 \leq q, t, s \leq 1} \mathfrak{g} \left(qst \frac{e^\xi}{2 \sinh \xi} + (1-q)(1-s)t \frac{e^\xi}{2 \cosh \xi} \right).$$

Using [17, Eq. (16.6)], we find

$$\max_{0 \leq q, t, s \leq 1} \mathfrak{g} \left(qst \frac{e^\xi}{2 \sinh \xi} + (1-q)(1-s)t \frac{e^\xi}{2 \cosh \xi} \right) = \max_{0 \leq q, s \leq 1} \mathfrak{g} \left(qs \frac{e^\xi}{2 \sinh \xi} + (1-q)(1-s) \frac{e^\xi}{2 \cosh \xi} \right).$$

By [17, Eqs. (16.7) and (16.17)], this further simplifies to

$$\max_{0 \leq u \leq \frac{\pi}{2}} \mathfrak{g} \left(\frac{\sin^2 u}{1 + \sin(2u)} \frac{e^\xi}{2 \sinh \xi} + \frac{\cos^2 u}{1 + \sin(2u)} \frac{e^\xi}{2 \cosh \xi} \right),$$

which proves the bound (2.26). The case $\alpha = \beta = 0$ follows from a continuity argument. Note that the condition $\operatorname{Re}(\alpha'), \operatorname{Re}(\beta') < \frac{1}{2}$ is equivalent to $|\operatorname{Re}(\alpha)|, |\operatorname{Re}(\beta)| < \frac{1}{2}$. The bound (2.27) can be established in a completely analogous manner by combining (7.1) and (6.25). This completes the proof of Theorem 2.6.

For Theorem 2.7, note that $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) = \widehat{R}_N^{(P_2)}(\nu, \xi, \alpha, \beta)$. Consequently, the estimate (2.28) follows directly from (2.27), which is also valid for $\xi > 0$.

8. PROOF OF THEOREM 2.8

In this section, we establish the error bounds presented in Theorem 2.8. Suppose that $0 < \zeta < \frac{\pi}{2}$. Assuming that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$, and noting that $\widehat{R}_N^{(Q)}(\nu, \xi, \alpha, \beta) = \widehat{R}_N^{(P_2)}(\nu, \xi, \alpha, \beta)$, we substitute (2.24) into the right-hand side of (3.6). This results in (2.29) with

$$\widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) = \frac{1}{2} e^{-((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i} \lim_{\varepsilon \rightarrow 0^+} \widehat{R}_N^{(Q)}(\nu, \varepsilon + i\zeta, \alpha, \beta) \\ + \frac{1}{2} e^{((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i} \lim_{\varepsilon \rightarrow 0^+} \widehat{R}_N^{(Q)}(\nu, \varepsilon - i\zeta, \alpha, \beta).$$

Applying the representation (6.23) and passing to the limit, we obtain

$$\begin{aligned}
(8.1) \quad \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) &= \frac{\Gamma(2\nu + \alpha + \beta + 2)}{2\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} e^{-((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i} \\
&\times \int_0^1 \int_0^1 t^{\alpha' + \beta'} (1-t)^{2\nu + \alpha + \beta - \alpha' - \beta'} s^{\alpha' - 1/2} (1-s)^{\beta' - 1/2} \widehat{R}_N^-(t, s, 1-s, i\zeta, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') ds dt \\
&+ \frac{\Gamma(2\nu + \alpha + \beta + 2)}{2\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} e^{((2\nu + \alpha + \beta + 1)\zeta - (\alpha + \frac{1}{2})\frac{\pi}{2})i} \\
&\times \int_0^1 \int_0^1 t^{\alpha' + \beta'} (1-t)^{2\nu + \alpha + \beta - \alpha' - \beta'} s^{\alpha' - 1/2} (1-s)^{\beta' - 1/2} \widehat{R}_N^-(t, s, 1-s, -i\zeta, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') ds dt.
\end{aligned}$$

A straightforward estimation yields the bound

$$\begin{aligned}
(8.2) \quad \left| \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) \right| &\leq \left| \frac{\Gamma(2\nu + \alpha + \beta + 2)}{2\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} \right| e^{\operatorname{Im}(\zeta_{\nu, N, 0}^{(3)})} \\
&\times \int_0^1 \int_0^1 t^{\operatorname{Re}(\alpha' + \beta')} (1-t)^{\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta')} s^{\operatorname{Re}(\alpha') - 1/2} (1-s)^{\operatorname{Re}(\beta') - 1/2} \\
&\times \left| \widehat{R}_N^-(t, s, 1-s, i\zeta, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') \right| ds dt \\
&+ \left| \frac{\Gamma(2\nu + \alpha + \beta + 2)}{2\Gamma(2\nu + \alpha + \beta - \alpha' - \beta' + 1)\Gamma(\alpha' + \frac{1}{2})\Gamma(\beta' + \frac{1}{2})} \right| e^{-\operatorname{Im}(\zeta_{\nu, N, 0}^{(3)})} \\
&\times \int_0^1 \int_0^1 t^{\operatorname{Re}(\alpha' + \beta')} (1-t)^{\operatorname{Re}(2\nu + \alpha + \beta - \alpha' - \beta')} s^{\operatorname{Re}(\alpha') - 1/2} (1-s)^{\operatorname{Re}(\beta') - 1/2} \\
&\times \left| \widehat{R}_N^-(t, s, 1-s, -i\zeta, \frac{1}{2} - \alpha', \frac{1}{2} - \beta') \right| ds dt.
\end{aligned}$$

From Lemma 6.1, we can deduce that

$$\begin{aligned}
\left| \widehat{R}_N^-(u, v_1, v_2, \pm i\zeta, A, B) \right| &\leq 2 \left| \frac{\sin(\pi A) \sin(\pi B)}{\sin(\pi \operatorname{Re}(A)) \sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(A))_N}{N!} \left(\frac{v_1}{2 \sin \zeta} \right)^N \right| u^N \\
&+ \sum_{\ell=0}^{N-1} \left| \frac{\sin(\pi B)}{\sin(\pi \operatorname{Re}(B))} \frac{(A)_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left(\frac{v_1}{2 \sin \zeta} \right)^\ell \left(\frac{v_2}{2 \cos \zeta} \right)^{N-\ell} \right| u^N \\
&\times \begin{cases} 2 \cos \zeta, & \text{if } 0 < \zeta < \frac{\pi}{4}, \\ \csc \zeta, & \text{if } \frac{\pi}{4} \leq \zeta < \frac{\pi}{2}, \end{cases}
\end{aligned}$$

provided that $-N < \operatorname{Re}(A) < 1$ and $0 < \operatorname{Re}(B) < 1$, and

$$\begin{aligned} \left| \widehat{R}_N^-(u, v_1, v_2, \pm i\zeta, A, B) \right| &\leq 2 \left| \frac{\sin(\pi A) \sin(\pi B)}{\sin(\pi \operatorname{Re}(A)) \sin(\pi \operatorname{Re}(B))} \frac{(\operatorname{Re}(B))_N}{N!} \left(\frac{v_2}{2 \cos \zeta} \right)^N \right| u^N \\ &\quad + \sum_{\ell=1}^N \left| \frac{\sin(\pi A)}{\sin(\pi \operatorname{Re}(A))} \frac{(\operatorname{Re}(A))_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left(\frac{v_1}{2 \sin \zeta} \right)^\ell \left(\frac{v_2}{2 \cos \zeta} \right)^{N-\ell} \right| u^N \\ &\quad \times \begin{cases} \sec \zeta, & \text{if } 0 < \zeta < \frac{\pi}{4}, \\ 2 \sin \zeta, & \text{if } \frac{\pi}{4} \leq \zeta < \frac{\pi}{2}, \end{cases} \end{aligned}$$

provided that $0 < \operatorname{Re}(A) < 1$ and $-N < \operatorname{Re}(B) < 1$. If A or B is an integer, then the limiting values must be taken in these bounds. Similarly, from Lemma 7.1 and [17, Eq. (16.19)], we obtain

$$\begin{aligned} \left| \widehat{R}_N^-(u, v_1, v_2, \pm i\zeta, A, B) \right| &\leq 2 \frac{\Gamma(\operatorname{Re}(A)) \Gamma(\operatorname{Re}(B))}{|\Gamma(A) \Gamma(B)|} \frac{|\Gamma(A+B+N)|}{\Gamma(\operatorname{Re}(A+B)+N)} \\ &\quad \times \sum_{\ell=0}^N \frac{(\operatorname{Re}(A))_\ell (\operatorname{Re}(B))_{N-\ell}}{\ell! (N-\ell)!} \left| \left(\frac{v_1}{2 \sin \zeta} \right)^\ell \left(\frac{v_2}{2 \cos \zeta} \right)^{N-\ell} \right| u^N, \end{aligned}$$

under the conditions $\operatorname{Re}(A), \operatorname{Re}(B) > 0$ and $\operatorname{Re}(A+B) < 1$. Substituting these estimates into (8.2) and observing that $\operatorname{Im} \left(\zeta_{\nu, N, 0}^{(3)} \right) = \operatorname{Im} \left(\zeta_{\nu, N, \ell}^{(3)} \right)$, $a_\ell(\alpha') = a_\ell(\alpha)$, $a_\ell(\operatorname{Re}(\alpha')) = a_\ell(\operatorname{Re}(\alpha))$, $a_\ell(\beta') = a_\ell(\beta)$, and $a_\ell(\operatorname{Re}(\beta')) = a_\ell(\operatorname{Re}(\beta))$ for all $0 \leq \ell \leq N$, we obtain the bounds (2.32), (2.33), and (2.35), respectively. The case $\alpha = \beta = 0$ in (2.35) follows by a continuity argument. Note that, for instance, in deriving (2.32), the conditions $-\frac{1}{2} < \operatorname{Re}(\alpha') < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta')| < \frac{1}{2}$ are required. However, these are equivalent to $-\frac{1}{2} < \operatorname{Re}(\alpha) < N + \frac{1}{2}$ and $|\operatorname{Re}(\beta)| < \frac{1}{2}$. A similar observation holds for the other bounds. The estimate (2.34) can be derived from (2.32) by applying the inequality (6.26).

The bounds for the remainder terms $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ and $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ in the truncated factorial series (2.30) and (2.31) can be derived in a similar fashion by combining (2.24) with (3.7) and (3.8), respectively. The details are left for the reader to verify.

The integral representation of $\widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta)$, along with the analogous representations for $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ and $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$, as well as the discussion at the end of Section 6, demonstrate that the factorial expansions (1.20), (1.21), and (1.22) terminate, providing an exact representation of the corresponding functions when both 2α and 2β are odd integers.

Finally, note that when

$$\frac{1}{|2 \sin \zeta|}, \quad \frac{1}{|2 \cos \zeta|} < 1 \quad \iff \quad \frac{\pi}{6} < \zeta < \frac{\pi}{3},$$

the convergence of the binomial expansion implies $\lim_{N \rightarrow +\infty} \widehat{R}_N^-(u, v_1, v_2, \pm i\zeta, A, B) = 0$. Consequently, from (8.1), we have $\lim_{N \rightarrow +\infty} \widehat{R}_N^{(P)}(\nu, \zeta, \alpha, \beta) = 0$, provided that $\operatorname{Re}(2\nu + \alpha + \beta + 1) > \operatorname{Re}(\alpha' + \beta')$. Thus, under these conditions, the infinite series in (1.20) converges to its left-hand side. The convergence of the series (1.21) and (1.22) follows in a similar manner, based on the analogous integral representations of the remainder terms $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$ and $\widehat{R}_N^{(Q)}(\nu, \zeta, \alpha, \beta)$.

REFERENCES

- [1] T. Aoki, T. Takahashi, M. Tanda, Exact WKB analysis and Jacobi polynomials, *RIMS Kôkyûroku Bessatsu* **B75** (2019), Article 11, pp. 123–137.
- [2] X.-X. Bai, Y.-Q. Zhao, A uniform asymptotic expansion for Jacobi polynomials via uniform treatment of Darboux's method, *J. Approx. Theory* **148** (2007), no. 1, pp. 1–11.

- [3] P. Baratella, L. Gatteschi, Remarks on asymptotics for Jacobi polynomials, *Calcolo* **28** (1991), pp. 129–137.
- [4] L.-C. Chen, M. E. H. Ismail, On asymptotics of Jacobi polynomials, *SIAM J. Math. Anal.* **22** (1991), no. 5, pp. 1442–1449.
- [5] H. S. Cohl, R. S. Costas-Santos, Multi-integral representations for Jacobi functions of the first and second kind, *Arab J. Basic Appl. Sci.* **30** (2023), no. 1, pp. 583–592.
- [6] G. Darboux, Mémoire sur l’approximation des fonctions de très-grandes nombres et sur une classe étendue de développements en série, *J. Math. Pures Appl.* **4** (1878), no. 3, pp. 5–57, 377–416.
- [7] E. Delabaere, J.-M. Rasoamanana, Somme effective d’une somme de Borel par séries de factorielles, *Ann. Inst. Fourier (Grenoble)* **57** (2007), no. 2, pp. 421–456.
- [8] T. M. Dunster, Asymptotic approximations for the Jacobi and ultraspherical polynomials, and related functions, *Methods Appl. Anal.* **6** (1999), no. 3, pp. 281–315.
- [9] L. Durand, Product formulas and Nicholson-type integrals for Jacobi functions. I. Summary of results, *SIAM J. Math. Anal.* **9** (1978), no. 1, pp. 76–86.
- [10] L. Durand, Asymptotic Bessel-function expansions for Legendre and Jacobi functions, *J. Math. Phys.* **60** (2019), no. 1, Article 013501, 14 pp.
- [11] D. Elliott, Uniform asymptotic expansions of the Jacobi polynomials and an associated function, *Math. Comp.* **25** (1971), no. 114, pp. 309–315.
- [12] A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, *Higher Transcendental Functions*, Vol. II, McGraw-Hill Book Company, Inc., New York, 1953.
- [13] A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, *Tables of Integral Transforms*, Vol. I, McGraw-Hill Book Company, Inc., New York, 1954.
- [14] C. L. Frenzen, R. Wong, A uniform asymptotic expansion of the Jacobi polynomials with error bounds, *Can. J. Math.* **37** (1985), no 5, pp. 979–1007.
- [15] L. Gatteschi, New error bounds for asymptotic approximations of Jacobi polynomials and their zeros, *Rend. Mat. Appl.* **14** (1994), no. 7, pp. 177–198.
- [16] A. Gil, J. Segura, N. M. Temme, Asymptotic expansions of Jacobi polynomials and of the nodes and weights of Gauss-Jacobi quadrature for large degree and parameters in terms of elementary functions, *J. Math. Anal. Appl.* **494** (2021), no. 2, Article 124642, 14 pp.
- [17] E. Hahn, Asymptotik bei Jacobi-Polynomen und Jacobi-Funktionen, *Math. Z.* **171** (1980), no. 3, pp. 201–226.
- [18] D. S. Jones, Asymptotics of the hypergeometric function, *Math. Methods Appl. Sci.* **24** (2001), no. 6, pp. 369–389.
- [19] A. B. J. Kuijlaars, A. Martínez-Finkelshtein, Strong asymptotics for Jacobi polynomials with varying non-standard parameters, *J. D’Anal. Math.* **94** (2004), no. 1, pp. 195–234.
- [20] P. McCabe, J. N. L. Connor, Nearside-farside analysis of differential cross sections: Diffraction and rainbow scattering in atom-atom and atom-molecule rotationally inelastic sudden collisions, *J. Chem. Phys.* **104** (1996), no. 6, pp. 2297–2311.
- [21] G. Nemes, Error bounds for the large-argument asymptotic expansions of the Hankel and Bessel functions, *Acta Appl. Math.* **150** (2017), no. 1, pp. 141–177.
- [22] G. Nemes, A. B. Olde Daalhuis, Large-parameter asymptotic expansions for the Legendre and allied functions, *SIAM J. Math. Anal.* **52** (2020), no. 1, pp. 437–470.
- [23] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.2 of 2024-09-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [24] A. B. Olde Daalhuis, Inverse factorial-series solutions of difference equations, *Proc. Edinb. Math. Soc.* **47** (2004), no. 2, pp. 421–448.
- [25] G. Szegő, Asymptotische Entwicklungen der Jacobischen Polynome, *Schr. der Konig. Gelehr. Gesell. Naturwiss. Kl.* **10** (1933), pp. 33–112.
- [26] G. Szegő, *Orthogonal Polynomials*, American Mathematical Society, New York, 1939.
- [27] O. Szehr, R. Zarouf, On the asymptotic behavior of Jacobi polynomials with first varying parameter, *J. Approx. Theory* **277** (2022), Article 105702, 31 pp.
- [28] G. N. Watson, Asymptotic expansions of hypergeometric functions, *Trans. Cambridge Philos. Soc.* **22** (1918), pp. 277–308.
- [29] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge University Press, Cambridge, England; The Macmillan Company, New York, 1944.

- [30] R. Wong, Q.-Q. Wang, On the asymptotics of the Jacobi function and its zeros, *SIAM J. Math. Anal.* **23** (1992), no. 6, pp. 1637–1649.
- [31] R. Wong, J.-M. Zhang, A uniform asymptotic expansion for the Jacobi polynomials with explicit remainder, *Appl. Anal.* **61** (1996), no. 1–2, pp. 17–29.
- [32] R. Wong, Y.-Q. Zhao, Estimates for the error term in a uniform asymptotic expansion of the Jacobi polynomials. *Anal. Appl. (Singap.)* **1** (2003), no. 2, pp. 213–241.
- [33] R. Wong, Y.-Q. Zhao, Uniform asymptotic expansion of the Jacobi polynomials in a complex domain, *Proc. R. Soc. Lond. A* **460** (2004), no. 2049, pp. 2569–2586.
- [34] X. F. Zhu, X. C. Li, Asymptotic approximations of Jacobi functions and their zeros with error bounds, *Acta Math. Sinica* **22** (2006), no. 3, pp. 729–740.

SCHOOL OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, XIDAZHI STREET, HARBIN 150001,
HEILONGJIANG, PEOPLE'S REPUBLIC OF CHINA

Email address: g.nemes@hit.edu.cn