

Local well-posedness for the Schrödinger-KdV system in $H^{s_1} \times H^{s_2}$, II

Yingzhe Ban¹, Jie Chen², Ying Zhang^{3,1}

¹*The Graduate School of China Academy of Engineering Physics, Beijing, 100088, P.R. China*

²*School of Sciences, Jimei University, Xiamen 361021, P.R. China*

³*Academy of Mathematics and Systems Science, CAS, Beijing 100190, P.R. China*

Abstract

In this paper, we continue the study of the local well-posedness theory for the Schrödinger-KdV system in the Sobolev space $H^{s_1} \times H^{s_2}$. We show the local well-posedness in $H^{-3/16} \times H^{-3/4}$ for $\beta = 0$. Combining our work [2], we also have the local well-posedness for $\max\{-3/4, s_1 - 3\} \leq s_2 \leq \min\{4s_1, s_1 + 2\}$. The result is sharp by using the contraction mapping argument.

1 Introduction

We study the Cauchy problem for Schrödinger-KdV system

$$\begin{cases} i\partial_t u + \partial_{xx} u = \alpha u v + \beta |u|^2 u, \\ \partial_t v + \partial_{xxx} v = \partial_x (\gamma |u|^2 - v^2/2), \\ (u, v)|_{t=0} = (u_0, v_0) \in H^{s_1} \times H^{s_2}. \end{cases} \quad (\text{S-KdV})$$

In this paper we concern the local well-posedness theory for (S-KdV) with $\beta = 0$. The constants $\alpha, \gamma \neq 0$ in the model is not essential. We would assume $\alpha = \gamma = 1$ for brevity.

This model appears in the study of resonant interaction between solitary wave of Langmuir and solitary wave of ion-acoustic. See Appert-Vaclavik [1]. It also appears in plasma physics and a diatomic lattice system. See [6, 4] and reference therein.

We recall some early results for this model. Tsutsumi showed the well-posedness in $H^{s+1/2} \times H^s$, $s \in \mathbb{N}$ in [9]. In [5], Guo-Miao showed the well-posedness in $H^s \times H^s$, $s \in \mathbb{N}$. Guo-Wang [6] obtained the well-posedness in $H^{s_1} \times H^{-3/4}$, $s_1 > -1/16$ and Wang-Cui showed the well-posedness in $H^{s_1} \times H^{-3/4}$, $s_1 > -3/16$ in [11]. See also [3, 4, 2] for other results.

In this paper, we obtain the following result.

Theorem 1.1. (**S-KdV**) is local well-posed in $H^{-3/16} \times H^{-3/4}$.

In fact combining the results in our work [2] we have

Corollary 1.2. (**S-KdV**) is local well-posed in $H^{s_1} \times H^{s_2}$ for

$$\max\{-3/4, s_1 - 3\} \leq s_2 \leq \min\{4s_1, s_1 + 2\}.$$

Recall the proof of Theorem 1.1 in [2] one has

Theorem 1.3. The data-to-solution mapping of (**S-KdV**) can not be C^2 except $\max\{-3/4, s_1 - 3\} \leq s_2 \leq \min\{4s_1, s_1 + 2\}$.

Thus it is reasonable to say than our local well-posedness result is sharp by using contraction mapping argument.

To show the local well-posedness in $H^{-3/16} \times H^{-3/4}$, there are two difficulties which come from the terms $\partial_x(|u|^2)$ and $\partial_x(v^2)$. In some sense, $H^{-3/16} \times H^{-3/4}$ is the “double critical” space for this model. We need some techniques that are different from before.

The paper is organized as follows. In Section 2, we rewrite (**S-KdV**) into a new form and construct the workspace. Then we show linear estimates, bilinear estimates related the Schrödinger equation, and trilinear estimate related the KdV equation. In Section 3, we estimate $F[u_0]$ (see the definition (2.1)). In Section 4, we give some refinement for the bilinear estimates related to the KdV equation and finally conclude the proof.

Notations. . For $a, b \in \mathbb{R}^+$, $a \lesssim b$ means that there exists $C > 0$ such that $a \leq Cb$ and $a \sim b$ means $a \lesssim b \lesssim a$ where C is a constant. We use $\langle \xi \rangle$ to denote $(1 + \xi^2)^{1/2}$. Let φ be an even, smooth function and $\varphi|_{[-1,1]} = 0$, $\varphi|_{[-2,2]^c} = 0$. $\varphi_N = \varphi(\cdot/N)$, $\psi_N = \varphi_N - \varphi_{N/2}$. We use inhomogeneous Littlewood-Paley projection: $P_N := \mathcal{F}^{-1}\psi_N \mathcal{F}$, $N \geq 2$; $P_1 = \mathcal{F}^{-1}\varphi \mathcal{F}$; $P_{>N} = \sum_{M>N} P_M$, etc.. We always use N, L to denote a dyadic number larger than 1.

Let $S_\lambda(t) = e^{i\lambda t \partial_{xx}}$, $K(t) = e^{-t \partial_{xxx}}$,

$$\mathcal{A}_\lambda(f) := i\lambda \int_0^t S_\lambda(t-s)f(s) ds, \quad \mathcal{B}(f) := \int_0^t K(t-s)\partial_x f(s) ds.$$

We always assume $0 < \lambda \ll 1$ in this paper. We define the modulation decomposition $Q_L^{S_\lambda}$, Q_L^K by $\mathcal{F}_{\tau,\xi}^{-1}\psi_L(\tau + \lambda\xi^2)\mathcal{F}_{t,x}$, $\mathcal{F}_{\tau,\xi}^{-1}\psi_L(\tau - \xi^3)\mathcal{F}_{t,x}$ respectively. Similarly, we define $Q_{>L}^{S_\lambda}$, $Q_{>L}^K$, $Q_1^{S_\lambda}$, and Q_1^K .

2 Function spaces and multilinear estimates

To show the well-posedness of (**S-KdV**) with $s_2 = -3/4$, we use the argument in [6]. By rescaling, we consider the equation

$$\begin{cases} i\partial_t u + \lambda\partial_{xx}u = \lambda uv, \\ \partial_t v + \partial_{xxx}v = \partial_x(|u|^2 - v^2), \\ (u, v)|_{t=0} = (u_{0,\lambda}, v_{0,\lambda}) \in H^{-3/16} \times H^{-3/4} \end{cases}$$

where $0 < \lambda \ll 1$, $\|u_{0,\lambda}\|_{H^{-3/16}} \lesssim \lambda^{21/16}$, $\|v_{0,\lambda}\|_{H^{3/4}} \lesssim \lambda^{3/4}$. Recall the definition of $U^p - V^p$ spaces in [7]. We define

$$\|u\|_{X_\lambda} = \|N^{-3/16} P_N u\|_{l_N^2 U_{S_\lambda}^2}, \quad \|v\|_Y = \|P_1 v\|_{L_x^2 L_t^\infty} + \|N^{-3/4} P_N v\|_{l_{N \geq 2}^2 U_K^2}.$$

For the brevity, we still denote $(u_{0,\lambda}, v_{0,\lambda})$ by (u_0, v_0) . The corresponding integral equation is

$$\begin{cases} u(t) = S_\lambda(t)u_0 - \mathcal{A}_\lambda(uv)(t), \\ v(t) = K(t)v_0 + \mathcal{B}(|u|^2 - v^2)(t). \end{cases}$$

We replace the function u in second equation with the first one and obtain

$$\begin{cases} u(t) = S_\lambda(t)u_0 - i\mathcal{A}_\lambda(uv)(t), \\ v(t) = K(t)v_0 + \mathcal{B}(|S_\lambda(t)u_0 - \mathcal{A}_\lambda(uv)|^2 - v^2)(t). \end{cases}$$

Let $\eta, \tilde{\eta} \in C_0^\infty$ with $\eta|_{[-1,1]} = 1$, $\tilde{\eta}|_{\text{supp}(\eta)} = 1$. Define

$$F[u_0](t) := \eta(t)\mathcal{B}(|S_\lambda(t)u_0|^2)(t). \quad (2.1)$$

In fact we can not show $F[u_0] \in Y$. Let $w(t) = v(t) - F[u_0](t)$. We consider the equation

$$\begin{cases} u(t) = \eta S_\lambda(t)u_0 - \eta \mathcal{A}_\lambda(u(w + F[u_0]))(t), \\ w(t) = \eta K(t)v_0 - \eta \mathcal{B}((w + F[u_0])^2)(t) + \eta \mathcal{B}(T_\lambda(u, w, u_0))(t) \end{cases} \quad (2.2)$$

where $T_\lambda(u, w, u_0) = 2\text{Im}(\mathcal{A}_\lambda(u(w + F[u_0]))\overline{S_\lambda(t)u_0}) - |\mathcal{A}_\lambda(u(w + F[u_0]))|^2$. Define

$$\|v\|_Z := \|P_{\lesssim 1} v\|_{L_x^2 L_t^\infty} + \|N^{-3/4} P_N v\|_{l_{N \geq 2}^2 V_K^2} + \|N^{1/4} P_N v\|_{l_N^2 L_x^\infty L_t^2}. \quad (2.3)$$

Note that for v supported on $[0, 1] \times \mathbb{R}$, $\|v\|_Z \lesssim \|v\|_Y$ by Strichartz, maximal function, and local smoothing estimates. We would show $F[u_0] \in Z$ in Section 3.

Directly, one has the following estimates due to the maximal function estimate and the definition of U^2 .

Lemma 2.1. $\|\eta S_\lambda(t)u_0\|_{X_\lambda} \lesssim \|u_0\|_{H^{-3/16}}$, $\|\eta K(t)v_0\|_Y \lesssim \|v_0\|_{H^{-3/4}}$.

Then, we show the bilinear estimate related to the Schrödinger equation.

Lemma 2.2. $\|\eta \mathcal{A}_\lambda(uv)\|_{X_\lambda} \lesssim \lambda^{1/2} \|u\|_{X_\lambda} \|v\|_Z$.

Proof. Firstly, for low frequency part of v we have

$$\begin{aligned} \left| \int_{\mathbb{R}^2} P_N(uP_1v)\tilde{\eta}^2 \bar{w} \, dxdt \right| &\lesssim \sum_{N_1 \sim N} \|\tilde{\eta} P_{N_1} u\|_{L_x^\infty L_t^2} \|P_1 v\|_{L_x^2 L_t^\infty} \|\tilde{\eta} w\|_{L_{t,x}^2} \\ &\lesssim (\lambda N)^{-1/2} \sum_{N_1 \sim N} \|P_{N_1} u\|_{U_{S_\lambda}^2} \|v\|_Z \|w\|_{V_{S_\lambda}^2}. \end{aligned} \quad (2.4)$$

By duality one has

$$\begin{aligned}\|\eta\mathcal{A}_\lambda(uP_1v)\|_{X_\lambda} &\lesssim \lambda \sum_N N^{-3/16} (\lambda N)^{-1/2} \sum_{N_1 \sim N} \|P_{N_1}u\|_{U_{S_\lambda}^2} \|v\|_Z \\ &\lesssim \lambda^{1/2} \|u\|_{X_\lambda} \|v\|_Z.\end{aligned}$$

Then by triangle inequality,

$$\|\eta\mathcal{A}_\lambda(uP_{>1}v)\|_{X_\lambda} \leq \sum_N \sum_{N_1} \sum_{N_2 \geq 2} N^{-3/16} \|\eta P_N \mathcal{A}_\lambda(P_{N_1}uP_{N_2}v)\|_{U_{S_\lambda}^2}.$$

If $N_2^2 \approx \lambda N_1$, for $\tau_1 + \tau_2 = \tau$, $\xi_1 + \xi_2 = \xi$, $|\xi_1| \sim N_1$, $|\xi_2| \sim N_2$, $|\xi| \sim N$ we have

$$|\tau_1 + \lambda\xi_1^2| + |\tau_2 - \xi_2^3| + |\tau + \lambda\xi^2| \gtrsim N_2 \max\{N_2^2, \lambda N_1\}.$$

Let $L = cN_2 \max\{N_2^2, \lambda N_1\}$. Typically we need to estimate

$$\begin{aligned}I &= \int_{\mathbb{R}^2} Q_{>L}^{S_\lambda} P_{N_1} u P_{N_2} v P_N \bar{w} \, dx dt, \\ II &= \int_{\mathbb{R}^2} P_{N_1} u Q_{>L}^K P_{N_2} v P_N \bar{w} \, dx dt\end{aligned}$$

where u, v, w are supported on $[-1, 1] \times \mathbb{R}$. By the Strichartz estimate one has

$$\|u\|_{L_{t,x}^6} \lesssim \lambda^{-1/6} \|u\|_{U_{S_\lambda}^2} \lesssim \lambda^{-1/6} \|u\|_{V_{S_\lambda}^2}, \quad \|u\|_{L_{t,x}^2} \lesssim \|u\|_{L_t^\infty L_x^2} \lesssim \|u\|_{V_{S_\lambda}^2}.$$

Thus $\|u\|_{L_{t,x}^4} \leq \|u\|_{L_{t,x}^6}^{3/4} \|u\|_{L_{t,x}^2}^{1/4} \lesssim \lambda^{-1/8} \|u\|_{V_{S_\lambda}^2}$. Similarly,

$$\|P_{N_2}v\|_{L_{t,x}^4} \lesssim N_2^{-1/8} \|v\|_{V_K^2}, \quad \|P_Nw\|_{L_{t,x}^4} \lesssim \lambda^{-1/8} \|w\|_{V_K^2}.$$

By the Hölder inequality we have

$$\begin{aligned}|I| &\lesssim \|Q_{>L}^{S_\lambda} P_{N_1} u\|_{L_{t,x}^2} \|P_{N_2} v\|_{L_{t,x}^4} \|P_N w\|_{L_{t,x}^4} \\ &\lesssim L^{-1/2} \lambda^{-1/8} \|u\|_{V_{S_\lambda}^2} N_2^{-1/8} \|v\|_{V_K^2} \|w\|_{V_K^2} \\ &\lesssim \lambda^{-1/8} N_2^{-5/8} \max\{N_2^2, \lambda N_1\}^{-1/2} \|u\|_{U_{S_\lambda}^2} \|v\|_{V_K^2} \|w\|_{V_K^2}.\end{aligned}$$

Also,

$$|II| \lesssim \lambda^{-1/4} N_2^{-1/2} \max\{N_2^2, \lambda N_1\}^{-1/2} \|u\|_{U_{S_\lambda}^2} \|v\|_{V_K^2} \|w\|_{V_K^2}.$$

Thus by the duality one has

$$\begin{aligned}&\|\eta P_N \mathcal{A}_\lambda(P_{N_1}uP_{N_2}v)\|_{U_{S_\lambda}^2} \\ &\lesssim \lambda \lambda^{-1/4} N_2^{-1/2} \max\{N_2^2, \lambda N_1\}^{-1/2} \|P_{N_1}u\|_{U_{S_\lambda}^2} \|P_{N_2}v\|_{V_K^2} \quad (2.5) \\ &\lesssim \lambda^{3/4} N_1^{3/16} N_2^{1/4} \max\{N_2^2, \lambda N_1\}^{-1/2} \|u\|_{X_\lambda} \|v\|_Z.\end{aligned}$$

Let N_{\max}, N_{med} be the maximum, medium of N_1, N_2, N . Note that

$$\sum_{N_{\max} \sim N_{\text{med}}} N^{-3/16} \lambda^{3/4} N_1^{3/16} N_2^{1/4} \max\{N_2^2, \lambda N_1\}^{-1/2} \lesssim \lambda^{3/4} \log \lambda^{-1}.$$

Thus we obtain

$$\sum_{N_2 \geq 2, N_2^2 \sim \lambda N_1, N_1, N} N^{-3/16} \|\eta P_N \mathcal{A}_\lambda(P_{N_1} u P_{N_2} v)\|_{U_{S_\lambda}^2} \lesssim \frac{\log \lambda^{-1}}{\lambda^{-3/4}} \|u\|_{X_\lambda} \|v\|_Z.$$

If $N_2^2 \sim \lambda N_1 \sim \lambda N$, by Lemma 4.12, (4.8) in [2], we have

$$\left| \int_{\mathbb{R}^2} P_{N_1} u P_{N_2} v P_N \bar{w} \, dx dt \right| \lesssim N_2^{-1} \|u\|_{U_{S_\lambda}^2} \|v\|_{U_K^2} \|w\|_{U_{S_\lambda}^2}.$$

On the other hand, by Hölder's inequality and Strichartz estimates, one has

$$\begin{aligned} \left| \int_{\mathbb{R}^2} P_{N_1} u P_{N_2} v P_N \bar{w} \, dx dt \right| &\lesssim \|u\|_{L_{t,x}^3} \|P_{N_2} v\|_{L_{t,x}^3} \|w\|_{L_{t,x}^3} \\ &\lesssim \|u\|_{L_t^{12} L_x^3} \|P_{N_2} v\|_{L_t^{12} L_x^3} \|w\|_{L_t^{12} L_x^3} \\ &\lesssim \lambda^{-1/6} N_2^{-1/12} \|u\|_{U_{S_\lambda}^{12}} \|v\|_{U_K^{12}} \|w\|_{U_{S_\lambda}^{12}}. \end{aligned}$$

Then by the interpolation (cf. Proposition 2.20 in [7]), we have

$$\left| \int_{\mathbb{R}^2} P_{N_1} u P_{N_2} v P_N \bar{w} \, dx dt \right| \lesssim N_2^{-1} \lambda^{-1/6} (\log N_2)^3 \|u\|_{U_{S_\lambda}^2} \|v\|_{V_K^2} \|w\|_{V_{S_\lambda}^2}.$$

By the duality one has

$$\begin{aligned} \|\eta P_N \mathcal{A}_\lambda(P_{N_1} u P_{N_2} v)\|_{U_{S_\lambda}^2} &\lesssim \lambda N_2^{-1} \lambda^{-1/6} (\log N_2)^3 \|P_{N_1} u\|_{U_{S_\lambda}^2} \|P_{N_2} v\|_{V_K^2} \\ &\lesssim \lambda^{5/6} N_2^{-1/4} N_1^{3/16} (\log N_2)^3 \|u\|_{X_\lambda} \|v\|_Z. \end{aligned} \tag{2.6}$$

Note that

$$\sum_{N_2^2 \sim \lambda N_1 \sim \lambda N} N^{-3/16} N_2^{-1/4} N_1^{3/16} \lambda^{5/6} (\log N_2)^3 \lesssim \lambda^{5/6}.$$

Overall one has

$$\sum_N \sum_{N_1} \sum_{N_2 \geq 2} N^{-3/16} \|\eta P_N \mathcal{A}_\lambda(P_{N_1} u P_{N_2} v)\|_{U_{S_\lambda}^2} \lesssim \lambda^{1/2} \|u\|_{X_\lambda} \|v\|_Z.$$

We finish the proof of this lemma. \square

Next we show the trilinear estimate related to the KdV equation.

Lemma 2.3. $\|\eta \mathcal{B}(\mathcal{A}_\lambda(uv)\bar{w})\|_Y \lesssim \lambda^{-1/2} \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}$.

Proof. Without loss of the generality we would assume that u, v, w are supported on $[-1, 1] \times \mathbb{R}$. For the low frequency we have

$$\begin{aligned}
\|\eta P_1 \mathcal{B}(\mathcal{A}_\lambda(uv)\bar{w})\|_{L_x^2 L_t^\infty} &\lesssim \|P_1(\mathcal{A}(uv)\bar{w})\|_{L_x^2 L_t^1} \\
&\lesssim \sum_{N_1 \sim N_2} \|P_1(P_{N_1} \mathcal{A}_\lambda(uv) P_{N_2} \bar{w})\|_{L_x^2 L_t^1} \\
&\lesssim \sum_{N_1 \sim N_2} \|\tilde{\eta} P_{N_1} \mathcal{A}_\lambda(uv)\|_{L_{t,x}^2} \|P_{N_2} w\|_{L_x^\infty L_t^2} \\
&\lesssim \sum_{N_1 \sim N_2} \|\tilde{\eta} P_{N_1} \mathcal{A}_\lambda(uv)\|_{U_{S_\lambda}^2} (\lambda N_2)^{-1/2} \|P_{N_2} w\|_{U_{S_\lambda}^2} \\
&\lesssim \lambda^{-1/2} \sum_{N_1 \sim N_2} N_1^{3/16} N_2^{-5/16} \|\tilde{\eta} \mathcal{A}_\lambda(uv)\|_{X_\lambda} \|w\|_{X_\lambda} \\
&\lesssim \lambda^{-1/2} \|\tilde{\eta} \mathcal{A}_\lambda(uv)\|_{X_\lambda} \|w\|_{X_\lambda}.
\end{aligned}$$

By Lemma 2.2 we obtain

$$\|\eta P_1 \mathcal{B}(\mathcal{A}_\lambda(uv)\bar{w})\|_{L_x^2 L_t^\infty} \lesssim \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}. \quad (2.7)$$

Since $\|P_N Q_L^K v\|_{L_x^\infty L_t^2} \lesssim N^{-1} L^{1/2} \|P_N Q_L^K v\|_{L_{t,x}^2}$, by the interpolation one has

$$\|P_N Q_L^K v\|_{L_x^{2/(1-\theta)} L_t^2} \lesssim (N^{-1} L^{1/2})^\theta \|P_N Q_L^K v\|_{L_{t,x}^2}, \quad \forall 0 < \theta < 1.$$

Thus (v is supported on $[-1, 1] \times \mathbb{R}$)

$$\begin{aligned}
\|P_N v\|_{L_x^{2/(1-\theta)} L_t^2} &\lesssim \sum_L (N^{-1} L^{1/2})^\theta \|P_N Q_L^K v\|_{L_{t,x}^2} \\
&\lesssim \sum_L (N^{-1} L^{1/2})^\theta L^{-1/2} \|P_N v\|_{V_K^2} \lesssim N^{-\theta} \|P_N v\|_{V_K^2}.
\end{aligned}$$

If $N_2 \lesssim N$ (which implies $N_1 \lesssim N$), by the duality, Minkowski, and Bernstein inequalities one has

$$\begin{aligned}
&\sum_{N_1, N_2 \lesssim N} \|\eta P_N \mathcal{B}(\mathcal{A}_\lambda(P_{N_1} u P_1 v) P_{N_2} \bar{w})\|_{U_K^2} \\
&\lesssim \sum_{N_1, N_2 \lesssim N} N N^{-\theta} \|P_{N_1} \mathcal{A}_\lambda(u P_1 v) P_{N_2} w\|_{L_x^{2/(1+\theta)} L_t^2} \\
&\lesssim \sum_{N_1, N_2 \lesssim N} N^{1-\theta} \|\tilde{\eta} P_{N_1} \mathcal{A}_\lambda(u P_1 v)\|_{L_x^{2/\theta} L_t^2} \|\tilde{\eta} P_{N_2} w\|_{L_x^2 L_t^\infty} \\
&\lesssim N^{3/2-\theta} \sum_{N_1, N_2 \lesssim N} (\lambda N_1)^{(\theta-1)/2} \|\tilde{\eta} P_{N_1} \mathcal{A}_\lambda(u P_1 v)\|_{U_{S_\lambda}^2} \|P_{N_2} w\|_{U_{S_\lambda}^2}.
\end{aligned}$$

By (2.4) one has $\|\tilde{\eta} P_{N_1} \mathcal{A}_\lambda(u P_1 v)\|_{U_{S_\lambda}^2} \lesssim \lambda (\lambda N_1)^{-1/2} N_1^{3/16} \|u\|_{X_\lambda} \|v\|_Z$. Thus we can control the former term by

$$N^{3/2-\theta} \sum_{N_1, N_2 \lesssim N} (\lambda N_1)^{(\theta-1)/2} \lambda (\lambda N_1)^{-1/2} N_1^{3/16} \|u\|_{X_\lambda} \|v\|_Z N_2^{3/16} \|w\|_{X_\lambda},$$

By choosing $15/16 < \theta < 1$, then

$$\lambda \sum_N N^{-3/4} N^{3/2-\theta} \sum_{N_1, N_2 \lesssim N} (\lambda N_1)^{(\theta-1)/2} (\lambda N_1)^{-1/2} N_1^{3/16} N_2^{3/16} \lesssim \lambda^{\theta/2}.$$

We obtain the desired estimate since we always assume $0 < \lambda \ll 1$.

If $N_2 \gg N$ (which implies $N_1 \sim N_2$) similar to the former argument, we have $\|P_N v\|_{L_{t,x}^{30/7}} \lesssim N^{-2/15} \|v\|_{V_K^2}$. Then by the duality one has

$$\begin{aligned} & \sum_{N_1 \sim N_2 \gg N} \|\eta P_N \mathcal{B}(\mathcal{A}_\lambda(P_{N_1} u P_1 v) P_{N_2} \bar{w})\|_{U_K^2} \\ & \lesssim \sum_{N_1 \sim N_2 \gg N} N N^{-2/15} \|\mathcal{A}_\lambda(P_{N_1} u P_1 v) P_{N_2} w\|_{L_{t,x}^{30/23}} \\ & \lesssim \sum_{N_1 \sim N_3 \gg N} N^{13/15} \|\tilde{\eta} \mathcal{A}_\lambda(P_{N_1} u P_1 v)\|_{L_{t,x}^2} \|P_{N_2} w\|_{L_{t,x}^{2/10}}^{3/10} \|P_{N_2} w\|_{L_{t,x}^6}^{7/10} \\ & \lesssim \lambda^{53/60} \sum_{N_1 \sim N_2 \gg N} N^{13/15} \|P_{N_1} u\|_{L_x^\infty L_t^2} \|P_1 v\|_{L_x^2 L_t^\infty} \|P_{N_2} w\|_{U_{S_\lambda}^2} \\ & \lesssim \lambda^{53/60} N^{13/15} \sum_{N_1 \sim N_2 \gg N} N_2^{3/16} (\lambda N_1)^{-1/2} N_1^{3/16} \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda} \\ & \lesssim \lambda^{23/60} N^{89/120} \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}. \end{aligned}$$

Note that $89/120 < 3/4$. Thus we have

$$\begin{aligned} & \sum_{N \geq 2} N^{-3/4} \sum_{N_1 \sim N_2 \gg N} \|\eta P_N \mathcal{B}(\mathcal{A}_\lambda(P_{N_1} u P_1 v) P_{N_2} \bar{w})\|_{U_K^2} \\ & \lesssim \lambda^{23/60} \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}. \end{aligned} \tag{2.8}$$

By (2.5), for $M_2^2 \asymp \lambda M_1$ we have

$$\|\eta P_{N_1} \mathcal{A}_\lambda(P_{M_1} u P_{M_2} v)\|_{U_{S_\lambda}^2} \lesssim \frac{\lambda^{3/4} M_1^{3/16} M_2^{1/4}}{\max\{M_2^2, \lambda M_1\}^{1/2}} \|u\|_{X_\lambda} \|v\|_Z.$$

By (2.6), for $M_2^2 \sim \lambda M_1$ we have

$$\|\eta P_{N_1} \mathcal{A}_\lambda(P_{M_1} u P_{M_2} v)\|_{U_{S_\lambda}^2} \lesssim \lambda^{5/6} M_2^{-1/4} M_1^{3/16} (\log(M_2/\lambda))^3 \|u\|_{X_\lambda} \|v\|_Z.$$

Thus,

$$\begin{aligned} & \|\eta P_{N_1} (\mathcal{A}_\lambda(u P_{>1} v))\|_{U_{S_\lambda}^2} \\ & \lesssim \sum_{M_2^2 \asymp \lambda M_1} \frac{\lambda^{3/4} M_1^{3/16} M_2^{1/4}}{\max\{M_2^2, \lambda M_1\}^{1/2}} \|u\|_{X_\lambda} \|v\|_Z \\ & + \sum_{M_2^2 \sim \lambda M_1 \sim \lambda N_1} \lambda^{5/6} M_2^{-1/4} M_1^{3/16} (\log(M_2/\lambda))^3 \|u\|_{X_\lambda} \|v\|_Z \\ & \lesssim (\lambda^{9/16} + \lambda^{17/24} N_1^{1/16} (\log N_1)^3) \|u\|_{X_\lambda} \|v\|_Z. \end{aligned} \tag{2.9}$$

Let $\tilde{u} = \eta \mathcal{A}_\lambda(u P_{>1} v)$. Now we estimate $\|\eta P_N \mathcal{B}(P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{U_K^2}$.

If $N^2 \ll \lambda N_1$ by Lemma 4.12, (4.7) in [2] one has

$$\left| \int_{\mathbb{R}^2} \partial_x (P_{N_1} \tilde{u} P_{N_2} \bar{w}) P_N \bar{v} \, dx dt \right| \lesssim \lambda^{-1} N_1^{-1/2} \|\tilde{u}\|_{U_{S_\lambda}^2} \|w\|_{U_{S_\lambda}^2} \|v\|_{U_K^2}.$$

On the other hand by the Hölder inequality we also have

$$\begin{aligned} \left| \int_{\mathbb{R}^2} \partial_x (P_{N_1} \tilde{u} P_{N_2} \bar{w}) P_N \bar{v} \, dx dt \right| &\lesssim N \|P_N (P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{L_{t,x}^2} \|\tilde{v}\|_{L_{t,x}^2} \\ &\lesssim \lambda^{-1/2} N^{1/2} \|\tilde{u}\|_{U_{S_\lambda}^2} \|w\|_{U_{S_\lambda}^2} \|v\|_{U_K^p}, \quad p > 2. \end{aligned}$$

Then by the interpolation theorem in [7] one has

$$\left| \int_{\mathbb{R}^2} \partial_x (P_{N_1} \tilde{u} P_{N_2} \bar{w}) P_N \bar{v} \, dx dt \right| \lesssim \lambda^{-1} N_1^{-1/2} \log(N_1/\lambda) \|\tilde{u}\|_{U_{S_\lambda}^2} \|w\|_{U_{S_\lambda}^2} \|v\|_{V_K^2}.$$

Combining Lemma 2.2 we have

$$\begin{aligned} &\sum_{N \geq 2} N^{-3/4} \sum_{N_1 \sim N_2 \gg N^2/\lambda} \|\eta P_N \mathcal{B}(P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{U_K^2} \\ &\lesssim \sum_{N \geq 2} N^{-3/4} \sum_{N_1 \sim N_2 \gg N^2/\lambda} \lambda^{-1} N_1^{-1/2} \log(N_1/\lambda) \|P_{N_1} \tilde{u}\|_{U_{S_\lambda}^2} \|P_{N_2} w\|_{U_{S_\lambda}^2} \\ &\lesssim \sum_{N_1 \sim N_2 \gg 1/\lambda} \lambda^{-1} N_1^{-1/2} \log(N_1/\lambda) \|P_{N_1} \tilde{u}\|_{U_{S_\lambda}^2} \|P_{N_2} w\|_{U_{S_\lambda}^2} \\ &\lesssim \lambda^{-7/8} \|\tilde{u}\|_{X_\lambda} \|w\|_{X_\lambda} \lesssim \lambda^{-3/8} \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}. \end{aligned} \tag{2.10}$$

Similarly if $N^2 \gg \lambda N_1$ by Lemma 4.12, (4.6) in [2] and the interpolation one has

$$\|\eta P_N \mathcal{B}(P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{U_K^2} \lesssim \lambda^{-1/2} N^{-1} \log(N/\lambda) \|\tilde{u}\|_{U_{S_\lambda}^2} \|w\|_{U_{S_\lambda}^2}.$$

Combining Lemma 2.2 we have

$$\begin{aligned} &\sum_{N \geq 2} N^{-3/4} \sum_{N_1, N_2 \ll N^2/\lambda} \|\eta P_N \mathcal{B}(P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{U_K^2} \\ &\lesssim \sum_{N \geq 2} N^{-3/4} \sum_{N_1, N_2 \ll N^2/\lambda} \lambda^{-1/2} N^{-1} \log(N/\lambda) \|P_{N_1} \tilde{u}\|_{U_{S_\lambda}^2} \|P_{N_2} w\|_{U_{S_\lambda}^2} \\ &\lesssim \sum_{N \geq 2} N^{-3/4} \sum_{N_1, N_2 \ll N^2/\lambda} \lambda^{-1/2} N^{-1} \log(N/\lambda) (N_1 N_2)^{3/16} \|\tilde{u}\|_{X_\lambda} \|w\|_{X_\lambda} \\ &\lesssim \lambda^{-7/8} \log(1/\lambda) \|\tilde{u}\|_{X_\lambda} \|w\|_{X_\lambda} \lesssim \lambda^{-3/8} \log(1/\lambda) \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}. \end{aligned} \tag{2.11}$$

If $N^2 \sim \lambda N_1$ similar to the former argument by Lemma 4.12, (4.8) one has

$$\|\eta P_N \mathcal{B}(P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{U_K^2} \lesssim \log(N/\lambda) \|\tilde{u}\|_{U_{S_\lambda}^2} \|w\|_{U_{S_\lambda}^2}.$$

Then by (2.9) we have

$$\begin{aligned}
& \sum_{N \geq 2} N^{-3/4} \sum_{N_1 \sim N_2 \sim N^2/\lambda} \|\eta P_N \mathcal{B}(P_{N_1} \tilde{u} P_{N_2} \bar{w})\|_{U_K^2} \\
& \lesssim \sum_{N \geq 2} N^{-3/4} \sum_{N_1 \sim N_2 \sim N^2/\lambda} \log(N/\lambda) \|P_{N_1} \tilde{u}\|_{U_{S_\lambda}^2} \|P_{N_2} w\|_{U_{S_\lambda}^2} \\
& \lesssim \sum_{N \geq 2} N^{-3/4} \sum_{N_1 \sim N_2 \sim N^2/\lambda} \log(N/\lambda) N_2^{3/16} \\
& \quad \cdot (\lambda^{9/16} + \lambda^{17/24} N_1^{1/16} (\log N_1)^3) \|u\|_{X_\lambda} \|v\|_Y \|w\|_{X_\lambda} \\
& \lesssim \lambda^{3/8} \|u\|_{X_\lambda} \|v\|_Z \|w\|_{X_\lambda}.
\end{aligned} \tag{2.12}$$

Combining (2.10), (2.11), and (2.12) we conclude the proof. \square

3 The estimates for $F[u_0]$

In this section, we show some estimates for $F[u_0]$. For the nonresonant case, we show that it belongs to U^2 type space. See Lemma 3.1. Also we show $F[u_0] \in Z$.

Lemma 3.1. *Recall the definition of $F[u_0]$ by (2.1). We have*

$$\|N^{-3/4} P_N (F[u_0] - F[P_{\sim N^2/\lambda} u_0])\|_{l_N^2 U_K^2} \lesssim \lambda^{-1} \|u_0\|_{H^{-3/16}}^2.$$

Proof. If $N \lesssim 1$, we claim that

$$\left| \int_{\mathbb{R}^2} S_\lambda(t) P_{N_1} u_0 \overline{S_\lambda(t) P_{N_2} u_0} P_N \bar{v} \, dx dt \right| \lesssim N_1^{-3/8} \|u_0\|_{L^2}^2 \|v\|_{V_K^2} \tag{3.1}$$

where v is supported on $[-1/2, 1/2] \times \mathbb{R}$. Let $q > 2$. By the Hölder inequality, local smoothing, maximal function, and Strichartz estimates one has

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} S_\lambda(t) P_{N_1} u_0 \overline{S_\lambda(t) P_{N_2} u_0} P_N \bar{v} \, dx dt \right| \\
& \lesssim \|\tilde{\eta} S_\lambda(t) P_{N_1} u_0\|_{L_x^{2q/(q-2)} L_t^2} \|\tilde{\eta} S_\lambda(t) P_{N_2} u_0\|_{L_{t,x}^2} \|P_N v\|_{L_x^q L_t^\infty} \\
& \lesssim \|\tilde{\eta} S_\lambda(t) P_{N_1} u_0\|_{L_x^\infty L_t^2}^{2/q} \|u_0\|_{L_x^2}^{1+(q-2)/q} \|v\|_{V_K^2} \\
& \lesssim (\lambda N_1)^{-1/q} \|u_0\|_{L^2}^2 \|v\|_{V_K^2}.
\end{aligned}$$

By choosing $q = 8/3$ we obtain (3.1). Then by the duality one has

$$\begin{aligned}
& \sum_{N \lesssim 1} \sum_{N_1 \sim N_2} N^{-3/4} \left\| \eta P_N \mathcal{B}(S_\lambda(t) P_{N_1} u_0 \overline{S_\lambda(t) P_{N_2} u_0}) \right\|_{U_K^2} \\
& \lesssim \sum_{N_1 \sim N_2} (\lambda N_1)^{-3/8} \|P_{N_1} u_0\|_{L^2} \|P_{N_2} u_0\|_{L^2} \lesssim \lambda^{-3/8} \|u_0\|_{H^{3/16}}^2.
\end{aligned}$$

If $\lambda N_1 \ll N^2$, by Lemma 4.12 in [2] one has

$$\left| \int_{\mathbb{R}^2} \partial_x (P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) P_N \bar{v} \, dx dt \right| \lesssim \lambda^{-1/2} N^{-1} \|u_0\|_{L^2}^2 \|v\|_{U_K^2}.$$

By the Hölder inequality and the Strichartz estimate one also has

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} \partial_x (P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) P_N \bar{v} \, dx dt \right| \\ & \lesssim \|\tilde{\eta} P_{N_1} S_\lambda(t) u_0\|_{L_{t,x}^3} \|\tilde{\eta} P_{N_2} S_\lambda(t) u_0\|_{L_{t,x}^2} \|\partial_x P_N v\|_{L_{t,x}^6} \\ & \lesssim \|P_{N_1} S_\lambda(t) u_0\|_{L_{t,x}^6}^{1/2} \|u_0\|_{L^2}^{3/2} N^{5/6} \|v\|_{U_K^6} \\ & \lesssim \lambda^{-1/12} N^{5/6} \|u_0\|_{L^2}^2 \|v\|_{U_K^6} \end{aligned}$$

where v is supported on $[-1/2, 1/2] \times \mathbb{R}$. Then by the interpolation in [7] we have

$$\left\| \eta P_N \mathcal{B}(P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) \right\|_{U_K^2} \lesssim \lambda^{-1/2} N^{-1} \log N \|u_0\|_{L^2}^2.$$

Since $\max\{N_1, N_2, N\} \sim \text{med}\{N_1, N_2, N\}$, thus

$$\begin{aligned} & \sum_{N \gg 1} \sum_{\lambda N_1 \ll N^2} N^{-3/4} \left\| \eta P_N \mathcal{B}(P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) \right\|_{U_K^2} \\ & \lesssim \sum_{N \gg 1} \sum_{\lambda \max\{N_1, N_2\} \ll N^2} N^{-3/4} \lambda^{-1/2} N^{-1} \log N \|P_{N_1} u_0\|_{L^2} \|P_{N_2} u_0\|_{L^2} \\ & \lesssim \lambda^{-7/8} \|u_0\|_{H^{-3/16}}^2. \end{aligned}$$

If $\lambda N_1 \gg N^2$, by Lemma 4.12 in [2] one has

$$\left| \int_{\mathbb{R}^2} \partial_x (P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) P_N \bar{v} \, dx dt \right| \lesssim \lambda^{-1} N_1^{-1/2} \|u_0\|_{L^2}^2 \|v\|_{U_K^2}.$$

By similar argument as $\lambda N_1 \ll N^2$ and the interpolation in [7] we have

$$\left\| \eta P_N \mathcal{B}(P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) \right\|_{U_K^2} \lesssim \lambda^{-1} N_1^{-1/2} \log N_1 \|u_0\|_{L^2}^2.$$

Thus,

$$\begin{aligned} & \sum_{N \gg 1} \sum_{\lambda N_1 \gg N^2} N^{-3/4} \left\| \eta P_N \mathcal{B}(P_{N_1} S_\lambda(t) u_0 \overline{S_\lambda(t) P_{N_2} u_0}) \right\|_{U_K^2} \\ & \lesssim \sum_{N \gg 1} \sum_{N_1 \sim N_2} N^{-3/4} \lambda^{-1} N_1^{-1/2} \log N_1 \|P_{N_1} u_0\|_{L^2} \|P_{N_2} u_0\|_{L^2} \\ & \lesssim \lambda^{-1} \|u_0\|_{H^{-3/16}}^2. \end{aligned}$$

We conclude the proof. \square

Although we can not show the desired estimate for $F[P_{\sim N^2/\lambda} u_0]$ in U_K^2 , one has the following estimate.

Lemma 3.2. $\|N^{-3/4} P_N F[P_{\sim N^2/\lambda} u_0]\|_{l_N^2 V_K^2} \lesssim \lambda^{-3/8} \|u_0\|_{H^{-3/16}}^2$.

Proof. By Lemma 4.12 in [2] one has

$$\left| \int_{\mathbb{R}^2} \partial_x(|S_\lambda(t)P_{\sim N^2/\lambda} u_0|^2) P_N \bar{v} \, dxdt \right| \lesssim \|u_0\|_{L^2}^2 \|v\|_{U_K^2}$$

where v is supported on $[-1, 1] \times \mathbb{R}$. Then by the duality one has

$$\begin{aligned} \|N^{-3/4} P_N F[P_{\sim N^2/\lambda} u_0]\|_{l_N^2 V_K^2} &\lesssim \|N^{-3/4} \|P_{\sim N^2/\lambda} u_0\|_{L^2}^2\|_{l_N^2} \\ &\lesssim \|(\lambda M)^{-3/16} P_M u_0\|_{l_M^2 L^2}^2 \\ &\lesssim \lambda^{-3/8} \|u_0\|_{H^{-3/16}}^2. \end{aligned}$$

To show the second inequality in the former argument we use that the cardinality of $\{N \in 2^{\mathbb{N}_0} : N^2 \sim \lambda M\}$ is uniformly bounded for M . \square

We give some refinement for the case $\lambda N_1 \sim \lambda N_2 \sim N^2$ in following two lemmas.

Lemma 3.3. *Let $N \gg 1$, $0 < c \ll 1 \ll C$ and I_1, I_2 be intervals included in $[-CN^2, -cN^2] \cup [cN^2, CN^2]$ with length less than cN . We assume $J := I_1 - I_2 \subset [-CN, cN] \cup [cN, CN]$ and $|\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2| \gtrsim N$ for any $\xi \in J, \xi_j \in I_j, \xi_1 + \xi_2 = \xi$. Then we have*

$$\left\| \eta P_J \mathcal{B}(S_\lambda(t)P_{I_1} u_0 \overline{S_\lambda(t)P_{I_2} u_0}) \right\|_{U_K^2} \lesssim \lambda^{-1/2} \|u_0\|_{L^2}^2.$$

Proof. Let $L = cN$. Since $|\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2| > L/2$ for $\xi_j \in I_j, \xi \in J, \xi_1 + \xi_2 = \xi$, we have

$$\int_{\mathbb{R}^2} \partial_x(S_\lambda(t)P_{I_1} u_0 S_\lambda(t)P_{I_2} u_0) Q_{\leq L}^K P_J \bar{v} \, dxdt = 0.$$

By the high modulation estimate and Lemma 4.2, (4.2) in [2] one has

$$\begin{aligned} &\left| \int_{\mathbb{R}^2} \partial_x(S_\lambda(t)P_{I_1} u_0 \overline{S_\lambda(t)P_{I_2} u_0}) Q_{>L}^K P_J \bar{v} \, dxdt \right| \\ &\lesssim \|P_J(S_\lambda(s)P_{I_1} u_0 \overline{S_\lambda(s)P_{I_2} u_0})\|_{L_{t,x}^2} \|\partial_x Q_{>L}^K P_J v\|_{L_{t,x}^2} \\ &\lesssim (\lambda N)^{-1/2} \|u_0\|_{L^2}^2 L^{-1/2} N \|v\|_{V_K^2} \sim \lambda^{-1/2} \|u_0\|_{L^2}^2 \|v\|_{V_K^2}. \end{aligned}$$

By the duality we conclude the proof. \square

Lemma 3.4. Let $N \gg 1$, $0 < c \ll 1 \ll C$ and I_1, I_2 be intervals included in $[-CN^2, -cN^2] \cup [cN^2, CN^2]$ with length $1/N$. We assume J included in $[-CN, -cN] \cup [cN, CN]$ with length $1/N^2$ and $|\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2| \lesssim 1$ for any $\xi \in J$, $\xi_j \in I_j$, $\xi_1 + \xi_2 = \xi$. Then we have

$$\left\| \eta P_J \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right\|_{U_K^2} \lesssim \|u_0\|_{L^2}^2.$$

Proof. Let $P_{I_1} u_0 = u_1$, $P_{I_2} u_0 = u_2$. By the definition of U_K^2 and the Taylor expansion one has

$$\begin{aligned} & \left\| \eta P_J \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right\|_{U_K^2} \\ &= \left\| \eta P_J \partial_x \int_0^t K(-s) (S_\lambda(s) u_1 \overline{S_\lambda(s) u_2}) ds \right\|_{U_t^2(L_x^2)} \\ &\sim \left\| \eta(t) \xi \chi_{\xi \in J} \int_0^t e^{-is\xi^3} \int_{\xi_1 + \xi_2 = \xi} e^{-i\lambda s\xi_1^2 + i\lambda s\xi_2^2} \widehat{u_1}(\xi_1) \overline{\widehat{u_2}}(-\xi_2) ds \right\|_{U_t^2(L_\xi^2)} \\ &\lesssim \sum_{k=0}^{\infty} \frac{1}{k!} \left\| \eta(t) \xi \chi_{\xi \in J} \int_0^t \int_{\xi_1 + \xi_2 = \xi} \widehat{u_1}(\xi_1) \overline{\widehat{u_2}}(-\xi_2) s^k (\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2)^k ds \right\|_{U_t^2 L_\xi^2} \\ &\lesssim \sum_{k=0}^{\infty} \frac{C^k N}{k!} \left\| \chi_{\xi \in J} \int_{\xi_1 + \xi_2 = \xi} |\widehat{u_1}(\xi_1) \overline{\widehat{u_2}}(-\xi_2)| ds \right\|_{L_\xi^2} \left\| \eta(t) \frac{t^{k+1} - 1}{k+1} \right\|_{U_t^2} \\ &\lesssim N \|u_1\|_{L^2} \|u_2\|_{L^2} |J|^{1/2} \lesssim \|u_0\|_{L^2}^2. \end{aligned}$$

Here we use $\|\eta(t)t^{k+1}\|_{U_t^2} \lesssim \|\eta(t)t^{k+1}\|_{H^1} \lesssim k$. \square

Lemma 3.5. $\|N^{1/4} P_N F[u_0]\|_{l_N^2 L_x^\infty L_t^2} \lesssim \lambda^{-1} \|u_0\|_{H^{-3/16}}^2$.

Proof. Firstly by Lemma 3.1 and the local smoothing estimate one has

$$\|N^{1/4} P_N (F[u_0] - F[P_{\sim N^2/\lambda} u_0])\|_{l_N^2 L_x^\infty L_t^2} \lesssim \lambda^{-1} \|u_0\|_{H^{-3/16}}^2$$

Thus we only need to show

$$\|N^{1/4} P_N F[P_{\sim N^2/\lambda} u_0]\|_{l_N^2 L_x^\infty L_t^2} \lesssim \lambda^{-1} \|u_0\|_{H^{-3/16}}^2.$$

If $N \lesssim 1$, by the Sobolev inequality, then

$$N^{1/4} \|P_N F[P_{\sim N^2/\lambda} u_0]\|_{L_x^\infty L_t^2} \lesssim \|P_{\sim N^2/\lambda} u_0\|_{L_x^4}^2 \lesssim \lambda^{-7/8} \|u_0\|_{H^{-3/16}}^2.$$

If $N \gg 1$, we claim

$$\|P_N F[P_{\sim N^2/\lambda} u_0]\|_{L_x^\infty L_t^2} \lesssim \lambda^{-1/2} N^{-1} \|u_0\|_{L^2}^2. \quad (3.2)$$

Let I_j , $j = 1, 2$ be intervals included in $[-CN^2/\lambda, cN^2/\lambda] \cup [cN^2/\lambda, CN^2/\lambda]$ with length cN . Also we assume for any $\xi_j \in I_j$, $j = 1, 2$ one has $|\eta_1 - \eta_2| \sim N$. We only need to show

$$\left\| \eta P_N \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} v_0}) \right\|_{L_x^\infty L_t^2} \lesssim \lambda^{-1/2} N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2}.$$

There exists a interval J included in $[-N, -N/2] \cup [N/2, N]$ with length 1 such that $|\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2| \gtrsim N^2$ for any $\xi \in [-N, -N/2] \cup [N/2, N] \setminus J := \hat{J}$, $\xi_j \in I_j$, $\xi = \xi_1 + \xi_2$. Then by Lemma 3.3, the local smoothing estimate, and the orthogonality we only need to show

$$\left\| \eta P_J \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} v_0}) \right\|_{L_x^\infty L_t^2} \lesssim \lambda^{-1/2} N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2}$$

where I_1, I_2, J are intervals with length 1, $|\xi_j| \sim N^2$ for any $\xi_j \in I_j$, $|\xi| \sim N$ for any $\xi \in J$.

There exists a interval J_1 included in J with length $1/N$ such that $|\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2| \gtrsim N$ for any $\xi \in J \setminus J_1$, $\xi_j \in I_j$, $\xi = \xi_1 + \xi_2$. Then again by Lemma 3.3, the local smoothing estimate, and the orthogonality we only need to show

$$\left\| \eta P_{J_1} \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} v_0}) \right\|_{L_x^\infty L_t^2} \lesssim \lambda^{-1/2} N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2}$$

where I_1, I_2, J_1 are intervals with length $1/N$.

There exists a interval J_2 included in J_1 with length $1/N^2$ such that one has $|\xi^3 + \lambda\xi_1^2 - \lambda\xi_2^2| \lesssim 1$ for any $\xi \in J_2$, $\xi_j \in I_j$, $\xi = \xi_1 + \xi_2$. Then by Lemma 3.4 and the local smoothing estimate one has

$$\begin{aligned} & \left\| \eta P_{J_2} \mathcal{B}(S_\lambda(t) P_I u_0 \overline{S_\lambda(t) P_J v_0}) \right\|_{L_x^\infty L_t^2} \\ & \lesssim N^{-1} \left\| \eta P_{J_2} \mathcal{B}(S_\lambda(t) P_I u_0 \overline{S_\lambda(t) P_J v_0}) \right\|_{U_K^2} \\ & \lesssim N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2}. \end{aligned}$$

Note that

$$\begin{aligned} & \mathcal{F}_x \left(\eta(t) P_{J_1 \setminus J_2} \mathcal{B}(S_\lambda(t) P_I u_0 \overline{S_\lambda(t) P_J v_0}) \right) \\ & \sim \eta(t) \xi \int_0^t \int_{\xi_1 + \xi_2 = \xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} e^{i(t-s)\xi^3} e^{-i\lambda s \xi_1^2 + i\lambda s \xi_2^2} \widehat{u_0}(\xi_1) \overline{\widehat{v_0}}(-\xi_2) \\ & \sim \eta(t) \int_{\xi_1 + \xi_2 = \xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} \frac{e^{it\xi^3} - e^{i\lambda t \xi (\xi_2 - \xi_1)}}{\xi^2 + \lambda \xi_1 - \lambda \xi_2} \widehat{u_0}(\xi_1) \overline{\widehat{v_0}}(-\xi_2) \\ & := \Lambda_1 + \Lambda_2. \end{aligned}$$

By the local smoothing estimate we have

$$\begin{aligned}
\|\mathcal{F}_\xi^{-1}\Lambda_1\|_{L_x^\infty L_t^2} &\lesssim N^{-1} \left\| \int_{\xi_1+\xi_2=\xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} \frac{\widehat{u}_0(\xi_1) \overline{\widehat{v}_0}(-\xi_2)}{\xi^2 + \lambda\xi_1 - \lambda\xi_2} \right\|_{L_\xi^2} \\
&\lesssim N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2} \left\| \frac{\chi_{|\xi| \sim J_1 \setminus J_2, \xi_1 \in I_1, \xi - \xi_1 \in I_2}}{\xi^2 + 2\lambda\xi_1 - \lambda\xi} \right\|_{L_{\xi_1}^\infty L_\xi^2} \\
&\lesssim N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2}.
\end{aligned}$$

Let ξ_j^0 be the center of I_j respectively. By the Taylor expansion we have

$$\begin{aligned}
&\|\mathcal{F}_\xi^{-1}\Lambda_2\|_{L_x^\infty L_t^2} \\
&\lesssim \sum_{k=0}^{\infty} \frac{1}{k!} \left\| \eta(t) \mathcal{F}_\xi^{-1} \int_{\xi_1+\xi_2=\xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} \frac{e^{i\lambda t \xi (\xi_2^0 - \xi_1^0)}}{\xi^2 + \lambda\xi_1 - \lambda\xi_2} \widehat{u}_0(\xi_1) \overline{\widehat{v}_0}(-\xi_2) \right. \\
&\quad \cdot (i\lambda t \xi (\xi_2 - \xi_2^0 - \xi_1 + \xi_1^0))^k \Bigg\|_{L_x^\infty L_t^2} \\
&\lesssim \sum_{k=0}^{\infty} \frac{1}{k!} \left\| \mathcal{F}_\xi^{-1} \int_{\xi_1+\xi_2=\xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} \frac{e^{i\lambda t \xi (\xi_2^0 - \xi_1^0)}}{\xi^2 + \lambda\xi_1 - \lambda\xi_2} \widehat{u}_0(\xi_1) \overline{\widehat{v}_0}(-\xi_2) \right. \\
&\quad \cdot \xi^k (\xi_2 - \xi_2^0 - \xi_1 + \xi_1^0)^k \Bigg\|_{L_x^\infty L_t^2}.
\end{aligned}$$

Note that $|\xi_2^0 - \xi_1^0| \sim N^2$. By the local smoothing estimate one has

$$\begin{aligned}
\|\mathcal{F}_\xi^{-1}\Lambda_2\|_{L_x^\infty L_t^2} &\lesssim N^{-1} \sum_{k=0}^{\infty} \frac{1}{k!} \left\| \int_{\xi_1+\xi_2=\xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} \frac{\widehat{u}_0(\xi_1) \overline{\widehat{v}_0}(-\xi_2)}{\xi^2 + \lambda\xi_1 - \lambda\xi_2} \right. \\
&\quad \cdot \xi^k (\xi_2 - \xi_2^0 - \xi_1 + \xi_1^0)^k \Bigg\|_{L_\xi^2} \\
&\lesssim \sum_{k=0}^{\infty} \frac{C^k}{k!} N^{-1} \left\| \int_{\xi_1+\xi_2=\xi} \chi_{|\xi| \sim J_1 \setminus J_2, \xi_j \in I_j} \frac{\widehat{u}_0(\xi_1) \overline{\widehat{v}_0}(-\xi_2)}{\xi^2 + \lambda\xi_1 - \lambda\xi_2} \right\|_{L_\xi^2} \\
&\lesssim N^{-1} \|u_0\|_{L^2} \|v_0\|_{L^2}.
\end{aligned}$$

We conclude the proof of (3.2). Then

$$\begin{aligned}
\|N^{1/4} P_N F[P_{\sim N^2/\lambda} u_0]\|_{l_{N \gg 1}^2 L_x^\infty L_t^2} &\lesssim \lambda^{-1/2} \|N^{1/4} N^{-1} \|P_{\sim N^2/\lambda} u_0\|_{L^2}^2\|_{l_N^2} \\
&\lesssim \lambda^{-7/8} \|u_0\|_{H^{-3/16}}^2.
\end{aligned}$$

Combining the nonresonant and low frequency cases, we finish the proof. \square

Recall the definition for the norm Z by (2.3). We have:

Corollary 3.6. *Combining Lemmas 3.1, 3.2, and 3.5 then*

$$\|F[u_0]\|_Z \lesssim \lambda^{-1} \|u_0\|_{H^{-3/16}}^2.$$

4 Bilinear estimates for KdV equation

We mainly use the argument in [10], §5.3.

Lemma 4.1. *Let $N_1, N_2 \geq 2$, $N_1 \sim N_2$. Then*

$$\|\eta P_{\lesssim 1} \mathcal{B}(P_{N_1} u P_{N_2} v)\|_{L_x^2 L_t^\infty} \lesssim N_1^{-3/2} \|u\|_{V_K^2} \|v\|_{V_K^2}.$$

Proof. If $N_1 \lesssim 1$ we have

$$\begin{aligned} \|\eta P_{\lesssim 1} \mathcal{B}(P_{N_1} u P_{N_2} v)\|_{L_x^2 L_t^\infty} &\lesssim \|\tilde{\eta} P_{N_1} u P_{N_2} v\|_{L_t^1 L_x^2} \\ &\lesssim \|P_{N_1} u\|_{L_t^\infty L_x^4} \|P_{N_2} v\|_{L_t^\infty L_x^4} \\ &\lesssim N_1^{-3/2} \|u\|_{L_t^\infty L_x^2} \|v\|_{L_t^\infty L_x^2}. \end{aligned}$$

Thus we consider the case $N_1 \sim N_2 \gg 1$. Let $L_{\min} = \min\{L_1, L_2\}$, $L_{\max} = \max\{L_1, L_2\}$. We decompose $P_{\lesssim 1}$ into $\sum_{K \lesssim 1} P_K$. By maximal function estimate, Lemma 5.7, (5.37) and (5.38) in [10] one has

$$\begin{aligned} &\sum_{K \lesssim N_1^{-2/3}} \|\eta P_K(\mathcal{B}(P_{N_1} u P_{N_2} v))\|_{L_x^2 L_t^\infty} \\ &\lesssim \sum_{K \lesssim N_1^{-2/3}} \sum_{L_1, L_2, L} K L^{-1/2} \|Q_{L_1}^K P_{N_1} u Q_{L_2}^K P_{N_2} v\|_{L_{t,x}^2} \\ &\lesssim \sum_{K \lesssim N_1^{-2/3}} K \sum_{L_1, L_2, L} L^{-1/2} \min\{L_{\min}^{1/2} K^{1/2}, L_{\min}^{1/2} L^{1/2} N_1^{-1}\} \\ &\quad \cdot \|P_{N_1} Q_{L_1}^K u\|_{L_{t,x}^2} \|P_{N_2} Q_{L_2}^K v\|_{L_{t,x}^2} \\ &\lesssim \sum_{K \lesssim N_1^{-2/3}} K \sum_{L_1, L_2, L} (L L_{\max})^{-1/2} \min\{K^{1/2}, L^{1/2} N_1^{-1}\} \|u\|_{V_K^2} \|v\|_{V_K^2} \\ &\lesssim \sum_{K \lesssim N_1^{-2/3}} K \sum_L L^{-1/2} \min\{K^{1/2}, L^{1/2} N_1^{-1}\} \|u\|_{V_K^2} \|v\|_{V_K^2} \\ &\lesssim N_1^{-3/2} \|u\|_{V_K^2} \|v\|_{V_K^2}. \end{aligned}$$

If $K \geq N_1^{-2/3}$, we firstly manipulate the case $L_{\max} \geq N_1^{5/4}$. Similar to the former argument one has

$$\begin{aligned} &\sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} \geq N_1^{5/4}} \|\eta P_K(\mathcal{B}(P_{N_1} Q_{L_1}^K u P_{N_2} Q_{L_2}^K v))\|_{L_x^2 L_t^\infty} \\ &\lesssim \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} \geq N_1^{5/4}, L} K (L L_{\max})^{-1/2} \min\{K^{1/2}, L^{1/2} N_1^{-1}\} \|u\|_{V_K^2} \|v\|_{V_K^2} \\ &\lesssim \sum_{L_{\max} \geq N_1^{5/4}} L_{\max}^{-1/2} N_1^{-1} \log(N_1) \|u\|_{V_K^2} \|v\|_{V_K^2} \\ &\lesssim N_1^{-3/2} \|u\|_{V_K^2} \|v\|_{V_K^2}. \end{aligned} \tag{4.1}$$

Thus we only need to estimate $L_{\max} < N_1^{5/4}$ for $N_1^{-2/3} \leq K \lesssim 1$. Let

$$\begin{aligned} f_{N_1, L_1}(\tau, \xi) &= \mathcal{F}_{t,x}(P_{N_1} Q_{L_1}^K u)(\tau + \xi^3, \xi), \\ g_{N_2, L_2}(\tau, \xi) &= \mathcal{F}_{t,x}(P_{N_2} Q_{L_2}^K v)(\tau + \xi^3, \xi). \end{aligned}$$

Then

$$\begin{aligned} &\mathcal{F}_x \left(\sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} P_K(\eta \mathcal{B}(P_{N_1} Q_{L_1}^K u P_{N_2} Q_{L_2}^K v)) \right) \\ &\sim \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} \eta(t) \int_0^t \psi(\xi/K) \xi e^{i(t-s)\xi^3} \\ &\quad \cdot \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \int_{\xi = \xi_1 + \xi_2} f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2) e^{it(\tau_1 + \xi_1^3) + it(\tau_2 + \xi_2^3)} \\ &\sim \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} \eta(t) \psi(\xi/K) \xi \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \int_{\xi = \xi_1 + \xi_2} \\ &\quad \cdot \frac{e^{it(\tau_1 + \tau_2 + \xi_1^3 + \xi_2^3)} - e^{it\xi^3}}{\tau_1 + \tau_2 + \xi_1^3 + \xi_2^3 - \xi^3} f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2) \\ &:= \mathcal{F}_x I - \mathcal{F}_x II. \end{aligned}$$

Note that on the support set of $f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2)$ one has $|\tau_1 + \tau_2 + \xi_1^3 + \xi_2^3 - \xi^3| \sim N_1^2 |\xi|$ since $L_{\max} < N_1^{5/4}$, $|\xi| \sim K \geq N_1^{-2/3}$. By the maximal function estimate and the Plancherel identity one has

$$\begin{aligned} &\|II\|_{L_x^2 L_t^\infty} \\ &\sim \left\| \int_{\substack{\xi = \xi_1 + \xi_2, \\ \tau_1, \tau_2}} \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} \frac{\psi(\xi/K) \xi f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2)}{\tau_1 + \tau_2 + \xi_1^3 + \xi_2^3 - \xi^3} \right\|_{L_\xi^2} \\ &\lesssim N_1^{-2} \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \sum_{L_{\max} < N_1^{5/4}} \|f_{N_1, L_1}(\tau_1, \cdot)\|_{L^2} \|g_{N_2, L_2}(\tau_2, \cdot)\|_{L^2} \\ &\lesssim N_1^{-2} \sum_{L_{\max} < N_1^{5/4}} L_1^{1/2} L_2^{1/2} \|f_{N_1, L_1}(\tau, \xi)\|_{L_{\tau, \xi}^2} \|g_{N_2, L_2}(\tau, \xi)\|_{L_{\tau, \xi}^2} \\ &\lesssim N_1^{-2} \sum_{L_{\max} < N_1^{5/4}} \|u\|_{V_K^2} \|v\|_{V_K^2} \lesssim N_1^{-3/2} \|u\|_{V_K^2} \|v\|_{V_K^2}. \end{aligned}$$

Define

$$\begin{aligned}
\mathcal{F}_x I' &:= \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} \eta(t) \psi(\xi/K) \xi \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \int_{\xi=\xi_1+\xi_2} \\
&\quad \frac{e^{it(\tau_1+\tau_2+\xi_1^3+\xi_2^3)}}{\xi_1^3 + \xi_2^3 - \xi^3} f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2) \\
&= \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} \eta(t) \psi(\xi/K) \int_{\xi=\xi_1+\xi_2} \\
&\quad \frac{1}{-3\xi_1\xi_2} \mathcal{F}_x(P_{N_1} Q_{L_1}^K u)(t, \xi_1) \mathcal{F}_x(P_{N_2} Q_{L_2}^K v)(t, \xi_2).
\end{aligned}$$

Let $f := \sum_{L_1 < N_1^{5/4}} f_{N_1, L_1}$, $g := \sum_{L_2 < N_1^{5/4}} g_{N_2, L_2}$. Then

$$\begin{aligned}
\mathcal{F}_x I - \mathcal{F}_x I' &= \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} \sum_{k=1}^{\infty} \eta(t) \psi(\xi/K) \xi \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \int_{\xi=\xi_1+\xi_2} \\
&\quad \frac{(-1)^k (\tau_1 + \tau_2)^k e^{it(\tau_1+\tau_2+\xi_1^3+\xi_2^3)}}{(\xi_1^3 + \xi_2^3 - \xi^3)^{k+1}} f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2).
\end{aligned}$$

Thus by the Strichartz estimate [8] one has

$$\begin{aligned}
&\|I - I'\|_{L_x^2 L_t^\infty} \\
&\lesssim \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \sum_{k=1}^{\infty} \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} |\tau_1 + \tau_2|^k K^{-k} \\
&\quad \cdot \left\| \eta(t) \mathcal{F}_\xi^{-1} \int_{\xi=\xi_1+\xi_2} \left(\frac{e^{it(\xi_1^3+\xi_2^3)}}{\xi_1^{k+1} \xi_2^{k+1}} f_{N_1, L_1}(\tau_1, \xi_1) g_{N_2, L_2}(\tau_2, \xi_2) \right) \right\|_{L_x^2 L_t^\infty} \\
&\lesssim \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \sum_{k=1}^{\infty} \sum_{N_1^{-2/3} \leq K \lesssim 1} \sum_{L_{\max} < N_1^{5/4}} N_1^{5k/4} K^{-k} \\
&\quad \cdot \|\mathcal{F}_\xi^{-1}(e^{it\xi^3} f_{N_1, L_1}(\tau_1, \xi)/\xi^{k+1})\|_{L_x^4 L_t^\infty} \|\mathcal{F}_\xi^{-1}(e^{it\xi^3} f_{N_2, L_2}(\tau_2, \xi)/\xi^{k+1})\|_{L_x^4 L_t^\infty} \\
&\lesssim \int_{\mathbb{R}_{\tau_1, \tau_2}^2} \sum_{k=1}^{\infty} \sum_{L_{\max} < N_1^{5/4}} N_1^{23k/12 - 2(k+1)+1/2} \|f_{N_1, L_1}(\tau_1, \cdot)\|_{L^2} \|f_{N_2, L_2}(\tau_2, \cdot)\|_{L^2} \\
&\lesssim \sum_{L_{\max} < N_1^{5/4}} N_1^{-37/12} (L_1 L_2)^{1/2} \|Q_{L_1}^K P_{N_1} u\|_{L_{t,x}^2} \|Q_{L_2}^K P_{N_2} v\|_{L_{t,x}^2} \\
&\lesssim \sum_{L_{\max} < N_1^{5/4}} N_1^{-37/12} \|u\|_{V_K^2} \|v\|_{V_K^2} \lesssim N^{-3/2} \|u\|_{V_K^2} \|v\|_{V_K^2}.
\end{aligned}$$

Note that there exists constant \tilde{c} such that

$$I' = \tilde{c} \eta(t) P_{N^{-2/3} \leq \cdot \lesssim 1} ((\partial_x^{-1} P_{N_1} Q_{< N_1^{5/4}}^K u) (\partial_x^{-1} P_{N_1} Q_{< N_1^{5/4}}^K v)).$$

By the Hölder inequality we have

$$\begin{aligned}
\|I'\|_{L_x^2 L_t^\infty} &\lesssim \|\partial_x^{-1} P_{N_1} Q_{< N_1^{5/4}}^K u\|_{L_x^4 L_t^\infty} \|\partial_x^{-1} P_{N_1} Q_{< N_1^{5/4}}^K v\|_{L_x^4 L_t^\infty} \\
&\lesssim N^{-2} \|P_{N_1} Q_{< N_1^{5/4}}^K u\|_{L_x^4 L_t^\infty} \|P_{N_2} Q_{< N_1^{5/4}}^K v\|_{L_x^4 L_t^\infty} \\
&\lesssim N^{-3/2} \|Q_{< N_1^{5/4}}^K u\|_{U_K^4} \|Q_{< N_1^{5/4}}^K v\|_{U_K^4} \\
&\lesssim N^{-3/2} \|Q_{< N_1^{5/4}}^K u\|_{V_K^2} \|Q_{< N_1^{5/4}}^K v\|_{V_K^2} \lesssim N^{-3/2} \|u\|_{V_K^2} \|v\|_{V_K^2}.
\end{aligned}$$

We conclude the proof. \square

Proposition 4.2. $\|\eta \mathcal{B}(uv)\|_Y \lesssim \|u\|_Y \|v\|_Y$.

Proof. By Lemma 4.1 we have

$$\begin{aligned}
&\|\eta P_1 \mathcal{B}(uv)\|_{L_x^2 L_t^\infty} \\
&\leq \|\eta P_1 \mathcal{B}(P_{\lesssim 1} u P_{\lesssim 1} v)\|_{L_x^2 L_t^\infty} + \sum_{N_1 \sim N_2 \gg 1} \|\eta P_1 \mathcal{B}(P_{N_1} u P_{N_2} v)\|_{L_x^2 L_t^\infty} \\
&\lesssim \|\tilde{\eta} P_{\lesssim 1} u P_{\lesssim 1} v\|_{L_t^1 L_x^2} + \sum_{N_1 \sim N_2 \gg 1} N_1^{-3/2} \|P_{N_1} u\|_{V_K^2} \|P_{N_2} v\|_{V_K^2} \\
&\lesssim \|P_{\lesssim 1} u\|_{L_t^\infty L_x^4} \|P_{\lesssim 1} v\|_{L_t^\infty L_x^4} + \|u\|_Z \|v\|_Z \\
&\lesssim \|u\|_Z \|v\|_Z.
\end{aligned}$$

Let $N \geq 2$. By the triangle inequality we have

$$\begin{aligned}
&\|\eta P_N \mathcal{B}(uv)\|_{U_K^2} \\
&\lesssim \sum_{N_1 \sim N} \|\eta P_N \mathcal{B}(P_{N_1} u P_{\lesssim 1} v)\|_{U_K^2} + \sum_{N_2 \sim N} \|\eta P_N \mathcal{B}(P_{\lesssim 1} u P_{N_2} v)\|_{U_K^2} \\
&\quad + \sum_{N_1, N_2 \gg 1} \|\eta P_N \mathcal{B}(P_{N_1} u P_{N_2} v)\|_{U_K^2} := I_N + I'_N + II_N.
\end{aligned}$$

For the term I we use Proposition 5.12 in [10].

$$\begin{aligned}
\|I_N\|_{U_K^2} &\lesssim \sum_{N_1 \sim N} \|\tilde{\eta}^2 \partial_x (P_{N_1} u P_{\lesssim 1} v)\|_{L_{t,x}^2} \\
&\lesssim \sum_{N_1 \sim N} N \|\tilde{\eta} P_{N_1} u\|_{L_x^\infty L_t^2} \|\tilde{\eta} P_{\lesssim 1} v\|_{L_x^2 L_t^\infty}.
\end{aligned}$$

Then by the local smoothing estimate one has

$$\begin{aligned}
\|N^{-3/4} I_N\|_{l_{N \geq 2}^2 U_K^2} &\lesssim \|\tilde{\eta} N^{1/4} P_N u\|_{l_N^2 L_x^\infty L_t^2} \|\tilde{\eta} P_{\lesssim 1} v\|_{L_x^2 L_t^\infty} \\
&\lesssim \|u\|_Z \|v\|_Z
\end{aligned}$$

By symmetry we control I'_N similarly.

For II_N we decompose the summation into four parts.

$$II_N = \sum_{N_1 \sim N_2 \sim N \gg 1} + \sum_{N_1 \sim N \gg N_2 \gg 1} + \sum_{N_2 \sim N \gg N_1 \gg 1} + \sum_{N_1 \sim N_2 \gg N} \dots \\ := II_N^{(1)} + II_N^{(2)} + II_N^{(3)} + II_N^{(4)}.$$

For $N_1 \sim N_2 \sim N \gg 1$ we show

$$\left| \int_{\mathbb{R}^2} \partial_x(P_{N_1}uP_{N_2}v)P_N\bar{w} dxdt \right| \lesssim N^{-3/4} \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2}$$

where u, v, w are supported on $[-1, 1] \times \mathbb{R}$. Let $L_{\min}, L_{\text{med}}, L_{\max}$ be the maximum, medium and minimum among L_1, L_2, L . By Lemma 5.7 in [10] we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} \partial_x(P_{N_1}uP_{N_2}v)P_N\bar{w} dxdt \right| \\ & \leq \sum_{L_{\max} \gtrsim cN^3} \left| \int_{\mathbb{R}^2} \partial_x(Q_{L_1}^K P_{N_1}u Q_{L_2}^K P_{N_2}v) \overline{P_N Q_L^K w} dxdt \right| \\ & \lesssim \sum_{L_{\max} \gtrsim cN^3} NL_{\min}^{1/2} L_{\text{med}}^{1/4} N^{-1/4} \|Q_{L_1}^K P_{N_1}u\|_{L_{t,x}^2} \|Q_{L_2}^K P_{N_2}v\|_{L_{t,x}^2} \|Q_L^K P_N w\|_{L_{t,x}^2} \\ & \lesssim \sum_{L_{\max} \gtrsim cN^3} L_{\min}^{1/2} L_{\text{med}}^{1/4} N^{3/4} (L_1 L_2 L)^{-1/2} \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2} \\ & \lesssim N^{-3/4} \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2} \end{aligned}$$

Then by the duality we have

$$\begin{aligned} \|N^{-3/4} II_N^{(1)}\|_{l_{N \geq 2}^2 V_K^2} & \lesssim \left\| N^{-3/2} \sum_{N_1 \sim N_2 \sim N \gg 1} \|P_{N_1}u\|_{V_K^2} \|P_{N_2}v\|_{V_K^2} \right\|_{l_{N \geq 2}^2} \\ & \lesssim \|N^{-3/4} P_N u\|_{l_{N \geq 2}^2 V_K^2} \|N^{-3/4} P_N v\|_{l_{N \geq 2}^2 V_K^2} \\ & \lesssim \|u\|_Z \|v\|_Z. \end{aligned}$$

For $N_1 \sim N \gg N_2 \gg 1$ we show

$$\left| \int_{\mathbb{R}^2} \partial_x(P_{N_1}uP_{N_2}v)P_N\bar{w} dxdt \right| \lesssim N^{-1/2} N_2^{1/4} \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2}.$$

In fact by Lemma 5.7 (5.38) in [10] we have

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} \partial_x(P_{N_1} u P_{N_2} v) P_N \bar{w} \, dx dt \right| \\
& \leq \sum_{L_1, L_2, L} \left| \int_{\mathbb{R}^2} \partial_x(Q_{L_1}^K P_{N_1} u Q_{L_2}^K P_{N_2} v) \overline{P_N Q_L^K w} \, dx dt \right| \\
& \lesssim \sum_{L_1, L_2, L} N(L_1 L_2 L)^{1/3} N^{-1/2} (N^2 N_2)^{-1/2} \\
& \quad \cdot \|Q_{L_1}^K P_{N_1} u\|_{L_{t,x}^2} \|Q_{L_2}^K P_{N_2} v\|_{L_{t,x}^2} \|Q_L^K P_N w\|_{L_{t,x}^2} \\
& \lesssim \sum_{L_1, L_2, L} (L_1 L_2 L)^{1/3} N^{-1/2} N_2^{-1/2} (L_1 L_2 L)^{-1/2} \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2} \\
& \lesssim N^{-1/2} N_2^{-1/2} \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2}.
\end{aligned}$$

Then by the duality we have

$$\begin{aligned}
\|N^{-3/4} II_N^{(2)}\|_{l_{N \geq 2}^2 U_K^2} & \lesssim \sum_{N_1 \sim N \gg N_2 \gg 1} N^{-5/4} N_2^{-1/2} \|P_{N_1} u\|_{V_K^2} \|P_{N_2} v\|_{V_K^2} \\
& \lesssim \|N^{-3/4} P_N u\|_{l_{N \geq 2}^2 V_K^2} \|N^{-3/4} P_N v\|_{l_{N \geq 2}^2 V_K^2} \\
& \lesssim \|u\|_Z \|v\|_Z.
\end{aligned}$$

By the symmetry we can control $II_N^{(3)}$ similarly.

Now, we estimate the term $II_N^{(4)}$. Let $N_1 \sim N_2 \gg N \geq 2$, $L = c N_1^2 N$. Firstly we estimate

$$\|\eta P_N \mathcal{B}(Q_{\leq L}^K (P_{N_1} u P_{N_2} v))\|_{U_K^2}.$$

Typically we control

$$\int_{\mathbb{R}^2} \partial_x(Q_{>L}^K P_{N_1} u P_{N_2} v) \overline{Q_{\leq L}^K P_N w} \, dx dt.$$

By high modulation and transversal estimates, we have

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} \partial_x(Q_{>L}^K P_{N_1} u P_{N_2} v) \overline{Q_{\leq L}^K P_N w} \, dx dt \right| \\
& \lesssim \|Q_{>L}^K P_{N_1} u\|_{L_{t,x}^2} \|P_{N_2} v \partial_x Q_{\leq L}^K P_N w\|_{L_{t,x}^2} \\
& \lesssim L^{-1/2} \|u\|_{V_K^2} N_2^{-1} N \|v\|_{U_K^2} \|w\|_{U_K^2} \\
& \sim N_1^{-2} N^{1/2} \|u\|_{V_K^2} \|v\|_{U_K^2} \|w\|_{U_K^2}.
\end{aligned}$$

On the other hand by the Strichartz estimate we have

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} \partial_x(Q_{>L}^K P_{N_1} u P_{N_2} v) \overline{Q_{\leq L}^K P_N w} \, dx dt \right| \\
& \lesssim \|Q_{>L}^K P_{N_1} u\|_{L_{t,x}^4} \|P_{N_2} v\|_{L_{t,x}^4} \|\partial_x Q_{\leq L}^K P_N w\|_{L_{t,x}^2} \\
& \lesssim \|Q_{>L}^K P_{N_1} u\|_{L_{t,x}^2}^{1/4} \|Q_{>L}^K P_{N_1} u\|_{L_{t,x}^6}^{3/4} \|P_{N_2} v\|_{L_{t,x}^2}^{1/4} \|P_{N_2} v\|_{L_{t,x}^6}^{3/4} N \|w\|_{U_K^6} \\
& \sim N_1^{-1/4} N \|u\|_{V_K^2} \|v\|_{U_K^6} \|w\|_{U_K^6}.
\end{aligned}$$

Then, by the interpolation [7], one has

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} \partial_x(Q_{>L}^K P_{N_1} u P_{N_2} v) \overline{Q_{\leq L}^K P_N w} \, dx dt \right| \\
& \lesssim N_1^{-2} N^{1/2} (\log N_1)^2 \|u\|_{V_K^2} \|v\|_{V_K^2} \|w\|_{V_K^2}.
\end{aligned}$$

Thus,

$$\begin{aligned}
& \sum_{N_1 \sim N_2 \gg N \geq 2} N^{-3/4} \|\eta P_N \mathcal{B}(Q_{\leq L}^K (P_{N_1} u P_{N_2} v))\|_{U_K^2} \\
& \lesssim \sum_{N_1 \sim N_2 \gg N \geq 2} N^{-3/4} N_1^{-2} N^{1/2} (\log N_1)^2 \|P_{N_1} u\|_{V_K^2} \|P_{N_2} v\|_{V_K^2} \\
& \lesssim \|u\|_Z \|v\|_Z.
\end{aligned}$$

Secondly we estimate $\|\eta P_N \mathcal{B}(Q_{\leq L}^K (P_{N_1} u P_{N_2} v))\|_{U_K^2}$. By high modulation and transversal estimates, we have

$$\begin{aligned}
& \left| \int_{\mathbb{R}^2} \partial_x(P_{N_1} u P_{N_2} v) \overline{Q_{>L}^K P_N w} \, dx dt \right| \\
& \lesssim \|P_N(P_{N_1} u P_{N_2} v)\|_{L_{t,x}^2} \|\partial_x Q_{>L}^K P_N w\|_{L_{t,x}^2} \\
& \lesssim (NN_1)^{-1/2} \|u\|_{U_K^2} \|v\|_{U_K^2} L^{-1/2} N \|w\|_{V_K^2} \\
& \sim N_1^{-3/2} \|u\|_{U_K^2} \|v\|_{U_K^2} \|w\|_{V_K^2}.
\end{aligned}$$

Then by the duality one has

$$\begin{aligned}
& \sum_{N_1 \sim N_2 \gg N \geq 2} \|N^{-3/4} \eta P_N \mathcal{B}(Q_{\leq L}^K (P_{N_1} u P_{N_2} v))\|_{U_K^2} \\
& \lesssim \sum_{N_1 \sim N_2 \gg N \geq 2} N^{-3/4} N_1^{-3/2} \|P_{N_1} u\|_{U_K^2} \|P_{N_2} v\|_{U_K^2} \\
& \lesssim \|u\|_Y \|v\|_Y.
\end{aligned}$$

We finish the proof of this proposition. \square

In the proof of Proposition 4.2, we control many parts by $\|u\|_Z \|v\|_Z$ except $II_N^{(4)}$. Fortunately, we have the following lemma which can help us to control this case when $v = F[u_0]$.

Lemma 4.3. $2 \leq N \ll N_1 \sim N_2$.

$$\begin{aligned} \|\eta P_N \mathcal{B}(P_{N_1} v P_{N_2} F[P_{\sim N_2^2/\lambda} u_0])\|_{U_K^2} &\lesssim (\lambda N_1^3/N)^{-1/2} \|v\|_{U_K^2} \|u_0\|_{L^2}^2, \\ \|\eta P_N \mathcal{B}(P_{N_1}(F[P_{\sim N_1^2/\lambda} u_0]) P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0]))\|_{U_K^2} &\lesssim (\lambda^2 N_1^3/N)^{-1/2} \|u_0\|_{L^2}^4. \end{aligned}$$

Proof. By the duality, we only need to show

$$\begin{aligned} &\left| \int_{\mathbb{R}^2} P_{N_1} v P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0]) \partial_x P_N w \, dx dt \right| \\ &\lesssim (\lambda N_1^3/N)^{-1/2} \|v\|_{U_K^2} \|u_0\|_{L^2}^2 \|w\|_{V_K^2} \end{aligned}$$

and

$$\begin{aligned} &\left| \int_{\mathbb{R}^2} P_{N_1}(F[P_{\sim N_1^2/\lambda} u_0]) P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0]) \partial_x P_N w \, dx dt \right| \\ &\lesssim (\lambda^2 N_1^3/N)^{-1/2} \|u_0\|_{L^2}^4 \|w\|_{V_K^2} \end{aligned}$$

where v, w are supported on $[-1, 1] \times \mathbb{R}$. Let $L = cNN_1N_2$. Firstly, we control

$$\left| \int_{\mathbb{R}^2} P_{N_1} v P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0]) Q_{\leq L}^K \partial_x P_N w \, dx dt \right|.$$

Since for $|\xi_1| \sim N_1$, $|\xi_1| \sim N_2$, $|\xi| \sim N$, $\xi_1 + \xi_2 = \xi$, one has $|\xi_1^3 + \xi_2^3 - (\xi_1 + \xi_2)^3| \gtrsim L$. Typically, we need to control

$$\left| \int_{\mathbb{R}^2} P_{N_1} v Q_{>L}^K P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0]) Q_{\leq L}^K \partial_x P_N w \, dx dt \right|.$$

By transversal and high modulation estimates we can control this term by

$$\begin{aligned} &\|P_{N_1} v Q_{\leq L}^K \partial_x P_N w\|_{L_{t,x}^2} \|Q_{>L}^K P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0])\|_{L_{t,x}^2} \\ &\lesssim N_1^{-1} N \|v\|_{U_K^2} \|w\|_{U_K^2} L^{-1/2} \|P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0])\|_{V_K^2} \\ &\sim N_1^{-2} N^{1/2} \|v\|_{U_K^2} \|w\|_{U_K^2} \|P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0])\|_{V_K^2}. \end{aligned}$$

Recall that v is supported on $[-1, 1] \times \mathbb{R}$. Thus for any $p \geq 1$, one has $\|v\|_{L_{t,x}^2} \lesssim \|v\|_{L_t^\infty L_x^2} \lesssim \|v\|_{U^p}$. By the Hölder inequality and Strichartz estimates, we can also control this term by

$$\begin{aligned} &\|P_{N_1} v\|_{L_{t,x}^3} \|Q_{\leq L}^K \partial_x P_N w\|_{L_{t,x}^6} \|Q_{>L}^K P_{N_2}(F[P_{\sim N_2^2} u_0])\|_{L_{t,x}^2} \\ &\lesssim \|P_{N_1} v\|_{L_{t,x}^2}^{1/2} \|P_{N_1} v\|_{L_{t,x}^6}^{1/2} N \|P_N w\|_{U_K^6} L^{-1/2} \|P_{N_2}(F[P_{\sim N_2^2} u_0])\|_{V_K^2} \\ &\sim N_1^{-13/12} N^{1/2} \|v\|_{U_K^6} \|w\|_{U_K^6} \|P_{N_2}(F[P_{\sim N_2^2} u_0])\|_{V_K^2}. \end{aligned}$$

Then, by the interpolation [7] and Lemma 3.2, one has

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} P_{N_1} v P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]) Q_{\leq L}^K \partial_x P_N w \, dx dt \right| \\ & \lesssim N_1^{-2} N^{1/2} (\log N_1)^2 \|v\|_{V_K^2} \|w\|_{V_K^2} \|P_{N_2} (F[P_{\sim N_2^2} u_0])\|_{V_K^2} \\ & \lesssim \lambda^{-3/8} N_1^{-2} N^{1/2} (\log N_1)^2 \|v\|_{V_K^2} \|w\|_{V_K^2} \|P_{N_2} (F[P_{\sim N_2^2} u_0])\|_{V_K^2}. \end{aligned}$$

Similarly, one has

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} P_{N_1} (F[P_{\sim N_1^2/\lambda} u_0]) P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]) Q_{\leq L}^K \partial_x P_N w \, dx dt \right| \\ & \lesssim N_1^{-2} N^{1/2} (\log N_1)^2 \|P_{N_1} (F[P_{\sim N_1^2/\lambda} u_0])\|_{V_K^2} \|P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0])\|_{V_K^2} \|w\|_{V_K^2} \\ & \lesssim \lambda^{-3/4} N_1^{-2} N^{1/2} (\log N_1)^2 \|u_0\|_{L^2}^4 \|w\|_{V_K^2}. \end{aligned}$$

For the other part, by the Hölder inequality,

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} P_{N_1} v P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]) Q_{>L}^K \partial_x P_N w \, dx dt \right| \\ & \lesssim \|P_N (P_{N_1} v P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]))\|_{L_{t,x}^2} \|Q_{>L}^K \partial_x P_N w\|_{L_{t,x}^2} \\ & \lesssim NL^{-1/2} \|P_N (P_{N_1} v P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]))\|_{L_{t,x}^2} \|w\|_{V_K^2}. \end{aligned}$$

To conclude the proof of this lemma, we show

$$\begin{aligned} & \|P_N (P_{N_1} v P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]))\|_{L_{t,x}^2} \lesssim (\lambda N N_1)^{-1/2} \|v\|_{U_K^2} \|u_0\|_{L^2}^2, \\ & \|P_N (P_{N_1} (F[P_{\sim N_1^2/\lambda} u_0]) P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]))\|_{L_{t,x}^2} \lesssim (N N_1)^{-1/2} \|u_0\|_{L^2}^4. \end{aligned}$$

To show the first one, we only need to show

$$\|P_N (P_{N_1} K(t) v_0 P_{N_2} (F[P_{\sim N_2^2/\lambda} u_0]))\|_{L_{t,x}^2} \lesssim (\lambda N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2.$$

Let I_1, I_2 be intervals included in $[-CN_2^2/\lambda, -cN_2^2/\lambda] \cup [cN_2^2/\lambda, CN_2^2/\lambda]$ with length cN_2 . Also, we can assume that for any $\eta_1 \in I_1, \eta_2 \in I_2$ one has $|\eta_1 - \eta_2| \sim N_2$. It is equivalent to

$$\begin{aligned} & \left\| P_N \left(P_{N_1} K(t) v_0 \tilde{\eta} P_{N_2} \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right) \right\|_{L_{t,x}^2} \\ & \lesssim (\lambda N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2. \end{aligned}$$

There exists a interval J included in $[-CN_2, -cN_2] \cup [cN_2, CN_2]$ with length 1 such that one has $|\xi_2^3 + \lambda\eta_1^2 - \lambda\eta_2^2| \gtrsim N_2^2$ for any $\xi_2 \in [-CN_2, -cN_2] \cup [cN_2, CN_2] \setminus J := \hat{J}$, $\eta_1 \in I_1, \eta_2 \in I_2, \eta_1 + \eta_2 = \xi_2$. Then by Lemma 3.3 and

$$\|P_N (P_{N_1} K(t) v_0 P_{N_2} K(t) v_1)\|_{L_{t,x}^2} \lesssim (N N_1)^{-1/2} \|v_0\|_{L^2} \|v_1\|_{L^2}, \quad (4.2)$$

we can control this part. Thus we only need to show

$$\begin{aligned} & \left\| P_N \left(P_{N_1} K(t) v_0 \tilde{\eta} P_J \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right) \right\|_{L^2_{t,x}} \\ & \lesssim (\lambda N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2. \end{aligned}$$

By the orthogonality, we can assume that I_1, I_2 are intervals with length 1.

There exists a interval J_1 included in J with length at most $1/N_2$ such that one has $|\xi_2^3 + \lambda\eta_1^2 - \lambda\eta_2^2| \gtrsim N_2$ for any $\eta \in J \setminus J_1$, $\eta_1 \in I_1$, $\eta_2 \in I_2$, $\eta_1 + \eta_2 = \eta$. By Lemma 3.3 and the former argument we only need to show

$$\begin{aligned} & \left\| P_N \left(P_{N_1} K(t) v_0 \tilde{\eta} P_{J_1} \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right) \right\|_{L^2_{t,x}} \\ & \lesssim (\lambda N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2 \end{aligned}$$

where I_1, I_2 are intervals with length $1/N_2$.

There exists a interval J_2 included in J_1 with length $1/N_2^2$ such that one has $|\xi_2^3 + \lambda\eta_1^2 - \lambda\eta_2^2| \lesssim 1$ for any $\xi_2 \in J_2$, $\eta_1 \in I_1$, $\eta_2 \in I_2$, $\eta_1 + \eta_2 = \xi_2$. By Lemma 3.4 and following the former argument we only need to show

$$\begin{aligned} & \left\| P_N \left(P_{N_1} K(t) v_0 \tilde{\eta} P_{J_1 \setminus J_2} \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right) \right\|_{L^2_{t,x}} \\ & \lesssim (\lambda N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2 \end{aligned}$$

where I_1, I_2 are intervals with length $1/N_2$. Note that

$$\begin{aligned} & \mathcal{F}_x \left(P_N \left(P_{N_1} K(t) v_0 \tilde{\eta} P_{J_1 \setminus J_2} \mathcal{B}(S_\lambda(t) P_{I_1} u_0 \overline{S_\lambda(t) P_{I_2} u_0}) \right) \right) \\ & \sim \int_{\substack{\xi_1 + \xi_2 = \xi, \\ \eta_1 + \eta_2 = \xi_2}} M(\xi, \xi_1, \eta_1) \widehat{v_0}(\xi_1) \widehat{u_0}(\eta_1) \overline{\widehat{u_0}(-\eta_2)} \tilde{\eta} (e^{it(\xi_1^3 + \xi_2^3)} - e^{it(\xi_1^3 + \lambda\xi_2(\eta_2 - \eta_1))}) \\ & := \Lambda_1 + \Lambda_2. \end{aligned}$$

where $M(\xi, \xi_1, \eta_1) = \chi_{|\xi| \sim N, |\xi_1| \sim N_1, \xi_2 \in J_1 \setminus J_2, \eta_1 \in I_1, \eta_2 \in I_2} / (\xi_2^2 + \lambda\eta_1 - \lambda\eta_2)$. Note that $M(\xi, \xi_2, \eta_1) \lesssim N_2$. For Λ_1 , by (4.2) and the Plancherel identity we have

$$\begin{aligned} \|\Lambda_1\|_{L_t^2 L_\xi^2} & \lesssim (N N_1)^{-1/2} \|v_0\|_{L^2} \\ & \quad \cdot \left\| \int_{\eta_1 + \eta_2 = \xi_2} \frac{\widehat{u_0}(\eta_1) \overline{\widehat{u_0}(-\eta_2)}}{\xi_2^2 + \lambda\eta_1 - \lambda\eta_2} \chi_{\xi_2 \in J_1 \setminus J_2, \eta_1 \in I_1, \eta_2 \in I_2} \right\|_{L_{\xi_2}^2} \\ & \lesssim (N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2 \left\| \frac{\chi_{\xi_2 \in J_1 \setminus J_2, \eta_1 \in I_1, \eta_2 \in I_2}}{\xi_2^2 + 2\lambda\eta_1 - \lambda\xi_2} \right\|_{L_{\eta_1}^\infty L_{\xi_2}^2} \\ & \lesssim (N N_1)^{-1/2} \|v_0\|_{L^2} \|u_0\|_{L^2}^2. \end{aligned}$$

Let η_j^0 be the center of I_j , $j = 1, 2$. Note that

$$\begin{aligned}\partial_{\xi_1}(\xi_1^3 + \lambda(\xi - \xi_1)(\eta_2^0 - \eta_1^0)) &= 3\xi_1^2 - \lambda(\eta_2^0 - \eta_1^0) \\ &= 3\xi_1^2 - \xi_2^2 + \xi_2^2 - \lambda(\eta_2^0 - \eta_1^0) \\ &= 2\xi_1^2 + \xi(\xi_1 - \xi_2) + \xi_2^2 - \lambda(\eta_2^0 - \eta_1^0) \sim N_1^2.\end{aligned}$$

Thus by the Taylor expansion and the transversal estimate one has

$$\begin{aligned}&\|\Lambda_2\|_{L_t^2 L_\xi^2} \\ &\lesssim \sum_{k=0}^{\infty} \left\| \int_{\substack{\xi_1+\xi_2=\xi, \\ \eta_1+\eta_2=\xi_2}} M(\xi, \xi_1, \eta_1) \widehat{v}_0(\xi_1) \widehat{u}_0(\eta_1) \overline{\widehat{u}_0}(-\eta_2) e^{it(\xi_1^3 + \lambda \xi_2(\eta_2^0 - \eta_1^0))} \right. \\ &\quad \cdot \tilde{\eta} \frac{(i\lambda t \xi_2(\eta_2 - \eta_2^0 - \eta_1 + \eta_1^0))^k}{k!} \left. \right\|_{L_t^2 L_\xi^2} \\ &\lesssim \sum_{k=0}^{\infty} \frac{1}{k!} \left\| \int_{\substack{\xi_1+\xi_2=\xi, \\ \eta_1+\eta_2=\xi_2}} M(\xi, \xi_1, \eta_1) \widehat{v}_0(\xi_1) \widehat{u}_0(\eta_1) \overline{\widehat{u}_0}(-\eta_2) e^{it(\xi_1^3 + \lambda \xi_2(\eta_2^0 - \eta_1^0))} \right. \\ &\quad \cdot (\xi_2(\eta_2 - \eta_2^0 - \eta_1 + \eta_1^0))^k \left. \right\|_{L_t^2 L_\xi^2} \\ &\lesssim \sum_{k=0}^{\infty} \frac{1}{k!} N_1^{-1} \|v_0\|_{L^2} \left\| \int_{\eta_1+\eta_2=\xi_2} \frac{\widehat{u}_0(\eta_1) \overline{\widehat{u}_0}(-\eta_2)}{\xi_2^2 + \lambda \eta_1 - \lambda \eta_2} \right. \\ &\quad \cdot \chi_{\xi_2 \in J_1 \setminus J_2, \eta_1 \in I_1, \eta_2 \in I_2} (\xi_2(\eta_2 - \eta_2^0 - \eta_1 + \eta_1^0))^k \left. \right\|_{L_{\xi_2}^2} \\ &\lesssim \sum_{k=0}^{\infty} \frac{C^k}{k!} N_1^{-1} \|v_0\|_{L^2} \left\| \int_{\eta_1+\eta_2=\xi_2} \frac{|\widehat{u}_0(\eta_1) \overline{\widehat{u}_0}(-\eta_2)|}{\xi_2^2 + \lambda \eta_1 - \lambda \eta_2} \chi_{\xi_2 \in J_1 \setminus J_2, \eta_1 \in I_1, \eta_2 \in I_2} \right. \\ &\quad \cdot (e^{it(\zeta_1 + \zeta_2)^2} - e^{i\lambda t(\zeta_1 + \zeta_2)(\zeta_2 - \zeta_1)}) (e^{it(\eta_1 + \eta_2)^2} - e^{i\lambda t(\eta_1 + \eta_2)(\eta_2 - \eta_1)}) \left. \right\|_{L_{\xi_2}^2} \\ &\lesssim N_1^{-1} \|v_0\|_{L^2} \|u_0\|_{L^2}^2.\end{aligned}$$

Thus we conclude the proof of the first inequality.

For the second one, similar to the former argument, we reduce it to

$$\begin{aligned}&\left\| \int_{\zeta_1+\zeta_2+\eta_1+\eta_2=\xi} \frac{\tilde{\eta} \chi_{|\xi| \sim N} M_1(\zeta_1, \zeta_2) M_2(\eta_1, \eta_2) \widehat{u}_0(\zeta_1) \widehat{u}_0(\zeta_2) \widehat{u}_0(\eta_1) \widehat{u}_0(\eta_2)}{((\zeta_1 + \zeta_2)^2 + \lambda \zeta_1 - \lambda \zeta_2)((\eta_1 + \eta_2)^2 + \lambda \eta_1 - \lambda \eta_2)} \right. \\ &\quad \cdot (e^{it(\zeta_1 + \zeta_2)^2} - e^{i\lambda t(\zeta_1 + \zeta_2)(\zeta_2 - \zeta_1)}) (e^{it(\eta_1 + \eta_2)^2} - e^{i\lambda t(\eta_1 + \eta_2)(\eta_2 - \eta_1)}) \left. \right\|_{L_t^2 L_\xi^2} \\ &\lesssim (N N_1)^{-1/2} \|u_0\|_{L^2}^4\end{aligned}$$

where $M_1(\zeta_1, \zeta_2) = \chi_{\zeta_1 + \zeta_2 \in J, \zeta_1 \in I_1, \zeta_2 \in I_2}$, $M_2(\eta_1, \eta_2) = \chi_{\eta_1 + \eta_2 \in \tilde{J}, \eta_1 \in \tilde{I}_1, \eta_2 \in \tilde{I}_2}$, $I_1, I_2, \tilde{I}_1, \tilde{I}_2$ are intervals included in $[-CN_1^2, -cN_1^2] \cup [cN_1^2, CN_1^2]$, J, \tilde{J} are

intervals included in $[-CN_1, cN_1] \cup [cN_1, CN_1]$ and

$$N_1^{-1} \leq |(\zeta_1 + \zeta_2)^2 + \lambda\zeta_1 - \lambda\zeta_2|, |(\eta_1 + \eta_2)^2 + \lambda\eta_1 - \lambda\eta_2| \ll N_1. \quad (4.3)$$

Recall the former argument. Let $\zeta_1 + \zeta_2 = \zeta, \eta_1 + \eta_2 = \eta$, we only need to show

$$\begin{aligned} & \left\| \int_{\zeta+\eta=\xi} \frac{\tilde{\eta}\chi_{|\xi|\sim N} M_1(\zeta_1, \zeta_2)M_2(\eta_1, \eta_2)\widehat{u_0}(\zeta_1)\widehat{u_0}(\zeta_2)\widehat{u_0}(\eta_1)\widehat{u_0}(\eta_2)}{(\zeta^2 + \lambda\zeta_1 - \lambda\zeta_2)(\eta^2 + \lambda\eta_1 - \lambda\eta_2)} \right. \\ & \quad \cdot e^{i\lambda t(\zeta(\zeta_2-\zeta_1)+\eta(\eta_2-\eta_1))} \left. \right\|_{L_t^2 L_\xi^2} \\ & \lesssim (NN_1)^{-1/2} \|u_0\|_{L^2}^4 \end{aligned}$$

Let ζ_j^0, η_j^0 be the center of I_j, \tilde{I}_j respectively. Note that

$$\begin{aligned} & \partial_\zeta(\lambda\zeta(\zeta_2^0 - \zeta_1^0) + \lambda(\xi - \zeta)(\eta_2^0 - \eta_1^0)) \\ &= \lambda\zeta_2^0 - \lambda\zeta_1^0 - \lambda(\eta_2^0 - \eta_1^0) \\ &= \zeta^2 - \eta^2 + \lambda\zeta_2^0 - \lambda\zeta_1^0 - \zeta^2 + \eta^2 - \lambda(\eta_2^0 - \eta_1^0) \\ &= \xi(\zeta - \eta) - (\zeta^2 + \lambda\zeta_1^0 - \lambda\zeta_2^0) + (\eta^2 - \lambda\eta_2^0 + \eta_1^0). \end{aligned}$$

By (4.3), $|\xi| \sim N, |\zeta - \eta| \sim N_1$, we obtain

$$|\partial_\zeta(\lambda\zeta(\zeta_2^0 - \zeta_1^0) + \lambda(\xi - \zeta)(\eta_2^0 - \eta_1^0))| \sim NN_1.$$

Following the same argument for the first inequality, by the Taylor expansion and the transversal estimate, we conclude the proof. \square

Proposition 4.4. $\|\eta\mathcal{B}(uF[u_0])\|_Y \lesssim \lambda^{-1} \|u\|_Y \|u_0\|_{H^{-3/16}}^2$.

Proof. By the proof of Proposition 4.2 and Lemma 3.6, we only need to control the following term

$$\sum_{N_1 \sim N_2 \gg N \geq 2} \|\eta P_N \mathcal{B}(P_{N_1} u P_{N_2} F[P_{\sim N_2^2/\lambda} u_0])\|_{U_K^2}.$$

By Lemma 4.3, we can control it by

$$\begin{aligned} & \sum_{N_1 \sim N_2 \gg N \geq 2} N^{-3/4} N_1^{-3/2} N^{1/2} \lambda^{-1/2} \|P_{N_1} u\|_{U_K^2} \|P_{\sim N_2^2/\lambda} u_0\|_{L^2}^2 \\ & \lesssim \lambda^{-7/8} \|u\|_Y \|u_0\|_{H^{-3/16}}^2. \end{aligned}$$

Combining Corollary 3.6 and Proposition 4.2, we conclude the proof. \square

Proposition 4.5. $\|\eta\mathcal{B}(F[u_0]^2)\|_Y \lesssim \lambda^{-2} \|u_0\|_{H^{-3/16}}^4$.

Proof. Similar to the argument for Proposition 4.4, we only need to control the term

$$\sum_{N_1 \sim N_2 \gg N \geq 2} N^{-3/4} \|\eta P_N \mathcal{B}(P_{N_1}(F[P_{\sim N_1^2/\lambda} u_0]) P_{N_2}(F[P_{\sim N_2^2/\lambda} u_0]))\|_{U_K^2}.$$

By Lemma 4.3 we can control it by

$$\sum_{N_1 \sim N_2 \gg N \geq 2} N^{-3/4} N_1^{-3/2} N^{1/2} \lambda^{-1} \|P_{\sim N_2^2/\lambda} u_0\|_{L^2}^4 \lesssim \lambda^{-7/4} \|u_0\|_{H^{-3/16}}^4.$$

By Corollary 3.6 and Proposition 4.2, we conclude the proof. \square

Combining Lemmas 2.1–2.3, Corollary 3.6, Propositions 4.2, 4.4, and 4.5, one can construct the solution (u, w) of (2.2). Let $v = w + F[u_0]$. By rescaling we obtain the local solution $(u, v) \in C([0, T]; H^{-3/16} \times H^{-3/4})$ of initial system (S-KdV) with $(s_1, s_2) = (-3/16, -3/4)$. For general (s_1, s_2) , see the argument in [2]. One can use the method in this paper to manipulate the case $s_1, s_2 < 0$ with some modification. However there exists easier argument. For example if $s_2 > -3/4$, we do not need rescaling. One can use the norms

$$\|u\|_{X_\lambda^{s_1}} := \|N^{s_1} P_N u\|_{l_N^2 U_{S_\lambda}^2}, \quad \|v\|_{Y^{s_2}} := \|P_1 v\|_{L_x^2 L_t^\infty} + \|N^{s_2} P_N v\|_{l_{N \geq 2}^2 V_K^2}.$$

Then we can show the contraction for the iteration of the initial integral equation. If $s_2 = -3/4$, $s_1 > -3/16$, we can show $\|\eta \mathcal{B}(\partial_x |u|^2)\|_Y \lesssim \|u\|_{X_\lambda^{s_1}}$. The argument is also much easier than the case $(s_1, s_2) = (-3/16, -3/4)$. See also [11]. We omit the details.

Acknowledgments: The first author is supported in part by the NSFC, grant 12301116. The authors would like to thank Professors Boling Guo and Baoxiang Wang for their invaluable support and encouragement.

References

- [1] K. Appert and J. Vaclavik, *Dynamics of coupled solitons*, Phys. Fluids **20** (1977), no. 11, 1845–1849.
- [2] Y. Ban, J. Chen, and Y. Zhang, *Local well-posedness for Schrödinger-KdV system in $H^{s_1} \times H^{s_2}$* , submitted (2024).
- [3] D. Bekiranov, T. Ogawa, and G. Ponce, *Weak solvability and well-posedness of a coupled Schrödinger-Korteweg-de Vries equation for capillary-gravity wave interactions*, Proc. Amer. Math. Soc. **125** (1997), no. 10, 2907–2919.

- [4] A. Corcho and F. Linares, *Well-posedness for the Schrödinger-Korteweg-de Vries system*, Tran. Amer. Math. Soc. **359** (2007), no. 9, 4089–4106.
- [5] B. Guo and C. Miao, *Well-posedness of the Cauchy problem for the coupled system of the Schrödinger-KdV equations*, Acta Math. Sin. **15** (1999), 215–224.
- [6] Z. Guo and Y. Wang, *On the well-posedness of the Schrödinger-Korteweg-de Vries system*, J. Differ. Equ. **249** (2010), no. 10, 2500–2520.
- [7] M. Hadac, S. Herr, and H. Koch, *Well-posedness and scattering for the KP-II equation in a critical space*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **26** (2009), no. 3, 917–941.
- [8] C. E. Kenig, G. Ponce, and L. Vega, *Oscillatory integrals and regularity of dispersive equations*, Indiana Univ. Math. J. **40** (1991), no. 1, 33–69.
- [9] M. Tsutsumi, *Well-posedness of the Cauchy problem for a coupled Schrödinger-KdV equation*, Math. Sci. Appl **2** (1993), 513–528.
- [10] B. Wang, Z. Huo, C. Hao, and Z. Guo, *Harmonic Analysis Method for Nonlinear Evolution Equations, I*, World Scientific, 2011.
- [11] H. Wang and S. Cui, *The Cauchy problem for the Schrödinger-KdV system*, J. Differ. Equ. **250** (2011), no. 9, 3559–3583.

YINGZHE BAN: THE GRADUATE SCHOOL OF CHINA ACADEMY OF ENGINEERING PHYSICS,
BEIJING, 100088, P.R. CHINA

E-mail address: banyingzhe22@gscaep.ac.cn

JIE CHEN: SCHOOL OF SCIENCES, JIMEI UNIVERSITY, XIAMEN 361021, P.R. CHINA

E-mail address: jiechern@163.com

YING ZHANG: THE GRADUATE SCHOOL OF CHINA ACADEMY OF ENGINEERING PHYSICS,
BEIJING, 100088, P.R. CHINA

E-mail address: zhangying21@gscaep.ac.cn