EIGENVALUE ESTIMATES FOR THE POLY-LAPLACE OPERATOR ON LATTICE SUBGRAPHS

BOBO HUA AND RUOWEI LI

ABSTRACT. We introduce the discrete poly-Laplace operator on a subgraph with Dirichlet boundary condition. We obtain upper and lower bounds for the sum of the first k Dirichlet eigenvalues of the poly-Laplace operators on a finite subgraph of lattice graph \mathbb{Z}^d extending classical results of Li-Yau and Kröger. Moreover, we prove that the Dirichlet 2l-order poly-Laplace eigenvalues are at least as large as the squares of the Dirichlet *l*-order poly-Laplace eigenvalues.

CONTENTS

1. Introduction	1
2. Preliminaries	7
2.1. Basic notations and facts of graphs	7
2.2. The Dirichlet poly-Laplace operator.	8
2.3. The integer lattice \mathbb{Z}^d and the Fourier transform on \mathbb{Z}^d	10
3. Estimates of Dirichlet poly-Laplace eigenvalues	14
3.1. Upper bound	14
3.2. Lower bound	16
3.3. Comparison between eigenvalues of l and 2l-order poly-Laplace	18
4. Appendix	21
References	21

1. INTRODUCTION

Let Ω be a connected bounded domain with a smooth boundary in a *d*dimensional Euclidean space \mathbb{R}^d and let ν be the outward unit normal vector field of $\partial\Omega$. Denote by Δ the Laplace operator on \mathbb{R}^d . Solutions of $\Delta f = 0$ on Ω are the classical harmonic functions that describe the equilibrium state of an elastic homogeneous membrane. Solutions of $\Delta^2 f = 0$ are called biharmonic, and they model equilibria of homogeneous plates. Similarly, solutions of $\Delta^l f = 0$, where $l \in \mathbb{N}_+$, are called poly-harmonic. One naturally considers the eigenvalue problem

(1.1)
$$\begin{cases} (-\Delta)^l f = \lambda^l f & \text{in } \Omega, \\ f|_{\partial\Omega} = \frac{\partial f}{\partial\nu}|_{\partial\Omega} = \dots = \frac{\partial^{l-1} f}{\partial^{l-1}\nu}|_{\partial\Omega} = 0 \end{cases}$$

We can order the eigenvalues as

$$0 < \lambda_1^l(\Omega) \leqslant \lambda_2^l(\Omega) \leqslant \ldots \leqslant \lambda_k^l(\Omega) \leqslant \ldots$$

For the case l = 1, the eigenvalue problem (1.1) is called a fixed membrane problem which has been well-studied. Beginning with the work of Weyl in 1912 [34], he states that as $k \to \infty$,

$$\lambda_k^1(\Omega) \sim C_d(\frac{k}{V(\Omega)})^{\frac{2}{d}},$$

where $V(\Omega)$ is the volume of the domain Ω and $C_d = (2\pi)^2 V_d^{-\frac{2}{d}}$ with V_d being the volume of the unit ball in \mathbb{R}^d .

For a planar domain Ω that tiles \mathbb{R}^2 , Pólya [29] provided the lower bound estimate for d = 2 as

$$\lambda_k^1(\Omega) \ge C_d(\frac{k}{V(\Omega)})^{\frac{2}{d}}.$$

His proof applies to any dimension $d \ge 2$, and he conjectured that this estimate holds for all bounded domains in \mathbb{R}^d . A partial solution to this conjecture was obtained by Lieb [25]

$$\lambda_k^1(\Omega) \ge D_d(\frac{k}{V(\Omega)})^{\frac{2}{d}},$$

where the constant $D_d < C_d$ and is proportional to C_d . Later Li-Yau [24] and Berezin [3] derived the sharp estimate known as the Berezin-Li-Yau inequality

$$\frac{1}{k}\sum_{i=1}^{k}\lambda_{i}^{1}(\Omega) \geq \frac{d}{d+2}C_{d}(\frac{k}{V(\Omega)})^{\frac{2}{d}}.$$

Then Li and Yau's estimate on the sum of the eigenvalues was improved by Melas [26] whose result was further improved in [19, 36]. The upper bound estimate on the sum of the eigenvalues was proved by Kröger [21] in 1994, which includes an extra term depending on the geometry of Ω . Moreover in the same spirit, Kröger [20, 21] obtained upper and lower bounds for the sum of the first k Neumann eigenvalues. Finally, Strichartz [31] developed a general framework for obtaining upper and lower bounds for the sum of the first k Dirichlet and Neumann eigenvalues.

For $l \ge 2$, poly-Laplace operators have various applications in physics. For example, for l = 2 the eigenvalue problem (1.1) is called a clamped plate problem [14, Chapter 11]. Agmon [1] and Pleijel [28] established the Weyl's asymptotics for the eigenvalues of the clamped plate problem. And Birman and Solomyak [4, 5] generalized the Weyl's asymptotics to arbitrary l. Later, Levine and Protter [23] extended the Berezin-Li-Yau inequality for arbitrary order $l \ge 1$,

(1.2)
$$\frac{1}{k} \sum_{i=1}^{k} \lambda_{i}^{l}(\Omega) \ge \frac{d}{d+2l} C_{d}^{l} (\frac{k}{V(\Omega)})^{\frac{2l}{d}} + \frac{d}{d+2l} C_{d}^{l} (\frac{k}{V(\Omega)})^{\frac{2(l-1)}{d}},$$

which is sharp in the sense of Weyl's asymptotics. Later [6] gave a significant improvement of (1.2) by adding l lower-order terms than $k^{2l/d}$ to its righthand side, and it was further improved by [33, 35]. For the upper bound of the sum of the first k-th eigenvalues of the Dirichlet bi-Laplace operator, Cheng and Wei in [7] prove that there exists a constant $r_0 > 0$ such that for $k \ge V(\Omega)r_0^n$,

(1.3)
$$\frac{1}{k} \sum_{j=1}^{k} \lambda_j^2(\Omega) \leqslant \frac{1 + \frac{4(d+4)(d^2+2d+6)}{d+2} \frac{V(\Omega_{r_0})}{V(\Omega)}}{\left(1 - \frac{V(\Omega_{r_0})}{V(\Omega)}\right)^{\frac{d+4}{d}}} \frac{dC_d^2}{d+4} \left(\frac{k}{V(\Omega)}\right)^{\frac{4}{d}}.$$

where $\Omega_r := \{x \in \Omega : \operatorname{dist}(x, \partial \Omega) < r^{-1}\}$. The estimate is sharp in the sense of the Weyl's asymptotics since $V(\Omega_{r_0}) \to 0$ as $r_0 \to +\infty$. The eigenvalue problem (1.1) for arbitrary order case has recently received more attention. For instance, look at [17] for universal bounds for eigenvalues, [10, 22, 27] for Lieb-Thirring type inequalities and [11] for positiving preserving problems.

In recent years, increasing attention has been paid to the analysis of graphs. The graph Laplacian has been extensively studied in the literature, e.g. [12, 13, 16, 2]. Hirschler and Woess in [15] studied bi-Laplace equations for directed networks and Markov chains. And Duffin considered the difference equations of the poly-Laplace type [8]. Other works [9, 30, 18] mainly focus on the numerical framework of bi-Laplace. However, there are no references available studies on the eigenvalue problem of the discrete poly-Laplace operators. The motivation comes from studying the discretization of the corresponding physical models and providing theoretical results for future numerical calculations and algorithms. In this paper, we first introduce the definition of the poly-Laplace operator on a locally finite connected graph G with Dirichlet boundary condition, and moreover, we prove the upper and lower estimates for the sum of the first k Dirichlet eigenvalues of poly-Laplace operators on a lattice graph.

Recall the setting of graphs. A locally finite, simple, undirected and connected graph G = (V, E) consists of the set of vertices V and the set of edges E. In the following $x \in G$ means that $x \in V$, and let |G| be the number of vertices in G. Two vertices x, y are called neighbors, denoted by $x \sim y$, if there exists $e \in E$ connecting x and y. The degree of $x \in V$ is defined as $\deg(x) \coloneqq |\{y : y \sim x\}|$. The graph distance on G is defined by

$$d(x,y) := \inf\{n | x = z_0 \sim \cdots \sim z_n = y\}.$$

Define balls $B(x,r) = \{y \in V : d(x,y) \leq r\}$ on G. A subgraph $G_1 = (V_1, E_1)$ of G means $V_1 \subset V$ and $E_1 \subset E$. For a subgraph Ω of G, recall the notion of the boundary $\delta\Omega$

$$\delta\Omega \coloneqq \{y \in V \setminus \Omega : \exists x \in \Omega, \text{s.t. } y \sim x\}.$$

Denote $\delta\Omega$ by $\delta_1\Omega$, and for a positive integer *l* define the *l*-th layer boundary as

$$\delta_l \Omega \coloneqq \delta(\Omega \cup \delta_1 \Omega \cup \ldots \cup \delta_{l-1} \Omega).$$

Recall that the graph Laplace operator is defined for any $f: V \to \mathbb{C}$ as

$$\Delta f(x) = \sum_{y \sim x} (f(y) - f(x))$$

For a positive integer l, the poly-Laplace operator can be regarded as iterations of the Laplace operator defined inductively

$$\Delta^l = \underbrace{\Delta \circ \Delta \circ \cdots \circ \Delta}_{l-\mathrm{th}}.$$

For a finite graph G = (V, E) with $V = \{x_1, \dots, x_n\}$, the Laplace matrix is defined as $\Delta = A - D$, where $D = \text{diag}(\text{deg}(x_1), \dots, \text{deg}(x_n))$ is the degree matrix and A is the adjacency matrix whose element A_{ij} is one when there is an edge from vertex x_i to vertex x_j , and zero when there is no edge. For any $f: V \to \mathbb{C}$,

$$(-\Delta)^{l} f(x) = (D - A)^{l} f(x) = \sum_{m=0}^{l} \binom{l}{m} D^{l-m} (-1)^{m} A^{m} f(x)$$
$$=: \sum_{y \in B(x,l)} a_{xy}^{l} f(y),$$

where

(1.4)
$$a_{xy}^{l} \coloneqq \sum_{m=0}^{l} \binom{l}{m} (-1)^{m} (\deg(y))^{l-m} p_{m}(x,y)$$

and $p_m(x,y) = \# \{ \text{paths } x = z_0 \sim \cdots \sim z_m = y \}$ is the number of paths of length *m* from vertex *x* to vertex *y*.

Given a finite subgraph $\Omega \subseteq G$, we introduce the poly-Laplace operator with Dirichlet boundary, denoted by $\Delta_{\Omega}^{l,\mathcal{D}}$, which is defined as

$$\Delta_{\Omega}^{l,\mathcal{D}} f \coloneqq \Delta^l f^*|_{\Omega}.$$

Here $f: \Omega \to \mathbb{C}$, and $f^*: G \to \mathbb{C}$ is the zero extension of u. One readily sees that $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$ is a self-adjoint and positive definite operator on Ω . The Dirichlet poly-Laplace eigenvalue problem is given by

(1.5)
$$(-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f(x) = \lambda^l f(x) \text{ in } \Omega.$$

There are $|\Omega|$ corresponding real eigenvalues:

$$0 < \lambda_1^l(\Omega) \leqslant \lambda_2^l(\Omega) \leqslant \ldots \leqslant \lambda_{|\Omega|}^l(\Omega).$$

See subsection 2.2 for more properties of this operator.

Consider the *d*-dimensional integer lattice \mathbb{Z}^d as a graph. We study the Dirichlet poly-Laplace problem on a finite subgraph Ω of \mathbb{Z}^d using the Fourier transform. In 2022, Bauer and Lippner [2] first studied the graph Laplacian eigenvalues with Dirichlet boundary condition on Ω of \mathbb{Z}^d . Later Wang [32] introduced the Dirichlet problems of graph fractional Laplacian and developed a framework for estimating the Dirichlet eigenvalues, they derived Laplacian. For the average of the first k Dirichlet eigenvalues, they derived

the upper estimate in the spirit of Kröger [21] and the lower estimate in the spirit of Li-Yau [24]. In this paper, we follow the proof strategies in [21, 24, 2, 32] to derive the upper and lower bounds for the Dirichlet eigenvalues of the discrete poly-Laplace operator.

The following are the main results of this paper.

Theorem 1.1. For a positive integer l, consider the Dirichlet poly-Laplace eigenvalue problem (1.5) on a finite subgraph Ω of \mathbb{Z}^d . Then

(a) for all $1 \leq k \leq \min\{1, \frac{V_d}{2^d}\} |\Omega|$, we have the upper bound estimate

$$\frac{1}{k}\sum_{j=1}^{k}\lambda_{j}^{l}(\Omega) \leqslant (2\pi)^{2l}\frac{d}{d+2l}\left(\frac{k}{V_{d}|\Omega|}\right)^{\frac{2l}{d}} + \frac{|\partial^{l}\Omega|}{|\Omega|}.$$

For $1 \leq k \leq \min\{1, \frac{V_d}{2^{d+1}}\}|\Omega|$, there holds

$$\lambda_{k+1}^{l}(\Omega) \leqslant (2\pi)^{2l} \frac{d \cdot 2^{\frac{d+2l}{d}}}{d+2l} \left(\frac{k}{V_d|\Omega|}\right)^{\frac{2l}{d}} + 2\frac{|\partial^{l}\Omega|}{|\Omega|}.$$

Recall that V_d is the volume of the unit ball in \mathbb{R}^d and

$$\begin{aligned} |\partial^{l}\Omega| &= \sum_{x \in \delta\Omega \cup \ldots \cup \delta_{l}\Omega} \sum_{y \in B(x,l) \cap \Omega} |a_{xy}^{l}| \\ &\leqslant 4^{l} d^{l} \left(|\delta\Omega| + \ldots + |\delta_{l}\Omega| \right), \end{aligned}$$

where a_{xy}^l is defined by (1.4).

(b) For all $1 \leq k \leq \min\{1, (\frac{\sqrt{6}}{2\pi})^d V_d\} |\Omega|$, the lower bound estimate is given as

$$\begin{split} \lambda_k^l(\Omega) &\ge \frac{1}{k} \sum_{j=1}^k \lambda_j^l(\Omega) \\ &\ge \sum_{m=0}^l \left(\begin{array}{c}l\\m\end{array}\right) \left(-\frac{1}{12}\right)^m (2\pi)^{2(l+m)} \frac{d}{d+2(l+m)} (\frac{k}{V_d|\Omega|})^{\frac{2(l+m)}{d}} > 0. \end{split}$$

Remark 1.2. (1) Compared with the continuous upper bound estimates (1.3) and lower bound estimates (1.2), the terms $(2\pi)^{2l} \frac{d}{d+2l} \left(\frac{k}{V_d|\Omega|}\right)^{\frac{2l}{d}}$ are the same and consistent with the Weyl's asymptotics. Moreover, the upper bound estimates (1.3) also contain a geometric quantity $V(\Omega_{r_0})$ that depends on the domain boundary.

(2) The boundary term $|\partial^l \Omega|$ can be bounded by $4^l d^l (|\delta \Omega| + \ldots + |\delta_l \Omega|)$. In particular, for l = 1 and l = 2, we can bound $|\partial^1 \Omega|$ and $|\partial^2 \Omega|$ by (2.7) and (2.8) respectively.

The ideas behind the proof are as follows. The upper bound estimate is based on Lemma 3.1 (was proved by the Rayleigh-Ritz formula in [2]), which gives an estimate (3.1) for eigenvalues of general self-adjoint and positive semidefinite operators. Let $g(x) = h_z(x) = e^{i\langle x, z \rangle}$ and integrate (3.1) in a measurable subset B of $[-\pi,\pi]^d$. By the properties of the Fourier transform we can prove Lemma 3.2. Finally, the upper bound estimate (a) follows from the proper choice for set B.

For the lower bound estimate, the key is to prove Lemma 3.3 which is an adaption of Li and Yau's method [24]. To prove the lemma, for a realvalued function F with $0 \leq F \leq M$ and $\int_{[-\pi,\pi]^d} F(z) dz \geq K$, first construct a radially symmetric function $\varphi(z): [-\pi,\pi]^d \to \mathbb{C}$ which satisfies $0 \leq \varphi(z) \leq$ $(\Phi(z))^l$ and

$$\int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz \ge \int_{[-\pi,\pi]^d} \varphi(z) F(z) dz.$$

Note that $\widetilde{F} = M \mathbb{1}_{B_R}$ minimizes the integral $\int_{[-\pi,\pi]^d} \varphi(z) F(z) dz$, and gives a lower bound of $\int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz$. Choose $F(z) = \sum_{j=1}^k |\langle \phi_j^l, h_z \rangle_{\Omega}|^2$, where $\{\phi_i^l\}_{1 \leq i \leq |\Omega|}$ are corresponding eigenfunctions of $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$, and (b) is derived from L. derived from Lemma 3.3.

Notably, on a graph G without the boundary, the eigenvalues of $(-\Delta)^l$ are actually the *l*-th power of the corresponding eigenvalues of the Laplace operator $-\Delta$. However, this relationship generally does not hold for the Dirichlet eigenvalue problems. The following theorem shows that the Dirichlet eigenvalues of lower-order poly-Laplace can provide a rough lower bound for the higher-order poly-Laplace. Therefore, it is necessary to establish a more precise estimate for the poly-Laplace eigenvalue, as stated in Theorem 1.1.

Theorem 1.3. For a finite subgraph Ω of G = (V, E) and a positive integer *l*, let λ_k^l , λ_k^{2l} be the k-th eigenvalues of the Dirichlet poly-Laplace $\Delta_{\Omega}^{l,\mathcal{D}}$ and $\Delta_{\Omega}^{2l,\mathcal{D}}$ respectively. Then for $1 \leq k \leq |\Omega|$,

$$\left(\lambda_k^l\right)^2 \leqslant \lambda_k^{2l}.$$

Moreover, if Ω is a finite subgraph of \mathbb{Z}^d , then

$$\left(\lambda_k^l\right)^2 < \lambda_k^{2l}$$

Remark 1.4. (1) We obtain the same estimate $(\lambda_k^l)^2 \leq \lambda_k^{2l}$ for the continuous settings, see Theorem 3.6.

(2) The strict inequality follows from Lemma 3.5, which states that there is no ℓ^2 -eigenfunction of poly-Laplace on \mathbb{Z}^d , while the strictness is usually not true on general graphs. For example, on the subgraph $\Omega = \{v_1, v_2\}$ of the 3-cycle $C_3 = \{v_1, v_2, v_3\}$, we have $(\lambda_2^1)^2 = (-3)^2 = 9 = \lambda_2^2$. (3) On a path graph $[0, n] \subseteq \mathbb{Z}$, we have $\frac{(\lambda_k^1)^2}{\lambda_k^2} \to c_k < 1$ as $n \to +\infty$, see

numerical experiments in Appendix.

For an infinite graph G = (V, E), the k-th eigenvalue of the poly-Laplace on G is defined as

$$\lambda_k^l(G) = \inf_{\substack{M \subseteq C_0(V) \\ \dim M = k}} \max_{\substack{f \in M \\ \|f\|_2 = 1}} \langle (-\Delta)^l f, f \rangle_G.$$

And it can be estimated by the exhaustion trick, see Definition 2.6.

Theorem 1.5. Let $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$ be an exhaustion of infinite graph G = (V, E) with bounded degree $\sup_{x \in V} \deg x < +\infty$, then

$$\lambda_k^l(G) = \lim_{i \to +\infty} \lambda_k^l(W_i).$$

In particular, for k = 1 we have

$$(\lambda_1^1(G))^l = \lambda_1^l(G) = \lim_{i \to +\infty} \lambda_1^l(W_i).$$

Remark 1.6. For k = 1, the bottom of the spectrum of poly-Laplace $(-\Delta)^{l}$ is the *l*-th power of the bottom of the spectrum of Laplace operator $-\Delta$.

The paper is organized as follows. In next section, we first recall the basic setting of graphs, and introduce the discrete poly-Laplace operator and some important lemmas. In Section 3, we prove Theorem 1.1 and Theorem 1.3.

2. Preliminaries

2.1. Basic notations and facts of graphs. We recall the setting of graphs. A locally finite, simple, undirected and connected graph G = (V, E) consists of the set of vertices V and the set of edges E. We denote the space of functions on G by $C(G) := \{f : V \to \mathbb{C}\}$, and let $C_0(G)$ be the set of all functions on G with finite support. For a finite graph G = (V, E), then C(G) is a finite dimensional complex vector space of functions defined on G, and equip it with the natural Hermitian scalar product

$$\langle f,g \rangle_G = \sum_{x \in G} f(x)g(x), \, \forall f,g \in C(G).$$

For $f \in C(G)$ and $p \ge 1$, define the ℓ^p -norm of f as

$$||f||_{\ell^p(G)} = \left(\sum_{x \in G} |f(x)|^p\right)^{\frac{1}{p}}.$$

And let

$$\ell^p(G) \coloneqq \left\{ f \in C(G) : \|f\|_{\ell^p(G)} < +\infty \right\}$$

be the space of ℓ^p -summable functions on G. For any edge $(x, y) \in E$ define the gradient operator as

$$\nabla_{xy}f = f(y) - f(x), \,\forall f \in C(G).$$

The Laplace operator on G is defined as

$$\Delta_G f(x) = \sum_{y \sim x} \nabla_{xy} f, \, \forall f \in C(G),$$

which is self-adjoint with respect to the above scalar product. And we write Δ instead of Δ_G when it causes no confusion.

For subgraphs $G_1, G_2 \subseteq G$, define the edge set of connecting G_1 and G_2 as

$$E(G_1,G_2) \coloneqq \left\{ (x,y) \in E : x \in G_1, y \in G_2 \right\}.$$

Let Ω be a finite subgraph of G. Give the notions of the boundary $\delta\Omega$ and the closure $\overline{\Omega}$ of the subgraph Ω as

$$\begin{split} \delta\Omega \coloneqq \left\{ y \in V \backslash \Omega : \exists x \in \Omega, \text{s.t. } y \sim x \right\}, \\ \bar{\Omega} \coloneqq \Omega \cup \delta\Omega. \end{split}$$

Denote $\delta\Omega$ by $\delta_1\Omega$, and for a positive integer *l* define the *l*-th layer boundary as

$$\delta_l \Omega \coloneqq \delta(\Omega \cup \delta_1 \Omega \cup \ldots \cup \delta_{l-1} \Omega).$$

For $f \in C(\Omega)$, the Dirichlet Laplace operator is defined as

$$\Delta_{\Omega}^{\mathcal{D}} f \coloneqq \Delta f^*|_{\Omega},$$

where $f^* \in C(G)$ obtained by extending functions to be 0 on Ω^c . Note that $\Delta_{\Omega}^{\mathcal{D}}$ is self-adjoint with respect to the Hermitian scalar product [2, Proposition 2.2]. For $f \in C(\overline{\Omega})$ and $x \in \delta\Omega$, define the discrete outward normal derivative of f at x as

$$\frac{\partial f}{\partial n}(x) \coloneqq \sum_{x \sim y \in \Omega} \nabla_{yx} f.$$

2.2. The Dirichlet poly-Laplace operator. In this subsection, we introduce the Dirichlet poly-Laplace operator and give the Dirichlet poly-Laplace eigenvalue problem.

Definition 2.1. Let Ω be a finite subgraph of G = (V, E), then for all functions $f \in C(\Omega)$ define the Dirichlet poly-Laplace operator $\Delta_{\Omega}^{l,\mathcal{D}} : C(\Omega) \to C(\Omega)$ as

$$\Delta_{\Omega}^{l,\mathcal{D}} f \coloneqq \Delta^l f^*|_{\Omega},$$

where $f^* \in C(G)$ obtained by extending functions to be 0 on Ω^c . In particular, we know $\Delta_{\Omega}^{1,\mathcal{D}}$ is the Dirichlet Laplace operator $\Delta_{\Omega}^{\mathcal{D}}$.

Lemma 2.2. Let Ω be a finite subgraph of G = (V, E), then $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$ is self-adjoint and positive definite.

Proof. For all functions $f, g \in C(\Omega)$, we have

$$\left\langle (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f, g \right\rangle_{\Omega} = \left\langle (-\Delta_G)^l f^* |_{\Omega}, g \right\rangle_{\Omega} = \left\langle (-\Delta_G)^l f^*, g^* \right\rangle_V.$$

Since Δ_G is self-adjoint, then

$$\left\langle \left(-\Delta_G\right)^l f^*, g^*\right\rangle_V = \left\langle f^*, \left(-\Delta_G\right)^l g^*\right\rangle_V = \left\langle f, \left(-1\right)^l \Delta_\Omega^{l,\mathcal{D}} g\right\rangle_\Omega.$$

And the positive definiteness follows from the positive definiteness of $-\Delta_G$.

In this paper, we consider the Dirichlet poly-Laplace eigenvalue problem on a finite subgraph Ω

$$(-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f = \lambda^l f, \text{ in } \Omega.$$

Since $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$ is self-adjoint, positive definite and a finite dimensional operator, then there exist $|\Omega|$ eigenvalues (with multiplicities), all of which are positive real numbers. We label them in increasing order

$$0 < \lambda_1^l(\Omega) \leqslant \lambda_2^l(\Omega) \leqslant \ldots \leqslant \lambda_{|\Omega|}^l(\Omega).$$

Denote the corresponding eigenfunctions by $\{\phi_i^l\}_{1 \le i \le |\Omega|}$ respectively.

Lemma 2.3. (Domain monotonicity) For finite subgraphs Ω_1 and Ω_2 of G = (V, E), if $\Omega_1 \subseteq \Omega_2$, then for all $1 \leq k \leq |\Omega_1|$

$$\lambda_k^l(\Omega_1) \ge \lambda_k^l(\Omega_2).$$

Proof. Since the poly-Laplace $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$ can be written in the min-max form

(2.1)
$$\lambda_k^l(\Omega) = \min_{\substack{M \subseteq C(\Omega) \\ \dim M = k}} \max_{\substack{f \in M \\ \|f\|_{\ell^2(\Omega)} = 1}} \langle (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f, f \rangle_{\Omega}.$$

Definition 2.4. For an infinite-dimensional Hilbert space \mathcal{H} , the spectrum of a bounded linear operator $T : \mathcal{H} \to \mathcal{H}$ is defined as

 $\sigma(T) \coloneqq \{\lambda \in \mathbb{C} : \lambda I - T \text{ not invertible}\}.$

And $\inf \sigma(T)$ is the bottom of the spectrum of T.

For an infinite graph G = (V, E), the k-th eigenvalue of the poly-Laplace on G is defined as

(2.2)
$$\lambda_k^l(G) = \inf_{\substack{M \subseteq C_0(V) \\ \dim M = k}} \max_{\substack{f \in M \\ \|f\|_{\ell^2(G)} = 1}} \langle (-\Delta)^l f, f \rangle_G.$$

Fact 2.5. On an infinite graph G = (V, E) with bounded degree $\sup_{x \in V} \deg x < +\infty$, the poly-Laplace $(-\Delta)^l$ is a bounded linear operator from $\ell^2(G)$ to $\ell^2(G)$, then (1) $\lambda_k^l(G) \subseteq \sigma((-\Delta)^l)$, (2) $\sigma((-\Delta)^l) = (\sigma(-\Delta))^l$, (3) $\lambda_1^l(G) = \inf \sigma((-\Delta)^l) = (\inf \sigma(-\Delta))^l = (\lambda_1^1(G))^l$. **Definition 2.6.** Let G = (V, E) be an infinite graph. A sequence of subsets of vertices $\mathcal{W} = \{W_i\}_{i=1}^{\infty}$ is called an exhaustion of G, if it satisfies

- (1) $W_1 \subseteq W_2 \subseteq \cdots \subseteq W_i \subseteq \cdots \subseteq V$,
- (2) $|W_i| < \infty$, for all $i = 1, 2, \cdots$,
- (3) $V = \bigcup_{i=1}^{\infty} W_i$.

By the exhaustion method we can estimate the eigenvalues of the poly-Laplace on G as Theorem 1.5.

Proof of Theorem 1.5. By the domain monotonicity $\lambda_k^l(W_i) \ge \lambda_k^l(W_{i+1})$, then the limit exists follows from the definition (2.1) and (2.2).

2.3. The integer lattice \mathbb{Z}^d and the Fourier transform on \mathbb{Z}^d . Let \mathbb{Z}^d be the *d*-dimensional integer lattice, and choose $S = \{e_1, \ldots, e_d, -e_1, \ldots, -e_d\}$ as the generating set. Here $e_i \in \mathbb{Z}^d$ is the vector whose *i*-th component is 1 and the rest are 0. Thus consider \mathbb{Z}^d as the Cayley graph generated by S. For any $s \in S$, the gradient operator $\nabla s : C(\mathbb{Z}^d) \to C(\mathbb{Z}^d)$ can be written for a function $f \in C(\mathbb{Z}^d)$ as

$$\nabla sf(x) = f(x+s) - f(x).$$

Note that the gradient operator on \mathbb{Z}^d satisfies the integration by parts and commutative law as follows.

Fact 2.7. (Integration by parts) For all $1 \leq i \leq d$ and $f, g \in C_0(\mathbb{Z}^d)$, we have

 $\langle \nabla_{e_i} f, g \rangle_{\mathbb{Z}^d} = \langle f, \nabla_{-e_i} g \rangle_{\mathbb{Z}^d}.$ (Commutative law) For all $1 \leq i, j \leq d$ and $f \in C(\mathbb{Z}^d)$, we have $\nabla_{e_i} \nabla_{e_i} f = \nabla_{e_i} \nabla_{e_i} f.$

Next, we introduce the Fourier transform on \mathbb{Z}^d . For $z \in [-\pi, \pi]^d$, we define $h_z(x) : \mathbb{Z}^d \to \mathbb{C}$ as $h_z(x) = e^{i\langle x, z \rangle}$, where $\langle x, z \rangle = \sum_{i=1}^d x_i z_i$. Thus, for all $f \in \ell^1(\mathbb{Z}^d)$, the Fourier transform on \mathbb{Z}^d is defined by

$$\mathscr{F}: \quad \mathbb{C}^{\mathbb{Z}^d} \to \quad \mathbb{C}^{[-\pi,\pi]}$$
$$f \quad \mapsto \quad \widehat{f},$$

and

$$\widehat{f}(z) = \sum_{x \in \mathbb{Z}^d} e^{-i\langle x, z \rangle} f(x) = \langle f, h_z \rangle_{\mathbb{Z}^d},$$
$$f(x) = (2\pi)^d \int_{[-\pi,\pi]^d} e^{i\langle x, z \rangle} \widehat{f}(z) dz.$$

Let Ω be a finite subgraph of \mathbb{Z}^d , and write $\Phi(z) = \sum_{i=1}^d (2 - 2\cos z_i)$ for $z \in [-\pi, \pi]^d$. Then we prove several critical lemmas as follows.

Lemma 2.8. Let *l* be a positive integer, then for all $f \in C_0(\mathbb{Z}^d)$ we have

$$\widehat{(-\Delta)^l}f(z) = (\Phi(z))^l \widehat{f}(z), \, \forall z \in [-\pi,\pi]^d.$$

10

Proof. By the definition of Fourier transform, for all $z \in [-\pi, \pi]^d$, we have

$$\begin{split} (-\Delta)^{l}f(z) &= \sum_{x \in \mathbb{Z}^{d}} e^{-i\langle x, z \rangle} (-\Delta)^{l} f(x) \\ &= \sum_{x \in \mathbb{Z}^{d}} (-\Delta) e^{-i\langle x, z \rangle} (-\Delta)^{l-1} f(x) \\ &= \Phi(z) \sum_{x \in \mathbb{Z}^{d}} e^{-i\langle x, z \rangle} (-\Delta)^{l-1} f(x) \\ &= (\Phi(z))^{l} \sum_{x \in \mathbb{Z}^{d}} e^{-i\langle x, z \rangle} f(x) = (\Phi(z))^{l} \widehat{f}(z). \end{split}$$

Lemma 2.9. Let Ω be a finite subgraph of \mathbb{Z}^d . For all $f \in C(\Omega)$, the following equalities holds:

(2.3)
$$\langle f, f \rangle_{\Omega} = \frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} |\langle f, h_z \rangle_{\Omega}|^2 dz,$$

(2.4)
$$\langle f, (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f \rangle_{\Omega} = \frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} (\Phi(z))^l |\langle f, h_z \rangle_{\Omega}|^2 dz.$$

Proof. The result of (2.3) follows from the Plancherel formula. For (2.4), letting $1 \le i_1 \le d$, we compute

$$\begin{split} &\frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} (2 - 2\cos z_{i_1}) |\langle f, h_z \rangle_{\Omega}|^2 dz \\ &= \frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} |e^{-i\langle e_{i_1},z \rangle} - 1|^2 |\langle f, h_z \rangle_{\Omega}|^2 dz \\ &= \frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} |\langle f, (e^{-i\langle e_{i_1},z \rangle} - 1) h_z \rangle_{\Omega}|^2 dz \\ &= \frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} |\langle f^*, \nabla_{-e_{i_1}} h_z \rangle_{\mathbb{Z}^d}|^2 dz \\ &= \frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} |\langle \nabla_{e_{i_1}} f^*, h_z \rangle_{\mathbb{Z}^d}|^2 dz \\ &= \langle \nabla_{e_{i_1}} f^*, \nabla_{e_{i_1}} f^* \rangle_{\mathbb{Z}^d} = \langle f^*, \nabla_{-e_{i_1}} \nabla_{e_{i_1}} f^* \rangle_{\mathbb{Z}^d} \end{split}$$

which follows from (2.3) and the integration by parts, see Fact 2.7. Then by summing over all coordinate directions for $1 \leq i_1, \ldots, i_l \leq d$ we get

$$\frac{1}{(2\pi)^d} \int_{[-\pi,\pi]^d} (\Phi(z))^l |\langle f, h_z \rangle_{\Omega}|^2 dz = \sum_{i_1,\dots,i_l=1}^d \langle f^*, \nabla_{-e_{i_l}} \nabla_{e_{i_l}} \cdots \nabla_{-e_{i_1}} \nabla_{e_{i_1}} f^* \rangle_{\mathbb{Z}^d}$$
$$= \langle f^*, (-\Delta)^l f^* \rangle_{\mathbb{Z}^d} = \langle f, (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f \rangle_{\Omega}.$$

Lemma 2.10. Let Ω be a finite subgraph of \mathbb{Z}^d , then the following holds:

(2.5)
$$\langle h_z, h_z \rangle_{\Omega} = |\Omega|.$$

(2.6)
$$|\langle h_z, (-1)^l \Delta_{\Omega}^{l,\mathcal{D}}(h_z|_{\Omega}) \rangle_{\Omega}| \leq (\Phi(z))^l |\Omega| + |\partial^l \Omega|_{\Omega}$$

where

$$\begin{aligned} |\partial^{l}\Omega| &= \sum_{x \in \delta\Omega \cup \ldots \cup \delta_{l}\Omega} \sum_{y \in B(x,l) \cap \Omega} |a_{xy}^{l}| \\ &\leqslant 4^{l} d^{l} \left(|\delta\Omega| + \ldots |\delta_{l}\Omega| \right), \end{aligned}$$

and a_{xy}^l is defined by (1.4). In particular, for p = 1 and p = 2, the terms $|\partial^1 \Omega|$ and $|\partial^2 \Omega|$ can be bounded by (2.7) and (2.8) respectively.

Proof. The first result (2.5) follows from

$$\langle h_z, h_z \rangle_{\Omega} = \sum_{x \in \Omega} h_z(x) \overline{h_z}(x) = \sum_{x \in \Omega} 1 = |\Omega|.$$

Note that

$$|\langle h_z, (-1)^l \Delta_{\Omega}^{l,\mathcal{D}}(h_z|_{\Omega}) \rangle_{\Omega}| = |\langle h_z^*, (-\Delta)^l (h_z|_{\Omega})^* \rangle_{\mathbb{Z}^d}|.$$

Since

$$\begin{aligned} |\langle h_z, (-\Delta)^l (h_z|_{\Omega})^* \rangle_{\mathbb{Z}^d}| &= |(-\Delta)^l (h_z|_{\Omega})^* (z)| \\ &= |(\Phi(z))^l (\widehat{h_z|_{\Omega}})^* (z)| \\ &= (\Phi(z))^l |\Omega|, \end{aligned}$$

and by the definition (1.4)

$$\begin{aligned} |\langle h_z, (-\Delta)^l (h_z|_{\Omega})^* \rangle_{\Omega^c}| &\leq \sum_{x \in \Omega^c} |e^{-i\langle x, z \rangle} (-\Delta)^l (h_z|_{\Omega})^* (x)| \\ &\leq \sum_{x \in \delta\Omega \cup \dots \cup \delta_l \Omega} |\sum_{y \in B(x,l) \cap \Omega} a_{xy}^l h_z(y)| \\ &\leq \sum_{x \in \delta\Omega \cup \dots \cup \delta_l \Omega} \sum_{y \in B(x,l) \cap \Omega} |a_{xy}^l| \eqqcolon |\partial^l \Omega|. \end{aligned}$$

Then we can estimate the left-hand side of (2.6) as

$$\begin{aligned} |\langle h_z, (-1)^l \Delta_{\Omega}^{l,\mathcal{D}}(h_z|_{\Omega}) \rangle_{\Omega}| &\leq |\langle h_z, (-\Delta)^l (h_z|_{\Omega})^* \rangle_{\mathbb{Z}^d}| + |\langle h_z, (-\Delta)^l (h_z|_{\Omega})^* \rangle_{\Omega^c}| \\ &\leq (\Phi(z))^l |\Omega| + |\partial^l \Omega|, \end{aligned}$$

Similar to (1.4), we define

$$(D+A)^l f(x) = \sum_{m=0}^l \begin{pmatrix} l \\ m \end{pmatrix} D^{l-m} A^m f(x) =: \sum_{y \in B(x,l)} b_{xy}^l f(y),$$

where D is the degree matrix and A is the adjacency matrix. Then

$$\sum_{y \in B(x,l)} |a_{xy}^l| \leqslant \sum_{y \in B(x,l)} b_{xy}^l \leqslant \sum_y \left((D+A)^l \right)_{xy} \leqslant 4^l d^l,$$

Hence we can get a upper estimate

$$|\partial^{l}\Omega| = \sum_{x \in \delta\Omega \cup \ldots \cup \delta_{l}\Omega} \sum_{y \in B(x,l) \cap \Omega} |a_{xy}^{l}| \leq 4^{l} d^{l} \left(|\delta\Omega| + \ldots |\delta_{l}\Omega| \right).$$

In particular, for l = 1 we have

(2.7)
$$|\langle h_z, -\Delta_{\Omega}^{\mathcal{D}}(h_z|_{\Omega})\rangle_{\Omega}| = \sum_{(x,y)\in E(\Omega)} |\nabla_{xy}h_z|^2 + \sum_{(x,y)\in E(\Omega,\delta\Omega)} |h_z(x)|^2 \\ \leqslant \Phi(z)|\Omega| + |E(\Omega,\delta\Omega)|.$$

And for l = 2 we have

$$\sum_{x \in \Omega^{c}} |\Delta^{2} (h_{z}|_{\Omega})^{*} (x)| \leq \sum_{x \in \delta\Omega} |\Delta^{2} (h_{z}|_{\Omega})^{*} (x)| + \sum_{x \in \delta_{2}\Omega} |\Delta^{2} (h_{z}|_{\Omega})^{*} (x)|$$
$$=: I + II.$$

For convenience, denote $E_1 := |E(\delta\Omega, \Omega)|, E_2 := |E(\delta\Omega)|, E_3 := |E(\delta\Omega, \delta_2\Omega)|.$ Then

$$\begin{split} \mathbf{I} &= \sum_{x \in \delta\Omega} |\sum_{y \sim x} \left(\Delta \left(h_z |_{\Omega} \right)^* (y) - \Delta \left(h_z |_{\Omega} \right)^* (x) \right) | \\ &\leqslant \sum_{(x,y) \in E(\delta\Omega,\Omega)} + \sum_{(x,y) \in E(\delta\Omega)} + \sum_{(x,y) \in E(\delta\Omega,\delta_2\Omega)} |\Delta \left(h_z |_{\Omega} \right)^* (y) - \Delta \left(h_z |_{\Omega} \right)^* (x) | \\ &= \sum_{(x,y) \in E(\delta\Omega,\Omega)} \left| \Delta \left(h_z |_{\Omega} \right)^* (y) + \frac{\partial \left(h_z |_{\Omega} \right)^*}{\partial n} (x) \right| + \sum_{(x,y) \in E(\delta\Omega,\delta_2\Omega)} \left| \frac{\partial \left(h_z |_{\Omega} \right)^*}{\partial n} (x) \right| \\ &\lesssim 4dE_1 + (E_1 + 2E_2 + E_3) \max_{x \in \delta\Omega} \left| \frac{\partial \left(h_z |_{\Omega} \right)^*}{\partial n} (x) \right| . \\ &\qquad \mathbf{II} \leqslant \sum_{x \in \delta_2\Omega} |\sum_{y \sim x} \left(\Delta \left(h_z |_{\Omega} \right)^* (y) - \Delta \left(h_z |_{\Omega} \right)^* (x) \right) | \\ &\leqslant \sum_{(x,y) \in E(\delta_2\Omega,\delta\Omega)} |\Delta \left(h_z |_{\Omega} \right)^* (y)| \\ &\leqslant \sum_{(x,y) \in E(\delta_2\Omega,\delta\Omega)} \sum_{(y,y') \in E(\delta\Omega,\Omega)} |h_z(y')| \leqslant E_3E_1. \end{split}$$

Hence

(2.8)
$$\left|\partial^{2}\Omega\right| \leq 4dE_{1} + (E_{1} + 2E_{2} + E_{3})\max_{x \in \delta\Omega} \left|\frac{\partial (h_{z}|_{\Omega})^{*}}{\partial n}(x)\right| + E_{3}E_{1}.$$

3. Estimates of Dirichlet Poly-Laplace eigenvalues

In this section, we follow the methods in [24, 2, 21] to derive the upper and lower bounds of Dirichlet poly-Laplace eigenvalues on a finite subgraph Ω of \mathbb{Z}^d . Recall that

$$(-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f = \lambda^l f, \text{ in } \Omega.$$

There are $|\Omega|$ positive real eigenvalues (with multiplicities)

$$0 < \lambda_1^l \leqslant \lambda_2^l \leqslant \ldots \leqslant \lambda_{|\Omega|}^l.$$

and denote the corresponding eigenfunctions by $\{\phi_i^l\}_{1 \le i \le |\Omega|}$ respectively.

3.1. **Upper bound.** We begin with a general lemma (proved in [2]) on eigenvalues.

Lemma 3.1. Let L be a self-adjoint, positive semidefinite operator on a finite dimensional, Hermitian, complex vector space W with Hermitian inner product \langle,\rangle . Let $0 \leq \gamma_1 \leq \ldots \leq \gamma_s$ be its eigenvalues, and choose an orthonormal basis of eigenfunctions $\{f_i\}_{1 \leq i \leq |\Omega|}$, where f_i corresponds to γ_i . Then for any $1 \leq k \leq s$ and any vector $g \in W$ one has

(3.1)
$$\gamma_{k+1}\langle g,g\rangle \leqslant \langle g,Lg\rangle + \sum_{j=1}^{k} (\gamma_{k+1} - \gamma_j) |\langle g,f_j\rangle|^2.$$

By choosing $g = h_z|_{\Omega}$ and $f_j = \phi_j^l$, we can derive an inequality between λ^l_{k+1} and $\sum_{j=1}^k \lambda_j^l$ as follows.

Lemma 3.2. Let Ω be a finite subgraph of \mathbb{Z}^d . For any $1 \leq k \leq |\Omega|$ and a measurable set $B \subset [-\pi, \pi]^d$, we have

$$\lambda_{k+1}^l(|\Omega||B| - (2\pi)^d k) \leqslant |\Omega| \cdot \int_{z \in B} (\Phi(z))^l dz - (2\pi)^d \sum_{j=1}^{\kappa} \lambda_j^l + |B| \cdot |\partial^l \Omega|,$$

where $\Phi(z) = \sum_{i=1}^{d} (2 - 2\cos z_i)$, and $\lambda_{k+1}^{l} = 0$ if $k = |\Omega|$.

 $\mathit{Proof.}\,$ In Lemma 3.1, we choose

$$W = \mathbb{C}^{\Omega}, \quad L = (-1)^l \Delta_{\Omega}^{l, \mathcal{D}}, \quad g = h_z|_{\Omega}, \quad \gamma_j = \lambda_j^l, \quad f_j = \phi_j^l,$$

which satisfying the conditions of Lemma 3.1 since $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$ is self-adjoint and positive definite. Integrating both sides of the inequality (3.1) over a measurable set $B \subset [-\pi, \pi]^d$ yields

$$\lambda_{k+1}^l \int_{z \in B} \langle h_z, h_z \rangle_{\Omega} \leqslant \int_{z \in B} \langle h_z, (-1)^l \Delta_{\Omega}^{l, \mathcal{D}}(h_z|_{\Omega}) \rangle_{\Omega} + \sum_{j=1}^k (\lambda_{k+1}^l - \lambda_j^l) \int_{z \in B} |\langle h_z, \phi_j^l \rangle_{\Omega}|^2$$

14

By Lemma 2.10, the left-hand side is equal to $\lambda_{k+1}^{l}|B||\Omega|$, and the right-hand side

$$\leq \int_{z\in B} ((\Phi(z))^l |\Omega| + |\partial^l \Omega|) dz + (2\pi)^d \sum_{j=1}^k (\lambda_{k+1}^l - \lambda_j^l) \langle \phi_j^l, \phi_j^l \rangle_{\Omega}$$
$$= |\Omega| \cdot \int_{z\in B} (\Phi(z))^l dz + |B| \cdot |\partial^l \Omega| + (2\pi)^d \sum_{j=1}^k (\lambda_{k+1}^l - \lambda_j^l),$$

which completes the proof of this lemma.

Then we can prove the upper bound estimate (a) of Theorem 1.1.

Proof of Theorem 1.1 (a). In Lemma 3.2, for $1 \leq k \leq \min\{1, \frac{V_d}{2^d}\} |\Omega|$, we choose the measurable set B as a ball of radius $2\pi (\frac{k}{V_d |\Omega|})^{\frac{1}{d}}$ centered at the origin in \mathbb{R}^d , so that the radius $R = 2\pi (\frac{k}{V_d |\Omega|})^{\frac{1}{d}} \leq \pi$ and $|B| = \frac{k(2\pi)^d}{|\Omega|}$. Thus we derive an inequality

(3.2)
$$(2\pi)^d \sum_{j=1}^k \lambda_j^l \leqslant |\Omega| \cdot \int_{z \in B} (\Phi(z))^l dz + |B| \cdot |\partial^l \Omega|.$$

In the rest of the proof, we need to estimate $\int_{z\in B} (\Phi(z))^l dz$. And by calculation,

$$(\Phi(z))^{l} = \left(\sum_{i=1}^{d} (2 - 2\cos z_{i})\right)^{l} = \left(\sum_{i=1}^{d} 4\sin^{2}\frac{z_{i}}{2}\right)^{l} \leqslant \left(\sum_{i=1}^{d} z_{i}^{2}\right)^{l} = |z|^{2l}.$$

This yields that

$$\int_{z\in B} (\Phi(z))^l dz \leqslant \int_{z\in B} |z|^{2l} dz = dV_d \int_0^R r^{2l} \cdot r^{d-1} dr = \frac{dV_d}{d+2l} R^{2l+d}.$$

Applying the above inequality to (3.2), we obtain

$$\frac{1}{k} \sum_{j=1}^{k} \lambda_j^l \leqslant \frac{dV_d}{d+2l} \frac{|\Omega| R^{d+2l}}{k(2\pi)^d} + \frac{|\partial^l \Omega|}{|\Omega|}$$
$$= (2\pi)^{2l} \frac{d}{d+2l} \left(\frac{k}{V_d |\Omega|}\right)^{\frac{2l}{d}} + \frac{|\partial^l \Omega|}{|\Omega|}$$

Similarly, for $1 \leq k \leq \min\{1, \frac{V_d}{2^{d+1}}\} |\Omega|$, we choose $R' = 2^{\frac{1}{d}}R$, which implies $R' = 2\pi (\frac{2k}{V_d |\Omega|})^{\frac{1}{d}} \leq \pi$ and $|B'| = \frac{2k(2\pi)^d}{|\Omega|}$. Then we repeat the above proof and ignore the term $\sum_{j=1}^k \lambda_j^l$, obtaining that

$$k(2\pi)^d \lambda_{k+1}^l \leq |\Omega| \cdot \int_{z \in B'} (\Phi(z))^l dz + |B'| \cdot |\partial^l \Omega|$$
$$\leq |\Omega| \cdot \frac{dV_d}{d+2l} R'^{d+2l} + \frac{2k(2\pi)^d |\partial^l \Omega|}{|\Omega|},$$

which implies that

$$\lambda_{k+1}^{l} \leqslant (2\pi)^{2l} \frac{d \cdot 2^{\frac{d+2l}{d}}}{d+2l} \left(\frac{k}{V_{d}|\Omega|}\right)^{\frac{2l}{d}} + 2\frac{|\partial^{l}\Omega|}{|\Omega|}.$$

3.2. Lower bound. To prove the lower bound estimate ((b) of Theorem 1.1), we follow the method of Li and Yau [24].

Lemma 3.3. (Modification of Lemma 1 from [24]) Let F be a real-valued function on \mathbb{R}^{d} such that $0 \leq F \leq M$ and

$$\int_{[-\pi,\pi]^d} F(z) dz \ge K.$$

Assume

$$(\frac{K}{MV_d})^{\frac{1}{d}} \leqslant \sqrt{6} < \pi.$$

Then we have

$$\int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz \ge \sum_{m=0}^l \binom{l}{m} \left(-\frac{1}{12} \right)^m \frac{dK}{d+2(l+m)} \left(\frac{K}{MV_d} \right)^{\frac{2(l+m)}{d}} > 0,$$
where $\Phi(z) = \sum^d (2 - 2 \cos z)$

where $\Phi(z) = \sum_{i=1}^{a} (2 - 2\cos z_i).$

Proof. Without loss of generality, assume that $\int_{[-\pi,\pi]^d} F(z) dz = K$ by decreasing F as necessary. Then prove the lemma in two steps.

Step 1: First we are going to construct a radially symmetric function $\varphi(z): [-\pi,\pi]^d \to \mathbb{R}$ such that $0 \leq \varphi(z) \leq (\Phi(z))^l$. Since $2-2\cos x \geq x^2 - x^4/12$ for all x. Then

$$\sum_{i=1}^{d} (2 - 2\cos z_i) \ge \sum_{i=1}^{d} (z_i^2 - \frac{1}{12}z_i^4) \ge \sum_{i=1}^{d} z_i^2 - \frac{1}{12} \left(\sum_{i=1}^{d} z_i^2\right)^2 = |z|^2 - \frac{1}{12}|z|^4,$$

which means for $|z| \leq \pi < 2\sqrt{3}$

$$(\Phi(z))^{l} = \left(\sum_{i=1}^{d} (2 - 2\cos z_{i})\right)^{l} \ge \left(|z|^{2} - \frac{1}{12}|z|^{4}\right)^{l} \eqqcolon H(|z|).$$

Note that H(|z|) is monotone increasing on $[0, \sqrt{6}]$. Then we define

$$\varphi(z) = \begin{cases} H(|z|), & |z| \le \sqrt{6}, \\ H(\sqrt{6}) = 3^l, & \sqrt{6} < |z| \le \pi. \end{cases}$$

Since $\Phi(z)$ is monotone increasing in $[-\pi,\pi]^d$ along each half line starting

at 0, then we get that $(\Phi(z))^l \ge \varphi(z)$ for all $z \in [-\pi, \pi]^d$. **Step 2:** Since $F \ge 0$, we can give a lower bound of $\int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz$ as

(3.3)
$$\int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz \ge \int_{[-\pi,\pi]^d} \varphi(z) F(z) dz.$$

16

Let $R = \left(\frac{K}{MV_d}\right)^{\frac{1}{d}} \leq \sqrt{6} < \pi$ such that $MR^dV_d = K$, and define

$$\widetilde{F}(z) = \begin{cases} M & , \ |z| \leq R \\ 0 & , \ |z| > R \end{cases}$$

Then we have

$$\int_{[-\pi,\pi]^d} \widetilde{F}(z) dz = \int_{[-\pi,\pi]^d} F(z) dz = K.$$

We claim that \widetilde{F} minimizes the integral $\int_{[-\pi,\pi]^d} \varphi(z)F(z)dz$ for all functions F satisfying $0 \leq F \leq M$ and $\int_{[-\pi,\pi]^d} F(z)dz = K$. Since $\varphi(z)$ is monotonic and radially symmetric, then

$$\begin{split} &\int_{[-\pi,\pi]^d} \varphi(z)(F(z) - \widetilde{F}(z))dz \\ &= \int_{[-\pi,\pi]^d \setminus B_R} \varphi(z)F(z)dz - \int_{B_R} \varphi(z)(M - F(z))dz \\ &\geqslant \varphi(R) \int_{[-\pi,\pi]^d \setminus B_R} F(z)dz - \varphi(R) \int_{B_R} (M - F(z))dz \\ &= \varphi(R)(\int_{[-\pi,\pi]^d} F(z)dz - \int_{B_R} Mdz) = 0. \end{split}$$

Thus we have the estimate

$$\begin{split} &\int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz \geqslant \int_{[-\pi,\pi]^d} \varphi(z) F(z) dz \geqslant \int_{[-\pi,\pi]^d} \varphi(z) \widetilde{F}(z) dz \\ &= M \int_{B_R} \varphi(z) dz = M dV_d \int_0^R H(r) r^{d-1} dr \\ &= M dV_d \left(\sum_{m=0}^l \binom{l}{m} \left(-\frac{1}{12} \right)^m \frac{R^{d+2(l+m)}}{d+2(l+m)} \right) \\ &= \sum_{m=0}^l \binom{l}{m} \left(-\frac{1}{12} \right)^m \frac{dK}{d+2(l+m)} \left(\frac{K}{MV_d} \right)^{\frac{2(l+m)}{d}}, \end{split}$$

which is positive when $R = \left(\frac{K}{MV_d}\right)^{\frac{1}{d}} \leq \sqrt{6}$.

Then we can prove the lower bound estimate (b) of Theorem 1.1.

Proof of Theorem 1.1 (b). Suppose eigenfunctions $\{\phi_j^l\}_{j=1}^k$ form a standardized orthogonal basis in $\ell^2(\Omega)$. We define P_j as the projection operator on the space spanned by ϕ_j^l , and P as the projection operator on the space spanned by $\{\phi_j^l\}_{j=1}^k$. Let $h_z(x) = e^{i\langle z, x \rangle}$ as before. To apply Lemma 3.3, let

$$F_j(z) = \langle P_j h_z, P_j h_z \rangle_{\Omega} = \|P_j h_z\|_{\ell^2(\Omega)}^2,$$

$$F(z) = \sum_{j=1}^{k} F_j(z) = \|Ph_z\|_{\ell^2(\Omega)}^2.$$

By Lemma 2.10, we have

$$\begin{split} F(z) &= \|Ph_z\|_{\ell^2(\Omega)}^2 = \sum_{j=1}^k |\langle \phi_j^l, h_z \rangle_{\Omega}|^2 \leqslant \|h_z\|_{\ell^2(\Omega)}^2 = |\Omega|, \\ \int_{[-\pi,\pi]^d} F(z) dz &= \sum_{j=1}^k \int_{[-\pi,\pi]^d} |\langle \phi_j^l, h_z \rangle_{\Omega}|^2 dz = (2\pi)^d \sum_{j=1}^k \langle \phi_j^l, \phi_j^l \rangle_{\Omega} = k(2\pi)^d, \\ \int_{[-\pi,\pi]^d} (\Phi(z))^l F(z) dz &= \sum_{j=1}^k \int_{[-\pi,\pi]^d} (\Phi(z))^l |\langle \phi_j^l, h_z \rangle_{\Omega}|^2 dz \\ &= (2\pi)^d \sum_{j=1}^k \langle \phi_j^l, (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} \phi_j^l \rangle_{\Omega} = (2\pi)^d \sum_{j=1}^k \lambda_j^l. \end{split}$$

By Lemma 3.3 with $M = |\Omega|, K = k(2\pi)^d$, we get

$$\begin{split} \lambda_k^l(\Omega) &\ge \frac{1}{k} \sum_{j=1}^k \lambda_j^l(\Omega) \\ &\ge \sum_{m=0}^l \binom{l}{m} \left(-\frac{1}{12}\right)^m (2\pi)^{2(l+m)} \frac{d}{d+2(l+m)} \left(\frac{k}{|\Omega|V_d}\right)^{\frac{2(l+m)}{d}}, \end{split}$$

which is positive when $k \leq \min\{1, (\frac{\sqrt{6}}{2\pi})^d V_d\} |\Omega|$.

3.3. Comparison between eigenvalues of l and 2l-order poly-Laplace. Next we want to compare the eigenvalue of $\Delta_{\Omega}^{2l,\mathcal{D}}$ and the square of eigenvalue of $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$. First we prove a lemma as follows.

Lemma 3.4. Let Ω be a finite subgraph of G and $l \in \mathbb{N}_+$, then for all $f \in C(\Omega)$,

$$\langle \left(\Delta_{\Omega}^{l,\mathcal{D}}\right)^2 f, f \rangle_{\Omega} \leqslant \langle \Delta_{\Omega}^{2l,\mathcal{D}} f, f \rangle_{\Omega}.$$

Proof. For all $f \in C(\Omega)$, we have

$$\begin{split} \langle \left(\Delta_{\Omega}^{l,\mathcal{D}} \right)^2 f, f \rangle_{\Omega} &= \langle \Delta_{\Omega}^{l,\mathcal{D}} f, \Delta_{\Omega}^{l,\mathcal{D}} f \rangle_{\Omega} = \langle \Delta^{l} f^* |_{\Omega}, \Delta^{l} f^* |_{\Omega} \rangle_{\Omega} \\ &\leqslant \langle \Delta^{l} f^*, \Delta^{l} f^* \rangle_{G} = \langle \Delta^{2l} f^*, f^* \rangle_{G} \\ &= \langle \Delta^{2l} f^* |_{\Omega}, f \rangle_{\Omega} = \langle \Delta_{\Omega}^{2l,\mathcal{D}} f, f \rangle_{\Omega}. \end{split}$$

The following lemma states that there is no ℓ^2 -eigenfunction of poly-Laplace on \mathbb{Z}^d .

Lemma 3.5. For $l \in \mathbb{N}_+$ and $\lambda^l > 0$, if $f \in \ell^2(\mathbb{Z}^d)$ satisfies

(3.4)
$$(-\Delta)^l f(x) = \lambda^l f(x) \text{ on } \mathbb{Z}^d.$$

Then $f \equiv 0$.

Proof. Since $\ell^2(\mathbb{Z}^d)$ is the completion of $C_0(\mathbb{Z}^d)$ in ℓ^2 norm. Assume that there exists a function $f \in C_0(\mathbb{Z}^d)$ satisfying (3.4). By the Fourier transform of both sides of (3.4), we have

$$(\Phi(z))^l \hat{f}(z) = \lambda^l \hat{f}(z)$$
 on $[-\pi, \pi]^d$.

Then

$$\left\{z \in \left[-\pi, \pi\right]^d | \, \widehat{f}(z) \neq 0\right\} \subseteq \left\{z \in \left[-\pi, \pi\right]^d | \, (\Phi(z))^l = \lambda^l\right\},$$

which is a set of Lebesgue measure zero. Hence $\hat{f}(z) \stackrel{a.e.}{=} 0$, and then f(x) = 0 on \mathbb{Z}^d .

Then we are ready to prove that the eigenvalues of $\Delta_{\Omega}^{2l,\mathcal{D}}$ are at least as large as the squares of the eigenvalues of $(-1)^l \Delta_{\Omega}^{l,\mathcal{D}}$.

Proof of Theorem 1.3. By the Min-Max-Theorem and Lemma 3.4, we know

$$\begin{split} \left(\lambda_k^l\right)^2 &= \min_{\substack{M \subseteq C(\Omega) \\ \dim M = k}} \max_{\substack{\|f\|_{\ell^2} = 1}} \langle (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f, f \rangle_{\Omega}^2 \\ &\leq \min_{\substack{M \subseteq C(\Omega) \\ \dim M = k}} \max_{\substack{\|f\|_{\ell^2} = 1}} \langle \Delta_{\Omega}^{l,\mathcal{D}} f, \Delta_{\Omega}^{l,\mathcal{D}} f \rangle_{\Omega} \langle f, f \rangle_{\Omega} \\ &= \min_{\substack{M \subseteq C(\Omega) \\ \dim M = k}} \max_{\substack{\|f\|_{\ell^2} = 1}} \langle \left(\Delta_{\Omega}^{l,\mathcal{D}}\right)^2 f, f \rangle_{\Omega} \\ &\leq \min_{\substack{M \subseteq C(\Omega) \\ \dim M = k}} \max_{\substack{\|f\|_{\ell^2} = 1}} \langle \Delta_{\Omega}^{2l,\mathcal{D}} f, f \rangle_{\Omega} = \lambda_k^{2l}, \\ &\leq \min_{\substack{M \subseteq C(\Omega) \\ \dim M = k}} \max_{\substack{\|f\|_{\ell^2} = 1}} \langle \Delta_{\Omega}^{2l,\mathcal{D}} f, f \rangle_{\Omega} = \lambda_k^{2l}, \end{split}$$

Moreover, if Ω is a finite subgraph of \mathbb{Z}^d , and the equality $(\lambda_k^l)^2 = \lambda_k^{2l}$ holds, then

$$\begin{split} \lambda_k^{2l} &= \max_{\substack{f \in M_0 \\ \|f\|_{\ell^2} = 1}} \langle \Delta_{\Omega}^{2l,\mathcal{D}} f, f \rangle_{\Omega} = \max_{\substack{f \in M_0 \\ \|f\|_{\ell^2} = 1}} \langle (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f, f \rangle_{\Omega}^2 = \left(\lambda_k^l\right)^2. \end{split}$$

Suppose $f_0 \in M_0$ such that $||f_0||_{\ell^2} = 1$ and

$$\left(\lambda_k^l\right)^2 = \langle (-1)^l \Delta_{\Omega}^{l,\mathcal{D}} f_0, f_0 \rangle_{\Omega}^2.$$

Then

$$\begin{split} \left(\lambda_k^l\right)^2 &= \langle (-\Delta)^l f_0^*, f_0^* \rangle_{\mathbb{Z}^d}^2 \leqslant \langle (-\Delta)^l f_0^*, (-\Delta)^l f_0^* \rangle_{\mathbb{Z}^d} \langle f_0^*, f_0^* \rangle_{\mathbb{Z}^d} \\ &= \langle (-\Delta)^{2l} f_0^*, f_0^* \rangle_{\mathbb{Z}^d} = \langle \Delta_{\Omega}^{2l, \mathcal{D}} f_0, f_0 \rangle_{\Omega} \leqslant \lambda_k^{2l}, \end{split}$$

where the first inequality follows from the Cauchy-Schwarz inequality. Since both sides of the above formula are equal, then there is a positive constant μ such that $(-\Delta)^l f_0^* = \mu f_0^*$. By Lemma 3.5 we have $f_0^* \equiv 0$, which contradicts to $\|f_0\|_{\ell^2} = 1$.

We also have the same estimate in the continuous settings.

Theorem 3.6. Let Ω be a connected bounded domain with smooth boundary in \mathbb{R}^d and ν be the outward unit normal vector field of $\partial\Omega$. For a positive integer l, if λ_k^{il} (i = 1, 2) are the k-th eigenvalues of the Dirichlet poly-Laplace problems

$$\begin{cases} (-\Delta)^{il}f = \lambda^{il}f, & \text{in }\Omega, \\ f = \frac{\partial f}{\partial \nu} = \ldots = \frac{\partial^{il-1}f}{\partial^{il-1}\nu} = 0, & \text{on }\partial\Omega, \end{cases}$$

respectively. Then

$$\left(\lambda_k^l\right)^2 \leqslant \lambda_k^{2l}.$$

Proof. Recall

$$H_0^{il}(\Omega) = \left\{ f : \|f\|_{H^{il}} = \sum_{|\alpha| \le il} \|D^{\alpha} f\|_{L^2} < +\infty, \ D^{\alpha} f|_{\partial\Omega} = 0 \text{ for } |\alpha| \le il - 1 \right\},$$

and note that $H_0^{il}(\Omega)$ is the completion of $C_0^{\infty}(\Omega)$ in H^{il} norm. Then

$$\begin{split} \left(\lambda_k^l\right)^2 &= \inf_{\substack{M \subseteq H_0^l(\Omega) \\ \dim M = k}} \sup_{\substack{\|f\|_{L^2} = 1}} \langle \nabla^l f, \nabla^l f \rangle_{\Omega}^2 \\ &= \inf_{\substack{M \subseteq C_0^{\infty}(\Omega) \\ \dim M = k}} \sup_{\substack{\|f\|_{L^2} = 1}} \langle (-\Delta)^l f, f \rangle_{\Omega}^2 \\ &\leq \inf_{\substack{M \subseteq C_0^{\infty}(\Omega) \\ \dim M = k}} \sup_{\substack{\|f\|_{L^2} = 1}} \langle (-\Delta)^l f, (-\Delta)^l f \rangle_{\Omega} \langle f, f \rangle_{\Omega} \\ &\leq \inf_{\substack{M \subseteq C_0^{\infty}(\Omega) \\ \dim M = k}} \sup_{\substack{\|f\|_{L^2} = 1}} \langle (-\Delta)^l f, (-\Delta)^l f \rangle_{\Omega} = \lambda_k^{2l}. \\ &= \inf_{\substack{M \subseteq H_0^{2l}(\Omega) \\ \dim M = k}} \sup_{\substack{\|f\|_{L^2} = 1}} \langle (-\Delta)^l f, (-\Delta)^l f \rangle_{\Omega} = \lambda_k^{2l}. \end{split}$$

4. Appendix

The following numerical experiment illustrates that on the path graph $[0, n] \subseteq \mathbb{Z}$, the ratio of the squares of discrete Dirichlet Laplace eigenvalues and Dirichlet bi-Laplace eigenvalues in Remark 1.4 (3) approximates the continuous case $(0, n) \subseteq \mathbb{R}$, that is,

$$\frac{\left(\lambda_k^1([0,n])\right)^2}{\lambda_k^2([0,n])} \to c_k = \frac{\left(\lambda_k^1((0,1))\right)^2}{\lambda_k^2((0,1))} < 1 \text{ as } n \to +\infty,$$

suggesting that the definition (1.5) is reasonable.



Figure 1. This figure shows the convergence of ratio of eigenvalues.

Acknowledgements. The authors would like to thank Florentin Münch for helpful discussions. B. Hua is supported by NSFC, No. 12371056, and by Shanghai Science and Technology Program [Project No. 22JC1400100].

Conflicts of Interests. The authors declared no potential conflicts of interests with respect to this article.

Ethics Approval. This study did not involve any human participants or animals, and therefore, ethical approval was not required.

Data Availability. Data availability is not applicable to this article as no new data were created or analyzed in this study.

References

- Shmuel Agmon. On kernels, eigenvalues, and eigenfunctions of operators related to elliptic problems. Comm. Pure Appl. Math., 18:627–663, 1965.
- [2] Frank Bauer and Gabor Lippner. Eigenvalue sum estimates for lattice subgraphs. Pure and Applied Mathematics Quarterly, 18(6):2339–2353, 2022. 1, 1, 1, 2.1, 3, 3.1
- [3] F. A. Berezin. Covariant and contravariant symbols of operators. Izv. Akad. Nauk SSSR Ser. Mat., 36:1134–1167, 1972. 1
- [4] M Sh Birman and MZ Solomyak. Asymptotic behavior of the spectrum of differential equations. Journal of Soviet Mathematics, 12:247–283, 1979. 1

- [5] M. Š. Birman and M. Z. Solomjak. Quantitative analysis in Sobolev imbedding theorems and applications to spectral theory, volume 114 of American Mathematical Society Translations, Series 2. American Mathematical Society, Providence, RI, 1980. Translated from the Russian by F. A. Cezus. 1
- [6] Qing-Ming Cheng, Xuerong Qi, and Guoxin Wei. A lower bound for eigenvalues of the poly-Laplacian with arbitrary order. *Pacific J. Math.*, 262(1):35–47, 2013. 1
- [7] Qing-Ming Cheng and Guoxin Wei. Upper and lower bounds for eigenvalues of the clamped plate problem. J. Differential Equations, 255(2):220–233, 2013. 1
- [8] RJ Duffin and EP Shelly. Difference equations of polyharmonic type. 1958. 1
- [9] Powei Feng and Joe Warren. Discrete bi-laplacians and biharmonic b-splines. ACM Transactions on Graphics (TOG), 31(4):1–11, 2012.
- [10] Clemens Förster and Jörgen Östensson. Lieb-Thirring inequalities for higher order differential operators. Math. Nachr., 281(2):199–213, 2008. 1
- [11] Filippo Gazzola, Hans-Christoph Grunau, and Guido Sweers. Polyharmonic boundary value problems, volume 1991 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2010. Positivity preserving and nonlinear higher order elliptic equations in bounded domains. 1
- [12] Robert Grone, Russell Merris, and Viakalathur Shankar Sunder. The laplacian spectrum of a graph. SIAM Journal on matrix analysis and applications, 11(2):218–238, 1990. 1
- [13] Stephen Guattery and Gary L Miller. Graph embeddings and laplacian eigenvalues. SIAM Journal on Matrix Analysis and Applications, 21(3):703–723, 2000. 1
- [14] Antoine Henrot. Extremum problems for eigenvalues of elliptic operators. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2006. 1
- [15] Thomas Hirschler and Wolfgang Woess. Laplace and bi-laplace equations for directed networks and markov chains. *Expositiones Mathematicae*, 39(2):271–301, 2021. 1
- [16] Bobo Hua and Matthias Keller. Harmonic functions of general graph laplacians. Calculus of Variations and Partial Differential Equations, 51(1):343–362, 2014. 1
- [17] Jürgen Jost, Xianqing Li-Jost, Qiaoling Wang, and Changyu Xia. Universal bounds for eigenvalues of the polyharmonic operators. *Trans. Amer. Math. Soc.*, 363(4):1821– 1854, 2011. 1
- [18] Ahmad Khalkhuzhaev, Shokhrukh Yu Kholmatov, and Mardon Pardabaev. Expansion of eigenvalues of rank-one perturbations of the discrete bilaplacian. arXiv preprint arXiv:1910.01369, 2019. 1
- [19] Hynek Kova rík, Semjon Vugalter, and Timo Weidl. Two-dimensional Berezin-Li-Yau inequalities with a correction term. *Comm. Math. Phys.*, 287(3):959–981, 2009.
- [20] P. Kröger. Upper bounds for the Neumann eigenvalues on a bounded domain in Euclidean space. J. Funct. Anal., 106(2):353–357, 1992. 1
- [21] Pawel Kröger. Estimates for sums of eigenvalues of the Laplacian. J. Funct. Anal., 126(1):217–227, 1994. 1, 1, 3
- [22] Ari Laptev and Timo Weidl. Recent results on Lieb-Thirring inequalities. In Journées "Équations aux Dérivées Partielles" (La Chapelle sur Erdre, 2000), pages Exp. No. XX, 14. Univ. Nantes, Nantes, 2000. 1
- [23] H. A. Levine and M. H. Protter. Unrestricted lower bounds for eigenvalues for classes of elliptic equations and systems of equations with applications to problems in elasticity. *Math. Methods Appl. Sci.*, 7(2):210–222, 1985. 1
- [24] Peter Li and Shing Tung Yau. On the Schrödinger equation and the eigenvalue problem. Comm. Math. Phys., 88(3):309–318, 1983. 1, 1, 1, 3, 3.2, 3.3
- [25] Elliott H. Lieb. The number of bound states of one-body Schroedinger operators and the Weyl problem. In Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), volume XXXVI of Proc. Sympos. Pure Math., pages 241–252. Amer. Math. Soc., Providence, RI, 1980. 1

- [26] Antonios D. Melas. A lower bound for sums of eigenvalues of the Laplacian. Proc. Amer. Math. Soc., 131(2):631–636, 2003. 1
- [27] Y. Netrusov and T. Weidl. On Lieb-Thirring inequalities for higher order operators with critical and subcritical powers. *Comm. Math. Phys.*, 182(2):355–370, 1996. 1
- [28] Å ke Pleijel. On the eigenvalues and eigenfunctions of elastic plates. Comm. Pure Appl. Math., 3:1–10, 1950. 1
- [29] G. Pólya. On the eigenvalues of vibrating membranes. Proc. London Math. Soc. (3), 11:419–433, 1961. 1
- [30] Florian Martin Schweiger. On the membrane model and the discrete Bilaplacian. PhD thesis, Universitäts-und Landesbibliothek Bonn, 2021. 1
- [31] Robert S. Strichartz. Estimates for sums of eigenvalues for domains in homogeneous spaces. J. Funct. Anal., 137(1):152–190, 1996. 1
- [32] Jiaxuan Wang. Eigenvalue estimates for the fractional laplacian on lattice subgraphs. arXiv preprint arXiv:2303.15766, 2023. 1
- [33] Guoxin Wei and Lingzhong Zeng. Estimates for eigenvalues of poly-harmonic operators. Adv. Nonlinear Stud., 16(1):31–44, 2016. 1
- [34] Hermann Weyl. Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung). Math. Ann., 71(4):441–479, 1912. 1
- [35] Selma Yıldırı m Yolcu and Türkay Yolcu. Eigenvalue bounds for the poly-harmonic operators. *Illinois J. Math.*, 58(3):847–865, 2014. 1
- [36] Selma Yıldırı m Yolcu and Türkay Yolcu. Multidimensional lower bounds for the eigenvalues of Stokes and Dirichlet Laplacian operators. J. Math. Phys., 53(4):043508, 17, 2012. 1

BOBO HUA: SCHOOL OF MATHEMATICAL SCIENCES, LMNS, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA; SHANGHAI CENTER FOR MATHEMATICAL SCIENCES, JIANG-WAN CAMPUS, FUDAN UNIVERSITY, NO. 2005 SONGHU ROAD, SHANGHAI 200438, CHINA.

Email address: bobohua@fudan.edu.cn

Ruowei Li: School of Mathematical Sciences, Fudan University, Shanghai 200433, China; Shanghai Center for Mathematical Sciences, Jiangwan Campus, Fudan University, No. 2005 Songhu Road, Shanghai 200438, China; MPI MIS Leipzig, 04103 Leipzig, Germany.

Email address: rwli19@fudan.edu.cn