

# INVARIANT THEORY AND COEFFICIENT ALGEBRAS OF LIE ALGEBRAS

YIN CHEN AND RUNXUAN ZHANG

**ABSTRACT.** The coefficient algebra of a finite-dimensional Lie algebra with respect to a faithful representation is defined as the subalgebra generated by all coefficients of the corresponding characteristic polynomial. We establish a connection between classical invariant theory and the coefficient algebras of finite-dimensional complex Lie algebras. Specifically, we prove that, with respect to any symmetric power of the standard representation: (1) the coefficient algebra of the upper triangular solvable complex Lie algebra is isomorphic to the algebra of symmetric polynomials; (2) the coefficient algebra of the general linear complex Lie algebra is the invariant ring of the general linear group with the conjugacy action on the full space of matrices; and (3) the coefficient algebra of the special linear complex Lie algebra can be generated by classical trace functions. As an application, we exactly exhibit the characteristic polynomial of the special linear complex Lie algebra.

## 1. INTRODUCTION

**1.1. Background.** The characteristic polynomial of a square matrix undoubtedly occupies a central position among the various branches of modern algebra. The theory of characteristic polynomial of multiple square matrices dates back to the works of Dedekind and Frobenius on group determinants of finite nonabelian groups in the 19th century. More precisely, the group determinant of a finite group  $G = \{1, g_1, \dots, g_n\}$  is a homogeneous polynomial in  $x_0, x_1, \dots, x_n$  defined as

$$(1.1) \quad \varphi_G := \det(x_0 \cdot I + x_1 \cdot \varphi(g_1) + \dots + x_n \cdot \varphi(g_n))$$

where  $I$  denotes the identity matrix and  $\varphi$  denotes the left regular representation of  $G$ . Factoring the group determinant into a product of irreducible polynomials and establishing connections between group determinants and the group structure theory are two fundamental tasks in this area, which paved the way for the representation theory of finite groups.

The group determinant of a finite group  $G$  can be generalized to the notion of characteristic polynomial of  $G$  with respect to any finite-dimensional faithful representation  $V$  via replacing the left regular representation  $\varphi$  in (1.1) by the group homomorphism from  $G$  to  $\mathrm{GL}(V)$ . Recently, an analogue of characteristic polynomials in Lie theory has been introduced by [HZ19] and has captured the attention of many researchers; see for example, [CCD19, JL22], and [GLW24]. One of our motivational examples was [HZ19, Theorem 2] which provides a sufficient and necessary condition for when a Lie algebra is solvable via its characteristic polynomial.

To better understand the characteristic polynomial of a Lie algebra  $\mathfrak{g}$ , we introduce a concept of the coefficient algebra of  $\mathfrak{g}$  with respect to a faithful representation  $V$  and the primary objective in this article is to compute the coefficient algebras for various Lie algebras. Our approach exposes

---

*Date:* November 19, 2024.

*Key words and phrases.* Lie algebras; coefficient algebras; invariant rings; characteristic polynomials.

*2020 Mathematics Subject Classification.* 13A50; 17B05; 15A15.

some unexpected connections between classical invariant theory and the characteristic polynomials of Lie algebras.

**1.2. Characteristic polynomial and coefficient algebra.** Let  $k$  be a field of any characteristic and  $\mathfrak{g}$  be a finite-dimensional Lie algebra over  $k$ . Suppose that  $V$  denotes a faithful  $m$ -dimensional representation of  $\mathfrak{g}$  and let's fix a basis  $\{e_1, e_2, \dots, e_n\}$  of  $\mathfrak{g}$ . We may use  $E_i$  to denote the image of  $e_i$  on  $V$  for all  $i = 1, 2, \dots, n$ . The *characteristic polynomial* of  $\mathfrak{g}_V$  with respect to the basis  $\{e_1, e_2, \dots, e_n\}$  is the following homogeneous polynomial of degree  $m$  in  $n + 1$  variables  $x_0, x_1, \dots, x_n$ :

$$(1.2) \quad \varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n) := \det(x_0 \cdot I_m + x_1 \cdot E_1 + \dots + x_n \cdot E_n).$$

Note that this definition is well-defined because it is independent of the choice of basis in  $V$ . Thus after choosing any basis of  $V$ , each  $E_i$  could be regarded as a square matrix of size  $m$ .

We take a new perspective to understand the characteristic polynomial  $\varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n)$ . In fact, since  $\varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n)$  also can be reviewed as a monic polynomial of degree  $m$  in the single variable  $x_0$  over the polynomial ring  $k[x_1, \dots, x_n]$ , we may write

$$(1.3) \quad \varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n) = x_0^m + c_1 \cdot x_0^{m-1} + \dots + c_{m-1} \cdot x_0 + c_m$$

for some homogeneous polynomials  $c_1, c_2, \dots, c_m \in k[x_1, \dots, x_n]$ . In a sense, understanding the polynomial  $\varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n)$  is equivalent to determining these coefficients  $c_1, c_2, \dots, c_m$ . This philosophy leads us to introduce the following concept.

**DEFINITION 1.1.** The  $k$ -subalgebra generated by  $\{c_1, c_2, \dots, c_m\}$  in  $k[x_1, \dots, x_n]$  is called the *coefficient algebra* of the characteristic polynomial  $\varphi_{\mathfrak{g}_V}$  and denoted as  $B_{\mathfrak{g}_V} := k[c_1, c_2, \dots, c_m]$ .

**1.3. Main results.** The main purpose of this article is to determine the structure of the coefficient algebra  $B_{\mathfrak{g}_V}$  for some well-known finite-dimensional linear Lie algebras over the complex field with respect to the symmetric powers of their standard representations. The structure of a coefficient algebra  $B_{\mathfrak{g}_V}$  would further enhance our understanding of the characteristic polynomial  $\varphi_{\mathfrak{g}_V}$  and the Lie representation  $(\mathfrak{g}, V)$ .

To articulate our main results, we suppose  $n, d \in \mathbb{N}^+$  in this subsection and use  $\mathfrak{u}_n(\mathbb{C})$  to denote the solvable Lie algebra of all upper triangular  $n \times n$ -matrices over the complex field  $\mathbb{C}$ . Let  $S^d(V)$  denote the  $d$ -th symmetric power of a vector space  $V$ . In particular, we identify  $S^1(V)$  with  $V$  itself. The first main result is the following.

**Theorem 1.2.** *The coefficient algebra of  $\mathfrak{u}_n(\mathbb{C})$  with respect to any symmetric power  $S^d(V)$  of the standard representation  $V$  is isomorphic to the symmetric invariant ring  $\mathbb{C}[x_1, x_2, \dots, x_n]^{S_n}$ , where  $S_n$  denotes the symmetric group of degree  $n$  with the standard permutation action on  $\{1, 2, \dots, n\}$ .*

We denote by  $\mathfrak{gl}_n(\mathbb{C})$  the general linear Lie algebra of degree  $n$  over  $\mathbb{C}$  and suppose that the general linear group  $\mathrm{GL}_n(\mathbb{C})$  of degree  $n$  acts on  $\mathfrak{gl}_n(\mathbb{C})$  in the conjugation way. Our second result below determines the coefficient algebra of  $\mathfrak{gl}_n(\mathbb{C})$ .

**Theorem 1.3.** *The coefficient algebra of  $\mathfrak{gl}_n(\mathbb{C})$  with respect to any symmetric power  $S^d(V)$  of the standard representation  $V$  is equal to the invariant ring*

$$\mathbb{C}[\mathfrak{gl}_n(\mathbb{C})]^{\mathrm{GL}_n(\mathbb{C})} = \mathbb{C}[x_{ij} \mid 1 \leq i, j \leq n]^{\mathrm{GL}_n(\mathbb{C})} = \mathbb{C}[s_1, s_2, \dots, s_n],$$

where  $s_r$  denote the classical conjugacy invariant of degree  $r$  in  $x_{ij}$ .

The third result computes the coefficient algebra of the special linear Lie algebra  $\mathfrak{sl}_n(\mathbb{C})$ .

**Theorem 1.4.** *The coefficient algebra of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to any symmetric power  $S^d(V)$  of the standard representation  $V$  is equal to the invariant ring  $\mathbb{C}[\mathfrak{sl}_n(\mathbb{C})]^{\mathrm{SL}_n(\mathbb{C})}$ , and so it is a polynomial algebra generated by the trace functions  $\mathrm{tr}_2, \mathrm{tr}_3, \dots, \mathrm{tr}_n$ .*

This theorem has an immediate application (Corollary 4.1) that determines the characteristic polynomial of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to the standard representation.

**1.4. Layout and conventions.** In Section 2, we begin with computing the coefficient algebra of  $\mathfrak{sl}_2(\mathbb{C})$  with respect to any representation  $V$  (Corollary 2.2) and provide a perspective from invariant theory to understand the characteristic polynomial of a Lie algebra with respect to a self-dual representation. We also prove that a Lie algebra is nilpotent if and only if the coefficient algebra with respect to the adjoint representation is zero; see Proposition 2.5. Section 3 is devoted the proofs of Theorems 1.2 and 1.3 in which we will see that how classical invariant theory plays a key role in determining characteristic polynomials and coefficient algebras. In Section 4, we prove Theorem 1.4, calculating the coefficient algebra of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to any symmetric power of the standard representation.

Throughout this paper,  $k$  denotes a field of any characteristic and  $\mathbb{C}$  denotes for the complex field. The symbol  $k[V]^G = k[x_1, x_2, \dots, x_n]^G$  is standard in invariant theory and denotes the invariant ring of a group  $G$  with respect to a representation  $V$ , where  $x_1, x_2, \dots, x_n$  form a basis of the dual space  $V^*$ ; see for example, [CW11] or [DK15] for a general reference on invariant theory and [CDG20, CT19, CZ23b] and [CZ24] for some recent applications of invariant theory.

## 2. $\mathfrak{sl}_2(\mathbb{C})$ AND NILPOTENT LIE ALGEBRAS

A new theory (concept) in Lie algebras usually begins with the study on  $\mathfrak{sl}_2(\mathbb{C})$ ; see [CCZ21, Che24], and [CZ23b, Section 5]. The characteristic polynomial of  $\mathfrak{sl}_2(\mathbb{C})$  with respect to any representation  $V$  has been computed by [CCD19, Theorem 4] and [JL22]. Based on their calculations, this section completely determines the coefficient algebras of  $\mathfrak{sl}_2(\mathbb{C})$  (for any representation) and nilpotent Lie algebras (for the adjoint representation). Additionally, we provide a viewpoint from invariant theory to understand the characteristic polynomials of a Lie algebra with respect to self-dual representations.

2.1.  $\mathfrak{sl}_2(\mathbb{C})$ . Let's begin with setting  $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$  and fixing a basis:

$$(2.1) \quad e_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, e_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Suppose  $V$  denotes the standard 2-dimensional representation of  $\mathfrak{sl}_2(\mathbb{C})$  on  $\mathbb{C}^2$ . Then the characteristic polynomial

$$\begin{aligned} \varphi_{\mathfrak{g}_V}(x_0, x_1, x_2, x_3) &= \det \left( x_0 \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + x_1 \cdot \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + x_2 \cdot \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} + x_3 \cdot \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right) \\ &= \det \begin{pmatrix} x_0 - x_2 & x_1 \\ x_3 & x_0 + x_2 \end{pmatrix} \\ &= x_0^2 - (x_2^2 + x_1 x_3). \end{aligned}$$

Hence,  $B_{\mathfrak{g}_V} = \mathbb{C}[x_2^2 + x_1x_3]$  is a polynomial subalgebra of  $\mathbb{C}[x_1, x_2, x_3]$  with Krull dimension 1. Moreover, this statement remains to hold if  $V$  is any irreducible representation of  $\mathfrak{sl}_2(\mathbb{C})$ .

**Proposition 2.1.** *Let  $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$  with the basis (2.1) and  $V$  be any irreducible representation of  $\mathfrak{sl}_2(\mathbb{C})$ . Then  $B_{\mathfrak{g}_V}$  is a polynomial subalgebra of  $\mathbb{C}[x_1, x_2, x_3]$  generated by  $x_2^2 + x_1x_3$ .*

*Proof.* By [CCD19, Theorem 4], we see that the characteristic polynomial  $\varphi_{\mathfrak{g}_V}(x_0, x_1, x_2, x_3)$  can be written as a product of either  $x_0$  or  $x_0^2 - a \cdot (x_2^2 + x_1x_3)$  for some  $a \in \mathbb{N}$ . In other words,

$$\varphi_{\mathfrak{g}_V}(x_0, x_1, x_2, x_3) = x_0^{\ell_0} \cdot \left(x_0^2 - a_1 \cdot (x_2^2 + x_1x_3)\right) \cdots \left(x_0^2 - a_s \cdot (x_2^2 + x_1x_3)\right)$$

for some  $s \in \mathbb{N}^+$  and  $\ell_0, a_1, a_2, \dots, a_s \in \mathbb{N}$ . For  $1 \leq i \leq s$ , let us define

$$\theta_i := a_i \cdot (x_2^2 + x_1x_3).$$

Recall that the coefficients of a polynomial equation can be expressed as elementary symmetric polynomials in their roots. Thus

$$\varphi_{\mathfrak{g}_V}(x_0, x_1, x_2, x_3) = x_0^{\ell_0} \cdot \left(x_0^{2s} - f_1(\theta_1, \dots, \theta_s) \cdot x_0^{2s-2} + \cdots + (-1)^s f_s(\theta_1, \dots, \theta_s)\right)$$

where  $f_i$  denotes the elementary symmetric polynomial of degree  $i$  for all  $i \in \{1, \dots, s\}$ . Note that each  $\theta_i \in \mathbb{C}[x_2^2 + x_1x_3]$ . Therefore,  $B_{\mathfrak{g}_V} = \mathbb{C}[x_2^2 + x_1x_3]$ .  $\square$

**Corollary 2.2.** *Let  $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$  with the basis (2.1) and  $V$  be any finite-dimensional faithful representation of  $\mathfrak{sl}_2(\mathbb{C})$ . Then  $B_{\mathfrak{g}_V}$  is generated by  $x_2^2 + x_1x_3$  in  $\mathbb{C}[x_1, x_2, x_3]$ .*

*Proof.* Note that for two faithful representations  $V$  and  $W$  of  $\mathfrak{g}$ , it follows from the definition of characteristic polynomial that  $\varphi_{\mathfrak{g}_{V \oplus W}} = \varphi_{\mathfrak{g}_V} \cdot \varphi_{\mathfrak{g}_W}$ . Thus combining Proposition 2.1 with the fact that  $\mathfrak{sl}_2(\mathbb{C})$  is completely reducible (i.e., each representation can be written as the direct sum of some irreducible representations), we see that  $I_{\mathfrak{g}_V} = \mathbb{C}[x_2^2 + x_1x_3]$ .  $\square$

More generally, recall that the base change in a Lie algebra  $\mathfrak{g}$  would induce a degree-preserving algebraic action on the polynomial ring  $k[x_0, x_1, \dots, x_n]$ . Specifically, if  $P$  denotes the transition matrix between two bases in  $\mathfrak{g}$ , then

$$\tilde{P} := \begin{pmatrix} 1 & 0 \\ 0 & P^\top \end{pmatrix}$$

can linearly transform  $\{x_0, x_1, \dots, x_n\}$  to a new generating set  $\{x_0, x'_1, \dots, x'_n\}$  of  $k[x_0, x_1, \dots, x_n]$  and then can be extended to a  $k$ -algebraic action on  $k[x_0, x_1, \dots, x_n]$ . Under this action, the characteristic polynomial with respect to the second basis of  $\mathfrak{g}$  is exactly equal to the image of the characteristic polynomial with respect to the first basis.

Together with Proposition 2.1 and Corollary 2.2 obtains the following result.

**Corollary 2.3.** *Let  $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$  with any basis and  $V$  be any finite-dimensional faithful representation of  $\mathfrak{sl}_2(\mathbb{C})$ . Then  $B_{\mathfrak{g}_V}$  is generated by an irreducible quadratic polynomial in  $\mathbb{C}[x_1, x_2, x_3]$ .*

**2.2. Self-dual representations.** A representation  $V$  of a Lie algebra and the dual  $V^*$  correspond to the same coefficient algebra, through their characteristic polynomials are different.

**Proposition 2.4.** *Let  $\mathfrak{g}$  be an  $n$ -dimensional Lie algebra with any basis and  $V$  be a finite-dimensional faithful representation of  $\mathfrak{g}$ . If  $V^*$  is the dual representation of  $V$ , then  $B_{\mathfrak{g}_{V^*}} = B_{\mathfrak{g}_V}$ .*

*Proof.* By [KKS24, Proposition 2.2], we see that the characteristic polynomial  $\varphi_{\mathfrak{g}_{V^*}}(x_0, x_1, \dots, x_n)$  is equal to  $(-1)^{\dim(V)} \cdot \varphi_{\mathfrak{g}_V}(-x_0, x_1, \dots, x_n)$ . This means that up to sign,  $\varphi_{\mathfrak{g}_{V^*}}$  and  $\varphi_{\mathfrak{g}_V}$  have the same coefficients. Hence,  $B_{\mathfrak{g}_{V^*}} = B_{\mathfrak{g}_V}$ .  $\square$

We would like to provide a point of view from invariant theory to better understand the characteristic polynomial of a Lie algebra  $\mathfrak{g}$  for self-dual representations.

Suppose the characteristic of the ground field  $k$  is not 2 and that  $V$  is a faithful self-dual representation of  $\mathfrak{g}$ , i.e.,  $V^* \cong V$ . Then

$$\varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n) = \varphi_{\mathfrak{g}_{V^*}}(x_0, x_1, \dots, x_n) = (-1)^{\dim(V)} \cdot \varphi_{\mathfrak{g}_V}(-x_0, x_1, \dots, x_n).$$

SUBCASE 1. If  $\dim(V)$  is even, then

$$(2.2) \quad \varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n) = \varphi_{\mathfrak{g}_V}(-x_0, x_1, \dots, x_n).$$

Namely,  $\varphi_{\mathfrak{g}_V}$  is an invariant polynomial under the action of the group  $\langle \sigma \rangle$  on  $k[x_0, x_1, \dots, x_n]$ , where

$$\sigma := \text{diag}\{-1, 1, \dots, 1\}.$$

Note that the order of  $\sigma$  is 2. From invariant theory, it is not difficult to see that the invariant ring  $k[x_0, x_1, \dots, x_n]^{\langle \sigma \rangle}$  is generated by  $\{x_0^2, x_1, \dots, x_n\}$ . This means that  $\varphi_{\mathfrak{g}_V}$  must be a polynomial in  $x_0^2, x_1, \dots, x_n$ .

SUBCASE 2. Suppose  $\dim(V)$  is odd and recall that the degree of  $\varphi_{\mathfrak{g}_V}$  equals  $\dim(V)$ , we have

$$(2.3) \quad \varphi_{\mathfrak{g}_V}(x_0, x_1, \dots, x_n) = -\varphi_{\mathfrak{g}_V}(-x_0, x_1, \dots, x_n) = \varphi_{\mathfrak{g}_V}(x_0, -x_1, \dots, -x_n).$$

In other words, the characteristic polynomial  $\varphi_{\mathfrak{g}_V}$ , in this case, is an invariant polynomial under the action of the group  $\langle -\sigma \rangle$  on  $k[x_0, x_1, \dots, x_n]$ . Since the invariant ring  $k[x_0, x_1, \dots, x_n]^{\langle -\sigma \rangle}$  is generated by

$$\{x_0, x_i x_j \mid 1 \leq i, j \leq n\},$$

it follows that  $\varphi_{\mathfrak{g}_V}$  is a polynomial in  $x_0$  and some  $x_i x_j$ , where  $i, j \in \{1, 2, \dots, n\}$ .

**2.3. Nilpotent Lie algebras.** We may obtain a sufficient and necessary description for when a complex Lie algebra is nilpotent via the coefficient algebra corresponding to the adjoint representation.

**Proposition 2.5.** *Let  $\mathfrak{g}$  be an  $n$ -dimensional complex Lie algebra with any basis and  $V$  be the adjoint representation of  $\mathfrak{g}$ . Then  $\mathfrak{g}$  is nilpotent if and only if  $B_{\mathfrak{g}_V} = 0$ .*

*Proof.* This immediately follows from [KKS24, Theorem 4.8], which states that  $\mathfrak{g}$  is nilpotent if and only if the corresponding characteristic polynomial is  $x_0^n$ .  $\square$

REMARK 2.6. Proposition 2.5 also can be obtained from the Engel's theorem. In fact, by Engel's theorem,  $\mathfrak{g}$  is nilpotent if and only if  $\mathfrak{g}$  has a basis with respect to which each element of  $\mathfrak{g}$  on the adjoint representation  $V$  is an upper triangular matrix along with diagonal entries 0; see, for example, [Car05, Corollary 2.11]. The latter statement is equivalent to saying that  $B_{\mathfrak{g}_V} = 0$ .  $\diamond$

Moreover, the Lie's theorem (see [Car05, Corollary 2.3]) is also greatly helpful in computing the characteristic polynomials and the coefficient algebras of some solvable Lie algebras, and we believe, especially for the cases of high-dimensional Lie algebras and representations.

Let's close this section with the following example that illustrates how we can use the Lie's theorem to calculate the coefficient algebra of a solvable Lie algebra.

EXAMPLE 2.7. Suppose that  $\mathfrak{g}$  denotes the 3-dimensional subalgebra of  $\mathfrak{gl}_3(\mathbb{C})$  consisting of all matrices of the following form

$$\begin{pmatrix} 0 & a & b \\ -a & 0 & c \\ 0 & 0 & 0 \end{pmatrix}$$

where  $a, b, c \in \mathbb{C}$ . Then  $\mathfrak{g}$  is a solvable but not nilpotent Lie algebra. Let  $V = \mathbb{C}^3$  be the standard representation of  $\mathfrak{g}$ . We define

$$(2.4) \quad A := \begin{pmatrix} 0 & x_1 & x_2 \\ -x_1 & 0 & x_3 \\ 0 & 0 & 0 \end{pmatrix}$$

The direct computing approach shows that

$$\varphi_{\mathfrak{g}_V}(x_0, x_1, x_2, x_3) = \det(x_0 \cdot I + A) = \det \begin{pmatrix} x_0 & x_1 & x_2 \\ -x_1 & x_0 & x_3 \\ 0 & 0 & x_0 \end{pmatrix} = x_0^3 + x_1^2 \cdot x_0.$$

Hence, the coefficient algebra  $B_{\mathfrak{g}_V} = \mathbb{C}[x_1^2]$ .

On the other hand, using Lie's theorem we may prove that there exists an upper triangular matrix in  $\mathfrak{gl}_3(\mathbb{C}[x_1, x_2, x_3])$  such that it is similar with  $A$ . By [Bro93, Corollary 16.4], the matrix  $x_0 \cdot I + A$  must be equivalent to an upper triangular matrix over  $\mathbb{C}[x_1, x_2, x_3][x_0]$ . In other words, after some elementary row/column transformations over  $\mathbb{C}[x_0, x_1, x_2, x_3]$ , the matrix  $x_0 \cdot I + A$  could be upper triangularized. Suppose that  $B \in \mathfrak{gl}_3(\mathbb{C}[x_0, x_1, x_2, x_3])$  denotes an equivalent upper triangular matrix of  $x_0 \cdot I + A$ . Then there exist two products  $P, Q$  of elementary matrices such that

$$B = P \cdot (x_0 \cdot I + A) \cdot Q.$$

Hence,

$$\det(x_0 \cdot I + A) = \frac{\det(B)}{\det(P) \cdot \det(Q)}.$$

In our case, we let  $P_1$  be the transformation of multiplying the first row with  $x_1$ ,  $P_2$  the transformation of multiplying the second row with  $x_0$ , and  $P_3$  be the transformation of adding the first row to the second row. Setting  $P = P_3 P_2 P_1$  and  $Q = I$ , we obtain

$$B = \begin{pmatrix} x_0 x_1 & x_1^2 & x_1 x_2 \\ 0 & x_0^2 + x_1^2 & x_0 x_3 + x_1 x_2 \\ 0 & 0 & x_0 \end{pmatrix}$$

Therefore,  $\det(x_0 \cdot I + A) = \frac{\det(B)}{\det(P) \cdot \det(Q)} = x_0^3 + x_1^2 \cdot x_0$ , and  $B_{\mathfrak{g}_V} = \mathbb{C}[x_1^2]$ .  $\diamond$

### 3. COEFFICIENT ALGEBRAS FOR $\mathfrak{u}_n(\mathbb{C})$ AND $\mathfrak{gl}_n(\mathbb{C})$

This section will work over the complex field  $\mathbb{C}$  and we will compute the coefficient algebras of the upper triangular matrix algebra  $\mathfrak{u}_n(\mathbb{C})$  and the general linear Lie algebra  $\mathfrak{gl}_n(\mathbb{C})$ , with respect to the symmetric powers of their standard representations. These two algebras play extremely significant roles in the complex representation theory of Lie algebras, because any solvable Lie algebra  $\mathfrak{h}$  could be viewed as a subalgebra of  $\mathfrak{u}_n(\mathbb{C})$  for some  $n$  (Lie's Theorem), and any Lie algebra  $\mathfrak{g}$  also could be viewed as a subalgebra of  $\mathfrak{gl}_n(\mathbb{C})$  under some representations (Ado's Theorem).

Suppose that  $V = \mathbb{C}^n$  denotes the  $n$ -dimensional standard representation of  $\mathfrak{gl}_n(\mathbb{C})$  and  $\mathfrak{u}_n(\mathbb{C})$  and  $S^d(V) = S^d(\mathbb{C}^n)$  denotes the  $d$ th symmetric power of  $\mathbb{C}^n$  for all  $d \in \mathbb{N}^+$ . We define

$$\{E_{ij} \mid 1 \leq i, j \leq n\}$$

to be the standard basis for  $\mathfrak{gl}_n(\mathbb{C})$ , where  $E_{ij}$  denotes the  $n \times n$ -matrix whose the  $(i, j)$ -entry is 1 and others 0. Recall that the symmetric invariant ring  $\mathbb{C}[x_1, x_2, \dots, x_n]^{S_n}$  is a polynomial generated by the elementary symmetric polynomials  $\epsilon_1, \epsilon_2, \dots, \epsilon_n$ ; see for example, [Stu08, Section 1.1].

We are ready to prove our first main result.

*Proof of Theorem 1.2.* Let's first consider the case of  $d = 1$ , i.e., the case of standard representation  $\mathbb{C}^n$ . In this case, recall that the dimension of  $\mathfrak{u}_n(\mathbb{C})$  is  $\frac{n(n+1)}{2}$  and the standard basis of  $\mathfrak{u}_n(\mathbb{C})$  could be chosen as  $\{E_{ij} \mid 1 \leq i \leq j \leq n\}$ . Thus we may assume that the characteristic polynomial is

$$\varphi(x_0, x_{ij}) = \det \left( x_0 \cdot I_n + \sum_{1 \leq i \leq j \leq n} x_{ij} \cdot E_{ij} \right)$$

for some variables  $x_0, x_{ij}$ . Hence,

$$\varphi(x_0, x_{ij}) = \det \left( x_0 \cdot I + \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ 0 & x_{22} & \ddots & \vdots \\ \vdots & \ddots & \ddots & x_{n-1,n} \\ 0 & \cdots & 0 & x_{nn} \end{pmatrix} \right) = \prod_{i=1}^n (x_0 + x_{ii}).$$

Therefore, the coefficient algebra of  $\mathfrak{u}_n(\mathbb{C})$  we are looking for is a polynomial algebra generated by  $n$  elementary polynomials in  $x_{11}, x_{22}, \dots, x_{nn}$ , which is isomorphic to  $\mathbb{C}[x_1, x_2, \dots, x_n]^{S_n}$ .

Suppose that  $d \geq 2$ . The resulting matrix of each  $E_{ij}$  (when  $i < j$ ) on the symmetric power  $S^d(\mathbb{C}^n)$  is a strictly upper triangular elementary matrix (i.e., upper triangular with diagonals 0 and in which there exists only one nonzero entry 1 above the diagonal) as well as the resulting matrices of all  $E_{ii}$  ( $1 \leq i \leq n$ ) are diagonal matrices with only one nonzero diagonal entry 1 and these matrices are distinct to each other. Hence, the characteristic polynomial in this case can be written as

$$\varphi(x_0, x_1, \dots, x_n) = x_0^r \cdot \prod_{i=1}^n (x_0 + x_i)$$

where  $r = \dim(S^d(\mathbb{C}^n)) - n = \binom{n+d-1}{d} - n$ . Therefore, the coefficient algebra is also isomorphic to  $\mathbb{C}[x_1, x_2, \dots, x_n]^{S_n}$ .  $\square$

**REMARK 3.1.** Computing invariants of the symmetric group  $S_n$  is a classical topic in the invariant theory. If the characteristic of  $k$  is zero, the invariant ring  $k[x_1, x_2, \dots, x_n]^{S_n}$  also can be generated by the power sums  $p_1, p_2, \dots, p_n$ , where

$$(3.1) \quad p_j := x_1^j + x_2^j + \cdots + x_n^j$$

for  $j \in \mathbb{N}^+$ ; see for example, [Stu08, Proposition 1.1.2] for more details.  $\diamond$

**EXAMPLE 3.2.** (1) For the Lie algebra  $\mathfrak{u}_2(\mathbb{C})$  and the standard representation  $\mathbb{C}^2$ , the coefficient algebra is  $\mathbb{C}[x_{11} + x_{22}, x_{11}x_{22}]$  and the characteristic polynomial  $\varphi(x_0, x_{11}, x_{22}) = (x_0 + x_{11})(x_0 + x_{22})$ .



(2) For the Lie algebra  $\mathfrak{u}_3(\mathbb{C})$  and the standard representation  $\mathbb{C}^3$ , the coefficient algebra is

$$\mathbb{C}[x_{11} + x_{22} + x_{33}, x_{11}x_{22} + x_{11}x_{33} + x_{22}x_{33}, x_{11}x_{22}x_{33}]$$

and the characteristic polynomial  $\varphi(x_0, x_{11}, x_{22}, x_{33}) = (x_0 + x_{11})(x_0 + x_{22})(x_0 + x_{33})$ .

(3) For the Lie algebra  $\mathfrak{u}_2(\mathbb{C})$  and the 2nd symmetric power of the standard representation  $\mathbb{C}^2$ , the coefficient algebra is also  $\mathbb{C}[x_{11} + x_{22}, x_{11}x_{22}]$  but the characteristic polynomial  $\varphi(x_0, x_{11}, x_{22}) = x_0^2 \cdot (x_0 + x_{11})(x_0 + x_{22})$ .  $\diamond$

To prove Theorem 1.3, let's consider the conjugation action of the general linear group  $\mathrm{GL}_n(\mathbb{C})$  on the underlying space of  $\mathfrak{gl}_n(\mathbb{C})$  defined by

$$(3.2) \quad (g, A) \mapsto g \cdot A \cdot g^{-1}$$

for all  $g \in \mathrm{GL}_n(\mathbb{C})$  and  $A \in \mathfrak{gl}_n(\mathbb{C})$ . It is well-known from classical invariant theory that the invariant ring

$$(3.3) \quad \mathbb{C}[\mathfrak{gl}_n(\mathbb{C})]^{\mathrm{GL}_n(\mathbb{C})} = \mathbb{C}[x_{ij} \mid 1 \leq i, j \leq n]^{\mathrm{GL}_n(\mathbb{C})} = \mathbb{C}[s_1, s_2, \dots, s_n]$$

is a polynomial algebra over  $\mathbb{C}$ ; see for example, [Kra17, Proposition 4.1.2].

*Proof of Theorem 1.3.* We still consider the case of  $d = 1$  first. Given  $n^2 + 1$  new variables  $x_0, x_{ij}$  for  $1 \leq i, j \leq n$  and setting a generic matrix  $X = (x_{ij})_{n \times n}$ , we may assume that the characteristic polynomial of  $\mathfrak{gl}_n(\mathbb{C})$  with respect to the standard basis of  $\mathbb{C}^n$  is

$$\begin{aligned} \varphi(x_0, x_{ij}) &= \det \left( x_0 \cdot I_n + \sum_{1 \leq i, j \leq n} x_{ij} \cdot E_{ij} \right) \\ &= \det(x_0 \cdot I_n + X) \\ &= x_0^n + \sum_{i=1}^n s_i \cdot x_0^{n-i} \end{aligned}$$

where  $s_i$  denotes the sum of all principal  $i \times i$ -minors of  $X$ ; see [Car17, Theorem 8.3]. Therefore, the coefficient algebra of  $\mathfrak{gl}_n(\mathbb{C})$  with respect to the standard representation  $\mathbb{C}^n (= S^1(\mathbb{C}^n))$  is exactly equal to  $\mathbb{C}[s_1, s_2, \dots, s_n] = \mathbb{C}[\mathfrak{gl}_n(\mathbb{C})]^{\mathrm{GL}_n(\mathbb{C})}$ .

Now suppose  $d \geq 2$  and consider the symmetric power  $S^d(\mathbb{C}^n)$ . We need to examine the resulting matrix  $[X]$  of the generic matrix  $X = (x_{ij})_{n \times n}$  on  $S^d(\mathbb{C}^n)$  and so the characteristic polynomial of  $\mathfrak{gl}_n(\mathbb{C})$  with respect to  $S^d(\mathbb{C}^n)$  is  $\det(x_0 \cdot I + [X])$ .

Let  $X_u$  be the matrix obtained from  $X$  by setting  $x_{ij} = 0$  for all  $i > j$  and let  $[X_u]$  be the resulting matrix of  $X_u$  on  $S^d(\mathbb{C}^n)$ . Clearly,  $X_u$  is an upper triangular matrix of size  $n$ . We have seen from the proof of Theorem 1.2 that for all  $i \in \{1, 2, \dots, \dim(S^d(\mathbb{C}^n))\}$ , if the  $i$ -th diagonal entry of the resulting matrix  $[X_u]$  is zero, then all entries on the  $i$ -th row and the  $i$ -th column in  $[X_u]$  are zero. We define

$$m := \dim(S^d(\mathbb{C}^n)) = \binom{n+d-1}{d}.$$

Then we have  $m - n$  zeros on the diagonal of  $[X_u]$ . Symmetrically, we may obtain the same conclusion by considering the low triangular part  $X_\ell$  of  $X$ . The total number of zero entries in  $[X]$  is

$$(2m - 1) + (2m - 1 - 2) + \dots + (2m - 1 - 2(m - n - 1)) = m^2 - n^2.$$



This means that each  $x_{ij}$  will occur in  $[X]$  and only occurs once. By Laplace's expansion, we obtain the characteristic polynomial of  $\mathfrak{gl}_n(\mathbb{C})$  with  $S^d(\mathbb{C}^n)$ :

$$(3.4) \quad \varphi(x_0, x_{ij}) = \det(x_0 \cdot I + [X]) = x_0^{m-n} \cdot \det(x_0 \cdot I + X).$$

Hence, the coefficient algebra in this case is also equal to  $\mathbb{C}[s_1, s_2, \dots, s_n] = \mathbb{C}[\mathfrak{gl}_n(\mathbb{C})]^{\mathrm{GL}_n(\mathbb{C})}$ .  $\square$

The following two examples illustrate how to compute the coefficients  $s_1, s_2, \dots, s_n$ .

EXAMPLE 3.3. The coefficient algebra for  $\mathfrak{gl}_2(\mathbb{C})$  can be generated by  $s_1$  and  $s_2$ , where

$$X = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}$$

and  $s_1 = \mathrm{Tr}(X) = x_{11} + x_{22}$  and  $s_2 = \det(X) = x_{11}x_{22} - x_{12}x_{21}$ . Thus the characteristic polynomial of  $\mathfrak{gl}_2(\mathbb{C})$  is  $x_0^2 + s_1 \cdot x_0 + s_2 = x_0^2 + (x_{11} + x_{22}) \cdot x_0 + (x_{11}x_{22} - x_{12}x_{21})$ .  $\diamond$

EXAMPLE 3.4. The coefficient algebra for  $\mathfrak{gl}_3(\mathbb{C})$  is generated by  $s_1, s_2$ , and  $s_3$ , where

$$X = \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix}.$$

and

$$\begin{aligned} s_1 &= \mathrm{Tr}(X) = x_{11} + x_{22} + x_{33} \\ s_2 &= \sum(\text{all } 2 \times 2 \text{ minors of } X) = \det \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} + \det \begin{pmatrix} x_{11} & x_{13} \\ x_{31} & x_{33} \end{pmatrix} + \det \begin{pmatrix} x_{22} & x_{23} \\ x_{32} & x_{33} \end{pmatrix} \\ &= (x_{11}x_{22} - x_{12}x_{21}) + (x_{11}x_{33} - x_{13}x_{31}) + (x_{22}x_{33} - x_{23}x_{32}) \\ s_3 &= \det(X). \end{aligned}$$

Therefore, the characteristic polynomial of  $\mathfrak{gl}_3(\mathbb{C})$  is  $x_0^3 + s_1 \cdot x_0^2 + s_2 \cdot x_0 + s_3$ .  $\diamond$

REMARK 3.5. The invariant ring  $\mathbb{C}[\mathfrak{gl}_n(\mathbb{C})]^{\mathrm{GL}_n(\mathbb{C})} = \mathbb{C}[x_{ij} \mid 1 \leq i, j \leq n]^{\mathrm{GL}_n(\mathbb{C})}$  also can be generated by the trace functions  $\mathrm{Tr}_1, \mathrm{Tr}_2, \dots, \mathrm{Tr}_n$ , where

$$(3.5) \quad \mathrm{Tr}_i : \mathfrak{gl}_n(\mathbb{C}) \longrightarrow \mathbb{C}, \quad A \mapsto \mathrm{Tr}(A^i)$$

for  $i \in \mathbb{N}^+$ . Namely,  $\mathrm{Tr}_i = \mathrm{Tr}(X^i)$  for all  $i \in \mathbb{N}^+$ . For example, if  $n = 2$ , then

$$\mathrm{Tr}_1 = \mathrm{Tr} \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} = x_{11} + x_{22} \text{ and } \mathrm{Tr}_2 = \mathrm{Tr} \left[ \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}^2 \right] = x_{11}^2 + 2x_{12}x_{21} + x_{22}^2;$$

see [KP96, Section 2.4] for more details. Note that these trace functions defined in (3.5) can be used to characterize the coefficient algebra of the special linear Lie algebra  $\mathfrak{sl}_n(\mathbb{C})$  in next section.  $\diamond$

#### 4. COEFFICIENT ALGEBRA FOR $\mathfrak{sl}_n(\mathbb{C})$

Suppose that  $n \geq 2$  and  $\mathfrak{sl}_n(\mathbb{C})$  denotes the special complex Lie algebra of degree  $n$ . Let  $V$  be the standard  $(n^2 - 1)$ -dimensional representation of  $\mathfrak{sl}_n(\mathbb{C})$  with the following basis:

$$\{E_{ij}, E_{s,s} - E_{s+1,s+1} \mid 1 \leq i \neq j \leq n, 1 \leq s \leq n-1\}.$$

Hence, the characteristic polynomial is

$$\begin{aligned}\varphi_{\mathfrak{g}_V}(x_0, x_{ij}, x_1, x_2, \dots, x_{n-1}) &= \det \left( x_0 \cdot I + \sum_{1 \leq i \neq j \leq n} x_{ij} \cdot E_{ij} + \sum_{s=1}^{n-1} x_s \cdot (E_{s,s} - E_{s+1,s+1}) \right) \\ &= x_0^n + c_1 \cdot x_0^{n-1} + c_2 \cdot x_0^{n-2} + \dots + c_{n-1} \cdot x_0 + c_n\end{aligned}$$

for some  $c_1, c_2, \dots, c_n \in \mathbb{C}[x_{ij}, x_1, x_2, \dots, x_{n-1} \mid 1 \leq i \neq j \leq n]$ .

Consider the generic matrix

$$(4.1) \quad \begin{aligned}X &:= \sum_{1 \leq i \neq j \leq n} x_{ij} \cdot E_{ij} + \sum_{s=1}^{n-1} x_s \cdot (E_{s,s} - E_{s+1,s+1}) \\ &= \begin{pmatrix} x_1 & x_{12} & x_{13} & \cdots & x_{1n} \\ x_{21} & x_2 - x_1 & x_{23} & \cdots & x_{2n} \\ \vdots & \cdots & \ddots & \ddots & \vdots \\ \vdots & \cdots & \cdots & x_{n-1} - x_{n-2} & x_{n-1,n} \\ x_{n1} & \cdots & \cdots & x_{n,n-1} & -x_{n-1} \end{pmatrix}.\end{aligned}$$

Note that  $c_1$  is equal to the trace of  $X$  up to sign, thus  $c_1 = 0$ , and the coefficient algebra we are going to compute is  $\mathbb{C}[c_2, \dots, c_n]$ .

For  $i \in \mathbb{N}^+$ , we suppose that  $\text{tr}_i$  denotes the restriction of  $\text{Tr}_i$  defined in (3.5) on  $\mathfrak{sl}_n(\mathbb{C})$ . Namely,  $\text{tr}_i$  is the polynomial obtained from  $\text{Tr}_i$  by substituting

$$x_{11} = x_1, x_{22} = x_2 - x_1, \dots, x_{n-1,n-1} = x_{n-1} - x_{n-2}, x_{nn} = -x_{n-1}.$$

For example, if  $n = 2$ , then

$$\text{tr}_1 = \text{Tr}_1(x_1, -x_1) = 0, \text{ and } \text{tr}_2 = \text{Tr}_2(x_1, x_{12}, x_{21}, -x_1) = 2(x_{11}^2 + x_{12}x_{21}).$$

More generally, we see that  $\text{tr}_1 = c_1 = 0$  for all  $n \in \mathbb{N}^+$ .

*Proof of Theorem 1.4.* We first suppose  $d = 1$  and in this case, we *claim* that  $\mathbb{C}[c_2, \dots, c_n] = \mathbb{C}[\text{tr}_2, \dots, \text{tr}_n]$ . Recall that

$$\mathbb{C}[x_{ij} \mid 1 \leq i, j \leq n]^{\text{GL}_n(\mathbb{C})} = \mathbb{C}[s_1, s_2, \dots, s_n] = \mathbb{C}[\text{Tr}_1, \text{Tr}_2, \dots, \text{Tr}_n].$$

Thus for each  $s_i$ , there exists a polynomial  $f_i \in \mathbb{C}[y_1, y_2, \dots, y_n]$  such that

$$s_i = f_i(\text{Tr}_1, \text{Tr}_2, \dots, \text{Tr}_n).$$

Note that up to sign,  $c_i$  is equal to the restriction of  $s_i$  on  $\mathfrak{sl}_n(\mathbb{C})$ , thus  $c_i$  can be expressed algebraically by  $\{\text{tr}_1, \text{tr}_2, \dots, \text{tr}_n\}$  (or  $\{\text{tr}_2, \dots, \text{tr}_n\}$  because of  $\text{tr}_1 = 0$ ). Hence,  $\mathbb{C}[c_2, \dots, c_n] \subseteq \mathbb{C}[\text{tr}_2, \dots, \text{tr}_n]$ . Conversely, as each  $\text{Tr}_i$  can be expressed algebraically by  $\{s_1, s_2, \dots, s_n\}$ , so restricting into  $\mathfrak{sl}_n(\mathbb{C})$  gives us an algebraic expression of  $\text{tr}_i$  in  $\{c_2, \dots, c_n\}$ . Thus  $\mathbb{C}[\text{tr}_2, \dots, \text{tr}_n] \subseteq \mathbb{C}[c_2, \dots, c_n]$  and the claim holds.

On the other hand, the conjugation action of the special linear group  $\text{SL}_n(\mathbb{C})$  on the underlying space of  $\mathfrak{sl}_n(\mathbb{C})$  induces a  $\mathbb{C}$ -algebraic action of  $\text{SL}_n(\mathbb{C})$  on the polynomial ring  $\mathbb{C}[\mathfrak{sl}_n(\mathbb{C})]$ , and the invariant ring  $\mathbb{C}[\mathfrak{sl}_n(\mathbb{C})]^{\text{SL}_n(\mathbb{C})}$  is minimally generated by

$$\{\text{tr}_2, \text{tr}_3, \dots, \text{tr}_n\};$$

see for example, [Bou05, Chapter VIII, Exercise 13]. Hence, the coefficient algebra of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to  $V$  is actually equal to  $\mathbb{C}[\mathfrak{sl}_n(\mathbb{C})]^{\mathrm{SL}_n(\mathbb{C})}$ .

To complete the proof, we need to consider the general case of  $d \geq 2$ . According to the argument in the proof of Theorem 1.3, the resulting matrix of the generic matrix  $X$  defined in (4.1) on  $S^d(V)$  is the restriction of the resulting matrix of  $(x_{ij})_{n \times n}$  on  $S^d(\mathbb{C}^n)$  to  $S^d(V)$ . This means that the characteristic polynomial of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to  $S^d(V)$  is the product of  $x_0^{m-n}$  and the characteristic polynomial of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to  $V$ , where  $m$  denotes the dimension of  $S^d(V)$ . Therefore, the coefficient algebra of  $\mathfrak{sl}_n(\mathbb{C})$  with respect to  $S^d(V)$  is also equal to  $\mathbb{C}[\mathfrak{sl}_n(\mathbb{C})]^{\mathrm{SL}_n(\mathbb{C})}$ .  $\square$

Theorem 1.4, together with the relationship between  $\{s_1, s_2, \dots, s_n\}$  and  $\{\mathrm{Tr}_1, \mathrm{Tr}_2, \dots, \mathrm{Tr}_n\}$  (see for example [Car17, Section 8.1.4]), obtains the following result.

**Corollary 4.1.** *The characteristic polynomial of the special linear Lie algebra  $\mathfrak{sl}_n(\mathbb{C})$  with respect to the standard representation is*

$$(4.2) \quad \varphi(x_0, x_{ij}, x_1, \dots, x_{n-1}) = x_0^n + \sum_{r=2}^n c_r \cdot x_0^{n-r}$$

where  $1 \leq i \neq j \leq n$  and

$$(4.3) \quad c_r := \frac{1}{r!} \det \begin{pmatrix} 0 & r-1 & 0 & \cdots & 0 \\ \mathrm{tr}_2 & 0 & r-2 & \ddots & \vdots \\ \mathrm{tr}_3 & \mathrm{tr}_2 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & 1 \\ \mathrm{tr}_r & \cdots & \mathrm{tr}_3 & \mathrm{tr}_2 & 0 \end{pmatrix}.$$

We provide the following examples to explain how to find the characteristic polynomial of  $\mathfrak{sl}_n(\mathbb{C})$ .

EXAMPLE 4.2. Suppose  $n = 2$ . Then  $X = \begin{pmatrix} x_1 & x_{12} \\ x_{21} & -x_1 \end{pmatrix}$ . Clearly,  $\mathrm{tr}_1 = \mathrm{Tr}(X) = 0$  and since

$$X^2 = \begin{pmatrix} x_1 & x_{12} \\ x_{21} & -x_1 \end{pmatrix} \begin{pmatrix} x_1 & x_{12} \\ x_{21} & -x_1 \end{pmatrix} = \begin{pmatrix} x_1^2 + x_{12}x_{21} & 0 \\ 0 & x_1^2 + x_{12}x_{21} \end{pmatrix}$$

it follows that  $\mathrm{tr}_2 = \mathrm{Tr}(X^2) = 2(x_1^2 + x_{12}x_{21})$ . Therefore, the characteristic polynomial is

$$\varphi(x_0, x_{12}, x_{21}, x_1) = x_0^2 - \frac{\mathrm{tr}_2}{2!} = x_0^2 - (x_1^2 + x_{12}x_{21}).$$

Compare this result with Section 2.1.  $\diamond$

EXAMPLE 4.3. Suppose  $n = 3$ . Then  $X = \begin{pmatrix} x_1 & x_{12} & x_{13} \\ x_{21} & x_2 - x_1 & x_{23} \\ x_{31} & x_{32} & -x_2 \end{pmatrix}$  and  $\mathrm{tr}_1 = \mathrm{Tr}(X) = 0$ . A direct calculation shows that

$$X^2 = \begin{pmatrix} x_{12}x_{21} + x_{13}x_{31} + x_1^2 & x_{13}x_{32} + x_{12}x_2 & x_{13}x_1 - x_{13}x_2 + x_{12}x_{23} \\ x_{23}x_{31} + x_{21}x_2 & x_{23}x_{32} + x_{12}x_{21} + x_1^2 - 2x_1x_2 + x_2^2 & x_{13}x_{21} - x_{23}x_1 \\ x_{31}x_1 - x_{31}x_2 + x_{32}x_{21} & x_{12}x_{31} - x_{32}x_1 & x_{13}x_{31} + x_{23}x_{32} + x_2^2 \end{pmatrix}.$$

Thus  $\mathrm{tr}_2 = \mathrm{Tr}(X^2) = 2(x_{12}x_{21} + x_{13}x_{31} + x_{23}x_{32} + x_1^2 - x_1x_2 + x_2^2)$ . Similarly, computing  $X^3$  obtains  $\mathrm{tr}_3 = \mathrm{Tr}(X^3) = 3(x_1x_{13}x_{31} - x_2x_{13}x_{31} + x_{12}x_{23}x_{31} + x_{13}x_{21}x_{32} - x_{23}x_{32}x_1 + x_{12}x_{21}x_2 + x_1^2x_2 - x_1x_2^2)$ .

Hence, the characteristic polynomial is

$$\varphi(x_0, x_{12}, \dots, x_{32}, x_1, x_2) = x_0^3 + \frac{1}{2!} \det \begin{pmatrix} 0 & 1 \\ \text{tr}_2 & 0 \end{pmatrix} \cdot x_0 + \frac{1}{3!} \det \begin{pmatrix} 0 & 2 & 0 \\ \text{tr}_2 & 0 & 1 \\ \text{tr}_3 & \text{tr}_2 & 0 \end{pmatrix}$$

which could be simplified as  $x_0^3 - \frac{\text{tr}_2}{2} \cdot x_0 + \frac{\text{tr}_3}{3}$ . ◇

**Acknowledgements.** This research was partially supported by the University of Saskatchewan under grant No. APEF-121159.

#### REFERENCES

- [Bou05] Nicolas Bourbaki, *Lie groups and Lie algebras. Chapters 7–9*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2005.
- [Bro93] William C. Brown, *Matrices over commutative rings*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 169, Marcel Dekker, Inc., New York, 1993.
- [CW11] H. E. A. Campbell and David L. Wehlau, *Modular invariant theory*, Encyclopaedia of Mathematical Sciences, vol. 139, Springer-Verlag, Berlin, 2011.
- [Car17] James B. Carrell, *Groups, matrices, and vector spaces*, Springer, New York, 2017.
- [Car05] Roger W. Carter, *Lie algebras of finite and affine type*, Cambridge Studies in Advanced Mathematics, vol. 96, Cambridge University Press, Cambridge, 2005.
- [CCZ21] Hongliang Chang, Yin Chen, and Runxuan Zhang, *A generalization on derivations of Lie algebras*, Electron. Res. Arch. **29** (2021), no. 3, 2457–2473.
- [CCD19] Zhiqi Chen, Xueqing Chen, and Ming Ding, *On the characteristic polynomial of  $\mathfrak{sl}(2, \mathbb{F})$* , Linear Algebra Appl. **579** (2019), 237–243.
- [Che24] Yin Chen, *Some Lie algebra structures on symmetric powers*, Amer. Math. Monthly (published online) (2024), <https://doi.org/10.1080/00029890.2024.2416375>.
- [CDG20] Yin Chen, Rong Du, and Yun Gao, *Modular quotient varieties and singularities by the cyclic group of order  $2p$* , Comm. Algebra **48** (2020), no. 12, 5490–5500.
- [CT19] Yin Chen and Zhongming Tang, *Vector invariant fields of finite classical groups*, J. Algebra **534** (2019), 129–144.
- [CZ23a] Yin Chen and Xinxin Zhang, *A class of quadratic matrix equations over finite fields*, Algebra Colloq. **30** (2023), no. 1, 169–180.
- [CZ23b] Yin Chen and Runxuan Zhang, *A commutative algebra approach to multiplicative Hom-Lie algebras*, Linear Multilinear Algebra **71** (2023), no. 7, 1127–1144.
- [CZ24] ———, *Shape enumerators of self-dual NRT codes over finite fields*, SIAM J. Discrete Math. **38** (2024), no. 4, 2841–2854.
- [DK15] Harm Derksen and Gregor Kemper, *Computational invariant theory*, Second enlarged edition, Encyclopaedia of Mathematical Sciences, vol. 130, Springer, Heidelberg, 2015.
- [GLW24] Amin Geng, Shoumin Liu, and Xumin Wang, *Characteristic polynomials and finite dimensional representations of simple Lie algebras*, New York J. Math. **30** (2024), 24–37.
- [HZ19] Zhiguang Hu and Philip B. Zhang, *Determinants and characteristic polynomials of Lie algebras*, Linear Algebra Appl. **563** (2019), 426–439.
- [JL22] Tianyi Jiang and Shoumin Liu, *Characteristic polynomials and finitely dimensional representations of  $\mathfrak{sl}(2, \mathbb{C})$* , Linear Algebra Appl. **647** (2022), 78–88.
- [KKS24] Korkeat Korkeathikhun, Borworn Khuhirun, Songpon Sriwongsa, and Keng Wiboonton, *More on characteristic polynomials of Lie algebras*, J. Algebra **643** (2024), 294–310.
- [Kra17] Hanspeter Kraft, *Algebraic transformation groups - an introduction*, Lecture notes for courses, 2017. available at: <https://kraftadmin.wixsite.com/hpkraft>.
- [KP96] Hanspeter Kraft and Claudio Procesi, *Classical invariant theory - a primer*, Lecture notes for courses, 1996. available at: <https://kraftadmin.wixsite.com/hpkraft>.
- [Stu08] Bernd Sturmfels, *Algorithms in invariant theory*, 2nd ed., Texts and Monographs in Symbolic Computation, SpringerWienNewYork, Vienna, 2008.

DEPARTMENT OF FINANCE AND MANAGEMENT SCIENCE, UNIVERSITY OF SASKATCHEWAN, SASKATOON, SK,  
CANADA, S7N 5A7

*Email address:* `yin.chen@usask.ca`

DEPARTMENT OF MATHEMATICS AND INFORMATION TECHNOLOGY, CONCORDIA UNIVERSITY OF  
EDMONTON, EDMONTON, AB, CANADA, T5B 4E4

*Email address:* `runxuan.zhang@concordia.ab.ca`