SUMS OF S-UNITS IN X-COORDINATES OF PELL EQUATIONS PARVATHI S NAIR AND S. S. ROUT

ABSTRACT. Let S be a fixed set of primes and let $(X_l)_{l\geq 1}$ be the X-coordinates of the positive integer solutions (X, Y) of the Pell equation $X^2 - dY^2 = 1$ corresponding to a non-square integer d > 1. We show that there are only a finite number of nonsquare integers d > 1 such that there are at least two different elements of the sequence $(X_l)_{l\geq 1}$ that can be represented as a sum of S-units with a fixed number of terms. Furthermore, we solve explicitly a particular case in which two of the X-coordinates are product of power of two and power of three.

1. INTRODUCTION

The problem of finding specific terms of a linear recurrence sequence of some particular form has a very rich history. Pethő [18] and Shorey-Stewart [21] independently studied the perfect powers in a non-degenerate binary recurrence sequence. In particular, they have considered the Diophantine equation

$$U_n = x^z \tag{1}$$

in integers n, x, z with $z \ge 2$, where $(U_n)_{n\ge 0}$ is a non-degenerate binary recurrence sequence and proved under certain natural assumptions that (1) contain only finitely many perfect powers. For example, Fibonacci and Lucas numbers, respectively, of the form x^z , with z > 1 has been recently proved by Bugeaud et al. [5]. Pethő [19] proved that there are no non-trivial perfect powers in Pell sequence (see also [7]). Pethő and Tichy [20] proved that there are only finitely many Fibonacci numbers of the form $p^a + p^b + p^c$, with prime p and integers $a > b > c \ge 0$. Marques and Togbé [16] found all Fibonacci and Lucas numbers of the form $2^a + 3^b + 5^c$ under the condition that $c \ge \max\{a, b\} \ge 0$.

Recently, Bertók et al., [3] under some mild assumptions gave finiteness result for the solutions of the Diophantine equation

$$U_n = b_1 p_1^{z_1} + \dots + b_s p_s^{z_s} \tag{2}$$

²⁰²⁰ Mathematics Subject Classification: 11B37 (Primary), 11D45, 11J86 (Secondary).

Keywords: Pell equations, Diophantine equations, S-units, linear forms in logarithms, reduction method.

Diophantine equations combining both S-units and recurrence sequences has been well studied by many authors (see [2, 3, 10, 13]). Suppose S is the set of distinct primes p_1, \ldots, p_l . Then a rational integer z is an S-unit if z can be written as

$$z = \pm p_1^{e_1} \cdots p_l^{e_l},\tag{3}$$

where e_1, \ldots, e_l are non-negative integers and we denote the set of S-units by \mathcal{U}_S . Let $(U_n)_{n\geq 0}$ be a recurrence sequence of order k with $k \geq 2$. Bérczes et al. [2] considered the equation

$$U_n = z_1 + \dots + z_r,\tag{4}$$

with some arbitrary but fixed $r \ge 1$, in unknown $n \ge 0$ and $z_1, \ldots, z_r \in \mathcal{U}_S$ and establish finiteness results for the solutions of (4).

Let d, t be nonzero integers with d > 1 square-free. Consider the Pell equation

$$x^2 - dy^2 = 1\tag{5}$$

in integers x, y. Recently, several mathematicians have investigated the following type of problem related to solution sets (i.e., X and Y-coordinates) of Pell equation (5). Assume that $\mathbb{U} := (U_n)_{n\geq 0}$ is some interesting sequence of positive integers and $\{(X_m, Y_m)\}_{m\geq 1}$ are sequence of solutions of Pell equation (5). What can one say about the number of solutions of the containment $X_m \in \mathbb{U}$ for a generic d? What about the number of solutions of the containment $Y_m \in \mathbb{U}$? For most of the binary recurrent sequences(Fibonacci numbers [4, 15], tribonacci numbers [14], rep-digits in some given integer base $b \geq 2$ [8, 11, 12]), the equation $X_m \in \mathbb{U}$ has at most one positive integer solution m for any given d except for finitely many values of d. Erazo et al., [10] showed under certain conditions that there are only a finite number non-square integers d > 1 such that there are at least two different elements of the sequence $(X_m)_{m\geq 1}$ that can be represented as a linear combination of prime powers with fixed primes, i.e.,

$$X_m = c_1 p_1^{n_1} + \dots + c_s p_s^{n_s}$$

In this paper, we extend this result to sums of S-units. In particular, we will prove that there are only a finite number non-square integers d > 1 such that there are at least two different elements of the sequence $(X_m)_{m\geq 1}$ that can be represented as a sum of S-units. Firstly, we prove the general case when elements of the sequence $(X_m)_{m\geq 1}$ equal to sum of S-units, that is,

$$X_l = z_1 + \dots + z_r,\tag{6}$$

where $z_1, \ldots, z_r \in \mathcal{U}_S$ and l are positive integers. Then in next result, we solve a particular case of (6) with $S = \{2, 3\}$.

2. NOTATION AND MAIN RESULTS

Let d > 1 be an integer which is not a square. The Pell equation

$$X^2 - dY^2 = 1, (7)$$

where $X, Y \in \mathbb{Z}_{>0}$ has infinitely many positive integer solutions (X, Y) and have the form

$$X + Y\sqrt{d} = X_l + Y_l\sqrt{d} = (X_1 + Y_1\sqrt{d})^l$$

for some $k \in \mathbb{Z}_{>0}$ and (X_1, Y_1) is the smallest positive integer solution of (7). The sequence $(X_l)_{l\geq 1}$ is a binary recurrence sequence satisfying the recurrence relation

$$X_l = 2X_1 X_{l-1} - X_{l-2} \tag{8}$$

for all $l \geq 2$. Setting

$$\gamma := X_1 + Y_1 \sqrt{d}$$
 and $\eta := X_1 - Y_1 \sqrt{d} = \gamma^{-1}$, (9)

so $\gamma \cdot \eta = X_1^2 - dY_1^2 = 1$. The Binet formula for X_l and Y_l are

$$X_l = \frac{\gamma^l + \eta^l}{2} \quad \text{and} \quad Y_l = \frac{\gamma^l - \eta^l}{2\sqrt{d}} \tag{10}$$

holds for all non-negative integers l. Let $s \in \mathbb{Z}_{>0}$ be fixed. We are interested to determine for which positive integers d > 1, the sequence $(X_l)_{l\geq 1}$ of X-coordinates of (7) has at least two different terms that can be represented as in (6). By denoting $z_i = p_1^{n_{i1}} \cdots p_s^{n_{is}}$ for all $1 \leq i \leq r$, we write (6) as

$$X_l = p_1^{n_{11}} p_2^{n_{12}} \cdots p_s^{n_{1s}} + \dots + p_1^{n_{r1}} p_2^{n_{r2}} \cdots p_s^{n_{rs}}.$$
 (11)

such that,

$$n_{rs} = \max_{\substack{1 \le i \le r \\ 1 \le j \le s}} \{n_{ij}\}.$$
 (12)

Now we are ready to state our main theorem.

Theorem 2.1. Let s be a fixed positive integer. Let $p_1 \leq \cdots \leq p_s$ be fixed primes with p_s odd. Let X_l be the X-coordinate of the Pell equation (7) with d > 1 non-square. Let $\epsilon > 0$ be arbitrary. Consider (6) in $l \geq 1$ and z_1, \ldots, z_r satisfying $|z_i|^{1+\epsilon} < |z_r|$, $(i = 1, \ldots, r-1)$ and (12). Let T_d be the set of solutions (l, z_1, \ldots, z_r) of (6). Then there exists effectively computable constants $c_8(s)$ and $c_9(s)$ depend only on the parameters s, p_s, r and ϵ such that

(i) if $d < p_s^{2c_8} r^2$, then $\#T_d \le c_9(s)$,

(11) if
$$d \ge p_s^{2c_8}r^2$$
, then $\#T_d \le 1$.

Note that Theorem 2.1 extends the corresponding statement from [10] to this more general situation. Our next theorem illustrates Theorem 2.1, in which we explicitly solve (11) for the case r = 1, s = 2 and $S = \{2, 3\}$.

4

Theorem 2.2. Let X_l be the X-coordinate of the Pell equation (7) with d > 1 non-square. Then there exist no d for which the equation

$$X_l = 2^{n_1} 3^{n_2}, \qquad n_1 \le n_2 \tag{13}$$

has at least two positive integer solutions.

3. Auxiliary results

In this section, we will prove some preliminary results to prove main theorems.

Lemma 3.1. Let $\gamma > 0$ be the fundamental solution of (7) for d > 1 non-square. Then

$$\left(\frac{1}{1+\sqrt{2}}\right)\gamma^{l} \leq X_{l} \leq (2-\sqrt{2})\gamma^{l}, \quad \text{for all } l \geq 1.$$
(14)

Proof. See Lemma 1 in [10].

Let α be an algebraic number of degree d. Then the *logarithmic height* of the algebraic number α is given by

$$h(\alpha) = \frac{1}{d} \left(\log |a| + \sum_{i=1}^{d} \log \max\{1, |\alpha^{(i)}|\} \right),$$

where a is the leading coefficient of the minimal polynomial of α over \mathbb{Z} and the $\alpha^{(i)}$'s are the conjugates of α in \mathbb{C} . In particular, if z = p/q is a rational number with gcd(p,q) = 1, and q > 0 then $h(z) = \log \max\{|p|, q\}$.

To prove Theorem 2.1, we use a lower bound of linear forms in logarithms to bound the index l, z_1, \ldots, z_s appearing in (6). In particular, we need the following result due to Matveev [17].

Lemma 3.2 (Matveev's Theorem). Let η_1, \ldots, η_t be real algebraic numbers and let d_1, \ldots, d_t be rational integers. Let $d_{\mathbb{L}}$ be the degree of the number field $\mathbb{L} = \mathbb{Q}(\eta_1, \ldots, \eta_t)$ over \mathbb{Q} . Let A_j be real numbers satisfying

 $A_j \ge \max \{ d_{\mathbb{L}} h(\eta_j), |\log \eta_j|, 0.16 \}, \quad j = 1, \dots, t.$

Assume that $B \ge \max\{|d_1|, \ldots, |d_t|\}$ and $\Lambda := \eta_1^{d_1} \cdots \eta_t^{d_t} - 1$. If $\Lambda \ne 0$, then

$$|\Lambda| \ge \exp\left(-1.4 \cdot 30^{t+3} \cdot t^{4.5} \cdot d_{\mathbb{L}}^2 (1 + \log d_{\mathbb{L}})(1 + \log B)A_1 \cdots A_t\right)$$

Lemma 3.3. Let B be a non-negative integer such that $\delta B \leq \alpha \log B + \beta$. If $\alpha \geq e\delta$, then

$$B \le (2/\delta)(\alpha \log(\alpha/\delta) + \beta).$$

Proof. See [6, Lemma 12.2.4]

We need some more results from the theory of continued fractions. The following result will be useful for the treatment of linear forms in logarithms.

Lemma 3.4. Let τ be an irrational number, M be a positive integer and $p_0/q_0, p_1/q_1, \ldots$, be all the convergents of the continued fraction $[a_0, a_1, \ldots]$ of τ . Let N be such that $q_N > M$. Then putting $a(M) := \max\{a_t : t = 0, 1, \ldots, N\}$, the inequality

$$|m\tau - n| > \frac{1}{(a(M) + 2)m}$$

holds for all pairs (n, m) of integers with 0 < m < M.

After getting the upper bound of n_2 , which is generally too large, the next step is to reduce it. For this reduction purpose, we use a variant of the Baker–Davenport result [1]. Here, for a real number x, let $||x|| := \min\{|x - n| : n \in \mathbb{Z}\}$ denote the distance from x to the nearest integer.

Lemma 3.5 (See [9]). Let τ be an irrational number, M be a positive integer and p/q be a convergent of the continued fraction of τ such that q > 6M. Let A, B, μ be some real numbers with A > 0 and B > 1. Put $\epsilon_1 := ||\mu q|| - M||\tau q||$, then there is no solution to the inequality

$$|m\tau - n + \mu| < AB^{-k},$$

in positive integers m, n and k with

$$m \le M$$
 and $k \ge \frac{\log(Aq/\epsilon_1)}{\log B}$.

4. Proof of Theorem 2.1

Suppose $1 \leq l_1 < l_2$, such that

$$X_{l_1} = p_1^{a_{11}} p_2^{a_{12}} \cdots p_s^{a_{1s}} + \dots + p_1^{a_{r1}} p_2^{a_{r2}} \cdots p_s^{a_{rs}}$$

and

$$X_{l_2} = p_1^{b_{11}} p_2^{b_{12}} \cdots p_s^{b_{1s}} + \dots + p_1^{b_{r1}} p_2^{b_{r2}} \cdots p_s^{b_{rs}}.$$
(15)

Denote

$$(l, n_{11}, \dots, n_{1s}, n_{21}, \dots, n_{2s}, \dots, n_{r1}, \dots, n_{rs}) = (l_1, a_{11}, \dots, a_{1s}, a_{21}, \dots, a_{2s}, \dots, a_{r1}, \dots, a_{rs})$$

or

$$(l, n_{11}, \dots, n_{1s}, n_{21}, \dots, n_{2s}, \dots, n_{r1}, \dots, n_{rs}) = (l_2, b_{11}, \dots, b_{1s}, b_{21}, \dots, b_{2s}, \dots, b_{r1}, \dots, b_{rs}).$$

By Lemma 3.1,

$$\frac{\gamma^l}{2.5} < X_l < \gamma^l. \tag{16}$$

From (11) and (12), it follows that,

$$p_s^{n_{rs}} < X_l < r p_s^{sn_{rs}}.$$
(17)

As $(X_l)_{l\geq 1}$ is an increasing sequence,

$$p_s^{a_{rs}} \le X_{l_1} \le X_{l_2} \le r p_s^{sb_{rs}}$$

and this implies

$$a_{rs} \le sb_{rs} + \frac{\log r}{\log p_s}.$$
(18)

From (16) and (17) we get

$$p_s^{n_{rs}} < \gamma^l < 2.5 X_l < 2.5 r p_s^{s n_{rs}},$$

and this further implies,

$$n_{rs}\log p_s < l\log\gamma < sn_{rs}\log p_s + \log 2.5r.$$
⁽¹⁹⁾

Thus,

$$\frac{n_{rs}}{\log\gamma} < \frac{l}{\log p_s} < l. \tag{20}$$

Since for $\epsilon > 0$, $|z_i|^{1+\epsilon} < |z_r|$, for i = 1, ..., r-1, then from (6) it follows that

$$X_l| = |z_1 + \dots + z_r| \le (r-1)|z_r|^{\frac{1}{1+\epsilon}} + |z_r| \le r|z_r|,$$
(21)

which implies

$$|z_r| \ge \frac{|X_l|}{r}.\tag{22}$$

If
$$|X_l| \leq \frac{\gamma^{(n_{rs}/\log \gamma)}}{4}$$
, then
$$\frac{1}{2}\gamma^l < |X_l| = \frac{1}{2}\gamma^l + \frac{1}{2}\eta^l \leq \frac{\gamma^{(n_{rs}/\log \gamma)}}{4} < \frac{\gamma^l}{4}$$

i.e.,

$$\frac{1}{2}\gamma^l < \frac{\gamma^l}{4},$$

which is not true. So assume that,

$$|X_l| > \frac{\gamma^{(n_{rs}/\log\gamma)}}{4}.$$
(23)

We rewrite (6) using (10) as

$$\frac{1}{2}\gamma^{l} - z_{r} = z_{1} + \dots + z_{r-1} - \frac{1}{2}\eta^{l}.$$
(24)

Dividing throughout by z_r ,

$$\left|2^{-1}\gamma^{l}z_{r}^{-1}-1\right| \leq \left|\frac{\sum_{i=1}^{r-1}z_{i}}{z_{r}}\right| + \left|\frac{\eta^{l}}{2z_{r}}\right|.$$
(25)

Next we will find bounds for each term in the right hand side of (25). From (22) and (23),

$$\frac{\sum_{i=1}^{r-1} |z_i|}{|z_r|} \le (r-1) \frac{|z_r|^{\frac{1}{1+\epsilon}}}{|z_r|} \le (r-1) \frac{1}{|z_r|^{\frac{\epsilon}{1+\epsilon}}} \le (r-1) \frac{(4r)^{\frac{\epsilon}{1+\epsilon}}}{\gamma^{\frac{(n_{rs}/\log\gamma)\epsilon}{1+\epsilon}}} \le c_1(s) \frac{1}{\gamma^{\frac{(n_{rs}/\log\gamma)\epsilon}{1+\epsilon}}},$$

where $c_1(s) = (r-1)(4r)^{\frac{\epsilon}{1+\epsilon}}$. To find a bound for $\left|\frac{\eta^l}{2z_r}\right|$,

$$\left|\frac{\eta^l}{2z_r}\right| < \frac{1}{2} \cdot \frac{4r\eta^l}{\gamma^{(n_{rs}/\log\gamma)}} < 2r \cdot \frac{\eta^l}{\gamma^{(n_{rs}/\log\gamma)}} < 2r \cdot \left(\frac{\eta}{\gamma}\right)^{n_{rs}/\log\gamma}$$

Substituting these bounds in (25),

$$\begin{aligned} \left| 2^{-1} \gamma^{l} z_{r}^{-1} - 1 \right| &\leq c_{1}(s) \frac{1}{\gamma^{\frac{(n_{rs}/\log\gamma)\epsilon}{1+\epsilon}}} + 2r \cdot \left(\frac{\eta}{\gamma}\right)^{n_{rs}/\log\gamma} \\ &\leq c_{2}(s) \cdot \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{n_{rs}/\log\gamma}. \end{aligned}$$

$$(26)$$

We know that, if $|x - 1| < \frac{1}{2}$, then $|\log x| < 2|x - 1|$. Set

$$\Lambda := 2^{-1} \gamma^l z_r^{-1} - 1 \quad \text{and} \quad \Gamma := -\log 2 + l \log \gamma - \log z_r.$$

Using the fact that $z_r = p_1^{n_{r1}} \cdots p_s^{n_{rs}}$,

$$|\Lambda| = \left|\frac{1}{2}\gamma^l p_1^{-n_{r1}} \cdots p_s^{-n_{rs}} - 1\right|.$$

If $|\Lambda| = 0$, then $\gamma^l = 2z_r$ is an integer, which is false for any $l \ge 1$. Hence $|\Lambda| \ne 0$. So applying Matveev's theorem (Lemma 3.2), with

 $(\eta_1, d_1) = (2, -1), \ (\eta_2, d_2) = (\gamma, l), \text{ and } (\eta_{i+2}, d_{i+2}) = (p_i, -n_{ri}) \text{ with } i = 1, \dots, s.$ Furthermore,

$$d_{\mathbb{L}} = 2, \ h(2) = \log 2, \quad h(\gamma) = \frac{1}{2} \log \gamma, \quad h(p_j) = \log p_j, \text{ where } j = 1, \dots, s.$$

We choose A_i 's as follows,

$$A_1 = 0.16,$$
 $A_2 = \log \gamma,$ $A_{i+2} = 2 \log p_i,$ with $i = 1, \dots, s.$

Since p_s is odd, without loss of generality we assume that $\min\{\gamma, n_{rs}\} \ge 2.5rp_s > 7$. From (19),

$$l < \frac{sn_{rs}\log p_s + \log 2.5r}{\log \gamma}.$$

Now if $\min\{\gamma, n_{rs}\} = \gamma$, then $2.5rp_s < \gamma$, which implies that, $\log p_s < \log \gamma - \log 2.5r$. Substituting this in above inequality,

$$l < \frac{sn_{rs}(\log \gamma - \log 2.5r) + \log 2.5r}{\log \gamma} < sn_{rs}.$$

If $\min\{\gamma, n_{rs}\} = n_{rs}$, then $n_{rs} < \gamma$, so as in the previous case $l < sn_{rs}$. Hence, in both cases $l < sn_{rs}$. So we can take $B = sn_{rs}$. Applying Matveev's theorem with t = s + 2,

$$\log |\Lambda| > -1.4 \cdot 30^{s+5} (s+2)^{4.5} 2^2 (1+\log 2) (1+\log(sn_{rs})) \cdot 0.16 (\log \gamma) (2\log p_1) \cdots (2\log p_s) > -c_3(s) (\log n_{rs}) (\log \gamma).$$
(27)

It is clear that, $\gamma^{\frac{\epsilon}{1+\epsilon}} < \gamma^2$ for $0 < \epsilon < 1$. Now taking logarithm on both sides of (26) and comparing with (27),

$$-c_3(s)(\log n_{rs})(\log \gamma) < \log c_2(s) - \left(\frac{\epsilon n_{rs}}{(1+\epsilon)\log \gamma}\right)\log \gamma$$

which further gives

$$n_{rs} < \left(\frac{1+\epsilon}{\epsilon}\right) \left(\log c_2(s) + c_3(s)(\log n_{rs})(\log \gamma)\right)$$

$$\leq c_4(s)(\log n_{rs})(\log \gamma).$$
(28)

This gives an upper bound for n_{rs} in terms of $\log \gamma$. For i = 1, 2 let

$$\Gamma_1^{(i)} := -\log 2 + l_i \log \gamma - \log(p_1^{m_{r1}} \cdots p_s^{m_{rs}})$$

= $-\log 2 + l_i \log \gamma - \sum_{k=1}^s m_{rk} \log p_k.$

Note that $m_{rk} = a_{rk}$ or b_{rk} for $k = 1, \ldots, s$. Next our aim is to eliminate term involving $\log \gamma$ from $\Gamma_1^{(i)}$. For that, consider

$$\Gamma_2 := l_1 \Gamma_1^{(2)} - l_2 \Gamma_1^{(1)} = (l_2 - l_1) \log 2 + (l_2 - l_1) \sum_{k=1}^s m_{rk} \log p_k.$$
(29)

Since $l_1 < l_2$, then from (26),

$$\begin{aligned} |\Gamma_{2}| &\leq l_{1} |\Gamma_{1}^{(2)}| + l_{2} |\Gamma_{1}^{(1)}| \\ &\leq l_{2} c_{2}(s) \cdot \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{b_{rs}/\log\gamma} + l_{2} c_{2}(s) \cdot \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{a_{rs}/\log\gamma} \\ &\leq l_{2} c_{2}(s) \cdot \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{a_{rs}/\log\gamma} \left(1 + \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{b_{rs}/\log\gamma - a_{rs}/\log\gamma}\right) \\ &\leq l_{2} c_{5}(s) \cdot \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{a_{rs}/\log\gamma} . \end{aligned}$$
(30)

Assume that, $c_5(s) \cdot l_2 \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{a_{rs}/\log \gamma} < 1/2$. Otherwise, we can derive a bound which is weaker than the bound with this assumption. Set

$$\Lambda_{2}| := |2^{(l_{2}-l_{1})} p_{1}^{n_{r1}(l_{2}-l_{1})} \cdots p_{s}^{n_{rs}(l_{2}-l_{1})} - 1|$$

= $|e^{\Gamma_{2}} - 1| < 2|\Gamma_{2}| < 2c_{5}(s) \cdot l_{2} \max\left\{\frac{1}{\gamma^{\frac{\epsilon}{1+\epsilon}}}, \frac{\eta}{\gamma}\right\}^{a_{rs}/\log\gamma}.$ (31)

We will apply Matveev's theorem for Λ_2 with

 $(\eta_1, d_1) := (2, l_2 - l_1), \qquad (\eta_{i+1}, d_{i+1}) := (p_i, n_{ri}(l_2 - l_1)), \text{ where } i = 1, \dots, s.$ Again, $d_{\mathbb{L}} = 1, h(2) = \log 2, \quad h(p_i) = \log p_i, \text{ with } i = 1, \dots, s$. Choose $A_1 = 2 \log 2, \qquad A_{i+1} = 2 \log p_i, \text{ with } i = 1, \dots, s$.

Note that $|n_{ri}(l_2 - l_1)| \le n_{rs}(l_2 + l_1)$ and $l_2 < sn_{rs}$. Here $B \ge \max_{1 \le i \le s} \{|l_2 - l_1|, |n_{ri}(l_2 - l_1)|\}$. So, we can set $B = 2s(n_{rs})^2$. Applying Matveev's theorem,

$$\log |\Lambda_2| > -1.4 \cdot 30^{s+4} \cdot (s+1)^{4.5} (1 + \log(2s(n_{rs})^2))(2\log 2)$$

(2 log p₁) · · · (2 log p_s)
> -c_6(s) log n_{rs}.

We have, $\gamma^{\frac{\epsilon}{1+\epsilon}} < \gamma^2$ for $0 < \epsilon < 1$. Now comparing the above inequality with (31), we get

$$-c_{6}(s)\log n_{rs} < \log(2l_{2}c_{5}(s)) - \frac{a_{rs}}{\log\gamma}\log \max\left\{\gamma^{\frac{\epsilon}{1+\epsilon}}, \gamma^{2}\right\}$$
$$\leq \log(2l_{2}c_{5}(s)) - \frac{a_{rs}}{\log\gamma}\log\gamma^{\frac{\epsilon}{1+\epsilon}}.$$

i.e.,

$$\frac{a_{rs}}{\log \gamma} \cdot \frac{\epsilon}{1+\epsilon} \log \gamma < c_6(s) \log n_{rs} + \log(2l_2c_5(s)),$$

which implies that,

$$a_{rs} < \left(\frac{1+\epsilon}{\epsilon}\right) \left(\log(2l_2c_5(s)) + c_6(s)\log n_{rs}\right).$$
(32)

Considering (28) for $n_{rs} = b_{rs}$,

 $b_{rs} < c_4(s)(\log b_{rs})(\log \gamma).$

By Lemma 3.3,

$$b_{rs} < 2(c_4(s)\log\gamma \cdot \log(c_4(s)\log\gamma)).$$
(33)

Using (19) for $n_{rs} = a_{rs}$ we get,

$$a_{rs}\log p_s < l_1\log \gamma < sa_{rs}\log p_s + \log 2.5r$$

and this implies,

$$\log \gamma < l_1 \log \gamma < sa_{rs} \log p_s + \log 2.5r. \tag{34}$$

Since $\sqrt{d-1} \le X_{l_1} \le r p_s^{sa_{rs}}$, so assume that $\log \gamma < 2a_{rs}s \log p_s$. Then from (33), $b_{rs} < 2(c_4(s)2a_{rs}s \log p_s \cdot \log(c_4(s)2a_{rs}s \log p_s)).$ (35)

Since $l_2 < sb_{rs}$, so from (32) and (35) we infer,

$$a_{rs} < \left(\frac{1+\epsilon}{\epsilon}\right) \left(\log(2sb_{rs}c_{5}(s)) + c_{6}(s)\log b_{rs}\right)$$

$$< \left(\frac{1+\epsilon}{\epsilon}\right) \left(\log(2s2(c_{4}(s)2a_{rs}s\log p_{s} \cdot \log(c_{4}(s)2a_{rs}s\log p_{s}))c_{5}(s))\right)$$

$$+ \left(\frac{1+\epsilon}{\epsilon}\right) \left(c_{6}(s)\log(2(c_{4}(s)2a_{rs}s\log p_{s} \cdot \log(c_{4}(s)2a_{rs}s\log p_{s})))\right).$$
mma 2.2

By Lemma 3.3,

$$a_{rs} < c_7(s)$$

Thus (34) implies that,

$$\log \gamma < sc_7(s) \log p_s + \log 2.5r,\tag{36}$$

where $c_7(s)$ is an effectively computable constant that depends only on s, p_s and ϵ . So,

$$\log \gamma < \log(p_s^{sc_7(s)}2.5r),$$

i.e.,

Since
$$\gamma > 2\sqrt{d-2}$$
, then
$$d < p_s^{2sc_7(s)} (2.5r)^2 < p_s^{2c_8(s)} r^2.$$

Also,

$$l_2 < b_{rs}s < s2(c_4(s)2c_7(s)s\log p_s \cdot \log(c_4(s)2c_7(s)s\log p_s)) =: c_9(s)$$

Thus, we conclude that there are only finitely many possibilities for d under the assumption that $\min\{\gamma, n_{rs}\} \ge 2.5rp_s$. If $\min\{\gamma, n_{rs}\} < 2.5rp_s$, then again we can obtain bounds on l, n_{rs} and d. In particular, if $n_{rs} = \min\{\gamma, n_{rs}\} < 2.5rp_s$, then by (19), we have upper bounds on $l, \log \gamma$ and then on d. However, if $\gamma = \min\{\gamma, n_{rs}\} < 2.5rp_s$, then there exists an upper bound on d and then on l and n_{rs} . This completes the proof of Theorem 2.1.

5. Proof of Theorem 2.2

Suppose that for $1 \leq l_1 < l_2$,

$$X_{l_1} = 2^{a_1} 3^{a_2}$$
 and $X_{l_2} = 2^{b_1} 3^{b_2}$, (37)

with $a_1 \leq a_2$ and $b_1 \leq b_2$. We denote

$$(l, n_1, n_2) = (l_1, a_1, a_2)$$
 or $(l, n_1, n_2) = (l_2, b_1, b_2).$

Assume $n_2 > n_1$. From the proof of Theorem 2.1,

$$3^{n_2} < X_l = 2^{n_1} 3^{n_2} < 3^{2n_2}. aga{38}$$

10

Also

$$3^{a_2} \le X_{l_1} \le X_{l_2} \le 3^{2b_2}$$

and this implies $a_2 \leq 2b_2$. From (38) and (16), we get $3^{n_2} < \gamma^l < 2.5 X_l < 2.5 \cdot 3^{2n_2}$. This gives,

$$n_2 \log 3 < l \log \gamma < 2n_2 \log 3 + \log 2.5. \tag{39}$$

Using (10), rewrite (13) as

$$\frac{\gamma^l}{2^{(n_1+1)}3^{n_2}} - 1 = \frac{\eta^l}{2^{(n_1+1)}3^{n_2}}$$

Setting $\Lambda := \gamma^{l} 2^{-(n_1+1)} 3^{-n_2} - 1$, we get

$$|\Lambda| \le \left| \frac{\eta^l}{2^{(n_1+1)} 3^{n_2}} \right| \le \frac{1}{3^{n_2}}.$$
(40)

Put

$$\Gamma = l \log \gamma - (n_1 + 1) \log 2 - n_2 \log 3 \tag{41}$$

and since $|\Lambda| = |e^{\Gamma} - 1| < \frac{1}{2}$, so $|\Gamma| < 2|\Lambda| < 2 \cdot 3^{-n_2}$. Since γ is not an integer, so $\Lambda \neq 0$. Hence we can use Matveev's theorem with t = 3 and

$$(\eta_1, d_1) = (2, -(n_1 + 1)),$$
 $(\eta_2, d_2) = (\gamma, l),$ $(\eta_3, d_3) = (3, -n_2).$

Further, $d_{\mathbb{L}} = 2, h(2) = \log 2, h(\gamma) = \frac{1}{2} \log \gamma, h(3) = \log 3, A_1 = 2 \log 2, A_2 = \log \gamma, A_3 = 2 \log 3$. Assume that $\min\{\gamma, n_2\} \ge 7.5$. Then one can see that $l < 2n_2$. Applying Matveev's theorem,

$$\log |\Lambda| > -c_{10}(\log n_2)(\log \gamma) \log 3,$$

where $c_{10} = 1.33 \cdot 10^{14}$. Comparing above inequality with (40), we get

$$n_2 < c_{11}(\log n_2) \log \gamma$$
, with $c_{11} = 1.34 \cdot 10^{14}$. (42)

To eliminate $\log \gamma$, let

$$\Gamma_1^{(i)} := -(m_1 + 1)\log 2 - m_2\log 3 + l_i\log\gamma,$$

where $m_i = a_i$ or b_i with i = 1, 2. Define

$$\Gamma_2 := l_1 \Gamma_1^{(2)} - l_2 \Gamma_1^{(1)} = (l_2 - l_1 + l_2 a_1 - l_1 b_1) \log 2 + (l_2 a_2 - l_1 b_2) \log 3.$$
(43)

Then proceeding as in the proof of Theorem 2.1, we get

$$|\Gamma_2| \le c_{12} \cdot l_2 3^{-\frac{a_2}{2}}$$
 with $c_{12} = 4$, (44)

 $b_2 < 2 \cdot 4c_{11}a_2 \log 3 \cdot \log(4c_{11}a_2 \log 3) < 8.62 \cdot 10^{28}$ and $l_2 < 2b_2 < 17.2 \cdot 10^{28}$. These bounds are very large for a computational search. Our next aim is to reduce these bounds to smaller numbers using Lemma 3.4 and 3.5. From (43) and (44),

$$\left|\frac{\log 2}{\log 3} - \frac{(l_1b_2 - l_2a_2)}{l_2(1+a_1) - l_1(1+b_1)}\right| \le \frac{c_{12}l_2}{3^{\frac{a_2}{2}} \cdot (l_2(1+a_1) - l_1(1+b_1))(\log 3)}$$

To use Lemma 3.4, put $\tau := \frac{\log 2}{\log 3}$ and $M_1 := 8.62 \cdot 10^{28}$. With the help of Mathematica finding q such that $q > M_1$ as $q_{60} > 8.62 \cdot 10^{28}$ and $a(M_1) = 55$. Now applying Lemma 3.4,

$$3^{\frac{a_2}{2}} < \frac{c_{12}l_2(l_2(1+a_1)-l_1(1+b_1))57}{\log 3} < 1824 \cdot M_1^3,$$

which implies that,

$$a_2 < \frac{2\log(1824 \cdot M_1^3)}{\log 3} < 377.55$$

i.e., $a_2 \leq 377$ and this gives $b_2 < 1.78 \cdot 10^{19}$. To get better bounds repeat this process with new *M* value. Next put $M_2 = 1.78 \cdot 10^{19}$. Finding the value of *q* as, $q_{41} > 1.78 \cdot 10^{19}$ and $a(M_2) = 55$. Hence $a_2 \leq 255$ and $b_2 < 1.2 \cdot 10^{19}$. Using Lemma 3.4 again with $M_3 = 1.2 \cdot 10^{19}$ to get $a_2 \leq 253$. This way will not give any better bounds further. So we will use another method.

Consider the following polynomial with integer coefficients:

$$X_{l} = \frac{1}{2} \left(\gamma^{l} + \eta^{l} \right) = \frac{1}{2} \left((X_{1} + Y_{1}\sqrt{d})^{l} + (X_{l} - Y_{l}\sqrt{d})^{l} \right)$$
$$= \frac{1}{2} \left(\left(X_{1} + \sqrt{X_{1}^{2} - 1} \right)^{l} + \left(X_{1} - \sqrt{X_{1}^{2} - 1} \right)^{l} \right) := P_{l}(X_{1}).$$

Now from (39) finding a bound for l_1 as,

$$l_1 < \frac{(2 \cdot 253)\log 3 + \log 2.5}{\log \gamma}.$$
(45)

We know that $X_1 \ge \sqrt{d-1}$ and $\gamma > 2\sqrt{d-2}$. Assume that d > 401, which gives $X_1 > 20$ and $l_1 \le 150$. Now from (37),

$$P_{l_1}(X_1) := 2^{a_1} 3^{a_2} \tag{46}$$

with $a_2 \in [0, 253]$, $a_1 \in [0, a_2]$, $l_1 \in [2, 150]$ and $X_1 > 20$. Note that $l_1 \ge 2$, because if we are taking $l_1 = 1$ we get $P_{l_1}(X_1) = X_1$. In this case there is nothing to solve.

For the case $l_1 > 1$ and d > 401, a quick computer search on (46) yields nothing; so if $l_1 > 1$ then $1 < d \le 401$. In this case also, we are not getting any solutions.

Since l_1 to be minimal, we must have $l_1 = 1$ for all d. Also $\gamma = X_1 + \sqrt{X_1^2 - 1}$. From (41),

$$|l\log\gamma - (n_1+1)\log 2 - n_2\log 3| < \frac{2}{3^{n_2}}$$

Rearranging and comparing with Lemma 3.5,

$$m := n_2, \tau := \frac{\log 3}{\log \gamma}, \mu := \frac{254 \cdot \log 2}{\log \gamma}, A := \frac{2}{\log \gamma}, B := 3, k := n_2.$$

We have an upper bound for b_2 , which is $M_4 := 1.179 \cdot 10^{19}$. Reduce this bound again and denote the reduced bound by $M_5(X_1)$ for each X_1 . Using Mathematica we get, max $M_5(X_1) = 52$. The X-coordinates of (7) satisfies the recurrence relation,

$$X_l = 2X_1 X_{l-1} - X_{l-2}$$

and for each X_1 a bound for l_2 is obtained using (39) denoting it by $l(X_1)$:

$$l_2 < \frac{2 \cdot M_5(X_1) \log 3 + \log 2.5}{\log \gamma} := l(X_1).$$

So, $\max_{X_1} l(X_1) = 130$. Now finding the minimum of $\epsilon_1(X_1)$ of Lemma 3.5 as $\min_{X_1} \epsilon_1(X_1) = \mathbf{v}$

0.0011. Now it remains to verify whether $\frac{X_{l_2}}{3^{b_2}}$ is of the form 2^{b_1} . Checking this for each value of X_1 in the range

$$l_2 \in [2, 130]$$
 and $b_2 \in [0, 52]$

using Mathematica, we find no solutions. This completes the proof of the Theorem 2.2.

Acknowledgment: First author's research is supported by UGC Fellowship (Ref No. 221610077314). S.S.R. work is supported by grant from Anusandhan National Research Foundation (ANRF) (File No.:CRG/2022/000268) and from National Board for Higher Mathematics (NBHM), Sanction Order No: 14053.

References

- [1] A. Baker and H. Davenport, The equations $3x^2 2 = y^2$ and $8x^2 7 = z^2$, Quart. J. Math. Oxford Ser. **20** (1969), 129–137.
- [2] A. Bérczes, L. Hajdu, I. Pink and S. S. Rout, Sums of S-units in recurrence sequences, J. Number Theory 196 (2019), 353-363.
- [3] Cs. Bertók, L. Hajdu, I. Pink and Zs. Rábai, *Linear combinations of prime powers in binary recurrence sequences*, Int. J. Number Theory **13** (2017), 261–271.
- [4] K. Bir, F. Luca and A. Togbé, On the X-coordinates of Pell equations which are Fibonacci numbers II, Colloq. Math. 149 (2018), 75-85.
- [5] Y. Bugeaud, M. Mignotte and S. Siksek, Classical and modular approaches to exponential Diophantine equations I. Fibonacci and Lucas perfect powers, Ann Math. 163 (2006), 969–1018.
- [6] Cohen, H. Number Theory. Volume II: Analytic and Modern Tools. Graduate Texts in Mathematics. Springer, New York (2007).
- [7] J. H. E. Cohn, Perfect Pell powers, Glasg. Math. J. 38 (1996), 19–20.
- [8] A. Dossavi-Yovo, F. Luca and A. Togbé, On the x-coordinates of Pell equations which are repdigits, Publ. Math. Debrecen 88 (2016), 381-399.
- [9] A. Dujella and A. Pethő, A generalization of a theorem of Baker and Davenport, Quart. J. Math. Oxford Ser. 49 (1998), 291–306.
- [10] H. Erazo, C. A. Gómez, and F. Luca, Linear combinations of prime powers in X-coordinates of Pell equations, Ramanujan J. 53 (2020), 123-137.
- [11] B. Faye and F. Luca, On the X-coordinates of Pell equations which are rep-digits, Fibonacci Q. 56 (2018), 52-62.
- [12] C. A. Gómez, F. Luca and F. S. Zottor, On X-coordinates of Pell equations which are rep-digits, Res. Number Theory 6(4) (2020), paper no. 41.

- [13] L. Hajdu and P. Sebestyén, Sums of S-units in the solution sets of generalized Pell equations, Arch. Math.115 (2020), 279-287.
- [14] F. Luca, A. Montejano, L. Szalay and A. Togbé, On the X-coordinates of Pell equations which are tribonacci numbers, Acta Arith. 179 (2017), 25-35.
- [15] F. Luca and A. Togbé, On the X-coordinates of Pell equations which are Fibonacci numbers, Math. Scand. 122 (2018), 18-30.
- [16] D. Marques and A Togbé, Fibonacci and Lucas numbers of the form 2^a + 3^b + 5^c, Proc. Japan Acad. Ser. A Math. Sci. 89 (2013), 47 - 50.
- [17] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in the logarithms of algebraic numbers, Izv. Math. 64 (2000), 1217–1269.
- [18] A. Pethő, Perfect powers in second order linear recurrences, J. Number Theory 15 (1982), 5-13.
- [19] A. Pethő, The Pell sequence contains only trivial perfect powers, Coll. Math. Soc. J. Bolyai, 60 sets, Graphs and Numbers, Budapest (1991), 561-568.
- [20] A. Pethő and R. F. Tichy, S-unit equations, linear recurrences and digit expansions, Publ. Math. Debrecen 42 (1993), 145-154.
- [21] T. N. Shorey and C. L. Stewart, On the Diophantine equation $ax^{2t} + bx^ty + cy^2 = d$ and pure powers in recurrence sequences, Math. Scand. **52** (1983), 24-36.

PARVATHI S NAIR, DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY CALICUT, KOZHIKODE-673 601, INDIA.

Email address: parvathisnair60@gmail.com; parvathi_p220245ma@nitc.ac.in

SUDHANSU SEKHAR ROUT, DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOL-OGY CALICUT, KOZHIKODE-673 601, INDIA.

Email address: sudhansu@nitc.ac.in; lbs.sudhansu@gmail.com