
Infinite Width Limits of Self Supervised Neural Networks

Maximilian Fleissner
Technical University of Munich

Gautham Govind Anil
IIT Madras

Debaghya Ghoshdastidar
Technical University of Munich

Abstract

The NTK is a widely used tool in the theoretical analysis of deep learning, allowing us to look at supervised deep neural networks through the lenses of kernel regression. Recently, several works have investigated kernel models for self-supervised learning, hypothesizing that these also shed light on the behaviour of wide neural networks by virtue of the NTK. However, it remains an open question to what extent this connection is mathematically sound — it is a commonly encountered misbelief that the kernel behaviour of wide neural networks emerges irrespective of the loss function it is trained on. In this paper, we bridge the gap between the NTK and self-supervised learning, focusing on two-layer neural networks trained under the Barlow Twins loss. We prove that the NTK of Barlow Twins indeed becomes constant as the width of the network approaches infinity. Our analysis technique is different from previous works on the NTK and may be of independent interest. Overall, our work provides a first rigorous justification for the use of classic kernel theory to understand self-supervised learning of wide neural networks. Building on this result, we derive generalization error bounds for kernelized Barlow Twins and connect them to neural networks of finite width.

1 Introduction

In recent years, self-supervised learning (SSL) has emerged as a powerful paradigm, building the foundation of several modern machine learning models. At

its core, SSL relies on the idea of using augmentations to encode a notion of similarity in otherwise unlabeled data. As a typical example, consider an image dataset. Even though labels may not be known, it is reasonable to believe that randomly cropping, slightly rotating, or blurring the images will not change the true underlying class information. Therefore, two augmented versions (x, x^+) of the same image should receive similar representations $f(x), f(x^+)$ in a lower-dimensional ambient space. This constitutes the basic intuition behind non-contrastive SSL, and several loss functions that capture this idea have emerged.¹ Among the most popular losses is Barlow Twins (Zbontar et al., 2021), a loss function that pushes the cross-correlation of the embeddings $f(x), f(x^+)$ towards the identity matrix. This aims to prevent a phenomenon known as dimension collapse, where the learned representations collapse to a single point in the embedding space.

Despite the empirical success of SSL on a range of tasks (Radford et al., 2021; Bachman et al., 2019), it has taken quite some time for deep learning theory to catch up with this innovation. Arguably one of the most promising avenues towards understanding the fundamental principles of SSL is by connecting it to kernel methods (Smola and Schölkopf, 1998). This of course is reminiscent of the supervised setting, where the NTK (Jacot et al., 2018; Lee et al., 2019) provides a powerful framework to understand several phenomena of deep learning with wide neural networks, including generalization (Simon et al., 2021), benign overfitting (Mallinar et al., 2022), and robustness (Bombari et al., 2023). While a significant number of researchers are currently looking at SSL from a kernel perspective (for an overview, see related works), and while almost all of these works are motivated with the NTK, the connection is left implicit. However, it is not a priori clear that neural networks trained under SSL actually behave like kernel machines in the infinite width limit. In fact, Anil et al. (2024) have recently demonstrated that for certain contrastive loss functions, the NTK is in fact **not** constant at infinite width. This casts a

¹In contrastive SSL, one additionally incorporates a notion of dissimilarity into the model by including negative examples x^- into the training procedure.

shadow of doubt on the validity of kernel approximations to SSL, urging us to take a closer look at the matter.

In this paper, we bridge the gap between SSL and the NTK for the Barlow Twins loss. We prove that for neural networks with one hidden layer, the neural tangent kernel indeed becomes constant as the width of the network approaches infinity. The proof technique is different from previous works on the NTK, and leverages Grönwall’s inequality (see Appendix F.1). This is necessitated by the training dynamics of the Barlow Twins loss, which make an extension of existing methods difficult. Our work confirms the hypothesized connection between kernel methods and neural network based SSL, and proves that Barlow Twins is akin to one of the most prominent representation learning methods, Kernel PCA (Schölkopf et al., 1997). Building on these insights, we use classic tools from learning theory to derive generalization error bounds for kernel versions of Barlow Twins, and then connect them to neural networks of finite width.

This paper is structured as follows. We discuss related works in Section 2 and present our formal setup in Section 3. Section 4 contains our main result, stated in the general setting of multi-dimensional embeddings $f : \mathbb{R}^d \rightarrow \mathbb{R}^K$ for the Barlow twins loss. In Section 5 we give a detailed proof sketch of the one-dimensional case, which is computationally less involved but contains the main technical ideas. Section 6 derives generalization error bounds for the Barlow twins loss in an abstract Hilbert space setting. Using the kernel trick and our newly established validity of the NTK approximation, we then relate these bounds to finite neural networks. Finally, we empirically verify our findings through experiments in Section 7.

2 Related Work

In comparison to supervised learning, the theoretical understanding of self-supervised learning is still at an early stage. Nonetheless, several works have investigated SSL using classic tools from statistical learning theory (Arora et al., 2019; Wei et al., 2020). Furthermore, there have been several successful attempts to look at SSL through the lenses of classic spectral and kernel methods (Kiani et al., 2022; Johnson et al., 2022; HaoChen et al., 2021; Cabannes et al., 2023; Esser et al., 2024). These works provide a number of useful insights into both theoretical as well as practical aspects of training SSL models, but leave the formal connection between the kernel regime and deep learning based SSL implicit.

The idea that neural networks behave like (neural tangent) kernel models in the infinite width limit was first

investigated by Jacot et al. (2018). Several later works further explored this connection in various contexts (Arora et al., 2019; Chizat et al., 2019; Lee et al., 2019; Liu et al., 2020a), mostly for the squared (or hinge) loss. In particular, Liu et al. (2020b) develop a general framework for looking at convergence to the NTK using the Hessian. The Barlow Twins loss does not fall under the umbrella of previous analysis, since it is a fourth order loss with very different training dynamics.

Thus, even though Simon et al. (2023) motivate their investigation of kernel versions of Barlow Twins precisely with the NTK, we are not aware of any derivation that proves this analogy is valid. Ziyin et al. (2022) investigate the landscape of several SSL losses, but their theory is stated in the linear setting. Closest to our work is Anil et al. (2024), who manage to bound the evolution of the NTK for certain contrastive loss functions, but do not quite prove constancy until convergence of the loss: Their bounds hold until a time that grows with network width, leaving the possibility that as wider networks are trained, the time till convergence also grows.

3 Formal Setup

In this work, we analyze an idealized version of the popular Barlow Twins loss (Zbontar et al., 2021), as considered in previous theoretical works on SSL (Simon et al., 2023). Training data consists of positive pairs, denoted $\mathcal{X} = \{x_1, \dots, x_N\} \subset \mathbb{R}^d$ and $\mathcal{X}_+ = \{x_1^+, \dots, x_N^+\} \subset \mathbb{R}^d$. We assume data lies on the unit sphere, that is $\|x_n\| = \|x_n^+\| = 1$ for all $n \in [N]$. For example, each pair (x_n, x_n^+) could consist of two (normalized) augmentations of the same underlying image, randomly cropped or blurred. The goal of Barlow Twins is now to learn the parameters of a neural network $f : \mathbb{R}^d \rightarrow \mathbb{R}^K$ that embeds each point into a lower-dimensional space. Typically, $K \ll d$. To encourage useful representations, the Barlow Twins loss function pushes the cross-moment matrix of $f(\mathcal{X})$ and $f(\mathcal{X}_+)$ to the identity matrix in \mathbb{R}^K . Defining

$$C = \frac{1}{2N} \sum_{n=1}^N (f(x_n)f(x_n^+)^T + f(x_n^+)f(x_n)^T) \in \mathbb{R}^{K \times K} \quad (1)$$

we minimize the loss function $\mathcal{L}(f) = \|C - I\|_F^2$ over the parameters of a neural network f . In this work, we restrict our analysis to two-layer neural networks, that is

$$f(x) = \frac{1}{\sqrt{M}} \sum_{m=1}^M w_m \phi(v_m^T x) \quad (2)$$

where $w_m \in \mathbb{R}^K$ and $v_m \in \mathbb{R}^d$ for all $m \in [M]$, and ϕ is a bounded, smooth activation function, and has bounded first derivative. For example, we could have $\phi(t) = \tanh(t)$. The weights are initialized as random independent Gaussians with constant variance, and collected in a vector $\theta \in \mathbb{R}^{M(d+K)}$ that is trained under gradient flow

$$\frac{\partial \theta}{\partial t} = \dot{\theta}(t) = -\frac{\partial \mathcal{L}}{\partial \theta} \quad (3)$$

We will later also discuss the ReLU activation, whose non-differentiability prevents us from formally defining gradient flow. The neural tangent kernel is defined as a time-varying, matrix-valued map $K_t : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^{K \times K}$, where

$$K_t(x, x') = \left(\left(\frac{\partial f_k(x)}{\partial \theta(t)} \right)^T \left(\frac{\partial f_l(x')}{\partial \theta(t)} \right) \right)_{k,l=1}^K \quad (4)$$

for all $x, x' \in \mathbb{R}^d$. To underline the dependence of the NTK on the parameters $\theta(t)$ that evolve during training, we sometimes also denote it as K_θ . The key insight of the NTK literature is that the NTK does not change during training if the width of the neural network approaches infinity. Consequently, the training dynamics of f approach those of kernel regression with respect to the (vector-valued) kernel at initialization K_0 . In general, the constancy of the NTK in the infinite width limit is essentially a consequence of two things: Firstly, the Hessian of the network scales as $\mathcal{O}(\frac{1}{\sqrt{M}})$, and secondly, the weights θ remain in a ball of bounded, width-independent radius R around θ_0 until convergence. Provided both of these conditions hold, one may then rely on the following result by Liu et al. (2020a).

Theorem 3.1. *Consider an initial parameter $\theta_0 \in \mathbb{R}^p$ and a ball $B(\theta_0, R)$ around θ_0 with radius $R > 0$. If the spectral norm of the Hessian of f satisfies $\|H(\theta)\|_2 = \mathcal{O}(\frac{1}{\sqrt{M}})$ for all $\theta \in B(\theta_0, R)$, then the neural tangent kernel satisfies*

$$|K_{\theta_0}(a, b) - K_\theta(a, b)| = \mathcal{O}(R/\sqrt{M}) \quad (5)$$

for all $\theta \in B(\theta_0, R)$ and all inputs $a, b \in \mathbb{R}^d$.

The neural networks we consider in the context of Barlow Twins directly inherit the condition $\|H(\theta)\|_2 = \mathcal{O}(\frac{1}{\sqrt{M}})$ from previous works on the NTK (Liu et al., 2020a). Indeed, the scaling of the Hessian in parameter space is independent of the the loss function that governs the training dynamics, and we therefore do not go into details here. But unless R stays independent of M , this by itself does not imply constancy of the NTK at infinite width. For the squared error, Liu et al. (2020b) use the PL^* condition to ensure that the

weights remain in a bounded ball of width-independent radius R . It is not obvious how to extend this idea to Barlow Twins loss. In fact, a direct extension of Liu et al. (2020b) does not work with Barlow Twins loss, since R depends on the learned representations as well. In this paper, we therefore take a different approach, and look at the evolution of the loss and the parameters *in time*.

4 Main Result

In light of the aforementioned discussion, we focus on proving that the weights of the network stay in a ball of fixed radius R , with high probability over random initialization. To this end, we fix a global constant $\delta > 0$, below which the loss is considered to be zero, and training is stopped. Our strategy is to verify the following two statements, both of which hold at large width M .

1. For any finite time $T > 0$, there exists a width-independent $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}(t)\| \leq \kappa$ with high probability for any network of sufficiently large width.
2. There exists a finite time T such that $\mathcal{L}(T) \leq \delta$ for any network of sufficiently large width.

Together, both imply that for large enough M , the weights θ remain within a ball of width-independent radius R around the initial θ_0 until T , with high probability. The reason is the following: Writing $R(t) = \|\theta(t) - \theta(0)\|^2$, we obtain

$$\begin{aligned} \frac{\partial}{\partial t} (R(t) + 1) &= 2 \sum_{m=1}^M (\theta_m(t) - \theta_m(0)) \dot{\theta}_m(t) \\ &\leq 2 \|\theta(t) - \theta(0)\| \|\dot{\theta}(t)\| \\ &= 2\sqrt{R(t)} \|\dot{\theta}(t)\| \\ &\leq 2\kappa(R(t) + 1) \end{aligned} \quad (6)$$

Since $R(0) = 0$, Grönwall's inequality (see Lemma F.1) implies that $R(t) + 1 \leq \exp(2\kappa t)$ for all $t \leq T$. Thus, the weights θ remain in a ball of radius $R = \exp(2\kappa T)$ until convergence.

Theorem 4.1. (Gradient of weights is bounded) *Fix any $T \geq 0$ and any $\epsilon > 0$. Then, there exists $M_0 \in \mathbb{N}$ and some $\kappa > 0$ such that, for all networks of width $M > M_0$, with probability at least $1 - \epsilon$, the weights θ satisfy $\|\dot{\theta}(t)\| \leq \kappa$ for all $t \leq T$ when trained under gradient flow.*

The proof is included in Appendix A.1. The statement is probabilistic because it only holds if the weights are not too large at initialization (which is certainly true

with high probability). It remains to show that for some sufficiently large width M , the convergence of the loss indeed happens within some finite time T , up to our fixed tolerance $\delta > 0$. We therefore turn to the evolution of $\mathcal{L}(t)$ over time. Defining

$$u(t) = \begin{bmatrix} \left(\frac{\partial \mathcal{L}}{\partial f_k(x_n)}\right)_{n,k} \\ \left(\frac{\partial \mathcal{L}}{\partial f_k(x_n^+)}\right)_{n,k} \end{bmatrix} \in \mathbb{R}^{2NK}$$

$$\mathbf{K}(t) = \begin{bmatrix} K_t(x_1, x_1) & \dots & K_t(x_1, x_N^+) \\ \dots & \dots & \dots \\ K_t(x_N^+, x_1) & \dots & K_t(x_N^+, x_N^+) \end{bmatrix} \in \mathbb{R}^{2NK \times 2NK} \quad (7)$$

we express the time evolution of the Barlow Twins loss in a more concise manner.

Lemma 4.2. (Time evolution of the loss) *Under gradient flow, the evolution of $\mathcal{L}(t)$ can be expressed as*

$$\frac{\partial}{\partial t} \mathcal{L}(t) = -u(t)^T \mathbf{K}(t) u(t) \quad (8)$$

The proof is included in Appendix A.2. Our main theorem requires an assumption on the loss at initialization.

Definition 4.3. (Definitions and Assumptions at initialization) *We assume that, under the initialization that ensures Theorem 4.1 to hold, the matrix $\mathbf{K}(0)$ is positive definite, with smallest eigenvalue $\lambda_{\min}(\mathbf{K}(0)) \geq \lambda > 0$, and that $\mathcal{L}(0) < 1$. Define*

$$\eta = \frac{4\lambda(1 - \sqrt{\mathcal{L}(0)})}{N} \quad (9)$$

and choose T such that the solution to the autonomous ODE $\frac{\partial}{\partial t} L = -\eta L$ satisfies $L(T) \leq \delta$, under the initial condition $L(0) = \mathcal{L}(0)$.

We state the main result.

Theorem 4.4. (Convergence of the loss in finite time) *Under the setting in Definition 4.3, there exists $M_1 \in \mathbb{N}$ such that, for all networks of width $M \geq M_1$, the loss under gradient flow satisfies $\frac{\partial}{\partial t} \mathcal{L}(t) < -\eta \mathcal{L}(t)$ for all time $t \leq T$. This implies that $\mathcal{L}(T) < \delta$.*

The proof of Theorem 4.4 is included in Appendix A.3. As a direct consequence of this result and of Theorem 3.1, we obtain the following corollary.

Corollary 4.5. (Convergence of the NTK) *Under the conditions of Theorem 4.4, there exists a radius $R > 0$ such that, with probability at least $1 - \epsilon$, the change of the NTK until convergence is $\mathcal{O}(R/\sqrt{M})$, with R depending only on δ, η , but independent of the network width.*

For the ReLU activation $\phi(x) = \max(0, x)$, gradient flow itself is ill-defined, due to the non-differentiability of the ReLU at zero. However, if we define the weak derivative $\frac{\partial \phi(0)}{\partial x} = 0$, and equate $\dot{\theta}(t) = -\frac{\partial \mathcal{L}}{\partial \theta}$ as per usual, then it is possible to prove Theorem 4.1 nonetheless, and all other results remain true as well. Of course, this is not entirely rigorous, because $\dot{\theta}(t) = -\frac{\partial \mathcal{L}}{\partial \theta}$ is not actually gradient flow. See Appendix C for details.

Remark 4.6. (Dropping the assumption on the loss) *The condition $\mathcal{L}(0) < 1$ is most likely not necessary. In Appendix E, we prove that in the linearized regime, the loss converges exponentially to zero if all eigenvalues of the embedding cross-moment matrix $C(0)$ are contained in $(0, 1)$. This suggests that “small” initialization is sufficient to enter the kernel regime at large width. Our experiments also do not impose it, and yet show convergence to the NTK (see Section 7). For wide ReLU networks, $\mathcal{L}(0) < 1$ can however be guaranteed by suitably scaling the Gaussian weights $v_m \in \mathbb{R}^d$ of the first layer by a data-dependent constant (see Appendix A.4).*

5 Proof Sketch for One Dimensional Embeddings

To give a better intuition on our proof strategy, we present a more detailed proof sketch for the simplest possible case of one-dimensional embeddings $f : \mathbb{R}^d \rightarrow \mathbb{R}$. We consider a neural network with one hidden layer and d -dimensional inputs x . For one-dimensional embeddings, the Barlow Twins loss function equates to

$$\mathcal{L}(f) = \left(\frac{1}{N} \sum_{n=1}^N f(x_n) f(x_n^+) - 1 \right)^2 \quad (10)$$

For ease of notation we define the cross-moment matrix of the embeddings

$$C = \frac{1}{N} \sum_{n=1}^N f(x_n) f(x_n^+) \quad (11)$$

As discussed in Section 4, we aim to verify that at sufficiently large width (i) there exists a width-independent $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}(t)\| \leq \kappa$ with high probability over random initialization of weights and (ii) $\mathcal{L}(T) \leq \delta$ at a finite time T independent of M . We begin by checking the first condition.

Theorem 5.1. (Gradient of weights is bounded) *Fix any $T \geq 0$ and any $\epsilon > 0$. Then, there exists $M_0 \in \mathbb{N}$ and some $\kappa > 0$ such that, for all networks of width $M > M_0$, with probability at least $1 - \epsilon$, the weights θ satisfy $\|\dot{\theta}(t)\| \leq \kappa$ for all $t \leq T$ when trained under gradient flow.*

The proof is included in Appendix B.1. For simplicity, the proof assumes $\mathcal{L}(0) < 1$, although this is not necessary here yet. Essentially, the proof proceeds in three steps.

1. Firstly, we show that $\frac{\partial}{\partial t} \left(\sum_{m=1}^M w_m^2(t) \right) \leq 8$ for all $t \leq T$. This implies that there exists some $\kappa_1 > 0$ such that for all $t \leq T$, we have $\frac{1}{M} \left(\sum_{m=1}^M w_m^2(t) \right) \leq \kappa_1$ with probability $\geq 1 - \epsilon$.

2. From there, we bound the maximum squared value that any representation takes until time T , that is

$$|f|^2 = \max_{n \in [N]} \sup_{t \leq T} \max (|f(x_n)|^2, |f(x_n^+)|^2) \quad (12)$$

by some $\kappa_2 > 0$ (again independent of M). Doing so requires the first part of the proof as well as Grönwall's inequality.

3. Finally, we show that

$$\|\dot{\theta}(t)\|^2 \leq \frac{16|f|^2}{M} \left(c_\phi^2 + dc_{\phi'}^2 \left(\sum_{m=1}^M w_m^2 \right) \right) \quad (13)$$

for all $t \leq T$. Combining this with the first and second part we conclude that there indeed exists $\kappa > 0$ such that

$$\sup_{t \leq T} \|\dot{\theta}(t)\| \leq \kappa \quad (14)$$

with probability $\geq 1 - \epsilon$.

With Theorem 5.1 established, it remains to prove convergence of the loss in finite time, for wide neural networks. To this end, we derive the evolution of $\mathcal{L}(t)$.

Lemma 5.2. (Time evolution of the loss) *The loss $\mathcal{L}(t)$ evolves over time as*

$$\frac{\partial}{\partial t} \mathcal{L}(t) = -\frac{4\mathcal{L}(t)}{N^2} \cdot \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T \mathbf{K}(t) \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \quad (15)$$

where we define

$$\mathbf{K}(t) = \begin{pmatrix} K_\theta(\mathcal{X}_+, \mathcal{X}_+) & K_\theta(\mathcal{X}_+, \mathcal{X}) \\ K_\theta(\mathcal{X}, \mathcal{X}_+) & K_\theta(\mathcal{X}, \mathcal{X}) \end{pmatrix} \quad (16)$$

which is just the kernel matrix of $(\mathcal{X}_+, \mathcal{X})$ at $\theta(t)$.

The proof is included in Appendix B.2. This is a non-autonomous ODE, linear in \mathcal{L} but multiplied with a time-dependent scalar function

$$g(t) = \frac{4}{N^2} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T \mathbf{K}(t) \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \quad (17)$$

on the right side. Suppose now that we can find a constant $\eta > 0$ such that $\forall t \leq T : g(t) > \eta$. Then, the solution to the non-autonomous ODE can be upper bounded by the solution to the autonomous ODE

$$\frac{\partial}{\partial t} L = -\eta L \quad (18)$$

which converges to zero up to our previously fixed $\delta > 0$ up to T , by choice of T . To ensure existence of such η constitutes the proof idea.

Theorem 5.3. (Convergence of the loss in finite time) *Under the setting in Definition 4.3, there exists $M_1 \in \mathbb{N}$ such that, for all networks of width $M \geq M_1$, the loss under gradient flow satisfies $\frac{\partial}{\partial t} \mathcal{L}(t) < -\eta \mathcal{L}(t)$ for all time $t \leq T$, by virtue of $g(t) > \eta$ for all $t \leq T$. This implies that $\mathcal{L}(T) < \delta$.*

The proof is included in Appendix B.3. Essentially, it relies on two observations: Firstly, the kernel matrix $\mathbf{K}(t)$ does not change much up to time T , provided M is large. This is due to Theorem 5.1, which guarantees that the weights remain in a bounded ball of radius R around their initialization (until time T), and Theorem 3.1, which then bounds the change in NTK as $\mathcal{O}(R/\sqrt{M})$. Therefore, the smallest eigenvalue of $\mathbf{K}(t)$ stays larger than $\lambda/2$ when M is sufficiently large. Thus,

$$\begin{aligned} g(t) &> \frac{2\lambda}{N^2} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \\ &\geq \frac{2\lambda}{N^2} \left(\begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right)^T \left(\begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \right) \end{aligned} \quad (19)$$

for all $t \leq T$, where we used Cauchy-Schwartz inequality. This expression can in turn be bounded from below, due to the non-increasing nature of the loss \mathcal{L} under gradient flow, which forces $C(t)$ to stay safely bounded from below by $1 - \sqrt{\mathcal{L}(0)}$.

6 Implications for the Generalization Error of Barlow Twins

Having established the validity of the NTK approximation for neural networks trained under the Barlow Twins loss, we leverage our newly found connection into generalization error bounds for (finite) neural networks. We wish to bound $\tilde{\mathcal{L}}(f) = \|\bar{C}(f) - I\|_F^2$ where the population cross-moment matrix is

$$\bar{C}(f) = \mathbb{E}_{x, x^+} \left[\frac{f(x)f(x^+)^T + f(x^+)f(x)^T}{2} \right] \quad (20)$$

Here, f denotes a neural network with M neurons, and p weights collected in a vector $\theta \in \mathbb{R}^p$. Whenever M is

large enough to ensure Corollary 4.5 kicks in, we know that

$$f_k(x, \theta_T) = f_k(x; \theta_0) + \langle \theta_T - \theta_0, \nabla f_k(x, \theta_0) \rangle + \mathcal{O}(R/\sqrt{M}) \quad (21)$$

with probability at least $1 - \epsilon$, where R is the radius of the ball within which the weights evolve until convergence (with high probability over random initialization). Therefore, under small $f_k(x, \theta_0)$, the trained neural network $f_k(x, \theta_T)$ can be approximated by

$$g_k(x) = \langle \theta_T - \theta_0, \nabla f_k(x, \theta_0) \rangle \quad (22)$$

up to some small positive $\zeta = f_k(x; \theta_0) + \mathcal{O}(R/\sqrt{M})$. Note that each g_k is a kernel model w.r.t the feature map $\psi_k(x) = \nabla f_k(x, \theta_0)$. In this section, we therefore derive generalization error bounds for kernel versions of Barlow Twins.

Generalization error for Barlow Twins in Hilbert spaces. We first place ourselves in an abstract Hilbert space framework. Denoting z, z^+ for positive pairs residing in some Hilbert space \mathcal{H} , we wish to learn a bounded linear operator $W : \mathcal{H} \rightarrow \mathbb{R}^K$ such as to minimize

$$\begin{aligned} \bar{\mathcal{L}}(W) &= \|W\bar{\Gamma}W^* - I_K\|_F^2, \text{ where} \\ \bar{\Gamma} &= \frac{1}{2}\mathbb{E}_{z, z^+} [z(z^+)^* + z^+z^*] \end{aligned} \quad (23)$$

Here $*$ denotes the adjoint, $\|\cdot\|_{\text{HS}}$ is the Hilbert-Schmidt norm of linear operators on \mathcal{H} (in the finite-dimensional setting, the Frobenius norm $\|\cdot\|_F$), and $\|\cdot\|$ is the operator norm.

Theorem 6.1. (Generalization error for Barlow Twins in Hilbert spaces) *Let $N \in \mathbb{N}$ and $N' \leq N$. Assume that pairs (z, z^+) are almost surely contained in a ball of radius $S > 0$ in a Hilbert space \mathcal{H} . Moreover, assume that for any set of training data, $\mathcal{L}(W) \leq \delta$ for some $\|W\| \leq B$. Then, with probability $\geq 1 - 2\epsilon$, it holds that*

$$\begin{aligned} \bar{\mathcal{L}}(W) &\leq 3\delta + \frac{3B^4}{N} \left(\hat{\mathbf{V}} + \exp\left(-\frac{(N' - 1)^2 \epsilon^2}{8S^8 N'}\right) \right) \\ &\quad + 3 \exp\left(-\frac{N\epsilon^2}{B^4 S^4}\right) \end{aligned} \quad (24)$$

where we define

$$\begin{aligned} \hat{\mathbf{V}} &= \frac{1}{N'(N' - 1)} \sum_{\substack{i < j \\ i, j \in [N']}} \|\Gamma_i - \Gamma_j\|_{\text{HS}}^2 \\ \Gamma_i &= \frac{1}{2} (z_i(z_i^+)^* + z_i^+z_i^*) \end{aligned} \quad (25)$$

and the randomness is over independently sampled positive pairs $(z_1, z_1^+), \dots, (z_N, z_N^+)$.

The proof relies on McDiarmid's inequality (McDiarmid et al., 1989) and is included in Appendix D.1. The quantity $\hat{\mathbf{V}}$ is purely empirical, and may be estimated from fewer samples by choosing $N' < N$. Of course, estimation is a strong word here — in an abstract Hilbert space setting, we may not be able to compute $\|\Gamma_i - \Gamma_j\|_{\text{HS}}^2$. However, when the inner product in \mathcal{H} takes the form of a kernel, $\hat{\mathbf{V}}$ can be expressed more explicitly.

Connecting to neural networks. Such is the case for the NTK approximation of the neural network (22). Defining $\psi(x) = (\psi_1(x), \dots, \psi_K(x))$ as the concatenation of all feature maps, we obtain the kernel inner product

$$\hat{K}_{\theta_0}(x, x') = \sum_{k=1}^K \langle \nabla f_k(x, \theta_0), \nabla f_k(x', \theta_0) \rangle \quad (26)$$

Then, defining

$$W = \begin{pmatrix} (\theta_T - \theta_0)^T & 0 & \dots \\ 0 & (\theta_T - \theta_0)^T & \dots \\ \dots & \dots & \dots \end{pmatrix} \in \mathbb{R}^{K \times Kp} \quad (27)$$

we obtain $g(x) = W\psi(x)$. Actually, multivariate functions such as g reside in a vector-valued RKHS. This is also clear from the fact that the NTK in the multivariate setting is a matrix-valued kernel K_{θ_0} . In a vector-valued RKHS, we do not have feature maps, but rather feature operators ψ . For the neural network, the feature operator is the Jacobian, that is $\psi(x) = D_{\theta}f(x)$. Under the trace inner product $\langle \psi(x), \psi(x') \rangle := \text{Trace}(\psi(x)^*\psi(x')) = \text{Trace}(K_{\theta_0}(x, x'))$ we see that this construction is identical to the one given above, in that it gives the same inner product between $\psi(x), \psi(x')$. By virtue of the kernel trick, we can now estimate $\hat{\mathbf{V}}$ without moving into the Hilbert space.

Lemma 6.2. (Estimating $\hat{\mathbf{V}}$ with kernels) *Consider the setting of Theorem 6.1. Suppose that there exists a feature map $\psi : \mathbb{R}^d \rightarrow \mathcal{H}$ such that $z_n = \psi(x_n)$ and $z_n^+ = \psi(x_n^+)$. Moreover, assume there exists a kernel $K_{\theta_0} : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ with $\langle \psi(x), \psi(x') \rangle = K_{\theta_0}(x, x')$ for all $x, x' \in \mathbb{R}^d$. Then, for all $i \neq j$, it holds that*

$$\begin{aligned} \|\Gamma_i - \Gamma_j\|_{\text{HS}}^2 &= \\ &0.5 (K_{\theta_0}(x_i, x_i)K_{\theta_0}(x_i^+, x_i^+) + K_{\theta_0}(x_i, x_i^+)^2) + \\ &0.5 (K_{\theta_0}(x_j, x_j)K_{\theta_0}(x_j^+, x_j^+) + K_{\theta_0}(x_j, x_j^+)^2) - \\ &K_{\theta_0}(x_i, x_j)K_{\theta_0}(x_i^+, x_j^+) - K_{\theta_0}(x_i, x_j^+)K_{\theta_0}(x_i^+, x_j) \end{aligned} \quad (28)$$

This result is proven in Appendix D.2. Furthermore, combining Theorem 6.1 with the fact that the neural




Figure 1: For a fixed sample size N , we plot different quantities for varying network width M . We then vary N and plot: (a) NTK change till convergence (b) Training Epochs till convergence (c) Squared norm of difference between representations of neural network and corresponding kernel model

network f can be approximated by a kernel model (22), we immediately obtain the following corollary.

Corollary 6.3. *Fix $\epsilon > 0$. Assume f is a neural with M neurons in the hidden layer, where M is large enough for Corollary 4.5 to hold. Suppose f is trained until the Barlow Twins loss is smaller than δ . Then, with probability at least $1 - 3\epsilon$, it holds that*

$$\bar{\mathcal{L}}(f_{NN}) \leq K\zeta(BS + \zeta) \left(K\zeta(BS + \zeta) + \sqrt{\nu(N, \epsilon, \delta')} \right) \quad (29)$$

where $\delta' = 2\delta + 2K^2\zeta^2(BS + \zeta)^2$ and $\nu(N, \epsilon, \delta')$ is the slack term from Theorem 6.1. B can be taken as the norm of $\|\theta_T - \theta_0\|$, and S is the radius of a ball that contains all $\psi(x)$ where $x \in S^{d-1}$.

The proof is included in Appendix D.3. It uses the fact that f is approximated up to ζ by a kernel model with probability $\geq 1 - \epsilon$. This allows bounding the difference between the loss of f and the loss of g . Then, we use Theorem 6.1, which holds with probability at least $1 - 2\epsilon$.

Remark 6.4. *Our analysis only bounds the pretraining loss. However, this can be passed on to guarantees on the classification error on downstream tasks, provided the augmentations elucidate enough of the underlying class structures. For example, see Section 5.2 of Cabannes et al. (2023).*

7 Experiments

In this Section, we verify our theoretical claims on the MNIST dataset (Deng, 2012). Optimization is done using gradient descent with a learning rate of 0.5. The threshold for loss convergence δ is set at 10^{-5} . We use a single-hidden layer neural network with tanh activation unless stated otherwise. Furthermore, for illustration purposes, we focus only on one-dimensional

embeddings. All experiments are run 10 times with different random seeds; their means along with standard deviations are plotted in the figures.

We first verify near-constancy of the NTK for Barlow Twins at large width, while not restricting ourselves to the setting of $\mathcal{L}(0) < 1$. Varying the sample size from 10 to 1000, we plot the norm of the NTK deviation by varying the hidden layer width M from 10 to 10000. Recall that in our theoretical results, η still depends on the sample size N , suggesting that as N grows, a larger width is necessary for the kernel regime to kick in. However, as Figure 1 (a) clearly displays, the neural network enters the kernel regime irrespective of the sample size N as M grows, with the change in the NTK near zero at 10^3 neurons in the hidden layer.

Next, we look at the number of training epochs required for the loss to converge below δ . As per Theorem 5.3, we expect this to be independent of the width of the neural network, provided the network is sufficiently large. This is empirically verified in Figure 1 (b).

With the NTK nearly constant at large width, we expect the representations learned by the finite neural network to be close to those learned by a corresponding kernel model, via gradient descent under the NTK at initialization $\mathbf{K}(0)$. We verify this in Figure 1 (c), where we see that irrespective of the number of samples, the representations at convergence are closer for wider neural networks. Additional experiments are included in Appendix G. There, we also check the variation of these quantities for networks with ReLU activation.

8 Discussion

In this paper, we connect self-supervised learning with neural networks to the neural tangent kernel regime. We prove that at infinite width, the NTK of a neural network trained under the Barlow Twins loss becomes constant. This is the first result that rigorously justifies the use of traditional kernel methods to understand SSL through the lenses of the NTK. Furthermore, the kernel connection enables us to bound the population Barlow Twins loss in terms of the empirical loss and a slack term that decays with the number of samples. This opens a number of interesting avenues and challenges for future work.

Extension to deep networks, and other losses.

It is desirable to extend the convergence results to other, in particular deep, architectures. This requires an extension of Theorem 4.1 to multiple layers. In addition, verifying the validity of the NTK approximation for other commonly used non-contrastive loss functions such as VIC-Reg (Bardes et al., 2021) is a natural next step. Moreover, our proof for the ReLU activations is not entirely rigorous, since $\frac{\partial \mathcal{L}}{\partial \theta}$ can only be understood in the weak sense. Finally, as discussed earlier, we believe the condition $\mathcal{L}(0) < 1$ can be dropped, but a proof remains to be established.

Improving generalization error bounds. Theorem 6.1 still relies on uniform bounds, despite closed-form expressions for kernel versions of Barlow Twins being available. To be precise, Simon et al. (2023) show that the minimum norm W^* that achieves zero loss lies in the top eigenspace of the cross-moment operator Γ . This W^* is referred to as the *spectral solution*. We believe that spectral perturbation bounds provide tighter guarantees on the generalization error.

Does gradient flow approach the spectral solution? Even with improved generalization bounds for the spectral solution W^* (as noted above), there is still a missing link: It is not yet known whether gradient flow actually approaches the spectral solution for linearized networks. Simon et al. (2023) prove it for “aligned” initialization (starting in the top eigenspace), and give heuristics for why it would also hold under “small” initialization.

Conclusion. Our work establishes the significance of the NTK to self-supervised learning, and confirms numerous recent works that have headed into this direction. As Belkin et al. (2018) put it, *to understand deep learning, we need to understand kernel learning*. In light of our results, we are tempted to add: To understand self-supervised learning, we need to understand representation learning with kernels.

Acknowledgements

This paper is supported by the DAAD programme Konrad Zuse Schools of Excellence in Artificial Intelligence, sponsored by the Federal Ministry of Education and Research.

References

- Anil, G. G., Esser, P., and Ghoshdastidar, D. (2024). When can we approximate wide contrastive models with neural tangent kernels and principal component analysis? *arXiv preprint arXiv:2403.08673*.
- Arora, S., Khandeparkar, H., Khodak, M., Plevrakis, O., and Saunshi, N. (2019). A theoretical analysis of contrastive unsupervised representation learning. *arXiv preprint arXiv:1902.09229*.
- Bachman, P., Hjelm, R. D., and Buchwalter, W. (2019). Learning representations by maximizing mutual information across views. *Advances in neural information processing systems*, 32.
- Bardes, A., Ponce, J., and LeCun, Y. (2021). Vicreg: Variance-invariance-covariance regularization for self-supervised learning. *arXiv preprint arXiv:2105.04906*.
- Belkin, M., Ma, S., and Mandal, S. (2018). To understand deep learning we need to understand kernel learning. In *International Conference on Machine Learning*, pages 541–549. PMLR.
- Blanchard, G., Bousquet, O., and Zwald, L. (2007). Statistical properties of kernel principal component analysis. *Machine Learning*, 66:259–294.
- Bombari, S., Kiyani, S., and Mondelli, M. (2023). Beyond the universal law of robustness: Sharper laws for random features and neural tangent kernels. In *International Conference on Machine Learning*, pages 2738–2776. PMLR.
- Cabannes, V., Kiani, B., Balestrieri, R., LeCun, Y., and Bietti, A. (2023). The ssl interplay: Augmentations, inductive bias, and generalization. In *International Conference on Machine Learning*, pages 3252–3298. PMLR.
- Chizat, L., Oyallon, E., and Bach, F. (2019). On lazy training in differentiable programming. *Advances in neural information processing systems*, 32.
- Deng, L. (2012). The mnist database of handwritten digit images for machine learning research [best of the web]. *IEEE signal processing magazine*, 29(6):141–142.
- Esser, P., Fleissner, M., and Ghoshdastidar, D. (2024). Non-parametric representation learning with kernels. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 38, pages 11910–11918.

- HaoChen, J. Z., Wei, C., Gaidon, A., and Ma, T. (2021). Provable guarantees for self-supervised deep learning with spectral contrastive loss. *Advances in Neural Information Processing Systems*, 34:5000–5011.
- Jacot, A., Gabriel, F., and Hongler, C. (2018). Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31.
- Johnson, D. D., El Hanchi, A., and Maddison, C. J. (2022). Contrastive learning can find an optimal basis for approximately view-invariant functions. In *The Eleventh International Conference on Learning Representations*.
- Kiani, B. T., Balestrieri, R., Chen, Y., Lloyd, S., and LeCun, Y. (2022). Joint embedding self-supervised learning in the kernel regime. *arXiv preprint arXiv:2209.14884*.
- Lee, J., Xiao, L., Schoenholz, S., Bahri, Y., Novak, R., Sohl-Dickstein, J., and Pennington, J. (2019). Wide neural networks of any depth evolve as linear models under gradient descent. *Advances in neural information processing systems*, 32.
- Liu, C., Zhu, L., and Belkin, M. (2020a). On the linearity of large non-linear models: when and why the tangent kernel is constant. *Advances in Neural Information Processing Systems*, 33:15954–15964.
- Liu, C., Zhu, L., and Belkin, M. (2020b). Toward a theory of optimization for over-parameterized systems of non-linear equations: the lessons of deep learning. *arXiv preprint arXiv:2003.00307*, 7.
- Mallinar, N., Simon, J., Abedsoltan, A., Pandit, P., Belkin, M., and Nakkiran, P. (2022). Benign, tempered, or catastrophic: Toward a refined taxonomy of overfitting. *Advances in Neural Information Processing Systems*, 35:1182–1195.
- McDiarmid, C. et al. (1989). On the method of bounded differences. *Surveys in combinatorics*, 141(1):148–188.
- Radford, A., Kim, J. W., Hallacy, C., Ramesh, A., Goh, G., Agarwal, S., Sastry, G., Askell, A., Mishkin, P., Clark, J., et al. (2021). Learning transferable visual models from natural language supervision. In *International conference on machine learning*, pages 8748–8763. PMLR.
- Schölkopf, B., Smola, A., and Müller, K.-R. (1997). Kernel principal component analysis. In *International conference on artificial neural networks*, pages 583–588. Springer.
- Simon, J. B., Dickens, M., Karkada, D., and DeWeese, M. R. (2021). The eigenlearning framework: A conservation law perspective on kernel regression and wide neural networks. *arXiv preprint arXiv:2110.03922*.
- Simon, J. B., Knutins, M., Ziyin, L., Geisz, D., Fetterman, A. J., and Albrecht, J. (2023). On the stepwise nature of self-supervised learning. In *International Conference on Machine Learning*, pages 31852–31876. PMLR.
- Smola, A. J. and Schölkopf, B. (1998). *Learning with kernels*, volume 4. Citeseer.
- Wei, C., Shen, K., Chen, Y., and Ma, T. (2020). Theoretical analysis of self-training with deep networks on unlabeled data. *arXiv preprint arXiv:2010.03622*.
- Zbontar, J., Jing, L., Misra, I., LeCun, Y., and Deny, S. (2021). Barlow twins: Self-supervised learning via redundancy reduction. In *International conference on machine learning*, pages 12310–12320. PMLR.
- Ziyin, L., Lubana, E. S., Ueda, M., and Tanaka, H. (2022). What shapes the loss landscape of self-supervised learning? *arXiv preprint arXiv:2210.00638*.

Supplementary Materials

A Proofs from Section 4

Appendix B contains a more detailed and computationally less involved derivation of all results for the one-dimensional case. It may be convenient to refer to it first. The proof strategy is identical.

A.1 Proof of Theorem 4.1

Proof. Recall that $T > 0$ is a fixed, width-independent point in time. We show that there exists $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}\| \leq \kappa$ for all networks of width $M > 8T$, with high probability over randomly initialized weights. Specifically, for any $\epsilon > 0$, we know that there exists $\kappa_1 > 0$ such that

$$\frac{\|\theta(0)\|^2}{M} \leq \kappa_1 - 1 \quad (30)$$

with probability at least $1 - \epsilon$, because the weights are independent Gaussians at initialization. We condition everything on this event. In this proof, we also assume $\mathcal{L}(0) < 1$ under said event. We need this condition anyway for Theorem 4.4 and therefore include it into our bounds here as well to simplify matters. However, any width-independent bound on $\mathcal{L}(0)$ is sufficient. Denote $c_\phi = \max_{t \in \mathbb{R}} |\phi(t)|$ and $c_{\phi'} = \max_{t \in \mathbb{R}} \left| \frac{\partial}{\partial t} \phi(t) \right|$. We will show the following three statements.

1. Firstly, we show that

$$\frac{\partial}{\partial t} \left(\sum_{m=1}^M \|w_m(t)\|^2 \right) \leq 8 \quad (31)$$

for all $t \leq T$. This immediately implies that there exists some $\kappa_1 > 0$ such that for all $t \leq T$

$$\frac{1}{M} \left(\sum_{m=1}^M \|w_m(t)\|^2 \right) \leq \kappa_1 \quad (32)$$

holds with probability $\geq 1 - \epsilon$ over random initialization, because

$$\frac{1}{M} \left(\sum_{m=1}^M \|w_m(t)\|^2 \right) \leq \frac{1}{M} \left(\sum_{m=1}^M \|w_m(0)\|^2 \right) + \frac{8T}{M} \leq \frac{1}{M} \left(\sum_{m=1}^M \|w_m(0)\|^2 \right) + 1 \quad (33)$$

where we used $M > 8T$ and the fact that

$$\frac{1}{M} \left(\sum_{m=1}^M \|w_m(0)\|^2 \right) \leq \frac{\|\theta(0)\|^2}{M} \leq \kappa_1 - 1 \quad (34)$$

with probability at least $1 - \epsilon$.

2. Using part 1, we bound the maximum squared value that any representation takes until time T , that is

$$|f|^2 = \max_{n \in [N]} \sup_{t \leq T} \max(\|f(x_n)\|^2, \|f(x_n^+)\|^2) \quad (35)$$

by some $\kappa_2 > 0$ (again independent of M).

3. We combine both statements to show that there exists $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}\| \leq \kappa$ with probability at least $1 - \epsilon$.

Part 1. First of all, we derive the evolution of all weights over time. Using the matrix chain rule,

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial w_{mk}} &= \text{Trace} \left(\left(\frac{\partial \mathcal{L}}{\partial C} \right)^T \frac{\partial C}{\partial w_{mk}} \right) \\ &= 2 \text{Trace} \left((C - I) \left(\frac{1}{2N} \frac{\partial}{\partial w_{mk}} \left(\sum_{n=1}^N f(x_n) f(x_n^+)^T + f(x_n^+) f(x_n)^T \right) \right) \right) \end{aligned} \quad (36)$$

and similarly,

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial v_{mr}} &= \text{Trace} \left(\left(\frac{\partial \mathcal{L}}{\partial C} \right)^T \frac{\partial C}{\partial v_{mr}} \right) \\ &= 2 \text{Trace} \left((C - I) \left(\frac{1}{2N} \frac{\partial}{\partial v_{mr}} \left(\sum_{n=1}^N f(x_n) f(x_n^+)^T + f(x_n^+) f(x_n)^T \right) \right) \right) \end{aligned} \quad (37)$$

For any $n \in [N]$ and any $i, j, k \in [K]$

$$\begin{aligned} &\frac{\partial}{\partial w_{mk}} (f_i(x_n) f_j(x_n^+) + f_j(x_n) f_i(x_n^+)) \\ &= \begin{cases} 2f_k(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) + 2f_k(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+), & \text{if } k = i = j \\ f_j(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) + f_j(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+), & \text{if } k = i \neq j \\ f_i(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) + f_i(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+), & \text{if } k = j \neq i \\ 0, & \text{if } k \notin \{i, j\} \end{cases} \end{aligned} \quad (38)$$

Equating $\frac{\partial}{\partial t} w_{mk} = -\frac{\partial \mathcal{L}}{\partial w_{mk}}$ gives

$$\begin{aligned} \frac{\partial}{\partial t} \left(\sum_{m=1}^M \|w_m\|^2 \right) &= 2 \sum_{m=1}^M \sum_{k=1}^K w_{mk} \left(\frac{\partial}{\partial t} w_{mk} \right) \\ &= 4 \text{Trace} \left((I - C) \left(\frac{1}{2N} \sum_{n=1}^N \sum_{k=1}^K A_{k,i,j}(n) \right)_{i,j=1}^K \right) \end{aligned} \quad (39)$$

where we define

$$\begin{aligned} A_{k,i,j}(n) &= \sum_{m=1}^M w_{mk} \cdot \frac{\partial}{\partial w_{mk}} (f_i(x_n) f_j(x_n^+) + f_i(x_n^+) f_j(x_n)) \\ &= \begin{cases} 2f_k(x_n^+) f_k(x_n) + 2f_k(x_n) f_k(x_n^+), & \text{if } k = i = j \\ f_j(x_n^+) f_k(x_n) + f_j(x_n) f_k(x_n^+), & \text{if } k = i \neq j \\ f_i(x_n^+) f_k(x_n) + f_i(x_n) f_k(x_n^+), & \text{if } k = j \neq i \\ 0, & \text{if } k \notin \{i, j\} \end{cases} \end{aligned} \quad (40)$$

Noticing that

$$\sum_{k=1}^K A_{k,i,j}(n) = \begin{cases} 2f_i(x_n^+) f_i(x_n) + 2f_i(x_n) f_i(x_n^+), & \text{if } i = j \\ 2f_i(x_n^+) f_j(x_n) + 2f_j(x_n) f_i(x_n^+), & \text{if } i \neq j \end{cases} \quad (41)$$

we obtain

$$\frac{\partial}{\partial t} \left(\sum_{m=1}^M \|w_m\|^2 \right) = 8 \text{Trace} ((I - C)C) \quad (42)$$

This is certainly bounded by some $\kappa_0 > 0$, because the loss is non-increasing under gradient flow, which forces all entries of C to stay bounded. In our case, with $\mathcal{L}(0) = \|I - C\|_F^2 < 1$, we may choose $\kappa_0 = 8$.

Regarding the evolution of the first layer weights v_{mr} , we note that for any $n \in [N]$, $i, j \in [K]$ and $r \in [d]$

$$\begin{aligned}
 \frac{\partial}{\partial v_{mr}} (f_i(x_n)f_j(x_n^+) + f_j(x_n)f_i(x_n^+)) &= \\
 \frac{1}{\sqrt{M}} f_j(x_n^+) w_{mj} \phi'(v_m^T x_n) x_n^{(r)} &+ \\
 \frac{1}{\sqrt{M}} f_i(x_n) w_{mi} \phi'(v_m^T x_n^+) (x_n^+)^{(r)} &+ \\
 \frac{1}{\sqrt{M}} f_j(x_n) w_{mi} \phi'(v_m^T x_n^+) (x_n^+)^{(r)} &+ \\
 \frac{1}{\sqrt{M}} f_i(x_n^+) w_{mi} \phi'(v_m^T x_n) x_n^{(r)} &
 \end{aligned} \tag{43}$$

Thus, we obtain

$$v_{mr}(t) = \frac{1}{N} \text{Trace} \left((C - I) \left(\frac{\partial}{\partial v_{mr}} (f_i(x_n)f_j(x_n^+) + f_j(x_n)f_i(x_n^+)) \right)_{i,j} \right) \tag{44}$$

Part 2. In this part we bound $|f|^2$. For any x and any $k \in [K]$, the k -th output f_k evolves over time as follows.

$$\begin{aligned}
 \frac{\partial}{\partial t} f_k(x) &= \left\langle \frac{\partial f_k(x)}{\partial \theta}, \frac{\partial}{\partial t} \theta \right\rangle \\
 &= - \sum_{m=1}^M \frac{\partial f_k(x)}{\partial \theta_m} \frac{\partial \mathcal{L}}{\partial \theta_m} \\
 &= - \sum_{m=1}^M \frac{\partial f_k(x)}{\partial \theta_m} \sum_{n=1}^N \sum_{l=1}^K \frac{\partial \mathcal{L}}{\partial f_l(x_n)} \frac{\partial f_l(x_n)}{\partial \theta_m} + \frac{\partial \mathcal{L}}{\partial f_l(x_n^+)} \frac{\partial f_l(x_n^+)}{\partial \theta_m} \\
 &= - \sum_{n=1}^N \sum_{l=1}^K \frac{\partial \mathcal{L}}{\partial f_l(x_n)} (K_\theta(x, x_n))_{k,l} + \frac{\partial \mathcal{L}}{\partial f_l(x_n^+)} (K_\theta(x, x_n^+))_{k,l}
 \end{aligned} \tag{45}$$

So we need to derive all $\frac{\partial \mathcal{L}}{\partial f_k(x_i)}$. For any $k \in [K]$, and any $i \in [N]$, we see that $\frac{\partial \mathcal{L}}{\partial f_k(x_i)} = \frac{1}{2N} A(i, k)$, where we define the matrix

$$A(i, k) = \begin{pmatrix} 0 & \dots & 0 & f_1(x_i^+) & 0 & \dots & 0 \\ 0 & \dots & 0 & f_2(x_i^+) & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ f_1(x_i^+) & f_2(x_i^+) & \dots & 2f_k(x_i^+) & \dots & \dots & f_K(x_i^+) \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & f_{K-1}(x_i^+) & 0 & \dots & 0 \\ 0 & \dots & 0 & f_K(x_i^+) & 0 & \dots & 0 \end{pmatrix} \tag{46}$$

and also let

$$A(i) = \begin{pmatrix} f_1(x_i^+) & \dots & f_1(x_i^+) \\ \dots & \dots & \dots \\ f_K(x_i^+) & \dots & f_K(x_i^+) \end{pmatrix} \in \mathbb{R}^{K \times K} \tag{47}$$

Notice that $A(i, k) = e_k e_k^T A(i)^T + A(i) e_k e_k^T$. Using the matrix chain rule, and the fact that $\frac{\partial \mathcal{L}}{\partial C} = 2(C - I)$, we

therefore obtain

$$\begin{aligned}
 \frac{\partial \mathcal{L}}{\partial f_k(x_i)} &= 2 \text{Trace} \left((C - I)^T \frac{\partial C}{\partial f_k(x_i)} \right) \\
 &= \frac{1}{N} \text{Trace} ((C - I)A(i, k)) \\
 &= \frac{1}{N} \left(\sum_{l=1}^K (C - I)_{l \bullet} A(i, k)_{\bullet l} \right) \\
 &= \frac{1}{N} \left(\sum_{l \neq k} (C - I)_{lk} f_l(x_i^+) + (C - I)_{\bullet k}^T f(x_i^+) \right) \\
 &= \frac{1}{N} \left(\sum_{l \neq k} C_{lk} f_l(x_i^+) + (C - I)_{\bullet k}^T f(x_i^+) \right) \\
 &= \frac{2}{N} (C_{1k} f_1(x_i^+) + C_{2k} f_2(x_i^+) + \dots + (C_{kk} - 1) f_k(x_i^+) + \dots + C_{Kk} f_K(x_i^+)) \\
 &= \frac{2}{N} (C - I)_{k \bullet} f(x_i^+) \\
 &= \frac{2}{N} e_k^T (C - I) f(x_i^+)
 \end{aligned} \tag{48}$$

where we exploited the symmetry of $A(i, k)$ in the sixth step. Overall, we have

$$\begin{aligned}
 \frac{\partial}{\partial t} f_k(x) &= -\frac{2}{N} \sum_{n=1}^N \sum_{l=1}^K K_\theta(x, x_n)_{k,l} \cdot e_l^T (C - I) f(x_n^+) + \\
 &\quad K_\theta(x, x_n^+)_{k,l} \cdot e_l^T (C - I) f(x_n) \\
 &= -\frac{2}{N} \sum_{l=1}^K K_\theta(x, [\mathcal{X}, \mathcal{X}_+])_{k,l} \cdot \mathbf{F}([\mathcal{X}_+, \mathcal{X}])^T (C - I) e_l
 \end{aligned} \tag{49}$$

where we write $\mathbf{F}([\mathcal{X}_+, \mathcal{X}]) \in \mathbb{R}^{K \times 2N}$ for the matrix that contains the representations at time t as columns. Moreover, with $\mathbf{K}_{k,l}$ the kernel matrix of $\mathcal{X}, \mathcal{X}_+$ at output k, l , we have

$$\frac{\partial}{\partial t} f_k([\mathcal{X}, \mathcal{X}_+]) = \frac{2}{N} \sum_{l=1}^K \mathbf{K}_{k,l} \cdot \mathbf{F}([\mathcal{X}_+, \mathcal{X}])^T (I - C) e_l \tag{50}$$

Therefore, the squared norm of the representations evolve as

$$\begin{aligned}
 \frac{\partial}{\partial t} \left(\sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\|^2 \right) &= 2 \sum_{k=1}^K (f_k([\mathcal{X}, \mathcal{X}_+]))^T \left(\frac{\partial}{\partial t} f_k([\mathcal{X}, \mathcal{X}_+]) \right) \\
 &= \frac{4}{N} \sum_{k=1}^K (f_k([\mathcal{X}, \mathcal{X}_+]))^T \sum_{l=1}^K \mathbf{K}_{k,l} \cdot \mathbf{F}([\mathcal{X}_+, \mathcal{X}])^T (C - I) e_l \\
 &\leq \frac{4}{N} \sum_{k,l=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\| \|\mathbf{K}_{k,l} \mathbf{F}([\mathcal{X}_+, \mathcal{X}])^T (C - I) e_l\| \\
 &\leq \frac{4}{N} \sum_{k,l=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\| \cdot \|\mathbf{K}_{k,l}\|_2 \cdot \|\mathbf{F}([\mathcal{X}_+, \mathcal{X}])^T (C - I) e_l\| \\
 &\leq \frac{4}{N} \sum_{k,l=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\| \cdot \|\mathbf{K}_{k,l}\|_2 \cdot \|\mathbf{F}([\mathcal{X}_+, \mathcal{X}])\|_2
 \end{aligned} \tag{51}$$

where we have used Cauchy-Schwartz inequality, and the fact that $\|I - C\|_2 \leq 1$ due to $\mathcal{L}(0) < 1$ in the last line. Continuing,

$$\begin{aligned}
 & \frac{4}{N} \sum_{k,l=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\| \cdot \|\mathbf{K}_{k,l}\|_2 \cdot \|\mathbf{F}([\mathcal{X}_+, \mathcal{X}])\|_2 \leq \\
 & \frac{4}{N} \max_{k,l} \|\mathbf{K}_{k,l}\|_2 \cdot \left(\sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\| \left(\sum_{l=1}^K \|f_l([\mathcal{X}, \mathcal{X}_+])\|^2 \right)^{1/2} \right) \leq \\
 & \frac{4}{N} \max_{k,l} \|\mathbf{K}_{k,l}\|_2 \cdot \left(\sum_{k,l=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\| \cdot \|f_l([\mathcal{X}, \mathcal{X}_+])\| \right) \leq \\
 & \frac{4K}{N} \max_{k,l} \|\mathbf{K}_{k,l}\|_2 \cdot \left(\sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\|^2 \right)
 \end{aligned} \tag{52}$$

where in the second line we bound the spectral norm of $\mathbf{F}([\mathcal{X}_+, \mathcal{X}])$ by its Frobenius norm, in the third line we bound the 2-norm by the 1-norm, and in the fourth line we use Cauchy-Schwartz again, which introduces a factor of K . We proceed to bound the spectral norm of the kernel matrices $\mathbf{K}_{k,l}$.

Note that for any k, l , and any $(a, b) \in [\mathcal{X}, \mathcal{X}_+] \times [\mathcal{X}, \mathcal{X}_+]$, we have

$$\begin{aligned}
 K_{k,l}(a, b) &= \left(\frac{\partial f_k(a)}{\partial \theta} \right)^T \left(\frac{\partial f_l(a)}{\partial \theta} \right) \\
 &= \frac{1}{M} \left(\sum_{m=1}^M \mathbf{1}_{k=l} \cdot \phi(v_m^T a) \phi(v_m^T b) \right) + \frac{1}{M} \left(\sum_{m=1}^M w_{mk} w_{ml} \phi'(v_m^T a) \phi'(v_m^T b) a^T b \right) \\
 &\leq c_\phi^2 + \frac{1}{M} \left(\sum_{m=1}^M |w_{mk} w_{ml}| \cdot c_{\phi'}^2 \right) \\
 &\leq c_\phi^2 + \frac{c_{\phi'}^2}{M} \cdot \left(\sum_{m=1}^M w_{mk}^2 + w_{ml}^2 \right) \\
 &\leq c_\phi^2 + c_{\phi'}^2 \kappa_1
 \end{aligned} \tag{53}$$

where the final line uses Part 1 of this proof, which bounds $\frac{1}{M} \sum_{m=1}^M \|w_m(t)\|^2$ uniformly in time by κ_1 . Also, recall that data lies on the unit sphere, so $a^T b \leq 1$. Overall, we see that the spectral norm of each kernel matrix $\mathbf{K}_{k,l}$ is bounded by

$$\|\mathbf{K}_{k,l}\|_2 \leq 2N (c_\phi^2 + c_{\phi'}^2 \kappa_1) \tag{54}$$

Plugging this in back into our evolution of the representations,

$$\frac{\partial}{\partial t} \sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\|^2 \leq (8Kc_\phi^2 + 8Kc_{\phi'}^2 \kappa_1) \left(\sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\|^2 \right) \tag{55}$$

We are thus in the setting of Grönwall's inequality, and obtain

$$\sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])\|^2 \leq \left(\sum_{k=1}^K \|f_k([\mathcal{X}, \mathcal{X}_+])(0)\|^2 \right) \cdot \exp(T (8Kc_\phi^2 + 8Kc_{\phi'}^2 \kappa_1)) =: \kappa_2 \tag{56}$$

for all time $t \leq T$. When $\frac{\|\theta(0)\|^2}{M} \leq \kappa_1 - 1$ this also gives a bound on $\|f_k([\mathcal{X}, \mathcal{X}_+])(0)\|^2$.

Part 3. We now control $\|\dot{\theta}(t)\|^2$. Using the time derivatives for w_{mk} and v_{mr} derived in Part 1,

$$\begin{aligned}
 |w_{mk}(t)|^2 &= \frac{1}{N^2} \text{Trace} \left((C - I) \left(\sum_{n=1}^N \frac{\partial}{\partial w_{mk}} (f_i(x_n) f_j(x_n^+) + f_j(x_n) f_i(x_n^+)) \right)_{i,j} \right)^2 \\
 &\leq \frac{1}{N^2} \left\| \sum_{n=1}^N \left(\frac{\partial}{\partial w_{mk}} (f_i(x_n) f_j(x_n^+) + f_j(x_n) f_i(x_n^+)) \right)_{i,j} \right\|_F^2 \\
 &= \mathcal{O} \left(\frac{\kappa_2 c_\phi^2}{M} \right)
 \end{aligned} \tag{57}$$

where we used Cauchy-Schwartz for the trace inner product and $\|C - I\|_F < 1$ in the second line, and $|f|^2 \leq \kappa_2$ from Part 2 in the third line. Summing up,

$$\sum_{m=1}^M \|w_m(t)\|^2 = \mathcal{O}(\kappa_2 c_\phi^2) \tag{58}$$

Similarly, we find that

$$\begin{aligned}
 |v_{mr}(t)|^2 &= \frac{1}{N^2} \text{Trace} \left((C - I) \left(\sum_{n=1}^N \frac{\partial}{\partial v_{mr}} (f_i(x_n) f_j(x_n^+) + f_j(x_n) f_i(x_n^+)) \right)_{i,j} \right)^2 \\
 &\leq \frac{1}{N^2} \left\| \sum_{n=1}^N \left(\frac{\partial}{\partial v_{mr}} (f_i(x_n) f_j(x_n^+) + f_j(x_n) f_i(x_n^+)) \right)_{i,j} \right\|_F^2 \\
 &= \mathcal{O} \left(\frac{\kappa_2 \|w_m\|^2 c_{\phi'}^2}{M} \right)
 \end{aligned} \tag{59}$$

Summing up and exploiting part 1 to bound $\frac{1}{M} \sum_{m=1}^M \|w_m(t)\|^2$ uniformly in time by κ_1 , we find that

$$\sum_{m=1}^M \|v_m\|^2 = \mathcal{O}(\kappa_1 \kappa_2 c_{\phi'}^2) \tag{60}$$

Combining both results, we see that there exists $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}(t)\| \leq \kappa$ with probability $\geq 1 - \epsilon$ over random initialization. \square

A.2 Proof of Lemma 4.2

Proof. Recall that for any x and any $k \in [K]$, the k -th output f_k evolves over time as follows.

$$\begin{aligned}
 \frac{\partial}{\partial t} f_k(x) &= \left\langle \frac{\partial f_k(x)}{\partial \theta}, \frac{\partial}{\partial t} \theta \right\rangle \\
 &= - \sum_{m=1}^M \frac{\partial f_k(x)}{\partial \theta_m} \frac{\partial \mathcal{L}}{\partial \theta_m} \\
 &= - \sum_{m=1}^M \frac{\partial f_k(x)}{\partial \theta_m} \sum_{n=1}^N \sum_{l=1}^K \frac{\partial \mathcal{L}}{\partial f_l(x_n)} \frac{\partial f_l(x_n)}{\partial \theta_m} + \frac{\partial \mathcal{L}}{\partial f_l(x_n^+)} \frac{\partial f_l(x_n^+)}{\partial \theta_m} \\
 &= - \sum_{n=1}^N \sum_{l=1}^K \frac{\partial \mathcal{L}}{\partial f_l(x_n)} (K_\theta(x, x_n))_{k,l} + \frac{\partial \mathcal{L}}{\partial f_l(x_n^+)} (K_\theta(x, x_n^+))_{k,l}
 \end{aligned} \tag{61}$$

From here, we write down the time evolution of the loss. We denote $*$, $+$ for the symbols that refer to an anchor sample $x = x^*$, or an augmentation x^+ .

$$\begin{aligned}
 \frac{\partial}{\partial t} \mathcal{L}(t) &= \sum_{i=1}^N \frac{\partial \mathcal{L}}{\partial f(x_i)} \frac{\partial}{\partial t} f(x_i) + \frac{\partial \mathcal{L}}{\partial f(x_i^+)} \frac{\partial}{\partial t} f(x_i^+) \\
 &= - \sum_{i,n=1}^N \sum_{k,l=1}^K \sum_{\alpha,\beta=*,+} \frac{\partial \mathcal{L}}{\partial f_k(x_i^\alpha)} K_{k,l}(x_i^\alpha, x_n^\beta) \frac{\partial \mathcal{L}}{\partial f_k(x_i^\beta)} \\
 &= - \sum_{i,n=1}^N \left(\frac{\partial \mathcal{L}}{\partial f(x_i)} \right)^T K(x_i, x_n) \left(\frac{\partial \mathcal{L}}{\partial f(x_n)} \right) + \\
 &\quad \left(\frac{\partial \mathcal{L}}{\partial f(x_i^+)} \right)^T K(x_i^+, x_n) \left(\frac{\partial \mathcal{L}}{\partial f(x_n)} \right) + \\
 &\quad \left(\frac{\partial \mathcal{L}}{\partial f(x_i)} \right)^T K(x_i, x_n^+) \left(\frac{\partial \mathcal{L}}{\partial f(x_n^+)} \right) + \\
 &\quad \left(\frac{\partial \mathcal{L}}{\partial f(x_i^+)} \right)^T K(x_i^+, x_n^+) \left(\frac{\partial \mathcal{L}}{\partial f(x_n^+)} \right)
 \end{aligned} \tag{62}$$

this can be expressed as

$$\frac{\partial}{\partial t} \mathcal{L}(t) = -u(t)^T \mathbf{K}(t) u(t) \tag{63}$$

where we have defined

$$u(t) = \left(\frac{\partial \mathcal{L}}{\partial f_1(x_1)} \quad \cdots \quad \frac{\partial \mathcal{L}}{\partial f_K(x_1)} \quad \cdots \quad \frac{\partial \mathcal{L}}{\partial f_K(x_N)} \quad \frac{\partial \mathcal{L}}{\partial f_1(x_1^+)} \quad \cdots \quad \frac{\partial \mathcal{L}}{\partial f_K(x_N^+)} \right)^T \tag{64}$$

$$\mathbf{K}(t) = \begin{pmatrix} K(x_1, x_1) & \cdots & K(x_1, x_N) & \cdots & K(x_1, x_N^+) \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ K(x_N, x_1) & \cdots & K(x_N, x_N) & \cdots & K(x_N, x_N^+) \\ K(x_1^+, x_1) & \cdots & K(x_1^+, x_N) & \cdots & K(x_1^+, x_N^+) \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ K(x_N^+, x_1) & \cdots & K(x_N^+, x_N) & \cdots & K(x_N^+, x_N^+) \end{pmatrix} \tag{65}$$

□

A.3 Proof of Theorem 4.4

Proof. Observe that

$$\frac{\partial}{\partial t} \mathcal{L}(t) = -u(t)^T \mathbf{K}(t) u(t) \leq -\|u(t)\|^2 \lambda_{\min}(\mathbf{K}(t)) \tag{66}$$

with \mathbf{K} and $u(t)$ as defined earlier. Recall from Appendix A.1 (part 3 of the proof) that

$$\frac{\partial \mathcal{L}}{\partial f_k(x_i)} = \frac{2}{N} e_k^T (C - I) f(x_i^+) \tag{67}$$

Thus, we can lower bound the squared Euclidean norm of $u(t)$ as

$$\begin{aligned}
 \|u(t)\|^2 &= \sum_{k=1}^K \sum_{i=1}^N \left(\frac{\partial \mathcal{L}}{\partial f_k(x_i)} \right)^2 + \left(\frac{\partial \mathcal{L}}{\partial f_k(x_i^+)} \right)^2 \\
 &= \sum_{k=1}^K \sum_{i=1}^N f(x_i^+)^T (C - I) e_k e_k^T (C - I) f(x_i^+) + f(x_i)^T (C - I) e_k e_k^T (C - I) f(x_i) \\
 &= \frac{4}{N^2} \sum_{i=1}^N f(x_i^+)^T (C - I)^2 f(x_i^+) + f(x_i)^T (C - I)^2 f(x_i) \\
 &\geq \frac{4}{N^2} \sum_{i=1}^N f(x_i^+) (C - I)^2 f(x_i) + f(x_i) (C - I)^2 f(x_i^+) \\
 &= \frac{4}{N^2} \text{Trace} \left((C - I)^2 \left(\sum_{i=1}^N f(x_i) f(x_i^+)^T + f(x_i^+) f(x_i)^T \right) \right) \\
 &= \frac{8}{N} \text{Trace}((C - I)^2 C)
 \end{aligned} \tag{68}$$

The inequality used above relies on the following fact: For any positive semi-definite matrix $A = Q^T \Lambda Q \in \mathbb{R}^K$ (where Q is orthonormal and Λ is diagonal with non-negative entries), and for any $v, w \in \mathbb{R}^K$, Cauchy-Schwartz yields

$$\begin{aligned}
 v^T A w + w^T A v &= (Qv)^T \Lambda (Qw) + (Qw)^T \Lambda (Qv) \\
 &= 2 \sum_{k=1}^K \lambda_k (Qv)_k (Qw)_k \\
 &\leq \sum_{k=1}^K \lambda_k ((Qv)_k^2 + (Qw)_k^2) \\
 &= v^T Q^T \Lambda Q v + w^T Q^T \Lambda Q w \\
 &= v^T A v + w^T A w
 \end{aligned} \tag{69}$$

All that remains to be done is to show that

$$\|u(t)\|^2 \geq \kappa \|C - I\|_F^2 \tag{70}$$

for some κ that is time-independent. By Von Neumann's trace inequality we can certainly choose $\kappa = \lambda_{\min}(C(t))$ at any given t . Thus, as long as the smallest eigenvalue of $C(t)$ is lower bounded until convergence, we are fine.

Recall that we assume $\mathcal{L}(0) \in (0, 1)$. Now assume that at some time t' we have $\lambda_{\min}(C(t')) < 1 - \sqrt{\mathcal{L}(0)}$. Then, denoting $\lambda_1, \dots, \lambda_K$ for the eigenvalues of $C(t')$, it follows that

$$\mathcal{L}(t') = \text{Trace}((C - I)^2) = \sum_{k=1}^K (\lambda_k - 1)^2 \geq (\lambda_{\min}(C(t')) - 1)^2 > \mathcal{L}(0) \tag{71}$$

This is a contradiction to the non-decreasing nature of the loss under gradient flow. Hence, as long as $\mathcal{L}(0) < 1$, we can lower bound $\lambda_{\min}(C)$ uniformly in time by $1 - \sqrt{\mathcal{L}(0)}$ from below. Thus, it holds that

$$\|u(t)\|^2 > \frac{8}{N} (1 - \sqrt{\mathcal{L}(0)}) \cdot \mathcal{L}(t) \tag{72}$$

and hence

$$\frac{\partial}{\partial t} \mathcal{L}(t) < -\frac{8(1 - \sqrt{\mathcal{L}(0)}) \lambda_{\min}(\mathbf{K}(t))}{N} \cdot \mathcal{L}(t) \tag{73}$$

holds for all t . We now take M to be so large that the entries of the kernel matrix change by less than some $\gamma > 0$ for all $t \leq T$, where $\gamma > 0$ is small enough to ensure that $\lambda_{\min}(\mathbf{K}(t)) \geq \lambda/2$ for all $t \leq T$. Such M certainly exists, because Theorem 3.1 guarantees that the change in the entries of the kernel matrix $\mathbf{K}(t)$ are no more than $\mathcal{O}(R/\sqrt{M})$ until time T , where $R = \exp(2\kappa T)$ following the discussion precluding Theorem 4.1. Overall, this implies that whenever $M \geq M_1$ for some $M_1 \in \mathbb{N}$, we have $\frac{\partial}{\partial t} \mathcal{L}(t) < -\eta \mathcal{L}(t)$ for all $t \leq T$. \square

A.4 Ensuring $\mathcal{L}(0) < 1$ for ReLU

Lemma A.1. (Scaling the first layer ensures small loss) *Consider the ReLU activation function. For any dataset $\mathcal{X}, \mathcal{X}_+$, there exist $s > 0$ and $M_0 \in \mathbb{N}$ such that for all $M \geq M_0$ and under $w_{mk} \sim \mathcal{N}(0, 1)$, $v_{mr} \sim \mathcal{N}(0, s^2)$, the loss at initialization satisfies $\mathcal{L}(0) < 1$ with high probability.*

Proof. In expectation, we have the following expression for the cross-moments at initialization.

$$\begin{aligned} \mathbb{E}[C_{kl}] &= \frac{1}{2N} \sum_{n=1}^N \mathbb{E}[f_k(x_n) f_l(x_n^+)] + \mathbb{E}[f_k(x_n^+) f_l(x_n)] \\ &= \frac{1}{2N} \sum_{n=1}^N \frac{1}{M} \sum_{m, m'=1}^M \mathbb{E} [w_{mk} (\phi(v_m^T x_n) \phi(v_{m'}^T x_n^+) + \phi(v_m^T x_n^+) \phi(v_{m'}^T x_n)) w_{m'l}] \end{aligned} \quad (74)$$

For $k \neq l$, the fact that all w_{mk} are independent and zero mean shows that $\mathbb{E}[C_{kl}] = 0$. When $k = l$, we have $\mathbb{E}[w_{mk} w_{m'l}] = \mathbf{1}(m = m')$, since the weights w_{mk} are standard Gaussians. Therefore,

$$\mathbb{E}[C_{kk}] = \frac{1}{N} \sum_{n=1}^N \frac{1}{M} \sum_m \mathbb{E} [\phi(v_m^T x_n) \phi(v_m^T x_n^+)] \quad (75)$$

For ReLU, the expectation is strictly positive and scales as s^2 . Thus, there certainly exists $s > 0$ such that

$$\mathbb{E}[C_{kk}] = \frac{2K - 1}{2K} \quad (76)$$

For large M , the variance of each C_{kl} is $\mathcal{O}(1/M)$ because the weights are independent. Thus, the matrix $I - C$ concentrates around $(2K)^{-1} I_K$, so $\mathcal{L}(0) = \|C(0) - I\|_F^2 < 1$ with high probability. \square

B Proofs from Section 5

B.1 Proof of Theorem 5.1

Proof. Recall that $T > 0$ is a fixed, width-independent point in time. We show that there exists $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}\| \leq \kappa$ for all networks of width $M > 8T$, with high probability over randomly initialized weights. Specifically, for any $\epsilon > 0$, we know that there exists $\kappa_1 > 0$ such that

$$\frac{\|\theta(0)\|^2}{M} \leq \kappa_1 - 1 \quad (77)$$

with probability at least $1 - \epsilon$, because the weights are independent Gaussians at initialization. We condition everything on this event. In this proof, we also assume $\mathcal{L}(0) < 1$ under said event. We need this condition anyway for Theorem 4.4 and therefore include it into our bounds here as well to simplify matters. However, any width-independent bound on $\mathcal{L}(0)$ is sufficient. Denote $c_\phi = \max_{t \in \mathbb{R}} |\phi(t)|$ and $c_{\phi'} = \max_{t \in \mathbb{R}} \left| \frac{\partial}{\partial t} \phi(t) \right|$. We will show the following three statements.

1. Firstly, we show that

$$\frac{\partial}{\partial t} \left(\sum_{m=1}^M w_m^2(t) \right) \leq 8 \quad (78)$$

for all $t \leq T$. This immediately implies that there exists some $\kappa_1 > 0$ such that for all $t \leq T$

$$\frac{1}{M} \left(\sum_{m=1}^M w_m^2(t) \right) \leq \kappa_1 \quad (79)$$

holds with probability $\geq 1 - \epsilon$ over random initialization, because

$$\frac{1}{M} \left(\sum_{m=1}^M w_m^2(t) \right) \leq \frac{1}{M} \left(\sum_{m=1}^M w_m^2(0) \right) + \frac{8T}{M} \leq \frac{1}{M} \left(\sum_{m=1}^M w_m^2(0) \right) + 1 \quad (80)$$

where we used $M > 8T$ and the fact that

$$\frac{1}{M} \left(\sum_{m=1}^M w_m^2(0) \right) \leq \kappa_1 - 1 \quad (81)$$

whenever $\frac{\|\theta(0)\|^2}{M} \leq \kappa_1 - 1$.

2. From there, we bound the maximum squared value that any representation takes until time T , that is

$$|f|^2 = \max_{n \in [N]} \sup_{t \leq T} \max (|f(x_n)|^2, |f(x_n^+)|^2) \quad (82)$$

by some $\kappa_2 > 0$ (again independent of M).

3. We combine both statements to show that there exists $\kappa > 0$ such that $\sup_{t \leq T} \|\dot{\theta}\| \leq \kappa$ with probability at least $1 - \epsilon$.

Part 1. We begin by computing

$$\frac{\partial \mathcal{L}}{\partial w_m} = 2(C-1) \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+) + f(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) \right] \right) \quad (83)$$

which equates to $-\frac{\partial}{\partial t} w_m$ under gradient flow, and

$$\frac{\partial \mathcal{L}}{\partial v_{mj}} = 2(C-1) \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} w_m \phi'(v_m^T x_n^+) (x_n^+)^{(j)} + f(x_n^+) \frac{1}{\sqrt{M}} w_m \phi'(v_m^T x_n) (x_n)^{(j)} \right] \right) \quad (84)$$

which equates to $-\frac{\partial}{\partial t}v_{mj}$ under gradient flow. Clearly,

$$\frac{\partial}{\partial t}(w_m^2) = 4(1-C) \left(\frac{1}{N} \sum_{n=1}^N [f(x_n)f_m(x_n^+) + f(x_n^+)f_m(x_n)] \right) \quad (85)$$

where we write $f_m(x) = \frac{1}{\sqrt{M}}w_m\phi(v_m^T x)$. Noticing that $f(x) = \sum_{m=1}^M f_m(x)$, we arrive at

$$\begin{aligned} \frac{\partial}{\partial t} \left(\sum_{m=1}^M w_m^2 \right) &= 4(1-C) \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \left(\sum_{m=1}^M f_m(x_n^+) \right) + f(x_n^+) \left(\sum_{m=1}^M f_m(x_n) \right) \right] \right) \\ &= 8(1-C)C \end{aligned} \quad (86)$$

The loss $\mathcal{L}(t) = (C-1)^2$ is non-increasing under gradient flow. Thus, since $\mathcal{L}(0) < 1$, we have $C \in (0, 1)$. We obtain $\frac{\partial}{\partial t} \left(\sum_{m=1}^M w_m^2(t) \right) \leq 8$ for all $t \leq T$ as desired.

Part 2. We consider the evolution of the representations $f(x)$ over time, where $x \in \mathcal{X}, \mathcal{X}_+$. Note that for any x ,

$$\begin{aligned} \frac{\partial}{\partial t} f(x) &= \left\langle \frac{\partial}{\partial \theta} f(x), \frac{\partial}{\partial t} \theta \right\rangle \\ &= - \sum_{m=1}^M \frac{\partial f(x)}{\partial \theta_m} \frac{\partial \mathcal{L}}{\partial \theta_m} \\ &= - \sum_{m=1}^M \frac{\partial}{\partial \theta_m} f(x) \left[\sum_{n=1}^N \left(\frac{\partial \mathcal{L}}{\partial f(x_n)} \frac{\partial f(x_n)}{\partial \theta_m} + \frac{\partial \mathcal{L}}{\partial f(x_n^+)} \frac{\partial f(x_n^+)}{\partial \theta_m} \right) \right] \\ &= - \sum_{n=1}^N \left[K_\theta(x, x_n) \frac{\partial \mathcal{L}}{\partial f(x_n)} + K_\theta(x, x_n^+) \frac{\partial \mathcal{L}}{\partial f(x_n^+)} \right] \end{aligned} \quad (87)$$

In our setting

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial f(x_n)} &= 2(C-1) \left(\frac{1}{N} f(x_n^+) \right) \\ \frac{\partial \mathcal{L}}{\partial f(x_n^+)} &= 2(C-1) \left(\frac{1}{N} f(x_n) \right) \end{aligned} \quad (88)$$

Hence, for any $i \in [N]$, we have

$$\begin{aligned} \frac{\partial}{\partial t} f(x_i) &= \frac{2(1-C)}{N} \cdot \sum_{n=1}^N [K_\theta(x_i, x_n) f(x_n^+) + K_\theta(x_i, x_n^+) f(x_n)] \\ &= \frac{2(1-C)}{N} \cdot K_\theta(x_i, [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \end{aligned} \quad (89)$$

and therefore we can write

$$\frac{\partial}{\partial t} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} = \frac{2(1-C)}{N} \cdot K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \quad (90)$$

Thus,

$$\frac{\partial}{\partial t} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 = 2 \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T \left(\frac{\partial}{\partial t} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right) \quad (91)$$

$$= \frac{4(1-C)}{N} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \quad (92)$$

By Cauchy-Schwartz inequality, for any positive semi-definite matrix A , it holds that

$$|y^T A z|^2 \leq (y^T A y) \cdot (z^T A z) \leq \|y\|^2 \|z\|^2 \|A\|_2^2 \quad (93)$$

In our case, this implies that

$$\begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \leq \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \cdot \|K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+])\|_2 \quad (94)$$

where we used the fact that

$$\left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\| = \left\| \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \right\| \quad (95)$$

since both vectors are just permutations of one another. It remains to show that the spectral norm of the time-dependent kernel matrix remains bounded until time T . Note that for any pair $(a, b) \in [\mathcal{X}, \mathcal{X}_+] \times [\mathcal{X}, \mathcal{X}_+]$, and for any time $t \leq T$, we have $|a^T b| \leq 1$. Therefore, every entry of the kernel matrix is bounded via

$$\begin{aligned} K_\theta(a, b) &= \left(\frac{\partial f(a)}{\partial \theta} \right)^T \left(\frac{\partial f(b)}{\partial \theta} \right) \\ &= \sum_{m=1}^M \left(\frac{\partial f(a)}{\partial w_m} \frac{\partial f(b)}{\partial w_m} + \sum_{j=1}^d \frac{\partial f(a)}{\partial v_{mj}} \frac{\partial f(b)}{\partial v_{mj}} \right) \\ &= \frac{1}{M} \sum_{m=1}^M \phi(v_m^T a) \phi(v_m^T b) + w_m^2 \phi'(v_m^T a) \phi'(v_m^T b) a^T b \\ &\leq \frac{1}{M} \sum_{m=1}^M c_\phi^2 + w_m^2 c_{\phi'}^2 \\ &\leq c_\phi^2 + c_{\phi'}^2 \left(\frac{1}{M} \sum_{m=1}^M w_m^2(t) \right) \\ &\leq c_\phi^2 + c_{\phi'}^2 \kappa_1 \end{aligned} \quad (96)$$

where we used Part 1 in the final inequality. Since the spectral norm is upper bounded by the trace, we obtain

$$\|K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+])\|_2 \leq 2N (c_\phi^2 + c_{\phi'}^2 \kappa_1) \quad (97)$$

Consequently, going back to the evolution of the representations, we see that

$$\frac{\partial}{\partial t} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \leq 8(1 - C) (c_\phi^2 + c_{\phi'}^2 \kappa_1) \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \leq 8 (c_\phi^2 + c_{\phi'}^2 \kappa_1) \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \quad (98)$$

Grönwall's inequality now ensures that there exists $\kappa_2 > 0$ such that

$$\left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t) \right\|^2 \leq \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (0) \right\|^2 \exp(8T (c_\phi^2 + c_{\phi'}^2 \kappa_1)) =: \kappa_2 \quad (99)$$

for all time $t \leq T$, where we used the fact that $\left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (0) \right\|^2$ can be bounded in terms of $\frac{\|\theta(0)\|^2}{M}$. Since

$$|f|^2 = \max_{n \in [N]} \sup_{t \leq T} \max (|f(x_n)|^2, |f(x_n^+)|^2) \leq \sup_{t \leq T} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t) \right\|^2 \quad (100)$$

this concludes part 2, giving us a bound κ_2 on $|f|^2$.

Part 3. We finally control $\|\dot{\theta}(t)\|^2$. Using the time derivatives for w_m, v_{mj} we derived in Part 1,

$$\begin{aligned}
 |w_m(t)|^2 &= 4(1-C)^2 \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+) + f(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) \right] \right)^2 \\
 &\leq 4(1-C)^2 \left(2|f|c_\phi \frac{1}{\sqrt{M}} \right)^2 \\
 &= 16(1-C)^2 \frac{|f|^2 c_\phi^2}{M} \\
 &\leq \frac{16\kappa_2 c_\phi^2}{M}
 \end{aligned} \tag{101}$$

for all $m \in [M]$. Similarly, it holds that

$$\begin{aligned}
 |v_{mj}(t)|^2 &= 4(1-C)^2 \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} w_m \phi'(v_m^T x_n^+) (x_n^+)^{(j)} + f(x_n^+) \frac{1}{\sqrt{M}} w_m \phi'(v_m^T x_n) (x_n)^{(j)} \right] \right)^2 \\
 &\leq 4(1-C)^2 \left(\frac{2|f||w_m|c_{\phi'}}{\sqrt{M}} \right)^2 \\
 &\leq \frac{16\kappa_2 w_m^2 c_{\phi'}^2}{M}
 \end{aligned} \tag{102}$$

for all $m \in [M], j \in [d]$. Thus, we obtain

$$\begin{aligned}
 \|\dot{\theta}(t)\|^2 &= \sum_{m=1}^M \sum_{j=1}^d (w_m(t))^2 + (v_{mj}(t))^2 \\
 &\leq \frac{1}{M} \left(\sum_{m=1}^M 16\kappa_2 c_\phi^2 + 16d\kappa_2 w_m^2 c_{\phi'}^2 \right)
 \end{aligned} \tag{103}$$

From part 1, $\frac{1}{M} \sum_{m=1}^M w_m^2(t) \leq \kappa_1$ holds for all $t \leq T$. Thus, defining

$$\kappa = 4\sqrt{\kappa_2 c_\phi^2 + d\kappa_1 \kappa_2 c_{\phi'}^2} \tag{104}$$

we obtain the desired high-probability bound on $\sup_{t \leq T} \|\dot{\theta}(t)\|$ and this concludes the proof. \square

B.2 Proof of Lemma 5.2

Proof. First recall what we showed in Appendix B.1. The function representations $f(x)$ evolve as

$$\frac{\partial}{\partial t} f(x) = - \sum_{n=1}^N K_\theta(x, x_n) \frac{\partial \mathcal{L}}{\partial f(x_n)} + K_\theta(x, x_n^+) \frac{\partial \mathcal{L}}{\partial f(x_n^+)} \tag{105}$$

for any $i \in [N]$. From there, it is apparent that

$$\begin{aligned}
 \frac{\partial}{\partial t} \mathcal{L}(t) &= \sum_{i=1}^N \frac{\partial \mathcal{L}}{\partial f(x_i)} \frac{\partial}{\partial t} f(x_i) + \frac{\partial \mathcal{L}}{\partial f(x_i^+)} \frac{\partial}{\partial t} f(x_i^+) \\
 &= - \sum_{i,n=1}^N \frac{\partial \mathcal{L}}{\partial f(x_i)} \frac{\partial \mathcal{L}}{\partial f(x_n)} K(x_n, x_i) + \frac{\partial \mathcal{L}}{\partial f(x_i^+)} \frac{\partial \mathcal{L}}{\partial f(x_n)} K(x_n, x_i^+) + \\
 &\quad \frac{\partial \mathcal{L}}{\partial f(x_i)} \frac{\partial \mathcal{L}}{\partial f(x_n^+)} K(x_n^+, x_i) + \frac{\partial \mathcal{L}}{\partial f(x_i^+)} \frac{\partial \mathcal{L}}{\partial f(x_n^+)} K(x_n^+, x_i^+) \\
 &= - \frac{4(C-1)^2}{N^2} \sum_{i,n=1}^N f(x_i^+) f(x_n^+) K(x_n, x_i) + f(x_i) f(x_n^+) K(x_n, x_i^+) + \\
 &\quad f(x_i^+) f(x_n) K(x_n^+, x_i) + f(x_i) f(x_n) K(x_n^+, x_i^+) \\
 &= - \frac{4}{N^2} \cdot \mathcal{L}(t) \cdot \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T \mathbf{K}(t) \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}
 \end{aligned} \tag{106}$$

where the kernel matrix $\mathbf{K}(t)$ is defined as

$$\mathbf{K}(t) = \begin{pmatrix} K_\theta(\mathcal{X}_+, \mathcal{X}_+) & K_\theta(\mathcal{X}_+, \mathcal{X}) \\ K_\theta(\mathcal{X}, \mathcal{X}_+) & K_\theta(\mathcal{X}, \mathcal{X}) \end{pmatrix} \tag{107}$$

□

B.3 Proof of Theorem 5.3

Proof. Note that the entries of the kernel matrix \mathbf{K} change by no more than $\mathcal{O}(\frac{\exp(2\kappa T)}{\sqrt{M}})$ up to time T . This is a consequence of Theorem 3.1, which states that the change is $\mathcal{O}(\frac{R}{\sqrt{M}})$ where R is the radius of the ball within which the weights evolve, and Theorem 5.1. Since the spectral norm is upper bounded by the trace of a matrix, for any $\gamma > 0$ there exists some large M such that

$$\|\mathbf{K}(t) - \mathbf{K}(0)\|_2 \leq \text{Trace}(\mathbf{K}(t) - \mathbf{K}(0)) \leq \gamma \tag{108}$$

for all $t \leq T$. We pick $\gamma = \frac{\lambda}{2}$, which ensures that

$$\lambda_{\min}(\mathbf{K}(t)) \geq \frac{\lambda}{2} \tag{109}$$

Define

$$z(t) = \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t) \tag{110}$$

We claim that $\frac{\|z(t)\|^2}{2N} \geq C(0)$ for all time $t \geq 0$. Indeed, assume this was not the case at a certain time $t' > 0$, where instead

$$\frac{\|z(t')\|^2}{2N} < C(0) \tag{111}$$

Then,

$$\begin{aligned}
 C(t') &= \frac{1}{2N} \sum_{n=1}^N f(x_n) f(x_n^+) + f(x_n^+) f(x_n) \\
 &= \frac{1}{2N} \left(\begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t') \right)^T \left(\begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} (t') \right) \\
 &\leq \frac{1}{2N} \|z(t')\|^2 \\
 &< C(0)
 \end{aligned} \tag{112}$$

The first inequality is Cauchy-Schwartz, using the fact that

$$\left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t') \right\| = \left\| \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} (t') \right\| = \|z(t')\| \quad (113)$$

The second inequality plugs in our assumption (111). But if $C(t') < C(0)$, then this leads to a contradiction. Since $C(0) \in (0, 1)$, we would obtain

$$\mathcal{L}(t') = (1 - C(t'))^2 > (1 - C(0))^2 = \mathcal{L}(0) \quad (114)$$

This is impossible, because gradient flow monotonically decreases \mathcal{L} . Thus, for all $t \leq T$, we know that

$$\frac{\|z(t)\|^2}{2N} \geq C(0) \quad (115)$$

From there, we complete the proof.

$$\begin{aligned} g(t) &= \frac{4}{N^2} \cdot z(t)^T \mathbf{K}(t) z(t) \\ &\geq \frac{4}{N^2} \cdot \|z(t)\|^2 \cdot \lambda_{\min}(\mathbf{K}(t)) \\ &> \frac{2\lambda}{N^2} \cdot \|z(t)\|^2 \\ &\geq \frac{2\lambda}{N^2} \cdot (2NC(0)) \\ &= \frac{4\lambda C(0)}{N} \\ &= \frac{4\lambda(1 - \sqrt{\mathcal{L}(0)})}{N} \\ &= \eta \end{aligned} \quad (116)$$

as desired. □

C Extending Theorem 5.1 to ReLU Activations

In this section, we prove that the weights remain in a ball of bounded radius even for ReLU activations, until any fixed time $T > 0$. The proof is not entirely rigorous, because gradient flow w.r.t. the weights is ill-defined due to the non-differentiability of the ReLU function. We restrict ourselves to an analysis of the one-dimensional setting.

Proof. Similar to the case for smooth bounded activations with bounded first derivatives (Appendix B.1), our proof consists of three parts. Remember that $T > 0$ is a fixed, width-independent point in time. We will show the following three statements, and assume $\mathcal{L}(0) < 1$ throughout. However, as before, all that we really need is boundedness of the loss at initialization. Note that our arguments are slightly reshuffled compared to the proof of Theorem 5.1. This is due to unboundedness of the ReLU function.

1. Firstly, we show that $\frac{\partial}{\partial t} \|\theta(t)\|^2 \leq 16$ for all $t \leq T$.
2. Secondly, we verify that for any $t \leq T$, it holds that

$$\|\dot{\theta}(t)\|^2 \leq \frac{16|f|^2 d \|\theta\|^2}{M} \quad (117)$$

where we denote

$$|f|^2 = \max_{n \in [N]} \sup_{t \leq T} \max(|f(x_n)|^2, |f(x_n^+)|^2) \quad (118)$$

for the maximum squared value that any representation takes until time T .

3. Thirdly, we show that there exists a constant κ_1 (again depending only on T) such that $|f|^2 \leq \kappa_1$.

Together, this will be enough. Part 1 implies that for all $t \leq T$, we have

$$\frac{\|\theta(t)\|^2}{M} \leq \frac{\|\theta(0)\|^2 + 16T}{M} \leq \frac{\|\theta(0)\|^2}{M} + 2 \quad (119)$$

since $M > 8T$. Then, combining part 2 and part 3, we arrive at

$$\|\dot{\theta}(t)\|^2 \leq \frac{16d\kappa_1(\|\theta(0)\|^2 + 16T)}{M} \leq 16d\kappa_1 \left(\frac{\|\theta(0)\|^2}{M} + 2 \right) \quad (120)$$

for all $t \leq T$. Note that for any $\epsilon > 0$ we can choose κ such that this expression is smaller than κ with probability at least $1 - \epsilon$. This is possible because the weights at initialization are independent Gaussians.

Part 1. We begin by computing

$$\frac{\partial \mathcal{L}}{\partial w_m} = 2(C-1) \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+) + f(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) \right] \right) \quad (121)$$

which equates to $-\frac{\partial}{\partial t} w_m$ under gradient flow, and

$$\frac{\partial \mathcal{L}}{\partial v_{mj}} = 2(C-1) \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} w_m \phi'(v_m^T x_n^+) (x_n^+)^{(j)} + f(x_n^+) \frac{1}{\sqrt{M}} w_m \phi'(v_m^T x_n) (x_n)^{(j)} \right] \right) \quad (122)$$

which equates to $-\frac{\partial}{\partial t} v_{mj}$ under gradient flow. Thus, writing $f_m(x) = \frac{1}{\sqrt{M}} w_m \phi(v_m^T x)$, it holds that

$$\frac{\partial}{\partial t} (w_m^2) = \frac{\partial}{\partial t} (\|v_m\|^2) = 4(1-C) \left(\frac{1}{N} \sum_{n=1}^N [f(x_n) f_m(x_n^+) + f(x_n^+) f_m(x_n)] \right) \quad (123)$$

where we used properties of the ReLU function ϕ , namely that

$$\phi(v_m^T x) = \phi'(v_m^T x) \cdot v_m^T x \quad (124)$$

Hence, noticing that $f(x) = \sum_{m=1}^M f_m(x)$, we arrive at

$$\begin{aligned} \frac{\partial}{\partial t} (\|\theta\|^2) &= \frac{\partial}{\partial t} \left(\sum_{m=1}^M (w_m^2 + \|v_m\|^2) \right) \\ &= 8(1-C) \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \left(\sum_{m=1}^M f_m(x_n^+) \right) + f(x_n^+) \left(\sum_{m=1}^M f_m(x_n) \right) \right] \right) \\ &= 16(1-C)C \end{aligned} \quad (125)$$

The loss is decreasing under gradient flow. Due to $\mathcal{L}(0) < 1$, we have $C \in (0, 1)$. We obtain $\frac{\partial}{\partial t} (\|\theta\|^2) \leq 16$ as claimed.

Part 2. We now control $\|\dot{\theta}(t)\|$. First note that $|\phi(v_m^T x)| \leq \|v_m\|$ due to $\|x\| = 1$ for all $x \in (\mathcal{X}, \mathcal{X}_+)$. Moreover,

$$\begin{aligned} |w_m(t)|^2 &= 4(1-C)^2 \left(\frac{1}{N} \sum_{n=1}^N \left[f(x_n) \frac{1}{\sqrt{M}} \phi(v_m^T x_n^+) + f(x_n^+) \frac{1}{\sqrt{M}} \phi(v_m^T x_n) \right] \right)^2 \\ &\leq 4(1-C)^2 \left(\frac{\|v_m\|}{N\sqrt{M}} \sum_{n=1}^N [|f(x_n)| + |f(x_n^+)|] \right)^2 \\ &\leq 4(1-C)^2 \left(\frac{2\|v_m\|\|f\|}{\sqrt{M}} \right)^2 \\ &= \frac{16(1-C)^2 \|v_m\|^2 \|f\|^2}{M} \\ &\leq \frac{16\|v_m\|^2 \|f\|^2}{M} \end{aligned} \quad (126)$$

holds for all $m \in [M]$. Similarly, it holds that

$$|v_{mj}(t)|^2 \leq \frac{16w_m^2 \|f\|^2}{M} \quad (127)$$

for all $m \in [M], j \in [d]$. Thus, we obtain

$$\begin{aligned} \|\dot{\theta}(t)\|^2 &= \sum_{m=1}^M \left[|w_m(t)|^2 + \sum_{j=1}^d |v_{mj}(t)|^2 \right] \\ &\leq \frac{16\|f\|^2}{M} \left[\sum_{m=1}^M (\|v_m\|^2 + dw_m^2) \right] \\ &\leq \frac{16\|f\|^2 d \|\theta\|^2}{M} \end{aligned} \quad (128)$$

as desired.

Part 3. Just as in Appendix B.1, we look at the evolution of the representations over time. We obtain

$$\frac{\partial}{\partial t} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} = \frac{2(1-C)}{N} \cdot K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \quad (129)$$

and thus

$$\begin{aligned} \frac{\partial}{\partial t} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 &= 2 \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T \left(\frac{\partial}{\partial t} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right) \\ &= \frac{4(1-C)}{N} \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \end{aligned} \quad (130)$$

By Cauchy-Schwartz inequality,

$$\begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix}^T K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+]) \begin{pmatrix} f(\mathcal{X}_+) \\ f(\mathcal{X}) \end{pmatrix} \leq \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \cdot \|K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+])\|_2 \quad (131)$$

Again, we must bound the spectral norm of the time-dependent kernel matrix until time T . This is now slightly different for ReLU. Note that for any pair $(a, b) \in [\mathcal{X}, \mathcal{X}_+] \times [\mathcal{X}, \mathcal{X}_+]$, we have

$$\begin{aligned} K_\theta(a, b) &= \left(\frac{\partial f(a)}{\partial \theta} \right)^T \left(\frac{\partial f(b)}{\partial \theta} \right) \\ &= \sum_{m=1}^M \frac{\partial f(a)}{\partial w_m} \frac{\partial f(b)}{\partial w_m} + \sum_{j=1}^d \frac{\partial f(a)}{\partial v_{mj}} \frac{\partial f(b)}{\partial v_{mj}} \\ &= \frac{1}{M} \sum_{m=1}^M \phi(v_m^T a) \phi(v_m^T b) + w_m^2 \phi'(v_m^T a) \phi'(v_m^T b) a^T b \\ &\leq \frac{1}{M} \sum_{m=1}^M \|v_m\|^2 + w_m^2 \\ &= \frac{\|\theta(t)\|^2}{M} \\ &\leq \frac{\|\theta(0)\|^2}{M} + 1 \end{aligned} \quad (132)$$

using part 1. Since the spectral norm is upper bounded by the trace, we obtain

$$\|K_\theta([\mathcal{X}, \mathcal{X}_+], [\mathcal{X}, \mathcal{X}_+])\|_2 \leq N \left(\frac{\|\theta(0)\|^2}{M} + 1 \right) \quad (133)$$

which is $\mathcal{O}(1)$. Consequently, going back to the evolution of the representations, we see that

$$\frac{\partial}{\partial t} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \leq 4 \left(\frac{\|\theta(0)\|^2}{M} + 1 \right) \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} \right\|^2 \quad (134)$$

Grönwall's inequality now ensures that

$$\left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t) \right\|^2 \leq \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (0) \right\|^2 \exp \left(4T \left(\frac{\|\theta(0)\|^2}{M} + 1 \right) \right) =: \kappa_1 \quad (135)$$

for all time $t \leq T$. Since

$$|f|^2 = \max_{n \in [N]} \sup_{t \leq T} \max (|f(x_n)|^2, |f(x_n^+)|^2) \leq \sup_{t \leq T} \left\| \begin{pmatrix} f(\mathcal{X}) \\ f(\mathcal{X}_+) \end{pmatrix} (t) \right\|^2 \quad (136)$$

this concludes part 3 and finishes the proof. \square

D Proofs from Section 6

D.1 Proof of Theorem 6.1

Proof. Our proof is based on previous works on the theoretical analysis of Kernel PCA (Blanchard et al., 2007). Denote by $\bar{L}(W) = \|W\bar{\Gamma}W^* - I\|_F$ the square root of the population Barlow Twins loss. Denote by $L(W) = \|W\Gamma W^* - I\|_F$ the empirical version, computed with respect to the empirical cross-moment matrix

$$\Gamma = \frac{1}{2N} \sum_{n=1}^N z_n(z_n^+)^* + z_n^+ z_n^* \quad (137)$$

Given independent positive pairs $(z_1, z_1^+), \dots, (z_N, z_N^+)$, define the map

$$\psi(x_1, \dots, x_N) = \sup_{\|W\| \leq B} |\bar{L}(W) - L(W)| \quad (138)$$

where $W : \mathcal{H} \rightarrow \mathbb{R}^K$ is a bounded linear operator. The function ψ satisfies a bounded differences inequality, because for any (y_n, y_n^+) (and with Γ' denoting the cross-moment matrix w.r.t the new sample) it holds that

$$\begin{aligned} & |\psi(z_1, \dots, z_n, \dots, z_N) - \psi(z_1, \dots, y_n, \dots, z_N)| \\ & \leq \sup_{\|W\| \leq B} \left| \|W\Gamma W^* - I\|_F - \|W\Gamma' W^* - I\|_F \right| \\ & \leq \sup_{\|W\| \leq B} \|W(\Gamma - \Gamma')W^*\|_F \\ & \leq B^2 \left\| \frac{1}{2N} (z_n(z_n^+)^* + (z_n^+)z_n^T - y_n(y_n^+)^* - (y_n^+)y_n^*) \right\|_{\text{HS}} \\ & \leq \frac{B^2}{N} (\|z_n(z_n^+)^*\|_{\text{HS}} + \|y_n(y_n^+)^*\|_{\text{HS}}) \\ & \leq \frac{2B^2 S^2}{N} \end{aligned} \quad (139)$$

Here, we used the fact that $\|AB\|_{\text{HS}} \leq \|A\|\|B\|_{\text{HS}}$ for operators A, B , and moreover that

$$\|z(z^+)^*\|_{\text{HS}}^2 = \|z\|\|z^+\| \leq S^2 \quad (140)$$

for any z, z^* (recall that they lie in a ball of radius no more than S in \mathcal{H}). Overall, McDiarmid's inequality yields that for any $\epsilon > 0$

$$\mathbb{P}(\psi(z_1, \dots, z_N) - \mathbb{E}_{z_1, \dots, z_N} [\psi(z_1, \dots, z_N)] > \epsilon) \leq \exp\left(-\frac{N\epsilon^2}{2B^4 S^4}\right) \quad (141)$$

The expectation of ψ (w.r.t samples z_1, \dots, z_N) can be bounded as follows.

$$\begin{aligned} \mathbb{E}_{z_1, \dots, z_N} [\psi(z_1, \dots, z_N)] & = \mathbb{E}_{z_1, \dots, z_N} \left[\sup_{\|W\| \leq B} |L(W) - \hat{L}(W)| \right] \\ & \leq \mathbb{E}_{z_1, \dots, z_N} \left[\sup_{\|W\| \leq B} \left| \|W\bar{\Gamma}W^* - I\|_F - \|W\Gamma W^* - I\|_F \right| \right] \\ & \leq B^2 \mathbb{E}_{z_1, \dots, z_N} [\|\Gamma - \bar{\Gamma}\|_{\text{HS}}] \end{aligned} \quad (142)$$

Let us write

$$\Gamma_i = \frac{1}{2} (z_i(z_i^+)^* + z_i^+ z_i^*) \quad (143)$$

so that $\Gamma = \frac{1}{N} \sum_{i=1}^N \Gamma_i$. Continuing with Jensen's inequality,

$$\begin{aligned}
 B^2 \mathbb{E}_{z_1, \dots, z_N} [\|\Gamma - \bar{\Gamma}\|_{\text{HS}}] &\leq B^2 \left(\mathbb{E}_{z_1, \dots, z_N} \left[\|\Gamma - \bar{\Gamma}\|_{\text{HS}}^2 \right] \right)^{1/2} \\
 &= B^2 \left(\mathbb{E}_{z_1, \dots, z_N} \left[\left\| \left(\frac{1}{N} \sum_{i=1}^N \Gamma_i \right) - \bar{\Gamma} \right\|_{\text{HS}}^2 \right] \right)^{1/2} \\
 &= B^2 \left(\mathbb{E}_{z_1, \dots, z_N} \left[\frac{1}{N^2} \sum_{i,j=1}^N \text{Trace} \left((\Gamma_i - \bar{\Gamma})^* (\Gamma_j - \bar{\Gamma}) \right) \right] \right)^{1/2} \\
 &= B^2 \left(\frac{1}{N^2} \sum_{i=1}^N \mathbb{E}_{\Gamma_i} \left[\|\Gamma_i - \bar{\Gamma}\|_{\text{HS}}^2 \right] \right)^{1/2} \\
 &= \frac{B^2}{\sqrt{N}} \left(\mathbb{E}_{\Gamma_i} \left[\|\Gamma_i - \bar{\Gamma}\|_{\text{HS}}^2 \right] \right)^{1/2}
 \end{aligned} \tag{144}$$

In the third line, we rewrote the Hilbert-Schmidt norm in terms of the trace inner product, then used independence and zero mean of all $\Gamma_i - \bar{\Gamma}, \Gamma_j - \bar{\Gamma}$ to drop the cross terms, and finally wrote everything in terms of a single expectation.

$$\mathbf{V} = \mathbb{E} \left[\|\Gamma_i - \bar{\Gamma}\|_{\text{HS}}^2 \right] \tag{145}$$

is the variance of the random operator Γ_i w.r.t the Hilbert-Schmidt norm, and therefore estimate it from the empirical variance

$$\hat{\mathbf{V}} = \frac{1}{N'(N' - 1)} \sum_{\substack{i < j \\ i, j \in [N']}} \|\Gamma_i - \Gamma_j\|_{\text{HS}}^2 \tag{146}$$

This is a function of independent random operators $\Gamma_1, \dots, \Gamma_{N'}$, and again satisfies a bounded differences inequality. For any collection of $\Gamma_1, \dots, \Gamma_n$ and any ‘‘new’’ element Γ'_n , it holds that

$$\begin{aligned}
 &\left| \hat{\mathbf{V}}(\Gamma_1, \dots, \Gamma_n, \dots, \Gamma_{N'}) - \hat{\mathbf{V}}(\Gamma_1, \dots, \Gamma'_n, \dots, \Gamma_{N'}) \right| \\
 &\leq \frac{1}{N'(N' - 1)} \sum_{i=1}^{N'} \left| \|\Gamma_i - \Gamma_n\|_{\text{HS}}^2 - \|\Gamma_i - \Gamma'_n\|_{\text{HS}}^2 \right| \\
 &\leq \frac{1}{(N' - 1)} \|\Gamma_n - \Gamma'_n\|_{\text{HS}}^2 \\
 &\leq \frac{4S^4}{N' - 1}
 \end{aligned} \tag{147}$$

where we again used the fact that data is contained in a ball of radius S . Using McDiarmid's inequality once again, we conclude that with probability $\geq 1 - \epsilon$ over random training data,

$$\mathbf{V} \leq \hat{\mathbf{V}} + \exp \left(-\frac{(N' - 1)^2 \epsilon^2}{8S^8 N'} \right) \tag{148}$$

Overall, this shows that with probability $\geq 1 - \epsilon$, the expected value of $\phi(z_1, \dots, z_N)$ is bounded as

$$\mathbb{E}[\phi(z_1, \dots, z_n)] \leq \frac{B^2}{\sqrt{N}} \left(\hat{\mathbf{V}} + \exp \left(-\frac{(N' - 1)^2 \epsilon^2}{8S^8 N'} \right) \right)^{1/2} \tag{149}$$

By a union bound, we conclude that with probability $\geq 1 - 2\epsilon$ over training samples

$$|\bar{L}(W) - L(W)| \leq \frac{B^2}{\sqrt{N}} \left(\hat{\mathbf{V}} + \exp \left(-\frac{(N' - 1)^2 \epsilon^2}{8S^8 N'} \right) \right)^{1/2} + \exp \left(-\frac{N\epsilon^2}{2B^4 S^4} \right) \tag{150}$$

holds uniformly over all W with $\|W\| \leq B$. Using the fact that $\mathcal{L}(W) \leq \delta$ almost surely over randomly drawn samples, we push it to the right and square both sides to arrive at

$$\bar{\mathcal{L}}(W) \leq 3\delta + \frac{3B^4}{N} \left(\hat{V} + \exp \left(-\frac{(N' - 1)^2 \epsilon^2}{8S^8 N'} \right) \right) + 3 \exp \left(-\frac{N\epsilon^2}{B^4 S^4} \right) \quad (151)$$

with probability at least $1 - 2\epsilon$. This concludes the proof. \square

D.2 Proof of Lemma 6.2

Proof. For all $i \neq j$, we can expand the Hilbert-Schmidt norm and use the cyclic property of the trace to obtain

$$\begin{aligned} \|\Gamma_i - \Gamma_j\|_{\text{HS}}^2 &= \|\Gamma_i\|_{\text{HS}}^2 + \|\Gamma_j\|_{\text{HS}}^2 - 2 \text{Trace}(\Gamma_i^* \Gamma_j) \\ &= \frac{1}{2} (\|\psi(x_i)\|^2 \|\psi(x_i^+)\|^2 + \langle \psi(x_i), \psi(x_i^+) \rangle^2 + \|\psi(x_j)\|^2 \|\psi(x_j^+)\|^2 + \langle \psi(x_j), \psi(x_j^+) \rangle^2) \\ &\quad - (\langle \psi(x_i), \psi(x_j) \rangle \cdot \langle \psi(x_i^+), \psi(x_j^+) \rangle) + \langle \psi(x_i), \psi(x_j^+) \rangle \cdot \langle \psi(x_i^+), \psi(x_j) \rangle \end{aligned} \quad (152)$$

which can be written purely in terms of the kernel, giving

$$\begin{aligned} \|\Gamma_i - \Gamma_j\|_{\text{HS}}^2 &= 0.5 (K(x_i, x_i)K(x_i^+, x_i^+) + K(x_i, x_i^+)^2) + \\ &\quad 0.5 (K(x_j, x_j)K(x_j^+, x_j^+) + K(x_j, x_j^+)^2) - \\ &\quad K(x_i, x_j)K(x_i^+, x_j^+) - K(x_i, x_j^+)K(x_i^+, x_j) \end{aligned} \quad (153)$$

which concludes the proof. \square

D.3 Proof of Corollary 6.3

Proof. Denote f for the neural network and $g = W\psi(x)$ for the corresponding NTK model from Equation (22). We first note that

$$\begin{aligned} & \left| \bar{\mathcal{L}}(f)^{1/2} - \bar{\mathcal{L}}(g)^{1/2} \right| \leq \\ & \left\| \mathbb{E} [f(x)f(x^+)^T + f(x^+)f(x)^T - g(x)g(x^+)^T - g(x^+)g(x)^T] \right\|_F = \\ & \left(\sum_{i,j=1}^K \mathbb{E} [|f_i(x)f_j(x^+) + f_i(x^+)f_j(x) - g_i(x)g_j(x^+) - g_i(x^+)g_j(x)|^2] \right)^{1/2} \leq \\ & \left(\sum_{i,j=1}^K \mathbb{E} [|f_i(x)f_j(x^+) - g_j(x^+)| + |g_i(x) - f_i(x)||g_j(x^+)| + |f_i(x^+)f_j(x) - g_j(x)| + |g_i(x^+) - f_i(x^+)||g_j(x)|]^2 \right)^{1/2} \leq \\ & \left(\sum_{i,j=1}^K \zeta^2 \mathbb{E} [|f_i(x)| + |g_j(x^+)| + |f_j(x^+)| + |g_j(x)|]^2 \right)^{1/2} \leq \\ & \left(\sum_{i,j=1}^K \zeta^2 \mathbb{E} [|g_i(x)| + |g_j(x^+)| + |g_j(x^+)| + |g_j(x)| + 2\zeta]^2 \right)^{1/2} \leq \\ & \sqrt{\sum_{i,j=1}^K \zeta^2 (BS + 2\zeta)^2} = \\ & K\zeta(BS + \zeta) \end{aligned} \quad (154)$$

where we used the fact that $\|f_i - g_i\|_\infty \leq \zeta$ for all $i \in [K]$ and that $g_i(x) = \langle w_i, \phi(x) \rangle \leq \|w_i\| \|\psi(x)\| \leq BS$ since the data is contained in a ball of radius S in the Hilbert space, and $\|W\| \leq B$.

The same derivation works for the empirical loss, so

$$\left| \mathcal{L}(f)^{1/2} - \mathcal{L}(g)^{1/2} \right| \leq K\zeta(BS + \zeta) \quad (155)$$

Now with $\mathcal{L}(f) < \delta$, we have

$$\mathcal{L}(g)^{1/2} < \sqrt{\delta} + K\zeta(BS + \zeta) \implies \mathcal{L}(g) \leq 2\delta + 2K^2\zeta^2(BS + \zeta)^2 =: \delta' \quad (156)$$

From here, Theorem 6.1 implies that

$$\bar{\mathcal{L}}(g) \leq \nu(N, \epsilon, \delta') \quad (157)$$

with probability at least $1 - 2\epsilon$, where $\nu'(N, \epsilon, \tilde{\delta})$ is the slack term in the corresponding theorem. Thus, with the same probability, it holds that

$$\begin{aligned} |\bar{\mathcal{L}}(f) - \bar{\mathcal{L}}(g)| &\leq \left| \bar{\mathcal{L}}(f)^{1/2} - \bar{\mathcal{L}}(g)^{1/2} \right| \cdot \left| \bar{\mathcal{L}}(f)^{1/2} + \bar{\mathcal{L}}(g)^{1/2} \right| \\ &\leq \left| \bar{\mathcal{L}}(f)^{1/2} - \bar{\mathcal{L}}(g)^{1/2} \right| \cdot \left(\left| \bar{\mathcal{L}}(f)^{1/2} - \bar{\mathcal{L}}(g)^{1/2} \right| + \bar{\mathcal{L}}(g)^{1/2} \right) \\ &\leq \left| \bar{\mathcal{L}}(f)^{1/2} - \bar{\mathcal{L}}(g)^{1/2} \right| \cdot \left(\left| \bar{\mathcal{L}}(f)^{1/2} - \bar{\mathcal{L}}(g)^{1/2} \right| + \sqrt{\nu(N, \epsilon, \delta')} \right) \\ &\leq K\zeta(BS + \zeta) \left(K\zeta(BS + \zeta) + \sqrt{\nu(N, \epsilon, \delta')} \right) \end{aligned} \quad (158)$$

which concludes the proof. □

E Dynamics of Linear Barlow Twins — Convergence from Small Initialization

We write the empirical linearized Barlow Twins loss as

$$\mathcal{L}(W) = \|W\Gamma W^T - I_K\|_F^2 \quad (159)$$

where Γ is the empirical cross-moment matrix between the anchors and the augmentations, either in \mathbb{R}^d or mapped into some RKHS.

It is known from recent works (Simon et al., 2023) that $\frac{\partial \mathcal{L}}{\partial W} = 4(W\Gamma W^T - I)W\Gamma$ and therefore, under gradient flow, it holds that $\frac{\partial}{\partial t} W = 4(I - W\Gamma W^T)W\Gamma$. We decompose the matrix W into $W = Q + W_0$, where the rows of W_0 are in the nullspace of Γ , and the rows of Q are in its orthogonal complement. Since $W\Gamma = Q\Gamma$, we have $\mathcal{L}(W) = \mathcal{L}(Q)$, and $\frac{\partial}{\partial t} W_0 = 0$. Therefore, the part of W that starts in the nullspace Γ stays completely unchanged. We may therefore w.l.o.g. restrict ourselves to the setting where Γ has only nonzero eigenvalues. We obtain

$$\frac{\partial}{\partial t} C = 4(I - C)W\Gamma^2 W^T + 4W\Gamma^2 W^T(I - C) \quad (160)$$

For an eigendecomposition $\Gamma = UDU^T$ with diagonal D and orthogonal U , we write $|\Gamma| = U|D|U^T$ where $|D|_{ii} = |D_{ii}|$. Moreover, denote by $\mu_\Gamma > 0$ the smallest nonzero eigenvalue of Γ in absolute value. Then,

$$\begin{aligned} \lambda_{\min}(W\Gamma^2 W^T) &= \min_{\|u\|=1} u^T W|\Gamma|^{1/2} |\Gamma| |\Gamma|^{1/2} W^T u \\ &\geq \mu_\Gamma \| |\Gamma|^{1/2} W^T u \|^2 \\ &\geq \mu_\Gamma \lambda_{\min}(W|\Gamma|W^T) \\ &\geq \mu_\Gamma \lambda_{\min}(C) \end{aligned} \quad (161)$$

Therefore, for any x having unit norm, the quadratic form $x^T C x$ satisfies

$$\begin{aligned} \frac{\partial}{\partial t} (x^T C x) &\geq 8\mu_\Gamma \lambda_{\min}(C)(1 - x^T C x), \\ \frac{\partial}{\partial t} (x^T C x) &\leq 8\lambda_{\max}(\Gamma)\lambda_{\max}(C)(1 - x^T C x) \end{aligned} \quad (162)$$

where we used Von-Neumann's trace inequality on the product of the two matrices $I - C$ and $W\Gamma^2 W^T$. This shows that as long as all eigenvalues of $C(0)$ are contained in $(0, 1)$, the map $t \mapsto x^T C(t)x$ is strictly increasing for all x , and has an equilibrium at 1. Consequently, we see that the smallest eigenvalue of C is strictly increasing, and that no eigenvalue of C can ever exceed 1. In addition,

$$\begin{aligned} \frac{\partial}{\partial t} \mathcal{L}(t) &= 2 \text{Trace} \left((C - I) \frac{\partial C}{\partial t} \right) \\ &= -16 \text{Trace} \left((C - I)^2 W\Gamma^2 W^T \right) \\ &\leq -16 \lambda_{\min}(C) \mu_\Gamma \cdot \mathcal{L}(t) \\ &\leq -16 \lambda_{\min}(C(0)) \mu_\Gamma \cdot \mathcal{L}(t) \\ &= -\eta \mathcal{L}(t) \end{aligned} \quad (163)$$

where $\eta = 16 \lambda_{\min}(C(0)) \mu_\Gamma$. Therefore, $\mathcal{L}(t) \leq \mathcal{L}(0) \exp(-\eta t)$ and we see that after $T = \frac{-\log \delta}{\eta}$ time, the loss is smaller than δ .

F Inequalities

F.1 Grönwall's Inequality

Lemma F.1. (*Grönwall's inequality*) Let $u(t)$ and $\beta(t)$ be two continuous functions on an interval $[a, b]$. Suppose $u(t)$ is differentiable on (a, b) and satisfies

$$\frac{\partial}{\partial t} u(t) \leq \beta(t)u(t) \tag{164}$$

for all $t \in (a, b)$. Then,

$$u(t) \leq u(0) \exp\left(\int_a^t \beta(s) ds\right) \tag{165}$$

for all $t \in [a, b]$.

F.2 McDiarmid's inequality

Lemma F.2. (*McDiarmid's inequality*) Let $(Z_1, \dots, Z_N) = Z$ be a finite sequence of independent random variables, each with values in \mathcal{Z} and let $\phi : \mathcal{Z}^N \rightarrow \mathbb{R}$ be a measurable function such that $|\phi(Z) - \phi(Z')| \leq \nu_n$ whenever Z, Z' differ only in n -th coordinate. Then, for every $\epsilon > 0$, it holds that

$$\mathbb{P}(\phi(Z) - \mathbb{E}[\phi(Z)] > \epsilon) \leq \exp\left(-\frac{2\epsilon^2}{\|\nu\|_2^2}\right) \tag{166}$$

G Experimental Details & Additional Experiments

All reported experiments have been run 10 times, across random seeds 0-9. The mean as well as standard deviation have been reported through plots for all experiments. All experiments were run using the publicly available Google Colaboratory. The experiments were run using CUDA enabled PyTorch on a T4 GPU with 15 GB memory.

Further, in this section, we also include some additional experiments. We first recreate the three plots provided in the main paper (NTK change till convergence, epochs till convergence and L_2 norm of representation difference) for the case of a single hidden layer neural network with ReLU activation:




Figure 2: For a fixed sample size N , we plot different quantities for varying network width M . We then vary N and plot: (a) NTK change till convergence (b) Training Epochs till convergence (c) Squared norm of difference between representations of neural network and corresponding kernel model.

As we can see in Figure 2, all three plots closely resemble the corresponding plots with TanH activation. This empirically validates that our claims hold for ReLU activation function as well, for which our derivation was not entirely rigorous (due to the non-differentiability of ReLU at zero). Next, we repeat the same for a 3 hidden layer neural network with ReLU activation:




Figure 3: For a fixed sample size N , we plot different quantities for varying network width M . We then vary N and plot: (a) NTK change till convergence (b) Training Epochs till convergence (c) Squared norm of difference between representations of neural network and corresponding kernel model.

While our proof is only for single-hidden layer neural networks, Figure 3 suggests that the analysis holds for deeper networks as well. We leave this for future work.