

**THE NONLOCAL HARNACK INEQUALITY
FOR ANTISYMMETRIC FUNCTIONS:
AN APPROACH VIA BOCHNER'S RELATION AND HARMONIC ANALYSIS**

SERENA DIPIERRO, MATEUSZ KWAŚNICKI, JACK THOMPSON, ENRICO VALDINOCI

Abstract. We revisit a Harnack inequality for antisymmetric functions that has been recently established for the fractional Laplacian and we extend it to more general nonlocal elliptic operators.

The new approach to deal with these problems that we propose in this paper leverages Bochner's relation, allowing one to relate a one-dimensional Fourier transform of an odd function with a three-dimensional Fourier transform of a radial function.

In this way, Harnack inequalities for odd functions, which are essentially Harnack inequalities of boundary type, are reduced to interior Harnack inequalities.

1. Introduction

One of the characterizing properties of classical harmonic functions is their “rigidity”: in spite of the fact that harmonic functions may exhibit, in general, different patterns, a common feature is that if, at a given point, a harmonic function “bends up” in a certain direction, then necessarily it has to “bend down” in another direction. This observation is typically formalised by the so-called maximum principle. Furthermore, the maximum principle is, in turn, quantified by the Harnack inequality [20] which asserts that the values of a positive harmonic function in a given ball are comparable (see [23] for a thorough introduction to the topic).

The growing interest recently surged in the study of nonlocal and fractional equations, especially in relation to fractional powers of the Laplace operator, has stimulated an intense investigation on the possible validity of Harnack-type inequalities in a nonlocal framework. Several versions of the Harnack inequality have been obtained for the fractional Laplacian as well as for more general nonlocal operators, see [3, 4, 31, 2, 30, 6, 33, 8, 18].

A striking difference between the classical and the fractional settings is that in the former a sign assumption on the solution is taken only in the region of interest, while in the latter such an assumption is known to be insufficient. Indeed, counterexamples to the validity of the nonlocal Harnack inequality in the absence of a global sign assumptions have been put forth in [22] (see also [5, Theorem 3.3.1] and [24, Theorem 2.2]). In fact, these counterexamples are just particular cases of the significant effect that the faraway oscillations may produce on the local patterns of solutions to nonlocal equations, see [10, 11] (in case of solutions which change sign, the nonlocal framework however allows for a Harnack inequality with a suitable integral remainder, see [24, Theorem 2.3]).

A rather prototypical situation in which the sign condition is violated is that of odd antisymmetric functions. In the nonlocal world, this situation frequently occurs, especially when dealing with moving planes and reflection methods. On these occasions, the lack of sign assumption needs to be replaced by bespoke maximum principles which carefully take into account the additional antisymmetrical structure of the problem under consideration, see [15, 21, 7, 16, 17, 9]. In particular, in the antisymmetric setting, the following nonlocal Harnack inequality has been established in [12]:

Theorem 1.1. *Suppose that Ω is an open set of \mathbb{R}^n , and let u be a function on \mathbb{R}^n such that u is $C^{2s+\delta}$ in Ω and antisymmetric, with*

$$\int_{\mathbb{R}^n} \frac{|x_1| |u(x)|}{(1 + |x|)^{n+2+2s}} dx < +\infty$$

and $x_1 u(x) \geq 0$ for every $x \in \mathbb{R}^n$.

Let c be a bounded Borel function on Ω . If

$$(-\Delta)^s u + cu = 0$$

in $\Omega_+ = \{x \in \Omega : x_1 > 0\}$, then for every compact subset K of Ω we have

$$\sup_{x \in K_0} \frac{u(x)}{x_1} \leq C(K, \Omega, \|c\|_{L^\infty(\Omega_+)}) \inf_{x \in K_0} \frac{u(x)}{x_1}, \quad (1.1)$$

where $K_0 = \{x \in K : x_1 \neq 0\}$.

In this article, we revisit the nonlocal Harnack inequality under a different perspective, providing a new, shorter proof of Theorem 1.1. This proof relies on the so-called Bochner's relation (or Hecke–Bochner identity), which links a one-dimensional Fourier transform of an odd function with a three-dimensional Fourier transform of a radial function: roughly speaking, for any function f on the positive reals (under a natural integrability condition), if $F(|Z|)$ denotes the 3-D Fourier transform of the 3-D radial function $f(|X|)$, then the 1-D Fourier transform of the 1-D function $xf(|x|)$ is equal to $izF(|z|)$. The complete statement of Bochner's relation is actually even more general (it works in higher dimensions too, and the variable x can be replaced by a homogeneous harmonic polynomial), see [32, page 72].

The gist of the argument that we present here is that Bochner's relation extends to the case of antisymmetric functions dealt with in Theorem 1.1. More specifically, the Fourier transform of an n -dimensional antisymmetric function $x_1 f(|x_1|, x')$ is equal to $iz_1 F(|z_1|, z')$, where $F(|Z|, z')$ is the $(n+2)$ -dimensional Fourier transform of $f(|X|, x')$, where X and Z are 3-dimensional vectors, x_1 and z_1 are real numbers, and x' and z' are $(n-1)$ -dimensional vectors.

Using the above property, one finds that the n -dimensional fractional Laplacian applied to $x_1 f(|x_1|, x')$ is equal to $x_1 g(|x_1|, x')$ if and only if the $(n+2)$ -dimensional fractional Laplacian applied to $f(|X|, x')$ is equal to $g(|X|, x')$ (in fact, this is true for an arbitrary function of the Laplacian, not just for its fractional powers).

This, in turn, enables one to apply the usual Harnack inequality in dimension $n+2$ to deduce a Harnack inequality (or, in a sense, a “boundary” Harnack inequality) in dimension n for antisymmetric functions. In this framework, Theorem 1.1 follows from a more classical result for positive solutions in dimension $n+2$.

For other applications of Bochner's relation to antisymmetric functions, see [13, Theorem 1], [26, Theorem 1.5 and Proposition 3.1] and [14].

Besides its intrinsic elegance and the conciseness of the techniques involved, one of the advantages of the methods leveraging Bochner's relation consists of the broad versatility of the arguments employed. In particular, the methodology employed allows us to extend the previous result in Theorem 1.1 to a more general class of nonlocal operators. To this end, given a positive integer n , we consider an operator L_n , acting on an appropriate class of functions $u : \mathbb{R}^n \rightarrow \mathbb{R}$, and taking the form

$$L_n u(x) = \int_{\mathbb{R}^n \setminus \{0\}} (u(x) - u(x+y) - \nabla u(x) \cdot y \chi_{B_1}(y)) j_n(|y|) dy, \quad (1.2)$$

where the kernel j_n satisfies the usual integrability condition

$$\int_{\mathbb{R}^n \setminus \{0\}} \min\{1, |x|^2\} j_n(|x|) dx < +\infty.$$

To proceed, we will need the following additional assumption:

$$j_n \text{ is differentiable and nonincreasing on } (0, \infty). \quad (1.3)$$

This allows us to introduce the operator L_{n+2} , defined by the same formula with kernel

$$j_{n+2}(r) = -\frac{j'_n(r)}{2\pi r}. \quad (1.4)$$

In this setting, we establish that:

Theorem 1.2. *Let L_n and L_{n+2} be given by (1.2), and suppose that conditions (1.3) and (1.4) are satisfied. Suppose that Ω is an open set of \mathbb{R}^n , and let u be a function on \mathbb{R}^n such that $u \in C^2(\Omega)$, $x_1^{-1}u \in L^\infty(\mathbb{R}^n)$, and u is antisymmetric with $x_1 u(x) \geq 0$ for all $x \in \mathbb{R}^n$.*

Let c be a bounded Borel function on Ω . If

$$L_n u + cu = 0$$

in $\Omega_+ = \{x \in \Omega : x_1 > 0\}$ and the operator L_{n+2} satisfies the Harnack inequality, then for every compact subset K of Ω we have

$$\sup_{x \in K_0} \frac{u(x)}{x_1} \leq C(K, \Omega, \|c\|_{L^\infty(\Omega_+)}) \inf_{x \in K_0} \frac{u(x)}{x_1}, \quad (1.5)$$

where $K_0 = \{x \in K : x_1 \neq 0\}$ and $C(K, \Omega, M)$ is the same as the constant in the Harnack inequality for L_{n+2} .

For the precise definition of an operator satisfying the Harnack inequality, see Definition 5.3. The Harnack inequality (with a scale-invariant constant) is known to hold for a large class of operators L_{n+2} , at least when $c = 0$; we refer to Remark 5.4 in Section 5 for further discussion.

The rest of the paper is organized as follows. In Section 2, we give some heuristic comments about the idea of reducing boundary Harnack inequalities to interior ones, to motivate the main strategy adopted in this paper, while, in Section 3, we recall Bochner's relation and frame it into a setting convenient for our purposes.

The techniques thus introduced are then applied to the case of the fractional Laplace operator in Section 4, leading to the proof of Theorem 1.1.

The case of general nonlocal operators will be explained in detail in Section 5.

2. Reducing boundary estimates to interior estimates: a heuristic discussion

The gist of the arguments that we exploit can be easily explained in the classical case of the Laplacian, in which all computations are straightforward, but already reveal a hidden higher dimensional geometric feature of the problem.

For this, we take coordinates $y \in \mathbb{R}^k$ and $z \in \mathbb{R}^m$. We use the notation $\rho = |y|$ and, given a smooth function u in a given domain of \mathbb{R}^{m+1} , for $\lambda \in \mathbb{R}$ we set

$$\tilde{v}(y, z) := |y|^\lambda u(|y|, z) = \rho^\lambda u(\rho, z).$$

Since, for every $j \in \{1, \dots, m\}$,

$$\rho \partial_{y_j} \rho = \frac{1}{2} \partial_{y_j} \rho^2 = \frac{1}{2} \partial_{y_j} (y_1^2 + \dots + y_k^2) = y_j,$$

and therefore

$$\nabla_y \rho = \frac{y}{\rho},$$

computing the Laplacian in cylindrical coordinates we find that

$$\begin{aligned} \Delta \tilde{v} &= \Delta_{y,z} \rho^\lambda u + \rho^\lambda \Delta_{y,z} u + 2 \nabla_{y,z} \rho^\lambda \cdot \nabla_{y,z} u \\ &= \Delta_y \rho^\lambda u + \rho^\lambda (\Delta_y u + \Delta_z u) + 2 \nabla_y \rho^\lambda \cdot \nabla_y u \\ &= (\lambda + k - 2) \lambda \rho^{\lambda-2} u + \rho^\lambda (\partial_\rho^2 u + (k-1) \rho^{-1} \partial_\rho u + \Delta_z u) \\ &\quad + 2(\lambda \rho^{\lambda-1} \nabla_y \rho) \cdot (\partial_\rho u \nabla_y \rho) \\ &= (\lambda + k - 2) \lambda \rho^{\lambda-2} u + (2\lambda + k - 1) \rho^{\lambda-1} \partial_\rho u + \rho^\lambda \partial_\rho^2 u + \rho^\lambda \Delta_z u. \end{aligned}$$

The choice $2\lambda + k - 1 = 0$ allows one to cancel the first order term. The additional choice $\lambda + k - 2 = 0$ makes the zero order term vanish. All in all, if $k = 3$ and $\lambda = -1$, we find that

$$\Delta \tilde{v} = \frac{\Delta u}{\rho}, \tag{2.1}$$

hence \tilde{v} is harmonic if and only if so is u .

For us, the interest of (2.1) is that, for instance, it allows us to recover a boundary Harnack inequality for the Laplace operator straightforwardly from the interior Harnack inequality. Specifically, *if c is a bounded Borel function on $B_1^+ := B_1 \cap \{x_1 > 0\} \subset \mathbb{R}^n$ and u is a nonnegative solution of $-\Delta u + cu = 0$ in B_1^+ , then*

$$\sup_{x \in B_{1/2}^+} \frac{u(x)}{x_1} \leq C \inf_{x \in B_{1/2}^+} \frac{u(x)}{x_1}, \tag{2.2}$$

where C is a positive constant depending only on n and $\|c\|_{L^\infty(B_1^+)}$.

In our setting, the proof of (2.2) can be obtained directly combining (2.1) and the classical Harnack inequality: namely, if we use the notation $\tilde{c}(y, z) := c(|y|, z)$, we deduce from (2.1) that

$$(-\Delta \tilde{v} + \tilde{c} \tilde{v})|y| = -\Delta u + cu = 0$$

whenever $(|y|, z) \in B_1$ and so, in particular, for all $(y, z) \in \mathbb{R}^3 \times \mathbb{R}^{n-1}$ such that $|(y, z)| < 1$.

This and the classical Harnack inequality lead to

$$\sup_{|(y,z)| < 1/2} \tilde{v}(y, z) \leq C \inf_{|(y,z)| < 1/2} \tilde{v}(y, z),$$

from which (2.2) plainly follows.

The strategy that we follow in this note is precisely to adapt this method to more general settings of nonlocal type. In this situation, additional symmetry structures have to be imposed on the solution, due to the nonlocal features of the operator. Moreover, the analog of (2.1) requires a series of more subtle arguments, since cylindrical coordinates are typically not easy to handle in a nonlocal framework, due to remote point interactions: this difficulty will be overcome by exploiting some classical tools from harmonic analysis, as described in detail in the forthcoming Section 3.

3. Bochner's relation and its consequences

3.1. Notation. Let $\ell \in \mathbb{N}$. As usual, a *solid harmonic polynomial* of degree ℓ is a homogeneous polynomial V on \mathbb{R}^n of degree ℓ (that is, $V(rx) = r^\ell V(x)$) satisfying the Laplace equation $\Delta V = 0$.

Below, we will also use symbols with a tilde to denote elements of $\mathbb{R}^{n+2\ell}$ and functions on $\mathbb{R}^{n+2\ell}$, while symbols without a tilde for vectors in \mathbb{R}^n and functions on \mathbb{R}^n . That is, for typographical convenience, we will usually (i.e., unless differently specified) write $x \in \mathbb{R}^n$ and $\tilde{x} \in \mathbb{R}^{n+2\ell}$, and similarly $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\tilde{f} : \mathbb{R}^{n+2\ell} \rightarrow \mathbb{R}$.

We use \mathcal{F}_n to denote the n -dimensional *Fourier transform*: if f is an integrable function on \mathbb{R}^n , then

$$\mathcal{F}_n f(\xi) = \int_{\mathbb{R}^n} f(x) e^{2\pi i \xi \cdot x} dx.$$

When we want to emphasize that we are taking the Fourier transform with respect to the variable x we also use the notation $\mathcal{F}_n^{(x)}$. For instance, if $f : \mathbb{R}^{2n} = \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$, to be written as $f(x, y)$, we have that, for each $y \in \mathbb{R}^n$,

$$\mathcal{F}_n^{(x)} f(\xi, y) = \int_{\mathbb{R}^n} f(x, y) e^{2\pi i \xi \cdot x} dx.$$

We are following here the convention on the Fourier transform on page 28 in [32].

We also consider L_n to be a Fourier multiplier on \mathbb{R}^n with radial symbol $\psi(|\xi|)$, that is

$$\mathcal{F}_n L_n f(\xi) = \psi(|\xi|) \mathcal{F}_n f(\xi) \tag{3.1}$$

for every Schwartz function f .

The case in which $\psi(r) = r^{2s}$ corresponds to the fractional Laplacian $L_n = (-\Delta)^s$, but the contents of this section work for general ψ with at most polynomial growth.

Let \mathcal{R} be a rotation on \mathbb{R}^n (that is, an orthogonal transformation of \mathbb{R}^n), and for a function f on \mathbb{R}^n denote $\mathcal{R}f(\xi) = f(\mathcal{R}\xi)$. By definition, we have the following transformation rule for the Fourier transform:

$$\mathcal{F}_n(\mathcal{R}f)(\xi) = \mathcal{R}(\mathcal{F}_n f)(\xi).$$

Radial functions are invariant under \mathcal{R} , and so

$$\psi(|\xi|) \mathcal{F}_n(\mathcal{R}f)(\xi) = \mathcal{R}[\psi(|\xi|) \mathcal{F}_n f(\xi)].$$

The inverse Fourier transform satisfies a similar transformation rule, so that

$$\mathcal{F}_n^{-1}[\psi(|\xi|) \mathcal{F}_n \mathcal{R}f(\xi)] = \mathcal{R}[\mathcal{F}_n^{-1}[\psi(|\xi|) \mathcal{F}_n f(\xi)]]$$

whenever f is, say, a Schwartz function. In view of (3.1), this means that

$$L_n \mathcal{R}f(\xi) = \mathcal{R} L_n f(\xi), \tag{3.2}$$

that is, L_n commutes with rotations.

Let us call a function f on \mathbb{R}^n isotropic with respect to the first k coordinates, or briefly k -isotropic, if $f(x) = f(y)$ whenever

$$|(x_1, x_2, \dots, x_k)| = |(y_1, y_2, \dots, y_k)| \quad \text{and} \quad (x_{k+1}, x_{k+2}, \dots, x_n) = (y_{k+1}, y_{k+2}, \dots, y_n).$$

We will only use this notion for $k = 3$: the proof of our main result involves 3-isotropic functions in \mathbb{R}^{n+2} .

A function f on \mathbb{R}^n is said to be *symmetric* (with respect to the first variable) if $f(-x_1, x_2, \dots, x_n) = f(x_1, x_2, \dots, x_n)$ (that is, if f is even with respect to the first variable; this coincides with the notion of a 1-isotropic function). Similarly, f is said to be *antisymmetric* (with respect to the first variable) if $f(-x_1, x_2, \dots, x_n) = -f(x_1, x_2, \dots, x_n)$. Clearly, f is symmetric if and only if $x_1 f(x)$ is antisymmetric.

If f is a k -isotropic function on \mathbb{R}^n , then f is invariant under every rotation \mathcal{R} which only acts on the first k coordinates and leaves the remaining $n-k$ coordinates unchanged: $\mathcal{R}f = f$. In view of (3.2), we have $\mathcal{R}(L_n f) = L_n(\mathcal{R}f) = L_n f$, that is, $L_n f$ is invariant under \mathcal{R} . In other words, L_n maps k -isotropic Schwartz functions into k -isotropic functions.

3.2. Fourier transforms of radial functions. Now we recall the tools from harmonic analysis which will come in handy for the development of the theory. The gist is the link of the Fourier transform of an n -dimensional radial function (multiplied by an appropriate harmonic polynomial) with that of its $(n + 2\ell)$ -dimensional counterpart.

For instance, roughly speaking, for any even function f on \mathbb{R} , if $\tilde{f} : \mathbb{R}^3 \rightarrow \mathbb{R}$ has the same profile of f (i.e., $\tilde{f}(\tilde{x}) = f(|\tilde{x}|)$ for each $\tilde{x} \in \mathbb{R}^3$), then the Fourier transform of the one-dimensional function $xf(x)$ coincides with $i\xi$ times the Fourier transform of the three-dimensional radial function \tilde{f} at $\tilde{\xi}$, as long as $\tilde{\xi} \in \mathbb{R}^3$ is such that $|\xi| = |\tilde{\xi}|$.

This fact holds true in higher dimensions as well, with suitable modifications, and goes under the name of Bochner's relation (or Hecke–Bochner identity). This will be our crucial tool and we now recall its precise statement:

Theorem 3.1 (Bochner's relation; Corollary on page 72 in [32]). *Let f and \tilde{f} be two radial Schwartz functions on \mathbb{R}^n and $\mathbb{R}^{n+2\ell}$, respectively, with the same profile function: $f(x) = \tilde{f}(\tilde{x})$ whenever $|x| = |\tilde{x}|$.*

Let V be a solid harmonic polynomial on \mathbb{R}^n of degree ℓ .

Then,

$$\mathcal{F}_n(Vf)(\xi) = i^\ell V(\xi) \mathcal{F}_{n+2\ell} \tilde{f}(\tilde{\xi}),$$

for every $\xi \in \mathbb{R}^n$ and $\tilde{\xi} \in \mathbb{R}^{n+2\ell}$ such that $|\tilde{\xi}| = |\xi|$.

In the statement of Theorem 3.1, the assumption on the profile function of f and \tilde{f} simply means that there exists a suitable $f_0 : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = f_0(|x|)$ and $\tilde{f}(\tilde{x}) = f_0(|\tilde{x}|)$ for all $x \in \mathbb{R}^n$ and $\tilde{x} \in \mathbb{R}^{n+2\ell}$ (the function f_0 is called in jargon “profile function”).

In our setting, we will only need Theorem 3.1 for $n = \ell = 1$ and $V(x) = x$. In this case Theorem 3.1 states that if f is an even function on \mathbb{R} and $\tilde{f}(\tilde{x}) = f(|x|)$ is the corresponding radial function on \mathbb{R}^3 , then

$$\mathcal{F}_1[xf(x)](\xi) = i\xi \mathcal{F}_3 \tilde{f}(\xi, 0, 0). \quad (3.3)$$

We have the following immediate extension of Theorem 3.1.

Lemma 3.2. Consider a function $f : \mathbb{R}^n = \mathbb{R} \times \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, to be written as $f(x, y)$, where $x \in \mathbb{R}$ and $y \in \mathbb{R}^{n-1}$.

Assume that f is a Schwartz function on \mathbb{R}^n which is symmetric with respect to the variable x (i.e., $f(-x, y) = f(x, y)$ for all $x \in \mathbb{R}$ and $y \in \mathbb{R}^{n-1}$).

Let $\tilde{f} : \mathbb{R}^{n+2} = \mathbb{R}^3 \times \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, to be written as $\tilde{f}(\tilde{x}, y)$, where $\tilde{x} \in \mathbb{R}^3$ and $y \in \mathbb{R}^{n-1}$, be a 3-isotropic function given by

$$\tilde{f}(\tilde{x}, y) = f(|\tilde{x}|, y).$$

Then,

$$\mathcal{F}_n[xf(x, y)](\xi, \eta) = i\xi \mathcal{F}_{n+2}\tilde{f}((\xi, 0, 0), \eta).$$

Proof. Given $y \in \mathbb{R}^{n-1}$, we apply Bochner's relation (3.3) with respect to $x \in \mathbb{R}$ to find that

$$\mathcal{F}_1^{(x)}[xf(x, y)](\xi, y) = i\xi \mathcal{F}_3^{(\tilde{x})}\tilde{f}(\xi, y).$$

It is now sufficient to apply the Fourier transform with respect to $y \in \mathbb{R}^{n-1}$ for each fixed $\xi \in \mathbb{R}$. \square

From Lemma 3.2 we obtain the following result.

Lemma 3.3. In the assumptions of Lemma 3.2, for every $x \in \mathbb{R}$ and $y \in \mathbb{R}^{n-1}$ we have that

$$L_n[xf(x, y)] = xL_{n+2}\tilde{f}((x, 0, 0), y).$$

Proof. Let us denote

$$\tilde{g}(\tilde{x}, y) = L_{n+2}\tilde{f}(\tilde{x}, y)$$

and

$$g(x, y) = \tilde{g}((x, 0, 0), y) = L_{n+2}\tilde{f}((x, 0, 0), y).$$

Since L_{n+2} maps 3-invariant functions into 3-invariant functions, \tilde{g} is 3-invariant.

Let now $(\xi, \eta) \in \mathbb{R} \times \mathbb{R}^{n-1}$ and let $\tilde{\xi} \in \mathbb{R}^3$ be such that $|\xi| = |\tilde{\xi}|$. Using, in order, the definition of L_n in (3.1), Lemma 3.2, once again the definition of L_{n+2} in (3.1), the definition of \tilde{g} , and once again Lemma 3.2, we find that

$$\begin{aligned} \mathcal{F}_n L_n[xf(x, y)](\xi, \eta) &= \psi(|(\xi, \eta)|) \mathcal{F}_n[xf(x, y)](\xi, \eta) \\ &= \psi(|(\xi, \eta)|) i\xi \mathcal{F}_{n+2}\tilde{f}(\xi, \eta) \\ &= i\xi \psi(|(\tilde{\xi}, \eta)|) \mathcal{F}_{n+2}\tilde{f}(\tilde{\xi}, \eta) \\ &= i\xi \mathcal{F}_{n+2}L_{n+2}\tilde{f}(\tilde{\xi}, \eta) \\ &= i\xi \mathcal{F}_{n+2}\tilde{g}(\tilde{\xi}, \eta) \\ &= \mathcal{F}_n[xg(x, y)](\xi, \eta). \end{aligned}$$

Thus, by inverting the Fourier transform,

$$L_n[xf(x, y)] = xg(x, y).$$

Combining this and the definition of g , we conclude that

$$L_n[xf(x, y)] = xL_{n+2}\tilde{f}((x, 0, 0), y),$$

as desired. \square

In the same vein, one shows the following more general result (we omit the proof since it can be obtained via the same argument as above, by utilizing the general identity in Theorem 3.1 instead of the ones specialized for the case $V(x) = x$).

Lemma 3.4. *Let V be a solid harmonic polynomial of degree ℓ on \mathbb{R}^k . Let $f(x, y)$, where $x \in \mathbb{R}^k$ and $y \in \mathbb{R}^{n-k}$, be a Schwartz function on \mathbb{R}^n which is isotropic with respect to the first variable x : $f(x, y) = f(x', y)$ whenever $|x| = |x'|$. Let $\tilde{f}(\tilde{x}, y)$, where $\tilde{x} \in \mathbb{R}^{k+2\ell}$ and $y \in \mathbb{R}^{n-k}$, be the corresponding Schwartz function on $\mathbb{R}^{n+2\ell}$, given by*

$$\tilde{f}(\tilde{x}, y) = f(x, y)$$

whenever $|\tilde{x}| = |x|$.

Then

$$L_n[V(x)f(x, y)] = V(x)L_{n+2\ell}\tilde{f}((x, 0), y),$$

where $(x, 0) \in \mathbb{R}^{k+2\ell}$ stands for the vector $x \in \mathbb{R}^k$ padded with 2ℓ zeroes.

4. Fractional Laplacians

4.1. Notation. Now we specialize the result of Section 3 to the case of the fractional Laplacian of order $s \in (0, 1)$, corresponding to the choice $\psi(r) = r^{2s}$ in Lemma 3.4. To this end, we introduce some further notation.

If f and g are functions on \mathbb{R}^n , we write

$$\langle f, g \rangle_{\mathbb{R}^n} = \int_{\mathbb{R}^n} f(x)g(x) dx.$$

We use the notation $f \in C^\alpha$ for non-integer $\alpha > 0$, meaning that $f \in C^{k,\beta}$ where $\alpha = k + \beta$ with $k \in \mathbb{N}$ and $\beta \in (0, 1)$.

4.2. Fractional Laplacian identities and a Harnack inequality. We now translate the results of Section 3 into the notation that we have just introduced.

Corollary 4.1. *Let f be a Schwartz function on \mathbb{R}^n , symmetric with respect to the first variable. Let \tilde{f} be the corresponding 3-isotropic Schwartz function on \mathbb{R}^{n+2} , given by*

$$\tilde{f}(\tilde{x}) = f\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right).$$

Finally, let $g(x) = x_1 f(x)$. Then

$$(-\Delta)^s g(x) = x_1 (-\Delta)^s \tilde{f}((x_1, 0, 0), x_2, \dots, x_n).$$

Proof. This follows directly from Lemma 3.3: indeed, choosing $\psi(r) = r^{2s}$ in (3.1) gives that L_n is the fractional Laplacian acting on functions of n variables and accordingly, for every $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$,

$$(-\Delta)^s g(x) = (-\Delta)^s [x_1 f(x)] = x_1 (-\Delta)^s \tilde{f}((x_1, 0, 0), x_2, \dots, x_n). \quad \square$$

Remark 4.2. The identity in Corollary 4.1 can be alternatively proved by a rather straightforward direct calculation. This involves the formula

$$\begin{aligned} r^2 \int_{\mathbb{S}^2} (\alpha^2 + (rz_1 - \beta)^2 + (rz_2)^2 + (rz_3)^2)^{-1-\gamma} \mathcal{H}^2(dz) \\ = \frac{1}{2\beta\gamma} ((\alpha^2 + (r - \beta)^2)^{-\gamma} - (\alpha^2 + (r + \beta)^2)^{-\gamma}), \end{aligned}$$

where \mathbb{S}^2 is the unit sphere in \mathbb{R}^3 and \mathcal{H}^2 is the surface measure. One substitutes $\alpha = \sqrt{(x_2 - y_2)^2 + \dots + (x_n - y_n)^2}$, $\beta = x_1$, $r = |y_1|$, and $\gamma = \frac{n}{2} + s$ and then integrates with respect to $y \in \mathbb{R}^n$; we omit the details.

An extension of Corollary 4.1 to arbitrary antisymmetric functions is rather straightforward: in particular, in item (c) below we consider the extension of the fractional Laplace operator on the class of antisymmetric functions, as described in [12].

Theorem 4.3. *Suppose that Ω is an open set of \mathbb{R}^n and $\delta > 0$. Let u be a function on \mathbb{R}^n such that u is $C^{2s+\delta}$ in Ω and antisymmetric, with*

$$\int_{\mathbb{R}^n} \frac{|x_1| |u(x)|}{(1+|x|)^{n+2+2s}} dx < +\infty. \quad (4.1)$$

Let \tilde{v} be the corresponding function on \mathbb{R}^{n+2} given by

$$\tilde{v}(\tilde{x}) = \frac{u(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}.$$

Then:

(a) if $\Omega_0 = \{x \in \Omega : x_1 \neq 0\}$, then the function \tilde{v} is $C_{\text{loc}}^{2s+\delta}$ in

$$\tilde{\Omega}_0 = \left\{ \tilde{x} \in \mathbb{R}^{n+2} : \left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2} \right) \in \Omega_0 \right\};$$

(b) we have that

$$\int_{\mathbb{R}^{n+2}} \frac{|\tilde{v}(\tilde{x})|}{(1+|\tilde{x}|)^{n+2+2s}} d\tilde{x} < +\infty;$$

(c) for every $x \in \Omega_0$, we have that¹

$$(-\Delta)^s u(x) = x_1 (-\Delta)^s \tilde{v}((x_1, 0, 0), x_2, \dots, x_n); \quad (4.2)$$

(d) for an arbitrary C_c^∞ function g on Ω , we have

$$\langle u, (-\Delta)^s g \rangle_{\mathbb{R}^n} = (2\pi)^{-1} \langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}}, \quad (4.3)$$

where

$$\tilde{f}(\tilde{x}) = \frac{g_A(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}$$

and

$$g_A(x) = \frac{1}{2} (g(x_1, \dots, x_n) - g(-x_1, x_2, \dots, x_n))$$

is the anti-symmetric part of g ;

(e) for an arbitrary C_c^∞ function \tilde{f} on

$$\tilde{\Omega} = \left\{ \tilde{x} \in \mathbb{R}^{n+2} : \left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2} \right) \in \Omega \right\},$$

we have

$$\langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}} = 2\pi \langle u, (-\Delta)^s g \rangle_{\mathbb{R}^n}, \quad (4.4)$$

where $g(x) = x_1 f(x)$ and f is given in terms of the 3-isotropic symmetrization of \tilde{f} by

$$f(x) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \tilde{f}(|x_1|z, x_2, \dots, x_n) \mathcal{H}^2(dz); \quad (4.5)$$

here \mathbb{S}^2 is the unit sphere in \mathbb{R}^3 and \mathcal{H}^2 is the surface measure.

¹In general, without the antisymmetry assumption, (4.1) is not a strong enough assumption on the decay of u for the fractional Laplacian to be well-defined. To rectify this issue, we have to use the antisymmetric fractional Laplacian, defined according to [12, Definition 1.2].

Remark 4.4. Note that, in Theorem 4.3 (a), the conclusion $\tilde{v} \in C_{\text{loc}}^{2s+\delta}(\tilde{\Omega}_0)$ cannot be relaxed to $\tilde{v} \in C^{2s+\delta}(\tilde{\Omega}_0)$. Indeed, consider $\Omega = (0, 1) \times \mathbb{R}^{n-1}$ and $u(x) = x_1^{2s+\delta}$ for $x_1 \geq 0$ extended antisymmetrically to all of \mathbb{R}^n . Then $\Omega_0 = \Omega$, $\tilde{\Omega}_0 = (B_1^3 \setminus \{0\}) \times \mathbb{R}^{n-1}$ (here B_1^3 denotes the unit ball in \mathbb{R}^3), and $u \in C^{2s+\delta}(\Omega_0)$, but

$$\tilde{v}(\tilde{x}) = (\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2)^{\frac{2s+\delta-1}{2}}$$

which is not in $C^{2s+\delta}(\tilde{\Omega}_0)$.

Proof of Theorem 4.3. To prove assertion (a), observe that, for all $\tilde{x} \in \tilde{\Omega}_0$,

$$\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} \neq 0.$$

Hence, the map from $\tilde{\Omega}_0 \rightarrow \mathbb{R}$ defined by $\tilde{x} \mapsto \frac{1}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}$, is in $C_{\text{loc}}^\infty(\tilde{\Omega}_0)$. Moreover,

$$\tilde{x} \mapsto u\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right)$$

is in $C^{2s+\delta}(\tilde{\Omega}_0)$ which implies $\tilde{v} \in C_{\text{loc}}^{2s+\delta}(\tilde{\Omega}_0)$. This proves (a).

Additionally,

$$\begin{aligned} \mathcal{I} &:= \int_{\mathbb{R}^{n+2}} \frac{|\tilde{v}(\tilde{x})|}{(1 + |\tilde{x}|)^{n+2+2s}} d\tilde{x} = \int_{\mathbb{R}^{n+2}} \frac{\left|u\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right)\right|}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} (1 + |\tilde{x}|)^{n+2+2s}} d\tilde{x} \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^{n-1}} \frac{|u(|a|, b)|}{|a| (1 + \sqrt{|a|^2 + |b|^2})^{n+2+2s}} da db, \end{aligned} \quad (4.6)$$

Now we observe that if $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ then, integrating in spherical coordinates, we get

$$\int_{\mathbb{R}^3} \Phi(|x|) dx = 4\pi \int_0^{+\infty} \tau^2 \Phi(\tau) d\tau. \quad (4.7)$$

Now, gathering (4.6) and (4.7), we see that

$$\mathcal{I} = \int_0^{+\infty} \int_{\mathbb{R}^{n-1}} \frac{\tau |u(\tau, b)|}{(1 + \sqrt{|\tau|^2 + |b|^2})^{n+2+2s}} db d\tau = \int_{\mathbb{R}^n} \frac{|x_1| |u(x)|}{(1 + |x|)^{n+2+2s}} dx,$$

from which (b) plainly follows.

To prove the claim in (c), we notice that if $f(x) = x_1^{-1}u(x)$ is a Schwartz function, item (c) reduces to Corollary 4.1. The extension to general u follows by mollification and cutting off. This approximation procedure is not completely standard in our scenario, so we have provided the details.

Let

$$\begin{aligned} \mathcal{A}_s(\mathbb{R}^n) &= \{u : \mathbb{R}^n \rightarrow \mathbb{R} \text{ s.t. } u \text{ is measurable, antisymmetric, and } \|u\|_{\mathcal{A}_s(\mathbb{R}^n)} < +\infty\}, \\ \mathcal{L}_s(\mathbb{R}^{n+2}) &= \{\tilde{v} : \mathbb{R}^{n+2} \rightarrow \mathbb{R} \text{ s.t. } \tilde{v} \text{ is measurable and } \|\tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})} < +\infty\}, \end{aligned}$$

where

$$\begin{aligned} \|u\|_{\mathcal{A}_s(\mathbb{R}^n)} &= \int_{\mathbb{R}^n} \frac{|x_1| |u(x)|}{(1 + |x|)^{n+2+2s}} dx, \\ \|\tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})} &= \int_{\mathbb{R}^{n+2}} \frac{|\tilde{v}(\tilde{x})|}{(1 + |\tilde{x}|)^{n+2+2s}} d\tilde{x}. \end{aligned}$$

Let η_ε be a standard mollifier on \mathbb{R}^{n+2} , that is η_1 is a smooth, nonnegative, compactly supported function with integral 1, and $\eta_\varepsilon(\tilde{x}) = \varepsilon^{-n-2}\eta_1(\tilde{x}/\varepsilon)$. We have $u \in \mathcal{A}_s(\mathbb{R}^n)$, and $\tilde{v} \in \mathcal{L}_s(\mathbb{R}^{n+2})$ by (b). Define

$$\tilde{v}_k(\tilde{x}) = (\tilde{v} * \eta_{1/k})(\tilde{x})\eta_k(\tilde{x}),$$

and

$$u_k(x) = x_1\tilde{v}_k((x_1, 0, 0), x_2, x_3, \dots, x_n).$$

Note that \tilde{v}_k and u_k are smooth and have compact support, so, in particular, they are Schwartz functions. By a standard estimate², we have $\tilde{v}_k \in \mathcal{L}_s(\mathbb{R}^{n+2})$ and

$$\|\tilde{v}_k - \tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})} \rightarrow 0$$

as $k \rightarrow \infty$. We have seen in the proof of (b) that for an arbitrary antisymmetric function u on \mathbb{R}^n and the corresponding function \tilde{v} on \mathbb{R}^{n+2} , $u \in \mathcal{A}_s(\mathbb{R}^n)$ if and only if $\tilde{v} \in \mathcal{L}_s(\mathbb{R}^{n+2})$, and in this case $\|u\|_{\mathcal{A}_s(\mathbb{R}^n)} = \|\tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})}$. In particular, since $\tilde{v}_k \in \mathcal{L}_s(\mathbb{R}^{n+2})$, we have $u_k \in \mathcal{A}_s(\mathbb{R}^n)$, and additionally

$$\|u_k - u\|_{\mathcal{A}_s(\mathbb{R}^n)} = \|\tilde{v}_k - \tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})}.$$

Furthermore, since $x_1^{-1}u \in C_{\text{loc}}^{2s+\delta}(\Omega_0)$, we have that $u_k \rightarrow u$ in $C_{\text{loc}}^{2s+\delta}(\Omega_0)$. These two properties and [12, Lemma 2.3] imply that $(-\Delta)^s u_k \rightarrow (-\Delta)^s u$ in Ω_0 . Next, $\tilde{v}_k \rightarrow v$ in $C_{\text{loc}}^{2s+\delta}(\tilde{\Omega}_0)$ using (a). Moreover,

$$\|\tilde{v}_k - \tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})} \rightarrow 0$$

as $k \rightarrow \infty$, so standard estimates for the fractional Laplacian imply $(-\Delta)^s \tilde{v}_k \rightarrow (-\Delta)^s \tilde{v}$ in $\tilde{\Omega}_0$. Using (c) for Schwartz functions u_k and \tilde{v}_k , and combining it with the above two convergence results, we conclude that (c) holds in the general case.

We now prove assertions (d) and (e). For (d), since \tilde{f} is 3-isotropic, also $(-\Delta)^s \tilde{f}$ is 3-isotropic:

$$(-\Delta)^s \tilde{f}(\tilde{x}) = (-\Delta)^s \tilde{f}\left(\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, 0, 0\right), \tilde{x}_4, \dots, \tilde{x}_{n+2}\right),$$

so Corollary 4.1 implies that

$$(-\Delta)^s \tilde{f}(\tilde{x}) = \frac{(-\Delta)^s g_A(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}.$$

Hence, using the definition of \tilde{v} , we have

$$\begin{aligned} & \langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}} \\ &= \int_{\mathbb{R}^{n+2}} \frac{u(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}) (-\Delta)^s g_A(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} d\tilde{x} \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^{n-1}} \frac{u(|a|, b) (-\Delta)^s g_A(|a|, b)}{|a|^2} db da. \end{aligned}$$

Integration in spherical coordinates leads to

$$\langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}} = 4\pi \int_0^{+\infty} \int_{\mathbb{R}^{n-1}} u(\tau, b) (-\Delta)^s g_A(\tau, b) db d\tau.$$

²The proof is very similar to the one for standard L^1 spaces. When \tilde{v} is continuous and compactly supported, an explicit calculation gives a bound for $\|\tilde{v}_k - \tilde{v}\|_{\mathcal{L}_s(\mathbb{R}^{n+2})}$ in terms of the modulus of continuity of \tilde{v} . For a general \tilde{v} , one approximates \tilde{v} by a continuous, compactly supported function in $\mathcal{L}_s(\mathbb{R}^{n+2})$. For the sake of completeness, we have included the proof in Appendix (A).

Then, since both u and $(-\Delta)^s g_A$ are antisymmetric,

$$\begin{aligned} \langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}} &= 2\pi \int_{-\infty}^{+\infty} \int_{\mathbb{R}^{n-1}} u(\tau, b) (-\Delta)^s g_A(\tau, b) db d\tau \\ &= 2\pi \langle u, (-\Delta)^s g \rangle_{\mathbb{R}^n}. \end{aligned}$$

For (e), we begin by assuming that \tilde{f} is 3-isotropic and deal with the general case later. Since \tilde{f} is 3-isotropic, Corollary 4.1 applies to \tilde{f} , f and g : we have

$$(-\Delta)^s \tilde{f}(\tilde{x}) = \frac{(-\Delta)^s g(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}.$$

Using spherical coordinates,

$$\begin{aligned} \langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}} &= \int_{\mathbb{R}^{n+2}} \frac{u(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}) (-\Delta)^s g(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} d\tilde{x} \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^{n-1}} \frac{u(|a|, b) (-\Delta)^s g(|a|, b)}{|a|^2} db da \\ &= 4\pi \int_0^{+\infty} \int_{\mathbb{R}^{n-1}} u(\tau, b) (-\Delta)^s g(\tau, b) db d\tau. \end{aligned}$$

Since both u and $(-\Delta)^s g$ are symmetric, we have

$$\langle \tilde{v}, (-\Delta)^s \tilde{f} \rangle_{\mathbb{R}^{n+2}} = 2\pi \int_{-\infty}^{+\infty} \int_{\mathbb{R}^{n-1}} u(\tau, b) (-\Delta)^s g(\tau, b) db d\tau = 2\pi \langle u, (-\Delta)^s g \rangle_{\mathbb{R}^n}.$$

Now, if \tilde{f} is not 3-isotropic, let

$$\tilde{F}(\tilde{x}) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \tilde{f}\left(z\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right) \mathcal{H}^2(dz)$$

be the 3-isotropic projection of \tilde{f} . By the definition of f ,

$$\tilde{F}(\tilde{x}) = f\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right),$$

and so, since we already proved the result for 3-isotropic functions, we have

$$\langle \tilde{v}, (-\Delta)^s \tilde{F} \rangle_{\mathbb{R}^{n+2}} = 2\pi \langle u, (-\Delta)^s g \rangle_{\mathbb{R}^n}.$$

Therefore, it remains to prove that

$$\langle \tilde{v}, (-\Delta)^s (\tilde{f} - \tilde{F}) \rangle_{\mathbb{R}^{n+2}} = 0. \quad (4.8)$$

Denote $\tilde{G} = \tilde{f} - \tilde{F}$. Then, by the definition of \tilde{F} ,

$$\int_{\mathbb{S}^2} \tilde{G}(\tau z, \tilde{x}_4, \dots, \tilde{x}_{n+2}) \mathcal{H}^2(dz) = 0$$

for every $\tau > 0$ and $(\tilde{x}_4, \dots, \tilde{x}_{n+2}) \in \mathbb{R}^{n-1}$. Thus, \tilde{G} is orthogonal to every 3-isotropic function: if \tilde{u} is 3-isotropic, then integration in spherical coordinates gives

$$\begin{aligned} \langle \tilde{u}, \tilde{G} \rangle &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^{n-1}} \tilde{u}(a, b) \tilde{G}(a, b) db da \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^{n-1}} \tilde{u}(|a|, 0, 0, b) \tilde{G}(a, b) db da \\ &= \int_0^{+\infty} \int_{\mathbb{S}^2} \int_{\mathbb{R}^{n-1}} \tau^2 \tilde{u}(\tau, 0, 0, b) \tilde{G}(\tau z, b) db \mathcal{H}^2(dz) d\tau \\ &= \int_0^{+\infty} \int_{\mathbb{R}^{n-1}} \tau^2 \tilde{u}(\tau, 0, 0, b) \left(\int_{\mathbb{S}^2} \tilde{G}(\tau z, b) \mathcal{H}^2(dz) \right) db d\tau = 0. \end{aligned}$$

We already know that $(-\Delta)^s$ maps 3-isotropic Schwartz functions into 3-isotropic functions. By approximation, the class of 3-isotropic functions in $L^2(\mathbb{R}^{n+2})$ forms an invariant subspace of $(-\Delta)^s$. Therefore, also its orthogonal complement in $L^2(\mathbb{R}^n)$ is invariant. Since \tilde{G} is orthogonal to the class of 3-isotropic functions, we conclude that $(-\Delta)^s \tilde{G}$ has the same property, and (4.8) follows. \square

In our framework, a natural application³ of Theorem 4.3 leads to the proof of Theorem 1.1:

Proof of Theorem 1.1. With the notation of Theorem 4.3, by item (c) of this result we have

$$(-\Delta)^s u(x) = x_1 (-\Delta)^s \tilde{v}((x_1, 0, 0), x_2, \dots, x_n)$$

for $x \in \Omega_0$, or equivalently⁴

$$\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} (-\Delta)^s \tilde{v}(\tilde{x}) = (-\Delta)^s u \left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2} \right), \quad (4.9)$$

for $\tilde{x} \in \tilde{\Omega}_0$ (recall that $(-\Delta)^3 \tilde{v}$ is 3-isotropic).

Therefore, if $x = (\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})$, we have

$$\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} \left((-\Delta)^s \tilde{v}(\tilde{x}) + c(x) \tilde{v}(\tilde{x}) \right) = (-\Delta)^s u(x) + c(x) u(x) = 0,$$

that is, setting $\tilde{c}(\tilde{x}) = c(x)$,

$$(-\Delta)^s \tilde{v}(\tilde{x}) + \tilde{c}(\tilde{x}) \tilde{v}(\tilde{x}) = 0 \quad (4.10)$$

in $\tilde{\Omega}_0$. In fact, the above identity holds in the weak sense in $\tilde{\Omega}$: this follows directly from item (e) of Theorem 4.3.

Since \tilde{v} is nonnegative in \mathbb{R}^{n+2} , the standard Harnack inequality for weak solutions of the Schrödinger equation (4.10) implies that for every compact subset \tilde{K} of $\tilde{\Omega}$, we have

$$\operatorname{ess\,sup}_{\tilde{K}} \tilde{v} \leq C(\tilde{K}, \tilde{\Omega}, \|\tilde{c}\|_{L^\infty(\tilde{\Omega})}) \operatorname{ess\,inf}_{\tilde{K}} \tilde{v}.$$

Applying this to $\tilde{K} = \{ \tilde{x} \in \mathbb{R}^{n+2} : (\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}) \in K \}$ with K a compact subset of Ω , we obtain

$$\operatorname{ess\,sup}_{x \in K} \frac{u(x)}{x_1} \leq C(K, \Omega, \|c\|_{L^\infty(\Omega_+)}) \operatorname{ess\,inf}_{x \in K} \frac{u(x)}{x_1},$$

³When $s = 1$, Theorem 1.1 boils down to (2.2).

⁴One can compare (2.1) with (4.9). Namely, (4.9) recovers (2.1) when $s = 1$ with the notation $y = (\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ and $z = (\tilde{x}_4, \dots, \tilde{x}_{n+2})$.

which is equivalent to (1.1). \square

5. More general nonlocal operators

In this section, we generalise the results obtained in the specific setting of the fractional Laplacian to the more general class of operators L_n presented in (1.2).

Given an open set $\Omega \subset \mathbb{R}^n$ and a function in $u \in C_{\text{loc}}^2(\Omega) \cap L^\infty(\mathbb{R}^n)$, by a standard computation (see, for example, the computation in [5, p. 9]), $L_n u(x)$ is defined for all $x \in \Omega$ and, moreover, $L_n u$ is bounded on compact subsets of Ω . Furthermore, the Fourier symbol of L_n is

$$\begin{aligned} \Psi_n(\xi) &= \int_{\mathbb{R}^n} (1 - e^{i\xi \cdot x} + \xi \cdot x \chi_{(0,1)}(|x|)) j_n(|x|) dx \\ &= \int_0^\infty \int_{\mathbb{S}^{n-1}} (1 - e^{ir\xi \cdot \omega} + ir(\xi \cdot \omega) \chi_{(0,1)}(r)) j_n(r) r^{n-1} \mathcal{H}^{n-1}(d\omega) dr \\ &= \frac{1}{2} \int_0^\infty \int_{\mathbb{S}^{n-1}} (2 - e^{ir\xi \cdot \omega} - e^{-ir\xi \cdot \omega}) j_n(r) r^{n-1} \mathcal{H}^{n-1}(d\omega) dr \\ &= \int_0^\infty \int_{\mathbb{S}^{n-1}} (1 - \cos(r\xi \cdot \omega)) j_n(r) r^{n-1} \mathcal{H}^{n-1}(d\omega) dr \\ &= \mathcal{H}^{n-2}(\mathbb{S}^{n-2}) \int_0^\infty \int_{-1}^1 (1 - \cos(rs|\xi|)) (1 - s^2)^{\frac{n-3}{2}} j_n(r) r^{n-1} ds dr \\ &= (2\pi)^{n/2} \int_0^\infty \left(\frac{1}{2^{n/2-1} \Gamma(\frac{n}{2})} - (r|\xi|)^{1-n/2} J_{n/2-1}(r|\xi|) \right) r^{n-1} j_n(r) dr, \end{aligned}$$

where J_ν is the Bessel function of the first kind; see [27, Chapter 10]. Hence, it follows that Ψ_n is rotationally invariant, that is $\Psi_n(\xi) = \psi(|\xi|)$, with ψ given as the Hankel transform

$$\psi(\tau) = (2\pi)^{n/2} \int_0^\infty \left(\frac{1}{2^{n/2-1} \Gamma(\frac{n}{2})} - (r\tau)^{1-n/2} J_{n/2-1}(r\tau) \right) r^{n-1} j_n(r) dr, \quad (5.1)$$

It is no mistake that ψ has no subscript n : under our assumptions (1.3) and (1.4) it turns out that the Fourier symbol Ψ_{n+2} of the operator L_{n+2} has the same profile function ψ . There are many different ways to prove this fact; for instance, it is a relatively simple consequence of Bochner's relation. A direct proof involving well-known properties of Bessel functions is perhaps even shorter: since $j_n(r) = 2\pi \int_r^\infty t j_{n+2}(t) dt$, we have

$$\begin{aligned} \psi(\tau) &= (2\pi)^{n/2+1} \int_0^\infty \int_r^\infty \left(\frac{1}{2^{n/2-1} \Gamma(\frac{n}{2})} - (r\tau)^{1-n/2} J_{n/2-1}(r\tau) \right) r^{n-1} t j_{n+2}(t) dt dr \\ &= (2\pi)^{n/2+1} \int_0^\infty \int_0^t \left(\frac{1}{2^{n/2-1} \Gamma(\frac{n}{2})} - (r\tau)^{1-n/2} J_{n/2-1}(r\tau) \right) r^{n-1} t j_{n+2}(t) dr dt \\ &= (2\pi)^{n/2+1} \int_0^\infty \left(\frac{1}{2^{n/2} \Gamma(\frac{n}{2} + 1)} - (t\tau)^{-n/2} J_{n/2}(t\tau) \right) t^{n+1} j_{n+2}(t) dt, \end{aligned}$$

which is exactly the same expression as (5.1) with n replaced by $n + 2$; here we have used Fubini's theorem and [27, Equation 10.22.1].

Therefore, Bochner's relation and its consequences proved in Section 3 apply to this setting. The previously established theory will give us information about L_n given information about L_{n+2} , namely that if L_{n+2} admits the Harnack inequality then L_n admits the antisymmetric Harnack inequality.

Corollary 5.1. *Let f be a symmetric Schwartz function on \mathbb{R}^n . Let \tilde{f} be the corresponding 3-isotropic Schwartz function on \mathbb{R}^{n+2} , given by*

$$\tilde{f}(\tilde{x}) = f\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right).$$

Finally, let $g(x) = x_1 f(x)$. Then

$$L_n g(x) = x_1 L_{n+2} \tilde{f}((x_1, 0, 0), x_2, \dots, x_n).$$

Proof. The proof is a consequence of Lemma 3.3. Indeed, recall that the symbol ψ given in (5.1) is rotationally invariant, and operators L_n and L_{n+2} satisfy formula (3.1). Accordingly, for every $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$,

$$L_n g(x) = L_n(x_1 f(x)) = x_1 L_{n+2} \tilde{f}((x_1, 0, 0), x_2, \dots, x_n),$$

as desired. \square

Recall that L_{n+2} maps 3-isotropic functions to 3-isotropic functions. We now provide an extension of Corollary 5.1 to antisymmetric functions. This is the counterpart of Theorem 4.3 for general operators.

Theorem 5.2. *Suppose that Ω is an open set of \mathbb{R}^n . Let u be an antisymmetric function on \mathbb{R}^n such that $u \in C_{\text{loc}}^2(\Omega)$, $x_1^{-1}u \in L^\infty(\mathbb{R}^n)$, and let \tilde{v} be the corresponding function on \mathbb{R}^{n+2} given by*

$$\tilde{v}(\tilde{x}) = \frac{u(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}.$$

Then:

(a) if $\Omega_0 = \{x \in \Omega : x_1 \neq 0\}$ and

$$\tilde{\Omega}_0 = \left\{ \tilde{x} \in \mathbb{R}^{n+2} : \left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2} \right) \in \Omega_0 \right\};$$

then the function \tilde{v} is $C_{\text{loc}}^2(\tilde{\Omega}_0) \cap L^\infty(\mathbb{R}^{n+2})$.

(b) for every $x \in \Omega_0$, we have that

$$L_n u(x) = x_1 L_{n+2} \tilde{v}((x_1, 0, 0), x_2, \dots, x_n); \quad (5.2)$$

(c) for an arbitrary C_c^∞ function g on Ω , we have

$$\langle u, L_n g \rangle_{\mathbb{R}^n} = (2\pi)^{-1} \langle \tilde{v}, L_{n+2} \tilde{f} \rangle_{\mathbb{R}^{n+2}}, \quad (5.3)$$

where

$$\tilde{f}(\tilde{x}) = \frac{g_A(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})}{\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}}$$

and

$$g_A(x) = \frac{1}{2} (g(x_1, \dots, x_n) - g(-x_1, x_2, \dots, x_n));$$

(d) for an arbitrary C_c^∞ function \tilde{f} on

$$\tilde{\Omega} = \left\{ \tilde{x} \in \mathbb{R}^{n+2} : \left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2} \right) \in \Omega \right\},$$

we have

$$\langle \tilde{v}, L_{n+2} \tilde{f} \rangle_{\mathbb{R}^{n+2}} = 2\pi \langle u, L_n g \rangle_{\mathbb{R}^n}, \quad (5.4)$$

where $g(x) = x_1 f(x)$ and f is given in terms of the 3-isotropic symmetrization of \tilde{f} by

$$f(x) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \tilde{f}(|x_1|z, x_2, \dots, x_n) \mathcal{H}^2(dz);$$

here \mathbb{S}^2 is the unit sphere in \mathbb{R}^3 and \mathcal{H}^2 is the surface measure.

Proof. Since the map $\tilde{x} \mapsto (\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})$ is smooth from $\tilde{\Omega}_0$ to Ω_0 and $x \mapsto x_1^{-1}u(x)$ is in $C_{\text{loc}}^2(\Omega_0)$, it follows that $v \in C_{\text{loc}}^2(\tilde{\Omega}_0)$. Moreover, the assumption $x_1^{-1}u \in L^\infty(\mathbb{R}^n)$ immediately implies that $\tilde{v} \in L^\infty(\mathbb{R}^{n+2})$, which gives (a).

To prove the claim in (b), we notice that if $f(x) = x_1^{-1}u(x)$ is a Schwartz function, item (b) reduces to Corollary 5.1. The extension to general u follows by mollification and cutting off. This is essentially the same as the argument applied in the proof of Theorem 4.3(c), and so we omit the details.

The proofs of (c) and (d) are analogous to the proofs of Theorem 4.3 (d) and (e) respectively. \square

In preparation for the proof of Theorem 1.2, we now give the definition of an operator satisfying the Harnack inequality.

Definition 5.3. Let \mathcal{A} be the collection of quadruples (Ω, K, c, v) such that $\Omega \subset \mathbb{R}^n$ is open; $K \subset \Omega$ is compact and connected; $c \in L^\infty(\Omega)$; and $v \in C_{\text{loc}}^2(\Omega) \cap L^\infty(\mathbb{R}^n)$ satisfies, in the weak sense,

$$\begin{cases} L_n v + cv = 0, & \text{in } \Omega \\ v \geq 0, & \text{in } \mathbb{R}^n. \end{cases} \quad (5.5)$$

We say that an operator L_n given by (1.2) admits the *Harnack inequality* if, for all $(\Omega, K, c, v) \in \mathcal{A}$, there holds that

$$\inf_K v \leq C \sup_K v$$

for some $C = C(\Omega, K, \|c\|_{L^\infty(\Omega)}) > 0$.

Remark 5.4. When $c = 0$ in (5.5), that is, when u is a harmonic function in Ω for the operator L_n , then Harnack inequality holds under fairly general conditions on the kernel j_n of L_n (see (1.2)). Indeed: Theorem 4.1 in [19] provides a Harnack inequality (in fact: a boundary Harnack inequality; see Remark 1.10(e) in that paper) when K and Ω are balls, under the following assumptions: j_n is positive and nonincreasing (so that the kernel of L_n is isotropic and *unimodal*), and there is a constant $C > 0$ such that $j_n(r+1) \geq C j_n(r)$ for $r \geq 1$ (so that in particular $j_n(r)$ goes to zero as $r \rightarrow \infty$ no faster than some exponential).

By Theorem 1.9 in [19], the constant in the Harnack inequality is *scale-invariant* if j_n is positive and decreasing and it satisfies the following scaling condition:

$$\frac{j_n(r)}{j_n(R)} \leq C \left(\frac{r}{R} \right)^{-n-\alpha}$$

for some constants $C > 0$ and $\alpha \in (0, 2)$.

While it is natural to expect that the results described above also hold for non-zero c , such extension does not seem to be available in literature. More restrictive results are given in [25, 1, 28] and references therein.

We have now developed the machinery to give the proof of Theorem 1.2.

Proof of Theorem 1.2. From Theorem 5.2 (b), we have that

$$L_n u(x) = x_1 L_{n+2} \tilde{v}((x_1, 0, 0), x_2, \dots, x_n)$$

for all $x \in \Omega_0$; or equivalently,

$$\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} L_{n+2} \tilde{v}(\tilde{x}) = L_n u\left(\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}\right)$$

for all $\tilde{x} \in \tilde{\Omega}_0$ (recall that $L_{n+2} u$ is 3-isotropic). It follows that if we denote $x = (\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2})$, then

$$\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2} (L_{n+2} \tilde{v}(\tilde{x}) + c(x) \tilde{v}(\tilde{x})) = L_n u(x) + c(x) u(x) = 0.$$

Setting $\tilde{c}(\tilde{x}) = c(x)$, we have that

$$L_{n+2} \tilde{v} + \tilde{c} \tilde{v} = 0 \quad \text{in } \tilde{\Omega}_0.$$

In fact, by Theorem 5.2 (d), the above equality holds in the weak sense in $\tilde{\Omega}$. By assumption, L_{n+2} admits the Harnack inequality, so from the definition applied to the quadruple $(\tilde{\Omega}, \tilde{K}, \tilde{c}, \tilde{v})$ with $\tilde{K} = \{\tilde{x} \in \mathbb{R}^{n+2} : (\sqrt{\tilde{x}_1^2 + \tilde{x}_2^2 + \tilde{x}_3^2}, \tilde{x}_4, \dots, \tilde{x}_{n+2}) \in K\}$, we have that

$$\sup_{\tilde{K}} \tilde{v} \leq C(\tilde{\Omega}, \tilde{K}, \|\tilde{c}\|_{L^\infty(\tilde{\Omega})}) \inf_{\tilde{K}} \tilde{v}.$$

We conclude that

$$\sup_{x \in K_0} \frac{u(x)}{x_1} \leq C(\tilde{\Omega}, \tilde{K}, \|c\|_{L^\infty(\Omega_+)}) \inf_{x \in K_0} \frac{u(x)}{x_1},$$

as desired. \square

Acknowledgments

SD has been supported by the Australian Future Fellowship FT230100333 “New perspectives on nonlocal equations”. JT is supported by an Australian Government Research Training Program Scholarship. EV has been supported by the Australian Laureate Fellowship FL190100081 “Minimal surfaces, free boundaries and partial differential equations.” MK has been supported by National Science Centre, Poland, 2023/49/B/ST1/04303.

This research was funded in whole or in part by National Science Centre, Poland, 2023/49/B/ST1/04303. For the purpose of Open Access, the authors have applied a CC-BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

Appendix A. Approximation of functions in a class of weighted L^1 spaces

In this section, we give a result that is possibly well-known to experts, but we decided to include since we were unable to find a reference that gave the result explicitly. The result is as follows.

Proposition A.1. *Let η_ε be a standard mollifier on \mathbb{R}^n , that is η_1 is a smooth, nonnegative function with integral 1 and support contained in the unit ball, and $\eta_\varepsilon(x) = \varepsilon^{-n} \eta_1(x/\varepsilon)$. Moreover, let $\alpha > 0$ and μ be the positive measure on \mathbb{R}^n given by*

$$d\mu = \frac{dx}{(1 + |x|)^\alpha}.$$

If $u \in L^1(\mu)$ and $u^\varepsilon = u * \eta_\varepsilon$ then

$$\|u^\varepsilon - u\|_{L^1(\mu)} \rightarrow 0$$

as $\varepsilon \rightarrow 0^+$.

Proof. First, suppose that u is a continuous, compactly supported function. Then

$$\|u^\varepsilon - u\|_{L^1(\mu)} \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y) - u(x)|\eta_\varepsilon(y-x)}{(1+|x|)^\alpha} dy dx.$$

Choosing $R > 1$ so that $u(x) = 0$ when $|x| \geq R$, we have

$$\begin{aligned} \|u^\varepsilon - u\|_{L^1(\mu)} &\leq \int_{\mathbb{R}^n} \int_{B_\varepsilon(x)} \frac{|u(y) - u(x)|\eta_\varepsilon(y-x)}{(1+|x|)^\alpha} dy dx \\ &= \int_{B_{2R}} \int_{B_\varepsilon(x)} \frac{|u(y) - u(x)|\eta_\varepsilon(y-x)}{(1+|x|)^\alpha} dy dx, \end{aligned}$$

where to obtain the second line above we use that if $|x| \geq 2R$ and $|x-y| < \varepsilon$ then $|y| \geq 2R - \varepsilon > R$ for ε small, so $u(x) = u(y) = 0$.

Next, if ω_u denotes the modulus of continuity of u , we have

$$\begin{aligned} \|u^\varepsilon - u\|_{L^1(\mu)} &\leq \omega_u(\varepsilon) \int_{B_{2R}} \int_{B_\varepsilon(x)} \frac{\eta_\varepsilon(y-x)}{(1+|x|)^\alpha} dy dx \\ &\leq \omega_u(\varepsilon) \int_{B_{2R}} \frac{dx}{(1+|x|)^\alpha} \rightarrow 0 \end{aligned}$$

as $\varepsilon \rightarrow 0^+$.

For a general $u \in L^1(\mu)$, using e.g. [29, Theorem 3.14] we may approximate u by a continuous, compactly supported function u_k satisfying

$$\|u - u_k\|_{L^1(\mu)} \leq \frac{1}{k}.$$

Then

$$\|u^\varepsilon - (u_k)^\varepsilon\|_{L^1(\mu)} \leq \int_{\mathbb{R}^n} \int_{B_\varepsilon(x)} \frac{|u(x) - u_k(y)|\eta_\varepsilon(x-y)}{(1+|x|)^\alpha} dy dx.$$

Since $1 + |y| \leq (1 + \varepsilon)(1 + |x|)$ whenever $|x - y| < \varepsilon$, it follows that

$$\begin{aligned} \|u^\varepsilon - (u_k)^\varepsilon\|_{L^1(\mu)} &\leq (1 + \varepsilon)^\alpha \int_{\mathbb{R}^n} \int_{B_\varepsilon(x)} \frac{|u(x) - u_k(y)|\eta_\varepsilon(x-y)}{(1+|y|)^\alpha} dy dx \\ &= (1 + \varepsilon)^\alpha \int_{\mathbb{R}^n} \frac{|u(x) - u_k(y)|}{(1+|y|)^\alpha} dy \\ &\leq \frac{(1 + \varepsilon)^\alpha}{k}. \end{aligned}$$

Since

$$\begin{aligned} \|u^\varepsilon - u\|_{L^1(\mu)} &\leq \|u^\varepsilon - (u_k)^\varepsilon\|_{L^1(\mu)} + \|(u_k)^\varepsilon - u_k\|_{L^1(\mu)} + \|u_k - u\|_{L^1(\mu)} \\ &\leq \frac{(1 + \varepsilon)^\alpha}{k} + \|(u_k)^\varepsilon - u_k\|_{L^1(\mu)} + \frac{1}{k} \end{aligned}$$

and

$$\limsup_{\varepsilon \rightarrow 0^+} \|(u_k)^\varepsilon(x) - u_k(x)\| = 0,$$

we conclude that

$$\limsup_{\varepsilon \rightarrow 0^+} \|u^\varepsilon - u\|_{L^1(\mu)} \leq \frac{2}{k}$$

for all $k \geq 1$, so $\lim_{\varepsilon \rightarrow 0^+} \|u^\varepsilon - u\|_{L^1(\mu)}$ exists and equals zero. \square

References

- [1] S. Athreya, K. Ramachandran, *Harnack inequality for non-local Schrödinger operators*, Potential Anal. 48 (2018), no. 4, 515–551.
- [2] R. Bass, M. Kassmann, *Harnack inequalities for non-local operators of variable order*. Trans. Amer. Math. Soc. 357 (2005), no. 2, 837–850.
- [3] R. F. Bass, D. A. Levin, *Harnack inequalities for jump processes*. Potential Anal. 17 (2002), no. 4, 375–388.
- [4] K. Bogdan, A. Stós, P. Sztonyk, *Harnack inequality for stable processes on d -sets*. Studia Math. 158 (2003), no. 2, 163–198.
- [5] C. Bucur, E. Valdinoci, *Nonlocal diffusion and applications*. Lecture Notes of the Unione Matematica Italiana, 20. Springer, Unione Matematica Italiana, Bologna, 2016.
- [6] L. Caffarelli, L. Silvestre, *An extension problem related to the fractional Laplacian*. Comm. Partial Differential Equations 32 (2007), no. 7-9, 1245–1260.
- [7] M. Cozzi, J. Dávila, M. del Pino, *Long-time asymptotics for evolutionary crystal dislocation models*. Adv. Math. 371 (2020), 107242, 109 pp.
- [8] A. Di Castro, T. Kuusi, G. Palatucci, *Nonlocal Harnack inequalities*. J. Funct. Anal. 267 (2014), no. 6, 1807–1836.
- [9] S. Dipierro, G. Poggesi, J. Thompson, E. Valdinoci, *The role of antisymmetric functions in non-local equations*. Trans. Amer. Math. Soc. 377 (2024), 1671–1692.
- [10] S. Dipierro, O. Savin, E. Valdinoci, *All functions are locally s -harmonic up to a small error*. J. Eur. Math. Soc. (JEMS) 19 (2017), no. 4, 957–966.
- [11] S. Dipierro, O. Savin, E. Valdinoci, *Local approximation of arbitrary functions by solutions of nonlocal equations*. J. Geom. Anal. 29 (2019), no. 2, 1428–1455.
- [12] S. Dipierro, J. Thompson, E. Valdinoci, *On the Harnack inequality for antisymmetric s -harmonic functions*. J. Funct. Anal. 285 (2023), no. 1, Paper No. 109917, 49 pp.
- [13] B. Dyda, *Fractional calculus for power functions and eigenvalues of the fractional Laplacian*. Fract. Calc. Appl. Anal. 15 (2012), no. 4, 536–555.
- [14] B. Dyda, A. Kuznetsov, M. Kwaśnicki, *Fractional Laplace operator and Meijer G -function*. Constr. Approx. 45 (2017), no. 3, 427–448.
- [15] M. M. Fall, S. Jarohs, *Overdetermined problems with fractional Laplacian*. ESAIM Control Optim. Calc. Var. 21 (2015), no. 4, 924–938.
- [16] J.-C. Felipe-Navarro, T. Sanz-Perela, *Semilinear integro-differential equations, I: Odd solutions with respect to the Simons cone*. J. Funct. Anal. 278 (2020), no. 2, 108309, 48 pp.
- [17] J.-C. Felipe-Navarro, *Uniqueness for linear integro-differential equations in the real line and applications*. Calc. Var. Partial Differential Equations 60 (2021), no. 6, Paper No. 220, 25 pp.
- [18] F. Ferrari, B. Franchi, *Harnack inequality for fractional sub-Laplacians in Carnot groups*. Math. Z. 279 (2015), no. 1-2, 435–458.
- [19] T. Grzywny, M. Kwaśnicki, *Potential kernels, probabilities of hitting a ball, harmonic functions and the boundary Harnack inequality for unimodal Lévy processes*. Stoch. Proc. Appl. 128 (2018), no. 1, 1–38.
- [20] A. Harnack, *Die Grundlagen der Theorie des logarithmischen Potentials und der eindeutigen Potentialfunktion in der Ebene*. V. G. Teubner, Leipzig, 1887.
- [21] S. Jarohs, T. Weth, *Symmetry via antisymmetric maximum principles in nonlocal problems of variable order*. Ann. Mat. Pura Appl. (4) 195 (2016), no. 1, 273–291.
- [22] M. Kassmann, *The classical Harnack inequality fails for non-local operators*. Sonderforschungsbereich 611, 2007.
- [23] M. Kassmann, *Harnack inequalities: an introduction*. Bound. Value Probl. 2007, Art. ID 81415, 21 pp.

- [24] M. Kassmann, *A new formulation of Harnack's inequality for nonlocal operators*. C. R. Math. Acad. Sci. Paris 349 (2011), no. 11-12, 637–640.
- [25] Y.-C. Kim, *Nonlocal Harnack inequalities for nonlocal Schrödinger operators with A_1 -Muckenhoupt potentials*, J. Math. Anal. Appl. 507 (2022), no. 1, 125746.
- [26] T. Kulczycki, M. Ryznar, *Gradient estimates of harmonic functions and transition densities for Lévy processes*. Trans. Amer. Math. Soc. 368 (2016), no. 1, 281–318.
- [27] F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Mille, B. V. Saunders, H. S. Cohl, M. A. McClain, eds., *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov/>, Release 1.1.6 of 2022-06-30
- [28] X. Ros-Oton, J. Serra, *The boundary Harnack principle for nonlocal elliptic operators in non-divergence form*, Potential Anal. 51 (2019), no. 3, 315–331.
- [29] W. Rudin, *Real and complex analysis*, McGraw-Hill Book Co., New York, 1987.
- [30] L. Silvestre, *Hölder estimates for solutions of integro-differential equations like the fractional Laplace*. Indiana Univ. Math. J. 55 (2006), no. 3, 1155–1174.
- [31] R. Song, Z. Vondraček, *Harnack inequality for some classes of Markov processes*. Math. Z. 246 (2004), no. 1-2, 177–202.
- [32] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*. Princeton University Press, 1970.
- [33] P. R. Stinga, J. L. Torrea, *Extension problem and Harnack's inequality for some fractional operators*. Comm. Partial Differential Equations 35 (2010), no. 11, 2092–2122.

Serena Dipierro: Department of Mathematics and Statistics, The University of Western Australia, 35 Stirling Highway, Crawley, Perth, WA 6009, Australia.
Email address: serena.dipierro@uwa.edu.au

Mateusz Kwaśnicki: Department of Pure Mathematics, Wrocław University of Science and Technology, Wyb. Wyspiańskiego 27, 50-370 Wrocław, Poland.
Email address: mateusz.kwasnicki@pwr.edu.pl

Jack Thompson: Department of Mathematics and Statistics, The University of Western Australia, 35 Stirling Highway, Crawley, Perth, WA 6009, Australia.
Email address: jack.thompson@research.uwa.edu.au

Enrico Valdinoci: Department of Mathematics and Statistics, The University of Western Australia, 35 Stirling Highway, Crawley, Perth, WA 6009, Australia.
Email address: enrico.valdinoci@uwa.edu.au