

Non-equilibrium steady states with a spatial Markov structure

Frank Redig ^{*} ¹ and Berend van Tol [†] ²

^{1,2}Institute of Applied Mathematics, Delft University of Technology, Delft, The Netherlands

November 19, 2024

Abstract

We investigate the structure of non-equilibrium steady states (NESS) for a class of exactly solvable models in the setting of a chain with left and right reservoirs. Inspired by recent results on the harmonic model [4], [5], we focus on models in which the NESS is a mixture of equilibrium product measures, and where the probability measure which describes the mixture has a spatial Markovian property. We completely characterize the structure of such mixture measures, and show that under natural scaling and translation invariance properties, the only possible mixture measures are coinciding with the Dirichlet process found in [4] in the context of the harmonic model.

1 Introduction

The structure of the stationary state of stochastic interacting particle systems driven out of equilibrium, for instance via boundary driving, is a challenging problem, and very few exactly solvable models are available. A famous example of an exactly solvable model in this context is the boundary driven exclusion process, solvable via the matrix ansatz solution, see [8], [9].

Recently, new explicit representations of non-equilibrium steady states (NESS) were discovered in the context of the boundary driven KMP model [14], [2], [7], and in the context of the boundary driven harmonic model [4], [5]. For a general class of mass transport models, including both the KMP and the harmonic model on general graphs, such representations were found via intertwining in [13]. The integrable structure of the harmonic model was discovered in [11], and explicit formulas for the factorial moments in the non-equilibrium stationary state are given.

An important further development is the discovery that in all these models the NESS can be written as a stochastic mixture of product measures with equilibrium marginals. E.g. for the KMP model the equilibrium measures are products of exponentials with identical parameters. If we then consider the KMP model on a chain with n sites and with left reservoir parameter θ_L and right reservoir parameter θ_R the NESS reads

$$\mu_{\theta_L, \theta_R}^{(n)} = \int (\otimes_{i=1}^n \nu_{\theta_i}) \Lambda_{\theta_L, \theta_R}^n(d\theta_1, \dots, d\theta_n). \quad (1)$$

^{*}Email: F.H.J.Redig@tudelft.nl

[†]Email: B.T.vanTol@tudelft.nl

Here ν_{θ_i} is an exponential distribution with parameter (i.e., expectation) θ_i , and $\Lambda_{\theta_L, \theta_R}^n$, the so-called mixture measure, is a probability measure on $[0, \infty)^n$. This mixture measure describes the distribution of the “hidden parameters” $(\theta_1, \dots, \theta_n)$. We then recover the equilibrium measures of the KMP model when $\theta_L = \theta_R$ because in that case the mixture measure $\Lambda_{\theta_L, \theta_R}^n$ becomes a Dirac measure. Moreover, it turns out that the mixture measure $\Lambda_{\theta_L, \theta_R}^n$ in (1) is itself the stationary measure of a Markovian dynamics on the space of parameters $(\theta_1, \dots, \theta_n)$ which is called the “hidden temperature” model [7] or in more general setting, the “hidden parameter” model [13]. Other models where the NESS can be written as a stochastic mixture of product measures include the boundary driven symmetric exclusion process [10], and the boundary driven inclusion process [13]. In both these models (as well as in the KMP model), the mixture measure cannot be obtained explicitly as it is the case for the harmonic models (on the chain), but it can only be characterized implicitly as the stationary measure of the hidden parameter model.

In the context of the (both discrete and continuous) harmonic model, a representation as in (1) exists, and in that context, the measure $\Lambda_{\theta_L, \theta_R}^n$ is available in closed form and given by the joint distribution of the order statistics of n independent uniforms on the interval $[\theta_L, \theta_R]$ (where we assume $\theta_L < \theta_R$). In [4], a class of generalized harmonic models is introduced, where also the mixture measure $\Lambda_{\theta_L, \theta_R}^n$ is explicit and given by an ordered Dirichlet distribution.

In [13] we provide a probabilistic understanding of the fact that for those harmonic models the mixture measure can be obtained explicitly, by showing that under the measure $\Lambda_{\theta_L, \theta_R}^n$, the conditional distribution of θ_i , given all $\theta_j, j \neq i$ is uniform on $[\theta_{i-1}, \theta_{i+1}]$, which is exactly the Markov structure of the joint distribution of order statistics of independent uniforms. As we showed in [4], this spatial Markov structure of the measure $\Lambda_{\theta_L, \theta_R}^n$ also leads naturally to the additivity principle for the pressure and the corresponding large deviation rate function (cf. [3]). Therefore, we can think of the spatial Markov structure of the mixture measure $\Lambda_{\theta_L, \theta_R}^n$ as a microscopic counterpart of the additivity principle [3].

In this paper, we want to focus on the precise consequences of the Markov structure of the mixture measure $\Lambda_{\theta_L, \theta_R}^n$. As a first result, we prove that in such a setting the measures $\Lambda_{\theta_L, \theta_R}^n$ are completely determined by the system consisting of a single site with left and right reservoirs, i.e., the measures $\Lambda_{\theta_L, \theta_R}^1$. As a second result we characterise those single site systems which are “extendable”, i.e., give rise to a Markov family $\{\Lambda_{\theta_L, \theta_R}^n, n \in \mathbb{N}\}$. Finally, as a third result, under natural assumptions of scaling and translation invariance, we prove that the only possible measures $\Lambda_{\theta_L, \theta_R}^n$ are the ordered Dirichlet distributions which appeared in the generalized harmonic models [4].

The measures $\Lambda_{\theta_L, \theta_R}^n$ are concentrated on the set of ordered n -tuples $\theta_L < \theta_1 < \theta_2 < \dots < \theta_n < \theta_R$. Because of this ordering restriction, the standard theory of Markov specifications from [12] chapter 10 and 11, or the Hammersley-Clifford theorem [6] is not directly applicable as both assume non-null specifications. We can think of the measures $\Lambda_{\theta_L, \theta_R}^n$ as the joint distribution of a Markov bridge, i.e., a joint distribution of the form

$$X_1^{(n)} = \cdot, \dots, X_n^{(n)} = \cdot | X_0^{(n)} = \theta_L, X_{n+1}^{(n)} = \theta_R$$

where $X_1^{(n)} = \cdot, \dots, X_n^{(n)}$ are the first n steps of a Markov process. However, because of the ordering restriction, this Markov bridge depends on n and therefore cannot be thought of as the conditioning of an infinite-volume Gibbs measure with nearest neighbor interaction.

The rest of our paper is organized as follows. In section 2 we define the two-sided Markov property and a notion of symmetry for the measures $\Lambda_{\theta_L, \theta_R}^n$. In section 3 we study their support properties. In section 4 we provide a way to reconstruct the measure $\Lambda_{\theta_L, \theta_R}^n$ corresponding to n sites from the measure corresponding to a single site. Finally, we find two properties concerning the shift and scale invariance of the measures to characterize the Dirichlet distributions.

2 Structure of the n -site densities

In this section we first introduce the necessary notation and give a formal definition of the Markovian structure of the mixture measure $\Lambda_{a,b}^n$. We also define a notion of symmetry which we will assume throughout this text. Next we prove that the Markov property implies a product structure for the n -site density and subsequently investigate consequences of the symmetry property.

Let $a, b \in \mathbb{R}^+$. We consider a family of probability density functions $\{\Lambda_{a,b}^n\}_{n \geq 1}$ corresponding to random vectors taking values in $(I_{a,b})^n$, where $I_{a,b} = (a \wedge b, a \vee b)$. That is, for fixed a, b and n we have random variables $\Theta_{1,n}, \dots, \Theta_{n,n}$ with joint density $\Lambda_{a,b}^n$:

$$\mathbb{P}_n((\Theta_{1,n}, \dots, \Theta_{n,n}) \in A) = \int_A \Lambda_{a,b}^n d\lambda_n, \quad (2)$$

for each Borel set $A \in \mathcal{B}((I_{a,b})^n)$ and $n \geq 1$ and where λ_n denotes the Lebesgue measure on \mathbb{R}^n . We assume that, for all values of $a, b \in \mathbb{R}^+$ and $n \geq 1$, we have a corresponding density $\Lambda_{a,b}^n$. This then defines the map $\Lambda^n : (a, b) \rightarrow \Lambda_{a,b}^n$ and we refer to this map as the n -site density. Moreover, we refer to $\Lambda_{a,b}^n$ as the the n -site density with left parameter a and right parameter b .

Throughout this text we often use the densities of marginals and conditional densities of the random variables $\Theta_{i,n}$, $1 \leq i \leq n$, which we denote as follows,

$$\Lambda_{a,b}^{n,i}(\theta_i) := \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{a,b}^n(\theta_1, \dots, \theta_i, \dots, \theta_n) d\theta_1 \dots d\theta_{i-1} d\theta_{i+1} \dots d\theta_n \quad (3)$$

for the i -th marginal and, for $\theta_i \in \text{supp}(\Lambda_{a,b}^{n,i})$,

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n | \Theta_i = \theta_i) := \frac{\Lambda_{a,b}^n(\theta_1, \dots, \theta_i, \dots, \theta_n)}{\Lambda_{a,b}^{n,i}(\theta_i)}. \quad (4)$$

We now define what it means for the family Λ^n to be two-sided Markov. For notational simplicity we omit the right index n in $\Theta_{1,n}, \dots, \Theta_{n,n}$ whenever n is fixed, i.e. we write $\Theta_1, \dots, \Theta_n$.

Definition 2.1. *The family $\{\Lambda^n\}_{n \geq 1}$ is called*

1. **Two-sided Markov** if the corresponding densities $\Lambda_{a,b}^n$, $n \geq 1$ and $a, b \in \mathbb{R}^+$, satisfy the following two properties

(i) *Two-sided Markov property:*

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n | \Theta_j = \theta_j) = \Lambda_{a,\theta_j}^{j-1}(\theta_1, \dots, \theta_{j-1}) \cdot \Lambda_{\theta_j,b}^{n-j}(\theta_{j+1}, \dots, \theta_n) \quad (5)$$

for all $\theta_1, \dots, \theta_n \in I_{a,b}$. Here we define $\Lambda_{a,b}^0 := 1$.

(ii) *Support restriction:* $\Lambda_{a,b}^1$ has support $I_{a,b}$. Here we define the support of a function $f : X \rightarrow \mathbb{R}$ as $\text{supp}(f) := \{x \in X : f(x) \neq 0\}$.

2. **Symmetric** if, for all $n \geq 1$, $a, b \in \mathbb{R}^+$ and $\theta_1, \dots, \theta_n \in I_{a,b}$

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \Lambda_{b,a}^n(\theta_n, \dots, \theta_1). \quad (6)$$

Remark 2.2. In this paper we are interested in densities $\Lambda_{a,b}^n$ which concentrate on the set of ordered n -tuples $a < \theta_1 < \dots < \theta_n < b$. This is guaranteed via the requirement on the support of Λ^1 , as we will prove in proposition 3.2.

Example 2.3. Consider the joint probability density of the order statistics of n i.i.d. uniform random variables on (a, b) with $a \leq b$. More precisely, for $n \geq 1$ we consider $X_1, X_2, \dots, X_n \sim U(a, b)$ and denote the corresponding order statistics by $\Theta_1 = X_{1:n}, \Theta_2 = X_{2:n}, \dots, \Theta_n = X_{n:n}$. The joint distribution of $\Theta_1, \dots, \Theta_n$ is given by

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = n! \cdot |b - a|^{-n} \mathbf{1}(a < \theta_1 < \theta_2 < \dots < \theta_n < b). \quad (7)$$

The marginals are given by

$$\Lambda_{a,b}^{n,i}(\theta) = \frac{n!}{(i-1)!(n-i)!} |b - a|^{-n} |b - \theta|^{n-i} |\theta - a|^{i-1} \mathbf{1}(\theta \in I_{a,b}). \quad (8)$$

The two-sided Markov property now follows from

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n | \Theta_i = \theta_i) = \frac{\Lambda_{a,b}^n(\theta_1, \dots, \theta_n)}{\Lambda_{a,b}^{n,i}(\theta_i)} = \frac{(i-1)!(n-i)!}{|b - \theta_i|^{n-i} |\theta_i - a|^{i-1}} \mathbf{1}(a < \theta_1 < \theta_2 < \dots < \theta_n < b) \quad (9)$$

and

$$\Lambda_{a,\theta_i}^{i-1}(\theta_1, \dots, \theta_{i-1}) \cdot \Lambda_{\theta_i,b}^{n-i}(\theta_{i+1}, \dots, \theta_n) = \frac{(i-1)!(n-i)!}{|b - \theta_i|^{n-i} |\theta_i - a|^{i-1}} \mathbf{1}(a < \theta_1 < \theta_2 < \dots < \theta_n < b). \quad (10)$$

We generalize the definition of the order statistics for the case $b \leq a$ in such a way that the family $\{\Lambda^n\}_{n \geq 1}$ becomes symmetric. I.e. we define

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \begin{cases} n! \cdot |b - a|^{-n} \mathbf{1}(a < \theta_1 < \theta_2 < \dots < \theta_n < b) & \text{if } a < b \\ n! \cdot |b - a|^{-n} \mathbf{1}(a > \theta_1 > \theta_2 > \dots > \theta_n > b) & \text{if } a > b \end{cases} \quad (11)$$

One can show the two-sided Markov property for the case $a > b$ in the same way as is done above for $a < b$.

2.1 Product form for two-sided Markov families

In this paper we aim to fully characterize the symmetric families which satisfy the two-sided Markov property. A first step in this direction is given by the proposition 2.4 below. It states the equivalence between the two-sided Markov property and the density being a product of marginals in two distinct ways. These product forms can directly be derived by repeatedly applying the two-sided Markov property. The converse can be derived by combining the two product representations.

Proposition 2.4. For notational convenience we write $\theta_0 := a$ and $\theta_{n+1} := b$. The following statements are equivalent:

1. $\{\Lambda^n\}_{n \geq 1}$ is two-sided Markov.

2. For all $n \geq 1$, $a, b \in \mathbb{R}^+$ and $\theta_1, \dots, \theta_n \in I_{a,b}$ we have

$$\begin{cases} \Lambda_{a,b}^n(\theta_1, \dots, \theta_n | \Theta_n = \theta_n) = \Lambda_{a,\theta_n}^{n-1}(\theta_1, \dots, \theta_{n-1}) \\ \Lambda_{a,b}^n(\theta_1, \dots, \theta_n | \Theta_1 = \theta_1) = \Lambda_{\theta_1,b}^{n-1}(\theta_2, \dots, \theta_n). \end{cases} \quad (12)$$

3. For all $n \geq 1$, $a, b \in \mathbb{R}^+$ and $\theta_1, \dots, \theta_n \in I_{a,b}$,

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \prod_{i=0}^{n-1} \Lambda_{\theta_i,b}^{n-i,1}(\theta_{i+1}) = \prod_{i=0}^{n-1} \Lambda_{a,\theta_{n-i+1}}^{n-i,n-i}(\theta_{n-i}). \quad (13)$$

Proof. 1 implies 2: This follows immediately from the definition of the two-sided Markov property.

2 implies 3: We use induction. (13) obviously holds for $n = 1$. Assume (13) holds for $n = k$, then we have, using (12),

$$\Lambda_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1}) = \Lambda_{a,b}^{k+1,k+1}(\theta_{k+1}) \cdot \Lambda_{a,\theta_{k+1}}^k(\theta_1, \dots, \theta_k) = \Lambda_{a,b}^{k+1,k+1}(\theta_{k+1}) \cdot \prod_{i=0}^{k-1} \Lambda_{a,\theta_{k-i+1}}^{k-i,k-i}(\theta_{k-i}) \quad (14)$$

and

$$\Lambda_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1}) = \Lambda_{a,b}^{k+1,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^k(\theta_2, \dots, \theta_{k+1}) = \Lambda_{a,b}^{k+1,1}(\theta_1) \cdot \prod_{i=1}^k \Lambda_{\theta_i,b}^{k-i,1}(\theta_{i+1}). \quad (15)$$

3 implies 1: We have the following two expressions for $\Lambda_{a,b}^n(\theta_1, \dots, \theta_n)$,

$$\begin{aligned} & \Lambda_{a,b}^n(\theta_1, \dots, \theta_n) \\ &= [\Lambda_{a,b}^{n,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^{n-1,1}(\theta_2) \dots \Lambda_{\theta_{j-1},b}^{n-j+1,1}(\theta_j)] \cdot [\Lambda_{\theta_j,b}^{j,j}(\theta_{j+1}) \cdot \Lambda_{\theta_{j+1},b}^{n-j-1,1}(\theta_{j+2}) \dots \Lambda_{\theta_{n-1},b}^{1,1}(\theta_n)] \end{aligned} \quad (16)$$

and

$$\begin{aligned} & \Lambda_{a,b}^n(\theta_1, \dots, \theta_n) \\ &= [\Lambda_{a,\theta_2}^{1,1}(\theta_1) \cdot \Lambda_{a,\theta_3}^{2,2}(\theta_2) \dots \Lambda_{a,\theta_j}^{j-1,j-1}(\theta_j)] \cdot [\Lambda_{a,\theta_j}^{j,j}(\theta_{j+1}) \cdot \Lambda_{a,\theta_{j+1}}^{j+1,j+1}(\theta_{j+2}) \dots \Lambda_{a,b}^{n,n}(\theta_n)]. \end{aligned} \quad (17)$$

Notice both expressions factor into a part which does not depend on $\theta_1, \dots, \theta_{j-1}$ and a part which does not depend on $\theta_{j+1}, \dots, \theta_n$. As a consequence

$$\frac{\Lambda_{a,b}^{n,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^{n-1,1}(\theta_2) \dots \Lambda_{\theta_{j-1},b}^{n-j+1,1}(\theta_j)}{\Lambda_{a,\theta_2}^{1,1}(\theta_1) \cdot \Lambda_{a,\theta_3}^{2,2}(\theta_2) \dots \Lambda_{a,\theta_j}^{j-1,j-1}(\theta_j)} = C(a, \theta_j, b) = \frac{\Lambda_{a,\theta_j}^{j,j}(\theta_{j+1}) \cdot \Lambda_{a,\theta_{j+1}}^{j+1,j+1}(\theta_{j+2}) \dots \Lambda_{a,b}^{n,n}(\theta_n)}{\Lambda_{a,\theta_j}^{j,j}(\theta_{j+1}) \cdot \Lambda_{a,\theta_{j+1}}^{j+1,j+1}(\theta_{j+2}) \dots \Lambda_{a,b}^{n,n}(\theta_n)} \quad (18)$$

for some function C . We tacitly used that the denominators above are non-zero. However if one of the factors in the denominator of the equation above is zero, then the two-sided Markov property holds automatically because both left and right hand side of the defining property (5) are equal to zero. One can now relate the factors of the first and the second expression via C . We use this to show the two-sided Markov property.

$$\begin{aligned}
\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) &= [\Lambda_{a,b}^{n,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^{n-1,1}(\theta_2) \dots \Lambda_{\theta_{j-1},b}^{n-j+1,1}(\theta_j)] \cdot [\Lambda_{\theta_j,b}^{n-j,1}(\theta_{j+1}) \cdot \Lambda_{\theta_{j+1},b}^{n-j-1,1}(\theta_{j+2}) \dots \Lambda_{\theta_{n-1},b}^{1,1}(\theta_n)] \quad (19) \\
&= C(a, \theta_j, b) \cdot [\Lambda_{a,\theta_2}^{1,1}(\theta_1) \cdot \Lambda_{a,\theta_3}^{2,2}(\theta_2) \dots \Lambda_{a,\theta_j}^{j-1,j-1}(\theta_j)] \cdot [\Lambda_{\theta_j,b}^{n-j,1}(\theta_{j+1}) \cdot \Lambda_{\theta_{j+1},b}^{n-j-1,1}(\theta_{j+2}) \dots \Lambda_{\theta_{n-1},b}^{1,1}(\theta_n)] \\
&= C(a, \theta_j, b) \cdot \Lambda_{a,\theta_j}^{j-1}(\theta_1, \dots, \theta_{j-1}) \cdot \Lambda_{\theta_j,b}^{n-j}(\theta_{j+1}, \dots, \theta_n).
\end{aligned}$$

Integrating both sides of this equation with respect to $\theta_1, \dots, \theta_{j-1}, \theta_{j+1}, \dots, \theta_n$ yields $C(a, \theta_j, b) = \Lambda_{a,b}^{n,j}(\theta_j)$, which then gives us the two-sided Markov property in (19). \square

Remark 2.5. Proposition (2.4) will play an important role in later applications since it shows the whole family is determined by its left and right-side marginals, i.e. $\{\Lambda^{n,1}\}_{n \geq 1}$ and $\{\Lambda^{n,n}\}_{n \geq 1}$. More precisely, suppose one has a family up to some $m \geq 0$, $\{\Lambda^n\}_{m \geq n \geq 1}$. Then we can construct a valid density Λ^{m+1} if we can find $\Lambda^{m+1,1}$ and $\Lambda^{m+1,m+1}$ such that

$$\Lambda_{a,b}^{m+1}(\theta_1, \dots, \theta_{m+1}) := \Lambda_{a,b}^{m+1,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^m(\theta_2, \dots, \theta_{m+1}) = \Lambda_{a,\theta_{m+1}}^m(\theta_1, \dots, \theta_m) \cdot \Lambda_{a,b}^{m+1,m+1}(\theta_{m+1}). \quad (20)$$

for all $a, b \in \mathbb{R}^+$ and $\theta_1, \dots, \theta_{m+1} \in I_{a,b}$. Indeed, one can substitute the product forms in (13) for $\Lambda_{a,b}^m$. Then $\Lambda_{a,b}^{m+1}$ again has the product form

$$\Lambda_{a,b}^{m+1}(\theta_1, \dots, \theta_{m+1}) = \prod_{i=0}^m \Lambda_{\theta_i,b}^{m+1-i,1}(\theta_{i+1}) = \prod_{i=0}^m \Lambda_{a,\theta_{m+2-i}}^{m+1-i,m+1-i}(\theta_{m+1-i}), \quad (21)$$

and hence satisfies the two-side Markov property.

2.2 Implications of the symmetry property

In this subsection we examine some of the consequences of the symmetry property defined in (6). First, we will relate the j -th and the $n - j + 1$ -th marginal. This is done in proposition 2.6. Next we show that the symmetry of a family $\{\Lambda^n\}_{n \geq 1}$ is completely determined by the symmetry of Λ^1 , provided that the family is two-sided Markov. This result is the content of 2.7.

Proposition 2.6. Let $\{\Lambda^n\}_{n \geq 1}$ be symmetric. Assume $a, b \in \mathbb{R}^+$, $\theta \in \mathbb{R}$ and $n \geq 1$, then

$$\Lambda_{a,b}^{n,j}(\theta) = \Lambda_{b,a}^{n,n-j+1}(\theta) \quad (22)$$

for all $j \in \{1, \dots, n\}$.

Proof.

$$\begin{aligned}
\Lambda_{a,b}^{n,j}(\theta) &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{a,b}^n(\theta_1, \dots, \theta, \dots, \theta_n) d\theta_1 \dots d\theta_{j-1} d\theta_{j+1} \dots d\theta_n \quad (23) \\
&= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{b,a}^n(\theta_n, \dots, \theta, \dots, \theta_1) d\theta_1 \dots d\theta_{j-1} d\theta_{j+1} \dots d\theta_n = \Lambda_{b,a}^{n,n-j+1}(\theta)
\end{aligned}$$

\square

Proposition 2.7. Let $\{\Lambda_{a,b}^n\}_{n \geq 1}$ be two-sided Markov with $\Lambda_{a,b}^1 = \Lambda_{b,a}^1$, then the whole family $\{\Lambda_{a,b}^n\}_{n \geq 1}$ is symmetric.

Proof. We use induction. The symmetry of $\Lambda_{a,b}^n$ holds for $n = 1$ by assumption. Suppose the symmetry holds for $n = k$ and that $(\theta_1, \dots, \theta_{k+1}) \in \text{supp } \Lambda_{a,b}^{k+1}$. Notice that, due to the two-sided Markov property,

$$1 = \frac{\Lambda_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1})}{\Lambda_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1})} = \frac{\Lambda_{a,b}^{k+1,1}(\theta_1) \cdot \Lambda_{\theta_1, \theta_k}^{k-1}(\theta_2, \dots, \theta_k) \cdot \Lambda_{\theta_1, b}^{k,k}(\theta_{k+1})}{\Lambda_{a,b}^{k+1,k+1}(\theta_{k+1}) \cdot \Lambda_{\theta_1, \theta_k}^{k-1}(\theta_2, \dots, \theta_k) \cdot \Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1)}. \quad (24)$$

We obtain

$$\frac{\Lambda_{a,b}^{k+1,1}(\theta_1)}{\Lambda_{a,b}^{k+1,k+1}(\theta_{k+1})} = \frac{\Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1)}{\Lambda_{\theta_1, b}^{k,k}(\theta_{k+1})}. \quad (25)$$

We will use this equation to show the symmetry for $n = k + 1$. Notice that we have symmetry if

$$\frac{\Lambda_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1})}{\Lambda_{b,a}^{k+1}(\theta_{k+1}, \dots, \theta_1)} = \frac{\Lambda_{a,b}^{k+1,1}(\theta_1) \cdot \Lambda_{\theta_1, b}^k(\theta_2, \dots, \theta_{k+1})}{\Lambda_{b,a}^{k+1,k+1}(\theta_1) \cdot \Lambda_{b, \theta_1}^k(\theta_{k+1}, \dots, \theta_2)} = 1. \quad (26)$$

By the induction hypothesis this is equivalent to

$$\frac{\Lambda_{a,b}^{k+1,1}(\theta_1)}{\Lambda_{b,a}^{k+1,k+1}(\theta_1)} = 1 \quad (27)$$

We will prove (27) to show symmetry for $n = k + 1$. We start from (25) and observe that

$$\frac{\Lambda_{a,b}^{k+1,1}(\theta_1)}{\Lambda_{a,b}^{k+1,k+1}(\theta_{k+1})} = \frac{\Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1)}{\Lambda_{\theta_1, b}^{k,k}(\theta_{k+1})} = \left(\frac{\Lambda_{\theta_1, b}^{k,k}(\theta_{k+1})}{\Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1)} \right)^{-1} = \left(\frac{\Lambda_{b, \theta_1}^{k,1}(\theta_{k+1})}{\Lambda_{\theta_{k+1}, a}^{k,k}(\theta_1)} \right)^{-1} = \frac{\Lambda_{b,a}^{k+1,k+1}(\theta_1)}{\Lambda_{b,a}^{k+1,1}(\theta_{k+1})}. \quad (28)$$

For the third equality we use the induction hypothesis and for the last equality we use (25). One can now separate the equation into a θ_1 -dependent and a θ_{k+1} -dependent side. That is

$$\frac{\Lambda_{a,b}^{k+1,1}(\theta_1)}{\Lambda_{b,a}^{k+1,k+1}(\theta_1)} = \frac{\Lambda_{a,b}^{k+1,k+1}(\theta_{k+1})}{\Lambda_{b,a}^{k+1,1}(\theta_{k+1})} = c \quad (29)$$

for some c independent from both θ_1 and θ_{k+1} . Integrating $\Lambda_{a,b}^{k+1,1}(\theta_1) = c \cdot \Lambda_{b,a}^{k+1,k+1}(\theta_1)$ over θ_1 yields $c = 1$. This yields (27) and hence concludes the proof. \square

3 Support of the densities

In this section we study the support of the densities $\{\Lambda_{a,b}^n\}_{n \geq 1}$. More precisely, we show that the support of the first marginal $\Lambda_{a,b}^{n,1}$ is given by $I_{a,b}$ for all $n \in \mathbb{N}$. The product form from the previous section then implies that $\Lambda_{a,b}^n$ is supported on the set of ordered n -tuples in the interval $I_{a,b}$.

For two-sided Markov families $\{\Lambda^n\}_{n \geq 1}$ one can find transition operators transforming on the marginals. More precise, for suitable i , one can find a raising operator $P_{n,i+1}$ and a lowering operator $Q_{n,i-1}$ transforming $\Lambda_{a,b}^{n,i}$ to $\Lambda_{a,b}^{n,i+1}$ and $\Lambda_{a,b}^{n,i-1}$ respectively. The proposition below states these transition operators explicitly.

Proposition 3.1. Let $\{\Lambda^n\}_{n \geq 1}$ be two-sided Markov. For $a, b \in \mathbb{R}^+$ and $n \geq 1$, $i \in \{1, \dots, n\}$ we have

$$\begin{cases} \Lambda_{a,b}^{n,i+1} = P_{n,i+1} \Lambda_{a,b}^{n,i} & \text{for } i \in \{1, \dots, n-1\} \\ \Lambda_{a,b}^{n,i-1} = Q_{n,i-1} \Lambda_{a,b}^{n,i} & \text{for } i \in \{2, \dots, n\} \end{cases} \quad (30)$$

Here $P_{n,i+1} : L^1(-\infty, \infty) \rightarrow L^1(-\infty, \infty)$ and $Q_{n,i-1} : L^1(-\infty, \infty) \rightarrow L^1(-\infty, \infty)$ are given by

$$\begin{cases} P_{n,i+1} f(x) = \int_{-\infty}^{\infty} f(y) \cdot \Lambda_{y,b}^{n-i,1}(x) dy \\ Q_{n,i-1} f(x) = \int_{-\infty}^{\infty} f(y) \cdot \Lambda_{a,y}^{i-1,i-1}(x) dy \end{cases} \quad (31)$$

Proof. It is straightforward to show that transition operators $P_{n,i+1}$ and $Q_{n,i-1}$ map $L^1(-\infty, \infty)$ functions to $L^1(-\infty, \infty)$ functions. Indeed, let $f \in L^1(-\infty, \infty)$, then

$$\int_{-\infty}^{\infty} |P_{n,i+1} f(x)| dx \leq \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |f(y)| \cdot |\Lambda_{y,b}^{n-i,1}(x)| dy dx \leq \|f\|_1 < \infty. \quad (32)$$

A similar argument can be used to show $Q_{n,i-1}$ map $L^1(-\infty, \infty)$ functions to $L^1(-\infty, \infty)$.

We now show that $\Lambda_{a,b}^{n,i+1} = P_{n,i+1} \Lambda_{a,b}^{n,i}$. One can use an analogous proof to show $\Lambda_{a,b}^{n,i-1} = Q_{n,i-1} \Lambda_{a,b}^{n,i}$.

We use the notation $\Lambda_{a,b}^{n,(i,i+1)}$ to denote the joint distribution of Θ_i and Θ_{i+1} , i.e.

$$\Lambda_{a,b}^{n,(i,i+1)}(\theta_i, \theta_{i+1}) = \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{a,b}^n(\theta_1, \dots, \theta_n) d\theta_1, \dots, d\theta_{i-1} d\theta_{i+2} \dots d\theta_n. \quad (33)$$

We write $\Lambda_{a,b}^{n,(i,i+1)}(y, \theta_{i+1} | \Theta_i = y)$ for the joint distribution under the condition $\Theta_i = y$. Suppose we have

$$\Lambda_{a,b}^{n,(i,i+1)}(y, \theta_{i+1} | \Theta_i = y) = \Lambda_{y,b}^{n-i,1}(\theta_{i+1}), \quad (34)$$

then the result follows directly:

$$\begin{aligned} \Lambda_{a,b}^{n,i+1}(\theta_{i+1}) &= \int_{-\infty}^{\infty} \Lambda_{a,b}^{n,i}(y) \cdot \Lambda_{a,b}^{n,(i,i+1)}(y, \theta_{i+1} | \Theta_i = y) dy \\ &= \int_{-\infty}^{\infty} \Lambda_{a,b}^{n,i}(y) \cdot \Lambda_{y,b}^{n-i,1}(\theta_{i+1}) dy \\ &= P_{n,i+1} \Lambda_{a,b}^{n,i}(\theta_{i+1}). \end{aligned} \quad (35)$$

To prove (34) we compute $\Lambda_{a,b}^{n,(i,i+1)}(y, \theta_{i+1} | \Theta_i = y)$,

$$\begin{aligned} &\Lambda_{a,b}^{n,(i,i+1)}(y, \theta_{i+1} | \Theta_i = y) \\ &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{a,b}^n(\theta_1, \dots, \theta_{i-1}, y, \theta_{i+1}, \dots, \theta_n | \Theta_i = y) d\theta_1 \dots d\theta_{i-1} d\theta_{i+2} \dots d\theta_n \\ &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{a,y}^{i-1}(\theta_1, \dots, \theta_{i-1}) \cdot \Lambda_{y,b}^{n-i}(\theta_{i+1}, \dots, \theta_n) d\theta_1 \dots d\theta_{i-1} d\theta_{i+2} \dots d\theta_n \\ &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \Lambda_{y,b}^{n-i}(\theta_{i+1}, \dots, \theta_n) d\theta_{i+2} \dots d\theta_n \\ &= \Lambda_{a,b}^{n-i,1}(\theta_{i+1}). \end{aligned} \quad (36)$$

□

In the following proposition we make use of the transition operators introduced in 3.1 in order to prove that the support of $\Lambda_{a,b}^1$ determines the support of the marginals $\Lambda_{a,b}^{n,1}$ and hence via the product structure the support $\Lambda_{a,b}^n$.

Proposition 3.2. *Let $\{\Lambda^n\}_{n \geq 1}$ be two-sided Markov and $a, b \in \mathbb{R}^+$. Then*

$$\text{supp}(\Lambda_{a,b}^1) = I_{a,b} \quad \text{implies} \quad \text{supp}(\Lambda_{a,b}^{n,1}) = I_{a,b} \quad (37)$$

for all $n \geq 1$.

Proof. We use induction. For $n = 1$ the statement holds by assumption. Assume $\forall m \leq k$: $\text{supp} \Lambda_{a,b}^{m,1} = I_{a,b}$. First we show that $\text{supp} \Lambda_{a,b}^{k+1,1} \subset I_{a,b}$. This part of the argument doesn't rely on the transition operators.

Step 1: the support of $\Lambda_{a,b}^{k+1,1}$ is a subset of $I_{a,b}$

Suppose $\theta_1 \in \mathbb{R}$ is in the support of $\Lambda_{a,b}^{k+1,1}$, i.e. $\Lambda_{a,b}^{k+1,1}(\theta_1) \neq 0$. Then, there exist $\theta_2, \dots, \theta_{k+1}$ such that

$$\begin{aligned} \Lambda_{a,b}^{k+1}(\theta_1, \theta_2, \dots, \theta_{k+1}) &= \Lambda_{a,b}^{k+1,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^{k,1}(\theta_2) \dots \Lambda_{\theta_k,b}^{1,1}(\theta_{k+1}) \\ &= \Lambda_{a,b}^{k+1,k+1}(\theta_{k+1}) \cdot \Lambda_{a,\theta_{k+1}}^{k,k}(\theta_k) \dots \Lambda_{a,\theta_2}^{1,1}(\theta_1) \neq 0. \end{aligned} \quad (38)$$

Suppose $\theta_1 < b$, then $\theta_1 < \theta_2 < \dots < \theta_{k+1} \leq b$ by the induction hypothesis and the first line of (38). The second line of (38) tells us that either $a < \theta_1 < \theta_2 < \dots < \theta_{k+1}$ or $a > \theta_1 > \theta_2 > \dots > \theta_{k+1}$. Combining the information from line one and line two gives $a < \theta_1 < b$. Similarly we can assume $\theta_1 < b$ and derive $b < \theta_1 < a$. We conclude $\theta_1 \in I_{a,b}$.

Step 2: $I_{a,b}$ is the support of $\Lambda_{a,b}^{k+1,1}$

We notice that $\Lambda_{a,b}^{k+1,1}$ is a fixed point for $Q_{k+1,1}P_{k+1,2}$ by virtue of proposition 3.1,

$$Q_{k+1,1} \left(P_{k+1,2} \Lambda_{a,b}^{k+1,1} \right) = Q_{k+1,1} \Lambda_{a,b}^{k+1,2} = \Lambda_{a,b}^{k+1,1}. \quad (39)$$

We simplify notation by writing f for $\Lambda_{a,b}^{k+1,1}$, g for $P_{k+1,2} \Lambda_{a,b}^{k+1,1}$, i.e. $f := \Lambda_{a,b}^{k+1,1}$ and $g := P_{k+1,2} \Lambda_{a,b}^{k+1,1}$.

Assume, without loss of generality, that $a \leq b$. Then,

$$g(x) = \int_{-\infty}^{\infty} f(y) \cdot \Lambda_{y,b}^{k,1}(x) dy = \int_a^b f(y) \cdot \Lambda_{y,b}^{k,1}(x) dy. \quad (40)$$

Note that under the induction hypothesis $\text{supp} \Lambda_{y,b}^{k,1} = [y, b]$ for $y \in [a, b]$ and that $\text{supp} f \subset I_{a,b} = [a, b]$. We see that

$$\text{supp} g = \{x \in \mathbb{R} : \text{ess inf}(\text{supp} f) < x < b\}, \quad (41)$$

where $\text{ess inf}(\text{supp} f) := \inf \{z \in \mathbb{R} : \lambda((-\infty, z] \cap \text{supp} f) \neq 0\}$ with λ the Lebesgue measure. Analogously, we have

$$f(x) = Q_{k+1,1}g(x) = \int_{-\infty}^{\infty} g(y) \cdot \Lambda_{a,y}^1(x) dy = \int_a^b g(y) \cdot \Lambda_{a,y}^1(x) dy \quad (42)$$

and

$$\text{supp}(f) = \{x \in \mathbb{R} : a < x < \text{ess sup}(\text{supp } g)\} \quad (43)$$

with $\text{ess sup}(\text{supp } g) := \sup \{z \in \mathbb{R} : \lambda([z, \infty) \cap \text{supp } g) \neq 0\}$. Notice that $\text{supp } g$ can not be empty since f is a fixed point of $Q_{k+1,1}P_{k+1,2}$. Indeed, if $\text{supp } g$ would be empty then f would be zero, which is not possible since $f = \Lambda_{a,b}^{k+1,1}$ is a density function. This implies $\text{ess sup}(\text{supp } g) = b$ since the support of g is the interval in (41). We conclude $\text{supp } f = \text{supp } \Lambda_{a,b}^{k+1,1} = (a, b) = I_{a,b}$. \square

Remark 3.3. *The support of the first marginals determines directly the support of the joint density. Indeed, the product expressions for $\Lambda_{a,b}^n$ in proposition 2.4 directly yields that*

$$\text{supp } \Lambda_{a,b}^n = \begin{cases} \{\theta_1, \dots, \theta_n \in \mathbb{R} : a < \theta_1 < \dots < \theta_n < b\} & \text{for } a < b \\ \{\theta_1, \dots, \theta_n \in \mathbb{R} : a > \theta_1 > \dots > \theta_n > b\} & \text{for } b < a \end{cases}. \quad (44)$$

4 Uniqueness and Recursive construction

In this subsection we attempt to construct families $\{\Lambda^n\}_{n \geq 1}$ satisfying the two-sided Markov property. In particular, we are interested in identifying which “one-site systems” Λ^1 extend to a full two-sided Markov family $\{\Lambda^n\}_{n \geq 1}$. We call those *extendable*. So far, in section 3, we have seen that both symmetry and the support of the first marginal $\Lambda^{n,1}$ are determined by Λ^1 . One of the results in this section gives uniqueness for extensions of Λ^1 which has support on $I_{a,b}$ for $a, b \in \mathbb{R}$. Moreover, we will use proposition 2.4 to identify the form of the Λ^1 which are extendable.

4.1 Uniqueness

In the previous subsection we found transition operators with the marginals as fixed points. Since these operators are defined in terms of marginals of smaller order, one could be tempted to show uniqueness recursively via a fixed point theorem (e.g the Krein-Rutman theorem). However, in our setting it appears to be non-trivial to satisfy the condition for such theorems. Therefore we give a more direct approach below, relying strongly on our knowledge about the support of the marginals.

Theorem 4.1. *Let $\{\Lambda^n\}_{n \geq 1}$ be a two-sided Markov family. Then Λ^1 uniquely determines $\{\Lambda^n\}_{n \geq 1}$.*

Proof. Suppose we have two families $\{\Lambda^n\}_{n \geq 1}$ and $\{\tilde{\Lambda}^n\}_{n \geq 1}$ which are both two-sided Markov. Moreover assume

$$\Lambda_{a,b}^1 = \tilde{\Lambda}_{a,b}^1. \quad (45)$$

We show

$$\{\Lambda^n\}_{n \geq 1} = \{\tilde{\Lambda}^n\}_{n \geq 1}. \quad (46)$$

using induction. For $n = 1$, the density functions are equal by assumption. Let $a, b \in \mathbb{R}^+$, $a < b$, be arbitrary and assume that $\Lambda^k = \tilde{\Lambda}^k$. For any $a < \theta_1 < \dots < \theta_{k+1} < b$ we have

$$\begin{aligned}
0 \neq \Lambda_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1}) &= \Lambda_{a,b}^{k+1,1}(\theta_1) \cdot \Lambda_{\theta_1, \theta_k}^{k-1}(\theta_2, \dots, \theta_k) \cdot \Lambda_{\theta_1, b}^{k,k}(\theta_{k+1}) \\
&= \Lambda_{a,b}^{k+1,k+1}(\theta_{k+1}) \cdot \Lambda_{\theta_1, \theta_k}^{k-1}(\theta_2, \dots, \theta_k) \cdot \Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1)
\end{aligned} \tag{47}$$

and

$$\begin{aligned}
0 \neq \tilde{\Lambda}_{a,b}^{k+1}(\theta_1, \dots, \theta_{k+1}) &= \tilde{\Lambda}_{a,b}^{k+1,1}(\theta_1) \cdot \Lambda_{\theta_1, \theta_k}^{k-1}(\theta_2, \dots, \theta_k) \cdot \Lambda_{\theta_1, b}^{k,k}(\theta_{k+1}) \\
&= \tilde{\Lambda}_{a,b}^{k+1,k+1}(\theta_{k+1}) \cdot \Lambda_{\theta_1, \theta_k}^{k-1}(\theta_2, \dots, \theta_k) \cdot \Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1).
\end{aligned} \tag{48}$$

We find

$$\frac{\Lambda_{a,b}^{k+1,1}(\theta_1)}{\Lambda_{a,b}^{k+1,k+1}(\theta_{k+1})} = \frac{\Lambda_{a, \theta_{k+1}}^{k,1}(\theta_1)}{\Lambda_{\theta_1, b}^{k,k}(\theta_{k+1})} = \frac{\tilde{\Lambda}_{a,b}^{k+1,1}(\theta_1)}{\tilde{\Lambda}_{a,b}^{k+1,k+1}(\theta_{k+1})}. \tag{49}$$

Let $\epsilon > 0$ be arbitrary and pick $\theta_{k+1} = b - \epsilon$. For all $\theta_1 \in (a, b - \epsilon]$

$$\frac{\Lambda_{a,b}^{k+1,1}(\theta_1)}{\Lambda_{a,b}^{k+1,k+1}(b - \epsilon)} = \frac{\tilde{\Lambda}_{a,b}^{k+1,1}(\theta_1)}{\tilde{\Lambda}_{a,b}^{k+1,k+1}(b - \epsilon)}. \tag{50}$$

Integration of both sides over θ_1 yields that $\Lambda_{a,b}^{k+1,k+1}(b - \epsilon) = \tilde{\Lambda}_{a,b}^{k+1,k+1}(b - \epsilon)$ and hence we have $\Lambda_{a,b}^{k+1,1}(x) = \tilde{\Lambda}_{a,b}^{k+1,1}(x)$ for $x \in (a, b - \epsilon)$. Since this holds for arbitrary ϵ , we can conclude that $\Lambda_{a,b}^{k+1,1} = \tilde{\Lambda}_{a,b}^{k+1,1}$. Moreover, we have $\Lambda_{a,b}^{k+1} = \tilde{\Lambda}_{a,b}^{k+1}$. For $b \leq a$ a similar argument holds. \square

Remark 4.2. *The theorem above essentially says that it is impossible for multiple two-sided Markov families to have the same single site density. However, which Λ^1 correspond to a two-sided Markov family, i.e., which Λ^1 are extendable is at this point an open question. This will be the subject of the next subsection, in particular in theorem 4.5 below we give a full characterization.*

4.2 Recursive construction

By virtue of proposition 2.4 we can attempt to recursively construct a family $\{\Lambda^n\}_{n \geq 1}$ from a suitable, symmetric, Λ^1 . In the proof of the theorem below we find a construction as described in remark 2.5, i.e. we find suitable left and right marginals to construct Λ^{n+1} from Λ^n . before we state this theorem, we first prove two preliminary results.

Lemma 4.3. *Let $\{\Lambda^n\}_{n \geq 1}$ be a symmetric two-sided Markov family. For arbitrary $n \geq 1$ and $y \in \mathbb{R}^+$, there exist y -dependent functions f_n, g_n and h_n on $(\mathbb{R}^+)^2$ such that*

$$\Lambda_{a,b}^{n,1}(x) = f_n(a, b) \cdot g_n(b, x) \cdot h_n(x, a) \cdot \mathbb{1}(x \in I_{a,b}) \tag{51}$$

for all $a, x, b \in \mathbb{R}^+$ such that $x \in I_{a,b}$ and $b \in I_{a,y}$. If a density is of this form we will refer to it as factorizable.

Proof. By conditioning on θ_1 and θ_{n+1} we find

$$\Lambda_{a,y}^{n+1,1}(\theta_1) \cdot \Lambda_{\theta_1, \theta_{n+1}}^{n-1}(\theta_2, \dots, \theta_n) \cdot \Lambda_{\theta_1, y}^{n,n}(\theta_{n+1}) = \Lambda_{a, \theta_{n+1}}^{n,1}(\theta_1) \cdot \Lambda_{\theta_1, \theta_{n+1}}^{n-1}(\theta_2, \dots, \theta_n) \cdot \Lambda_{a,y}^{n+1,n+1}(\theta_{n+1}), \tag{52}$$

where both sides of the equation are equal to $\Lambda_{a,y}^{n+1}(\theta_1, \dots, \theta_{n+1})$. To proceed we need information on the support of $\Lambda_{a,y}^{n+1}$. To this end we assume $a < y$. As a consequence, we can pick $a < \theta_1 < \dots < \theta_{n+1} < y$, then

$$\frac{\Lambda_{a,y}^{n+1,1}(\theta_1)}{\Lambda_{y,a}^{n+1,1}(\theta_{n+1})} = \frac{\Lambda_{a,y}^{n+1,1}(\theta_1)}{\Lambda_{a,y}^{n+1,n+1}(\theta_{n+1})} = \frac{\Lambda_{a,\theta_{n+1}}^{n,1}(\theta_1)}{\Lambda_{\theta_1,y}^{n,n}(\theta_{n+1})} = \frac{\Lambda_{a,\theta_{n+1}}^{n,1}(\theta_1)}{\Lambda_{y,\theta_1}^{n,1}(\theta_{n+1})}. \quad (53)$$

The first and third equality use the symmetry property. The second equality follows from (52). The same equation can be derived for $y < a$ and $\theta_1 > \theta_{n+1}$, hence we continue with general $a, y \in \mathbb{R}^+$ and $\theta_1 \in I_{a,\theta_{n+1}}, \theta_{n+1} \in I_{a,y}$. Notice that in the case $a < y$ we have $a < x < b < y$, whereas in the case $y < a$ we have $y < b < x < a$. That is, in the first case y acts as an upper bound for the interval $I_{a,b}$ and in the second case it acts like a lower bound. We observe that

$$\Lambda_{a,\theta_{n+1}}^{n,1}(\theta_1) = [\Lambda_{y,a}^{n+1,1}(\theta_{n+1})]^{-1} \cdot \Lambda_{y,\theta_1}^{n,1}(\theta_{n+1}) \cdot \Lambda_{a,y}^{n+1,1}(\theta_1). \quad (54)$$

One can now take $\theta_1 = x$ and $\theta_{n+1} = b$ to obtain

$$\Lambda_{a,b}^{n,1}(x) = [\Lambda_{y,a}^{n+1,1}(b)]^{-1} \cdot \Lambda_{y,x}^{n,1}(b) \cdot \Lambda_{a,y}^{n+1,1}(x). \quad (55)$$

We can then define the functions f_n, g_n and h_n as follows

$$f_n(a, b) := [\Lambda_{y,a}^{n+1,1}(b)]^{-1}, \quad g_n(b, x) := \Lambda_{y,x}^{n,1}(b), \quad h_n(x, a) := \Lambda_{a,y}^{n+1,1}(x). \quad (56)$$

□

Lemma 4.4. *Let $\{\Lambda^n\}_{n \geq 1}$ be a symmetric two-sided Markov family. For arbitrary $z_1, z_2 \in \mathbb{R}^+$, $z_1 \leq z_2$, there exist functions f, g and h on \mathbb{R} such that*

$$\Lambda_{a,b}^1(x) = f(a, b) \cdot g(b, x) \cdot h(x, a) \cdot \mathbf{1}(x \in I_{a,b}) \quad (57)$$

for all $a, b \in (z_1, z_2)$. Moreover, the functions f and g can be chosen to be symmetric.

Proof. Let $Z_1 > 0$ and $Z_2 > 0$ be such that $Z_1 < z_1$ and $z_2 < Z_2$. Let $a, b \in (Z_1, Z_2)$ and assume $a \leq b$. Then $b \in I_{a,Z_2}$, hence proposition 4.3 gives us functions \widehat{f}, \widehat{g} and \widehat{h} such that

$$\Lambda_{a,b}^1(x) = \widehat{f}(a, b) \cdot \widehat{g}(b, x) \cdot \widehat{h}(x, a) \cdot \mathbf{1}(x \in I_{a,b}). \quad (58)$$

Similarly we can assume $b \leq a$, then $b \in I_{Z_1,a}$, hence there are functions $\widetilde{f}, \widetilde{g}$ and \widetilde{h} such that

$$\Lambda_{a,b}^1(x) = \widetilde{f}(a, b) \cdot \widetilde{g}(b, x) \cdot \widetilde{h}(x, a) \cdot \mathbf{1}(x \in I_{a,b}). \quad (59)$$

hence we can define f, g and h as

$$f(a, b) = \begin{cases} \widehat{f}(a, b) & \text{if } a < b \\ \widetilde{f}(a, b) & \text{if } b < a \end{cases} \quad g(b, x) = \begin{cases} \widehat{g}(b, x) & \text{if } x < b \\ \widetilde{g}(b, x) & \text{if } b < x \end{cases} \quad h(x, a) = \begin{cases} \widehat{h}(x, a) & \text{if } a < x \\ \widetilde{h}(x, a) & \text{if } x < a \end{cases}$$

This show the first part of the lemma, we proceed with general $a, b \in I_{Z_1, Z_2}$ and $x \in I_{a,b}$. Now we only have to show that we can choose f and g symmetric. We define $\psi(a, x) = \sqrt{g(a, x) \cdot h(x, a)}$ and $\phi(a, b) = \sqrt{f(a, b) \cdot f(b, a)}$. Notice that, using the symmetry of the family $\{\Lambda^n\}_{n \geq 1}$,

$$\begin{aligned}
\Lambda_{a,b}^1(x) &= \sqrt{\Lambda_{a,b}^1(x) \cdot \Lambda_{b,a}^1(x)} \\
&= \sqrt{f(a,b) \cdot f(b,a)} \cdot \sqrt{g(a,x) \cdot h(x,a)} \cdot \sqrt{g(b,x) \cdot h(x,b)} \\
&= \phi(a,b) \cdot \psi(a,x) \cdot \psi(b,x).
\end{aligned} \tag{60}$$

We now restrict a, b to (z_1, z_2) and claim that

$$\frac{\psi(y,x)}{\psi(x,y)} = \frac{c_2(y)\mathbf{1}(x < y) + c_1(y)\mathbf{1}(y < x)}{c_1(x)\mathbf{1}(x < y) + c_2(x)\mathbf{1}(y < x)} \tag{61}$$

for some functions c_1 and c_2 and $x \neq y$, $x, y \in (z_1, z_2)$. Before we prove this claim we demonstrate how we will use it. If the claim holds, then we have

$$\begin{aligned}
\Psi(x,y) &:= [c_1(x)\mathbf{1}(x < y) + c_2(x)\mathbf{1}(y < x)] \cdot \psi(y,x) \\
&= [c_2(y)\mathbf{1}(x < y) + c_1(y)\mathbf{1}(y < x)] \cdot \psi(x,y) = \Psi(y,x)
\end{aligned} \tag{62}$$

for all $x, y \in (z_1, z_2)$ with $x \neq y$. As a consequence

$$\Lambda_{a,b}^1(x) = \phi(a,b) \cdot \psi(a,x) \cdot \psi(b,x) \cdot \mathbf{1}(x \in I_{a,b}) = \phi(a,b) \cdot \Psi(b,x) \cdot \gamma(a,x) \cdot \mathbf{1}(x \in I_{a,b}). \tag{63}$$

Here we define γ , for $x \in I_{a,b}$, as

$$\begin{aligned}
\gamma(a,x) &:= [c_1(x) \cdot \mathbf{1}(a < x) + c_2(x) \cdot \mathbf{1}(x < a)]^{-1} \psi(a,x) \\
&= [c_1(x) \cdot \mathbf{1}(x < b) + c_2(x) \cdot \mathbf{1}(b < x)]^{-1} \psi(a,x).
\end{aligned} \tag{64}$$

Notice that the last equality relies on the fact that $x \in I_{a,b}$. Indeed, for $x \in I_{a,b}$: $a < x \iff x < b$ and $b < x \iff x < a$. We can pick $f = \phi$, $g = \Psi$ and $h = \gamma$. Since ϕ and Ψ are symmetric, this completes the proof.

Finally, we prove the claim (61). Recall that (60) holds on the whole interval (Z_1, Z_2) rather than only on the subinterval (z_1, z_2) . Let $x, y \in I_{z_1, z_2}$. If we have $x < y$ then

$$\frac{\Lambda_{z_1, z_2}^{2,1}(x)}{\Lambda_{z_2, z_1}^{2,1}(y)} = \frac{\Lambda_{z_1, y}^1(x)}{\Lambda_{z_2, x}^1(y)} = \frac{\phi(z_1, y) \cdot \psi(z_1, x) \cdot \psi(y, x)}{\phi(z_2, x) \cdot \psi(z_2, y) \cdot \psi(x, y)}. \tag{65}$$

Define $C_{1,2}(x) := \frac{\psi(z_1, x)}{\Lambda_{z_1, z_2}^{2,1}(x) \cdot \phi(z_2, x)}$ and $C_{2,1}(y) := \frac{\psi(z_2, y)}{\Lambda_{z_2, z_1}^{2,1}(y) \cdot \phi(z_1, y)}$. The equation above reads then

$$\frac{C_{2,1}(y)}{C_{1,2}(x)} = \frac{\psi(y, x)}{\psi(x, y)}. \tag{66}$$

Now assume $y < x$, similarly we find

$$\frac{C_{1,2}(y)}{C_{2,1}(x)} = \frac{\psi(y, x)}{\psi(x, y)}. \tag{67}$$

Combining these two expressions yields

$$\frac{\psi(y, x)}{\psi(x, y)} = \frac{C_{2,1}(y) \cdot \mathbf{1}(x < y) + C_{1,2}(y) \cdot \mathbf{1}(y < x)}{C_{1,2}(x) \cdot \mathbf{1}(x < y) + C_{2,1}(x) \cdot \mathbf{1}(y < x)} \tag{68}$$

We define $c_1 = C_{1,2}$ and $c_2 = C_{2,1}$ to obtain the required form. This proves (61). \square

We can now state a characterization of the Λ^1 which generate a symmetric two-sided Markov family.

Theorem 4.5. *The following two statements are equivalent,*

1. *The family $\{\Lambda^n\}_{n \geq 0}$ is symmetric and two-sided Markov*
2. *$\forall z_1, z_2 \in \mathbb{R}^+, a, b \in I_{z_1, z_2}$ there exists a symmetric $g : \mathbb{R}^2 \rightarrow \mathbb{R}^+$ such that*

$$\Lambda_{a,b}^1(x) = f(a, b) \cdot g(b, x) \cdot g(x, a) \cdot \mathbf{1}(x \in I_{a,b}) \quad (69)$$

with normalization

$$f(a, b) = \left(\int_{a \wedge b}^{a \vee b} g(b, y) \cdot g(y, a) dy \right)^{-1}. \quad (70)$$

Proof. 1 implies 2. Lemma 4.4 states that we can write

$$\Lambda_{a,b}^1(x) = \tilde{f}(a, b) \cdot \tilde{g}(b, x) \cdot \tilde{h}(x, a) \cdot \mathbf{1}(x \in I_{a,b}) \quad (71)$$

with \tilde{f} and \tilde{g} symmetric. Using the symmetry of Λ^1 , we can deduce that \tilde{h} is of the form $\tilde{h}(x, a) = c(x) \cdot \tilde{g}(x, a)$ for some function c . Indeed, the symmetry gives

$$\tilde{f}(a, b) \cdot \tilde{g}(b, x) \cdot \tilde{h}(x, a) = \tilde{f}(b, a) \cdot \tilde{g}(a, x) \cdot \tilde{h}(x, b) \quad (72)$$

for $x \in I_{a,b}$. Using the symmetry of \tilde{f} , we obtain

$$\frac{\tilde{g}(b, x)}{\tilde{h}(x, b)} = \frac{\tilde{g}(a, x)}{\tilde{h}(x, a)} =: c^{-1}(x). \quad (73)$$

This then gives us $\tilde{h}(x, b) = c(x) \cdot \tilde{g}(b, x)$. We define $g(a, x) := \sqrt{c(x) \cdot c(a)} \cdot \tilde{g}(a, x)$ and $f(a, b) := \tilde{f}(a, b) / \sqrt{c(a) \cdot c(b)}$. Notice that f and g are symmetric. We can now write Λ^1 in the form

$$\Lambda_{a,b}^1(x) = f(a, b) \cdot g(a, x) \cdot g(x, b). \quad (74)$$

Since $\Lambda_{a,b}^1$ is a probability density, f must indeed have the form (70).

2 implies 1. We define the family $\{\Lambda^n\}_{n \geq 1}$ recursively. Let $z_1, z_2 \in \mathbb{R}^+$ and assume that for some $m \geq 1$, the first marginals $\Lambda^{1,1}, \dots, \Lambda^{m,1}$ have the form

$$\Lambda_{a,b}^{k,1}(x) = f_k(a, b) \cdot g_k(b, x) \cdot h_k(x, a) \cdot \mathbf{1}(x \in I_{a,b}) \quad (75)$$

for $a, b \in (z_1, z_2)$, and $1 \leq k \leq m$ with f_k and g_k symmetric. Notice that this is true for Λ^1 with $f_1(a, b) := f(a, b)$, $g_1(b, x) := g(b, x)$ and $h_1(x, a) = g(x, a)$. We define $\Lambda_{a,b}^{m+1,1}$ and $\Lambda_{a,b}^{m+1,m+1}$ as

$$\Lambda_{a,b}^{m+1,1}(x) = f_{m+1}(a, b) \cdot [f_m(b, x)]^{-1} \cdot h_m(x, a) \cdot \mathbf{1}(x \in I_{a,b}) \quad (76)$$

and

$$\Lambda_{a,b}^{m+1,m+1}(x) = \Lambda_{b,a}^{m+1,1}(x) = f_{m+1}(b, a) \cdot [f_m(a, x)]^{-1} \cdot h_m(x, b) \cdot \mathbf{1}(x \in I_{a,b}) \quad (77)$$

with f_{m+1} the normalization

$$f_{m+1}(a, b) = \left(\int_{a \wedge b}^{a \vee b} [f_m(b, x)]^{-1} \cdot h_m(x, a) dx \right)^{-1}. \quad (78)$$

We show that f_{m+1} is symmetric. Assume $a \leq b$. Then, using the recursion,

$$\begin{aligned} & [f_{m+1}(a, b)]^{-1} \\ &= \int_a^b g(a, x_1) \int_{x_1}^b g(x_1, x_2) \dots \int_{x_m}^b g(x_{m+1}, b) dx_{m+1} \dots dx_2 dx_1 \\ &= \int_a^b \int_{x_1}^b \dots \int_{x_m}^b g(a, x_1) \cdot g(x_1, x_2) \cdot \dots \cdot g(x_{m+1}, b) dx_{m+1} \dots dx_2 dx_1 \\ &= \int_a^b \int_a^b \dots \int_a^b \mathbf{1}(a \leq x_1 \leq x_2 \leq \dots \leq x_{m+1} \leq b) \prod_{i=0}^n g(x_i, x_{i+1}) dx_1 dx_2 \dots dx_{m+1} \end{aligned} \quad (79)$$

and

$$\begin{aligned} & [f_{m+1}(b, a)]^{-1} \\ &= \int_a^b g(b, x_{m+1}) \int_a^{x_{m+1}} g(x_{m+1}, x_m) \dots \int_a^{x_2} g(x_1, a) dx_1 dx_2 \dots dx_{m+1} \\ &= \int_a^b \int_a^{x_{m+1}} \dots \int_a^{x_2} g(b, x_{m+1}) \cdot g(x_{m+1}, x_m) \cdot \dots \cdot g(x_1, a) dx_1 dx_2 \dots dx_{m+1} \\ &= \int_a^b \int_a^b \dots \int_a^b \mathbf{1}(a \leq x_1 \leq x_2 \leq \dots \leq x_{m+1} \leq b) \prod_{i=0}^n g(x_i, x_{i+1}) dx_1 dx_2 \dots dx_{m+1} \end{aligned} \quad (80)$$

where it is understood that $x_0 = a$ and $x_{m+2} = b$. This immediately gives the symmetry of f_{m+1} . We proceed to show that $\Lambda_{a,b}^{m+1,1}$ and $\Lambda_{a,b}^{m+1,m+1}$ indeed yield a density $\Lambda_{a,b}^{m+1}$ which is two sided Markov. For arbitrary $\theta_1, \dots, \theta_{m+1} \in \mathbb{R}^+$ we calculate

$$\begin{aligned} & \Lambda_{a,b}^{m+1,1}(\theta_1) \cdot \Lambda_{\theta_1,b}^m(\theta_2, \dots, \theta_{m+1}) = \Lambda_{a,b}^{m+1,1}(\theta_1) \cdot \Lambda_{\theta_1,\theta_{m+1}}^{m-1}(\theta_2, \dots, \theta_m) \cdot \Lambda_{\theta_1,b}^{m,m}(\theta_{m+1}) \\ &= f_{m+1}(a, b) \cdot [f_m(b, \theta_1)]^{-1} \cdot h_m(\theta_1, a) \cdot \Lambda_{\theta_1,\theta_{m+1}}^{m-1}(\theta_2, \dots, \theta_m) \\ &\quad \cdot f_m(b, \theta_1) \cdot g_m(\theta_1, \theta_{m+1}) \cdot h_m(\theta_{m+1}, b) \cdot \mathbf{1}(\theta_1 \in I_{a,b}, \theta_{m+1} \in I_{\theta_1,b}) \\ &= f_{m+1}(a, b) \cdot h_m(\theta_1, a) \cdot \Lambda_{\theta_1,\theta_{m+1}}^{m-1}(\theta_2, \dots, \theta_m) \\ &\quad \cdot g_m(\theta_1, \theta_{m+1}) \cdot h_m(\theta_{m+1}, b) \cdot \mathbf{1}(\theta_1 \in I_{a,b}, \theta_{m+1} \in I_{\theta_1,b}) \end{aligned} \quad (81)$$

and

$$\begin{aligned} & \Lambda_{a,\theta_{m+1}}^m(\theta_1, \dots, \theta_m) \cdot \Lambda_{a,b}^{m+1,m+1}(\theta_{m+1}) = \Lambda_{a,\theta_{m+1}}^{m,1}(\theta_1) \cdot \Lambda_{\theta_1,\theta_{m+1}}^{m-1}(\theta_2, \dots, \theta_m) \cdot \Lambda_{a,b}^{m+1,m+1}(\theta_{m+1}) \\ &= f_m(a, \theta_{m+1}) \cdot g_m(\theta_{m+1}, \theta_1) \cdot h_m(\theta_1, a) \cdot \Lambda_{\theta_1,\theta_{m+1}}^{m-1}(\theta_2, \dots, \theta_m) \\ &\quad \cdot f_{m+1}(b, a) \cdot [f_m(a, \theta_{m+1})]^{-1} \cdot h_m(\theta_{m+1}, b) \cdot \mathbf{1}(\theta_1 \in I_{a,\theta_{m+1}}, \theta_1 \in I_{a,b}) \\ &= f_{m+1}(a, b) \cdot h_m(\theta_1, a) \cdot \Lambda_{\theta_1,\theta_{m+1}}^{m-1}(\theta_2, \dots, \theta_m) \\ &\quad \cdot g_m(\theta_1, \theta_{m+1}) \cdot h_m(\theta_{m+1}, b) \cdot \mathbf{1}(\theta_1 \in I_{a,b}, \theta_{m+1} \in I_{\theta_1,b}). \end{aligned} \quad (82)$$

Observe that these two are equal. As is explained in remark 2.5, one can use proposition 2.4 to construct a Λ^{m+1} for which the two-sided Markov property holds. Since Λ^1 is clearly symmetric, the whole family $\{\Lambda^n\}_{n \geq 1}$ is symmetric as is shown in proposition 2.7. This concludes the proof. \square

We explicitly state the recursion from the proof of theorem 4.5. This is done in the definition below, alongside the introduction of some terminology.

Definition 4.6. Λ^1 is called a generating density if it is of the form in (69) for all $z_1, z_2 \in \mathbb{R}^+$ such that $z_1 < z_2$ and $a, b \in (z_1, z_2)$. The unique symmetric two-sided Markov family $\{\Lambda^n\}_{n \geq 1}$ with first marginals

$$\Lambda_{a,b}^{n,1}(x) = f_n(a, b) \cdot g_n(b, x) \cdot h_n(x, a) \cdot \mathbb{1}(x \in I_{a,b}) \quad (83)$$

is called the generated family. The functions f_n, g_n and h_n on \mathbb{R}^+ are defined via the recursion

$$\begin{cases} f_{n+1}(a, b) = \left(\int_{a \wedge b}^{a \vee b} [f_n(b, x)]^{-1} \cdot h_1(x, a) dx \right)^{-1} \\ g_{n+1}(b, x) = [f_n(a, x)]^{-1} \\ h_{n+1}(x, a) = h(x, a) = g(x, a) \end{cases} \quad (84)$$

with $f_1 = f$ and $g_1 = g$. For fixed z_1, z_2 we call (f_n, g_n, h_n) the n -th order factors generated by Λ^1 . The functions f_n and g_n are symmetric for all $n \geq 1$, as is shown in the proof of theorem 4.5.

Remark 4.7. Notice that in the definition above, the n -th order factors are all constructed from just one function. Indeed, given only g_1 , we know h_1 and f_1 since h_1 is simply defined to be equal to g_1 and $f_1(a, b)$ is the normalization of $g_1(b, x) \cdot h_1(x, a)$. The recursive construction gives the higher order factors.

Example 4.8. We verify that the density of the order statistics are indeed recursively generated according to (84). (8) evaluated in $i = 1$ yields

$$\Lambda_{a,b}^{n,1}(x) = n \cdot |b - a|^{-n} |b - x|^{n-1} \mathbb{1}(x \in I_{a,b}). \quad (85)$$

This expression is clearly factorizable for $x \in I_{a,b}$, we can take

$$f_n(a, b) = n! |b - a|^{-n}, \quad g_n(b, x) = \frac{|b - x|^{n-1}}{(n-1)!}, \quad h_n(x, a) = 1. \quad (86)$$

We immediately see that $f_1(a, b) = |b - a|^{-1}$ and $g_1(b, x) = 1$ are symmetric, hence Λ^1 is indeed generating. Moreover f_n and g_n are symmetric and satisfy the recursion,

$$f_{n+1}^{-1}(a, b) = \int_{a \wedge b}^{a \vee b} h_1(x, a) \cdot f_n^{-1}(b, x) dx = \int_{a \wedge b}^{a \vee b} \frac{|b - x|^n}{n!} dx = \frac{|b - a|^{n+1}}{(n+1)!} \quad (87)$$

and

$$g_{n+1}(b, x) = [f_n(b, x)]^{-1} = \frac{|b - x|^n}{n!}. \quad (88)$$

We conclude that f_n, g_n and h_n are indeed the n -th order factors generated by Λ^1 .

Example 4.9 (gapped order statistics). In example 4.8 we reconstructed the joint density corresponding to the order statistics of uniform random variables. In this case we had $g_1(a, b) = 1$, defining the whole family. Since g_1 is indeed symmetric strictly positive, we could have directly seen that the order statistics are symmetric and two-sided Markov. In fact we recover related families if we take $g_1(a, b) = [(s-1)!]^{-1} \cdot |b - a|^{s-1}$ for any integer $s \geq 1$. For this choice of g_1 we obtain the so-called *gapped* order statistics of the uniform random variables. These are defined as follows. For fixed $n \geq 1$ and $a, b \in \mathbb{R}^+$ we define $N = s(n+1) - 1$ and consider

$X_1, X_2, \dots, X_N \sim U(a \wedge b, a \vee b)$. Then $\Theta_1 := X_{s:N}, \Theta_2 := X_{2s:N}, \dots, \Theta_n := X_{ns:N}$ are the gapped order statistics with gap size s . For $a < b$, their joint densities and first marginals are known [15] to be

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \frac{N!}{[(s-1)!]^n} \cdot \frac{1}{|b-a|^n} \prod_{i=1}^{n+1} (\theta_i - \theta_{i-1})^{s-1} \cdot \mathbf{1}(a < \theta_1 < \dots < \theta_n < b) \quad (89)$$

and

$$\Lambda_{a,b}^{n,1}(\theta) = \frac{(s(n+1)-1)!}{(s-1)!(sn-1)!} \cdot \frac{1}{|b-a|^{s(n+1)-1}} \cdot |b-\theta|^{sn-1} \cdot |b-\theta|^{s-1} \cdot |\theta-a|^{s-1} \mathbf{1}(\theta \in I_{a,b}) \quad (90)$$

where it is understood that $\theta_0 = a$ and $\theta_{n+1} = b$. Akin to the density for $s = 1$ in example 2.3, the indicator is replaced by $\mathbf{1}(b < \theta_n < \dots < \theta_1 < a)$ in the case where $b < a$. One can check that this family is symmetric and two-sided Markov by directly verifying definition 2.1. We demonstrate the recursion in (84). Notice that $\Lambda_{a,b}^{n,1}$ is indeed factorizable for

$$f_n(a, b) = \frac{(s(n+1)-1)!}{|b-a|^{s(n+1)-1}}, \quad g_n(b, \theta) = \frac{|b-\theta|^{sn-1}}{(sn-1)!}, \quad h_n(\theta, a) = \frac{|\theta-a|^{s-1}}{(s-1)!}. \quad (91)$$

Take $n = 1$. We can immediately see that f_1, g_1 and h_1 define a generating density. f_n also follows from the recursion since $g_n = f_{n-1}^{-1}$.

Example 4.10 (Dirichlet processes). *Notice that we can in fact generalize the example above by taking $s \in \mathbb{R}^+$ and defining*

$$f_n(a, b) = \frac{\Gamma(s(n+1))}{|b-a|^{s(n+1)-1}}, \quad g_n(b, \theta) = \frac{|b-\theta|^{sn-1}}{\Gamma(sn)}, \quad h_n(\theta, a) = \frac{|\theta-a|^{s-1}}{\Gamma(s)}. \quad (92)$$

It is again easy to check that the recursion in (84) works. These generalized distributions are known in literature under the name ordered Dirichlet distribution [1]. The joint density is then given by the analogue of (89):

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \frac{\Gamma(s(n+1))}{[\Gamma(s)]^n} \cdot \frac{1}{|b-a|^n} \prod_{i=1}^{n+1} (\theta_i - \theta_{i-1})^{s-1} \cdot \mathbf{1}(a < \theta_1 < \dots < \theta_n < b) \quad (93)$$

5 Characterizing properties of Dirichlet densities

In this section we find characterizing properties for $\{\Lambda^n\}_{n \geq 1}$ corresponding to an ordered Dirichlet distribution, i.e., the family given in (93). In theorem 4.5, we provided a characterization of those Λ^1 which are part of a two-sided symmetric Markov family $\{\Lambda^n, n \in \mathbb{N}\}$. Due to this characterization, it is easy to generate a large class of symmetric two-sided families. Indeed, we can perform the construction in definition 4.6 for an arbitrary integrable and symmetric function $g : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$. However, we will prove that under natural assumptions such as scaling and shift invariance (detailed below), the only two-sided Markov families are the Dirichlet processes of example 4.10.

Definition 5.1. *The family $\{\Lambda^n\}_{n \geq 1}$ is*

1. **Scale invariant** if for all $a, b, \gamma \in \mathbb{R}^+, n \geq 1$ and $\theta_1 \in I_{a,b}, \theta_2 \in I_{\theta_1,b}, \dots, \theta_n \in I_{\theta_{n-1},b}$,

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \gamma^n \cdot \Lambda_{\gamma \cdot a, \gamma \cdot b}^n(\gamma \cdot \theta_1, \dots, \gamma \cdot \theta_n). \quad (94)$$

2. **Shift invariant** if, for all $a, b \in \mathbb{R}^+$, $\gamma > -(a \wedge b)$, $n \geq 1$ and $\theta_1 \in I_{a,b}, \theta_2 \in I_{\theta_1,b}, \dots, \theta_n \in I_{\theta_{n-1},b}$

$$\Lambda_{a,b}^n(\theta_1, \dots, \theta_n) = \Lambda_{a+\gamma, b+\gamma}^n(\theta_1 + \gamma, \dots, \theta_n + \gamma). \quad (95)$$

We start with a useful lemma.

Lemma 5.2. *Let $\phi : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be a continuous symmetric function which satisfies the following scale and shift properties*

$$\begin{cases} \phi(\gamma \cdot x, \gamma \cdot y) = \tilde{\lambda}_1(\gamma) \cdot e^{\tilde{\lambda}_2(\gamma)|x-y|} \cdot \phi(x, y) \\ \phi(x + \gamma, y + \gamma) = \hat{\lambda}_1(\gamma) \cdot e^{\hat{\lambda}_2(\gamma)|x-y|} \cdot \phi(x, y) \end{cases} \quad (96)$$

for some continuous functions $\tilde{\lambda}_1, \tilde{\lambda}_2, \hat{\lambda}_1, \hat{\lambda}_2 : \mathbb{R}^+ \rightarrow \mathbb{R}^+$. Then ϕ is of the form

$$\phi(x, y) = t \cdot |x - y|^\sigma \quad (97)$$

with $\sigma, t \in \mathbb{R}$, i.e., $\tilde{\lambda}_1(\gamma) = \gamma^\sigma$, $\hat{\lambda}_1 = 1$ and $\tilde{\lambda}_2 = \hat{\lambda}_2 = 0$.

Proof. First we will show that $\tilde{\lambda}_1(a) = a^\sigma$ and $\tilde{\lambda}_2(a) = u(a - 1)$.

We have the following restriction on $\tilde{\lambda}_1$ and $\tilde{\lambda}_2$. Let $a, b \in \mathbb{R}^+$

$$\tilde{\lambda}_1(a) \cdot \tilde{\lambda}_1(b) \cdot e^{-[\tilde{\lambda}_2(a) \cdot b + \tilde{\lambda}_2(b)] \cdot |x-y|} \cdot \phi(x, y) = \phi(a \cdot b \cdot x, a \cdot b \cdot y) = \tilde{\lambda}_2(a \cdot b) \cdot e^{-\tilde{\lambda}_2(a \cdot b) |x-y|} \cdot \phi(x, y). \quad (98)$$

Since this expression must hold for all x and y , we have the two equations for $\tilde{\lambda}_1$ and $\tilde{\lambda}_2$,

$$\tilde{\lambda}_1(a \cdot b) = \tilde{\lambda}_1(a) \cdot \tilde{\lambda}_1(b) \quad \text{and} \quad \tilde{\lambda}_2(a) \cdot b + \tilde{\lambda}_2(b) = \tilde{\lambda}_2(a \cdot b). \quad (99)$$

This shows that $\tilde{\lambda}_1$ is of the form $\tilde{\lambda}_1(\gamma) = \gamma^\sigma$, for some $\sigma \in \mathbb{R}$. The second equation tells us that $\tilde{\lambda}_2(\gamma) = u \cdot (\gamma - 1)$ for $u \in \mathbb{R}$. To see this, use (99) to write

$$\tilde{\lambda}_2(a) \cdot b + \tilde{\lambda}_2(b) = \tilde{\lambda}_2(a \cdot b) = \tilde{\lambda}_2(b) \cdot a + \tilde{\lambda}_2(a). \quad (100)$$

Assume that $\tilde{\lambda}_2(a) \cdot \tilde{\lambda}_2(b) \neq 0$. We divide by $\tilde{\lambda}_2(a) \cdot \tilde{\lambda}_2(b)$ and separate the equation in a and b ,

$$u^{-1} := \frac{a}{\tilde{\lambda}_2(a)} - \frac{1}{\tilde{\lambda}_2(a)} = \frac{b}{\tilde{\lambda}_2(b)} - \frac{1}{\tilde{\lambda}_2(b)}. \quad (101)$$

This implies that $\tilde{\lambda}_2$ is of the form $\tilde{\lambda}_2(\gamma) = u \cdot (\gamma - 1)$ or $\tilde{\lambda}_2(\gamma) = 0$. Since $\tilde{\lambda}_2(\gamma)$ is continuous, we have $\tilde{\lambda}_2(\gamma) = u \cdot (\gamma - 1)$ for some $u \in \mathbb{R}$, which might be zero. We can perform a similar procedure as we did above for the shift property. That is, we start from the requirement

$$\hat{\lambda}_1(a) \cdot \hat{\lambda}_1(b) \cdot e^{-[\hat{\lambda}_2(a) + \hat{\lambda}_2(b)] \cdot |x-y|} \cdot \phi(x, y) = \phi(a + b + x, a + b + y) = \hat{\lambda}_2(a + b) \cdot e^{-\hat{\lambda}_2(a+b) |x-y|} \cdot \phi(x, y). \quad (102)$$

Again, we can argue that this equation should hold for all x and y to obtain

$$\hat{\lambda}_1(a + b) = \hat{\lambda}_1(a) \cdot \hat{\lambda}_1(b) \quad \text{and} \quad \hat{\lambda}_2(a) + \hat{\lambda}_2(b) = \hat{\lambda}_2(a + b) \quad (103)$$

for arbitrary $a, b \in \mathbb{R}^+$. The first equation shows that $\hat{\lambda}_1(\gamma) = \exp(v \cdot \gamma)$, $v \in \mathbb{R}$ while the second equation gives $\hat{\lambda}_2(\gamma) = w \cdot \gamma$, $w \in \mathbb{R}$. Now that we know the general form of $\tilde{\lambda}_1, \tilde{\lambda}_2, \hat{\lambda}_1$ and $\hat{\lambda}_2$, we find an explicit form for ϕ and argue that $u, v, w = 0$. Assume $y > x$, then

$$\begin{aligned}
\phi(x, y) &= \widehat{\lambda}_1(x) \cdot e^{\widehat{\lambda}_2(x)|y-x|} \cdot \phi(0, y-x) \\
&= \widehat{\lambda}_1(x) \cdot e^{\widehat{\lambda}_2(x)|y-x|} \cdot \widetilde{\lambda}_1(y-x) \cdot e^{\widetilde{\lambda}_2(y-x)|y-x|} \cdot \phi(0, 1) \\
&= t \cdot e^{v \cdot x} \cdot e^{w \cdot x(y-x)} \cdot (y-x)^\sigma \cdot e^{u(y-x-1)(y-x)},
\end{aligned} \tag{104}$$

where $t := \phi(0, 1)$. We will then show that $u = v = w = 0$.

By the scaling property,

$$\begin{aligned}
&\phi(\gamma \cdot x, \gamma \cdot y) \\
&= t \cdot \gamma^\sigma \cdot (y-x)^\sigma \cdot \exp[\gamma(vx - u(y-x)) + \gamma^2(wx(y-x) + u(y-x)^2)]
\end{aligned} \tag{105}$$

and

$$\begin{aligned}
\phi(\gamma \cdot x, \gamma \cdot y) &= \widetilde{\lambda}_1(\gamma) \cdot \exp[\widetilde{\lambda}_2(\gamma)(y-x)] \cdot \phi(x, y) \\
&= \gamma^\sigma \cdot \exp[u(\gamma-1) \cdot (y-x)] \cdot \phi(x, y) \\
&= t \cdot \gamma^\sigma \cdot \exp[u(\gamma-1) \cdot (y-x)] \cdot (y-x)^\sigma \cdot \exp[(vx - u(y-x)) + (wx(y-x) + u(y-x)^2)].
\end{aligned} \tag{106}$$

These two expressions should be equal for all $x, y \in \mathbb{R}$, which is only possible if u, v and w are zero. In that case we find $\phi(x, y) = t \cdot (y-x)^\sigma$. Clearly this expression satisfies the shift property. Due to the symmetry assumption on ϕ we have, for general x, y , we finally obtain

$$\phi(x, y) = t \cdot |y-x|^\sigma. \tag{107}$$

□

Theorem 5.3. Let $\{\Lambda^n\}_{n \geq 1}$ be a symmetric, two-sided Markov family, i.e., Λ^1 is of the form (69)

$$\Lambda_{a,b}^1(x) = f(a, b) \cdot g(b, x) \cdot g(a, x) \tag{108}$$

with

$$f(a, b)^{-1} = \int_{a \wedge b}^{a \vee b} g(a, x) \cdot g(b, x) dx \tag{109}$$

for some symmetric function $g : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$. We assume $g(x, \cdot)$ to be differentiable on $\mathbb{R}^+ \setminus \{x\}$ for all $x \in \mathbb{R}^+$.

If $\{\Lambda^n\}_{n \geq 1}$ is both scale and shift invariant, then, for some $s, t \in \mathbb{R}^+$

$$\Lambda_{a,b}^1(x) = f(a, b) \cdot (t \cdot |b-x|^{s-1}) \cdot (t \cdot |x-a|^{s-1}). \tag{110}$$

That is, we recover the density corresponding to the Dirichlet process with parameter s from example 4.10.

Proof. Notice that for $\theta_1, \theta_2 \in I_{a,b}$

$$\frac{\Lambda_{a,b}^1(\theta_1)}{\Lambda_{a,b}^1(\theta_2)} = \frac{\Lambda_{\gamma \cdot a, \gamma \cdot b}^1(\gamma \cdot \theta_1)}{\Lambda_{\gamma \cdot a, \gamma \cdot b}^1(\gamma \cdot \theta_2)}. \tag{111}$$

Bringing all the factors containing θ_1 to the left hand side and all the terms containing θ_2 to the right hand side shows that for any $\theta \in I_{a,b}$,

$$\frac{g(a, \theta) \cdot g(b, \theta)}{g(\gamma \cdot a, \gamma \cdot \theta) \cdot g(\gamma \cdot b, \gamma \cdot \theta_1)} = \frac{\Lambda_{a,b}^1(\theta)}{\Lambda_{\gamma \cdot a, \gamma \cdot b}^1(\gamma \cdot \theta)} = \tilde{\kappa}_1(a, b, \gamma). \quad (112)$$

Define $\tilde{g}(a, \theta) := \log[g(a, \theta)/g(\gamma \cdot a, \gamma \cdot \theta)]$. We differentiate with respect to θ and obtain

$$\tilde{g}^{(0,1)}(a, \theta) + \tilde{g}^{(0,1)}(b, \theta) = 0, \quad (113)$$

where $\tilde{g}^{(0,1)}$ denotes the derivative of \tilde{g} in the second entry. As a consequence

$$\begin{cases} \tilde{g}^{(0,1)}(a, \theta) = -\tilde{g}^{(0,1)}(b, \theta) & \text{for } \theta \in I_{a,b} \\ \tilde{g}^{(0,1)}(a, \theta) = \tilde{g}^{(0,1)}(b, \theta) & \text{for } \theta \notin I_{a,b}. \end{cases} \quad (114)$$

This is easily seen as follows. Consider the case $a, b < \theta$ and let $c > \theta$. Then $\tilde{g}^{(0,1)}(a, \theta) = -\tilde{g}^{(0,1)}(c, \theta)$ and $\tilde{g}^{(0,1)}(b, \theta) = -\tilde{g}^{(0,1)}(c, \theta)$, so we indeed have $\tilde{g}^{(0,1)}(a, \theta) = \tilde{g}^{(0,1)}(b, \theta)$. A similar argument can be made for the case where $a, b > \theta$. We can now directly infer that

$$\tilde{g}^{(0,1)}(a, \theta) = \begin{cases} -\tilde{\kappa}_2(\gamma) & \text{for } \theta < a \\ \tilde{\kappa}_2(\gamma) & \text{for } \theta > a. \end{cases} \quad (115)$$

As a consequence,

$$\tilde{g}(a, \theta) = \begin{cases} -\tilde{\kappa}_2(\gamma) \cdot \theta + \tilde{\kappa}_3(a, \gamma) & \text{for } \theta < a \\ \tilde{\kappa}_2(\gamma) \cdot \theta + \tilde{\kappa}_4(a, \gamma) & \text{for } \theta > a. \end{cases} \quad (116)$$

We now use the explicit form of \tilde{g} to find a scaling property for g . If $a < \theta$, then

$$\log \frac{g(a, \theta)}{g(\gamma \cdot a, \gamma \cdot \theta)} = \tilde{\kappa}_4(a, \gamma) + \tilde{\kappa}_2(\gamma) \cdot \theta. \quad (117)$$

Likewise, if $\theta < a$, then

$$\log \frac{g(a, \theta)}{g(\gamma \cdot a, \gamma \cdot \theta)} = \tilde{\kappa}_3(a, \gamma) - \tilde{\kappa}_2(\gamma) \cdot \theta. \quad (118)$$

For now we assume $a < \theta$. Using the symmetry of g we obtain the following relation

$$\tilde{\kappa}_3(a, \gamma) - \tilde{\kappa}_2(\gamma) \cdot \theta = \tilde{\kappa}_4(\theta, \gamma) + \tilde{\kappa}_2(\gamma) \cdot a. \quad (119)$$

We do separation of variables in a and θ . This yields the following expression which is independent of θ and a

$$\tilde{\kappa}_5(\gamma) := \tilde{\kappa}_3(a, \gamma) - \tilde{\kappa}_2(\gamma) \cdot a = \tilde{\kappa}_4(\theta, \gamma) + \tilde{\kappa}_2(\gamma) \cdot \theta. \quad (120)$$

We rewrite $\log g(a, \theta)/g(\gamma \cdot a, \gamma \cdot \theta)$ as

$$\log \frac{g(a, \theta)}{g(\gamma \cdot a, \gamma \cdot \theta)} = \tilde{\kappa}_5(\gamma) + \tilde{\kappa}_2(\gamma)(\theta - a). \quad (121)$$

Under the assumption $\theta < a$ we find

$$\log \frac{g(a, \theta)}{g(\gamma \cdot a, \gamma \cdot \theta)} = \tilde{\kappa}_5(\gamma) + \tilde{\kappa}_2(\gamma)(a - \theta). \quad (122)$$

Now we have the following relation between $g(a, \theta)$ and $g(\gamma \cdot a, \gamma \cdot \theta)$

$$g(\gamma \cdot a, \gamma \cdot \theta) = e^{-\tilde{\kappa}_5(\gamma)} \cdot e^{-\tilde{\kappa}_2(\gamma) \cdot |a - \theta|} \cdot g(a, \theta) \quad (123)$$

Notice this scaling property for g is derived from just the scale invariance. Moreover, one can mutatis mutandis repeat the procedure above to derive a shift property. This leaves us with the following expressions for scaling and shifts

$$\begin{cases} g(a + \gamma, \theta + \gamma) = e^{-\widehat{\kappa}_5(\gamma)} \cdot e^{-\widehat{\kappa}_2(\gamma) \cdot |a - \theta|} \cdot g(a, \theta) \\ g(\gamma \cdot a, \gamma \cdot \theta) = e^{-\widetilde{\kappa}_5(\gamma)} \cdot e^{-\widetilde{\kappa}_2(\gamma) \cdot |a - \theta|} \cdot g(a, \theta) \end{cases} \quad (124)$$

Here $\widehat{\kappa}_5$ and $\widehat{\kappa}_2$ are some continuous function of γ analogous to $\widetilde{\kappa}_5$ and $\widetilde{\kappa}_2$. By virtue of lemma 5.2, we can conclude the proof. \square

Remark 5.4. A close inspection of the proofs of lemma 5.2 and theorem 5.3 reveals that it is not enough to only assume one of the properties in 5.1. That is, if one only assumes either shift or scale invariance, then there is a much wider class of symmetric two-sided Markov families. To illustrate this, we recover the specific form of the generating densities which give rise to scale, respectively shift, invariance.

Scale invariance: Let $\{\Lambda^n\}_{n \geq 1}$ be a symmetric two-sided Markov family as introduced in the statement of theorem 5.3 which only satisfies scale invariance. The proof of the theorem then shows that g satisfies (123) i.e.,

$$g(\gamma \cdot a, \gamma \cdot \theta) = e^{-\widetilde{\kappa}_5(\gamma)} \cdot e^{-\widetilde{\kappa}_2(\gamma) \cdot |a - \theta|} \cdot g(a, \theta) \quad (125)$$

for some continuous functions $\widetilde{\kappa}_5$ and $\widetilde{\kappa}_2$. The proof of lemma 5.2 gives candidates for the form of these two functions. Indeed, (99) gives $\exp\{-\widetilde{\kappa}_5(\gamma)\} = \gamma^s$ and $\widetilde{\kappa}_2(\gamma) = u \cdot (\gamma - 1)$ for some $s, u \in \mathbb{R}$. Now we define $\varphi(x) := g(1, x) = g(x, 1)$, for $x \in (0, 1]$ and find

$$g(a, \theta) = e^{-\widetilde{\kappa}_5(a \vee \theta)} \cdot e^{-\widetilde{\kappa}_2(a \vee \theta) \cdot |1 - \frac{a \wedge \theta}{a \vee \theta}|} \cdot g\left(\frac{a}{a \vee \theta}, \frac{\theta}{a \vee \theta}\right) = (a \vee \theta)^s \cdot e^{u(a \vee \theta - 1) \cdot |1 - \frac{a \wedge \theta}{a \vee \theta}|} \cdot \varphi\left(\frac{a \wedge \theta}{a \vee \theta}\right) \quad (126)$$

This yields the following generating density

$$\Lambda_{a,b}^1(\theta) = f(a, b) \cdot \theta^s \cdot e^{u(\frac{a \wedge b}{\theta} + \frac{\theta}{a \vee b})} \cdot \varphi\left(\frac{a \wedge b}{\theta}\right) \varphi\left(\frac{\theta}{a \vee b}\right) \cdot \mathbb{1}(\theta \in I_{a,b}). \quad (127)$$

Here, $f(a, b)$ is the normalization of $\Lambda_{a,b}^1$. It is straightforward to check that this density is indeed scale invariant.

Shift invariance: Let $\{\Lambda^n\}_{n \geq 1}$ be a symmetric two-sided Markov family as introduced in the statement of theorem 5.3 which only satisfies shift invariance. As was mentioned in the proof of theorem 5.3, g also obeys a shift property of a form similar to(123). That is,

$$g(a + \gamma, \theta + \gamma) = e^{-\widehat{\kappa}_5(\gamma)} \cdot e^{-\widehat{\kappa}_2(\gamma) \cdot |a - \theta|} \cdot g(a, \theta). \quad (128)$$

(103) in the proof of lemma 5.2 yields that $\widehat{\kappa}_5(\gamma) = -v \cdot \gamma$ and $\widehat{\kappa}_2(\gamma) = -w \cdot \gamma$, where $v, w \in \mathbb{R}$. Define $\varphi(x) := g(0, x) = g(x, 0)$ for $x \in \mathbb{R}^+$, then

$$g(a, \theta) = e^{-\widehat{\kappa}_5(a \wedge \theta)} \cdot e^{-\widehat{\kappa}_2(a \wedge \theta)} g(a - a \wedge \theta, \theta - a \wedge \theta) = e^{v \cdot a \wedge \theta} \cdot e^{w \cdot a \wedge \theta \cdot |a - \theta|} \cdot \varphi(|\theta - a|). \quad (129)$$

This gives the following generating density

$$\Lambda_{a,b}^1(\theta) = f(a, b) \cdot e^{v\theta + w(a+b)\theta - w\theta^2} \cdot \varphi(|\theta - a|) \varphi(|\theta - b|). \quad (130)$$

A short calculation reveals that Λ^1 is indeed shift invariant.

5.1 Densities depending on distances

In this subsection we discuss a special case of shift and scale invariant densities where the factor $g_1(b, x)$ is of the form $g_1(b, x) = \Phi(|b - x|)$ for some differentiable function $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$. Notice that one automatically has shift invariance for densities of this form, whereas the scale invariance imposes an extra restriction on Φ . We provide then a simpler proof to show that such $\{\Lambda^n\}_{n \geq 1}$ correspond to the Dirichlet processes of example 4.10. We start with the a lemma which is a direct consequence of theorem 4.5.

Lemma 5.5. *Let $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be integrable. Suppose Λ^1 is of the form*

$$\Lambda_{a,b}^1(x) = f_1(a, b) \cdot \Phi(|b - x|) \cdot \Phi(|x - a|) \quad (131)$$

with

$$f_1(a, b)^{-1} = \int_{a \wedge b}^{a \vee b} \Phi(|b - x|) \cdot \Phi(|x - a|) dx \quad (132)$$

for all $a, b \in \mathbb{R}^+$. Then there exists a symmetric, two-sided Markov family $\{\tilde{\Lambda}^n\}_{n \geq 1}$ with $\tilde{\Lambda}^1 = \Lambda^1$.

Proof. Clearly $\Lambda^1(a, b)$ is symmetric and factorizable for f_1 , g_1 and h_1 , where the latter two are given by

$$g_1(x, y) = h_1(x, y) = \Phi(|x - y|) \quad (133)$$

and f_1 is as defined in the statement of the lemma. Theorem 4.5 yields the existence of $\{\tilde{\Lambda}^n\}_{n \geq 1}$. \square

Theorem 5.6. *Let $\{\Lambda^n\}_{n \geq 1}$ be a symmetric, two-sided Markov family where Λ^1 is of the form*

$$\Lambda_{a,b}^1(x) = f_1(a, b) \cdot \Phi(|b - x|) \cdot \Phi(|x - a|) \quad (134)$$

with

$$f_1(a, b)^{-1} = \int_{a \wedge b}^{a \vee b} \Phi(|b - x|) \cdot \Phi(|x - a|) dx \quad (135)$$

for some differentiable function $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$. Moreover, assume that $\{\Lambda^n\}_{n \geq 1}$ is scale-invariant. Then, for some $s, t \in \mathbb{R}^+$

$$\Lambda_{a,b}^1(x) = f_1(a, b) \cdot (t \cdot |b - x|^s) \cdot (t \cdot |x - a|^s). \quad (136)$$

Proof. We start from the following observation, for $a \leq \theta_1, \theta_2 \leq b$,

$$\frac{\Phi(\theta_1 - a) \cdot \Phi(b - \theta_1)}{\Phi(\theta_2 - a) \cdot \Phi(b - \theta_2)} = \frac{\Lambda_{a,b}^1(\theta_1)}{\Lambda_{a,b}^1(\theta_2)} = \frac{\Lambda_{\gamma \cdot a, \gamma \cdot b}^1(\gamma \cdot \theta_1)}{\Lambda_{\gamma \cdot a, \gamma \cdot b}^1(\gamma \cdot \theta_2)} = \frac{\Phi(\gamma \cdot (\theta_1 - a)) \cdot \Phi(\gamma \cdot (b - \theta_1))}{\Phi(\gamma \cdot (\theta_2 - a)) \cdot \Phi(\gamma \cdot (b - \theta_2))}. \quad (137)$$

For the second equality we use the scale invariance. We bring the θ_1 and θ_2 dependent factors to the left- and right hand side respectively. This yields

$$\frac{\Phi(\theta_1 - a) \cdot \Phi(b - \theta_1)}{\Phi(\gamma \cdot (\theta_1 - a)) \cdot \Phi(\gamma \cdot (b - \theta_1))} = \frac{\Phi(\theta_2 - a) \cdot \Phi(b - \theta_2)}{\Phi(\gamma \cdot (\theta_2 - a)) \cdot \Phi(\gamma \cdot (b - \theta_2))} = c_{a,b}(\gamma). \quad (138)$$

Here $c_{a,b}(\gamma)$ is a constant which depends only on a, b and γ . We introduce a new function $\tilde{\Phi} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ as $\tilde{\Phi}(x) = \log(\Phi(x)/\Phi(\gamma \cdot x))$. The equation above can be rewritten as

$$\tilde{\Phi}(\theta - a) + \tilde{\Phi}(b - \theta) = \log(c_{a,b}(\gamma)). \quad (139)$$

Differentiation with respect to θ yields

$$\tilde{\Phi}'(\theta - a) - \tilde{\Phi}'(b - \theta) = 0. \quad (140)$$

Given arbitrary $x, y \in \mathbb{R}^+$ we can find a, b and $\theta \in \mathbb{R}^+$ such that $a \leq \theta \leq b$ and $x = \theta - a$, $y = b - \theta$. We find that $\tilde{\Phi}'(x) = \tilde{\Phi}'(y)$. Since x and y are arbitrary, this means that $\tilde{\Phi}'$ is constant and hence that $\tilde{\Phi}$ is of the form

$$\tilde{\Phi}(x) = \alpha(\gamma) \cdot x + \beta(\gamma) \quad (141)$$

which gives

$$\Phi(\gamma \cdot x) = e^{-\alpha(\gamma) \cdot x - \beta(\gamma)} \Phi(x). \quad (142)$$

In what follows we absorb the minuses in $\alpha(\gamma)$ and $\beta(\gamma)$. Next we differentiate with respect to γ , this gives

$$[\alpha'(\gamma) \cdot x + \beta'(\gamma)] \cdot \Phi(\gamma \cdot x) = x \cdot \Phi'(\gamma \cdot x) \quad (143)$$

or, substituting x/γ for x ,

$$[\alpha'(\gamma) \cdot x + \gamma \cdot \beta'(\gamma)] \cdot \Phi(x) = x \cdot \Phi'(x). \quad (144)$$

The expression above is an ODE which has solution

$$\Phi(x) = k \cdot e^{\alpha'(\gamma) \cdot x} \cdot x^{\gamma \cdot \beta'(\gamma)}, \quad (145)$$

with $k \in \mathbb{R}$ an arbitrary constant. Since Φ can not depend on γ , we must have that $\alpha'(\gamma) = \alpha$ and $\gamma \cdot \beta'(\gamma)$ are constants. Hence we must have the following form for Φ ,

$$\Phi(x) = k \cdot e^{\alpha \cdot x} \cdot x^\beta. \quad (146)$$

It is straightforward to check that for a function Φ of this form the whole generated family satisfies the homogeneity condition. One should notice that we can set $\alpha = 0$ because

$$\Lambda_{a,b}^1(x) = f_1(a, b) \cdot \Phi(|x - a|) \cdot \Phi(|b - x|) = f_1(a, b) \cdot k^2 \cdot |x - a|^\beta \cdot |b - x|^\beta \quad (147)$$

□

References

- [1] Barry C Arnold, Narayanaswamy Balakrishnan, and Haikady Navada Nagaraja. *A first course in order statistics*. SIAM, 2008.
- [2] Lorenzo Bertini, Alberto De Sole, Davide Gabrielli, Giovanni Jona-Lasinio, and C233569507120473 Landim. Stochastic interacting particle systems out of equilibrium. *Journal of Statistical Mechanics: Theory and Experiment*, 2007(07):P07014, 2007.
- [3] Thierry Bodineau and Bernard Derrida. Current fluctuations in nonequilibrium diffusive systems: an additivity principle. *Physical review letters*, 92(18):180601, 2004.
- [4] Gioia Carinci, Chiara Franceschini, Rouven Frassek, Cristian Giardinà, and Frank Redig. Large deviations and additivity principle for the open harmonic process, 2023.
- [5] Gioia Carinci, Chiara Franceschini, Davide Gabrielli, Cristian Giardinà, and Dimitrios Tsagkarogiannis. Solvable stationary non equilibrium states. *Journal of Statistical Physics*, 191(1):10, 2024.
- [6] P Clifford and JM Hammersley. Markov fields on finite graphs and lattices. 1971.
- [7] Anna De Masi, Pablo A Ferrari, and Davide Gabrielli. Hidden temperature in the kmp model. *arXiv preprint arXiv:2310.01672*, 2023.
- [8] Bernard Derrida. An exactly soluble non-equilibrium system: the asymmetric simple exclusion process. *Physics Reports*, 301(1-3):65–83, 1998.
- [9] Bernard Derrida, JL Lebowitz, and ER Speer. Free energy functional for nonequilibrium systems: an exactly solvable case. *Physical Review Letters*, 87(15):150601, 2001.
- [10] Simone Floreani and Adrián González Casanova. Non-equilibrium steady state of the symmetric exclusion process with reservoirs. *arXiv preprint arXiv:2307.02481*, 2023.
- [11] Rouven Frassek and Cristian Giardinà. Exact solution of an integrable non-equilibrium particle system. *Journal of Mathematical Physics*, 63(10), 2022.
- [12] Hans-Otto Georgii. *Gibbs measures and phase transitions*. Walter de Gruyter GmbH & Co. KG, Berlin, 2011.
- [13] Cristian Giardinà, Frank Redig, and Berend van Tol. Intertwining and propagation of mixtures for generalized kmp models and harmonic models. *arXiv preprint arXiv:2406.01160*, 2024.
- [14] Claude Kipnis, Carlo Marchioro, and Errico Presutti. Heat flow in an exactly solvable model. *Journal of Statistical Physics*, 27:65–74, 1982.
- [15] Tomoya Yamada and Tadashi Matsunawa. Quantitative approximation to the ordered dirichlet distribution under varying basic probability spaces. *Annals of the Institute of Statistical Mathematics*, 52:197–214, 2000.