

Whittaker spaces for reducible unitary principal series representations of $\widetilde{SL}_2(F)$

Dani Szpruch

November 19, 2024

Abstract

Let F be a p -adic field containing the full group of n^{th} roots of 1 and let \widetilde{G} be the n -fold cover of $SL_2(F)$ constructed by Kubota. In this paper we compute the dimension of the space of Whittaker functionals of the two irreducible summands inside a reducible unitary genuine principal series representation of \widetilde{G} . We also show how these dimensions change when the Whittaker character is modified. As an application we determine the action of the twisted Kazhdan-Patterson n -fold cover of $GL_2(F)$ on the two summands. We emphasize that our main results addresses both ramified and unramified representations and do not rely on the assumption that the cover is tame.

1 Introduction

Let G be a quasi-split reductive group defined over a p -field F . Fix a Borel subgroup $B = H \ltimes N$ and a non-degenerate character ψ of N . Let M be a Levi subgroup of G containing H and let ψ' be the restriction of ψ to $N \cap M$.

Let σ be a smooth admissible irreducible representation of M and let $Wh_{\psi'}(\sigma)$ be the space of ψ' -Whittaker functionals on σ . σ is called ψ' -generic if $Wh_{\psi'}(\sigma)$ is non-trivial. The well known uniqueness of Whittaker model states that if σ is ψ' -generic then $\dim Wh_{\psi'}(\sigma) = 1$. Let $\pi = I(\sigma)$ be the representation of G parabolically induced from σ . Rodier Heredity implies that if σ is ψ' -generic then $\dim Wh_{\psi}(\pi) = 1$. In particular, if π is reducible then exactly one of its irreducible constituents is ψ -generic.

This situation is changed dramatically when we move to covering groups. For these groups, uniqueness of Whittaker model rarely holds, see [5]. In recent years the problem of determining the space of Whittaker functionals of various representations of covering groups has been studied extensively. See for example [1], [3], [4] [7], [20].

If the uniqueness of Whittaker model fails for a certain covering group, it fails already for its genuine principal series representations. The goal of this paper is to determine the dimensions of the space of Whittaker functionals for the irreducible constituents of reducible unitary genuine principal series representations of $\widetilde{G} = \widetilde{SL}_2(F)$, the n -fold cover of $SL_2(F)$ defined in [11]. We emphasize that our results apply to both ramified and unramified representations and to any Whittaker character. We also emphasize that we do not rely on the assumption that the cover is tame. Namely, we do not assume that $\gcd(n, p) = 1$.

Such reducible representations of \tilde{G} exists only if n is odd, see [17], Proposition 5.5 and Section 5.7. Similar to the linear case, the set of reducible unitary genuine principal series representations of \tilde{G} is parameterized by the set of non-trivial quadratic characters of F^* , see Section 3.1 below.

Let π be a reducible genuine unitary principal series representation of \tilde{G} . It is a sum of two non-isomorphic irreducible representations. Since $\dim Wh_\psi(\pi)$ is odd we may write $\pi = \pi_\psi^+ \oplus \pi_\psi^-$ where the irreducible summands π_ψ^\pm are defined by the relation

$$\dim Wh_\psi(\pi_\psi^+) > \dim Wh_\psi(\pi_\psi^-).$$

Our main result given in Section 3.4, Theorem 1, states that

$$\dim Wh_\psi(\pi_\psi^+) = \|n\|^{-\frac{1}{2}} \frac{n+1}{2}, \quad \dim Wh_\psi(\pi_\psi^-) = \|n\|^{-\frac{1}{2}} \frac{n-1}{2}.$$

Here $\|\cdot\|$ is the usual p -adic absolute value. Moreover, if one changes the Whittaker character to ψ_c where $c \in F^*$ (see the body of the paper for details) then

$$\dim Wh_{\psi_c}(\pi_\psi^+) = \|n\|^{-\frac{1}{2}} \frac{n+\theta(c)}{2}, \quad \dim Wh_{\psi_c}(\pi_\psi^-) = \|n\|^{-\frac{1}{2}} \frac{n-\theta(c)}{2}.$$

Here θ is the non-trivial quadratic character of F^* associated with π .

We now give an outline of the proof of Theorem 1: first we normalize the standard intertwining operator by a suitable γ -factor so that the normalized operator is a \tilde{G} isomorphism of order 2. We then use the normalized operator to construct the projections from $Wh_\psi(\pi)$ to $Wh_\psi(\pi_\psi^\pm)$. The last step in the proof is a computation of the rank of these projections. The computation ultimately boils down to a computation of the trace of a certain scattering matrix. A by-product of our proof is that π_ψ^\pm are identified as the ± 1 eigenspaces of the normalized intertwining operator.

Our results in the unramified case are among Gao's results in [3]. The normalization factor in that work of Gao is the one coming from the Gindikin-Karpelevich formula and his computation of the trace uses the explicit description of the scattering matrices obtained by McNamara in [14]. Both Gindikin-Karpelevich formula and McNamara's work are applicable for unramified representations only. In fact McNamara assumes that the Whittaker character is normalized.

Our first novelty is the usage the of the explicit computation of the Plancherel measure in [17], Theorem 5.7 to obtain the normalization factor, see Section 3.1 below. In the unramified case it agrees with Gao's. Our second novelty is the usage of the coarse description of the local coefficients matrices in [17], Theorem 4.12 to compute the trace. Using the formalism from [17], this last computation turns out to be very simple, in fact it is nothing but Fourier inversion for finite abelian groups, see Section 3.3.

This paper is organized as follows. First, in Section 2 we introduce some known notions and results regarding the p -adic field F and the covering group \tilde{G} along with some material on its principal series representations. Our main result is presented in Section 3. We have included two remarks in the end of Section 3.4: in Remark 3 we discuss the analogy to the linear case and the relation to the theory of R -groups. Remark 4 addresses the relation between the normalization factor and the Shahidi-type invariant defined in [5].

In Section 4 we use our main result to determine the action of the Kazhdan-Patterson twisted n -fold cover of $GL_2(F)$ on π_ψ^\pm . Finally, in Section 5 we use this action along with our main result to study the unramified case.

2 Preliminaries

2.1 The p -adic field

Let F be a finite extension of \mathbb{Q}_p . Denote by q the cardinality of its residue field. Denote by \mathbb{O}_F its ring of integers. Fix ϖ , a generator of \mathbb{P}_F , the maximal ideal of \mathbb{O}_F . We normalize the absolute value on F such that $\|\varpi\| = q^{-1}$.

Let ψ be a non-trivial character of F . We define $e(\psi)$, the conductor of ψ , to be the smallest integer k such that ψ is trivial on \mathbb{P}_F^k . For $c \in F^*$ let ψ_c be the character of F given by $x \mapsto \psi(cx)$. The map $\psi \mapsto \psi_c$ is an isomorphism from F to its Pontryagin dual \widehat{F} . We denote by d_ψ the ψ -self dual Haar measure on F .

Let χ be a character of F^* . If χ is ramified we define its conductor, $e(\chi)$, to be the smallest $k \in \mathbb{N}$ such that χ is trivial on $1 + \mathbb{P}_F^k$. If χ is unramified we set $e(\chi) = 0$. For $s \in \mathbb{C}$ we set $\chi_s = \|\cdot\|^s \chi$. In this paper θ will denote a non-trivial quadratic character of F^* .

Tate γ -factor associated with χ , ψ and $s \in \mathbb{C}$ is defined by

$$\gamma(s, \chi, \psi) = \epsilon(s, \chi, \psi) \frac{L(1-s, \chi^{-1})}{L(s, \chi)}$$

where $\epsilon(s, \chi, \psi)$ is a monomial function in q^{-s} satisfying

$$\epsilon(1-s, \chi^{-1}, \psi) = \chi(-1) \epsilon(s, \chi, \psi)^{-1}, \quad (1)$$

$$\epsilon(s, \chi, \psi_c) = \chi(c) \|c\|^{s-\frac{1}{2}} \epsilon(s, \chi, \psi), \quad (2)$$

$$\epsilon(s+t, \chi, \psi) = q^{e(\psi)-e(\chi)t} \epsilon(s, \chi, \psi), \quad (3)$$

We note that if $e(\psi) = 0$ and χ is unramified then $\epsilon(s, \chi, \psi) = 1$. For all these assertions see Section 1 in [15] for example. Combining (1) and (3) one obtains

$$\epsilon(1-s, \chi^{-1}, \psi) \epsilon(1+s, \chi, \psi) = \chi(-1) q^{e(\psi)-e(\chi)}. \quad (4)$$

Fix an odd integer $n \in \mathbb{N}$. We shall assume in this paper that F^* contains the full group of n^{th} roots of 1. Denote this cyclic group by μ_n . We identify μ_n with the group of n^{th} roots of 1 in \mathbb{C}^* and suppress this identification. Let

$$(\cdot, \cdot) : F^* \times F^* \rightarrow \mu_n$$

be the n^{th} power Hilbert symbol. Recall that it is an anti-symmetric bilinear form and that its kernel in each argument is F^{*n} . Hence, it gives rise to a non-degenerate bilinear form on $F^*/F^{*n} \times F^*/F^{*n}$. In particular, it identifies F^*/F^{*n} with $\widehat{F^*/F^{*n}}$. The group $\widehat{F^*/F^{*n}}$ may also be identified with the group of characters of F^* whose order divide n . For $x \in F^*$ let η_x be the character of F^* defined by

$$y \mapsto \eta_x(y) = (x, y).$$

When convenient we shall also think of η_x as an element in $\widehat{F^*/F^{*n}}$.

A subgroup \overline{J} and of F^*/F^{*n} is called a maximal isotropic subgroup if

$$\overline{J} = \bigcap_{x \in \overline{J}} \ker(\eta_x).$$

By [17], Lemma 2.3, there exist two maximal isotropic subgroups \overline{J} and \overline{K} of F^*/F^{*n} such that

1. $\overline{J} \times \overline{K} = F^*/F^{*n}$.
2. The map $k \mapsto \eta_k|_{\overline{J}}$ is an isomorphism from \overline{K} to the Pontryagin dual of \overline{J} .

Remark 1. In the case where $\gcd(n, p) = 1$, the standard choice in the literature is

$$(\overline{J}, \overline{K}) = (\mathbb{O}_F^* F^{*n} / F^{*n}, \langle \varpi \rangle F^{*n} / F^{*n}).$$

We shall not use this.

Fix $k \in \overline{K}$. The following partial γ -factor was defined in [17], Page 127,

$$\gamma_J(s, \chi, \psi, k) = (\#\overline{J})^{-1} \sum_{j \in \overline{J}} \gamma(s, \chi \eta_j, \psi) \eta_k(j).$$

See [17], Theorem 2.12 for the motivation for this definition. By Fourier inversion for finite abelian groups we have

$$\sum_{k \in \overline{K}} \gamma_J(s, \chi, \psi, k) = \gamma(s, \chi, \psi). \quad (5)$$

2.2 The covering of $SL_2(F)$

Most of the material in this section and in Sections 2.3, 2.4 and 2.5 is well known. Full details and references can be found in [17].

Let $G = SL_2(F)$ be the group of two by two matrices with entries in F whose determinants equal 1. Let $N \simeq F$ be the group of upper triangular unipotent matrices in G and let $H \simeq F^*$ be the group of diagonal elements inside G . Denote $B = H \rtimes N$. For $x \in F$, and $a \in F^*$ we shall write

$$u(x) = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, \quad h(a) = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}.$$

The map $u(x) \mapsto \psi(x)$ defines a character of N we shall continue to denote it by ψ .

Let $\widetilde{G}^{(n)} = \widetilde{SL_2(F)^n}$ be the topological central extension of $SL_2(F)$ by μ_n constructed by Kubota in [11]. We have the short exact sequence

$$1 \rightarrow \mu_n \rightarrow \widetilde{G} \rightarrow G \rightarrow 1.$$

We shall denote by \widetilde{A} the inverse image in \widetilde{G} of a subgroup A of G . A (complex) representation of \widetilde{G} or any of its subgroups containing μ_n is called genuine if the central subgroup μ_n acts by the previously fixed injective character $\mu_n \hookrightarrow \mathbb{C}^*$.

2.3 The metaplectic torus

If $n > 2$, \tilde{H} is not abelian. Rather it is a 2-step nilpotent group whose center has a finite index. We note that $Z(\tilde{H})$, the center of \tilde{H} , is \tilde{C} where

$$C = \{h(a) \mid a \in F^{*n}\}.$$

By the Stone-Von Neumann Theorem, see Theorem 3.1 in [19] for example, the isomorphism class of a genuine smooth irreducible representation σ of \tilde{H} is determined by its central character χ_σ . Moreover, a realization of σ is given by inducing from χ'_σ , a character of a maximal abelian subgroup of \tilde{H} which extends χ_σ . The set of maximal abelian subgroups of \tilde{H} is in bijection with the set of maximal isotropic subgroups of F^*/F^{*n} . Let J be the pull-back of \tilde{J} to F^* . Set

$$M = \{h(a) \mid a \in J\}.$$

Then \tilde{M} is a maximal abelian subgroup of \tilde{H} and

$$[\tilde{H} : \tilde{M}] = [J : F^{*n}] = \sqrt{[F^* : F^{*n}]} = n\|n\|^{-\frac{1}{2}}$$

See Section 3 in [17]. This implies that $\dim(\sigma) = n\|n\|^{-\frac{1}{2}}$.

Since n is odd $Z(\tilde{H})$ is canonically isomorphic to $C \times \mu_n$ and \tilde{M} is canonically isomorphic to $M \times \mu_n$. This implies that set of genuine characters of $Z(\tilde{H})$ is canonically parameterized by the set of characters of F^{*n} and that the set of characters of \tilde{M} is canonically parameterized by the set of characters of J , see Section 4.2 in [17].

The set of characters of F^{*n} is naturally identified with the set of n^{th} powers of characters of F^* . Thus, given a genuine smooth irreducible representation σ of \tilde{H} we may associate to it a unique character χ^n where χ is a character of F^* . We may think of χ^n as the restriction of χ to F^{*n} . Note that σ is unitary if and only if χ is unitary. We shall identify the restriction of χ to J with a genuine character of \tilde{M} and we continue to denote this genuine character by χ .

For $t \in \tilde{H}$, an inverse image of $h(a)$ we set $\|t\| = \|a\|$. We now define σ_s to be the representation of \tilde{H} given by $t \mapsto \|t\|^s \sigma(t)$. If χ^n is the character of F^* associated with σ then $(\chi_s)^n$ is the character of F^* associated with σ_s .

Fix w , an inverse image of $w_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ in \tilde{G} . Since w normalizes \tilde{H} and μ_n is central in \tilde{G} , the automorphism of \tilde{H} defined by $t \mapsto wtw^{-1}$ is independent of the choice of an inverse image of w_0 . We define σ^w to be a representation of \tilde{H} by setting

$$t \mapsto \sigma^w(t) = \sigma(wtw^{-1}).$$

We note that $(\sigma^w)^w = \sigma$, $(\sigma_s)^w = (\sigma^w)_{-s}$ and that if χ^n is associated σ then χ^{-n} is associated with σ^w .

2.4 Genuine Principal series representations

\tilde{G} splits uniquely over N . We shall identify N with its image in \tilde{G} . Moreover, $\tilde{B} = \tilde{H} \ltimes N$. Therefore, as in the linear case, any representation (σ, V_σ) of \tilde{H} can be extended to a

representation of \tilde{B} by defining it to be trivial on N . We shall not distinguish between representations of \tilde{H} and those of \tilde{B} . We consider the normalized parabolic induction

$$I(\sigma, s) = \text{Ind}_{\tilde{B}}^{\tilde{G}} \sigma_{s+1}$$

and we set $I(\sigma) = I(\sigma, 0)$.

Fix $f \in I(\sigma)$ and $s \in \mathbb{C}$. Given $g \in \tilde{G}$ we pick $t \in \tilde{H}$, $n \in N$ and k an inverse image inside \tilde{G} of an element of $SL_2(\mathbb{O}_F)$ such that $g = tnk$ and we set

$$f_s(g) = \|t\|^s f(g). \quad (6)$$

As in the linear case one verifies, using the Iwasawa decomposition of G , that f_s is a well defined element in $I(\sigma, s)$ and that $f \mapsto f_s$ is an isomorphism of vector spaces. Similar to the linear case we have the following:

Lemma 1. *For $f \in I(\sigma)$ and $g \in \tilde{G}$ the integral*

$$\int_F f_s(wu(x)g) d_\psi x$$

converges absolutely to a V_σ valued rational function in q^{-s} provided that $\|\chi^n(\varpi)\| < q^{\text{Re}(s)n}$. Here χ^n is the character of F^ associated with σ . We shall denote its meromorphic continuation by $(A(\sigma, s, \psi)(f_s))(g)$. Away from its poles,*

$$A(\sigma, s, \psi) \in \text{Hom}_{\tilde{G}}(I(\sigma, s), I(\sigma^w, -s)).$$

For the proof see Section 7 of [13] for example.

We note that $A(\sigma, s, \psi)$ depends on ψ only via the choice of the Haar-measure and that since $d_{\psi_c} = \|c\|^{\frac{1}{2}} d_\psi$ we have

$$A(\sigma, s, \psi_c) = \|c\|^{\frac{1}{2}} A(\sigma, s, \psi). \quad (7)$$

While this dependence is typically suppressed in the literature we chose to indicate it in this paper.

2.5 The space of Whittaker functionals

A ψ -Whittaker functional on a representation (π, V_π) of \tilde{G} is a functional on V_π satisfying

$$\xi(\pi(u)v) = \psi(u)\xi(v)$$

for all $v \in V_\pi$, $u \in N$. Denote by $Wh_\psi(\pi) = \text{Hom}_N(\pi, \psi)$ the space of ψ -Whittaker functionals on π . If τ is another representation of \tilde{G} then any $A \in \text{Hom}_{\tilde{G}}(\pi, \tau)$ gives rise to a linear map

$$A^\psi : Wh_\psi(\tau) \rightarrow Wh_\psi(\pi)$$

defined by $\xi \mapsto \xi \circ A$. We note that if $\pi = \tau$ and A is the identity map then A^ψ is also the identity map.

As noted in the introduction, unlike the linear case, the space of Whittaker functionals on smooth admissible irreducible genuine representations of \widetilde{G} may be greater than 1. By Rodier Heredity, regardless of the question of reducibility, we have

$$\dim Wh_\psi(I(\sigma)) = n\|n\|^{-\frac{1}{2}},$$

see Lemma 3.5 in [17] for example.

3 Main result

3.1 Plancherel measure

Let σ be a smooth genuine irreducible representation of \widetilde{H} . There exists a rational function in q^{-ns} , $\mu_\psi(\sigma^w, s)$, such that

$$A(\sigma^w, -s, \psi) \circ A(\sigma, s, \psi) = \chi(-1)\mu_\psi^{-1}(\sigma, s)Id. \quad (8)$$

(here Id is the identity map). See [17], Section 5.1 for exact details. By Theorem 5.1 of [8] if $gcd(p, n) = 1$ we have

$$\mu_\psi(\sigma, s) = \chi^n(-1)\gamma(1 + ns, \chi^n, \psi_n)\gamma(1 - ns, \chi^{-n}, \psi_n) \quad (9)$$

Here χ^n is the character of F^* associated with σ (one needs to use (4)). By Theorem 5.7 of [17], if χ^n is unramified then (9) holds also for the case where $gcd(p, n) > 1$ (note that $q^{e(\psi_n)} = \|n\|q^{e(\psi)}$).

The Knapp-Stein dimension theorem for covering groups proven in [12] states that given that σ is unitary, $I(\sigma)$ is reducible if and only if $\sigma \simeq \sigma^w$ and $\mu_n^{-1}(\sigma, s)$ is analytic at $s = 0$. By studying the analytical properties of the Plancherel measure it was proven in Proposition 5.5 of [17] that if σ is unitary then $I(\sigma)$ is reducible if and only if χ^n is a non-trivial quadratic character of F^*/F^{*n} . Since n is odd, any non-trivial quadratic character of F^*/F^{*n} may be extended to a unique non-trivial quadratic character of F^* .

For θ , a non-trivial quadratic character of F^* , let $\sigma = \sigma_\theta$ be (the isomorphism class of) the genuine unitary representation of \widetilde{H} whose associated character is $\theta^n = \theta$. By the above, as in the linear case, the map $\theta \mapsto I(\sigma_\theta)$ is a bijection from the set of non-trivial quadratic characters of F^* to the set of reducible genuine principal series representations of \widetilde{G} induced from a unitary data.

Proposition 1. *Equation (9) holds for the case where $\chi^n = \chi = \theta$ is a non-trivial quadratic character of F^* .*

Proof. We only need to consider the case where $gcd(n, p) > 1$ and θ is ramified. Note that in this case $p \neq 2$ as n is odd. Consequently, $e(\theta) = 1$ (see Lemma 4.15 in [17] for example). Since $gcd(2, n) = 1$, for any ramified element in $\eta \in \widehat{F^*/F^{*n}}$ we have $e(\theta\eta) = e(\eta)$. By Theorem 5.7 in [17] we only need to prove that

$$[F^* : F^{*n}]^{-1} \sum_{\eta \in \widehat{F^*/F^{*n}}} q^{-e(\theta\eta)} = \|n\|q^{-1}.$$

Indeed, the n unramified elements in $\widehat{F^*/F^{*n}}$ contribute nq^{-1} the sum above. Thus,

$$[F^* : F^{*n}]^{-1} \sum_{\eta \in \widehat{F^*/F^{*n}}} q^{-e(\theta\eta)} = n^{-2} \|n\| \left(nq^{-1} + \sum_{\eta \in \widehat{F^*/F^{*n}}, \eta \text{ is ramified}} q^{-e(\eta)} \right).$$

From the identify in last line of Page 156 of [17] it follows that

$$\sum_{\eta \in \widehat{F^*/F^{*n}}, \eta \text{ is ramified}} q^{-e(\eta)} = n(n-1)q^{-1}.$$

The proof is now complete. \square

3.2 Induction by stages

Recall that we may realize σ as $\text{Ind}_{\widetilde{M}}^{\widetilde{H}} \chi'_\sigma$ where χ'_σ is a character of \widetilde{M} whose restriction to $Z(\widetilde{H})$ equals χ_σ . Recall also that if χ^n is the character of F^* associated with σ then we may identify χ'_σ with the restriction of χ to J . Using induction by stages we may realize $I(\sigma, s)$ as

$$I(\chi, s) = \text{Ind}_{\widetilde{NM}}^{\widetilde{G}} \chi_{s+1}.$$

Note that by Stone-Von Neumann theorem, if η is a character of F^* whose restriction to F^{*n} is trivial (equivalently, $\eta \in \widehat{F^*/F^{*n}}$) then $I(\chi, s) \simeq (\chi\eta, s)$. When $s = 0$ we write $I(\chi)$ instead of $I(\chi, 0)$.

Using this realization, away from its poles, the intertwining operator $A(\chi, s, \psi)$ takes the form

$$A(\chi, s, \psi) \in \text{Hom}_{\widetilde{G}}(I(\chi, s), I((\chi^{-1}, -s))).$$

See the commutative diagram in Page 137 of [17] for a proof. That diagram also implies that

$$A(\chi^{-1}, -s, \psi) \circ A(\chi, s, \psi) = \chi(-1) \mu_\psi^{-1}(\sigma, s) \text{Id}.$$

In the special case where $\chi = \theta$ is a non-trivial quadratic character of F^* , $A(\theta, 0, \psi)$ is a self intertwining operator. Moreover, by the theory of R -groups, it is not a scalar operator, see [12]. Using Proposition 1 and Equation (2) we obtain

$$A(\theta, 0, \psi)^2 = \|n\| \gamma(1, \theta, \psi)^2 \text{Id}. \quad (10)$$

3.3 A scattering matrix

In Section 4.2 of [17] we have introduced a basis $\{\lambda_{b, \chi, \psi, s} \mid b \in \overline{K}\}$ for $Wh_\psi(I(\chi, s))$ and a function

$$\tau(\cdot, \cdot, \chi, s, \psi) : \overline{K} \times \overline{K} \rightarrow \mathbb{C}[q^{-s}].$$

defined by the relation

$$A^\psi(\chi, s, \psi)(\lambda_{a^{-1}, \chi^{-1}, \psi, -s}) = \sum_{b \in \overline{K}} \tau(a, b, \chi, s, \psi) \lambda_{b, \chi, \psi, s}. \quad (11)$$

See [17], Proposition 4.9. The appearance of a^{-1} rather than a on the left hand side is a manifestation of the fact that both Whittaker spaces $Wh_\psi(I(\chi, s))$ and $Wh_\psi(I(\chi^{-1}, -s))$ were identified in a canonical way with the space of functionals on σ . This identification gave rise to the local coefficients matrix whose characteristic polynomial is an invariant of σ and ψ , see Section 3.2 of [6].

However, if $\chi = \theta$ is a non-trivial quadratic character of F^* then $A^{\psi_c}(\theta, 0, \psi)$ is an endomorphism on the space $Wh_{\psi_c}(I(\theta, 0))$ and we wish to change our point of view. In this special case it follows from the above that if we define

$$M(\cdot, \cdot, \chi, \psi) : \overline{K} \times \overline{K} \rightarrow \mathbb{C}[q^{-s}]$$

by

$$M(a, b, \theta, \psi) = \tau(a^{-1}, b, \theta, 0, \psi).$$

then

$$A^\psi(\theta, 0, \psi)(\lambda_{a, \theta, \psi, 0}) = \sum_{b \in \overline{K}} M(a, b, \theta, \psi) \lambda_{b, \theta, \psi, 0}. \quad (12)$$

In other words, $M(\cdot, \cdot, \theta, \psi)$ represents the endomorphism $A^\psi(\theta, 0, \psi)$.

Remark 2. *Suppose that $\gcd(n, p) = 1$ and that θ is the unique non-trivial quadratic unramified character of F^* . If one fixes \overline{J} as in Remark 1 then the conjugacy class of $M(\cdot, \cdot, \chi, \psi)$ is the same as the conjugacy class of the scattering matrix computed in [14] as both matrices represents the same map.*

Lemma 2. *Let θ be a non-trivial quadratic character of F^* . We have*

$$M(a, b, \theta, \psi) = \gamma_J(1, \theta \eta_{a^{-1}b}, \psi, (ab)^{-1})$$

Proof. This follows at once from the coarse description of $\tau(a, b, \chi, 0, \psi)$ given in [17], Theorem 4.12. \square

Proposition 2. *Assume that θ is a non-trivial quadratic character of F^* . We have*

$$\text{tr } A^{\psi_c}(\theta, 0, \psi) = \theta(c) \gamma(1, \theta, \psi).$$

Proof. Assume first that $c = 1$. From (12) it follows that

$$\text{Tr } A^\psi(\theta, 0, \psi) = \sum_{a \in \overline{K}} M(a, a, \theta, \psi).$$

Lemma 2 gives

$$\sum_{a \in \overline{K}} M(a, a, \theta, \psi) = \sum_{a \in \overline{K}} \gamma_J(1, \theta, \psi, a^{-2}) = \sum_{k \in \overline{K}} \gamma_J(1, \theta, \psi, k).$$

For the last equality we have used the fact that since \overline{K} is an abelian group of odd order, $a \mapsto a^{-2}$ is an automorphism of \overline{K} . With (5) the proof of for the case where $c = 1$ is completed.

We now move to the general case. From (7) it follows that $\|c\|^{-\frac{1}{2}} M(\cdot, \cdot, \theta, \psi_c)$ represents $A^{\psi_c}(\theta, 0, \psi)$. Thus,

$$\text{Tr } A^{\psi_c}(\theta, 0, \psi) = \|c\|^{-\frac{1}{2}} \sum_{a \in \overline{K}} M(a, a, \theta, \psi_c) = \|c\|^{-\frac{1}{2}} \gamma(1, \theta, \psi_c).$$

With (2) we finish. \square

3.4 Whittaker dimensions

From this point we shall write $\pi = I(\sigma_\theta)$. We define a normalized intertwining operator

$$\mathbb{A}(\theta, \psi) = \|n\|^{-\frac{1}{2}} \gamma(1, \theta, \psi)^{-1} A(\theta, 0, \psi). \quad (13)$$

From (10) it follows that $\mathbb{A}(\theta, \psi)^2 = Id$. Since $\mathbb{A}(\theta, \psi)$ is not a scalar operator we may write $I(\theta)$ as a sum of two irreducible representations of \tilde{G}

$$I(\theta) = \pi_\psi^+ \oplus \pi_\psi^-$$

where π_ψ^\pm is the eigenspace of $\mathbb{A}(\theta, \psi)$ corresponding to ± 1 . We note that since $\mathbb{A}(\theta, \psi)^2$ is the identity map it follows that $\frac{1}{2}(I \pm \mathbb{A}(\theta, \psi))$ are the projections from $I(\theta)$ to π_ψ^\pm .

Theorem 1. *Let θ be a non-trivial quadratic character of F^* . Denote $\pi = I(\sigma_\theta)$. We have*

$$\dim Wh_{\psi_c}(\pi_\psi^+) = \|n\|^{-\frac{1}{2}} \frac{n + \theta(c)}{2}, \quad \dim Wh_{\psi_c}(\pi_\psi^-) = \|n\|^{-\frac{1}{2}} \frac{n - \theta(c)}{2}.$$

In particular,

$$\dim Wh_\psi(\pi_\psi^+) = \|n\|^{-\frac{1}{2}} \frac{n + 1}{2}, \quad \dim Wh_\psi(\pi_\psi^-) = \|n\|^{-\frac{1}{2}} \frac{n - 1}{2}.$$

Proof. Since $\frac{1}{2}(Id \pm \mathbb{A}(\theta, \psi))^{\psi_c}$ are the projections from $Wh_{\psi_c}(I(\theta))$ to $Wh_{\psi_c}(\pi_\psi^\pm)$ we obtain

$$\begin{aligned} \dim Wh_{\psi_c}(\pi_\psi^\pm) &= \text{rank} \frac{1}{2}(Id \pm \mathbb{A}(\theta, \psi))^{\psi_c} = \text{tr} \frac{1}{2}(Id \pm \mathbb{A}(\theta, \psi))^{\psi_c} \\ &= \frac{1}{2}(n \|n\|^{-\frac{1}{2}} \pm \|n\|^{-\frac{1}{2}} \gamma(1, \theta, \psi)^{-1} \text{tr} A^{\psi_c}(\theta, 0, \psi)) \end{aligned}$$

With Proposition 2 we finish. □

Remark 3. *Since $\dim Wh_\psi(\pi)$ is odd we know a-priori that the ψ -Whittaker dimensions of the two irreducible summands inside π are not equal. From the last Theorem it follows that we can define π_ψ^\pm by stating that*

$$\dim Wh_\psi(\pi_\psi^+) > \dim Wh_\psi(\pi_\psi^-). \quad (14)$$

This second characterization of π_ψ^\pm shows that the definition of these sub-representations as eigenspaces is free of the choice of \bar{J} giving rise to the realization $I(\theta)$. The action of the intertwining operator on the two summands inside π is analogous to the formula for the action of the intertwining operator in the case of quasi-split reductive algebraic groups given in [10]. That formula involved the R -groups and had the generic summand as its base point. Here the R -group is the group of two elements and due to (14), π_ψ^+ plays the role of the generic summand.

Remark 4. *Let σ be an irreducible genuine representation of \tilde{H} and let χ^n the character of F^* associated with σ . In Section 3.5.1 of [5] we have defined an invariant $S_w(\sigma, s, \psi)$ using the determinant of the local coefficient matrix and the Plancherel measure and made the*

point that it is an analog of the reciprocal of Shahidi local coefficients defined for quasi-split reductive algebraic groups, [16]. In particular, it follows at once from the definition that

$$S_w(\sigma, s, \psi)S_w(\sigma^w, -s, \psi) = \chi^n(-1)\mu_\psi^{-1}(\sigma, s). \quad (15)$$

Note that Equation (15) is unchanged if we replace $S_w(\sigma, s, \psi)$ by

$$S'_w(\sigma, s, \psi) = \chi^n(n)S_w(\sigma, s, \psi).$$

In the case where $\gcd(n, p) = 1$ and σ is unramified it was shown in Theorem 3.14 of [5] that $S_w(\sigma, s, \psi) = \gamma(1 - ns, \chi^{-n}, \psi_n)$. This last result was recently extended to all σ under the additional assumption that p is odd, see the introduction of [18]. Thus, if p is odd and relatively prime to n then from (2) it follows that the normalization factor $\|n\|^{\frac{1}{2}}\gamma(1, \theta, \psi)$ we have used in (13) for $A(\theta, 0, \psi)$ equals $S'_w(\sigma_\theta, 0, \psi)$. It is expected of course that these factors are equal in the other cases as well: note that from (15), Proposition 1 and Equation (2) it follows that $S'_w(\sigma_\theta, 0, \psi)^2 = \|n\|\gamma(1, \theta, \psi)^2$. Thus, the factors are equal at least up to a sign for all p .

4 The action of $\widetilde{GL}_2(F)$

Let $\widetilde{GL}_2(F)$ be the c -twisted central extension of $GL_2(F)$ by μ_n introduced in [9]. The derived group of $\widetilde{GL}_2(F)$ is \widetilde{G} . Therefore, $\widetilde{GL}_2(F)$ is acting on \widetilde{G} by conjugations. For $h \in \widetilde{G}$ and $g \in \widetilde{G}$ set $h^g = g^{-1}hg$.

Since μ_n is central in $\widetilde{GL}_2(F)$ this action factors through $GL_2(F)$ so for $h \in \widetilde{G}$ and $g \in GL_2(F)$ we can write h^g with no ambiguity. We also have an action of \widetilde{G} on the set of isomorphism classes of representations of \widetilde{G} . For a representation τ of \widetilde{G} and $g \in GL_2(F)$ we define a representation τ^g of \widetilde{G} by $h \mapsto \tau(h^g)$.

The centralizer of \widetilde{G} in $\widetilde{GL}_2(F)$ is \widetilde{S} , the inverse image in $\widetilde{GL}_2(F)$ of the group of scalar matrices in $GL_2(F)$, see Lemma 8.2 in [6]. Thus, the action of $\widetilde{GL}_2(F)$ on the set of isomorphism classes of representations of \widetilde{G} reduces to an action of $\widetilde{GL}_2(F)/\widetilde{G}^+$ where \widetilde{G}^+ is the inverse image in $\widetilde{GL}_2(F)$ of

$$G^+ = SG = \{g \in G \mid \det(g) \in F^{*2}\}.$$

Let $g \in \widetilde{GL}_2(F)$ be an inverse image in $\widetilde{GL}_2(F)$ of $g' \in GL_2(F)$. Set $\det(g) = \det(g')$. The map $g \mapsto \det(g)$ gives rise to an isomorphism from $\widetilde{G}/\widetilde{G}^+$ to F^*/F^{*2} . As a set of representatives of $\widetilde{G}/\widetilde{G}^+$ we can take a set of inverse images of

$$\{\text{diag}(1, c) \mid c \in F^*/F^{*2}\}.$$

We note that the group of inverse images in $\widetilde{GL}_2(F)$ of the diagonal subgroup in $GL_2(F)$ normalizes N . In particular for $u \in N$ we have

$$\psi(u^{\text{diag}(1, c)}) = \psi_c(u). \quad (16)$$

Lemma 3. *Let τ be a genuine principal series representation of \widetilde{G} . For any $g \in \widetilde{GL}_2(F)$, $\tau^g \simeq \tau$.*

Proof. We realize τ as $I(\chi)$, an induction from \widetilde{MN} . Let $g \in \widetilde{GL_2(F)}$ be an inverse image of $g' = \text{diag}(1, c)$ and let $m \in \widetilde{M}$ be an inverse image of $m' \in M$. Since g' commutes with m' there exists a character $\eta : M \rightarrow \mu_n$ such that $m^g = \eta(m')m$. The map $T : I(\chi)^g \rightarrow I(\chi\eta)$ defined by $T(f)(h) = f(h^g)$ is a \widetilde{G} isomorphism. Finally, by Lemma 8.3 in [6], since n is odd, $Z(\widetilde{H}) = Z(\widetilde{T}) \cap \widetilde{G}$. Consequently, the restriction of η to $Z(\widetilde{H})$ is trivial. From the Stone-Von Neumann theorem it follows that $I(\chi\eta) \simeq I(\chi)$. \square

Remark 5. Lemma 3 is false for even fold covers of G since in this case $Z(\widetilde{T}) \cap \widetilde{G} \subsetneq Z(\widetilde{H})$.

Theorem 2. Let θ be a non-trivial quadratic character of F^* . Denote $\pi = I(\sigma_\theta)$. Fix $g \in \widetilde{GL_2(F)}$.

1. If $\theta(\det g) = 1$ then $(\pi_\psi^+)^g \simeq \pi_\psi^+$ and $(\pi_\psi^-)^g \simeq \pi_\psi^-$.
2. If $\theta(\det g) = -1$ then $(\pi_\psi^+)^g \simeq \pi_\psi^-$ and $(\pi_\psi^-)^g \simeq \pi_\psi^+$.

Proof. From Lemma 3 it follows that either g fixes each of the 2 irreducible sub representations of π or it swaps them. Thus, we only need to determine the isomorphism class of $(\pi_\psi^+)^g$.

As before we may assume that g is an inverse image of $\text{diag}(1, c)$. In light of Theorem 1 and Remark 3 it is sufficient to show that $Wh_\psi(\pi^+) = Wh_{\psi_c}((\pi_\psi^+)^g)$. Indeed if $\xi \in Wh_\psi(\pi^+)$ then from (16) we deduce that for $u \in N$ we have

$$\xi \circ \pi^g(u) = \psi_c(u)\xi.$$

This shows that $Wh_\psi(\pi^+) \subseteq Wh_{\psi_c}((\pi_\psi^+)^g)$. The other inclusion is proven in the same way. \square

5 The unramified case

In this section we shall use Theorem 1 and Theorem 2 to prove and slightly extend the results in Section 5.4.1 of [7].

Assume that $\gcd(n, p) = 1$. In this case \widetilde{G} splits over $SL_2(\mathbb{O}_F)$ uniquely. Denote by K_1 the image of this group in \widetilde{G} . Denote also $K_2 = K_1^{\text{diag}(1, \varpi)}$.

Let σ be an irreducible genuine representation of \widetilde{H} . Suppose that σ is an irreducible representation associated with a character χ^n of F^* . $I(\sigma)$ is called unramified with respect to K_i if it has a non-zero fixed K_i vector. As in the linear case, due to the Iwasawa decomposition, if $I(\sigma)$ is unramified then it has a one dimensional K_i fixed subspace. We note that $I(\sigma)$ is unramified with respect to K_1 if and only if χ^n is unramified, see Section 8.6 in [6]. Since $I(\sigma)$ is ramified with respect to K_1 if and only if $I(\sigma)^{\text{diag}(1, \varpi)}$ is ramified with respect to K_2 it follows from Lemma 3 that $I(\sigma)$ is unramified with respect to K_1 if and only if it is unramified with respect to K_2 . We remark here that for even fold covers of G , a genuine principal series representation can not be unramified with respect to both K_1 and K_2 .

Note that F^* has exactly one non-trivial unramified quadratic character, namely $x \mapsto \theta_u(x) = (-1)^{\text{val}(x)}$. Thus, there exists a unique genuine reducible unramified principal

series representation π of \tilde{G} induced from a unitary representation. For $i = 1, 2$ let v_i be the unique (up to normalization) K_i spherical vector and let V_i be the \tilde{G} invariant subspace of π generated by v_i .

If $e(\psi)=0$ then from Proposition 7.3 in [2] we deduce that $\mathbb{A}(\theta, \psi)v_1 = v_1$. Hence if $e(\psi)=0$ then $V_1 = \pi_\psi^+$.

Theorem 3. *Assume that $\gcd(n, p) = 1$. Let π be the unique genuine unitary reducible unramified principal series representation of \tilde{G} . For $i = 1, 2$ let v_i be the unique (up to normalization) K_i unramified vector and let V_i be the \tilde{G} sub-representation of π generated by v_i . We have*

1. *If $e(\psi)$ is even then $V_1 = \pi_\psi^+$ and $V_2 = \pi_\psi^-$.*
2. *If $e(\psi)$ is odd then $V_1 = \pi_\psi^-$ and $V_2 = \pi_\psi^+$.*

Proof. Since we already know that if $e(\psi) = 0$ then $V_1 = \pi_\psi^+$ and since $e(\psi) \equiv e(\psi_c) \pmod{2}$ if and only if $\theta_u(c) = 1$ it follows from Theorem 1 and Remark 3 that if $e(\psi)$ is even then $V_1 = \pi_\psi^+$ and that if $e(\psi)$ is odd then $V_1 = \pi_\psi^-$. To finish the proof we note that $v_2 \in V_1^{diag(1, \varpi)}$ and that since $\theta_u(\varpi) = -1$ it now follows from Theorem 2 that if $e(\psi)$ is even then $V_2 = \pi_\psi^-$ and that if $e(\psi)$ is odd then $V_2 = \pi_\psi^+$. \square

Corollary 1.

$$\mathbb{A}(\theta_u, \psi)v_1 = (-1)^{e(\psi)}v_1, \quad \mathbb{A}(\theta_u, \psi)v_2 = (-1)^{e(\psi)+1}v_2.$$

Acknowledgment. I would like to thank Fan Gao and Nadya Gurevich for numerous enlightening communications.

Funding. This research was supported by the Israel Science Foundation (grant No. 1643/23).

References

- [1] GAO, F. Distinguished theta representations for certain covering groups. *Pacific J. Math.* 290, 2 (2017), 333–379.
- [2] GAO, F. The Langlands-Shahidi L -functions for Brylinski-Deligne extensions. *Amer. J. Math.* 140, 1 (2018), 83–137.
- [3] GAO, F. R-group and whittaker space of some genuine representations. *Journal of the Institute of Mathematics of Jussieu* (2021), 61.
- [4] GAO, F., GUREVICH, N., AND KARASIEWICZ, E. Genuine whittaker functor and the quantum affine schur–weyl duality. *arXiv preprint* (2022).
- [5] GAO, F., SHAHIDI, F., AND SZPRUCH, D. On the local coefficients matrix for coverings of SL_2 . In *Geometry, algebra, number theory, and their information technology applications*, vol. 251 of *Springer Proc. Math. Stat.* Springer, Cham, 2018, pp. 207–244.

- [6] GAO, F., SHAHIDI, F., AND SZPRUCH, D. Local coefficients and gamma factors for principal series of covering groups. *Mem. Amer. Math. Soc.* 283, 1399 (2023), v+135.
- [7] GAO, F., SHAHIDI, F., AND SZPRUCH, D. Restrictions, l-parameters, and local coefficients for genuine representations. *arXiv preprint arXiv:2102.08859* (accepted for publication in *Mémoires de la Société Mathématique de France*, 2023).
- [8] GOLDBERG, D., AND SZPRUCH, D. Plancherel measures for coverings of p -adic $SL_2(F)$. *Int. J. Number Theory* 12, 7 (2016), 1907–1936.
- [9] KAZHDAN, D. A., AND PATTERSON, S. J. Metaplectic forms. *Inst. Hautes Études Sci. Publ. Math.*, 59 (1984), 35–142.
- [10] KEYS, C. D., AND SHAHIDI, F. Artin L -functions and normalization of intertwining operators. *Ann. Sci. École Norm. Sup. (4)* 21, 1 (1988), 67–89.
- [11] KUBOTA, T. *On automorphic functions and the reciprocity law in a number field*. Lectures in Mathematics, Department of Mathematics, Kyoto University, No. 2. Kinokuniya Book-Store Co., Ltd., Tokyo, 1969.
- [12] LUO, C. Knapp-Stein dimension theorem for finite central covering groups. *Pacific J. Math.* 306, 1 (2020), 265–280.
- [13] MCNAMARA, P. J. Principal series representations of metaplectic groups over local fields. In *Multiple Dirichlet series, L-functions and automorphic forms*, vol. 300 of *Progr. Math.* Birkhäuser/Springer, New York, 2012, pp. 299–327.
- [14] MCNAMARA, P. J. The metaplectic Casselman-Shalika formula. *Trans. Amer. Math. Soc.* 368, 4 (2016), 2913–2937.
- [15] SCHMIDT, R. Some remarks on local newforms for $GL(2)$. *J. Ramanujan Math. Soc.* 17, 2 (2002), 115–147.
- [16] SHAHIDI, F. *Eisenstein series and automorphic L-functions*, vol. 58 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2010.
- [17] SZPRUCH, D. On Shahidi local coefficients matrix. *Manuscripta Math.* 159, 1-2 (2019), 117–159.
- [18] SZPRUCH, D. A p -adic analog of hasse–davenport product relation involving ϵ -factors. *Forum Mathematicum* (2024).
- [19] WEISSMAN, M. H. Metaplectic tori over local fields. *Pacific J. Math.* 241, 1 (2009), 169–200.
- [20] ZOU, J. Local metaplectic correspondence and applications. *Math. Z.* 305, 3 (2023), Paper No. 43, 33.