Classification of quadratic forms over finite fields with maximal and minimal Artin-Schreier curves

Ruikai Chen

School of Mathematical Sciences, South China Normal University, Guangzhou 510631, China Email: chen.rk@outlook.com

Abstract. This paper explores quadratic forms over finite fields with associated Artin-Schreier curves. Specifically, we investigate quadratic forms of $\mathbb{F}_{q^n}/\mathbb{F}_q$ represented by polynomials over \mathbb{F}_{q^n} with q odd, characterizing them using certain matrices defined by coefficients of the polynomials. In particular, a comprehensive treatment will be given for those polynomials whose coefficients all lie in \mathbb{F}_q . Afterwards, the results on quadratic forms will be applied to get maximal and minimal Artin-Schreier curves explicitly. Keywords. Artin-Schreier curve, finite field, polynomial, quadratic form

1 Introduction

Given $\mathbb{F}_{q^n}/\mathbb{F}_q$, an extension of a finite field of odd characteristic, and the polynomial ring $\mathbb{F}_{q^n}[x]$, define

$$
\text{Tr}(x) = \sum_{i=0}^{n-1} x^{q^i}.
$$

Let ℓ be a nonzero q-linear polynomial over \mathbb{F}_{q^n} of the form

$$
\ell(x) = \sum_{i=0}^{n-1} a_i x^{q^i}.
$$

Here a q-linear polynomial over \mathbb{F}_{q^n} is also viewed as a linear endomorphism of $\mathbb{F}_{q^n}/\mathbb{F}_q$. Then Tr($x\ell(x)$) as a map from \mathbb{F}_{q^n} to \mathbb{F}_q is a quadratic form of $\mathbb{F}_{q^n}/\mathbb{F}_q$ (in fact, every quadratic form of $\mathbb{F}_{q^n}/\mathbb{F}_q$ can be written in this form), and it induces a homogeneous polynomial over \mathbb{F}_q of degree 2 in *n* indeterminates or a zero polynomial. Quadratic forms over finite fields have attracted significant attention in number theory, with applications in various areas including coding theory, cryptography and combinatorics. Furthermore,

they are directly related to the number of rational points of some Artin-Schreier curves over finite fields. More precisely, consider the projective curve over \mathbb{F}_{q^n} defined by

$$
y^{q} - y - x\ell(x) \in \mathbb{F}_{q^n}[x, y].
$$

If the degree of ℓ is q^m , then the genus of this curve is $\frac{(q-1)q^m}{2}$ (see [\[14,](#page-31-0) Proposition 3.7.10]). For the number N of rational points of the above curve, the well-known Hasse-Weil bound asserts that

$$
q^{n} + 1 - (q - 1)q^{\frac{n}{2} + m} \le N \le q^{n} + 1 + (q - 1)q^{\frac{n}{2} + m}.
$$

The curve is called a maximal curve if N attains the upper bound, and called a minimal curve if N attains the lower bound. We have special interests in such curves for their unique properties.

There has been a lot of research concerning quadratic forms over finite fields and related maximal or minimal Artin-Schreier curves. To name a few, in [\[7\]](#page-30-0) and [\[6\]](#page-30-1), the quadratic forms represented by monomials are classified. Also, one may refer to [\[5\]](#page-30-2) for those curves defined by monomials, and to [\[1\]](#page-30-3) for those curves defined by polynomials with coefficients in prime fields. Some other results can be found in [\[3,](#page-30-4) [11,](#page-31-1) [12,](#page-31-2) [13,](#page-31-3) [2,](#page-30-5) [10,](#page-30-6) [9\]](#page-30-7).

In this paper, we will present a general approach in the language of quadratic forms to determine the number of rational points of those Artin-Schreier curves. Given a basis $\alpha_1, \ldots, \alpha_n$ of $\mathbb{F}_{q^n}/\mathbb{F}_q$, for $w = w_1\alpha_1 + \cdots + w_n\alpha_n$ with $w_1, \ldots, w_n \in \mathbb{F}_q$ we have

$$
\operatorname{Tr}(w\ell(w)) = (w_1 \quad \cdots \quad w_n) C \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix},
$$

where C is an $n \times n$ matrix whose (i, j) entry is $\frac{1}{2} \text{Tr}(\alpha_i \ell(\alpha_j) + \alpha_j \ell(\alpha_i))$. We call C the associated matrix of the quadratic form $\text{Tr}(x\ell(x))$ with respect to the basis $\alpha_1, \ldots, \alpha_n$. Since C is symmetric, it can be diagonalized as $C = P^TDP$ for some nonsingular matrix P and some diagonal matrix D over \mathbb{F}_q (cf. [\[8,](#page-30-8) Theorem 6.21]). Let $r = \text{rank } C$ be the rank of C, and rearrange the entries of D so that its first r diagonal entries d_1, \ldots, d_r are nonzero. Then

$$
\operatorname{Tr}(w\ell(w)) = \begin{pmatrix} w_1 & \cdots & w_n \end{pmatrix} P^{\mathrm{T}} D P \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix} = d_1 w_1^2 + \cdots + d_r w_r^2.
$$
 (1)

Let χ be the canonical additive character of \mathbb{F}_q , and η be the quadratic character of \mathbb{F}_q^* . For $a \in \mathbb{F}_q^*$, we have

$$
\sum_{u \in \mathbb{F}_q} \chi(au^2) = \eta(a)\mathcal{G},
$$

 $\mathcal{G} = \sum$ $u \in \mathbb{F}_q^*$ $\chi(u)\eta(u)$

is the quadratic Gauss sum of \mathbb{F}_q with $\mathcal{G}^2 = \eta(-1)q$. Replacing $(w_1 \cdots w_n) P^T$ by $(w_1 \cdots w_n)$ in [\(1\)](#page-1-0), we get

$$
\sum_{w \in \mathbb{F}_{q^n}} \chi(\text{Tr}(w\ell(w)))
$$
\n
$$
= \sum_{w_1, \dots, w_n \in \mathbb{F}_q} \chi(d_1 w_1^2 + \dots + d_r w_r^2)
$$
\n
$$
= \sum_{w_1 \in \mathbb{F}_q} \chi(d_1 w_1^2) \dots \sum_{w_r \in \mathbb{F}_q} \chi(d_r w_r^2) \sum_{w_{r+1}, \dots, w_n \in \mathbb{F}_q} 1
$$
\n
$$
= \eta(d_1 \cdots d_r) \mathcal{G}^r q^{n-r}.
$$

Here the numbers r, called the rank of the quadratic form, and $\eta(d_1 \cdots d_r)$, denoted by ϵ , act as invariants of the quadratic form $\text{Tr}(x\ell(x))$, which do not depend on the choice of basis. In particular, if C is nonsingular, then

$$
\epsilon = \eta(d_1 \cdots d_r) = \eta(\det D) = \eta(\det C)
$$

since $C = P^T D P$ with $\det P \in \mathbb{F}_q^*$.

Now we are ready for the number of roots of $\text{Tr}(x\ell(x)) + \lambda$ in \mathbb{F}_{q^n} for $\lambda \in \mathbb{F}_q$, which is, by the orthogonality relations for characters,

$$
q^{-1} \sum_{w \in \mathbb{F}_{q^n}} \sum_{u \in \mathbb{F}_q} \chi(u \text{Tr}(w\ell(w)) + u\lambda)
$$

= $q^{-1} \sum_{u \in \mathbb{F}_q} \sum_{w_1, \dots, w_n \in \mathbb{F}_q} \chi(u(d_1 w_1^2 + \dots + d_r w_r^2)) \chi(u\lambda)$
= $q^{n-1} + q^{-1} \sum_{u \in \mathbb{F}_q^*} \eta(u)^r \epsilon \mathcal{G}^r q^{n-r} \chi(u\lambda);$

that is

$$
\begin{cases} q^{n-1} + (-1)^{\frac{(q-1)r}{4}} \epsilon(q-1) q^{n-1-\frac{r}{2}} & \text{if } r \text{ is even,} \\ q^{n-1} & \text{if } r \text{ is odd} \end{cases}
$$

when $\lambda = 0$, and is

$$
\begin{cases} q^{n-1} - (-1)^{\frac{(q-1)r}{4}} \epsilon q^{n-1-\frac{r}{2}} & \text{if } r \text{ is even,} \\ q^{n-1} + (-1)^{\frac{(q-1)(r+1)}{4}} \eta(\lambda) \epsilon q^{n-1-\frac{r-1}{2}} & \text{if } r \text{ is odd} \end{cases}
$$

where

when $\lambda \neq 0$. For the Artin-Schreier curve over \mathbb{F}_{q^n} defined by $y^q - y - x\ell(x)$, it has exactly one point at infinity. Given $x_0 \in \mathbb{F}_{q^n}$, $y_0^q - y_0 = x_0 \ell(x_0)$ for some $y_0 \in \mathbb{F}_{q^n}$ if and only if $Tr(x_0\ell(x_0)) = 0$, in which case the number of such y_0 in \mathbb{F}_{q^n} is q. Therefore, we obtain the following conclusion (cf. [\[1,](#page-30-3) Corollary 3.7]).

Lemma 1. The Artin-Schreier curve over \mathbb{F}_{q^n} defined by $y^q - y - x\ell(x)$ with $\deg \ell = q^m$ is maximal or minimal if and only if n is even and $r = n - 2m$. In this case, it is maximal if $\epsilon = (-1)^{\frac{(q-1)r}{4}}$, and is minimal otherwise.

We have seen that the invariants r and ϵ carry essential information about the quadratic form, which comprise the central topic of this paper. In Section [2,](#page-4-0) we propose a general method to determine the invariants so as to classify all quadratic forms represented by polynomials over finite fields. In Section [3,](#page-10-0) we focus on a special case where all the coefficients of the polynomials belong to \mathbb{F}_q , providing a complete characterization. In Section [4,](#page-17-0) we study some polynomials of simple forms inducing maximal or minimal Artin-Schreier curves.

The following notation will be used throughout. For $\mathbb{F}_{q^n}/\mathbb{F}_q$ with $n > 1$ and q odd, let l be a nonzero q-linear polynomial over \mathbb{F}_{q^n} of degree less than q^n . Denote by Tr and N the trace map and the norm map of $\mathbb{F}_{q^n}/\mathbb{F}_q$ respectively. The invariant ϵ of the quadratic form Tr($x\ell(x)$) is defined above, and η is the quadratic character of \mathbb{F}_q^* .

Consider \mathbb{F}_{q^n} as a vector space over \mathbb{F}_q . For a subspace W of \mathbb{F}_{q^n} , denote by W^{\perp} the subspace consisting of all elements $\alpha \in \mathbb{F}_{q^n}$ such that $\text{Tr}(\alpha x)$ vanishes on W (one can check that $(W^{\perp})^{\perp} = W$ and $\dim_{\mathbb{F}_q} W + \dim_{\mathbb{F}_q} W^{\perp} = n$. Given a q-linear polynomial L over \mathbb{F}_{q^n} of the form

$$
L(x) = \sum_{i=0}^{n-1} a_i x^{q^i},
$$

define L' as

$$
L'(x) = a_0 x + \sum_{i=1}^{n-1} (a_i x)^{q^{n-i}},
$$

and let

$$
M_L = \begin{pmatrix} a_0 & a_1 & \cdots & a_{n-1} \\ a_{n-1}^q & a_0^q & \cdots & a_{n-2}^q \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{q^{n-1}} & a_2^{q^{n-1}} & \cdots & a_0^{q^{n-1}} \end{pmatrix}.
$$

For a positive integer k and a matrix M over some field, let $M^{(k)}$ denote the submatrix of M formed by its first k rows and k columns.

2 Classification of quadratic forms by matrices

For L and L' defined above, it is clear that $\text{Tr}(\alpha L(\beta)) = \text{Tr}(L'(\alpha)\beta)$ for all $\alpha, \beta \in \mathbb{F}_{q^n}$. In addition, $L'(\alpha) = 0$ if and only if $\text{Tr}(\alpha L(\beta)) = 0$ for all $\beta \in \mathbb{F}_{q^n}$, if and only if $\alpha \in (\text{im } L)^{\perp}$. This means ker $L' = (\text{im } L)^{\perp}$.

Let $\alpha_1, \ldots, \alpha_n$ be a basis of $\mathbb{F}_{q^n}/\mathbb{F}_q$ with a matrix

$$
A = \begin{pmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ \alpha_1^q & \alpha_2^q & \cdots & \alpha_n^q \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{q^{n-1}} & \alpha_2^{q^{n-1}} & \cdots & \alpha_n^{q^{n-1}} \end{pmatrix} . \tag{2}
$$

Then the (i, j) entry of A^TM_LA is

$$
\begin{pmatrix}\n\alpha_i & \alpha_i^q & \cdots & \alpha_i^{q^{n-1}}\n\end{pmatrix} M_L\n\begin{pmatrix}\n\alpha_j \\
\alpha_j^q \\
\vdots \\
\alpha_j^{q^{n-1}}\n\end{pmatrix} =\n\begin{pmatrix}\n\alpha_i & \alpha_i^q & \cdots & \alpha_i^{q^{n-1}}\n\end{pmatrix}\n\begin{pmatrix}\nL(\alpha_j) \\
L(\alpha_j)^q \\
\vdots \\
L(\alpha_j)^{q^{n-1}}\n\end{pmatrix}
$$
\n
$$
= \text{Tr}(\alpha_i L(\alpha_j)).
$$

For the quadratic form $\text{Tr}(x\ell(x))$, its associated symmetric matrix C with respect to the basis $\alpha_1, \ldots, \alpha_n$ has (i, j) entry

$$
\frac{1}{2} \text{Tr}(\alpha_i \ell(\alpha_j) + \alpha_j \ell(\alpha_i)) = \frac{1}{2} \text{Tr}(\alpha_i \ell(\alpha_j) + \alpha_i \ell'(\alpha_j)).
$$

If $L = \frac{\ell + \ell'}{2}$ $\frac{1}{2}$, then clearly $C = A^{T} M_L A$, and the rank of the quadratic form is rank $C =$ rank M_L since A is nonsingular. This will be employed to characterize the quadratic form in terms of the coefficients of ℓ . For rank M_L , we have the following results.

Theorem 2. Let

$$
L(x) = \sum_{i=0}^{n-1} a_i x^{q^i}
$$

with coefficients in \mathbb{F}_{q^n} . For an integer k with $0 < k \leq n$, the following conditions are equivalent:

- (i) dim_{\mathbb{F}_q} ker $L \geq k$;
- (ii) rank $M_L \leq n - k$;
- (iii) the first $n k + 1$ rows of M_L are linearly dependent over \mathbb{F}_{q^n} .

As a consequence, $\dim_{\mathbb{F}_q} \ker L + \operatorname{rank} M_L = n$, and the first $\operatorname{rank} M_L$ number of rows of M_L are linearly independent over \mathbb{F}_{q^n} while the first rank $M_L + 1$ number of rows are not.

Proof. First, suppose that $\dim_{\mathbb{F}_q} \ker L \geq k$. Take a basis of ker L/\mathbb{F}_q and extend it to a basis of $\mathbb{F}_{q^n}/\mathbb{F}_q$, denoted by $\alpha_1, \ldots, \alpha_n$, so that

$$
M_L \begin{pmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ \alpha_1^q & \alpha_2^q & \cdots & \alpha_n^q \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{q^{n-1}} & \alpha_2^{q^{n-1}} & \cdots & \alpha_n^{q^{n-1}} \end{pmatrix} = \begin{pmatrix} L(\alpha_1) & L(\alpha_2) & \cdots & L(\alpha_n) \\ L(\alpha_1)^q & L(\alpha_2)^q & \cdots & L(\alpha_n)^q \\ \vdots & \vdots & \ddots & \vdots \\ L(\alpha_1)^{q^{n-1}} & L(\alpha_2)^{q^{n-1}} & \cdots & L(\alpha_n)^{q^{n-1}} \end{pmatrix},
$$

where the matrix on the right side has at least k columns being zero. Then it has rank at most $n - k$, and so does M_L .

Next, [\(ii\)](#page-4-1) implies [\(iii\)](#page-4-2) since the rank of M_L is the maximal number of its rows that are linearly independent over \mathbb{F}_{q^n} .

Suppose now that the first $n - k + 1$ rows of M_L are linearly dependent over \mathbb{F}_{q^n} . Then

$$
\begin{pmatrix} b_0 & b_1 & \cdots & b_{n-k} & 0 & \cdots & 0 \end{pmatrix} M_L
$$

is zero for some $b_0, b_1, \ldots, b_{n-k} \in \mathbb{F}_{q^n}$. Note that

$$
(b_0 \quad b_1 \quad \cdots \quad b_{n-k} \quad 0 \quad \cdots \quad 0) \ M_L
$$
\n
$$
= (b_0 \quad b_1 \quad \cdots \quad b_{n-k} \quad 0 \quad \cdots \quad 0) \begin{pmatrix} a_0 & a_1 & \cdots & a_{n-1} \\ a_{n-1}^q & a_0^q & \cdots & a_{n-2}^q \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{q^{n-1}} & a_2^{q^{n-1}} & \cdots & a_0^{q^{n-1}} \end{pmatrix}
$$
\n
$$
= \left(\sum_{i=0}^{n-k} b_i a_{-i}^{q^i} \sum_{i=0}^{n-k} b_i a_{1-i}^{q^i} \quad \cdots \quad \sum_{i=0}^{n-k} b_i a_{n-1-i}^{q^i} \right),
$$
\n(3)

where the subscripts are taken modulo *n*. Therefore, for $L_0(x) = \sum_{i=0}^{n-k} b_i x^{q^i}$, one has

$$
L_0(L(x)) = \sum_{i=0}^{n-k} b_i \sum_{j=0}^{n-1} a_j^{q^i} x^{q^{i+j}} \equiv \sum_{j=0}^{n-1} \sum_{i=0}^{n-k} b_i a_{j-i}^{q^i} x^{q^j} \equiv 0 \pmod{x^{q^n} - x},\tag{4}
$$

 \Box

and then im $L \subseteq \ker L_0$. As

$$
\dim_{\mathbb{F}_q} \text{im } L \le \dim_{\mathbb{F}_q} \ker L_0 \le n - k,
$$

it turns out that $\dim_{\mathbb{F}_q} \ker L \geq k$. This indicates that [\(iii\)](#page-4-2) implies [\(i\).](#page-4-3)

For $L = \frac{\ell + \ell'}{2}$ $\frac{1}{2}$, we will see that ϵ depends on some submatrix of M_L . In fact, for $r = \dim_{\mathbb{F}_q} \text{im } L$, the submatrix $M_L^{(r)}$ $L^{(r)}$ is nonsingular and its determinant will be used to determine ϵ .

Theorem 3. Let $L = \frac{\ell + \ell'}{2}$ $\frac{e^{+\ell}}{2}$, and suppose that $\dim_{\mathbb{F}_q} \text{im } L = r$ for some integer r with $0 < r \leq n$. If $r = n$, then det $M_L \in \mathbb{F}_q$ and $\epsilon = (-1)^{n-1}\eta(\det M_L)$. Suppose that $r < n$. Then det $M_L^{(r)}$ $L^{(r)} \neq 0$, and

- (i) $\int \det M_L^{(r)}$ $\binom{r}{L}^{q+1} = \left(\det M_L^{(r,1)}\right)$ $\binom{m}{L}^2$, where $M_L^{(r,1)}$ $L^{(1,1)}$ is the submatrix formed by the r consecutive rows starting from the second row of M_L along with the first r columns of M_L , and $\epsilon = 1$ if and only if $\left(\det M_L^{(r)}\right)$ $\binom{r}{L}^{\frac{q+1}{2}} = \det M_L^{(r,1)}$ $\stackrel{(T,1)}{L};$
- (ii) $\epsilon = 1$ if and only if det $M_L^{(r)}$ $L^{(r)}$ is a square in \mathbb{F}_{q^n} , when *n* is odd;
- (iii) $a^{2q^r \frac{q^m-1}{q-1}}$ $\frac{m_{-1}}{q-1}$ det $M_L^{(r)} \in \mathbb{F}_q$ and $\epsilon = (-1)^{n-1} \eta \left(a^{2q^r \frac{q^m-1}{q-1}} \right)$ $\frac{m-1}{q-1}$ det $M_L^{(r)}$ $\binom{r}{L}$, when $r = n - 2m$ and ℓ has degree q^m with leading coefficient a.

Proof. Take a basis $\alpha_1, \ldots, \alpha_n$ of $\mathbb{F}_{q^n}/\mathbb{F}_q$ such that $\alpha_{r+1}, \ldots, \alpha_n \in \text{ker } L$, and define A as in [\(2\)](#page-4-4). By the discussion preceding Theorem [2,](#page-4-5) the quadratic form $Tr(x\ell(x))$ has an associated symmetric matrix written as

$$
\begin{pmatrix} C & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} = A^{\mathrm{T}} M_L A,
$$

where C is an $r \times r$ submatrix over \mathbb{F}_q and other entries are all zero. Let $C = P^T D P$ for some nonsingular matrix P and some diagonal matrix D over \mathbb{F}_q , so that

$$
\begin{pmatrix} C & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} = \begin{pmatrix} P^{\mathrm{T}} & \mathbf{0} \\ \mathbf{0} & I_{n-r} \end{pmatrix} \begin{pmatrix} D & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} P & \mathbf{0} \\ \mathbf{0} & I_{n-r} \end{pmatrix},
$$

where I_{n-r} is the identity matrix of order $n-r$ over \mathbb{F}_q . It follows from Theorem [2](#page-4-5) that rank $C = \text{rank } M_L = r$, so the diagonal entries of D are all nonzero and $\epsilon = \eta(\det D)$ $\eta(\det C)$.

Observe that

$$
(\det A)^q = \begin{vmatrix} \alpha_1^q & \alpha_2^q & \cdots & \alpha_n^q \\ \alpha_1^{q^2} & \alpha_2^{q^2} & \cdots & \alpha_n^{q^2} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1 & \alpha_2 & \cdots & \alpha_n \end{vmatrix} = (-1)^{n-1} \det A,
$$

so $(\det A)^2 \in \mathbb{F}_q$. If $r = n$, then $C = A^T M_L A$, and

$$
\epsilon = \eta((\det A)^2 \det M_L) = (-1)^{n-1} \eta(\det M_L).
$$

Assume now that $r < n$ and let β_1, \ldots, β_n be the dual basis of $\alpha_1, \ldots, \alpha_n$ in $\mathbb{F}_{q^n}/\mathbb{F}_q$, so that

$$
A^{-1} = \begin{pmatrix} \beta_1 & \beta_1^q & \cdots & \beta_1^{q^{n-1}} \\ \beta_2 & \beta_2^q & \cdots & \beta_2^{q^{n-1}} \\ \vdots & \vdots & \ddots & \vdots \\ \beta_n & \beta_n^q & \cdots & \beta_n^{q^{n-1}} \end{pmatrix}.
$$

Partition it as

$$
A^{-1} = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix},
$$

where B_{11} is an $r\times r$ submatrix. Then

$$
M_L = (A^{\mathrm{T}})^{-1} \begin{pmatrix} C & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} A^{-1}
$$

= $\begin{pmatrix} B_{11}^{\mathrm{T}} & B_{21}^{\mathrm{T}} \\ B_{12}^{\mathrm{T}} & B_{22}^{\mathrm{T}} \end{pmatrix} \begin{pmatrix} C & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$
= $\begin{pmatrix} B_{11}^{\mathrm{T}} C & \mathbf{0} \\ B_{12}^{\mathrm{T}} C & \mathbf{0} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$
= $\begin{pmatrix} B_{11}^{\mathrm{T}} C B_{11} & B_{11}^{\mathrm{T}} C B_{12} \\ B_{12}^{\mathrm{T}} C B_{11} & B_{12}^{\mathrm{T}} C B_{12} \end{pmatrix}$,

and

$$
\det M_L^{(r)} = (\det B_{11})^2 \det C.
$$
 (5)

Note that

$$
\begin{vmatrix} \beta_1 & \beta_1^q & \cdots & \beta_1^{q^{r-1}} & \beta_1^{q^r} \\ \beta_2 & \beta_2^q & \cdots & \beta_2^{q^{r-1}} & \beta_2^{q^r} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \beta_r & \beta_r^q & \cdots & \beta_r^{q^{r-1}} & \beta_r^{q^r} \\ x & x^q & \cdots & x^{q^{r-1}} & x^{q^r} \end{vmatrix}
$$

= det $B_{11}x^{q^r} + \cdots + (-1)^r (\det B_{11})^q x$,

as a linear endomorphism of $\mathbb{F}_{q^n}/\mathbb{F}_q$ whose kernel is exactly the subspace spanned by β_1, \ldots, β_r , as is easily verified. Also, det B_{11} is nonzero, and so is det $M_L^{(r)}$ $L^{(r)}$. Divide the above polynomial by $\det B_{11}$ to get a monic polynomial

$$
L_0(x) = x^{q^r} + \dots + (-1)^r (\det B_{11})^{q-1} x.
$$

The statement [\(ii\)](#page-6-0) follows immediately from [\(5\)](#page-7-0) since $(\det C)^{\frac{q^n-1}{2}} = (\det C)^{\frac{q-1}{2}}$ when n is odd. Since $L = L'$, we have $\text{im } L = (\ker L)^{\perp} = \ker L_0$, and

$$
L_0(L(x)) \equiv 0 \pmod{x^{q^n} - x}.
$$

By the same argument as in [\(3\)](#page-5-0) and [\(4\)](#page-5-1), we find that

$$
((-1)^r (\det B_{11})^{q-1} \cdots 1) \begin{pmatrix} M_L^{(r)} \\ v \end{pmatrix}
$$

is zero, where v is the $1 \times r$ submatrix formed by the first r entries of the $(r + 1)$ -st row of M_L . Cramer's rule leads to

$$
(-1)^r (\det B_{11})^{q-1} = (-1)^r \frac{\det M_L^{(r,1)}}{\det M_L^{(r)}}.
$$

This, coupled with

$$
\Big(\det M_L^{(r)}\Big)^{\frac{q-1}{2}}=(\det B_{11})^{q-1}(\det C)^{\frac{q-1}{2}},
$$

implies [\(i\).](#page-6-1)

To prove [\(iii\),](#page-6-2) suppose that $\dim_{\mathbb{F}_q}\ker L=2m$ and

$$
\ell(x)+\ell'(x)=\cdots+(ax)^{q^{n-m}}+ax^{q^m}+\cdots.
$$

Let L_1 be the polynomial over \mathbb{F}_{q^n} of degree less than q^n such that

$$
L'_1(x) \equiv 2a^{-q^{n-2m}} L(x)^{q^{n-m}} \pmod{x^{q^n} - x};
$$

that is

$$
L'_1(x) = \dots + x^{q^{n-2m}} + a^{q^{n-m}-q^{n-2m}}x,
$$

with

$$
L_1(x) = x^{q^{2m}} + \dots + a^{q^{n-m}-q^{n-2m}}x.
$$

Then ker $L'_1 = \ker L$, spanned by $\alpha_{r+1}, \ldots, \alpha_n$. Accordingly, $\operatorname{im} L_1 = (\ker L'_1)^{\perp} = \ker L_0$ and

$$
L_0(L_1(x)) \equiv 0 \pmod{x^{q^n} - x}.
$$

It has been seen that L_0 is monic of degree q^r , and L_1 is monic of degree q^{2m} by a simple investigation, which means

$$
L_0(L_1(x)) = x^{q^n} - x.
$$

Equating the constant terms gives

$$
(-1)^r (\det B_{11})^{q-1} a^{q^r (q^m - 1)} = -1.
$$

Finally, taking [\(5\)](#page-7-0) into account we obtain

$$
a^{2q^r\frac{q^m-1}{q-1}}\det M_L^{(r)}=a^{2q^r\frac{q^m-1}{q-1}}(\det B_{11})^2\det C\in \mathbb{F}_q,
$$

where

$$
\left(a^{2q^r \frac{q^m-1}{q-1}} (\det B_{11})^2\right)^{\frac{q-1}{2}} = a^{q^r (q^m-1)} (\det B_{11})^{q-1} = (-1)^{r-1},
$$

so

$$
\eta\left(a^{2q^r \frac{q^m-1}{q-1}} \det M_L^{(r)}\right) = (-1)^{r-1} \eta(\det C) = (-1)^{n-1} \epsilon,
$$

as desired.

Example 4. We classify all those quadratic forms of $\mathbb{F}_{q^4}/\mathbb{F}_q$ as follows (the results for $\mathbb{F}_{q^2}/\mathbb{F}_q$ and $\mathbb{F}_{q^3}/\mathbb{F}_q$ can be found in [\[4,](#page-30-9) Section 3]). Let $\ell(x) = 2ax^{q^2} + 2bx^q + cx$ and $L(x) = \frac{\ell(x) + \ell'(x)}{2} = (bx)^{q^3} + (a^{q^2} + a)x^{q^2} + bx^q + cx$ be nonzero polynomials over \mathbb{F}_{q^4} , so that 2 3

$$
M_L = \begin{pmatrix} c & b & a^{q^2} + a & b^{q^3} \\ b & c^q & b^q & a^{q^3} + a^q \\ a^{q^2} + a & b^q & c^{q^2} & b^{q^2} \\ b^{q^3} & a^{q^3} + a^q & b^{q^2} & c^{q^3} \end{pmatrix}.
$$

Then

• rank $M_L = 1$ if and only if

$$
\begin{vmatrix} c & b \\ b & c^q \end{vmatrix} = \begin{vmatrix} b & a^{q^2} + a \\ c^q & b^q \end{vmatrix} = \begin{vmatrix} a^{q^2} + a & b^{q^3} \\ b^q & a^{q^3} + a^q \end{vmatrix} = 0,
$$

in which case $\epsilon = 1$ if and only if $c^{\frac{q+1}{2}} = b$;

• rank $M_L = 2$ if and only if c b b c^q $\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array}\\ \end{array} \end{array} \end{array}$ $\neq 0$ and c b $a^{q^2} + a$ b c^q b^q = b $a^{q^2} + a$ b^{q^3} c^q b^q $a^{q^3} + a^q$

 $a^{q^2}+a \quad b^q \qquad c^{q^2}$

in which case $\epsilon = 1$ if and only if

$$
\begin{vmatrix} c & b \\ b & c^q \end{vmatrix}^{\frac{q+1}{2}} = \begin{vmatrix} b & c^q \\ a^{q^2} + a & b^q \end{vmatrix};
$$

 b^q c^{q^2} b^{q^2}

 $= 0,$ \Box

• rank $M_L = 3$ if and only if

$$
\begin{vmatrix} c & b & a^{q^2} + a \\ b & c^q & b^q \\ a^{q^2} + a & b^q & c^{q^2} \end{vmatrix} \neq 0
$$

and det $M_L = 0$, in which case $\epsilon = 1$ if and only if

$$
\begin{vmatrix} c & b & a^{q^2} + a \\ b & c^q & b^q \\ a^{q^2} + a & b^q & c^{q^2} \end{vmatrix} \stackrel{q+1}{=} \begin{vmatrix} b & c^q & b^q \\ a^{q^2} + a & b^q & c^{q^2} \\ b^{q^3} & a^{q^3} + a^q & b^{q^2} \end{vmatrix};
$$

• rank $M_L = 4$ if and only if det $M_L \neq 0$, in which case $\epsilon = -\eta(\det M_L)$.

3 Quadratic forms represented by polynomials with coefficients in \mathbb{F}_q

For $L = \frac{\ell + \ell'}{2}$ $\frac{1}{2}$, consider the special case that L has all coefficients in \mathbb{F}_q . Some Artin-Schreier curves induced from this type of polynomials have been studied in, e.g., [\[1,](#page-30-3) Theorem 5.3] and [\[9,](#page-30-7) Theorem 3.1], with some restrictions such as $gcd(n, q) = 1$. In fact, the matrix M_L is a circulant matrix that can be easily diagonalized, provided that there exists a primitive n-th root of unity over \mathbb{F}_q . However, this is not necessarily the case. In this section, we will address the problem, providing a complete characterization of all those quadratic forms.

For a polynomial f in the polynomial ring $\mathbb{C}[z]$ written as

$$
f(z) = \sum_{i=0}^{n-1} c_i z^i,
$$

let C_f be the circulant matrix defined by

$$
C_f = \begin{pmatrix} c_0 & c_1 & \cdots & c_{n-1} \\ c_{n-1} & c_0 & \cdots & c_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ c_1 & c_2 & \cdots & c_0 \end{pmatrix}
$$

.

The rank of C_f and the determinant of some submatrix of it can be determined as follows.

Lemma 5. Let ζ_1, \ldots, ζ_n be the distinct n-th roots of unity in \mathbb{C} . Let f and g be polynomials over $\mathbb C$ such that $\prod_{j=1}^r g(\zeta_j) \neq 0$ and

$$
f(z) = \sum_{i=0}^{n-1} c_i z^i \equiv g(z) \prod_{i=r+1}^{n} (z - \zeta_i) \pmod{z^n - 1}
$$

with $0 < r \leq n$. Then rank $C_f = r$ and $\det C_f^{(r)} = \prod_{j=1}^r \zeta_j^{-r} g(\zeta_j)$.

Proof. Let Z be the $n \times n$ matrix whose (i, j) entry is ζ_j^{i-1} . A simple investigation yields

$$
Z^{-1} = \frac{1}{n} \begin{pmatrix} 1 & \zeta_1^{-1} & \cdots & \zeta_1^{-(n-1)} \\ 1 & \zeta_2^{-1} & \cdots & \zeta_2^{-(n-1)} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \zeta_n^{-1} & \cdots & \zeta_n^{-(n-1)} \end{pmatrix}.
$$

Note that

$$
C_{f}Z = \begin{pmatrix} c_{0} & c_{1} & \cdots & c_{n-1} \\ c_{n-1} & c_{0} & \cdots & c_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ c_{1} & c_{2} & \cdots & c_{0} \end{pmatrix} \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \zeta_{1} & \zeta_{2} & \cdots & \zeta_{n} \\ \vdots & \vdots & \ddots & \vdots \\ \zeta_{1}^{n-1} & c_{2}^{n-1} & \cdots & c_{n}^{n-1} \end{pmatrix}
$$

$$
= \begin{pmatrix} f(\zeta_{1}) & f(\zeta_{2}) & \cdots & f(\zeta_{n}) \\ \zeta_{1}f(\zeta_{1}) & \zeta_{2}f(\zeta_{2}) & \cdots & \zeta_{n}f(\zeta_{n}) \\ \vdots & \vdots & \ddots & \vdots \\ \zeta_{1}^{n-1}f(\zeta_{1}) & \zeta_{2}^{n-1}f(\zeta_{2}) & \cdots & c_{n}^{n-1}f(\zeta_{n}) \end{pmatrix}
$$

$$
= \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \zeta_{1} & \zeta_{2} & \cdots & \zeta_{n} \\ \vdots & \vdots & \ddots & \vdots \\ \zeta_{1}^{n-1} & \zeta_{2}^{n-1} & \cdots & c_{n}^{n-1} \end{pmatrix} \begin{pmatrix} f(\zeta_{1}) & & & \\ & f(\zeta_{2}) & & \\ & & \ddots & \\ & & & & f(\zeta_{n}) \end{pmatrix}
$$

$$
= ZD,
$$

where D is the above diagonal matrix. This proves rank $C_f = r$, as $\prod_{j=1}^r f(\zeta_j)$ = $\prod_{j=1}^r g(\zeta_j) \neq 0.$

Partition D into four blocks where the upper-left one is $D^{(r)}$, so that the others are all zero. Since $C_f = ZDZ^{-1}$, partitioning C_f , Z and Z^{-1} in the same manner gives

$$
\det C_f^{(r)} = \frac{1}{n^r} \begin{vmatrix} 1 & 1 & \cdots & 1 \\ \zeta_1 & \zeta_2 & \cdots & \zeta_r \\ \vdots & \vdots & \ddots & \vdots \\ \zeta_1^{r-1} & \zeta_2^{r-1} & \cdots & \zeta_r^{r-1} \end{vmatrix} \begin{vmatrix} 1 & \zeta_1^{-1} & \cdots & \zeta_1^{-(r-1)} \\ 1 & \zeta_2^{-1} & \cdots & \zeta_2^{-(r-1)} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \zeta_r^{-1} & \cdots & \zeta_r^{-(r-1)} \end{vmatrix} \det D^{(r)}.
$$

Calculating the Vandermonde determinants, we get

$$
n^{r} \det C_{f}^{(r)} = \prod_{1 \leq j < i \leq r} (\zeta_{i} - \zeta_{j}) \prod_{1 \leq j < i \leq r} (\zeta_{i}^{-1} - \zeta_{j}^{-1}) \prod_{j=1}^{r} f(\zeta_{j})
$$
\n
$$
= \prod_{1 \leq i < j \leq r} (\zeta_{j} - \zeta_{i}) \prod_{1 \leq j < i \leq r} (\zeta_{j} - \zeta_{i}) \zeta_{i}^{-1} \zeta_{j}^{-1} \prod_{j=1}^{r} f(\zeta_{j})
$$
\n
$$
= \prod_{j=1}^{r} \prod_{\substack{1 \leq i \leq r \\ i \neq j}} (\zeta_{j} - \zeta_{i}) \prod_{1 \leq j < i \leq r} \zeta_{i}^{-1} \zeta_{j}^{-1} \prod_{j=1}^{r} \left(g(\zeta_{j}) \prod_{i=r+1}^{n} (\zeta_{j} - \zeta_{i}) \right)
$$
\n
$$
= \prod_{j=1}^{r} \prod_{\substack{1 \leq i \leq n \\ i \neq j}} (\zeta_{j} - \zeta_{i}) \prod_{j=1}^{r} \zeta_{j}^{1-r} \prod_{j=1}^{r} g(\zeta_{j})
$$
\n
$$
= \prod_{j=1}^{r} n \zeta_{j}^{-1} \prod_{j=1}^{r} \zeta_{j}^{1-r} g(\zeta_{j})
$$
\n
$$
= n^{r} \prod_{j=1}^{r} \zeta_{j}^{-r} g(\zeta_{j}),
$$

where the equality

$$
\prod_{\substack{1 \le i \le n \\ i \ne j}} (\zeta_j - \zeta_i) = n\zeta_j^{-1}
$$

can be obtained by taking formal derivatives of both sides of

$$
(z - \zeta_j) \prod_{\substack{1 \le i \le n \\ i \ne j}} (z - \zeta_i) = z^n - 1
$$

at ζ_j .

Remark 6. In this lemma, $\prod_{j=1}^{r} \zeta_j^{-r} g(\zeta_j)$ can be calculated from the coefficients of f using the resultant of $\frac{f(z)}{\gcd(f(z),z^n-1)}$ and $\frac{z^n-1}{\gcd(f(z),z^n-1)}$.

 \Box

For $L(x) = \sum_{i=0}^{n-1} a_i x^{q^i}$ with all coefficients in \mathbb{F}_q , define

$$
L^*(x) = \sum_{i=0}^{n-1} a_i x^i.
$$

Here is the finite field version of the preceding result (for fields of even characteristic as well).

Theorem 7. Let $\omega_1, \ldots, \omega_n$ be the (not necessarily distinct) n-th roots of unity over \mathbb{F}_q such that $x^n - 1 = \prod_{i=1}^n (x - \omega_i)$. Let L be a nonzero q-linear polynomial over \mathbb{F}_q of degree $less than qⁿ$ and

$$
\gcd(L^*(x), x^n - 1) = \prod_{i=r+1}^n (x - \omega_i)
$$

with $0 < r \leq n$, so that

$$
L^*(x) \equiv G(x) \prod_{i=r+1}^n (x - \omega_i) \pmod{x^n - 1}
$$

for some polynomial G over \mathbb{F}_q (for instance, let $G(x) = \frac{L^*(x)}{\gcd(L^*(x), x^n - 1)}$). Then rank $M_L =$ r, and det $M_L^{(r)} = \prod_{j=1}^r \omega_j^{-r} G(\omega_j)$. In particular, if $r = n$, then det $M_L = \prod_{j=1}^n L^*(\omega_j)$.

Proof. Let $n = p^e \nu$ with $gcd(\nu, p) = 1$, where p is the characteristic of \mathbb{F}_q . Denote by K the $(q-1)$ -st cyclotomic field with ring of integers \mathcal{O}_K , and let $E = K(\zeta)$ for a primitive n-th root of unity ζ in C. The ring of integers of E is $\mathcal{O}_E = \mathcal{O}_K[\zeta]$. For a prime P of K lying above $p\mathbb{Z}$, the residue field \mathcal{O}_K/P has q elements and there is a surjective ring homomorphism σ from \mathcal{O}_K to \mathbb{F}_q with kernel P. Take the minimal polynomial of ζ over K, which has coefficients all in \mathcal{O}_K , and apply σ to each of these coefficients to get a polynomial over \mathbb{F}_q . Let ω be a root of this polynomial, so that $\mathbb{F}_q(\omega)$ is a fintie extension of \mathbb{F}_q . By setting $\sigma(\zeta) = \omega$, we extend σ to a well-defined homomorphism (still denoted by σ) from $\mathcal{O}_K[\zeta]$ onto $\mathbb{F}_q(\omega)$, whose kernel is a prime \mathfrak{P} of E containing P. Note that ζ^{p^e} is a primitive ν -th root of unity and

$$
\prod_{i=1}^{\nu-1} (1 - \zeta^{p^{e_i}}) = \nu \notin \mathfrak{P},
$$

for otherwise $\nu \in \mathfrak{P} \cap \mathbb{Z} = p\mathbb{Z}$. Then $\omega^i \neq 1$ for $1 \leq i < \nu$ since $(1 - \omega^i)^{p^e} = \sigma(1 - \zeta^{p^{e_i}}) \neq 0$. Furthermore, $\omega^{\nu} = 1$ as $({\omega}^{\nu})^{p^e} = \sigma(\zeta)^n = 1$, so ω is a primitive *v*-th root of unity over \mathbb{F}_q . We conclude that σ maps the group of *n*-th roots of unity in $\mathbb C$ onto the group of ν -th roots of unity over \mathbb{F}_q , and thus

$$
x^n - 1 = \prod_{i=1}^n (x - \sigma(\zeta_i)),
$$

where ζ_1,\ldots,ζ_n are distinct *n*-th roots of unity in $\mathbb C$ such that $\sigma(\zeta_i) = \omega_i$ for $1 \leq i \leq n$.

In what follows, for a polynomial f over \mathcal{O}_K , denote by f the polynomial over \mathbb{F}_q obtained by applying σ to each coefficient of f. Then $G = \overline{g}$ for some polynomials g over \mathcal{O}_K , and

$$
L^*(x) = \bar{g}(x) \prod_{i=r+1}^n (x - \omega_i) + \bar{h}(x)(x^n - 1)
$$

for some polynomials h over \mathcal{O}_K , as $x^n - 1$ divides $L^*(x) - G(x) \prod_{i=r+1}^n (x - \omega_i)$ in $\mathbb{F}_q[x]$. Let

$$
f(z) = g(z) \prod_{i=r+1}^{n} (z - \zeta_i) + h(z)(z^n - 1),
$$

so that $\bar{f} = L^*$. Here $\prod_{j=1}^r g(\zeta_j) \neq 0$; otherwise $\prod_{j=1}^r \bar{g}(\omega_j) = 0$, but

$$
gcd(L^*(x), x^n - 1) = gcd\left(\bar{g}(x) \prod_{i=r+1}^n (x - \omega_i), x^n - 1\right)
$$

$$
= gcd\left(\bar{g}(x), \prod_{i=1}^r (x - \omega_i)\right) \prod_{i=r+1}^n (x - \omega_i).
$$

It follows from Lemma [5](#page-10-1) that

$$
\det C_f^{(r)} = \prod_{j=1}^r \zeta_j^{-r} g(\zeta_j).
$$

Calculating the determinants of matrices by definition, one gets

$$
\det M_L^{(r)} = \sigma(\det C_f^{(r)}) = \prod_{j=1}^r \sigma(\zeta_j^{-r} g(\zeta_j)) = \prod_{j=1}^r \omega_j^{-r} \overline{g}(\omega_j),
$$

and

$$
\det M_L = \prod_{j=1}^n \omega_j^{-n} G(\omega_j) = \prod_{j=1}^n L^*(\omega_j)
$$

when $r = n$. Moreover, every singular submatrix (of zero determinant) of C_f induces a singular submatrix of M_L , so rank $M_L \le$ rank $C_f = r$. This implies rank $M_L = r$. \Box

Theorem 8. Continue with the notation in the last theorem, and suppose that $L = \frac{\ell + \ell'}{2}$ 2 be nonzero with all coefficients in \mathbb{F}_q . Then

- (i) $\epsilon = 1$ if and only if $\prod_{j=1}^r G(\omega_j)^{\frac{q-1}{2}} = (-1)^{r-1} c^{\frac{q-1}{2}r+1}$, where c is the constant term of $gcd(L^*(x), x^n - 1)$ satisfying $c^2 = 1$;
- (ii) $\epsilon = (-1)^{(n-1-\mu)(\frac{q-1}{2}r+1)}\eta(\prod_{j=1}^r G(\omega_j)),$ where μ is the multiplicity of -1 as a root $of \gcd(L^*(x), x^n - 1);$

(iii)
$$
\epsilon = (-1)^{n-1} \eta \big(\prod_{j=1}^{n} L^*(\omega_j) \big)
$$
, when $r = n$;

(iv) $\epsilon = (-1)^{(n-1)(\frac{q-1}{2}m+1)}\eta(2a)^n$, when $r = n-2m < n$ and ℓ has degree q^m with leading coefficient a.

Proof. The first three statements are apparent in the case $r = n$. Suppose that $0 < r < n$ and let

$$
L^*(x) = G(x) \prod_{i=r+1}^n (x - \omega_i) + H(x)(x^n - 1)
$$

for some polynomial H over \mathbb{F}_q . Let k be a sufficiently large integer such that $x^{k-n+r}G(x^{-1})$ and $x^{k-n}H(x^{-1})$ are both polynomials in $\mathbb{F}_q[x]$. Then

$$
x^{k}L^{*}(x^{-1}) = x^{k}G(x^{-1}) \prod_{i=r+1}^{n} (x^{-1} - \omega_{i}) + x^{k}H(x^{-1})(x^{-n} - 1)
$$

= $x^{k-n+r}G(x^{-1}) \prod_{i=r+1}^{n} \omega_{i}(\omega_{i}^{-1} - x) + x^{k-n}H(x^{-1})(1 - x^{n}),$

which means

$$
x^{k} L^{*}(x^{-1}) \equiv x^{k-n+r} G(x^{-1}) \prod_{i=r+1}^{n} \omega_{i}(\omega_{i}^{-1} - x) \pmod{x^{n} - 1}.
$$
 (6)

For $L(x) = \sum_{i=0}^{n-1} a_i x^{q^i}$, one has

$$
x^{k}L^{*}(x^{-1}) = \sum_{i=0}^{n-1} a_{i}x^{k-i} \equiv \sum_{i=0}^{n-1} a_{i}x^{k+n-i} \equiv x^{k}L'^{*}(x) \pmod{x^{n}-1},
$$

so

$$
\gcd(x^{k} L^{*}(x^{-1}), x^{n} - 1) = \gcd(x^{k} L^{*}(x), x^{n} - 1) = \prod_{i=r+1}^{n} (x - \omega_{i})
$$

as $L = L'$. It follows from [\(6\)](#page-15-0) that $\prod_{i=r+1}^{n} (x - \omega_i^{-1})$ divides $x^k L^*(x^{-1})$, and consequently it divides $\prod_{i=r+1}^{n}(x - \omega_i)$. Then

$$
\prod_{i=r+1}^{n} (x - \omega_i) = \prod_{i=r+1}^{n} (x - \omega_i^{-1}),
$$

where ω_i and ω_i^{-1} have the same multiplicity. Hence, $(-1)^{\mu} \prod_{j=1}^{r} \omega_j = \prod_{j=1}^{n} \omega_j = (-1)^{n-1}$. Moreover,

$$
c = (-1)^{n-r} \prod_{i=r+1}^{n} \omega_i = (-1)^{n-r} \prod_{i=r+1}^{n} \omega_i^{-1} = c^{-1},
$$

and thus $\prod_{j=1}^{r} \omega_j = (-1)^{r-1}c$ with $c^2 = 1$.

Let L_1 be the polynomial of degree less than q^n with

$$
L_1(x) \equiv L(x)^q \pmod{x^{q^n} - x},
$$

so that

$$
L_1^*(x) \equiv xL^*(x) \equiv xG(x) \prod_{i=r+1}^n (x - \omega_i) \pmod{x^n - 1}.
$$

Since

$$
\gcd(L_1^*(x), x^n - 1) = \gcd(xL^*(x), x^n - 1) = \prod_{i=r+1}^n (x - \omega_i),
$$

the last theorem can be applied to get

$$
\det M_{L_1}^{(r)} = \prod_{j=1}^r \omega_j^{1-r} G(\omega_j).
$$

Observe that the submatrix of M_L defined in [\(i\)](#page-6-1) of Theorem [3](#page-6-3) is, in this case, exactly $M_{L_1}^{(r)}$ $L_1^{(r)}$. Then $\epsilon=1$ if and only if

$$
\prod_{j=1}^r (\omega_j^{-r} G(\omega_j))^{\frac{q+1}{2}} = \prod_{j=1}^r \omega_j^{1-r} G(\omega_j);
$$

that is,

$$
\prod_{j=1}^r G(\omega_j)^{\frac{q-1}{2}} = \prod_{j=1}^r \omega_j^{\frac{q-1}{2}r+1}.
$$

The results [\(i\)](#page-14-0) and [\(ii\)](#page-14-1) can be easily verified.

Finally, by the assumption of [\(iv\)](#page-14-2) we have

$$
x^m L^*(x) \equiv \frac{a}{2} x^{2m} + \dots + \frac{a}{2} \pmod{x^n - 1},
$$

where the polynomial on the right side has degree $2m$, while

$$
\gcd(x^m L^*(x), x^n - 1) = \gcd(L^*(x), x^n - 1) = \prod_{i=r+1}^n (x - \omega_i)
$$

has degree $n - r = 2m$. Therefore,

$$
x^{2m}+\cdots+1=\prod_{i=r+1}^n(x-\omega_i),
$$

which implies $\prod_{j=1}^{r} \omega_j = (-1)^{r-1}c = (-1)^{n-1}$ and

$$
L^*(x) \equiv x^{n-m} x^m L^*(x) \equiv \frac{a}{2} x^{n-m} \prod_{i=r+1}^n (x - \omega_i) \pmod{x^n - 1}.
$$

Then

$$
\epsilon = (-1)^{n-1} \eta \left(\det M_L^{(r)} \right)
$$

= $(-1)^{n-1} \eta \left(\prod_{j=1}^r \frac{a}{2} \omega_j^m \right)$
= $(-1)^{n-1} \eta \left((-1)^{(n-1)m} \left(\frac{a}{2} \right)^r \right)$
= $(-1)^{(n-1)(\frac{q-1}{2}m+1)} \eta (2a)^n$,

as a consequence of [\(iii\)](#page-6-2) in Theorem [3.](#page-6-3)

4 Maximal and minimal Artin-Schreier curves from some quadratic forms

To find a maximal or minimal curve as in Lemma [1,](#page-3-0) it is necessary and sufficient to let n be even and ℓ have degree q^m such that $\dim_{\mathbb{F}_q} \ker L = 2m$ for $L = \frac{\ell + \ell'}{2}$ $\frac{e^{i\theta}}{2}$. The trivial cases include

- $\ell(x) = ax^{q^{\frac{n}{2}}}$ with $a \in \mathbb{F}_{q^n}^*$ such that $a^{q^{\frac{n}{2}}} + a = 0$;
- $\ell(x) = ax$ with $a \in \mathbb{F}_{q^n}^*$.

Using the obtained classification results of quadratic forms, we can derive more such curves explicitly and decide whether they are maximal or minimal, which depends on $(-1)^{\frac{(q-1)(\hat{n}-2m)}{4}}\epsilon$. The case of odd *n* will be considered as well.

Example 9. Consider $\ell(x) = 2ax^{q^2} + 2bx^q + cx$ with $a \in \mathbb{F}_q^*$ and $b, c \in \mathbb{F}_q$, and suppose that $n > 4$. For $L = \frac{\ell + \ell'}{2}$ $\frac{e}{2}$, we have

$$
x^{2}L^{*}(x) \equiv ax^{4} + bx^{3} + cx^{2} + bx + a \pmod{x^{n} - 1},
$$

and $\dim_{\mathbb{F}_q} \ker L = 4$ if and only if $gcd(ax^4 + bx^3 + cx^2 + bx + a, x^n - 1)$ has degree 4. For an *n*-th root of unity ω over \mathbb{F}_q , it is clear that ω and ω^{-1} have the same multiplicity as roots of $ax^4 + bx^3 + cx^2 + bx + a$. In addition, the multiplicity of -1 , as well as of 1, must be even. In order for $a^{-1}(ax^4 + bx^3 + cx^2 + bx + a)$ to divide $x^n - 1$, it is necessarily

- the product of two polynomials of the form $(x \omega)(x \omega^{-1})$ with $\omega \in \mathbb{F}_q$ or $\omega^{q+1} = 1$, or
- $(x \omega)(x \omega^q)(x \omega^{-1})(x \omega^{-q})$ with $\omega \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ such that $\omega^{q+1} \neq 1$, or
- $(x \omega)(x \omega^q)(x \omega^{q^2})(x \omega^{q^3})$ with $\omega \in \mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}$.

 \Box

The sufficiency is routine to check. If $\dim_{\mathbb{F}_q} \ker L = 4$, then $\epsilon = (-1)^{n-1} \eta(a)^n$ by [\(iv\)](#page-14-2) of Theorem [8.](#page-14-3)

Next, we investigate certain binomials that may induce maximal or minimal Artin-Schreier curves.

Proposition 10. Let $L(x) = (ax)^{q^{n-1}} + ax^q + bx$ for $a, b \in \mathbb{F}_{q^n}^*$ with $n \geq 4$. Define a sequence of polynomials in $\mathbb{F}_{q^n}[x]$ as

$$
f_k(x) = -f_{k-1}(x) - x^{q^{k-2}} f_{k-2}(x)
$$
 for $k \ge 2$,

with $f_0 = 1$ and $f_1 = -1$. Then $\dim_{\mathbb{F}_q} \ker L = 2$ if and only if

$$
f_{n-1}(a^2b^{-1-q}) = N(a) - N(b)f_n(a^2b^{-1-q}) = 0,
$$

if and only if

$$
a^{2q^{n-3}} \det M_L^{(n-3)} - b^{q^{n-2}} \det M_L^{(n-2)} = \mathcal{N}(a) + (-1)^n a^{2q^{n-2}} \det M_L^{(n-2)} = 0.
$$

Proof. The kernel of L has at most q^2 elements, since

$$
L(x)^q \equiv a^q x^{q^2} + b^q x^q + ax \pmod{x^{q^n} - x}.
$$

By Theorem [2,](#page-4-5) $\dim_{\mathbb{F}_q} \ker L = 2$ if and only if

$$
(\beta_0 \cdots \beta_{n-2} 0) \begin{pmatrix} b & a & & & a^{q^{n-1}} \\ a & b^q & \ddots & & \\ & a^q & \ddots & a^{q^{n-4}} & \\ & & \ddots & b^{q^{n-3}} & a^{q^{n-3}} \\ & & & a^{q^{n-3}} & b^{q^{n-2}} & a^{q^{n-2}} \\ a^{q^{n-1}} & & & a^{q^{n-2}} & b^{q^{n-1}} \end{pmatrix}
$$

is zero for some $\beta_0, \ldots, \beta_{n-2} \in \mathbb{F}_{q^n}$ one of which is nonzero. Assume that this is the case. If $\beta_0 = 0$, then clearly $\beta_1, \ldots, \beta_{n-2}$ are all zero. Hence, let $\beta_0 = 1$ without loss of generality. Now we have

$$
\begin{cases}\nb + a\beta_1 = 0, \\
a^{q^{k-2}}\beta_{k-2} + b^{q^{k-1}}\beta_{k-1} + a^{q^{k-1}}\beta_k = 0 \quad \text{for } 2 \le k \le n-2, \\
a^{q^{n-3}}\beta_{n-3} + b^{q^{n-2}}\beta_{n-2} = 0, \\
a^{q^{n-1}} + a^{q^{n-2}}\beta_{n-2} = 0,\n\end{cases}
$$

where $\beta_1, \ldots, \beta_{n-2}$ are uniquely determined by all but the last two equations. We conclude that $\dim_{\mathbb{F}_q} \ker L = 2$ only if for the sequence of elements in \mathbb{F}_{q^n} defined by

$$
\beta_k = -a^{-q^{k-1}} \left(b^{q^{k-1}} \beta_{k-1} + a^{q^{k-2}} \beta_{k-2} \right) \text{ for } k \ge 2,
$$

with $\beta_0 = 1$ and $\beta_1 = -a^{-1}b$, one has

$$
a^{q^{n-3}} \beta_{n-3} + b^{q^{n-2}} \beta_{n-2} = a^{q^{n-1}} + a^{q^{n-2}} \beta_{n-2} = 0.
$$
 (7)

The converse is apparent.

We show by induction that with β_k defined above,

$$
a^{q^{k-1}+\dots+q+1}\beta_k = b^{q^{k-1}+\dots+q+1}f_k(a^2b^{-1-q}) = (-1)^k \det M_L^{(k)}
$$
(8)

for $1 \leq k \leq n-2$. For $k \in \{1,2\}$, the claim is obvious. Let $3 \leq k \leq n-2$ and assume that it holds for $k-1$ and $k-2$. Then

$$
a^{q^{k-1}+\cdots+q+1}\beta_k
$$

= $-a^{q^{k-2}+\cdots+q+1} (b^{q^{k-1}}\beta_{k-1} + a^{q^{k-2}}\beta_{k-2})$
= $-b^{q^{k-1}}a^{q^{k-2}+\cdots+q+1}\beta_{k-1} - a^{2q^{k-2}}a^{q^{k-3}+\cdots+q+1}\beta_{k-2}$
= $-b^{q^{k-1}+\cdots+q+1}f_{k-1}(a^2b^{-1-q}) - a^{2q^{k-2}}b^{q^{k-3}+\cdots+q+1}f_{k-2}(a^2b^{-1-q})$
= $b^{q^{k-1}+\cdots+q+1}(-f_{k-1}(a^2b^{-1-q}) - (a^2b^{-1-q})^{q^{k-2}}f_{k-2}(a^2b^{-1-q}))$
= $b^{q^{k-1}+\cdots+q+1}f_k(a^2b^{-1-q}),$

and

$$
a^{q^{k-1}+\cdots+q+1}\beta_k
$$

= $-b^{q^{k-1}}a^{q^{k-2}+\cdots+q+1}\beta_{k-1} - a^{2q^{k-2}}a^{q^{k-3}+\cdots+q+1}\beta_{k-2}$
= $(-1)^k b^{q^{k-1}}$ det $M_L^{(k-1)} - (-1)^k a^{2q^{k-2}}$ det $M_L^{(k-2)}$
= $(-1)^k$ det $M_L^{(k)}$,

where the last equality follows from

$$
\begin{vmatrix}\nb & a & & & & \\
a & b^{q} & \cdots & a^{q^{k-3}} & & \\
& a^{q} & \cdots & a^{q^{k-2}} & a^{q^{k-2}} & \\
& & a^{q^{k-2}} & b^{q^{k-1}} & \\
& & & a^{q^{k-2}} & b^{q^{k-1}}\n\end{vmatrix}
$$
\n
$$
= b^{q^{k-1}} \det M_L^{(k-1)} - a^{q^{k-2}} \begin{vmatrix}\nb & a & & & \\
a & b^{q} & \cdots & a^{q^{k-4}} & \\
& a^q & \cdots & a^{q^{k-3}} & \\
& & & \ddots & b^{q^{k-3}} & \\
& & & & a^{q^{k-3}} & a^{q^{k-2}}\n\end{vmatrix}
$$
\n
$$
= b^{q^{k-1}} \det M_L^{(k-1)} - a^{2q^{k-2}} \det M_L^{(k-2)}.
$$

Finally, by [\(8\)](#page-19-0) we have

$$
a^{q^{n-4}+\cdots+q+1} (a^{q^{n-3}} \beta_{n-3} + b^{q^{n-2}} \beta_{n-2})
$$

= $a^{q^{n-3}} b^{q^{n-4}+\cdots+q+1} f_{n-3} (a^2 b^{-1-q}) + a^{-q^{n-3}} b^{q^{n-2}+\cdots+q+1} f_{n-2} (a^2 b^{-1-q})$
= $a^{-q^{n-3}} b^{q^{n-2}+\cdots+q+1} ((a^2 b^{-1-q})^{q^{n-3}} f_{n-3} (a^2 b^{-1-q}) + f_{n-2} (a^2 b^{-1-q}))$
= $-a^{-q^{n-3}} b^{q^{n-2}+\cdots+q+1} f_{n-1} (a^2 b^{-1-q}),$

and

$$
a^{q^{n-3}+\cdots+q+1} (a^{q^{n-3}} \beta_{n-3} + b^{q^{n-2}} \beta_{n-2})
$$

= $(-1)^{n-1} a^{2q^{n-3}} \det M_L^{(n-3)} + (-1)^n b^{q^{n-2}} \det M_L^{(n-2)},$

with

$$
a^{q^{n-2}+\cdots+q+1} (a^{q^{n-1}} + a^{q^{n-2}} \beta_{n-2})
$$

= N(a) + N(b)(a²b^{-1-q})^{qⁿ⁻²} f_{n-2}(a²b^{-1-q})
= N(a) + (-1)ⁿa^{2qⁿ⁻²} det M_L⁽ⁿ⁻²⁾.

Given the condition in [\(7\)](#page-19-1), this completes the proof.

Corollary 11. Continue with the notation in the last proposition, and suppose that $a^2b^{-1-q} \in$ $\mathbb{F}_q.$ Define a sequence of polynomials in $\mathbb{F}_q[x]$ as

 \Box

$$
g_k(x) = (-1)^k \sum_{i=0}^{\lfloor \frac{k}{2} \rfloor} (-1)^i {k-i \choose i} x^i
$$

for $k \geq 0$. Then $\dim_{\mathbb{F}_q} \ker L = 2$ if and only if

$$
g_{n-1}(a^2b^{-1-q}) = N(a) - N(b)g_n(a^2b^{-1-q}) = 0.
$$

Proof. For $k \geq 2$, one has

$$
\begin{split}\n&(-1)^{k-1}(g_{k-1}(x)+xg_{k-2}(x)) \\
&= \sum_{i=0}^{\lfloor \frac{k-1}{2} \rfloor} (-1)^i {k-1-i \choose i} x^i - \sum_{i=0}^{\lfloor \frac{k-2}{2} \rfloor} (-1)^i {k-2-i \choose i} x^{i+1} \\
&= 1 + \sum_{i=1}^{\lfloor \frac{k-1}{2} \rfloor} (-1)^i {k-1-i \choose i} x^i + \sum_{i=1}^{\lfloor \frac{k}{2} \rfloor} (-1)^i {k-1-i \choose i-1} x^i \\
&= 1 + \sum_{i=1}^{\lfloor \frac{k-1}{2} \rfloor} (-1)^i {k-i \choose i} x^i + \sum_{i=\lfloor \frac{k-1}{2} \rfloor+1}^{\lfloor \frac{k}{2} \rfloor} (-1)^i {k-1-i \choose i-1} x^i \\
&= (-1)^k g_k(x) - \sum_{i=\lfloor \frac{k-1}{2} \rfloor+1}^{\lfloor \frac{k}{2} \rfloor} \left((-1)^i {k-i \choose i} x^i - (-1)^i {k-1-i \choose i-1} x^i\right).\n\end{split}
$$

If k is odd, then $\lfloor \frac{k-1}{2} \rfloor$ $\frac{-1}{2}$] = $\lfloor \frac{k}{2}$ $\frac{k}{2}$]; otherwise, $\lfloor \frac{k-1}{2} \rfloor$ $\frac{-1}{2}$] + 1 = $\lfloor \frac{k}{2}$ $\frac{k}{2}$] = $\frac{k}{2}$ $\frac{k}{2}$ and

$$
\binom{k-\frac{k}{2}}{\frac{k}{2}} - \binom{k-1-\frac{k}{2}}{\frac{k}{2}-1} = 1 - 1 = 0.
$$

Consequently,

$$
g_k(x) = -g_{k-1}(x) - xg_{k-2}(x)
$$
 for $k \ge 2$

with $g_0 = 1$ and $g_1 = -1$ by definition. Note that

$$
f_k(x) = -f_{k-1}(x) - x^{q^{k-2}} f_{k-2}(x) \equiv -f_{k-1}(x) - xf_{k-2}(x) \pmod{x^q - x}
$$

for $k \ge 2$ with $f_0 = 1$ and $f_1 = -1$. Therefore,

$$
f_k(x) \equiv g_k(x) \pmod{x^q - x}
$$

and $f_k(a^2b^{-1-q}) = g_k(a^2b^{-1-q}).$

Corollary 12. Let $\ell(x) = 2ax^q + bx$ and $L(x) = (ax)^{q^{n-1}} + ax^q + bx$ for $a, b \in \mathbb{F}_{q^n}^*$ with $n \geq 4$. If $\dim_{\mathbb{F}_q} \ker L = 2$, then $\epsilon = (-1)^{\frac{q+1}{2}(n-1)} \eta(N(a))$.

Proof. It follows from Proposition [10](#page-18-0) that $N(a) + (-1)^n a^{2q^{n-2}} \det M_L^{(n-2)} = 0$. Therefore,

$$
\epsilon = (-1)^{n-1} \eta((-1)^{n-1} \mathcal{N}(a)) = (-1)^{\frac{q+1}{2}(n-1)} \eta(\mathcal{N}(a)),
$$

as a result of [\(iii\)](#page-6-2) of Theorem [3.](#page-6-3)

 \Box

 \Box

Consider the binomial $\ell(x) = ax^{q^m} + bx^{q^l}$ for $a, b \in \mathbb{F}_{q^n}^*$ with positive integers m and l. It is easy to see that $\dim_{\mathbb{F}_q} \ker \ell$ is either 0 or $gcd(m-l,n)$. Denote by v_2 the 2-adic valuation of Q, and by Tr_k the trace map of $\mathbb{F}_{q^n}/\mathbb{F}_{q^k}$ for a positive divisor k of n. We have the following results concerning the composition of two linear endomorphisms of $\mathbb{F}_{q^n}/\mathbb{F}_q$.

Theorem 13. Let $\ell(x) = ax^{q^m} + bx^{q^l}$ for $a, b \in \mathbb{F}_{q^n}^*$ with $0 < l < m < \frac{n}{2}$, $d = \gcd(m, m - l)$ and $e = \text{lcm}(m + l, m - l)$. Then

- (i) $\alpha \ell(x)^{q^{n-m-l}} = \ell'(x)$ for some $\alpha \in \mathbb{F}_{q^n}^*$ if and only if $a^{q^l+1} = b^{q^m+1}$;
- (ii) if both $m+l$ and $m-l$ divide $n, a = \gamma^{q^m+1}\delta$ and $b = -\gamma^{q^l+1}\delta$ for $\gamma \in \mathbb{F}_{q^n}^*$ and $\delta \in \mathbb{F}_{q^{m-l}}^*$ such that
	- $v_2(l) \ge v_2(m-l)$ and $gcd\left(\frac{n}{e}\right)$ $(\frac{n}{e}, q) \neq 1$, or • $v_2(l) < v_2(m-l)$ and δ $q^{m-l}-1$ $\frac{q^{n-l}-1}{q^d+1}$ gcd($\frac{n}{e}, q^d+1$) = 1 with either $\delta^{\frac{q^{m-l}-1}{q^d+1}}$ $\frac{q^d+1}{r^d} \neq 1$ or $\gcd\left(\frac{n}{e}\right)$ $(\frac{n}{e}, q) \neq 1,$

then dim_{\mathbb{F}_q} ker $L = 2m$ for $L = \frac{\ell + \ell'}{2}$ $\frac{e^{\mu}}{2}$;

(iii) the converse of [\(ii\)](#page-22-0) is true provided that $a^{q^l+1} = b^{q^m+1}$.

Proof. First of all, we claim that

• if
$$
v_2(l) \ge v_2(m-l)
$$
, then $(q^m - 1) \frac{q^e - 1}{q^{m+l} - 1}$ is divisible by $q^{m-l} - 1$;

• if $v_2(l) < v_2(m-l)$, then

$$
\gcd\left((q^m-1)\frac{q^e-1}{q^{m+l}-1}, q^{m-l}-1\right)=\frac{q^{m-l}-1}{q^d+1}
$$

and

$$
\gcd\left((q^m-1)\frac{q^e-1}{q^{m+l}-1}\frac{n}{e}, q^{m-l}-1\right) = \frac{q^{m-l}-1}{q^d+1} \gcd\left(\frac{n}{e}, q^d+1\right),\,
$$

when n is divisible by e .

For $d = \gcd(m, m - l) = \gcd(l, m - l)$, if $v_2(l) \ge v_2(m - l)$, then $\gcd(m + l, m - l) =$ $gcd(2l, m-l) = d$ and

$$
(q^m - 1)\frac{q^e - 1}{q^{m+l} - 1} = \frac{q^m - 1}{q^d - 1}\frac{(q^e - 1)(q^d - 1)}{q^{m+l} - 1}
$$

is divisible by $q^{m-l}-1$. To see this, note that $(z^{m+l}-1)(z^{m-l}-1)$ divides $(z^e-1)(z^d-1)$ in $\mathbb{C}[z]$; both of them are monic with coefficients in Z, which means $\frac{(z^e-1)(z^d-1)}{(z^{m+l}-1)(z^{m-l}-1)}$ is a polynomial over Z. Assume that $v_2(l) < v_2(m - l)$. Then $gcd(m + l, m - l) = 2d$ and $\left(z^{\frac{m+l}{2d}}-1\right)\left(z^{\frac{m-l}{2d}}-1\right)$ divides $\left(z^{\frac{e}{2d}}-1\right)\left(z-1\right)$ over \mathbb{Z} . Moreover,

$$
\frac{\left(z^{\frac{e}{2d}}-1\right)(z-1)}{\left(z^{\frac{m+l}{2d}}-1\right)\left(z^{\frac{m-l}{2d}}-1\right)}=\frac{z^{\frac{e}{2d}-1}}{z^{\frac{m+l}{2d}}-1}\frac{z^{\frac{e}{2d}}}{z^{\frac{m-l}{2d}}-1}\equiv\frac{\frac{e}{2d}}{\frac{m+l}{2d}\frac{m-l}{2d}}\equiv1\pmod{z-1},
$$

with $\left(z^{\frac{e}{2d}}-1\right)(z-1)$ $\frac{\frac{z^{2a}-1}{z^{a}+1}}{\frac{z^{m+1}-1}{z^{a}-1}\left(\frac{z-a}{z-a-1}\right)}-1$ divisible by $z-1$ over \mathbb{Z} . Substitute q^{2d} for z to get

$$
\frac{(q^e - 1)(q^{2d} - 1)}{(q^{m+l} - 1)(q^{m-l} - 1)} \equiv 1 \pmod{q^{2d} - 1}.
$$
\n(9)

Now

$$
(qm - 1)\frac{qe - 1}{qm+l - 1} = \frac{qm-l - 1}{qd + 1}\frac{qm - 1}{qd - 1}\frac{(qe - 1)(q2d - 1)}{(qm+l - 1)(qm-l - 1)}.
$$

Noting that $v_2(m) = v_2(l + m - l) = \min\{v_2(l), v_2(m - l)\} = v_2(d)$ and

$$
\frac{q^m - 1}{q^d - 1} = \frac{(q^d)^{\frac{m}{d}} - 1}{q^d - 1} = \sum_{i=0}^{\frac{m}{d} - 1} q^{di} \equiv 1 \pmod{q^d + 1},
$$

with

$$
\frac{(q^e - 1)(q^{2d} - 1)}{(q^{m+l} - 1)(q^{m-l} - 1)} \equiv 1 \pmod{q^d + 1}
$$

from [\(9\)](#page-23-0), we get

$$
\gcd\left((q^m-1)\frac{q^e-1}{q^{m+l}-1}, q^{m-l}-1\right)
$$

=
$$
\frac{q^{m-l}-1}{q^d+1} \gcd\left(\frac{q^m-1}{q^d-1}\frac{(q^e-1)(q^{2d}-1)}{(q^{m+l}-1)(q^{m-l}-1)}, q^d+1\right)
$$

=
$$
\frac{q^{m-l}-1}{q^d+1}.
$$

A similar argument gives

$$
\gcd\left((q^m-1)\frac{q^e-1}{q^{m+l}-1}\frac{n}{e}, q^{m-l}-1\right) = \frac{q^{m-l}-1}{q^d+1} \gcd\left(\frac{n}{e}, q^d+1\right),\,
$$

when e divides n . The claim is now valid.

For $\alpha \in \mathbb{F}_{q^n}$, we have

$$
\alpha \ell(x)^{q^{n-m-l}} = \alpha a^{q^{n-m-l}} x^{q^{n-l}} + \alpha b^{q^{n-m-l}} x^{q^{n-m}},
$$

and

$$
\ell'(x) = (ax)^{q^{n-m}} + (bx)^{q^{n-l}}.
$$

Then $\alpha \ell(x)^{q^{n-m-l}} = \ell'(x)$ if and only if

$$
\alpha a^{q^{n-m-l}} - b^{q^{n-l}} = \alpha b^{q^{n-m-l}} - a^{q^{n-m}} = 0,
$$

which is equivalent to

$$
\alpha^{q^{m+l}} = b^{q^m} a^{-1} = a^{q^l} b^{-1},\tag{10}
$$

and [\(i\)](#page-22-1) follows. Suppose that this is the case. Let $\ell_0(x) = \alpha x^{q^{n-m-1}} + x$ so that

$$
\dim_{\mathbb{F}_q} \ker(\ell_0 \circ \ell) = \dim_{\mathbb{F}_q} (\ker \ell_0 \cap \operatorname{im} \ell) + \dim_{\mathbb{F}_q} \ker \ell
$$

\n
$$
\leq \dim_{\mathbb{F}_q} \ker \ell_0 + \dim_{\mathbb{F}_q} \ker \ell
$$

\n
$$
\leq \gcd(m + l, n) + \gcd(m - l, n)
$$

\n
$$
\leq m + l + m - l
$$

\n
$$
= 2m.
$$
\n(11)

Here, all the equalities hold if and only if ker $\ell_0 \cap \text{im } \ell = \text{ker } \ell_0$, $\dim_{\mathbb{F}_q} \text{ker } \ell_0 = m + l$ and $\dim_{\mathbb{F}_q} \ker \ell = m - l$ with both $m + l$ and $m - l$ dividing n.

Next, we show [\(iii\)](#page-22-2) with the assumption that $\alpha \ell(x)^{q^{n-m-l}} = \ell'(x)$ and $\dim_{\mathbb{F}_q} \ker(\ell_0 \circ \ell) =$ $\dim_{\mathbb{F}_q} \ker L = 2m$. Given $\dim_{\mathbb{F}_q} \ker \ell_0$, we see that $\alpha = -\beta^{q^{n-m-l}-1}$ for some $\beta \in \mathbb{F}_{q^n}^*$, and

$$
\ell_0(x) = -\beta^{q^{n-m-l}-1} x^{q^{n-m-l}} + x = -\beta^{-1} ((\beta x)^{q^{n-m-l}} - \beta x)
$$

with ker $\ell_0 = \beta^{-1} \mathbb{F}_{q^{m+l}}$. Likewise, $b = -a \gamma^{q^l - q^m}$ for some $\gamma \in \mathbb{F}_{q^n}^*$, so that

$$
\ell(x) = ax^{q^m} - a\gamma^{q^l - q^m} x^{q^l} = a\gamma^{-q^m} ((\gamma x)^{q^m} - (\gamma x)^{q^l})
$$

with $\text{im } \ell = a\gamma^{-q^m} \text{ ker } \text{Tr}_{m-l}$. It follows from [\(10\)](#page-24-0) that

$$
a^{q^l-q^m} = \frac{a^{q^l+1}}{a^{q^m+1}} = \frac{b^{q^m+1}}{a^{q^m+1}} = \gamma^{(q^l-q^m)(q^m+1)},
$$

and thus $a = \gamma^{q^m+1} \delta$ for some $\delta \in \mathbb{F}_{q^{m-l}}^*$ with $b = -a\gamma^{q^l-q^m} = -\gamma^{q^l+1} \delta$. Furthermore,

$$
-\beta^{1-q^{m+l}} = \alpha^{q^{m+l}} = b^{q^m} a^{-1} = -\gamma^{q^{m+l}-1} \delta^{q^m-1},
$$

so

$$
(\beta \gamma)^{1 - q^{m+l}} = \delta^{q^m - 1}.
$$
\n(12)

Recall that ker $\ell_0 \subseteq \text{im } \ell$, which is

$$
\beta^{-1} \mathbb{F}_{q^{m+l}} \subseteq a\gamma^{-q^m} \ker \operatorname{Tr}_{m-l} = \gamma \delta \ker \operatorname{Tr}_{m-l} = \gamma \ker \operatorname{Tr}_{m-l}.
$$

In other words, Tr_{m-l} vanishes on $(\beta \gamma)^{-1} \mathbb{F}_{q^{m+l}}$. Clearly, $e = \text{lcm}(m+l, m-l)$ divides n and

$$
\sum_{i=0}^{\frac{e}{m-l}-1} (\text{Tr}_e((\beta \gamma)^{-1})x)^{q^{(m-l)i}}
$$
\n
$$
\equiv \sum_{i=0}^{\frac{e}{m-l}-1} \text{Tr}_e((\beta \gamma)^{-1}x)^{q^{(m-l)i}}
$$
\n
$$
\equiv \text{Tr}_{m-l}((\beta \gamma)^{-1}x)
$$
\n
$$
\equiv 0 \pmod{x^{q^{m+l}} - x}.
$$
\n(13)

Here $x^{q^{(m-l)i}} \equiv x \pmod{x^{q^{m+l}} - x}$ if and only if $(m-l)i \equiv 0 \pmod{m+l}$; i.e., $i \equiv 0$ (mod $\frac{e}{m-l}$). Then $\text{Tr}_e((\beta \gamma)^{-1})=0$, which, combined with [\(12\)](#page-24-1), leads to

$$
0 = \sum_{i=0}^{\frac{n}{e}-1} (\beta \gamma)^{-q^{ei}} = (\beta \gamma)^{-1} \sum_{i=0}^{\frac{n}{e}-1} (\beta \gamma)^{1-q^{ei}} = (\beta \gamma)^{-1} \sum_{i=0}^{\frac{n}{e}-1} (\delta^{q^{m}-1})^{\frac{q^{ei}-1}{q^{m+1}-1}}.
$$

Let $\theta = \delta^{(q^m - 1)} \frac{q^e - 1}{q^{m+l}}$ $\frac{q^{m+l}-1}{q^m}$, which belongs to $\mathbb{F}_{q^e}^*$, so that

$$
\sum_{i=0}^{\frac{n}{e}-1} \theta^i = \sum_{i=0}^{\frac{n}{e}-1} \theta^{\frac{q^{ei}-1}{q^e-1}} = 0.
$$
 (14)

Then $\theta^{\frac{n}{e}} = 1$; in addition, $\theta \neq 1$ when gcd $\left(\frac{n}{e}\right)$ $\left(\frac{n}{e}, q\right) = 1$. If $v_2(l) \ge v_2(m-l)$, then $\theta = 1$ by the claim, and necessarily gcd $\left(\frac{n}{e}\right)$ $(\frac{n}{e}, q) \neq 1$. Otherwise, $\theta = 1$ if and only if δ $q^{m-l}-1$ $\overline{q^{d+1}} = 1$, and $\theta^{\frac{n}{e}} = 1$ if and only if δ $q^{m-l}-1$ $\frac{a^{n-i}-1}{q^d+1}$ gcd $(\frac{n}{e},q^d+1)$ = 1. This proves [\(iii\).](#page-22-2)

Assume the conditions in [\(ii\).](#page-22-0) Then

$$
\ell(x) = ax^{q^m} + bx^{q^l} = \gamma \delta((\gamma x)^{q^m} - (\gamma x)^{q^l}),
$$

with $\dim_{\mathbb{F}_q} \ker \ell = m - l$ and $\operatorname{im} \ell = \gamma \ker \mathrm{Tr}_{m-l}$. It is straightforward to verify that $a^{q^l+1} = b^{q^m+1}$, and $\alpha \ell(x)^{q^{n-m-l}} = \ell'(x)$ with

$$
\alpha^{q^{m+l}} = b^{q^m} a^{-1} = -\gamma^{q^{m+l}-1} \delta^{q^m-1}
$$

from [\(10\)](#page-24-0). Now [\(11\)](#page-24-2) applies and it remains to show that $\dim_{\mathbb{F}_q} \ker \ell_0 = m + l$ with $\ker \ell_0 \subseteq \text{im } \ell, \text{ where } \ell_0(x) = \alpha x^{q^{n-m-l}} + x. \text{ For } \theta = \delta^{(q^m-1)\frac{q^e-1}{q^{m+l}}}.$ q^{m+l-1} , the condition on δ together with the claim implies $\sum_{i=0}^{\frac{n}{e}-1} \theta^i = 0$, so

$$
\left(-\alpha\right)^{q^{m+l}\frac{q^n-1}{q^{m+l}-1}} = \left(\gamma^{q^{m+l}-1}\delta^{q^m-1}\right)^{\frac{q^n-1}{q^{m+l}-1}} = \delta^{\left(q^m-1\right)\frac{q^e-1}{q^{m+l}-1}\frac{q^n-1}{q^e-1}} = \theta^{\frac{n}{e}} = 1.
$$

Then $\alpha = -\beta^{q^{n-m-l}-1}$ for some $\beta \in \mathbb{F}_{q^n}^*$, so that $(\beta \gamma)^{1-q^{m+l}} = \delta^{q^m-1}$ and ker $\ell_0 =$ $\beta^{-1}\mathbb{F}_{q^{m+l}}, \text{ with}$

$$
\mathrm{Tr}_e((\beta\gamma)^{-1}) = (\beta\gamma)^{-1} \sum_{i=0}^{\frac{n}{e}-1} (\delta^{q^{m}-1})^{\frac{q^{ei}-1}{q^{m+l}-1}} = (\beta\gamma)^{-1} \sum_{i=0}^{\frac{n}{e}-1} \theta^i = 0.
$$

By the same argument as in [\(13\)](#page-25-0), we have ker $\ell_0 \subseteq \text{im } \ell$ and establish [\(ii\).](#page-22-0)

Recall that

$$
\sum_{w \in \mathbb{F}_{q^n}} \chi(\text{Tr}(w\ell(w))) = \epsilon \mathcal{G}^r q^{n-r},
$$

where r is the rank of the quadratic form, χ is the canonical additive character of \mathbb{F}_q and G is the quadratic Gauss sum of \mathbb{F}_q . The quadratic form from one of the polynomials in the above theorem can be further investigated.

Proposition 14. Let $\ell(x) = \gamma^{q^m+1} \delta x^{q^m} - \gamma^{q^l+1} \delta x^{q^l}$ for $\gamma \in \mathbb{F}_{q^m}^*$ and $\delta \in \mathbb{F}_{q^{m-l}}^*$ with $0 < l < l$ $m < \frac{n}{2}$, $v_2(l) \ge v_2(m-l)$, n divisible by both $m+l$ and $m-l$, and $\gcd\left(\frac{n}{\text{lcm}(m+l)}\right)$ $\frac{n}{\text{lcm}(m+l,m-l)}, q \neq 1.$ Then $\epsilon = (-1)^{n-1+\frac{q-1}{2}(\frac{n}{d}-1)m} \eta(N(2\delta))$ for $d = \gcd(m, m-l)$.

Proof. We show that there exists some $c \in \mathbb{F}_{q^{m-l}}^*$ with $\delta c^{q^m+1} \in \mathbb{F}_{q^d}$, so that

$$
x\ell(x) = \gamma^{q^m+1} \delta c^{q^m+1} (c^{-1}x)^{q^m+1} - \gamma^{q^l+1} \delta c^{q^l+1} (c^{-1}x)^{q^l+1}
$$

=
$$
\delta c^{q^m+1} ((\gamma c^{-1}x)^{q^m+1} - (\gamma c^{-1}x)^{q^l+1}).
$$
 (15)

Note that $gcd(q^m + 1, q^{m-l} - 1)$ divides

$$
\gcd(q^{2m}-1, q^{m-l}-1) = q^{\gcd(2m, m-l)} - 1 = q^d - 1,
$$

so $\gcd(q^m + 1, q^{m-l} - 1)$ divides $\gcd(q^m + 1, q^d - 1) = 2$ as $q^m + 1 \equiv 2 \pmod{q^d - 1}$. Then

$$
\gcd(q^m + 1, q^{m-l} - 1) = 2,
$$

and it suffices to find an element $c_0 \in \mathbb{F}_{q^{m-l}}^*$ such that $\delta c_0^2 \in \mathbb{F}_{q^d}$. Take $\alpha \in \mathbb{F}_q^*$ $_{q^d}^*$ that is a nonsquare element in $\mathbb{F}_{q^{m-l}}$, so that one may let either $c_0^{-2} = \delta$ or $c_0^{-2} = \delta \alpha$ to get δc_0^2 lying in \mathbb{F}_{q^d} . Such α exists in \mathbb{F}_q^* $_{q^{d}}^{*}$ since α $q^{m-l}-1$ $\frac{c^{l-1}-1}{2} = \alpha^{\frac{q^d-1}{2}\frac{m-l}{d}}$, while $v_2(d) = \min\{v_2(l), v_2(m-l)\}$ $v_2(m-l)$ and $\frac{m-l}{d}$ is odd.

For the canonical additive character χ_d of \mathbb{F}_{q^d} , it follows from [\(15\)](#page-26-0) that

$$
\epsilon \mathcal{G}^{n-2m} q^{2m} = \sum_{w \in \mathbb{F}_{q^n}} \chi(\text{Tr}(w\ell(w))) = \sum_{w \in \mathbb{F}_{q^n}} \chi_d(\text{Tr}_d(\delta c^{q^m+1} w(w^{q^m} - w^{q^l}))).
$$

 \Box

Now $\text{Tr}_d\big(\delta c^{q^m+1}x\big(x^{q^m}-x^{q^l}\big)\big)$ is a quadratic form of $\mathbb{F}_{q^n}/\mathbb{F}_{q^d}$ with coefficients in \mathbb{F}_{q^d} . Taking the absolute value into account, we have

$$
\sum_{w \in \mathbb{F}_{q^n}} \chi_d \big(\text{Tr}_d \big(\delta c^{q^m+1} w \big(w^{q^m} - w^{q^l} \big) \big) \big) = \epsilon_d \mathcal{G}_d^{\frac{n-2m}{d}} q^{2m},
$$

where $\mathcal{G}_d = (-1)^{d-1} \mathcal{G}^d$ ([\[8,](#page-30-8) Theorem 5.15]) and

$$
\epsilon_d = (-1)^{\left(\frac{n}{d} - 1\right)\left(\frac{q^d - 1}{2}\frac{m}{d} + 1\right)} \eta\left(\mathcal{N}\left(2\delta c^{q^m + 1}\right)\right) = (-1)^{\left(\frac{n}{d} - 1\right)\left(\frac{q - 1}{2}m + 1\right)} \eta\left(\mathcal{N}\left(2\delta\right)\right),
$$

as a consequence of [\(iv\)](#page-14-2) in Theorem [8.](#page-14-3) Altogether, it gives

$$
\epsilon \mathcal{G}^{n-2m} q^{2m}
$$

= $(-1)^{\left(\frac{n}{d}-1\right)\left(\frac{q-1}{2}m+1\right)} \eta(N(2\delta))((-1)^{d-1} \mathcal{G}^d)^{\frac{n-2m}{d}} q^{2m}$
= $(-1)^{n-1+\frac{q-1}{2}\left(\frac{n}{d}-1\right)m} \eta(N(2\delta)) \mathcal{G}^{n-2m} q^{2m},$

 \Box

which completes the proof.

So far, we have been studying Artin-Schreier curves of the form $y^q - y - x\ell(x)$ from quadratic forms of $\mathbb{F}_{q^n}/\mathbb{F}_q$. One may be interested in curves defined by $y^{q^k} - y - x\ell(x)$ for some positive divisor k of n, even if $\text{Tr}_k(x\ell(x))$ is not a quadratic form of $\mathbb{F}_{q^n}/\mathbb{F}_{q^k}$. Here is an example.

Corollary 15. Continue with the notation and assumptions in the last proposition. For a positive divisor k of $m - l$, the number of rational points of the curve over \mathbb{F}_{q^n} defined by $y^{q^k} - y - x\ell(x)$ is

$$
\begin{cases} q^n + 1 - (-1)^{\frac{(q-1)n}{4}} \eta(N(\delta))(q^k - 1)q^{\frac{n}{2} + m} & \text{if } \frac{n}{k} \text{ is even,} \\ q^n + 1 & \text{if } \frac{n}{k} \text{ is odd.} \end{cases}
$$

It is a maximal or minimal curve if $\frac{n}{k}$ is even.

Proof. As is easily checked, the curve has genus $\frac{(q^k-1)q^m}{2}$ $\frac{1}{2}$, with exactly one point at infinity. The number of its other rational points over \mathbb{F}_{q^n} is

$$
\sum_{w \in \mathbb{F}_{q^n}} \sum_{u \in \mathbb{F}_{q^k}} \chi_k(u \text{Tr}_k(w\ell(w)))
$$
\n
$$
= \sum_{u \in \mathbb{F}_{q^k}} \sum_{w \in \mathbb{F}_{q^n}} \chi(\text{Tr}(uw\ell(w)))
$$
\n
$$
= q^n + \sum_{u \in \mathbb{F}_{q^k}^*} (-1)^{n-1+\frac{q-1}{2}(\frac{n}{d}-1)m} \eta(N(2u\delta)) \mathcal{G}^{n-2m} q^{2m},
$$

where $\mathcal{G}^{n-2m} = (-1)^{\frac{(q-1)(n-2m)}{4}} q^{\frac{n}{2} - m}$ when *n* is even, and

$$
\sum_{u \in \mathbb{F}_{q^k}^*} \eta(\mathcal{N}(2u)) = \begin{cases} q^k - 1 & \text{if } \frac{n}{k} \text{ is even,} \\ 0 & \text{if } \frac{n}{k} \text{ is odd.} \end{cases}
$$

A simple calculation yields the result.

As a final remark, we introduce an alternative approach towards maximal and minimal Artin-Schreier curves.

Proposition 16. Let L be a q-linear polynomial over \mathbb{F}_{q^n} of degree less than q^n . Then $L = L'$ with $\dim_{\mathbb{F}_q} \text{im } L = r$ for some positive integer r if and only if

$$
L(x) = \sum_{i=1}^{r} d_i \alpha_i \text{Tr}(\alpha_i x)
$$

for some $d_i \in \mathbb{F}_q^*$ and $\alpha_1, \ldots, \alpha_r$ that are linearly independent in $\mathbb{F}_{q^n}/\mathbb{F}_q$. In this case, $\epsilon = \eta(d_1 \cdots d_r)$ for $L = \frac{\ell + \ell'}{2}$ $\frac{e^{\epsilon}}{2}$.

Proof. The sufficiency is obvious. If L has the above form, then

$$
\operatorname{Tr}(x\ell(x)) = \operatorname{Tr}(xL(x)) \equiv \sum_{i=1}^r d_i \operatorname{Tr}(\alpha_i x)^2 \pmod{x^{q^n} - x},
$$

where $(\text{Tr}(\alpha_1 x), \dots, \text{Tr}(\alpha_r x))$ maps \mathbb{F}_{q^n} onto \mathbb{F}_q^r , and thus $\epsilon = \eta(d_1 \cdots d_r)$ by definition.

Conversely, suppose that $L = L'$ and $\dim_{\mathbb{F}_q} \text{im } L = r$. Let β_1, \dots, β_r be a basis of im L over \mathbb{F}_q . Then there exist $\alpha_1, \ldots, \alpha_r \in \mathbb{F}_{q^n}$ such that

$$
L(x) = \beta_1 \text{Tr}(\alpha_1 x) + \cdots + \beta_r \text{Tr}(\alpha_r x),
$$

for every linear form of $\mathbb{F}_{q^n}/\mathbb{F}_q$ is $\text{Tr}(\alpha x)$ for some $\alpha \in \mathbb{F}_{q^n}$. Let W be the subspace of $\mathbb{F}_{q^n}/\mathbb{F}_q$ spanned by $\alpha_1, \ldots, \alpha_r$. Clearly, ker $L = W^{\perp}$ and then $\dim_{\mathbb{F}_q} W = r$, so $\alpha_1, \ldots, \alpha_r$ is a basis of $(\ker L)^{\perp}$ over \mathbb{F}_q , where $(\ker L)^{\perp} = (\ker L')^{\perp} = \mathrm{im} L$. Hence, there exists a nonsingular matrix C over \mathbb{F}_q such that

$$
(\beta_1 \cdots \beta_r) = (\alpha_1 \cdots \alpha_r) C.
$$

Then

$$
L(x) = (\alpha_1 \quad \cdots \quad \alpha_r) C \begin{pmatrix} Tr(\alpha_1 x) \\ \vdots \\ Tr(\alpha_r x) \end{pmatrix}.
$$

 \Box

Observe that

$$
L'(x) = (\alpha_1 \quad \cdots \quad \alpha_r) C^{\mathrm{T}} \begin{pmatrix} \text{Tr}(\alpha_1 x) \\ \vdots \\ \text{Tr}(\alpha_r x) \end{pmatrix},
$$

and there exist $x_1, \ldots, x_r \in \mathbb{F}_{q^n}$ such that

$$
\text{Tr}(\alpha_i x_j) = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}
$$

Since $L(x_j) = L'(x_j)$ for $j = 1, ..., n$, one has

$$
(\alpha_1 \cdots \alpha_r) C = (\alpha_1 \cdots \alpha_r) C^{\mathrm{T}}.
$$

This implies $C = C^T$, and $C = P^TDP$ for a nonsingular matrix P over \mathbb{F}_q and a diagonal matrix with nonzero entries $d_1, \ldots, d_r \in \mathbb{F}_q$. As a result,

$$
L(x) = (\alpha_1 \quad \cdots \quad \alpha_r) P^{\mathrm{T}}DP \begin{pmatrix} \text{Tr}(\alpha_1 x) \\ \vdots \\ \text{Tr}(\alpha_r x) \end{pmatrix}.
$$

Substituting $(\alpha_1 \cdots \alpha_r)$ for $(\alpha_1 \cdots \alpha_r) P^T$, we get the desired basis of im L. \Box

As we have seen, if $L = \frac{\ell + \ell'}{2}$ $\frac{f^{(1)}f^{(2)}}{2}$ for a q-linear polynomial ℓ over \mathbb{F}_{q^n} of degree less than q^n , then $L = L'$, and vice versa. Now, assume that n is even. Except the trivial cases mentioned at the beginning of this section, every maximal or minimal curve over \mathbb{F}_{q^n} defined by $y^q - y - x\ell(x)$ arises from

$$
L(x) = \sum_{i=1}^{r} d_i \alpha_i \text{Tr}(\alpha_i x) = \sum_{i=0}^{n-1} a_i x^{q^i},
$$

where r is even with $d_i \in \mathbb{F}_q^*$ and $\alpha_1, \ldots, \alpha_r$ linearly independent in $\mathbb{F}_{q^n}/\mathbb{F}_q$, such that $a_{\frac{n-r}{2}+1}, a_{\frac{n-r}{2}+2}, \ldots, a_{\frac{n}{2}}$ are all zero. Then we may take the part of degree $q^{\frac{n-r}{2}}$ in L. Whether it is maximal or minimal depends on $(-1)^{\frac{(q-1)r}{4}}\eta(d_1\cdots d_r)$. For example, let

$$
L(x) = d_1 \alpha_1 \text{Tr}(\alpha_1 x) + d_2 \alpha_2 \text{Tr}(\alpha_2 x) = \sum_{i=0}^{n-1} \left(d_1 \alpha_1^{q^i+1} + d_2 \alpha_2^{q^i+1} \right) x^{q^i},
$$

with $d_1, d_2 \in \mathbb{F}_q^*$, $\alpha_1 \alpha_2^{-1} \notin \mathbb{F}_q$ and $d_1 \alpha_1^{q^{\frac{n}{2}}+1} + d_2 \alpha_2^{q^{\frac{n}{2}}+1} = 0$, and let

$$
\ell(x) = 2 \sum_{i=0}^{\frac{n}{2}-1} \left(d_1 \alpha_1^{q^i+1} + d_2 \alpha_2^{q^i+1} \right) x^{q^i},
$$

so that $L = \frac{\ell + \ell'}{2}$ $\frac{1}{2}e^{\frac{\mu}{2}}$. Here $\deg \ell = q^{\frac{n}{2}-1}$ since $L(x)$, $q^{\frac{n}{2}-1}$ is congruent to a polynomial of degree at most q^{n-2} modulo $x^{q^n} - x$, while $\dim_{\mathbb{F}_q} \ker L = n-2$. That describes all the maximal and minimal Artin-Schreier curves from quadratic forms of $\mathbb{F}_{q^n}/\mathbb{F}_q$ of rank 2 (cf. [\[1,](#page-30-3) Theorem 5.5] and [\[2\]](#page-30-5)).

Acknowledgement

The author would like to express his sincere gratitude to Prof. Sihem Mesnager for her advice.

References

- [1] N. Anbar and W. Meidl. Quadratic functions and maximal Artin–Schreier curves. Finite Fields and Their Applications, 30:49–71, 2014.
- [2] D. Bartoli, L. Quoos, Z. Saygı, and E. S. Yılmaz. Explicit maximal and minimal curves of Artin–Schreier type from quadratic forms. Applicable Algebra in Engineering, Communication and Computing, 32(4):507–520, 2021.
- [3] E. Çakçak and F. Özbudak. Curves related to Coulter's maximal curves. Finite Fields and Their Applications, 14(1):209–220, 2008.
- [4] R. Chen and S. Mesnager. Evaluation of weil sums for some polynomials and associated quadratic forms. Cryptography and Communications, 15(3):661–673, 2023.
- [5] R. S. Coulter. The number of rational points of a class of Artin–Schreier curves. Finite Fields and Their Applications, 8(4):397–413, 2002.
- [6] R. W. Fitzgerald and Y. Kottegoda. Invariants of trace forms in odd characteristic and authentication codes. Finite Fields and Their Applications, 48:1–9, 2017.
- [7] A. Klapper. Cross-correlations of quadratic form sequences in odd characteristic. Designs, Codes and Cryptography, 11(3):289–305, 1997.
- [8] R. Lidl and H. Niederreiter. Finite fields. Cambridge University Press, 1997.
- [9] D. Oliveira and F. B. Martínez. Artin-Schreier curves given by \mathbb{F}_q -linearized polynomials. Discrete Mathematics, 346(12):113630, 2023.
- [10] J. A. Oliveira, H. Borges, and F. B. Mart´ınez. On the number of rational points on Artin-Schreier hypersurfaces. Finite Fields and Their Applications, 90:102229, 2023.
- [11] F. Özbudak, E. Saygı, and Z. Saygı. Quadratic forms of codimension 2 over certain finite fields of even characteristic. Cryptography and Communications, 3(4):241–257, 2011.
- [12] F. Özbudak, E. Saygı, and Z. Saygı. Quadratic forms of codimension 2 over finite fields containing \mathbb{F}_4 and Artin–Schreier type curves. Finite Fields and Their Applications, 18(2):396–433, 2012.
- [13] F. Özbudak and Z. Saygı. Explicit maximal and minimal curves over finite fields of odd characteristics. Finite Fields and Their Applications, 42:81–92, 2016.
- [14] H. Stichtenoth. Algebraic function fields and codes. Springer Science & Business Media, 2009.