

Gravity and Unification: Insights from $SL(2N, C)$ Gauge Symmetries

J. L. Chkareuli^{1,2}

¹*Institute of Theoretical Physics, Ilia State University, 0162 Tbilisi, Georgia*

²*Andronikashvili Institute of Physics, Tbilisi State University, 0177 Tbilisi, Georgia*

Abstract

The perspective that gravity governs the unification of all known elementary forces calls for an extension of the gauge gravity symmetry group $SL(2, C)$ to the broader local symmetry $SL(2N, C)$, where N reflects the internal $SU(N)$ symmetry subgroup. This extension is shown to lead to a consistent hyperunification framework, provided that the tetrad fields of $SL(2, C)$ retain their invertibility condition in the extended theory, thus maintaining their connection to gravity. As a result, while the full gauge multiplet of $SL(2N, C)$ typically comprises vector, axial-vector, and tensor field submultiplets of $SU(N)$, only the vector submultiplet and the singlet tensor field manifest in the observed particle spectrum. The axial-vector submultiplet remains decoupled from ordinary matter, while the tensor submultiplet acquires the Planck scale order masses. Consequently, the effective symmetry of the theory reduces to $SL(2, C) \times SU(N)$, bringing together $SL(2, C)$ gauge gravity and $SU(N)$ grand unification. As all states in $SL(2N, C)$ are also classified by their spin, some $SU(N)$ grand unified models, including the standard $SU(5)$, appear unsuitable for the standard spin-1/2 quarks and leptons. However, applying $SL(2N, C)$ symmetry to a model of composite quarks and leptons, where constituent chiral preons form the fundamental representations, identifies $SL(16, C)$ with its effective $SL(2, C) \times SU(8)$ symmetry accommodating all three quark-lepton families, as the most compelling candidate for hyperunification of the existing fundamental forces.

1 Introduction

A deepening recognition has long emerged that gravity exhibits notable similarities to the other three fundamental forces when examined within the framework of conventional gauge theory [1, 2, 3]. Specifically, the spin-connection fields that gauge the Lorentz symmetry arise much like photons and gluons in the Standard Model. This opens the intriguing possibility that these spin-connections could be unified with the Standard Model gauge bosons within some non-compact symmetry group, potentially leading to a unification of all known gauge forces. If we accept that gravity itself governs the symmetry structure of this unification, then the search for such a symmetry naturally extends the gauge gravity group $SL(2, C)$ to the broader local symmetry $SL(2N, C)$, where N defines the degree of the internal $SU(N)$ subgroup. We focus on $SL(2, C)$ and its extensions rather than the more conventional $SO(1, 3)$, as $SL(2, C)$ more fundamentally captures the spinorial nature of fermions in spacetime. In this context, we refer to such frameworks as hyperunified theories (HUTs), with particular emphasis on the $SL(2N, C)$ model examined here in detail. This unification integrates $SL(2, C)$ gauge gravity with $SU(N)$ grand unified theory (GUT). The $SU(N)$ subgroup within $SL(2N, C)$ is designated as a "hyperflavor" symmetry, with fields associated with its representations termed "hyperflavored" fields, contrasting with the "neutral" spin-connection field of local $SL(2, C)$ symmetry, which corresponds to the graviton. Remarkably, the total gauge hypermultiplet of the $SL(2N, C)$ HUT may encompass both spin-1 and spin-2 fields, offering a potential unified framework for describing all fundamental forces – including gravity – should some key challenges, typically encountered in such theories, be successfully navigated.

Numerous models in the literature propose unifying gravity with other interactions by merging local Lorentz and internal symmetries into a non-compact covering group [4, 5, 6, 7]. Their difficulties are well known and, to varying degrees, they generally appear in the $SL(2N, C)$ HUT as well [8]. Firstly, the vector fields in the total gauge hypermultiplet of this group are always accompanied by the axial-vector fields which must be somehow excluded from the theory as there is no direct indication of their existence. Then, while vector fields are proposed to mediate ordinary gauge interactions, the tensor fields from the same hypermultiplet must provide the subtle gravitational interactions required to align with reality. The crucial point lies in the fact that, whereas in pure gravity case, one can solely consider the action being linear in the curvature (R) constructed from a tensor field, the unification with other interactions necessitates the inclusion of quadratic curvature (R^2) terms as well. Consequently, tensor fields in these terms will induce interactions comparable to those of the gauge vector fields in the Standard Model. Moreover, the tensor fields, akin to the vector ones, exhibit now the internal $SU(N)$ symmetry features implying the existence of the multiplet of hyperflavored gravitons rather than a single neutral one. Apart from that, such $R + R^2$ Lagrangians for gravity are generally known to contain ghosts and tachyons rendering them essentially unstable. And lastly, but perhaps most importantly, a potential pitfall in hyperunified theories stems from the Coleman-Mandula theorem [9] concerning the impossibility of merging spacetime and internal symmetries. It is worth noting that this theorem initially surfaced precisely in connection with one of the special cases of $SL(2N, C)$, specifically the $SL(6, C)$ symmetry [10], used a long time

ago as a possible relativistic version of the global $SU(6)$ symmetry model describing the spin-unitary spin symmetry classification of mesons and baryons [11].

In contrast, we aim to demonstrate here how, in the $SL(2N, C)$ HUT framework, these difficulties can be naturally overcome in the way as yet unexplored. The key idea is that the interactions of the extended gauge multiplet I_μ in the $SL(2N, C)$ theory – comprising generally the vector, axial-vector and tensor field submultiplets – are suitably shaped by the associated tetrad fields. Crucially, these tetrads are assumed to retain their form in the extended theory, continuing to satisfy the invertibility condition as they do in pure $SL(2, C)$ gravity. Consequently, from the entire gauge multiplet I_μ only the $SU(N)$ hyperflavor-charged vector fields and hyperflavor-neutral tensor field emerge in the observed particle spectrum. The axial-vector fields remain sterile to ordinary matter, while the hyperflavor-charged tensor fields acquire the Planck scale order masses through the linear curvature terms so as to remain only with the effective $SL(2, C) \times SU(N)$ symmetry in the theory. In this sense, the tetrads not only define the geometric structure of spacetime but also govern which local internal symmetries and associated gauge field interactions are really operative in it.

Next, the extension of $SL(2, C)$ gauge gravity to $SL(2N, C)$ naturally calls for the inclusion of some "safe" quadratic curvature terms in the gravitational sector of the extended theory, alongside the standard quadratic strength-tensor terms for the vector fields. This requirement uniquely identifies, from among all possible candidates, the ghost-free curvature-squared gravity Lagrangian initially proposed by Neville [12] (see also [13]), as the most appropriate model for such an extension. Consequently, the resulting theory, after reduction, includes the $SL(2, C)$ symmetric $R + R^2$ Einstein-Cartan type gravity action, which remains free from ghosts and tachyons, in addition to the conventional $SU(N)$ gauge vector field theory regarded as the GUT candidate. Eventually, the theory effectively manifests a local $SL(2, C) \times SU(N)$ symmetry, rather than an entire $SL(2N, C)$ symmetry, which solely serves to determine the structure of the gauge and matter multiplets. In this way, the restrictions imposed by the Coleman-Mandula theorem appear naturally circumvented.

The paper is organized as follows. Section 2 revisits the $SL(2, C)$ gauge gravity in an updated form. Section 3 presents the $SL(2N, C)$ HUT, introducing an invertibility constraint on the tetrads, which guides its reduction to the effective $SL(2, C) \times SU(N)$ symmetry framework detailed in the subsequent sections. Section 4 explores the hyperunified linear and quadratic curvature Lagrangians for gravity and other fundamental forces. Section 5 highlights specific HUT models with a particular focus on the $SL(16, C)$ theory that gives rise to the $SU(8)$ GUT, accommodating all three families of composite quarks and leptons. The paper concludes with a summary in Section 6.

2 $SL(2, C)$ gauge gravity

We begin by presenting the $SL(2, C)$ gravity model, drawing in part from the pioneering work [3]. At any point in spacetime, we assume a local frame where the global $SL(2, C)$

symmetry group acts. Under this symmetry, the fundamental fermions transform as

$$\Psi \rightarrow \Omega\Psi, \quad \Omega = \exp\left\{\frac{i}{4}\theta_{ab}\gamma^{ab}\right\} \quad (1)$$

where the matrix Ω satisfies a pseudounitariness condition, $\Omega^{-1} = \gamma_0\Omega^\dagger\gamma_0$, with the transformation parameters θ_{ab} taken to be constant for now. To maintain the invariance of the kinetic terms, $i\bar{\Psi}\gamma^\mu\partial_\mu\Psi$, the gamma matrices must be replaced by a set of tetrad matrices e^μ which transform like

$$e^\mu \rightarrow \Omega e^\mu \Omega^{-1} \quad (2)$$

In general, the tetrad matrices e^μ and their conjugates e_μ incorporate the appropriate tetrad fields e_a^μ and e_μ^a , respectively,

$$e^\mu = e_a^\mu \gamma^a, \quad e_\mu = e_\mu^a \gamma_a \quad (3)$$

which transforms infinitesimally as

$$\delta e^{\mu c} = \frac{1}{2}\theta_{ab}(e^{\mu a}\eta^{bc} - e^{\mu b}\eta^{ac}) \quad (4)$$

They, as usual, satisfy the orthonormality or invertibility conditions

$$e_\mu^a e_a^\nu = \delta_\mu^\nu, \quad e_\mu^a e_b^\mu = \delta_b^a \quad (5)$$

and determine the metric tensors in the theory

$$g_{\mu\nu} = \frac{1}{4}\text{Tr}(e_\mu e_\nu) = e_\mu^a e_\nu^b \eta_{ab}, \quad g^{\mu\nu} = \frac{1}{4}\text{Tr}(e^\mu e^\nu) = e_a^\mu e_b^\nu \eta^{ab} \quad (6)$$

Turning now to the case where the $SL(2, C)$ transformations (1) become local, $\theta_{ab} \equiv \theta_{ab}(x)$, one must introduce the spin-connection gauge field multiplet I_μ which transforms as usual

$$I_\mu \rightarrow \Omega I_\mu \Omega^{-1} - \frac{1}{ig}(\partial_\mu \Omega)\Omega^{-1} \quad (7)$$

This defines the covariant derivative for the fermion field

$$\partial_\mu \Psi \rightarrow D_\mu \Psi = \partial_\mu \Psi + ig I_\mu \Psi \quad (8)$$

where g is the gauge coupling constant. The multiplet I_μ gauging the $SL(2, C)$ takes the following form by definition

$$I_\mu = \frac{1}{4}T_{\mu[ab]}\gamma^{ab} \quad (9)$$

where the flat spacetime tensor field components $T_{\mu[ab]}$ transform according to

$$\delta T_\mu^{[ab]} = \frac{1}{2}\theta_{[cd]}[(T_\mu^{[ac]}\eta^{bd} - T_\mu^{[ad]}\eta^{bc}) - (T_\mu^{[bc]}\eta^{ad} - T_\mu^{[bd]}\eta^{ac})] - \frac{1}{g}\partial_\mu\theta^{[ab]} \quad (10)$$

The tensor field $T_{\mu[ab]}$ may in principle propagate, while the tetrad e^μ is not treated as a dynamical field. The invariant Lagrangian built from the tensor field strength

$$T_{\mu\nu}^{[ab]} = \partial_{[\nu} T_{\mu]}^{[ab]} + g\eta_{cd} T_{[\mu}^{[ac]} T_{\nu]}^{[bd]} \quad (11)$$

can be written in a conventional form

$$e\mathcal{L}_G = \frac{1}{2\kappa} e_{[a}^\mu e_{b]}^\nu T_{\mu\nu}^{[ab]}, \quad e \equiv [-\det Tr(e^\mu e^\nu)/4]^{-1/2} \quad (12)$$

where κ stands for the modified Newtonian constant $8\pi/M_{Pl}^2$. This form arises after using the commutator for tetrads and some standard relations for γ matrices¹. In fact, this is the simplest pure gravity Lagrangian in the Palatini-type formulation. Its variation with respect to the tensor field imposes a constraint that expresses the tensor field in terms of the tetrads and their derivatives, reducing $e\mathcal{L}_G$ to the standard Einstein Lagrangian. The factor e , while not relevant for $SL(2, C)$ gauge invariance, introduces an extra invariance of the action under general four-coordinate transformations of $GL(4, R)$ [3].

In the presence of fermions, the gauge-invariant matter coupling, expressed via the covariant derivative (8, 9) leads to an additional interaction between the tensor field and the spin-current density

$$e\mathcal{L}_M = -\frac{1}{2} g\epsilon^{abcd} T_{\mu[ab]} e_c^\mu \bar{\Psi} \gamma_d \gamma_5 \Psi \quad (13)$$

This is a key feature of the Einstein-Cartan type gravity [2] resulting, beyond standard General Relativity, in the tiny four-fermion (spin current-current) interaction in the matter sector

$$\kappa (\bar{\Psi} \gamma_d \gamma_5 \Psi) (\bar{\Psi} \gamma^d \gamma_5 \Psi) \quad (14)$$

3 Extending gravity: gauge $SL(2N, C)$ theories

The view that gravity governs the symmetry structure underlying the unification of all known elementary forces necessitates extending the gauge gravity symmetry group $SL(2, C)$ to the broader local symmetry $SL(2N, C)$, where N defines the degree of the internal $SU(N)$ symmetry as a subgroup. We refer to this as hyperflavor symmetry, which encompasses all known quantum numbers associated with quarks and leptons – such as color, weak isospin, and family numbers. Indeed, this hyperflavor $SU(N)$ symmetry underpins the grand unification of the three other fundamental forces acting on quarks and leptons. By proceeding along these lines, we aim to explore the new insights that may arise from such a formulation.

¹We give here some of them used throughout the paper

$$\begin{aligned} \gamma^{ab} &= i[\gamma^a, \gamma^b]/2, \quad \gamma^a \gamma^b = \gamma^{ab}/i + \eta^{ab} \hat{1}, \quad \gamma_c \gamma^{[ab]} \gamma^c = 0 \\ [\gamma^{ab}, \gamma^{a'b'}] &= 2i(\eta^{ab'} \gamma^{ba'} + \eta^{ba'} \gamma^{ab'} - \eta^{aa'} \gamma^{bb'} - \eta^{bb'} \gamma^{aa'}) \\ Tr(\gamma^{ab} \gamma^{a'b'}) &= 4(\eta^{aa'} \eta^{bb'} - \eta^{ab'} \eta^{ba'}), \quad Tr(\gamma^{ab} \gamma_{cd}) = 4(\delta_c^a \delta_d^b - \delta_c^b \delta_d^a) \\ Tr(\gamma^{ab} \gamma^{a'b'} \gamma^{a''b''}) &= 4i[\eta^{aa'} (\eta^{a''b'} \eta^{bb''} - \eta^{a''b} \eta^{b'b''}) + \eta^{ab'} (\eta^{a''b} \eta^{a'b''} - \eta^{a'a''} \eta^{bb''}) \\ &\quad + \eta^{a'b} (\eta^{aa''} \eta^{b'b''} - \eta^{a''b'} \eta^{ab''}) + \eta^{bb'} (\eta^{a'a''} \eta^{ab''} - \eta^{a'b''} \eta^{aa''})] \end{aligned}$$

where $\hat{1}$ in the above is the 4×4 unit matrix.

3.1 Basics of $SL(2N, C)$

As mentioned, the $SL(2N, C)$ symmetry group encompasses, among its primary subgroups, the gravity $SL(2, C)$ symmetry which covers the orthochronous Lorentz group, and the internal hyperflavor $SU(N)$ symmetry proposed to unify all quarks and leptons. Indeed, the $8N^2 - 2$ generators of $SL(2N, C)$ are formed from the tensor products of the generators of $SL(2, C)$ and generators of $SU(N)$ so that the basic transformation applied to the fermions looks as follows

$$\Omega = \exp \left\{ \frac{i}{2} \left[\left(\theta^k + i\theta_5^k \gamma_5 \right) \lambda^k + \frac{1}{2} \theta_{ab}^K \gamma^{ab} \lambda^K \right] \right\} \quad (K = 0, k) \quad (15)$$

Here, among the λ^K matrices, λ^k ($k = 1, \dots, N^2 - 1$) represent the $SU(N)$ Gell-Mann matrices, while λ^0 is the unit matrix $\hat{1}$ corresponding to the $U(1)$ generator involved (the parameters θ parameters may either be constant or, in general, functions of spacetime coordinate). Henceforth, we use uppercase Latin letters (I, J, K) for the $U(1) \times SU(N)$ symmetry, while the lowercase letters (i, j, k) for the case of pure $SU(N)$ symmetry².

For description of the fermion matter in the theory one needs to reintroduce the tetrad multiplet which generally has a form

$$e_\mu = (e_\mu^{aK} \gamma_a + e_{\mu 5}^{aK} \gamma_a \gamma_5) \lambda^K \quad (16)$$

It transforms, as before, according to (2), though the transformation matrix is now given by equation (15). This tetrad structure can be simplified by excluding its axial-vector component, which can be achieved by imposing gauge-invariant constraints on the tetrads. To facilitate this, a special nondynamical $SL(2N, C)$ scalar multiplet can be introduced into the theory.

$$S = \exp \{ i [(s^k + ip^k \gamma_5) \lambda^k + t_{ab}^K \gamma^{ab} \lambda^K / 2] \} \quad (17)$$

which transforms like as $S \rightarrow \Omega S$. With this scalar multiplet one can form a new tetrad in terms of the gauge invariant construction, $S^{-1} e S$. So, choosing appropriately the flat space components in the S field one can turn the tetrad axial part to zero and establish symmetry between Greek and Latin spacetime indices [10]. Nonetheless, even these reduced tetrads can be regarded at best as some dynamical fields [10] rather than standard vielbeins that satisfy the invertibility conditions (5) as in the pure gravity case.

²Some relations for λ matrices used below are given here

$$\begin{aligned} [\lambda^k, \lambda^l] &= 2i f^{klm} \lambda^m, \quad \{\lambda^k, \lambda^l\} = 2(\delta^{kl} \hat{1} + d^{klm} \lambda^m) \\ \lambda^k \lambda^l \lambda^k &= -\lambda^l, \quad \lambda^K \lambda^l \lambda^K = 0, \quad Tr(\lambda^k \lambda^l) = N \delta^{kl} \end{aligned}$$

The connections with a standard choice of the $SU(N)$ matrices are given by the links

$$\lambda^K = \sqrt{2N} T^K, \quad f^{ijk} = \sqrt{N/2} F^{ijk}, \quad d^{ijk} = \sqrt{N/2} D^{ijk}$$

3.2 Tetrads linked to gravity

Meanwhile, some challenge lies in the fact that the tetrads in (16), along with a neutral component, also incorporate $SU(N)$ hyperflavored components which may generally hinder the preservation of the invertibility conditions. Consequently, this imposes a stringent restriction on the permissible form of tetrads. To illustrate, let us assume the tetrads adopt the general $SL(2N, C)$ covariant form

$$e_\mu = e_\mu^{aK} \gamma_a \lambda^K, \quad e_\mu^{aK} e_b^{\mu K'} = \Delta_b^{aKK'}, \quad e_\mu^{aK} e_a^{\nu K'} = \Delta_\mu^{\nu KK'} \quad (18)$$

with certain yet unspecified constructions for $\Delta_b^{aKK'}$ and $\Delta_\mu^{\nu KK'}$, which in the pure gravity case are expected to satisfy the standard arrangement

$$\Delta_b^{a00} = \delta_b^a, \quad \Delta_\mu^{\nu 00} = \delta_\mu^\nu \quad (19)$$

Then multiplying the conditions (18) by the tetrad multipliers $e_\sigma^{bK''}$ and $e_a^{\sigma K''}$, respectively, one come after simple calculations to

$$e_\mu^{aK} = \Delta_b^{aK0} e_\mu^{b0}, \quad e_a^{\mu K'} = \Delta_a^{bK'0} e_b^{\mu 0} \quad (20)$$

that finally gives

$$\Delta_b^{aKK'} = \Delta_c^{aK0} \Delta_b^{cK'0} \quad (21)$$

and correspondingly

$$\Delta_\mu^{\nu KK'} = \Delta_\sigma^{\nu K0} \Delta_\mu^{\sigma K'0} \quad (22)$$

For the constant and multiplicative forms of these functions, the only viable solution arises as

$$e_\mu^{aK} e_b^{\mu K'} = \Delta_b^{aKK'} = \delta_b^a \delta^{K0} \delta^{K'0}, \quad e_\mu^{aK} e_a^{\nu K'} = \Delta_\mu^{\nu KK'} = \delta_\mu^\nu \delta^{K0} \delta^{K'0} \quad (23)$$

which essentially mirrors the pure gravity case. Therefore, the invertibility condition for tetrads holds only if they predominantly belong to the $SL(2, C)$ subgroup rather than the entire $SL(2N, C)$ group

$$e_\mu^{aK} = e_\mu^a \delta^{K0} \quad (24)$$

Otherwise, their connection to general relativity is lost.

Note that tetrads (24) may arise from a spontaneous-like breakdown of $SL(2N, C)$ in the tetrad sector, though they are not treated as dynamical fields in the theory, a premise adopted henceforth. This ansatz implies that the initial symmetry breaks as

$$SL(2N, C) \rightarrow SL(2, C) \times SU(N) \quad (25)$$

ultimately defining the effective symmetry of the theory. One could, in principle, start directly with this symmetry bringing together $SL(2, C)$ gauge gravity and $SU(N)$ grand unification. However, it is particularly insightful to examine how the hyperunified theory organizes itself in this broken symmetry phase, driven by the tetrads, while maintaining its connection to gravity. This may, in general, yield new and distinctive effects associated with hyperunification, as discussed later.

3.3 Gauging $SL(2N, C)$

Once the $SL(2N, C)$ transformation (15) becomes local one needs, as usual, to introduce the gauge field multiplet I_μ transforming as

$$I_\mu \rightarrow \Omega I_\mu \Omega^{-1} - \frac{1}{ig} (\partial_\mu \Omega) \Omega^{-1} \quad (26)$$

whose strength-tensor takes the form

$$I_{\mu\nu} = \partial_{[\mu} I_{\nu]} + ig [I_\mu, I_\nu] \quad (27)$$

This provides the fermion multiplet with the covariant derivative

$$\partial_\mu \Psi \rightarrow D_\mu \Psi = \partial_\mu \Psi + ig I_\mu \Psi \quad (28)$$

where g is the universal gauge coupling constant of the proposed hyperunification. The I_μ multiplet includes in general the vector, axial-vector and tensor field submultiplets of $SU(N)$, and also the neutral tensor field

$$I_\mu = V_\mu + A_\mu + T_\mu = \frac{1}{2} \left(V_\mu^k + i A_\mu^k \gamma_5 \right) \lambda^k + \frac{1}{4} T_{\mu[ab]}^K \gamma^{ab} \lambda^K \quad (K = 0, k) \quad (29)$$

as follows from its decomposition to the flat spacetime component fields. Thereby, the corresponding total strength tensor written in the component fields comes to

$$\begin{aligned} I_{\mu\nu} = & \frac{1}{2} \partial_{[\mu} \left(V^k - i A^k \gamma_5 \right)_{\nu]} \lambda^k - \frac{1}{2} f^{ijk} g (V^i - i A^i \gamma_5)_\mu (V^j - i A^j \gamma_5)_\nu \lambda^k \\ & + \frac{1}{4} \left(\partial_{[\mu} T_{\nu]}^{[ab]K} \gamma_{ab} \lambda^K + i \frac{g}{4} T_\mu^{[ab]K} T_\nu^{[a'b']K'} [\lambda^K \gamma_{ab}, \lambda^{K'} \gamma_{a'b'}] \right) \end{aligned} \quad (30)$$

Similarly, the gauge invariant fermion matter couplings, when given in terms of the I_μ submultiplets, take the form

$$e\mathcal{L}_M = -\frac{g}{2} \bar{\Psi} \left\{ e^\mu, \left[\frac{1}{2} \left(V_\mu^k - i A_\mu^k \gamma_5 \right) \lambda^k + \frac{1}{4} T_{\mu[ab]}^K \gamma^{ab} \lambda^K \right] \right\} \Psi \quad (31)$$

where, as one can readily observe, the vector, axial-vector and tensor fields interact everywhere in (30) and (31) with the universal gauge coupling constant g of $SL(2N, C)$. Just the tensor fields in these equations provide gravitational interaction in the $SL(2N, C)$ HUTs that, aside from the standard linear curvature Lagrangian for gravity (12), includes the conventional quadratic strength terms for all gauge field submultiplets involved.

As mentioned, this raises a crucial issue: how to ensure that only the hyperflavored submultiplet of vector fields V_μ^k and a neutral tensor field $T_{\mu[ab]}^0$ (underlying the $SU(N)$ GUTs and Einstein-Cartan gravity, respectively) appear in the observed particle spectrum, while the hyperflavored submultiplets of axial-vector and tensor fields, A_μ^k and $T_{\mu[ab]}^k$, are discriminated in the theory. Remarkably, if the tetrads retain their pure gravity form discussed above, the submultiplet of axial-vector fields typically remains sterile to ordinary matter, while the submultiplet of tensor fields acquires the Planck scale order masses. Furthermore, they can be completely excluded from the theory provided that the gauge hypermultiplet I_μ is properly constrained through the tetrads involved, as we demonstrate below.

3.4 Axial-vector fields

Let us now focus more closely on the vector and axial-vector fields which are the basic spin-1 carriers of the hyperflavor $SU(N)$ symmetry in the $SL(2N, C)$ theory. Their own sector stemming from the common strength tensor (30) looks as

$$e\mathcal{L}^{V,A} = -\frac{1}{4}[\partial_{[\mu}V_{\nu]}^k - gf^{ijk}(V_{\mu}^iV_{\nu}^j + A_{\mu}^iA_{\nu}^j)]^2 - \frac{1}{4}[\partial_{[\mu}A_{\nu]}^k]^2 \quad (32)$$

where, as one can see, the vector fields acquire a conventional gauge theory form, while the axial-vector field couplings break this gauge invariance³. At the same time, as follows from the matter sector of the theory (31), the vector fields interact with ordinary matter fermions

$$e\mathcal{L}_M^V = -\frac{g}{2}V_{\mu}^i\bar{\Psi}e_a^{\mu}\gamma^a\lambda^i\Psi \quad (33)$$

while axial-vector fields do not, thus being sterile to them.

One might attempt to reconcile axial-vector fields with reality, despite the lack of direct evidence for their existence. Meanwhile, according to the Lagrangian (32), they could give rise to various processes, including the decay of vector fields into invisible axial-vector modes. Current data, in principle, do not rule out such decays for the W and Z bosons in the Standard Model, whose total width fractions into invisible modes remain relatively large [14]. A traditional approach to rendering these modes unobservable at low energies involves making them superheavy via an extensively expanded Higgs sector, while keeping the vector fields, which gauge the Standard Model, massless or sufficiently light. However, this approach faces significant challenges, as the axial-vector fields typically follow the same mass-generating mechanism as their vector counterparts.

Notably, despite the local $SL(2N, C)$ invariance in the theory, the very presence of the axial-vector fields breaks the special gauge symmetry $SU(N)$ associated solely to the vector fields. In this context, an intriguing possibility arises if the axial-vector fields condense at some Planck-scale order \mathcal{M} , thus providing a true vacuum in the theory, $\langle A_{\mu}^k \rangle = \mathbf{n}_{\mu}^k \mathcal{M}$, whose direction is given by the unit Lorentz vector \mathbf{n}_{μ}^k ($\mathbf{n}_{\mu}^{\mu}\mathbf{n}_{\mu}^k = 1$). Remarkably, in such a vacuum, as is directly seen, gauge invariance for the vector fields is fully restored, though a tiny spontaneous breaking of the Lorentz invariance at the scale \mathcal{M} may appear [8].

In particular, for the condensation of axial-vector fields, instead of introducing a conventional polynomial potential for the entire gauge hypermultiplet, one can impose a covariant constraint of the form

$$\frac{1}{4}Tr(I_{\mu}I^{\mu}) = \mathcal{M}^2 \quad (34)$$

which, in terms of field components, reads

$$(V_{\mu}^k)^2 + (A_{\mu}^k)^2 + (T_{\mu}^{abK})^2 = \mathcal{M}^2 \quad (35)$$

³Certainly, there are also the intersecting terms of vector and axial-vector with hyperflavored tensor field submultiplet $T_{\mu}^{[ab]k}$ which we discuss later.

The typical solution to this constraint might be related to a special Goldstone-type configuration

$$A_\mu^k = \mathbf{a}_\mu^k + \mathbf{n}_\mu^k \sqrt{\mathcal{M}^2 - (V_\mu^k)^2 - (T_\mu^{abK})^2 - \mathbf{a}^2}, \quad \mathbf{n}_\mu^k \mathbf{a}_\mu^k = 0 \quad (\mathbf{a}^2 \equiv (\mathbf{a}_\mu^k)^2) \quad (36)$$

This parametrization shows that, while the axial multiplet condenses, represented by an effective Higgs mode formed from the vector and tensor field invariants, its massless excitations are orthogonal to the vacuum direction, aligned along the unit Lorentz vector \mathbf{n}_μ^k . We have further employed the factorized form, $\mathbf{n}_\mu^k \equiv \mathbf{n}_\mu \epsilon^k$, where \mathbf{n}_μ is the unit Lorentz vector ($\mathbf{n}_\mu^2 = \pm 1$, while ϵ^k is associated with the internal $SU(N)$ symmetry ($\epsilon^k \epsilon^k = 1$).

Now, turning back to the spin-1 field Lagrangian (32) and substituting the A_μ^i expression (36) one can confirm that first-order terms in the zero modes \mathbf{a}_μ^i do not appear, provided the orthogonality conditions

$$\mathbf{n}_\mu \mathbf{a}_\mu^k = 0, \quad \mathbf{n}_\mu V_\mu^k = 0, \quad (\partial \mathbf{n}) \mathbf{a}_\nu^k = 0 \quad (37)$$

hold. These can be viewed as gauge conditions for the zero modes \mathbf{a}_μ^k and vector field V_μ^k , respectively, while the last condition implies that the zero modes are independent of the x coordinate along the direction where Lorentz symmetry is broken. Neglecting higher-order terms in the zero modes, one arrives at the Lagrangian

$$e\mathcal{L}^{V,A} = -\frac{1}{4} \left(\partial_{[\mu} V_{\nu]}^k - f^{ijk} V_\mu^i V_\nu^j / 2 \right)^2 + \frac{1}{\mathcal{M}^2} \left\{ \left[\mathbf{n}_{[\mu}^k (V_\sigma^i \partial_{\nu]} V_\sigma^i) \right]^2 + \left[\mathbf{n}_{[\mu}^k (T_\sigma^{Kab} \partial_{\nu]} T_{\sigma ab}^K) \right]^2 \right\} \quad (38)$$

which consists of the conventional vector field gauge-invariant Lagrangian, plus small non-invariant and Lorentz-violating terms arising from the square root in (36) when the lowest-order terms in $(V_\mu^i)^2/\mathcal{M}^2$ and $(T_\mu^{abK})^2/\mathcal{M}^2$ are considered.

At higher orders, the kinetic and interaction terms of the axial-vector zero modes emerge, subtly influencing the observable sector. Notably, some of these modes acquire large masses due to the fact that the symmetry of the constraint (35) is significantly higher than that of the Lagrangian, making certain zero modes pseudo-Goldstone states. From their mass term in (32)

$$(\mathbf{a}_\mu^i \mathbf{a}^{\mu i'}) (\mathbf{n}_\nu^j \mathbf{n}^{\nu j'}) f^{ijk} f^{i'j'k} \mathcal{M}^2 / 2 \quad (39)$$

it follows that modes associated with the "non-diagonal" generators of $SU(N)$ acquire superheavy masses, whereas those corresponding to the "diagonal" generators remain massless, though only at the tree level. Consequently, some distinct processes involving these modes may arise, while the modes themselves remain decoupled from ordinary matter, as previously noted.

A more radical approach, discussed below, would be to impose a specific tetrad filtering condition that automatically excludes the axial-vector field from the total gauge multiplet (29).

3.5 Constraints by tetrad filtering

We begin by proposing some tetrad filtering condition applied directly to the general gauge multiplet (29) that, instead of being imposed by postulate, can be incorporated into the theory through the Lagrange multiplier type term

$$C \left(I_\mu - \frac{1}{4} e_\sigma I_\mu e^\sigma \right)^2 \quad (40)$$

where $C(x)$ is some multiplier function. This term, upon variation under C , yields the constraint condition

$$I_\mu = e_\sigma I_\mu e^\sigma / 4 \quad (41)$$

using the "neutral" tetrads, e_σ and e^σ (24), which are only permitted in the theory. This yields

$$I_\mu = \frac{1}{4} e_\sigma^{aK} e_b^{\sigma K'} (\gamma_a \lambda^K I_\mu \gamma^b \lambda^{K'}) \quad (42)$$

which, upon employing the invertibility conditions of the tetrad (23), results in the equality

$$\left(V_\mu^k + i A_\mu^k \gamma_5 \right) \lambda^k + T_{\mu[ab]}^K \gamma^{ab} \lambda^K / 2 = \left(V_\mu^k - i A_\mu^k \gamma_5 \right) \lambda^k$$

This implies that the reduced gauge multiplet (44) comprises solely the vector fields

$$I_\mu = V_\mu^k \lambda^k / 2 \quad (43)$$

while the axial-vector and tensor field submultiplets vanish identically. Remarkably, by imposing the covariant constraint (41) the starting $SL(2N, C)$ symmetry group is effectively reduced to the pure unitary $SU(N)$ symmetry case. In a sense, the constraint acts as a symmetry-breaking mechanism, but unlike typical scenarios, nothing remains of the original $SL(2N, C)$ gauge sector except its $SU(N)$ part.

Further, to also only keep the axial-vector field submultiplet in the theory one could use another tetrad filtering instead

$$I_\mu = e_\rho e_\sigma I_\mu e^\sigma e^\rho / 16 \quad (44)$$

which goes to

$$\left(V_\mu^k + i A_\mu^k \gamma_5 \right) \lambda^k + T_{\mu[ab]}^K \gamma^{ab} \lambda^K / 2 = \left(V_\mu^k + i A_\mu^k \gamma_5 \right) \lambda^k$$

resulting in the gauge multiplet

$$I_\mu = \left(V_\mu^k + i A_\mu^k \gamma_5 \right) \lambda^k / 2 \quad (45)$$

with the vanished tensor field submultiplet.

And, finally, to include the tensor field submultiplet instead of the axial-vector one, we could require the "less stringent" filtering constraint instead

$$e_\rho I_\mu e^\rho = e_\rho e_\sigma I_\mu e^\sigma e^\rho / 4 \quad (46)$$

that removes the tensor field submultiplet in the both sides. Specifically, this leads to

$$\left(V_\mu^k - iA_\mu^k\gamma_5\right)\lambda^k = e_\rho\left(V_\mu^k - iA_\mu^k\gamma_5\right)\lambda^k e^\rho/4 = \left(V_\mu^k + iA_\mu^k\gamma_5\right)\lambda^k \quad (47)$$

resulting in the vanishing of the axial-vector multiplet, $A_\mu^k = 0$, while the tensor field multiplet remains unaffected. Consequently, one could in principle use the constraint (46) and the properly reduced gauge field hypermultiplet

$$I_\mu = V_\mu^l\lambda^l/2 + T_{\mu[ab]}^K\gamma^{ab}\lambda^K/4 \quad (48)$$

that would solely lead to the theory of vector and tensor fields.

When examining the structure of the total gauge multiplet I_μ in the $SL(2N, C)$ HUT, it becomes evident that while its vector submultiplet persists under any tetrad filtering constraint, the inclusion of axial-vector and tensor field submultiplets in the theory depends crucially on the specific filtering applied. We do not impose these constraints here, though some were employed in earlier works [8, 15].

4 Lagrangians

Below, we present the linear and quadratic stress-tensor Lagrangians for all the field submultiplets comprising the total gauge multiplet I_μ . Our primary focus will be on the Lagrangians for the tensor fields, as the common Lagrangian for the vector and axial-vector submultiplets has already been discussed in (32). Furthermore, considering that with neutral tetrads taken, the axial-vector fields decouple from ordinary matter, we can temporarily set them aside. One could choose the vacuum where they are condensed. This, as was mentioned above, effectively strengthens the gauge nature of the vector fields in the theory. Another, more radical approach might involve their complete exclusion from the theory through an appropriate tetrad filtering constraint (46), ultimately leading to a properly reduced total gauge multiplet (48).

4.1 Linear strength-tensor theories

We are taking the linear hyperunified theory Lagrangian in the Palatini type form

$$e\mathcal{L}_H^{(1)} \sim Tr\{[e^\mu, e^\nu]I_{\mu\nu}\} \quad (49)$$

In the $SL(2N, C)$ case the strength tensor $I_{\mu\nu}$ (30), apart from tensor submultiplet underlying the gravity sector, comprises the vector and axial-vector submultiplets as well. However, due to the neutral tetrad chosen (24) satisfying the commutator

$$[e^\mu, e^\nu] = -2ie_a^\mu e_b^\nu \gamma^{ab} \quad (50)$$

one can easily confirm that they do not contribute to the Lagrangian (49).

Eventually, for the tensor field strength in (30) one has, after taking the necessary traces of products involving γ and λ matrices, the following linear curvature gravity Lagrangian

$$e\mathcal{L}_H^{(1)T} = \frac{1}{2\kappa} \left(\partial_{[\mu} T_{\nu]}^{[ab]0} + g\eta_{cd} T_{[\mu}^{[ac]K} T_{\nu]}^{[bd]K} \right) e_a^\mu e_b^\nu \quad (51)$$

Similarly, beyond the fermion matter coupling with vector fields (33), there are also couplings with the tensor submultiplet, as derived from the general matter Lagrangian (31)

$$e\mathcal{L}_M^T = -\frac{g}{2}\epsilon^{abcd}T_{\mu[ab]}^K\bar{\Psi}e_c^\mu\gamma_d\lambda^K\gamma^5\Psi \quad (52)$$

where couplings of the tensor fields with the neutral and hyperflavored spin density currents appear with the same coupling constant g .

Notably, in the Lagrangian (51), only the neutral component $T_\mu^{[ab]0}$ of the total tensor field multiplet $T_{[\mu}^{[ab]K}$ of $U(1) \times SU(N)$ contains derivative terms, while the hyperflavored components $T_\mu^{[ab]k}$ are solely involved through interaction terms. This indicates that the neutral tensor field alone gauges gravity, while the hyperflavored components are simply reduced to spin currents, given by $\epsilon^{abcd}\bar{\Psi}e_{\mu c}\gamma_d\lambda^k\gamma^5\Psi$. When they both, $T_{[\mu}^{[ab]0}$ and $T_\mu^{[ab]k}$, are independently eliminated from the entire linear tensor field Lagrangian $e\mathcal{L}_G^{(1)} + e\mathcal{L}_M^{(T)}$, one arrives at the Einstein-Cartan type gravity containing, besides the usual GR, the tiny 4-fermion spin density interaction

$$\kappa(\bar{\Psi}\gamma_c\gamma^5\lambda^K\Psi)(\bar{\Psi}\gamma^c\gamma^5\lambda^K\Psi) \quad (53)$$

which in contrast to the standard case [2] includes the hyperflavored four-fermion interaction terms as well.

4.2 Quadratic strength-tensor theories

4.2.1 Pure gravity case

As previously noted, extending the $SL(2, C)$ gauge gravity to the hyperunified $SL(2N, C)$ theory naturally implies the inclusion of some "safe" quadratic curvature terms in its gravitational sector, alongside the standard quadratic terms for the vector field strength tensors. This criterion distinctly identifies the ghost-free curvature-squared gravity model, originally proposed by Neville [12], as the most fitting for such an extension.

Accordingly, we employ the corresponding Lagrangian.

$$\mathcal{L}_G^{(2)} = \lambda e T_{abcd} (T^{abcd} - 4T^{acbd} + T^{cdab}) \quad (54)$$

where the curvature tensor $T_{\mu\nu}^{ab}$ is contracted with tetrads, $T^{abcd} = T_{\mu\nu}^{ab}e^{\mu c}e^{\nu d}$, and properly (anti)symmetrized. In the $SL(2, C)$ gravity case this curvature is constructed from tensor fields involved, while the constant λ , though currently arbitrary, may potentially be determined within the $SL(2N, C)$ unification framework. The Lagrangian $\mathcal{L}_G^{(2)}$, when properly expressed through the tensor field strengths and tetrad components takes the form

$$e\mathcal{L}_G^{(2)} = \lambda(T_{ab}^{\mu\nu}T_{\mu\nu}^{ab} - 4T_{ab}^{\mu\nu}T_{\rho\nu}^{ac}e_{\mu c}e^{\rho b} + T_{ab}^{\mu\nu}T_{\rho\sigma}^{cd}e_{\mu c}e_{\nu d}e^{\rho a}e^{\sigma b}) \quad (55)$$

In view of the extension to the $SL(2N, C)$ hyperunified theory, we suggest that all terms in the Lagrangian (55) may be understood as originating from the corresponding traces of general matrix couplings

$$e\mathcal{L}_G^{(2)} = aTr(I_{\mu\nu}I^{\mu\nu}) + b^{(r)}Tr(I_{\mu\nu}, I^{\rho\nu}, e^\mu, e_\rho)^{(r)} + c^{(s)}Tr(I_{\mu\nu}, I^{\rho\sigma}, e^\mu, e^\nu, e_\rho, e_\sigma)^{(s)} \quad (56)$$

They include all possible mutual placements of the strength-tensors and tetrads inside of traces taken with arbitrary coefficients a , $b^{(r)}$ and $c^{(s)}$ ($r = 1, 2, \dots$; $s = 1, 2, \dots$; here and below, summation is implied over repeated indices r and s). The strength tensor $I_{\mu\nu}$ in the pure gravity case is solely related to the tensor field strength $T_{\mu\nu}^{[ab]}$, as outlined above in (11). Thus, we have for some basic couplings in (56)

$$16e\mathcal{L}_G^{(2)} = aT_{\mu\nu}^{ab}T_{cd}^{\mu\nu}U_{[ab]}^{[cd]} + T_{\mu\nu}^{ab}T_{cd}^{\rho\nu}e^{\mu f}e_{\rho g}\left(b^{(r)}U_{[ab]f}^{(r)[cd]g}\right) \\ + T_{\mu\nu}^{ab}T_{cd}^{\rho\sigma}e^{\mu f}e^{\nu f'}e_{\rho g}e_{\sigma g'}\left(c^{(s)}U_{[ab]ff'}^{(s)[cd]gg'}\right) \quad (57)$$

where the above U tensors are related to the traces of the corresponding products of γ matrices

$$U_{[ab]}^{[cd]} = \text{Tr}(\gamma_{ab}\gamma^{cd}) = 4(\delta_a^c\delta_b^d - \delta_b^c\delta_a^d) \\ U_{[ab]f}^{(1)[cd]g} = \text{Tr}(\gamma_{ab}\gamma^{cd}\gamma_f\gamma^g), \quad U_{[ab]f}^{(2)[cd]g} = \text{Tr}(\gamma_{ab}\gamma_f\gamma^{cd}\gamma^g), \quad \dots \\ U_{[ab]ff'}^{(1)[cd]gg'} = \text{Tr}(\gamma_{ab}\gamma^{cd}\gamma_f\gamma_{f'}\gamma^g\gamma^{g'}), \quad U_{[ab]ff'}^{(2)[cd]gg'} = \text{Tr}(\gamma_{ab}\gamma^{cd}\gamma_f\gamma_{f'}\gamma^g\gamma^{g'}), \dots \quad (58)$$

including all their mutual placements. The sum of terms in the Lagrangian (57), with appropriately justified coefficients a , $b^{(r)}$ and $c^{(s)}$ will inevitably lead to the ghost-free Lagrangian (55). Notably, the second term in (57) not only contributes on its own but also influences the first term, while the third term, in addition to generating its own contribution, similarly affects both the first and second terms. As a result, the Lagrangian (57) takes the form

$$e\mathcal{L}_G^{(2)} = \left(A'T_{ab}^{\mu\nu}T_{\mu\nu}^{ab} + B'T_{ab}^{\mu\nu}T_{\rho\nu}^{ac}e_{\mu c}e^{\rho b} + C'T_{ab}^{\mu\nu}T_{\rho\sigma}^{cd}e_{\mu c}e_{\nu d}e^{\rho a}e^{\sigma b}\right)/16 \quad (59)$$

The new numerical factors A' , B' and C' arise due to the trace contributions to the terms with the original factors a , $b^{(r)}$ and $c^{(s)}$, and their appropriate summation

$$A' = a + b^{(r_a)}t_b^{(r_a)} + c^{(s_a)}t_c^{(s_a)} \\ B' = b + c^{(s_b)}t_c^{(s_b)} \\ C' = c^{(s_c)}t_c^{(s_c)} \quad (60)$$

where the appropriate parameters $b^{(r_a)}$, $c^{(s_a, s_b, s_c)}$ and corresponding trace values $t_b^{(r_a)}$, $t_c^{(s_a, s_b, s_c)}$ are explicitly indicated. While these connections between the new and old factors are theoretically calculable, they are not relevant for what follows. The only requirement is that gravity remains free from ghosts and tachyons, imposing the following conditions on the new parameters

$$A' = -B'/4 = C' = \lambda \quad (61)$$

that directly leads to the acceptable quadratic curvature Lagrangian (55).

Consequently, the spin-connection field T_μ^{ab} becomes truly dynamic. In the particle sector, besides the ordinary massless graviton, there exists a scalar torsion excitation $S(0^-)$ which could, in principle, normally propagate. However, it typically possesses a Planck-scale order mass, $m_S^2 \sim M_P^2/\lambda$, making it unlikely to have any observable significance unless the numerical parameter λ is exceedingly large.

4.2.2 Extension to $SL(2N, C)$ theory

The extension of quadratic curvature terms to the $SL(2N, C)$ HUT follows directly from the general Lagrangian (56) where the strength-tensor $I_{\mu\nu}$ now relates to the total gauge multiplet of $SL(2N, C)$ rather than only the gauge tensor field of $SL(2, C)$.

Let us first focus on the vector field submultiplet in this extension and examine how the internal symmetry in the hyperunified theory is organized. This can be analyzed by considering the Lagrangian (56) term by term using the reduced gauge multiplet (48) and neutral tetrads (the axial-vector fields are now excluded from consideration). As follows, for the pure vector field part all three terms in (56) results in a conventional gauge-invariant form for the vector field interaction.

Indeed, from the structure of the vector field part in the general strength-tensor $I_{\mu\nu}$ (30) one obtains the following expressions for the first, second, and third terms in (56), respectively

$$\begin{aligned} Tr\{V_{\mu\nu}V^{\rho\nu}\} &= N(V_{\mu\nu}^k V^{\mu\nu k})/4, \\ Tr\{V_{\mu\nu}V^{\rho\nu}e^\mu e_\rho\} &= N(V_{\mu\nu}^k V^{\mu\nu k}), \\ Tr\{V_{\mu\nu}V^{\rho\sigma}e^\mu e^\nu e_\rho e_\sigma\} &= 2N(V_{\mu\nu}^k V^{\mu\nu k}) \end{aligned} \quad (62)$$

where the tetrad invertibility conditions (23) and corresponding traces for γ and λ matrices have been applied. Summing these terms with their respective coefficients, the resulting Lagrangian for the vector fields takes the form

$$e\mathcal{L}_H^{(2)V}/N = (V_{\mu\nu}^k V^{\mu\nu k})(a/4 + B + 2C), \quad B = \sum_r b^{(r)}, \quad C = \sum_s c^{(s)}$$

that remains the $SU(N)$ gauge invariant form for their interactions, while the factors a , B and C satisfy the standard condition for the vector field multiplet

$$N(a/4 + B + 2C) = -1/4 \quad (63)$$

As to the tensor field interactions, they emerge from the direct $SL(2N, C)$ extension of the pure gravity Lagrangian in (57)

$$\begin{aligned} 16e\mathcal{L}_H^{(2)T}/N &= aT_{\mu\nu}^{abK}T_{cd}^{\mu\nu K}U_{[ab]}^{[cd]} + T_{\mu\nu}^{abK}T_{cd}^{\rho\nu K}e^{\mu f}e_{\rho g}\left(b^{(r)}U_{[ab]f}^{(r)[cd]g}\right) \\ &\quad + T_{\mu\nu}^{abK}T_{cd}^{\rho\sigma K}e^{\mu f}e^{\nu f'}e_{\rho g}e_{\sigma g'}\left(c^{(s)}U_{[ab]ff'}^{(s)[cd]gg'}\right) \end{aligned} \quad (64)$$

So, collecting both vector and tensor field terms one finally comes to the $SL(2N, C)$ invariant quadratic Lagrangian

$$\begin{aligned} e\mathcal{L}_H^{(2)} &= -1/4V_{\mu\nu}^k V^{\mu\nu k} \\ &\quad + \lambda(T_{ab}^{\mu\nu K}T_{\mu\nu}^{abK} - 4T_{ab}^{\mu\nu K}T_{\rho\nu}^{acK}e_{\mu c}e^{\rho b} + T_{ab}^{\mu\nu K}T_{\rho\sigma}^{cdK}e_{\mu c}e_{\nu d}e^{\rho a}e^{\sigma b}) \end{aligned} \quad (65)$$

where the λ parameter is related again to the factors A' , B' and C' (60) in the pure gravity case. They meet the ghost-free gravity conditions which have practically the same form as in (61)

$$A' = -B'/4 = C' = \lambda/N \quad (66)$$

These conditions, encompassing all the initial parameters a , $b^{(r)}$ and $c^{(s)}$ are readily satisfied, while still allowing the parameter λ to remain arbitrary, though it may be expected of order 1, similar to other parameters in the hyperunified theory.

Note that we do not consider here the intersecting terms between the vector and tensor fields which necessarily arise for the hyperflavored tensor field submultiplet $T_\mu^{[ac]k}$ in the general gauge multiplet couplings (56). As follows from the general covariant derivative (30), these couplings will include the intersecting terms of the form

$$gf^{ijk}D_{[\mu}V^i{}_{\nu]}T_\mu^{[ab]j}T_\nu^{[a'b']k}(\eta_{aa'}\eta_{bb'} - \eta_{ab'}\eta_{a'b}) \quad (67)$$

and some similar ones. They directly break $SU(N)$ gauge invariance because, although the vector fields appear with covariant derivatives, the tensor fields in these terms cannot be treated as matter fields in a way that would preserve such invariance. Fortunately, however, these tensor fields seem to acquire masses at the Planck scale, rendering them irrelevant at lower energies, as argued below.

Combining the linear and quadratic curvature terms for the tensor fields, as presented in (51) and (65), respectively, one leads to a remarkable conclusion concerning the heavy mass origin for the $SU(N)$ tensor field submultiplet. Actually, as noted earlier, within the linear curvature terms, only the neutral tensor field component contains derivative terms, while the hyperflavored components appear solely through polynomial couplings. It can now be readily seen that these couplings

$$e\frac{1}{2\kappa}g\eta_{cd}\left(T_\mu^{[ac]K}T_\nu^{[bd]K} - T_\nu^{[ac]K}T_\mu^{[bd]K}\right)e_a^\mu e_b^\nu \quad (\mu, \nu; c, d = 0, 1, 2, 3) \quad (68)$$

may generate the Planck scale order masses for the $SU(N)$ tensor field submultiplet. Actually, in flat spacetime, where $e_a^\mu = \delta_a^\mu$ and $e_b^\nu = \delta_b^\nu$, these couplings yield a Fierz-Pauli type mass term for the 2-tensor field multiplet

$$-\frac{g}{2\kappa}\eta_{cd}\left(T_\nu^{[\mu c]K}T_\mu^{[\nu d]K} - T_\mu^{[\mu c]K}T_\nu^{[\nu d]K}\right) \quad (69)$$

which ultimately leads to the mass

$$M_T^2 \sim (g/\lambda)M_P^2 \quad (70)$$

This implies that, alongside the massless graviton and the heavy scalar excitation associated with the neutral tensor field ($K = 0$), as seen in the pure gravity case [12], the hyperunified theory also incorporates the $SU(N)$ hyperflavor multiplet of tensor fields having superheavy masses. These fields may become active at the Planck scale, potentially influencing the cosmology of the early universe.

5 Application to GUTs

5.1 Symmetry breaking scenario

It is evident that the hyperunification of all elementary forces implies that, while gravity is basically governed by its unique linear tensor field strength Lagrangian (51), the quadratic strength terms for all components of gauge multiplet I_μ are naturally unified in a common $SL(2N, C)$ invariant Lagrangian (65). Consequently, the gauge sector of the hyperunified Lagrangian takes the form

$$e\mathcal{L}_H = e\mathcal{L}_H^{(1)T} + e\mathcal{L}_H^{(2)T} - \frac{1}{4}V_{\mu\nu}^k V^{\mu\nu k} \quad (71)$$

containing, besides the unified Einstein-Cartan type gravity given by the Lagrangians (51) and (65), the standard $SU(N)$ invariant vector field part. Meanwhile, the axial-vector fields are effectively excluded from the theory, either due to their sterility, condensation, or the tetrad filtering constraints discussed earlier.

A conventional scenario for breaking the $SL(2N, C)$ invariance in the theory typically relies on an appropriate set of scalar fields capable of first reducing this invariance to the intermediate $SL(2, C) \times SU(N)$ symmetry, and subsequently to the Standard Model. However, in our framework, there is no necessity to induce the initial stage of symmetry breaking explicitly. Indeed, due to the neutral tetrad pattern proposed, the gauge submultiplets related to the "nondiagonal" generators of $SL(2N, C)$ either are generically superheavy, as observed above with hyperflavored tensor fields due to the relevant linear curvature terms, or remain sterile to ordinary matter, as is the case with the axial-vector fields (alternatively, they can be entirely filtered out of the theory).

As to the internal $SU(N)$ symmetry violation down to the Standard Model one actually need to have the adjoint scalar multiplets of the type

$$\Phi = (\phi^k + i\phi_5^k \gamma_5)\lambda^k + \phi_{ab}^K \gamma^{ab} \lambda^K / 2 \quad (72)$$

which transform under $SL(2N, C)$ as

$$\Phi \rightarrow \Omega \Phi \Omega^{-1} \quad (73)$$

It generally contains, apart the scalar components, the pseudoscalar and "tensor" components as well. However, as in the above gauge multiplet case, one can use again the tetrad projection mechanism to filter away these "superfluous" components, just like as it was done in (41)

$$\Phi = e_\sigma \Phi e^\sigma / 4$$

As a result, with tetrads satisfying the invertibility conditions (23) there is only left the pure scalar components in the $SU(N)$ symmetry breaking multiplet Φ

$$\Phi = \phi^k \lambda^k \quad (k = 1, \dots, N^2 - 1)$$

providing (with other similar scalar multiplets) the breaking of the $SU(N)$ GUT down to the Standard Model. The final symmetry breaking to $SU(3)_c \times U(1)_{em}$ is provided by extra scalar multiplets whose assignment depends on which multiplets are chosen for quarks and leptons.

5.2 $SU(5)$ and its direct extension

As discussed, the $SL(2N, C)$ HUT effectively exhibits a local $SL(2, C) \times SU(N)$ symmetry, rather than an entire $SL(2N, C)$ symmetry, which primarily serves to determine the structure of the gauge and matter multiplets. This results in the $SL(2, C)$ gauge gravity on one side and $SU(N)$ grand unified theory on the other. Given that all states involved in the $SL(2N, C)$ theories are additionally classified according to their spin values, many potential $SU(N)$ GUTs, including the conventional $SU(5)$ theory [16], appear to be irrelevant for standard spin 1/2 quarks and leptons. However, the application of $SL(2N, C)$ to their proposed preon constituents turns out both natural and exceptionally promising.

Note first that the $SL(2N, C)$ symmetry is presently applied to the chiral fermions, leading to the decomposition of a general transformation (15) into distinct transformations for lefthanded and righthanded fermions

$$\begin{aligned}\Omega_L &= \exp \left\{ \frac{i}{2} \left[\left(\theta^k - i\theta_5^k \right) \hat{1}\lambda^k + \frac{1}{2}\theta_{ab}^K \Sigma^{ab} \lambda^K \right] \right\}, \\ \Omega_R &= \exp \left\{ \frac{i}{2} \left[\left(\theta^k + i\theta_5^k \right) \hat{1}\lambda^k + \frac{1}{2}\theta_{ab}^K \bar{\Sigma}^{ab} \lambda^K \right] \right\}\end{aligned}\quad (74)$$

in the chiral basis for γ matrices. Here Σ^{ab} and $\bar{\Sigma}^{ab}$ are given by

$$\Sigma^{ab} = \frac{i}{2}(\sigma^a \bar{\sigma}^b - \sigma^b \bar{\sigma}^a), \quad \bar{\Sigma}^{ab} = \frac{i}{2}(\bar{\sigma}^a \sigma^b - \bar{\sigma}^b \sigma^a) \quad (75)$$

where the two-dimensional matrices σ^a and $\bar{\sigma}^a$

$$\gamma^a = \begin{pmatrix} 0 & \sigma^a \\ \bar{\sigma}^a & 0 \end{pmatrix} \quad (76)$$

are expressed in terms of the unit and Pauli matrices as

$$\sigma^a = (\hat{1}, \boldsymbol{\sigma}), \quad \bar{\sigma}^a = (\hat{1}, -\boldsymbol{\sigma}) \quad (77)$$

Accordingly, the gauge multiplets of $SL(2N, C)$ associated with both lefthanded and righthanded fermions are appropriately specified.

We begin with the familiar $SU(5)$, which could naturally emerge from the $SL(10, C)$ hyperunification. In this context, some of its low-dimensional chiral fermion multiplets (lefthanded for certainty) can be represented in terms of the $SU(5) \times SL(2, C)$ components as

$$\Psi_L^{i\mathbf{a}}, \quad 10 = (\bar{\mathbf{5}}, 2) \quad (78)$$

and

$$\Psi_{L[ai, jb]} = \Psi_{L[ij]\{\mathbf{ab}\}} + \Psi_{L\{ij\}[\mathbf{ab}]}, \quad 45 = (10, 3) + (15, 1) \quad (79)$$

Here, we have used that any common antisymmetry across two or more combined $SL(10, C)$ indices ($i\mathbf{a}, j\mathbf{b}, k\mathbf{c}$) implies antisymmetry in the $SU(5)$ indices ($i, j, k = 1, \dots, 5$) and symmetry in the spinor indices ($\mathbf{a}, \mathbf{b}, \mathbf{c} = 1, 2$), and the reverse holds as well (the dimensionality of the representations is also indicated). Notably, while the fermionic $SU(5)$ antiquintet is

easily constructed (78), the fermionic decuplet does not arise from the purely antisymmetric $SL(10, C)$ representation (79). Instead, the tensor in (79) corresponds to a collection of vector and scalar multiplets rather than fermionic ones. This effectively rules out the standard $SU(5)$ GUT, along with its supersymmetric or string-inspired extensions [17, 18] from consideration.

Note in this connection, that all GUTs where fermions are assigned to the pure antisymmetric representations seem to be also irrelevant since the spin magnitude of appearing states are not in conformity with what we have in reality. The most known example of this kind is the $SU(11)$ GUT [19] with all three quark-lepton families collected in its one-, two-, three-, and four-index antisymmetric representations. No doubt this GUT should also be excluded in the framework of the considered $SL(2N, C)$ theories. Actually, for the right 1/2 spin value of ordinary quarks and leptons these theories should include more complicated fermion multiplets having in general the upper and lower indices rather than the pure asymmetric ones. The point is, however, that such multiplets appear enormously large and contain in general lots of exotic states which never been detected. This may prompt further exploration into the composite structure of quarks and leptons, for whose constituents – preons – the $SL(2N, C)$ unification might look much simpler. As we demonstrate below, applying the $SL(2N, C)$ symmetry to the model of composite quarks and leptons – comprising constituent chiral preons in its fundamental representations and the masslessness condition for their composites – leads to the identification of the $SL(16, C)$ HUT as the most likely candidate for the hyperunification of existing elementary forces, accommodating all three quark-lepton families [8, 15].

5.3 $SU(8)$ with composite quarks and leptons

Following the recent discussion [20], we introduce N lefthanded and N righthanded preons being the fundamental multiplets $P_{Li\mathbf{a}}^\alpha$ and $P_{Ri\mathbf{a}}^{\alpha'}$ of the vectorlike "metaflavor" $SL(2N, C)$ HUT symmetry ($i = 1, \dots, N$; $\mathbf{a} = 1, 2$) times some local left-right "metacolor" $SO(n)_L \times SO(n)_R$ symmetry ($\alpha = 1, \dots, n$; $\alpha' = 1, \dots, n$) binding preons inside quarks and leptons⁴. Both of these symmetries are obviously anomaly-free and the numbers of metaflavors (N) and metacolors (n) are not yet determined. The metaflavor symmetry describes preons at small distances as well as their composites at large ones. They are produced individually from the lefthanded and righthanded preons due to confining forces of the above metacolor symmetry. Some of these composites, including the observed quarks and leptons, are expected to be much lighter than their composition scale. For that, the accompanying chiral symmetry $SU(N)_L \times SU(N)_R$ of the preons should be preserved at large distances in a way that – when it is considered as the would-be local symmetry group with some spectator gauge fields and fermions – the corresponding triangle anomaly matching conditions [21] are satisfied. Namely, the $SU(N)_L^3$ and $SU(N)_R^3$ anomalies related to N lefthanded and N righthanded preons have to individually match those for lefthanded and righthanded composite fermions being produced by the $SO(n)_L$ and $SO(n)_R$ metacolor forces, respectively.

⁴By tradition, we call them the "metaflavor" and "metacolor" symmetry, while still referring to the $SU(N)$ subgroup of $SL(2N, C)$ as the hyperflavor symmetry.

Moreover, as is turned out, just this condition, when being properly strengthened, can determine the particular metaflavor symmetry $SL(2N, C)$ in the theory. Indeed, we first assume that all composites, both lefthanded and righthanded, have just the three-preon configuration ($n = 3$), thus fixing the metacolor symmetry to $SO(3)_L \times SO(3)_R$. And second and most importantly, they belong to a single representation of their chiral symmetries $SU(N)_L$ and $SU(N)_R$, respectively, rather than to some set of representations. Then it turns out that among all their third-rank representations the anomaly matching condition holds individually only for multiplets of the type $\psi_{[i j]L}^k$ and $\psi_{[i j]R}^k$ ($i, j, k = 1, 2, \dots, N$), that gives the unique solution to the number of preons N , both lefthanded and righthanded,

$$N^2/2 - 7N/2 - 1 = 3, \quad N = 8 \quad (80)$$

This means that among all possible chiral symmetries only the $SU(8)_L \times SU(8)_R$ symmetry can in principle provide masslessness of lefthanded and righthanded fermion composites at large distances. This in turn identifies – among all metaflavor $SL(2N, C)$ symmetries – just $SL(16, C)$ as the most likely candidate for hyperunification. Note that, in contrast to the above global chiral symmetry, in the local $SL(16, C)$ metaflavor theory, being as yet vectorlike, all metaflavor triangle anomalies are automatically cancelled out.

Turning now from the chiral symmetry multiplets $\psi_{[i j]L,R}^k$ to the corresponding $SL(16, C)$ composite multiplets $\Psi_{[ia, jb]L,R}^{kc}$ one can write them in terms of the $SU(8) \times SL(2, C)$ components as the collection

$$\Psi_{[ia, jb]}^{kc} = \Psi_{[ij]\{ab\}}^{kc} + \Psi_{\{ij\}[ab]}^{kc}, \quad 1904 = (216, 2) + (216 + 8, 4) + (280 + 8, 2) \quad (81)$$

which contains some spin 1/2 and 3/2 lefthanded and righthanded composite fermion submultiplets. Meanwhile, as one can easily confirm, among all submultiplets in (81) only the $(216, 2)_{L,R}$ ones satisfy individually the anomaly matching condition for the chiral $SU(8)_L$ and $SU(8)_R$ symmetries, respectively. As a result, all the other submultiplets there have then to acquire superheavy masses. This actually means that only the $SU(8) \times SL(2, C)$ subgroup of the $SL(16, C)$ HUT symmetry survives at large distances where the composite fermions emerge. Surprisingly enough, this is consistent with what we had above, albeit from a different perspective. Namely, the filtered $SL(2N, C)$ gauge theory, in which only neutral tensor field and vector field multiplet remain, turns out to be effectively reduced to the $SU(N) \times SL(2, C)$ invariant theory. Now, in the composite model, for the particular case of metaflavored symmetry $SL(16, C)$, this independently follows from the preservation of the accompanying chiral symmetry $SU(8)_L \times SU(8)_R$ at large distances, thus leading to the theory with the residual metaflavor symmetry $SU(8) \times SL(2, C)$.

Remarkably, the above $(216, 2)_{L,R}$ submultiplets being decomposed into the standard $SU(5)$ GUT and family symmetry $SU(3)_F$ looks as

$$(216, 2)_{L,R} = [(\bar{5} + 10, \bar{3}) + (45, 1) + (5, 8 + 1) + (24, 3) + (1, 3) + (1, \bar{6})]_{L,R} \quad (82)$$

where the first term in the squared brackets, when taken for lefthanded states in 216_L , describes all three quark-lepton families being the family symmetry triplets. However, there are also the similar righthanded states in 216_R in our still vectorlike $SL(16, C)$ theory.

This means that, while preons are left massless being protected by their own metacolors, the composites (82) being metacolor singlets could in principle pair up and acquire the heavy Dirac masses.

To avoid this for the submultiplet of physical quarks and leptons in (82), $(\bar{5} + 10, \bar{3})_L$, one may propose, following the scenario developed in [20], some spontaneous breaking of the basic L - R symmetry in the theory. This is assumed to follow from the sector of righthanded preons that reduces the chiral symmetry of their composites down to $[SU(5) \times SU(3)]_R$. Actually, such a breaking may readily appear due to a possible condensation of massive composite scalars which unavoidably appear in the theory together with composite fermions. This means that, though the massless righthanded preons still possess the $SU(8)_R$ symmetry, the masslessness of their composites at large distances is now solely controlled by its remained $[SU(5) \times SU(3)]_R$ part. Thus, while nothing really happens with the lefthanded preon composites still completing the total multiplet $(216, 2)_L$ in (82), the righthanded preon composites with their residual chiral symmetry no longer include all submultiplets given in $(216, 2)_R$. Very remarkably, the corresponding anomaly matching condition "organizes" their composite spectrum in such a way that the submultiplet $(\bar{5} + 10, \bar{3})_R$ is absent among the righthanded preon composites. As a result, all the lefthanded submultiplets in $(216, 2)_L$, except the $(\bar{5} + 10, \bar{3})_L$, will then pair up, thus becoming heavy and decoupling from laboratory physics [20].

Accordingly, once the L - R symmetry is violated in the theory, the vectorlike metaflavor symmetry $SU(8) \times SL(2, C)$, while still working for preons, will also break down to its subgroup $[SU(5) \times SU(3)_F] \times SL(2, C)$ for their large-distance composites. So, one eventually comes to the conventional $SU(5)$ GUT [16] together with the extra local $SU(3)_F$ family symmetry [22] describing just three standard families of composite quarks and leptons. Both types of the triangle anomalies, $SU(5)^3$ and $SU(3)_F^3$, emerging at this stage are properly cancelled out in the theory.

The further symmetry violation is related, as was mentioned above, to the adjoint scalar field multiplet Φ (72) which in the present context breaks the $SU(5)$ to the Standard Model. As to the final breaking of the SM and accompanied family symmetry $SU(3)_F$, it appears through the extra multiplets $H^{[ia,jb,kc,ld]}$, and $\chi_{[ia,jb]}$ and $\chi_{\{ia,jb\}}$ of $SL(16, C)$, respectively. These multiplets contain, among others, the true scalar components which develop the corresponding VEVs and give masses to the weak bosons, as well as the flavor bosons of the $SU(3)_F$. They also generate masses to quarks and leptons located in the lefthanded fermion multiplet (81, 82) through the $SL(16, C)$ invariant Yukawa couplings

$$\begin{aligned} & \frac{1}{\mathcal{M}} \left[\Psi_{[ja, kb]L}^{ic} C \Psi_{[me, nf]L}^{ld} \right] H^{\{[ja, kb], [me, nf]\}} (a_u \chi_{[ic, ld]} + b_u \chi_{\{ic, ld\}}) \\ & \frac{1}{\mathcal{M}} \left[\Psi_{[ja, kb]L}^{ic} C \Psi_{[ic, me]L}^{ld} \right] H^{\{[ja, kb], [me, nf]\}} (a_d \chi_{[ld, nf]} + b_d \chi_{\{ld, nf\}}) \end{aligned} \quad (83)$$

with different index contraction for the up quarks, and down quarks and leptons, respectively ($i, j, k, l, m, n = 1, \dots, 8$; $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \mathbf{f} = 1, 2$). The mass \mathcal{M} stands for some effective scale in the theory that in the composite model of quarks and leptons can be related to their compositeness scale, while $a_{u,d}$ and $b_{u,d}$ are some dimensionless constants of the order of 1. Actually, these couplings contain two types of scalar multiplets with the following

$SU(8) \times SL(2, C)$ components – the H multiplet $H^{\{[ja, kb], [me, nf]\}}$ containing the true scalar components

$$H^{[jkmn]\{[ab], [ef]\}}(70, 1) \quad (84)$$

and symmetric and antisymmetric χ multiplets, $\chi_{\{ic, ld\}}$ and $\chi_{[ic, ld]}$, whose scalar components look as

$$\chi_{[il][cd]}(28, 1), \quad \chi_{[cd]\{il\}}(36, 1) \quad (85)$$

Decomposing them into the components of the final $SU(5) \times SU(3)_F$ symmetry one finds the full set of scalars

$$\begin{aligned} 70 &= (5, 1) + (\bar{5}, 1) + (10, \bar{3}) + (\bar{10}, 3) \\ 28 &= (5, 3) + (10, 1) + (1, \bar{3}) \\ 36 &= (5, 3) + (15, 1) + (1, 6) \end{aligned} \quad (86)$$

containing the $SU(5)$ quintets $(5, 1)$ and $(\bar{5}, 1)$ to break the Standard Model at the electroweak scale M_{SM} and the the $SU(3)_F$ triplet and sextet, $(1, \bar{3})$ and $(1, 6)$, to properly break the family symmetry at some large scale M_F . One may refer to the scalars (84) and (85) as the "vertical" and "horizontal" ones, respectively, which are actually the simplest choice to form the above Yukawa couplings. Working in pairs in them, they presumably determine masses and mixings of all quarks and leptons. And the last but not the least, they may be indeed composed, in the model considered, from the same preons as quarks and leptons [20].

6 Conclusion

We have investigated the potential of the local $SL(2N, C)$ symmetry to unify all fundamental forces, including gravity. The view that gravity alone dictates the symmetry structure required for this hyperunification naturally leads to extending the $SL(2, C)$ gauge gravity group to the broader local symmetry $SL(2N, C)$, where N defines the internal $SU(N)$ symmetry as a subgroup. This extension yields a consistent hyperunification framework, provided that the tetrad fields of $SL(2, C)$ gravity retain their form in the extended theory, thereby preserving the crucial invertibility condition that would otherwise be lost. Consequently, while the full gauge multiplet of $SL(2N, C)$ encompasses vector, axial-vector and tensor field submultiplets of the $SU(N)$ symmetry, only the vector field submultiplet and the singlet tensor field emerge in the observed particle spectrum. The axial-vector fields remain decoupled from ordinary matter, while the tensor field submultiplet acquires mass at the Planck scale. As a result, the effective symmetry reduces to $SL(2, C) \times SU(N)$, representing $SL(2, C)$ gauge gravity and $SU(N)$ grand unified theory, thereby naturally circumventing the restrictions imposed by the Coleman-Mandula theorem.

For the gravitational part, the extension of $SL(2, C)$ gauge gravity to $SL(2N, C)$ HUT necessitates the inclusion of some "safe" quadratic curvature terms in the gravitational sector of the extended theory, alongside the standard quadratic terms for the strength-tensors of the vector fields. This requirement uniquely identifies, from among all possible

candidates, the ghost-free curvature-squared gravity Lagrangian [12, 13], as the most appropriate model for such an extension. Consequently, the resulting theory, after reduction, includes the $SL(2, C)$ symmetric $R + R^2$ Einstein-Cartan type gravity action, which remains free from ghosts and tachyons. The theory also contains the properly suppressed four-fermion (spin current-current) interaction (53) which, in contrast to the standard case [2], includes the hyperflavor depending interaction terms as well.

For the grand unification part, in turn, since all states involved in $SL(2N, C)$ theories are additionally classified by spin magnitude, the $SU(N)$ GUTs with purely antisymmetric matter multiplets, including the usual $SU(5)$ theory, turn out to be irrelevant for the standard 1/2-spin quarks and leptons. Meanwhile, the $SU(8)$ grand unification for all three families of composite quarks and leptons, arising from the $SL(16, C)$ theory formulated for preon constituents, appears to be particularly interesting.

Meanwhile, it is important to clarify that $SL(2N, C)$ hyperunification does not imply a single universal coupling constant for gravity and other interactions, as is usually assumed in unified theories. Instead, it suggests that all these forces are provided by vector and tensor fields being the members of the same $SL(2N, C)$ gauge hypermultiplet. A universal constant is indeed necessary for the standard quadratic strength terms of vector and tensor fields. However, the pure gravitational interaction has a fundamentally different coupling, linear in the tensor field strength (51). This unique coupling arises solely due to the presence of tetrads, which are essential ingredients for an $SL(2N, C)$ invariant theory. It comes with its own independent coupling constant ($1/2\kappa$), conventionally related to the Planck mass. Significantly, the vector (and axial-vector) fields cannot have these linear strength terms alongside the standard quadratic ones. Furthermore, we should emphasize the special role of tetrads within the entire $SL(2N, C)$ theory framework. The key point is that tetrads should be truly neutral, devoid of any $SU(N)$ hyperflavored components. Otherwise, they cannot be treated as standard vielbein fields satisfying the invertibility conditions (23).

Finally, axial-vector fields may present challenges within the theory, though, as previously discussed, these could be addressed through their condensation [8] or via the tetrad filtering constraint (46), which fully eliminates them. However, there exists a fundamentally different approach that may resolve this issue specifically for the tetrads considered. Unlike vector and tensor fields, axial-vector fields do not directly couple to fermionic matter, as shown earlier. This points to a novel scenario for hyperunification, where all gauge fields associated with the $SL(2N, C)$ symmetry arise as composite bosons formed from fermion pairs, rather than being elementary fields. This approach, which has long been considered a viable alternative in quantum electrodynamics [23], gravity [24] and Yang-Mills theories [25, 26, 27], has yet to be applied to noncompact unified symmetries. In this framework, where only the global $SL(2N, C)$ symmetry is initially imposed on the pure fermionic Lagrangian with suitably constrained fermion currents, one might expect that only composite vector and tensor fields emerge in the effective gauge theory, while axial-vector fields are never generated.

Further study may also focus on the phenomenological aspects of the theory. The spontaneous breaking of the $SL(2N, C)$ HUT through the effective $SL(2, C) \times SU(N)$ symmetry down to the Standard Model and beyond will give rise to numerous new processes. These

processes arise from the generalization of both the gravity and Standard Model sectors, leading to new particles and interactions. As partially discussed earlier, this includes axial-vector fields, which may introduce corrections to the decays of electroweak bosons in the Standard Model, and superheavy tensor fields, which could have a significant impact on the cosmology of the early universe.

These significant issues will be addressed elsewhere.

Acknowledgments

I would like to thank Colin Froggatt and Holger Nielsen for interesting discussions.

References

- [1] R. Utiyama, *Phys. Rev.* **101** (1956) 1597.
- [2] T.W. Kibble, *J. Math. Phys.* **2** (1960) 212.
- [3] C.J. Isham, A. Salam and J. Strathdee, *Lett. Nuovo Cim.* **5** (1972) 969.
- [4] Y.M. Cho, *J. Math. Phys.* **16** (1975) 2029, *Phys. Rev. D* **14** (1976) 3335;
Y.M. Cho and P.G.O. Freund, *Phys. Rev. D* **12** (1975) 1711.
- [5] J.C. Huang and P.W. Dennis, *Phys. Rev. D* **15** (1977) 983, *D* **24** (1981) 3125
- [6] R. Percacci, *Phys. Lett. B* **144** (1984) 37, *Nucl. Phys. B* **353** (1991) 271;
F. Nesti and R. Percacci, *J. Phys. A* **41** (2008) 075405, *Phys. Rev. D* **81** (2010) 025010.
- [7] A.H. Chamseddine, *Phys. Rev. D* **70** (2004) 084006, *Int. J. Geom. Meth. Mod. Phys.* **3** (2006) 149.
- [8] J.L. Chkareuli, *Phys. Lett. B* **834** (2022) 137417.
- [9] S. Coleman and J. Mandula, *Phys. Rev.* **159** (1967) 1251.
- [10] C.J. Isham, A. Salam and J. Strathdee, *Phys. Rev. D* **8** (1973) 2600, *Phys. Rev. D* **9** (1974) 1702.
- [11] F. Gursev and L. A. Radicati, *Phys. Rev. Lett.* **13** (1964) 173;
B. Sakita, *Phys. Rev.* **137** (1964) B 1756.
- [12] D.E. Neville, *Phys. Rev. D* **18** (1978) 3535.
- [13] E. Sezgin and P. van Nieuwenhuizen, *Phys. Rev. D* **21** (1980) 3269.

- [14] R.L. Workman et al. (Particle Data Group), Prog. Theor. Exp. Phys. **2022** (2022) 083C01 .
- [15] J.L. Chkareuli, Eur. Phys. J. C **84** (2024) 11, 1212; arXiv: 2310.11847 [hep-th].
- [16] H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32** (1974) 438.
- [17] J. Hisano, H. Murayama, T. Yanagida, Nucl. Phys. B **402** (1993) 46 .
- [18] J.L. Chkareuli, I.G. Gogoladze, Phys. Rev. D **58** (1998) 055011, arXiv: hep-ph/9803335 [hep-ph].
- [19] H. Georgi, Nucl. Phys. B **156** (1979) 126.
- [20] J.L. Chkareuli, Nucl. Phys. B **941** (2019) 425, arXiv:1901.07428 [hep-ph].
- [21] G. 't Hooft, in *Recent Developments in Gauge Theories*, edited by G.'t Hooft et al (Plenum, New-York, 1980).
- [22] J.L. Chkareuli, JETP Lett. **32** (1980) 671;
Z.G. Berezhiani and J.L. Chkareuli, Sov. J. Nucl. Phys. **37** (1983) 618;
Z.G. Berezhiani, Phys. Lett. B **129** (1983) 99; ibid B **150** (1985) 177.
- [23] J.D. Bjorken, Ann. Phys. (N.Y.) **24** (1963) 174.
- [24] P.R. Phillips, Phys. Rev. **146** (1966) 966.
- [25] T. Eguchi, Phys.Rev. D **14** (1976) 2755;
H. Terazava, Y. Chikashige, K. Akama, Phys. Rev. D **15** (1977) 480.
- [26] M. Suzuki, Phys. Rev. D **37** (1988) 210; Phys. Rev. D **82** (2010) 045026.
- [27] J.L. Chkareuli, Phys. Lett. B **817** (2021) 136281, arXiv: 2102.11217 [hep-ph].