

# On the geometric fixed points of the real topological cyclic homology of $\mathbb{Z}/4$

Thomas Read

thomas.read@warwick.ac.uk

## Abstract

We study the homotopy groups of the geometric fixed points of the real topological cyclic homology of  $\mathbb{Z}/4$ . We relate these groups to the values of the non-abelian derived functors of the functor  $M \mapsto (M \otimes_{\mathbb{Z}/4} M)^{C_2}$  at the  $\mathbb{Z}/4$ -module  $\mathbb{Z}/2$ , which we precisely calculate with computer assistance up to degree 6, and calculate in general up to slight remaining ambiguity. Using these results we compute  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  exactly for  $i \leq 1$ , up to an extension problem for  $2 \leq i \leq 5$ , and describe the asymptotic growth of this group for large  $i$ . A consequence of these computations is that there exists some  $0 \leq i \leq 5$  such that the canonical map comparing the genuine symmetric and symmetric  $L$ -theory spectra of  $\mathbb{Z}/4$  is not an isomorphism on degree  $i$  homotopy, and moreover this comparison map is never an isomorphism on homotopy in sufficiently large degrees.

## Contents

<b>Introduction</b>	<b>2</b>
Relation to $L$ -theory . . . . .	5
Computer code and results . . . . .	7
Acknowledgements . . . . .	7
<b>1 Preliminaries and conventions</b>	<b>7</b>
1.1 Point-set models and model structures . . . . .	7
1.2 Dold-Kan equivalence . . . . .	9
1.3 Real topological Hochschild homology . . . . .	10
<b>2 Low degree homotopy of <math>\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}</math></b>	<b>11</b>
2.1 Decomposition of the equaliser formula for $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ . . . . .	11
2.2 Homotopy groups of $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ and $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$ . . . . .	14
2.3 Action of the maps $f$ and $r$ on homotopy . . . . .	19
2.4 Resolving the extension problem in degree 1 . . . . .	27

<b>3</b>	<b>Asymptotic growth estimates</b>	<b>32</b>
3.1	Non-abelian derived functor computations . . . . .	34
3.2	Application to size estimates . . . . .	41
<b>References</b>		<b>43</b>

## Introduction

Trace methods, introduced in [MHB93], study the  $K$ -theory spectrum of a ring spectrum  $A$  by mapping it to other invariants that are more tractable to compute. The most basic such invariant is the topological Hochschild homology spectrum  $\mathrm{THH}(A)$ , from which one can obtain further invariants such as the  $p$ -typical topological cyclic homology spectrum  $\mathrm{TC}(A; p)$ . If  $A$  is a ring spectrum with anti-involution,  $\mathrm{THH}(A)$  refines to a real cyclotomic spectrum  $\mathrm{THR}(A)$ , the real topological Hochschild homology (introduced in [HM15], [Dot12] and [Høg16]). This receives a trace map from the real algebraic  $K$ -theory  $\mathrm{KR}(A)$  of [HM15]. From  $\mathrm{THR}(A)$  one can obtain further approximations to real  $K$ -theory, in particular the  $p$ -typical real topological cyclic homology, a genuine  $\mathbb{Z}/2$ -spectrum denoted  $\mathrm{TCR}(A; p)$  (defined in [Høg16]). We will study the geometric fixed points of  $\mathrm{TCR}(A; p)$ .

Dotto, Moi and Patchkoria analyse the dihedral structure of  $\mathrm{THR}(A; p)$  in [DMP24], and use this to obtain a formula for the geometric fixed points of  $\mathrm{TCR}(A; p)$ . In particular when  $p = 2$  they show that for  $A$  a connective ring spectrum with anti-involution,  $\mathrm{TCR}(A; 2)^{\phi_{\mathbb{Z}/2}}$  can be expressed as a homotopy equaliser

$$\mathrm{TCR}(A; 2)^{\phi_{\mathbb{Z}/2}} \simeq \mathrm{hoeq} \left( (\mathrm{THR}(A; 2)^{\phi_{\mathbb{Z}/2}})^{C_2} \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[r]{} \end{array} \mathrm{THR}(A; 2)^{\phi_{\mathbb{Z}/2}} \right) \quad (1)$$

where  $C_2$  denotes the Weyl group of  $\mathbb{Z}/2$  in  $O(2)$ . We can consider  $A^{\phi_{\mathbb{Z}/2}}$  as an  $A$ -module via the Frobenius module structure

$$A \xrightarrow{\Delta} (N_{\{e\}}^{\mathbb{Z}/2}(A))^{\phi_{\mathbb{Z}/2}} \xrightarrow{\mu^{\phi_{\mathbb{Z}/2}}} A^{\phi_{\mathbb{Z}/2}}$$

(where  $N_{\{e\}}^{\mathbb{Z}/2}(-)$  is the Hill-Hopkins-Ravenel norm of [HHR16], see Section 1.1 for more details). Then there is an equivalence of  $C_2$ -spectra

$$\mathrm{THR}(A; 2)^{\phi_{\mathbb{Z}/2}} \simeq A \otimes_{N_{\{e\}}^{C_2} A} N_{\{e\}}^{C_2}(A^{\phi_{\mathbb{Z}/2}}),$$

(where  $A$  is a  $N_{\{e\}}^{C_2} A$ -spectrum via the multiplication map  $\mu : N_{\{e\}}^{C_2} A \rightarrow A$  and  $N_{\{e\}}^{C_2}(A^{\phi_{\mathbb{Z}/2}})$  is a  $N_{\{e\}}^{C_2} A$ -spectrum via applying the norm to the Frobenius module structure on  $A^{\phi_{\mathbb{Z}/2}}$ ). If  $A^{\phi_{\mathbb{Z}/2}}$  splits as an  $A$ -module into the direct sum of its homotopy groups

$$A^{\phi_{\mathbb{Z}/2}} \simeq \bigoplus_{n \geq 0} \Sigma^n H(\pi_n A^{\phi_{\mathbb{Z}/2}})$$

then we get corresponding decompositions of the terms  $(\mathrm{THR}(A; 2)^{\phi_{\mathbb{Z}/2}})^{C_2}$  and  $\mathrm{THR}(A; 2)^{\phi_{\mathbb{Z}/2}}$  appearing in the equaliser (1), and [DMP24] identifies the maps  $f$  and

$r$  under these decompositions. This approach is a key ingredient in the identification of the homotopy type of  $\mathrm{TCR}(\mathbb{Z}; 2)^{\phi_{\mathbb{Z}/2}}$  ([DMP24]) and the computation of the homotopy groups of  $\mathrm{TCR}(k; 2)^{\phi_{\mathbb{Z}/2}}$  for  $k$  a field ([DMP24] for fields of odd characteristic and perfect fields of characteristic 2, extended in [Dot24] by Dotto to all fields).

The purpose of the present paper is to apply this decomposition to study  $\mathrm{TCR}(\mathbb{Z}/4; 2)^{\phi_{\mathbb{Z}/2}}$ . From this point onwards we will write  $\mathrm{TCR}(A)$  for  $\mathrm{TCR}(A; 2)$ , leaving the prime  $p = 2$  implicit. In Section 2.1 we show that  $(\underline{H\mathbb{Z}/4})^{\phi_{\mathbb{Z}/2}}$  (with the Frobenius module structure, and where the underline indicates we are considering  $\mathbb{Z}/4$  as having trivial involution) splits as an  $\underline{H\mathbb{Z}/4}$ -module into the direct sum of its homotopy groups, and so following the above approach we obtain a decomposition of the equaliser diagram (1) for  $A = \underline{H\mathbb{Z}/4}$ . The homotopy groups of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  are immediate from the decomposition, while to calculate the homotopy groups of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$  we need to analyse summands of the form

$$\left( \underline{H\mathbb{Z}/4} \otimes_{N_{\{e\}}^{C_2}(\underline{H\mathbb{Z}/4})} N_{\{e\}}^{C_2}(\Sigma^n \underline{H\mathbb{Z}/2}) \right)^{C_2}$$

for  $n \in \mathbb{N}$ . In Section 2.2 we show that the homotopy groups of these summands can be expressed using non-abelian derived functors. Non-abelian derived functors were introduced by Dold and Puppe in [DP58] and developed in [DP61]. Just as derived functors are computed using free resolutions in chain complexes, non-abelian derived functors are computed using free resolutions in simplicial modules; we later recall this in Definition 2.8. The key result is the following (proved in Corollary 2.11):

**Lemma A.** *Let  $F_R : \mathrm{Mod}_R \rightarrow \mathrm{Ab}$  defined by  $F_R(M) = (M \otimes_R M)^{C_2}$ . Then*

$$\pi_i \left( \left( \underline{H\mathbb{Z}/4} \otimes_{N_{\{e\}}^{C_2}(\underline{H\mathbb{Z}/4})} N_{\{e\}}^{C_2}(\Sigma^n \underline{H\mathbb{Z}/2}) \right)^{C_2} \right) \cong L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$$

where  $L_i^{(n)} F_R : \mathrm{Mod}_R \rightarrow \mathrm{Ab}$  is the  $i$ th non-abelian derived functor of type  $n$  of  $F_R$ .

To compute these values we initially turn to brute force calculation, using computer assistance to evaluate  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  for  $i, n \leq 6$ .

**Theorem B.** *The low degree homotopy groups of the terms in the equaliser (1) for  $A = \underline{H\mathbb{Z}/4}$  are as follows:*

$i$	$\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})$	$\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$
0	$\mathbb{Z}/4$	$\mathbb{Z}/2$
1	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^3$
2	$(\mathbb{Z}/2)^7$	$(\mathbb{Z}/2)^6$
3	$(\mathbb{Z}/2)^{13}$	$(\mathbb{Z}/2)^{10}$
4	$(\mathbb{Z}/2)^{15} \oplus (\mathbb{Z}/4)^3$	$(\mathbb{Z}/2)^{15}$
5	$(\mathbb{Z}/2)^{27}$	$(\mathbb{Z}/2)^{21}$
6	$(\mathbb{Z}/2)^{34}$	$(\mathbb{Z}/2)^{28}$

In Section 2.3 we continue by computing the action on homotopy of the above-mentioned maps  $f$  and  $r$  up to degree 6. In the case of  $f$  this is a straightforward extension of our computer program; for the map  $r$  we do a more involved analysis. As a result we can write down the long exact sequence associated to the homotopy equaliser of  $f$  and  $r$ , and so obtain short exact sequences describing  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  for  $-1 \leq i \leq 5$ . That is, we compute the desired homotopy groups up to solving certain extension problems. For  $i = -1$  and  $i = 0$  these extension problems are trivial, and in Section 2.4 we solve the extension problem for  $i = 1$ . Our results are summarised in the following:

**Theorem C.** *All non-trivial homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  are in degrees  $\geq -1$ . We compute*

$$\begin{aligned}\pi_{-1}(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) &\cong \mathbb{Z}/2 \\ \pi_0(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) &\cong \mathbb{Z}/4 \\ \pi_1(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) &\cong (\mathbb{Z}/2)^2,\end{aligned}$$

and we have short exact sequences

$$\begin{aligned}0 &\rightarrow \mathbb{Z}/2 \rightarrow \pi_2(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \rightarrow (\mathbb{Z}/2)^2 \rightarrow 0 \\ 0 &\rightarrow (\mathbb{Z}/2)^2 \rightarrow \pi_3(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \rightarrow (\mathbb{Z}/2)^4 \rightarrow 0 \\ 0 &\rightarrow (\mathbb{Z}/2)^4 \rightarrow \pi_4(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \rightarrow (\mathbb{Z}/2)^6 \oplus \mathbb{Z}/4 \rightarrow 0 \\ 0 &\rightarrow (\mathbb{Z}/2)^9 \rightarrow \pi_5(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \rightarrow (\mathbb{Z}/2)^{10} \rightarrow 0.\end{aligned}$$

Moreover the groups  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  are 4-torsion for all  $i$ .

In particular note  $\pi_1$  and  $\pi_5$  cannot be isomorphic. Demonstrating this was part of our original motivation for computing these homotopy groups. As we explain in the following subsection, via showing that the homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  are not 4-periodic in positive degrees, we can conclude that the homotopy groups of the genuine symmetric and symmetric  $L$ -theories of  $\mathbb{Z}/4$  are not canonically isomorphic in non-negative degrees, resolving a question of Calmès et. al from [Cal+21] Example 1.3.11.

In Section 3.1 we return to a more theoretical approach, studying the non-abelian derived functors  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  for general  $i$  and  $n$ . Note that the non-abelian derived functors of  $F_R$  are isomorphic to those of J. H. C. Whitehead's functor  $\Lambda_R$  of [Whi50] (since  $F_R$  and  $\Lambda_R$  agree on free modules); the non-abelian derived functors of  $\Lambda_R$  were previously studied in [ST74] and [Sim74] but most of their results are only valid when 2 is not a zero-divisor in  $R$ . We almost precisely narrow down the value of  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  for all  $i$  and  $n$ , with just slight uncertainties remaining.

**Theorem D.** *For  $i \geq 0$  fixed,  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  takes the same value for all  $0 \leq n \leq \lfloor i/2 \rfloor$ . Moreover we determine this constant value, up to two possibilities in some*

cases:

$$L_i^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong L_i^{(1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong \dots \cong L_i^{\lfloor i/2 \rfloor} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \\ \cong \begin{cases} (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 1} \text{ or } (\mathbb{Z}/2)^{\lfloor i/2 \rfloor} \oplus \mathbb{Z}/4 & \text{if } i \equiv 0 \pmod{4}, \\ (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 1} \text{ or } (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 2} & \text{if } i \equiv 1 \pmod{4}, \\ (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 1} & \text{if } i \equiv 2 \pmod{4}, \\ (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 2} & \text{if } i \equiv 3 \pmod{4}. \end{cases}$$

For each  $k$  we have  $|L_{4k}^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)| = |L_{4k+1}^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)|$ . That is, if we know the value for some  $i$  congruent to 0 mod 4 then this tells us the value for  $i + 1$ .

For  $\lfloor i/2 \rfloor < n \leq i$  we have

$$L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{i-n+1}.$$

And  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) = 0$  for  $n > i$ .

This result, proved in Theorem 3.16, is enough to put quite tight bounds on the order of  $\pi_i((\text{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})$ . In Section 3.2 we use the long exact sequence in homotopy for the equaliser formula (1) to get upper and lower bounds for the orders of the groups  $\pi_i(\text{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ . Despite Theorem C showing that these groups are not 4-periodic in all positive degrees, one might still have imagined that beyond a certain point they could start repeating with some period. The following result (proved in Theorem 3.19) shows that this is not the case; rather, the homotopy groups continue to grow in size.

**Theorem E.** *The logarithm of the order of the group*

$$\pi_i(\text{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$$

*exhibits asymptotically quadratic growth with respect to  $i$ .*

*More precisely, for  $i \geq 0$  we have upper and lower bounds*

$$2^{(i-1)(i+1)/8} \leq |\pi_i(\text{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})| \leq 2^{(i+2)(9i+20)/8}.$$

## Relation to $L$ -theory

Our original motivation for studying the geometric fixed points of TCR was due to connections to  $L$ -theory via a result of Harpaz, Nikolaus and Shah [HNS]. In this subsection we describe the  $L$ -theoretic consequences of our calculations.

Let us first recall the relevant flavours of  $L$ -theory. Let  $R$  be a commutative ring, possibly equipped with an involution. Under the perspective of Calmès et al. [Cal+23b] we can study Poincaré structures on the perfect derived  $\infty$ -category  $\mathcal{D}^p(R)$ , and their associated  $L$ -theory spectra. In particular we have the symmetric structure  $\mathcal{Q}_R^s$  and the genuine symmetric structure  $\mathcal{Q}_R^{gs}$  (see [Cal+23a] 4.2.12, 4.2.13, 4.2.23). The  $L$ -theory spectrum ([Cal+23b] 4.4.4)  $L(\mathcal{D}^p(R), \mathcal{Q}_R^s)$  associated to the symmetric

structure is Ranicki's 4-periodic symmetric  $L$ -theory spectrum, while the spectrum  $L(\mathcal{D}^p(R), \mathcal{Q}_R^{gs})$  associated to the genuine symmetric structure has homotopy groups given by the non-periodic symmetric  $L$ -groups defined by Ranicki in [Ran92].

We would like to understand the relation between these two different  $L$ -theories. There is a canonical comparison map

$$L(\mathcal{D}^p(R), \mathcal{Q}_R^{gs}) \rightarrow L(\mathcal{D}^p(R), \mathcal{Q}_R^s) \quad (2)$$

from the genuine symmetric  $L$ -theory to the symmetric  $L$ -theory. Theorem 6 of [Cal+21] says that if  $R$  is a coherent ring with finite global dimension  $d$ , then this map is an injection on homotopy groups in degree  $d - 2$  and an isomorphism on homotopy in degrees  $\geq d - 1$ . That is, for a sufficiently nice ring  $R$  these  $L$ -theories agree in sufficiently high degrees. In [Cal+21] Example 1.3.11 they show that this bound is tight when allowing rings with involution: for the  $d$ -dimensional ring  $R = F_2[\mathbb{Z}^d]$  with involution induced by the inversion of the group  $\mathbb{Z}^d$ , the map (2) is not surjective in degree  $d - 2$ . However in general there are not many rings  $R$  where both  $L$ -theory spectra have been computed, and after giving the above example Calmès et. al pose the question of identifying any commutative ring with trivial involution such that it is not the case that the map (2) is an isomorphism on all non-negative homotopy groups.

We use our results to answer this question, showing that in the case of  $R = \mathbb{Z}/4$  (considered as having trivial involution) the two  $L$ -theories are far from always coinciding in non-negative degrees.

**Corollary F.** *The comparison map from genuine symmetric to symmetric  $L$ -theories*

$$\pi_i \left( L(\mathcal{D}^p(\mathbb{Z}/4), \mathcal{Q}_{\mathbb{Z}/4}^{gs}) \right) \rightarrow \pi_i \left( L(\mathcal{D}^p(\mathbb{Z}/4), \mathcal{Q}_{\mathbb{Z}/4}^s) \right)$$

*fails to be an isomorphism for some  $0 \leq i \leq 5$ , and moreover for  $i$  sufficiently large it is never an isomorphism.*

*Proof.* We use the result of [HNS] relating  $L$ -theories to TCR. Let  $\mathcal{Q}_R^q$  denote the quadratic Poincaré structure on  $\mathcal{D}^p(R)$ , whose associated  $L$ -theory  $L(\mathcal{D}^p(R), \mathcal{Q}_R^q)$  is Ranicki's 4-periodic quadratic  $L$ -theory spectrum. The Poincaré structure  $\mathcal{Q}_R^q$  is the initial Poincaré structure on  $\mathcal{D}^p(R)$  compatible with the duality  $\text{Hom}(-, R)$  ([Cal+23a] 1.3.5), so there is a canonical map  $L(\mathcal{D}^p(R), \mathcal{Q}_R^q) \rightarrow L(\mathcal{D}^p(R), \mathcal{Q}_R^{gs})$ . The cofibre of this map is known as normal  $L$ -theory. Harpaz, Nikolaus and Shah show that in fact normal  $L$ -theory is equivalent to the geometric fixed points of TCR, so we have a cofibre sequence of spectra

$$L(\mathcal{D}^p(R), \mathcal{Q}_R^q) \rightarrow L(\mathcal{D}^p(R), \mathcal{Q}_R^{gs}) \rightarrow \text{TCR}(R)^{\phi_{\mathbb{Z}/2}}.$$

Consider the map of cofibre sequences

$$\begin{array}{ccccc} L(\mathcal{D}^p(R), \mathcal{Q}_R^q) & \longrightarrow & L(\mathcal{D}^p(R), \mathcal{Q}_R^{gs}) & \longrightarrow & \text{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}} \\ \parallel & & \downarrow & & \downarrow \\ L(\mathcal{D}^p(R), \mathcal{Q}_R^q) & \longrightarrow & L(\mathcal{D}^p(R), \mathcal{Q}_R^s) & \longrightarrow & C, \end{array}$$

where  $C$  is the cofibre of  $L(\mathcal{D}^p(R), \Omega_R^q) \rightarrow L(\mathcal{D}^p(R), \Omega_R^s)$ .

Suppose the map  $L(\mathcal{D}^p(R), \Omega_R^{qs}) \rightarrow L(\mathcal{D}^p(R), \Omega_R^s)$  is an isomorphism on homotopy groups  $\pi_i$  for all  $0 \leq i \leq 5$ . Consider the map of long exact sequences in homotopy groups induced by the above map of cofibre sequences. By the 5-lemma the map  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2} \rightarrow C$  is an isomorphism on  $\pi_i$  for  $1 \leq i \leq 5$ . Observe  $L(\mathcal{D}^p(R), \Omega_R^q) \rightarrow L(\mathcal{D}^p(R), \Omega_R^s)$  is a map of  $L(\mathcal{D}^p(R), \Omega_R^s)$ -modules and  $L(\mathcal{D}^p(R), \Omega_R^s)$  is a 4-periodic ring spectrum, so  $C$  is 4-periodic and hence we conclude that  $\pi_1(\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \cong \pi_5(\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$ . However Theorem C contradicts this, and so our earlier supposition must be false; that is, the comparison map between the homotopy groups of the genuine symmetric and symmetric  $L$ -groups must fail to be an isomorphism somewhere in degrees 0 to 5.

Similarly suppose the map  $L(\mathcal{D}^p(R), \Omega_R^{qs}) \rightarrow L(\mathcal{D}^p(R), \Omega_R^s)$  induces an isomorphism on  $\pi_i$  for some particular  $i$ . Then the 4-lemma shows that  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow \pi_i(C)$  is injective. But  $C$  is 4-periodic and the homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  grow according to Theorem E, so for  $i$  sufficiently large  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$  cannot inject into  $\pi_i(C)$ . We conclude that beyond a certain point the comparison map is never an isomorphism on homotopy.  $\square$

## Computer code and results

The computer program used to support the computations in this paper is available at [https://github.com/thjread/tcr\\_geo\\_fixed](https://github.com/thjread/tcr_geo_fixed), together with the raw program output. The computer code is written in Python, utilising the GAP computational group theory library [Gap24] via SageMath [Sag24].

## Acknowledgements

I would like to thank Emanuele Dotto and Irakli Patchkoria for encouraging me to study these homotopy groups, and for many helpful conversations during the course of the project.

The author is supported by the Warwick Mathematics Institute Centre for Doctoral Training, and gratefully acknowledges funding from the University of Warwick.

# 1 Preliminaries and conventions

We first establish some notation, and review some technical details and background material.

## 1.1 Point-set models and model structures

In a few places our argument depends on using a particular point-set model of spectra. We will work with orthogonal  $\mathbb{Z}/2$ -equivariant spectra, generally following Schwede's lecture notes [Sch23] for the basic constructions described as follows.

We denote the Eilenberg-MacLane spectrum of an abelian group  $A$  by  $HA$ , and model this as an orthogonal spectrum via letting the  $n$ th level  $(HA)_n$  be the reduced  $A$ -linearisation of the  $n$ -sphere  $A[S^n]$ . Similarly we denote the Eilenberg-MacLane spectrum of an abelian group  $A$  with involution  $w$  by  $H(A, w)$ , and model this as an orthogonal  $\mathbb{Z}/2$ -spectrum via letting  $H(A, w)_n = A[S^n]$  with the  $\mathbb{Z}/2$ -action induced by  $w$ . Sometimes for  $A$  an abelian group we want to work with the  $\mathbb{Z}/2$ -spectrum  $H(A, \text{id}_A)$ , the Eilenberg-MacLane spectrum of  $A$  considered as having trivial involution; for convenience we still denote this  $HA$ , and whether we mean a spectrum or a  $\mathbb{Z}/2$ -spectrum should be clear from context. We define the geometric fixed points of a  $\mathbb{Z}/2$ -spectrum  $X$  via  $X^{\phi_{\mathbb{Z}/2}}(M) := X(M \otimes_{\mathbb{S}} \rho_{\mathbb{Z}/2})^{\mathbb{Z}/2}$ , and we define the non-derived version of the norm of a spectrum  $Y$  via  $\tilde{N}_{\{e\}}^{\mathbb{Z}/2}(Y) := Y \otimes_{\mathbb{S}} Y$  where  $\mathbb{Z}/2$  acts by swapping the factors.

We work with the flat model structure of [Sto11], [BDS22]. This in particular means that given a strictly commutative ring spectrum we can choose a strictly commutative and flat cofibrant replacement. Except when explicitly stated we will always want to work with derived constructions on spectra. We let  $N_{\{e\}}^{C_2}$  denote the derived Hill-Hopkins-Ravenel norm (i.e. taking a cofibrant replacement followed by applying  $\tilde{N}_{\{e\}}^{C_2}$ ) and write  $\otimes_R$  for the derived smash product relative to a ring spectrum  $R$ . Note that the above geometric fixed point construction does not need to be derived: it gives the correct homotopy type for any  $\mathbb{Z}/2$ -spectrum  $X$  without needing to take a cofibrant replacement.

For a  $\mathbb{Z}/2$ -spectrum  $X$  we denote the homotopy groups of the genuine  $C_2$ -fixed points by

$$\pi_i^{C_2}(X) := \pi_i(X^{C_2}).$$

We now describe the Frobenius module structure on the geometric fixed points of a commutative ring spectrum ([DMPR21] Section 2.5). There is a canonical map

$$Y \xrightarrow{\Delta} (\tilde{N}_{\{e\}}^{C_2}(Y))^{\phi_{C_2}},$$

which is an equivalence of  $\mathbb{Z}/2$ -spectra whenever  $Y$  is cofibrant. If  $R$  is a commutative ring spectrum then there is a canonical multiplication map

$$\tilde{N}_{\{e\}}^{C_2}(R) \xrightarrow{\mu} R.$$

Using these maps we define the Frobenius module structure as follows.

**Definition 1.1.** The Frobenius  $R$ -module structure on  $R^{\phi_{\mathbb{Z}/2}}$  is defined by the map

$$R \xrightarrow{\Delta} (N_{\{e\}}^{C_2}(R))^{\phi_{C_2}} \xrightarrow{\mu^{\phi_{\mathbb{Z}/2}}} R^{\phi_{\mathbb{Z}/2}}.$$

In fact although this definition uses the derived norm, we could have equally used the underived norm.

**Lemma 1.2.** *The map defining the Frobenius module structure is homotopic to*

$$R \xrightarrow{\Delta} (\tilde{N}_{\{e\}}^{C_2}(R))^{\phi_{C_2}} \xrightarrow{\mu^{\phi_{\mathbb{Z}/2}}} R^{\phi_{\mathbb{Z}/2}}.$$

*Proof.* Taking a commutative cofibrant replacement  $R^{\text{cof}}$  for  $R$ , we have a commutative diagram

$$\begin{array}{ccccc}
R^{\text{cof}} & \xrightarrow{\Delta} & \left( \tilde{N}_{\{e\}}^{\mathbb{Z}/2}(R^{\text{cof}}) \right)^{\phi_{\mathbb{Z}/2}} & \xrightarrow{\mu^{\phi_{\mathbb{Z}/2}}} & (R^{\text{cof}})^{\phi_{\mathbb{Z}/2}} \\
\downarrow \simeq & & \downarrow & & \downarrow \simeq \\
R & \xrightarrow{\Delta} & \left( \tilde{N}_{\{e\}}^{\mathbb{Z}/2} R \right)^{\phi_{\mathbb{Z}/2}} & \xrightarrow{\mu^{\phi_{\mathbb{Z}/2}}} & R^{\phi_{\mathbb{Z}/2}}.
\end{array}$$

Now consider the two paths around the circumference from  $R$  to  $R^{\phi_{\mathbb{Z}/2}}$ . □

## 1.2 Dold-Kan equivalence

Let  $R$  a commutative ring. The Dold-Kan equivalence [Dol58] is an equivalence between the category of connective chain complexes of  $R$ -modules  $\text{Ch}_{\geq 0}(R)$  and the category of simplicial  $R$ -modules  $\text{sMod}_R$ . Let us briefly recall the definitions of these functors, since these constructions will be an essential part of our calculations and we will need to reason about the concrete details later on.

We denote the functor from chain complexes to simplicial modules by

$$\sigma : \text{Ch}_{\geq 0}(R) \rightarrow \text{sMod}_R.$$

Let  $M_*$  be a connective chain complex. Then the simplicial  $R$ -module  $\sigma M_\bullet$  is given level-wise by

$$\sigma M_n = \bigoplus_{[n] \twoheadrightarrow [k]} M_k$$

where the sum is taken over all surjections out of  $[n]$  in the simplex category  $\Delta$ ; that is, all surjective weakly increasing maps  $[n] \twoheadrightarrow [k]$  for  $0 \leq k \leq n$ . The simplicial structure is as follows. Given a map  $[m] \rightarrow [n]$  in  $\Delta$ , we need to produce a map  $\sigma M_n \rightarrow \sigma M_m$ . It suffices to describe how to map out of each direct summand of  $\sigma M_n$ , and so for each surjection  $[n] \twoheadrightarrow [k]$  we will give a map  $M_k \rightarrow \sigma M_m$ . Factor the composite  $[m] \rightarrow [n] \twoheadrightarrow [k]$  as a surjection followed by an injection  $[m] \twoheadrightarrow [m'] \hookrightarrow [k]$ , then we take the map  $M_k \rightarrow M_{m'}$  from the chain complex structure of  $M$ , and compose it with the summand inclusion  $M_{m'} \leq \sigma M_m$  corresponding to  $[m] \twoheadrightarrow [m']$ .

The inverse functor from simplicial modules to chain complexes is the normalised chain complex functor

$$N : \text{sMod}_R \rightarrow \text{Ch}_{\geq 0}(R).$$

Given a simplicial  $R$ -module  $M_\bullet$ , we define  $NM_*$  as follows: let  $NA_n$  be the subgroup of  $A_n$  that is killed by all the face maps  $d_i$  for  $i < n$ , and let the differential  $NA_n \rightarrow NA_{n-1}$  be given by  $(-1)^n d_n$ .

**Theorem 1.3** (Dold-Kan equivalence [Dol58]). *The functors  $\sigma$  and  $N$  give an equivalence of categories*

$$\text{Ch}_{\geq 0}(R) \simeq \text{sMod}_R.$$

*Proof.* See [GJ09] Section III.2 (and note the correction to the proof in the errata). □

Additionally recall the Moore complex of a simplicial  $R$ -module. Given a simplicial  $R$ -module  $A_\bullet$ , the Moore complex is the chain complex  $A_*$  which is given by  $A_n$  in degree  $n$ , and has boundary map  $\partial : A_n \rightarrow A_{n-1}$  given by the alternating sum of the face maps  $\sum_{i=0}^n (-1)^i d_i$ . The Moore complex  $A_*$  and the normalised chain complex  $NA_*$  are naturally homotopy equivalent, and so in practice we can use either in situations where we only care about the result up to homotopy.

### 1.3 Real topological Hochschild homology

In this subsection we briefly recall the construction of  $\mathrm{THR}(A)$  and  $\mathrm{TCR}(A; p)$  in the special case where  $A$  is a discrete commutative ring with trivial involution (this is the only case we will need). These constructions were introduced in [HM15], [Dot12], [Høg16]; good sources for full details include [DMPR21] and [DMP24].

Let  $HA$  be a flat model for the Eilenberg-MacLane spectrum of  $A$ , in orthogonal spectra. The cyclic nerve of  $HA$  becomes a dihedral spectrum if we give it an appropriate involution. We denote this construction  $N^{di}HA$ . In level  $n$  we have  $(N^{di}HA)_n = HA^{\otimes n+1}$ , which we visualise as  $n+1$  copies of  $A$  tensored together in a circle, equipped with rotation and reflection actions. We define the real topological Hochschild homology of  $A$  to be the geometric realisation

$$\mathrm{THR}(A) := |N^{di}A|.$$

This is an orthogonal spectrum with an  $O(2)$ -action, so we can regard it as a genuine  $O(2)$ -spectrum. Note the underlying cyclotomic spectrum of  $\mathrm{THR}(A)$  is  $\mathrm{THH}(A)$ , while the underlying genuine  $\mathbb{Z}/2$ -spectrum is

$$A \otimes_{N_e^{\mathbb{Z}/2}A} A$$

(where  $A$  has  $N_e^{\mathbb{Z}/2}A$ -module structure given by  $N_e^{\mathbb{Z}/2}A \xrightarrow{\mu} A$ ). In particular we find that the geometric fixed points are

$$\mathrm{THR}(A)^{\phi\mathbb{Z}/2} \simeq A^{\phi\mathbb{Z}/2} \otimes_A A^{\phi\mathbb{Z}/2}.$$

Let  $p$  a prime. We define the  $p$ -typical truncated real topological restriction homology of  $A$  by

$$\mathrm{TRR}^{n+1}(A; p) := \mathrm{THR}(A)^{C_{p^n}},$$

where  $C_{p^n} \leq D_{p^n} \leq O(2)$ . We can consider this as a  $\mathbb{Z}/2$ -spectrum, since the Weyl group of  $C_{p^n}$  inside  $D_{p^n}$  is  $\mathbb{Z}/2$ . There is a  $\mathbb{Z}/2$ -equivariant map  $R : \mathrm{TRR}^{n+1}(A; p) \rightarrow \mathrm{TRR}^n(A; p)$  coming from the real cyclotomic structure of  $\mathrm{THR}(A)$ ; using this we define the real topological restriction homology

$$\mathrm{TRR}(A; p) := \mathrm{holim} \left( \cdots \xrightarrow{R} \mathrm{TRR}^{n+1}(A; p) \xrightarrow{R} \mathrm{TRR}^n(A; p) \xrightarrow{R} \cdots \xrightarrow{R} \mathrm{TRR}^1(A; p) \right).$$

The inclusion  $C_{p^{n-1}} \leq C_{p^n}$  induces a  $\mathbb{Z}/2$ -equivariant restriction map

$$F : \mathrm{TRR}^n(A; p) = \mathrm{THR}(A)^{C_{p^n}} \rightarrow \mathrm{THR}(A)^{C_{p^{n-1}}} = \mathrm{TRR}^{n-1}(A; p),$$

known as the Frobenius. This in turn induces an endomorphism  $F$  on  $\mathrm{TRR}(A; p)$ . Finally we define the real topological cyclic homology of  $A$  as the  $\mathbb{Z}/2$ -equivariant spectrum

$$\mathrm{TCR}(A; p) := \mathrm{hoeq}(\mathrm{TRR}(A; p) \underset{\mathrm{id}}{\overset{F}{\rightrightarrows}} \mathrm{TRR}(A; p)).$$

This construction of TCR from THR is analogous to the usual construction of TC from THH, and indeed the underlying spectrum of  $\mathrm{TCR}(A; p)$  is  $\mathrm{TC}(A; p)$ .

As mentioned in the introduction, we will not work directly with the above construction of TCR, but instead use the following formula of Dotto, Moi and Patchkoria.

**Theorem 1.4** ([DMP24] 2.14). *We can describe topological cyclic restriction homology by the following homotopy equaliser*

$$\mathrm{TCR}(A)^{\phi_{\mathbb{Z}/2}} \simeq \mathrm{hoeq} \left( (\mathrm{THR}(A)^{\phi_{\mathbb{Z}/2}})^{C_2} \underset{r}{\overset{f}{\rightrightarrows}} \mathrm{THR}(A)^{\phi_{\mathbb{Z}/2}} \right).$$

(see [DMP24] for descriptions of the maps, and we will later give an explicit description in the case  $A = H\mathbb{Z}/4$ ).

## 2 Low degree homotopy of $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$

In this section we compute the low degree terms of the long exact sequence in homotopy groups corresponding to the equaliser formula of Theorem 1.4, and make some progress solving the resulting extension problems. As a result we compute  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  for  $i \leq 1$  and obtain short exact sequences for these homotopy groups for  $2 \leq i \leq 5$ .

### 2.1 Decomposition of the equaliser formula for $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$

In order to compute the homotopy groups of the terms in the equaliser, we want to apply Lemma 4.3 of [DMP24] to decompose the copies of  $\mathrm{THR}(\mathbb{Z}/4^{\phi_{\mathbb{Z}/2}})$ . But to satisfy the conditions of the lemma we first need to show that  $H\mathbb{Z}/4^{\phi_{\mathbb{Z}/2}}$ , endowed with the Frobenius  $H\mathbb{Z}/4$ -module structure, splits as the direct sum of its homotopy groups.

**Lemma 2.1.** *For an abelian group  $A$  we have*

$$\pi_i(HA^{\phi_{\mathbb{Z}/2}}) \cong \begin{cases} \mathrm{coker}(A \xrightarrow{2} A) & i \geq 0, i \text{ even}, \\ \mathrm{ker}(A \xrightarrow{2} A) & i \geq 0, i \text{ odd}, \\ 0 & i < 0. \end{cases}$$

*Proof.* The spectrum  $HA$  is connective, so its geometric fixed points are too. Now consider the cofibre sequence

$$HA_{h\mathbb{Z}/2} \rightarrow HA \rightarrow HA^{\phi_{\mathbb{Z}/2}}.$$

Since  $HA$  is concentrated in degree 0, the long exact sequence in homotopy groups gives us an exact sequence

$$0 \rightarrow \pi_1(HA^{\phi\mathbb{Z}/2}) \rightarrow \pi_0(HA_{h\mathbb{Z}/2}) \rightarrow \pi_0(HA) \rightarrow \pi_0(HA^{\phi\mathbb{Z}/2}) \rightarrow 0 \quad (3)$$

and isomorphisms

$$\pi_i(HA^{\phi\mathbb{Z}/2}) \cong \pi_{i-1}(HA_{h\mathbb{Z}/2})$$

for  $i \geq 2$ .

But we have

$$\pi_i(HA_{h\mathbb{Z}/2}) \cong H_i(\mathbb{RP}^\infty; A) \cong \begin{cases} A & i = 0, \\ \text{coker}(A \xrightarrow{2} A) & i > 0, i \text{ odd}, \\ \text{ker}(A \xrightarrow{2} A) & i > 0, i \text{ even}. \end{cases}$$

And the map  $A \cong \pi_0(HA_{h\mathbb{Z}/2}) \rightarrow \pi_0(HA) \cong A$  appearing in the exact sequence (3) is the multiplication map  $A \xrightarrow{2} A$ . The result follows.  $\square$

**Lemma 2.2.** *We have a splitting of  $H\mathbb{Z}/4$ -modules*

$$H\mathbb{Z}/4^{\phi\mathbb{Z}/2} \simeq \bigoplus_{n \geq 0} \Sigma^n H\pi_n(H\mathbb{Z}/4^{\phi\mathbb{Z}/2}) \cong \bigoplus_{n \geq 0} \Sigma^n H\mathbb{Z}/2.$$

where we consider  $H\mathbb{Z}/4^{\phi\mathbb{Z}/2}$  as an  $H\mathbb{Z}/4$ -module via the Frobenius module structure of Definition 1.1 and consider  $H\mathbb{Z}/2$  as an  $H\mathbb{Z}/4$ -module via the quotient map  $H\mathbb{Z}/4 \rightarrow H\mathbb{Z}/2$ .

*Proof.* First note that by Lemma 2.1 we indeed have  $\pi_n(H\mathbb{Z}/4^{\phi\mathbb{Z}/2}) \cong \mathbb{Z}/2$  for  $n \geq 0$ .

Recall Lemma 1.2, which says that the Frobenius module structure on  $H\mathbb{Z}/4^{\phi\mathbb{Z}/2}$  can be defined by the map

$$\varphi : H\mathbb{Z}/4 \xrightarrow{\Delta} (\tilde{N}_{\{e\}}^{\mathbb{Z}/2}(H\mathbb{Z}/4))^{\phi\mathbb{Z}/2} \xrightarrow{\mu^{\phi\mathbb{Z}/2}} H\mathbb{Z}/4^{\phi\mathbb{Z}/2}. \quad (4)$$

We claim that  $\varphi$  factors along the quotient  $H\mathbb{Z}/4 \twoheadrightarrow H\mathbb{Z}/2$ . This then implies the lemma, since it means the Frobenius  $H\mathbb{Z}/4$ -module structure is the pullback of a  $H\mathbb{Z}/2$ -module structure, and  $\mathbb{Z}/2$  is a field so all  $H\mathbb{Z}/2$ -modules split uniquely as the sum of their homotopy groups.

Let the ideal  $I = 2\mathbb{Z}/4$  be the kernel of the quotient  $\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2$ . If we were just working with discrete rings, then to show that a ring homomorphism out of  $\mathbb{Z}/4$  factors through the quotient  $\mathbb{Z}/2$  it would suffice to show that the homomorphism is zero on  $I$ . We will use the technology of Hovey [Hov14] to apply a similar argument to ring spectra.

Considering  $I$  as a non-unital ring spectrum, naturality with respect to  $I \hookrightarrow \mathbb{Z}/4$  gives a commutative diagram

$$\begin{array}{ccccc} HI & \xrightarrow{\Delta} & (\tilde{N}_{\{e\}}^{\mathbb{Z}/2}(HI))^{\phi\mathbb{Z}/2} & \xrightarrow{\mu^{\phi\mathbb{Z}/2}} & HI^{\phi\mathbb{Z}/2} \\ \downarrow & & \downarrow & & \downarrow \\ H\mathbb{Z}/4 & \xrightarrow{\Delta} & (\tilde{N}_{\{e\}}^{\mathbb{Z}/2}(H\mathbb{Z}/4))^{\phi\mathbb{Z}/2} & \xrightarrow{\mu^{\phi\mathbb{Z}/2}} & H\mathbb{Z}/4^{\phi\mathbb{Z}/2}. \end{array}$$

The multiplication map  $\tilde{N}_{\{e\}}^{\mathbb{Z}/2}(HI) \xrightarrow{\mu} HI$  can be expressed as

$$\tilde{N}_{\{e\}}^{\mathbb{Z}/2}(HI) \rightarrow H(I \otimes I) \xrightarrow{H(\nu)} HI$$

where the first map is an equivalence of underlying spectra and  $\nu : I \otimes I \rightarrow I$  is the multiplication map on  $I$ . But the ring multiplication of  $I$  is zero, so we conclude  $\mu$  is also the zero map. Taking the geometric fixed points of the zero map gives the zero map, so in the top row  $\mu^{\phi_{\mathbb{Z}/2}} = 0$ . By commutativity of the diagram we see that the composite  $HI \hookrightarrow H\mathbb{Z}/4 \xrightarrow{\varphi} H\mathbb{Z}/4^{\phi_{\mathbb{Z}/2}}$  is zero (on the nose, not just null-homotopic).

Now we apply Theorem 1.4 of [Hov14]. We have shown there is a map of Smith ideals

$$\begin{array}{ccc} HI & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ H\mathbb{Z}/4 & \xrightarrow{\varphi} & (H\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}} \end{array}$$

and so taking cofibres we get a commutative square of ring spectra

$$\begin{array}{ccc} H\mathbb{Z}/4 & \xrightarrow{\varphi} & (H\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}} \\ \downarrow & & \parallel \\ H\mathbb{Z}/2 & \longrightarrow & (H\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}, \end{array}$$

giving the desired factorisation.  $\square$

Now by Lemma 4.3 of [DMP24] we get decompositions of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  and  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$ .

**Corollary 2.3.** *We have an equivalence of  $C_2$ -spectra*

$$\begin{aligned} \mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}} &\simeq \bigoplus_{n \geq 0} \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \\ &\quad \oplus \bigoplus_{0 \leq n < m} \Sigma^{n+m} (C_2)_+ \otimes_{\mathbb{S}} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \end{aligned}$$

(where  $\rho$  is the regular representation of  $C_2$  and  $(C_2)_+ \otimes_{\mathbb{S}} (-)$  denotes the left adjoint of the restriction functor from  $\mathbb{Z}/2$ -spectra to spectra) and an equivalence of spectra

$$\begin{aligned} (\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2} &\simeq \bigoplus_{n \geq 0} \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2} \\ &\quad \oplus \bigoplus_{0 \leq n < m} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2. \end{aligned}$$

Note that we are using the  $N_{\{e\}}^{C_2}(H\mathbb{Z}/4)$ -module structure on  $H\mathbb{Z}/4$  defined by the multiplication map  $\mu : N_{\{e\}}^{C_2}(H\mathbb{Z}/4) \rightarrow H\mathbb{Z}/4$ .  $\square$

*Remark 2.4.* Observe that on underlying spectra the first equivalence gives

$$\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}} \simeq \bigoplus_{n,m \geq 0} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2.$$

Again by [DMP24] Lemma 4.3, we get descriptions in terms of these decompositions for the two maps  $f, r : (\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2} \rightarrow \mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  whose homotopy equaliser computes  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ .

**Corollary 2.5.** *The map  $f : (\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2} \rightarrow \mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/4}}$  splits up according to the direct sum decomposition of Corollary 2.3. It is defined on the  $n$ -indexed summands by restriction maps*

$$\mathrm{res}_e^{C_2} : \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2} \rightarrow \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2),$$

and on the  $(n, m)$ -indexed summands by diagonal maps

$$\Delta : \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \rightarrow \Sigma^{n+m}(C_2)_+ \otimes_{\mathbb{S}} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2.$$

The map  $r$  kills the  $(n, m)$ -summands of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$ , and is defined on the  $n$ -summands by

$$\begin{aligned} \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2} &\rightarrow \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{\phi_{C_2}} \\ &\simeq \Sigma^n (H\mathbb{Z}/4)^{\phi_{C_2}} \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \\ &\simeq \bigoplus_{m \geq 0} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \\ &\rightarrow \mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}} \end{aligned}$$

where the first map is the canonical map from genuine to geometric fixed points, the second map comes from monoidality of geometric fixed points, the third map is the splitting of Lemma 2.2, and the final map is the inclusion of a direct summand according to the decomposition of Remark 2.4.  $\square$

## 2.2 Homotopy groups of $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ and $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$

Now let us discuss the computation of the homotopy groups of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  and  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$ .

**Proposition 2.6.** *We have*

$$\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \cong \bigoplus_{\substack{n,m,k \geq 0 \\ n+m+k=i}} \mathbb{Z}/2.$$

*Proof.* Considering a free resolution of  $H\mathbb{Z}/2$  as an  $H\mathbb{Z}/4$ -module, then we can compute

$$H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \simeq \bigoplus_{k \geq 0} \Sigma^k H\mathbb{Z}/2.$$

So by Corollary 2.3 and Remark 2.4 we get an equivalence of underlying spectra

$$\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2} \simeq \bigoplus_{n,m,k \geq 0} \Sigma^{n+m+k} H\mathbb{Z}/2$$

and the result follows.  $\square$

**Proposition 2.7.** *Looking at just the off-diagonal part of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$  under the decomposition of Corollary 2.3, we get*

$$\pi_i \left( \bigoplus_{0 \leq n < m} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \right) \cong \bigoplus_{\substack{n,m,k \geq 0 \\ n < m \\ n+m+k=i}} \mathbb{Z}/2.$$

*Proof.* Similarly to Proposition 2.6, we have an equivalence of spectra

$$\bigoplus_{0 \leq n < m} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \simeq \bigoplus_{0 \leq n < m} \bigoplus_{k \geq 0} \Sigma^{n+m+k} H\mathbb{Z}/2$$

giving the result.  $\square$

So we just need to focus on calculating the homotopy groups of the diagonal summands of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$ , which are of the form

$$\left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2}$$

for  $n \geq 0$ . We approach this using the concept of non-abelian derived functors, as defined in [DP58]. First let us recall the definition.

Let  $R$  be a commutative ring and let  $T : \mathrm{Mod}_R \rightarrow \mathrm{Ab}$  be a functor with  $T(0) = 0$ . Let  $i \geq 0$  and  $n \geq 0$  be natural numbers. Given an  $R$ -module  $M$ , let  $M_\bullet$  be a level-wise free simplicial  $R$ -module with geometric realisation  $\Sigma^n T M$  (for example take a free resolution for  $M[n]$  in chain complexes of  $R$ -modules, then apply the Dold-Kan equivalence of Theorem 1.3). Then we define a functor

$$L_i^{(n)} T : \mathrm{Mod}_R \rightarrow \mathrm{Ab}$$

via

$$L_i^{(n)} T(M) := \pi_i(|T(M_\bullet)|)$$

where  $T(M_\bullet)$  means the simplicial abelian group produced by applying  $T$  level-wise to  $M_\bullet$ .

**Definition 2.8** (Non-abelian derived functor). We call  $L_i^{(n)} T$  the  $i$ th non-abelian derived functor of type  $n$  of  $T$ .

*Remark 2.9.* Observe that  $H(M_\bullet)$  is zero in degrees less than  $n$ . So  $L_i^{(n)}H(M) = 0$  for  $i < n$ .

**Lemma 2.10.** *Let  $R$  be a commutative ring and  $M$  an  $R$ -module. Let  $M_\bullet$  a free simplicial  $R$ -module with realisation  $\Sigma^n HM$ . Then*

$$\Sigma^{n\rho} HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM) \simeq |H(M_\bullet \otimes_R M_\bullet, w)|,$$

where  $w$  is the involution on  $M_i \otimes_R M_i$  given by swapping the two copies of  $M_i$ . This equivalence is natural in  $M_\bullet$  (if we take  $M$  to depend functorially on  $M_\bullet$ ).

*Proof.* First rewrite as

$$\begin{aligned} \Sigma^{n\rho} HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM) &\simeq HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(\Sigma^n HM) \\ &\simeq \left| HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM_\bullet) \right|, \end{aligned}$$

where the first equivalence comes from the interaction of suspension and the norm construction, and the second equivalence uses the fact that geometric realisation commutes with the norm and tensor products.

Let  $B_i$  index a basis of the free module  $M_i$ , i.e.  $M_i \cong \bigoplus_{B_i} R$ . We first establish some notation. Consider the set  $(B_i \times B_i)_{C_2}$ , where  $C_2$  acts on  $B_i \times B_i$  by swapping the two factors. Denote the diagonal part by  $B_i^d = \{(u, u) \in (B_i \times B_i)_{C_2}\} \cong B_i$  and denote the off-diagonal part by  $B_i^o = (B_i \times B_i)_{C_2} \setminus B_i^d$ . Then we have a series of equivalences of simplicial  $C_2$ -spectra

$$\begin{aligned} &HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM_\bullet) \\ &\simeq HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2} \left( \bigoplus_{B_\bullet} HR \right) \\ &\simeq HR \otimes_{N_{\{e\}}^{C_2}(HR)} \left( \bigoplus_{B_\bullet^d} N_{\{e\}}^{C_2}(HR) \oplus \bigoplus_{B_\bullet^o} \left( \Sigma^\infty(C_2)_+ \otimes_{\mathbb{S}} N_{\{e\}}^{C_2}(HR) \right) \right) \\ &\simeq \bigoplus_{B_\bullet^d} HR \oplus \bigoplus_{B_\bullet^o} \left( \Sigma^\infty(C_2)_+ \otimes_{\mathbb{S}} HR \right) \\ &\simeq H \left( \bigoplus_{B_\bullet \times B_\bullet} R, w' \right) \\ &\simeq H(M_\bullet \otimes_R M_\bullet, w), \end{aligned}$$

where  $w'$  is the involution on  $\bigoplus_{B_i \times B_i} R$  given by swapping the two copies of  $B_i$ , and which under  $\bigoplus_{B_i \times B_i} R \cong M_i \otimes_R M_i$  becomes the involution  $w$  described in the lemma statement.

To see naturality, consider the map of spectra

$$HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM_i) \rightarrow H(M_i \otimes M_i)$$

obtained as the canonical map from the left hand side to the Eilenberg-MacLane spectrum of its degree zero homotopy, and observe that taking geometric realisations gives the map in the statement of the lemma; from this definition of the map it is clear that it is natural in  $M_\bullet$ .  $\square$

The immediate consequence of this theorem is that we can describe some homotopy groups of interest using non-abelian derived functors. We present these generally but note the case we will be interested in is when  $R = \mathbb{Z}/4$  and  $M = \mathbb{Z}/2$ . First we consider the homotopy groups of the genuine  $C_2$ -fixed points; this appeared as Lemma A in the introduction. Define  $F_R : \text{Mod}_R \rightarrow \text{Ab}$  on an  $R$ -module  $M$  by  $F_R(M) = (M \otimes_R M)^{C_2}$ , where  $M \otimes_R M$  has the  $C_2$ -action given by swapping the factors.

**Corollary 2.11.** *We have*

$$\pi_i^{C_2} \left( \Sigma^{n\rho} HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM) \right) \cong L_i^{(n)} F_R(M)$$

where  $L_i^{(n)} F_R$  denotes the  $i$ th non-abelian derived functor of type  $n$  of  $F_R$ .

*Proof.* We have

$$\begin{aligned} \pi_i^{C_2} \left( \Sigma^{n\rho} HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM) \right) &\cong \pi_i^{C_2} (|H(M_\bullet \otimes_R M_\bullet, w)|) \\ &\cong \pi_i (|H F_R(M_\bullet)|) \\ &\cong L_i^{(n)} F_R(M). \end{aligned} \quad \square$$

Next we consider the homotopy groups of the underlying non-equivariant spectrum. Define  $G_R : \text{Mod}_R \rightarrow \text{Ab}$  on an  $R$ -module  $M$  by  $G_R(M) = M \otimes_R M$ .

**Corollary 2.12.** *We have*

$$\pi_i \left( \Sigma^{n\rho} HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM) \right) \cong L_i^{(n)} G_R(M)$$

where  $L_i^{(n)} G_R$  denotes the  $i$ th non-abelian derived functor of type  $n$  of  $G_R$ .

*Proof.* We have

$$\begin{aligned} \pi_i \left( \Sigma^{n\rho} HR \otimes_{N_{\{e\}}^{C_2}(HR)} N_{\{e\}}^{C_2}(HM) \right) &\cong \pi_i (|H(M_\bullet \otimes_{\mathbb{Z}/4} M_\bullet)|) \\ &\cong L_i^{(n)} G_R(M). \end{aligned} \quad \square$$

However this time when  $R = \mathbb{Z}/4$  and  $M = \mathbb{Z}/2$  we can also compute the homotopy groups directly, giving us a computation of  $L_i^{(n)} G_{\mathbb{Z}/4}(\mathbb{Z}/2)$ .

**Lemma 2.13.** *We have*

$$L_i^{(n)} G_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong \begin{cases} 0 & \text{if } i < 2n \\ \mathbb{Z}/2 & \text{if } i \geq 2n. \end{cases}$$

*Proof.* On underlying spectra we have

$$\begin{aligned} \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) &\cong \Sigma^{2n}(H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2) \\ &\cong \bigoplus_{k \geq 0} \Sigma^{2n+k} H\mathbb{Z}/2 \end{aligned}$$

and observe

$$\pi_i \left( \bigoplus_{k \geq 0} \Sigma^{2n+k} H\mathbb{Z}/2 \right) \cong \begin{cases} 0 & \text{if } i < 2n \\ \mathbb{Z}/2 & \text{if } i \geq 2n. \end{cases}$$

Now by Corollary 2.12 the result follows.  $\square$

For small  $i$  and  $n$  we can compute  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  directly with computer assistance. We work with the free resolution of  $\mathbb{Z}/2[n]$  in chain complexes given by

$$\cdots \xrightarrow{2} \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \rightarrow 0 \rightarrow \cdots,$$

where the rightmost  $\mathbb{Z}/4$  is in degree  $n$ . Applying the Dold-Kan equivalence (recalled in Theorem 1.3) between chain complexes and simplicial modules gives a level-wise free simplicial  $\mathbb{Z}/4$ -module  $M_\bullet$  with geometric realisation  $\Sigma^n H\mathbb{Z}/2$ . We use this to compute the non-abelian derived functor exactly as defined in Definition 2.8: apply  $F_{\mathbb{Z}/4}$  level-wise, then compute homology groups of the Moore complex.

Without any particularly sophisticated optimisation and using a normal laptop computer, we managed to compute just far enough to prove the non-periodicity result mentioned in the introduction. However the simplicial structure coming from Dold-Kan gets exponentially larger as the degree increases, so even with more serious computing power we wouldn't be able to go much farther without significant algorithmic improvements.

Combining the computations from the computer program with our earlier discussion, we obtain the following. These results appeared as Theorem B in the introduction.

**Theorem 2.14.** *The low degree homotopy groups of the terms in the equaliser of Theorem 1.4 computing  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  are as follows:*

$i$	$\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})$	$\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$
0	$\mathbb{Z}/4$	$\mathbb{Z}/2$
1	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^3$
2	$(\mathbb{Z}/2)^7$	$(\mathbb{Z}/2)^6$
3	$(\mathbb{Z}/2)^{13}$	$(\mathbb{Z}/2)^{10}$
4	$(\mathbb{Z}/2)^{15} \oplus (\mathbb{Z}/4)^3$	$(\mathbb{Z}/2)^{15}$
5	$(\mathbb{Z}/2)^{27}$	$(\mathbb{Z}/2)^{21}$
6	$(\mathbb{Z}/2)^{34}$	$(\mathbb{Z}/2)^{28}$

*Proof.* Proposition 2.6 computed  $\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ . For  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$  we work with the decomposition of Corollary 2.3. Proposition 2.7 computed the homotopy

groups of the off-diagonal part of the decomposition. We use a computer program to compute the homotopy groups of the  $n$ -indexed diagonal summands of the decomposition, which by Corollary 2.11 are given by the values of the non-abelian derived functors  $L_i^{(n)}F_R$  at  $\mathbb{Z}/2$ .

For example at  $i = 1$  the computer program calculates

$$L_1^{(0)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^2 \quad L_1^{(1)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong \mathbb{Z}/2$$

(and  $L_1^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) = 0$  for  $n > 1$ ) so we get

$$\begin{aligned} \pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}) &\cong \bigoplus_{0 \leq n \leq 1} L_1^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) \oplus \bigoplus_{\substack{n,m,k \geq 0 \\ n < m \\ n+m+k=1}} \mathbb{Z}/2 \\ &\cong (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2 \\ &\cong (\mathbb{Z}/2)^4. \end{aligned}$$

See the code repository for the full output for all  $i$ . □

### 2.3 Action of the maps $f$ and $r$ on homotopy

We can go further and compute the maps  $f$  and  $r$  on the homotopy groups that we have calculated, allowing us to obtain short exact sequences describing the homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ . Recall the descriptions of  $f$  and  $r$  in Corollary 2.5, defined in terms of the decompositions in Corollary 2.3. The actions of the maps on the  $(n, m)$ -summands of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$  are straightforward:  $f$  acts by the diagonal map, and  $r$  is zero. This section is mostly occupied by analysing what happens on the diagonal  $n$ -indexed summands.

First let us consider how  $f$  acts on the homotopy groups of the diagonal summands. For each  $n \geq 0$ , the map  $f$  takes the  $n$ -indexed summand of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$  to the  $n$ -indexed summand of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  by the restriction map. Recall from Corollary 2.11 that the degree  $i$  homotopy of the  $n$ -indexed summand of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2}$  is isomorphic to  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$ , and recall from Corollary 2.12 and Lemma 2.13 that the degree  $i$  homotopy of the  $n$ -indexed summand of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  is isomorphic to  $L_i^{(n)}G_{\mathbb{Z}/4}(\mathbb{Z}/2)$ , which is 0 for  $i < 2n$  and  $\mathbb{Z}/2$  for  $i \geq 2n$ . Then we find:

**Proposition 2.15.** *By Corollary 2.5 the map  $f$  restricted to the  $n$ -indexed diagonal summands of the source and target is the restriction map*

$$\pi_i^{C_2} \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right) \rightarrow \pi_i \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right).$$

*Under the isomorphisms between these groups and non-abelian derived functors, we claim this becomes the map*

$$L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_i^{(n)}G_{\mathbb{Z}/4}(\mathbb{Z}/2)$$

*induced by the natural transformation  $F_R \Rightarrow G_R$  given at a module  $M$  by inclusion of fixed points  $(M \otimes_R M)^{C_2} \hookrightarrow M \otimes_R M$ .*

*Proof.* Let  $M_\bullet$  be a free simplicial  $\mathbb{Z}/4$ -module with realisation  $\Sigma^n \mathbb{Z}/2$ . Under the equivalence of Lemma 2.10 the restriction map on the  $n$ -indexed summands becomes the restriction map

$$\pi_i^{C_2}(|H(M_\bullet \otimes_{\mathbb{Z}/4} M_\bullet, w)|) \rightarrow \pi_i(|H(M_\bullet \otimes_{\mathbb{Z}/4} M_\bullet, w)|)$$

which is equivalently the map

$$\pi_i(|H(F_{\mathbb{Z}/4}(M_\bullet))|) \rightarrow \pi_i(|H(G_{\mathbb{Z}/4}(M_\bullet))|)$$

induced by the inclusion of fixed points natural transformation  $F_R \Rightarrow G_R$ . But this is precisely the definition of the map on non-abelian derived functors induced by  $F_R \Rightarrow G_R$ .  $\square$

We can straightforwardly extend our computer program to compute this map on non-abelian derived functors. Moreover there is no problem identifying the source and target with the appropriate summands of the groups computed in Theorem 2.14: we computed  $\pi_i((\text{THR}(\mathbb{Z}/4)^{\phi \mathbb{Z}/2})^{C_2})$  using non-abelian derived functors in the first place, and  $\mathbb{Z}/2$  has no non-trivial automorphisms so for  $i \geq 2n$  we can immediately identify  $L_i^{(n)} G_{\mathbb{Z}/4}(\mathbb{Z}/2)$  with the appropriate summand of  $\pi_i(\text{THR}(\mathbb{Z}/4)^{\phi \mathbb{Z}/2})$  (namely the  $(n, m, k) = (n, n, i - 2n)$ -indexed summand under the indexing of Proposition 2.6).

It remains to compute the action of the map  $r$ , which proves to be more involved. Recall our description of  $r$  in Corollary 2.5. Let  $M_*$  be the standard free resolution of  $\mathbb{Z}/2[n]$  by a chain complex of free  $\mathbb{Z}/4$ -modules

$$\cdots \rightarrow \mathbb{Z}/4 \xrightarrow{\cdot 2} \mathbb{Z}/4 \xrightarrow{\cdot 2} \mathbb{Z}/4 \rightarrow 0 \rightarrow \cdots,$$

and let  $\sigma : \text{Ch}_{\geq 0}(\mathbb{Z}/4) \rightarrow \text{sMod}_{\mathbb{Z}/4}$  be the Dold-Kan equivalence functor recalled in Theorem 1.3, so we have  $\sigma M_\bullet$  a simplicial  $\mathbb{Z}/4$ -module with geometric realisation  $\Sigma^n H\mathbb{Z}/2$ . Rewriting the description of  $r$  in terms of this free simplicial resolution of  $M$ , the action of  $r$  on the  $n$ -indexed diagonal component of  $(\text{THR}(\mathbb{Z}/4)^{\phi \mathbb{Z}/2})^{C_2}$  is given by

$$\begin{aligned} \left| \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet) \right)^{C_2} \right| &\rightarrow \left| \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet) \right)^{\phi C_2} \right| \\ &\simeq |H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} H\sigma M_\bullet| \\ &\simeq H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} |H\sigma M_\bullet| \\ &\simeq H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} \Sigma^n H\mathbb{Z}/2 \\ &\simeq \bigoplus_{m \geq 0} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2 \\ &\rightarrow \text{THR}(\mathbb{Z}/4)^{\phi \mathbb{Z}/2}, \end{aligned} \tag{5}$$

where the first map is the geometric realisation of the canonical map from genuine to geometric fixed points, the second map comes from monoidality of geometric fixed points, and the remaining maps are all the evident canonical maps. The source is

the geometric realisation of a (non-discrete) simplicial spectrum, not just a simplicial abelian group as before, so we can no longer straightforwardly plug it into the computer; we will need to use some tricks.

We start by simplifying the map from genuine to geometric fixed points.

**Lemma 2.16.** *Write  $\sigma M_\bullet = \bigoplus_{B_\bullet} \mathbb{Z}/4$ . Then we have a commutative diagram of simplicial spectra*

$$\begin{array}{ccc} \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet) \right)^{C_2} & \longrightarrow & \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet) \right)^{\phi C_2} \\ \downarrow \cong & & \downarrow \cong \\ \bigoplus_{(B_\bullet \times B_\bullet)_{C_2}} H\mathbb{Z}/4 & \longrightarrow & \bigoplus_{B_\bullet} H\mathbb{Z}/4^{\phi C_2} \end{array}$$

where the bottom horizontal map is defined by the projection

$$\bigoplus_{(B_\bullet \times B_\bullet)_{C_2}} H\mathbb{Z}/4 \rightarrow \bigoplus_{B_\bullet^d} H\mathbb{Z}/4 \cong \bigoplus_{B_\bullet} H\mathbb{Z}/4$$

onto the diagonal components of  $(B_\bullet \times B_\bullet)_{C_2}$  (recall that we split the set  $(B_i \times B_i)_{C_2}$  into diagonal and off-diagonal parts as  $B_i^d \amalg B_i^o$ ) composed with the map

$$\bigoplus_{B_\bullet} H\mathbb{Z}/4 \rightarrow \bigoplus_{B_\bullet} H\mathbb{Z}/4^{\phi C_2}$$

induced component-wise by the canonical map  $H\mathbb{Z}/4 \cong H\mathbb{Z}/4^{C_2} \rightarrow H\mathbb{Z}/4^{\phi C_2}$ . The simplicial structure on the bottom right group is such that the isomorphism

$$\bigoplus_{B_\bullet} H\mathbb{Z}/4^{\phi C_2} \cong H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} \bigoplus_{B_\bullet} H\mathbb{Z}/4 \cong H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} H\sigma M_\bullet$$

is an isomorphism of simplicial spectra, where we consider  $H\mathbb{Z}/4^{\phi C_2}$  as an  $H\mathbb{Z}/4$ -module via the Frobenius module structure.

*Proof.* Recalling the proof of Lemma 2.10 we have an equivalence of simplicial  $C_2$ -spectra

$$\begin{aligned} \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet) \right)^{C_2} &\cong \left( \bigoplus_{B_\bullet^d} H\mathbb{Z}/4 \oplus \bigoplus_{B_\bullet^o} (C_2)_+ \otimes_{\mathbb{S}} H\mathbb{Z}/4 \right)^{C_2} \\ &\cong H \left( \bigoplus_{B_\bullet \times B_\bullet} \mathbb{Z}/4, w' \right)^{C_2} \\ &\cong HF_{\mathbb{Z}/4}(\sigma M_\bullet), \end{aligned}$$

where  $w'$  is the involution on  $\bigoplus_{B_\bullet \times B_\bullet} \mathbb{Z}/4$  corresponding to interchanging the copies of  $B_\bullet$ , and  $F_{\mathbb{Z}/4}(N) := (N \otimes_{\mathbb{Z}/4} N)^{C_2}$ . We have

$$F_{\mathbb{Z}/4}((\sigma M)_i) \cong \bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/4 \cong \bigoplus_{B_i^d} \mathbb{Z}/4 \oplus \bigoplus_{B_i^o} \mathbb{Z}/4,$$

and observe that  $\bigoplus_{B_i^d} \mathbb{Z}/4$  is a sub-simplicial abelian group. In fact if we let  $M_\bullet^0$  be the chain complex

$$\cdots \mathbb{Z}/4 \xrightarrow{\cdot 0} \mathbb{Z}/4 \xrightarrow{\cdot 0} \mathbb{Z}/4 \rightarrow 0$$

(with the lowest degree  $\mathbb{Z}/4$  in degree  $n$ ) then  $\bigoplus_{B_i^d} \mathbb{Z}/4 \cong \sigma M_\bullet^0$ . However the level-wise subgroups  $\bigoplus_{B_i^o} \mathbb{Z}/4$  do not define a sub-simplicial abelian group, since face maps applied to off-diagonal terms can hit diagonal terms. That is, our level-wise decomposition of  $\bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/4$  does not give a decomposition of simplicial abelian groups.

Similarly analysing the geometric fixed points, we get an equivalence of simplicial spectra

$$\begin{aligned} \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet) \right)^{\phi_{C_2}} &\cong \left( \bigoplus_{B_\bullet^d} H\mathbb{Z}/4 \oplus \bigoplus_{B_\bullet^o} \Sigma^\infty(C_2)_+ \otimes_{\mathbb{S}} H\mathbb{Z}/4 \right)^{\phi_{C_2}} \\ &\cong \bigoplus_{B_\bullet^d} H\mathbb{Z}/4^{\phi_{C_2}} \\ &\cong H\mathbb{Z}/4^{\phi_{C_2}} \otimes_{H\mathbb{Z}/4} H\sigma M_\bullet, \end{aligned} \tag{6}$$

where in the last line we take  $H\mathbb{Z}/4^{\phi_{C_2}}$  with the Frobenius  $H\mathbb{Z}/4$ -module structure. Note that this time all the off-diagonal components vanish. We should briefly explain how the Frobenius module structure arises here. The copies of  $H\mathbb{Z}/4^{\phi_{C_2}}$  arise from the isomorphism

$$H\mathbb{Z}/4^{\phi_{C_2}} \cong \left( H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/4) \right)^{\phi_{C_2}}.$$

The simplicial structure maps involve acting (with the usual  $H\mathbb{Z}/4$ -module structure) on the copy of  $H\mathbb{Z}/4$  contained in the norm on the right hand side. But under the isomorphism this is exactly the action in the Frobenius module structure on  $H\mathbb{Z}/4^{\phi_{C_2}}$ .

Finally observe that under the above equivalences of spectra, the map from genuine to geometric fixed points of  $H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\sigma M_\bullet)$  becomes the map

$$\bigoplus_{(B_\bullet \times B_\bullet)_{C_2}} H\mathbb{Z}/4 \rightarrow \bigoplus_{B_\bullet} H\mathbb{Z}/4^{\phi_{C_2}}$$

as described in the statement of the Lemma.  $\square$

Now we need to understand how the map  $r$  acts on homotopy classes as computed by our computer program. Recall that our program works by computing the homology groups of the Moore complex of the simplicial abelian group

$$F_{\mathbb{Z}/4}(\sigma M_{\bullet}) \cong \bigoplus_{(B_{\bullet} \times B_{\bullet})_{C_2}} \mathbb{Z}/4.$$

Fix some  $i \geq n$ , and let  $\partial_i$  denote the degree  $i$  differential of the Moore complex. A homology class  $[x] \in \ker(\partial_i)/\text{im}(\partial_{i+1}) \cong \pi_i \left( \bigoplus_{(B_{\bullet} \times B_{\bullet})_{C_2}} H\mathbb{Z}/4 \right)$  is represented by some  $x \in \ker(\partial_i) \leq \bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/4$ . We want to determine the image  $r_*([x])$ .

**Lemma 2.17.** (i) *If we take a purely off-diagonal element  $x \in \ker(\partial_i) \cap \bigoplus_{B_i^{\circ}} \mathbb{Z}/4$ , then  $r_*([x]) = 0$ .*

(ii) *For  $x \in \ker(\partial_i)$ ,  $r_*([2x]) = 0$ .*

*Proof.* Part (i) is clear from the description of the map in Lemma 2.16. To be fully rigorous, consider the simplicial map from the  $i$ -simplex  $\Delta^i \rightarrow \bigoplus_{(B_{\bullet} \times B_{\bullet})_{C_2}} H\mathbb{Z}/4$  representing  $[x]$ , and observe that the postcomposition with the projection onto the diagonal  $\bigoplus_{B_i^{\circ}} H\mathbb{Z}/4$  is the zero map.

For (ii), observe that  $\pi_i(\text{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  is entirely 2-torsion, so for any  $x \in \ker(\partial_i)$  we have  $r_*([2x]) = 2r_*([x]) = 0$ .  $\square$

*Remark 2.18.* So for a general  $x \in \ker(\partial_i)$ , in order to compute  $r_*([x])$  it suffices to analyse the residue class of  $x$  in

$$\ker(\partial_i)/(\text{im}(\partial_{i+1}) + \ker(\partial_i) \cap \bigoplus_{B_i^{\circ}} \mathbb{Z}/4 + 2\ker(\partial_i)).$$

Let  $e_{(u,v)} \in \bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/4$  denote the basis element corresponding to  $(u, v) \in (B_i \times B_i)_{C_2}$ . Recalling the Dold-Kan equivalence of Theorem 1.3, explicitly  $B_i$  is the set of surjective maps  $[i] \twoheadrightarrow [j]$  where  $n \leq j \leq i$ , so there is an element  $\text{id}_{[i]} \in B_i$  corresponding to the identity map.

**Lemma 2.19.** *We have an isomorphism*

$$\ker(\partial_i)/(\text{im}(\partial_{i+1}) + \ker(\partial_i) \cap \bigoplus_{B_i^{\circ}} \mathbb{Z}/4 + 2\ker(\partial_i)) \cong \mathbb{Z}/2,$$

where the left hand side is generated by the residue class of  $e_{(\text{id}_{[i]}, \text{id}_{[i]})}$ .

More precisely, let  $\theta : \bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/4 \rightarrow \mathbb{Z}/2$  be the map that extracts the coefficient of  $e_{(\text{id}_{[i]}, \text{id}_{[i]})}$  and reduces it mod 2. Then  $\theta|_{\ker(\partial_i)} : \ker(\partial_i) \rightarrow \mathbb{Z}/2$  exhibits the above quotient.

*Proof.* Note that  $e_{(\text{id}_{[i]}, \text{id}_{[i]})} \in \ker(\partial_i)$ , so the restriction of  $\theta$  to  $\ker(\partial_i)$  remains surjective. We need to show that

$$\text{im}(\partial_{i+1}) + \ker(\partial_i) \cap \bigoplus_{B_i^{\circ}} \mathbb{Z}/4 + 2\ker(\partial_i) = \ker(\theta|_{\ker(\partial_i)}).$$

It is straightforward to check that the left hand side is contained in the right hand side. It remains to show that every element of the right hand side is contained in the left hand side.

First consider  $i = 0$ . Then  $\partial_i$  is the zero-map and  $\ker(\partial_i) = \bigoplus_{(B_0 \times B_0)_{C_2}} \mathbb{Z}/4 \cong \mathbb{Z}/4$ . We see that  $\ker(\theta|_{\ker(\partial_i)}) = \ker(\theta) = 2\mathbb{Z}/4$  and  $2\ker(\partial_i) = 2\mathbb{Z}/4$  so the right hand side is contained in the left hand side as desired.

Otherwise we have  $i \geq 1$ . Let  $x \in \ker(\theta|_{\ker(\partial_i)})$  i.e.  $x$  is an element of  $\ker(\partial_i)$  with an even coefficient of  $e_{(\text{id}_{[i]}, \text{id}_{[i]})}$ . We want to find  $y \in \bigoplus_{(B_{i+1} \times B_{i+1})_{C_2}} \mathbb{Z}/4$  such that the diagonal part of  $\partial_{i+1}(y)$  agrees with that of  $x$ , then  $x - \partial_{i+1}(y) \in \ker(\partial_i) \cap \bigoplus_{B_i^o} \mathbb{Z}/4$  so  $x \in \text{im}(\partial_{i+1}) + \ker(\partial_i) \cap \bigoplus_{B_i^o} \mathbb{Z}/4$  and we're done.

Write  $x = x^d + x^o$  where  $x^d \in \bigoplus_{B_i^d} \mathbb{Z}/4$  is the diagonal part and  $x^o \in \bigoplus_{B_i^o} \mathbb{Z}/4$  is the off-diagonal part. Consider what happens if we reduce everything modulo 2, which we denote with an overline. We get an element

$$\bar{x} = \bar{x}^d + \bar{x}^o \in \bigoplus_{B_i^d} \mathbb{Z}/2 \oplus \bigoplus_{B_i^o} \mathbb{Z}/2 \cong \bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/2.$$

Note the coefficient of  $\bar{e}_{(\text{id}_{[i]}, \text{id}_{[i]})}$  in  $\bar{x}$  is zero. Explicitly considering the face maps in  $\bigoplus_{(B_\bullet \times B_\bullet)_{C_2}} \mathbb{Z}/4$  shows that the on-diagonal part of  $\partial_i(x^o)$  is a multiple of 2, and we know  $\partial_i(x^o) + \partial_i(x^d) = \partial_i(x) = 0$ , so  $\partial_i(x^d)$  must also be zero modulo 2. That is,  $\bar{x}^d$  is in the kernel of the Moore boundary map for the simplicial abelian group  $\bigoplus_{B_i^d} \mathbb{Z}/2 \cong \sigma M_\bullet^0 / 2\sigma M_\bullet^0$ . However the Dold-Kan isomorphism  $\sigma$  commutes with reducing mod 2, so this is the simplicial abelian group  $\sigma(M_\bullet^0 / 2M_\bullet^0)$ , where we observe  $M_\bullet^0 / 2M_\bullet^0$  is the chain complex with  $\mathbb{Z}/2$  in every degree  $\geq n$  and zero boundary maps.  $M_\bullet^0 / 2M_\bullet^0$  has homology  $\mathbb{Z}/2$  in every degree  $\geq n$ , so by Dold-Kan the  $i$ th homology group of the Moore complex of  $\bigoplus_{B_i^d} \mathbb{Z}/2$  is also  $\mathbb{Z}/2$ . Moreover the quotient defining this homology group

$$\ker\left(\partial_i : \bigoplus_{B_i^d} \mathbb{Z}/2 \rightarrow \bigoplus_{B_{i-1}^d} \mathbb{Z}/2\right) / \text{im}\left(\partial_{i+1} : \bigoplus_{B_{i+1}^d} \mathbb{Z}/2 \rightarrow \bigoplus_{B_i^d} \mathbb{Z}/2\right) \cong \mathbb{Z}/2$$

is explicitly given by extracting the coefficient of  $\bar{e}_{(\text{id}_{[i]}, \text{id}_{[i]})}$ . Hence  $\bar{x}^d$  represents zero in homology, so there exists  $\bar{y} \in \bigoplus_{B_{i+1}^d} \mathbb{Z}/2$  such that  $\partial_{i+1}(\bar{y}) = \bar{x}$ .

Pick  $y'$  a lift of  $\bar{y}$  to  $\bigoplus_{B_{i+1}^d} \mathbb{Z}/4$ . Including  $\bigoplus_{B_{i+1}^d} \mathbb{Z}/4$  in  $\bigoplus_{(B_{i+1} \times B_{i+1})_{C_2}} \mathbb{Z}/4$ , we can consider  $y'$  an element of  $\bigoplus_{(B_{i+1} \times B_{i+1})_{C_2}} \mathbb{Z}/4$ . Then  $x^d$  and  $\partial_{i+1}(y')$  agree modulo 2, that is,  $x^d - \partial_{i+1}(y') = 2z$  for some  $z \in \bigoplus_{B_i^d} \mathbb{Z}/4$ .

Consider some element of  $B_i$ . An element of  $B_i$  corresponds to a surjective map  $[i] \rightarrow [j]$  for some  $j$ ; if this map sends  $k \mapsto a_k$  then let us denote the corresponding element of  $B_i$  by the tuple of values  $a = (a_0, \dots, a_i = j)$ . Let  $a' = (a_0, a_0, a_1, a_2, \dots, a_i) \in B_{i+1}$  and  $a'' = (a_0, a_1, a_1, a_2, \dots, a_i) \in B_{i+1}$  (recall we are considering the case where  $i \geq 1$ ), and note that  $a' \neq a''$ . Then  $\partial_{i+1}(e_{(a', a'')}) = 2e_{(a, a)} + h$  where  $h \in \bigoplus_{B_i^o} \mathbb{Z}/4$  is purely off-diagonal. By adding up elements of this form according to the coefficients of  $z$ , we can find  $y'' \in \bigoplus_{B_{i+1}^o} \mathbb{Z}/4$  such that the on-diagonal part of  $\partial_{i+1}(y'')$  equals  $2z$ .

Let  $y = y' + y''$ . Then  $x^d - \partial_{i+1}(y)$  is purely off-diagonal, so  $x - \partial_{i+1}(y) \in \ker(\partial_i) \cap \bigoplus_{B_i^o} \mathbb{Z}/4$  as desired and the claim is proved.  $\square$

*Remark 2.20.* We have used our computer program to double check this lemma for all values of  $n$  and  $i$  relevant for our later calculations (i.e.  $0 \leq n \leq i \leq 6$ ). Indeed this was how we originally realised that the lemma might be true.

**Proposition 2.21.** *Recall that  $x$  represents a homology class  $[x] \in \ker(\partial_i)/\text{im}(\partial_{i+1})$ , which by Corollary 2.11 we are identifying with the degree  $i$  homotopy of the  $n$ -indexed diagonal component of  $(\text{THR}(\mathbb{Z}/2)^{\phi\mathbb{Z}/2})^{C_2}$ . If the coefficient of  $e_{(\text{id}_{[i]}, \text{id}_{[i]})}$  in  $x$  is even then  $r_*([x]) = 0$ . Otherwise  $r_*([x])$  considered as an element of*

$$\pi_i(\text{THR}(\mathbb{Z}/2)^{\phi\mathbb{Z}/2}) \cong \bigoplus_{\substack{n', m, k \geq 0 \\ n' + m + k = i}} \mathbb{Z}/2$$

is the generator of the copy of  $\mathbb{Z}/2$  indexed by  $(n, 0, i - n)$ , where the above decomposition of  $\pi_i(\text{THR}(\mathbb{Z}/2)^{\phi\mathbb{Z}/2})$  is that of Proposition 2.6 (except using  $n'$  as an index instead of  $n$ , since we are already using  $n$ ).

*Proof.* By Remark 2.18 and Lemma 2.19 we deduce that for  $x \in \ker(\partial_i)$ , if the coefficient of  $e_{(\text{id}_{[i]}, \text{id}_{[i]})}$  in  $x$  is odd then  $r_*([x]) = r_*([e_{(\text{id}_{[i]}, \text{id}_{[i]})}])$  and otherwise  $r_*([x]) = 0$ . Hence it only remains to compute  $r_*([e_{(\text{id}_{[i]}, \text{id}_{[i]})}])$ .

Observe that the inclusion  $\bigoplus_{B_i^d} \mathbb{Z}/4 \hookrightarrow \bigoplus_{(B_i \times B_i)_{C_2}} \mathbb{Z}/4$  takes the basis element  $e_{\text{id}_{[i]}} \in \bigoplus_{B_i} \mathbb{Z}/4 \cong \bigoplus_{B_i^d} \mathbb{Z}/4$  to  $e_{(\text{id}_{[i]}, \text{id}_{[i]})}$ . This basis element  $e_{\text{id}_{[i]}}$  is in the kernel of the Moore differential on  $\bigoplus_{B_i^d} \mathbb{Z}/4 \cong \sigma M_{\bullet}^0$ , and in fact represents the generator of  $\pi_i(|H\sigma M_{\bullet}^0|) \cong H_i(M_{\bullet}^0) \cong \mathbb{Z}/4$ . So  $r_*([e_{(\text{id}_{[i]}, \text{id}_{[i]})}])$  is the image of  $1 \in \mathbb{Z}/4 \cong \pi_i\left(\bigoplus_{B_i^d} H\mathbb{Z}/4\right)$  under the map on  $\pi_i$  induced by

$$\begin{aligned} \left| \bigoplus_{B_i^d} H\mathbb{Z}/4 \right| &\rightarrow \left| \bigoplus_{(B_{\bullet} \times B_{\bullet})_{C_2}} H\mathbb{Z}/4 \right| \rightarrow \left| \bigoplus_{B_{\bullet}} H\mathbb{Z}/4^{\phi C_2} \right| \\ &\xrightarrow{\cong} \left| H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} H\sigma M_{\bullet} \right| \hookrightarrow \text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/4}, \end{aligned} \tag{7}$$

where as discussed in Lemma 2.16 the composite of the first two maps is the geometric realisation of the simplicial map

$$\bigoplus_{B_i^d} H\mathbb{Z}/4 \cong \bigoplus_{B_{\bullet}} H\mathbb{Z}/4 \rightarrow \bigoplus_{B_{\bullet}} (H\mathbb{Z}/4)^{\phi C_2}$$

given component-wise by  $H\mathbb{Z}/4 \rightarrow H\mathbb{Z}/4^{\phi C_2}$ , and the final inclusion is as described in (5).

Under the decomposition of  $H\mathbb{Z}/4$ -modules  $H\mathbb{Z}/4^{\phi C_2} \simeq \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2$  we see

$$\begin{aligned} H\mathbb{Z}/4^{\phi C_2} \otimes_{H\mathbb{Z}/4} H\sigma M_{\bullet} &\simeq \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\sigma M_{\bullet} \\ &\cong \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\sigma M_{\bullet}^0 \end{aligned}$$

where we use the fact that  $\sigma M_\bullet$  and  $\sigma M_\bullet^0$  are the same modulo 2. Now evaluating the geometric realisations we have

$$\left| \bigoplus_{B_\bullet^d} H\mathbb{Z}/4 \right| \cong \left| H\sigma M_\bullet^0 \right| \cong \bigoplus_{k \geq 0} \Sigma^{n+k} H\mathbb{Z}/4$$

and

$$\begin{aligned} \left| \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\sigma M_\bullet^0 \right| &\cong \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} \left| H\sigma M_\bullet^0 \right| \\ &\cong \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} \bigoplus_{k \geq 0} \Sigma^{n+k} H\mathbb{Z}/4 \\ &\cong \bigoplus_{k \geq 0} \Sigma^{n+k} \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2. \end{aligned}$$

Under these isomorphisms, the map (7) becomes

$$\bigoplus_{k \geq 0} \Sigma^{n+k} H\mathbb{Z}/4 \rightarrow \bigoplus_{k \geq 0} \Sigma^{n+k} \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/2 \hookrightarrow \bigoplus_{n', m, k \geq 0} \Sigma^{n'+m+k} H\mathbb{Z}/2 \simeq \text{THR}(\mathbb{Z}/2)^{\phi\mathbb{Z}/2}$$

where the first map is induced by  $H\mathbb{Z}/4 \rightarrow H\mathbb{Z}/4^{\phi C_2} \cong \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/4$ , the inclusion includes the  $n' = n$  components, and the final equivalence is as described in Section 2.2. The map  $H\mathbb{Z}/4 \rightarrow \bigoplus_{m \geq 0} \Sigma^m H\mathbb{Z}/4$  induces a quotient  $\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2$  on  $\pi_0$ . Under  $\left| \bigoplus_{B_\bullet^d} H\mathbb{Z}/4 \right| \cong \bigoplus_{k \geq 0} \Sigma^{n+k} H\mathbb{Z}/4$  the homotopy group  $[e_{\text{id}_{[i]}}]$  becomes the generator of  $\pi_i(\Sigma^{n+k} H\mathbb{Z}/4)$  for  $k = i - n$ . So its image under (7) is the generator of  $\pi_i(\Sigma^{n'+m+k} H\mathbb{Z}/2)$  for  $n' = n$ ,  $m = 0$  and  $k = i - n$ .  $\square$

We are now ready to exploit the long exact sequence for the homotopy equaliser of  $r - f$ .

**Theorem 2.22.** *We have short exact sequences*

$$\begin{aligned} 0 \rightarrow \mathbb{Z}/2 \rightarrow \pi_{-1}(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow 0 \\ 0 \rightarrow \pi_0(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow \mathbb{Z}/4 \rightarrow 0 \\ 0 \rightarrow \mathbb{Z}/2 \rightarrow \pi_1(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow \mathbb{Z}/2 \rightarrow 0 \\ 0 \rightarrow \mathbb{Z}/2 \rightarrow \pi_2(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow (\mathbb{Z}/2)^2 \rightarrow 0 \\ 0 \rightarrow (\mathbb{Z}/2)^2 \rightarrow \pi_3(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow (\mathbb{Z}/2)^4 \rightarrow 0 \\ 0 \rightarrow (\mathbb{Z}/2)^4 \rightarrow \pi_4(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow (\mathbb{Z}/2)^6 \oplus \mathbb{Z}/4 \rightarrow 0 \\ 0 \rightarrow (\mathbb{Z}/2)^9 \rightarrow \pi_5(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow (\mathbb{Z}/2)^{10} \rightarrow 0. \end{aligned}$$

*Proof.* The long exact sequence of homotopy groups for the fibre sequence

$$\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2} \rightarrow (\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2} \xrightarrow{r-f} \text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$$

splits into short exact sequences

$$0 \rightarrow \text{coker}((r - f)_{i+1}) \rightarrow \pi_i(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \rightarrow \ker((r - f)_i) \rightarrow 0,$$

so we just need to compute the kernels and cokernels of the maps  $(r - f)_i$ .

Using Propositions 2.15 and 2.21 we extend our computer program to compute the action of the maps  $r$  and  $f$  on homotopy groups. See the code repository for the full output; the computed kernels and cokernels are reproduced in the following table:

$i$	$\pi_i((\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2})$	$\pi_i(\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$	$\ker((r - f)_i)$	$\text{coker}((r - f)_i)$
0	$\mathbb{Z}/4$	$\mathbb{Z}/2$	$\mathbb{Z}/4$	$\mathbb{Z}/2$
1	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^3$	$\mathbb{Z}/2$	0
2	$(\mathbb{Z}/2)^7$	$(\mathbb{Z}/2)^6$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$
3	$(\mathbb{Z}/2)^{13}$	$(\mathbb{Z}/2)^{10}$	$(\mathbb{Z}/2)^4$	$\mathbb{Z}/2$
4	$(\mathbb{Z}/2)^{15} \oplus (\mathbb{Z}/4)^3$	$(\mathbb{Z}/2)^{15}$	$(\mathbb{Z}/2)^6 \oplus \mathbb{Z}/4$	$(\mathbb{Z}/2)^2$
5	$(\mathbb{Z}/2)^{27}$	$(\mathbb{Z}/2)^{21}$	$(\mathbb{Z}/2)^{10}$	$(\mathbb{Z}/2)^4$
6	$(\mathbb{Z}/2)^{34}$	$(\mathbb{Z}/2)^{28}$	$(\mathbb{Z}/2)^{15}$	$(\mathbb{Z}/2)^9$ .

Let us describe the program output in detail in the case  $i = 1$ , since we will use this in the next section. In Theorem 2.14 we already computed  $\pi_1((\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}) \cong (\mathbb{Z}/2)^4$  and  $\pi_1(\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \cong (\mathbb{Z}/2)^3$ . Let us denote the basis elements of  $\pi_1((\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2})$  by  $\{x_1, x_2, x_3, x_4\}$ , where  $\{x_1, x_2\}$  generates the 0-indexed diagonal component isomorphic to  $L_1^{(0)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$ ,  $\{x_3\}$  generates the 1-indexed diagonal component  $L_1^{(1)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  and  $\{x_4\}$  generates the off-diagonal component. Let the basis elements of  $\pi_1(\text{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$  be denoted by  $\{y_1, y_2, y_3\}$ , which corresponds to  $(n, m, k) = (0, 0, 1), (0, 1, 0), (1, 0, 0)$  respectively in the indexing of Proposition 2.6. Then the program computes that  $f$  takes  $x_1 \mapsto 0, x_2 \mapsto y_1, x_3 \mapsto 0$  and  $x_4 \mapsto y_2 + y_3$ , and  $r$  takes  $x_1 \mapsto y_1, x_2 \mapsto 0, x_3 \mapsto y_3$  and  $x_4 \mapsto 0$ . From this we calculate that  $(r - f)_1$  has kernel  $\mathbb{Z}/2$  and trivial cokernel.  $\square$

*Remark 2.23.* Note that all non-trivial homotopy groups of  $\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  are in degrees  $\geq -1$ . Indeed for  $A$  a ring spectrum with involution such that  $A$  and  $A^{\phi\mathbb{Z}/2}$  are connective, then  $(\text{THR}(A)^{\phi\mathbb{Z}/2})^{C_2}$  and  $\text{THR}(A)^{\phi\mathbb{Z}/2}$  are also connective, so the long exact sequence shows that  $\pi_i(\text{TCR}(A)^{\phi\mathbb{Z}/2}) = 0$  for  $i < -1$ .

**Corollary 2.24.** *The groups  $\pi_i(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$  are 4-torsion for all  $i$ .*

*Proof.* Since  $\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  is a ring spectrum, all the  $\pi_i(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$  are modules over the ring  $\pi_0(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \cong \mathbb{Z}/4$  hence must be 4-torsion.  $\square$

## 2.4 Resolving the extension problem in degree 1

We would prefer to actually compute the homotopy groups, not just fit them into exact sequences. The above shows  $\pi_{-1}(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \cong \mathbb{Z}/2$  and  $\pi_0(\text{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \cong \mathbb{Z}/4$ . The following analysis lets us additionally compute  $\pi_1$ , and makes some progress in simplifying the problem for higher homotopy groups.

The key observation is that the map  $r - f$  almost splits as a direct sum. Recall how the maps  $f$  and  $r$  are defined in Corollary 2.5 in terms of the decompositions of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  and  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$  described in Corollary 2.3. The map  $f$  sends the diagonal summand indexed by  $n$  to the diagonal summand indexed by  $n$ , and sends the off-diagonal summand indexed by  $(n, m)$  into the off-diagonal summands indexed by  $(n, m)$  and  $(m, n)$ . The map  $r$  is zero on the off-diagonal summands, and sends the diagonal summand indexed by  $n$  into the summands indexed  $(n, m)$  for  $m \geq 0$ . So  $r$  preserves the  $n$  index, and so does  $f$  if we restrict to the diagonal summands. We record this observation in the following lemma.

Let

$$D_n := \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2}$$

denote the  $n$ -indexed diagonal summand of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$ , and let

$$D := \bigoplus_{n \geq 0} D_n.$$

**Lemma 2.25.** *The restriction  $(r - f)|_D : D \rightarrow \mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  is a direct sum of maps*

$$\gamma_n : D_n \rightarrow \bigoplus_{m \geq 0} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2,$$

where the right hand side is the sum of the  $(n, m)$ -indexed summands of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  for  $m \geq 0$ .

*Proof.* Immediate from Corollary 2.5. □

Homotopy fibres preserve direct sums, so we have

$$\mathrm{hofib}((r - f)|_D) \cong \bigoplus_{n \geq 0} \mathrm{hofib}(\gamma_n).$$

Our computer program computes the action of  $r - f$  on homotopy in terms of the decomposition of Corollary 2.3, so for each  $n$  we can easily restrict to obtain the action of the map  $\gamma_n$ , and from this calculate a long exact sequence of homotopy groups for  $\mathrm{hofib}(\gamma_n)$ .

Let us record the resulting short exact sequences involving  $\pi_1(\mathrm{hofib}(\gamma_n))$ .

**Lemma 2.26.** *We have*

$$\begin{aligned} 0 \rightarrow (\mathbb{Z}/2)^2 &\rightarrow \pi_1(\mathrm{hofib}(\gamma_0)) \rightarrow \mathbb{Z}/2 \rightarrow 0 \\ 0 \rightarrow \mathbb{Z}/2 &\rightarrow \pi_1(\mathrm{hofib}(\gamma_1)) \rightarrow 0 \end{aligned}$$

and  $\pi_1(\mathrm{hofib}(\gamma_n)) = 0$  for  $n > 1$ .

*Proof.* From the long exact sequence of homotopy groups we get short exact sequences

$$0 \rightarrow \mathrm{coker}(\pi_{i+1}(\gamma_n)) \rightarrow \pi_i(\mathrm{hofib}(\gamma_n)) \rightarrow \ker(\pi_i(\gamma_n)) \rightarrow 0.$$

Immediate from the output of the computer program, we can read off the relevant restrictions of  $r - f$  and compute kernels and cokernels. For example, recall our description of the action of  $f$  and  $r$  on  $\pi_1$  in the proof of Theorem 2.22. We have  $\pi_1(D_0) \cong L_1^{(0)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^2$  corresponding to the subgroup generated by  $\{x_1, x_2\}$  in our previous computation, and  $\pi_1$  of the sum of the  $(0, m)$ -indexed components of  $\text{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  is  $(\mathbb{Z}/2)^2$  corresponding to the subgroup generated by  $\{y_1, y_2\}$  previously. The map  $f$  takes  $x_1 \mapsto 0$  and  $x_2 \mapsto y_1$  respectively, and the map  $r$  takes  $x_1 \mapsto y_1$  and  $x_2 \mapsto 0$ . So  $\gamma_0$  is given by  $x_1 \mapsto y_1$  and  $x_2 \mapsto -y_1$ , and we conclude that  $\pi_1(\gamma_0)$  has kernel  $\mathbb{Z}/2$ .

The full set of kernels and cokernels needed is as follows:

$$\begin{aligned} \text{coker}(\pi_2(\gamma_0)) &\cong (\mathbb{Z}/2)^2 & \ker(\pi_1(\gamma_0)) &\cong \mathbb{Z}/2 \\ \text{coker}(\pi_2(\gamma_1)) &\cong \mathbb{Z}/2 & \ker(\pi_1(\gamma_1)) &\cong 0. \end{aligned}$$

For  $n > i$  the short exact sequences are evidently all zero.  $\square$

Now we need to relate the homotopy groups of the restricted homotopy fibre  $\text{hofib}((r - f)|_D)$  to those of the original homotopy fibre  $\text{hofib}(r - f) \simeq \text{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ . Ultimately this will allow us to compute  $\pi_1(\text{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ .

We first note a small homological algebra lemma that we will need in the subsequent proposition.

**Lemma 2.27.** *Suppose we have a map of short exact sequences in an abelian category*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & X & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & C & \longrightarrow & D & \longrightarrow & X & \longrightarrow & 0 \end{array}$$

where the rightmost vertical map is equality. Then the left hand square is a pullback. If the map  $A \rightarrow C$  is an epimorphism then the left hand square is also a pushout.

*Proof.* Note that a square with the top horizontal map a monomorphism and the right vertical map an epimorphism is a pushout iff it is a pullback. Now consider the diagram

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \\ \downarrow & & \downarrow \\ * & \longrightarrow & X. \end{array}$$

Since  $X$  is the cokernel of  $C \rightarrow D$ , the bottom square is a pushout. As  $C \rightarrow D$  is monic and  $D \rightarrow X$  is epic, the bottom square is also a pullback. Similarly as  $X$  is the cokernel of  $A \rightarrow B$  the outer square is a pushout and a pullback. So by the pasting law for pullbacks the upper square is a pullback.

If  $A \rightarrow C$  is epic then so is  $B \rightarrow D$  by the 4-lemma applied to the original map of short exact sequences, so we deduce the upper square is also a pushout.  $\square$

**Proposition 2.28.** *Let  $(r - f)_i$  denote the map induced by  $r - f$  on homotopy groups  $\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}) \rightarrow \pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})$ . Then we have a pushout and pullback square*

$$\begin{array}{ccc} \mathrm{coker}((r - f)_{i+1}|_{\pi_{i+1}(D)}) & \longrightarrow & \pi_i(\mathrm{hofib}((r - f)|_D)) \\ \downarrow & & \downarrow \\ \mathrm{coker}((r - f)_{i+1}) & \longrightarrow & \pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \end{array}$$

where the horizontal arrows come from the long exact sequences for the homotopy fibres of  $(r - f)|_D$  and  $r - f$ , and the vertical maps are induced by the inclusion of  $D$  into  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$ .

*Proof.* We have a map of fibre sequences:

$$\begin{array}{ccccc} \mathrm{hofib}((r - f)|_D) & \longrightarrow & D & \xrightarrow{(r-f)|_D} & \mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2} \\ \downarrow & & \downarrow & & \parallel \\ \mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2} & \longrightarrow & (\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2} & \xrightarrow{r-f} & \mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}, \end{array}$$

giving a corresponding map of long exact sequences in homotopy, and so maps of the short exact sequences involving the homotopy groups of the homotopy fibres. Note that since  $D$  is a summand of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$ ,  $\pi_i(D)$  is a summand of  $\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2})$ , and so the map induced by  $(r - f)|_D$  on degree  $i$  homotopy groups is  $(r - f)_i|_{\pi_i(D)}$ . Then we have a map of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{coker}((r - f)_{i+1}|_{\pi_{i+1}(D)}) & \longrightarrow & \pi_i(\mathrm{hofib}((r - f)|_D)) & \longrightarrow & \ker((r - f)_i|_{\pi_i(D)}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathrm{coker}((r - f)_{i+1}) & \longrightarrow & \pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) & \longrightarrow & \ker((r - f)_i) \longrightarrow 0 \end{array}$$

The map  $\mathrm{coker}((r - f)_{i+1}|_{\pi_{i+1}(D)}) \rightarrow \mathrm{coker}((r - f)_{i+1})$  is evidently surjective. We claim that the map  $\ker((r - f)_i|_{\pi_i(D)}) \rightarrow \ker((r - f)_i)$  is an isomorphism; that is, we claim that  $\ker((r - f)_i)$  is contained within  $\pi_i(D)$ . Let  $0 \leq n < m$ . If we post-compose  $(r - f)_i$  with the projection map onto the off-diagonal component of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}$  indexed by  $(n, m)$

$$\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2}) \twoheadrightarrow \pi_i(\Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2)$$

then we get the projection map onto the off-diagonal component of  $(\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}$  indexed by  $(n, m)$

$$\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi\mathbb{Z}/2})^{C_2}) \twoheadrightarrow \pi_i(\Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2)$$

(observe that this is what happens if we post-compose  $f_i$  with the projection, and on homotopy  $r_i$  only hits non-diagonal components indexed by  $(n', 0)$  for some  $n'$ ). So  $\ker((r - f)_i)$  is entirely contained within the kernel of every projection onto an off-diagonal component, i.e. is contained in the diagonal part of the decomposition. We conclude the right hand vertical map is an isomorphism.

Now we are done by Lemma 2.27.  $\square$

*Remark 2.29.* Our earlier computer calculations suffice to compute the left hand vertical map of the pushout square, so we've reduced the problem of computing  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  to that of computing  $\pi_i(\mathrm{hofib}((r-f)|_D)) \cong \bigoplus_{n=0}^i \pi_i(\mathrm{hofib}(\gamma_n))$  and the pushout. In fact as we will see in the following calculation, even if while computing  $\pi_i(\mathrm{hofib}(\gamma_n))$  we run into extension problems that we are unable to solve, sometimes we can still compute the pushout.

**Theorem 2.30.** *We have*

$$\pi_1(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \cong (\mathbb{Z}/2)^2.$$

*Proof.* We apply the above proposition to compute  $\pi_1$ . Taking the direct sum of the  $n = 0$  and  $n = 1$  short exact sequences from Lemma 2.26, we see that

$$\begin{aligned} \mathrm{coker}((r-f)_2|_{\pi_2(D)}) &\cong (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/2, \\ \pi_1(\mathrm{hofib}((r-f)|_D)) &\cong \pi_1(\mathrm{hofib}(\gamma_0)) \oplus \pi_1(\mathrm{hofib}(\gamma_1)) \cong \pi_1(\mathrm{hofib}(\gamma_0)) \oplus \mathbb{Z}/2, \\ \ker((r-f)_1|_{\pi_1(D)}) &\cong \mathbb{Z}/2 \oplus 0. \end{aligned}$$

We saw earlier that  $\mathrm{coker}((r-f)_2) \cong \mathbb{Z}/2$ , and explicitly examining the action of  $r-f$  on  $\pi_2$  shows that the quotient

$$\mathrm{coker}((r-f)_2|_{\pi_2(D)}) \twoheadrightarrow \mathrm{coker}((r-f)_2)$$

is the projection of  $(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/2$  onto the second summand. Now Proposition 2.28 gives us a pushout square

$$\begin{array}{ccc} (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/2 & \hookrightarrow & \pi_1(\mathrm{hofib}(\gamma_0)) \\ \downarrow & & \downarrow \\ \mathbb{Z}/2 & \hookrightarrow & \pi_1(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}). \end{array}$$

Considering this pushout we find that  $\pi_1(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  is the cokernel of the composition

$$(\mathbb{Z}/2)^2 \rightarrow \pi_1(\mathrm{hofib}(\gamma_0)) \hookrightarrow \pi_1(\mathrm{hofib}(\gamma_0)) \oplus \mathbb{Z}/2,$$

where the first map is the map  $(\mathbb{Z}/2)^2 \cong \mathrm{coker}(\pi_2(\gamma_0)) \rightarrow \pi_1(\mathrm{hofib}(\gamma_0))$  from the exact sequence in Lemma 2.26, and the second map is the inclusion of the first factor. The exact sequence from Lemma 2.26 tells us that  $\pi_1(\mathrm{hofib}(\gamma_0))/(\mathbb{Z}/2)^2 \cong \mathbb{Z}/2$ , so we find

$$\pi_1(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \cong \pi_1(\mathrm{hofib}(\gamma_0))/(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/2 \cong (\mathbb{Z}/2)^2. \quad \square$$

*Remark 2.31.* Unfortunately the same approach is not enough on its own to compute higher homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ . For example we run into an extension problem when computing  $\pi_2$  of the homotopy kernel of  $\gamma_1$ , and this time we would actually need to solve this extension problem in order to compute  $\pi_2(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ . Nevertheless these techniques can potentially constrain the set of possible homotopy groups. For instance applying this analysis to  $\pi_4(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  gives a second proof that this group is 4-torsion (note we already knew this since it is a  $\pi_0(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ -module).

### 3 Asymptotic growth estimates

Our results so far show that the first few homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$  do not obey a 4-periodic pattern. We would like to further prove that the homotopy groups grow arbitrarily large, and so there is no point beyond which they start repeating with any period. To show this we will make asymptotic estimates that show that the logarithm of the size of the group  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  grows quadratically in  $i$ . In order to do so we almost completely compute the groups  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  for all  $i$  and  $n$ ; this is a result that may be of independent interest.

Recall as in Theorem 2.22 the long exact sequence associated to the homotopy fibre of  $r-f$  and the resulting short exact sequences. If we can estimate the sizes of the homotopy groups involved in the long exact sequence, then this gives us constraints on the sizes of the kernels and cokernels appearing in the short exact sequences, and so gives bounds on the size of  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ . In particular note we get these estimates without needing any control over the maps in the long exact sequence. We describe this general technique in the following lemma.

**Lemma 3.1.** *Let*

$$A \xrightarrow{u} B \rightarrow C \rightarrow D \xrightarrow{v} E$$

*be an exact sequence of abelian groups. Then we have bounds*

$$\max\left(1, \frac{|B|}{|A|}\right) \cdot \max\left(1, \frac{|D|}{|E|}\right) \leq |C| \leq |B| \cdot |D|.$$

*Proof.* We have a short exact sequence

$$0 \rightarrow \mathrm{coker}(u) \rightarrow C \rightarrow \mathrm{ker}(v) \rightarrow 0$$

so

$$|C| = |\mathrm{coker}(u)| \cdot |\mathrm{ker}(v)|.$$

And we have bounds

$$\max\left(1, \frac{|B|}{|A|}\right) \leq |\mathrm{coker}(u)| \leq |B|$$

and

$$\max\left(1, \frac{|D|}{|E|}\right) \leq |\mathrm{ker}(v)| \leq |D|,$$

where note that if  $|B| < |A|$  we take the trivial lower bound  $|\mathrm{coker}(u)| \geq 1$ , and similarly if  $|D| < |E|$  we take the trivial lower bound  $|\mathrm{ker}(v)| \geq 1$ .  $\square$

Applying this lemma to the long exact sequence for  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ , appropriate bounds on the sizes of  $\pi_j((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})$  and  $\pi_j(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  will give us upper and lower bounds for the size of  $\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$ .

First recall that we entirely understand the non-equivariant homotopy groups of  $\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ .

**Proposition 3.2.** *The group  $\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$  has order  $2^{\frac{1}{2}(i+1)(i+2)}$ .*

*Proof.* Recall from Proposition 2.6 that

$$\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}) \cong \bigoplus_{\substack{n,m,k \geq 0 \\ n+m+k=i}} \mathbb{Z}/2.$$

The number of  $\mathbb{Z}/2$ -summands is the number of ways of writing  $i$  as the sum of three non-negative integers  $i = n + m + k$ . For each  $0 \leq n \leq i$ , there are  $i - n + 1$  possible choices of non-negative integers  $m, k$  such that  $m + k = i - n$ . So in total we have

$$\sum_{n=0}^i i - n + 1 = \frac{1}{2}(i+1)(i+2)$$

summands. □

As before, analysing  $\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})$  takes more work.

**Lemma 3.3.** *The group  $\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})$  has order*

$$2^{\lfloor (i+1)^2/4 \rfloor} \cdot \prod_{0 \leq n \leq i} |L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)|.$$

*Proof.* Recall from Corollary 2.3 that

$$\begin{aligned} (\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2} &\simeq \bigoplus_{n \geq 0} \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2} \\ &\oplus \bigoplus_{0 \leq n < m} \Sigma^{n+m} H\mathbb{Z}/2 \otimes_{H\mathbb{Z}/4} H\mathbb{Z}/2. \end{aligned}$$

By Corollary 2.11 we know that the  $i$ th homotopy group of the  $n$ -indexed diagonal summand is computed using non-abelian derived functors as  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$ , and by general properties of non-abelian derived functors these groups are zero for  $n > i$ . So the diagonal part has order

$$\prod_{0 \leq n \leq i} |L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)|.$$

As we observed in Proposition 2.7 the homotopy of the off-diagonal terms is given by

$$\pi_i \left( \bigoplus_{0 \leq n < m} \bigoplus_{k \geq 0} \Sigma^{n+m+k} H\mathbb{Z}/2 \right) \cong \bigoplus_{\substack{n,m,k \geq 0 \\ n < m \\ n+m+k=i}} \mathbb{Z}/2.$$

Let us count the number of  $\mathbb{Z}/2$  summands appearing here. For each  $0 \leq k \leq i$ , there are  $\lfloor (i-k-1)/2 \rfloor + 1$  ways to pick non-negative integers  $n, m$  with  $n < m$  and  $n + m = i - k$ , so the total number of summands is

$$\begin{aligned} \sum_{k=0}^i \lfloor (i-k-1)/2 \rfloor + 1 &= \begin{cases} i(i+2)/4 & \text{if } i \text{ is even} \\ (i+1)^2/4 & \text{if } i \text{ is odd.} \end{cases} \\ &= \left\lfloor \frac{(i+1)^2}{4} \right\rfloor \end{aligned}$$

Multiplying the orders of the diagonal and off-diagonal parts gives the result. □

### 3.1 Non-abelian derived functor computations

To make further progress we need to study  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2)$ . As a warm-up, let us compute  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/4)$ ; this will come in useful later.

**Lemma 3.4.** *We have*

$$L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/4) \cong \pi_i^{C_2}(\Sigma^{n\rho}H\mathbb{Z}/4) \cong \begin{cases} \mathbb{Z}/4 & \text{if } n \text{ is even and } i = 2n \\ \mathbb{Z}/2 & \text{if } n \leq i < 2n, \text{ or if } n \text{ is odd and } i = 2n \\ 0 & \text{otherwise} \end{cases}$$

*Proof.* By the same reasoning as Corollary 2.11, Lemma 2.10 gives

$$L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/4) \cong \pi_i^{C_2} \left( \Sigma^{n\rho}H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/4) \right).$$

But  $\Sigma^{n\rho}H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/4) \cong \Sigma^{n\rho}H\mathbb{Z}/4$  so

$$L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/4) \cong \pi_i^{C_2}(\Sigma^{n\rho}H\mathbb{Z}/4).$$

This is precisely the Bredon homology of  $S^{n\rho}$  with coefficients in the Mackey functor of the abelian group  $\mathbb{Z}/4$  with trivial involution. For  $n$  even this is the homology of the chain complex

$$\cdots 0 \rightarrow \mathbb{Z}/4 \xrightarrow{0} \mathbb{Z}/4 \xrightarrow{2} \cdots \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \xrightarrow{0} \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \rightarrow 0 \cdots$$

where the non-zero groups are in degrees  $n$  to  $2n$ . For  $n$  odd the chain complex looks like

$$\cdots 0 \rightarrow \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \xrightarrow{0} \cdots \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \xrightarrow{0} \mathbb{Z}/4 \xrightarrow{2} \mathbb{Z}/4 \rightarrow 0 \cdots$$

with non-zero groups in the same degrees. The homology of these complexes gives the claimed result.  $\square$

Now we recall some of the theory of non-abelian derived functors from [DP58] and [ST74].

**Definition 3.5** (Cross effect, originally introduced in [EL53]). Let  $T : \text{Mod}_R \rightarrow \text{Ab}$  be a functor with  $T(0) = 0$ . The second cross-effect is the functor  $T_2 : (\text{Mod}_R)^2 \rightarrow \text{Ab}$  defined on objects by

$$T_2(A_1, A_2) := \text{Im}(\text{id}_{T(A_1 \oplus A_2)} - T(i_{A_1}p_{A_1}) - T(i_{A_2}p_{A_2}))$$

where  $i_{A_i}, p_{A_i}$  are the canonical injections and projections of  $A_i$  into and out of  $A_1 \oplus A_2$ . The functor  $T_2$  is defined on morphisms in a natural way; see [ST74] Section 4 for full details.

We define higher cross effects  $T_k : (\text{Mod}_R)^k \rightarrow \text{Ab}$  inductively via

$$T_k(A_1, \dots, A_k) := (T_{k-1}(A_1, \dots, A_{k-2}, -))_2(A_{k-1}, A_k).$$

**Lemma 3.6.** *The functor  $F_R(M) := (M \otimes_R M)^{C_2}$  has second cross-effect*

$$A_1, A_2 \mapsto A_1 \otimes_R A_2,$$

*and third cross-effect zero – accordingly we say that  $F_R$  is quadratic.*

*Proof.* We compute

$$(F_R)_2(A_1, A_2) := \text{Im}(\text{id}_{F_R(A_1 \oplus A_2)} - F_R(i_{A_1} p_{A_1}) - F_R(i_{A_2} p_{A_2})) \cong A_1 \otimes_R A_2.$$

For the third cross-effect we have

$$(F_R)_3(A_1, A_2, A_3) := ((F_R)_2(A_1, -))_2(A_2, A_3).$$

But  $(F_R)_2(A_1, -) = A_1 \otimes_R (-)$  is additive, so has second cross-effect zero. Hence  $F_R$  has third cross-effect zero.  $\square$

**Lemma 3.7.** *Let  $R$  be a commutative ring and let  $M$  be an  $R$ -module. We have a long exact sequence*

$$\dots \rightarrow L_i^{(n)} G_R(M) \rightarrow L_i^{(n)} F_R(M) \rightarrow L_{i+1}^{(n+1)} F_R(M) \rightarrow L_{i-1}^{(n)} G_R(M) \rightarrow L_{i-1}^{(n)} F_R(M) \rightarrow \dots$$

*where  $G_R(M) = M \otimes_R M$  is the tensor square functor.*

*We have  $L_i^{(n)} G_R(M) = 0$  for  $i < 2n$ , hence  $L_i^{(n)} F_R(M) \cong L_{i+1}^{(n+1)} F_R(M)$  for  $i < 2n$ . Explicitly this isomorphism is given by the suspension map  $\sigma_i^{(n)}(M)$  of [DP58] Equation 2.4.*

*Proof.* This follows by Corollary 3.9 of [DP58] and surrounding remarks, noting by the previous lemma that  $(F_R)_2(A, A) = G_R(A)$  and  $F_R$  is quadratic.  $\square$

*Remark 3.8.* We will use this lemma in the case  $R = \mathbb{Z}/4$  and  $M = \mathbb{Z}/2$ , where recall from Lemma 2.13 that

$$L_i^{(n)} G_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong \begin{cases} 0 & \text{if } i < 2n \\ \mathbb{Z}/2 & \text{if } i \geq 2n. \end{cases}$$

For  $i < 2n$  we have isomorphisms

$$L_i^{(n)} F_R(M) \xrightarrow[\cong]{\sigma_i^{(n)}(M)} L_{i+1}^{(n+1)} F_R(M) \xrightarrow[\cong]{\sigma_{i+1}^{(n+1)}(M)} L_{i+2}^{(n+2)} F_R(M) \xrightarrow[\cong]{\sigma_{i+2}^{(n+2)}(M)} \dots$$

Accordingly we can study these stable values. In fact this stabilisation occurs for the non-abelian derived functors of any functor ([DP58] Equation 3.11), so we can make the following general definition.

**Definition 3.9.** Let  $X : \text{Mod}_R \rightarrow \text{Ab}$  a functor with  $X(0) = 0$ . The left stable derived functors of  $X$  are defined by

$$L_q^s X(M) := \text{colim} \left( L_q^{(0)} X(M) \xrightarrow{\sigma_q^{(0)}(M)} L_{q+1}^{(1)} X(M) \xrightarrow{\sigma_{q+1}^{(1)} X(M)} L_{q+2}^{(2)} X(M) \rightarrow \dots \right).$$

*Remark 3.10.* Since by Remark 2.9  $L_i^{(n)}X(M) = 0$  for  $i < n$ , we see that  $L_q^s X(M) = 0$  for  $q < 0$ .

*Remark 3.11.* By Lemma 3.4 we have  $L_{q+n}^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/4) \cong \mathbb{Z}/2$  for  $n > q \geq 0$ . So

$$L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/4) \cong \mathbb{Z}/2$$

for  $q \geq 0$ .

Simson and Tyc study stable derived functors in [ST74]. In particular they compute the left stable derived functor of J. H. C. Whitehead's functor  $\Lambda_R : \text{Mod}_R \rightarrow \text{Mod}_R$  for certain rings  $R$ . This is of interest to us since  $\Lambda_R$  and  $F_R$  are naturally isomorphic on free modules, so have the same non-abelian derived functors. While Simson and Tyc do not consider the case  $R = \mathbb{Z}/4$ , we can still make use of some of their techniques.

**Proposition 3.12.** *For  $q \geq 0$  we have*

$$L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{q+1}.$$

*Proof.* Applying [ST74] Corollary 5.8 to the short exact sequence

$$0 \rightarrow \mathbb{Z}/2 \xrightarrow{2} \mathbb{Z}/4 \rightarrow \mathbb{Z}/2 \rightarrow 0$$

gives a long exact sequence

$$\cdots \rightarrow L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_q^s(\mathbb{Z}/4) \rightarrow L_q^s(\mathbb{Z}/2) \rightarrow L_{q-1}^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow \cdots.$$

As noted in Remark 3.11,  $L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/4) \cong \mathbb{Z}/2$  for  $q \geq 0$ . We claim that the maps  $L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/4)$  in the sequence, which come from applying  $L_q^s F_{\mathbb{Z}/4}$  to  $\mathbb{Z}/2 \xrightarrow{2} \mathbb{Z}/4$ , are zero.

Let us work through how to compute this map from the definitions. The map  $\mathbb{Z}/2[n] \xrightarrow{2} \mathbb{Z}/4[n]$  corresponds to the map of free resolutions

$$\begin{array}{ccccccc} \cdots & \xrightarrow{2} & \mathbb{Z}/4 & \xrightarrow{2} & \mathbb{Z}/4 & \xrightarrow{2} & \mathbb{Z}/4 & \longrightarrow & 0 & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \downarrow^2 & & \downarrow & & \\ \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \mathbb{Z}/4 & \longrightarrow & 0 & \longrightarrow & \cdots \end{array}$$

where the non-zero vertical map is in degree  $n$ . To compute  $L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_i^{(n)}F_{\mathbb{Z}/4}(\mathbb{Z}/4)$  we apply Dold-Kan to give a map of simplicial  $\mathbb{Z}/4$ -modules, apply  $F_{\mathbb{Z}/4}$  level-wise giving a map of simplicial abelian groups, then take degree  $i$  homotopy. Observe that after applying Dold-Kan, the level-wise maps are still all multiples of 2. However applying  $F_{\mathbb{Z}/4}$  to a map that is a multiple of 2 gives the zero map (since  $F_{\mathbb{Z}/4}(2\alpha) = 4F_{\mathbb{Z}/4}(\alpha) = 0$ ). So the resulting map of simplicial abelian groups is zero on the nose, and hence induces the zero map on homotopy.

As a result, for  $q \geq 0$  our long exact sequence splits up into short exact sequences of the form

$$0 \rightarrow \mathbb{Z}/2 \rightarrow L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_{q-1}^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow 0.$$

By Remark 3.10,  $L_{-1}^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) = 0$ . And by [ST74] Corollary 8.10,  $L_q^s F_{\mathbb{Z}/4}(M) \cong L_q^s \Lambda_{\mathbb{Z}/4}(M)$  is always 2-torsion. So we see by induction on  $q$  that

$$L_q^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{q+1}. \quad \square$$

*Remark 3.13.* We can now compute the value of the stable derived functor  $L_q^s F_{\mathbb{Z}/4}$  applied to any  $\mathbb{Z}/4$ -module. Indeed stable derived functors are additive, and every  $\mathbb{Z}/4$ -module is a direct sum of copies of  $\mathbb{Z}/4$  and  $\mathbb{Z}/2$ . So it suffices to compute the values of the functor at  $\mathbb{Z}/4$  and  $\mathbb{Z}/2$ , which we did in Remark 3.11 and Proposition 3.12.

So we have computed  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  in the stable range  $i < 2n$ . It remains to see what happens when  $i \geq 2n$ . To do so we borrow some techniques from Section 5 of [DMP24], but adapted to our current context of analysing  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ , whereas they were analysing  $\mathrm{TCR}(\mathbb{Z})^{\phi_{\mathbb{Z}/2}}$ .

**Lemma 3.14.** *We have a fibre sequence*

$$\begin{array}{c} \left( \Sigma^{n\rho+\sigma} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2} \oplus \Sigma^{2n} \mathbb{Z}/2 \\ \downarrow \\ (\Sigma^{n\rho} \mathbb{Z}/4)^{C_2} \\ \downarrow \\ \left( \Sigma^{n\rho} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2) \right)^{C_2} \end{array}$$

where  $\sigma$  is the sign representation of  $C_2$  (so  $\rho = 1 + \sigma$ ).

*Proof.* First apply Lemma 5.1 of [DMP24] to the map  $\mathbb{Z}/2 \xrightarrow{2} \mathbb{Z}/4$ . This map has cofibre  $\mathbb{Z}/2$ , so we get a square of  $N_{\{e\}}^{C_2}(H\mathbb{Z}/4)$ -modules

$$\begin{array}{ccc} \mathrm{Ind}_{\{e\}}^{C_2}(H\mathbb{Z}/2 \otimes_{\mathbb{S}} H\mathbb{Z}/2) & \xrightarrow{\mathrm{Ind}((H\mathbb{Z}/2) \otimes 2)} & \mathrm{Ind}_{\{e\}}^{C_2}(H\mathbb{Z}/2 \otimes_{\mathbb{S}} H\mathbb{Z}/4) \\ \downarrow \mathrm{id} & & \downarrow 2 \otimes (\widetilde{H\mathbb{Z}/4}) \\ N_{\{e\}}^{C_2}(H\mathbb{Z}/2) & \xrightarrow{N(2)} & N_{\{e\}}^{C_2}(H\mathbb{Z}/4) \end{array}$$

with total cofibre  $N_{\{e\}}^{C_2}(H\mathbb{Z}/2)$ . Here  $\mathrm{Ind}_{\{e\}}^{C_2}$  is left adjoint to the forgetful functor from  $N_{\{e\}}^{C_2}(H\mathbb{Z}/4)$ -modules in  $C_2$ -spectra to  $H\mathbb{Z}/4 \otimes_{\mathbb{S}} H\mathbb{Z}/4$ -modules in spectra, and the vertical maps are adjoint to the identity of  $H\mathbb{Z}/4 \otimes_{\mathbb{S}} H\mathbb{Z}/4$  and the map  $2 \otimes (\widetilde{H\mathbb{Z}/4})$  respectively.

Following the approach of [DMP24] Lemma 5.2, we apply the functor

$$H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} (-),$$

and note that this preserves the total cofibre. Simplifying the result somewhat, we get the square

$$\begin{array}{ccc} \mathrm{Ind}_{\{e\}}^{C_2}(HZ/2 \otimes_{HZ/4} HZ/2) & \xrightarrow{\mathrm{Ind}((HZ/2) \otimes 2)} & \mathrm{Ind}_{\{e\}}^{C_2}(HZ/2 \otimes_{HZ/4} HZ/4) \\ \downarrow \tilde{\mathrm{id}} & & \downarrow 2 \otimes \widetilde{(HZ/4)} \\ HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/2) & \xrightarrow{HZ/4 \otimes N(2)} & HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/4) \end{array}$$

with total cofibre  $HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/2)$ .

The top horizontal map is zero, since  $(HZ/2) \otimes 2 : HZ/2 \otimes_{HZ/4} HZ/2 \rightarrow HZ/2 \otimes_{HZ/4} HZ/4$  is evidently the zero map. The top right spectrum simplifies to  $\mathrm{Ind}_{\{e\}}^{C_2}(HZ/2)$ , and the bottom right spectrum simplifies to  $HZ/4$ . The left vertical map is given by the fold map

$$HZ/2 \otimes_{HZ/4} HZ/2 \oplus HZ/2 \otimes_{HZ/4} HZ/2 \rightarrow HZ/2 \otimes_{HZ/4} HZ/2$$

on underlying spectra, and has cofibre

$$\Sigma^\sigma HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/2)$$

(in general the cofibre sequence  $(C_2)_+ \rightarrow \mathbb{S}^0 \rightarrow \mathbb{S}^\sigma$  gives us a sequence  $\mathrm{Ind}_{\{e\}}^{C_2} X \xrightarrow{\tilde{\mathrm{id}}} X \rightarrow \Sigma^\sigma X$ ). Hence we find the pushout of the top horizontal and left vertical maps is

$$\left( \Sigma^\sigma HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/2) \right) \oplus \mathrm{Ind}_{\{e\}}^{C_2}(HZ/2).$$

So the (co)fibre sequence defining the total cofibre is

$$\begin{array}{c} \Sigma^\sigma HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/2) \oplus \mathrm{Ind}_{\{e\}}^{C_2}(HZ/2) \\ \downarrow \\ HZ/4 \\ \downarrow \\ HZ/4 \otimes_{N_{\{e\}}^{C_2}(HZ/4)} N_{\{e\}}^{C_2}(HZ/2). \end{array}$$

Taking the  $\Sigma^{n\rho}$  suspension followed by genuine  $C_2$ -fixed points gives the desired fibre sequence.  $\square$

**Corollary 3.15.** *We have a long exact sequence*

$$\begin{aligned} \cdots \rightarrow L_{i+1}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \oplus \pi_i(\Sigma^{2n} HZ/2) &\rightarrow \pi_i^{C_2}(\Sigma^{n\rho} HZ/4) \rightarrow L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow \\ L_i^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \oplus \pi_{i-1}(\Sigma^{2n} HZ/2) &\rightarrow \pi_{i-1}^{C_2}(\Sigma^{n\rho} HZ/4) \rightarrow L_{i-1}^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow \cdots \end{aligned}$$

*Proof.* Consider the long exact sequence in homotopy for the fibre sequence of Lemma 3.14. It remains to observe that for a  $C_2$ -spectrum  $X$ ,

$$\pi_i^{C_2}(\Sigma^{n\rho+\sigma} X) \cong \pi_{i+1}^{C_2}(\Sigma \Sigma^{n\rho+\sigma} X) \cong \pi_{i+1}^{C_2}(\Sigma^{(n+1)\rho} X)$$

so

$$\pi_i^{C_2}(\Sigma^{n\rho+\sigma} H\mathbb{Z}/4 \otimes_{N_{\{e\}}^{C_2}(H\mathbb{Z}/4)} N_{\{e\}}^{C_2}(H\mathbb{Z}/2)) \cong L_{i+1}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2).$$

Hence identifying terms in the long exact sequence as appropriate non-abelian derived functors gives the result.  $\square$

We can now use this long exact sequence, together with our earlier results, to almost completely compute  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  (this appeared as Theorem D in the introduction).

**Theorem 3.16.** *For  $i \geq 0$  fixed,  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  takes the same value for all  $0 \leq n \leq \lfloor i/2 \rfloor$ . Moreover we determine this constant value, up to two possibilities in some cases:*

$$\begin{aligned} L_i^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) &\cong L_i^{(1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong \dots \cong L_i^{\lfloor i/2 \rfloor} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \\ &\cong \begin{cases} (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 1} \text{ or } (\mathbb{Z}/2)^{\lfloor i/2 \rfloor} \oplus \mathbb{Z}/4 & \text{if } i \equiv 0 \pmod{4}, \\ (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 1} \text{ or } (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 2} & \text{if } i \equiv 1 \pmod{4}, \\ (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 1} & \text{if } i \equiv 2 \pmod{4}, \\ (\mathbb{Z}/2)^{\lfloor i/2 \rfloor + 2} & \text{if } i \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

For each  $k$  we have  $\left| L_{4k}^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \right| = \left| L_{4k+1}^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \right|$ . That is, if we know the value for some  $i$  congruent to 0 mod 4 then this tells us the value for  $i+1$  and vice versa.

For  $\lfloor i/2 \rfloor < n \leq i$  we have

$$L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{i-n+1}.$$

And  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) = 0$  for  $n > i$  (recall from Remark 2.9 that this is a general fact about non-abelian derived functors).

*Proof.* The claim for  $\lfloor i/2 \rfloor < n \leq i$  follows from Proposition 3.12 (see also Lemma 3.7), since for  $n \leq i < 2n$  we have

$$L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong L_{i-n}^s F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{i-n+1}.$$

The claim that  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  takes a constant value for  $0 \leq n \leq \lfloor i/2 \rfloor$  follows from the long exact sequence of Corollary 3.15, since in Lemma 3.4 we showed  $\pi_i^{C_2}(\Sigma^{n\rho} H\mathbb{Z}/4) = 0$  for  $i > 2n$ , so for  $i > 2n+1$  we get exact sequences

$$0 \rightarrow L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_i^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow 0.$$

To try to evaluate this constant value, we now analyse what happens near the boundary case  $i = 2n$ .

Consider the exact sequence of Lemma 3.7 and Remark 3.8, noting that

$$L_{2n+1}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{n+1},$$

so we get an exact sequence

$$\mathbb{Z}/2 \rightarrow L_{2n}^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow (\mathbb{Z}/2)^{n+1} \rightarrow 0.$$

This shows that

$$L_{2n}^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \in \{(\mathbb{Z}/2)^{n+1}, (\mathbb{Z}/2)^{n+2}, (\mathbb{Z}/2)^n \oplus \mathbb{Z}/4\}. \quad (8)$$

Next consider the following part of the sequence from Corollary 3.15:

$$\begin{aligned} & L_{2n}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \oplus \pi_{2n-1}(\Sigma^{2n} H\mathbb{Z}/2) \rightarrow \pi_{2n-1}^{C_2}(\Sigma^{n\rho} H\mathbb{Z}/4) \rightarrow L_{2n-1}^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \\ & \rightarrow L_{2n-1}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \oplus \pi_{2n-2}(\Sigma^{2n} H\mathbb{Z}/2). \end{aligned} \quad (9)$$

By Lemma 3.4 we have  $\pi_i^{C_2}(\Sigma^{n\rho} H\mathbb{Z}/4) \cong \mathbb{Z}/2$  for  $n \leq i < 2n$ , so  $\pi_{2n-1}^{C_2}(\Sigma^{n\rho} H\mathbb{Z}/4) \cong \mathbb{Z}/2$ . Also substituting in  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{i-n+1}$  for  $n \leq i < 2n$  and  $\pi_i(\Sigma^{2n} H\mathbb{Z}/4) = 0$  for  $i \neq 2n$ , we get

$$L_{2n}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow \mathbb{Z}/2 \rightarrow (\mathbb{Z}/2)^n \rightarrow (\mathbb{Z}/2)^{n-1}.$$

But any map  $(\mathbb{Z}/2)^n \rightarrow (\mathbb{Z}/2)^{n-1}$  has kernel of order at least 2, so the map  $\mathbb{Z}/2 \rightarrow (\mathbb{Z}/2)^n$  must be an injection, and hence the map  $L_{2n}^{(n+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow \mathbb{Z}/2$  is zero.

Now we look at the part of the exact sequence of Corollary 3.15 leading up to the section in (9), and split the argument according to the parity of  $n$ . Let  $n = 2k + 1$  be odd, then we get

$$\begin{aligned} 0 \rightarrow L_{4k+3}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_{4k+3}^{(2k+2)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \oplus \mathbb{Z}/2 \rightarrow \mathbb{Z}/2 \rightarrow \\ L_{4k+2}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow L_{4k+2}^{(2k+2)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow 0 \end{aligned}$$

using that  $\pi_i^{C_2}(\Sigma^{(2k+1)\rho} H\mathbb{Z}/4)$  is zero for  $i > 4k+2$  and isomorphic to  $\mathbb{Z}/2$  at  $i = 4k+2$  (by Lemma 3.4), and as we just observed the map into that group for  $i = 4k+1 = 2n-1$  is zero. Filling in the values of  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  that we have already computed gives

$$0 \rightarrow L_{4k+3}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow (\mathbb{Z}/2)^{2k+3} \rightarrow \mathbb{Z}/2 \rightarrow L_{4k+2}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow (\mathbb{Z}/2)^{2k+1} \rightarrow 0.$$

This shows  $L_{4k+2}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  has order at most  $2^{2k+2}$ , so considering the list of possibilities in (8) we must have

$$L_{4k+2}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{2k+2}.$$

Then the rest of the exact sequence shows

$$L_{4k+3}^{(2k+1)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{2k+3}.$$

Finally consider the case where  $n = 2k$  is even. We obtain a very similar exact sequence with the difference that now  $\pi_{4k}^{C_2}(\Sigma^{2k\rho} H\mathbb{Z}/4) \cong \mathbb{Z}/4$ , so we get

$$0 \rightarrow L_{4k+1}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow (\mathbb{Z}/2)^{2k+2} \rightarrow \mathbb{Z}/4 \rightarrow L_{4k}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \rightarrow (\mathbb{Z}/2)^{2k} \rightarrow 0$$

There is no exact sequence  $\mathbb{Z}/4 \rightarrow (\mathbb{Z}/2)^{2k+2} \rightarrow (\mathbb{Z}/2)^{2k} \rightarrow 0$ , so considering our list of possibilities we must have

$$L_{4k}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \in \{(\mathbb{Z}/2)^{2k+1}, (\mathbb{Z}/2)^{2k} \oplus \mathbb{Z}/4\}.$$

If  $L_{4k}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{2k+1}$  then the map from  $\mathbb{Z}/4$  has kernel  $\mathbb{Z}/2$ , so we find

$$L_{4k+1}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{2k+1}.$$

Otherwise  $L_{4k}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{2k} \oplus \mathbb{Z}/4$ ; in this case the map from  $\mathbb{Z}/4$  is injective and we find

$$L_{4k+1}^{(2k)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \cong (\mathbb{Z}/2)^{2k+2}. \quad \square$$

*Remark 3.17.* Note this result agrees with the output of our computer program, which we have used to compute  $L_i^{(n)} F_{\mathbb{Z}/4}$  for  $i, n \leq 6$ . In the cases where the above theorem does not give a definite answer, the program gives us

$$\begin{aligned} L_0^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) &\cong \mathbb{Z}/4, & L_1^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) &\cong (\mathbb{Z}/2)^2, \\ L_4^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) &\cong (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/4, & L_5^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) &\cong (\mathbb{Z}/2)^4. \end{aligned}$$

### 3.2 Application to size estimates

Combining Theorem 3.16 with Lemma 3.3 gives us lower and upper bounds for  $|\pi_i((\text{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})|$ .

**Theorem 3.18.** *We have bounds*

$$2^{(i+1)(5i+7)/8} \leq |\pi_i((\text{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})| \leq 2^{(i+2)(5i+8)/8}.$$

*Proof.* We work with the base 2 logarithms of the orders of the groups to make the notation easier. Recall Lemma 3.3, which says that

$$\log_2 |\pi_i((\text{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})| = \lfloor (i+1)^2/4 \rfloor + \sum_{0 \leq n \leq i} \log_2 |L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)|.$$

We apply the results of Theorem 3.16 to analyse the sum  $\sum_{0 \leq n \leq i} \log_2 |L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)|$ . For the range  $0 \leq n \leq \lfloor i/2 \rfloor$  we know  $L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)$  takes a constant value; we have

$$\sum_{0 \leq n \leq \lfloor i/2 \rfloor} \log_2 |L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)| = (\lfloor i/2 \rfloor + 1) \log_2 |L_i^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2)|$$

where  $\log_2 \left| L_i^{(0)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \right| \in \{ \lfloor i/2 \rfloor + 1, \lfloor i/2 \rfloor + 2 \}$ , giving lower and upper bounds for this term. For the range  $\lfloor i/2 \rfloor < n \leq i$  we have

$$\begin{aligned} \sum_{\lfloor i/2 \rfloor < n \leq i} \log_2 \left| L_i^{(n)} F_{\mathbb{Z}/4}(\mathbb{Z}/2) \right| &= \sum_{\lfloor i/2 \rfloor < n \leq i} (i - n + 1) \\ &= (\lfloor i/2 \rfloor)(\lfloor i/2 \rfloor + 1)/2. \end{aligned}$$

Combining these calculations and for convenience eliminating the floors and ceilings (at the cost of very slightly loosening the bounds) gives the result.  $\square$

Finally we have enough ingredients to use Lemma 3.1 get asymptotic bounds on the order of the homotopy groups of  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ . The following appeared as Theorem E in the introduction.

**Theorem 3.19.** *The groups*

$$\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})$$

*grow rapidly: the logarithm of the order asymptotically grows quadratically in  $i$ .*

*More precisely, for  $i \geq 0$  we have upper and lower bounds*

$$2^{(i-1)(i+1)/8} \leq |\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})| \leq 2^{(i+2)(9i+20)/8}.$$

*Proof.* Apply Lemma 3.1 to the long exact sequence for  $\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}}$ , giving

$$\begin{aligned} \frac{|\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})|}{|\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})|} &\leq |\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})| \\ &\leq |\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})| \cdot |\pi_{i+1}(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})|. \end{aligned}$$

Now by Proposition 3.2 computing  $|\pi_i(\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})|$  and Theorem 3.18 proving bounds on  $|\pi_i((\mathrm{THR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})^{C_2})|$ , we find

$$\begin{aligned} \log_2 (|\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})|) &\geq (i+1)(5i+7)/8 - (i+1)(i+2)/2 \\ &= (i-1)(i+1)/8 \end{aligned}$$

and

$$\begin{aligned} \log_2 (|\pi_i(\mathrm{TCR}(\mathbb{Z}/4)^{\phi_{\mathbb{Z}/2}})|) &\leq (i+2)(5i+8)/8 + (i+2)(i+3)/2 \\ &= (i+2)(9i+20)/8 \end{aligned} \quad \square$$

## References

- [BDS22] Morten Brun, Bjørn Ian Dundas, and Martin Stolz. “Equivariant Structure on Smash Powers” (2022). arXiv: 1604.05939 [math].
- [Cal+21] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. “Hermitian K-theory for stable  $\infty$ -categories III: Grothendieck-Witt groups of rings” (2021). arXiv: 2009.07225 [math].
- [Cal+23a] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. “Hermitian K-theory for stable  $\infty$ -categories I: Foundations”. *Selecta Mathematica* 29.1 (2023), p. 10.
- [Cal+23b] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. “Hermitian K-theory for stable  $\infty$ -categories II: Cobordism categories and additivity” (2023). arXiv: 2009.07224 [math].
- [DMP24] Emanuele Dotto, Kristian Moi, and Irakli Patchkoria. “On the Geometric Fixed Points of Real Topological Cyclic Homology”. *Journal of the London Mathematical Society* 109.2 (2024), e12862.
- [DMPR21] Emanuele Dotto, Kristian Moi, Irakli Patchkoria, and Sune Precht Reeh. “Real Topological Hochschild Homology”. *Journal of the European Mathematical Society* 23.1 (2021), pp. 63–152.
- [Dol58] Albrecht Dold. “Homology of Symmetric Products and Other Functors of Complexes”. *Annals of Mathematics* 68.1 (1958), pp. 54–80.
- [Dot12] Emanuele Dotto. “Stable Real K-theory and Real Topological Hochschild Homology” (2012). arXiv: 1212.4310 [math].
- [Dot24] Emanuele Dotto. “An Analogue of the Milnor Conjecture for the de Rham-Witt Complex in Characteristic 2” (2024). arXiv: 2405.02054 [math].
- [DP58] Albrecht Dold and Dieter Puppe. “Non-Additive Functors, Their Derived Functors, and the Suspension Homomorphism”. *Proceedings of the National Academy of Sciences* 44.10 (1958), pp. 1065–1068.
- [DP61] Albrecht Dold and Dieter Puppe. “Homologie nicht-additiver Funktoren. Anwendungen”. *Annales de l’institut Fourier* 11 (1961), pp. 201–312.
- [EL53] Samuel Eilenberg and Saunders Mac Lane. “On the Groups  $H(\Pi, n)$ , I”. *The Annals of Mathematics* 58.1 (1953), p. 55.
- [Gap24] The Gap Group. *GAP – Groups, Algorithms, and Programming, Version 4.13.1*. <https://www.gap-system.org>. 2024.
- [GJ09] Paul G. Goerss and John F. Jardine. *Simplicial Homotopy Theory*. Birkhäuser, 2009.

- [HHR16] Michael Hill, Michael Hopkins, and Douglas Ravenel. “On the Nonexistence of Elements of Kervaire Invariant One”. *Annals of Mathematics* 184.1 (2016), pp. 1–262.
- [HM15] Lars Hesselholt and Ib Madsen. “Real algebraic K-theory”. Accessed 2024-10-30: <https://web.math.ku.dk/~larsh/papers/s05/book.pdf>. 2015.
- [HNS] Yonatan Harpaz, Thomas Nikolaus, and Jay Shah. “Real Topological Cyclic Homology and Normal L-theory”. To appear.
- [Høg16] Amalie Høgenhaven. “Real Topological Cyclic Homology of Spherical Group Rings” (2016). arXiv: 1611.01204 [math].
- [Hov14] Mark Hovey. “Smith Ideals of Structured Ring Spectra” (2014). arXiv: 1401.2850 [math].
- [MHB93] I. Madsen, W. C. Hsiang, and M. Bökstedt. “The cyclotomic trace and algebraic K-theory of spaces.” *Inventiones mathematicae* 111.3 (1993), pp. 465–540.
- [Ran92] A.A. Ranicki. *Algebraic L-theory and Topological Manifolds*. Cambridge Tracts in Mathematics 102. Cambridge University Press, 1992.
- [Sag24] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 10.4)*. <https://www.sagemath.org>. 2024.
- [Sch23] Stefan Schwede. “Lectures on Equivariant Stable Homotopy Theory”. Accessed 2024-10-30: <https://www.math.uni-bonn.de/~schwede/equivariant.pdf>. 2023.
- [Sim74] Daniel Simson. “Stable Derived Functors of the Second Symmetric Power Functor, Second Exterior Power Functor and Whitehead Gamma Functor”. *Colloquium Mathematicae* 32.1 (1974), pp. 49–55.
- [ST74] Daniel Simson and Andrzej Tyc. “Connected Sequences of Stable Derived Functors and Their Applications”. *Dissertationes Mathematicae* 111 (1974).
- [Sto11] Martin Stolz. “Equivariant Structure on Smash Powers of Commutative Ring Spectra” (2011). arXiv: 1604.05939 [math].
- [Whi50] J. H. C. Whitehead. “A Certain Exact Sequence”. *Annals of Mathematics* 52.1 (1950), pp. 51–110.