

Edge universality of β -additions through Dunkl operators

David Keating¹ and Jiaming Xu²

¹Department of Mathematics, University of Illinois Urbana-Champaign, Champaign, USA,
dkeating@illinois.edu

²Department of Mathematics, KTH Royal Institut of Technology, Stockholm, Sweden,
jxu0800@gmail.com

Abstract

It is well known that the edge limit of Gaussian/Laguerre Beta ensembles is given by $\text{Airy}(\beta)$ point process. We prove an universality result that this also holds for a general class of additions of Gaussian and Laguerre ensembles. In order to make sense of β -addition, we introduce type A Bessel function as the characteristic function of our matrix ensemble following the line of [GM], [BCG]. Then we extract its moment information through the action of Dunkl operators, a class of differential operators originated from special function theory. We do the action explicitly on the Bessel generating functions of our additions, and after the asymptotic analysis, we obtain certain limiting functional in terms of conditional Brownian bridges of the Laplace transform of $\text{Airy}(\beta)$, which is universal up to proper rescaling among all our additions.

Contents

1	Introduction	2
1.1	Overview	2
1.2	Matrix additions for general β	3
1.3	Main result	4
1.4	Proof ingredients	6
2	Bessel generating functions	7
2.1	Dunkl operators	7
2.2	Multivariate Bessel functions	8
2.3	Bessel generating functions	8
2.4	Connection to free probability	10
3	Moment calculations	12
3.1	The Dunkl actions	12
3.2	Further grouping	16
3.3	A Simplification	16
4	Mapping to random walks	17
4.1	Discrete excursions and free cumulants	17
4.2	Conditional random walk	18
4.3	The partition function	19
4.4	Steepest descent	20
5	First moment	23
5.1	Explicit Functional expression	24
5.2	Scaling	28
5.3	Functional CLT	28
5.4	Universality	30

6	Properties of random walk bridges	31
6.1	Conditional functional CLT's	31
6.2	Time homogeneity of down steps	31
6.3	Tail estimate of the walks	33
7	Higher moments	34
7.1	Some technical lemmas	36
7.2	Piecewise walks and scaling	37
7.3	Blowing up terms and cancellations	39
7.4	Convergence of the functional	43
7.5	Renormalization	49
7.6	Tail estimate of the power sums	50
8	Proof of main theorem	55
9	Further studies	58
10	Appendix A: Ballot problem and more steepest descent	59
11	Appendix B: Proof of Lemma 6.5	63

1 Introduction

1.1 Overview

The focus of this article is the intersection of three topics in random matrix theory: addition of random matrices, β -ensembles, and edge universality. The study of addition of independent large random matrices dates back to Voiculescu, in which the author considers

$$C(N) = A(N) + B(N),$$

where $A(N)$, $B(N)$ are two independent $N \times N$ real/complex self-adjoint matrices with deterministic eigenvalues $a_1(N) \geq \dots \geq a_N(N)$, $b_1(N) \geq \dots \geq b_N(N)$ and uniformly random eigenvectors. Voiculescu [Vo] proves that if the empirical measures of $A(N)$, $B(N)$ have

$$\frac{1}{N} \sum_{i=1}^N \delta_{a_i(N)} \rightarrow \mu_A, \quad \frac{1}{N} \sum_{i=1}^N \delta_{b_i(N)} \rightarrow \mu_B$$

for some deterministic measures μ_A , μ_B on \mathbb{R} , then

$$\frac{1}{N} \sum_{i=1}^N \delta_{c_i(N)} \rightarrow \mu_C,$$

where

$$\mu_C = \mu_A \boxplus \mu_B$$

is also a deterministic measure on \mathbb{R} , known as the *free convolution* of μ_A and μ_B . This classical result was an early breakthrough in the subject called *free probability*, that has various interplay with different types of random matrix additions, see e.g [B1], [B2], [MSS].

In random matrix literature, β -ensembles are random N -tuples of real numbers $x_1 \geq x_2 \geq \dots \geq x_N$ with the density function

$$\frac{1}{Z_{n,\beta}} \prod_{1 \leq i < j \leq N} (x_i - x_j)^\beta \prod_{i=1}^N \exp[-V(x_i)], \quad (1)$$

where $Z_{n,\beta}$ is a normalizing constant, and $V(x)$ is the potential function satisfying certain analytic conditions. When $\beta = 1, 2, 4$, such density corresponds to invariant matrix ensembles with real/complex/quaternion entries, see e.g [D] for more details. For general $\beta > 0$, the parameter is understood as the inverse temperature. Two of the most classical objects in this context are Gaussian β -ensembles ($G\beta E$), when $V(x) = \frac{\beta}{2}x^2$, and Laguerre

β -ensembles ($L\beta E$), when $V(x) = \left[\frac{\beta}{2}(a+1) - 1\right] \log(x) - \frac{\beta}{2}x$. For $G\beta E$ and $L\beta E$, Dumitriu and Edelman [DE] give a concrete tridiagonal matrix realization of the density function. This construction is later extended to Jacobi ensembles by Killip and Nenciu [KN], then to a larger class of potentials $V(x)$ by Krishnapur, Rider, and Virag [KRV]. The tridiagonal structure is a main tool in studying β -ensembles.

To study the addition of invariant random matrices for general $\beta > 0$, rather than using the density in Eqn. (1), an alternate algebraic approach was developed. This is defined in [GM], where the authors use the type A Bessel function, a multivariate special function with representation theoretic background, to play the role of characteristic function of the matrix ensembles. While never formally written, it is believed that for any fixed $\beta > 0$, the asymptotic empirical measure of self-adjoint additions is still characterized by the free convolution as for $\beta = 1, 2$, see e.g [BCG, Appendix]. In addition, [GM], [Me], [MeP], [BCG] study the asymptotic empirical measures of the additions as $N \rightarrow \infty$ in low and high temperature regime (i.e, when $\beta \rightarrow \infty$ and $\beta \rightarrow 0$, resp.), and obtain two other versions of convolutions that differ from the classical free convolution. Another similar story happens with the additions of $N \times M$ rectangular matrices with certain invariant properties: [B1], [B2] study the asymptotic empirical measures of singular values for fixed $\beta = 1, 2$ when $N, M \rightarrow \infty$, [Xu] generalize the additions to $\beta > 0$ by using the type BC Bessel function, and [GrM], [Gri], [Xu] obtained several limiting results about the rectangular empirical measures in low and high temperatures.

One common feature of the literature of matrix additions mentioned above is that they only consider the global asymptotic results, while in this text we study the *edge behavior* of the random spectrum. For a single matrix ensemble without any additions, when $\beta = 1, 2$, there is a huge amount of known results. For example, it is known that the largest eigenvalue of a large class of random matrix ensembles, including $G\beta E$ and $L\beta E$, converges weakly to the Tracy-Widom distribution (TW_β) after proper rescaling. Alternatively, one can consider the largest k eigenvalues, where k is any fixed positive integer, the same arguments show that the rescaled k -tuples converge to the largest k particles of *Airy- β process*. The universal edge limit of large random matrices was also extended to Wigner matrices by [Sos], and to generalized real/complex self-adjoint matrices in [PS], [TV], [EYY], [LY], [BEY] and other works. For $\beta \neq 1, 2, 4$, when there are no invariant matrices, [RRV], [GSh], [DG], [PaSh], [Sh], [BEY] and other works show the edge universality for a large class of β -ensembles defined by Eqn.(1). We refer the readers to [BEY, Section 1] for a more detailed survey.

One of the main contributions of this article is to extend the edge universality of self-adjoint square matrices to a class of additions of finitely many $G\beta E$ and $L\beta E$, for general $\beta > 0$. This problem was studied in [Ahn], [JP] using different approaches. In these works the authors add invariant matrices of more general form, however, the results and proofs only hold for $\beta = 1, 2$. In the current work on β -additions, we use the same ensembles as in [GM], [BCG], whose definition rely on multivariate special functions.

Our proof of main result relies on the *moment method*. Heuristically, when taking power sums

$$\sum_{i=1}^N \lambda_i(N)^M$$

of all eigenvalues, where $M \in \mathbb{Z}_{>0}$ is large enough compared to N , the statistic is dominated by the $\lambda_i(N)$'s with largest absolute value. This allows one to study the edge behavior. A similar approach was initiated in [Sos] (see also [SS1], [SS2]), and later applied in [Pa], [Sod], [GSh] to study the edge limit of various random matrix ensembles. One common feature of the above work is that their eigenvalue power sums are obtained by directly calculating powers of the corresponding Wigner matrices/tridiagonal realization of β -ensembles, and then taking the trace. Such approach does not work in our setting, since to our best knowledge, a concrete matrix structure no longer exists for β -additions. Instead, we extract the moment information of the additions from their characteristic function by using *type A Dunkl operators*, a class of differential operators established in special function literature, see Section 2 for details. The applications of Dunkl operators to random matrices was initiated by [BCG], using the same type of operators as in this text, and later [Xu] extended such connection to rectangular matrices using Dunkl operators of type BC. Both these two work study the expectation of $\sum_{i=1}^N \lambda_i(N)^k$ for fixed $k \in \mathbb{Z}_{\geq 1}$, while in [GXZ], which is concurrent to this text, type A Dunkl operators are used to give explicit expressions of the Laplace transform of Airy $_\beta$ line ensemble, an infinite collection of continuous random curves, whose one-time marginal gives Airy(β).

1.2 Matrix additions for general β

We start from the additions of complex self-adjoint matrices, which corresponds to $\beta = 2$. Fix $\delta_1, \delta_2 \in \mathbb{R}$, $\delta_2 \geq 0$, and a sequence $\{\alpha_i\}_{i \geq 1}$ with $\alpha_i \in \mathbb{R}$ and $\sum_{i=1}^{\infty} \alpha_i^2 < \infty$. We consider the addition of matrices which

takes the form

$$\delta_1 I + \sqrt{\delta_2} \frac{X + X^*}{2} + \sum_{i=1}^{\infty} \alpha_i \left(\frac{1}{2} V_i V_i^* - I \right). \quad (2)$$

where X is an $N \times N$ matrix with i.i.d $\mathcal{N}(0, N) + i\mathcal{N}(0, N)$ entries, and V_i are i.i.d columns of i.i.d $\mathcal{N}(0, 1) + i\mathcal{N}(0, 1)$ entries with size N . The matrix $\frac{X+X^*}{2}$ is the Gaussian Unitary ensemble, and when $\alpha_1 = \dots = \alpha_L = a$,

$$\sum_{i=1}^L \frac{\alpha}{2} V_i V_i^* = \frac{\alpha}{2} Y Y^*,$$

where Y is a $N \times L$ rectangular matrix with i.i.d complex Gaussian entries, is the $N \times L$ Laguerre Unitary ensemble (with weight α). We also note that Eqn. (2) is studied in asymptotic representation theory: it is the $N \times N$ corner of a random infinite complex Hermitian matrix, whose distribution is invariant under conjugation of $U(\infty)$ and ergodic (see [OV]).

In this text we ignore the addition of constant by taking $\delta_1 = \sum_i \alpha_i$, and study the β -version of above addition. Here $\beta > 0$ is a parameter that is commonly viewed as *inverse temperature*, and GUE, LUE are replaced by Gaussian Beta ensemble (G β E) and Laguerre Beta ensemble (L β E) respectively. Since for $\beta \neq 1, 2, 4$ there is no concrete invariant matrix of G β E and L β E, the β -addition is defined in terms of multiplication of the type-A Bessel generating functions

$$G(z_1, \dots, z_N; \beta/2)$$

of the summands, as in [GM], [BCG]. We discuss this in more detail in Section 2.

For a $N \times N$ G β E and finitely many $N \times L_i$ L β E's, it is shown in [AN] that their (real-valued) linear combinations gives a legitimate probability measure P_N of N -tuples

$$\lambda_1 \geq \dots \geq \lambda_N$$

in \mathbb{R}^N . The same paper explains that these measures appear in a classification problem of ergodic Gibbs measures, see also [BR]. By classical free probability theory, the empirical measure of λ_i 's converges to a deterministic measure μ on \mathbb{R} as $N \rightarrow \infty$. The main focus of this text is the study of the upper edge behavior near μ_+ , the right endpoint of $\text{supp}(\mu)$. This corresponds to the fluctuation of the largest eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$, for a finite fixed k .

1.3 Main result

As a random point process on the real line parameterized by $\beta > 0$, the Airy- β process (abbreviated Airy(β)) is known as the universal object characterizing the scaling limit of the largest eigenvalues of various random matrix ensemble. While for $\beta = 1, 2, 4$ there have been several characterizations of the limiting point configuration and the corresponding Tracy-Widom distributions in the past literature, for general β the explicit expressions for Airy(β) are relatively less understood. In [RRV], the authors make sense of the stochastic Airy operator

$$SAO_\beta = -\frac{d^2}{dx^2} + x + \frac{2}{\sqrt{\beta}} B'(x) \quad (3)$$

on $L^2(\mathbb{R}_{\geq 0})$ with Dirichlet boundary condition at 0, where $B(x)$ denotes the Brownian motion and the differential makes sense as Schwartz distribution. See [RRV, Section 2] for the precise analytic issues to define such operator. SAO_β is known to have a orthonormal basis of eigenfunctions with eigenvalues $-\eta_i$ ($i = 1, 2, \dots$). The points in the Airy(β) process are given by these eigenvalues.

On the other hand, [GSh] introduces integral operators $\mathcal{U}(T)$ ($T \geq 0$), called the *stochastic Airy semigroup*, such that by [GSh, Corollary 2.12],

$$\mathcal{U}(T) \stackrel{d}{=} \exp\left(-\frac{T}{2} SAO_\beta\right) \quad (4)$$

for each T . The right side above is the unique stochastic operator with the same eigenfunctions as SAO_β and eigenvalues $\exp\left(-\frac{T\eta_i}{2}\right)$'s. In addition, $\mathcal{U}(T)$ is almost surely trace class (see [GSh, Proposition 2.4]), and the same proposition also gives an alternate characterization of Airy(β) in terms of its Laplace transform. In particular, we have the following expression.

Proposition 1.1. [GSh, Proposition 2.14]

$$\begin{aligned} \mathcal{E}[\text{Trace}(\mathcal{U}(T))] &= \mathbb{E} \left[\sum_{i=1}^{\infty} e^{\frac{T\eta_i}{2}} \right] \\ &= \frac{2}{\pi} T^{-\frac{3}{2}} \mathbb{E} \left[\exp \left(-\frac{T^{\frac{3}{2}}}{2} \int_0^1 \mathfrak{E}(t) dt + \frac{T^{\frac{3}{2}}}{2\beta} \int_0^{\infty} (l_y)^2 dy \right) \right], \end{aligned} \quad (5)$$

where \mathfrak{E} is a standard Brownian excursion on $[0, 1]$, and each l_y is the total local time of \mathfrak{E} at level y .

Note that Proposition 1.1 gives an explicit formula for the first moment of

$$\sum_{i=1}^{\infty} e^{\frac{T\eta_i}{2}}, \quad (6)$$

the Laplace transform of $\text{Airy}(\beta)$. Moreover, [GSh] provides an explicit kernel for $\mathcal{U}(T)$, which is given in terms of Brownian bridges conditioned to stay nonnegative. It might also be possible to extract explicit expressions for higher moments of Eqn. (6) from the kernel although we do not pursue that here.

Our main result verifies that the upper edge limit of the following class of β -additions is given by $\text{Airy}(\beta)$. Now we introduce the following notions necessary for stating the theorem.

Fix $\beta > 0$. Let $X(N)$ be a $N \times N$ Gaussian β -ensemble,

$$V_{N, L_1}(N), \dots, V_{N, L_k}(N)$$

be independent $N \times L_i$ ($L_i \geq N$) Laguerre β -ensembles, and $\alpha_1, \dots, \alpha_k \in \mathbb{R}$ be the coefficients of $V_{N, L_1}(N), \dots, V_{N, L_k}(N)$, respectively. Assume that there exists $\gamma_i > 0$, as $N \rightarrow \infty$,

$$\frac{L_i}{N} \rightarrow \gamma_i \geq 1 \quad (7)$$

for each $i = 1, 2, \dots, k$.

Fix $\delta \geq 0$, consider the β -addition

$$\sqrt{\delta} X(N) \boxplus_N^{\beta} \alpha_1 V_{N, L_1}(N) \boxplus_N^{\beta} \dots \boxplus_N^{\beta} \alpha_k V_{N, L_k}(N). \quad (8)$$

For $\beta = 1, 2, 4$ this is the usual additions of independent Gaussian and Wishart matrices, and for general $\beta > 0$ we give a generalized definition in Section 2.

Without loss of generality set

$$|\alpha_1| > |\alpha_2| > \dots > |\alpha_k| > 0. \quad (9)$$

For $l \in \mathbb{Z}_{\geq 1}$, let

$$\kappa_2 = \delta + \alpha_1^2 \gamma_1 + \dots + \alpha_k^2 \gamma_k, \quad \kappa_2(N) = \delta + \alpha_1^2 \frac{L_1(N)}{N} + \dots + \alpha_k^2 \frac{L_k(N)}{N}, \quad (10)$$

and

$$\kappa_l = \alpha_1^l \gamma_1 + \dots + \alpha_k^l \gamma_k, \quad \kappa_l(N) = \alpha_1^l \frac{L_1(N)}{N} + \dots + \alpha_k^l \frac{L_k(N)}{N}, \quad (11)$$

for $l \neq 2$. For each $N = 1, 2, \dots$, assume that

$$\kappa_l(N) \geq 0 \quad (12)$$

for all l 's. Let

$$V(z) = \frac{1}{z} + \sum_{l=1}^{\infty} \kappa_l z^{l-1} \quad (13)$$

be a meromorphic function on \mathbb{C} locally near 0.

Theorem 1.2 (Edge universality of β -ensemble additions). *Let $\lambda_i(N)$ ($i = 1, 2, \dots, N$) denote the eigenvalues of our $N \times N$ addition in Eqn. (8). Assume that (7), (9) and (12) hold, then there exists a positive constant C_0 , a sequence of positive real numbers $\mu_+(N)$, such that*

$$\lim_{N \rightarrow \infty} \left\{ \lambda_i^l := \frac{\lambda_i(N) - \mu_+(N)N}{N^{\frac{1}{3}}} \right\}_{i=1}^N \stackrel{d}{=} C_0 \cdot \text{Airy}(\beta), \quad (14)$$

the Airy- β process.

The following statement gives explicit descriptions of the parameters in Theorem 1.2.

Proposition 1.3. *Let $z_c > 0$ be the unique root of $V'(z)$ on $(0, \frac{1}{a_1})$. Then $V(z_c) > 0$, and*

$$\begin{aligned}\mu_+(N) &\rightarrow \mu_+ = V(z_c) \text{ as } N \rightarrow \infty, \\ C_0 &= 2^{-\frac{1}{3}} V''(z_c)^{\frac{1}{3}}.\end{aligned}\tag{15}$$

Remark 1.4. *A key restriction of our ensembles is that $\kappa_l \geq 0$ for all l 's. This is guaranteed when we have only additions of Laguerre ensembles, i.e., all $\alpha_i > 0$. In general, by taking $a_1 > 0$, it's not hard to see that $\kappa_l > 0$ for all l large enough. However, what is not covered in our result is when the terms being subtracted contribute enough to make $\kappa_l < 0$ for some l . We conjecture our result to still be true in that case, see Section 9 for more details.*

Remark 1.5. *Note that in the addition (8) we allow the absence of the Gaussian ensemble in the sum. It is possible that if $\delta > 0$ is large enough, the result in Theorem 1.2 can be (partially) obtained by the approach in [AH], which relies on the analysis of Dyson Brownian motion. However, their approach doesn't cover the cases $\beta \in (0, 1)$, or $\delta = 0$.*

Example 1.6. *A special case of Theorem 1.2 gives the classical edge limit of a single Laguerre β -ensemble of variance 1. In this case we have*

$$V(z) = \frac{1}{z} + \frac{\gamma}{1-z}, \quad z_c = \frac{1}{\sqrt{\gamma} + 1}, \quad \mu_+ = V(z_c) = (\sqrt{\gamma} + 1)^2,\tag{16}$$

and

$$C_0 = \frac{(\sqrt{\gamma} + 1)^{\frac{4}{3}}}{\gamma^{\frac{1}{6}}}.\tag{17}$$

Plugging in these parameters gives exactly [RRV, Theorem 1.4].

1.4 Proof ingredients

Define $\lambda_i^l = N^{-\frac{1}{3}}(\lambda_i(N) - \mu_+(N)N)$ as in Theorem 1.2. We calculate the asymptotic distribution of λ_i^l 's in terms of the Laplace transform

$$\sum_{i=1}^N e^{\frac{T\lambda_i^l}{\mu_+}},$$

which is a collection of random variables parameterized by $T > 0$.

We compute the l^{th} moments of the Laplace transform for all $l \in \mathbb{Z}_{\geq 1}$ by taking asymptotic of

$$\mathbb{E} \left[\left(\sum_{i=1}^N \lambda_i(N)^M \right)^l \right],$$

where $M \approx TN^{\frac{2}{3}}$. More precisely, we show that for each element in the class of additions we consider, the limiting moment expressions are all the same, after rescaling λ_i^l by C_0 . This establishes the edge universality, and since we know a single Laguerre ensemble converges to $\text{Airy}(\beta)$, so do the other additions.

The way we make sense of β -additions for general $\beta > 0$ is based on the symmetric function

$$B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \beta/2),$$

which is known as *type-A multivariate Bessel function* and plays the role of characteristic function of our ensembles. Moreover, we take the actions of *type-A Dunkl operator*, a class of differential operators $\mathcal{D}_i (i = 1, 2, \dots, N)$ on $\mathbb{E}[B_{(\lambda_1, \dots, \lambda_N)}(z_1, \dots, z_N)]$. Since the power sums of \mathcal{D}_i 's have type-A Bessel functions as eigenfunctions, such actions extract the moment information of λ_i 's.

Such approach using Dunkl action in random matrices follows the same line as [BCG] and [Xu]. But compared to these work, which take the global limit of the empirical measures, we focus on edge behavior and therefore we take the large powers of $\lambda_i(N)$'s, whose power grows with order $O(N^{\frac{2}{3}})$. This makes the asymptotic analysis of the explicit actions much more involved. When $l = 1$, we observe the appearance of

certain nonnegative walk bridge that starts from 0 and ends at 0 with length M . Moreover, the increments of this walk bridge are characterized by the free cumulants arising from the β -free convolution. We view it as a random walk excursion, and show that when $N, M \rightarrow \infty$, it converges weakly to a Brownian excursion on $[0, 1]$. In Section 5, we show the explicit expressions under the Dunkl action then converges to certain functional of Brownian excursion in the $l = 1$ case, postponing some technical toolkits to Section 6. When $l > 1$, the proof follows a similar flavor, but there are extra subtleties arising that makes the argument much more technical. We resolve these separately in Section 7, and again give explicit expressions for high moments of the Laplace transforms where Brownian excursion is upgraded to certain conditional nonnegative Brownian bridges. Section 8 gives the final steps to pass from moment convergence of λ_i^l 's to weak convergence.

Besides proving the edge universality, the limiting expressions arising in our proof provide new formulas that characterize $\text{Airy}(\beta)$. In a concurrent paper [GXZ], the same action of Dunkl operators is applied on the Bessel generating function of β -Dyson Brownian motions and corners process of Gaussian β -ensemble, which gives a similar combinatorial structure in terms of conditional random walk bridges. Such expressions converge to the multi-time Laplace transform of the Airy_β line ensemble, which specializes to our limiting expression when taking one-time marginal (though with slightly different presentation).

A natural problem that we do not handle in this text is to identify our limiting first moment expression with Eqn. (5), given in [GSh], or even identify our higher moment expressions with some corresponding formulas of $\text{Airy}(\beta)$ given in the literature by separate combinatorial arguments.

Acknowledgements

We thank Vadim Gorin for lots of stimulating discussions, and Lingfu Zhang for his helpful comments during the preparation of the text. We are also grateful to Yun Li for pointing out useful references about Airy processes, to Julien Sohier and Francesco Caravenna for pointing out useful references about conditional functional CLT.

Notations. For any real numbers $a < b$, we denote $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{Z}$. We also use $A \sim B$ to express $c_1 \leq A/B \leq c_2$ for some constants $c_1 < c_2$, uniform in the variables of A, B according to the context. Similarly, $A \lesssim B$ expresses $A \leq c_3 B$ for some uniform constant $c_3 > 0$. For a real-valued function f whose domain is a subset of \mathbb{R} , we use $f(x-)/f(x+)$ to denote the left/right limit of f at $x \in \mathbb{R}$, when it exists. f is *Cádlág* if it is right continuous and has a left limit for each x in the interior of its domain.

2 Bessel generating functions

In this section we give a brief review of Dunkl operators of type A and Bessel generating functions. See [Ro], [A] and references therein for more details. In Section 2.3, we use these generating function to define the notion of ‘addition’ for general β . In what follows we fix the real parameter $\theta > 0$.

2.1 Dunkl operators

Consider functions of N variables, z_1, \dots, z_N . Define an operator acting on these function, known as the *Dunkl operator* (of type A), by

$$\mathcal{D}_i := \frac{\partial}{\partial z_i} + \theta \sum_{j:j \neq i} \frac{1}{z_i - z_j} \circ (1 - s_{i,j}), \quad i = 1, \dots, N \quad (18)$$

where $s_{i,j}$ permutes the variables x_i and x_j . It is well-known that the Dunkl operators commute, that is,

$$\mathcal{D}_i \mathcal{D}_j = \mathcal{D}_j \mathcal{D}_i, \quad i, j = 1, \dots, N.$$

We also define

$$\mathcal{P}_k := \mathcal{D}_1^k + \dots + \mathcal{D}_N^k, \quad k > 0,$$

known as the *symmetric Dunkl operators*.

2.2 Multivariate Bessel functions

Let (a_1, a_2, \dots, a_N) be an N -tuple of real numbers satisfying $a_1 \leq a_2 \leq \dots \leq a_N$. The *multivariate Bessel function* $B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta)$ has many definitions. For example, it can be seen as the expectation of a certain observable of the θ -corner process or as a limit of Jack symmetric polynomials, see e.g [OO], [BCG, Section 2.2].

Here we list several well-known properties of these functions:

- They are symmetric in z_1, \dots, z_N . This can be seen immediately from the limit of the Jack polynomials.
- They can be extended to an entire function on $2N$ variables $a_1, \dots, a_N, z_1, \dots, z_N$.
- They have the normalization $B_{(a_1, \dots, a_N)}(0, \dots, 0; \theta) = 1$.
- They are eigenfunctions of the \mathcal{P}_k operators. In particular,

$$\mathcal{P}_k B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta) = \left(\sum_{i=1}^N a_i^k \right) B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta).$$

2.3 Bessel generating functions

Let $W_N := \{(a_1, a_2, \dots, a_N) \in \mathbb{R}^N \mid a_1 \geq a_2 \geq \dots \geq a_N\}$. Given a probability measure μ on W_N , we define the *Bessel generating function* of μ by

$$G_N(z_1, \dots, z_N; \theta, \mu) = \int_{a_1 \leq \dots \leq a_N} B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta) d\mu(a_1, \dots, a_N). \quad (19)$$

It follows from the symmetry of the Bessel functions that $G_N(z_1, \dots, z_N; \theta)$ is also symmetric in the variables z_1, \dots, z_N , and from the normalization of the Bessel functions we see that $G_N(0, \dots, 0; \theta) = 1$.

We will be interested in measures μ that satisfy a certain decay condition.

Definition 2.1. [BCG, Definition 2.7] We say that a measure μ on W_n is exponentially decaying with exponent $R > 0$ if

$$\int_{a_1 \leq \dots \leq a_n} e^{NR \max_i |a_i|} d\mu(a_1, \dots, a_N) < \infty.$$

This guarantees that the integral (19) converges for all (z_1, \dots, z_N) in the domain

$$\Omega_R := \{(z_1, \dots, z_N) \in \mathbb{C}^N : |\Re z_i| < R, i = 1, \dots, N\},$$

see [BCG, Lemma 2.9]. For such measures, we have the following proposition.

When $\theta = 1/2, 1, 2$, the Bessel function can be seen as an example of a spherical transform (see [Hel]). For example, when $\theta = 1$, we have

$$B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta) = \int_U \exp \operatorname{Tr}(UAU^*Z) dU \quad (20)$$

where $A = \operatorname{diag}(a_1, \dots, a_N)$, $Z = \operatorname{diag}(z_1, \dots, z_N)$ and dU is the Haar measure on $N \times N$ Unitary matrices. One can see

$$M_A = UAU^*$$

as a random hermitian matrix with fixed eigenvalues given by $a_1 \leq \dots \leq a_N$.

Suppose we take the sum of two such random matrices, M_A and M_B , with fixed eigenvalues $a_1 \leq \dots \leq a_N$ and $b_1 \leq \dots \leq b_N$, respectively. Consider the eigenvalues c_1, \dots, c_N of the result, by Eqn. (20) and the linearity of the trace, we have

$$\mathbb{E}[B_{(c_1, \dots, c_N)}(z_1, \dots, z_N; \theta) | \vec{a}, \vec{b}] = B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta) B_{(b_1, \dots, b_N)}(z_1, \dots, z_N; \theta)$$

It follows that

$$G_N(z_1, \dots, z_N; \theta, \mu_C) = G_N(z_1, \dots, z_N; \theta, \mu_A) G_N(z_1, \dots, z_N; \theta, \mu_B).$$

When $\theta \neq 1/2, 1, 2$ and there is no underlying invariant matrices to sum, so we define addition in terms of the generating functions.

Definition 2.2. Let μ_A and μ_B be exponentially decaying measures on W_N . Define $\mu_C := \mu_A \boxplus_N^\beta \mu_B$ to be the measure whose Bessel generating function is given by

$$G_N(z_1, \dots, z_N; \theta, \mu_C) = G_N(z_1, \dots, z_N; \theta, \mu_A) G_N(z_1, \dots, z_N; \theta, \mu_B). \quad (21)$$

We note that it is a well-known open problem in the general case to construct a probability measure on N -tuples (c_1, \dots, c_N) for which Eqn. (21) holds. However, for the sums we will be considering this has been done in [AN]. We denote this measure as $P_{N,\beta}$.

We will focus on two particular matrix ensembles: the Gaussian β ensembles and the Laguerre β ensembles.

Definition 2.3. For $\beta > 0$, a $N \times N$ Gaussian β -ensemble is a random N -tuples $(\lambda_1 \geq \dots \geq \lambda_N) \in \mathbb{R}^N$ with density function

$$\frac{1}{Z_{N,\beta}} \prod_{1 \leq i < j \leq N} (\lambda_i - \lambda_j)^\beta \prod_{i=1}^N \exp\left(-\frac{\lambda_i^2}{2N}\right), \quad (22)$$

and a $N \times L$ Laguerre β -ensemble ($N \leq L$) is a random N -tuples $(\lambda_1 \geq \dots \geq \lambda_N) \in \mathbb{R}^N$ with density function

$$\frac{1}{Z_{N,L,\beta}} \prod_{i=1}^N \left[\lambda_i^{\frac{\beta}{2}(L-N+1)-1} \exp\left(-\frac{\beta \lambda_i}{2}\right) \right] \prod_{1 \leq i < j \leq N} (\lambda_i - \lambda_j)^\beta. \quad (23)$$

$Z_{N,\beta}$ and $Z_{N,L,\beta}$ are certain normalizing constants.

Proposition 2.4. Let $\theta = \beta/2$. For the $N \times N$ $G\beta E$ the Bessel generating function is given by

$$G_N(z_1, \dots, z_N; \theta) = \exp\left(\frac{N(z_1^2 + \dots + z_N^2)}{2\theta}\right).$$

For the $N \times L$ $L\beta E$ the Bessel generating function is given by

$$G_N(z_1, \dots, z_N; \theta) = \prod_{i=1}^N \frac{\exp(-Lz_i)}{\left(1 - \frac{z_i}{\theta}\right)^{L\theta}}.$$

A sketch of proof of these two facts are given in Lemma 4.5 of [BCG] using Jack polynomials and [Ne] respectively.

Fix $\beta > 0$, $\delta \geq 0$, $\alpha_1, \dots, \alpha_k \in \mathbb{R}$. Consider the β -addition

$$\sqrt{\delta} X(N) \boxplus_N^\beta \alpha_1 V_{N,L_1}(N) \boxplus_N^\beta \dots \boxplus_N^\beta \alpha_k V_{N,L_k}(N).$$

If $\lambda_1 < \dots < \lambda_N$ are the random eigenvalues of this sum, then by Definition 2.2 and Proposition 2.4, its Bessel generating function is given by

$$G_N(z_1, \dots, z_N; \theta) = \mathbb{E}[B_{(\lambda_1, \dots, \lambda_N)}(z_1, \dots, z_N; \theta)] = \prod_{k=1}^N \mathcal{F}(z_k) \quad (24)$$

where

$$\begin{aligned} \mathcal{F}(z) &= \exp\left(\frac{\delta N}{2\theta} z^2\right) \prod_{i=1}^k \frac{\exp(-\alpha_i L_i z)}{\left(1 - \frac{\alpha_i z}{\theta}\right)^{L_i \theta}} \\ &= \exp\left(\frac{\delta N}{2\theta} z^2 + \theta \sum_{i=1}^k \sum_{l=1}^{\infty} \frac{1}{l} \frac{\alpha_i^l L_i + \dots + \alpha_k^l L_k}{\theta^l} z^l\right), \end{aligned} \quad (25)$$

so

$$G_N(z_1, \dots, z_N; \theta) = \exp\left(\frac{\delta N}{2\theta} p_2(z) + \theta \sum_{l=1}^{\infty} \frac{1}{l} \frac{\alpha_1^l L_1 + \dots + \alpha_k^l L_k}{\theta^l} p_l(z)\right), \quad (26)$$

where $p_k(z_1, \dots, z_N)$ is the k -th power sum symmetric function.

Remark 2.5. *This class of matrices was chosen in large part because the multivariate Bessel generating function factors into the product of single variable functions $\mathcal{F}(z)$. This greatly simplifies the computation in what follows. However, the measure $P_{N,\beta}$ supporting this BGF is also of independent interest, known as (up to shift by a constant) the extremal consistent distributions on the infinite families of interlacing arrays under Dixon-Anderson conditional probability distribution, see [AN]. The result in that paper generalizes the classification of ergodic unitary invariant infinite complex random matrices in [OV] to all $\beta > 0$.*

One of the key features of multivariate Bessel functions is that they are the (unique up to constant) eigenfunctions of Dunkl operators, see Proposition 2.6 below. A probabilistic version of this result is given in Proposition 2.7. This will be the starting point of our moment calculations in the remaining sections.

Let $p_k(x_1, \dots, x_N)$ be the degree k symmetric power sum of x_1, \dots, x_N , where $x_i (i = 1, 2, \dots, N)$ are formal variables that can be operators \mathcal{D}_i .

Proposition 2.6. *[Op] Fix $\theta > 0$, $N \in \mathbb{Z}_{\geq 1}$. For any $k = 1, 2, \dots$,*

$$p_k(\mathcal{D}_1, \dots, \mathcal{D}_N) B_{(a_1, \dots, a_N; \theta)}(z_1, \dots, z_N) = p_k(a_1, \dots, a_N) B_{(a_1, \dots, a_N)}(z_1, \dots, z_N; \theta). \quad (27)$$

Proposition 2.7. *Fix $\theta > 0$, $N \in \mathbb{Z}_{\geq 1}$. Define $G_N(z_1, \dots, z_N; \theta)$ as in Eqn. (26), and let $\lambda_1 \geq \dots \geq \lambda_N$ be its random eigenvalues. Then for any $k_1, \dots, k_m \in \mathbb{Z}_{\geq 1}$, we have*

$$\mathbb{E} \left[\prod_{i=1}^m p_{k_i}(\lambda_1, \dots, \lambda_N) \right] = \left(\prod_{i=1}^m p_{k_i}(\mathcal{D}_1, \dots, \mathcal{D}_N) \right) G_N(z_1, \dots, z_N; \theta). \quad (28)$$

Proof. This follows immediately from Proposition 2.6 and the fact $B_{(a_1, \dots, a_N)}(0, \dots, 0; \theta) = 1$, (see [BCG, Proposition 2.10] for more details), once we check that $P_{N,\beta}$ is exponentially decaying.

We give a tail estimate of $P_{N,\beta}$, that will also be used in Section 8. Consider the addition

$$C = A \boxplus_N^\beta B,$$

where A, B have deterministic eigenvalues $\lambda_1(A) \geq \dots \geq \lambda_N(A)$, $\lambda_1(B) \geq \dots \geq \lambda_N(B)$ and C has random eigenvalues $\lambda_1(C) \geq \dots \geq \lambda_N(C)$. By [A, Lemma 3.23],

$$\lambda_1(C) \leq \lambda_1(A) + \lambda_1(B).$$

Another ingredient is that for $G\beta E$ and $L\beta E$, the tail probability of their largest eigenvalue λ_1 is decaying exponentially fast, when λ_1 is going beyond the support of the global empirical measure, see e.g [LR, Section 1] for the detailed statements. By a simple union bound, for N large enough, as long as the empirical measure of P_N converges weakly to some deterministic measure on \mathbb{R} (which is the case, see Section 2.4), we have for $x > 1$,

$$\mathbb{P}_{N,\beta}[\lambda_1 \geq (1+x)\mu_+(N)N] \leq \exp(-CN \cdot x^{\frac{3}{2}}), \quad (29)$$

for some numerical constant $C(\beta) > 0$. This in particular finishes the proof. \square

2.4 Connection to free probability

We briefly recall some basic properties of free convolution, that corresponds to the asymptotic behavior of self-adjoint matrix additions. Given a measure μ on \mathbb{R} , the free cumulants $\{\kappa_l(\mu)\}_{l=1}^\infty \in \mathbb{R}^\infty$ are a collection of quantities that linearizes the free convolution, i.e, for $\mu_C = \mu_A \boxplus \mu_B$

$$\kappa_l(\mu_C) = \kappa_l(\mu_A) + \kappa_l(\mu_B)$$

for each $l = 1, 2, \dots$. For more detailed exposition, see e.g [No], [NS].

Since in this text we study the additions of β -Hermitian matrices that generalizes [Vo], it's not surprising that free cumulants again come into play. More precisely, they appears as the partial derivative at zero of the logarithm of the Bessel generating function.

Recall that for each $N = 1, 2, \dots$, we consider the addition

$$\sqrt{\delta} X(N) \boxplus_N^\beta \alpha_1 V_{N,L_1}(N) \boxplus_N^\beta \dots \boxplus_N^\beta \alpha_k V_{N,L_k}(N), \quad (30)$$

where $X(N)$ is a $N \times N$ Gaussian ensemble and $V_{N,L_i}(N)$ is a $N \times L_i(N)$ Laguerre ensemble. For simplicity, we will always assume a_1, \dots, a_k are independent of N .

Define the finite- N version of the free cumulants as follows: let

$$\begin{aligned}\kappa_1(N) &= \frac{1}{N} \left(\alpha_1 L_1(N) + \dots + \alpha_k L_k(N) \right) \\ \kappa_2(N) &= \frac{1}{N} \left(\delta N + \alpha_1^2 L_1(N) + \dots + \alpha_k^2 L_k(N) \right) \\ \kappa_l(N) &= \frac{1}{N} \left(\alpha_1^l L_1(N) + \dots + \alpha_k^l L_k(N) \right) \quad l \geq 3.\end{aligned}\tag{31}$$

We assume that $\gamma_1(N) := L_1(N)/N \rightarrow \gamma_1 > 0, \dots, \gamma_k(N) = L_k(N)/N \rightarrow \gamma_k > 0$ as $N \rightarrow \infty$, and without loss of generality assume that $a_1 > 0$ (the case $a_1 < 0$ only differ by a factor -1). Then we have

$$\begin{aligned}\kappa_1(N) &\longrightarrow \kappa_1 = \alpha_1 \gamma_1 + \dots + \alpha_k \gamma_k \\ \kappa_2(N) &\longrightarrow \kappa_2 = \delta + \alpha_1^2 \gamma_1 + \dots + \alpha_k^2 \gamma_k \\ \kappa_l(N) &\longrightarrow \kappa_l = \alpha_1^l \gamma_1 + \dots + \alpha_k^l \gamma_k \quad l \geq 3\end{aligned}\tag{32}$$

as $N \rightarrow \infty$. Since $\alpha_1 > 0$, $\kappa_l(N) > 0$ for all l 's large enough.

Under the setting above, as $N \rightarrow \infty$, the empirical measure

$$\frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i/N}$$

of our addition converges weakly in probability to a deterministic probability measure μ on \mathbb{R} , and μ_+ in Theorem 1.2 is the right endpoint of $\text{supp}(\mu)$. In addition, μ is uniquely determined by its free cumulants $\{\kappa_l\}_{l=1}^\infty$.

For $\beta = 1, 2$, the result is a special case of the more general classical statement in [Vo] presented at the beginning of the text, while for all $\beta > 0$ similar statement is also a well-believed to be true, but maybe never formally written down. We present it in a way adapted to [BCG, Claim 9.1] to highlight the connection to Bessel generating function.

Recall for our addition of matrices we have

$$G_N(z_1, \dots, z_N; \theta) = G_N(z_1, \dots, z_N; \theta) = \exp \left(\frac{\delta N}{2\theta} p_2(z) + \theta \sum_{l=1}^{\infty} \frac{1}{l} \frac{\alpha_1^l L_1 + \dots + \alpha_k^l L_k}{\theta^l} p_l(z) \right).$$

By direct calculation we see that as $N \rightarrow \infty$,

$$\frac{1}{N} \partial_i \ln G(z_1, \dots, z_N) \Big|_{z_1=\dots=z_N=0} = \kappa_1(N) \rightarrow \kappa_1\tag{33}$$

$$\frac{1}{N} \partial_i^2 \ln G(z_1, \dots, z_N) \Big|_{z_1=\dots=z_N=0} = \frac{\kappa_2(N)}{\theta} \rightarrow \frac{\kappa_2}{\theta}\tag{34}$$

$$\frac{1}{N} \partial_i^l \ln G(z_1, \dots, z_N) \Big|_{z_1=\dots=z_N=0} = (l-1)! \frac{\kappa_l(N)}{\theta^{l-1}} \rightarrow (l-1)! \frac{\kappa_l}{\theta^{l-1}}, \quad l \geq 3\tag{35}$$

$$\frac{1}{N} \partial_i \partial_j \ln G(z_1, \dots, z_N) \Big|_{z_1=\dots=z_N=0} = 0, \quad i \neq j.\tag{36}$$

By [BCG, Claim 9.1] we have the convergence of empirical measures (in the sense of moments), and κ_l 's are the free cumulants of μ since it is shown to satisfy the combinatorial relation Eqn. (72). Since $\{\kappa_l\}_{l=1}^\infty$ are explicitly one-one corresponding to the sequence of moments $\{m_k\}_{k=1}^\infty$, it completely characterizes μ (when the measure decays fast enough, which is the case in this text).

Example 2.8 (Gaussian ensemble). *Letting $\alpha_i = 0$ for all i 's and $\delta = 1$ results in $G\beta E$. Note that all the free cumulants are zero, except for $\kappa_2 = 1$. This implies that the limiting spectral measure is given by Wigner's semi-circle law which has density*

$$d\mu(x) = \frac{1}{2\pi} \sqrt{4 - x^2}, \quad |x| \leq 2\tag{37}$$

and 0 otherwise. The non-zero moments are given by

$$m_{2n} = \int_{-2}^2 x^{2n} d\mu(x) = C_n \quad (38)$$

where $C_n = \frac{1}{n+1} \binom{2n}{n}$ is the n -th Catalan number.

Example 2.9 (Laguerre ensemble). Suppose we take $\alpha_1 = 1$ and $\delta, \alpha_2, \alpha_3, \dots$ to be 0, then we get a $N \times L$ $L\beta E$. It is known that if $N, L \rightarrow \infty$ such that $L/N \rightarrow \gamma > 1$, then the limiting spectral measure is given by the Marchenko-Pastur distribution

$$d\mu(x) = \frac{1}{2\pi} \frac{\sqrt{(x - \mu_-)(\mu_+ - x)}}{x}, \quad \mu_- \leq x \leq \mu_+ \quad (39)$$

where $\mu_{\pm} = (\sqrt{\gamma} \pm 1)^2$. It is known that the Voiculescu R -transform characterizes the random variable, and for μ it is

$$R(z) = \frac{\gamma}{1-z} = \sum_{n=0}^{\infty} \gamma z^n. \quad (40)$$

Recall that $R(z)$ is the generating function of the free cumulants. We see that $\kappa_l = \gamma$ for each l .

3 Moment calculations

In this section we compute the Dunkl action on our Bessel generating function, defined in the previous section. When applying the Dunkl operator a large number of times to our Bessel generating function, the resulting expression can be written the original BGF times a polynomial in the variables z_1, \dots, z_N with a large number of terms. The main goal of this section is to organize and group these terms in a way that will allow us to give them a probabilistic interpretation that is amenable to further analysis.

3.1 The Dunkl actions

Let $\lambda_1 \geq \dots \geq \lambda_N$ be the random eigenvalues. We define the rescaled eigenvalue fluctuations as

$$\lambda'_i := N^{-1/3}(\lambda_i - \mu_+(N)N), \quad (41)$$

where we use the same scaling as the edge limit of a single Gaussian or Laguerre ensemble, see e.g [RRV].

Our goal is to characterize the distribution of λ'_i 's by the (random) Laplace transform on empirical measure

$$\sum_{i=1}^N e^{\frac{T\lambda'_i}{\mu_+}} \quad (42)$$

for $T > 0$. Note that by Eqn. (29), Eqn. (42) has finite moments when T is small enough.

We approximate the Laplace transform by rescaled power sums of the λ_i 's, more precisely,

$$e^{T\lambda'_i/\mu_+} \approx \left(1 + \frac{\lambda'_i}{\mu_+(N)N^{2/3}} \right)^{TN^{2/3}} \quad (43)$$

$$= \left(1 + \frac{N^{-1/3}(\lambda_i - \mu_+N)}{\mu_+(N)N^{2/3}} \right)^{TN^{2/3}} \quad (44)$$

$$= \left(\frac{\lambda_i}{\mu_+(N)N} \right)^{TN^{2/3}}. \quad (45)$$

Remark 3.1. We present the details showing how the above approximation leads to the weak convergence of extreme eigenvalues in Section 8. Heuristically, the estimation works well for the largest λ_i 's which are close to μ_+N , the upper edge of the limiting empirical measure. For λ_i 's that are smaller, $e^{T\lambda'_i/\mu_+}$ is always close to

0, while $|\frac{\lambda_i}{\mu_+ N}|$ is close to 0 only if $\lambda_i \gg -\mu_+ N$. This holds for almost all λ_i 's if and only if $|\mu_-| < |\mu_+| = \mu_+$, which is the case as long as there exists at least one Laguerre component in the sum.

For a single Gaussian ensemble, note that each λ_i is equal in distribution to $-\lambda_i$ and therefore

$$\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+ N} \right)^M = 0$$

when M is odd, and is approximately $2 \sum_{i=1}^N e^{T\lambda_i/\mu_+}$ when M is even. So the right moment expression for Gaussian β -ensemble is

$$\sum_{i=1}^N e^{T\lambda_i/\mu_+} \approx \frac{1}{2} \cdot \left[\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+ N} \right)^{\lfloor TN^{\frac{2}{3}} \rfloor} + \sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+ N} \right)^{\lfloor TN^{\frac{2}{3}} \rfloor + 1} \right], \quad (46)$$

see e.g [GSh]. As we only consider the case when there is at least one Laguerre component, we avoid this subtlety.

Our goal reduces to characterizing

$$\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+(N)N} \right)^{\lfloor TN^{2/3} \rfloor}, \quad (47)$$

which we do by calculating its l^{th} moments, for all $l \in \mathbb{Z}_{\geq 1}$. Assume that $\mu_+ > |\mu_-| \geq 0$, which will always be the case under our assumption in Theorem 1.2, and consider an operator of the form

$$\frac{1}{(\mu_+(N)N)^M} \left(\sum_{i=1}^N \mathcal{D}_i^M \right) \Big|_{z_1=\dots=z_N=0} \quad (48)$$

acting on $G(z_1, \dots, z_n; \theta)$, where

$$M = \lfloor TN^{2/3} \rfloor, \quad (49)$$

and \mathcal{D}_i is the Dunkl operator. By Proposition 2.7, we have

$$\frac{1}{(\mu_+(N)N)^M} \sum_{i=1}^N \mathbb{D}_i^M G(z_1, \dots, z_N; \theta) \Big|_{z_1=\dots=z_N=0} = \mathbb{E} \left[\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+(N)N} \right)^M \right]. \quad (50)$$

This gives the first moment of Eqn. (47). We will discuss this in detail in Section 5. The higher moments can be obtained by similar actions, which we will discuss in detail in Section 7.

Remark 3.2. For fixed $T > 0$, $M = \lfloor TN^{\frac{2}{3}} \rfloor$ is growing with N . Throughout the remainder of the text we will switch between using the index N and M depending on which is more relevant in context.

For now we focus on the Dunkl action giving the first moment. Recall that $G_N(\cdot)$ is given by

$$G_N(z_1, \dots, z_N; \theta) = \exp \left(\frac{\delta N}{2\theta} p_2(z) + \theta \sum_{l=1}^{\infty} \frac{1}{l} \frac{\alpha_1^l L_1(N) + \dots + \alpha_k^l L_k(N)}{\theta^l} p_l(z) \right).$$

for the ensembles we are interested in. Since G_N is symmetric in the z 's, and we are acting on it by a symmetric power sum of \mathcal{D}_i 's, it suffices to consider only the action of \mathcal{D}_1^M on G_N , since

$$\frac{1}{(\mu_+(N)N)^M} \left(\sum_{i=1}^N \mathcal{D}_i^M \right) G_N(z_1, \dots, z_N; \theta) \Big|_{z_1=\dots=z_N=0} = \frac{N}{(\mu_+(N)N)^M} \cdot \mathcal{D}_1^M G_N(z_1, \dots, z_N; \theta) \Big|_{z_1=\dots=z_N=0} \quad (51)$$

To simplify the expression, we pair each \mathcal{D}_i with $1/N$ and consider the action of $N^{-1}\mathcal{D}_1$. Recall from Eqn. (18) that

$$\frac{1}{N} \mathcal{D}_1 = \frac{1}{N} \partial_1 + \frac{\theta}{N} \sum_{j=2}^N \frac{1 - s_{1j}}{z_1 - z_j}. \quad (52)$$

Note that the operator consists of two parts: a partial derivative term ∂_1 and the swapping term $\frac{1-s_{1j}}{z_1-z_j}$.

From now on we will treat the parameter θ as constant and omit it from notation. Let us compute the action of $N^{-1}\mathcal{D}_1$ on functions of the form

$$h(z) \cdot G_N(z),$$

where $G_N(z)$ is the Bessel generating function and $h(z)$ is a polynomial of z_1, \dots, z_N .

By straightforward calculation, we have that the action of the partial derivative can be written

$$\begin{aligned} (N^{-1}\partial_1)[h(z)G_N(z)] &= \left[\left(\frac{\alpha_1 L_1(N) + \dots + \alpha_k L_k(N)}{N} + \frac{\delta + \alpha_1^2 L_1(N) + \dots + \alpha_k^2 L_k(N)}{N\theta} z_1 \right. \right. \\ &\quad \left. \left. + \sum_{l=2}^{\infty} \frac{\alpha_1^{l+1} L_1(N) + \dots + \alpha_k^{l+1} L_k(N)}{N\theta^l} z_1^l + \frac{1}{N} \partial_1 \right) h(z) \right] \cdot G_N(z), \end{aligned} \quad (53)$$

while, for $j = 2, 3, \dots, N$, the action of the swapping term can be written

$$\left(\frac{\theta}{N} \cdot \frac{1 - s_{1j}}{z_1 - z_j} \right) [h(z)G_N(z)] = \left[\left(\frac{\theta}{N} \hat{d}_1(j) + \frac{\theta}{N} \Delta d_j \right) h(z) \right] \cdot G_N(z), \quad (54)$$

where, for each j , $\hat{d}_1(j)$, Δd_j are linear operators on polynomials of z_1, z_j whose action on monomials $z_1^p z_j^q$, $p, q \in \mathbb{Z}_{\geq 0}$, is given by

$$\hat{d}_1(j) (z_1^p z_j^q) = \begin{cases} z_1^{p-1} z_j^q, & \text{if } p \geq 1, p > q \\ 0, & \text{otherwise} \end{cases} \quad (55)$$

and

$$\Delta d_j (z_1^p z_j^q) = \begin{cases} z_1^{p-2} z_j^{q+1} + z_1^{p-3} z_j^{q+2} + \dots + z_1^{q+1} z_j^{p-2} + z_1^q z_j^{p-1}, & \text{if } p > q, p > 1 \\ (-1)(z_1^p z_j^{q-1} + z_1^{p+1} z_j^{q-2} + \dots + z_1^{q-2} z_j^{p-1} + z_1^{q-1} z_j^p), & \text{if } p < q \\ 0, & \text{if } p = q. \end{cases} \quad (56)$$

Note that when summing over the other indices j ,

$$\sum_{j=2}^N \hat{d}_1(j) = (N - Y_1) \hat{d}_1$$

where we define Y_1 to be the operator that, when acting on a monomial, it multiply the monomial by the number of $j \in \{2, 3, \dots, N\}$ such that the degree of z_j is larger or equal to the degree of z_1 (e.g. $Y_1[x_1^3 x_3^3 x_4^2 x_7^5] = 2x_1^3 x_3^2 x_4^5 x_7$), and

$$\hat{d}_1(z_1^p) = \begin{cases} z_1^{p-1}, & \text{if } p \geq 1 \\ 0, & \text{otherwise} \end{cases} \quad (57)$$

We start with the expression $G_N(z)$ with $h(z) = 1$. From the above calculation, we see that

$$\begin{aligned} &(N^{-1}\mathcal{D}_1)^M G_N(z) \\ &= \left[\left(\frac{\alpha_1 L_1(N) + \dots + \alpha_k L_k(N)}{N} + \frac{\delta N + \alpha_1^2 L_1(N) + \dots + \alpha_k^2 L_k(N)}{N\theta} z_1 + \sum_{l=2}^{\infty} \frac{\alpha_1^{l+1} L_1(N) + \dots + \alpha_k^{l+1} L_k(N)}{N\theta^l} z_1^l \right. \right. \\ &\quad \left. \left. + \frac{\theta}{N} \sum_{j=2}^N \hat{d}_1(j) + \frac{\theta}{N} \Delta d_j + \frac{1}{N} \partial_1 \right)^M 1 \right] G_N(z) \\ &= \left[\left(\sum_{l=1}^{\infty} \kappa_l(N) \theta^{-(l-1)} z_1^{l-1} + \theta \frac{N - Y_1}{N} \hat{d}_1 + \frac{\theta}{N} \Delta d_j + \frac{1}{N} \partial_1 \right)^M 1 \right] G_N(z), \end{aligned} \quad (58)$$

where we have used Equation (31) to rewrite the expression in the second line in terms of the free cumulants.

Remark 3.3. Identify Y_1 with the scalar coming out from its action, which is changing from time to time. When N is very large, one should have the idea that in the “generic” case Y_1 is always small enough so that $\frac{N-Y_1}{N} \approx 1$, and we can think the corresponding term as $\theta \hat{d}_1$ without affecting the asymptotic of the action.

So far we have rewritten our action as the large product of operators acting on the polynomial $h(z) = 1$. Each of these operators is the sum of simpler operators

$$\kappa_l(N)\theta^{-(l-1)}z_1^{l-1} \quad (l = 1, 2, \dots), \quad \theta \frac{N - Y_1}{N} \hat{d}_1, \quad \frac{\theta}{N} \Delta d_j, \quad \text{and} \quad \frac{1}{N} \partial_1. \quad (59)$$

By expanding the expression in the last line of (58), we get a huge sum of the form

$$\left(\sum S_M \cdots S_2 S_1(1) \right) G_N(z), \quad (60)$$

where each operator S_i is chosen from those listed in Eqn. (59) and the sum is over all possible sequences of the S_i .

We may view each term in the sum as a discrete-time process $\{P(t)\}_{t=0}^M$ of polynomials in the variables z_1, \dots, z_N . That is, for any sequence S_1, \dots, S_M of operators, we get a sequence of polynomials such that $P(0) = 1$ and the polynomial at time t is given by $P(t) = S_t \cdots S_2 S_1(1)$. Note that at after step i , if $P(i) \neq 0$, in this process the total degree either

- increase by $l - 1$ if $S_i = \kappa_l(N)\theta^{-(l-1)}z_1^{l-1}$, or
- decreases by 1 otherwise.

Recall that our goal is to understand Eqn. (51), in which we set $z_1 = \dots = z_N = 0$. Since $G_N(0) = 1$,

$$(S_M \cdots S_2 S_1(1)) G_N(z) \Big|_{z_1 = \dots = z_N = 0} = (S_M \cdots S_2 S_1(1)) \Big|_{z_1 = \dots = z_N = 0}, \quad (61)$$

we see that only those $\{P(t)\}_{t=0}^M$ for which $P(M)$ is a constant will make a nontrivial contribution to the sum in (59). We make the following definition.

Definition 3.4. Let $\mathcal{P}(M)$ to be the collection of sequences of polynomials $\{P(t)\}_{t=0}^M$ in the variables z_1, \dots, z_N such that

- $P(0) = 1$,
- $P(t) = S_t \cdots S_2 S_1(1)$, where the S_i are chosen from the operators listed in Eqn. (59),
- and $P(M)$ is a constant.

We will see later (Sections 4 and 5) how to interpret these polynomial sequences from a probabilistic point of view.

Example 3.5. Let's consider the case $M = 3$ with two variables z_1 and z_2 . In this case, there are seven possible sequences (S_1, S_2, S_3) of operators that result in $\{P(t)\}_{t=0}^3 \in \mathcal{P}(3)$ with $P(3) \neq 0$. We list these as well as the corresponding sequences of polynomials below. Here we write $S = l, -1, \Delta, \partial$ as shorthand for the operators in (59).

(S_1, S_2, S_3)	$\{1, P(1), P(2), P(3)\}$
$(2, -1, -1)$	$\left\{1, \frac{\kappa_3(N)}{\theta^2} z_1^2, \frac{\kappa_3(N)}{\theta} z_1, \kappa_3(N)\right\}$
$(2, \partial, \partial)$	$\left\{1, \frac{\kappa_3(N)}{\theta^2} z_1^2, \frac{2\kappa_3(N)}{N\theta^2} z_1, \frac{2\kappa_3(N)}{N^2\theta^2}\right\}$
$(2, \Delta, \Delta)$	$\left\{1, \frac{\kappa_3(N)}{\theta^2} z_1^2, \frac{\kappa_3(N)}{N\theta} z_2, -\frac{\kappa_3(N)}{N^2}\right\}$
$(1, -1, 0)$	$\left\{1, \frac{\kappa_2(N)}{\theta} z_1, \kappa_2(N), \kappa_2(N)\kappa_1(N)\right\}$
$(1, 0, -1)$	$\left\{1, \frac{\kappa_2(N)}{\theta} z_1, \frac{\kappa_2(N)\kappa_1(N)}{\theta} z_1, \kappa_2(N)\kappa_1(N)\right\}$
$(0, 1, -1)$	$\left\{1, \kappa_1(N), \frac{\kappa_2(N)\kappa_1(N)}{\theta} z_1, \kappa_2(N)\kappa_1(N)\right\}$
$(0, 0, 0)$	$\left\{1, \kappa_1(N), \kappa_1(N)^2, \kappa_1(N)^3\right\}$

3.2 Further grouping

We now describe how we will group the different sequences $\{P(t)\}_{t=0}^M$. Recall that, among the four types of choices of operator S_t , $\kappa_l(N)\theta^{-(l-1)}z_1^{l-1}$ increases the total degree of $P(t-1)$ by $l-1 \in \mathbb{Z}_{\geq 0}$, while each of $\theta \frac{N-Y_1}{N} \hat{d}_1$, $\frac{\theta}{N} \Delta d_j$, and $\frac{1}{N} \partial_1$ decrease the total degree by 1. Of the three ways to decrease the total degree, we will think of $\theta \frac{N-Y_1}{N} \hat{d}_1$ as the “standard” choice, and think of the other two as so-called “weird” choices.

For a given $\{P(t)\}_{t=0}^M$, let $\vec{k} = (k_2, \dots, k_N)$, p denote the number of uses of the operators $\frac{\theta}{N} \Delta d_j$ ($j = 2, 3, \dots, N$), and $\frac{1}{N} \partial_1$, respectively, in $S_M \cdots S_2 S_1$. Denote the set of all sequences $\{P(t)\}_{t=0}^M$ with (\vec{k}, p) such choices of operators by $\mathcal{P}_{\vec{k}, p}$, where we suppress the dependence on M in the notation.

Now, by the above discussion, we have

Lemma 3.6. *We may write the Dunkl action on our BGF as*

$$\begin{aligned} \left(\frac{\mathcal{D}_1}{N}\right)^M G_N(z_1, \dots, z_N; \theta) \Big|_{z_1=\dots=z_N=0} &= \left(\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}(M)} P(M) \right) G_N(0, \dots, 0; \theta) \\ &= \left(\sum_{\vec{k}, p} \sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k}, p}} P(M) \right) \end{aligned} \quad (62)$$

where, by Definition 3.4, $P(M)$ is a constant for all $\{P(t)\}_{t=0}^M \in \mathcal{P}(M)$.

In subsequent sections, our approach when taking the asymptotic of the power sum will be as follows: fix $T > 0$ and take $M = \lfloor TN^{\frac{2}{3}} \rfloor$, we will show that for each fixed (\vec{k}, p) , the contribution of $\mathcal{P}_{\vec{k}, p}$ to

$$\frac{N}{(\mu_+(N)N)^M} \cdot \mathcal{D}_1^M G(z_1, \dots, z_N; \theta) \Big|_{z_1=\dots=z_N=0}$$

converges. Together with a tail estimate given in Section 7.6, this will verify that

$$\mathbb{E} \left[\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+(N)N} \right)^M \right]$$

converges as $M \rightarrow \infty$.

3.3 A Simplification

Throughout the remaining of the text, we replace the operator

$$\theta \frac{N-Y_1}{N} \hat{d}_1$$

that appears in Eqn. (59) by

$$\theta \hat{d}_1.$$

This is possible, as mentioned in Remark 3.3, since in generic case, Y_1 is of order $O(1)$, and as there are $M \sim N^{\frac{2}{3}}$ steps in the action, the difference caused by such replacement is of order

$$1 - \left(1 - \frac{1}{N}\right)^{N^{\frac{2}{3}}} \sim o(1).$$

The non-generic case occurs when the number of weird jumps given by $\frac{\theta}{N} \Delta d_j$'s is of order at least $N^{\frac{1}{3}} \sim M^{\frac{1}{2}}$. Since $|\frac{N-Y_1}{N}| \leq 1$, and we will show in Section 7.6, Lemma 7.28, that the total contribution such terms is exponentially small, such replacement still does not change the limit.

4 Mapping to random walks

In this section, we introduce our main tool for understanding the collection of discrete processes $\{P(t)\}_{t=0}^M$ of polynomials which have no “weird jumps”, that is, those $\{P(t)\} \in \mathcal{P}_{\bar{0},0}$. We will show that one can identify each such $\{P(t)\}_{t=0}^M$ with a discrete-time walk $\{W(t)\}_{t=0}^M$ on $\mathbb{Z}_{\geq 0}$. We will construct a probability measure on the space of all such walks by viewing them as certain random walks with iid steps conditioned to remain nonnegative. The distribution of the iid steps depends on the free cumulants of empirical measure of the eigenvalues of our ensemble. Along the way, we prove some combinatorial results showing that the partition function of the walks can be written in terms of the moment m_M of the spectral measure. Using this we get a contour integral expression for the partition function. Finally, we do a steepest descent analysis using this contour integral expression for the partition function of the walks as their length goes to infinity.

4.1 Discrete excursions and free cumulants

First, we will show that by considering the total degree of the polynomials in a sequence $\{P(t)\}_{t=0}^M \in \mathcal{P}_{\bar{0},0}$, we have a bijection with certain discrete-time excursion, that is, discrete-time walks on $\mathbb{Z}_{\geq 0}$ of fixed length M that begin and end at 0.

Proposition 4.1. *Fix $M \in \mathbb{Z}_{\geq 2}$, and define the discrete-time processes $\{P(t)\}_{t=0}^M$ as in Section 3.1. Then each sequence in $\mathcal{P}_{\bar{0},0}$ is in one-to-one correspondence to a discrete-time walk $\{W(t)\}_{t=0}^M$ on $\mathbb{Z}_{\geq 0}$, such that*

- $W(0) = W(M) = 0$.
- $W(t) \geq 0$ for all t 's.
- For $t = 0, 1, 2, \dots, M-1$,

$$W(t+1) - W(t) \in \{-1, 0, 1, 2, \dots\}.$$

Proof. Since we prohibit the use of “weird jumps”, by (59) the process $P(t)$ is driven by the actions of

$$\kappa_l(N)\theta^{-(l-1)}z_l^{l-1} \quad (l = 1, 2, \dots), \quad \text{or} \quad \theta \hat{d}_1. \quad (63)$$

Consider only the degree of z_1, \dots, z_N at time $t = 0, 1, 2, \dots, M$, one immediately has the following observations:

- The degrees of z_2, \dots, z_N are always 0 as the above operator can only add or remove powers of z_1 ,
- from time t to $t+1$, the degree of z_1 increases by $l-1 = 0, 1, 2, \dots$, when taking the a step of the first type of action, and decreases by 1 when taking the second type,
- the degree of z_1 is nonnegative throughout and must be zero at time M .

Define the height of the walk W at time t by

$$W(t) = \text{total degree of } P(t) = \text{degree of } z_1 \text{ of } P(t). \quad (64)$$

As the increment in the walk is completely determined by the increment in the total degree of $\{P(t)\}_{t=0}^M$ and vice-versa, this map is clearly bijective. The properties listed in the proposition follow immediately from the observations above, giving the result. \square

The above proposition motivates the following definition.

Definition 4.2. *Given $M \in \mathbb{Z}_{\geq 2}$, let $\mathcal{W}(M)$ be the space of all discrete walks $\{W(t)\}_{t=0}^M$ satisfying the three properties listed in Proposition 4.1.*

Next we consider the contribution of each $\{P(t)\}_{t=0}^M \in \mathcal{P}_{\bar{0},0}$ to the sum given in Lemma 3.6. By the lemma, each sequence of polynomials in $\mathcal{P}(M)$ contributes $P(M)$ to the sum, where by construction $P(M)$ is a constant. In the case that the sequence is in $\mathcal{P}_{\bar{0},0}$, this constant is particularly simple to describe.

First, we note that by the observations in the proof of Proposition 4.1, if $\{P(t)\}_{t=0}^M \in \mathcal{P}_{\bar{0},0}$ then each polynomial $P(t)$ is in fact a monomial in the variable z_1 . Using this, we give the following definition.

Definition 4.3. For each $\{P(t)\}_{t=0}^M \in \mathcal{P}_{\bar{0},0}$, let the weight $w_t(\cdot)$ of $\{P(t)\}_{t=0}^M$ be the coefficient of the monomial $P(t)$. Write $w(\cdot) = w_M(\cdot)$.

Because of the bijection given in Proposition 4.1, we assign weight to elements in $\mathcal{W}(M)$ in the natural way.

Remark 4.4. Note that from time t to $t+1$, by (59), w_t gets a new factor $\theta^{-(l-1)}$ when the total degree of $P(t)$ increases by $l-1$, and w_t gets multiplied by θ when the total degree decreases by 1 when we choose \hat{d}_1 . As the total degree starts and ends at zero, the degree of θ must be zero at time M . Therefore, in the remainder of this section, it's convenient to replace the choice of operators S_i as defined in Eqn. (59) by

$$\kappa_l(N)z_l^{l-1} \quad (l = 1, 2, \dots), \quad \hat{d}_1. \quad (65)$$

4.2 Conditional random walk

Here we give an alternative way to describe the excursions $\mathcal{W}(M)$. Given a sequence $\{\kappa_l(N)\}_{l=1}^\infty$ where $\kappa_l(N) \geq 0$ for all l 's, define a random variable $X(N)$ taking values in $\{-1, 0, 1, 2, \dots\}$ by

$$\begin{aligned} \mathbb{P}(X(N) = -1) &= \frac{1}{V_N(z)z}, \\ \mathbb{P}(X(N) = l) &= \frac{\kappa_{l+1}(N)z^l}{V_N(z)}, \quad l \geq 0 \end{aligned} \quad (66)$$

where $z > 0$ is a parameter we will specify shortly, and $V_N(z)$ is a normalizing factor so that these are probabilities. Note the normalizing factor

$$V_N(z) = \frac{1}{z} + \sum_{l=1}^{\infty} \kappa_l(N)z^{l-1}$$

is exactly the Voiculescu transform of $\{\kappa_l(N)\}_{l=1}^\infty$.

Suppose that $V_N'(z)$ has a unique positive zero $z_c(N)$ in its radius of convergence and that $V_N''(z_c(N)) > 0$. This will be true in the specific cases we are interested in. We have

$$\mathbb{E}[X(N)] = \frac{zV_N'(z)}{V_N(z)}, \quad \mathbb{E}[X(N)^2] = \frac{z^2V_N''(z)}{V_N(z)}. \quad (67)$$

Setting $z = z_c(N)$, we see that $\mathbb{E}[X(N)] = 0$ and

$$\text{Var}(X(N)) = \frac{z_c(N)^2V_N''(z_c(N))}{V(z_c(N))} := \sigma_N^2.$$

We have the following important observation relating the weight $w(\cdot)$ given in Definition 4.3 to these random variables $X(N)$.

Proposition 4.5. Treat $\mathcal{W}(M)$ as a probability space such that

$$\mathbb{P}[\{W(t)\}_{t=0}^M] = \frac{w(\{W(t)\}_{t=0}^M)}{\sum_{P \in \mathcal{W}(M)} w(P)}. \quad (68)$$

Then for $t = 0, 1, 2, \dots, M$, we may write

$$W(t) = \sum_{i=1}^t X_i$$

becomes a random walk where $X_i = -1, 0, 1, 2, \dots$

We have an equality in distribution between the walks $\{W(t)\}$ and the random walk

$$W'(t) = \sum_{i=1}^t X'_i,$$

such that the X'_i are independent with $X'_i \stackrel{d}{=} X(N)$ defined in Eqn. (66), and we condition on the event $W'(0) = W'(M) = 0$ and $W'(t) \geq 0$ for all $0 \leq t \leq M$.

Proof. This follows directly from Eqn. (66) and Section 4.1, once we observe that the degree of $z_c(N)$ of all $\mathbb{P}[W'(t)]$ in the conditional event is 0, because $W'(t)$ has no height change after time M . \square

Because of Proposition 4.5, we obtain a probabilistic point of view of $\{W(t)\}_{t=0}^M$, and we use it throughout the remaining of this text. We also observe that since

$$\sigma_N^2 \rightarrow \sigma^2 := \frac{z_c^2 V''(z_c)}{V(z_c)} \quad (69)$$

as indicated in Section 2.4, the sequence of unconditional walk $W'(t)$ (whose increments change with N) has

$$\frac{1}{\sigma\sqrt{M}} \sum_{i=1}^M X_i' \implies \mathcal{N}(0, 1), \quad (70)$$

as $N \rightarrow \infty$. In other words, $W'(t)$ lies in the domain of attraction of the standard Gaussian law.

4.3 The partition function

Our next goal is to calculate the partition function of $\mathcal{W}(M)$, given by

$$Z_{\mathcal{W}(M)} = \sum_{\{W(t)\}_{t=0}^M} w(\{W(t)\}_{t=0}^M). \quad (71)$$

As a first step, we recall the relation between moments $\{m_k\}_{k=1}^\infty$ and free cumulants $\{\kappa_l\}_{l=1}^\infty$ of some measure μ on \mathbb{R} . It is well-known that the n -th moment m_n can be expressed in terms as a sum of products of the free cumulants $\{\kappa_l\}_{l=1}^\infty$.

Proposition 4.6. (*[No, Section 1.6]*) *For any $M = 1, 2, \dots$, The M -th moment m_M can be expressed as*

$$m_M = \sum_{\pi \in NC(M)} \prod_{B \in \pi} \kappa_{|B|} \quad (72)$$

where the sum is over non-crossing partitions $\pi = B_1 \sqcup \dots \sqcup B_n$ of the set $\{1, 2, \dots, M\}$, $B \in \pi$ is a block of the partition, and $|B|$ is the size of the block.




Figure 1: The graph on the left represents a noncrossing partition π of $[6]$, where $B_1 = \{1, 4, 5, 6\}$, $B_2 = \{2, 3\}$, and the graph on the right represents a crossing partition π' of $[6]$, where $B_1 = \{1, 3, 5, 6\}$, $B_2 = \{2, 4\}$.

Next, we prove the following combinatorial result, that allows us to connect our discrete walks $\mathcal{W}(M)$ with free cumulants.

Proposition 4.7. Fix $M \in \mathbb{Z}_{\geq 2}$. There's an explicit bijection between $\mathcal{W}(M)$ and $NC(M)$, the set of all non-crossing partitions of $\{1, 2, \dots, M\}$.

Moreover, for $\{W(t)\}_{t=0}^M$ corresponding to $\pi \in NC(M)$,

$$w(\{W(t)\}_{t=0}^M) = \prod_{B \in \pi} \kappa_{|B|}(N). \quad (73)$$

Proof. The result is well-known, see e.g [NS, Proposition 9.8], but for completeness we provide a proof here. Given $\pi = B_1 \sqcup \dots \sqcup B_n \in NC(M)$, let $b_j = \min(B_j)$ be the leftmost element in block B_j . We construct a walk $\{W(t)\}_{t=0}^M$ by

- $W(0) = 0$.
- Let $W(t) - W(t-1) = l - 1$ if $t = b_j$ for some $j = 1, 2, \dots, n$, where $l = |B_j|$.
- Let $W(t) - W(t-1) = -1$ otherwise, for $t = 1, 2, \dots, M$.

We leave it to readers to check that $\{W(t)\}_{t=0}^M \in \mathcal{W}(M)$ under this construction. It's clear that $\pi \mapsto \{W(t)\}_{t=0}^M$ is injective. To see it's surjective, note that given any $\{W(t)\}_{t=0}^M \in \mathcal{W}(M)$, we may construct a partition $\pi = B_1 \sqcup \dots \sqcup B_n$ such that $\{\min(B_1), \dots, \min(B_n)\}$ is fixed by the times in which the walk goes up by $l_1 - 1, \dots, l_n - 1 \in \{0, 1, 2, \dots\}$. Let $|B_j| = l_j$, then the the starting position and size of each block fixes a unique non-crossing partition $\pi \in NC(M)$.

The weight in (73) follows from the above argument and (65). \square

By Propositions 4.6 and 4.7 we immediately see that

Corollary 4.8. Fix $M \in \mathbb{Z}_{\geq 2}$. The partition function of $\mathcal{W}(M)$ is equal to m_M , the M^{th} moment corresponding to free cumulants $\{\kappa_l(N)\}_{l=1}^{\infty}$. That is,

$$Z_{\mathcal{W}(M)} = \sum_{\{W(t)\}_{t=0}^M} w(\{W(t)\}_{t=0}^M) = m_M(N). \quad (74)$$

The following result is classical in free probability theory, and can be found in for example [BCG, Claim 9.1].

Proposition 4.9. Given a sequence of free cumulants $\{\kappa_l\}_{l=1}^{\infty}$, let $V(z) = \frac{1}{z} + \sum_{l=1}^{\infty} \kappa_l z^{l-1}$ be its Voiculescu transform. Then the corresponding moments

$$\begin{aligned} m_k &= \frac{1}{k+1} [z^{-1}] (V(z)^{k+1}) \\ &= \frac{1}{k+1} \frac{1}{2\pi i} \oint_{C_z} V(z)^{k+1} dz, \end{aligned} \quad (75)$$

for any $k = 1, 2, \dots$. The second equality follows from basic complex analysis, and it holds in the sense that $V(z)$ is naturally extended as a meromorphic function on \mathbb{C} , and C_z is a properly chosen contour that surrounds only one pole $z = 0$ of $V(z)$.

Proof. The result follows from the combinatorial description of the moments as partition functions of a random walk (Corollary 4.8) and Proposition 10.1 in Appendix A. \square

4.4 Steepest descent

According to Corollary 4.8 and Proposition 4.9, the partition function of $\mathcal{W}(M)$ is given by

$$m_M(N) = \frac{1}{M+1} \frac{1}{2\pi i} \oint_{C_z} V_N(z)^{M+1} dz. \quad (76)$$

To calculate its asymptotic when $N, M \rightarrow \infty$, we do steepest descent of the above contour integral. Recall that

$$\begin{aligned} V_N(z) &= \frac{1}{z} + \sum_{l=1}^{\infty} \kappa_l(N) z^{l-1} \\ &= \frac{1}{z} + \delta z + \frac{\gamma_1(N) a_1}{1 - a_1 z} + \dots + \frac{\gamma_k(N) a_k}{1 - a_k z} \end{aligned} \quad (77)$$

where the second equality holds when $-1/a_1 < z < 1/a_1$. Define $V(z)$ in a similar way by replacing each coefficient $\kappa_l(N)$ by its limit when $N \rightarrow \infty$. From now on we denote $V_N(z), V(z)$ as the meromorphic extension on \mathbb{C} given in the second line of Eqn. (77).

By direct calculation,

$$V'_N(z) = -\frac{1}{z^2} + \delta + \frac{\gamma_1(N) a_1^2}{(1 - a_1 z)^2} + \dots + \frac{\gamma_k(N) a_k^2}{(1 - a_k z)^2}, \quad (78)$$

$$V''_N(z) = \frac{2}{z^3} + \frac{2\gamma_1(N) a_1^3}{(1 - a_1 z)^3} + \dots + \frac{2\gamma_k(N) a_k^3}{(1 - a_k z)^3}, \quad (79)$$

One can check that V'_N is monotonically increasing from $-\infty$ to ∞ in $(0, \frac{1}{a_1})$ and therefore there's a unique zero of $V'_N(z)$ in $(0, \frac{1}{a_1})$, which we denote by $z_c(N)$. We choose our contour $C_z(N)$ in Eqn. (76) to be the circle centered at 0 with radius $z_c(N)$. Since on such $C_z(N)$, each $\kappa_l(N) z^{l-1}$ has constant modulus, and they have the same argument only if $z > 0$, $z_c(N)$ is the unique point that maximizes $|V_N(z)|$ on the contour.

Remark 4.10. Note that $V_N(z_c(N)) = \frac{1}{z_c} + \sum_{l=1}^{\infty} \kappa_l(N) z_c^{l-1}$ gives a convergent power series, where each term in the sum is positive.

Denote $f_N(z) = \log V_N(z)$, which is analytic near $z = z_c(N)$ since $V_N(z) > 0$ on $(0, \frac{1}{a_1})$. Let $C_z^\epsilon(N)$ be the arc of the circle $C_z(N)$ with argument $(-\frac{\epsilon}{\sqrt{M+1}}, \frac{\epsilon}{\sqrt{M+1}})$. Then we have

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{C_z(N)} V_N(z)^{M+1} dz \stackrel{(a)}{\approx} \frac{1}{2\pi i} \oint_{C_z^\epsilon(N)} V_N(z)^{M+1} dz \\ &= \frac{1}{2\pi i} \oint_{C_z^\epsilon(N)} \exp[(M+1)f_N(z)] dz \\ &\stackrel{(b)}{\approx} \frac{1}{2\pi i} \oint_{C_z^\epsilon(N)} \exp[(M+1)f_N(z_c(N))] \exp\left[(M+1) \frac{f''_N(z_c(N))}{2} (z - z_c(N))^2\right] dz \\ &\stackrel{(c)}{\approx} V_N(z_c(N))^{M+1} \cdot \frac{1}{2\pi i} \oint_{C_z^\epsilon} \exp\left[(M+1) \frac{f''(z_c)}{2} (z - z_c)^2\right] dz \\ &\stackrel{(d)}{\approx} \frac{V_N(z_c(N))^{M+1}}{\sqrt{M+1}} \cdot \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp\left[-\frac{|f''(z_c)|}{2} t^2\right] dt \\ &= \frac{V(z_c(N))^{M+1}}{\sqrt{M+1}} \cdot \frac{1}{\sqrt{2\pi}} \frac{1}{\sqrt{|f''(z_c)|}}. \end{aligned} \quad (80)$$

We are a bit sketchy here since the above calculation is quite standard: the integral on $c_z(N)$ is close to the integral on $c_z^\epsilon(N)$ since the modulus of the integrand decays exponentially fast away from it, then the rest follows from Taylor expansion and a change of variables. To make this rigorous, and to take its asymptotic ratio with $\mu_+(N)^M$, we prove the following lemma, such that the part (i) in which corresponds to the part (i) in Eqn. (80).

Lemma 4.11. (a). There exists constants $\delta_\epsilon(1) > \delta_\epsilon(2) > 0$, $K > 0$ independent of N , such that

$$|V_N(z_c(N))| - \sup_{z \in C_z(N) \setminus C_z^\epsilon(N)} |V_N(z)| \geq \delta_\epsilon(1) \cdot \frac{1}{\sqrt{M}}, \quad (81)$$

$$\sup_{z \in C_z^{\epsilon/K}(N)} |V_N(z_c(N)) - V_N(z)| \leq \delta_\epsilon(2) \cdot \frac{1}{\sqrt{M}}. \quad (82)$$

(b).

$$\sup_{z, z_c \in C_z^\epsilon(N)} \exp \left[(M+1) \frac{f^{(3)}(z_c(N))}{6} (z - z_c(N))^3 \right] \longrightarrow 1$$

as $N \rightarrow \infty$.

(c).

$$\begin{aligned} z_c(N) &\longrightarrow z_c, \\ f_N''(z_c(N)) &\longrightarrow f''(z_c), \end{aligned}$$

as $N \rightarrow \infty$, and

(d).

$$f''(z_c) > 0.$$

Proof. Eqn. (82) follows from estimating the difference of each corresponding term of $V_N(z_c(N))$ and $V_N(z)$ in the expression given by the second line of Eqn. (77). To prove Eqn. (81), we use the expression in the first line of (77) instead. It's not hard to see that

$$\begin{aligned} \left| V_N(z_c(N)) \right| - \sup_{z \in C_z(N) \setminus C_z^\epsilon(N)} |V_N(z)| &\geq \left| \min(1, \kappa_2(N)) \cdot \left(\frac{1}{z} + z \right) - \min(1, \kappa_2(N)) \cdot \left(\frac{1}{z_c(N)} + z_c(N) \right) \right| \\ &\geq \delta_\epsilon(1) \cdot \frac{1}{\sqrt{M}} \end{aligned} \quad (83)$$

for some $\delta_\epsilon(1) > 0$. To make $\delta_\epsilon(1) > \delta_\epsilon(2)$, just take K large enough.

(b) holds since $f^{(3)}(z_c(N))$ is clearly bounded and $z - z_c(N) \sim \frac{1}{\sqrt{M}}$ on $C_z^\epsilon(N)$. (c) and (d) hold by definition. \square

The asymptotic of our partition function follows from the above lemma.

Proposition 4.12. *We have that*

$$\mu_+(N) = V(z_c(N)), \quad (84)$$

and as $N \rightarrow \infty$,

$$\begin{aligned} \mu_+(N) &\rightarrow \mu_+, \\ N \cdot \frac{m_M(N)}{\mu_+(N)^M} &\rightarrow \frac{1}{\sqrt{2\pi}} \frac{V(z_c)}{\sqrt{f''(z_c)}} T^{-\frac{3}{2}}, \end{aligned} \quad (85)$$

equivalently,

$$\frac{m_M(N)}{\mu_+(N)^M} \sim \frac{1}{\sqrt{2\pi}} \frac{V(z_c)^{\frac{3}{2}}}{V''(z_c)^{\frac{1}{2}}} M^{-\frac{3}{2}}. \quad (86)$$

Proof. By (75) and (80) it remains to check the steps in (80) using Lemma 4.11.

To show (a), notice that

$$V_N(z) \approx V(z_c) > 0$$

on $C_z^\epsilon(N)$, and $C_z^\epsilon(N)$ is of length $\sim \frac{1}{\sqrt{M}}$. Given the two estimates in Lemma 4.11. (a), we have that

$$\left| \frac{1}{2\pi i} \oint_{C_z^\epsilon(N)} V_N(z)^{M+1} dz \right| \geq \text{const} \cdot \left(|V(z_c)| - \delta_\epsilon(2) \cdot \frac{1}{\sqrt{M}} \right)^{M+1} \cdot \frac{1}{\sqrt{M}}, \quad (87)$$

and

$$\begin{aligned} \left| \frac{1}{2\pi i} \oint_{C_z(N) \setminus C_z^\epsilon(N)} V_N(z)^{M+1} dz \right| &\leq \frac{1}{2\pi} \oint_{C_z(N) \setminus C_z^\epsilon(N)} |V_N(z)|^{M+1} dz \\ &\leq \frac{1}{2\pi} \oint_{C_z(N) \setminus C_z^\epsilon(N)} \left(|V_N(z_c(N))| - \delta_\epsilon(1) \cdot \frac{1}{\sqrt{M}} \right)^{M+1} dz \\ &\sim \text{const} \cdot \left(|V_N(z_c(N))| - \delta_\epsilon(1) \cdot \frac{1}{\sqrt{M}} \right)^{M+1}. \end{aligned} \quad (88)$$

Then clearly

$$\left| \frac{1}{2\pi i} \oint_{C_z(N) \setminus C_z^e(N)} V_N(z)^{M+1} dz \right| \ll \left| \frac{1}{2\pi i} \oint_{C_z^e(N)} V_N(z)^{M+1} dz \right| \quad (89)$$

when $M \rightarrow \infty$.

Eqn. (b) follows since we can uniformly bound the remainder term in Taylor expansion of $f_N(z)$ on $C_z^e(N)$. (c) follows since $c_z^e(N)$ and c_z^e are the same in shape, then by applying a change of variables

$$\frac{z_1 - z_c(N)}{z_c(N)} = \frac{z_2 - z_c}{z_c},$$

it's a direct consequence of dominated convergence theorem. (d) follows from another change of variables

$$z - z_c = i \cdot \frac{1}{\sqrt{M+1}} t,$$

and $f''(z_c) > 0$ gives the right sign of the exponent. \square

Remark 4.13. We will see the later that we may write

$$\frac{m_M(N)}{\mu_+(N)^M} \sim \frac{1}{\sqrt{2\pi}} \frac{V(z_c)^{\frac{3}{2}}}{V''(z_c)^{\frac{1}{2}}} M^{-\frac{3}{2}} = \frac{1}{\sqrt{2\pi}} \frac{1}{\sigma P_{-1}} M^{-3/2}. \quad (90)$$

where σ^2 is given in Eqn. (69) and P_{-1} is the limiting fraction of down steps (see Lemma 6.2).

5 First moment

Recall from Eqn. (51) that we have

$$\frac{N}{(\mu_+(N)N)^M} \cdot (\mathcal{D}_1^M) G_N(z_1, \dots, z_N; \theta) \Big|_{z_1=\dots=z_N=0} = \mathbb{E} \left[\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+(N)N} \right)^M \right]$$

To understand the action of the Dunkl operators on the BGF, we denoted the set of all sequences $\{P(t)\}_{t=0}^M$ with (\vec{k}, p) weird jumps respectively by $\mathcal{P}_{\vec{k}, p}$. Here k_j, p gives the number of jumps given by

$$\frac{1}{N} \Delta d_j (j = 2, 3, \dots, N), \quad \frac{1}{N\theta} \partial_1 \quad (91)$$

respectively.

Remark 5.1. Note that these differ from the form given in Eqn. (59) by a factor of θ^{-1} . We make this change to simplify our notation, following the same idea as Remark 4.4.

Fixing (\vec{k}, p) , we may write the contributions of $\mathcal{P}_{\vec{k}, p}$ to the Dunkl action above as

$$\left(\frac{N}{(\mu_+(N)N)^M} \sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{0}, 0}} P(M) \right) \cdot \frac{\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k}, p}} P(M)}{\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{0}, 0}} P(M)} = \left(\frac{N}{(\mu_+(N)N)^M} m_M(N) \right) \cdot \frac{\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k}, p}} P(M)}{Z_{\mathcal{W}(M)}} \quad (92)$$

where $Z_{\mathcal{W}(M)}$ is the partition function of the conditional random walks defined in Definition. 4.2 and the equality follows from Corollary 4.8.

In the remainder of this section we will show that

$$\frac{\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k}, p}} P(M)}{Z_{\mathcal{W}(M)}} = \frac{1}{Z_{\mathcal{W}(M)}} \sum_{W \in \mathcal{W}(M)} I_{\vec{k}, p; M}(W) w(\{W(t)\}_{t=0}^M)$$

where $I_{\vec{k}, p}$ is certain functional of the random walk excursion whose asymptotic expression we will show takes a universal form as a functional of a Brownian excursion. We separate the first moment calculation done in this section from the higher moment calculation done in Section 7 because, although the technique is similar for both cases, there are some subtleties in the higher moment case that could distract from the general argument.

5.1 Explicit Functional expression

5.1.1 The block process

First, let us consider the operators Δd_j whose action on monomials is given by Eqn. (56).

Recall from Section 4 that, the operator decreases the total degree of P by one. Due to this, we still have a mapping between the sequence of polynomials in which we allow the use of the operators Δd_j , $j = 2, \dots, N$, and the walks $\mathcal{W}(M)$ by mapping the total degree of the polynomial $P(t)$ to the height of the walk $W(t)$. In this case, however, $P(t)$ may be made up of several monomials, each with different powers of the variables z_i , $i \geq 1$ (but the same total degree). In order to keep track of the powers each of the variables z_1, \dots, z_N contribute to the total degree, we introduce the block process.

First, we define a piecewise continuous version of the walks in $\mathcal{W}(M)$.

Definition 5.2. Given $\{W(t)\}_{t=0}^M \in \mathcal{W}(M)$ define a piecewise continuous walk $E_M(s)$ by

$$E_M(s) = W(\lfloor s \rfloor) \quad 0 \leq s \leq M. \quad (93)$$

Let $\mathcal{E}(M)$ be the set of all such walks.

Definition 5.3. Given a sequence of polynomials $\{P(t)\}_{t=0}^M$ of z_1, \dots, z_N as in Section 3.1, define $q_i(s)$ ($i = 2, \dots, N$) such that

$$q_i(s) = \text{degree of } z_i \text{ in } P(\lfloor s \rfloor) \quad (94)$$

Clearly $q_j(s) \in \mathbb{Z}_{\geq 0}$ for all s , and is flat on each interval $[i-1, i)$ for $i = 1, 2, \dots, M-1$. Define the block process as

$$\vec{q}(s) = (q_2(s), \dots, q_N(s)) \quad (95)$$

and let

$$q(s) = \sum_{j=2}^N q_j(s) \quad (96)$$

The walk $E_M(s)$ as well as the block process $q(s)$ uniquely determines a choice of monomial in each polynomial in the sequence $\{P(t)\}_{t=0}^M$. We will view each component of the block process themselves as walks on the nonnegative integers by letting $q_i(t)$ be the height of the walk at time t . See Example 5.4. In fact, if we fix a walk E_M , then the paths $\vec{q}(t) = (q_2(t), \dots, q_N(t))$ come from a valid block process iff

- Each q_i is piecewise constant, right continuous, and has a left limit. Moreover, the jumps in q_i can only occur at the same time as down steps in E_M and only one of the q_i can have a jump at any given time.
- $E_M(t) \geq q(t)$ for all $t \in [0, M]$.
- If at time t there is a weird jump of height h in q_j then we must have

$$h \in \begin{cases} \{1, 2, \dots, E_M(t-) - q(t-) - q_j(t-) - 1\}, & \text{if } E_M(t-) - q(t-) > q_j(t-) \\ \{-1, -2, \dots, E_M(t-) - q(t-) - q_j(t-)\}, & \text{if } E_M(t-) - q(t-) < q_j(t-) \end{cases}$$



This essentially follows from Eqn. (56) and the definitions of E_M and the block process. Recall that, from Eqn. (56), a weird jump on variable z_j always comes with a factor of $\frac{1}{N}$; moreover, jumps that lowers the degree of z_j multiply the coefficient of the polynomial by a factor of -1 . We can replicate this by giving a weight to the block process of N^{-1} to the number of jumps and -1 to the number of down jumps in the block process \vec{q} .

Using the above we will be able to compute the sum

$$\frac{\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{k,p}} P(M)}{Z_{\mathcal{W}(M)}}$$

by first fixing a walk and taking the weighted sum over all possible block processes compatible with the walk, then summing over all walks. We will view the contribution of this sum for a fix walk as the desired functional of the walk.

Example 5.4. Below is an example with $M = 28$. On top, the solid black line is the walk $E_M(s)$. Below, the dashed blue line is $q_2(s)$, the dashed orange line is $q_3(s)$, and the dashed black line is $q(s)$, the sum of the other two dashed lines.



For this example, we have that the down step at $t = 6$ comes from the action of Δd_2 . The heights of the walk and the blocks at this time picks out the monomial $z_1^2 z_2^2$ in the polynomial $P(6)$. Similarly, the down step at $t = 11$ comes from the action of Δd_3 and the heights pick out the monomial $z_1^2 z_2^2 z_3^3$ from the polynomial $P(11)$. As there are three down jumps in the block process, it is given weight $(-1)^3$.

To that end, now consider the case when $|\vec{k}| > 1$, $p = 0$. As a first step, suppose $|\vec{k}| = 2$. Recall that we only consider sequences of polynomials such that $P(M)$ is a constant and note that Δd_j is the only operator that effects the power of z_j . As such, if $k_j \neq 0$, we must have $k_j \geq 2$, since the first action of the operator will increase the power of z_j and the last action must decrease the power back to zero. Thus, in this case, we can only have weird jumps corresponding to a single variable, say z_j .

Remark 5.5. This constraint will no longer be present in the higher moment case, Section 7.

Given $\vec{v} = (v_2, \dots, v_N) \in \mathbb{Z}_{\geq 0}^{N-1}$, let $v = \sum_{i=2}^N v_i$. For a fix excursion E_M , define

$$A_{\vec{v},h}^j(s,t) := \mathbf{1}(E_M(\tau) \geq v \text{ for all } \tau \in [s,t]) \cdot \mathbf{1}(|h| \leq |E_M(t-) - v - v_j - (\text{sgn}(h) + 1)/2|) \\ \cdot \mathbf{1}(h(E_M(t-) - v - v_j) > 0).$$

One should think of this as follows: if at time s , we have that the height of the blocks in the block process are given by $\vec{q}(s-) = \vec{v}$, then $A_{\vec{v},h}^j(s,t)$ is 1 iff it is possible that the next weird jump is in variable j and occurs at time t with height h , otherwise the product of indicators is 0.

Returning to the case $|\vec{k}| = 2$, let $t_1 < t_2$ be the times of the weird jumps and h_1, h_2 be their size, then the weighted sum over all possible block processes compatible with E_M is given by

$$\begin{aligned} I_{2,0;M}^j(E_M) &= \frac{1}{N^2} \sum_{\substack{h_1, h_2 \\ h_1+h_2=0}} \sum_{\substack{\text{down steps} \\ t_1 < t_2}} \left(\prod_{i=1}^2 \text{sgn}(h_i) \right) A_{\vec{0}, h_1}^j(0, t_1) A_{k_1 \vec{e}_j, h_2}^j(t_1, t_2) \\ &= -\frac{1}{N^2} \sum_{\text{down step } t_1} \sum_{h_1 > 0} \mathbf{1}(E_M(t_1) \geq h_1). \end{aligned}$$

where \vec{e}_j is the j -th standard basis vector. The second equality follows from the fact that $A_{\vec{0}, h_1}^j(0, t_1)$ is only nonzero if $h_1 > 0$ and, moreover, h_2 and t_2 are completely determined as $A_{k_1 \vec{e}_j, h_2}^j(t_1, t_2)$ is only nonzero if $E_M(t_2-) = h_1$ since $h_2 = -h_1$. In particular, t_2 must be the first time after t_1 in which $E_M(t_2) = h_1 - 1$.

Let $I_{2,0;M}(E_M) = \sum_{j=2}^N I_{2,0;M}^j(E_M)$. From the above, we have

$$\begin{aligned} \sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k}, 0}, |\vec{k}|=2} P(M) &= \sum_{E_M \in \mathcal{E}(M)} I_{2,0;M}(E_M) w(E_M) \\ &= (N-1) \sum_{E_M \in \mathcal{E}(M)} I_{2,0;M}^2(E_M) w(E_M) \end{aligned}$$

where the factor of $N-1$ comes from the $N-1$ choices of index j .

If $|\vec{k}| = m > 0, p = 0$ and only one k_j is non-zero, we have that the weighted sum over all possible block processes compatible with E_M can be written

$$\begin{aligned} I_{m,0;M}^j(E_M) &= \frac{1}{N^m} \sum_{\substack{h_1, \dots, h_m \\ h_1 + \dots + h_m = 0}} \sum_{\substack{\text{down steps} \\ t_1 < \dots < t_m}} \left(\prod_{i=1}^m \text{sgn}(h_i) \right) \left(A_{\vec{0}, h_1}^j(0, t_1) A_{h_1 \vec{e}_j, h_2}^j(t_1, t_2) A_{(h_1+h_2) \vec{e}_j, h_3}^j(t_2, t_3) \dots \right. \\ &\quad \left. A_{(h_1+\dots+h_{m-1}) \vec{e}_j, h_m}^j(t_{m-1}, t_m) \right). \end{aligned}$$

Remark 5.6. *While in the case $|\vec{k}| = 2$ we were able to give a relatively simple expression for the functional, in the general case it is unclear if such a simplifications can be done. Note, however, that our later arguments will only require control of the scaling of the functional and not on its precise expression.*

5.1.2 The partial derivative operator

Now, let us consider the case $\vec{k} = \vec{0}$. In order to collect the contribution of each application of ∂_1 operator to Eqn. (92) we recall that each partial derivative returns the height of the walk at that time and that these jumps always correspond to down steps. For a fixed walk, we may think of marking p of the down steps and multiplying the weight of the walk by its height at those points.

To that end, define a functional

$$I_{0,p;M}(E_M) = \frac{1}{(N\theta)^p} \sum_{\substack{\text{down steps} \\ s_1 < s_2 < \dots < s_p}} \prod_{i=1}^p E_M(s_i-) \quad (97)$$

where, from (91), each use of the partial derivative comes with a factor of $\frac{1}{N}$. We see that we may write

$$\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{0}, p}} P(M) = \sum_{E_M \in \mathcal{E}(M)} I_{0,p;M}(E_M) w(E_M).$$

Note that the partial derivative is in the variable z_1 while the height of the walk corresponds to the total degree. Thus, in general, an application of the partial derivative at time s brings down a factor of $E_M(s-)$ minus the total height of the blocks in the block process, rather than just $E_M(s-)$.

5.1.3 Full Functional Expression

Now let us consider the general case. Although here we write an explicit expression for the functional, its exact form will not be important in what follows. As mentioned in Remark 5.6, our arguments will only require control of the scaling of the functional and not on its precise expression.

For this case we can have weird jumps in several of the variables z_j or, equivalently, in several $q_j(t)$ of the block process. Let $n(\vec{k}) = n$ be the number of nonzero k_i 's. Note that there are $\binom{N-1}{n}$ ways to choose which k_i are nonzero and that the resulting functional does not depend on what choice we make. Thus we can always choose k_2, \dots, k_{n+1} to be nonzero and multiply the functional by an overall factor of $\binom{N-1}{n}$ to handle the other possible choices.

For a fixed vector \vec{k} , define the set $\mathcal{M}_{\vec{k}}$ of all possible pairings

$$\mu_{\vec{k}} : \{1, \dots, |\vec{k}|\} \rightarrow \underbrace{\{2, \dots, 2\}}_{k_2} \underbrace{\{3, \dots, 3\}}_{k_3} \dots \underbrace{\{N, \dots, N\}}_{k_N}$$

in which $\mu_{\vec{k}}(j) = i$ means the j -th jump in the block process is a jump in q_i . Each $\mu_{\vec{k}} \in \mathcal{M}_{\vec{k}}$ gives a choice of ordering of the jumps. Note that, as before, the last jump for each of the q_i is completely determined by the previous jumps.

Finally, for a fixed walk E_M and fixed choice of \vec{k} with $n(\vec{k}) = n$, we may write the weighted sum over all compatible block processes as the functional

$$\begin{aligned} I_{\vec{k}, p; M}(E_M) &= \frac{1}{N^{|\vec{k}|+p} \theta^p} \binom{N-1}{n} \\ &\cdot \sum_{\mu \in \mathcal{M}_{\vec{k}}} \sum_{\substack{h_1^{(2)}, \dots, h_{k_2}^{(2)} \\ h_1^{(2)} + \dots + h_{k_2}^{(2)} = 0}} \dots \sum_{\substack{h_1^{(N)}, \dots, h_{k_N}^{(N)} \\ h_1^{(N)} + \dots + h_{k_N}^{(N)} = 0}} \sum_{\substack{\text{down steps} \\ t_1 < \dots < t_{|\vec{k}|}}} \sum_{\substack{\text{down steps} \\ s_1 < \dots < s_p}} \left(\mathbf{1}(t_i \neq s_j, i = 1, \dots, |\vec{k}|, j = 1, \dots, p) \right) \\ &\cdot \left(\prod_{i=2}^N \prod_{j=1}^{k_i} \text{sgn}(h_j^{(i)}) \right) \left(\prod_{i=1}^p (E_M(s_i -) - q(s_i -)) A_{\vec{h}}^\mu(t_1, \dots, t_{|\vec{k}|}) \right) \end{aligned} \quad (98)$$

where $\vec{h} = (h_*^{\mu(1)}, \dots, h_*^{\mu(|\vec{k}|)})$ with the subscripts starting from 1 and increasing to the right for each superscript,

$$\begin{aligned} A_{\vec{h}}^\mu(t_1, \dots, t_{|\vec{k}|}) &= \left(A_{\vec{0}, h_1}^{\mu(1)}(0, t_1) A_{h_1 \vec{e}_{\mu(1)}, h_2}^{\mu(2)}(t_1, t_2) A_{h_1 \vec{e}_{\mu(1)} + h_2 \vec{e}_{\mu(2)}, h_3}^{\mu(3)}(t_2, t_3) \dots \right. \\ &\quad \left. A_{h_1 \vec{e}_{\mu(1)} + h_2 \vec{e}_{\mu(2)} + \dots + h_{|\vec{k}|-1} \vec{e}_{\mu(|\vec{k}|-1)}}^{\mu(|\vec{k}|)}(t_{|\vec{k}|-1}, t_{|\vec{k}|}) \right), \end{aligned}$$

and $q(s) = \sum_{i=1}^{|\vec{k}|} h_i \mathbf{1}(t_i < s)$.

With this we have

$$\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k}, p}} P(M) = \sum_{E_M \in \mathcal{E}(M)} I_{\vec{k}, p; M}(E_M) w(E_M).$$

Example 5.7. Continuing Example 5.4, we have $\vec{k} = (3, 2)$ and $p = 0$.

The weird jump heights are given by

$$\begin{aligned} h_1^{(2)} &= 2, & h_2^{(2)} &= -1, & h_3^{(2)} &= -1 \\ h_1^{(3)} &= 3, & h_2^{(3)} &= -3 \end{aligned}$$

They occur at times

$$t_1 = 6, t_2 = 11, t_3 = 16, t_4 = 18, t_5 = 25$$

with pairing $\mu : \{1, 2, 3, 4, 5\} \rightarrow \{2, 2, 2, 3, 3\}$ given by

$$\mu(1) = 2, \mu(2) = 3, \mu(3) = 2, \mu(4) = 3, \mu(5) = 2.$$

To conclude this section, note that we can view $w(E_M)/Z_{\mathcal{W}(M)}$ as a probability measure on the set of conditioned random walks. Thus we may write

$$\begin{aligned} \frac{\sum_{\{P(t)\}_{t=0}^M \in \mathcal{P}_{\vec{k},p}} P(M)}{Z_{\mathcal{W}(M)}} &= \frac{\sum_{E_M \in \mathcal{E}(M)} I_{\vec{k},p;M}(E_M) w(E_M)}{Z_{\mathcal{W}(M)}} \\ &= \mathbb{E}_{\mathcal{E}(M)}[I_{\vec{k},p;M}] \end{aligned} \quad (99)$$

where the expectation is over the conditioned random walks $\mathcal{E}(M)$.

5.2 Scaling

Here we will show that

$$\frac{\sum_{E_M \in \mathcal{E}(M)} I_{\vec{k},p;M}(E_M) w(E_M)}{Z_{\mathcal{W}(M)}} \sim O(1)$$

as $M \rightarrow \infty$ for each choice of fixed \vec{k} and p . Suppose $n(\vec{k}) = n$. First note that the contribution of the partial derivative operators has a simple bound

$$\prod_{i=1}^p (E_M(s_{i-}) - q(s_{i-})) \leq \prod_{i=1}^p (E_M(s_{i-}) \leq (\max_s E_M(s))^p \sim (\sigma\sqrt{M})^p,$$

where the last line holds with high probability as $M \rightarrow \infty$ by the functional CLT given in the next section. As each of the derivative operators must occur at a down step and, as we will see in Lemma 6.2, the limiting fraction of down steps is given by $P_{-1} = \frac{1}{z_c V(z_c)}$, we have

$$\frac{1}{(N\theta)^p} \sum_{\substack{\text{down steps} \\ s_1 < s_2 < \dots < s_p}} \prod_{i=1}^p (E_M(s_{i-}) - q(s_{i-})) \lesssim \frac{1}{(N\theta)^p} (P_{-1}M)^p (\sigma\sqrt{M})^p \sim \left(\frac{\sigma P_{-1} T^{3/2}}{\theta} \right)^p$$

where we recall that $M = \lfloor TN^{\frac{2}{3}} \rfloor$.

We can similarly bound the number of ways to choose the heights and the times of the jumps corresponding to the operators Δd_j . For these, however, recall that the height and time of the last jump for each index j is completely determined by the previous jumps. Thus we have

$$\begin{aligned} &\frac{1}{N^{|\vec{k}|}} \sum_{\mu \in \mathcal{M}_{\vec{k}}} \sum_{\substack{h_1^{(2)}, \dots, h_{k_2}^{(2)} \\ h_1^{(2)} + \dots + h_{k_2}^{(2)} = 0}} \dots \sum_{\substack{h_1^{(N)}, \dots, h_{k_N}^{(N)} \\ h_1^{(N)} + \dots + h_{k_N}^{(N)} = 0}} \sum_{\substack{\text{down steps} \\ t_1 < \dots < t_{|\vec{k}|}}} |A_h^\mu(t_1, \dots, t_{|\vec{k}|})| \\ &\lesssim \frac{1}{N^{|\vec{k}|}} \binom{|\vec{k}|}{k_1, k_2, \dots} (P_{-1}M)^{|\vec{k}|-n} (\sigma\sqrt{M})^{|\vec{k}|-n} \\ &\sim \frac{1}{N^n} \binom{|\vec{k}|}{k_1, k_2, \dots} (\sigma P_{-1} T^{3/2})^{|\vec{k}|-n} \end{aligned}$$

where the multinomial coefficient counts the number of ways to order the weird jumps coming from \vec{k} .

Altogether we have

$$|I_{\vec{k},p;M}(E_M)| \lesssim \frac{1}{N^n} \binom{N-1}{n} \binom{|\vec{k}|}{k_1, k_2, \dots} \frac{(\sigma P_{-1} T^{3/2})^{|\vec{k}|+p-n}}{\theta^p} \sim O(1).$$

5.3 Functional CLT

Our next goal is to show that under proper rescaling, $I_{\vec{k},p;M}$, the functional on $E_M(s)$, converges to $I_{\vec{k},p}$, a limiting functional on $\mathfrak{E}(s)$, where $\mathfrak{E}(s)$ is a standard Brownian excursion on $[0, 1]$. This follows from a limit transition of the functional, and some technical estimates that will be proved in Section 6.

Here we state a general result on the convergence of random walks to Brownian excursions due to [CC] (see also [S]). Let $S(t) = u_1(M) + \dots + u_t(M)$, $t = 0, 1, 2, \dots, M$ be an integer valued aperiodic random walk with

i.i.d step $u_i(M)$ whose distribution depends on M , and $\mathbb{E}[u_i(M)] = 0$. We assume that S lies in the domain of attraction of the standard Gaussian law, i.e, there exists $\sigma > 0$ such that

$$\frac{S_M}{\sigma\sqrt{M}} \implies \mathbf{N}(0, 1). \quad (100)$$

Let $P_M^{*,x,y}$ be the law of $S(t)$ conditioned to be non-negative and satisfy $S(0) = x \geq 0, S(m) = y \geq 0$. Define Ω to be the space of càdlàg functions on $[0, 1]$ endowed with the standard Skorohod topology and define $X^M : \mathbb{Z}^M \rightarrow \Omega$ by

$$X^M(u_1, \dots, u_M)(s) = \left(\frac{1}{\sigma\sqrt{M}} \sum_{i=1}^{\lfloor tM \rfloor} u_i \right)_{s \in [0,1]}. \quad (101)$$

Theorem 5.8. [CC, Theorem. 2.4 and Corollary 2.5] As $M \rightarrow \infty$, the following convergence holds in Ω :

$$P_n^{*,0,0} \circ (X^M)^{-1} \implies \mathfrak{E} \quad (102)$$

where \mathfrak{E} is the law of the standard Brownian excursion.

By the discussion at the end of Section 4.2, the unconditional walk $W^l(t)$ lies in the domain of attraction of standard Gaussian law, with $\sigma^2 = \frac{z_c^2 V''(z_c)}{V(z_c)}$. Let

$$\mathfrak{E}_M(s) = \frac{1}{\sigma\sqrt{M}} E_M(sM)$$

for $s \in [0, 1]$. Then by the Theorem 5.8, $\mathfrak{E}_M(t)$ converges weakly to \mathfrak{E} as $M \rightarrow \infty$.

Theorem 5.9. As $M \rightarrow \infty$,

$$I_{\vec{k},p;M}(\mathfrak{E}_M) \rightarrow I_{\vec{k},p}(\mathfrak{E}), \quad (103)$$

where

$$\begin{aligned} I_{\vec{k},p}(\mathfrak{E}) &= (\sigma P_{-1} T^{3/2})^{|\vec{k}|+p-n} \\ &\cdot \sum_{\mu \in \mathfrak{M}_{\vec{k}}} \int d\vec{H} \int_{u_1 < \dots < u_{|\vec{k}|-n+1}} d\vec{u} \int_{v_1 < \dots < v_p} d\vec{v} \left(\prod_{i,j} \text{sgn}(H_j(i)) \right) \left(\prod_{i=1}^p \mathfrak{E}(v_i) - y(v_i) \right) A^\mu(\vec{u}, \vec{H}; \mathfrak{E}) \\ &= (\sigma P_{-1} T^{3/2})^{|\vec{k}|+p-n} \\ &\cdot \sum_{\mu \in \mathfrak{M}_{\vec{k}}} \int d\vec{H} \int_{u_1 < \dots < u_{|\vec{k}|-n+1}} d\vec{u} \left(\prod_{i,j} \text{sgn}(H_j(i)) \right) \frac{1}{p!} \left(\int_0^1 \mathfrak{E}(s) - y(s) ds \right)^p A^\mu(\vec{u}, \vec{H}; \mathfrak{E}) \end{aligned} \quad (104)$$

for a particular function $A^\mu(\vec{u}, \vec{H}; \mathfrak{E})$. Here $\vec{u} \in \mathbb{R}^{|\vec{k}|-n+1}$, $\vec{v} \in \mathbb{R}^p$, and $\vec{H} \in \mathbb{R}^{|\vec{k}|-n+1}$ has $k_i - 1$ components for each $i = 2, \dots, n+1$ which we denote $H_j(i)$ for $j = 1, \dots, k_i - 1$, and $y(s)$ is the sum of all heights in \vec{H} that have occurred before time s .

Moreover,

$$\mathbb{E}_{\mathcal{E}(M)}[I_{\vec{k},p;M}] \rightarrow \mathbb{E}[I_{\vec{k},p}], \quad (105)$$

where the expectation on the right side is taken on the standard Brownian excursion \mathfrak{E} .

Proof. By the above theorem the random walks $\mathfrak{E}_M(t)$ converges weakly to \mathfrak{E} as $M \rightarrow \infty$. In Section 6.3 we show that we have good control on the tail of the random walk, therefore the convergence in expectation follows from the weak convergence.

After rescaling all heights by $\frac{1}{\sigma\sqrt{M}}$ the sums

$$\sum_{\substack{h_1^{(2)}, \dots, h_{k_2}^{(2)} \\ h_1^{(2)} + \dots + h_{k_2}^{(2)} = 0}} \dots \sum_{\substack{h_1^{(N)}, \dots, h_{k_N}^{(N)} \\ h_1^{(N)} + \dots + h_{k_N}^{(N)} = 0}}$$

become Riemann sums for the integrals

$$(\sigma\sqrt{M})^{|\vec{k}|-n} \int d\vec{H}$$

where $\vec{H} = \{H_j(i)\}_{j=1,\dots,k_i-1, i=2,\dots,n+1}$, $d\vec{H} = \prod_{i,j} dH_j(i)$, and the integral is over $\mathbb{R}^{|\vec{k}|-n}$. Note here we used that the last height in each sum is fixed by the previous heights.

Similarly, after rescaling time by $\frac{1}{M}$, the sums

$$\sum_{\substack{\text{down steps} \\ t_1 < \dots < t_{|\vec{k}|}}} \sum_{\substack{\text{down steps} \\ s_1 < \dots < s_p}}$$

become Riemann sums for the integrals

$$(P_{-1}M)^{|\vec{k}|+p-n} \int_{u_1 < \dots < u_{|\vec{k}|-n}} d\vec{u} \int_{v_1 < \dots < v_p} d\vec{v}$$

where $\vec{v} \in \mathbb{R}^{|\vec{k}|-n+1}$ and $\vec{u} \in \mathbb{R}^p$. Here we again using that the last jump times of each z_i are completely determined by the previous jumps. The factor P_{-1} takes into account that the fraction of down steps is asymptotically P_{-1} , on each local region of $\mathfrak{E}_M(s)$, see Lemma 6.2.

Note that the product of indicator functions $A_h^\mu(t_1, \dots, t_{|\vec{k}|})$ is still well-defined when we replace E_M with \mathfrak{E} and the jump heights by there rescaled versions. Call the corresponding product of indicators $A^\mu(\vec{u}, \vec{H}; \mathfrak{E})$. Similarly, by replacing E_M with \mathfrak{E} and q by y , we can replace

$$\left(\prod_{i=1}^p (E_M(s_i-) - q(s_i-)) \right) \text{ by } (\sigma\sqrt{M})^p \left(\prod_{i=1}^p (\mathfrak{E}(s_i) - y(s_i)) \right).$$

Finally, we are left with a factor

$$\frac{(\sigma P_{-1} M^{3/2})^{|\vec{k}|+p-n}}{N^{|\vec{k}|+p}} \binom{N-1}{n} \rightarrow (\sigma P_{-1} T^{3/2})^{|\vec{k}|+p-n}$$

□

5.4 Universality

Recall that in Theorem 1.2 we would like to show that

$$\left\{ \lambda_i' := \frac{\lambda_i(N) - \mu_+(N)N}{N^{\frac{1}{3}}} \right\}_{i=1}^N \implies C_0 \cdot \text{Airy}(\beta).$$

In other words, after proper rescaling, the limiting distribution of each $\{\lambda_i'\}$ is the same regardless of choice of ensemble. For a single $G\beta E$ we know from the classical result in [RRV, Theorem 1.1] that $C_0 = 1$, $\mu_+ = 2$, and clearly $\sigma = 1$, $P_{-1} = \frac{1}{2}$.

As we calculated the first moment of $\sum_i e^{\frac{T\lambda_i'}{\mu_+}}$, for the edge universality to hold, we must have that the limiting expression in Theorem 5.9 is the same as in the $G\beta E$ case, after replacing T by $\frac{\mu_+}{2C_0}T$ for a general ensemble.

For the first moment and fixed \vec{k} and p , note that the only place where information about the specific matrix summation appears in the limiting functional (104) is through the factor $(\sigma P_{-1} T^{3/2})^{|\vec{k}|+p-n}$. The universality then follows if for a general ensemble

$$\sigma P_{-1} \left(\frac{\mu_+}{2C_0} \right)^{3/2} = 1 \cdot \frac{1}{2}.$$

But this follows immediately after writing each quantity on the LHS in terms of z_c and $V(z_c)$.

6 Properties of random walk bridges

Consider the (conditional) random walks with increments given by Section 4.2. For $x, y \in \mathbb{Z}_{\geq 0}$, $L \in \mathbb{Z}_{>1}$, let $\mathcal{W}(x, y, L; M)$ be the set of walk bridges from x to y of length L , with increments and weights given by Eqn. (66), conditioning on staying nonnegative. In particular, $\mathcal{W}(0, 0, M; M)$ gives the walk excursion $\{W(t)\}_{t=0}^M$ arising in the first moment calculation in Section 5. We use $|\mathcal{W}(x, y, L; M)|$ to denote the partition function of $\mathcal{W}(x, y, L; M)$.

In this section, we provide several results regarding the walks sampled from $\mathcal{W}(x, y, L; M)$.

6.1 Conditional functional CLT's

We introduce two more results from [CC], that deal with the conditional random walk bridge with different starting and ending heights. These will be used later in Section 7 when we prove the convergence in high moments.

Fix $c > 0$. Define $S(t)(t = 0, 1, 2, \dots, \lfloor cM \rfloor)$, $X^{\lfloor cM \rfloor} : \mathbb{Z}^{\lfloor cM \rfloor} \rightarrow \Omega$, $P^{*,x,y}$ similarly as in Section 5.3.

Theorem 6.1. *Suppose x_M, y_M are integers such that*

$$\frac{x_M}{\sigma\sqrt{M}} \rightarrow u > 0, \quad \frac{y_M}{\sigma\sqrt{M}} \rightarrow v > 0$$

as $M \rightarrow \infty$. Then one has the convergence

$$P_{\lfloor cM \rfloor}^{*,0,x_M} \circ (X^{\lfloor cM \rfloor})^{-1} \Longrightarrow \mathfrak{E}_u(t), \quad (106)$$

$$P_{\lfloor cM \rfloor}^{*,y_M,0} \circ (X^{\lfloor cM \rfloor})^{-1} \Longrightarrow \tilde{\mathfrak{E}}_v(t), \quad (107)$$

as $M \rightarrow \infty$, where $\mathfrak{E}_u(t)(t \in [0, c])$ is a Brownian bridge from 0 to u conditioned on staying nonnegative, $\tilde{\mathfrak{E}}_v(t)(t \in [0, c])$ is a Brownian bridge from v to 0 conditioned on staying nonnegative.

Suppose x_M, y_M are integers such that

$$\frac{x_M}{\sigma\sqrt{M}} \rightarrow u > 0, \quad \frac{y_M}{\sigma\sqrt{M}} \rightarrow v > 0$$

as $M \rightarrow \infty$. Then one has the convergence

$$P_{\lfloor cM \rfloor}^{*,x_M,y_M} \circ (X^{\lfloor cM \rfloor})^{-1} \Longrightarrow \mathfrak{E}_{u,v}(t), \quad (108)$$

as $M \rightarrow \infty$, where $\mathfrak{E}_{u,v}(t)(t \in [0, c])$ is a Brownian bridge from u to v conditioned on staying nonnegative.




Figure 2: The three cases for the paths in Theorem 6.1.

6.2 Time homogeneity of down steps

Lemma 6.2. *Suppose we are given a sequence in M of random walks $W_M(t)$, sampled from $\mathcal{W}(x_M, y_M, L; M)$, such that there are constants $C > 0, c_2 > c_1 > 0$ uniform in M , where $0 \leq x_M, y_M \leq C\sqrt{M}$, $c_1M \leq L \leq c_2M$. Then for any $0 < t_1 < t_2 < 1$, define $\rho_{-1}(t_1, t_2)$ to be the proportion of down-steps in $\llbracket t_1L, t_2L \rrbracket$. Then we have*

$$\rho_{-1}(t_1, t_2) \rightarrow P_{-1} \quad (109)$$

as $M \rightarrow \infty$, where $P_{-1} = \frac{1}{z_c V(z_c)}$.

Remark 6.3. *By the definition of $X_1(N)$, we recall from Section 4.2 that*

$$P_{-1} = \lim_{N \rightarrow \infty} \mathbb{P}[X(N) = -1]. \quad (110)$$

Proof. Since $t_1 > 0, t_2 < 1$, $x_M, y_M \sim \sqrt{M}$, under the scaling of $W_M(s)$ according to Theorem 6.1, we have that with high probability,

$$0 < \tilde{c}_1 \leq \frac{1}{\sqrt{L}} W_M(\lfloor t_1 L \rfloor) \leq \tilde{c}_2, \quad 0 < \tilde{c}_1 \leq \frac{1}{\sqrt{M}} W_M(\lfloor t_2 L \rfloor) \leq \tilde{c}_2$$

for some constants $\tilde{c}_1, \tilde{c}_2 > 0$. Therefore, it suffices to prove the convergence of the statement for a walk $W'_M(s)$ sampled from $\mathcal{W}(x_M, y_M, L; M)$, uniformly for any fixed integer-valued $x_M, y_M \in \llbracket \tilde{c}_1 \sqrt{L}, \tilde{c}_2 \sqrt{L} \rrbracket$.

Let $\tilde{W}(t) (t = 0, 1, 2, \dots, L)$ be a walk with the same increments and weights as $W(t)$, except that it doesn't have the conditioning to stay nonnegative. Observe that in the regime of x_M, y_M we are considering,

$$\mathbb{P}[\tilde{W}(t) \geq 0 \text{ for all } t \mid \tilde{W}(0) = x_M, \tilde{W}(L) = y_M] > \epsilon \quad (111)$$

for some $\epsilon > 0$, uniformly in x_M, y_M, L, M .

The idea of remaining of the proof is to first remove the conditioning that the walk stays nonnegative, then calculate the probability distribution of ρ_{-1} , defined as the proportion of down steps of $\tilde{W}(t)$, and show that with probability going to 1 we have $\rho_{-1} \approx P_{-1}$. More precisely, we have for any $\delta > 0$,

$$\begin{aligned} \mathbb{P} \left[|\rho_{-1} - P_{-1}| > \delta \mid \tilde{W}(t) \geq 0 \text{ for all } t \right] &= \frac{\mathbb{P} \left[|\rho_{-1} - P_{-1}| > \delta, \tilde{W}(t) \geq 0 \text{ for all } t \right]}{\mathbb{P} \left[\tilde{W}(t) \geq 0 \text{ for all } t \right]} \\ &\leq \frac{1}{\epsilon} \mathbb{P} \left[|\rho_{-1} - P_{-1}| > \delta \right]. \end{aligned} \quad (112)$$

It remains to show the last line above goes to 0 as $M \rightarrow \infty$, uniformly in $y_1(M), y_2(M)$. This is done by a straightforward calculation.

Abbreviate $\mathbb{P}[X_1(M) = i]$ by $p_i(M)$ and $y_M - x_M$ by $H(M) \in [(c_1 - c_2 - 1)\sqrt{L}, (c_2 - c_1 + 1)\sqrt{L}]$, $H(M)'$ by $L + H(M)$. Then for $J_{-1}, J_0, J_1, J_2, \dots \in \mathbb{Z}_{\geq 0}$,

$$J_{-1} + J_1 + \dots + J_{H(M)'} = L \quad (113)$$

where J_i is the number of jumps of size i . We have

$$\begin{aligned} &\mathbb{P} \left[\text{there are } J_i \text{ jumps of size } i \text{ in the walk, for each } i = -1, 0, 1, \dots, H(M)' \right] \\ &= \binom{L}{J_{-1} \ J_0 \ J_1 \ \dots \ J_{H(M)'}} p_{-1}(M)^{J_{-1}} p_0(M)^{J_0} p_1(M)^{J_1} \dots p_{H(M)'}(M)^{J_{H(M)'}} \\ &\approx (2\pi)^{-\frac{H(M)'}{2}} (H(M)')^{H(M)'+\frac{1}{2}} J_{-1}^{-(J_{-1}+\frac{1}{2})} J_0^{-(J_0+\frac{1}{2})} \dots J_{H(M)'}^{-(J_{H(M)'}+\frac{1}{2})} \\ &\quad \cdot p_{-1}(M)^{J_{-1}} p_0(M)^{J_0} \dots p_{H(M)'}(M)^{J_{H(M)'}} \end{aligned} \quad (114)$$

conditioning on the event that

$$-J_{-1} + J_1 + 2J_2 + \dots + H(M)' J_{H(M)'} = H(M). \quad (115)$$

The third line of Eqn. (114) follows from Stirling's formula.

By Eqn. (113) and (115), we rewrite J_{-1}, J_1 in terms of the other terms, such that

$$\begin{aligned} J_{-1} &= \frac{1}{2} \left[H(M)' - J_0 + J_2 + 2J_3 + \dots + (H(M)' - 1) J_{H(M)'} \right] - \frac{1}{2} H(M) \\ &\approx \frac{1}{2} \left[H(M)' - J_0 + J_2 + 2J_3 + \dots + (H(M)' - 1) J_{H(M)'} \right], \\ J_1 &= \frac{1}{2} \left[H(M)' - J_0 - 3J_2 - 4J_3 - \dots - (H(M)' + 1) J_{H(M)'} \right] + \frac{1}{2} H(M) \\ &\approx \frac{1}{2} \left[H(M)' - J_0 - 3J_2 - 4J_3 - \dots - (H(M)' + 1) J_{H(M)'} \right]. \end{aligned} \quad (116)$$

where the \approx makes sense since $H(M) \sim O(\sqrt{M})$ and $J_{-1}, J_1 \sim O(M)$.

Our goal is to find $(J_{-1}, J_0, J_1, \dots)$ that maximize the probability in Eqn. (114). We take the logarithm of the right side of Eqn. (114), plug in the expression in Eqn. (116), and take the partial derivative of the resulting expression in J_i 's. Then we have after taking $\frac{\partial}{\partial J_i}$ ($i = 0, 2, 3, \dots$):

$$\begin{aligned}
& \log(p_i(M)) - \frac{i+1}{2} \log(p_1(M)) + \frac{i-1}{2} \log(p_{-1}(M)) \\
& - \left[\log(J_i) + \frac{J_i + \frac{1}{2}}{J_i} - \frac{i+1}{2} \log(J_1) - \frac{J_1 - \frac{1}{2}i + 1}{J_1} + \frac{i-1}{2} \log(J_{-1}) + \frac{J_{-1} + \frac{1}{2}i - 1}{J_{-1}} \right] \\
& = \log(p_i(M)) - \frac{i+1}{2} \log(p_1(M)) + \frac{i-1}{2} \log(p_{-1}(M)) \\
& - \left[\log\left(\frac{J_i}{H(M)'}\right) + \frac{J_i + \frac{1}{2}}{J_i} - \frac{i+1}{2} \log\left(\frac{J_1}{H(M)'}\right) - \frac{J_1 - \frac{1}{2}i + 1}{J_1} + \frac{i-1}{2} \log\left(\frac{J_{-1}}{H(M)'}\right) + \frac{J_{-1} + \frac{1}{2}i - 1}{J_{-1}} \right] \quad (117) \\
& = 0 \\
& \approx \log(p_i(M)) - \frac{i+1}{2} \log(p_1(M)) + \frac{i-1}{2} \log(p_{-1}(M)) \\
& - \left[\log(J_i) - \frac{i+1}{2} \log(J_1) + \frac{i-1}{2} \log(J_{-1}) \right],
\end{aligned}$$

where in the approximation in the last line we treat J_i as a positive real number, such that $\frac{J_i + \frac{1}{2}}{J_i} \approx 1$ (this is the case for the generic step i 's that dominate the walk), and similarly $\frac{J_1 - \frac{1}{2}}{J_1}, \frac{J_{-1} + \frac{1}{2}}{J_{-1}} \approx 1$.

There are $H(M)' - 1$ (approximately linear) equations with $H(M)' - 1$ unknowns $\log(J_2), \dots, \log(J_{H(M)'})$. We observe that for each M ,

$$\frac{J_i}{L} = \frac{J_i}{J_{-1} + J_0 + \dots + J_{H(M)'}} \approx p_i(M), \quad (118)$$

hence $J_i \approx [p_i(M)L]$ (close to 0 when i is large) are the unique local region that maximize

$$\mathbb{P}[\text{there are } J_i \text{ jumps of size } +i \text{ in the walk}].$$

It's direct to see from Eqn. (114) that as $M \rightarrow \infty$ the distribution function concentrates near its maximum, which in particular implies that for any $\delta > 0$,

$$\mathbb{P}\left[\left|\frac{J_{-1}}{H(M)'} - \mathbb{P}[X_1(M) = -1]\right| > \delta\right] \rightarrow 0.$$

Since $\rho_{-1} = \frac{J_{-1}}{L}$, $X_1(M) \rightarrow X_1$ in distribution, the result follows. \square

6.3 Tail estimate of the walks

We give two results of the extreme values of such walks, that are used in the tail estimate argument in Section 7.6.

Let $\mathcal{W}_1(x, L; M), \mathcal{W}_2(x, L; M)$ be the walk with the same increments conditioning on staying nonnegative, with $W(0) = x/W(L) = x$ respectively.

Lemma 6.4. (a). For fixed $x \in \mathbb{Z}_{\geq 0}, L \in \mathbb{Z}_{\geq 1}$, let $W(t)$ be a walk sampled from $\mathcal{W}_2(x, L; M)$, where $W(0)$ is undetermined. Then there exists constant $C > 0, c > 0$, which are uniform in M, L and x , such that for all $h > 0$,

$$\mathbb{P}[W(0) - x > h\sqrt{L}] \leq C \exp(-ch^2).$$

(b). For fixed $x, y \in \mathbb{Z}_{\geq 0}, L \in \mathbb{Z}_{\geq 1}$, $W(t)(t = 0, 1, 2, \dots, L)$ be sampled from $\mathcal{W}(x, y, L; M)$. Then there exists constant $C > 0, c > 0$, which are uniform in M, L, x and y , such that for all $h > 0$,

$$\mathbb{P}\left[\max_{0 \leq t \leq L} W(t) - \max(x, y) > h\sqrt{L}\right] \leq C \exp(-ch^2).$$

Proof. By taking the constant C large enough, it suffices to consider $h > 100$.

Proof of (a): For brevity we only consider the case $x = 0$, and the general cases can be argued similarly. When $y = 0$, we are sampling $W(t)$ from $\mathcal{W}_2(0, L; M)$, and

$$\mathbb{P}[W(0) = \lfloor h\sqrt{L} \rfloor] = \frac{|\mathcal{W}(\lfloor h\sqrt{L} \rfloor, 0, L; M)|}{|\mathcal{W}_2(0, L; M)|}.$$

Note that the inequality is trivial when $\lfloor h\sqrt{L} \rfloor > L$, since our walk has increments ≥ -1 . And for $\lfloor h\sqrt{L} \rfloor \leq L$, the result can be obtained from a steepest descent using Lemma 10.4 and Lemma 10.5.

Proof of (b): If $x > y$, let $\bar{\tau} = \max_{t \in \llbracket 0, L \rrbracket} \{t : W(t) = x\}$, and it suffices to show the statement for $y = x$ conditioning on $\bar{\tau}$, which is included in the case $x \leq y$. So we assume $x \leq y$.

For a walk $W(t)$ sampled from $\mathcal{W}(x, y, L; M)$ such that $W(t) \geq y + \lfloor h\sqrt{L} \rfloor$ for some t , let $\tau = \max_{t \in \llbracket 0, L \rrbracket} \{t : W(t) = y + \lfloor h\sqrt{L} \rfloor\}$ be the last time $W(t)$ is above $y + \lfloor h\sqrt{L} \rfloor$. We split the walk as a concatenation of two walk bridges on $\llbracket 0, \tau \rrbracket$ and $\llbracket \tau, L \rrbracket$ respectively, and count the partition function of such concatenation. Compared to $\mathcal{W}(x, y, L; M)$, $W(t)$ is fixed at $t = L - \tau$, which gives a factor $\frac{1}{\sqrt{L-\tau}}$. Moreover, the unconditional walk bridge from $y + \lfloor h\sqrt{L} \rfloor$ to y with length $L - \tau$ has a weight $\sim \exp\left(-c\frac{(h\sqrt{L})^2}{L-\tau}\right)$, and the conditioning event that such walk is below $\lfloor y + h\sqrt{L} \rfloor$ gives a factor $\leq \frac{h\sqrt{L}}{L-\tau}$. Summing over at most L choices of τ gives the result. \square

Lemma 6.5. For $x, y \in \mathbb{Z}_{\geq 0}$, $L, M \in \mathbb{Z}_{\geq 1}$, sample the walk bridge $W(t)$ in $\mathcal{W}(x, y, L; M)$. Take discrete intervals $\llbracket X_1, Y_1 \rrbracket, \dots, \llbracket X_k, Y_k \rrbracket$ contained in $\llbracket 0, L \rrbracket$. Assume that $Y_i - X_i \geq 1$, and the intervals are distinct except for at endpoints. For each $i = 1, 2, \dots, k$, denote $V_i = W(X_i) - \min_{\llbracket X_i, Y_i \rrbracket} W + 1$. Take $a_1, \dots, a_k \in \mathbb{Z}_{>0}$ and let $a = a_1 + \dots + a_k$. Then there exists constant $C > 0$ uniform in x, y, L, M , such that

$$\mathbb{E} \left[\prod_{i=1}^k \frac{V_i^{a_i}}{a_i!} \right] < C^a \log(a)^a (a^{-\frac{a}{2}} L^{\frac{a}{2}} + a^{-a} \max(x-y, 0)^a), \quad (119)$$

and such expectation has a Gaussian tail, in the sense that there exists another constant $c > 0$ uniform in x, y, L, M , such that

$$\begin{aligned} \mathbb{E} \left[\prod_{i=1}^k \frac{V_i^{a_i}}{a_i!} (\mathbf{1}_{\max W > \max(x, y) + h\sqrt{L}} + \mathbf{1}_{\min W < \min(x, y) - h\sqrt{L}}) \right] \\ < C^a \log(a)^a (a^{-\frac{a}{2}} L^{\frac{a}{2}} + a^{-a} (\max(x-y, 0) + h\sqrt{L})^a) \exp(-ch^2) \end{aligned} \quad (120)$$

for all $h > 0$.

We postpone its proof to Appendix B.

7 Higher moments

In this section, we present a more general way to express the l^{th} moments of

$$\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+(N)N} \right)^M$$

for any fixed $T > 0$, $l \geq \mathbb{Z}_{\geq 1}$. Recall from Proposition 2.7 and discussion in Section 3.1 that for $l \in \mathbb{Z}_{\geq 1}$, the l^{th} moment is obtained from the action of

$$\frac{1}{\mu_+(N)^{lM}} \cdot \left[\left(\frac{\mathcal{D}_1}{N} \right)^M + \dots + \left(\frac{\mathcal{D}_N}{N} \right)^M \right]^l, \quad (121)$$

which we may rewrite as

$$\frac{1}{\mu_+(N)^{lM}} \cdot \left[\sum_{n_0=1}^l \binom{N}{n_0} \sum_{l_1+\dots+l_{n_0}=l} \binom{l}{l_1, \dots, l_{n_0}} \left(\frac{\mathcal{D}_{n_0}}{N} \right)^{l_{n_0}M} \dots \left(\frac{\mathcal{D}_1}{N} \right)^{l_1M} \right] \quad (122)$$

where, because of the symmetry of $G(z_1, \dots, z_N; \theta)$ and the commutativity of D_j 's, we may assume that only the first n_0 Dunkl operators appear in each term. Note that the combinatorial sum

$$\sum_{n_0=1}^l \sum_{l_1+\dots+l_{n_0}=l}$$

is finite for each fixed l , hence it suffices to study for fixed $n_0 \in \mathbb{Z}_{\geq 1}$, $(l_1, \dots, l_{n_0}) \in \mathbb{Z}_{\geq 1}^{n_0}$, the action

$$\left(\frac{D_{n_0}}{N}\right)^{l_{n_0}M} \cdots \left(\frac{D_1}{N}\right)^{l_1M} \quad (123)$$

on $G(z_1, \dots, z_N; \theta)$. By almost the same arguments as in Section 3, we can again write our expression as a huge sum of sequences of monomials $\{P(t)\}_{t=0}^{lM}$, each of which corresponds to a (random) walk bridge $\{W(t)\}_{t=0}^{lM}$ in which the height of the walk $W(t)$ equals the total degree of the polynomial $P(t)$. Note that the length of $\{P(t)\}_{t=0}^{lM}$ and $\{W(t)\}_{t=0}^{lM}$ is now lM instead of M . Moreover, similar to Section 5, after proper modification each $\{W(t)\}_{t=0}^{lM}$ is obtained from the sequence of actions chosen from one of the four operators

$$\kappa_l(N)z_i^{l-1} \quad (l = 1, 2, \dots), \quad \hat{d}_i, \quad \frac{1}{N}\Delta d_j (j \neq i), \quad \text{and} \quad \frac{1}{N}\theta^{-1}\partial_i. \quad (124)$$

We call z_i the *main variable* if up-steps in the walk correspond to increases in the degree of z_i . Unlike Section 5, for which z_1 was always the main variable, in this case the main variable depends on where we are in the walk. We have that the main variable is z_1 for $t = 1, 2, \dots, l_1M$, z_2 for $t = l_1M + 1, \dots, (l_1 + l_2)M, \dots$, and z_{n_0} for $t = (l_1 + \dots + l_{n_0-1})M + 1, \dots, lM$. The weird jumps on the i^{th} time segment are given by $\Delta_j d_j (j \neq i)$, which changes the degree of z_i and z_j just as described in Eqn. (56) with z_1 replaced with z_i .

As in Section 5, we construct a piecewise continuous version of $W(t)$. Let

$$E_M(t) = W(\lfloor t \rfloor) \quad 1 \leq t \leq lM + 1. \quad (125)$$

For $i = 1, 2, \dots, n_0 - 1$, let $C_i = (l_1 + \dots + l_i)M$ be the splitting time between when z_i is the main variable and when z_{i+1} is the main variable (such a time does not exist for $l = 1$). Let $C_{n_0} = lM$.

Define the block processes $q_i(t) (i = 1, 2, \dots, N)$ in the similar way as Definition 5.3, such that they give the degree of z_i on $[0, C_{n_0}] \setminus [C_{i-1}, C_i]$ if $i = 1, 2, \dots, n_0$, and on $[0, C_{n_0}]$ if $i = n_0 + 1, \dots, n$, throughout the actions. Note, however, that we may now have blocks q_1 corresponding to the degree of z_1 , when z_1 is not the main variable. Let $q(t) = \sum_{j \neq i} q_j(t)$, for $t \in [C_{i-1}, C_i]$ ($i = 1, 2, \dots, n_0$). From this definition, we have each $q_i(t)$ and $q(t)$ is Cádrlág on $[0, lM + 1]$, and all the discontinuity points are given by the occurring time of the weird jumps of z_1, \dots, z_n and the splitting point C_i 's. Moreover, $E_M(t) - q(t)$ gives the degree of the main variable at time t .

Fix n_0 and (l_1, \dots, l_{n_0}) . As before, we fix the number of weird jumps $k_1, \dots, k_{n_0} \geq 0, k_{n_0+1}, \dots, k_n \geq 1$, where k_i gives the number of Δd_i 's changing the degree of z_i and the current main variable. The block process $\vec{q} = (q_1(s), \dots, q_n(s))$ is determined by the jumps

$$\{(s_m(i), h_m(i))\}_{m=1,2,\dots,k_i, i=1,2,\dots,n},$$

such that $(s_m(i), h_m(i))$ gives the occurring time $s_m(i) \in \llbracket 1, C_{i-1} \rrbracket \cup \llbracket C_i + 1, C_{n_0} \rrbracket$, and height

$$h_m(i) = q_i(s_m(i)) - q_i(s_m(i)-) \in \mathbb{Z} \setminus \{0\}$$

of the m^{th} weird jump changing the degree of z_i and the current main variable.

Let p be the number of ∂ 's from time 1 to lM . We split further the ∂ actions by

$$\vec{p} = (p_1, \dots, p_{n_0}),$$

where $p = p_1 + \dots + p_{n_0}$, p_i is the number of ∂_i 's at times when z_i is the main variable.

Remark 7.1. For each weird jump we have $1 \leq s_m(i) \leq lM$ and $s_m(i), h_m(i) \in \mathbb{Z}$. Unlike the first moment case, in which the heights of the weird jumps for a given variable must sum to zero, for general $l > 1$, $h_1(i) + \dots + h_{k_i}(i)$ is not necessarily 0 as we will see in Section 7.2.

As in the first moment case, we will write the Dunkl action as the sum of certain functionals of the random walks, one for every choice of \vec{k} and \vec{p} . In fact, we will show that the action

$$\binom{N}{n_0} \binom{l}{l_1, \dots, l_{n_0}} \frac{\left(\frac{D_1}{N}\right)^{l_1 M} \dots \left(\frac{D_{n_0}}{N}\right)^{l_{n_0} M}}{\mu_+(N)^{lM}}$$

on $G(z_1, \dots, z_N; \theta)$ can be written as the sum of $C(N, n_0, \vec{k}) \cdot I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M)$ over the parameters \vec{k}, \vec{p} . Here $C(N, n_0, \vec{k})$ is a combinatorial prefactor and $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M)$ is the contribution to the Dunkl action coming from terms with (\vec{k}, \vec{p}) weird jumps divided by the partition function of the conditioned random walk (see Eqn. (133)).

We will show that as $N \rightarrow \infty$ these functionals converge to functionals on a Brownian excursion. Showing this convergence is more subtle than in the first moment case. We will see that there are some walks/functional combinations for which the limiting value does not converge to a finite value; however, such terms will appear in pairs with opposite sign and cancel up to a finite order term.

7.1 Some technical lemmas

We begin by proving two results about the integer-valued random walk with i.i.d steps. The first result considers the contribution of the configurations when the walk goes below 0 in a small time, and the second considers the probability it stays nonnegative for time $O(M)$, where $M \rightarrow \infty$.

Lemma 7.2. *For $x \in \mathbb{Z}_{\geq 1}$, $\tau \in \mathbb{Z}_{\geq 1}$, consider a random walk bridge $W(t)$ ($t = 0, 1, 2, \dots, M$), that is sampled from $\mathcal{W}(x, x, \tau - 1; M)$, and $W(\tau) - W(\tau - 1) = -1$.*

Fix some $0 < \epsilon < 1$, we have that there is a constant $C > 0$ uniform in x, ϵ, M ,

$$\sum_{\tau=1}^{\epsilon M} \tau \cdot \frac{|\mathcal{W}(x, x, \tau - 1; M)| z_c(N)^{-1}}{\mu_+(N)^\tau} \leq C\sqrt{\epsilon M}, \quad (126)$$

for all $0 \leq t \leq \epsilon M$.

Proof. Since $z_c(N)$ and $\mu_+(N)$ converges as $M \rightarrow \infty$, it suffices to estimate

$$\tau \cdot \frac{|\mathcal{W}(x, x, \tau - 1; M)|}{\mu_+(N)^{\tau-1}}$$

for $0 \leq \tau - 1 \leq \epsilon M - 1$. By Proposition 4.12,

$$\frac{|\mathcal{W}(x, x, \tau - 1; M)|}{\mu_+(N)^{\tau-1}} = \frac{m_{\tau-1}(N)}{\mu_+(N)^{\tau-1}} \sim \tau^{-\frac{3}{2}}$$

when τ is large. Then by choosing the constant C to be large enough, one can bound $\frac{|\mathcal{W}(x, x, \tau-1; M)|}{\mu_+(N)^{\tau-1}}$ by $C\tau^{-\frac{3}{2}}$. Summing over all τ 's gives the result. \square

Let $\mathcal{W}_1(x, L; M)$, $\mathcal{W}_2(x, L; M)$ be the walk with the same increments conditioning on staying nonnegative, with $W(0) = x/W(L) = x$ respectively. We use $|\cdot|$ to denote the partition function of these two types of walks as for $\mathcal{W}(x, y, L; M)$.

Let $W_1(t)$, $W_2(t)$ be walks sampled from $\mathcal{W}_1(0, L; M)/\mathcal{W}_2(0, L; M)$ respectively. Here one can think $W_2(t)$ as a walk running backward in time.

Lemma 7.3. *Fix parameter $c > 0$. Then as $L \rightarrow \infty, M \rightarrow \infty$, we have*

$$\sqrt{L} \cdot \mathbb{P}[W_2(t) \geq 0 \text{ for all } t] \rightarrow \frac{1}{\sqrt{2\pi}} \frac{\sigma}{P_{-1}}.$$

Proof. By Lemma 10.4, we have that the partition function for walks starting at a height H and ending at a height 0 after L steps restricted to stay weakly above zero is given by

$$\frac{H+1}{L+1} \frac{1}{2\pi i} \oint \left(\frac{z}{z_c(N)} \right)^H V_N(z)^{L+1} dz. \quad (127)$$

Let $W'(t) = W'_M(t)$ ($t = 0, 1, 2, \dots, L$) be a walk with the same ending point 0, same increments and weights as $W_2(t)$, but without conditioning to stay nonnegative. We may write

$$\begin{aligned}
& \mathbb{P}(W'(t) \geq 0 \text{ for all } t \in \{0, \dots, L\}) \\
&= \sum_{H=0}^{\infty} \mathbb{P}(W'(0) = H, W'(t) \geq 0 \text{ for all } t \in \{0, \dots, L\}) \\
&= \frac{1}{V_N(z_c(N))^L} \sum_{H=0}^{\infty} \frac{H+1}{L+1} \frac{1}{2\pi i} \oint \left(\frac{z}{z_c(N)} \right)^H V_N(z)^{L+1} dz \\
&= \frac{1}{V_N(z_c(N))^L (L+1)} \frac{1}{2\pi i} \oint \left(\sum_{H=0}^{\infty} (H+1) \left(\frac{z}{z_c(N)} \right)^H \right) V_N(z)^{L+1} dz \\
&= \frac{1}{V_N(z_c(N))^L (L+1)} \frac{1}{2\pi i} \oint \frac{1}{(1-z/z_c(N))^2} V_N(z)^{L+1} dz,
\end{aligned}$$

where our contour contains only the pole at zero. Now by Lemma 10.5, using steepest descent as $L \rightarrow \infty$, we have

$$\frac{1}{2\pi i} \oint \frac{1}{(1-z/z_c(N))^2} V_N(z)^{L+1} dz \approx \frac{\sqrt{L+1} z_c(N)^2 V_N(z_c(N))^{L+1}}{\sqrt{2\pi}} \left(\frac{V_N''(z_c(N))}{V_N(z_c(N))} \right)^{1/2}. \quad (128)$$

(There are some subtleties that must be dealt with due to the critical point $z_c(N)$ of $V_N(z)$ being a pole of $\frac{1}{(1-z/z_c(N))^2}$, see the proof of Lemma 10.5 for details.) It follows that

$$\mathbb{P}(W'(t) \geq 0 \text{ for all } t \in \{0, \dots, L\}) \approx \frac{1}{\sqrt{L+1}} \frac{z_c(N)^2 V_N(z_c(N))}{\sqrt{2\pi}} \left(\frac{V_N''(z_c(N))}{V_N(z_c(N))} \right)^{1/2}. \quad (129)$$

Multiplying both sides by \sqrt{L} , then taking $L \rightarrow \infty$, $M \rightarrow \infty$ gives the limit $\frac{1}{\sqrt{2\pi}} \frac{\sigma}{P_{-1}}$, where the last expression follows from the fact that $\sigma^2 = \frac{z_c^2 V''(z_c)}{V(z_c)}$ and $P_{-1} = \frac{1}{z_c V(z_c)}$. \square

Lemma 7.3 gives us the following weight of $W_1(t)$, $W_2(t)$. For any given $y \sim \sqrt{M}$, by Theorem 6.1 and local limit theorem for our conditional walk (see e.g [BD, Proposition 1] and [S, Lemma 3.1 and 3.2]), we have for $L = \lfloor cM \rfloor$ for some $c > 0$,

$$\mathbb{P}\left(\frac{1}{\sigma\sqrt{M}} W_2(0) = y\right) \sim \frac{1}{P_{-1}} \frac{1}{M}, \quad (130)$$

where we hide the constant depending on c and y . Recall also that the random walk bridge from 0 to 0 of length $2cM$ has partition function $m_{2cM}(N) \sim \frac{1}{P_{-1}\sigma} V_N(z_c(N))^{2cM} M^{-\frac{3}{2}}$. Then again by Theorem 5.8 and local limit theorem, the walk bridge of length $2cM$ (rescaled by $\sigma\sqrt{M}$) arriving at height y at time cM has weight $\sim \frac{1}{P_{-1}\sigma^2} V_N(z_c(N))^{2cM} M^{-2}$. Let $\bar{W}(t) = \bar{W}_M(t)$ ($t = 0, 1, 2, \dots, L$) be a walk with the same ending point 0, same increments and weights as $W_1(t)$, but without conditioning to stay nonnegative. By the Markov property and local limit theorem, we get that for $L = \lfloor cM \rfloor$,

$$\mathcal{P}(\bar{W}(t) \geq 0 \text{ for all } t) = \frac{|\mathcal{W}(0, \lfloor cM \rfloor; M)|}{\mu_+(N)^{\lfloor cM \rfloor}} \sim \frac{1}{\sigma} \frac{1}{\sqrt{M}}, \quad (131)$$

where we hide some different constant depending on c and y .

7.2 Piecewise walks and scaling

Consider the space of Cadlag functions $\mathbb{D}([0, lM + 1])$ with the usual topology. For a given block process \vec{q} , we define a set $A_M(\vec{q})$ of $f \in \mathbb{D}([0, lM + 1])$ such that $f \in A_M(\vec{q})$ if all the followings hold:

- $f(t) \geq q(t)$ for $t \in [0, lM + 1]$.
- For the jump $(s_m(i), h_m(i))$,

$$|h_m(i)| \leq |f(t-) - q(t-) - q_i(t-) - (\text{sgn}(h_m(i)) + 1)|.$$

- For the jump $(s_m(i), h_m(i))$,

$$h_m(i) \cdot (f(t-) - q(t-) - q_i(t-)) > 0.$$

Specify this to the process $E_M(s)$ which is C adl ag on $[0, lM + 1]$. We say $E_M \in B_M(\vec{q})$ if all jumps in \vec{q} occur at down-steps in E_M , that is, times s such that $E_M(s + 1) - E_M(s) = -1$.

$E_M \in A_M(\vec{q})$, $E_M \in B_M(\vec{q})$ comprise all the necessary conditions for the appearance of the pair (E_M, \vec{q}) of walk/block configuration in the action of (123). But to understand the contribution to the terms to the action in the general $l > 1$ case is considerably more complicated than the case $l = 1$.

For a given (E_M, \vec{q}) , we consider the block process $q_i(t)$ for $i = 1, 2, \dots, n$. The behavior of the last weird jump of the variable z_i depends on whether or not that variable is one of the possible main variables. We list the possible cases below.

- (i) If $i = n_0 + 1, \dots, n$ then, as in first moment calculation, the last weird jump of z_i sends $deg(z_i)$ back to zero, so we must have that $E_M(t-) - q(t-) = 0$ at the occurring time of the jump.
- (ii) Similarly, if $i = n_0$, $deg(z_{n_0})$ is sent back to 0 by the walk by the last weird jump in z_{n_0} .

If $i = 1, \dots, n_0 - 1$, then there are two other possible cases.

- (iii) If there is at least one weird jump of z_i after time C_i , then the last such weird jump sends $deg(z_i)$ back to zero, with $E_M(t-) - q(t-) = 0$ at that time.
- (iv) If there is no weird jump of z_i after time C_i then $deg(z_i)$ must be zero at time C_i , thus we must have $E_M(C_i-) - q(C_i-) = 0$.

We now consider the scaling of the action. We first choose n_0 variables from z_1, \dots, z_N which we may assume are the variables z_1, \dots, z_{n_0} appearing in increasing order. We then choose $n - n_0$ variables from z_{n_0+1}, \dots, z_N which we may assume are the variables z_{n_0+1}, \dots, z_n , **whose ‘‘first jumps’’ (which by definition must exist) appear in increasing order**. We record the number of weird jumps of each z_i as a vector \vec{k} .

Definition 7.4. Let \vec{k} be a integer-valued vector of length $n_0 \cdot n$, that records the number of weird jump of each z_j ($j = 1, 2, \dots, n$) between each C_{i-1} and C_i ($i = 1, 2, \dots, n_0$). That is, $k_{i,j}$, $i = 1, 2, \dots, n_0$, $j = 1, \dots, n$ is the number of weird jumps in variable z_j between C_{i-1} and C_i . We call \vec{k} the jump data of a walk/block configuration.

In particular, we require that for each \vec{k} , each $i = 1, 2, \dots, n_0$, $n_0 + 1 \leq j_1 < j_2 \leq n$, $k_{1,j_1} + \dots + k_{i,j_1} = 0$ implies $k_{1,j_2} + \dots + k_{i,j_2} = 0$. In other words, the jump data should not contradict the restriction of the first jumps

$$s_1(n_0 + 1) < \dots < s_1(n). \quad (132)$$

Then for each fixed $n_0, n, (l_1, \dots, l_{n_0})$, and \vec{k} , number of ∂ 's p , the action of Eqn. (123) on $G(z_1, \dots, z_N; \theta)$ gives an expression of the form

$$\binom{N}{n_0} \binom{l}{l_1, \dots, l_{n_0}} \binom{N - n_0}{1!, \dots, 1!, N - n} \left(\frac{1}{N} \right)^{|\vec{k}|+p} \left(\frac{N}{\mu_+(N)^{lM}} m_{lM}(N) \right) \frac{1}{N} \quad (133)$$

. $\frac{\text{sum over occurring time of } \partial\text{'s, walks and blocks with } n \text{ nontrivial variables}}{m_{lM}(N)}$.

The scaling of the sums of terms corresponding to walk/block configurations with only situation (i)-(iii) behaves like in the first moment calculation. However, in case (iv) subtleties arise at the splitting times C_1, \dots, C_{n_0-1} . It will be useful for us to consider the local behavior of the walk/block configuration near such C_i . See Figure 3. On the left of C_i , there are also three sub-cases:

1. $q_{i+1}(C_i) > 0$, which means that $deg(z_{i+1})$ is nonzero at the splitting time C_i .
2. $q_{i+1}(C_i) = 0$, which means that $deg(z_{i+1}) = 0$ at the splitting time C_i , and there's no jump of z_{i+1} before time C_i .
3. $q_{i+1}(C_i) = 0$, and there are at least one jumps of z_{i+1} before time C_i . In particular the last of such jump must send $deg(z_{i+1})$ back to 0.




Figure 3: Here we give a schematic for the example when $n_0 = 2$. In the top figure, the blue block corresponding to q_2 is nonzero at time C_1 , so at the splitting time the degree of z_2 is nonzero. In the bottom two figures, $q_2(C_1) = 0$ and the degree of z_2 is zero at the splitting time. Note that we could have the immediate creation of a block q_1 at the splitting time (bottom-left) or the total degree can be zero at the splitting time (bottom-right).

We will see that the sum of the terms in case (iv.1) or (iv.2) blows up, because of the local behavior on the left of time C_i in their walk/block configurations. To handle this, we will show that these divergent cases cancel by developing a coupling between terms in (iv.1) and (iv.2) with opposite signs. See Examples 7.5 and 7.9. After these cancellations the sum of terms in case (iv) will also be of order $O(1)$. In Section 7.3, below, we give the precise details of this cancellation. The sum of terms in the third case will have order $o(1)$ and do not need a coupling (see Lemma 7.15).

7.3 Blowing up terms and cancellations

Let τ_i be a stopping time such that

$$\tau_i = \begin{cases} \min_{C_{i+1} \leq t \leq C_{i+1}} \{t : E_M(t) = E_M(C_i) - 1\}, & \text{if such } t \text{ exists;} \\ \infty, & \text{otherwise.} \end{cases} \quad (134)$$

Consider the case at some C_i ($i = 1, 2, \dots, n_0 - 1$) that case (iv) occurs. Let $\bar{\tau}_i$ be the maximal time before C_i that $E_M(t)$ is less than $E_M(C_i)$. Because we are in case (iv), we must have $E_M(C_i) = q(C_i -)$. In order for $q(\bar{\tau}_i) \leq E_M(\bar{\tau}_i)$, there must be at least one weird jump between times $\bar{\tau}_i$ and C_i .

There are at most $C_i - \bar{\tau}_i$ choices of occurring time of this weird and note that the height of this jump is fixed by the data of all its previous weird jumps and the condition that $E_M(C_i) = q(C_i)$. Therefore, the scaling factor contributed by the local behavior of walk/block configuration at time $[\bar{\tau}_i, C_i]$ is bounded by

$$\mathbb{E}[C_i - \bar{\tau}_i] = \sum_{t \geq 1} 1 \cdot t \cdot \mathbb{P}[C_i - \bar{\tau}_i = t]. \quad (135)$$

Intuitively, since $E_M(t)$ behaves like Brownian motion for M large, $C_i - \bar{\tau}_i$ is small with high probability. More precisely, by Lemma 7.3 (from its proof one can see that the convergence holds uniformly for L large enough), we have

$$\sum_{t=1}^{\epsilon M} t \cdot \mathbb{P}[C_i - \bar{\tau}_i = t] = \sum_{t=0}^{\epsilon M} \mathbb{P}[C_i - \bar{\tau}_i \geq t] \sim \sum_{t \leq \epsilon M} \frac{1}{\sqrt{t}} \sim \sqrt{\epsilon M}.$$


In particular, the main contribution to the sum is from the configurations that $C_i - \bar{\tau}_i \sim O(M)$. Note that it is enough to consider just one weird jump in $[\bar{\tau}_i, C_i]$ as additional weird jumps at worst contribute an $O(1)$ term to the scaling factor.

It will be more clear why contributions from such configurations give a blowing-up partial sum by looking at the following example.

Example 7.5. Let us consider the case when $n_0 = 2$. We are interested in the actions of

$$\frac{N(N-1)}{\mu_+(N)^{2M}} \left(\frac{\mathcal{D}_2}{N}\right)^M \left(\frac{\mathcal{D}_1}{N}\right)^M$$

on the BGF. First, let's consider the case in which there are no weird jumps. In this case, corresponding walk must hit zero at the M -th step as shown in the example below with $M = 14$.




The partition function for such walks is given by $m_M(N)^2$ and we have

$$\frac{N(N-1)}{\mu_+(N)^{2M}} m_M(N)^2 = O(1)$$

by Proposition 4.12.

Now consider the case in which we have exactly 1 weird jump. There are two possibilities. We could have that the weird jump adds powers of z_2 in the section where z_1 is the main variable, in which case the height of the z_2 block must be equal to the height of the walk at step M . Or we could have a weird jump removing left-over powers of z_1 in the section where z_2 is the main variable. Examples of the two possibilities are given below with the block drawn dashed in blue.



One can check that the corresponding functional on walks of length $2M$ has

$$\frac{N(N-1)m_{2M}(N)}{\mu_+^{2M}(N)} \frac{1}{N} (I_{(7,7)}((0, 0, 1, 0), \vec{0}; M) + I_{(7,7)}((0, 1, 0, 0), \vec{0}; M)) = O(\sqrt{M})$$

due to contributions from the first case (the second case contributes $O(1)$ since the occurring time and height of the jump have no freedom given the walk).

To resolve this issue, let X_i denote the occurring time of the closest weird jump on the right of C_i and let $Y_i = \min(X_i, C_{i+1})$.

Locally on the right of C_i , we further classify the following possible situations.

- (a) If $Y_i > \tau_i$, or if $Y_i = \tau_i$ and the weird jump at time τ_i sends $\text{deg}(z_j)$ to 0 for some $j \neq i, i + 1$.
- (b) If $Y_i < \tau_i$, or if $Y_i = \tau_i$ and the weird jump at time τ_i does not send its corresponding $\text{deg}(z_j)$ back to 0.

Remark 7.6. *In case (b), if there is one weird jump of some z_j between time C_i and τ_i , then that jump doesn't send $\text{deg}(z_j)$ back to 0 no matter whether it's occurring at time τ_i or before. The addition of a further jump needed to send $\text{deg}(z_j)$ back to 0 makes the terms vanish in the scaling limit. Only case (a) have terms that are problematic in the scaling limit.*

To deal with case (a), we make a coupling between those configurations in case (a1) with those that fall into (a2). For (E_M, \vec{q}) that fall in case (a1) on the right of some C_i , fix one of such i , and consider all (E'_M, \vec{q}') such that $E'_M = E_M$, and \vec{q} and \vec{q}' have the same jump times, heights and jumping variables, except that before time C_i , we replace all the jumps of z_{i+1} by jumps of z_j (given by Δd_j), $j \in \{n + 1, \dots, N\}$, with the same time and heights as before. In addition, we add one more jump from Δd_j in (E'_M, \vec{q}') at time τ_i , the earliest time after C_i that a jump can send $\text{deg}(z_j)$ back to 0. Then the resulting (E'_M, \vec{q}') falls in case (2a).

Remark 7.7. *There are some terms in case (2a) that are not coupled: the first jump after C_i at τ_i is a jump of z_1, \dots, z_{n_0} . We don't worry about these terms since their total contribution is of order $O(1)$, see Section 7.4.*


Fixing (E_M, \vec{q}) and (E'_M, \vec{q}') , we couple the terms from these two configurations with same number of partial derivatives in the corresponding actions. If there's no ∂ , then the one extra jump that sends $\text{deg}(z_j)$ back to 0 gives an extra factor $-\frac{1}{N}$ while we have $N - n$ choices of variable z_j , so the sum of the coupled terms is equal to the (single) term from (E_M, \vec{q}) multiplied by a factor

$$1 - \frac{N - n}{N} \sim O\left(\frac{1}{N}\right) \tag{136}$$

for each fixed n . The discussion around Eqn. (135) shows this vanishes in the limit, since the scaling factor given by the sum of terms of case (iv) behavior locally on the left of time C_i is of order at most $O(\sqrt{M})$. Hence for simplicity, as $n \leq N^{\frac{1}{4}} \ll \sqrt{M}$, we can ignore the effect given by $1 - \frac{N-n}{N}$ below.

Remark 7.8. *When $n \geq N^{\frac{1}{4}}$, in Section 7.6 we do the tail estimation of the contribution of those walk/block configurations, and show that it is decaying in the speed of $C^{-n} \lesssim C^{-N^{\frac{1}{4}}}$ for some $C > 1$ uniformly in N . Hence the error of the coupling is again negligible.*

Example 7.9. *The blowing-up terms coming from configurations one weird jump will cancel with certain configurations with two weird jumps. The configuration shown in Example 7.5 above will cancel with*



where the red dashed line indicates the block for some z_i , $i \neq 1, 2$. Note to the left of C_1 we have kept the same weird jump location and height (but have changed the index) and we have added a new weird jump to the right of C_1 at the unique location that can send the block to zero.

Introduce a parameter $\epsilon \in (0, 1]$, and for the following discussion we assume that $\tau_i - C_i \leq \epsilon M$. If there is one ∂ , we sum over the choices of time $1 \leq s \leq lM$ that ∂ can take place. Then since, up to some constant, for a fixed walk sum over the possible locations of ∂ gives the area between the block process and the walk, and since the block configuration from (E'_M, \vec{q}') has one more rectangular block with height at most $E'_M(C_i)$ and width $\tau_i - C_i$ after the cancellation of the coupled terms, we have an error bounded by $(\tau_i - C_i)E'_M(C_i)$.

We know from Section 6.3 that $E_M(C_i)$ is of order \sqrt{M} with exponentially decaying tail probability, and by Lemma 7.2, $\mathbb{E}[\tau_i \mathbf{1}_{1 \leq \tau_i \leq \epsilon M}] \lesssim \sqrt{\epsilon M}$. Again the scaling factor given by the sum of terms of case (iv) behavior locally on the left of time C_i is of order at most $O(\sqrt{M})$ due to the discussion around Eqn. (135). Also recall that the operator ∂ comes with a factor $\frac{1}{N}$, hence the error after cancellation bounded up to some uniform constant by

$$\frac{\sqrt{\epsilon M} \sqrt{M} \sqrt{M}}{N} \lesssim \sqrt{\epsilon}. \quad (137)$$

For general $p \geq 1$, let A denote the area between walk and blocks of a given walk/block configuration in case (1), say on interval $[[C_i, C_{i+1}]]$, and let \square denote the area of the added block in the coupled configuration, then up to negligible errors and neglecting any overall minus signs, the sum of the coupled terms can be written as

$$\sum_{p=1,2,\dots} \frac{1}{p!} [A^p - (A - \square)^p] \approx \exp(A) - \exp(A - \square), \quad (138)$$

when M is large. Again by Section 6.3, generically each $A \lesssim M\sqrt{M}$ and $\frac{\square}{A} \lesssim \epsilon$. Hence the expectation of the sum over local behavior on the left of time C_i of Eqn. (137) is still bounded by $O(\sqrt{\epsilon})$. Therefore we get the following result.

Lemma 7.10. *Assume that $\tau_i - C_i \leq \epsilon M$ for some $\epsilon \in (0, 1]$. For each C_i , the sum over terms corresponding to case (a) behavior locally on the left of time C_i and at time $[C_i + 1, \tau_i]$, is of order bounded by $O(\sqrt{\epsilon})$.*

In general, for a walk/block configuration there can be $\tilde{n} \in \{0, 1, 2, \dots, n_0 - 1\}$ splitting points, C_i , with case (a) behavior, denoted as $C_{j_1}, \dots, C_{j_{\tilde{n}}}$ where $j_1 < j_2 < \dots < j_{\tilde{n}}$. We start from C_{j_1} and couple as above, then identify the configuration before time C_{j_1} of the coupled two terms, then go to time C_{j_2} , and couple the two types of configuration with behavior (1a) and (2a) near C_{j_2} respectively, and iterate until the $2^{\tilde{n}}$ terms are combined as a whole. One can check by definition that such iteration is well-defined, and since in case (a) we have $\tau_i - C_i \leq M$, by repeatedly using Lemma 7.10, the total contribution of terms in case (a) is bounded uniformly in N .

Remark 7.11. *In this group with $2^{\tilde{n}}$ elements, the number of nontrivial variables n differ by at most $\tilde{n} \leq n_0 - 1$, which is a constant uniformly bounded in N .*

Now we consider case (b), which has similar scaling behavior as case (a). Note that in this case, $0 \leq \tau_i - C_i \leq lM$.

Lemma 7.12. *Assume that $\tau_i - C_i \leq \epsilon M$ for some $\epsilon \in (0, 1]$. For each C_i , the sum over terms corresponding to case (b) behavior locally on the left of time C_i and in the time interval $[C_i, C_i + \epsilon M]$, is of order bounded by $O(\sqrt{\epsilon})$.*

Proof. Since there are m weird jump at time $t \in [C_i + 1, \tau_i]$, ($m \geq 1$), the number of choices of such jumps is bounded by τ_i^m . Each weird jump introduces a factor $\frac{1}{N}$, and with high probability has $\lesssim O(\sqrt{M})$ choices of the height. Besides this, the walk/block configuration behave in the same way on the right of τ_i . Recall again that the sum of terms in the action can be thought as expectation of some functional (characterized by drawing blocks and taking area between walks and block processes) of the random walk bridge. Hence the sum over the terms with local behavior on the right of time C_i of case (b) is bounded by

$$\sum_{t=1}^{cM} \frac{t^m \sqrt{M}^m}{N^m} \cdot \mathbb{P}[\tau_i = t] \lesssim \sqrt{\epsilon} \frac{1}{\sqrt{M}}$$

by Lemma 7.2. Locally on the left of time C_i , again the sum gives a scaling factor \sqrt{M} . By the Markov property of the walk we can multiply these two scaling, which gives $O(\sqrt{\epsilon})$ bound as in case (a). \square

By the discussions above, we see that in case (a) or (b), the sum over terms with $\tau_i - C_i \leq \epsilon M$ for $c \leq 1$ gives at most order $\sqrt{\epsilon}$ contribution. We will show in Section 7.4 that the case (b) when $\tau_i > \epsilon M$ gives contribution of order $O(1)$. This together with the fact that away from splitting point C_i 's, the sum is always of order $O(1)$, shows that the total sum of terms with fixed number of weird jumps (up to cancellations in case (a)) is of order $O(1)$, and sum of terms with $\tau_i - C_i \leq \epsilon M$ vanish when taking $\epsilon \rightarrow 0$.

7.4 Convergence of the functional

We give the functional expression of the higher moment action in this section. Because of the case (iv) behavior for $l > 1$ discussed in Section 7.3, that gives $O(1)$ contribution (after cancellations) with only walks of vanishing weight in the limit, we need a slightly more complicated presentation compared to Section 5.1.

Recall the notion of \vec{k} from Definition 7.4. For a given vector \vec{k} , which gives the number of weird jumps of z_1, \dots, z_n in each domain $[C_{i-1}, C_i]$, ($i = 1, 2, \dots, n_0$), as in Section 5.2, we first determine the scaling of the total contribution of terms given by such \vec{k} .

Definition 7.13. For $i = 1, 2, \dots, n$, we call the jump of z_i with the latest occurring time $s_{k_i}(i)$ the last jump of z_i , if

- $i = 1, 2, \dots, n_0$, $k_i \geq 1$, and this jump sends $\deg(z_i)$ back to 0, i.e. $q_i(s_{k_i}(i)-) > 0$, $q_i(s_{k_i}(i)) = 0$; or
- $i = n_0 + 1, \dots, n$.

Note that for $i = 1, 2, \dots, n_0$, the last jump might not exist.

Definition 7.14. Let K be a set of vector \vec{k} , such that $\vec{k} \in K$ if there exists any orientation of the order of weird jumps so that the following holds: for each $j = 1, 2, \dots, n_0$ where the last weird jump of z_j is after time C_j , we have that the last weird jump of z_j is the first weird jump after some C_i , ($j \leq i \leq n_0 - 1$), and we force that there is no jump of z_{i+1} before C_i ¹. In particular, we include the case with no weird jumps in K .

Lemma 7.15. For $\vec{k} \in K$, up to coupling in Section 7.3, the sum of terms given by \vec{k} are of order $O(1)$, and the total contribution of terms given by $\vec{k} \notin K$ vanish as $N, M \rightarrow \infty$.

Proof. We first note the non-vanishing contribution is given by the ‘‘canonical’’ walk-block configurations: configurations such that there is at most one jump of each z_i that sends its degree back to 0, and therefore before such jump, generically $\deg(z_i) > 0$.

Consider the scaling factor given by z_{n_0+1}, \dots, z_n : each time introducing a new variable into the blocks gives a factor N . As in the first moment case, the last jump is ‘‘fixed’’ by the earlier jumps and the walk, and all other weird jumps have freedom of order M and \sqrt{M} horizontally and vertically. Each jump has a factor $\frac{1}{N}$. These together gives order $O(1)$.

Now consider $i = 1, 2, \dots, n_0$, where we recall that there are $O(N^{n_0})$ choices of these ‘‘main variables’’. Assume that there are no weird jumps of the z_i ’s, then the walk is forced to return to $q(C_i-)$ at time C_i . This gives factor $O(\frac{1}{\sqrt{M}})$ at each C_1, \dots, C_{n_0-1} . By Lemma 7.2, generically $E_M(t) \geq q(C_i-)$ for $[C_i - \epsilon M, C_i]$, which by Lemma 7.3 gives factor $O(\frac{1}{\sqrt{M}})$ at each C_1, \dots, C_{n_0-1} . Locally on the right of C_i ’s, if $\tau_i \leq \epsilon M$, the walk is ‘‘free’’ in the sense that it no longer needs to stay above $E_M(C_i)$ for long enough time. There must be a jump at τ_i , which is generically the last jump of z_j ($j \geq n_0 + 1$), so we are in case (2a) and do the coupling to resolve the blow up. If $\tau_i > \epsilon M$, by Eqn (131) this gives a factor $O(\frac{1}{\sqrt{M}})$. So the total order locally near each C_i is $O(\frac{N}{M^{\frac{3}{2}}}) = O(1)$.

Now suppose there are weird jumps in the z_j , $j = 2, \dots, n_0$. Fix one of such j ’s. If there are jumps of z_j before C_{j-1} , the walk is ‘‘free’’ locally on the right of C_{j-1} as above. By Eqn. (131) this give factor $O(\sqrt{M})$. On the right of C_{j-1} , if $\tau_{j-1} \leq \epsilon M$, which is the generic case, we are in case (1a) and do the coupling in Section 7.3 to resolve the blow up. If $\tau_{j-1} > \epsilon M$, then the walk is still under the restriction to stay above $E(C_{j-1})$ locally on the right of C_{j-1} as if there’s no jump of z_j before C_{j-1} .

If there are jumps of z_j after C_j , we have one ‘‘fixed jump’’ that gives factor $O(\frac{1}{N})$, the walk locally on the left of C_j is ‘‘free’’ in its height and no longer staying above $E_M(C_j)$, which gives factor $O(\sqrt{M}^2)$. Therefore, in order for the sum over such terms to be not vanishing, the last jump of z_j needs to ‘‘release’’ the walk locally on the right of some C_i ($i \geq j$), i.e. it needs to be the first jump after C_i where $\deg(z_{i+1})$ is forced to be 0 at time C_i , as in Figure 4. For $\vec{k} \in K$ this is indeed the case, otherwise the contribution of the terms vanish. \square

In the remaining of this section we consider only $\vec{k} \in K$. In order for taking the limit, we update the interpretation of ‘‘fixed’’ jumps, and consider the block process \vec{q} as a collection of free variables $\{(s_m(i), h_m(i))\}_{i=1}^n$ that deforms ‘‘continuously’’, as follows.

¹It follows, that $\deg(z_{i+1}) = 0$ at time C_i




Figure 4: In the above example, we see that at time C_i the path hits the top of the block, corresponding to the variable z_i , so $\deg(z_{i+1})$ must be zero. A short time after C_i the block is forced to zero, and the path is “released”.

Definition 7.16. Given the block process \vec{q} , for $j = 1, 2, \dots, n$, we view $(s_{k_j}(j), h_{k_j}(j))$, the last jump of z_j , as **fixed** by the previous jumps of z_j and the walk, if

- (i). $j = 1, 2, \dots, n_0$ and $s_{k_j}(j) > C_j$;
- (ii). $j = n_0 + 1, \dots, n$.

More precisely, we must have that after the last jump, the block is at height zero, so $h_{k_j}(j) = -\sum_{m=1}^{k_j-1} h_m(j)$, and $s_{k_j}(j)$ is at the earliest time after $s_{k_{j-1}}(j)$ where $E_M(t)$ jumps from $E_M(s_{k_{j-1}}(j))$ to $E_M(s_{k_{j-1}}(j)) - 1$.

Furthermore, we say for $j = 1, 2, \dots, n_0 - 1$, its last jump is **strongly fixed**, if² it is fixed and it is the first weird jump after some C_i ($j \leq i \leq n_0 - 1$), where $\deg(z_{i+1})$ is forced to be 0 at time C_i . A fixed jump that is not strongly fixed is **weakly fixed**.

Examples of such fixed jumps are given in Figure 5.

Remark 7.17. Intuitively, a weakly fixed jump only has freedom in its occurring time, while a strongly fixed jump is completely determined by the walk/block configuration.




Figure 5: Examples of fixed jumps of the block process as in Definition 7.16. In the top figure, the block corresponds to variable z_j with $j \in \{1, 2, \dots, n_0\}$ and its last jump is occurring as the first weird jump after the time C_j , so strongly fixed. In the bottom figure the block belongs to a variable z_i with $i \in \{n_0 + 1, \dots, n\}$.

For a fixed \vec{q} , according to the discussion in the proof of Lemma 7.15, we now determine the appropriate events conditioning the walk $E_M(t)$ depends on \vec{q} such that $(E_M(t), \vec{q}(t))$ is a valid walk/block pair. Given \vec{q} , we first **deform** it according to the following rule, abuse the notation by still denoting the deformed block process by \vec{q} :

- For $j = 1, 2, \dots, n_0 - 1$, the last jump of z_j is strongly fixed and occurs at time $\tilde{\tau}_i \in \llbracket C_i, C_{i+1} \rrbracket$ for some $i = j, j + 1, \dots, n_0$, then let $q_j(t) = 0$ for $t \geq C_i$ ³.

Then the conditioning of $E_M(t)$ must obey the following rules, given the **deformed block process** \vec{q} :

²Then by definition, if $\vec{k} \in K$, all the last jumps of z_1, \dots, z_{n_0-1} are strongly fixed, and the last jump of z_{n_0} does not exist.

³Note that $q_j(C_i-)$ might still be nonzero.

- $E_M(t) \geq q(t) \geq 0$ for $0 \leq t \leq lM$.
- For each $i = 1, 2, \dots, n_0 - 1$, if there is no jump of z_i after time C_i , let $X_{i,1}$ denote the occurring time of the closest weird jump on the left of (i.e. \leq) C_i , and take $X_{i,1} = \emptyset$ when it doesn't exist. Let $Y'_{i,1} = \max(X_{i,1}, C_{i-1})$, we condition on the event that $E_M(C_i) = q(C_i-)$, $E_M(t) \geq q(C_i-)$ for $t \in [Y'_{i,1}, C_i)$. If there is one more jump of z_i after time C_i (which is strongly fixed and is removed from the block process in the above deformation), we make no extra conditioning. If there are two or more jumps of z_i after time C_i , we condition on the event that $E_M(C_i) = q(C_i-) + q_i(C_i)$.
- On the right of time C_i , let $X_{i,2}$ denote the occurring time of the closest weird jump on the right of (i.e. $>$) C_i , and take $X_{i,2} = \emptyset$ when it doesn't exist. If we are in case (2), let $Y'_{i,2} = \min(X_{i,2}, C_{i+1})$, and condition on the event that $\tau_i = Y'_{i,2}$ when we are in case (2a) and require coupling,⁴ or on the event that $\tau_i \geq Y'_{i,2}$ when we are in case (2b)⁵.
- On the right of time C_i , if we are in case (1), let $Y'_{i,2} \in \llbracket C_i + 1, \min(X_{i,2}, C_{i+1}) \rrbracket$. We add an extra “virtual block” with height $q_{i+1}(C_i)$ extending from time $t = C_i$ to $Y'_{i,2}$. We condition on the event that $\tau_i = Y'_{i,2}$ when $Y'_{i,2} \leq \min(X_{i,2}, C_{i+1}) - 1$, on the event that $\tau_i \geq \min(X_{i,2}, C_{i+1})$ when $Y'_{i,2} = \min(X_{i,2}, C_{i+1})$.
- At all remaining weakly fixed jump $s_{k_j}(j)$, condition on the event that $E_M(s_{k_j}(j)-) = q(s_{k_j}(j)-)$, $E_M(t) \geq q(t)$ between $s_{k_j}(j)$, and the rightmost weird jump/the splitting point on its left.

See Figure 6.

Definition 7.18. For a given block process \vec{q} , denote the conditional process by $E'_M(t)$, such that $E'_M(t) = E'_M(\lfloor t \rfloor)$ as before, and denote the updated block process possibly with extra virtual blocks by \vec{q}' .

Definition 7.19. Fix $M, n_0, l_1, \dots, l_{n_0}$. Fix a parameter $\epsilon > 0$. For a given \vec{q}' , we say \vec{q}' is in the ϵ -domain, if for each $i = 1, 2, \dots, n_0 - 1$, $Y'_{i,2} \geq \epsilon M$.

Remark 7.20. The above detailed classification of conditioning of the event $\{\tau_i > \epsilon M\}$ is designed for the limiting expression $I_{(l_1, \dots, l_{n_0})}(\vec{k}, p)$ when taking $\epsilon \rightarrow 0$, see Theorem 7.23, 7.27. In this event the “virtual block” always exists.




Figure 6: **Left:** If there are no weird jumps of z_i after C_i then we condition the walk on the event that $E_M(C_i-) = q(C_i-)$ and $E_M(t) \geq q(C_i)$ for $t \in [Y'_{i,1}, C_i)$. **Center:** If we are in case (2) in which $\deg(z_{i+1}) = 0$ at time C_i , then we condition the walk on the event $E_M(C_i) = q(C_i)$. Further, if we are in case (2a) then we condition on the event that $Y'_{i,2} = \tau_i$. Note that after time $Y'_{i,2}$ there can be more blocks. **Right:** If we are in case (1a) we add a virtual block as shown by the dashed block in the figure.

Note that $E'_M(t)$ is piecewise independent split by the conditioning times, that deforms “continuously” with \vec{q}' , and each \vec{q}' might correspond to more than one \vec{q}' 's.

Example 7.21. When $\vec{q} = 0$, i.e. there's no weird jump, then $\vec{q}' = 0$, and $E'_M(t)$ is a combination of n_0 independent random walks starting and ending at 0 and staying nonnegative from time C_{j-1} to C_j .

⁴ $\deg(z_j)$ is sent to 0 at $Y'_{i,2}$ for some $j = n_0 + 1, \dots, N$

⁵ No extra conditioning if we are in case (2a) but the jump at $Y'_{i,2}$ is strongly fixed.

Recall the Eqn. (133) deals with the scaling of our higher moment action. After the above conditioning, we upgrade it to be the following:

$$\binom{N}{n_0} \binom{l}{l_1, \dots, l_{n_0}} \binom{N - n_0}{1!, \dots, 1!, N - n} \left(\frac{1}{N} \right)^{|k|+p} \left(\frac{\text{partition function of the conditional walk}}{\mu_+^{lM}} \right) \cdot \frac{\text{sum over occurring time of } \partial\text{'s, walks and blocks with } n \text{ nontrivial variables}}{\text{partition function of the conditional walk}} \quad (139)$$

After all the preparations, by omitting the coupled terms in case (a) and (b) that will vanish when $\epsilon \rightarrow 0$ as discussed in Lemma 7.10, 7.12, we give the functional expression under action of (123) that converges in the limit. It is nothing but a rephrase and slight generalization of the functional in Section 7.2.

We now split the sum of terms in the Dunkl action according to \vec{k}, \vec{p} .

Proposition 7.22. *Fix $\epsilon > 0$ small enough, n_0, n and (l_1, \dots, l_{n_0}) . The action*

$$\binom{N}{n_0} \binom{l}{l_1, \dots, l_{n_0}} \frac{\left(\frac{D_1}{N} \right)^{l_1 M} \dots \left(\frac{D_{n_0}}{N} \right)^{l_{n_0} M}}{\mu_+(N)^{lM}}$$

on $G(z_1, \dots, z_N; \theta)$ is a sum of $C(N, n_0, \vec{k}) \cdot I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M)$ over the parameters \vec{k}, \vec{p} , such that

$$C(N, n_0, \vec{k}) = \binom{N}{n_0} \binom{l}{l_1, \dots, l_{n_0}} \binom{N - n_0}{1!, \dots, 1!, N - n} N^{-n} \approx \binom{l}{l_1, \dots, l_{n_0}} \frac{1}{n_0!} \quad (140)$$

as N is large. Moreover, one can replace each $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M)$ by $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M, \epsilon)$ up to error of order $O(\sqrt{\epsilon})$, where

$$I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M, \epsilon) = \sum_{\vec{q}'} F_{(l_1, \dots, l_{n_0})}(\vec{q}', \vec{p}; M) \cdot \frac{N^{n-|k|-p} \cdot \text{Par}(E_M^l(\vec{q}'))}{\mu_+(N)^{lM}}, \quad (141)$$

$\text{Par}(E_M^l(\vec{q}'))$ is the partition function of conditional walk $E_M^l(\vec{q}')$ in Eqn. (139), \vec{q}' is summed over unfixed and weakly fixed jumps of all possible updated block processes in the ϵ -domain, under the restrictions of \vec{k} , and in particular the constraint that

$$s_1(n_0 + 1) < \dots < s_1(n); \quad (142)$$

and

$$\begin{aligned} F_{(l_1, \dots, l_{n_0})}(\vec{q}', \vec{p}; M) &= \sum_f \mathbf{1}_{f \in A_M(\vec{q}')} \mathbf{1}_{f \in B_M(\vec{q}')} \left(\prod_{i=1}^n \prod_{m=1}^{k_i} \text{sgn}(h_m(i)) \right) \\ &\quad \sum_{v_1 < \dots < v_p \in D(f)} \theta^{-p} \prod_{j=1}^p [f(v_j) - q(v_j)] \cdot \text{Prob}(f; E_M^l(\vec{q}')) \\ &= \mathbb{E}_{\vec{q}'} \left[\mathbf{1}_{E_M^l \in A_M(\vec{q}')} \mathbf{1}_{E_M^l \in B_M(\vec{q}')} \left(\prod_{i=1}^n \prod_{m=1}^{k_i} \text{sgn}(h_m(i)) \right) \right. \\ &\quad \left. \prod_{i=1}^{n_0} \left[\sum_{C_{i-1} < v_1 < \dots < v_{p_i} \leq C_i \in D(E_M^l)} \theta^{-p_i} \prod_{j=1}^{p_i} [E_M^l(v_j) - q(v_j)] \right] \right], \end{aligned} \quad (143)$$

where $q(s) = \sum_{j \neq i} q_j(s)$, and z_i is the main variable at time s . The expectation is taken on E_M^l that depends on \vec{q}' , and $D(E_M^l)$ denotes the set of all down steps in E_M^l .

Now we pass from discrete to continuous. For $l_1, \dots, l_{n_0} \in \mathbb{Z}_{>0}$, $l = l_1 + \dots + l_{n_0}$, let \vec{y} be a block process on $[0, l]$, with splitting points $l_1, \dots, l_1 + \dots + l_i$ ($i = 1, 2, \dots, n_0 - 1$), defined in almost the same way as the \vec{q} . As the case for \vec{q} , \vec{y} is a Cádłág function, whose discontinuity points are given by the jump times and the splitting points (in particular $y_i(t) = 0$ on $[l_1 + \dots + l_{i-1}, l_1 + \dots + l_i]$ for $i = 1, 2, \dots, n_0$). Denote the jumps of \vec{y} by

$$\{(S_m(i), H_m(i))\}_{m=1,2,\dots,k_i, i=1,2,\dots,n},$$

$S_m(i) \in [0, l] \setminus [l_1 + \dots + l_{i-1}, l_1 + \dots + l_i]$, $H_m(i) \in \mathbb{R} \setminus \{0\}$. Let $y(s) = \sum_{j \neq i} y_j(s)$. For a given \vec{y} , we define a set $A(\vec{y})$ for $f \in \mathbb{D}([0, l])$ such that $f \in A(\vec{y})$ if all the followings hold:

- $f(t) \geq y(t)$ for $t \in [0, l]$.
- For the jump $(S_m(i), H_m(i))$,

$$|H_m(i)| \leq |f(t-) - y(t-) - y_i(t-) - (\text{sgn}(H_m(i)) + 1)|.$$

- For the jump $(S_m(i), H_m(i))$,

$$H_m(i) \cdot (f(t-) - y(t-) - y_i(t-)) > 0$$

Denote the deformed and updated block process of \vec{y} with possible extra virtual blocks by \vec{y}' . \vec{y}' and the notion of ϵ -domain for \vec{y}' are defined similar as \vec{q}' , except that we replace C_i by $l_1 + \dots + l_i$ and ϵM by ϵ .

Theorem 7.23. For each fixed $\vec{k} \in K, \vec{p}$, when $M \rightarrow \infty$,

$$C(N, n_0, \vec{k}) I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M, \epsilon) \rightarrow \binom{l}{l_1, \dots, l_{n_0}} \frac{1}{n_0!} I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; \epsilon) \quad (144)$$

where

$$I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; \epsilon) = \int d\vec{y}' g(\vec{y}') T^{\frac{3}{2}(|\vec{k}| - n + p)} \mathbb{E} \left[\mathbf{1}_{\mathfrak{E}'_{\vec{y}'}} \in A(\vec{y}') (P_{-1}\sigma)^{|\vec{k}| - n + p} \left(\prod_{i=1}^n \prod_{m=1}^{k_i} \text{sgn}(H_m(i)) \right) \left(\frac{1}{\theta^p} \prod_{i=1}^{n_0} \frac{1}{p_i!} \left[\int_{l_{i-1}}^{l_i} \mathfrak{E}'_{\vec{y}'}(s) - y(s) ds \right]^{p_i} \right) \right], \quad (145)$$

\vec{y}' is the updated block process, that record the time and height of all (not strongly fixed) weird jumps, and is integrated in the ϵ -domain, under the restrictions given by \vec{k} , in particular the constraint

$$S_1(n_0 + 1) < \dots < S_1(n); \quad (146)$$

and $g(\vec{y}')$ is a continuous density function only depending on \vec{y}' and l_1, \dots, l_{n_0} , and $\mathfrak{E}'_{\vec{y}'}$ is a piecewise independent conditional Brownian bridge, whose conditioning depends on \vec{y}' .

Remark 7.24. For each given $\vec{k} \in K$, if there is any strongly fixed jump of z_1, \dots, z_{n_0-1} right after some C_i , then the limiting occurring time of its corresponding jump in $\vec{y}(s)$ is fixed to be at $l_1 + \dots + l_i$, as described by the deformation of the block process.

Remark 7.25. One can see that the limiting expression $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; \epsilon)$ is universal for all ensembles we consider in this text, by the same reasoning as in Section 5.4. Moreover, although the domains of \vec{y}' and $\mathfrak{E}'_{\vec{y}'}$ have certain restrictions parameterized by ϵ , we prefer to leave this implicit since such effects vanish when $\epsilon \rightarrow 0$, see Section 7.5.

Remark 7.26. The density function $g(\vec{y}')$ can be written out explicitly, since it gives the distribution of a Brownian bridge, with explicit conditioning piecewisely. We choose to leave it as the current form for brevity, and for the reason that the key point of our result is the convergence and the universality of the limit rather than its specific expression. The interested readers can check [GXZ, Section 2] for a more explicit limiting expression.

Proof. First recall that

$$C(N, n_0, \vec{k}) \rightarrow \binom{l}{l_1, \dots, l_{n_0}} \frac{1}{n_0!} \quad \text{as } N \rightarrow \infty.$$

Again by Lemma 6.4, we have a good control on the maximum of the random walk bridge, so we can truncate the maximum at height $h\sqrt{M}$ for some arbitrarily large $h > 0$.

Rescale $E_M(t)$ horizontally by M and vertically by $\sigma\sqrt{M}$, so that the walk/block locates on $[0, l]$, and the Riemann sum $\sum_{\vec{q}'} \sum_{v_1 < \dots < v_p}$ converges to the corresponding integral as in Theorem 5.9. One can split the conditional walk $E_M'(t)$ into independent pieces (by conditioning on the height at the boundaries⁶ that correspond to one of the cases in Theorem 5.8 and 6.1. Then for fixed (rescaled) \vec{q}' , the walk $\mathfrak{E}'_M(s) = \frac{1}{\sigma\sqrt{M}} E_M'(sM)$ converges in distribution to the corresponding Brownian bridge. This provides a scaling factor $\sigma^{|\vec{k}| - (n - c_0) + |\vec{p}|}$. Note that Lemma 6.2 does not require the walk to start and end at the same height, so by exactly the same argument as before, the down steps in $\mathfrak{E}'_M(s)$ always distribute homogeneously with limiting proportion P_{-1} , and $\mathbf{1}_{\mathfrak{E}'_M \in B'_M(\vec{q}')} v_1, \dots, v_{|\vec{p}|} \in D(E'_M)$ gives $P_{-1}^{|\vec{k}| - (n - c_0) + |\vec{p}|}$ in the limit. This shows

$$\left(\frac{1}{M\sqrt{M}} \right)^p F_{(l_1, \dots, l_{n_0})}(\vec{q}, p; M) \rightarrow \mathbb{E}_{\vec{y}'} \left[\mathbf{1}_{\mathfrak{E}' \in A(\vec{y}')} \mathbf{1}_{\mathfrak{E}' \in C(\vec{y}')} (P_{-1}\sigma)^p \left(\prod_{i=1}^n \prod_{m=1}^{k_i} \text{sgn}(H_m(i)) \right) \left(\prod_{i=1}^{n_0} \frac{1}{p_i!} \left[\int_{l_{i-1}}^{l_i} \mathfrak{E}'_{\vec{y}'}(s) - y(s) \right]^{p_i} \right) \right] \quad (147)$$

as $M \rightarrow \infty$, where \vec{y}' is the rescaled limit of \vec{q}' , and it's clear that the right side above is continuous in \vec{y}' .

It remains to take the asymptotic of

$$(M\sqrt{M})^{|\vec{p}|} \frac{N^{n - |\vec{k}| - |\vec{p}|} \text{Par}(E'_M(\vec{q}'))}{\mu_+(N)^{lM}}.$$

We again split $E'_M(t)$ into bounded number of independent pieces of walks conditioning on the height at the splitting points, and consider

$$\frac{\text{Par}(\cdot)}{\mu_+(N)^L},$$

where the numerator is the total weight of a piece of walk with fixed starting and ending point of length L . Multiplying the expressions above then summing over the splitting heights gives $\frac{\text{Par}(E'_M(\vec{q}'))}{\mu_+^{lM}}$.

The generic regime for each piece of walk is that $L \sim O(M)$, so in particular $L \rightarrow \infty$. Each piece of walk has one of the three configurations in Figure 2, depending on whether there is a conditioning of touching the block at the starting or ending point. Recall that $\mu_+(N) = V_N(z_c(N))$, the partition function of one step in Lemma 7.3. Therefore we make use of Eqn. (130), (131) when seeing the first two configurations, since the corresponding expression differs to $\frac{\text{Par}(\cdot)}{\mu_+^L}$ only by a factor

$$z_c(N)^{y_M - x_M}.$$

When we see the third configuration, simply view $\frac{\text{Par}(\cdot)}{\mu_+^L}$ as

$$\mathbb{P}_{x_M}(\text{the walk ends at } y_M \text{ and staying nonnegative}) \cdot z_c(N)^{x_M - y_M}.$$

Also observe that after multiplying all segments this factor disappears.

We already see that the order of $(M\sqrt{M})^{|\vec{p}|} \frac{N^{n - |\vec{k}| - |\vec{p}|} \text{Par}(E'_M(\vec{q}'))}{\mu_+(N)^{lM}}$ is $O(1)$. To see the constant factors, recall by Eqn. (130) and (131) that only P_{-1} and σ that depend on the walk increments come into play. Moreover we have the following observations.

⁶The boundaries of the independent pieces of walk is a subset of C_i 's ($i = 0, 1, 2, \dots, n_0$), $\min(X_{i,1}, C_{i-1})$'s, $\min(\tau_i, C_{i+1})$'s ($i = 1, 2, \dots, n_0 - 1$), and the last jump $s_{k_i}(i)$ of each z_i ($i = 1, 2, \dots, n$), so the total number of such walk bridges is bounded by $3n_0 + n$, which is finite for each fixed \vec{k} .

- When there is no jump, $Par(E_M^l(\vec{q}')) = m_{l_1 M}(N) \cdots m_{l_{n_0} M}(N)$, which by Eqn. (90) gives a factor $(P_{-1}\sigma)^{-n_0} = (P_{-1}\sigma)^{-n}$.
- When there's a jump of some z_i for $i = 1, 2, \dots, n_0 - 1$ after C_i , by the proof of Lemma 7.15 it must "release" the walk locally on the left of C_i , and the walk locally on the right of some C_j , in order to be of order $O(1)$. Then by Eqn. (130) and (131) we obtain one copy of $P_{-1}\sigma$.
- Each weird jump that is not a "last jump" also asymptotically gives a factor $P_{-1}\sigma$.
- $N \sim T^{-\frac{3}{2}} M \sqrt{M}$.

Therefore we conclude that for \vec{y}' restricted in the ϵ -domain, when $M \rightarrow \infty$,

$$(M\sqrt{M})^{|\vec{p}|} \frac{N^{n-|\vec{k}|-|\vec{p}|} Par(E_M^l(\vec{q}'))}{\mu_+(N)^{lM}} \rightarrow (P_{-1}\sigma)^{|\vec{k}|-n} \cdot g(\vec{y}') T^{-\frac{3}{2}(n-|\vec{k}|-|\vec{p}|)}. \quad (148)$$

The result then follows from dominated convergence theorem. \square

7.5 Renormalization

Indeed, some of the integrals $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; \epsilon)$ diverges to $\pm\infty$ when $\epsilon \rightarrow 0$, because of the behavior (a) described in Section 7.3. More precisely, locally on the right of some C_i in case (a), as the leg $Y_{i,2}^l$ of the updated block approaches C_i with distance lower bounded by ϵM , we are conditioning on the events that for some $w_i > \epsilon$, there's a walk bridge from C_i to $C_i + \lfloor w_i M \rfloor$, staying above $q(C_i)$ and with height $q(C_i)$ at the starting and ending point, or equivalently a walk bridge sampled from $\mathcal{W}(0, 0, \lfloor w_i M \rfloor; M)$. By calculating its partition function $|\mathcal{W}(0, 0, \lfloor w_i M \rfloor; M)|$, we see that it contributes a factor $(w_i)^{-\frac{3}{2}}$ to $g(\vec{y}')$ in the limit. Therefore we have an integral of order (up to factors independent of ϵ)

$$\int_{\epsilon}^1 w_i^{-\frac{3}{2}} dw_i,$$

which is finite for all $\epsilon > 0$ but blows up in the limit.

Recall that for each $l \geq 1$, we split

$$\frac{1}{\mu_+(N)^{lM}} \left[\sum_{i=1}^N \left(\frac{\mathcal{D}_i}{N} \right)^M \right]^l G(z_1, \dots, z_N; \theta) \Big|_{z_1, \dots, z_N=0}$$

into sums of $C(N, n_0, \vec{k}) \cdot I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; M)$ according to the weird jump \vec{k}, \vec{p} .

For a given $\vec{k} \in K$, that has case (a) behavior at splitting points $C_{j_1}, \dots, C_{j_{\tilde{n}}}$, we handle the divergence of the sum over contributions of configurations with jump data \vec{k} , by the couplings of $2^{\tilde{n}}$ configurations that have either case (1a) or (2a) behavior at each C_{j_i} , see the end of Section 7.3 for more details. By the argument there, we already know that under coupling, the sum of contributions corresponding to $2^{\tilde{n}}$ jump data \vec{k} 's is uniformly bounded in M . More precisely, the couplings produce a factor of order w_i at C_{j_i} , coming from the area of the extra block from C_{j_i} to $C_{j_i} + w_i M$ in case (2a). So the integral becomes of order (up to factors independent of w_i 's, c.f. Eqn. (138))

$$\prod_{i=1}^{\tilde{n}} \int_{\epsilon}^1 w_i^{-\frac{3}{2}} w_i dw_i = \text{constant} - O(\sqrt{\epsilon}).$$

This is simply a rephrasing of Lemma 7.10. Because of Theorem 7.23, we have that the convergence holds outside of the ϵ -domain, so we have the following.

Theorem 7.27. *For a given $\vec{k} \in K$, that has case (a) behavior at splitting points $C_{j_1}, \dots, C_{j_{\tilde{n}}}$, summing up $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p}; \epsilon)$ for $2^{\tilde{n}}$ different \vec{k} 's that are grouped with \vec{k} by the coupling, then taking $\epsilon \rightarrow 0$ gives a collection of convergent integrals.*

We denote each of such convergent integral as the sum over a finite collection of \vec{k} 's of $I_{(l_1, \dots, l_{n_0})}(\vec{k}, \vec{p})$.

7.6 Tail estimate of the power sums

In order to show that (after renormalization in last section) the convergence of the expression of each (\vec{k}, \vec{p}) when $M \rightarrow \infty$ implies the convergence of the total sum, we prove the following lemma, which shows that the summands decay exponentially fast as the number of weird jumps or ∂ increase. Unlike when showing the convergence, instead of separating different \vec{k} 's, here we combine all the terms $C(N, n_0, \vec{k}) \cdot I_{l_1, \dots, l_{n_0}}(\vec{k}, \vec{p}; M)$ with the same data n , $|\vec{k}| - n + n_0$ and p_1, \dots, p_{n_0} .

Lemma 7.28. *Fix $n_0 \geq 1$. There exists a constant $C > 1$ uniformly in \vec{k}, \vec{p}, M and $\vec{l} = (l_1, \dots, l_{n_0})$, such that the total contribution of*

$$\left| \sum_{\vec{k}} C(N, n_0, \vec{k}) \cdot I_{l_1, \dots, l_{n_0}}(\vec{k}, \vec{p}; M) \right| \leq C^{-[n + (|\vec{k}| - n + n_0) + (p_1 + \dots + p_{n_0})]}, \quad (149)$$

where \vec{k} are summed over the group with the same n and $|\vec{k}| - n + n_0$.

Proof. The key of this proof is that one needs to enumerate the walk/block configurations from a different point of view compared to the previous sections. We start from the case when $n_0 = 1$ which corresponds to first moment tail estimate.

Like in Section 5, we first draw the random walk excursion, which is of length M from 0 to 0 conditioning on staying nonnegative. Then we draw the block process $\vec{q}(t)$ **backwards in time**. More precisely, we first fix all the jump time, without specifying their corresponding variables. Denote the “free jumps” and last jumps (defined in Definition 7.13, and free jumps are all the remaining weird jumps) of all these variables by $j_1, \dots, j_{|\vec{k}| - n + n_0 + 1}$, and l_1, \dots, l_{n-1} respectively, ordered reversely in the occurring time. For notation convenience, denote time M and time 0 as j_0 and $j_{|\vec{k}| - n + n_0}$ respectively.

We draw the blocks recursively. For $i = 1, 2, \dots$, On the time interval between j_{i-1} and j_i , if there are u ($u \geq 1$) last jumps, observe from their combinatorial feature (the block touches the walk at those points when there's a down step -1) that, the number of positions these jumps can locate is

$$E_M(j_i) - \min_{t \in \llbracket j_i + 1, j_{i-1} - 1 \rrbracket} E_M(t).$$

Therefore there are no more than

$$\frac{[E_M(j_i) - \min_{t \in \llbracket j_i + 1, j_{i-1} - 1 \rrbracket} E_M(t)]^u}{u!} \quad (150)$$

of the last jumps on $\llbracket j_i + 1, j_{i-1} - 1 \rrbracket$. These jumps, if they exist, will change $\vec{q}(t)$, so after specifying them (and equivalently knowing the value of $\vec{q}(j_i)$), we determine the value of $\vec{q}(j_{i-1})$.

If the jump at j_i is a “first jump” of some z_l , then $\deg(z_l)$ must be 0 before it and therefore there's no freedom of $\vec{q}(j_{i-1})$ conditioning on $\vec{q}(j_i)$. In other cases, we have the key observation that if the jump at j_i is of z_l , then given $q_l(j_i)$, the choice of $q_l(j_{i-1})$ is at most $2q_l(j_i)$. So the total number of choices of $\vec{q}(j_{i-1})$ is bounded by $\sum_l q_l(j_i) \leq \max_{0 \leq t \leq M} E_M(t)$.

Now we consider the case for general $\vec{l} = (l_1, \dots, l_{n_0})$. Recall that we need to do conditioning on the walk, and in this proof we do it in a slightly different order. For simplicity, we first consider the walk/block configurations given by $\vec{k} \in K$, and not in case (3) on the left of all C_1, \dots, C_{n_0-1} ⁷. Define $X_{i,1}, Y_{i,1}, X_{i,2}$ and τ_i as before. We count the total weight of configurations in the following steps:

0. Choose n_0 “main variables” and $n - n_0$ “auxillary variables” from z_1, \dots, z_N .

1. Fix $\tau_1, \dots, \tau_{n_0-1} \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$.

2. Fix all the weird jump times of z_i where $i = 1, 2, \dots, n_0$. Since $k \in K$, either there is no jump of z_i after time C_i , or the last jump of z_i is the first jump after some C_j where $i \leq j \leq n_0 - 1$. Then based on the data fixed in step 1, determine the local behavior type⁸ of the configuration on the right of each C_j ($1 \leq j \leq n_0 - 1$): (1a), (2a) (1b) or (2b), as defined in Section 7.2 and 7.3.

⁷For each fixed \vec{k} , the contribution given by configurations with $\vec{k} \notin K$ or with some case (3) behavior turn out to be of order $o(1)$, see the previous sections.

⁸If there are jumps of z_i after C_i , some C_j ($j \geq i$) must has case (2a) behavior on its right, and once such j is chosen, the last jump time of z_i is fixed at τ_j .

3. Recall that in case (2a), there is a last jump of z_j at τ_i that sends $q_j(t) = \deg(z_j)$ back to 0. We further restrict ourselves to the case that $j = 1, 2, \dots, n_0$, since the case $j = n_0 + 1, \dots, n$ will be coupled with configurations in case (1a) as described in Section 7.3.

4. For each $i = n_0 + 1, \dots, n$, determine the occurring time of the first jump of z_i . Then determine the occurring time of all the remaining free jumps of z_{n_0+1}, \dots, z_n , without specifying their corresponding variables⁹. Based on step 2 and 4, we have the restrictions that

- If we have case (1a), there's no jump on $\llbracket C_i + 1, \tau_i \rrbracket$. If for some C_i we have case (2a), there's no jump on $\llbracket C_i + 1, \tau_i - 1 \rrbracket$, and one jump of $z_i (i = n_0 + 1, \dots, n)$ at τ_i .

5. Based on step 1, if there's no jump of z_i after C_i , $\deg(z_i)$ is forced to be 0 at C_i , i.e., $E_M(C_i) - \lim_{t \rightarrow C_i^-} q(t) = 0$, fix $Y_{i,1}$, under the restrictions that

- There's no jump on $\llbracket Y_{i,1} + 1, C_i \rrbracket$.
- There's one jump at $Y_{i,1}$ when $Y_{i,1} > C_{i-1}$.

6. Split $\llbracket 0, lM \rrbracket$ into disjoint segments by C_i 's, $Y_{i,1}$'s, τ_i 's (when $\tau_i < \infty$). Denote the segments by $\tilde{I}_1, \dots, \tilde{I}_{\tilde{S}}$, ordered backwards in time, and the left endpoint of I_q by \tilde{t}_q . More precisely, all the above positions (except $C_0=0$) serve as the right boundary of the segment on their left. Note that $\tilde{S} \leq 3n_0$.

Split $\llbracket 0, lM \rrbracket$ into disjoint intervals by C_i 's, $Y_{i,1}$'s, τ_i 's (when $\tau_i < \infty$), all the jumps of z_1, \dots, z_{n_0} , and all the free jumps (which have determined occurring time by step 5). More precisely, all the above positions (except $C_0=0$) serve as the right boundary of the interval on their left. Note that the intervals are subdivisions of the segments. Denote the intervals defined in step 6 by $I_q (q = 1, 2, \dots, S)$, and the left endpoint of I_q by t_q . Note that since the total number of C_i 's, $Y_{i,1}$'s, τ_i 's is bounded by $3n_0$, we have

$$|\vec{k}| - n + n_0 \leq S \leq |\vec{k}| - n + 4n_0.$$

Then we determine two data of the last jump times¹⁰ l_{n_0+1}, \dots, l_n . First, we determine the number of last jumps in each interval, under the same restrictions given in step 4 and 5. Second, we determine the order of these $n - n_0$ last jumps.

7. For $p = 1, 2, \dots, \tilde{S}$, starting from $p = 1$ (the segment ending at lM), draw a walk $E_M(t)$ on \tilde{I}_p , that is "compatible" with all the "jump time data" determined in the previous steps¹¹.

8. Still on segment \tilde{I}_p , doing backwards in time, determine the height of all free jumps, and last jumps of z_1, \dots, z_n on each $I_q \subset \tilde{I}_p$, in the same way as in the case $n_0 = 1$.

9. Repeat step 7 and 8 on segment \tilde{I}_{p+1} until $p = \tilde{S}$. Note that the walks on each segments are mutually independent, conditioning on the boundary heights.

We now make a discussion of the factors arising from each of the steps in the above counting procedure. Recall that the contribution of each walk/block configuration can be both positive or negative, and by counting we get a summation of their absolute values. For simplicity, we omit the effects given by $\partial_1, \dots, \partial_{n_0}$, i.e., take $\vec{p} = 0$ for now.

0. The combinatorial factor

$$\binom{N}{n_0} \binom{N - n_0}{n - n_0} \sim \frac{N^n}{(n - n_0)!}.$$

Each weird jump and ∂ give a factor $\frac{1}{N}$, so in total $\left(\frac{1}{N}\right)^{|\vec{k}|+|\vec{p}|}$.

1. The choice of τ_i 's will give a weighted sum in terms of the corresponding walks, see the previous sections for more details.

⁹This turns out to be helpful for bounding the total weight of the configurations.

¹⁰They should be occurring after the corresponding first jumps, but we don't take this restriction into account in this proof.

¹¹More precisely, we have

- $E_M(t) \geq E_M(C_i)$ on $[Y_{i,1}, C_i]$;
- $E_M(C_i) = E_M(\tau_i - 1) = E_M(\tau_i) + 1$ when $\tau_i < \infty$, and $E_M(t) \geq E_M(C_i)$ on $[C_i, C_{i+1}]$ otherwise;
- for $j = 1, 2, \dots, n_0$, if l_j exists, $E_M(t) \geq E_M(l_j -)$ on the interval whose right boundary is l_j ; and
- $E_M(t) = E_M(t - 1) - 1$ at each jump time t .

2. The total number of choice of local behavior types is bounded by a constant. Conditioning on such choice, let $\tilde{k}_j = k_j$ if no jump of z_j is after C_j , and $\tilde{k}_j = k_j - 1$ otherwise. Then we have the combinatorial factor

$$\leq \binom{LM}{\tilde{k}_1, \dots, \tilde{k}_{n_0}, LM - \sum_{j=1}^{n_0} \tilde{k}_j} \leq \frac{(LM)^{\sum_{j=1}^{n_0} \tilde{k}_j}}{\tilde{k}_1! \dots \tilde{k}_{n_0}!}.$$

4. We have the combinatorial factor at most ¹²

$$\binom{LM - \sum_{j=1}^{n_0} k_j}{1!, \dots, 1!, |\vec{k}| - \sum_{j=1}^{n_0} k_j - 2n + 2n_0, LM - |\vec{k}| + n - n_0} \leq \frac{(LM)^{|\vec{k}| - \sum_{j=1}^{n_0} k_j - n + n_0}}{(|\vec{k}| - \sum_{j=1}^{n_0} k_j - 2n + 2n_0)!},$$

note that the factor on the right side of the inequality gives the correct order when $|\vec{k}|$ is fixed.

5. Fixing $Y_{i,1}$ gives some restrictions to both the walk and the jump times. Let δ and $\delta_1, \delta_2, \delta_3$ denote the number of $Y_{i,1} > C_{i-1}$, and the number of $Y_{i,1} > C_{i-1}$ where there is a free jump of z_{n_0+1}, \dots, z_n /last jump of z_{n_0+1}, \dots, z_n /jump of z_1, \dots, z_{n_0} at $Y_{i,1}$, respectively. Then by definition $\delta = \delta_1 + \delta_2 + \delta_3 \leq n_0 - 1$. Since the occurring time of the last jumps of z_{n_0+1}, \dots, z_n are not fixed yet, we get a factor

$$\sim (LM)^{\delta_2}.$$

6. We have the combinatorial factor (note that $|\vec{k}| - n + n_0 \geq n - n_0$ by definition, so choosing S among its possible values makes a $O(1)$ difference)

$$\binom{S - 1 + n - n_0}{n - n_0} (n - n_0)! \sim \frac{(|\vec{k}| - n + n_0 - 1 + n - n_0)!}{(|\vec{k}| - n + n_0 - 1)!}.$$

7 and 8. On each I_q , denote the number of last jumps of z_{n_0+1}, \dots, z_n by $u(q)$. Then as in $n_0 = 1$ case, the number of choices of last jump heights on each I_q is bounded by

$$\frac{[E_M(t_q) - \min_{t \in I_q} E_M(t)]^{u'(q)}}{u'(q)!},$$

where $u'(q) = u(q)$ or $u(q) - 1$ is the ‘‘degree of freedom’’ of the last jump height, where the latter case happens when in some of the configurations, one of the last jump heights is fixed by the restriction $E_M(C_i) = q(C_i) = \sum_j q_{j \neq i}(C_i)$, when C_i has case (2) behavior and $I_q \subset [C_i, C_{i+1}]$.

As for the choices of each free jump height and its corresponding variable, there are three cases:

- no freedom if it’s a first jump;
- if not a first jump, generically bounded by (and of the same order of) $\max_{0 \leq t \leq LM} E_M(t)$; and
- in some of the configurations, one of the free jump height is fixed by the same restriction as above, there are at most $n - n_0$ choices of this jump about its corresponding variables.

Also note each restriction at C_i can fix at most one free or last jump, and there are at most $n_0 - 1$ such restrictions. So the total loss of degree of freedom of free and last jumps are at most $n_0 - 1$.

Denote the degree of freedom loss of the free jumps by δ_4 , the number of z_1, \dots, z_{n_0} that doesn’t have a last jump (equivalently, no jump after C_i) + the number of z_1, \dots, z_{n_0} that doesn’t have a first jump (equivalently, no jump before $C_{i-1} + 1$) by δ_5 , then we obtain the factor

$$n^{\delta_4} \left[\max_{0 \leq t \leq LM} E_M(t) \right]^{|\vec{k}| - 2n - \delta_4 + \delta_5} \prod_{q=1}^S \frac{[E_M(t_q) - \min_{t \in I_q} E_M(t)]^{u'(q)}}{u'(q)!}, \quad (151)$$

To sum over all the walk compatible with the given ‘‘jump time data’’, we rely on the same probabilistic interpretation as in previous sections, such that each walk is a combination of two groups of (mutually independent) random objects: the walk heights at the endpoints of each segment $\tilde{I}_1, \dots, \tilde{I}_S$, and the walk bridges

¹²The key is that the number of available time positions of the jumps is of order $O(M)$.

on each segment conditioning on the boundaries (and staying nonnegative). We claim without recalling more details from previous sections that, for each given “jump time data”, the collection of N -factors together with the factor $\mu_+(N)^{lM}$ give the right order renormalization¹³ such that for all N large enough, up to a $O(1)$ factor¹⁴, one can view the weight of $E_M(t)$ as the probability measure of a conditional walk bridge. Because of this, we omit all the N -factors from now on.

We take the weighted sum over $E_M(t)$ as taking an expectation on the conditional walk bridge described above. Denote $\max_{t \in \tilde{I}_p} E_M(t) = h_p N^{\frac{1}{3}}$, then by Lemma 6.4, among any possible conditioning at C_i 's, $\text{Prob}[h \leq h_p < h + 1] \sim \exp(-ch^2)$ for some constant $c > 0$ uniform in N, M ¹⁵. As for the second part of Eqn. (151), we apply¹⁶ Lemma 6.5. Combining these two together gives an upper bound of order

$$\sim h^{|\vec{k}|-2n-\delta_4+\delta_5} \prod_{p=1}^{\tilde{S}} C^{\tilde{u}(p)} \log(\tilde{u}(p))^{\tilde{u}(p)} [\tilde{u}(p)^{-\tilde{u}(p)/2} + \tilde{u}(p)^{-\tilde{u}(p)} h^{\tilde{u}(p)}] \exp(-ch^2), \quad (152)$$

where we sum over the walks with $N^{-\frac{1}{3}} \cdot \max_{0 \leq t \leq lM} E_M(t) \in [h, h + 1)$, and $\tilde{u}(p)$ denotes the total degree of freedom of last jumps on \tilde{I}_p , given by a sum over $u(q)$ for $q = 1, 2, \dots, S$ such that $I_q \subset \tilde{I}_p$. Note that the total loss of degree of freedom on each \tilde{I}_p is 0 or 1.

To upper bound the product

$$\prod_{p=1}^{\tilde{S}} C^{\tilde{u}(p)} \log(\tilde{u}(p))^{\tilde{u}(p)} [\tilde{u}(p)^{-\tilde{u}(p)/2} + \tilde{u}(p)^{-\tilde{u}(p)} h^{\tilde{u}(p)}], \quad (153)$$

note that at least one $\tilde{u}(p) > \frac{n-n_0}{3n_0}$, and for this $\tilde{u}(p)$ we have¹⁷

$$\tilde{u}(p)^{-\tilde{u}(p)/2} + \tilde{u}(p)^{-\tilde{u}(p)} h^{\tilde{u}(p)} \leq \left(\frac{n-n_0}{3n_0}\right)^{-\frac{n-n_0}{6n_0}} + \left[\left(\frac{3n_0 h}{n-n_0}\right)^{\frac{n-n_0}{3n_0}} \text{ or } \left(\frac{3n_0 h}{n-n_0}\right)^n\right];$$

where for the first term we use the fact that $x \mapsto x^{-\frac{x}{2}}$ is decreasing and upper bound it at $x = \frac{n-n_0}{3n_0}$, and the two cases for the second term correspond to the regime $h < \frac{n-n_0}{3n_0}$ (so $\frac{h}{\tilde{u}(p)} \leq \frac{3n_0 h}{n-n_0} < 1$ and we minimize the exponent) or $h \geq \frac{n-n_0}{3n_0}$ (so $\max(\frac{h}{\tilde{u}(p)}, 1) \leq \frac{3n_0 h}{n-n_0}$ and we maximize the exponent at n).

For other $\tilde{u}(p)$'s, we use the fact that the function $x \mapsto x^{-x} h^x$ is increasing when $x < h/e$, and decreasing when $x > h/e$. So instead we have

$$\tilde{u}(p)^{-\tilde{u}(p)/2} + \tilde{u}(p)^{-\tilde{u}(p)} h^{\tilde{u}(p)} \leq 1 + e^{h/e}.$$

Therefore we can bound Eqn. (153) by (taking $h \geq \mathbb{Z}_{\geq 0}$)

$$C^n \log(n)^n [h/(n-n_0)]^{c(n-n_0)+n_0 \mathbf{1}_{h \geq n-n_0}},$$

where c is a positive constant between $\frac{1}{6n_0}$ and 1. The exponent on the right comes from the fact that

$$\sum_{p=1}^{\tilde{S}} \tilde{u}(p) \leq n.$$

Summing over $h \in \mathbb{Z}_{\geq 0}$, we have that for $u_1, u_2 \geq 0$,

$$\begin{aligned} \sum_{h \geq 0} h^{u_1+u_2} (n-n_0)^{-u_1} \exp(-ch^2) &\sim C^{u_1+u_2} \sqrt{(u_1+u_2)!} (n-n_0)^{-u_1} \\ &\sim C^{u_1+u_2} \left(\frac{u_1+u_2}{e}\right)^{\frac{u_1+u_2}{2}} (u_1+u_2)^{\frac{1}{4}} (n-n_0)^{-u_1}, \end{aligned}$$

¹³When there are $\tilde{n} \leq n_0$ C_i 's with case (1a) behavior, this is true up to a coupling of $2^{\tilde{n}}$ corresponding terms, or equivalently restricting to the configurations that all $\tau_i \geq \epsilon M$ for arbitrarily small $\epsilon > 0$.

¹⁴When $\vec{k} \in K$, otherwise the factor is $o(1)$.

¹⁵ c depends only on l, T and σ .

¹⁶When $|\vec{k}|$ is fixed as in the previous sections, the number of intervals S is bounded, so one can bound the second part of Eqn. (151) in the same way as the first part. The main purpose introducing this lemma is to get an upper bound uniformly in $|\vec{k}|$ and S , because the bound only depends on the number of segments \tilde{S} , which is still uniformly bounded.

¹⁷We get 1 when the exponent $n-n_0$ is 0.

where $u_1 = c(n - n_0) + n_0 \mathbf{1}_{h \geq n - n_0} \geq 0$ and $u_2 = |\vec{k}| - 2n - \delta_4 + \delta_5 \in \mathbb{Z}_{\geq 0}$, $u_1 + u_2 \approx |\vec{k}| - n + n_0$.

We now collect all the factors appearing in step 0 to 8, except that we omit the N -factor. This gives an upper bound of order

$$\frac{1}{(n - n_0)!} \frac{1}{(|\vec{k}| - \sum_{j=1}^{n_0} k_j - 2n + 2n_0)!} \frac{1}{k_1! \cdots k_{n_0}!} \frac{(|\vec{k}| - n + n_0 - 1 + n - n_0)!}{(|\vec{k}| - n + n_0 - 1)!} \log(n)^n n^{\delta_4} C^{|\vec{k}|} (u_1 + u_2)^{(u_1 + u_2)/2} (n - n_0)^{-u_1} (u_1 + u_2)^{\frac{1}{4}}. \quad (154)$$

Combining $\frac{1}{(n - n_0)!}$ and $\frac{(|\vec{k}| - n + n_0 - 1 + n - n_0)!}{(|\vec{k}| - n + n_0 - 1)!}$, as well as $\frac{1}{(|\vec{k}| - \sum_{j=1}^{n_0} k_j - 2n + 2n_0)!}$ and $\frac{1}{k_1! \cdots k_{n_0}!}$, we upper bound Eqn. (154) by (the constant C takes larger value than above)

$$\log(n)^n n^{\delta_4} C^{|\vec{k}|} \frac{1}{(|\vec{k}| - 2n + 2n_0)!} (u_1 + u_2)^{(u_1 + u_2)/2} (n - n_0)^{-u_1} (u_1 + u_2)^{\frac{1}{4}}. \quad (155)$$

Then replacing $|\vec{k}| - 2n + 2n_0$ by u_2 and $n - n_0$, n by u_1 , and changing the constant C again, the above can be bounded by

$$\log(u_1)^{u_1} u_1^{\delta_4} C^{u_1 + u_2} \frac{1}{u_2!} (u_1 + u_2)^{(u_1 + u_2)/2} u_1^{-u_1} (u_1 + u_2)^{\frac{1}{4}}, \quad (156)$$

which is clearly decaying exponentially fast as u_1 or $u_2 \rightarrow \infty$. This finishes the calculation for $\vec{p} = 0$.

Considering the contribution given by $\vec{p} \neq 0$. Indeed, for any fixed walk/block configuration, one can sum over all possible \vec{p} and get a factor bounded by

$$\left(1 + \frac{\max_{0 \leq t \leq lM} E_M(t)}{N} \right)^{lM} \leq \exp(Ch)$$

for some constant $C > 0$ independent¹⁸ of N , $N^{-\frac{1}{3}} \max_{0 \leq t \leq lM} E_M(t) \in [h, h + 1)$. Clearly this factor will be absorbed by the factor $\exp(-ch^2)$ in Eqn. (154).

Now we turn to the case $\vec{k} \notin K$. The walk/block configurations in this group have one or two of the following non-generic behaviors:

- For some $i = 1, 2, \dots, n_0 - 1$, there are more than one jumps of z_i after C_i , or the unique jump of z_i after C_i is not the first jump after any C_j ($j = i, i + 1, \dots, n$)¹⁹.
- For some $i = 2, 3, \dots, n_0$, there are jumps of z_i before C_{i-1} , but $\deg(z_i) = 0$ at time C_i . In other words, there is an (unnecessary) extra last jump of z_i before²⁰ C_i .

We claim (without giving more details) the counting procedure for configurations with such behaviors does not differ much compared to the one for $\vec{k} \in K$, and therefore the total weight of such configurations (with fixed N , n and $|\vec{k}|$) also decays exponentially fast as n and $|\vec{k}|$ grow. On the other hand, by Lemma 7.15 such weight is of order $o(1)$ as $N \rightarrow \infty$, so the result follows. \square

Remark 7.29. *The reason why we count the blocks backwards in time is that, if counting them forward in time, then given $\vec{q}(t - 1)$ ($t \in \llbracket 0, lM \rrbracket$), the number of choices of $\vec{q}(t)$ would only be bounded by $n \cdot E_M(t - 1)$: there are at most n choices of jump variables, and each has freedom bounded by the current height of the walk.*

We see from Theorem 7.23 that as in first moment calculation, each copy of $P_{-1}\sigma$ comes with $T^{\frac{3}{2}}$. Therefore we conclude with the following statement.

¹⁸ C depends only on l , T and σ .

¹⁹In other words, it is not “releasing the walk locally on the right of C_j ”, see Lemma 7.15 and its proof.

²⁰This is the case 3 behavior defined in Section 7.3

Theorem 7.30. Fix $T > 0$. For each $l \geq 1$, $M = \lfloor TN^{\frac{2}{3}} \rfloor$, as $M \rightarrow \infty$, the l^{th} moment of $\sum_{i=1}^N \left(\frac{\lambda_i}{\mu_+ N} \right)^M$ of each our target ensemble converges to

$$\begin{aligned} & \sum_{n_0=1}^l \sum_{l_1+\dots+l_{n_0}=l} \sum_{\vec{k} \in K} \sum_{p_1, \dots, p_{n_0}=0}^{\infty} \binom{l}{l_1, \dots, l_{n_0}} \frac{1}{n_0!} I_{(l_1, \dots, l_{n_0})}(\vec{k}, p) \\ &= \sum_{m=n-2n_0}^{\infty} \left(\sum_{n_0=1}^l \sum_{l_1+\dots+l_{n_0}=l} \sum_{\vec{k} \in K} \sum_{\vec{p} \in \mathbb{Z}_{\geq 0}^{n_0}} \mathbf{1}_{m=|\vec{k}|-n+|\vec{p}|} \binom{l}{l_1, \dots, l_{n_0}} \frac{1}{n_0!} I_{(l_1, \dots, l_{n_0})}(\vec{k}, p) \right) \end{aligned} \quad (157)$$

which is a universal expression²¹, in the sense that m gives the power of factor $T^{\frac{3}{2}} P_{-1} \sigma$ in the summand s . Here $m \geq n - 2n_0$ since $|\vec{k}| \geq 2(n - n_0)$. The sum over \vec{k} converges in the sense given in Theorem 7.27.

This finishes the higher moment calculation.

8 Proof of main theorem

In this section we finish the proof of Theorem 1.2, based on Theorem 7.23 and 7.30. By the above two theorems, we have the convergence of

$$\mathbb{E} \left[\prod_{j=1}^l \left(\sum_i \left(1 + \frac{\lambda'_i(N)}{\mu_+(N)N^{\frac{2}{3}}} \right)^{\lfloor T_j N^{\frac{2}{3}} \rfloor (+0 \text{ or } 1)} \right) \right], \quad (158)$$

for any fixed $T_1, \dots, T_l > 0$. This is because Eqn. (158) differs from

$$\left(\sum_{i=1}^N \left(\frac{\lambda_i(N)}{\mu_+(N)N} \right)^M \right)^l$$

only by choosing M to be either the closest odd or even to $\lfloor TN^{\frac{2}{3}} \rfloor$, and taking T 's to be distinct. From the proofs presented in Section 3 to Section 7 we see that the convergence follows from exactly the same arguments.

Lemma 8.1. Convergence of Eqn. (158) implies the convergence of Laplace transform

$$\mathbb{E} \left[\prod_{j=1}^l \left(\sum_i \exp \left(\frac{T_j \lambda'_i(N)}{\mu_+} \right) \right) \right], \quad (159)$$

to the same limit, for all $T_1, \dots, T_l > 0$.

Proof. Split Eqn. (158) as a finite sum of products of terms of the form

$$\mathbb{E} \left[\text{const} \cdot \prod_{j=1}^l \lambda_{i_j}(N)^{\lfloor T_j N^{\frac{2}{3}} \rfloor (+0 \text{ or } 1)} \right],$$

and split Eqn. (159) in similar way. Then we consider four different regime of $\lambda'_i(N)$ (equivalently of $\lambda_i(N)$):

1. $\lambda'_i(N) \leq -\mu_+(N)N^{\frac{2}{3}}$.
2. $-\mu_+(N)N^{\frac{2}{3}} \leq \lambda'_i(N) < -N^{\frac{1}{100}}$,
3. $-N^{\frac{1}{100}} \leq \lambda'_i(N) < N^{\frac{1}{100}}$,

²¹In the concurrent work [GXZ] that gives Laplace transform of Airy $_{\beta}$ line ensemble, when $\tau_1 = \dots = \tau_m$, the expression $\mathbf{L}_{\beta}(\vec{\mathbf{k}}, \vec{\tau})$ is identical to Eqn. (157) in our text (when $P_{-1} = \frac{1}{2}, \sigma = 1$). Compared with our exposition, [GXZ] gives explicit density function of the Brownian bridges, and moreover sums over \vec{p} to get an exponentiated area between Brownian bridge and blocks in $\mathbf{I}_{\beta, \vec{\mathbf{k}}}[\vec{\mathbf{p}}, \vec{\mathbf{v}}, \mathbf{H}]$, and does an extra cancellation between different (l_1, \dots, l_{n_0}) 's in Section 4.6. Both operations are designed to simplify the expression.

$$4. \lambda'_i(N) \geq N^{\frac{1}{100}},$$

In regime (1), (2), since there are at most N such $\lambda'_i(N)$'s, the contribution of products containing such $\exp\left(\frac{T_j \lambda'_i(N)}{\mu_+}\right)$ in Eqn. (159) is clearly small. Similarly the contribution of products containing $\lambda_i(N)^{\lfloor T_j N^{\frac{2}{3}} \rfloor}$ in regime (2) is small. The contribution of products containing $\lambda_i(N)^{\lfloor T_j N^{\frac{2}{3}} \rfloor}$ in regime (1) is also small, since when changing the exponent $\lfloor T_j N^{\frac{2}{3}} \rfloor$ by 1 the sign of such terms change, but the value of Eqn. (158) is almost unchanged for N large enough.

In regime (3), which is the “generic” regime, by elementary calculation we have

$$\begin{aligned} 0 &\leq \exp\left(\frac{T_j \lambda'_i(N)}{\mu_+(N)}\right) - \left(1 + \frac{\lambda'_i(N)}{\mu_+(N) N^{\frac{2}{3}}}\right)^{\lfloor T_j N^{\frac{2}{3}} \rfloor (+0 \text{ or } 1)} \\ &\leq \left[\exp\left(\frac{T_j}{\mu_+(N)^2} N^{\frac{1}{100} \cdot 2 - \frac{2}{3}}\right) - 1 \right] \left(1 + \frac{\lambda'_i(N)}{\mu_+(N) N^{\frac{2}{3}}}\right)^{\lfloor T_j N^{\frac{2}{3}} \rfloor (+0 \text{ or } 1)}. \end{aligned} \quad (160)$$

Since $\mu_+(N) \rightarrow \mu_+$ and there are at most N^l terms inside the bracket, the difference of above type goes to 0 as $N \rightarrow \infty$ as well.

Finally, the contribution of products containing terms in regime (4) is $o(1)$ in Eqn. (159), and therefore in Eqn. (158) as well, for all positive T_j 's, because of the exponential tail behavior of $\lambda'_1(N)$ outside of $\mu_+(N)$ given in Eqn. (29). \square

Denote the random point configuration of $\{\frac{1}{C_0} \lambda'_i(N)\}_{i=1}^N$ by \mathcal{X}_N , a sequence of point processes on \mathbb{R} . By the universality in Theorem 7.30 and Lemma 8.1, the Laplace transform of \mathcal{X}_N converges in moment to the Laplace transform of $\text{Airy}(\beta)$.

We claim that $\{\mathcal{X}_N\}$ is tight in the $W^\#$ topology of boundedly finite Borel measures on \mathbb{R} , see [DVe1, A2.6.] for the definition. By [DVe2, Proposition 11.1.VI.], this is equivalent to checking that for any compact interval $A \subset \mathbb{R}$, the random variable

$$N_A := \text{number of particles in } A$$

is uniformly tight in N . It suffices to check $A = [t, \infty)$ for any $t \in \mathbb{R}$ since such sets contain all compact intervals. We have

$$\sum_i \left(1 + \frac{\lambda'_i(N)}{\mu_+(N) N^{\frac{2}{3}}}\right)^{\lfloor T N^{\frac{2}{3}} \rfloor (+0 \text{ or } 1)} \geq \frac{N_A}{2} \exp\left(\frac{Tt}{\mu_+}\right)$$

when N is large and the exponent is even, and its expectation converges as $N \rightarrow \infty$ as a special case of Eqn. (158), hence the uniform tightness follows.

It remains to check that the moments of Laplace transform of $\text{Airy}(\beta)$ uniquely determines a point process on \mathbb{R} . If we have this, because of tightness of $\{\mathcal{X}_N\}$ and the moment convergence of Laplace transform, every subsequence of $\{\mathcal{X}_N\}$ converges to the same limit $\text{Airy}(\beta)$, and so does the whole sequence.

By definition of correlation measures ρ_l of point processes (see [DVe1], [DVe2]), for a point process \mathcal{X} on \mathbb{R} with particles $\Lambda_0 < \Lambda_1 < \Lambda_2 < \dots$

$$\mathbb{E} \left[\prod_{j=1}^l \left(\sum_i \exp(T_j \Lambda_i) \right) \right] = \int \cdots \int \rho_l(dx_1, \dots, dx_l) e^{T_1 x_1 + \dots + T_l x_l} + \text{lower order terms}, \quad (161)$$

where the lower order terms give a linear combination of Laplace transforms of $\rho_1, \dots, \rho_{l-1}$ of similar form.

Lemma 8.2. *All ρ_l 's are uniquely determined by the Laplace transform above.*

Proof. By iteration starting from $l = 1$ on Eqn. (161), it remains to show that for any $l = 1, 2, \dots$, the Laplace transform

$$\int \cdots \int \rho_l(dx_1, \dots, dx_l) e^{T_1 x_1 + \dots + T_l x_l} \quad (162)$$

$(T_1, \dots, T_l > 0)$ uniquely determines ρ_l . It's clear that Eqn. (162) is finite for all $(T_1, \dots, T_l) \in \mathbb{R}_{>0}^l$, so fix some $(S_1, \dots, S_l) \in \mathbb{R}_{>0}^l$, we have $\rho_l(dx_1, \dots, dx_l)e^{S_1x_1 + \dots + S_lx_l}$ is in L^1 , i.e a finite measure, and its Fourier transform is fixed by Eqn. (162) under analytic continuation. Then $\rho_l(dx_1, \dots, dx_l)e^{S_1x_1 + \dots + S_lx_l}$ and hence $\rho_l(dx_1, \dots, dx_l)$ is also fixed. \square

Recall also that by definition, the distribution of \mathcal{X} is determined by finite-dimensional distribution of N_{A_1}, \dots, N_{A_k} , for any compact sets A_1, \dots, A_k , and

$$\mathbb{E}[N_A(N_A - 1) \cdots (N_A - k + 1)] = \int_A \cdots \int_A \rho_k(dx_1, \dots, dx_k). \quad (163)$$

So the (joint) moments of N_A 's are fixed by correlation measures. It remains to check that for any compact set A , the tail probability of N_A is exponentially decaying, so moments of N_A determine the distribution of N_A . We check this for $A = (\lambda, \infty)$ for any $\lambda \in \mathbb{R}$, which will suffice.

Since $\mathcal{X} = \text{Airy}(\beta)$, we use the following result from [RRV, Theorem 1.2]: for $\lambda \in \mathbb{R}$,

$$\mathbb{P}[N_{(\lambda, +\infty)} \geq k] = \mathbb{P}[\eta_k > \lambda] = \int_{\mathbb{R}^k} \kappa(\lambda, dx_1) \kappa(x_1, dx_2) \cdots \kappa(x_{k-1}, dx_k), \quad (164)$$

where $\kappa(x, \cdot)$ is the distribution of first passage time to $-\infty$ of diffusion $p(t)$ ($B(t)$ is a standard Brownian motion)

$$dp(t) = (t - p(t)^2) dt + \frac{2}{\sqrt{\beta}} dB(t), \quad (165)$$

that starts from $+\infty$ at time x .

Then Eqn. (164) can be rewritten as

$$\mathbb{P}[N_{(\lambda, +\infty)} \geq k] = \mathbb{P}[p(t) \text{ reaches } -\infty \text{ for at least } k \text{ times when starting from time } \lambda].$$

We collect the facts about $\kappa(x, \cdot)$ we need in the next lemma.

- Lemma 8.3.** (1). $\kappa(x, y) = 0$ for $y < x$.
(2). For any $x \in \mathbb{R}$, $\kappa(x, \{+\infty\}) > 0$.
(3). $\kappa(x, \{+\infty\})$ increases with x .

Proof. (1) is clear from definition. To see (2) we set $k = 1$ in Eqn. (164) and get

$$\kappa(x, \{\infty\}) = 1 - \mathbb{P}[\eta_1 > x] = \mathbb{P}[\eta_1 \leq x], \quad (166)$$

which is clearly in $(0, 1)$ for all $x \in \mathbb{R}$ (see [RRV, Theorem 1.3] for a tail estimate of η_1). As for (3), it suffices to observe that the diffusion in Eqn. (165) starting at time x is equivalent to

$$dp_x(t) = (x + t - p_x(t)^2) dt + \frac{2}{\sqrt{\beta}} dB(t), \quad (167)$$

starting from time 0. By Eqn. (166) we also have $\kappa(x_1, \{\infty\}) \geq \kappa(x_2, \{\infty\})$ for $x_1 \geq x_2$. \square

It's clear from definition that each run of $p(t)$ after the restart from $+\infty$ is independent. Hence by the above properties of transition measure $\kappa(x, \cdot)$, we have

$$\mathbb{P}[p(x) \text{ reaches } -\infty \text{ for at least } k + 1 \text{ times when starting from time } \lambda] \leq (1 - \kappa(\lambda, \{+\infty\}))^{k+1},$$

which is exponentially decaying. This finishes the proof. \square

9 Further studies

Recall that in this text we assume that for the additions of our β -ensembles, the limiting free cumulants $\kappa_1, \kappa_2, \dots$ are all nonnegative, see Eqn. (32). We do this because in the proofs presented in Section 5 to Section 7, we use a probabilistic point of view, that is, we interpret the huge sum arising from the actions of Dunkl operators on $G(z_1, \dots, z_N; \theta)$ as certain functional of a conditional random walk $W_M(t)$. The steps $X_i(M)$ of $W_M(t)$ are i.i.d before conditioning, and as $M \rightarrow \infty$,

$$X_1(M) \implies X_1,$$

whose distribution function is given in terms of the free cumulants $\kappa_1, \kappa_2, \dots$ and the Voiculescu transform $V(z)$.

Recall also that this assumption includes all cases in which we add Gaussian ensembles with finitely many Laguerre ensembles, and a subset of the cases in which we have subtractions of Laguerre ensembles. We naturally conjecture that if we assume similar convergence of each summands as stated in Section 1.3, the edge limit of the additions will still be Airy β -process under proper rescaling, even when some κ_i 's are negative.

It seems that our current proof cannot be directly applied to this case. For simplicity, let us only consider the first moment calculation. There are two main ingredients of our proof that do not work as before: the steepest descent calculation in Section 4 and the functional CLT in Section 5.3.

Consider $\kappa_l(N)$'s and $V_N(z)$ defined as in Eqn. (10), (11) and (13), and take $\alpha_1 > 0$, $\alpha_2 < 0$. For the steepest descent of such $V_N(z)$, we claim without proof that there is one unique root of $V_N'(z)$ in $(-\frac{1}{\alpha_2}, 0)$ and another in $(0, \frac{1}{\alpha_1})$, which we denote as $z_{c,1}(N)$ and $z_{c,2}(N)$, respectively. By choosing a contour $C_z'(N)$ symmetric about real line that crosses both $z_{c,1}(N)$ and $z_{c,2}(N)$ and only surrounds the pole $z = 0$ of $V_N(z)$, one can check that

$$|m_M(N)| = \frac{1}{M+1} \left| \frac{1}{2\pi i} \oint_{C_z'(N)} V_N(z)^{M+1} dz \right| \sim \frac{1}{\sqrt{M}} |V_N(z_c(N))|^{M+1}, \quad (168)$$

where $z_c(N)$ is whichever of $z_{c,1}(N)$, $z_{c,2}(N)$ that maximizes $|V_N(z)|$ on $C_z'(N)$. Without loss of generality consider $V_N(z_c(N)) > 0$. Since the edge behavior is characterized by $m_M(N)$ for $M \sim N^{\frac{2}{3}}$, we believe that the critical point $z_c(N)$ in the steepest descent gives the upper edge $\mu_+(N) = V(z_c(N))$ as before.

Question: Is it always the case that $|z_{c,1}(N)| < \frac{1}{\alpha_1}$, so that $V_N(z_{c,1}(N))$ converges absolutely as $N \rightarrow \infty$?

The functional CLT argument becomes even more tricky. If we take the same interpretation of conditional random walk with i.i.d step $X_i(M)$'s as before, then $X_i(M)$ is a formal integer-valued random variable, such that for $l \geq 0$,

$$\mathbb{P}(X_i(M) = l) = \frac{\kappa_{l+1}(N) z_c(N)^l}{V_N(z_c(N))},$$

which is negative for some l . Everything can continue to make sense at the level of measure theory by replacing the probability measure with a signed measure to total mass 1. However, such formal random objects behave poorly from the probabilistic point of view as lots of elementary arguments break down. In particular, in Section 5, $P_{-1} = \mathbb{P}(X_i = -1)$ give the asymptotic proportion of down steps, but our arguments no longer hold when X_i is not supported by a probability measure, because even the classical weak LLN no longer hold. In addition, it's not clear to us whether one can state the analog for sign measures of Theorem 5.8 and 6.1 that deal with the weak convergence of conditional random walks to conditional Brownian motions. Another obstruction is showing the tail probability of the random walk decay exponentially as done in Section 6.3. We give a simple example to highlight the issue: if one instead take the i.i.d sum of formal step X_i such that $\mathbb{E}[X_i] = 0$, with distribution say

$$\mathbb{P}(X_i = 1) = \frac{3}{2}, \quad \mathbb{P}(X_i = 3) = -\frac{1}{2},$$

where \mathbb{P} denotes a signed measure of total mass 1, then the total variation of the tail "probability" of $X_1 + \dots + X_M$ is growing to infinity as $M \rightarrow \infty$. Therefore, the weak convergence of the signed measure does not imply the convergence of moments.

Question: Can one generalize the probabilistic interpretation of Dunkl action to formal random walks supported by signed measure, or is a different point of view needed?

Question: Suppose the probabilistic argument can be generalized. Recall also that the sum in Dunkl action does not directly involve the parameter $z_c(N)$ (the walk starts and ends at 0, so powers of $z_c(N)$ cancel out after conditioning). However, $z_c(N)$ and $V_N(z_c(N))$ determines the scaling factor/variance of the random walk σ_N^2 . Is this still the case for negative cumulants? If yes, among the two candidates, $z_{c,1}(N)$ and $z_{c,2}(N)$, what makes the functional “choose” $z_c(N)$ that maximizes $|V_N(z)|$?

10 Appendix A: Ballot problem and more steepest descent

Suppose we have a walk $W(t)$ ($t = 0, 1, 2, \dots, L$), for which the possible increments are $\{(1, k)\}_{k=-1,0,\dots}$. We assign a weight of w_k to a step of height k , and let the weight of each walk be the product of the weights of its increments. Fix the endpoint of all the walks to be starting at $(0, y_0)$ and ending at $(L, 0)$. Let $Z_{(y_0,0,L)}(\{w\})$ be the partition function for these walks, $Z_{(y_0,0,L)}^{good}(\{w\})$ be the partitions function for the walks constrained to stay strictly above the x -axis before the L^{th} step, and $Z_{(y_0,0,L)}^{bad}(\{w\})$ be the partition function for the walks that are not ‘good’.

We are interested in computing the partition function of the ‘good’ paths and its asymptotic growth rate. In the literature these types of paths are known as Lukasiewicz paths. See [BP], and references therein, for a study of these paths with uniform weights. As far as the authors are aware, the asymptotic results for weighted paths presented at the end of this appendix are new.

First, we compute the partition function.

Proposition 10.1. *The partition function for the ‘good’ paths can be written*

$$Z_{(y_0,0,L)}^{good}(\{w\}) = \frac{y_0}{L} Z_{(y_0,0,L)}(\{w\}). \quad (169)$$

Remark 10.2. *When $w_{-1} = w_1 = 1$ and all other $w_i = 0$ this is equivalent to Bertrand’s ballot problem [Ber]. Many other variations of the problem have been solved. The variation presented here in Proposition 10.1 is essentially done in [GS, Section 6] using a generalized version of the ‘reflection principle’. For completeness, we present a proof below.*

Proof. The ‘bad’ walks can either end with a down-, flat-, or up-step, where the up-step can be of height $k = 1, 2, \dots$. We have

$$Z_{(y_0,0,L)}^{bad}(\{w\}) = Z_{(y_0,0,L)}^{bad,-}(\{w\}) + Z_{(y_0,0,L)}^0(\{w\}) + \sum_{k=1}^{\infty} Z_{(y_0,0,L)}^{+,k}(\{w\}) \quad (170)$$

where $Z_{(y_0,0,L)}^{bad,-}(\{w\})$ is the partition function for walks that end with a down-step and are bad, $Z_{(y_0,0,L)}^0(\{w\})$ is the partition function for walks that end with a flat-step with no other constraints, and $Z_{(y_0,0,L)}^{+,k}(\{w\})$ is the partition function for walks that end with an up-step of height k with no other constraints.

Note that each bad walk that ends with a down step must have some last time (before step L) in which they hit the x -axis. We can partition the walks in $Z_{(y_0,0,L)}^{bad,-}(\{w\})$ by the size k of the up-step at that time and the height $i = 1, \dots, k$ of the walk after that step. For each walk in $(bad, -, k, i)$ we can take all the steps after (and including) the last time it hits the x -axis and rotate them 180 degrees to get a walk in $(+, k)$. Note that this is invertible: for a walk in $(+, k)$ find the last down step starting from height $-(k - i)$ and rotate all paths after this. See Example 10.3.

Since these rotations are weight preserving it follows that

$$Z_{(y_0,0,L)}^{bad,-,k,i}(\{w\}) = Z_{(y_0,0,L)}^{+,k}(\{w\})$$

for each k and i . As for each k , i ranges over $1, \dots, k$, we have

$$Z_{(y_0,0,L)}^{bad,-}(\{w\}) = \sum_k k Z_{(y_0,0,L)}^{+,k}(\{w\}). \quad (171)$$

From Eqn. (170) and Eqn. (171), we see that

$$\begin{aligned} Z_{(y_0,0,L)}^{good}(\{w\}) &= Z_{(y_0,0,L)}(\{w\}) - Z_{(y_0,0,L)}^{bad}(\{w\}) \\ &= Z_{(y_0,0,L)}^+(\{w\}) + Z_{(y_0,0,L)}^0(\{w\}) + Z_{(y_0,0,L)}^-(\{w\}) - Z_{(y_0,0,L)}^{bad}(\{w\}) \\ &= Z_{(y_0,0,L)}^-(\{w\}) - \sum_k k Z_{(y_0,0,L)}^{+,k}(\{w\}). \end{aligned} \quad (172)$$

Now, suppose we fix the number of steps of size $i = -1, 0, 1, \dots$ to be n_i . Note that we must have

$$\sum_i n_i = L \quad \text{and} \quad \sum_i i n_i = -y_0.$$

The weight of a walk with this distribution of steps is given by $\prod_i w_i^{n_i}$. The contribution that such walks make to $Z_{(y_0, 0, L)}^-(\{w\}) - \sum_k k Z_{(y_0, 0, L)}^{+, k}(\{w\})$ can be written


$$\begin{aligned} & \left(\binom{L-1}{n_{-1}-1, n_0, n_1, \dots} - \sum_k k \binom{L-1}{n_{-1}, n_1, \dots, n_k-1, \dots} \right) \prod_i w_i^{n_i} \\ &= \frac{n_{-1} - \sum_k k n_k}{L} \binom{L}{n_{-1}, n_0, n_1, \dots} \prod_i w_i^{n_i} \\ &= \frac{y_0}{L} \binom{L}{n_{-1}, n_0, n_1, \dots} \prod_i w_i^{n_i}. \end{aligned}$$

Finally, summing over all choices of the n_i , we have


$$Z_{(y_0, 0, L)}^{\text{good}}(\{w\}) = \frac{y_0}{L} Z_{(y_0, 0, L)}(\{w\}).$$

□

Example 10.3. Here we have $y_0 = 2$ and $L = 9$. Below is a walk in $(\text{bad}, -, 3, 2)$, that is the last step to hit the x -axis is a step of size 3 that ends at height 2:




After the rotation we get



a walk in $(+, 3)$.

Note that the walks



in $(\text{bad}, -, 3, 1)$ and $(\text{bad}, -, 3, 3)$, resp., also map to the same walk in $(+, 3)$ after rotating their highlighted blocks.




Figure 7: The contour used in the steepest descent analysis in Lemma 10.5.

Let us now restrict to the types of walks defined in Section 4.1. In this case, for notation simplicity omitting the index N , we have $w_{-1} = z_c^{-1}$ and $w_i = \kappa_{i+1} z_c^i$ for $i = 0, 1, \dots$, where z_c is the critical point of $V(z) = \frac{1}{z} + \sum_{i=1}^{\infty} \kappa_i z^{i-1}$. Note that the power of z_c in the weight of a walk is equal to the overall change in height.

Lemma 10.4. *The partition function for the walk bridges defined as above from $(0, H)$ to $(M, 0)$ is given by*

$$\frac{H+1}{M+1} \frac{1}{2\pi i} \oint \left(\frac{z}{z_c}\right)^H V(z)^{M+1} dz \quad (173)$$

where our contour only encloses the pole at zero. Taking $H = 0$ we recover Proposition 4.9.

Proof. The results follows as a corollary of Proposition 10.1 by taking $L = M + 1$ and $y_0 = H + 1$. \square

In Section 7, in particular Lemma 7.3, we are interested in walks that are free to start at any non-negative height. In this case, the partition function can be written

$$\sum_{H=0}^{\infty} \frac{H+1}{M+1} \frac{1}{2\pi i} \oint \left(\frac{z}{z_c}\right)^H V(z)^{M+1} dz = \frac{1}{2\pi i} \oint \frac{1}{(1-z/z_c)^2} V(z)^{M+1} dz \quad (174)$$

where the above holds as long as the contour is chosen so that $|z| < |z_c|$.

We would like to compute the asymptotic growth rate of

$$\frac{1}{2\pi i} \oint \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz$$

as $L \rightarrow \infty$.

Lemma 10.5.

$$\frac{1}{2\pi i} \oint \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz \sim \frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1}}{\sqrt{2\pi}} \left(\frac{V''(z_c)}{V(z_c)}\right)^{1/2}$$

as $L \rightarrow \infty$.

Proof. In order to do steepest descent as in Section 4.4, we would normally deform our contour to be a circle around the origin passing through z_c . However, this is problematic as z_c is a pole of $\frac{1}{(1-z/z_c)^2}$, so we must be more careful. Instead, consider a circle of radius ϵ around z_c , we take as our contour the usual contour until it intersects with the small circle, then we take the inner semicircle of the small circle (so that our contour does not contain z_c). Call the portion of our original contour C and the portion of the small circle C_ϵ . See Figure 7.

We have

$$\frac{1}{2\pi i} \oint \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz = \frac{1}{2\pi i} \int_C \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz + \frac{1}{2\pi i} \int_{C_\epsilon} \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz$$

Let $f(z) = \log V(z)$. The first integral can be computed as before.

$$\begin{aligned} \frac{1}{2\pi i} \int_C \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz &= \frac{1}{2\pi i} \int_C \frac{1}{(1-z/z_c)^2} e^{(L+1)f(z)} dz \\ &\approx \frac{1}{2\pi i} \int_C \frac{1}{(1-z/z_c)^2} e^{(L+1)[f(z_c) + \frac{f''(z_c)}{2}(z-z_c)^2]} dz \end{aligned}$$

Take $z - z_c = \frac{it}{\sqrt{L+1}}$. Note $t \in (-\infty, -\epsilon\sqrt{L+1}) \cup (\epsilon\sqrt{L+1}, \infty) =: I_{\epsilon,L}$, so we see that we need ϵ to be small compared to $\sqrt{L+1}$. Take $\epsilon = \frac{1}{L}$.

$$\begin{aligned} \frac{1}{2\pi i} \int_C \frac{1}{(1-z/z_c)^2} e^{(L+1)(f(z_c) + \frac{f''(z_c)}{2}(z-z_c)^2)} dz \\ \approx -\frac{1}{2\pi} \int_{I_{\epsilon,L}} \frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1}}{t^2} e^{-\frac{f''(z_c)}{2} t^2} dt \\ = -\frac{1}{\epsilon\sqrt{L+1}} \frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1}}{\pi} e^{-\frac{f''(z_c)}{2} \epsilon^2 (L+1)} + \frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1} f''(z_c)}{2\pi} \int_{I_{\epsilon,L}} e^{-\frac{f''(z_c)}{2} t^2} dt \end{aligned}$$

where the second line is an approximation similar as in the approximation (d) in Eqn. (80), and the third line follows by integration by parts.

The integral of the small semicircle can also be computed. Take $z = z_c + \epsilon e^{i\theta}$.

$$\begin{aligned} \frac{1}{2\pi i} \int_{C_\epsilon} \frac{1}{(1-z/z_c)^2} V(z)^{L+1} dz &= \frac{1}{2\pi} \int_{3\pi/2}^{\pi/2} \frac{z_c^2}{\epsilon} e^{-i\theta} V(z_c + \epsilon e^{i\theta})^{L+1} d\theta \\ &\approx \frac{1}{2\pi} \int_{3\pi/2}^{\pi/2} \frac{z_c^2}{\epsilon} e^{-i\theta} V(z_c)^{L+1} d\theta \\ &= \frac{1}{\epsilon} \frac{z_c^2 V(z_c)^{L+1}}{\pi} \end{aligned}$$

The second line follows from Taylor expanding $V(z_c + \epsilon e^{i\theta})^{L+1}$ around its critical point z_c with $\epsilon = \frac{1}{L}$,

$$\begin{aligned} V(z_c + \epsilon e^{i\theta})^{L+1} &= V(z_c)^{L+1} + \frac{1}{2}(L+1)V(z_c)^L V''(z_c) \epsilon^2 e^{2i\theta} + O(L^2 \epsilon^3) \\ &= V(z_c)^{L+1} + \frac{1}{2} \frac{L+1}{L^2} V(z_c)^L V''(z_c) e^{2i\theta} + O\left(\frac{1}{L}\right) \\ &\approx V(z_c)^{L+1} \end{aligned}$$

Looking only at the two terms that are proportional to $\frac{1}{\epsilon}$, we see that

$$\begin{aligned} &-\frac{1}{\epsilon\sqrt{L+1}} \frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1}}{\pi} e^{-\frac{f''(z_c)}{2} \epsilon^2 (L+1)} + \frac{1}{\epsilon} \frac{z_c^2 V(z_c)^{L+1}}{\pi} \\ &= -\frac{1}{\epsilon} \frac{z_c^2 V(z_c)^{L+1}}{\pi} e^{-\frac{f''(z_c)}{2} \epsilon^2 (L+1)} + \frac{1}{\epsilon} \frac{z_c^2 V(z_c)^{L+1}}{\pi} \\ &\approx -\frac{1}{\epsilon} \frac{z_c^2 V(z_c)^{L+1}}{\pi} \left(1 - \frac{f''(z_c)}{2} \epsilon^2 (L+1)\right) + \frac{1}{\epsilon} \frac{z_c^2 V(z_c)^{L+1}}{\pi} \\ &= \frac{z_c^2 V(z_c)^{L+1}}{\pi} \frac{f''(z_c)}{2} \epsilon (L+1). \end{aligned}$$

Note that with $\epsilon = \frac{1}{L}$ this is $O(V(z_c)^{L+1})$ and will be small compared to the remaining term which we compute below.

The other term gives

$$\frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1} f''(z_c)}{2\pi} \int_{I_{\epsilon,L}} e^{-\frac{f''(z_c)}{2} t^2} dt = \frac{\sqrt{L+1} z_c^2 V(z_c)^{L+1} \sqrt{f''(z_c)}}{\sqrt{2\pi}}$$

as $\epsilon \rightarrow 0$. This is the main contribution. Noting that $f''(z_c) = \frac{V''(z_c)}{V(z_c)}$ gives the desired result. \square

11 Appendix B: Proof of Lemma 6.5

Recall that for $x, y \in \mathbb{Z}_{\geq 0}$, $L \in \mathbb{Z}_{> 1}$, $\mathcal{W}(x, y, L; M)$ is the set of walk bridges from x to y of length L , with increments and weights given by Eqn. (66), conditioning on staying nonnegative.

As a preparation, we first prove the following Gaussian tail bound about a walk bridge, that is away from its mean at a certain point in the middle.

Lemma 11.1. *Let $H, G \in \mathbb{Z}_{\geq 0}$, $L \in \mathbb{Z}_{\geq 2}$. Sample a walk $W(t)$ from $\mathcal{W}(H, G, L; M)$, and let take $L' \in \llbracket 1, L-1 \rrbracket$. If $H \geq G$, then there exists constants $C', c > 0$ uniformly in H, G, L, L' and M , such that for all $h > 0$,*

(a).

$$\mathbb{P}[W(L') > G + (H - G)(L - L')/L + h(L - L')^{\frac{1}{2}}] \leq C' \exp(-ch^2);$$

(b).

$$\mathbb{P}[W(L') < H - (H - G)L'/L - hL'^{\frac{1}{2}}] \leq C' \exp(-ch^2).$$

If $H < G$, then there exists constants $C', c > 0$ uniformly in H, G, L, L' and M , such that for all $h > 0$,

(c).

$$\mathbb{P}[W(L') > G + h(L - L')^{\frac{1}{2}}] \leq C' \exp(-ch^2);$$

(d).

$$\mathbb{P}[W(L') < H - hL'^{\frac{1}{2}}] \leq C' \exp(-ch^2).$$

Proof. We focus on the proof of (a), and (b) can be proved in a similar way by a rotation of 180 degree (the restriction that $W(t) \geq 0$ further decreases the tail probability).

To prove (a), we only give detailed argument of the case $W(t)$ sampled from $\mathcal{W}(H - G, 0, L; M)$, and to simplify the notation we replace $H - G$ by $H' \in \mathbb{Z}_{\geq 0}$. The tail probability of the general $G > 0$ situation can be bounded following the same path, and gives at most the same order. After the transformation, our goal becomes proving that

$$\mathbb{P}[W(L') > H'(L - L')/L + h(L - L')^{\frac{1}{2}}] \leq C' \exp(-ch^2).$$

Denote $W(L') = \tilde{H}$. It then suffices to show that

$$\frac{|\mathcal{W}(H', \tilde{H}, L'; M)| \cdot |\mathcal{W}(\tilde{H}, 0, L - L', M)|}{|\mathcal{W}(H', 0, L; M)|} \lesssim \frac{1}{\sqrt{\min(L', L - L')}} \exp\left(-c \left(\frac{\tilde{H}^2}{L - L'}\right)\right). \quad (175)$$

Let $\tilde{\mathcal{W}}(H', 0, L; M)$ be the set of walks with the same increments (whose weights change with M) as above, starting at H' , ending at 0 and has length L , without any further conditioning. Let $|\tilde{\mathcal{W}}(H', 0, L; M)|$ be the partition function of such walk bridge, which can be expressed as

$$\frac{1}{2\pi i} \oint \left(\frac{z}{z_c(N)}\right)^{H'+1} V_N(z)^{L+1} dz, \quad (176)$$

where the contour is around 0.

Using such expression, by a steepest descent similar as in Section 4.4, we have that for some constants $0 < c_1 < c_2$, $c > 0$ uniform in L, M , $\mathbf{H} \in \llbracket 0, L \rrbracket$,

$$\begin{aligned} \frac{|\tilde{\mathcal{W}}(\mathbf{H}, 0, L; M)|}{V_N(z_c(N))^L} &\leq c_2 \frac{1}{\sqrt{L}} \exp\left(-c \left(\frac{\mathbf{H}^2}{L}\right)\right), \\ c_1 \frac{1}{\sqrt{L}} \exp\left(-c \left(\frac{\mathbf{H}^2}{L}\right)\right) &\leq \frac{|\tilde{\mathcal{W}}(\mathbf{H}, 0, L; M)|}{V_N(z_c(N))^L} \leq c_2 \frac{1}{\sqrt{L}} \exp\left(-c \left(\frac{\mathbf{H}^2}{L}\right)\right), \quad \text{when } 0 \leq \mathbf{H} \leq g(L), \end{aligned} \quad (177)$$

where $g(\cdot) : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ can be any function such that $\frac{g(L)}{L} \rightarrow 0$ as $L \rightarrow \infty$. For some constant $C > 0$ uniform in $L_1, L_2, M \in \mathbb{Z}_{> 1}$,

$$\frac{|\tilde{\mathcal{W}}(\lfloor \mathbf{a}L_1 \rfloor + \mathbf{H}, 0, L_1; M)| \cdot |\tilde{\mathcal{W}}(\lfloor \mathbf{a}L_2 \rfloor - \mathbf{H}, 0, L_2; M)|}{|\tilde{\mathcal{W}}(\lfloor \mathbf{a}L_1 \rfloor, 0, L; M)| \cdot |\tilde{\mathcal{W}}(\lfloor \mathbf{a}L_2 \rfloor, 0, L; M)|} \leq C \frac{|\tilde{\mathcal{W}}(\lfloor \mathbf{b}L_1 \rfloor + \mathbf{H}, 0, L_1; M)| \cdot |\tilde{\mathcal{W}}(\lfloor \mathbf{b}L_2 \rfloor - \mathbf{H}, 0, L_2; M)|}{|\tilde{\mathcal{W}}(\lfloor \mathbf{b}L_1 \rfloor, 0, L; M)| \cdot |\tilde{\mathcal{W}}(\lfloor \mathbf{b}L_2 \rfloor, 0, L; M)|} \quad (178)$$

for any $0 \leq \mathbf{b} \leq \mathbf{a}$, $\lfloor bL_2 \rfloor - \mathbf{H} \geq 0$. Also note that

$$|\tilde{\mathcal{W}}(\mathbf{H}, 0, L; M)| = |\tilde{\mathcal{W}}(\mathbf{H} + \mathbf{H}', \mathbf{H}', L; M)| \quad (179)$$

for any $\mathbf{H}' \in \mathbb{Z}$. Also by Lemma 10.4 we have,

$$\begin{aligned} |\mathcal{W}(\tilde{H}, 0, L - L', M)| &= \frac{\tilde{H} + 1}{L - L' + 1} |\tilde{\mathcal{W}}(\tilde{H} + 1, 0, L - L' + 1, M)| \\ |\mathcal{W}(H', 0, L; M)| &= \frac{H' + 1}{L' + 1} |\tilde{\mathcal{W}}(H' + 1, 0, L + 1; M)|. \end{aligned}$$

We estimate $|\mathcal{W}(H', \tilde{H}, L'; M)|$ as follows. If $H' \geq \tilde{H}$, we have

$$\frac{|\mathcal{W}(H', \tilde{H}, L'; M)|}{V_N(z_c(N))^{L'}} = \sum_{m=0}^{\tilde{H}} \mathbb{P} \left[\min_{\llbracket 0, L' \rrbracket} W = m \right] \lesssim \frac{(H' + 1)(\tilde{H} + 1)}{L'} \frac{|\tilde{\mathcal{W}}(H' - \tilde{H}, 0, L'; M)|}{V_N(z_c(N))^{L'}}. \quad (180)$$

To see this, first observe that the bound follows directly from Eqn. (177) to (179) when $\tilde{H} \geq \frac{1}{2}\sqrt{L'}$, so next we consider the case $0 \leq \tilde{H} < \frac{1}{2}\sqrt{L'}$ (which implies $(\tilde{H} - m)^2 \leq \frac{1}{4}L'$). We split $W(t)$ as the concatenation of two walk bridges at $\tilde{\tau}$, the first time $W(t) = m$, where $m = \min_{\llbracket 0, L' \rrbracket} W \in \llbracket 0, \tilde{H} \rrbracket$, then we have

$$\mathbb{P} \left[\min_{\llbracket 0, L' \rrbracket} W = m \right] \sim \sum_{0 \leq \tilde{\tau} \leq L'} \frac{|\mathcal{W}(H' - m, 0, \tilde{\tau}; M)|}{V_N(z_c(N))^{\tilde{\tau}}} \frac{|\mathcal{W}(0, \tilde{H} - m, L' - \tilde{\tau}; M)|}{V_N(z_c(N))^{L' - \tilde{\tau}}}.$$

By Theorem 6.1, the maximal possible order of the second fraction is given when $L' - \tilde{\tau} \geq (\tilde{H} - l)^2$, and similarly the maximal possible order of the first fraction is given when $\tilde{\tau} \geq \min((H' - m)^2, 0.1L')$. Therefore we assume that $(\tilde{H} - m)^2 \leq L' - \tilde{\tau} \leq L' - \min((H' - m)^2, 0.1L')$. In this region, we have

$$\begin{aligned} \frac{|\mathcal{W}(H' - m, 0, \tilde{\tau}; M)|}{V_N(z_c(N))^{\tilde{\tau}}} &= \frac{|\mathcal{W}(H' - m, 0, \tilde{\tau}; M)|}{|\tilde{\mathcal{W}}(H' - m, 0, \tilde{\tau}; M)|} \frac{|\tilde{\mathcal{W}}(H' - m, 0, \tilde{\tau}; M)|}{V_N(z_c(N))^{\tilde{\tau}}} \lesssim \frac{H' - m}{\tilde{\tau}^{\frac{3}{2}}} \frac{|\tilde{\mathcal{W}}(H' - \tilde{H}, 0, L'; M)|}{V_N(z_c(N))^{L'}}, \\ &\frac{|\mathcal{W}(0, \tilde{H} - m, L' - \tilde{\tau}; M)|}{V_N(z_c(N))^{L' - \tilde{\tau}}} \lesssim \frac{\tilde{H} - m}{(L' - \tilde{\tau})^{\frac{3}{2}}}, \end{aligned} \quad (181)$$

where in both lines we use local limit theorem, see e.g [BD, Proposition 1]. The result then follows by summing over $\tilde{\tau}$ in the region and summing over \tilde{H} choices of m 's.

If $H' < \tilde{H}$, we claim that

$$\frac{|\mathcal{W}(H', \tilde{H}, L'; M)|}{V_N(z_c(N))^{L'}} \lesssim \frac{H' \tilde{H}}{(L')^{\frac{3}{2}}}. \quad (182)$$

To see this, first note that the bound is trivial when $H' > \sqrt{L'}$, so assume $H' \leq \sqrt{L'}$. Then we again split the walk at $\tilde{\tau}$, the stopping time that $W(t)$ reaches its minimum m , and restrict $\tilde{\tau}$ to a subset of $\llbracket 0, L' \rrbracket$ similar as above. For each such $\tilde{\tau}$, we have (by local limit theorem, see e.g [BD, Proposition 1])

$$\frac{|\mathcal{W}(H' - m, 0, \tilde{\tau}; M)|}{V_N(z_c(N))^{\tilde{\tau}}} \lesssim \frac{H' - m}{\tilde{\tau}^{\frac{3}{2}}}, \frac{|\mathcal{W}(0, \tilde{H} - m, L' - \tilde{\tau}; M)|}{V_N(z_c(N))^{L' - \tilde{\tau}}} \lesssim \frac{\tilde{H} - m}{(L' - \tilde{\tau})^{\frac{3}{2}}}$$

when $\tilde{H} - m \leq 100\sqrt{L' - \tilde{\tau}}$, but the second inequality is indeed easier to reach when $\tilde{H} - m \geq 100\sqrt{L' - \tilde{\tau}}$, since the right side grows with \tilde{H} , while the left side does not. So the bound holds for all \tilde{H} . Summing over $\tilde{\tau}$'s and m 's gives Eqn. (182).

Finally, we estimate the ratio

$$\frac{|\tilde{\mathcal{W}}(H' - \tilde{H}, 0, L'; M)| \cdot |\tilde{\mathcal{W}}(\tilde{H} + 1, 0, L - L' + 1, M)|}{|\tilde{\mathcal{W}}(H' + 1, 0, L + 1; M)|} \quad \text{or} \quad \frac{\left(\frac{|\tilde{\mathcal{W}}(\tilde{H} + 1, 0, L - L' + 1, M)|}{V_N(z_c(N))^{L - L' + 1}} \right)}{\left(\frac{|\tilde{\mathcal{W}}(H' + 1, 0, L + 1; M)|}{V_N(z_c(N))^{L + 1}} \right)},$$

depending on whether $H' \geq \tilde{H}$ or not, by Eqn. (177) to (179), and obtain the Gaussian tail bound $\leq \exp\left(-c\left(\frac{\tilde{H}-H'}{\sqrt{L-L'}}\right)^2\right)$ for both cases. Combining all of the above estimates for the left side of Eqn. (175) gives the desired bound.

Next we consider (c), and (d) is proved in a similar way. Similar as in (a), we reduce to the case that $W(t)$ is sampled from $\mathcal{W}(0, H', L; M)$, and would like to prove

$$\mathbb{P}[W(L') > H' + h(L - L')^{\frac{1}{2}}] \leq C' \exp(-ch^2).$$

When $L' \leq L - L'$, this is a weaker statement than Lemma 6.4 and therefore follows from there. When $L - L' < L'$, denote $W(L') = \tilde{H} > H'$, we use similar argument as in (a) to show that

$$\frac{|\mathcal{W}(0, \tilde{H}, L'; M)| \cdot |\mathcal{W}(\tilde{H}, H', L - L', M)|}{|\mathcal{W}(0, H', L; M)|} \leq \frac{1}{\sqrt{\min(L', L - L')}} \exp\left(-c\left(\frac{\tilde{H} - H'}{\sqrt{L - L'}}\right)^2\right). \quad (183)$$

More precisely, by local limit theorem (see e.g [BD, Proposition 1]) have

$$\frac{\left(\frac{|\mathcal{W}(0, \tilde{H}, L'; M)|}{V_N(z_c(N))^{L'}}\right)}{\left(\frac{|\mathcal{W}(0, H', L; M)|}{V_N(z_c(N))^L}\right)} \leq \frac{\tilde{H} + 1}{H' + 1}.$$

Moreover, as already shown above, we again have

$$\begin{aligned} \frac{|\tilde{\mathcal{W}}(\tilde{H}, H', L - L'; M)|}{V_N(z_c(N))^{L-L'}} &\leq \frac{1}{\sqrt{L - L'}} \exp\left(-c\left(\frac{\tilde{H} - H'}{\sqrt{L - L'}}\right)^2\right), \\ \frac{|\mathcal{W}(\tilde{H}, H', L - L'; M)|}{V_N(z_c(N))^{L-L'}} &\leq \frac{(H' + 1)(\tilde{H} + 1)}{(L - L')^{\frac{3}{2}}} \exp\left(-c\left(\frac{\tilde{H} - H'}{\sqrt{L - L'}}\right)^2\right). \end{aligned} \quad (184)$$

If $H' + 1 \geq \sqrt{L - L'}$, then $\frac{\tilde{H}+1}{H'+1} = 1 + \frac{\tilde{H}-H'}{H'+1} \leq \frac{\tilde{H}-H'}{\sqrt{L-L'}}$, which will be absorbed by the Gaussian tail, so we use the first bound above and simply bound $\frac{|\mathcal{W}(\tilde{H}, H', L-L'; M)|}{|\tilde{\mathcal{W}}(\tilde{H}, H', L-L'; M)|}$ by 1. If $H' + 1 < \sqrt{L - L'}$, we use the second bound above and multiply it with $\frac{\tilde{H}+1}{H'+1}$, and the factor $\frac{\tilde{H}^2}{L-L'} < (1 + \frac{\tilde{H}-H'}{\sqrt{L-L'}})^2$ is again absorbed by the Gaussian tail. So the result follows. \square

Now we finish the proof of Lemma 6.5. A similar statement to this lemma is proved for Bernoulli walks, see [GXZ, Lemma A.7]. For our walks, whose increment is unbounded in the upward direction, as we define V_i to be $W(X_i) - \min_{[X_i, Y_i]} W + 1$ instead of the larger quantity $\max_{[X_i, Y_i]} W - \min_{[X_i, Y_i]} W + 1$, the statement holds in a similar way (by expanding the fraction on the left side and using the tail bound below), when we also replace $|x - y|$ by $\max(x - y, 0)$, as long as we can still prove for all $h > 0$, there exists constants $C' > 0$, $c > 0$ uniformly in x, y, L, M , such that

$$\mathbb{P}[Z_i > h] \leq C' \exp(-ch^2)$$

where

$$Z_i = \frac{V_i - \max(x - y, 0)(Y_i - X_i)/L}{(Y_i - X_i)^{\frac{1}{2}}}.$$

Under the event $\mathcal{E} = \{Z_i > h\}$, one of the following must happen

$$\begin{aligned} \mathcal{E}_1 : & W(X_i) - W(Y_i) > \max(x - y, 0)(Y_i - X_i)/L + \frac{h}{2}(Y_i - X_i)^{\frac{1}{2}} - 1; \text{ or} \\ \mathcal{E}_2 : & \min_{[X_i, Y_i]} W(t) < \min(W(X_i), W(Y_i)) - \frac{h}{2}(Y_i - X_i)^{\frac{1}{2}}, \end{aligned}$$

since one can check that $\mathcal{E}_1^c \cap \mathcal{E}_2^c \subset \mathcal{E}^c$.

We first show that \mathcal{E}_2 happens with probability $\leq \exp(-ch^2)$. Conditioning on the boundary values $W(X_i)$ and $W(Y_i)$, we are sampling a walk from the set $\mathcal{W}(W(X_i), W(Y_i), Y_i - X_i; M)$. And the result follows directly from Lemma 6.4, since a rotation of 180 degree of the walk bridge turn taking minimum into taking maximum, and the restriction that $W(t) \geq 0$ in $\mathcal{W}(W(X_i), W(Y_i), Y_i - X_i; M)$ further decreases $\mathbb{P}[\mathcal{E}_2]$.

As for \mathcal{E}_1 , If $x - y \leq 0$, we further reduce the event \mathcal{E}_1 to the union of the following two events

$$\begin{aligned} \mathcal{E}'_1 &: W(X_i) - W(Y_i) > h(Y_i - X_i)^{\frac{1}{2}}/4 - 1; \text{ or} \\ \mathcal{E}'_2 &: W(X_i) - W(Y_i) > h(Y_i - X_i)^{\frac{1}{2}}/4 - 1. \end{aligned}$$

We consider the event \mathcal{E}'_1 , and \mathcal{E}'_2 can be dealt with using similar argument (by instead using Lemma 11.1(b),(d)). We would like to show that for any $W(Y_i)$, for a walk sampled from $\mathcal{W}(x, W(Y_i), Y_i; M)$,

$$\mathbb{P}[W(X_i) > W(Y_i) + h(Y_i - X_i)^{\frac{1}{2}}/4 - 1] \leq C' \exp(-ch^2).$$

When $x > W(Y_i)$, it follows from Lemma 11.1(a) that we have a stronger estimate

$$\mathbb{P}[W(X_i) > W(Y_i) + (x - W(Y_i))(Y_i - X_i)/Y_i + h(Y_i - X_i)^{\frac{1}{2}}/4 - 1] \leq C' \exp(-ch^2).$$

Hence the result holds. When $x \leq W(Y_i)$, this follows from Lemma 11.1(c). This again gives an upper bound of the tail probability.

If $x - y > 0$, we further reduce the event \mathcal{E}_1 to the union of the following four events

$$\begin{aligned} \mathcal{E}'_1 &: W(Y_i) \leq x, W(X_i) - W(Y_i) > (x - W(Y_i))(Y_i - X_i)/Y_i + h(Y_i - X_i)^{\frac{1}{2}}/4 - 1; \\ \mathcal{E}'_2 &: W(X_i) \geq y, W(X_i) - W(Y_i) > (W(X_i) - y)(Y_i - X_i)/Y_i + h(Y_i - X_i)^{\frac{1}{2}}/4 - 1; \\ \mathcal{E}'_3 &: W(Y_i) > x, W(X_i) - W(Y_i) > (x - y)(Y_i - X_i)/L + h(Y_i - X_i)^{\frac{1}{2}}/2 - 1; \\ \mathcal{E}'_4 &: W(X_i) < y, W(X_i) - W(Y_i) > (x - y)(Y_i - X_i)/L + h(Y_i - X_i)^{\frac{1}{2}}/2 - 1; \end{aligned}$$

For \mathcal{E}'_1 , conditioning on $W(Y_i) \leq x$, $\mathbb{P}[\mathcal{E}'_1|W(Y_i)]$ is bounded using Lemma 11.1(a). Similarly $\mathbb{P}[\mathcal{E}'_2|W(X_i)]$ is bounded using Lemma 11.1(b). For \mathcal{E}'_3 , $\mathbb{P}[\mathcal{E}'_3|W(Y_i)]$ is bounded by $\mathbb{P}[W(X_i) - W(Y_i) > h(Y_i - X_i)^{\frac{1}{2}}/2 - 1]$, where $W(t)$ is sampled from $\mathcal{W}(W(X_i), W(Y_i), Y_i; M)$, and the latter probability can be bounded by Lemma 11.1(c). Similarly $\mathbb{P}[\mathcal{E}'_4|W(X_i)]$ is bounded using Lemma 11.1(d). This finishes the proof. \square

References

- [A] J.P. Anker, An introduction to Dunkl theory and its analytic aspects. *Analytic, Algebraic and Geometric Aspects of Differential Equations*. Birkhauser, Cham (2017), 3-58.
- [Ahn] A. Ahn, Airy point process via supersymmetric lifts, *Probability and Mathematical Physics*, vol. 3, no. 4, Feb. 2023, pp. 869-938.
- [AH] A. Aggarwal, J. Huang, Edge Rigidity of Dyson Brownian Motion with General Initial Data. <https://arxiv.org/abs/2308.04236>
- [AN] T. Assiotis, J. Najnudel, The boundary of the orbital beta process, *Moscow Mathematical Journal*, 21(4), 659-694. arXiv:1905.08684
- [B1] F. Benaych-Georges, Rectangular random matrices, related convolution, *Probab. Theory Relat. Fields*, 144, 471-515, (2009). arXiv:math/0507336
- [B2] F. Benaych-Georges, Rectangular R-Transform as the Limit of Rectangular Spherical Integrals, *Journal of Theoretical Probability*, 24, 969-987, (2011). arXiv:0909.0178
- [BCG] F. Benaych-Georges, C. Cuenca, V. Gorin, Matrix Addition and the Dunkl Transform at High Temperature, *Communications in Mathematical Physics*, 394, 735-795, (2022). arXiv:2105.03795

- [Ber] J. Bertrand, Solution d'un problème, *Comptes Rendus de l'Académie des Sciences*, Paris 105 (1887), 369.
- [BD] A. Bryn-Jones, R. A. Doney, A functional limit theorem for random walk conditioned to stay non-negative, *J. London Math Soc.* (2), 74(1):244-258, 2006.
- [BP] J.-L. Baril, H. Prodinger. Enumeration of partial Lukasiewicz paths, *Enumerative Combinatorics and Applications*, 2022.
- [BR] D. Brennecken, M. Roesler, Limits of Bessel functions for root systems as the rank tends to infinity, arXiv:2401.02515.
- [BEY] P. Bourgade, L. Erdos, H.T. Yau . Edge universality of beta ensembles. *Commun. Math. Phys.* 332, 261–353, (2014).
- [CC] F. Caravenna, L. Chaumont, An invariance principle for random walk bridges conditioned to stay positive, *Electron. J. Probab.*, 18: 1-32 (2013).
- [CC2] F. Caravenna, L. Chaumont, Invariance principles for random walks conditioned to stay positive, *Ann. Inst. H. Poincaré Probab. Statist.*, 44 (1) 170 - 190, (2008).
- [CH] I. Corwin, A. Hammond, Brownian Gibbs property for Airy line ensembles, *Invent. Math.*, 195(2), 441-508 (2014).
- [D] I. Dumitriu, Eigenvalue statistics for beta-ensembles. *PhD thesis, Massachusetts Institute of Technology*, (2003).
- [DE] I. Dumitriu and A. Edelman. Matrix models for beta ensembles. *J. Math. Phys.*, 43(11):5830-5847, 2002.
- [DG] P. Deift and D. Gioev, Universality at the edge of the spectrum for unitary, orthogonal, and symplectic ensembles of random matrices, *Comm. Pure Appl. Math.* 60 (2007), no. 6, 867–910.
- [DM] B. Davis, D. MacDonald, An elementary proof of the local central limit theorem, *Journal of Theoretical Probability*, 8, 693–701 (1995).
- [DL] L. Dumaz, C. Labbé, The stochastic Airy operator at large temperature, *Annals of Applied Probability*, 32(6): 4481-4534 (2022).
- [DLV] L. Dumaz, Y. Li, B. Valko. Operator level hard-to-soft transition for β -ensembles. *Electronic Journal of Probability*, 2021(26):1-28, (2021).
- [Du] C. Dunkl, Differential-Difference Operators Associated to Reflection Groups. *Transactions of the American Mathematical Society*, 311, no. 1 (1989), 167-183.
- [DVe1] D. Daley, D. Vere-Jones, An introduction to the theory of point processes: Vol. I. Elementary theory and methods, Springer-Verlag, New York, 2003.
- [DVe2] D. Daley, D. Vere-Jones, An introduction to the theory of point processes: Vol. II. Elementary theory and methods, Springer-Verlag, New York, 2008.
- [EYY] L. Erdos, H.-T. Yau, and J. Yin, Universality for generalized Wigner matrices with Bernoulli distribution, *J. Comb.* 2 (2011), no. 1, 15–81.
- [GM] V. Gorin, A. W. Marcus, Crystallization of random matrix orbits. *International Mathematics Research Notices* 2020, no. 3 (2020), 883-913.
- [GXZ] V. Gorin, J. Xu, L. Zhang, Airy $_{\beta}$ line ensemble and its Laplace transform, (2024+).
- [Gri] A. Gribinski, A theory of singular values for finite free probability. arXiv:2208.09768 (2022).
- [GrM] A. Gribinski, A. Marcus, A rectangular additive convolution for polynomials, *Journal of Combinatorial Theory*. arXiv:1904.11552

- [GSh] V. Gorin, M. Shkolnikov, Stochastic Airy semigroup through tridiagonal matrices, *Annals of probability*, 46(4): 2287-2344 (July 2018)
- [GS] I.P. Goulden and L.G. Serrano, Maintaining the spirit of the reflection principle when the boundary line has arbitrary integer slope, *J. Comb. Theory, Series A* 104 (2003), 317-326.
- [Hel] S. Helgason, Groups and Geometric Analysis, *American Mathematical Society*, (2012).
- [JP] H. C. Ji, J. Park, Tracy-Widom limit for free sum of random matrices. arXiv:2110.05147
- [Ke] D. P. Kennedy, The distribution of the maximum Brownian excursion, *Journal of Applied Probability*, Volume 13, Issue 2, (1976), 371-376.
- [KM] O. Khorunzhiy, J. F. Marckert, Uniform bounds for exponential moment of maximum of a Dyck path, *Electron Communication of Probability*, 14: 327-333 (2009).
- [KN] R. Killip, I. Nenciu. Matrix models for circular ensembles, *International Mathematics Research Notices*, no. 50 (2004): 2665-2701.
- [KRV] M. Krishnapur, B. Rider, B. Virag, Universality of the stochastic Airy operator, *Communications on Pure and Applied Mathematics*, Volume 69, Issue 1, (2016), 145-199.
- [LR] M. Ledoux and B. Rider. Small deviations for beta ensembles, *Electronic Journal of Probability* 15 (2009): 1319-1343.
- [LY] J.-O. Lee and J. Yin, A necessary and sufficient condition for edge universality of Wigner matrices, *Duke Math. J.* 163 (2014), no. 1, 117–173.
- [Me] P. Mergny, An explicit example for the high temperature convolution: crossover between the binomial law $B(2,1/2)$ and the arcsine law. arXiv:2204.07744
- [MeP] P. Mergny, M. Potters, Rank one HCIZ at high temperature: interpolating between classical and free convolutions, *SciPost Physics*, 12, 022 (2022).
- [MSS] A. Marcus, D. Spielman, N. Srivastava. Finite free convolutions of polynomials, *Probab. Theory Relat. Fields* 182, 807–848 (2022). arXiv:1504.00350
- [Ne] Y. A. Neretin, Rayleigh triangles and non-matrix interpolation of matrix beta-integrals, *Sbornik: Mathematics*, Volume 194, Number 4 (2003).
- [No] J. Novak, Three lectures on free probability, *Random matrix theory, interacting particle systems, and integrable systems* 65.309-383 (2014): 13. arXiv:1205.2097
- [NS] A. Nica, R. Speicher, Lectures on the Combinatorics of Free Probability, *London Mathematical Society Lecture Note Series* 335 (2006).
- [OO] A. Okounkov, G. Olshanski, Shifted Jack polynomials, binomial formula, and applications. *Mathematical Research Letters*, 4 (1997), 69-78, arXiv:q-alg/9608020.
- [Op] E. M. Opdam, Dunkl operators, Bessel functions and the discriminant of a finite Coxeter group, *Compositio Mathematica*, vol. 85, 3, 333-373 (1993).
- [OV] G. Olshanski, A. Vershik, Ergodic unitarily invariant measures on the space of infinite Hermitian matrices, *Contemporary Mathematical Physics. F. A. Berezin's memorial volume. Amer. Math. Transl. Ser. 2*, vol. 175 (R. L. Dobrushin et al., eds), (1996), pp. 137-175. arXiv:math/9601215
- [Pa] S. Péché, Universality results for the largest eigenvalues of some sample covariance matrix ensembles, *Probab. Theory Relat. Fields*, 143, 481–516 (2009).
- [PaSh] L. Pastur and M. Shcherbina, On the edge universality of the local eigenvalue statistics of matrix models, *Mat. Fiz. Anal. Geom.* 10 (2003), no. 3, 335–365.
- [PS] S. Péché and A. Soshnikov, Wigner random matrices with non-symmetrically distributed entries, *J. Stat. Phys.* 129 (2007), no. 5-6, 857–884.

- [Ro] M. Roesler, Dunkl Operators: Theory and Applications. *Lecture Notes in Mathematics* vol. 1817, (2003), 93-136.
- [RRV] J. Ramirez, B. Rider, B. Virag, Beta ensembles, stochastic Airy spectrum, and a diffusion. *J. Amer. Math. Soc.* 24, 919–944.
- [S] J. Sohler, A functional limit convergence towards Brownian excursion, Preprint (2010), arXiv:1012.0118 [math.PR].
- [Sh] M. Shcherbina, Edge universality for orthogonal ensembles of random matrices, *J. Stat. Phys.* 136 (2009), no. 1, 35–50.
- [Sod] S. Sodin, Several applications of the moment method in random matrix theory, arXiv:1406.3410.
- [Sos] A. Soshnikov, Universality at the edge of the spectrum in Wigner random matrices, *Comm. Math. Phys.* 207, (1999), 697–733.
- [SS1] Y. Sinai, A. Soshnikov, Central limit theorem for traces of large random symmetric matrices with independent matrix elements, *Boletim da Sociedade Brasileira de Matemática-Bulletin/Brazilian Mathematical Society*, 29(1):1–24, (1998).
- [SS2] Y. Sinai, A. Soshnikov, A refinement of Wigner’s semicircle law in a neighborhood of the spectrum edge for random symmetric matrices, *Functional Analysis and Its Applications*, 32(2):114–131, (1998).
- [TV] T. Tao and V. Vu, Random matrices: Universality of local eigenvalue statistics up to the edge, *Comm. Math. Phys.* 298 (2010), 549–572.
- [Vo] D. Voiculescu, Limit laws for random matrices and free products, *Inventiones Mathematicae*, 104 (1991), 201-220.
- [Xu] J. Xu, Rectangular matrix additions in low and high temperatures. <https://arxiv.org/abs/2303.13812>