$egin{aligned} Research in \ A stronomy and \ A strophysics \end{aligned}$ 

# **Galaxy Specific Star Formation Rate Is Independent of Halo Spin**

Zichen Hua<sup>1,2</sup>, Yu Rong<sup>1,2\*</sup>

<sup>1</sup> Department of Astronomy, University of Science and Technology of China, Hefei, Anhui 230026, China; *rongyua@ustc.edu.cn* 

<sup>2</sup> School of Astronomy and Space Sciences, University of Science and Technology of China, Hefei 230026, Anhui, China

Received 20XX Month Day; accepted 20XX Month Day

**Abstract** Utilizing ALFALFA HI data, we investigate the relationship between specific star formation rate (sSFR) and halo spin across various star-forming galaxies. Our analysis reveals no significant correlation between sSFR and halo spin, irrespective of the galactic environment. Previous research suggests that high-spin halos tend to harbor extended, low-density stellar distributions due to suppressed gas cooling and star formation. However, unlike galaxy size and density, sSFR may primarily reflect the current star-forming state rather than long-term history, indicating potential independence from halo spin.

Key words: galaxies: evolution - galaxies: formation - methods: statistical

# **1 INTRODUCTION**

In the standard galaxy formation model, halo spin is considered pivotal in galaxy formation and evolution, influencing morphology and regulating baryonic fraction (Mo et al. 1998; Guo et al. 2011; Rong et al. 2017; Amorisco & Loeb 2016; Benavides et al. 2023; van den Bosch 1998; Diemand et al. 2005; Desmond et al. 2017). However, hydrodynamical simulations (e.g., Kim & Lee 2013; Jiang et al. 2019; Yang et al. 2023; Di Cintio et al. 2019) have sparked debate on the role of halo spin in low-mass galaxies. For most dwarf galaxies, stellar distributions may be independent of or weakly dependent on halo spin, with concentrations and baryonic feedback exerting more significant influence on low-mass galaxy properties. Halos with shallower potential wells and lower concentrations exhibit stronger feedback effects, leading to reduced star formation efficiencies, lower stellar mass fractions, and more extended stellar distributions (Kravtsov et al. 2018; Sales et al. 2022; Hopkins et al. 2012; Sawala et al. 2015). Despite extensive research, the influence of halo spin on galaxy structure and evolution remains incompletely understood and lacks consensus.

The star formation rate, a crucial parameter defining galaxy evolutionary state, is influenced by internal factors such as feedback, fuel supply, metallicity, and dust content (Springel 2000; Hjorth et al. 2014; Grudić et al. 2018; Hayward et al. 2011). Environmental factors also play a significant role in shaping the star formation rate (González & Padilla 2009; Tinker et al. 2017). Among the myriad of galaxy properties, the impact of halo properties on the star formation rate, while not direct, is paramount. Halos' mass and surface density have been shown to significantly affect star formation rates in galaxies (Dahlem et al. 2006; Kimm et al. 2009), yet the influence of halo spin on star formation rate remains uncertain. Rong et al. (2024a) propose a link between spin and star formation rate, suggesting that increased halo spin could hinder gas accretion and slow star formation processes. However, this proposed scenario requires further investigation.

HI surveys conducted with single-dish telescopes, such as the Arecibo Legacy Fast Alfa Survey (ALFALFA; Giovanelli et al. 2005; Haynes et al. 2018) and the undergoing FAST All Sky HI survey (FASHI; Zhang et al. 2024), provide valuable HI spectra from numerous star-forming galaxies. These surveys offer crucial dynamical information on galaxies, facilitating the estimation of spin parameters for galaxies with varying star formation rates and enabling the study of halo spin's impact on star formation rate.

In this study, we utilize a semi-analytic approach to estimate halo spin for each HI-bearing galaxy cataloged in ALFALFA and investigate the relationship between halo spins and stellar densities of galaxies. Section 2 introduces the sample data and outlines the methodology for estimat-

<sup>\*</sup> corresponding author

ing halo spin. Section 3 presents a statistical analysis of the dependence of galaxy stellar densities on halo spins. Our findings are summarized in section 4.

### 2 DATA

# 2.1 Sample and specific star formation rate

We draw our galaxy sample from ALFALFA, a comprehensive HI survey covering 6,600 deg<sup>2</sup> at high Galactic latitudes. The ALFALFA ( $\alpha$ .100; Haynes et al. 2018) catalog, catalog, released by Haynes et al. (2018), includes ~31,500 sources with radial velocities below 18,000 km s<sup>-1</sup>. For each source, the catalog provides the HI spectrum signal-to-noise ratio (SNR), cosmological distance, 50% peak width of the HI line ( $W_{50}$ ) corrected for instrumental effects, HI mass ( $M_{\rm HI}$ ), among other properties.

We match ALFALFA galaxies with MPA-JHU DR7 SDSS measurements to obtain star formation rates (SFRs) (Brinchmann et al. 2004) and stellar mass ( $M_{\star}$ ) based on photometric fits. The specific SFR (sSFR) for each galaxy is then calculated as sSFR = SFR/ $M_{\star}$  (yr<sup>-1</sup>). To focus on star-forming galaxies, we select those with log sSFR >  $\log(1/3t_{H(z=0)}) \approx -10.62$  (Jing et al. 2021), where  $t_{H(z=0)}$  represents the Hubble time at redshift 0.

#### 2.2 Rotation velocity and halo spin

The rotation velocity is given by  $V_{\rm rot} = W_{50}/2/\sin\phi$ , with inclination  $\phi$  of the HI disk estimated via the optical axis ratio b/a (from Durbala et al. 2020) and  $\sin\phi = \sqrt{(1-(b/a)^2)/(1-q_0^2)}$ , setting  $\phi = 90^\circ$  for  $b/a \le q_0$ . We assume  $q_0 \sim 0.2$  for massive galaxies and  $q_0 \sim$ 0.4 (Rong et al. 2024b) for low-mass galaxies ( $M_{\star} < 10^{9.5} M_{\odot}$ ). We exclude galaxies with  $\phi < 50^\circ$  or low SNR (< 10) to ensure accurate  $V_{\rm rot}$  measurements.

Some HI-bearing galaxies exhibit velocity dispersiondominated kinematics. These galaxies, identified by their HI line profiles exhibiting a 'single-horned' shape (ElBadry et al. 2018), pose challenges in accurately estimating rotation velocities and, consequently, halo spins. Following Hua et al. 2024, we employ the kurtosis parameter  $k_4 < -1.0$  to restrict our analysis to robust subsets of isolated galaxies with double-horned HI profiles to exclude potential contamination from dispersion-dominated systems.

Assuming an isothermal halo model with negligible baryonic gravitational influence, the halo spin parameter  $\lambda_h$  is estimated as (Hernandez et al. 2007):

$$\lambda_{\rm h} \simeq 21.8 \frac{R_{\rm HI,d}/\rm kpc}{(V_{\rm rot}/\rm km s^{-1})^{3/2}},$$
 (1)

where  $R_{\rm HI,d}$  is the HI disk scale length, derived from:

$$\Sigma_{\rm HI}(R) = \Sigma_{\rm HI,0} \exp(-R/R_{\rm HI,d}), \qquad (2)$$

where  $\Sigma_{\rm HI,0}$  is the central surface density of the HI disk. The total HI mass  $M_{\rm HI}$  is linked to the scale length as

$$M_{\rm HI} = 2\pi \Sigma_{\rm HI,0} R_{\rm HI,d}^2. \tag{3}$$

Additionally, we introduce the HI radius  $r_{\rm HI}$ , defined as the radius at which the HI surface density reaches  $1 \text{ M}_{\odot} \text{pc}^{-2}$ .  $r_{\rm HI}$  is calculated using the observed  $r_{\rm HI}$ - $M_{\rm HI}$ relation (Wang et al. 2016; Gault et al. 2021):  $\log r_{\rm HI} =$  $0.51 \log M_{\rm HI} - 3.59$  (Wang et al. 2016; Gault et al. 2021). Therefore, at  $r_{\rm HI}$ , we have

$$\Sigma_{\rm HI,0} \exp(-r_{\rm HI}/R_{\rm HI,d}) = 1 \,\mathrm{M}_{\odot} \mathrm{pc}^{-2}.$$
 (4)

By using equations (3) and (4), we can compute the value of  $R_{\rm HI,d}$  for each galaxy in our sample, thereby enabling the estimation of the halo spin.

# **3 RESULTS**

Panels a and b of Fig. 1 illustrate the sSFR-halo spin relation for low-mass  $(M_{\star} < 10^{9.5} M_{\odot})$  and high-mass  $(M_{\star} > 10^{9.5} M_{\odot})$  galaxies. Correlation analysis reveals no dependence of sSFR on halo spin, with correlation coefficients close to 0 (correlation coefficients  $\simeq 0.10$  and -0.04 for the low-mass and high-mass regimes, respectively).

Since environment also affects galaxy properties (e.g., Moore et al. 1996; Mayer et al. 2001; Mastropietro et al. 2005; Smith et al. 2015; Kazantzidis et al. 2011), we control for environmental influences by using the galaxy group catalog by Saulder et al. (2016), applying a friends-offriends algorithm to SDSS DR12 (Alam et al. 2015) and 2MASS Redshift Survey data (Huchra et al. 2012), adjusted for biases (e.g., Malmquist bias). Isolated galaxiesthose located more than three virial radii from galaxy groups-are examined again in panels c and d of Fig. 1, yielding similarly weak correlations between sSFR and  $\lambda_h$ (with correlation coefficients of 0.08 for the low-mass sample and -0.03 for the high-mass sample).

#### **4 CONCLUSION**

Using ALFALFA HI data, we examine sSFR-halo spin relationships for a range of star-forming galaxies, finding no significant correlation regardless of environment. Previous work suggests that high-spin halos host extended, lowdensity stellar distributions due to inhibited gas cooling and star formation (Rong et al. 2024a). However, unlike size and density, sSFR may reflect only the current starforming state rather than long-term history, implying independence from halo spin. Future simulation-based studies are necessary to verify this scenario.

Acknowledgements Y.R. acknowledges supports from the CAS Pioneer Hundred Talents Program (Category B), and the NSFC grant 12273037, as well as the USTC Research Funds of the Double First-Class Initiative.



Fig. 1 sSFR versus halo spin parameter for low-mass (left) and high-mass (right) galaxies. The top panels show the full star-forming galaxy sample, while the bottom panels focus on isolated galaxies. Median sSFR values with  $1\sigma$  error bars for each bin in  $\log \lambda_h$  are represented in blue and red, respectively. Best-fit linear trends are indicated by the corresponding lines.

# References

Alam, M. P. et al. 2015, ApJS, 219, 12 2

- Amorisco, N. C. & Loeb, A. 2016, MNRAS, 459, L51 1
- Benavides, J. A., Sales, L. V., Abadi, M. G., Marinacci, F., Vogelsberger, M., Hernquist, L. 2023, MNRAS, 522, 1033 1
- Brinchmann, J., Charlot, S., White, S. D. M., Tremonti, C., Kauffmann, G., Heckman, T., Brinkmann, J. 2004,

MNRAS, 351, 1151 2 Dahlem, M., Lisenfeld, U., Rossa, J. 2006, A&A, 457, 121

- Desmond, H., Mao Y.-Y., Wechsler R. H., Crain R. A., Schaye J. 2017, MNRAS, 471, L11 1
- Di Cintio, A., Brook, C. B., Macciò, A. V., Dutton, A. V., Cardona-Barrero, S. 2019, MNRAS, 486, 2535 1
- Diemand, J., Madau, P., Moore, B. 2005, MNRAS, 364, 367 1

- Durbala, A., Finn, R. A., Crone Odekon, M., Haynes, M. P., Koopmann, R. A., O'Donoghue, A. A. 2020, AJ, 160, 271 2
- ElBadry, K. et al. 2018, MNRAS, 473, 1930 2
- Gault, L. et al. 2021, AJ, 909, 19 2
- Giovanelli, R. et al. 2005, AJ, 130, 6 1
- González, R. E., Padilla, N. D. 2009, MNRAS, 397, 1498
- Grudić, M. Y., Hopkins, P. F., Faucher-Giguère, C.-A., Quataert, E., Murray, N., Kereš, D. 2018, MNRAS, 475, 3511 1
- Guo, Q., et al. 2011, MNRAS, 413, 101 1
- Haynes, M. P. et al. 2018, ApJ, 861, 49 1, 2
- Hayward, C. C., Kereš, D., Jonsson, P., Narayanan, D., Cox, T. J., Hernquist, L. 2011, ApJ, 743, 159 1
- Hernandez, X., Park, C., Cervantes-Sodi, B., & Choi, Y.-Y. 2007, MNRAS, 375, 163 2
- Hjorth, J., Gall, C., Michalowski, M. J. 2014, ApJ, 872, L23 1
- Hopkins, P. F., Quataert, E., Murray, N. 2012, MNRAS, 421, 3488 1
- Hua, Z., Rong, Y., Hu, H.-J. 2024, eprint arXiv:2403.16754 2
- Huchra, J. P. et al. 2012, ApJS, 199, 26 2
- Jiang, F., et al. 2019, MNRAS, 488, 4801 1
- Jing, Y.-J., Rong, Y., Wang, J., Guo, Q., Gao, L. 2021, RAA, 21, 218 2
- Kazantzidis, S., Lokas, E., Callegari, S., Mayer, L., Moustakas, L. 2011, ApJ, 726, 98 2
- Kim, J.-h. & Lee, J. 2013, MNRAS, 432, 1701 1
- Kimm, T., et al. 2009, MNRAS, 394, 1131 1
- Kravtsov, A. V., Vikhlinin, A. A., Astronomy Letters, 44, 8 1
- Mastropietro, C., Moore, B., Mayer, L., Debattista, V., Piffaretti, R., Stadel, J. 2005, MNRAS, 364, 607 2
- Mayer, L., Governato, F., Colpi, M., Moore, B., Quinn, T., Wadsley, J., Stadel, J., Lake, G. 2001, ApJ, 547, L123 2
- Mayer, L., Kazantzidis, S., Mastropietro, C., Wadsley, J. 2007, Nature, 445, 738
- Mo, H. J., Mao, S. D. & White, S. D. M. 1998, MNRAS, 295, 319 1
- Moore, B., Katz, N., Lake, G., et al. 1996, Nature, 379, 613 2
  - MNRAS, 431, 2307-2316
- Peng, Y.-J., Renzini, A. 2020, MNRAS, 491L, 51
- Rong, Y., Guo, Q., Gao, L., Liao, S., Xie, L., Puzia, T. H., Sun, S., Pan, J. 2017, MNRAS, 470, 4231 1
- Rong, Y., Hu, H., He, M., Du, W., Guo, Q. Wang, H.-Y., Zhang, H.-X., Mo, H. 2024a, arXiv:2404.00555 1, 2
- Rong, Y., He, M., Hu, H., Zhang, H.-X., Wang, H.-Y. 2024b, arXiv:2409.00944 2

- Sales, L. V., Wetzel, A., Fattahi, A. 2022, Nature Astronomy, 6, 897 1
- Saulder, C., van Kampen, E., Chilingarian, I. V., Mikske, S., Zeilinger, W. W. 2016, A&A, 596, A14 2
- Sawala, T., et al. 2015, MNRAS, 448, 2941 1
- Smith, R., Sánchez-Janssen, R., Beasley, M. A., et al. 2015, MNRAS, 454, 2502 2
- Springel, V. 2000, MNRAS, 312, 859 1
- Tinker, J. L., Wetzel, A. R., Conroy, C. 2017, MNRAS, 472, 2504 1
- van den Bosch, F. C. 1998, ApJ, 507, 601 1
- Wang, J., Koribalski, B. S., Serra, P., van der Hulst, T., Roychowdhury, S., Kamphuis, P., Chengalur, J. N. 2016, MNRAS, 460, 2143 2
- Yang, H., Gao, L., Frenk, C. S., Grand, R. J. J., Guo, Q., Liao, S., Shao, S. 2023, MNRAS, 518, 5253 1
- Zhang, C.-P., Zhu, M., Jiang, P., et al. 2024, Science China Physics, Mechanics & Astronomy, 67, 219511 1