Cost of controllability of the Burgers' equation linearized at a steady shock in the vanishing viscosity limit

Vincent Laheurte

ABSTRACT. We consider the one-dimensional Burgers equation linearized at a stationary shock, and investigate its null-controllability cost with a control at the left endpoint. We give an upper and a lower bound on the control time required for this cost to remain bounded in the vanishing viscosity limit, as well as a rough description of an admissible control. The proof relies on complex analysis and adapts methods previously used to tackle the same issue with a constant transport term.

CONTENTS

1. Introduction	1
1.1. Shock profiles for Burgers equation	1
1.2. Vanishing viscosity controllability problem	2
1.3. Control of the limit system	3
1.4. Related problems	4
1.5. Main results	4
2. Spectral analysis of the operator	5
3. Proof of the upper bound on T_{unif}	9
3.1. Scheme of the proof	9
3.2. Controllability of the first mode. Proof of Lemma 3.1	10
3.3. Controllability of the other modes. Proof of Lemma 3.2	11
4. Proof of the lower bound on T_{unif}	14
Appendix A. Construction of the bi-orthogonal family, proof of Lemma 3.4	16
References	18

1. Introduction

1.1. Shock profiles for Burgers equation. When considering the one-dimensional inviscid Burgers' equation

$$\begin{cases} \partial_t u(t,x) + u(t,x)\partial_x u(t,x) = 0, & t \in \mathbb{R}^+, x \in \mathbb{R}, \\ u(0,x) = u_0(x), & x \in \mathbb{R}, \end{cases}$$
(1)

shocks may arise in finite time regardless of the initial datum's regularity. Indeed, by the method of characteristics, if u_0 is at least C^1 and there exists some $x \in \mathbb{R}$ such that $u'_0(x) < 0$, two distinct characteristics will meet in time

$$T = -\frac{1}{\inf_{\substack{x \in \mathbb{R} \\ 1}} u_0'(x)},$$

which causes a jump discontinuity to appear in the solution u for times greater than T. Such shocks are jumps from u^- to u^+ , with $u^- > u^+$, and propagate, by the Rankine-Hugoniot condition, at a speed

$$s = \frac{u^- + u^+}{2}.$$

Stationary shocks therefore exist and are jump discontinuities from a positive value to its opposite. To study the shocks, we may restrain the equation to an interval [-L, L], L > 0 and impose suitable Dirichlet conditions on the endpoints:

$$\begin{cases} \partial_t u + u \partial_x u = 0, \\ u(t, -L) = 1, \\ u(t, L) = -1, \\ u(0, x) = u_0(x). \end{cases}$$
(2)

This system admits infinitely many stationary solutions, which are all of the form

$$U_{\sigma}(x) = \begin{cases} 1 & \text{if } x < \sigma, \\ -1 & \text{if } x > \sigma. \end{cases}$$

Moreover, if the initial datum u_0 has bounded variations, then the corresponding solution converges in finite time to one of these stationary solutions.

If the initial datum has low regularity, the system becomes ill-posed, as initial solutions with a jump discontinuity from a smaller to a greater value lead to infinitely many weak solutions. To avoid this issue and define the "right" unique solution, the approach first used by Hopf [10] is to add a small viscosity term $\varepsilon \partial_x^2$, which makes the system well-posed, and make ε tend to 0. We are therefore interested in the viscous system

$$\begin{cases} \partial_t u^{\varepsilon} + u^{\varepsilon} \partial_x u^{\varepsilon} = \varepsilon \partial_x^2 u^{\varepsilon}, \\ u^{\varepsilon}(t, -L) = 1, \\ u^{\varepsilon}(t, L) = -1. \end{cases}$$
(3)

Contrarily to the inviscid system (2), this one admits a unique stationary shock, which is given by

$$U^{\varepsilon}(x) = -\kappa \tanh\left(\frac{\kappa x}{2\varepsilon}\right),\tag{4}$$

where $\kappa > 0$ is such that $U^{\varepsilon}(\pm L) = \mp 1$.

The L^2 stability of such viscous shocks in the context of Burgers equation was first studied by Il'in and Oleinik [11] by a maximum principle, as well as Sattinger [23] via spectral analysis tools. We refer to [12,21,22] for L^2 stability results with other flux functions f(u), convex or non-convex.

We wish to study some controllability aspects of the viscous Burgers equation linearized at the shock profile (4).

1.2. Vanishing viscosity controllability problem. We study the left-side null-controllability of the viscous Burgers' equation linearized around a stationary shock. Namely, we set an arbitrary control time T > 0, and consider the control system

$$\begin{cases} \partial_t u^{\varepsilon} + \partial_x (U^{\varepsilon} u^{\varepsilon}) = \varepsilon \partial_x^2 u^{\varepsilon}, & x \in (-L, L), t \in (0, T), \\ u^{\varepsilon} (t, -L) = h(t), & t \in (0, T), \\ u^{\varepsilon} (t, L) = 0, & t \in (0, T), \\ u^{\varepsilon} (0, x) = u_0, & x \in (-L, L), \end{cases}$$
(5)

where the initial datum u_0 lies in $H_0^1 \cap H^2(-L, L)$, and the shock profile U^{ε} is given by

$$U^{\varepsilon}(x) = -\tanh\left(\frac{x}{2\varepsilon}\right),$$

where we ignore the κ from (4) as it does not affect the calculations in a meaningful way.

Given $\varepsilon > 0, T > 0$, the issue of null-controllability is, for any initial datum, to find a control $h \in L^2(0, T)$ such that the solution of (5) satisfies $u^{\varepsilon}(T) = 0$. If this is possible, we may define the null-controllability cost:

$$\mathcal{C}(T,\varepsilon) := \sup_{u_0 \in H_0^1 \cap H^2, \|u_0\|_{L^2} = 1} \inf \left\{ \|h\|_{L^2(0,T)} : u^{\varepsilon}(T) = 0 \right\}.$$

This type of problems has been deeply studied [5] and the system (5) can be proved to be null-controllable for any $\varepsilon > 0$ and T > 0. In this paper, we will tackle the issue of uniform null-controllability in the vanishing viscosity limit. Namely, we want to find the minimal time T_{unif} such that, for any $T > T_{\text{unif}}$, the controllability cost $C(T, \varepsilon)$ remains bounded as $\varepsilon \to 0$.

1.3. Control of the limit system. Formally taking $\varepsilon = 0$, the limit control problem we obtain can be written as

$$\begin{cases} \partial_t u - \partial_x (\operatorname{sgn}(x)u) = 0, \\ u(t, -L) = h(t), \\ u(t, L) = 0, \\ u(0, x) = 0. \end{cases}$$
(6)

By the method of characteristics, for any time t > 0, the solution u(t) of this system is given by

$$\forall x \in (-L,0), \quad u(t,x) = \begin{cases} u_0(x-t) & t < x+L \\ h(t-x-L) & t \ge x-L \end{cases}$$
(7)

$$\forall x \in (0, L), \quad u(t, x) = \begin{cases} u_0(x+t) & t < L - x \\ 0 & t \ge x - L \end{cases},$$
(8)

and the solution has a Dirac in x = 0 of mass

$$m(t) = \begin{cases} \int_{-t}^{t} u_0(x) \, \mathrm{d}s & t < L\\ \int_{-L}^{L} u_0(x) \, \mathrm{d}x + \int_{0}^{t-L} h(s) \, \mathrm{d}s & t \ge L \end{cases}$$
(9)

From this expression, one easily obtains that the system (6) is only null-controllable if T > L, and $u(T) \equiv 0$ if and only if the two following constraints on h are verified:

$$h(t) = 0, \quad t \ge T - L,$$
 (10)

$$\int_{0}^{T-L} h(t) \,\mathrm{d}t + \int_{-L}^{L} u_0(x) \,\mathrm{d}x = 0 \tag{11}$$

We then immediately conclude that the optimal control associated to an initial datum u_0 is given by

$$h(t) = \begin{cases} -\frac{1}{T-L} \int_{-L}^{L} u_0(x) \, \mathrm{d}x, & \text{if } t < T-L, \\ 0 & \text{if } t \ge T-L. \end{cases}$$
(12)

Applying Cauchy-Schwarz' inequality, the null-controllability cost of the limit system is therefore given by

$$\mathcal{C}(T,0) = \frac{\sqrt{2L}}{T-L} \tag{13}$$

It may then be expected that the viscous problem (5) is uniformly null-controllable for times T > L, and that its null-controllability cost converges to $\lim_{\varepsilon \to 0} C(T, \varepsilon) = C(T, 0) = \frac{\sqrt{2L}}{T-L}$

1.4. Related problems. This kind of problem was first addressed by Coron and Guerrero [3], where they consider a transport-diffusion system with a constant transport:

$$\begin{cases} \partial_t u + M \partial_x u = \varepsilon \partial_x^2 u, \\ u(t,0) = h(t), \\ u(t,L) = 0. \end{cases}$$
(14)

This system highlights different behaviors depending on the sign of the transport M. By using a dissipation argument and Carleman estimates, Coron and Guerrero [3] obtained a first upper-bound on uniform null-controllability time, which was later improved by Glass [8] through a method of moments, and by Lissy [18] with a reduction to the control of heat equation in short time. This result was later improved by Dardé-Ervedoza [4] with sharper estimates on the heat equation. In [19], Lissy also provided a non-trivial lower bound on the uniform null-controllability time, with complex-analytic tools.

Similar results were then obtained for non-constant transport terms M(t, x) in several space dimensions by Guerrero and Lebeau [9], only proving the existence of a uniform time, without any estimate on it. Laurent and Léautaud [15, 16] later provided an upper bound on the uniform control time in similar contexts.

In the nonlinear case, some partial results were given by Glass and Guerrero [7] for the Burgers equation, and extended by Léautaud [17] to other conservation laws, by first bringing the solution to the neighborhood of a traveling wave.

1.5. Main results. We state below the main results, giving the existence and constraints on the control time required for the null-controllability cost to remain bounded in the vanishing viscosity limit.

THEOREM 1.1. There exists a minimal time $T_{unif} < +\infty$ such that the system (5) is uniformly nullcontrollable for any time $T > T_{unif}$. Moreover, this minimal time verifies

$$T_{\text{unif}} \in \left[(4\sqrt{2} - 2); 4\sqrt{3} \right] L. \tag{15}$$

REMARK 1.2. The upper bound on the uniform null-controllability time matches the one obtained in [18] with M = 1, as we work on an interval of length 2L. The lower bound is however slightly worse than the one from [19] due to the behavior of the eigenfunctions of our system.

We also provide a description of some admissible control function h for times large enough.

THEOREM 1.3. Let $T > T^* = 4\sqrt{3}L$. Then, there exists C > 0 such that, for any $u_0 \in H_0^1 \cap H^2$, $\varepsilon > 0$, there exists a control function $h_{u_0}^{\varepsilon} \in L^2(0,T)$ such that the solution u^{ε} of (5) satisfies $u^{\varepsilon}(T) = 0$ and

$$\|h_{u_0}^{\varepsilon}\|_{L^2(0,T)} \le \left(\frac{2\sqrt{2L}}{T - T^*} + Ce^{-\frac{C}{\varepsilon}}\right) \|u_0\|_{L^2(-L,L)}.$$
(16)

As ε goes to 0, this control function converges in $L^2(0,T)$ towards the limit control

$$h_{u_0}^0(t) = \begin{cases} -\frac{2}{T-T^*} \int_{-L}^{L} u_0(x) \, \mathrm{d}x, & t \le \frac{T-T^*}{2}, \\ 0 & t > \frac{T-T^*}{2}. \end{cases}$$
(17)

The proof of $T_{\text{unif}} \leq 4\sqrt{3}L$ and of (16) is given in Section 3.1. It uses two intermediate results which are proved, respectively in Section 3.2 and in Section 3.3. The proof of $T_{\text{unif}} \geq (4\sqrt{2} - 2)L$ is given in Section 4. These results rely on a the spectral analysis of the operator at stake in the system (5) which is performed in Section 2. The proof of Theorem (1.3) is done within the proof of the upper bound, as we give a constructive proof of the control cost. More precisely, the estimate (16) is proven in Section 48, and the limit behavior (17) is given in (56) and (77).

REMARK 1.4. The admissible control built in Theorem 1.3 has the same structure as the optimal control of the limit system (12). It morally cancels out the mean of the solution u^{ε} in an arbitrarily small time and then exploits the strong dissipation of the system.

It is currently unclear whether these results may improve or prove further controllability results in the nonlinear case, extending the work of [7], or if similar results can be obtained for other conservation laws or more generally for systems with other singularities.

2. Spectral analysis of the operator

We are interested in the eigenvalues and eigenfunctions of the linearized Burgers operator

$$\mathcal{L}^{\varepsilon}u(x) := -\partial_x \left(\tanh\left(\frac{x}{2\varepsilon}\right) u(x) \right) - \varepsilon \partial_x^2 u(x).$$

This operator, among other viscous conservation laws, has been studied in [14], with a deeper look at the metastability phenomenon in [6, 20]. In these papers, the authors highlighted that the eigenvalues $(\lambda_k^{\varepsilon})_{k\geq 0}$ of the operator $\mathcal{L}^{\varepsilon}$ are simple, real, positive, and are distributed as follows:

$$\lambda_0^{\varepsilon} = \mathcal{O}\left(\exp\left(-\frac{1}{\varepsilon}\right)\right), \quad \lambda_k^{\varepsilon} > \frac{1}{4\varepsilon}, k \ge 1.$$

The first notable difference with the operator of the Coron-Guerrero system (14) is the presence of an exponentially small eigenvalue. The results from [3, 8] make use of the strong dissipation of the system. If we wish to use a similar approach, we need to treat the term corresponding to the first eigenvalue separately. Moreover, In order to apply the method of moments, we need sharper estimates regarding the distribution of the eigenvalues, as well as some information about the eigenfunctions. We obtain the following result:

LEMMA 2.1. The eigenvalues $(\lambda_k^{\varepsilon})_{k\geq 0}$ of the operator $\mathcal{L}^{\varepsilon}$ acting on $H_0^1 \cap H^2(-L, L)$ verify the following estimates:

$$\exists C > 0: 0 < \lambda_0^{\varepsilon} < Ce^{-\frac{L}{2\varepsilon}},\tag{18}$$

$$\frac{1}{4\varepsilon} + k^2 \frac{\pi^2 \varepsilon}{4L^2} < \lambda_k^{\varepsilon} < \frac{1}{4\varepsilon} + (k+1)^2 \frac{\pi^2 \varepsilon}{4L^2}, \quad k \ge 1$$
(19)

$$|\lambda_k^{\varepsilon} - \lambda_j^{\varepsilon}| \ge |k^2 - j^2| \frac{\varepsilon \pi^2}{4L^2}, \quad j,k \ge 1.$$
⁽²⁰⁾

Moreover, the associated eigenfunctions $(\psi_k^{\varepsilon})_{k\geq 0}$ of $(\mathcal{L}^{\varepsilon})^*$ are such that

$$\frac{\|\psi_0^{\varepsilon}\|_{L^2(-L,L)}}{|\varepsilon(\psi_0^{\varepsilon})'(-L)|} \le 2\sqrt{2L},\tag{21}$$

$$\frac{\|\psi_k^{\varepsilon}\|_{L^2(-L,L)}}{|\varepsilon(\psi_k^{\varepsilon})'(-L)|} \le \frac{4L}{k\pi\sqrt{\varepsilon}}, \quad k \ge 1.$$
(22)

More precisely, the first eigenfunction satisfies, as $\varepsilon \to 0$,

$$\frac{\psi_0^k}{\varepsilon(\psi_0^k)'(-L)} \xrightarrow{L^2(-L,L)} 1 \tag{23}$$

Above the operator $(\mathcal{L}^{\varepsilon})^*$ denotes the adjoint operator of $\mathcal{L}^{\varepsilon}$ with homogeneous Dirichlet condition, so that an eigenfunction of $(\mathcal{L}^{\varepsilon})^*$ satisfies the eigenvalue problem

$$\begin{cases} (\mathcal{L}^{\varepsilon})^{*}\psi^{\varepsilon} = \lambda^{\varepsilon}\psi^{\varepsilon}, \\ \psi^{\varepsilon}(\pm L) = 0, \end{cases}$$
(24)

REMARK 2.2. The distribution of the eigenvalues (19) and the information about the gap (20) are important in the computations later, and are not directly obtainable to our knowledge by standard methods such as perturbation tools, Sturm-Liouville theory or min-max formulation. We also insist on the fact that the eigenvalues of our operator $(\mathcal{L}^{\varepsilon})^*$ strongly resemble the ones of the Coron-Guerrero operator, for $M = \pm 1$ and an interval of length 2L. PROOF. We proceed in four steps.

Step 1. Reduction to an self-adjoint operator. Following [6, 15] we start by observing that the operator $\mathcal{L}^{\varepsilon}$ is conjugated to the self-adjoint operator:

$$P^{\varepsilon} := -\varepsilon^2 \partial_x^2 - \frac{1}{4} + \frac{1}{2} \tanh^2 \left(\frac{x}{2\varepsilon}\right),$$
$$\mathcal{L}^{\varepsilon} = \operatorname{sech}\left(\frac{x}{2\varepsilon}\right) \left(\frac{1}{\varepsilon} P^{\varepsilon}\right) \cosh\left(\frac{x}{2\varepsilon}\right), \tag{25}$$

Therefore the eigenvalue problem (24) can then be reduced to the simpler problem

$$\begin{cases} \frac{P^{\varepsilon}}{\varepsilon}\varphi^{\varepsilon} = \lambda^{\varepsilon}\varphi^{\varepsilon}, \\ \varphi^{\varepsilon}(\pm L) = 0, \end{cases}$$
(26)

by taking

$$\varphi^{\varepsilon}(x) = \operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\psi^{\varepsilon}.$$

Step 2. Reduction to a constant potential. As seen in [1, Proposition 3.1], the operator P^{ε} can be factored under the form

$$P^{\varepsilon} = (a^{\varepsilon})^* a^{\varepsilon},$$

where the operator a^{ε} is given by

$$a^{\varepsilon} := \varepsilon \partial_x + \frac{1}{2} \tanh\left(\frac{x}{2\varepsilon}\right).$$

Moreover, switching a^{ε} and its conjugate gives the much simpler operator

$$a^{\varepsilon}(a^{\varepsilon})^* = -\varepsilon^2 \partial_x^2 + \frac{1}{4}.$$

The price we have to pay is that the boundary condition is no longer the Dirichlet condition but rather a Robin condition. Indeed we now have the equivalent eigenvalue problem

$$\begin{cases} \left(-\varepsilon^2 \partial_x^2 + \frac{1}{4}\right) f^{\varepsilon} = \varepsilon \lambda^{\varepsilon} f^{\varepsilon}, \\ (a^{\varepsilon})^* f^{\varepsilon}(\pm L) = 0. \end{cases}$$
(27)

If the eigenpair $(\lambda^{\varepsilon}, f^{\varepsilon})$ solves the system (27), then the pair $(\lambda^{\varepsilon}, (a^{\varepsilon})^* f^{\varepsilon})$ solves (26). Conversely, a pair $(\lambda^{\varepsilon}, \varphi^{\varepsilon})$ solving (26) leads to a solution to system (27) by taking $(\lambda^{\varepsilon}, a^{\varepsilon}\varphi^{\varepsilon})$. In the sequel we distinguish two cases.

Step 3. An exponentially small eigenvalue.

If an eigenvalue satisfies $\lambda_{\varepsilon} < \frac{1}{4\varepsilon}$, then there exists $A, B \in \mathbb{R}$ such that the associated eigenfunction f^{ε} is of the form

$$f^{\varepsilon}(x) = A \cosh\left(\frac{\sqrt{\frac{1}{4}} - \varepsilon\lambda^{\varepsilon}}{\varepsilon}x\right) + B \sinh\left(\frac{\sqrt{\frac{1}{4}} - \varepsilon\lambda^{\varepsilon}}{\varepsilon}x\right)$$

The boundary condition $(a^{\varepsilon})^* f^{\varepsilon}(\pm L) = 0$ then translates to

$$A = 0, \quad \sqrt{\frac{1}{4} - \varepsilon \lambda^{\varepsilon}} = \frac{1}{2} \tanh\left(\frac{L}{2\varepsilon}\right) \tanh\left(\frac{\sqrt{\frac{1}{4} - \varepsilon \lambda^{\varepsilon}}}{\varepsilon}L\right).$$
(28)

By a concavity argument, this constraint admits two solutions in the interval $(-\infty, \frac{1}{4\varepsilon}]$: $\frac{1}{4\varepsilon}$, which we discard as it does not give any nonzero function f^{ε} , and some $\lambda_0^{\varepsilon} \in [0, \frac{1}{4\varepsilon})$.

by

Rewriting the condition (28) immediately gives

$$4\varepsilon\lambda_0^{\varepsilon} = 1 - \tanh^2\left(\frac{L}{2\varepsilon}\right) \tanh^2\left(\frac{\sqrt{\frac{1}{4}} - \varepsilon\lambda_0^{\varepsilon}}{\varepsilon}L\right),\,$$

and performing rough upper bounds quickly yields

$$4\varepsilon\lambda_0^{\varepsilon} \le 2\left(-\exp\left(\frac{L}{2\varepsilon}\right) + \exp\left(-\frac{\sqrt{\frac{1}{4}-\varepsilon\lambda_0^{\varepsilon}}}{\varepsilon}L\right)\right).$$

We note that λ_0^{ε} must be small, in particular $\lambda_0^{\varepsilon} \leq \frac{3}{16\varepsilon}$, and immediately conclude that the first eigenvalue of $\mathcal{L}^{\varepsilon}$ verifies

$$0 < \lambda_0^{\varepsilon} < \frac{C}{\varepsilon} e^{-\frac{L}{2\varepsilon}}.$$

The corresponding eigenfunction is given by

$$f_0^{\varepsilon}(x) = \sinh\left(\frac{\sqrt{\frac{1}{4} - \varepsilon\lambda_0^{\varepsilon}}}{\varepsilon}x\right).$$
(29)

Therefore the corresponding eigenfunction of P^{ε} is

$$\varphi_0^{\varepsilon} = (a^{\varepsilon})^* f_0^{\varepsilon} = -\sqrt{\frac{1}{4} - \varepsilon \lambda_0^{\varepsilon}} \cosh\left(\frac{\sqrt{\frac{1}{4} - \varepsilon \lambda_0^{\varepsilon}}}{\varepsilon}x\right) + \frac{1}{2} \tanh\left(\frac{x}{2\varepsilon}\right) \sinh\left(\frac{\sqrt{\frac{1}{4} - \varepsilon \lambda_0^{\varepsilon}}}{\varepsilon}x\right). \tag{30}$$

Finally the corresponding eigenfunction for $(\mathcal{L}^{\varepsilon})^*$ is $\psi_0^{\varepsilon}(x) = \cosh\left(\frac{x}{2\varepsilon}\right)\varphi_0^{\varepsilon}(x)$. Direct estimates then give the upper bound

$$\|\psi_0^{\varepsilon}\|_{L^2(-L,L)} \le \frac{\sqrt{2L}}{2},\tag{31}$$

while taking the derivative in x = -L gives the lower bound

$$|\varepsilon\left(\psi_{0}^{\varepsilon}\right)'\left(-L\right)| \geq \frac{1}{4}.$$
(32)

Combining the bounds (31) and (32) then gives the desired estimates (21). We may refine this result in the limit $\varepsilon \to 0$, and obtain

$$-2\varepsilon(\psi_0^k)'(-L) \to 1, \tag{33}$$

$$-2\psi_0^k \xrightarrow{L^2(-L,L)} 1. \tag{34}$$

This finally yields (23).

Step 4. The other eigenvalues. Now, let us consider eigenvalues verifying $\lambda^{\varepsilon} \geq \frac{1}{4\varepsilon}$. Then an associated eigenfunction can be written as

$$f^{\varepsilon}(x) = A \cos\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right) + B \sin\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right).$$

The boundary condition then leads to one of the two following constraints:

$$\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}} \sin\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}L\right) - \frac{1}{2} \tanh\left(\frac{L}{2\varepsilon}\right) \cos\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}L\right) = 0, \quad (35)$$

or

$$\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}} \cos\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}L\right) + \frac{1}{2} \tanh\left(\frac{L}{2\varepsilon}\right) \sin\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}L\right) = 0.$$
(36)

We introduce the unknown

$$\theta^{\varepsilon} := \arctan\left(\frac{\sqrt{4\varepsilon\lambda^{\varepsilon}-1}}{\tanh\left(\frac{L}{2\varepsilon}\right)}\right).$$

Constraints (35) and (36) then read

$$\sin\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}L - \theta^{\varepsilon}\right) = 0 \quad \text{or} \quad \cos\left(\frac{\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}}}{\varepsilon}L - \theta^{\varepsilon}\right) = 0.$$
(37)

Therefore $\lambda^{\varepsilon} > \frac{1}{4\varepsilon}$ is an eigenvalue if and only if it satisfies

$$\sqrt{\varepsilon\lambda^{\varepsilon} - \frac{1}{4}} = \frac{\varepsilon}{L} \left(k\frac{\pi}{2} + \theta^{\varepsilon} \right), \quad \text{for some } k \ge 1.$$
(38)

The constraint (38) has a unique solution λ_k^{ε} for each k. Moreover, since $\theta^{\varepsilon} \in (0, \frac{\pi}{2})$ and θ^{ε} is an increasing function of λ^{ε} , we immediately deduce the properties (19) and (20). We distinguish two cases.

• If $k \ge 1$ is even, the constraint (35) holds and the associated eigenfunction of $a^{\varepsilon}(a^{\varepsilon})^*$ is

$$f_k^{\varepsilon}(x) = \cos\left(\frac{\sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right).$$

The corresponding eigenfunction for P^{ε} is then given by $\varphi_k^{\varepsilon}(x) = (a^{\varepsilon})^* f_k^{\varepsilon}(x)$, that is

$$\varphi_k^{\varepsilon}(x) = -\sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}}\sin\left(\frac{\sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right) + \frac{1}{2}\tanh\left(\frac{L}{2\varepsilon}\right)\cos\left(\frac{\sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right).$$
(39)

Finally the eigenfunction for $\mathcal{L}^{\varepsilon}$ is obtained by taking $\psi_k^{\varepsilon}(x) = \cosh\left(\frac{x}{2\varepsilon}\right)\varphi_k^{\varepsilon}(x)$. Direct estimates then give the upper bound

$$\|\psi_k^{\varepsilon}\|_{L^2(-L,L)} \le \varepsilon \sqrt{\lambda_k^{\varepsilon}} \cosh\left(\frac{L}{2\varepsilon}\right),\tag{40}$$

as well as the lower bound on the derivative

$$|(\psi_k^{\varepsilon})'(-L)| \ge \frac{1}{2}\sqrt{\lambda_k^{\varepsilon}}\sqrt{\lambda_k^{\varepsilon} - \frac{1}{4\varepsilon}}\cosh\left(\frac{L}{2\varepsilon}\right).$$
(41)

Combining (40) and (41) leads to the desired estimate (22).

• Likewise, when k is odd, we instead get the following formula for the eigenfunction φ_k^{ε} of P^{ε} :

$$\varphi_k^{\varepsilon}(x) = \sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}} \cos\left(\frac{\sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right) + \frac{1}{2} \tanh\left(\frac{L}{2\varepsilon}\right) \sin\left(\frac{\sqrt{\varepsilon\lambda_k^{\varepsilon} - \frac{1}{4}}}{\varepsilon}x\right).$$

The same analysis leads again to the estimate (22) and the proof of Lemma 2.1 is complete.

8

REMARK 2.3. The information on λ_0^{ε} is not new and does not improve the one obtained by [14] by using an eigenfunction of $\mathcal{L}^{\varepsilon}$ acting on $H^2(\mathbb{R})$ which is exponentially small in $\pm L$, yet we provide an alternative proof, as the Darboux transformation, see Step 2, allows us to do so, and gives an exact expression of the associated eigenfunction.

3. Proof of the upper bound on T_{unif}

3.1. Scheme of the proof. As mentioned in Section 2, we control the solution of system (5) to zero in two steps, splitting the imparted time interval (0, T) into two parts $(0, \tau)$ and (τ, T) , for an intermediate time $\tau \in (0, T)$ which will be determined later, see (47). In a first time, we eliminate the first mode of u^{ε} , namely its projection on the eigenfunction ψ_0^{ε} . Then, we control the rest by exploiting the strong dissipation, and concluding with the method of moments, similarly to the work of O. Glass in [8] and P. Lissy in [18]. This is the object of the two following preliminary lemmas. Recall that the functions $(\psi_k^{\varepsilon})_k$ are the eigenfunctions of the operator $(\mathcal{L}^{\varepsilon})^*$ defined in (30) and (39).

LEMMA 3.1. Let T > 0, $\tau \in (0,T)$. For any $\varepsilon > 0$ and any initial datum $u_0 \in H^2 \cap H_0^1$, there is a control function $h_1 \in L^2(0,\tau)$ verifying:

$$\|h_1\|_{L^2(0,\tau)} \le \frac{2\sqrt{2L}}{\tau} \|u_0\|_{L^2(-L,L)},\tag{42}$$

such that the solution u^{ε} of (5) satisfies at time τ , for any $k \in \mathbb{N}$,

$$\langle u^{\varepsilon}(\tau), \psi_{k}^{\varepsilon} \rangle = e^{-\lambda_{k}^{\varepsilon}\tau} \langle u_{0}, \psi_{k}^{\varepsilon} \rangle - \frac{(\psi_{k}^{\varepsilon})'(-L)}{(\psi_{0}^{\varepsilon})'(-L)} \cdot \frac{e^{-\lambda_{0}^{\varepsilon}\tau} \langle u_{0}, \psi_{0}^{\varepsilon} \rangle}{\tau} \int_{0}^{\tau} e^{-(\lambda_{k}^{\varepsilon} - \lambda_{0}^{\varepsilon})t} \,\mathrm{d}t, \tag{43}$$

where the brackets $\langle \cdot, \cdot \rangle$ denote the inner product in $L^2(-L, L)$. In particular, for k = 0,

$$\langle u^{\varepsilon}(\tau), \psi_0^{\varepsilon} \rangle = 0. \tag{44}$$

In the following lemma we continue from the state $u^{\varepsilon}(\tau)$ as a new initial data for the system (5) now considered on the time interval (τ, T) .

LEMMA 3.2. Let $T > 0, \tau \in (0, T)$ with

$$T - \tau > T^* = 4\sqrt{3L}.$$
 (45)

Then there exists C > 0 such that for any $u_0 \in H^2 \cap H^1_0$, for any $\varepsilon > 0$, there exists a control function $h_2 \in L^2(\tau, T)$ with

$$\|h_2\|_{L^2(\tau,T)} \le C e^{-\frac{C}{\varepsilon}} \|u_0\|_{L^2(-L,L)}$$
(46)

such that the solution u^{ε} of (5) on the time interval (τ, T) starting at time τ with the initial data $u^{\varepsilon}(\tau)$ given by the formula (43), for $k \in \mathbb{N}$, satisfies $u^{\varepsilon}(T) = 0$.

The proofs of Lemma 3.1 and 3.2 are respectively given in the next two sections. Let us take them as granted for the moment and conclude the proof of the upper bound on T_{unif} , that is $T_{\text{unif}} \leq 4\sqrt{3}L$ and (16). We fix a time $T > T^*$, and set

$$\tau = \frac{T - T^*}{2},\tag{47}$$

so that $T - \tau > T^*$. Let $\varepsilon > 0, u_0 \in H_0^1 \cap H^2$. We obtain our control $h \in L^2(0,T)$ by concatenating the two controls h_1, h_2 built in Lemmas 3.1 and 3.2, namely

$$h(t) := \begin{cases} h_1(t) & \text{if } 0 < t < \tau, \\ h_2(t) & \text{if } \tau < t < T. \end{cases}$$

This choice of control steers the initial datum u_0 to the final state $u^{\varepsilon}(T) = 0$. Moreover such a control h verifies

$$\|h\|_{L^{2}(0,T)} \leq \|h_{1}\|_{L^{2}(0,\tau)} + \|h_{2}\|_{L^{2}(\tau,T)}$$

$$\leq \left(\frac{2\sqrt{2L}}{T - T^{*}} + Ce^{-\frac{C}{\varepsilon}}\right) \|u_{0}\|_{L^{2}(-L,L)},$$
(48)

which proves the uniform controllability time verifies $T_{\text{unif}} \leq T^*$, and leads to (16).

3.2. Controllability of the first mode. Proof of Lemma 3.1. We begin with a preliminary lemma exploiting duality equalities to describe the behavior of each mode of the solution.

LEMMA 3.3. Let $\varepsilon > 0$, $t_1, t_2 \in [0, T]$, and $k \ge 0$. The evolution of the k-th mode of the solution u^{ε} of the control problem (5) is prescribed by the relation:

$$\langle u^{\varepsilon}(t_2), \psi_k^{\varepsilon} \rangle = e^{-\lambda_k^{\varepsilon}(t_2 - t_1)} \langle u^{\varepsilon}(t_1), \psi_k^{\varepsilon} \rangle + \varepsilon(\psi_k^{\varepsilon})'(-L) \int_{t_1}^{t_2} e^{-\lambda_k^{\varepsilon}(t - t_1)} h(t_1 + t_2 - t) \, \mathrm{d}t.$$
(49)

PROOF. We introduce $\zeta_k^{\varepsilon}(t, x)$ as the solution to the adjoint uncontrolled system

$$\begin{cases} \partial_t \zeta_k^{\varepsilon} + (\mathcal{L}^{\varepsilon})^* \zeta_k^{\varepsilon} = 0, \\ \zeta_k^{\varepsilon}(t, -L) = \zeta_k^{\varepsilon}(t, L) = 0, \\ \zeta_k^{\varepsilon}(t_1, x) = \psi_k^{\varepsilon}(x). \end{cases}$$
(50)

Using the definition of the operators $\mathcal{L}^{\varepsilon}, (\mathcal{L}^{\varepsilon})^*$ and integrating by parts, it holds

$$\frac{\mathrm{d}}{\mathrm{d}t}\langle u^{\varepsilon}(t), \zeta_k^{\varepsilon}(t_1+t_2-t)\rangle = \varepsilon \partial_x \zeta_k^{\varepsilon}(t_1+t_2-t,-L)u^{\varepsilon}(t,-L).$$
(51)

Therefore, integrating this relation from t_1 to t_2 , we have the duality equality

$$\langle u^{\varepsilon}(t_2), \zeta_k^{\varepsilon}(t_1) \rangle = \langle u^{\varepsilon}(t_1), \zeta_k^{\varepsilon}(t_2) \rangle + \int_{t_1}^{t_2} \varepsilon \partial_x \zeta_k^{\varepsilon}(t, -L) h(t_1 + t_2 - t) \, \mathrm{d}t$$
(52)

Then, we recall ψ_k^{ε} is an eigenfunction of $\mathcal{L}^{\varepsilon}$, so that ζ_k^{ε} is explicitly known as

$$\zeta_k^{\varepsilon}(t,x) = \psi_k^{\varepsilon}(x)e^{-\lambda_k^{\varepsilon}(t-t_1)}$$

Substituting this into (52) yields

$$\langle u^{\varepsilon}(t_2), \psi_k^{\varepsilon} \rangle = e^{-\lambda_k^{\varepsilon}(t_2 - t_1)} \langle u^{\varepsilon}(t_1), \psi_k^{\varepsilon} \rangle + \varepsilon(\psi_k^{\varepsilon})'(-L) \int_{t_1}^{t_2} e^{-\lambda_k^{\varepsilon}(t - t_1)} h(t_1 + t_2 - t) \, \mathrm{d}t,$$

which concludes the proof.

Applying Lemma 3.3 with $k = 0, t_1 = 0, t_2 = \tau$ gives the relation

$$\langle u^{\varepsilon}(\tau), \psi_{0}^{\varepsilon} \rangle = e^{-\lambda_{0}^{\varepsilon}\tau} \langle u_{0}, \psi_{0}^{\varepsilon} \rangle + \varepsilon (\psi_{0}^{\varepsilon})'(-L) \int_{0}^{\tau} e^{-\lambda_{0}^{\varepsilon}t} h(\tau - t) \,\mathrm{d}t.$$
(53)

We then choose

$$h(t) = -\frac{\langle u_0, \psi_0^{\varepsilon} \rangle}{\varepsilon(\psi_0^{\varepsilon})'(-L)} \cdot \frac{e^{-\lambda_0^{\varepsilon} t}}{\tau}.$$
(54)

It immediately follows that

$$\langle u^{\varepsilon}(\tau), \psi_0^{\varepsilon} \rangle = 0.$$

We have the upper bound

$$h(t)| \leq \frac{\|\psi_0^{\varepsilon}\|_{L^2(-L,L)}}{|\varepsilon(\psi_0^{\varepsilon})'(-L)|} \cdot \frac{\|u_0\|_{L^2(-L,L)}}{\tau}.$$

Using the estimate (21) then immediately yields

$$\|h\|_{L^2(0,\tau)} \le \frac{2\sqrt{2L}}{\tau} \|u_0\|_{L^2(-L,L)}.$$

Finally, to perform the desired identities on $u^{\varepsilon}(\tau)$, we apply Lemma 3.3 again, for $k \ge 1$ and $t_1 = 0, t_2 = \tau$. This leads to the duality equalities

$$\langle u^{\varepsilon}(\tau), \psi_{k}^{\varepsilon} \rangle = e^{-\lambda_{k}\tau} \langle u_{0}, \psi_{k}^{\varepsilon} \rangle + \varepsilon(\psi_{k}^{\varepsilon})'(-L) \int_{0}^{\tau} e^{-\lambda_{k}^{\varepsilon}t} h(\tau - t) \,\mathrm{d}t$$

Plugging the expression of the control h(t), the solution at final time then verifies, for $k \ge 1$,

$$\langle u^{\varepsilon}(\tau), \psi_{k}^{\varepsilon} \rangle = e^{-\lambda_{k}^{\varepsilon}\tau} \langle u_{0}, \psi_{k}^{\varepsilon} \rangle - \frac{(\psi_{k}^{\varepsilon})'(-L)}{(\psi_{0}^{\varepsilon})'(-L)} \cdot \frac{e^{-\lambda_{0}^{\varepsilon}\tau} \langle u_{0}, \psi_{0}^{\varepsilon} \rangle}{\tau} \int_{0}^{\tau} e^{-(\lambda_{k}^{\varepsilon}-\lambda_{0}^{\varepsilon})t} \,\mathrm{d}t.$$
(55)

This concludes the proof of Lemma 3.1.

We additionnally note that, as $\varepsilon \to 0$, we may compute the limit of the control h using the property (23) of the eigenfunction ψ_0^{ε} . It holds

$$e^{-\lambda_0^{\varepsilon}t} \xrightarrow{L^2(0,\tau)} 1,$$
$$\frac{\psi_0^{\varepsilon}}{\varepsilon(\psi_0^k)'(-L)} \xrightarrow{L^2(-L,L)} 1$$

Plugging those limits in the expression of the control (54), we obtain

$$h^{\varepsilon} \xrightarrow{L^{2}(0,\tau)} - \frac{\int_{-L}^{L} u_{0}(x) \,\mathrm{d}x}{\tau},\tag{56}$$

which proves the first half of (17)

3.3. Controllability of the other modes. Proof of Lemma 3.2. Let $T > 0, \tau \in (0, T)$ with

$$T - \tau > T^* = 4\sqrt{3}L.$$
 (57)

Let $u_0 \in H_0^1 \cap H^2$, and $u^{\varepsilon}(\tau)$ prescribed by the conditions (43). We perform the change of variable

$$w(t,x) := e^{\frac{t}{4\varepsilon}} u^{\varepsilon}(t+\tau,x), \tag{58}$$

so that the solution u^{ε} of (5) on the time interval (τ, T) starting at time τ with the initial data $u^{\varepsilon}(\tau)$ provides a solution w to the system

$$\begin{cases} \partial_t w + \left(\mathcal{L}^{\varepsilon} - \frac{1}{4\varepsilon}\right) w = 0, \\ w(t, -L) = \tilde{h}(t), \\ w(t, L) = 0, \\ w(0, x) = u^{\varepsilon}(\tau, x), \end{cases}$$
(59)

on $(0, \hat{T})$ with

 $\widehat{T} := T - \tau, \quad \widetilde{h}(t) = e^{\frac{t}{4\varepsilon}} h(t + \tau).$

The eigenvalues of the operator $\mathcal{L}^{arepsilon}-rac{1}{4arepsilon}$ acting on $H^1_0\cap H^2$ are given by

$$\mu_k^{\varepsilon} := \lambda_k^{\varepsilon} - \frac{1}{4\varepsilon}, \quad k \ge 0, \tag{60}$$

and the associated eigenbasis is the family $(\psi_k^{\varepsilon})_{k\geq 0}$. The interest of this new system is that, by Lemma 2.1, the eigenvalues $(\mu_k^{\varepsilon})_{k\geq 1}$ of the associated operator look like the ones of the heat operator, for which one has some precise controllability results, in particular from [24]. The eigenvalue μ_0^{ε} is negative of order $-\frac{1}{4\varepsilon}$, but this does not induce any problem as we already eliminated the corresponding eigenmode in the previous step.

Without control the system already dissipates and drives the system close to zero in finite time. However, driving the state exactly to zero in finite time will require a control which may be huge a priori. In order to exploit the dissipation of the system, we therefore split again the imparted time interval into two parts by introducing $m \in (0, 1)$ which we will determine later, to optimize the control cost, and our strategy is as follows. First, on the time interval $[0, m\hat{T}]$, we set h(t) = 0 and let the dissipation happen. Then, we apply a duality result similar to Lemma 3.3 to the operator $\mathcal{L}^{\varepsilon} - \frac{1}{4\varepsilon}$, to obtain that for any $k \ge 0$,

$$\langle w(\widehat{T}), \psi_k^{\varepsilon} \rangle = e^{-\mu_k^{\varepsilon} \widehat{T}} \langle w(0), \psi_k^{\varepsilon} \rangle + \varepsilon (\psi_k^{\varepsilon})' (-L) \int_{m\widehat{T}}^{\widehat{T}} \widetilde{h}(t) e^{-\mu_k^{\varepsilon} (\widehat{T}-t)} \, \mathrm{d}t.$$
(61)

Therefore, since the $(\psi_k^{\varepsilon})_{k\geq 0}$ form a Hilbert basis, the null-control problem, which by (58), reduces to $w(\hat{T}) = 0$, is equivalent to finding a control $\tilde{h}(t)$ so that for any $k \geq 0$,

$$\int_{m\widehat{T}}^{\widehat{T}} \widetilde{h}(t)e^{\mu_k^{\varepsilon}t} \,\mathrm{d}t = -\frac{\langle w(0), \psi_k^{\varepsilon} \rangle}{\varepsilon(\psi_k^{\varepsilon})'(-L)}.$$
(62)

Shifting the integral to center the time interval around 0, one rephrases this condition as

$$\int_{-(1-m)\hat{T}/2}^{(1-m)T/2} \widetilde{h}\left(t + \frac{m+1}{2}\widehat{T}\right) e^{\mu_k^{\varepsilon}t} dt = -e^{-\mu_k^{\varepsilon}\frac{m+1}{2}\widehat{T}} \frac{\langle w(0), \psi_k^{\varepsilon} \rangle}{\varepsilon(\psi_k^{\varepsilon})'(-L)}.$$
(63)

A classical method used for such moment problems, in particular for controllability results on the heat equation and other parabolic systems, is to look for a control $\tilde{h}(t)$ as a series of functions $(\tilde{q}_k)_{k\geq 1}$ which are bi-orthogonal to the functions $\left(e^{\mu_j^{\varepsilon}t}\right)_{j\geq 0}$ in $L^2\left(-\frac{(1-m)\hat{T}}{2}, \frac{(1-m)\hat{T}}{2}\right)$, see [5]. To determine such functions $(\tilde{q}_k)_{k\geq 1}$ we are going to use the following result regarding the construction of a family bi-orthogonal to the functions $\left(e^{\mu_k^{\varepsilon}t}\right)_{k\geq 1}$.

LEMMA 3.4. For any $\varepsilon > 0, \widetilde{T} > 0$, there exists a family of functions $(q_k^{\varepsilon})_{k\geq 1}$ in $L^2\left(-\frac{\widetilde{T}}{2}, \frac{\widetilde{T}}{2}\right)$ such that $\zeta \frac{\widetilde{T}}{2}$

$$\forall j \ge 0, k \ge 1, \int_{-\frac{\tilde{T}}{2}}^{\frac{1}{2}} e^{\mu_j^{\varepsilon} t} q_k^{\varepsilon}(t) \, \mathrm{d}t = \delta_{j,k},\tag{64}$$

Moreover, for any $\kappa > 1$, there exists such a family and a constant $c = c(\varepsilon, \kappa, L, \tilde{T})$ which is a rational function of its arguments, independent of $k \ge 1$, such that

$$\|q_k^{\varepsilon}\|_{L^2\left(-\frac{\tilde{T}}{2},\frac{\tilde{T}}{2}\right)} \le c \frac{1}{\mu_k^{\varepsilon}} e^{\frac{3\kappa L^2}{\varepsilon T}},\tag{65}$$

This result is proved in Appendix A and heavily relies on the construction of [24]. We then use Lemma 3.4 for $\tilde{T} = (1 - m)\hat{T}$ and for some $\kappa > 1$ to be determined later, see (76). We may now construct the control $\tilde{h} \in L^2(0, \hat{T})$, as

$$\widetilde{h}(t) = \begin{cases} 0, & t \in [0, mT], \\ -\sum_{k \ge 1} c_k q_k^{\varepsilon} \left(t - \frac{m+1}{2} \widehat{T} \right), & t \in [m\widehat{T}, \widehat{T}], \end{cases}$$
(66)

where the coefficients c_k are given by

$$c_k := e^{-\mu_k^{\varepsilon} \frac{m+1}{2}\hat{T}} \frac{\langle w(0), \psi_k^{\varepsilon} \rangle}{\varepsilon(\psi_k^{\varepsilon})'(-L)}.$$
(67)

We will prove that this control is well-defined, steers the solution of (59) to 0, and satisfies suitable L^2 estimates.

This choice of control clearly solves the moment problem (63), and this yields, for $k \ge 0$,

$$\langle w(T), \psi_k^{\varepsilon} \rangle = 0.$$
 (68)

Therefore, provided h is well-defined, since $y(T)=w(\widehat{T})e^{-\frac{\widehat{T}}{4\varepsilon}},$ the control

$$h(t) = e^{-\frac{t}{4\varepsilon}} \widetilde{h}(t-\tau)$$

steers $u^{\varepsilon}(\tau)$ to $u^{\varepsilon}(T) = 0$.

We may now estimate the norm of the control \tilde{h} . We start by bounding the coefficients c_k . Reminding that $w(0) = u^{\varepsilon}(\tau)$ is given by the conditions (43), we can rewrite

$$c_k = e^{-\mu_k^{\varepsilon} \frac{1+m}{2} \widehat{T}} \left(e^{-\lambda_k^{\varepsilon} \tau} \frac{\langle u_0, \psi_k^{\varepsilon} \rangle}{\varepsilon(\psi_k^{\varepsilon})'(-L)} - \frac{e^{-\lambda_0^{\varepsilon} \tau}}{\tau} \int_0^{\tau} e^{-(\lambda_k^{\varepsilon} - \lambda_0^{\varepsilon})t} \, \mathrm{d}t \frac{\langle u_0, \psi_0^{\varepsilon} \rangle}{\varepsilon(\psi_0^{\varepsilon})'(-L)} \right)$$
(69)

We estimate both terms separately. For the first term, we use the property (22) of the eigenfunctions ψ_k^{ε} , it gives:

$$\left| e^{-\mu_k^{\varepsilon} \frac{1+m}{2} \widehat{T}} e^{-\lambda_k^{\varepsilon} \tau} \frac{\langle u_0, \psi_k^{\varepsilon} \rangle}{\varepsilon(\psi_k^{\varepsilon})'(-L)} \right| \leq \frac{\|u_0\|_{L^2} \|\psi_k^{\varepsilon}\|_{L^2(-L,L)}}{|\varepsilon(\psi_k^{\varepsilon})'(-L)|} \leq \frac{4L}{k\pi\sqrt{\varepsilon}} \|u_0\|_{L^2(-L,L)}.$$

$$(70)$$

For the second term, we use the property (21) of the first eigenfunction ψ_0^{ε} .

$$\left| e^{-\mu_{k}^{\varepsilon} \frac{1+m}{2} \widehat{T}} \frac{e^{-\lambda_{0}^{\varepsilon} \tau}}{\tau} \int_{0}^{\tau} e^{-(\lambda_{k}^{\varepsilon} - \lambda_{0}^{\varepsilon})t} dt \frac{\langle u_{0}, \psi_{0}^{\varepsilon} \rangle}{\varepsilon(\psi_{0}^{\varepsilon})'(-L)} \right| \leq \frac{1}{\tau(\lambda_{k}^{\varepsilon} - \lambda_{0}^{\varepsilon})} \cdot \frac{\|u_{0}\|_{L^{2}(-L,L)} \|\psi_{0}^{\varepsilon}\|_{L^{2}(-L,L)}}{|\varepsilon(\psi_{0}^{\varepsilon})'(-L)|} \leq \frac{2\sqrt{2L}}{\tau(\lambda_{k}^{\varepsilon} - \lambda_{0}^{\varepsilon})} \|u_{0}\|_{L^{2}(-L,L)}.$$
(71)

Combining the estimates (70) and (71) we deduce from (69) that for any $k \ge 1$,

$$|c_k| \le \left(\frac{4L}{k\pi\sqrt{\varepsilon}} + \frac{2\sqrt{2L}}{\tau(\lambda_k^{\varepsilon} - \lambda_0^{\varepsilon})}\right) \|u_0\|_{L^2(-L,L)}.$$
(72)

Finally, the estimate (65) gives

$$\|\widetilde{h}\|_{L^{2}(0,\widehat{T})} \leq \sum_{k\geq 1} \frac{1}{\mu_{k}^{\varepsilon}} \left(\frac{4L}{k\pi\sqrt{\varepsilon}} + \frac{2\sqrt{2L}}{\tau(\lambda_{k}^{\varepsilon} - \lambda_{0}^{\varepsilon})} \right) c \exp\left(\frac{3\kappa L^{2}}{\varepsilon(1-m)\widehat{T}}\right) \|u_{0}\|_{L^{2}(-L,L)}.$$
(73)

By the distribution of the eigenvalues (19), the sum is finite, and there exists $c = c(\varepsilon, \kappa, L, T, m, \tau) > 0$ a rational function of its arguments such that,

$$\|\widetilde{h}\|_{L^2(0,\widehat{T})} \le c \exp\left(\frac{3\kappa L^2}{\varepsilon(1-m)\widehat{T}}\right) \|u_0\|_{L^2}.$$
(74)

Since \tilde{h} vanishes on $[0, m\hat{T}]$ and $h(t + \tau) = e^{-\frac{t}{4\varepsilon}}\tilde{h}(t)$, it immediately follows that

$$\|h\|_{L^{2}(\tau,T)} \leq e^{-\frac{m\hat{T}}{4\varepsilon}} \|\tilde{h}\|_{L^{2}(0,\hat{T})}$$
$$\leq c \exp\left(-\frac{m\hat{T}}{4\varepsilon} + \frac{3\kappa L^{2}}{\varepsilon(1-m)\hat{T}}\right) \|u_{0}\|_{L^{2}(-L,L)}$$
(75)

Then the term $-\frac{m\hat{T}}{4} + \frac{3\kappa L^2}{(1-m)\hat{T}}$ is negative whenever $\hat{T}^2 > \frac{12\kappa L^2}{m(1-m)}$. We take $m = \frac{1}{2}$ to minimise that bound, and obtain that the cost to steer $u^{\varepsilon}(\tau)$ to 0 is exponentially small as soon as

$$\widehat{T} > 4\sqrt{3}\sqrt{\kappa}L = \sqrt{\kappa}T^*,\tag{76}$$

which is possible for some $\kappa > 1$ by the assumption (45). This concludes the proof of Lemma 3.2. Moreover, it implies that

$$h^{\varepsilon} \xrightarrow{L^2(\tau,T)} 0. \tag{77}$$

Combining (56) and (77) concludes the proof of identity (17), and of Theorem 1.3

4. Proof of the lower bound on T_{unif}

In this section we prove that $T_{\text{unif}} \ge (4\sqrt{2}-2)L$. We follow the work from [19], improving [2, pp 106-109], to give a lower bound on the norm of any control function steering a well-chosen initial datum to zero. We set $T > 0, \varepsilon > 0$, and choose the initial datum of the control system (5) as $u_0^{\varepsilon} = \operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}$. Let h(t) the associated optimal control function. By definition of the null-controllability cost, it verifies

$$\|h\|_{L^2(0,T)} \le \mathcal{C}(T,\varepsilon) \left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}\right\|.$$
 (78)

Moreover, applying Lemma 3.3 on the whole time interval [0, T], we have, for $k \ge 0$

$$\langle u^{\varepsilon}(T), \psi_{k}^{\varepsilon} \rangle = e^{-\lambda_{k}^{\varepsilon}T} \langle \operatorname{sech}\left(\frac{x}{2\varepsilon}\right) \varphi_{1}^{\varepsilon}, \psi_{k}^{\varepsilon} \rangle + \varepsilon(\psi_{k}^{\varepsilon})'(-L) \int_{0}^{T} h(t) e^{-\lambda_{k}^{\varepsilon}(T-t)} \,\mathrm{d}t.$$

$$= e^{-\lambda_{k}^{\varepsilon}T} \langle \varphi_{1}^{\varepsilon}, \varphi_{k}^{\varepsilon} \rangle + \varepsilon(\psi_{k}^{\varepsilon})'(-L) \int_{0}^{T} h(t) e^{-\lambda_{k}^{\varepsilon}(T-t)} \,\mathrm{d}t.$$

$$(79)$$

We introduce the function

$$v(z) := \int_{-\frac{T}{2}}^{\frac{T}{2}} h\left(t + \frac{T}{2}\right) e^{-izt} \,\mathrm{d}t.$$
(80)

v is the Fourier transform of a compactly supported function, and is therefore entire.

Since the operator P^{ε} acting on $H_0^1 \cap H^2(-L, L)$ is self-adjoint, the family $(\varphi_k^{\varepsilon})_{k\geq 1}$ is an orthogonal family. It then follows from (79) and the constraint $u^{\varepsilon}(T) = 0$ that

$$v(i\lambda_k^{\varepsilon}) = \begin{cases} -e^{\frac{-\lambda_1^{\varepsilon}T}{2}} \frac{\|\varphi_1^{\varepsilon}\|^2}{\varepsilon(\psi_1^{\varepsilon})'(-L)}, & k = 1\\ 0, & k \neq 1. \end{cases}$$
(81)

Using the bound (78), we also have, for any $z \in \mathbb{C}$

$$|v(z)| \leq \mathcal{C}(T,\varepsilon)\sqrt{T}\exp\left(\frac{T|Im(z)|}{2}\right) \left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}\right\|$$

We defined the rescaled entire function

$$f(z) := v\left(\frac{z}{4\varepsilon}\right),$$

which immediately verifies, for $z \in \mathbb{C}$,

$$|f(z)| \le \mathcal{C}(T,\varepsilon)\sqrt{T} \exp\left(\frac{T|Im(z)|}{8\varepsilon}\right) \left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}\right\|.$$
(82)

Similarly to the proof of the upper bound, we rewrite the null-control problem in terms of moments of the control function h, and we express it as the inverse Fourier transform of some entire function with prescribed values at some key points. This point of view is well studied in complex analysis, notably with the Hadamard and Weyl factorization theorems. In this case, we will apply a representation theorem from [13, p.56] to write, for z in the upper half-plane,

$$\ln|f(z)| = \sum_{\ell \ge 1} \ln\left|\frac{z - a_\ell}{z - \overline{a_\ell}}\right| + \sigma Im(z) + \frac{Im(z)}{\pi} \int_{\mathbb{R}} \frac{\ln|f(s)|}{|s - z|^2} \,\mathrm{d}s,\tag{83}$$

where σ is the type of the entire function f and the $(a_\ell)_{\ell \ge 1}$ are the roots of f in the upper half-plane. We apply this equality in $z = 4i\varepsilon\lambda_1^{\varepsilon}$, and want to find an upper bound on the right-hand side.

For the second term, we know from the estimate (82) that the type of f verifies $\sigma \leq \frac{T}{8\varepsilon}$, so that

$$\sigma \operatorname{Im}(4i\varepsilon\lambda_1^\varepsilon) \le \frac{\lambda_1^\varepsilon T}{2}.$$
(84)

Applying the estimate (82) again with $s \in \mathbb{R}$, we get

$$\ln|f(s)| \le \ln\left(\mathcal{C}(T,\varepsilon)\sqrt{T} \left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}\right\|\right),$$

and we may directly deduce a bound for the third term:

$$\frac{Im(4i\varepsilon\lambda_{1}^{\varepsilon})}{\pi} \int_{\mathbb{R}} \frac{\ln|f(s)|}{|s-4i\varepsilon\lambda_{1}^{\varepsilon}|^{2}} \, \mathrm{d}s \leq \frac{4\varepsilon\lambda_{1}^{\varepsilon}}{\pi} \ln\left(\mathcal{C}(T,\varepsilon)\sqrt{T} \left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_{1}^{\varepsilon}\right\|\right) \int_{\mathbb{R}} \frac{1}{s^{2}+(4\varepsilon\lambda_{1}^{\varepsilon})^{2}} \, \mathrm{d}s$$
$$\leq \ln\left(\mathcal{C}(T,\varepsilon)\sqrt{T} \left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_{1}^{\varepsilon}\right\|\right) \tag{85}$$

Finally, for the first term, we only use the known roots of the form $4i\varepsilon\lambda_k^{\varepsilon}$, for $k\geq 2$, which provides the upper bound

$$\sum_{\ell \ge 1} \ln \left| \frac{4i\varepsilon\lambda_1^\varepsilon - a_\ell}{4i\varepsilon\lambda_1^\varepsilon - \overline{a_\ell}} \right| \le \sum_{k \ge 2} \ln \left(\frac{\lambda_k^\varepsilon - \lambda_1^\varepsilon}{\lambda_k^\varepsilon + \lambda_1^\varepsilon} \right).$$

With the distribution of the eigenvalues (19) and (20), this becomes

$$\sum_{k\geq 2} \ln\left(\frac{\lambda_k^{\varepsilon} - \lambda_1^{\varepsilon}}{\lambda_k^{\varepsilon} + \lambda_1^{\varepsilon}}\right) \leq \sum_{k\geq 2} \ln\left(\frac{((k+1)^2 - 1)\frac{\varepsilon\pi^2}{4L^2}}{\frac{1}{2\varepsilon} + (k^2 + 1)\frac{\varepsilon\pi^2}{4L^2}}\right)$$
$$\leq \sum_{k\geq 2} \left(\frac{(k^2 - 1)\frac{\varepsilon\pi^2}{4L^2}}{\frac{1}{2\varepsilon} + (k^2 + 1)\frac{\varepsilon\pi^2}{4L^2}}\right) - \ln\left(\frac{3\varepsilon\pi^2}{4L^2}\right).$$

Comparing the series with an integral and integrating by parts then yields the upper bound

$$\sum_{\ell \ge 1} \ln \left| \frac{4i\varepsilon\lambda_1^\varepsilon - a_\ell}{4i\varepsilon\lambda_1^\varepsilon - \overline{a_\ell}} \right| \le -\frac{\sqrt{2}L}{\varepsilon} + 2\ln\left(1 + \frac{2L^2}{\pi^2\varepsilon^2}\right) + \ln\left(\frac{4L^2}{3\varepsilon\pi^2}\right) + 2.$$
(86)

Replacing $f(4i\varepsilon\lambda_1^{\varepsilon})$ with its expression and combining the bounds (84), (85) and (86) then gives

$$-\frac{\lambda_1^{\varepsilon}T}{2}\ln\left(\frac{\|\varphi_1^{\varepsilon}\|^2}{\varepsilon(\psi_1^{\varepsilon})'(-L)}\right) \le -\frac{\sqrt{2}L}{\varepsilon} + 2\ln\left(1 + \frac{2L^2}{\pi^2\varepsilon^2}\right) + \ln\left(\frac{4L^2}{3\varepsilon\pi^2}\right) + 2 + \frac{\lambda_1T}{2} + \ln\left(\mathcal{C}(T,\varepsilon)\sqrt{T}\left\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}\right\|\right)$$

Rearranging the terms gives the lower bound on the null-controllability cost

$$\mathcal{C}(T,\varepsilon) \ge e^2 \cdot \frac{4L^2}{3\varepsilon\pi^2} \left(1 + \frac{2L^2}{\pi^2\varepsilon^2}\right)^2 \frac{\|\varphi_1^\varepsilon\|^2}{\varepsilon(\psi_1^\varepsilon)'(-L) \|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^\varepsilon\|} \exp\left(-\lambda_1^\varepsilon T + \frac{\sqrt{2}L}{\varepsilon}\right).$$
(87)

Finally, we use the expression of the eigenfunction φ_1^{ε} given by (39), which implies the existence of a constant c > 0 such that

$$\frac{\|\varphi_1^{\varepsilon}\|^2}{\varepsilon(\psi_1^{\varepsilon})'(-L)\|\operatorname{sech}\left(\frac{x}{2\varepsilon}\right)\varphi_1^{\varepsilon}\|} \ge c\exp\left(-\frac{L}{2\varepsilon}\right).$$
(88)

Plugging this estimate into the bound (87) leads to

$$\mathcal{C}(T,\varepsilon) \ge ce^2 \cdot \frac{4L^2}{3\varepsilon\pi^2} \left(1 + \frac{2L^2}{\pi^2\varepsilon^2}\right)^2 \exp\left(-\lambda_1^\varepsilon T + \frac{\sqrt{2}L}{\varepsilon} - \frac{L}{2\varepsilon}\right)$$
(89)

Since $\lambda_1^{\varepsilon} \leq \frac{1}{4\varepsilon} + 4\varepsilon \frac{\pi^2}{4L^2}$, the null-controllability cost $C(T, \varepsilon)$ explodes as ε approaches 0 if the quantity $-\frac{T}{4} + \sqrt{2}L - \frac{L}{2}$ is positive, namely whenever

$$T < (4\sqrt{2} - 2)L.$$
 (90)

The proof of the lower bound on T_{unif} is then concluded.

Appendix A. Construction of the bi-orthogonal family, proof of Lemma 3.4

We follow the construction of the bi-orthogonal family by Tenenbaum and Tucsnak [24], with a few small adaptations to fit the spectrum of our operator. Let $\tilde{T} > 0, \varepsilon > 0, \kappa > 1$. We wish to build a family $(q_k^{\varepsilon})_{k\geq 1}$ which satisfies the constraints (64), as well as the L^2 estimate (65). Rescaling in time, we set

 $S = \frac{\pi^2 \varepsilon}{4L^2} \widetilde{T}, \quad \eta_k^\varepsilon = \frac{4L^2}{\pi^2 \varepsilon} \mu_k^\varepsilon,$

where we recall (μ_k) is defined by (60), and we instead construct a family $(p_k^{\varepsilon})_{k\geq 1}$ of $L^2\left(-\frac{S}{2}, \frac{S}{2}\right)$ that verifies

$$\forall j \ge 0, k \ge 1, \int_{-\frac{S}{2}}^{\frac{S}{2}} e^{\eta_j^{\varepsilon} t} p_k^{\varepsilon}(t) \, \mathrm{d}t = \delta_{j,k},\tag{91}$$

and then take

$$q_k^{\varepsilon}(t) := \frac{4L^2}{\pi^2 \varepsilon} p_k^{\varepsilon} \left(\frac{4L^2 t}{\pi^2 \varepsilon} \right).$$
(92)

From Proposition 2.1, and more precisely estimates (18), (19) and (20), the family $(\eta_k^{\varepsilon})_{k>0}$ satisfies:

$$\eta_0^\varepsilon = -\frac{L^2}{\pi^2 \varepsilon^2} + O(e^{-1/\varepsilon}),\tag{93}$$

$$(k+1)^2 \ge \eta_k^{\varepsilon} \ge k^2, \quad k \ge 1, \tag{94}$$

$$|\eta_j^{\varepsilon} - \eta_k^{\varepsilon}| \ge |j^2 - k^2|, \quad j,k \ge 1.$$
(95)

We want to build a family of entire functions $g_k, k \ge 1$ that are integrable on the real axis, and verify, for any $j \ge 0, k \ge 1$,

$$g_k(i\eta_j^\varepsilon) = \delta_{jk}.\tag{96}$$

An approach to build such a family has been developed in [5], and consists in taking functions of the form

$$g_k(z) = \Phi_k(z) \frac{f(z)}{f(i\eta_k^\varepsilon)},\tag{97}$$

where Φ_k is a "natural" entire function with its poles in the $(i\eta_j)_{j\neq k}$, and the function f is a multiplier which aims at compensating the growth of Φ_k on the real axis.

We define, for any $k \ge 1$,

$$\Phi_k(z) = \prod_{j \ge 0, j \ne k} \frac{-iz - \eta_j^{\varepsilon}}{\eta_k^{\varepsilon} - \eta_j^{\varepsilon}}.$$
(98)

For any $j \ge 0, k \ge 1$, it immediately holds

$$\Phi_k(i\eta_j^\varepsilon) = \delta_{jk}.\tag{99}$$

Moreover, using the estimates (94) and (95), there holds, for $x \in \mathbb{R}$,

$$\begin{aligned} |\Phi_k(x)| &\leq \left| \frac{-ix - \eta_0^{\varepsilon}}{\eta_k^{\varepsilon} - \eta_0^{\varepsilon}} \right| \prod_{j\geq 1, j\neq k} \left| \frac{-ix - (j+1)^2}{k^2 - j^2} \right|, \\ &\leq \left| \frac{-ix - \eta_0^{\varepsilon}}{\eta_k^{\varepsilon} - \eta_0^{\varepsilon}} \right| \cdot \left| \frac{-ix - k^2}{-ix - 1} \right| \cdot \prod_{j\geq 1, j\neq k} \left| \frac{-ix - j^2}{k^2 - j^2} \right| \end{aligned}$$
(100)

Rewriting the last product using Hadamard's product formulation of the sine, it then leads, for any $x \in \mathbb{R}$, to

$$|\Phi_k(x)| \le c(x,k,\varepsilon,L) \exp\left(\pi\sqrt{|x|/2}\right),\tag{101}$$

where, here and in the rest of the proof, c is an abuse of notation to denote a rational fraction of its arguments. Moreover, for any $k \ge 1$ and $z \in \mathbb{C}$ we have

$$|\Phi_k(z)| = \prod_{j \ge 0, j \ne k} \left(1 + \frac{-iz - \eta_k^{\varepsilon}}{\eta_k^{\varepsilon} - \eta_j^{\varepsilon}} \right).$$

By the distribution of the spectrum (95), the sum $\sum_{j \neq k} |\eta_j^{\varepsilon} - \eta_k^{\varepsilon}|^{-1}$ converges, so that

$$\Phi_k(z) \le \exp\left(c(k,\varepsilon,L)| - iz - \eta_k^{\varepsilon}|\right)$$

It immediately follows, for $k \ge 1, z \in \mathbb{C}$,

$$|\Phi_k(z)| \le c(k,\varepsilon,L) \exp(c(k,\varepsilon,L)|z|).$$
(102)

We now introduce the same multiplier used in [24]. For $\beta > 0, \delta > 0$, we set

$$\nu = \frac{(\pi + \delta)^2}{\beta},$$

and we define the entire function H_{β} by

$$H_{\beta}(z) := C_{\nu} \int_{-1}^{1} \exp\left(-\frac{\nu}{1-t^2} - i\beta tz\right) dt,$$
(103)

where C_{ν} is the normalizing constant such that $H_{\beta}(0) = 1$. The entire H_{β} then verifies the following estimates:

$$|H_{\beta}(iy)| \ge \frac{1}{11\sqrt{\nu+1}} \exp\left(\frac{\beta|y|}{2\sqrt{\nu+1}}\right), \qquad \forall y \in \mathbb{R}, \qquad (104)$$

$$H_{\beta}(z)| \le e^{\beta|z|} \qquad \qquad \forall z \in \mathbb{C},, \tag{105}$$

$$|H_{\beta}(x)| \le c(\beta, \delta)\sqrt{\nu+1} \exp\left(\frac{3\nu}{4} - (\pi + \frac{\delta}{2})\sqrt{|x|}\right), \qquad \forall x \in \mathbb{R}.$$
 (106)

We choose the multiplier

$$f(z) := H_{\beta}\left(\frac{z}{2}\right).$$

Recalling the definition of g_k (97) and combining (101), (104) and (106), we obtain

$$|g_k(x)| \le c(x,k,\varepsilon,L,\nu) \exp\left(-\frac{\beta\eta_k^\varepsilon}{4\sqrt{\nu+1}} + \frac{3\nu}{4} - \frac{\delta}{2\sqrt{2}}\sqrt{|x|}\right)$$
(107)

We then set $\beta \in \left(\frac{S}{\kappa}, S\right)$, and $\delta > 0$ such that

$$\nu < \frac{(4-\delta)\pi^2\kappa}{4S}.$$

It immediately follows, for any $k \ge 1$, that

$$\|g_k\|_{L^1(\mathbb{R})} \le c(\varepsilon, \kappa, L, S) \frac{\exp\left(\frac{3\pi^2 \kappa}{4S}\right)}{\eta_k^{\varepsilon}}.$$
(108)

By growth estimates (102) and (105), we may apply the Paley-Wiener theorem, and there exists a family $(p_k^{\varepsilon})_{k\geq 1}$ in $L^2(-\frac{S}{2},\frac{S}{2})$ such that

$$g_k(z) = \int_{-\frac{S}{2}}^{\frac{S}{2}} p_k(t) e^{-itz} \,\mathrm{d}t, \tag{109}$$

and $||p_k||_{L^2\left(-\frac{S}{2},\frac{S}{2}\right)} \leq c(\varepsilon,\kappa,L,S)(\eta_k^{\varepsilon})^{-1} \exp\left(\frac{3\pi^2\kappa}{4S}\right)$. Moreover, the relation (99) translates to $g_k(i\eta_j^{\varepsilon}) = \delta_{jk}$ for $j \geq 0, k \geq 1$, and therefore the family $(p_k^{\varepsilon})_{k\geq 1}$ satisfies (91).

Finally, the family $(q_k)_{k>1}$ defined by (92) immediately verifies (64), as well as

$$\begin{aligned} \|q_{k}^{\varepsilon}\|_{L^{2}\left(-\frac{\tilde{T}}{2},\frac{\tilde{T}}{2}\right)} &= \frac{2L}{\pi\sqrt{\varepsilon}} \|p_{k}^{\varepsilon}\|_{L^{2}\left(-\frac{S}{2},\frac{S}{2}\right)} \\ &\leq c(\varepsilon,\kappa,L,S)(\eta_{k}^{\varepsilon})^{-1} \exp\left(\frac{3\pi^{2}\kappa}{4S}\right) \\ &\leq c(\varepsilon,\kappa,L,S)(\mu_{k}^{\varepsilon})^{-1} \exp\frac{3\kappa L^{2}}{\varepsilon\tilde{T}}, \end{aligned}$$
(110)

which proves the estimate (65). The Lemma 3.4 is now fully proven.

References

- [1] G. Carbou, *Stability of static walls for a three-dimensional model of ferromagnetic material*, Journal de mathématiques pures et appliquées **93** (2010), no. 2, 183–203.
- [2] J.-M. Coron, *Control and nonlinearity*, Mathematical Surveys and Monographs, vol. 136, American Mathematical Society, Providence, RI, 2007.
- [3] J-M Coron and S. Guerrero, Singular optimal control: a linear 1-d parabolic-hyperbolic example, Asymptotic Analysis 44 (2005), no. 3-4, 237–257.
- [4] J. Dardé and S. Ervedoza, On the cost of observability in small times for the one-dimensional heat equation, Analysis & PDE 12 (2019), no. 6, 1455–1488.
- [5] H. O Fattorini and D. L Russell, Uniform bounds on biorthogonal functions for real exponentials with an application to the control theory of parabolic equations, Quarterly of Applied Mathematics 32 (1974), no. 1, 45–69.
- [6] R. Folino, C. Lattanzio, C. Mascia, and M. Strani, *Metastability for nonlinear convection-diffusion equations*, Nonlinear Differential Equations and Applications NoDEA 24 (2017), no. 4, 35.
- [7] O. Glass and S. Guerrero, *On the uniform controllability of the burgers equation*, SIAM Journal on Control and Optimization **46** (2007), no. 4, 1211–1238.
- [8] O. Glass, A complex-analytic approach to the problem of uniform controllability of a transport equation in the vanishing viscosity limit, Journal of Functional Analysis **258** (2010), no. 3, 852–868.
- [9] S. Guerrero and G. Lebeau, Singular optimal control for a transport-diffusion equation, Communications in Partial Differential Equations 32 (2007), no. 12, 1813–1836.
- [10] E. Hopf, *The partial differential equation* $u_t + uu_x = \mu u_{xx}$, Communications on Pure and Applied Mathematics **3** (1950), no. 3, 201–230.
- [11] A. M. Il'in and O. A. Oleinik, Asymptotic behavior of solutions of the cauchy problem for some quasi-linear equations for large values of the time, Matematicheskii Sbornik 93 (1960), no. 2, 191–216.
- [12] S. Kawashima and A. Matsumura, Stability of shock profiles in viscoelasticity with non-convex constitutive relations, Communications on Pure and Applied Mathematics 47 (1994), no. 12, 1547–1569.
- [13] P. Koosis, The logarithmic integral, Vol. 1, Cambridge university press, 1988.
- [14] G. Kreiss and H.-O. Kreiss, Convergence to steady state of solutions of burgers' equation, Applied Numerical Mathematics 2 (1986), no. 3-5, 161–179.
- [15] C. Laurent and M. Léautaud, Uniform observability estimates for linear waves, ESAIM: Control, Optimisation and Calculus of Variations 22 (2016), no. 4, 1097–1136.
- [16] C. Laurent and M. Léautaud, On uniform controllability of 1d transport equations in the vanishing viscosity limit, Comptes Rendus. Mathématique 361 (2023), no. G1, 265–312.
- [17] M. Léautaud, Uniform controllability of scalar conservation laws in the vanishing viscosity limit, SIAM Journal on Control and Optimization 50 (2012), no. 3, 1661–1699.
- [18] P. Lissy, A link between the cost of fast controls for the 1-d heat equation and the uniform controllability of a 1-d transportdiffusion equation, Comptes Rendus Mathematique 350 (2012), no. 11-12, 591–595.
- [19] P. Lissy, Explicit lower bounds for the cost of fast controls for some 1-d parabolic or dispersive equations, and a new lower bound concerning the uniform controllability of the 1-d transport-diffusion equation, Journal of Differential Equations 259 (2015), no. 10, 5331–5352.
- [20] C. Mascia and M. Strani, Metastability for nonlinear parabolic equations with application to scalar viscous conservation laws, SIAM Journal on Mathematical Analysis 45 (2013), no. 5, 3084–3113.
- [21] A. Matsumura and M. Mei, Nonlinear stability of viscous shock profile for a non-convex system of viscoelasticity (1997).

- [22] A. Matsumura and K. Nishihara, *Asymptotic stability of traveling waves for scalar viscous conservation laws with non-convex nonlinearity*, Communications in Mathematical Physics **165** (1994), 83–96.
- [23] D. H Sattinger, On the stability of waves of nonlinear parabolic systems, Advances in Mathematics 22 (1976), no. 3, 312–355.
- [24] G. Tenenbaum and M. Tucsnak, *New blow-up rates for fast controls of schrödinger and heat equations*, Journal of Differential Equations **243** (2007), no. 1, 70–100.

IMB, UNIVERSITÉ DE BORDEAUX

Email address: vincent.laheurte@math.u-bordeaux.fr