

ON SHARP ANISOTROPIC HARDY INEQUALITIES

XIA HUANG AND DONG YE

ABSTRACT. Recently, Yanyan Li and Xukai Yan showed in [7, 8] the following interesting Hardy inequalities with anisotropic weights: Let $n \geq 2$, $p \geq 1$, $p\alpha > 1 - n$, $p(\alpha + \beta) > -n$, then there exists $C > 0$ such that

$$\| |x|^\beta |x'|^{\alpha+1} \nabla u \|_{L^p(\mathbb{R}^n)} \geq C \| |x|^\beta |x'|^\alpha u \|_{L^p(\mathbb{R}^n)}, \quad \forall u \in C_c^1(\mathbb{R}^n).$$

Here $x' = (x_1, \dots, x_{n-1}, 0)$ for $x = (x_i) \in \mathbb{R}^n$. In this note, we will determine the best constant for the above estimate when $p = 2$ or $\beta \geq 0$. Moreover, as refinement for very special case of Li-Yan's result in [8], we provide explicit estimate for the anisotropic L^p -Caffarelli-Kohn-Nirenberg inequality.

2020 MSC: 26D10, 35A23, 46E35

Keywords: Anisotropic Hardy inequality, best constant, anisotropic L^p -Caffarelli-Kohn-Nirenberg inequality

1. INTRODUCTION

Recently, Li-Yan studied in [7] the asymptotic stability of the (-1) -homogeneous axisymmetric stationary solutions to Navier-Stokes equations in \mathbb{R}^3 . A key estimate was

$$(1.1) \quad \int_{\mathbb{R}^3} |\nabla u|^2 dx \geq C \int_{\mathbb{R}^3} \frac{u^2}{|x'| |x|} dx, \quad \forall u \in C_c^1(\mathbb{R}^3)$$

where $x' = (x_1, x_2, 0)$. Li-Yan [7] obtained (1.1) by proving the following strengthened inequality:

$$(1.2) \quad \int_{\mathbb{R}^3} \frac{|x'|}{|x|} |\nabla u|^2 dx \geq C \int_{\mathbb{R}^3} \frac{u^2}{|x'| |x|} dx, \quad \forall u \in C_c^1(\mathbb{R}^3).$$

It is worthy to remark that (1.2) improves also the classical Hardy inequality

$$\int_{\mathbb{R}^3} |\nabla u|^2 dx \geq \frac{1}{4} \int_{\mathbb{R}^3} \frac{u^2}{|x|^2} dx, \quad \forall u \in C_c^1(\mathbb{R}^3).$$

This motivated them to show the following general anisotropic Hardy inequalities, see [7, Theorem 1.3] and [8, pages 6-7].

Theorem A. Let $n \geq 2$, $x' = (x_1, \dots, x_{n-1}, 0)$ for $x = (x_i) \in \mathbb{R}^n$. Assume that $p \geq 1$, $p\alpha > 1 - n$, $p(\alpha + \beta) > -n$, then there exists positive constant C depending on n, p, α and β , such that

$$(1.3) \quad \| |x|^\beta |x'|^{\alpha+1} \nabla u \|_{L^p(\mathbb{R}^n)} \geq C \| |x|^\beta |x'|^\alpha u \|_{L^p(\mathbb{R}^n)}, \quad \forall u \in C_c^1(\mathbb{R}^n).$$

The interesting estimates (1.3) were used by Li-Yan to establish a generalized and improved anisotropic version of Caffarelli-Kohn-Nirenberg's interpolation inequalities, see [8, Theorem 1.1], which established necessary and sufficient conditions to have

$$(1.4) \quad \| |x|^{\gamma_1} |x'|^\alpha u \|_{L^s(\mathbb{R}^n)} \leq C \| |x|^{\gamma_2} |x'|^\mu \nabla u \|_{L^p(\mathbb{R}^n)}^a \| |x|^{\gamma_3} |x'|^\beta u \|_{L^q(\mathbb{R}^n)}^{1-q}, \quad \forall u \in C_c^1(\mathbb{R}^n).$$

Our main purpose here is to study the best constant for the inequalities (1.3). A first remark is that for any $n \geq 2$, $p \geq 1$, $|x|^\beta |x'|^\alpha \in L_{loc}^p(\mathbb{R}^n)$ if and only if

$$(1.5) \quad p\alpha > 1 - n \quad \text{and} \quad p(\alpha + \beta) > -n.$$

This fact can be seen by spherical coordinates. Our first result gives a complete answer for best constant of (1.3) when $p = 2$.

Theorem 1.1. Let $n \geq 2$, $p = 2$, and $\alpha, \beta \in \mathbb{R}$ satisfy (1.5). Denote by $C_{n,\alpha,\beta}$ the sharp constant in (1.3) with $p = 2$, i.e. the best constant to claim

$$(1.6) \quad \int_{\mathbb{R}^n} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla u|^2 dx \geq C_{n,\alpha,\beta} \int_{\mathbb{R}^n} |x'|^{2\alpha} |x|^{2\beta} u^2 dx, \quad \forall u \in C_c^1(\mathbb{R}^n).$$

Then we have

$$(1.7) \quad C_{n,\alpha,\beta} = \frac{(n-1+2\alpha)^2 - [\sqrt{\max(K,1)} - 1]^2}{4}$$

where $K = -4\beta(n+2\alpha+\beta) = (n+2\alpha)^2 - (n+2\alpha+2\beta)^2$.

For more general $p \geq 1$, we obtain the following partial result where the best constant is determined when $\beta \geq 0$.

Theorem 1.2. *Let $n \geq 2$, $p \geq 1$ and $\alpha, \beta \in \mathbb{R}$ satisfy (1.5). Denote still by $C_{n,\alpha,\beta}$ the best constant in (1.3), then*

$$C_{n,\alpha,\beta} = \left(\frac{n-1+p\alpha}{p} \right)^p \quad \text{for any } \beta \geq 0.$$

Moreover, $C_{n,\alpha,\beta} \geq \left(\frac{n-1+p\alpha+p\beta}{p} \right)^p$ if $\beta < 0$ and $p(\alpha+\beta) > 1-n$.

Furthermore, we give an alternative proof for the anisotropic L^p -Caffarelli-Kohn-Nirenberg inequalities, i.e a very special case of Li-Yan's general result (1.4) with $s = p = q > 1$ and $a = \frac{1}{p}$. Let $n \geq 2$, $p \geq 1$, we know that $|x'|^\mu |x|^{\gamma_2}$, $|x'|^\beta |x|^{\gamma_3}$, $|x'|^\alpha |x|^{\gamma_1} \in L^p_{loc}(\mathbb{R}^n)$ if and only if

$$(1.8) \quad \min(\alpha, \beta, \mu) > \frac{1-n}{p}, \quad \min(\alpha + \gamma_1, \mu + \gamma_2, \beta + \gamma_3) > -\frac{n}{p}.$$

Notice also that in our special case, the assumptions [8, (1.10)-(1.13)] are equivalent to

$$(1.9) \quad \alpha + \gamma_1 = \frac{\mu + \gamma_2 - 1}{p} + \frac{(p-1)(\beta + \gamma_3)}{p}, \quad \gamma_1 \leq \frac{\gamma_2 - 1}{p} + \frac{(p-1)\gamma_3}{p}.$$

Theorem 1.3. *Let $n \geq 2$, $p > 1$. Assume that $\alpha, \beta, \mu, \gamma_1, \gamma_2, \gamma_3$ satisfy (1.8)-(1.9), then for any $u \in C_c^1(\mathbb{R}^n)$, there holds*

$$(1.10) \quad \| |x|^{\gamma_2} |x'|^\mu \nabla u \|_{L^p(\mathbb{R}^n)}^{\frac{1}{p}} \| |x|^{\gamma_3} |x'|^\beta u \|_{L^p(\mathbb{R}^n)}^{\frac{p-1}{p}} \geq \frac{n+p(\alpha+\gamma_1)}{p} \| |x|^{\gamma_1} |x'|^\alpha u \|_{L^p(\mathbb{R}^n)}.$$

If moreover $\alpha = \beta = \mu$ and $\gamma_3 - \gamma_2 + 1 > 0$, the constant $\frac{n+p(\alpha+\gamma_1)}{p}$ is sharp.

Our approach departs from an elementary identity: Let $\Omega \subset \mathbb{R}^n$ be an open set, $V \in C^1(\Omega)$ and $f \in C^2(\Omega)$ be positive, then

$$(1.11) \quad \int_{\Omega} V |\nabla u|^2 dx = - \int_{\Omega} \frac{\operatorname{div}(V \nabla f)}{f} u^2 dx + \int_{\Omega} V f^2 \left| \nabla \left(\frac{u}{f} \right) \right|^2 dx, \quad \forall u \in C_c^1(\Omega).$$

The above equality can be showed using integration by parts; or by taking $\vec{F} = -\frac{\nabla f}{f}$ in the more general equality

$$(1.12) \quad \int_{\Omega} V |\nabla u|^2 dx = \int_{\Omega} \left[\operatorname{div}(V \vec{F}) - V |\vec{F}|^2 \right] u^2 dx + \int_{\Omega} V |\nabla u + u \vec{F}|^2 dx.$$

These identities suggest to find weighted Hardy or Poincaré inequalities by testing suitable positive functions $f \in C_c^1(\Omega)$, and provide a natural way to study Hardy type inequalities. This idea has been used in many situations in the literature, and summarized in [6]. For example, the Bessel pair with radial potential (V, W) introduced by Ghoussoub-Moradifam [5] is a special case of (1.11) for radial function $f(x) = f(|x|)$, since

$$f''(r) + \left(\frac{n-1}{r} + \frac{V'}{V} \right) f'(r) + \frac{cW}{V} f(r) = 0 \quad \text{is equivalent to say } -\operatorname{div}(V \nabla f) = cW f.$$

Furthermore, we remark that the last integral in (1.11) is zero if and only if u/f is a constant. It is well known that the optimal Hardy inequality cannot be reached in general, that is, the *best* choice of f does not belong to the proper functional space. However, we can check eventually sharpness of the subsequent weight $W := -\frac{\operatorname{div}(V \nabla f)}{f}$ by choosing appropriate functions u to approximate f .

The last term in (1.11) can be interpreted also as a kind of stability, since it measures in some sense the *distance* between u and the eventual linear space generated by the optimal choice f .

To prove Theorem 1.1, according to V , we will apply (1.11) with $f(x) = |x'|^\theta |x|^\lambda$, and try to optimize the subsequent weight W with suitable choice of the parameters $\theta, \lambda \in \mathbb{R}$. As θ or λ are allowed to be

negative, the corresponding anisotropic Hardy inequalities are firstly proved in $C_c^1(\mathbb{R}^n \setminus \{x' = 0\})$, then extended to $C_c^1(\mathbb{R}^n)$ by density argument. Moreover, we study the sharpness by trying to approximate the optimal choice of f .

An equality similar to (1.11) exists for general $p > 1$, where we replace the last integral by a Piconé type term, see [6, section 10.2]. Let (\mathcal{M}, g) be a Riemannian manifold, consider $V \in C^1(\mathcal{M})$ and $\vec{F} \in C^1(\mathcal{M}, T_g\mathcal{M})$, then for any $u \in C_c^1(\mathcal{M})$, there holds

$$(1.13) \quad \int_{\mathcal{M}} [V|\nabla u|^p + (p-1)V|\vec{F}|^p|u|^p]dg = \int_{\mathcal{M}} \operatorname{div} \left(V|\vec{F}|^{p-2}\vec{F} \right) |u|^p dg + \int_{\mathcal{M}} V\mathcal{R}(\nabla u, u\vec{F})dg,$$

where

$$\mathcal{R}(\vec{X}, \vec{Y}) = (p-1)\|\vec{Y}\|^p + \|\vec{X}\|^p + p\|\vec{Y}\|^{p-2}\langle \vec{Y}, \vec{X} \rangle \geq 0.$$

In particular, let $\vec{F} = -\frac{\nabla f}{f}$ and $\mathcal{M} = \Omega$, we get that (see also [2, 3, 4] with f depending on one variable)

$$(1.14) \quad \int_{\Omega} V|\nabla u|^p dx = - \int_{\Omega} \frac{\operatorname{div} (V|\nabla f|^{p-2}\nabla f)}{f^{p-1}} |u|^p dx + \int_{\Omega} V\mathcal{R} \left(\nabla u, -u\frac{\nabla f}{f} \right) dx.$$

Hence we obtain the L^p -Hardy inequality (with suitable V , f and u)

$$(1.15) \quad \int_{\Omega} V|\nabla u|^p dx \geq - \int_{\Omega} \frac{\operatorname{div} (V|\nabla f|^{p-2}\nabla f)}{f^{p-1}} |u|^p dx.$$

Here again, we need not any symmetry assumption on V or f , again the residual term in (1.14) is zero if and only if u/f is constant. Therefore we can proceed similarly as for L^2 case to handle Theorem 1.2.

Remark 1.4. *The identity (1.15) holds also for $p = 1$, if $V \in C^1(\Omega)$, $f \in C^{1,1}(\Omega)$ satisfies $|\nabla f| > 0$. In that case,*

$$\int_{\Omega} V|\nabla u| dx \geq - \int_{\Omega} \operatorname{div} \left(V \frac{\nabla f}{|\nabla f|} \right) |u| dx, \quad \forall u \in C_c^1(\Omega).$$

Moreover, notice that for any $\kappa > 0$, $(\kappa^{-1}V, \kappa^{\frac{1}{p-1}}\vec{F})$ does not change the subsequent weight $W = \operatorname{div}(V|\vec{F}|^{p-2}\vec{F})$ on the right hand side of (1.13). Taking

$$\kappa_0^{\frac{p}{p-1}} = \frac{\int_{\Omega} V|\nabla u|^p dg}{\int_{\Omega} V|\vec{F}|^p |u|^p dg},$$

we obtain a special weighted L^p -Caffarelli-Kohn-Nirenberg inequality as follows.

$$(1.16) \quad \begin{aligned} & \left(\int_{\Omega} V|\nabla u|^p dg \right)^{\frac{1}{p}} \left(\int_{\Omega} V|\vec{F}|^p |u|^p dg \right)^{\frac{p-1}{p}} \\ &= \frac{1}{p} \int_{\Omega} \operatorname{div}(V|\vec{F}|^{p-2}\vec{F})|u|^p dg + \int_{\omega} \frac{V}{p\kappa_0} \mathcal{R} \left(\nabla u, u\kappa_0^{\frac{1}{p-1}}\vec{F} \right) dg \\ &\geq \frac{1}{p} \int_{\Omega} \operatorname{div}(V|\vec{F}|^{p-2}\vec{F})|u|^p dg. \end{aligned}$$

We will choose suitable \vec{F} to prove Theorem 1.3.

Remark 1.5. *For Li-Yan's inequality (1.4) with $p = 2$ and $q = 1$, Bao-Chen [1] considered recently the existence, the symmetry and the symmetry breaking region of extremal functions.*

2. PROOF OF THEOREM 1.1.

Let $n \geq 2$ and $x' = (x_1, \dots, x_{n-1}, 0)$ for $x = (x_i) \in \mathbb{R}^n$. Let $V = |x'|^{2\alpha+2}|x|^{2\beta}$ with α, β satisfying (1.5) with $p = 2$. Consider $f = f_1 f_2$, then

$$\frac{\operatorname{div}(V\nabla f)}{f} = \frac{\operatorname{div}(V\nabla f_1)}{f_1} + \frac{\operatorname{div}(V\nabla f_2)}{f_2} + 2V \frac{\nabla f_1 \cdot \nabla f_2}{f_1 f_2} =: I_1 + I_2 + I_3.$$

Choose now $f_1 = |x'|^\theta$, $f_2 = |x|^\lambda$, directly calculus yields that in $\mathbb{R}^n \setminus \{x' = 0\}$,

$$\nabla V = \left[(2\alpha + 2) \frac{x'}{|x'|^2} + 2\beta \frac{x}{|x|^2} \right] V, \quad \nabla f_1 = \theta |x'|^{\theta-2} x', \quad \nabla f_2 = \lambda |x|^{\lambda-2} x$$

and

$$\Delta f_1 = \theta(n-3+\theta)|x'|^{\theta-2}, \quad \Delta f_2 = \lambda(n-2+\lambda)|x|^{\lambda-2}.$$

Hence

$$\begin{aligned} I_1 &= \frac{\nabla V \cdot \nabla f_1 + V \Delta f_1}{f_1} = \left[\frac{\theta(2\alpha+2)}{|x'|^2} + \frac{2\theta\beta}{|x|^2} \right] V + V \frac{\theta(n-3+\theta)}{|x'|^2} \\ &= \left[\frac{\theta(n-1+\theta+2\alpha)}{|x'|^2} + \frac{2\theta\beta}{|x|^2} \right] V, \end{aligned}$$

and

$$I_2 = \frac{\nabla V \cdot \nabla f_2 + V \Delta f_2}{f_2} = \frac{\lambda(n+\lambda+2\alpha+2\beta)}{|x|^2} V, \quad I_3 = \frac{2\theta\lambda}{|x|^2} V.$$

Therefore we get

$$\begin{aligned} (2.1) \quad -\frac{\operatorname{div}(V\nabla f)}{f} &= -\left[\frac{\theta(n-1+\theta+2\alpha)}{|x'|^2} + \frac{\lambda(n+2\alpha+2\beta+2\theta+\lambda)+2\beta\theta}{|x|^2} \right] V \\ &= H_1(\theta, \lambda) \frac{V}{|x'|^2} + H_2(\theta, \lambda) \frac{V x_n^2}{|x'|^2 |x|^2} \end{aligned}$$

where

$$H_1(\theta, \lambda) = -\left[\theta(n-1+\theta+2\alpha+2\beta) + \lambda(n+2\alpha+2\beta+2\theta+\lambda) \right] := H(\theta) - H_2(\theta, \lambda)$$

with

$$H(\theta) = -\theta(n-1+2\alpha+\theta), \quad H_2(\theta, \lambda) = \lambda(n+2\alpha+2\beta+2\theta+\lambda) + 2\beta\theta.$$

Seeing Li-Yan's estimate (1.3), we aim to find the maximum value of H_1 under the constraint $H_2 \geq 0$. As $\lim_{|\theta| \rightarrow \infty} H(\theta) = -\infty$, and $\lim_{|\lambda| \rightarrow \infty} H_1(\theta, \lambda) = -\infty$ uniformly for bounded θ , $\max_{H_2 \geq 0} H_1$ exists.

It is easy to see that $\partial_\theta H_1 - \partial_\lambda H_1 \equiv 1$, hence H_1 has no critical point in \mathbb{R}^2 and $\max_{H_2 \geq 0} H_1$ is reached on the subset $\{H_2 = 0\}$, i.e. when

$$(2.2) \quad \lambda(n+2\alpha+2\beta+2\theta+\lambda) + 2\beta\theta = 0.$$

If $K = -4\beta(n+2\alpha+\beta) \leq 0$, then for any $\theta \in \mathbb{R}$, there exists λ such that $H_2(\theta, \lambda) = 0$, because the discriminant for the quadratic equation (2.2) of λ satisfies

$$(n+2\alpha+2\beta+2\theta)^2 - 8\beta\theta = 4\theta^2 + 4(n+2\alpha)\theta - (n+2\alpha+2\beta)^2 \geq 0 \quad \text{in } \mathbb{R}.$$

This means that

$$(2.3) \quad \max_{H_2=0} H_1 = \max_{\mathbb{R}} H(\theta) = H(\theta_0) = \frac{(n-1+2\alpha)^2}{4}, \quad \text{where } \theta_0 = \frac{1-n-2\alpha}{2}.$$

Let $K > 0$, then (2.2) holds true for $\lambda \neq -\beta$ and

$$\theta = \theta(\lambda) := -\frac{\lambda(n+2\alpha+2\beta+\lambda)}{2(\lambda+\beta)}.$$

Clearly,

$$(2.4) \quad \min_{\lambda < -\beta} \theta(\lambda) = \theta(\lambda_1) = -\frac{(n+2\alpha) - \sqrt{K}}{2} =: \theta_1, \quad \text{with } \lambda_1 = -\beta - \frac{\sqrt{K}}{2}$$

and

$$\lim_{\lambda \rightarrow -\infty} \theta(\lambda) = \lim_{\lambda \rightarrow -\beta^-} \theta(\lambda) = +\infty.$$

If $K \in (0, 1]$, then $\theta(\lambda_0) = \theta_0$ for $\lambda_0 = -\beta - \frac{1+\sqrt{1-K}}{2}$, which yields

$$(2.5) \quad \max_{H_2=0} H_1 = H_1(\theta_0, \lambda_0) = \theta_0^2, \quad \text{if } K \in (0, 1].$$

Consider now $K > 1$, we can check that

$$\max_{H_2=0, \lambda < -\beta} H_1 = H(\theta_1).$$

By the same, for any $K > 0$, there holds

$$\max_{H_2=0, \lambda > -\beta} H_1 = H(\theta_2) \quad \text{with } \theta_2 = -\frac{(n+2\alpha) + \sqrt{K}}{2}.$$

Notice that $H(\theta_2) < H(\theta_1)$ for any $K > 0$, hence for $K > 1$, there holds

$$\begin{aligned} \max_{H_2=0} H_1 &= H_1(\theta_1, \lambda_1) = H(\theta_1) = -\theta_1(n-1+2\alpha+\theta_1) \\ (2.6) \quad &= \frac{n+2\alpha-\sqrt{K}}{2} \times \frac{n-2+2\alpha+\sqrt{K}}{2} \\ &= \frac{(n-1+2\alpha)^2}{4} - \frac{(\sqrt{K}-1)^2}{4}. \end{aligned}$$

Finally, we obtain

$$\max_{H_2 \geq 0} H_1 = \max_{H_2=0} H_1 = \frac{(n-1+2\alpha)^2 - [\sqrt{\max(K, 1)} - 1]^2}{4} = C_{n, \alpha, \beta}$$

with $K = -4\beta(n+2\alpha+\beta)$. Seeing (2.1) and the equality (1.11),

$$(2.7) \quad \int_{\mathbb{R}^n} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla u|^2 dx \geq C_{n, \alpha, \beta} \int_{\mathbb{R}^n} |x'|^{2\alpha+2-2} |x|^{2\beta} u^2 dx, \quad \forall u \in C_c^1(\mathbb{R}^n \setminus \{x' = 0\}).$$

Under the assumption (1.5) with $p = 2$, i.e. $2\alpha > 1-n$, $2(\alpha+\beta) > -n$, there holds (using the spherical coordinates)

$$\lim_{\epsilon \rightarrow 0^+} \int_{|x'| \leq M\epsilon, |x| \leq R} |x'|^{2\alpha} |x|^{2\beta} dx = 0, \quad \forall M, R > 0.$$

For any $u \in C_c^1(\mathbb{R}^n)$, we consider $u_\epsilon(x) = u(x) - u(x)\eta(|x'|/\epsilon)$ for $\epsilon \in (0, 1)$, with a standard cut-off function $\eta \in C_c^1(\mathbb{R})$. Applying (2.7) to u_ϵ and tending $\epsilon \rightarrow 0^+$, we can claim the estimate (1.6).

Now we show the sharpness of the constants $C_{n, \alpha, \beta}$ in (1.7). We will use the spherical coordinates for $n \geq 2$, that is

$$\begin{cases} x_1 &= r \sin \varphi_1 \sin \varphi_2 \cdots \sin \varphi_{n-2} \sin \varphi_{n-1}, \\ x_2 &= r \sin \varphi_1 \sin \varphi_2 \cdots \sin \varphi_{n-2} \cos \varphi_{n-1}, \\ &\dots \\ x_{n-1} &= r \sin \varphi_1 \cos \varphi_2, \\ x_n &= r \cos \varphi_1 \end{cases}$$

where $r \in \mathbb{R}_+$, $\varphi_k \in [0, \pi]$ for $1 \leq k \leq n-2$ if $n \geq 3$ and $\varphi_{n-1} \in [0, 2\pi]$, so

$$dx = r^{n-1} (\sin \varphi_1)^{n-2} (\sin \varphi_2)^{n-3} \cdots \sin \varphi_{n-2} dr d\varphi_1 \cdots d\varphi_{n-1}.$$

Let $v(x) = h(s)g(r)$, $s = |x'|$ and $r = |x|$, then

$$\nabla v(x) = h'(s)g(r) \frac{x'}{s} + h(s)g'(r) \frac{x}{r},$$

and

$$|\nabla v(x)|^2 = h'(s)^2 g^2(r) + h^2(s) g'(r)^2 + 2h(s)h'(s)g(r)g'(r) \frac{s}{r}.$$

Hence

$$\begin{aligned} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla v(x)|^2 &= h'(s)^2 g^2(r) s^{2\alpha+2} r^{2\beta} + h^2(s) g'(r)^2 s^{2\alpha+2} r^{2\beta} + 2h(s)h'(s)g(r)g'(r) s^{2\alpha+3} r^{2\beta-1} \\ &= h'(s)^2 g^2(r) (\sin \varphi_1)^{2\alpha+2} r^{2\beta+2\alpha+2} + h^2(s) g'(r)^2 (\sin \varphi_1)^{2\alpha+2} r^{2\beta+2\alpha+2} \\ &\quad + 2h(s)h'(s)g(r)g'(r) (\sin \varphi_1)^{2\alpha+3} r^{2\beta+2\alpha+2} \\ &=: J_1 + J_2 + J_3. \end{aligned}$$

Denote $\Sigma = (0, \pi)$, we have

$$\begin{aligned} & \int_{\mathbb{R}^n} J_1 dx \\ &= \int_{\mathbb{R}_+ \times \Sigma^{n-2} \times (0, 2\pi)} h'(r \sin \varphi_1)^2 g^2(r) r^{n+2\alpha+2\beta+1} (\sin \varphi_1)^{n+2\alpha} (\sin \varphi_2)^{n-3} \dots \sin \varphi_{n-2} dr d\varphi_1 \dots d\varphi_{n-1} \\ &= \omega_{n-1} \int_{\mathbb{R}_+ \times \Sigma} h'(r \sin \varphi_1)^2 g^2(r) r^{n+2\alpha+2\beta+1} (\sin \varphi_1)^{n+2\alpha} dr d\varphi_1 \end{aligned}$$

where ω_{n-1} stands for the volume of the unit sphere in \mathbb{R}^{n-1} . For the estimates of J_i , we consider three subcases.

Case $K > 1$. Seeing (2.6), we choose the test function $v = hg$ with $h(s) = s^{\theta_1}$, $g(r) = (r^2 + \epsilon^2)^{\frac{\lambda}{2}} \eta(r)$ and $\eta \in C_c^1(\mathbb{R})$ a standard cut-off function. Then

$$h'(r \sin \varphi_1)^2 g^2(r) r^{n+2\alpha+2\beta+1} (\sin \varphi_1)^{n+2\alpha} = \theta_1^2 (\sin \varphi_1)^{\sqrt{K}-2} g^2(r) r^{2\beta-1+\sqrt{K}},$$

so

$$\int_{\mathbb{R}^n} J_1 dx = \omega_{n-1} \theta_1^2 \int_{\Sigma} (\sin \varphi_1)^{\sqrt{K}-2} d\varphi_1 \int_0^{+\infty} g^2(r) r^{2\beta-1+\sqrt{K}} dr.$$

For any $\lambda > -1$, there holds

$$(2.8) \quad \int_0^\pi (\sin s)^\lambda ds = B\left(\frac{\lambda+1}{2}, \frac{1}{2}\right),$$

where $B(\cdot, \cdot)$ stands for Euler's Beta function

$$B(t, \gamma) := \int_0^1 s^{t-1} (1-s)^{\gamma-1} ds, \quad \forall t, \gamma > 0.$$

We know also that

$$(2.9) \quad B(t+1, \gamma) = \frac{t}{t+\gamma} B(t, \gamma), \quad \forall t, \gamma > 0.$$

On the other hand, for any $\epsilon \in (0, 1)$,

$$\begin{aligned} \int_0^{+\infty} g^2(r) r^{2\beta-1+\sqrt{K}} dr &= \int_0^{+\infty} (r^2 + \epsilon^2)^{\lambda_1} r^{2\beta-1+\sqrt{K}} \eta^2(r) dr \\ &= \int_0^{+\infty} (t^2 + 1)^{-\beta - \frac{\sqrt{K}}{2}} t^{2\beta-1+\sqrt{K}} \eta^2(\epsilon t) dt \\ &= -\ln \epsilon + O(1). \end{aligned}$$

Here and after, $O(1)$ means a quantity uniformly bounded for $\epsilon \in (0, 1)$. Indeed, we used the following fact.

Lemma 2.1. *Assume that $\xi \in L_{loc}^1(\mathbb{R}_+)$ and $\xi(s) - s^{-1} \in L^1([1, \infty))$. Let $\zeta \in C_c^1(\mathbb{R})$ be a standard cut-off function, then there is $C > 0$ such that for any $\epsilon \in (0, 1)$,*

$$\left| \int_0^{+\infty} \xi(t) \zeta(\epsilon t) dt + \ln \epsilon \right| \leq C.$$

There holds then

$$\int_{\mathbb{R}^n} J_1 dx = \omega_{n-1} \theta_1^2 B\left(\frac{\sqrt{K}-1}{2}, \frac{1}{2}\right) |\ln \epsilon| + O(1), \quad \forall \epsilon \in (0, 1).$$

Similarly, as

$$g'^2(r) = \lambda_1^2 r^2 (r^2 + \epsilon^2)^{\lambda_1-2} \eta^2(r) + 2\lambda_1 r (r^2 + \epsilon^2)^{\lambda_1-1} \eta \eta'(r) + (r^2 + \epsilon^2)^{\lambda_1} \eta'^2(r),$$

for $\epsilon \in (0, 1)$ we have

$$\begin{aligned}
\int_0^{+\infty} g'^2(r) r^{2\beta+1+\sqrt{K}} dr &= \lambda_1^2 \int_0^{+\infty} (r^2 + \epsilon^2)^{\lambda_1-2} r^{2\beta+3+\sqrt{K}} \eta^2(r) dr \\
&\quad + 2\lambda_1 \int_0^{+\infty} (r^2 + \epsilon^2)^{\lambda_1-1} r^{2\beta+2+\sqrt{K}} \eta \eta'(r) dr \\
&\quad + \int_0^{+\infty} (r^2 + \epsilon^2)^{\lambda_1} r^{2\beta+1+\sqrt{K}} \eta'^2(r) dr \\
&= \lambda_1^2 \int_0^{+\infty} (t^2 + 1)^{-\beta - \frac{\sqrt{K}}{2} - 2} t^{2\beta+3+\sqrt{K}} \eta^2(\epsilon t) dt + O(1) \\
&= \lambda_1^2 |\ln \epsilon| + O(1).
\end{aligned}$$

We see that

$$\begin{aligned}
\int_{\mathbb{R}^n} J_2 dx &= \omega_{n-1} \int_{\Sigma} (\sin \varphi_1)^{\sqrt{K}} d\varphi_1 \int_0^{+\infty} g'(r)^2 r^{2\beta+1+\sqrt{K}} dr \\
&= \omega_{n-1} \lambda_1^2 B\left(\frac{\sqrt{K}+1}{2}, \frac{1}{2}\right) |\ln \epsilon| + O(1),
\end{aligned}$$

and also

$$\begin{aligned}
\int_{\mathbb{R}^n} J_3 dx &= 2\omega_{n-1} \times \theta_1 \int_{\Sigma} (\sin \varphi_1)^{\sqrt{K}} d\varphi_1 \int_0^{+\infty} g(r) g'(r) r^{2\beta+\sqrt{K}} dr \\
&= 2\omega_{n-1} \theta_1 B\left(\frac{\sqrt{K}+1}{2}, \frac{1}{2}\right) \int_0^{+\infty} g(r) g'(r) r^{2\beta+\sqrt{K}} dr \\
&= 2\omega_{n-1} \theta_1 \lambda_1 B\left(\frac{\sqrt{K}+1}{2}, \frac{1}{2}\right) |\ln \epsilon| + O(1).
\end{aligned}$$

Finally, using (2.9), we arrive at

$$(2.10) \quad \int_{\mathbb{R}^n} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla v|^2 dx = A \omega_{n-1} B\left(\frac{\sqrt{K}-1}{2}, \frac{1}{2}\right) |\ln \epsilon| + O(1)$$

where

$$A = \theta_1^2 + (\lambda_1^2 + 2\theta_1 \lambda_1) \frac{\sqrt{K}-1}{\sqrt{K}}.$$

Recall that $\theta_1 = \frac{-(n+2\alpha)+\sqrt{K}}{2}$, $\lambda_1 = -\beta - \frac{\sqrt{K}}{2}$ and $K = -4\beta(n+2\alpha+\beta)$, so $\lambda_1^2 = 2\beta\theta_1$ and

$$\begin{aligned}
A &= \theta_1^2 + \theta_1(2\beta + 2\lambda_1) \frac{\sqrt{K}-1}{\sqrt{K}} = \theta_1^2 + \theta_1(1 - \sqrt{K}) \\
&= \frac{(n-1+2\alpha)^2}{4} - \frac{(\sqrt{K}-1)^2}{4}.
\end{aligned}$$

Moreover,

$$\begin{aligned}
\int_{\mathbb{R}^n} |x'|^{2\alpha} |x|^{2\beta} v^2(x) dx &= \omega_{n-1} \int_{\mathbb{R}_+ \times \Sigma} (\sin \varphi_1)^{n+2\alpha+2\theta_1-2} g^2(r) r^{n+2\alpha+2\beta+2\theta_1-1} dr d\varphi_1 \\
&= \omega_{n-1} \int_{\mathbb{R}_+ \times \Sigma} (\sin \varphi_1)^{\sqrt{K}-2} g^2(r) r^{2\beta+\sqrt{K}-1} dr d\varphi_1 \\
&= \omega_{n-1} B\left(\frac{\sqrt{K}-1}{2}, \frac{1}{2}\right) \int_0^{+\infty} g^2(r) r^{2\beta+\sqrt{K}-1} dr \\
&= \omega_{n-1} B\left(\frac{\sqrt{K}-1}{2}, \frac{1}{2}\right) |\ln \epsilon| + O(1).
\end{aligned}$$

Therefore,

$$\lim_{\epsilon \rightarrow 0^+} \frac{\int_{\mathbb{R}^n} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla v(x)|^2 dx}{\int_{\mathbb{R}^n} |x'|^{2\alpha} |x|^{2\beta} v^2(x) dx} = \frac{(n-1+2\alpha)^2}{4} - \frac{(\sqrt{K}-1)^2}{4}.$$

Notice that the function $h(s) = s^{\theta_1}$ is not smooth at $s = 0$, however as all involved integrals converge, we may use eventually a family of smooth functions to approximate h , so we omit the details. This means that for $K > 1$, the constant $\frac{(n-1+2\alpha)^2}{4} - \frac{(\sqrt{K}-1)^2}{4}$ is sharp to claim (1.6).

The analysis for other cases are similar, we will go through quickly.

Case $K = 1$. Here we take $v(x) = h(s)g(r)$ with $h(s) = s^{\theta_0+\sigma}$, $g(r) = (r^2 + \epsilon^2)^{\frac{\lambda_0-\sigma}{2}} \eta(r)$, where $s = |x'|$, $r = |x|$, $\sigma > 0$ and $\lambda_0 = -\beta - \frac{1}{2}$. Remark that (1.5) with $p = 2$ yields

$$(2.11) \quad n + 2\alpha + \beta = \frac{n + 2\alpha + 2\beta}{2} + \frac{n + 2\alpha}{2} > \frac{1}{2}.$$

Therefore $K = 1$ implies $2\beta \in (-1, 0)$. We get then

$$\begin{aligned} \int_{\mathbb{R}^n} |x'|^{2\alpha} |x|^{2\beta} v^2(x) dx &= \omega_{n-1} \int_0^\pi (\sin \varphi_1)^{2\sigma-1} d\varphi_1 \times \int_0^{+\infty} r^{2\beta+2\sigma} (r^2 + \epsilon^2)^{-\beta-\frac{1}{2}-\sigma} \eta^2(r) dr \\ &= \omega_{n-1} B\left(\sigma, \frac{1}{2}\right) |\ln \epsilon| + O_\sigma(1). \end{aligned}$$

Here $\xi(r) = r^{2\beta+2\sigma} (r^2 + \epsilon^2)^{-\beta-\frac{1}{2}-\sigma}$ satisfies all assumptions of Lemma 2.1, and $O_\sigma(1)$ stands for a quantity uniformly bounded for $\epsilon \in (0, 1)$ when $\sigma > 0$ is fixed. By the same, there holds

$$\begin{aligned} \int_{\mathbb{R}^n} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla v|^2 dx &= \omega_{n-1} |\ln \epsilon| \left[(\theta_0 + \sigma)^2 B\left(\sigma, \frac{1}{2}\right) + (\lambda_0^2 + 2\lambda_0\theta_0 + 2\lambda_0\sigma) B\left(\sigma + 1, \frac{1}{2}\right) \right] + O_\sigma(1) \\ &= \omega_{n-1} |\ln \epsilon| \left[(\theta_0 + \sigma)^2 + (\lambda_0^2 + 2\lambda_0\theta_0 + 2\lambda_0\sigma) \frac{\sigma}{\sigma+1} \right] B\left(\sigma, \frac{1}{2}\right) + O_\sigma(1). \end{aligned}$$

Taking first $\epsilon \rightarrow 0^+$ and secondly $\sigma \rightarrow 0^+$, we see that $C_{n,\alpha,\beta} \leq \theta_0^2$ for $K = 1$.

Case $K < 1$. Let $v(x) = h(s)g(r)$ with $h(s) = s^{\theta_0+\sigma}$, $g(r) = (r^2 + \epsilon^2)^{\frac{\lambda_0}{2}} \eta(r)$. Remark that $\beta > -\frac{1}{2}$ by (2.11), let

$$0 < \sigma < \frac{\sqrt{1-K}}{2} \quad \text{and} \quad \lambda_0 = -\beta - \frac{1 + \sqrt{1-K}}{2}.$$

The above choice is motivated by (2.5). There holds then

$$\begin{aligned} \int_{\mathbb{R}^n} |x'|^{2\alpha} |x|^{2\beta} v^2(x) dx &= \omega_{n-1} \int_0^\pi (\sin \varphi_1)^{2\sigma-1} d\varphi_1 \times \int_0^{+\infty} r^{2\beta+2\sigma} (r^2 + \epsilon^2)^{-\beta-\frac{1+\sqrt{1-K}}{2}} \eta^2(r) dr \\ &= \omega_{n-1} B\left(\sigma, \frac{1}{2}\right) \epsilon^{2\sigma-\sqrt{1-K}} \times \int_0^{+\infty} t^{2\beta+2\sigma} (t^2 + 1)^{-\beta-\frac{1+\sqrt{1-K}}{2}} \eta^2(\epsilon t) dt \\ &= \omega_{n-1} B\left(\sigma, \frac{1}{2}\right) \epsilon^{2\sigma-\sqrt{1-K}} \times [A_1 + o_\sigma(1)]. \end{aligned}$$

Here $o_\sigma(1)$ stands for a quantity tending to zero as ϵ goes to 0 for fixed $\sigma > 0$, and

$$A_1 = \int_0^{+\infty} t^{2\beta+2\sigma} (t^2 + 1)^{-\beta-\frac{1+\sqrt{1-K}}{2}} dt < \infty.$$

Similarly, we get

$$\begin{aligned} \int_{\mathbb{R}^n} |x'|^{2\alpha+2} |x|^{2\beta} |\nabla v|^2 dx &= \omega_{n-1} \epsilon^{2\sigma-\sqrt{1-K}} \times [A_1 + o_\sigma(1)] (\theta_0 + \sigma)^2 B\left(\sigma, \frac{1}{2}\right) \\ &\quad + \omega_{n-1} \epsilon^{2\sigma-\sqrt{1-K}} \times [A_2 + o_\sigma(1)] B\left(\sigma + 1, \frac{1}{2}\right) \\ &= \omega_{n-1} \epsilon^{2\sigma-\sqrt{1-K}} [A_1 \theta_0^2 + O(\sigma) + o_\sigma(1)] B\left(\sigma, \frac{1}{2}\right) \end{aligned}$$

with

$$A_2 = \lambda_0^2 \int_0^{+\infty} t^{2\beta+2\sigma+4} (t^2+1)^{-\beta-\frac{1+\sqrt{1-K}}{2}-2} dt + 2\lambda_0\theta_0 \int_0^{+\infty} t^{2\beta+2\sigma+2} (t^2+1)^{-\beta-\frac{1+\sqrt{1-K}}{2}-1} dt < \infty.$$

Taking first $\epsilon \rightarrow 0^+$ and secondly $\sigma \rightarrow 0^+$, we see that $C_{n,\alpha,\beta} \leq \theta_0^2$ for $K < 1$. \square

Remark 2.2. Take $\alpha = \beta = -\frac{1}{2}$ and $n \geq 3$, we have

$$(2.12) \quad \int_{\mathbb{R}^n} \frac{|x'|}{|x|} |\nabla u|^2 dx \geq \left[\frac{n^2 - 6n + 6}{4} + \frac{\sqrt{2n-3}}{2} \right] \int_{\mathbb{R}^n} \frac{u^2}{|x'| |x|} dx, \quad \forall u \in C_c^1(\mathbb{R}^n).$$

Here the constant $\frac{n^2-6n+6}{4} + \frac{\sqrt{2n-3}}{2}$ is sharp. In particular, the best constant for (1.2) is $\frac{2\sqrt{3}-3}{4}$.

Remark 2.3. The estimate (2.12) is trivial when $n = 2$, but an anisotropic Leray type inequality exists. Let \mathbb{B}^2 denote the open unit disc in \mathbb{R}^2 , there holds

$$\int_{\mathbb{B}^2} \frac{|x_1|}{|x|} |\nabla u|^2 dx \geq \frac{1}{4} \int_{\mathbb{B}^2} \frac{|x_1|}{|x|^3 (\ln|x|)^2} u^2 dx, \quad \forall u \in C_c^1(\mathbb{B}^2).$$

Here we consider $f(x) = \sqrt{-\ln|x|}$ and check that

$$-\frac{\operatorname{div}(|x_1||x|^{-1}\nabla f)}{f} = \frac{|x_1|}{4|x|^3(\ln|x|)^2} \quad \text{in } \mathbb{B}^2 \setminus \{0\}.$$

3. PROOF OF THEOREM 1.2

Let $V = |x'|^{(\alpha+1)p}|x|^{\beta p}$ and $f = |x'|^\gamma$, then

$$\nabla V = \left[(\alpha+1)p \frac{x'}{|x'|^2} + \beta p \frac{x}{|x|^2} \right] V,$$

and $\nabla f = \gamma|x'|^{\gamma-2}x'$, $\Delta f = \gamma(n-3+\gamma)|x'|^{\gamma-2}$. There hold also

$$|\nabla f|^{p-2} = |\gamma|^{p-2}|x'|^{(\gamma-1)(p-2)}, \quad \nabla(|\nabla f|^{p-2}) = |\gamma|^{p-2}(\gamma-1)(p-2)|x'|^{\gamma p-2\gamma-p}x'.$$

According to the Hardy inequality (1.15), we will calculate

$$-\frac{\operatorname{div}(V|\nabla f|^{p-2}\nabla f)}{f^{p-1}} = -\left[\frac{\operatorname{div}(V\nabla f)|\nabla f|^{p-2}}{f^{p-1}} + \frac{V\nabla f \cdot \nabla(|\nabla f|^{p-2})}{f^{p-1}} \right] =: -(K_1 + K_2).$$

More precisely,

$$\frac{\operatorname{div}(V\nabla f)}{f} = \frac{\nabla V \cdot \nabla f + V\Delta f}{f} = \left[\frac{\gamma(n-3+(\alpha+1)p+\gamma)}{|x'|^2} + \frac{\gamma\beta p}{|x|^2} \right] V.$$

This yields

$$K_1 = \frac{\operatorname{div}(V\nabla f)}{f} \times \frac{|\nabla f|^{p-2}}{f^{p-2}} = V|\gamma|^{p-2} \left[\frac{\gamma(n-3+(\alpha+1)p+\gamma)}{|x'|^p} + \frac{\gamma\beta p}{|x'|^{p-2}|x|^2} \right].$$

On the other hand,

$$K_2 = V|\gamma|^{p-2} \frac{\gamma(\gamma-1)(p-2)}{|x'|^p}.$$

Hence

$$\begin{aligned} W &= -\frac{\operatorname{div}(V|\nabla f|^{p-2}\nabla f)}{f^{p-1}} = -V|\gamma|^{p-2} \left[\frac{\gamma(n-3+(\alpha+1)p+\gamma) + \gamma(\gamma-1)(p-2)}{|x'|^p} + \frac{\gamma\beta p}{|x'|^{p-2}|x|^2} \right] \\ &= V|x'|^{-p} \left\{ -|\gamma|^{p-2} \gamma [(p-1)\gamma + n-1 + p\alpha] - |\gamma|^{p-2} \gamma \beta p \frac{|x'|^2}{|x|^2} \right\} \\ &=: V|x'|^{-p} \left[\overline{H}_1(\gamma) + \overline{H}_2(\gamma) \frac{|x'|^2}{|x|^2} \right]. \end{aligned}$$

We consider respectively two cases according to the sign of β .

Case $\beta \geq 0$. Recall that $p\alpha > 1-n$. Let $n-1+p\alpha = -p\gamma_0$, then $\gamma_0 < 0$,

$$\max_{\mathbb{R}} \overline{H}_1(\gamma) = \overline{H}_1(\gamma_0) = |\gamma_0|^p \quad \text{and} \quad \overline{H}_2(\gamma_0) = \beta p |\gamma_0|^{p-1} \geq 0.$$

Thanks to (1.14) or (1.15), we have

$$\| |x|^\beta |x'|^{\alpha+1} \nabla u \|_{L^p(\mathbb{R}^n)} \geq |\gamma_0|^p \| |x|^\beta |x'|^\alpha u \|_{L^p(\mathbb{R}^n)}, \quad \forall u \in C_c^1(\mathbb{R}^n \setminus \{x' = 0\}).$$

Recall that $|x|^\beta |x'|^\alpha \in L_{loc}^p(\mathbb{R}^n)$ under the condition (1.5), similarly as for the case $p = 2$, we can extend the above estimate for $u \in C_c^1(\mathbb{R}^n)$ by approximation, so $C_{n,\alpha,\beta} \geq |\gamma_0|^p$.

Now we prove the sharpness of the above estimate. Consider $v = |x'|^\gamma g(r)$ with $g(r) = (r^2 + \epsilon^2)^{\frac{\lambda}{2}} \eta$, a cut-off function η and

$$\gamma = -\frac{n-1+p\alpha}{p} + \sigma = \gamma_0 + \sigma, \quad \lambda = -\beta - \frac{1}{p} - \sigma, \quad \epsilon, \sigma > 0.$$

Then for $\epsilon \in (0, 1)$, applying Lemma 2.1 and (2.8),

$$(3.1) \quad \begin{aligned} \int_{\mathbb{R}^n} |x'|^{p\alpha} |x|^{p\beta} |v|^p dx &= \omega_{n-1} \int_0^\pi (\sin \varphi_1)^{n-2+(\alpha+\gamma)p} d\varphi_1 \int_0^{+\infty} r^{\beta p} g^p(r) dr \\ &= \omega_{n-1} B\left(\frac{p\sigma}{2}, \frac{1}{2}\right) \times |\ln \epsilon| + O_\sigma(1) \end{aligned}$$

where $O_\sigma(1)$ stands for a quantity uniformly bounded for $\epsilon \in (0, 1)$ and fixed $\sigma \in (0, 1)$. On the other hand,

$$\begin{aligned} |\nabla v|^2 &= |\gamma |x'|^{\gamma-2} x' g(r) + |x'|^\gamma g'(r) \frac{x}{r}|^2 \\ &= |x'|^{2\gamma-2} \left[\gamma^2 g^2(r) + |x'|^2 g'^2(r) + 2\gamma \frac{|x'|^2}{r} g g'(r) \right] =: |x'|^{2\gamma-2} G(r). \end{aligned}$$

In \mathbb{B}^n the unit ball of \mathbb{R}^n , as $\eta \equiv 1$, we have

$$G(r) = \gamma^2 (r^2 + \epsilon^2)^\lambda + \lambda^2 (r^2 + \epsilon^2)^{\lambda-2} |x'|^2 r^2 + 2\gamma \lambda (r^2 + \epsilon^2)^{\lambda-1} |x'|^2.$$

Therefore

$$\begin{aligned} |x'|^{p(\alpha+1)} |x|^{p\beta} |\nabla v|^p &= |x'|^{p(\gamma-1)} G^{\frac{p}{2}}(r) \\ &= |x'|^{p(\alpha+\gamma)} |x|^{p\beta} (r^2 + \epsilon^2)^{\frac{p\lambda}{2}} \left(\gamma^2 + \lambda^2 \frac{|x'|^2 r^2}{(r^2 + \epsilon^2)^2} + 2\gamma \lambda \frac{|x'|^2}{r^2 + \epsilon^2} \right)^{\frac{p}{2}}. \end{aligned}$$

Let $\epsilon, \sigma \in (0, 1)$, clearly

$$\left| \lambda^2 \frac{|x'|^2 r^2}{(r^2 + \epsilon^2)^2} + 2\gamma \lambda \frac{|x'|^2}{r^2 + \epsilon^2} \right| \leq C \frac{|x'|^2}{r^2 + \epsilon^2} \quad \text{in } \mathbb{R}^n.$$

By mean value theorem, as $|x'| \leq r$, there holds

$$\left| \left(\gamma^2 + \lambda^2 \frac{|x'|^2 r^2}{(r^2 + \epsilon^2)^2} + 2\gamma \lambda \frac{|x'|^2}{r^2 + \epsilon^2} \right)^{\frac{p}{2}} - |\gamma|^p \right| \leq C \frac{|x'|^2}{r^2 + \epsilon^2} \quad \text{in } \mathbb{R}^n.$$

Using spherical coordinates, we see that

$$(3.2) \quad \begin{aligned} &\int_{\mathbb{B}^n} |x'|^{p(\alpha+1)} |x|^{p\beta} |\nabla v|^p dx \\ &= \omega_{n-1} \int_0^\pi (\sin \varphi_1)^{n-2+p(\alpha+\gamma)} \\ &\quad \times \int_0^1 r^{n-1+p(\alpha+\beta+\gamma)} (r^2 + \epsilon^2)^{\frac{p\lambda}{2}} \left(\gamma^2 + \lambda^2 \frac{|x'|^2 r^2}{(r^2 + \epsilon^2)^2} + 2\gamma \lambda \frac{|x'|^2}{r^2 + \epsilon^2} \right)^{\frac{p}{2}} dr \\ &=: \omega_{n-1} |\gamma|^p \int_0^\pi (\sin \varphi_1)^{p\sigma-1} \int_0^1 r^{p(\beta+\sigma)} (r^2 + \epsilon^2)^{\frac{p\lambda}{2}} dr + L_1 \\ &= \omega_{n-1} |\gamma|^p B\left(\frac{p\sigma}{2}, \frac{1}{2}\right) |\ln \epsilon| + O_\sigma(1) + L_1. \end{aligned}$$

Moreover,

$$(3.3) \quad \begin{aligned} |L_1| &\leq C \int_0^\pi (\sin \varphi_1)^{p\sigma+1} \int_0^1 r^{p(\beta+\sigma)+2} (r^2 + \epsilon^2)^{\frac{p\lambda}{2}-1} dr \leq C \left[B\left(\frac{p\sigma}{2} + 1, \frac{1}{2}\right) |\ln \epsilon| + O_\sigma(1) \right] \\ &= C \frac{p\sigma}{p\sigma+1} B\left(\frac{p\sigma}{2}, \frac{1}{2}\right) |\ln \epsilon| + O_\sigma(1). \end{aligned}$$

On $2\mathbb{B}^n \setminus \mathbb{B}^n$, directly calculation gives

$$|\nabla v|^2 = |x'|^{2\gamma-2} \left[\gamma^2 g^2(r) + |x'|^2 g'^2(r) + 2\gamma \frac{|x'|^2}{r} g g'(r) \right] \leq C |x'|^{2\gamma-2}.$$

Consequently

$$|x'|^{(\alpha+1)p} |x|^{\beta p} |\nabla v|^2 \leq C |x'|^{n-1+p\sigma} |x|^{\beta p},$$

and then

$$(3.4) \quad \int_{B_2 \setminus B_1} |x'|^{(\alpha+1)p} |x|^{\beta p} |\nabla v|^2 dx \leq CB \left(\frac{p\sigma}{2}, \frac{1}{2} \right).$$

Combining (3.2)-(3.4), for small enough $\sigma > 0$ and $\epsilon \in (0, 1)$, we can claim

$$\int_{\mathbb{R}^n} |x'|^{(\alpha+1)p} |x|^{\beta p} |\nabla v|^2 = \omega_{n-1} |\gamma|^p B\left(\frac{p\sigma}{2}, \frac{1}{2}\right) |\ln \epsilon| \times \left[1 + O(\sigma) \right] + B\left(\frac{p\sigma}{2}, \frac{1}{2}\right) \times O(1) + O_\sigma(1).$$

Tend first $\epsilon \rightarrow 0^+$, secondly set $\sigma \rightarrow 0^+$, we conclude that $C_{n,\alpha,\beta} \leq |\gamma_0|^p$ seeing (3.1). So $C_{n,\alpha,\beta} = |\gamma_0|^p$ for $p > 1$ and $\beta \geq 0$.

Case $\beta < 0$. Here we take still $f(x) = |x'|^\gamma$, but rewrite

$$\begin{aligned} W &= V |x'|^{-p} \left[-|\gamma|^{p-2} \gamma (n-1 + (\alpha+\beta)p + (p-1)\gamma) + |\gamma|^{p-2} \gamma \beta p \frac{x_n^2}{|x|^2} \right] \\ &=: V |x'|^{-p} \left[\tilde{H}_1(\gamma) + \tilde{H}_2(\gamma) \frac{x_n^2}{|x|^2} \right] \end{aligned}$$

where

$$\tilde{H}_1(\gamma) = -|\gamma|^{p-2} \gamma (n-1 + (\alpha+\beta)p + (p-1)\gamma), \quad \tilde{H}_2(\gamma) = |\gamma|^{p-2} \gamma \beta p.$$

Denote $\tilde{\gamma}_0 = -\frac{n-1+(\alpha+\beta)p}{p}$. If $p(\alpha+\beta) > 1-n$, there hold $\tilde{\gamma}_0 < 0$ and

$$\max_{\mathbb{R}} \tilde{H}_1(\gamma) = \tilde{H}_1(\tilde{\gamma}_0) = |\tilde{\gamma}_0|^p, \quad \tilde{H}_2(\tilde{\gamma}_0) \geq 0.$$

Seeing (1.15) and using approximation with functions in $C_c^1(\mathbb{R}^n \setminus \{x' = 0\})$, we claim $C_{n,\alpha,\beta} \geq |\tilde{\gamma}_0|^p$. \square

4. PROOF OF THEOREM 1.3

Thanks to (1.9), we need only to prove (1.10) for

$$(4.1) \quad \alpha p = \beta(p-1) + \mu, \quad \text{hence } \gamma_1 p = \gamma_3(p-1) + \gamma_2 - 1.$$

Let $V(x) = |x'|^{p\mu} |x|^{p\gamma_2}$ and $\vec{F}(x) = |x'|^{\beta-\mu} |x|^{\gamma_3-\gamma_2-1} x$, direct calculation yields that in $\mathbb{R}^n \setminus \{x' = 0\}$,

$$|\vec{F}| = |x'|^{\beta-\mu} |x|^{\gamma_3-\gamma_2}, \quad V |\vec{F}|^{p-2} = |x'|^{\beta(p-2)+2\mu} |x|^{\gamma_3(p-2)+2\gamma_2}$$

and

$$\operatorname{div}(\vec{F}) = (n-1 + \beta - \mu + \gamma_3 - \gamma_2) |x'|^{\beta-\mu} |x|^{\gamma_3-\gamma_2-1}.$$

Hence we have, with (4.1),

$$\begin{aligned} W &= \operatorname{div}(V |\vec{F}|^{p-2} \vec{F}) = V |\vec{F}|^{p-2} \operatorname{div}(\vec{F}) + \nabla(V |\vec{F}|^{p-2}) \cdot \vec{F} \\ &= [n + p(\alpha + \gamma_1)] |x'|^{\alpha p} |x|^{\gamma_1 p}. \end{aligned}$$

Applying (1.16), as $V |\vec{F}|^p = |x'|^{\beta p} |x|^{\gamma_3 p}$, we get immediately (1.10) for $u \in C_c^1(\mathbb{R}^n \setminus \{x' = 0\})$ with (4.1). Recall that under the condition (1.8), $|x'|^\alpha |x|^{\gamma_1}$, $|x'|^\beta |x|^{\gamma_3}$, $|x'|^\mu |x|^{\gamma_2} \in L_{loc}^p(\mathbb{R}^n)$. Similarly as for Theorem 1.1 and 1.2, we can extend the estimate for $u \in C_c^1(\mathbb{R}^n)$ by approximation.

Furthermore, if $\alpha = \beta = \mu$, then the above $\vec{F}(x) = |x|^{\gamma_3 - \gamma_2 - 1}x$. Then $u_0(x) = e^{-\kappa_0^{\frac{1}{p-1}}|x|^{\gamma_3 - \gamma_2 + 1}}$ (with suitable value $\kappa_0 > 0$) satisfies that the residual term in (1.16),

$$\mathcal{R}(\nabla u_0, u_0 \kappa_0^{\frac{1}{p-1}} \vec{F}) \equiv 0 \quad \text{in } \mathbb{R}^n \setminus \{0\}.$$

Hence with $\gamma_3 - \gamma_2 + 1 > 0$ and standard approximation, we can be convinced easily that the estimate (1.10) is sharp. \square

Remark 4.1. *The same proof provides also a generalization of [2, Corollary 1.2] to the anisotropic case. Let $p > 1$, $p\alpha > 1 - n$ and $\gamma_1 p = \gamma_3(p - 1) + \gamma_2 - 1$, there holds*

$$\| |x|^{\gamma_2} |x'|^\alpha \nabla u \|_{L^p(\mathbb{R}^n)}^{\frac{1}{p}} \| |x|^{\gamma_3} |x'|^\alpha u \|_{L^p(\mathbb{R}^n)}^{\frac{p-1}{p}} \geq \frac{|n + p(\alpha + \gamma_1)|}{p} \| |x|^{\gamma_1} |x'|^\alpha u \|_{L^p(\mathbb{R}^n)}, \quad \forall u \in C_c^1(\mathbb{R}^n \setminus \{0\}).$$

Moreover, for either $\gamma_3 - \gamma_2 + 1 > 0$, $(\gamma_2 + \alpha)p \geq p - n$; or $\gamma_3 - \gamma_2 + 1 < 0$, $(\gamma_2 + \alpha)p \leq p - n$, the above estimate is sharp.

Acknowledgements. The authors are partially supported by NSFC (No. 12271164) and Science and Technology Commission of Shanghai Municipality (No. 22DZ2229014).

Data availability. Data sharing is not applicable to this work as no new data were created or analyzed in the study.

REFERENCES

- [1] J. Bao; X. Chen, On the anisotropic Caffarelli-Kohn-Nirenberg type inequalities: existence, symmetry breaking region and symmetry of extremal functions. To appear in Commun. Contemp. Math. (2024).
- [2] A. Do; J. Flynn; N. Lam; G. Lu, L^p -Caffarelli-Kohn-Nirenberg inequalities and their stabilities, arXiv2310.07083 (2023).
- [3] N. Duy; N. Lam; G. Lu, p -Bessel pairs, Hardy's identities and inequalities and Hardy-Sobolev inequalities with monomial weights. J. Geom. Anal. 32(4) (2022), paper no. 109.
- [4] J. Flynn; N. Lam; G. Lu, L^p -Hardy identities and inequalities with respect to the distance and mean distance to the boundary. arXiv2310.18758, (2023).
- [5] N. Ghoussoub; A. Moradifam, Bessel pairs and optimal Hardy and Hardy-Rellich inequalities. Math. Ann. 349(1) (2011), 1-57.
- [6] X. Huang; D. Ye, First order Hardy inequalities revisited. Commun. Math. Res. 38(4) (2022), 535-559.
- [7] Y. Li; X. Yan, Asymptotic stability of homogeneous solutions of incompressible stationary Navier-Stokes equations. J. Diff. Equa. 297 (2021), 226-245.
- [8] Y. Li; X. Yan, Anisotropic Caffarelli-Kohn-Nirenberg type inequalities. Adv. Math. 419 (2023), paper no. 108958.

XIA HUANG

SCHOOL OF MATHEMATICAL SCIENCES AND SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, 200241, PEOPLE'S REPUBLIC OF CHINA

Email address: xhuang@cpde.ecnu.edu.cn

DONG YE

SCHOOL OF MATHEMATICAL SCIENCES AND SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, 200241, PEOPLE'S REPUBLIC OF CHINA

Email address: dye@math.ecnu.edu.cn