COMBINATORICS OF GENERALIZED ORTHOGONAL POLYNOMIALS OF TYPE R_{II}

JANG SOO KIM AND MINHO SONG

ABSTRACT. In 1995, Ismail and Masson introduced orthogonal polynomials of types R_I and R_{II} , which are defined by specific three-term recurrence relations with additional conditions. Recently, Kim and Stanton found a combinatorial interpretation for the moments of orthogonal polynomials of type R_I in the spirit of the combinatorial theory of orthogonal polynomials due to Flajolet and Viennot. In this paper, we push this combinatorial model further to orthogonal polynomials of type R_{II} . Moreover, we generalize orthogonal polynomials of type R_{II} by relaxing some of their conditions. We then prove a master theorem, which generalizes combinatorial models for moments of various types of orthogonal polynomials: classical orthogonal polynomials, Laurent biorthogonal polynomials, and orthogonal polynomials of types R_I and R_{II} .

1. INTRODUCTION

A (classical) orthogonal polynomial sequence $(P_n(x))_{n\geq 0}$ is a sequence of polynomials such that deg $P_n(x) = n$ and $\mathcal{L}(P_n(x)P_m(x)) = \kappa_n \delta_{n,m}$ with $\kappa_n \neq 0$, for some linear functional \mathcal{L} defined on the space of polynomials. It is well known that $(P_n(x))_{n\geq 0}$ is a (monic) orthogonal polynomial sequence if and only if it satisfies a three-term recurrence relation

(1.1)
$$P_{n+1}(x) = (x - b_n)P_n(x) - \lambda_n P_{n-1}(x), \quad n \ge 0,$$

where $P_{-1}(x) = 0, P_0(x) = 1$ and $\lambda_n \neq 0$ for $n \ge 1$.

In this paper, we focus on two quantities of orthogonal polynomials: moments and dual coefficients. The values $\mu_n := \mathcal{L}(x^n)$ are called the *moments* of the orthogonal polynomials. By the combinatorial theory of orthogonal polynomials due to Flajolet [4] and Viennot [12], the moment μ_n is a generating function for the Motzkin paths from (0,0) to (n,0). More generally, Viennot [12] showed that the *generalized moment*

(1.2)
$$\mu_{n,r,s} := \frac{\mathcal{L}(x^n P_r(x) P_s(x))}{\mathcal{L}(P_s(x)^2)}$$

is a generating function for the Motzkin paths from (0, r) to (n, s).

For a sequence $(P_n(x))_{n\geq 0}$ of any polynomials with deg $P_n(x) = n$, the *(generalized) dual* coefficients $\tau_{n,r,s}$ are defined by

$$x^n P_r(x) = \sum_{s \ge 0} \tau_{n,r,s} P_s(x).$$

If $(P_n(x))_{n\geq 0}$ is a sequence of orthogonal polynomials, then, by the orthogonality, the dual coefficient $\tau_{n,r,s}$ is equal to the generalized moment $\mu_{n,r,s}$.

In 1995, Ismail and Masson [5] introduced orthogonal polynomials of types R_I and R_{II} . Orthogonal polynomials $P_n^I(x)$ of type R_I are defined by the three-term recurrence

(1.3)
$$P_{n+1}^{I}(x) = (x - b_n)P_n^{I}(x) - (a_n x + \lambda_n)P_{n-1}^{I}(x), \quad n \ge 0,$$

where $P_{-1}^{I}(x) = 0$ and $P_{0}^{I}(x) = 1$, and $a_n \neq 0$ and $P_{n}^{I}(-\lambda_n/a_n) \neq 0$ for $n \ge 1$. Let

$$d_m^I(x) = \prod_{i=1}^m (a_i x + \lambda_i), \qquad Q_m^I(x) = \frac{P_m^I(x)}{d_m^I(x)}.$$

Ismail and Masson [5, Theorem 2.1] (see also [10, Theorem 2.1]) showed that there is a unique linear functional \mathcal{L}^{I} on the vector space $W^{I} := \operatorname{span}\{x^{n}Q_{m}^{I}(x): 0 \leq n < m\}$ such that $\mathcal{L}^{I}(1) = 1$ and $\mathcal{L}^{I}(x^{n}Q_{m}^{I}(x)) = 0$ for $0 \leq n < m$.

Kim and Stanton [10] presented a combinatorial theory of orthogonal polynomials of type R_I . In particular, they found combinatorial interpretations of the moments $\mu_n^I := \mathcal{L}^I(x^n)$ and the generalized moments $\mu_{n,r,s}^I := \mathcal{L}^I(x^n P_r^I(x) Q_s^I(x))$ using Motzkin–Schröder paths. This generalizes a result of Kamioka [8] that the moments of Laurent biorthogonal polynomials are generating functions for Schröder paths.

The main objective of this paper is to provide a combinatorial theory of orthogonal polynomials of type R_{II} that generalizes the previous results on classical orthogonal polynomials, Laurent biorthogonal polynomials, and orthogonal polynomials of type R_I . Orthogonal polynomials $P_n^{II}(x)$ of type R_{II} are defined by $P_{-1}^{II}(x) = 0$, $P_0^{II}(x) = 1$, and

(1.4)
$$P_{n+1}^{II}(x) = (x - b_n)P_n^{II}(x) - (c_n x^2 + a_n x + \lambda_n)P_{n-1}^{II}(x), \quad n \ge 0,$$

with the assumptions that $c_n \neq 0$, $P_n^{II}(\alpha_n) \neq 0$, and $P_n^{II}(\beta_n) \neq 0$ for $n \geq 1$, where α_n and β_n are the zeros of $c_n x^2 + a_n x + \lambda_n$. Let

$$d_m^{II}(x) = \prod_{i=1}^m (c_i x^2 + a_i x + \lambda_i), \qquad Q_m^{II}(x) = \frac{P_m^{II}(x)}{d_m^{II}(x)}.$$

Ismail and Masson [5, Theorem 3.5] showed that, for fixed N_0 and N_1 , there is a unique linear functional \mathcal{L}^{II} on the vector space

(1.5)
$$V^{II} := \operatorname{span}\{x^n Q_m^{II}(x) : 0 \le n < m\}$$

such that $\mathcal{L}^{II}(1) = N_0$, $\mathcal{L}^{II}(xQ_1^{II}(x)) = N_1$, and $\mathcal{L}^{II}(x^nQ_m^{II}(x)) = 0$ for $0 \le n < m$. See [6, 11] for recent work on specific orthogonal polynomials of type R_{II} .

There are two crucial differences between orthogonal polynomials of type R_{II} and classical orthogonal polynomials, as well as orthogonal polynomials of type R_I . The first difference is that, unlike $P_n(x)$ and $P_n^I(x)$, defined in (1.1) and (1.3), respectively, the orthogonal polynomials $P_n^{II}(x)$ of type R_{II} , defined in (1.4), are not monic. This induces a significantly different behavior in their dual coefficients, which will be explained shortly. The second difference is that the vector space W^{II} , given in (1.5), does not contain x^n for $n \ge 1$. This means that the moments $\mu_n^{II} = \mathcal{L}^{II}(x^n)$ are not defined, let alone the generalized moments $\mu_{n,r,s}^{II} = \mathcal{L}^{II}(x^n P_r^{II}(x) Q_s^{II}(x))$. Our strategy to overcome this limitation is to simply consider a linear functional defined on the larger space

(1.6)
$$W^{II} := \operatorname{span}\{x^n Q_m^{II}(x) : n, m \ge 0\}.$$

It may seem somewhat artificial to consider the space W^{II} only to include x^n for $n \ge 1$. However, this approach offers two benefits. First, it allows us to develop a combinatorial theory of orthogonal polynomials of type R_{II} that generalizes previous results on classical orthogonal polynomials, Laurent biorthogonal polynomials, and orthogonal polynomials of type R_I . Second, it enables us to discover a combinatorial interpretation for the dual coefficients $\tau_{n,r,s}^{II}$ defined below, which is similar to our combinatorial model for $\mu_{n,r,s}^{II}$.

Let $P_n(x)$ be the orthogonal polynomials defined in (1.1). Since $P_n(x)$ are monic, their dual coefficients are polynomials in variables $\mathbf{b} = (b_0, b_1, ...)$ and $\mathbf{\lambda} = (\lambda_1, \lambda_2, ...)$. Moreover, since $\tau_{n,r,s} = \mu_{n,r,s}$, the dual coefficients $\tau_{n,r,s}$ are, in fact, polynomials with nonnegative integer coefficients. See [3] and [2] for recent work on dual coefficients of orthogonal polynomials.

Now consider the polynomials $P_n^{II}(x)$ defined in (1.4) and let $\tau_{n,r,s}^{II}$ denote the their dual coefficients. Since $P_n^{II}(x)$ are not monic, we cannot expect their dual coefficients to be polynomials. Indeed, we have

$$x^{3} = \frac{P_{3}^{II}(x)}{1 - c_{1} - c_{2}} + \frac{A \cdot P_{2}^{II}(x)}{(1 - c_{1} - c_{2})(1 - c_{1})} + \frac{B \cdot P_{1}^{II}(x)}{(1 - c_{1} - c_{2})(1 - c_{1})} + \frac{C \cdot P_{0}^{II}(x)}{(1 - c_{1} - c_{2})(1 - c_{1})},$$

$$A = a_1 + a_2 + b_0 + b_1 + b_2 - b_2c_1 - b_0c_2,$$

$$B = a_2b_0c_1 + b_0b_1c_1 - a_1b_0c_2 - b_0^2c_2 - b_0b_1c_2 + a_1^2 + a_1a_2 + 2a_1b_0 + b_0^2 + 2a_1b_1 + a_2b_1 + b_0b_1 + b_1^2 - c_1\lambda_1 - c_1\lambda_2 + \lambda_1 + \lambda_2,$$

$$C = a_2b_0^2c_1 + b_0^2b_1c_1 - a_1b_0^2c_2 - b_0^3c_2 + a_1^2b_0 + a_1a_2b_0 + 2a_1b_0^2 + b_0^3 + a_1b_0b_1 - b_0c_1\lambda_1 - b_0c_2\lambda_1 + a_1\lambda_1 + a_2\lambda_1 + 2b_0\lambda_1 + b_1\lambda_1.$$

However, surprisingly, if we expand the dual coefficient $\tau_{n,r,s}^{II}$ as a formal power series, then the coefficients are nonnegative integers. For example, the dual coefficient $\tau_{3,0,1}^{II} = \frac{B}{(1-c_1-c_2)(1-c_1)}$ is equal to

$$\lambda_1 + \lambda_2 + a_1^2 + a_1a_2 + 2a_1b_0 + b_0^2 + 2a_1b_1 + a_2b_1 + b_0b_1 + b_1^2 + c_1\lambda_1 + c_2\lambda_1 + c_1\lambda_2 + c_2\lambda_2 + T,$$

where T is a formal power series in which every term has total degree greater than 2.

In this paper, we show that $\tau_{n,r,s}^{II}$ is a formal power series with nonnegative integer coefficients by giving its combinatorial interpretation. We note an interesting fact that, contrary to the case of classical orthogonal polynomials, for orthogonal polynomials of type R_{II} (and also for orthogonal polynomials of type R_I), the dual coefficient $\tau_{n,r,s}^{II}$ is not equal to the generalized moment $\mu_{n,r,s}^{II}$. We will find combinatorial models for $\tau_{n,r,s}^{II}$ and $\mu_{n,r,s}^{II}$, which are closely related. In fact, this is how the authors discovered the combinatorial model for $\tau_{n,r,s}^{II}$ in the first place; they found a combinatorial model for $\mu_{n,r,s}^{II}$ and an identity, see (5.2), which naturally suggests a combinatorial model for $\tau_{n,r,s}^{II}$.

The structure of the paper is as follows. In Section 2, we provide basic definitions and review some known results. In Section 3, we find some bases of the vector space $W^{II} = \text{span}\{x^n Q_m^{II}(x) :$ $n, m \ge 0\}$ and construct a linear functional on W^{II} . We then give a combinatorial interpretation for the generalized moments $\mu_{n,r,s}^{II}$. In Section 4, we generalize orthogonal polynomials of type R_{II} by relaxing some of their conditions. We prove a master theorem, which generalizes all combinatorial models for generalized moments of various types of orthogonal polynomials. In Section 5, we provide a combinatorial interpretation for the dual coefficient $\tau_{n,r,s}^{II}$. In Section 6, we propose two sufficient conditions for the convergence of $\mu_{n,r,s}$. In Section 7, we consider the moments of orthogonal polynomials of type R_{II} when $(b_n)_{n\ge 0}$, $(\lambda_n)_{n\ge 1}$, $(a_n)_{n\ge 1}$, and $(c_n)_{n\ge 1}$ are constant sequences. We show that, in this case, the moment of orthogonal polynomials of type R_{II} can be written as the moment of some classical orthogonal polynomials and also as the moment of some orthogonal polynomials of type R_I .

2. Preliminaries

In this section, we introduce basic definitions and known results. We start with the definition of orthogonal polynomials of type R_{II} , introduced by Ismail and Masson [5].

Definition 2.1. We say that $(P_n(x))_{n\geq 0}$ is a sequence of orthogonal polynomials of type R_{II} if $P_{-1}(x) = 0, P_0(x) = 1$, and

(2.1)
$$P_{n+1}(x) = (x - b_n)P_n(x) - (c_n x^2 + a_n x + \lambda_n)P_{n-1}(x), \quad n \ge 0,$$

for some sequences $\boldsymbol{b} = (b_0, b_1, \ldots), \boldsymbol{\lambda} = (\lambda_1, \lambda_2, \ldots), \boldsymbol{a} = (a_1, a_2, \ldots), \text{ and } \boldsymbol{c} = (c_1, c_2, \ldots)$ such that $c_n \neq 0, P_n(\alpha_n) \neq 0$, and $P_n(\beta_n) \neq 0$ for $n \geq 1$, where α_n and β_n are the roots of $c_n x^2 + a_n x + \lambda_n$. Given the polynomials $P_n(x)$, we will use the following notation:

$$d_n(x) = \prod_{i=1}^n (c_i x^2 + a_i x + \lambda_i), \qquad Q_n(x) = \frac{P_n(x)}{d_n(x)},$$
$$e_n(x) = \prod_{i=1}^n (x - \alpha_i), \qquad f_n(x) = \prod_{i=1}^n (x - \beta_i).$$



FIGURE 1. An R_{II} path $P = UUBUBUVUUVBDHU \in \mathsf{R}_{6,1}$ with wt $(P) = a_2 a_3 b_0 c_2^3 \lambda_1$. The thick segment from (1,1) to (2,2) represents UBUBU.

Here, we use the standard convention that the empty product is defined to be 1; for example, $d_0(x) = 1$. We also define the vector space

(2.2)
$$W = \operatorname{span}\{x^n Q_m(x) : n, m \ge 0\}.$$

Ismail and Masson [5] showed that orthogonal polynomials of type R_{II} have partial orthogonality as follows.

Theorem 2.2. [5, Theorem 3.5] For the polynomials $P_n(x)$ defined in Definition 2.1 and for any numbers N_0 and N_1 , there is a unique linear functional \mathcal{L} on W such that $\mathcal{L}(1) = N_0$, $\mathcal{L}(xQ_1(x)) = N_1$, and $\mathcal{L}(x^nQ_m(x)) = 0$ for $0 \le n < m$.

Now we review known combinatorial properties of these orthogonal polynomials, which will be generalized to orthogonal polynomials of type R_{II} in this paper. To put them all in one framework, we introduce R_{II} paths.

A lattice path is a sequence $p = (p_0, p_1, \ldots, p_n)$ of points $p_i \in \mathbb{Z} \times \mathbb{Z}_{\geq 0}$. For $i \geq 1$, the *i*th step s_i of p is defined by $s_i = p_i - p_{i-1}$. We will also consider the path p as the sequence $s_1 \ldots s_n$ of its steps (together with its starting point p_0).

Definition 2.3. An R_{II} path is a lattice path consisting of up steps U = (1,1), horizontal steps H = (1,0), down steps D = (1,-1), vertical-down steps V = (0,-1), and backward-down steps B = (-1,-1). Denote by $R_{n,r,s}$ the set of R_{II} paths from (0,r) to (n,s). We also write $R_{k,m} = R_{k,0,m}$ and $R_k = R_{k,0,0}$.

For later use, we also define restricted R_{II} paths.

Definition 2.4. A restricted R_{II} path is an R_{II} -path in $R_{n,r,s}$ that immediately ends when it touches the line x = n. We denote by $\tilde{R}_{n,r,s}$ the set of restricted R_{II} paths in $R_{n,r,s}$. The additional condition used to define restricted R_{II} paths from R_{II} paths will be referred to as the restricted condition.

For an R_{II} path $p = s_1 \dots s_k$, we define the *weight* wt(p) of p by wt(p) = $\prod_{t=1}^k \text{wt}(s_t)$, where

(2.3)
$$\operatorname{wt}(s_t) = \begin{cases} 1 & \text{if } s_t = U, \\ b_i & \text{if } s_t = H \text{ starting at height } i, \\ \lambda_i & \text{if } s_t = D \text{ starting at height } i, \\ a_i & \text{if } s_t = V \text{ starting at height } i, \\ c_i & \text{if } s_t = B \text{ starting at height } i. \end{cases}$$

See Figure 1. Note that steps U and B can be overlapped in an R_{II} path as shown in Figure 1. Thus, there can be infinitely many elements in $\mathsf{R}_{n,r,s}$ and $\sum_{p \in \mathsf{R}_{n,r,s}} \operatorname{wt}(p)$ is a formal power series in the variables in $\boldsymbol{b}, \boldsymbol{\lambda}, \boldsymbol{a}$, and \boldsymbol{c} .

A Motzkin path is an R_{II} path without vertical-down steps and backward-down steps. A Schröder path is an R_{II} path without down steps and backward-down steps. A Motzkin–Schröder path is an R_{II} path without backward-down steps. Let $Mot_{n,r,s}$ (resp. $Sch_{n,r,s}$ and $MS_{n,r,s}$) denote the set of Motzkin paths (resp. Schröder paths and Motzkin–Schröder paths) from (r, 0) to (n, s).

We are now ready to state known results on various orthogonal polynomials. Note that for the case of classical orthogonal polynomials, by the well-known fact $\mathcal{L}(P_n(x)^2) = \lambda_1 \cdots \lambda_n$, we have

$$Q_n(x) = \frac{P_n(x)}{d_n(x)} = \frac{P_n(x)}{\lambda_1 \cdots \lambda_n} = \frac{P_n(x)}{\mathcal{L}(P_n(x)^2)}.$$

Theorem 2.5. [12, Proposition 17, pp. I–15] Let $P_n(x)$ be the (classical) orthogonal polynomials given by (2.1) with $\mathbf{c} = \mathbf{a} = \mathbf{0}$. Then there is a linear functional \mathcal{L} on W such that

$$\mathcal{L}(x^n P_r(x) Q_s(x)) = \sum_{p \in \mathsf{Mot}_{n,r,s}} \mathrm{wt}(p).$$

Theorem 2.6. [8, Theorem 17] Let $P_n(x)$ be the Laurent biorthogonal polynomials given by (2.1) with $c = \lambda = 0$. Then there is a linear functional \mathcal{L} on W such that

$$\mathcal{L}(x^n P_r(x) Q_s(x)) = \sum_{p \in \mathsf{Sch}_{n,r,s}} \mathrm{wt}(p).$$

We note that Theorem 2.6 is an equivalent statement of [8, Theorem 17]; see [10, Theorem 4.7].

Theorem 2.7. [10, Theorem 3.12] Let $P_n(x)$ be the orthogonal polynomials of type R_I given by (2.1) with c = 0. Then there is a linear functional \mathcal{L} on W such that

$$\mathcal{L}(x^n P_r(x)Q_s(x)) = \sum_{p \in \mathsf{MS}_{n,r,s}} \mathrm{wt}(p).$$

Finally, we introduce dual coefficients of polynomials.

Definition 2.8. Let $(P_n(x))_{n\geq 0}$ be a sequence of polynomials such that deg $P_n(x) = n$ for all $n \geq 0$. The coefficients $\rho_{n,k}$ and dual coefficients $\tau_{n,k}$ of the polynomials $P_n(x)$ are defined by

(2.4)
$$P_n(x) = \sum_{k \ge 0} \rho_{n,k} x^k, \quad x^n = \sum_{k \ge 0} \tau_{n,k} P_k(x).$$

We also define the *(generalized)* dual coefficients $\tau_{n,r,s}$ by

(2.5)
$$x^{n}P_{r}(x) = \sum_{s \ge 0} \tau_{n,r,s}P_{s}(x).$$

Suppose that $P_n(x)$ be the (classical) orthogonal polynomials given by (2.1) with $\mathbf{c} = \mathbf{a} = \mathbf{0}$ and let \mathcal{L} be the corresponding linear functional. By the orthogonality $\mathcal{L}(P_r(x)Q_s(x)) = \delta_{r,s}$, multiplying $Q_s(x)$ both sides of (2.5) and taking \mathcal{L} yields

(2.6)
$$\mathcal{L}(x^n P_r(x) Q_s(x)) = \tau_{n,r,s}.$$

Thus, Theorem 2.5 gives the following corollary.

Corollary 2.9. Let $P_n(x)$ be the (classical) orthogonal polynomials given by (2.1) with c = a = 0. Then

$$\tau_{n,r,s} = \sum_{p \in \mathsf{Mot}_{n,r,s}} \mathrm{wt}(p).$$

We note that (2.6) no longer holds for Laurent biorthogonal polynomials or orthogonal polynomials of type R_I or R_{II} . In Section 5, we will find an analogous result of Corollary 2.9 for orthogonal polynomials of type R_{II} using restricted R_{II} paths.

3. A COMBINATORIAL INTERPRETATION FOR GENERALIZED MOMENTS

3.1. Vector space and linear functional. Throughout this section we follow the notation in Definition 2.1. Recall the vector space

$$W = \operatorname{span}\{x^n Q_m(x) : n, m \ge 0\}.$$

In this subsection, we provide a basis for W as follows.

Theorem 3.1. The vector space $W = \text{span}\{x^n Q_m(x) : n, m \ge 0\}$ has a basis

$$\mathcal{B} = \{x^n : n \ge 0\} \cup \{Q_n(x) : n \ge 1\} \cup \{xQ_n(x) : n \ge 1\}$$

To prove this theorem, we consider

$$W' = \operatorname{span}\{x^n/d_m(x) : n, m \ge 0\}$$

We will show that W = W'. Since W is clearly contained in W', it suffices to show that a basis of W' is contained in W.

Lemma 3.2. The vector space $W' = \operatorname{span}\{x^n/d_m(x) : n, m \ge 0\}$ has a basis

$$\mathcal{B}' = \{x^n : n \ge 0\} \cup \{1/d_m(x) : m \ge 1\} \cup \{x/d_m(x) : m \ge 1\}$$

Proof. Since $1/d_m(x) \sim x^{-2m}$ and $x/d_m(x) \sim x^{-2m+1}$ asymptotically up to constants, \mathcal{B}' is linearly independent. Since $\mathcal{B}' \subset W'$, it suffices to show that $\operatorname{span}(B') = W'$. To see this, we claim that $x^n/d_m(x) \in \operatorname{span}(B')$ for any $n, m \geq 0$. First, we write the fraction as

(3.1)
$$\frac{x^n}{d_m(x)} = q(x) + \frac{r(x)}{d_m(x)},$$

where q(x) and r(x) are polynomials and $\deg(r(x)) < 2m$. Now it suffices to show that $r(x)/d_m(x) \in \operatorname{span}(\mathcal{B}')$. If $\deg r(x) \leq 1$, then we have $r(x)/d_m(x) \in \operatorname{span}(\mathcal{B}')$. Otherwise, we can write

(3.2)
$$\frac{r(x)}{d_m(x)} = \frac{(c_m x^2 + a_m x + \lambda_m) r_1(x) + c_1(x)}{d_m(x)} = \frac{r_1(x)}{d_{m-1}(x)} + \frac{c_1(x)}{d_m(x)},$$

where $r_1(x)$ is a polynomial with deg $r_1(x) < 2m - 2$ and $c_1(x)$ is a polynomial with deg $c_1(x) \leq 1$. By iterating (3.2), we can express $r(x)/d_m(x)$ as a linear combination of the elements in \mathcal{B}' , which completes the proof.

Lemma 3.3. Let $W = \text{span}\{x^n Q_m(x) : n, m \ge 0\}$ and $W' = \text{span}\{x^n/d_m(x) : n, m \ge 0\}$. Then we have W = W'.

Proof. Since $W \subseteq W'$ and $x^n \in W$ for all $n \geq 0$, by Lemma 3.2, it suffices to show that $1/d_m(x), x/d_m(x) \in W$ for $m \geq 0$. We prove this by induction on m. First, the base case m = 0 is clear.

Now, let $m \ge 1$. We claim that $1/d_m(x) \in W$. If $r_m(x)$ is the quotient of $P_m(x)$ when divided by $x - \beta_m$, then we have

$$\frac{P_m(x)}{d_{m-1}(x)(x-\beta_m)} = \frac{r_m(x)}{d_{m-1}(x)} + \frac{P_m(\beta_m)}{d_{m-1}(x)(x-\beta_m)}.$$

Using (3.2) repeatedly, we can write $r_m(x)/d_{m-1}(x)$ as a linear combination of $1/d_i(x)$ and $x/d_i(x)$ for i < m, that is, $r_m(x)/d_{m-1}(x) \in W$ by the inductive hypothesis. Since $P_m(\beta_m) \neq 0$, we have

(3.3)
$$\frac{1}{d_{m-1}(x)(x-\beta_m)} = \frac{1}{P_m(\beta_m)} \left(c_m x Q_m(x) - c_m \alpha_m Q_m(x) - \frac{r_m(x)}{d_{m-1}(x)} \right) \in W.$$

We also have $1/(d_{m-1}(x)(x - \alpha_m)) \in W$ in the same way. If $\alpha_m \neq \beta_m$, then

$$\frac{1}{d_m(x)} = \frac{1}{c_m(\beta_m - \alpha_m)} \left(\frac{1}{d_{m-1}(x)(x - \beta_m)} - \frac{1}{d_{m-1}(x)(x - \alpha_m)} \right) \in W.$$

If $\alpha_m = \beta_m$, then we can write $Q_m(x)$ as

$$Q_m(x) = \frac{P_m(x)}{d_m(x)} = \frac{c_m(x - \alpha_m)^2 R_{m-2}(x) + \gamma c_m(x - \alpha_m) + P_m(\alpha_m)}{d_m(x)}$$
$$= \frac{R_{m-2}(x)}{d_{m-1}(x)} + \frac{\gamma}{d_{m-1}(x)(x - \alpha_m)} + \frac{P_m(\alpha_m)}{d_m(x)}$$

for some constant γ . Again using (3.2) repeatedly, we have $\frac{R_{m-2}(x)}{d_{m-1}(x)} \in W$ by the inductive hypothesis and hence

$$\frac{1}{d_m(x)} = \frac{1}{P_m(\alpha_m)} \left(Q_m(x) - \frac{R_{m-2}(x)}{d_{m-1}(x)} - \frac{\gamma}{d_{m-1}(x)(x-\alpha_m)} \right) \in W,$$

as claimed.

Now it remains to prove $x/d_m(x) \in W$. We write $Q_m(x)$ as

(3.4)
$$Q_m(x) = \frac{(c_m x^2 + a_m x + \lambda_m) U_m(x)}{d_m(x)} + \frac{c_1(x)}{d_m(x)} = \frac{U_m(x)}{d_{m-1}(x)} + \frac{c_1(x)}{d_m(x)},$$

where $c_1(x)$ is a polynomial of degree at most 1. Note that, if m = 1, we have $U_0(x) = 0$. Similarly to how we showed $r_m(x)/d_{m-1}(x) \in W$, we use the same induction argument to obtain $U_m(x)/d_{m-1}(x) \in W$. Together with $1/d_m(x) \in W$, we obtain $x/d_m(x) \in W$, as desired. \Box

Now we prove Theorem 3.1.

Proof of Theorem 3.1. Since $\mathcal{B} \subseteq W$, it suffices to show that \mathcal{B} contains $1/d_m(x)$ and $x/d_m(x)$ for any $m \ge 1$. It can be shown by induction. We suppose $1/d_k(x), x/d_k(x) \in \mathcal{B}$ for $k \le m-1$. By (3.2) and (3.3), we inductively obtain $1/d_m(x) \in \mathcal{B}$. Moreover, using (3.4), we obtain $x/d_m(x) \in \mathcal{B}$, as desired.

Remark 3.4. In [5, Theorem 3.5], Ismail and Masson implicitly use the fact (without providing a proof) that $\{Q_n(x) : n \ge 1\} \cup \{xQ_n(x) : n \ge 1\}$ is a basis for span $\{x^nQ_m(x) : 0 \le n \le m < \infty\}$. This fact can be established by following the same argument as in the proofs of the preceding two lemmas and Theorem 3.1.

3.2. Moments and generalized moments for orthogonal polynomials of type R_{II} . In this subsection, we construct a linear functional \mathcal{L} on $W = \operatorname{span}\{x^n Q_m(x) : n, m \ge 0\}$ that gives the partial orthogonality for orthogonal polynomials of type R_{II} . Moreover, we provide a combinatorial description for the generalized moments $\mathcal{L}(x^n P_r(x)Q_s(x))$.

By Theorem 3.1, we can define a linear functional on W as follows.

Definition 3.5. The linear functional \mathcal{L} on the vector space span $\{x^n Q_m(x) : n, m \ge 0\}$ is defined by

$$\mathcal{L}(x^n) = \sum_{P \in \mathsf{R}_n} \operatorname{wt}(P), \quad n \ge 0,$$
$$\mathcal{L}(Q_n(x)) = \mathcal{L}(xQ_{n+1}(x)) = 0, \quad n \ge 1$$
$$\mathcal{L}(xQ_1(x)) = \sum_{P \in \mathsf{R}_{1,1}} \operatorname{wt}(P).$$

Definition 3.6. The generalized moment $\mu_{n,r,s}$ of orthogonal polynomials of type R_{II} is given by

$$\mu_{n,r,s} = \mathcal{L}(x^n P_r(x) Q_s(x)).$$

We also write $\mu_{n,m} = \mu_{n,0,m}$ and $\mu_n = \mu_{n,0,0}$.

By multiplying $1/d_n(x)$ on both sides of (2.1), we have

(3.5)
$$(c_{n+1}x^2 + a_{n+1}x + \lambda_{n+1})Q_{n+1}(x) = (x - b_n)Q_n(x) - Q_{n-1}(x).$$

The next proposition provides a combinatorial interpretation for the values of \mathcal{L} on the set $\{x^n Q_m(x) : n, m \ge 0\}.$

Proposition 3.7. Suppose $c_n \neq 0$ for all $n \geq 0$. Then we have

$$\mu_{n,m} = \sum_{p \in \mathsf{R}_{n,m}} \operatorname{wt}(p).$$

Proof. Let $\mu'_{n,m}$ be the right-hand side of the equation. We will prove $\mu_{n,m} = \mu'_{n,m}$ using induction on (n,m). By Definition 3.5, the statement holds for m = 0, n = 0, or n = 1.

Now let $n \ge 2$ and $m \ge 1$, and suppose that the statement holds for all $(n', m') \ne (n, m)$ with $n' \le n$ and $m' \le m$. By (3.5),

$$(c_m x^2 + a_m x + \lambda_m)Q_m(x) = (x - b_{m-1})Q_{m-1}(x) - Q_{m-2}(x).$$

Multiply x^{n-2} on both sides and taking \mathcal{L} yields

$$(3.6) c_m \mu_{n,m} + a_m \mu_{n-1,m} + \lambda_m \mu_{n-2,m} = \mu_{n-1,m-1} - b_{m-1} \mu_{n-2,m-1} - \mu_{n-2,m-2}.$$

On the other hand, considering the last step of $p \in \mathsf{R}_{n-1,m-1}$, we have

$$\mu'_{n-1,m-1} = c_m \mu'_{n,m} + a_m \mu'_{n-1,m} + \lambda_m \mu'_{n-2,m} + b_{m-1} \mu'_{n-2,m-1} + \mu'_{n-2,m-2}$$

which is the same recurrence relation as (3.6). Thus, applying the inductive hypothesis to every term in (3.6) except $c_m \mu_{n,m}$ gives $c_m \mu_{n,m} = c_m \mu'_{n,m}$. Since $c_m \neq 0$, the statement also holds for (n,m), which completes the proof.

Remark 3.8. Using the recurrence (3.6) along with Definition 3.5, one can see that $\mu_{n,m} = 0$ for $0 \le n < m$. This shows the orthogonality of orthogonal polynomials of type R_{II} .

We are now ready to obtain a combinatorial description of $\mu_{n,r,s}$.

Theorem 3.9. For nonnegative integers n, r, and s, we have

$$\mu_{n,r,s} = \sum_{P \in \mathsf{R}_{n,r,s}} \operatorname{wt}(P).$$

Proof. Let $\mu'_{n,r,s} = \sum_{P \in \mathbb{R}_{n,r,s}} \operatorname{wt}(P)$. We will prove $\mu_{n,r,s} = \mu'_{n,r,s}$ by induction on r. The base step r = 0 follows from Proposition 3.7. For the inductive step, let $k \geq 0$ and suppose that $\mu_{n,r,s} = \mu'_{n,r,s}$ holds for all $r \leq k$. Replacing n by k in (2.1), multiplying $x^n Q_s(x)$ to both sides of the resulting equation, and taking \mathcal{L} , we obtain

(3.7)
$$\mu_{n,k+1,s} = \mu_{n+1,k,s} - b_k \mu_{n,k,s} - c_k \mu_{n+2,k-1,s} - a_k \mu_{n+1,k-1,s} - \lambda_k \mu_{n,k-1,s}.$$

On the other hand, considering the first step of $p \in \mathsf{R}_{n+1,k,s}$, we have

$$\mu'_{n+1,k,s} = \mu'_{n,k+1,s} + b_k \mu'_{n,k,s} + c_k \mu'_{n+2,k-1,s} + a_k \mu'_{n+1,k-1,s} + \lambda_k \mu'_{n,k-1,s},$$

which is the same recurrence as (3.7). Thus, by the inductive hypothesis, $\mu_{n,r,s} = \mu'_{n,r,s}$ also holds for r = k + 1. By induction, we obtain the theorem.

4. Generalized orthogonal polynomials of type R_{II}

In this section, we extend the definition of orthogonal polynomials of type R_{II} by relaxing some of their conditions.

Definition 4.1. We say that $(P_n(x))_{n\geq 0}$ is a sequence of generalized orthogonal polynomials of type R_{II} , or R_{II} polynomials for short, if $P_{-1}(x) = 0$, $P_0(x) = 1$, and

$$P_{n+1}(x) = (x - b_n)P_n(x) - (c_n x^2 + a_n x + \lambda_n)P_{n-1}(x), \quad n \ge 0,$$

where $\boldsymbol{a} = (a_1, a_2, \ldots), \boldsymbol{b} = (b_1, b_2, \ldots), \boldsymbol{c} = (c_1, c_2, \ldots), \text{ and } \boldsymbol{\lambda} = (\lambda_1, \lambda_2, \ldots)$ are sequences such that, for all $n \geq 1$, at least one of λ_n , a_n , or c_n is nonzero. We will use the following notation:

$$d_n(x) = \prod_{i=1}^n (c_i x^2 + a_i x + \lambda_i), \qquad Q_n(x) = \frac{P_n(x)}{d_n(x)}.$$

Note that R_{II} polynomials contain various types of orthogonal polynomials:

- Orthogonal polynomials of type R_{II} are R_{II} polynomials such that $c_n \neq 0, P_n(\alpha_n) \neq 0$, and $P_n(\beta_n) \neq 0$ for $n \ge 1$, where α_n and β_n are the roots of $c_n x^2 + a_n x + \lambda_n$.
- Orthogonal polynomials of type R_I are R_{II} polynomials such that $c_n = 0, a_n \neq 0$, and $P_n(-\lambda_n/a_n) \neq 0$ for $n \geq 1$.
- Laurent biorthogonal polynomials are R_{II} polynomials such that $c_n = \lambda_n = 0, a_n \neq 0$, and $P_n(0) \neq 0$ for $n \geq 1$.

• Classical orthogonal polynomials are R_{II} polynomials such that $c_n = a_n = 0$ and $\lambda_n \neq 0$ for $n \ge 1$.

Let

$$W = \operatorname{span}\{x^n Q_m(x) : n, m \ge 0\}.$$

Recall from (3.5) that, for $k \ge 0$,

(4.1)
$$(c_{k+1}x^2 + a_{k+1}x + \lambda_{k+1})Q_{k+1}(x) = (x - b_k)Q_k(x) - Q_{k-1}(x),$$

where $Q_{-1}(x) = 0$. If we multiply x^{ℓ} , we obtain

(4.2)
$$c_{k+1}x^{\ell+2}Q_{k+1}(x) + a_{k+1}x^{\ell+1}Q_{k+1}(x) + \lambda_{k+1}x^{\ell}Q_{k+1}(x) = x^{\ell+1}Q_k(x) - b_kx^{\ell}Q_k(x) - x^{\ell}Q_{k-1}(x).$$

Let $S \subseteq \mathbb{Z}_{\geq 0}^2$ and $B = \{x^n Q_m(x) : (n,m) \in S\}$. We say that $x^n Q_m(x)$ is obtained from Bby a single application of (4.1) if there exist $k, \ell \geq 0$ such that $x^n Q_m(x)$ appears in (4.2) with a nonzero coefficient and, for every term $x^{n'}Q_{m'}(x)$ in (4.2) with $(n',m') \neq (n,m)$, either its coefficient is zero or it is contained in B. We say that $x^n Q_m(x)$ is obtained from B by multiple applications of (4.1) if there is a sequence $(n_1,m_1), (n_2,m_2), \ldots, (n_t,m_t)$ with $(n_t,m_t) = (n,m)$ such that $x^{n_i}Q_{m_i}(x)$ is obtained from B_{i-1} by a single application of (4.1) for all $i = 1, 2, \ldots, t$, where $B_0 = B$ and $B_i = B_{i-1} \cup \{x^{n_i}Q_{m_i}(x)\}$.

Definition 4.2. A basis B of W is said to be good if $B = \{x^n Q_m(x) : (n,m) \in S\}$ for a subset $S \subseteq \mathbb{Z}^2_{>0}$ and, for every $n, m \ge 0, x^n Q_m(x)$ is obtained from B by multiple applications of (4.2).

Theorem 4.3. Suppose that there is a good basis of W. Then there is a linear functional \mathcal{L} on W such that, for all $n, r, s \ge 0$,

$$\mathcal{L}(x^n P_r(x) Q_s(x)) = \sum_{p \in \mathsf{R}_{n,r,s}} \mathrm{wt}(p).$$

Proof. Let $B = \{x^n Q_m(x) : (n,m) \in S\}$ be a good basis of W. Define \mathcal{L} on B by

(4.3)
$$\mathcal{L}(x^n Q_m(x)) = R_{n,m},$$

where $R_{n,m} = \sum_{p \in \mathsf{R}_{n,m}} \operatorname{wt}(p)$. Since *B* is a basis, we can extend the definition of \mathcal{L} to *W* by linearity. We claim that (4.3) holds for all $n, m \geq 0$.

To prove the claim, first suppose that $x^n Q_m(x)$ is obtained from B by a single application of (4.1). Applying \mathcal{L} to both sides of (4.2) gives

$$(4.4) c_{k+1}R'_{\ell+2,k+1} + a_{k+1}R'_{\ell+1,k+1} + \lambda_{k+1}R'_{\ell,k+1} = R'_{\ell+1,k} - b_kR'_{\ell,k} - R'_{\ell,k-1}$$

where $R'_{i,j} = \mathcal{L}(x^i Q_j(x))$. By the construction of \mathcal{L} , if $x^i Q_j(x) \in B$, then $R'_{i,j} = R_{i,j}$. Considering the last step of any path $p \in \mathsf{R}_{\ell+1,k}$, we have

$$(4.5) R_{\ell+1,k} = c_{k+1}R_{\ell+2,k+1} + a_{k+1}R_{\ell+1,k+1} + \lambda_{k+1}R_{\ell,k+1} + b_kR_{\ell,k} + R_{\ell,k-1}.$$

Since every term $R'_{i,j}$ with $(i,j) \neq (n,m)$ appearing in (4.4) satisfies $R'_{i,j} = R_{i,j}$, comparing (4.4) and (4.5) gives $R'_{n,m} = R_{n,m}$, hence $\mathcal{L}(x^n Q_m(x)) = R_{n,m}$.

Now consider $x^n Q_m(x)$ for arbitrary $n, m \ge 0$. Since B is a good basis, $x^n Q_m(x)$ is obtained from B by multiple applications of (4.2). Applying the previous argument iteratively, we obtain that $\mathcal{L}(x^n Q_m(x)) = R_{n,m}$. Therefore (4.3) holds for all $n, m \ge 0$, as claimed.

This shows the theorem for the case r = 0. The general case $r \ge 0$ can be proved by induction by the same arguments in the proof of Theorem 3.9.

Since $\mathsf{R}_{n,0,m} = \emptyset$ for n < m, we obtain the following orthogonality.

Corollary 4.4. Suppose that there is a good basis of W. Then there is a linear functional \mathcal{L} on W such that $\mathcal{L}(1)$ is a power series in $\mathbf{c} = (c_1, c_2, ...)$ with constant term 1 and, for all $0 \le n < m$,

$$\mathcal{L}(x^n Q_m(x)) = 0.$$

JANG SOO KIM AND MINHO SONG

polynomials	good bases
orthogonal polynomials	$\{(n,m):m=0\}$
Laurent biorthogonal polynomials	$\{(n,m): n = 0 \text{ or } m = 0\}$
orthogonal polynomials of type R_I	$\{(n,m): n = 0 \text{ or } m = 0\}$
orthogonal polynomials of type R_{II}	$\{(n,m): 0 \le n \le 1 \text{ or } m = 0\}$

TABLE 1. Orthogonal polynomials and their good bases.

Since there are good bases as in Table 1, Theorem 4.3 implies the combinatorial models of generalized moments for classical orthogonal polynomials, Laurent biorthogonal polynomials, and orthogonal polynomials of types R_I and R_{II} in Theorems 2.5, 2.6, and 2.7. Theorem 4.3 also includes orthogonal polynomials that do not belong to any of these four classes of orthogonal polynomials; for example, one can consider the sequence c with $c_{2n+1} = 0$ and $c_{2n} \neq 0$.

Note that in Theorem 4.3 we have the assumption that W has a good basis. It would be interesting to characterize the existence of a good basis.

Problem 4.5. Find a characterization of the sequences a, b, c, and λ such that W has a good basis.

5. A COMBINATORIAL INTERPRETATION FOR DUAL COEFFICIENTS

In this section, we provide a combinatorial interpretation of the (generalized) dual coefficients $\tau_{n,r,s}$ of R_{II} polynomials introduced in Section 4, which are orthogonal polynomials of type R_{II} with relaxed conditions.

We start with an observation on the combinatorial interpretation for generalized moments $\mu_{n,r,s}$ of R_{II} polynomials. Let $P_n(x)$ be R_{II} polynomials defined in the previous section. Recall that dual coefficients $\tau_{n,r,s}$ are defined by

(5.1)
$$x^n P_r(x) = \sum_{s>0} \tau_{n,r,s} P_s(x).$$

Multiply $Q_{\ell}(x)$ and take \mathcal{L} on both sides of (5.1) yields

(5.2)
$$\mu_{n,r,\ell} = \sum_{s \ge 0} \tau_{n,r,s} \mu_{0,s,\ell}$$

By Theorem 4.3, $\mu_{n,r,\ell}$ is the generating function for all R_{II} paths from (0, r) to (n, ℓ) . Hence, the identity (5.2) naturally suggests that $\tau_{n,r,s}$ is the generating function for the restricted R_{II} paths from (0, r) to (n, s) defined in Definition 2.4. Indeed, such a generating function satisfies (5.2). However, to conclude that $\tau_{n,r,s}$ is equal to this generating function, it must be shown that there is a unique solution for $\tau_{n,r,s}$ to the equation (5.2). Unfortunately, this is not the case. For example, if $\tau'_{0,0,1} = \mu_{0,1,0}^{-1} \mu_{0,0,0}$ and $\tau'_{0,0,s} = 0$ for $s \neq 1$, then we also have

$$\mu_{0,0,0} = \sum_{s \ge 0} \tau'_{0,0,s} \mu_{0,s,0}.$$

Interestingly, however, the suggested generating function turns out to be equal to $\tau_{n,r,s}$, which is the main result in this section.

Theorem 5.1. For nonnegative integers n, r, and s, we have

(5.3)
$$\tau_{n,r,s} = \sum_{p \in \widetilde{\mathsf{R}}_{n,r,s}} \operatorname{wt}(p).$$

Note that in order to prove Theorem 5.1, it suffices to show that the right-hand side of (5.3), denoted $\tau'_{n,r,s}$, satisfies the same relation (5.1) as $\tau_{n,r,s}$, that is,

(5.4)
$$x^{n}P_{r}(x) = \sum_{s\geq 0} \tau'_{n,r,s}P_{s}(x).$$



FIGURE 2. An example of $T \in DT_8$ with $wt(T) = a_2 b_3 c_5 \lambda_7$.

We will prove (5.4) by giving combinatorial interpretations for both sides and finding a sign-reversing involution. To do this we introduce the following definition.

Definition 5.2. Consider a $1 \times n$ square board whose boxes are labeled 1 through n from left to right. Let M_i (resp. D_i) be a monomino (resp. domino) with i dots inside. A *dotted tiling of size* n is a tiling of the board using tiles in $\{M_0, M_1, D_0, D_1, D_2\}$. The set of dotted tilings of size n is denoted by DT_n .

For a dotted tiling T, we define the weight wt(T) to be the product of the weights $wt(\tau)$ of the tiles τ in T, where

$$\operatorname{wt}(\tau) = \begin{cases} x & \text{if } \tau = M_1, \\ -b_{i-1} & \text{if } \tau = M_0 \text{ with label } i, \\ -\lambda_{i-1} & \text{if } \tau = D_0 \text{ with label } (i-1,i), \\ -a_{i-1}x & \text{if } \tau = D_1 \text{ with label } (i-1,i), \\ -c_{i-1}x^2 & \text{if } \tau = D_2 \text{ with label } (i-1,i). \end{cases}$$

See Figure 2 for an example.

By (2.1), it is immediate that

(5.5)
$$P_n(x) = \sum_{T \in DT_n} \operatorname{wt}(T)$$

We are now ready to prove Theorem 5.1.

Proof of Theorem 5.1. As discussed before, it suffices to show (5.4). By (5.5), we can rewrite (5.4) as

(5.6)
$$\sum_{k\geq 0} \tau'_{n,r,k} P_k(x) = \sum_{(p,T)\in X} \operatorname{wt}(p) \operatorname{wt}(T),$$

where X is the set of pairs (p, T) such that $p \in \tilde{\mathsf{R}}_{n,r,k}$ and $T \in \mathrm{DT}_k$ for some $k \ge 0$. To prove (5.4), we establish a sign-reversing involution on X such that the sum over its fixed points is equal to $x^n P_r(x)$.

Consider $(p,T) \in X$ with $p \in \widetilde{\mathsf{R}}_{n,r,k}$ and $T \in \mathrm{DT}_k$. For $m \ge 1$, we denote by p^m the *m*th-tolast step in p, and by T^m the *m*th-to-last tile in T. Let i (resp. j) be the maximum number of consecutive U's (resp. M_1 's) at the end of p (resp. T). Here, if $p^1 \ne U$, then i = 0, and if $T^1 \ne M_1$, then j = 0. We define $\phi(p,T) = (p',T')$ considering the two cases $i \le j$ and i > j.

Case 1: $i \leq j$. Note that p^{i+1} does not exist if and only if p consists only of up-steps, in which case i = n = k - r. We define p' and T' in the following subcases.

Case 1-1: $p^{i+1} = H$. Let p' be the path obtained from p by replacing p^{i+1} with U. Let T' be the dotted tiling obtained from T by inserting a new tile M_0 after T^{i+1} .

Case 1-2: $p^{i+1} = D$. Let p' be the path obtained from p by replacing p^{i+1} with U. Let T' be the dotted tiling obtained from T by inserting a new tile D_0 after T^{i+1} .

- **Case 1-3:** $p^{i+1} = V$. In this case, we have $i \ge 1$. Let p' be the path obtained from p by removing p^{i+1} . Let T' be the dotted tiling obtained from T by replacing T^i with D_1 .
- **Case 1-4:** $p^{i+1} = B$. In this case, we have $i \ge 2$. Let p' be the path obtained from p by removing the pair of steps (p^{i+1}, U) , where U is the step following p^{i+1} . Let T' be the dotted tiling obtained from T by replacing the pair of tiles (T^i, T^{i-1}) with a single tile D_2 .



FIGURE 3. An example of $(P,T) \in \widetilde{\mathsf{R}}_{5,2,5} \times \mathrm{DT}_5$ for Case 1-1 and its image $(\phi(P), \phi(T)) \in \widetilde{\mathsf{R}}_{5,2,6} \times \mathrm{DT}_6$ corresponding to Case 2-1, described in the proof of Theorem 5.1.



FIGURE 4. An example of $(P,T) \in \widetilde{\mathsf{R}}_{5,2,4} \times \mathrm{DT}_4$ for Case 1-2 and its image $(\phi(P), \phi(T)) \in \widetilde{\mathsf{R}}_{5,2,6} \times \mathrm{DT}_6$ corresponding to Case 2-2, described in the proof of Theorem 5.1.

Case 1-5: p^{i+1} does not exist. In this case, let (p', T') = (p, T).

Case 2: i > j. In this case, the starting height k - i of the step p^i is at least 1 because $k - i > k - j \ge 0$. We define p' and T' in the following subcases.

- **Case 2-1:** $T^{j+1} = M_0$. Let p' be the path obtained from p by replacing p^{i+1} with H. Let T' be the dotted tiling obtained from T by removing the tile T^{j+1} .
- **Case 2-2:** $T^{j+1} = D_0$. Let p' be the path obtained from p by replacing p^{i+1} with D. Let T' be the dotted tiling obtained from T by removing the tile T^{j+1} .
- **Case 2-3:** $T^{j+1} = D_1$. Let p' be the path obtained from p by inserting a step V before p^{i+1} . Let T' be the dotted tiling obtained from T by replacing T^{j+1} with M_1 .
- **Case 2-4:** $T^{j+1} = D_2$. Let p' be the path obtained from p by inserting a pair of steps (B, U) before p^{i+1} . Let T' be the dotted tiling obtained from T by replacing T^{j+1} with a pair of tiles (M_1, M_1) .

See Figures 3 to 6. It is straightforward to check that ϕ is a sign-reversing involution on X whose fixed points are the pairs (p, T), where p is the unique path in $\mathbb{R}_{n,r,n+r}$ consisting only of up-steps and T is a dotted tiling in DT_{n+r} such that the last n tiles are all equal to M_1 . Since $\mathrm{wt}(p) \mathrm{wt}(T) = x^n \mathrm{wt}(T_0)$, where T_0 is the dotted tiling obtained from T by removing the last n tiles, we have

(5.7)
$$\sum_{(p,T)\in X} \operatorname{wt}(p) \operatorname{wt}(T) = x^n \sum_{T_0\in \operatorname{DT}_r} \operatorname{wt}(T_0) = x^n P_r(x).$$

Then (5.4) follows from (5.6) and (5.7), which completes the proof.

If $c_n = 0$ in Theorem 5.1, we obtain the following corollary.



FIGURE 5. An example of $(P,T) \in \widetilde{\mathsf{R}}_{5,2,4} \times \mathrm{DT}_4$ for Case 1-3 and its image $(\phi(P), \phi(T)) \in \widetilde{\mathsf{R}}_{5,2,5} \times \mathrm{DT}_5$ corresponding to Case 2-3, described in the proof of Theorem 5.1.



FIGURE 6. An example of $(P,T) \in \widetilde{\mathsf{R}}_{5,2,4} \times \mathrm{DT}_4$ for Case 1-4 and its image $(\phi(P), \phi(T)) \in \widetilde{\mathsf{R}}_{5,2,4} \times \mathrm{DT}_4$ corresponding to Case 2-4, described in the proof of Theorem 5.1.

Corollary 5.3. Let $(P_n^I(x))_{n\geq 0}$ be the sequence of orthogonal polynomials of type R_I defined in (1.3) and let $\tau_{n,r,s}^I$ be their dual coefficients. Then we have

$$\tau^I_{n,r,s} = \sum_{p \in \widetilde{\mathsf{R}}^I_{n,r,s}} \operatorname{wt}(p),$$

where $\widetilde{\mathsf{R}}_{n,r,s}^{I}$ is the set of restricted R_{II} paths from (r,0) to (n,s) without backward-down steps.

Recall that a Motzkin path is an R_{II} path without vertical-down steps and backward-down steps. Thus, there is no difference between Motzkin paths and restricted Motzkin paths. Therefore, if $c_n = a_n = 0$ in Theorem 5.1, we obtain Corollary 2.9.

Remark 5.4. One may try to prove Theorem 5.1 by induction on n. Once the base steps n = 0 and n = 1 have been proved, we can easily apply the inductive argument as in the case of $\mu_{n,r,s}$. The case n = 0 is immediate from the definition. However, the other case n = 1 is not obvious. Note that, for the case of $\mu_{n,r,s}$, both cases n = 0 and n = 1 follow from the definition of \mathcal{L} .

6. Convergence of the moments and dual coefficients

In Section 3, we found a combinatorial description for generalized moments $\mu_{n,r,s}$ in terms of the weight sum of R_{II} paths. As there are infinitely many R_{II} paths from (0, r) to (n, s), we need to check whether the weight sum converges. In this section, we give sufficient conditions on the weights of steps for convergent weight sums of R_{II} paths and restricted R_{II} paths.

Let $C_n = \frac{1}{n+1} {2n \choose n}$ be the *n*th Catalan number.

Lemma 6.1. Suppose $b_m = b$, $\lambda_m = \lambda$, $a_m = a$, and $c_m = c$ for all m. Then

$$\mu_n = \sum_{i,j,k\geq 0} C_{i+j+k} \binom{n+j+2k}{n-2i-j} \binom{i+j+k}{j} \binom{i+k}{i} b^{n-2i-j} \lambda^i a^j c^k.$$

Proof. Suppose that $p \in \mathsf{R}_n$ has u up-steps, h horizontal steps, i down steps, j vertical-down steps, and k backward-down steps. Then

$$(n,0) = u(1,1) + h(1,0) + i(1,-1) + j(0,-1) + k(-1,-1)$$

which implies u = i + j + k and h = n - 2i - j. Therefore

(6.1)
$$\mu_n = \sum_{p \in \mathsf{R}_n} \operatorname{wt}(p) = \sum_{i,j,k \ge 0} f_n(i,j,k) b^{n-2i-j} \lambda^i a^j c^k,$$

where $f_n(i, j, k)$ is the number of paths $p \in \mathsf{R}_n$ with *i* down steps, *j* vertical-down steps, and *k* backward-down steps.

Given a path $p \in \mathsf{R}_n$, define $\phi(p)$ to be the labeled Motzkin path obtained from p by replacing every vertical-down step with a down step labeled a, every backward-down step with a down step labeled c. If p has the given number of steps of each type as above, then $\phi(p)$ is a labeled Motzkin path from (0,0) to (n+j+2k) with n-2i-j horizontal steps, i (unlabeled) down steps, j down steps labeled a, and k down steps labeled c. Such a labeled Motzkin path can be obtained from a Dyck path from (0,0) to (2i+2j+2k) by inserting n-2i-j horizontal steps in $\binom{n+j+2k}{n-2i-j}$ ways, labeling j down steps by a in $\binom{i+j+k}{j}$ ways, and labeling k down steps by c in $\binom{i+k}{k} = \binom{i+k}{i}$ ways. This implies that

(6.2)
$$f_n(i,j,k) = C_{i+j+k} \binom{n+j+2k}{n-2i-j} \binom{i+j+k}{j} \binom{i+k}{i}.$$

By (6.1) and (6.2), we obtain the desired formula.

Lemma 6.2. Suppose $b_m = b$, $\lambda_m = \lambda$, $a_m = a$, and $c_m = c$ for all m and |c| < 1/4. Then μ_n converges absolutely.

Proof. By Lemma 6.1, we have $\mu_n = \sum_{i,j,k\geq 0} f_n(i,j,k) b^{n-2i-j} \lambda^i a^j c^k$, where $f_n(i,j,k)$ is given in (6.2). Since $f_n(i,j,k) = 0$ unless $2i + j \leq n$, there are finitely many pairs (i,j) in this sum. Thus it suffices to show that, for fixed *i* and *j* with $2i + j \leq n$, the sum

(6.3)
$$\sum_{k\geq 0} f_n(i,j,k)c^k$$

converges.

By Stirling's formula, for fixed m and large N, we have the asymptotic behaviors

$$C_N \sim \frac{4^N}{N^{3/2}\sqrt{\pi}}, \qquad \binom{N}{m} \sim \frac{N^m}{m!}$$

Thus, for fixed i, j, n and large k,

$$f_n(i,j,k)c^k \sim \frac{4^{i+j+k}}{(i+j+k)^{3/2}\sqrt{\pi}} \cdot \frac{(n+j+2k)^{n-2i-j}}{(n-2i-j)!} \cdot \frac{(i+j+k)^j}{j!} \cdot \frac{(i+k)^i}{i!}c^k \sim D \cdot (4c)^k k^{n-i-3/2},$$

where D is a constant with respect to k. Since |4c| < 1, by the ratio test, (6.3) converges.

Theorem 6.3. Let $(a_m)_{m\geq 0}, (b_m)_{m\geq 0}, (c_m)_{m\geq 0}$ and $(\lambda_m)_{m\geq 0}$ be bounded sequences of complex numbers. Suppose $|c_m| < \frac{1}{4} - \epsilon$ for all m, where $\epsilon > 0$ is a fixed number. Then the generalized moments $\mu_{n,r,s}$ converge absolutely.

Proof. Suppose $|a_m| < a, |b_m| < b, |\lambda_m| < r$ for all m. Let $r = \max(\lambda, 1)$ and $c = 1/4 - \epsilon$. For $p \in \mathsf{R}_{n,r,s}$, let p' be the R_{II} path obtained from p by adding r up steps at the beginning and s down steps at the end. Then $p' \in \mathsf{R}_{n+r+s}$ and $|\operatorname{wt}(p)| = |\lambda^{-s} \operatorname{wt}(p')| \leq |\operatorname{wt}(p')|$. Thus

$$\sum_{p \in \mathsf{R}_{n,r,s}} |\operatorname{wt}(p)| \le \sum_{p \in \mathsf{R}_{n+r+s}} |\operatorname{wt}(p)| \le \sum_{p \in \mathsf{R}_{n+r+s}} |\widehat{\operatorname{wt}}(p)|,$$

where $\widehat{\mathrm{wt}}(p)$ is the weight $\mathrm{wt}(p)$ with substitution $(b_i, \lambda_i, a_i, c_i) \mapsto (b, r, a, c)$ for all *i*. By Lemma 6.2, the rightmost side in the above inequalities converges, which implies that $\mu_{n,r,s}$ converges absolutely.

$$\mu(x) = \frac{1}{\cdot} + \frac{bx(\mu(x))}{+} + \frac{x(\mu(x))}{x} + \frac{x(\mu(x))}{a}(\mu(x)) + \frac{x(\mu(x))}{\lambda x}(\mu(x))$$

FIGURE 7. A visualization of the functional equation for $\mu(x)$.

For the convergence of $\tau_{n,r,s}$, we only need the condition on c.

Corollary 6.4. Suppose that $|c_m| < \frac{1}{4} - \epsilon$ for all m, where $\epsilon > 0$ is a fixed number. Then the dual coefficients $\tau_{n,r,s}$ converge absolutely.

Proof. Recall that $\tau_{n,r,s} = \sum_{p \in \widetilde{\mathsf{R}}_{n,r,s}} \operatorname{wt}(p)$. Because of the restricted condition, each $p \in \widetilde{\mathsf{R}}_{n,r,s}$ cannot go over the line y = n + r. Thus

(6.4)
$$\sum_{p \in \widetilde{\mathsf{R}}_{n,r,s}} |\operatorname{wt}(p)| \le \sum_{p \in \mathsf{R}_{n,r,s}} |\operatorname{wt}'(p)|,$$

where wt'(p) is the weight of p using the sequences a', b', c', and λ' obtained from a, b, c, and λ by substituting $b_i = a_i = \lambda_i = c_i = 0$ for i > n + r. Then, by Theorem 6.3, we see that the right-hand side of (6.4) converges.

7. Moments with constant recurrence coefficients

In this section, we assume that $a_m = a$, $b_m = b$, $c_m = c$, and $\lambda_m = \lambda$, for all m. The goal is to find a formula for μ_n in this case.

Let

$$\mu(x) = \sum_{n \ge 0} \mu_n x^n.$$

Considering the first time each R_{II} path returns to the x-axis as shown in Figure 7, we obtain the following functional equation:

$$\mu(x) = 1 + bx\mu(x) + (c + ax + \lambda x^2)\mu(x)^2.$$

Solving the equation yields

$$\mu(x) = \frac{1 - bx - \sqrt{(1 - 4c) - (4a + 2b)x - (4\lambda - b^2)x^2}}{2(c + ax + \lambda x^2)}.$$

We aim to show that (up to a constant) μ_n can be expressed as the moment of certain classical or type R_I orthogonal polynomials. Recall that the moments of classical orthogonal polynomials (resp. orthogonal polynomials of type R_i) are equal to the weight sums of certain Motzkin paths (resp. Motzkin–Schröder paths). Thus, the goal is to reinterpret the values h_n in the framework of Motzkin or Motzkin–Schröder path models. While there are infinitely many R_{II} paths from (0,0) to (n,0), the number of such Motzkin or Motzkin–Schröder paths is finite. In this point of view, our results give useful ways of calculating μ_n . Note that this section generalizes the result on moments of R_I polynomials in [10, Section 6.2].

Let \mathcal{C} be the generating function for the Catalan numbers C_n :

$$\mathcal{C} = \sum_{n \ge 0} C_n c^n = \frac{1 - \sqrt{1 - 4c}}{2c}$$

Proposition 7.1. Suppose that $a_m = a$, $b_m = b$, $c_m = c$, and $\lambda_m = \lambda$, for all m. We have

$$\mu_n = \mathcal{C} \cdot \mu_n(\mathbf{B}, \mathbf{\Lambda})$$

Here, $\mu_n(\mathbf{B}, \mathbf{\Lambda})$ is the nth moment for classical orthogonal polynomials given by

$$p_{n+1}(x) = (x - B_n)p_n(x) - \Lambda_n p_{n-1}(x), n \ge 0,$$

where $p_{-1}(x) = 0, p_0(x) = 1$, and

$$B_{0} = \frac{1 - \sqrt{1 - 4c}}{2c\sqrt{1 - 4c}}a + \frac{b}{\sqrt{1 - 4c}}, \qquad \Lambda_{1} = \frac{(b^{2}c + a^{2} + ab - 4c\lambda + \lambda)(1 - \sqrt{1 - 4c})}{2c\sqrt{1 - 4c}},$$
$$B_{n} = \frac{2a + b}{1 - 4c}, \qquad \Lambda_{n+1} = \frac{b^{2}c + a^{2} + ab - 4c\lambda + \lambda}{(1 - 4c)^{2}}, \qquad n \ge 1.$$

Proof. By Flajolet's theory on continued fractions [4],

(7.1)
$$\sum_{n\geq 0} \mu_n(\mathbf{B}, \mathbf{\Lambda}) x^n = \frac{1}{1 - B_0 x - \Lambda_1 x^2 \cdot g(x)}$$

where

$$g(x) = \frac{1}{1 - B_1 x - \frac{\Lambda_2 x^2}{1 - B_2 x - \dots}}$$

Since $B_1 = B_2 = \cdots$ and $\Lambda_2 = \Lambda_3 = \cdots$, we see that

$$g(x) = \frac{1}{1 - B_1 x - \Lambda_2 x^2 \cdot g(x)}$$

Solving this function equation gives

(7.2)
$$g(x) = \frac{2}{1 - B_1 x - \sqrt{1 - 2B_1 x + (B_1^2 - 4\Lambda_2)x^2}}$$

By (7.1) and (7.2),

$$\mathcal{C}\sum_{n\geq 0}\mu_n(\mathbf{B}, \mathbf{\Lambda})x^n = \frac{\mathcal{C}}{1 - B_0 x - \frac{2\Lambda_1 x^2}{1 - B_1 x - \sqrt{1 - 2B_1 x + (B_1^2 - 4\Lambda_2)x^2}} = \frac{2}{1 - bx + \sqrt{(1 - 4c) - (4a + 2b)x - (4\lambda - b^2)x^2}} = \mu(x),$$

as desired.

As a corollary, we obtain a formula for the Hankel determinant $\det(\mu_{i+j})_{i,j=0}^n$.

Corollary 7.2. Suppose that $a_m = a$, $b_m = b$, $c_m = c$, and $\lambda_m = \lambda$, for all m. Then, we have

$$\det(\mu_{i+j})_{i,j=0}^{n} = \frac{\left(b^{2}c + a^{2} + ab - 4c\lambda + \lambda\right)^{\binom{n+1}{2}} \left(\frac{1 - \sqrt{1 - 4c}}{2c}\right)^{2n+1}}{\sqrt{1 - 4c}^{\binom{2n+1}{2}}}$$

Proof. By Proposition 7.1 and the fact that $\det(\mu_{i+j}(\mathbf{B}, \mathbf{\Lambda}))_{i,j=0}^n = \Lambda_1^n \Lambda_2^{n-1} \cdots \Lambda_n^1$, we have

$$\det(\mu_{i+j})_{i,j=0}^{n} = \mathcal{C}^{n+1} \det(\mu_{i+j}(\mathbf{B}, \mathbf{\Lambda}))_{i,j=0}^{n}$$
$$= \mathcal{C}^{n+1} \Lambda_{1}^{n} \Lambda_{2}^{n-1} \cdots \Lambda_{n}^{1}$$
$$= \frac{(b^{2}c + a^{2} + ab - 4c\lambda + \lambda)^{\binom{n+1}{2}} \left(\frac{1 - \sqrt{1 - 4c}}{2c}\right)^{2n+1}}{\sqrt{1 - 4c}^{\binom{2n+1}{2}}}.$$

Now we show that (up to a constant) μ_n can also be interpreted as the moment of orthogonal polynomials of type R_I .

Proposition 7.3. Suppose that $a_m = a$, $b_m = b$, $c_m = c$, and $\lambda_m = \lambda$, for all m. Then $\mu_n = \mathcal{C} \cdot \mu_n(\mathbf{B}, \mathbf{\Lambda}, \mathbf{A}).$

Here, $\mu_n(\mathbf{B}, \mathbf{\Lambda}, \mathbf{A})$ is the nth moment for orthogonal polynomials of type R_I given by

$$p_{n+1}(x) = (x - B_n)p_n(x) - (A_n x + \Lambda_n)p_{n-1}(x), \qquad n \ge 0,$$

where $p_{-1}(x) = 0, p_0(x) = 1$, and

$$B_{0} = \frac{b}{\sqrt{1 - 4c}}, \qquad B_{n} = \frac{b}{1 - 4c}, \qquad n \ge 1,$$

$$\Lambda_{1} = \frac{(b^{2}c - 4c\lambda + \lambda)(1 - \sqrt{1 - 4c})}{2c\sqrt{1 - 4c^{3}}}, \qquad \Lambda_{n+1} = \frac{b^{2}c - 4c\lambda + \lambda}{(1 - 4c)^{2}}, \qquad n \ge 1,$$

$$A_{1} = \frac{1 - \sqrt{1 - 4c}}{2c\sqrt{1 - 4c}}a, \qquad A_{n+1} = \frac{a}{1 - 4c}, \qquad n \ge 1.$$

Proof. The proof is similar to the proof of Proposition 7.1. By a direct computation, we obtain

$$\begin{split} \mathcal{C} \sum_{n \ge 0} \mu_n(\mathbf{B}, \mathbf{\Lambda}, \mathbf{A}) x^n &= \frac{\mathcal{C}}{1 - B_0 x - \frac{A_1 x + \Lambda_1 x^2}{1 - B_1 x - \frac{A_2 x + \Lambda_2 x^2}{1 - B_2 x - \dots}}}{\frac{\mathcal{C}}{1 - B_0 x - \frac{2A_1 x + 2\Lambda_1 x^2}{1 - B_1 x + \sqrt{1 - (4A_2 + 2B_1)x - (4\Lambda_2 - B_1^2)x^2}}} \\ &= \mu(x). \end{split}$$

The moment $\mu_n(\mathbf{B}, \mathbf{\Lambda})$ (resp. $\mu_n(\mathbf{B}, \mathbf{\Lambda}, \mathbf{A})$) of (classical) orthogonal polynomials (resp. orthogonal polynomials of type R_I) is the generating function for Motzkin paths (resp. Motzkin–Schröder paths) from (0,0) to (n,0), which is a finite sum. Hence, Propositions 7.1 and 7.3 provide practical methods for computing μ_n . At this point, we can pose natural questions:

- (1) Are the path models in Propositions 7.1 and 7.3 applicable for $\mu_{n,m}$ or $\mu_{n,r,s}$?
- (2) If not, are there Motzkin or Motzkin–Schröder paths with different weights that can be used for computing $\mu_{n,m}$ or $\mu_{n,r,s}$?

The answer to the first question is "no". For instance, one can verify that the coefficient $[c]\mu_{2,1}$ of c in $\mu_{2,1}$ is equal to 10a + 8b. In contrast, let $\hat{\mu}_{2,1}$ be the weight sum of Motzkin paths (resp. Motzkin–Schröder paths) from (0,0) to (2,1) with the weights given in Proposition 7.1 (resp. Proposition 7.3). Then $[c]\hat{\mu}_{2,1} = 11a + 6b$ (resp. $[c]\hat{\mu}_{2,1} = 7a + 6b$).

Regarding the second question, we observe that $\mu_{i,i} = \mu_{i,0,i} = C^{i+1}$. This implies that it is impossible to find classical or type R_I orthogonal polynomials whose generalized moments $\tilde{\mu}_{n,r,s}$ are the same as $\mu_{n,r,s}$, since $\tilde{\mu}_{i,0,i} = 1$. One might consider introducing a scaling factor F such that $\mu_{n,r,s} = F \cdot \tilde{\mu}_{n,r,s}$ as we have F = C when r = s = 0. However, we note that it is not possible to interpret $\mu_{n,r,s}$ as a weight sum of Motzkin–Schröder paths. In fact, one can verify that

(7.3)
$$\mu_{0,1,0} = \left(\frac{1-\sqrt{1-4c}}{2c}\right)^2 \cdot \frac{a+2bc}{\sqrt{1-4c}} \text{ and } \mu_{0,2,1} = \mu_{0,1,0} \cdot \frac{1-c-\sqrt{1-4c}}{c}.$$

To interpret these using Motzkin–Schröder paths, $\mu_{0,i,i-1}$ must be the weight A_i of a vertical down step (0, -1) starting at height *i*. This implies that $\mu_{0,2,0} = A_2A_1 = \mu_{0,2,1}\mu_{0,1,0}$, which does not contain any λ by (7.3). This contradicts the fact that $\mu_{0,2,0}$ must contain a term λc , since *DB* is an R_{II} path from (0, 2) to (0, 0) whose weight is λc .

Therefore, we need to consider combinatorial models different from Motzkin–Schröder paths. In the literature, there are other lattice path models for orthogonal polynomials. For example, Jang and Song [7] provide lattice paths called gentle Motzkin paths for their study of orthogonal polynomials on the unit circle.

We end this section with the following open problem.

Problem 7.4. Find a lattice path model that can be used to calculate $\mu_{n,m}$ or $\mu_{n,r,s}$ in a finite process.

References

- M. Bousquet-Mélou and M. Mishna. Walks with small steps in the quarter plane. In Algorithmic probability and combinatorics, volume 520 of Contemp. Math., pages 1–39. Amer. Math. Soc., Providence, RI, 2010.
- [2] S. Corteel, B. Jonnadula, J. P. Keating, and J. S. Kim. Lecture hall graphs and the Askey scheme. Preprint, arXiv:2311.12761, 2023.
- [3] S. Corteel and J. S. Kim. Lecture hall tableaux. Adv. Math., 371:107266, 35, 2020.
- [4] P. Flajolet. Combinatorial aspects of continued fractions. Discrete Math., 32(2):125–161, 1980.
- [5] M. E. H. Ismail and D. R. Masson. Generalized orthogonality and continued fractions. J. Approx. Theory, 83(1):1–40, 1995.
- [6] M. E. H. Ismail and A. Sri Ranga. R_{II} type recurrence, generalized eigenvalue problem and orthogonal polynomials on the unit circle. *Linear Algebra Appl.*, 562:63–90, 2019.
- [7] J. Jang and M. Song. Combinatorics of orthogonal polynomials on the unit circle. *Preprint*, arXiv:2407.07508, 2024.
- [8] S. Kamioka. A combinatorial representation with Schröder paths of biorthogonality of Laurent biorthogonal polynomials. *Electron. J. Combin.*, 14(1):Research Paper 37, 22, 2007.
- [9] S. Kamioka. A combinatorial derivation with Schröder paths of a determinant representation of Laurent biorthogonal polynomials. *Electron. J. Combin.*, 15(1):Research Paper 76, 20, 2008.
- [10] J. S. Kim and D. Stanton. Combinatorics of orthogonal polynomials of type R_I. Ramanujan J., 61(2):329–390, 2023.
- [11] V. Shukla and A. Swaminathan. Chain sequences and zeros of polynomials related to a perturbed R_{II} type recurrence relation. J. Comput. Appl. Math., 422:Paper No. 114916, 24, 2023.
- [12] G. Viennot. Une théorie combinatoire des polynômes orthogonaux généraux. Lecture Notes, UQAM, 1983.
- [13] A. Zhedanov. Biorthogonal rational functions and the generalized eigenvalue problem. J. Approx. Theory, 101(2):303–329, 1999.

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON, SOUTH KOREA *Email address*: jangsookim@skku.edu

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON, SOUTH KOREA *Email address*: smh3227@skku.edu