

## First-principle theory of the Casimir screening effect

Yury A. Budkov<sup>1,2,3, a)</sup> and Petr E. Brandyshev<sup>1,2</sup>

<sup>1)</sup>*Laboratory of Computational Physics, HSE University, Tallinskaya st. 34, 123458 Moscow, Russia*

<sup>2)</sup>*School of Applied Mathematics, HSE University, Tallinskaya st. 34, 123458 Moscow, Russia*

<sup>3)</sup>*Laboratory of Multiscale Modeling of Molecular Systems, G.A. Krestov Institute of Solution Chemistry of the Russian Academy of Sciences, 153045, Akademicheskaya st. 1, Ivanovo, Russia*

In this letter, we use the formalism of finite-temperature quantum field theory to investigate the Casimir force between flat, ideally conductive surfaces containing confined, but mobile ions. We demonstrate that in the Gaussian approximation, the contribution of ionic fluctuations is separate from the contribution of electromagnetic fluctuations that are responsible for the standard Casimir effect. This is in line with the "separation hypothesis", which was previously used on a purely intuitive basis. Our analysis demonstrates the significance of calculating the zero Matsubara frequency component in the electromagnetic contribution, using the formula developed by Schwinger et al., as opposed to other researchers based on the Lifshitz theory.

---

<sup>a)</sup>ybudkov@hse.ru

The Casimir effect, named after Hendrik Casimir, who first predicted it in 1948<sup>1</sup>, is a phenomenon that occurs when two neutral metal surfaces are brought very close together in vacuum at zero temperature. The extension of Casimir's result to arbitrary temperatures and general dielectric media was accomplished by Lifshitz et al.<sup>2,3</sup>, with subsequent refinements by Schwinger et al.<sup>4</sup> and others. Casimir effect is caused by the quantum fluctuations of electromagnetic fields in vacuum that generate a difference in the ground state energy density between the inside and outside of the metal surfaces, creating a net attraction between them. Although the Casimir force is very weak, it has been detected and experimentally quantified, and shown to play a crucial role in nanotechnology, particularly in the development of microelectromechanical systems (MEMS) and other nano devices<sup>5-7</sup>. Overall, the Casimir effect is an prime example of the complex and unintuitive nature of quantum physics<sup>8,9</sup>. It demonstrates how empty space is actually permeated by electromagnetic field fluctuations that generate long range interactions between bodies having a significant impact on many everyday phenomena.

Despite the significant progress made in the theoretical understanding of the Casimir effect<sup>8-11</sup>, there are still some unresolved issues that need to be addressed.

One of the unresolved problems in the theory of the Casimir effect is the Casimir force in the presence of mobile charged particles (ions) between flat perfectly conductive walls. In the early works<sup>12-15</sup> it was assumed that electromagnetic field fluctuations responsible for the attractive Casimir force can be treated separately from the pure image effects of the metallic slabs on the electrolyte ions. This assumption was referred to as the "separation hypothesis"<sup>16</sup>. Following the separation hypothesis, the two long-range forces cancel each other out exactly<sup>13-15</sup>. This canceling could be referred to as the "Casimir screening effect"<sup>17</sup>. Despite the plausibility of this hypothesis, it has not been confirmed by the fundamental principles of quantum electrodynamics.

We would like to emphasize that the screening of the Casimir force caused by the presence of ions in the space between metal walls is different from the screening related to the presence of mass within the quanta of the vector field within the framework of the Proca model<sup>18-20</sup>. Unlike 'ionic screening', 'mass screening', which is essentially a quantum effect, occurs for all terms in the Casimir force expansion in the Matsubara modes<sup>18</sup>.

The second issue is related to the behavior of the Casimir force at high temperatures. When the dielectric function of the material of the walls approaches infinity ( $\epsilon(\omega) \rightarrow \infty$ ),

the zero-Matsubara frequency term in the Lifshitz formula is not uniquely defined, as its value depends on the order in which the limits of the dielectric function approaching infinity and the Matsubara frequency approaching zero ( $\omega_n \rightarrow 0$ ) are taken. To obtain the ideal-conductor result based on electrostatic boundary conditions, Schwinger et al. postulated the following order<sup>4</sup>:  $\epsilon(\omega) \rightarrow \infty$  and then  $\omega_n \rightarrow 0$ . Thus they obtained twice as high value of the high temperature limit of the Casimir force than what was obtained by Lifshitz et al.<sup>3</sup>. Jancovici and Šamaj<sup>16,17</sup> and independently Buenzli and Martin<sup>21</sup> questioned the result of Schwinger et al., modeling the Casimir interaction of metal walls as the interaction of half-spaces containing equilibrium plasma comprised of classical ions. Using methods based on the formalism of correlation functions<sup>21</sup> and the theory of linear response<sup>16</sup>, the Casimir force was obtained, two times smaller than by Schwinger et al., in accordance with the Lifshitz limiting regime. Later, Buenzli and Martin<sup>22</sup> reproduced their result assuming that metals consist of mobile quantum charges in thermal equilibrium with the photon field at a fixed temperature. Recently, Brandyshev and Budkov<sup>23</sup> following the finite-temperature quantum field theory<sup>24</sup>, directly reproduced Schwinger's result for the Casimir force. It is important to note that their first-principle quantum electrodynamics-based approach does not require the assumption of the order of the aforementioned limits, which may indicate the correctness of the result obtained by Schwinger and coworkers. Below we provide more detailed analysis.

The aim of this study is to conduct a comprehensive analysis of these issues, considering their interconnections within the finite-temperature quantum field theory approach.

We assume that there are  $N_+$  positively charged ions (cations) and  $N_-$  negatively charged ions (anions) in a slit-like pore. The ions have charges of  $q_+$  and  $q_-$ , respectively, and the walls of the pore are ideally conductive and electrically neutral. The distance between the walls is  $L$ . We assume that the system, which consists of ions, conductive walls, and a quantum electromagnetic field, is in thermodynamic equilibrium with metal walls at a specific temperature  $T$ . The total ionic charge is zero, i.e.  $q_+N_+ + q_-N_- = 0$ , and the system is electroneutral.

Thus, we start from the partition function of the described system with a fixed gauge

(see Appendix A)

$$Z = \oint \mathcal{D}A |D| \oint d\sigma[\{\mathbf{r}_i^{(+)}\}] \oint d\sigma[\{\mathbf{r}_j^{(-)}\}] \exp \left[ S_{EM}[A] - i \int_0^\beta d\tau \int d^3\mathbf{x} (\hat{\rho}\varphi - \hat{\mathbf{j}} \cdot \mathbf{A}) \right], \quad (1)$$

where  $A_\mu = (-\varphi, \mathbf{A})$  is the 4-potential of the electromagnetic field with effective action

$$S_{EM}[A] = \int_0^\beta d\tau \int d^3\mathbf{x} \mathcal{L}_{\text{eff}}, \quad (2)$$

$$\mathcal{L}_{\text{eff}} = \frac{1}{2} A_\mu \square A_\mu + \frac{1-\alpha}{2} \left( \partial_\mu A_\mu \right)^2, \quad (3)$$

$|D| = \text{Det}(-\square)$  is the functional determinant and  $\alpha$  is the gauge parameter determining the family of similar gauges<sup>25</sup>; we have also introduced the integration measures over the paths,  $\mathbf{r}_i^{(a)}(\tau)$ , of ions in the imaginary 'time'  $\tau$ , i.e.

$$\oint d\sigma[\{\mathbf{r}_i^{(a)}\}](\dots) = \oint \prod_{i=1}^{N_+} \frac{\mathcal{D}\mathbf{r}_i^{(a)}}{C_a} \exp \left[ - \int_0^\beta d\tau \frac{m_a (\dot{\mathbf{r}}_i^{(a)}(\tau))^2}{2} \right] (\dots), \quad a = \pm, \quad (4)$$

where  $m_a$  is the mass of ion  $a$ , and  $C_a = \oint \mathcal{D}\mathbf{r}_i^{(a)} e^{-\int_0^\beta d\tau \frac{m_a (\dot{\mathbf{r}}_i^{(a)}(\tau))^2}{2}}$  are the normalization constants; the symbol  $\oint$  means the functional integration takes into account periodic boundary conditions for the integration variables,

$$A_\mu(0, \mathbf{x}) = A_\mu(\beta, \mathbf{x}), \quad \mu = 0, 1, 2, 3, \quad (5)$$

$$\mathbf{r}_i^{(a)}(0) = \mathbf{r}_i^{(a)}(\beta), \quad \beta = \frac{1}{T}. \quad (6)$$

We would like to point out that the partition function, which is the trace of the density matrix, can be formally obtained from the quantum amplitude of a system written in terms of a functional integral, by performing a Wick rotation from time to inverse temperature,  $\tau \rightarrow -i\tau$ <sup>26,27</sup>. Note also that we use the Euclidean metric for D'Alembert operator, i.e.  $\square = \partial_\tau^2 + \Delta$  and a unit system, where  $k_B = \hbar = c = 1$  ( $k_B$  is the Boltzmann constant,  $\hbar$  is the Planck constant, and  $c$  is the speed of light).

The microscopic charge density and current density of ions can be determined as follows

$$\hat{\rho}(\tau, \mathbf{x}) = \sum_{a=\pm} q_a \sum_{i=1}^{N_a} \delta \left( \mathbf{x} - \mathbf{r}_i^{(a)}(\tau) \right), \quad (7)$$

$$\hat{\mathbf{j}}(\tau, \mathbf{x}) = \sum_{a=\pm} q_a \sum_{j=1}^{N_a} \dot{\mathbf{r}}_j^{(a)}(\tau) \delta(\mathbf{x} - \mathbf{r}_j^{(a)}(\tau)). \quad (8)$$

It is easy to check that the electric charge and current density obey the conservation law,  $\partial_\tau \hat{\rho} + \nabla \cdot \hat{\mathbf{j}} = 0$ . Taking into account (7) and (8), we can rewrite the partition function (1) in the form

$$Z = \oint \mathcal{D}A |D| \exp[S_{EM}[A]] \prod_{a=\pm} \left[ \oint \frac{\mathcal{D}\mathbf{r}_a}{C_a} e^{-\int_0^\beta d\tau \left( \frac{m_a \dot{\mathbf{r}}_a^2(\tau)}{2} + i q_a \varphi(\tau, \mathbf{r}_a(\tau)) - i q_a \dot{\mathbf{r}}_a(\tau) \cdot \mathbf{A}(\tau, \mathbf{r}_a(\tau)) \right)} \right]^{N_a}. \quad (9)$$

Now, let us rewrite the 'paths' of ions as  $\mathbf{r}_a(\tau) = \mathbf{x}_a + \boldsymbol{\xi}_a(\tau)$ , where  $\mathbf{x}_a$  is the constant radius-vector and variables  $\boldsymbol{\xi}_a(\tau)$  describe the quantum 'smoothing' of particles and satisfy zero boundary conditions  $\boldsymbol{\xi}_a(0) = \boldsymbol{\xi}_a(\beta) = 0$ . Considering the ions as classical particles, that is, neglecting the terms containing the variables  $\boldsymbol{\xi}_a(\tau)$  and  $\dot{\boldsymbol{\xi}}_a(\tau)$ , we can obtain

$$Z = \oint \mathcal{D}A |D| \exp[S_{EM}[A]] \prod_{a=\pm} \left[ \int \frac{d\mathbf{x}}{V} e^{-i \int_0^\beta d\tau q_a \varphi(\tau, \mathbf{x})} \right]^{N_a}, \quad (10)$$

where  $V$  is the volume of system and we took into account that  $C_a = V \int \mathcal{D}\boldsymbol{\xi} e^{-\int_0^\beta d\tau \frac{m_a \dot{\boldsymbol{\xi}}^2(\tau)}{2}}$  with  $\boldsymbol{\xi}(0) = \boldsymbol{\xi}(\beta) = 0$ . The ions can be treated as classical particles if their thermal wavelengths,  $\lambda_a$ , are much smaller than the average inter-ionic distances, which is always the case.

In the thermodynamic limit  $N_a \rightarrow \infty$ ,  $V \rightarrow \infty$ ,  $N_a/V \rightarrow n_a$ , where  $n_a$  are the average ionic concentrations, we can obtain<sup>28,29</sup>

$$Z = \oint \mathcal{D}\varphi \mathcal{D}\mathbf{A} |D| \exp[S_{EM}[\varphi, \mathbf{A}] + S_{ion}[\varphi]], \quad (11)$$

where we have used that  $\mathcal{D}A = \mathcal{D}\varphi \mathcal{D}\mathbf{A}$  and introduced the contribution of the ions to the effective action

$$S_{ion}[\varphi] = \sum_{a=\pm} n_a \int d\mathbf{x} \left( e^{-i \int_0^\beta d\tau q_a \varphi(\tau, \mathbf{x})} - 1 \right). \quad (12)$$

As can be seen from eqs. (11) and (12), for pure classical ions, the vector potential drops out from the ion part of effective action. In other words, classical charged particles in equilibrium can only affect the fluctuations in the electric potential  $\varphi$ , which is of course in accordance with the Bohr-Van Leeuwen theorem<sup>30</sup>.

To proceed with the calculation of the partition function (11), we need to specify a gauge. The simplest one is the Feynman gauge, for which  $\alpha = 1$ . In this case, the kinetic terms for all the fields  $A_\mu$  are in canonical form, meaning they are orthogonalized and uncoupled. Let us calculate the partition function (11) in the Gaussian approximation. Expanding the ionic part of the effective action ( $S_{ion}$ ) in a power series of  $\varphi$  and neglecting terms higher than second order, we get

$$Z = \oint \mathcal{D}\varphi \mathcal{D}\mathbf{A} |D| e^{\frac{1}{2} \int_0^\beta d\tau \int d^3\mathbf{x} A_\mu \square A_\mu - \frac{1}{2} \sum_{a=\pm} q_a^2 n_a \int_0^\beta d\tau \int_0^\beta d\tau' \int d\mathbf{x} \varphi(\tau, \mathbf{x}) \varphi(\tau', \mathbf{x})}. \quad (13)$$

By calculating the Gaussian functional integral, taking into account the boundary conditions for an ideal metal (see Appendix B and ref.<sup>23</sup>), we can obtain the surface free energy density

$$f = -\frac{1}{\beta \mathcal{A}} \ln Z, \quad (14)$$

where  $\mathcal{A}$  is the total surface area of the bounding walls. Thus, we have

$$f = f_1 + f_2, \quad (15)$$

where

$$f_1 = \frac{1}{2\beta} \int \frac{d^2\mathbf{q}}{(2\pi)^2} \left( \sum_{l=1}^{\infty} \ln \lambda'(\mathbf{q}, 0, l) + \sum_{l=0}^{\infty} \ln \lambda(\mathbf{q}, 0, l) \right), \quad (16)$$

$$f_2 = \frac{2}{\beta} \int \frac{d^2\mathbf{q}}{(2\pi)^2} \sum_{l=0}^{\infty} \sum_{n=1}^{\infty} \ln \lambda(\mathbf{q}, n, l), \quad (17)$$

$$\lambda(\mathbf{q}, n, l) = \mathbf{q}^2 + q_l^2 + \omega_n^2, \quad \lambda'(\mathbf{q}, n, l) = \mathbf{q}^2 + q_l^2 + \omega_n^2 + \varkappa^2, \quad (18)$$

where  $\varkappa = r_D^{-1} = (\beta \sum_a q_a^2 n_a)^{1/2}$  is the inverse Debye length,  $q_l = \pi l/L$  is the lateral photon momentum, and  $\omega_n = 2\pi n/\beta$  is the  $n$ th Matsubara frequency. A dashed sum means that the term with the number  $l = 0$  is included with a multiplier  $1/2$ .

Using the same method as in<sup>23</sup>, we can sum over  $l$  and differentiate with respect to  $L$  to obtain the disjoining pressure – the total force per unit area acting on the bounding surfaces<sup>31</sup>,

$$\begin{aligned} \Pi &= - \left( \frac{\partial f}{\partial L} - \frac{\partial f}{\partial L} \Big|_{L \rightarrow \infty} \right) \\ &= -\frac{T}{4\pi} \int_0^\infty dq q \sqrt{q^2 + \varkappa^2} \left( \coth \left( \sqrt{q^2 + \varkappa^2} L \right) - 1 \right) - \frac{T}{4\pi} \int_0^\infty dq q^2 \left( \coth(qL) - 1 \right) \\ &\quad - \frac{T}{\pi} \sum_{n=1}^{\infty} \int_0^\infty dq q \sqrt{q^2 + \omega_n^2} \left( \coth \left( \sqrt{q^2 + \omega_n^2} L \right) - 1 \right). \quad (19) \end{aligned}$$

The first term on the right-hand side of equation (19) represents the contribution from electric field fluctuations. Based on the Bohr-Van Leeuwen theorem, only the pure electric component of the disjoining pressure is influenced by the presence of classical charged particles in a volume between metal walls. That is why this contribution depends on the inverse Debye length,  $\varkappa$ . The second and third terms in the expression for the electromagnetic pressure are due to fluctuations that are not affected by the presence of classical ions.

Eq. (19) can be rewritten as follows. Let us add and subtract the second term. This will result in a purely repulsive ion contribution to the disjoining pressure,

$$\Pi_{ion} = -\frac{T}{4\pi} \int_0^\infty dq q \sqrt{q^2 + \varkappa^2} \left( \coth \left( \sqrt{q^2 + \varkappa^2} L \right) - 1 \right) + \frac{T}{4\pi} \int_0^\infty dq q^2 \left( \coth(qL) - 1 \right), \quad (20)$$

which can be derived within the statistical field theory of the Coulomb gas<sup>15,28</sup> using image-charge concept and the negative contribution of the electromagnetic field fluctuations, which coincides exactly with the result of Schwinger and coworkers<sup>4</sup>:

$$\Pi_{EM} = -\frac{T}{\pi} \sum_{n=0}^{\infty} \int_0^\infty dq q \sqrt{q^2 + \omega_n^2} \left( \coth \left( \sqrt{q^2 + \omega_n^2} L \right) - 1 \right), \quad (21)$$

so that

$$\Pi = \Pi_{ion} + \Pi_{EM}. \quad (22)$$

Thus, our first-principle calculations justify the *separation hypothesis* discussed in the Introduction.

Note that  $\Pi_{ion}$  vanishes when  $\varkappa = 0$ . Note also that it can be simplified to

$$\Pi_{ion} = \frac{T}{8\pi L^3} \left( \zeta(3) - \frac{1}{2} \int_{2\varkappa L}^\infty \frac{dy y^2}{e^y - 1} \right). \quad (23)$$

The ion contribution to the disjoining pressure has the following asymptotic behavior

$$\Pi_{ion} \simeq \begin{cases} \frac{T\zeta(3)}{8\pi L^3}, & \varkappa L \gg 1 \\ \frac{T\varkappa^2}{4\pi L}, & \varkappa L \ll 1. \end{cases} \quad (24)$$

The electromagnetic contribution can be also simplified to<sup>4</sup>

$$\Pi_{EM} = -\frac{T}{4\pi L^3} \sum_{n=0}^{\infty} \int_{2\omega_n L}^\infty \frac{dy y^2}{e^y - 1}. \quad (25)$$

Now let us discuss which formula correctly describes the Casimir disjoining pressure in a high-temperature regime in the absence of ions - the Schwinger or Lifshitz formula. As discussed in the Introduction, Lifshitz's theory when applied to the limit of ideally conductive walls encounters an ambiguity that requires taking limits:  $\omega_n \rightarrow 0$  and  $\epsilon(\omega) \rightarrow \infty$ . As mentioned above, Schwinger et al. postulated the following order:  $\epsilon(\omega) \rightarrow \infty$  and then  $\omega_n \rightarrow 0$ . At first glance, there seem to be no fundamental reasons why this particular order should be preferred over the one discussed in the works of Lifshitz and co-workers<sup>3</sup>. However, there are two main arguments that support Schwinger's rule being correct.

First, from general physical considerations, in the limit of an ideal metal ( $\epsilon(\omega) \rightarrow \infty$ ), the specific properties of the boundary surfaces become insignificant, and we obtain a pure electromagnetic field decoupled from matter in equilibrium with perfectly reflecting flat surfaces. This pure quantum field system, in its dynamics, must obey the symmetry of four-dimensional spacetime with respect to the Poincare group, ISO(3,1). When in thermodynamic equilibrium, the symmetry of four-dimensional spacetime reduces to the symmetry of four-dimensional Euclidean space with respect to the ISO(4) group, where the time is replaced by the imaginary time (inverse temperature). As we have shown in our previous work<sup>23</sup>, the Schwinger's formula follows from the Helmholtz free energy (or partition function)<sup>23</sup> which is invariant under the ISO(4) group. At the same time, the expression obtained by Lifshitz et al. does not correspond to the partition function that is invariant under ISO(4). Thus, we conclude that the disjoining pressure obtained by Schwinger respects the symmetries of four-dimensional space. The same can be said about the tangential pressure and, therefore, surface tension that we have recently obtained<sup>23</sup>. It is important to note that, due to the ISO(4) symmetry of the partition function, we can obtain the low-temperature expansion of the disjoining pressure, as obtained by Mehra<sup>32</sup>, from a resummation of Schwinger's high-temperature expansion<sup>4,23</sup>.

The second argument is based on the fundamental principles of classical electrodynamics and statistical mechanics. The difference between the results obtained by Schwinger et al. and those derived from Lifshitz's theory is due to the fact that the latter does not account for classical radiation pressure, which occurs in the high-temperature limit. In contrast, Schwinger's approach does take this into consideration. Indeed, the zero Matsubara



frequency term in the total electromagnetic disjoining pressure<sup>4,23</sup>,

$$\Pi_{EM}^{(n=0)} = -\frac{T}{2\pi} \int_0^\infty dq q^2 (\coth(qL) - 1) = -\frac{T\zeta(3)}{4\pi L^3}, \quad (26)$$

is simply the disjoining pressure generated by equilibrium classical radiation. This is demonstrated using classical electrodynamics and the Gibbs statistical mechanical approach, as explained in Appendix C.

In conclusion, our justification of the separation hypothesis is rooted in the fundamental principles of quantum electrodynamics and quantum statistical physics. We have shown that in the Gaussian approximation the positive ionic contribution to the Casimir force, which is of electrostatic origin, can be separated from the electromagnetic fluctuations that give rise to the original Casimir effect. It is crucial to highlight the significance of calculating the zero Matsubara mode term in the electromagnetic contribution using the formula put forth by Schwinger et al. instead of Lifshitz et al. This choice is supported by the ISO(4) symmetry of the Helmholtz free energy, from which Schwinger's formula can be derived and the correspondence between the zero Matsubara mode in Schwinger's expansion and the pressure of classical electromagnetic radiation in equilibrium with conductive walls, as explained by classical electrodynamics and statistical physics.

## Appendix A: Gauge invariance of partition function

The partition function of the described system can be written as follows

$$\mathcal{Z} = \oint \mathcal{D}A \oint d\sigma[\{\mathbf{r}_i^{(+)}\}] \oint d\sigma[\{\mathbf{r}_j^{(-)}\}] e^{\mathcal{S}}, \quad (A1)$$

where

$$\mathcal{S} = \int_0^\beta d\tau \int d^3\mathbf{x} \left[ \mathcal{L}_{EM} - i(\hat{\rho}\varphi - \hat{\mathbf{j}} \cdot \mathbf{A}) \right], \quad (A2)$$

is the total action with the Lagrangian of the electromagnetic field

$$\mathcal{L}_{EM} = -\frac{1}{4} F_{\mu\nu} F_{\mu\nu}, \quad (A3)$$

where  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  is the electromagnetic tensor. Using the electric charge conservation law,

$$\partial_\tau \hat{\rho} + \nabla \cdot \hat{\mathbf{j}} = 0, \quad (A4)$$

we can easily check that (A1) is a gauge invariant.

To fix gauge, let us consider the identity

$$\oint \mathcal{D}\theta |D| \delta[\partial_\mu A_\mu^\theta(x) - B(x)] = 1, \quad (\text{A5})$$

where  $|D|$  is determined by (B5),  $A_\mu^\theta$  is defined by (B6), and  $B(x)$  is an arbitrary function;  $\delta[\cdot]$  is the functional generalization of Dirac's delta-function. Using the identity (A5), the partition function can be rewritten as

$$\mathcal{Z} = \oint \mathcal{D}\theta \oint \mathcal{D}A \oint d\sigma[\{\mathbf{r}_i^{(+)}\}] \oint d\sigma[\{\mathbf{r}_j^{(-)}\}] |D| \delta[\partial_\mu A_\mu^\theta - B] e^{\mathcal{S}}. \quad (\text{A6})$$

Calculation of the functional integral over all possible functions  $B(x)$  and dividing by gauge group volume  $\mathcal{V}_\theta = \oint \mathcal{D}\theta$ ,

$$Z = \mathcal{V}_\theta^{-1} \oint \mathcal{D}B \mathcal{Z} \exp\left(-\frac{\alpha B^2}{2}\right), \quad (\text{A7})$$

results in (1). From (A5) it follows that  $\mathcal{Z}$  in (A6) does not depend on  $B(x)$  thus the integration over  $B(x)$  is equivalent to multiplying by infinite constant<sup>23</sup>.

## Appendix B: Calculation of partition function

The partition function at  $\alpha = 1$  can be written as

$$Z = |D| (\Delta_0 \Delta_1 \Delta_2 \Delta_3)^{-\frac{1}{2}} \quad (\text{B1})$$

where  $\Delta_\mu$  is the determinant corresponding to the functional integrations over  $A_\mu$ . Let us consider the eigenfunctions of the D'Alembert operator

$$\begin{aligned} -\square v(x, \mathbf{q}, n, l) &= \lambda(\mathbf{q}, n, l) v(x, \mathbf{q}, n, l), \\ l &= 1, 2, \dots, \quad n = 0, \pm 1, \pm 2, \dots \end{aligned} \quad (\text{B2})$$

satisfying the boundary conditions at  $x_3 = 0$  and  $x_3 = L$

$$v = 0, \quad (\text{B3})$$

with eigenvalues

$$\lambda(\mathbf{q}, n, l) = \mathbf{q}^2 + q_l^2 + \omega_n^2, \quad (\text{B4})$$

The determinant  $|D|$  can be rewritten as

$$|D| = \text{Det} \left( - \frac{\delta \square \theta}{\delta \theta} \right), \quad (\text{B5})$$

where  $\theta(x)$  is the parameter of the gauge transformation<sup>23,24</sup>

$$A_\mu^\theta = A_\mu - \partial_\mu \theta. \quad (\text{B6})$$

It has been shown recently<sup>23</sup> that  $\theta(x)$  is satisfied by the same boundary conditions as  $v$ , so that

$$|D| = \exp \left( \sum_{l=1}^{\infty} \ln \bar{\lambda}_l \right), \quad (\text{B7})$$

where

$$\ln \bar{\lambda}_l = \mathcal{A} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \sum_{n \in \mathbb{Z}} \ln \lambda(\mathbf{q}, n, l) \quad (\text{B8})$$

and also  $A_0$ ,  $A_1$  and  $A_2$  are satisfied the same boundary conditions. Therefore

$$\int \mathcal{D}A_1 \exp \left( \frac{1}{2} \left( A_1, \square A_1 \right) \right) = \Delta_1^{-\frac{1}{2}}, \quad (\text{B9})$$

$$\Delta_1 = |D|, \quad (\text{B10})$$

and for  $A_2$  one gets the same result as for  $A_1$ , i.e.  $\Delta_1 = \Delta_2$ . Let us note that  $v(x, \mathbf{q}, 0, l)$  are eigenfunctions of other operator

$$(\varkappa^2 - \square) v(x, \mathbf{q}, 0, l) = \lambda'(\mathbf{q}, 0, l) v(x, \mathbf{q}, 0, l), \quad (\text{B11})$$

with other eigenvalues

$$\lambda'(\mathbf{q}, 0, l) = \mathbf{q}^2 + q_l^2 + \varkappa^2, \quad (\text{B12})$$

where

$$\varkappa^2 = \beta \sum_{a=\pm} n_a q_a^2. \quad (\text{B13})$$

From this, one can get

$$\int \mathcal{D}\varphi \exp \left( \frac{1}{2} \left( \varphi, \square \varphi \right) - \frac{1}{2} \sum_{a=\pm} q_a^2 n_a \int d^3 \mathbf{x} \left( \int_0^\beta d\tau \varphi(\tau, \mathbf{x}) \right)^2 \right) = \Delta_0^{-\frac{1}{2}}, \quad (\text{B14})$$

where

$$\varphi(x) = \varphi(\tau, \mathbf{x}) = \sum_{l=1}^{\infty} \sum_{n \in \mathbb{Z}} \int d^2 \mathbf{q} a_0(\mathbf{q}, n, l) v(x, \mathbf{q}, n, l) + h.c., \quad (\text{B15})$$

and after integrating  $\varphi(x)$  with respect to  $\tau$  in the expression (B14) only one non-zero term with  $n = 0$  remains in sum (B15) because the integral of the function  $v(x, \mathbf{q}, n, l) \sim e^{i\omega_n \tau}$  is equal to zero at  $n \neq 0$ . Thus, we have

$$\ln \Delta_0 = \mathcal{A} \sum_{l=1}^{\infty} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \left( \ln \lambda'(\mathbf{q}, 0, l) + 2 \sum_{n=1}^{\infty} \ln \lambda(\mathbf{q}, n, l) \right), \quad (\text{B16})$$

$A_3$  is satisfied the other boundary conditions at  $x_3 = 0$  and  $x_3 = L$

$$\partial_3 A_3 = 0, \quad (\text{B17})$$

due to that on the surfaces of plates the charge density is equal to zero, i.e.  $\nabla \cdot \mathbf{E} = 0$ . Thus, let us consider the eigenfunctions

$$\begin{aligned} -\square u(x, \mathbf{q}, n, l) &= \lambda(\mathbf{q}, n, l) u(x, \mathbf{q}, n, l), \\ l &= 0, 1, 2, \dots, \quad n = 0, \pm 1, \pm 2, \dots \end{aligned} \quad (\text{B18})$$

with the boundary conditions at  $x_3 = 0$  and  $x_3 = L$

$$\partial_3 u = 0, \quad (\text{B19})$$

then one can get

$$\int \mathcal{D}A_3 \exp \left( \frac{1}{2} \left( A_3, \square A_3 \right) \right) = \Delta_3^{-\frac{1}{2}}, \quad (\text{B20})$$

where

$$\Delta_3 = \exp \left( \sum_{l=0}^{\infty} \ln \bar{\lambda}_l \right). \quad (\text{B21})$$

that can be rewritten as

$$\ln \Delta_3 = \mathcal{A} \sum_{l=0}^{\infty} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \left( \ln \lambda(\mathbf{q}, 0, l) + 2 \sum_{n=1}^{\infty} \ln \lambda(\mathbf{q}, n, l) \right) \quad (\text{B22})$$

### Appendix C: Disjoining pressure of electromagnetic radiation in classical physics

Let us use the standing electromagnetic waves between perfect metal flat walls separated by distance  $L$ . We consider the Coulomb gauge, i.e. that  $\varphi = 0$  and  $\nabla \cdot \mathbf{A} = 0$ . The general solution of the D'Alembert equation,

$$\frac{\partial^2 \mathbf{A}}{\partial t^2} - \Delta \mathbf{A} = 0, \quad (\text{C1})$$

with boundary conditions on the walls corresponding to the ideal metal

$$\partial_3 A_3(x_1, x_2, 0, t) = \partial_3 A_3(x_1, x_2, L, t) = 0, \quad (\text{C2})$$

$$\mathbf{A}^{\parallel}(x_1, x_2, 0, t) = \mathbf{A}^{\parallel}(x_1, x_2, L, t) = 0, \quad \mathbf{A}^{\parallel} = (A_1, A_2, 0), \quad (\text{C3})$$

satisfying the gauge relation  $\nabla \cdot \mathbf{A} = 0$ , is determined by the well-known formula<sup>33</sup>

$$\mathbf{A} = \sum_{\mathbf{q}} \sum_{l=0}^{\infty} {}'' \left( a_{\mathbf{q}l}^{(1)} [\mathbf{e}_{\mathbf{q}}, \mathbf{e}_3] \sin(q_l x_3) + a_{\mathbf{q}l}^{(2)} \left( \mathbf{e}_{\mathbf{q}} \frac{i q_l}{\omega_{\mathbf{q}l}} \sin(q_l x_3) - \mathbf{e}_3 \frac{|\mathbf{q}|}{\omega_{\mathbf{q}l}} \cos(q_l x_3) \right) \right) e^{i\mathbf{q} \cdot \boldsymbol{\rho}} + c.c., \quad (\text{C4})$$

where  $\boldsymbol{\rho} = (x_1, x_2, 0)$ ,  $q_l = \pi l/L$ ,  $l = 0, 1, 2, \dots$ ,  $\mathbf{q} = (2\pi n_x/L_x, 2\pi n_y/L_y, 0)$ ,  $n_x, n_y \in Z$ ,  $\omega_{\mathbf{q}l} = \sqrt{\mathbf{q}^2 + q_l^2}$ ,  $\mathbf{e}_{\mathbf{q}} = \mathbf{q}/|\mathbf{q}|$ ,  $\mathbf{e}_3$  is the unit vector along the  $x_3$ -axis,  $[\mathbf{e}_{\mathbf{q}}, \mathbf{e}_3]$  is the vector product; the double prime on the sum means that there is an extra factor of  $1/\sqrt{2}$  for the term with  $l = 0$ .

Note that the coefficients  $a_{\mathbf{q}l}^{(\alpha)}$  (the index  $\alpha = 1, 2$  indicates the polarization direction) are the functions of time and are satisfied the equations

$$\ddot{a}_{\mathbf{q}l}^{(\alpha)} + \omega_{\mathbf{q}l}^2 a_{\mathbf{q}l}^{(\alpha)} = 0, \quad (\text{C5})$$

i.e.  $a_{\mathbf{q}l}^{(\alpha)} \sim e^{-i\omega_{\mathbf{q}l} t}$ .

The electric and magnetic fields are determined by relations

$$\mathbf{E} = -\dot{\mathbf{A}}, \quad \mathbf{B} = [\nabla, \mathbf{A}], \quad (\text{C6})$$

which yield

$$\mathbf{E} = \sum_{\mathbf{q}} \sum_{l=0}^{\infty} {}'' \left( -a_{\mathbf{q}l}^{(1)} i\omega_{\mathbf{q}l} [\mathbf{e}_{\mathbf{q}}, \mathbf{e}_3] \sin(q_l x_3) + a_{\mathbf{q}l}^{(2)} (\mathbf{e}_{\mathbf{q}} q_l \sin(q_l x_3) + i\mathbf{e}_3 |\mathbf{q}| \cos(q_l x_3)) \right) e^{i\mathbf{q} \cdot \boldsymbol{\rho}} + c.c., \quad (\text{C7})$$

$$\mathbf{B} = \sum_{\mathbf{q}} \sum_{l=0}^{\infty} {}'' \left( -a_{\mathbf{q}l}^{(2)} i\omega_{\mathbf{q}l} [\mathbf{e}_{\mathbf{q}}, \mathbf{e}_3] \cos(q_l x_3) + a_{\mathbf{q}l}^{(1)} (\mathbf{e}_{\mathbf{q}} q_l \cos(q_l x_3) - i\mathbf{e}_3 |\mathbf{q}| \sin(q_l x_3)) \right) e^{i\mathbf{q} \cdot \boldsymbol{\rho}} + c.c. \quad (\text{C8})$$

The total energy of the electromagnetic field is

$$H = \frac{1}{2} \int d^3 \mathbf{r} (\mathbf{E}^2 + \mathbf{B}^2). \quad (\text{C9})$$

With using eqs. (C4), (C7), (C8) integration yields

$$H = \frac{V}{2} \sum_{\alpha=1,2} \sum_{\mathbf{q}} \sum_{l=0}^{\infty} (q_l^2 + \mathbf{q}^2) a_{\mathbf{q}l}^{(\alpha)*} a_{\mathbf{q}l}^{(\alpha)} \quad (\text{C10})$$

where  $V = L_x L_y L$  is the total volume enclosed by the metal walls.

Then, introducing the canonical variables

$$Q_{\mathbf{q}l}^{(\alpha)} = \frac{\sqrt{V}}{2}(a_{\mathbf{q}l}^{(\alpha)} + a_{\mathbf{q}l}^{(\alpha)*}), \quad (\text{C11})$$

$$P_{\mathbf{q}l}^{(\alpha)} = -i\omega_{\mathbf{q}l} \frac{\sqrt{V}}{2}(a_{\mathbf{q}l}^{(\alpha)} - a_{\mathbf{q}l}^{(\alpha)*}), \quad (\text{C12})$$

where the symbol  $*$  means the complex conjugation, we can rewrite the total energy of the electromagnetic field in their terms as follows

$$H[P, Q] = \frac{1}{2} \sum_{\alpha=1,2} \sum_{\mathbf{q}} \sum_{l=0}^{\infty} \left( P_{\mathbf{q}l}^{(\alpha)2} + \omega_{\mathbf{q}l}^2 Q_{\mathbf{q}l}^{(\alpha)2} \right). \quad (\text{C13})$$

Note that these variables are canonical because they satisfy the Hamilton's equations:

$$\dot{Q}_{\mathbf{q}l}^{(\alpha)} = \frac{\partial H}{\partial P_{\mathbf{q}l}^{(\alpha)}}, \quad \dot{P}_{\mathbf{q}l}^{(\alpha)} = -\frac{\partial H}{\partial Q_{\mathbf{q}l}^{(\alpha)}}. \quad (\text{C14})$$

For the harmonic oscillators of the electromagnetic field which are in equilibrium with metal walls we have the following Gibbs distribution function

$$\mathcal{P}[P, Q] = \frac{1}{Z} e^{-\beta H[P, Q]}. \quad (\text{C15})$$

The Helmholtz free energy of the electromagnetic field is

$$F = -T \ln Z, \quad (\text{C16})$$

where the partition function is

$$Z = \int \mathcal{D}P \mathcal{D}Q e^{-\beta H[P, Q]}. \quad (\text{C17})$$

Calculating the Gaussian integrals, we obtain the Helmholtz free energy up to infinite constant which does not depend on  $L$ :

$$F = T \sum_{\mathbf{q}} \sum_{l=0}^{\infty} \ln(\mathbf{q}^2 + q_l^2). \quad (\text{C18})$$

The electromagnetic radiation pressure is

$$P = -\frac{\partial(F/\mathcal{A})}{\partial L}, \quad (\text{C19})$$

where  $\mathcal{A} = L_x L_y$  is the total area of the metal walls. Therefore, we obtain

$$P = -\frac{2T}{L} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \sum_{l=1}^{\infty} \frac{q_l^2}{\mathbf{q}^2 + q_l^2}. \quad (\text{C20})$$

where we have used that for very large walls we have

$$\sum_{\mathbf{q}}(\cdot) \rightarrow \mathcal{A} \int \frac{d^2 \mathbf{q}}{(2\pi)^2}(\cdot). \quad (\text{C21})$$

Using the same approach to regularizing divergent sums as in<sup>23</sup>, we obtain

$$P = -T \int \frac{d^2 \mathbf{q}}{(2\pi)^2} q \coth(qL). \quad (\text{C22})$$

In agreement with (26), the radiation disjoining pressure is

$$\Pi_{EM}^{(cl)} = P - P|_{L \rightarrow \infty} = \Pi_{EM}^{(n=0)} = -T \int \frac{d^2 \mathbf{q}}{(2\pi)^2} q (\coth(qL) - 1) = -\frac{T\zeta(3)}{4\pi L^3}. \quad (\text{C23})$$

## ACKNOWLEDGMENTS

The authors would like to express their gratitude to Rudolf Podgornik for his valuable discussions and insights on these issues. The authors thank the Russian Science Foundation (Grant No. 24-11-00096) for financial support.

## REFERENCES

- <sup>1</sup>H. B. Casimir, “On the attraction between two perfectly conducting plates,” in *Proc. Kon. Ned. Akad. Wet.*, Vol. 51 (1948) p. 793.
- <sup>2</sup>E. M. Lifshitz, M. Hamermesh, *et al.*, “The theory of molecular attractive forces between solids,” in *Perspectives in theoretical physics* (Elsevier, 1992) pp. 329–349.
- <sup>3</sup>I. E. Dzyaloshinskii, E. M. Lifshitz, and L. P. Pitaevskii, “The general theory of van der waals forces,” *Advances in Physics* **10**, 165–209 (1961).
- <sup>4</sup>J. Schwinger, L. L. DeRaad Jr, and K. A. Milton, “Casimir effect in dielectrics,” *Annals of Physics* **115**, 1–23 (1978).
- <sup>5</sup>H. B. Chan, V. A. Aksyuk, R. N. Kleiman, D. J. Bishop, and F. Capasso, “Quantum mechanical actuation of microelectromechanical systems by the casimir force,” *Science* **291**, 1941–1944 (2001).

- <sup>6</sup>G. Palasantzas, M. Sedighi, and V. B. Svetovoy, “Applications of casimir forces: Nanoscale actuation and adhesion,” *Applied Physics Letters* **117** (2020).
- <sup>7</sup>B. Elsaka, X. Yang, P. Kästner, K. Dingel, B. Sick, P. Lehmann, S. Y. Buhmann, and H. Hillmer, “Casimir effect in mems: Materials, geometries, and metrologies—A review,” *Materials* **17**, 3393 (2024).
- <sup>8</sup>Y. S. Barash and V. L. Ginzburg, “Electromagnetic fluctuations in matter and molecular (van-der-waals) forces between them,” *Soviet Physics Uspekhi* **18**, 305 (1975).
- <sup>9</sup>M. Bordag, U. Mohideen, and V. M. Mostepanenko, “New developments in the casimir effect,” *Physics reports* **353**, 1–205 (2001).
- <sup>10</sup>J. Mahanty and B. W. Ninham, “Dispersion forces,” (No Title) (1976).
- <sup>11</sup>V. A. Parsegian, *Van der Waals forces: a handbook for biologists, chemists, engineers, and physicists* (Cambridge university press, 2005).
- <sup>12</sup>E. Barouch, J. Perram, and E. Smith, “On the interdependence of van der waals and double layer forces,” *Chemical Physics Letters* **19**, 131–133 (1973).
- <sup>13</sup>V. Gorelkin and V. Smilga, “Calculation of intermolecular interaction forces between bodies separated by a film of a strong electrolyte solution,” *Soviet Journal of Experimental and Theoretical Physics* **36**, 761 (1973).
- <sup>14</sup>C. Barnes and B. Davies, “Statistical mechanics and lifshitz theory for electrolytes. part 1,” *Journal of the Chemical Society, Faraday Transactions 2: Molecular and Chemical Physics* **71**, 1667–1689 (1975).
- <sup>15</sup>B. W. Ninham and V. Yaminsky, “Ion binding and ion specificity: the hofmeister effect and onsager and lifshitz theories,” *Langmuir* **13**, 2097–2108 (1997).
- <sup>16</sup>B. Jancovici and L. Šamaj, “Screening of classical casimir forces by electrolytes in semi-infinite geometries,” *Journal of Statistical Mechanics: Theory and Experiment* **2004**, P08006 (2004).
- <sup>17</sup>B. Jancovici and L. Šamaj, “Casimir force between two ideal-conductor walls revisited,” *Europhysics Letters* **72**, 35 (2005).
- <sup>18</sup>G. Barton and N. Dombey, “Casimir effect for massive photons,” *Nature* **311**, 336–339 (1984).
- <sup>19</sup>G. Barton and N. Dombey, “The casimir effect with finite-mass photons,” *Annals of Physics* **162**, 231–272 (1985).



- <sup>20</sup>L. Teo, “Casimir effect of massive vector fields,” *Physical Review D-Particles, Fields, Gravitation, and Cosmology* **82**, 105002 (2010).
- <sup>21</sup>P. R. Buenzli and P. A. Martin, “Microscopic origin of universality in casimir forces,” *Journal of statistical physics* **119**, 273–307 (2005).
- <sup>22</sup>P. Buenzli and P. A. Martin, “Microscopic theory of the casimir force at thermal equilibrium: Large-separation asymptotics,” *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics* **77**, 011114 (2008).
- <sup>23</sup>P. Brandyshev and Y. A. Budkov, “Finite-temperature quantum field theory of casimir forces,” arXiv preprint arXiv:2409.14450 (2024).
- <sup>24</sup>J. I. Kapusta and C. Gale, *Finite-temperature field theory: Principles and applications* (Cambridge university press, 2007).
- <sup>25</sup>S. Weinberg, *The quantum theory of fields* (Cambridge university press, 1995).
- <sup>26</sup>R. P. Feynman, A. R. Hibbs, and D. F. Styer, *Quantum mechanics and path integrals* (Courier Corporation, 2010).
- <sup>27</sup>J. Zinn-Justin, *Path integrals in quantum mechanics* (OUP Oxford, 2010).
- <sup>28</sup>Y. A. Budkov and N. N. Kalikin, *Statistical Field Theory of Ion-Molecular Fluids: Fundamentals and Applications in Physical Chemistry and Electrochemistry* (Springer Nature).
- <sup>29</sup>Y. A. Budkov, “Statistical field theory of ion–molecular solutions,” *Physical Chemistry Chemical Physics* **22**, 14756–14772 (2020).
- <sup>30</sup>B. Savoie, “A rigorous proof of the bohr–van leeuwen theorem in the semiclassical limit,” *Reviews in Mathematical Physics* **27**, 1550019 (2015).
- <sup>31</sup>B. V. Derjaguin, N. V. Churaev, V. M. Muller, and V. Kisin, *Surface forces* (Springer, 1987).
- <sup>32</sup>J. Mehra, “Temperature correction to the casimir effect,” *Physica* **37**, 145–152 (1967).
- <sup>33</sup>G. Barton, “Quantum electrodynamics of spinless particles between conducting plates,” *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **320**, 251–275 (1970).