Regular subdivisions, bounds on initial ideals, and categorical limits

George Balla 1 , Daniel Corey 2 , Igor Makhlin 1 , and Victoria Schleis 34

1 *Institute of Mathematics, Technische Universität Berlin, Germany*

²*Department of Mathematical Sciences, University of Nevada, Las Vegas, Las Vegas, NV*

³*Department of Mathematical Sciences, Durham University, United Kingdom*

4 *Institute for Advanced Study, Princeton, NJ*

Abstract. We extend two known constructions that relate regular subdivisions to initial degenerations of projective toric varieties and Grassmannians. We associate a point configuration *A* with any homogeneous ideal *I*. We obtain upper and lower bounds on each initial ideal of *I* in terms of regular subdivisions of *A*. Both bounds can be interpreted categorically via limits over face posets of subdivisions. We also investigate when these bounds are exact. This is an extended abstract of a forthcoming paper.

Keywords: Grassmannians, initial degenerations, regular subdivisions, secondary fans, toric varieties

Introduction

Given an ideal $I \subset \mathbb{C}[x_1,\ldots,x_m]$, one may associate an initial ideal in_{*w*} *I* with any vector *w* ∈ **R***m*. This notion has its origins in computer algebra and plays a central role in the area of Gröbner theory. It has also established its importance in algebraic geometry, where it serves as a standard tool for constructing flat degenerations. We refer to [\[7](#page-11-0)] for an introduction to various aspects of initial ideals.

On the other hand, if *A* is a configuration of *m* points in real space, any $w \in \mathbb{R}^m$ determines a regular subdivision subd*^w A*. Defined in its modern form in [\[4](#page-11-1)], this concept can be traced to the work of Voronoy. It provides a standard method of building point-set triangulations and is widely applied in discrete and computational geometry, algebraic combinatorics and other fields. A detailed introduction can be found in [\[2](#page-11-2)].

A rather basic observation is that both of the above constructions depend on a vector $w \in \mathbb{R}^n$. A somewhat less trivial similarity is that (under mild assumptions) the set of w providing a given initial ideal of *I* is a polyhedral cone and so is the set of *w* providing a given regular subdivision of *A*. Moreover, in both cases the collection of all such cones is a complete polyhedral fan in \mathbb{R}^n : the Gröbner fan Gröb *I* and the secondary fan Sec *A*.

In certain settings these observations lead to deep connections between the two notions. We mention several such results. First, let *A* be a lattice point configuration and *I* be the respective toric ideal. In [\[7](#page-11-0), Chapter 8] it is shown that Gröb *I* refines Sec *A*, i.e.

the initial ideal determines the subdivision uniquely. Moreover, results in [\[7](#page-11-0), [9](#page-11-3)] provide a rather nice geometric interpretation: the irreducible components of the zero set of in*^w I* are the toric varieties of the maximal cells of subd_{*w*} A .

Now, let $I_{k,n}$ be the Plücker ideal defining the Grassmannian $Gr(k, n)$ and let A consist of the vertices of the hypersimplex $\Delta(k, n)$. A particularly important subfan of Gröb $I_{k,n}$ is the tropical Grassmannian Trop *Ik*,*n*. An important subfan of Sec *A* is the Dressian Dr(*k*, *n*), parametrizing matroid subdivisions of ∆(*k*, *n*). In [\[8](#page-11-4)] it is shown that every cone of Trop $I_{k,n}$ is contained in a cone of $Dr(k,n)$, providing a morphism between the two subfans. Subsequently, [\[1\]](#page-11-5) establishes a relationship between the initial degeneration and the matroid subdivision determined by $w \in \text{Trop } I_{k,n}$: the very affine scheme defined by $\int \ln w \, I_{k,n}$ admits a closed immersion into a certain inverse limit over the cells of subd_{*w*} *A*.

The broad motivation of our project is to search for a unifying and generalizing context for the above results. In this abstract we present extensions of the results in [\[1](#page-11-5)] and [\[9](#page-11-3)] to arbitrary projective schemes, also showing that these two settings should be viewed as dual to each other.

As a first key step we associate a point configuration $\mathcal{A}(I)$ with any homogeneous ideal *I*. One may view $A(I)$ as a projection of the unit simplex onto the lineality space of Gröb *I*. For the Plücker ideal this is the hypersimplex, for a toric ideal *I* it is the point configuration defining *I*.

The first main result we describe concerns bounds on in*^w I* by ideals associated to regular subdivisions of $A(I)$. To the subdivision $\Theta = \text{subd}_w A(I)$ we associate an ideal *I*_{*w*} defined as a sum of ideals over the cells of Θ . Dually, to the subdivision Θ^* = subd_{$-w$} $\mathcal{A}(I)$ an ideal *I^w* defined as an intersection of ideals over the cells of Θ^* .

Theorem 0.1 (cf. Theorems [5.3](#page-6-0) and [5.10\)](#page-7-0). In $\mathbb{C}[x_1, \ldots, x_n]$, we have the inclusion of ideals $I_w \subseteq \text{in}_w I \subseteq I^w$.

If $I = I_{k,n}$, then the inverse limit in [\[1\]](#page-11-5) may be realized as the very affine scheme cut out by the ideal I_w . If *I* is toric, then I^w is the radical of $\text{in}_w I$ as show in [\[9](#page-11-3)].

Next, with every cell Δ of Θ we associate a ring R_{Δ} . For faces $\Delta \subset \Gamma$ of Θ , we have an embedding $R_{\Delta} \to R_{\Gamma}$. Dually, we associate rings R^{Δ} with cells Δ of Θ^* and obtain surjections $R^{\overline{\Gamma}} \to R^{\Delta}$ for $\Delta \subset \Gamma$. Thus, the assignments $\Delta \mapsto R_{\Delta}$ and $\Delta \mapsto R^{\Delta}$ define diagrams indexed by the face posets of Θ and Θ[∗] respectively. We find the limits of these diagrams. Set $R_w = \mathbb{C}[x_1, \dots, x_m]/I_w$ and $R^w = \mathbb{C}[x_1, \dots, x_m]/I^w$.

Theorem 0.2 (cf. Theorems [6.2](#page-9-0) and [6.4\)](#page-9-1)**.** *We have canonical isomorphisms* $R_w \cong \lim_{\to \infty} R_{\Delta}$ $and R^w \cong \varprojlim_{\Theta^*} R^{\Delta}.$

The ideals I_w and I^w are algorithmically easier than in_{*w*} I to compute, and a natural question is when are these inclusions actually equalities. We quantify the accuracy of these bounds by the sets $\Omega(I) = \{w \in \mathbb{R}^n : I_w = in_w I\}$, and $\Omega^*(I) = \{w \in \mathbb{R}^n : I_w = I\}$ $\{in_{w} I\}$. Further, we prove a structural result about these sets.

Theorem 0.3 (cf. Theorems [7.1](#page-10-0) and [7.3\)](#page-10-1). *The set* $\Omega(I)$ *is the support of a subfan of* Sec (A) *, and* $\Omega^*(I)$ *is the support of a subfan of* $-$ Sec(*A*)*.*

Finally, we extend some of these results to the setting of very affine schemes. Suppose that the ideal *I* contains no monomials and hence defines a very affine scheme. We associate to each cell Δ of Θ a very affine scheme X_{Δ}° . Then, for faces $\Delta \subset \Gamma$ we have morphisms $X_{\Gamma}^{\circ} \to X_{\Delta}^{\circ}$. The assignment $\Delta \mapsto X_{\Delta}^{\circ}$ defines a finite diagram of very affine schemes and we may form their limit $\lim_{\Delta \to 0} X^{\circ}_{\Delta}$ in the category of affine schemes. Our final result establishes a naturally defined morphism of schemes

$$
\operatorname{in}_w X^\circ \to \varprojlim_{\Theta} X^\circ_{\Delta}
$$

which, under mild hypotheses, is a closed immersion. This generalizes the main theorem in [\[1\]](#page-11-5).

1 Point configurations and subspaces

In this section we define a simple correspondence between point configurations and vector subspaces that plays an important role in our results.

For the entirety of this paper we fix a finite labeling set *E*. An *E-labeled point configuration* in the **R**-vector space *V* is a sequence of points $A = (a_e)_{e \in E}$ in *V*. We denote by aff *A* the affine hull of *A*, the dimension of *A* is dim $A = \dim(\text{aff } A)$. Two point configurations $A = (a_e)_{e \in E}$ and $B = (b_e)_{e \in E}$ are *affinely equivalent* if there is an affine bijection φ : aff *A* → aff *B* such that φ (a_e) = b_e for all $e \in E$. Denote by [*A*]_{aff} the class of all point configurations affinely equivalent to *A*.

Definition 1.1. *Given a point configuration* $A = (a_e)_{e \in E}$ *in the* **R***-vector space V, define the following linear subspace of* **R***^E :*

 $\mathcal{L}(A) = \{w \in \mathbb{R}^E : \text{ there is an affine function } f \colon V \to \mathbb{R} \text{ such that } w_e = f(a_e) \text{ for all } e \in E\}.$

Evidently, dim $\mathcal{L}(A) = \dim A + 1$ and $\mathcal{L}(A)$ contains the all-ones vector $(1, \ldots, 1)$. Furthermore, if *A* and *B* are affinely equivalent, then $\mathcal{L}(A) = \mathcal{L}(B)$. In fact, the assignment $[A]_{\text{aff}} \mapsto \mathcal{L}(A)$ defines a bijection between affine equivalence classes and subspaces containing $(1, \ldots, 1)$. The inverse bijection is obtained as follows. Let $\{u_e\}_{e \in E}$ be the standard basis of (**R***^E*) ∗ .

Definition 1.2. *Given a linear subspace* $L ⊂ ℝ^E$, *let u*^{L} *denote the restriction of u_e to L. The corresponding E-labeled point configuration in L*[∗] *is*

$$
\mathcal{A}(L)=(u_e^L)_{e\in E}.
$$

The following is easy to check.

Proposition 1.3. The map $L \mapsto [\mathcal{A}(L)]_{\text{aff}}$ is a bijection from the set of subspaces in \mathbb{R}^E con*taining* (1, . . . , 1) *to the set of affine equivalence classes of E-labeled point configurations. The inverse bijection takes the affine equivalence class of A to* $\mathcal{L}(A)$ *, i.e. one has* $\mathcal{L}(\mathcal{A}(L)) = L$ *for any L containing* (1, . . . , 1)*.*

For a subspace $L \subset \mathbb{R}^E$ there is an alternative realization of the point configuration $\mathcal{A}(L)$ via orthogonal projections. The standard scalar product on \mathbb{R}^E induces an isomorphism $\iota_L\colon L\to L^*$. Let $\pi_L\colon \mathbb{R}^E\to L$ denote the orthogonal projection onto *L* and let $(\varepsilon_e)_{e \in E}$ be the standard basis in \mathbb{R}^E .

Proposition 1.4. The isomorphism ι_L takes the point configuration $(\pi_L(\varepsilon_e))_{e \in E}$ to $\mathcal{A}(L)$ *.* In *particular, the two point configurations are affinely equivalent.*

2 Regular subdivisions and secondary fans

We briefly overview regular subdivisions and secondary fans of point configurations. A comprehensive treatment can be found in [\[2](#page-11-2)].

For an *E*-labeled point configuration *A* in *V*, a *subpoint configuration* of *A* is any point configuration of the form $A' = (a_e)_{e \in E'}$ with labeling set $E' \subseteq E$; we write $A' \subseteq A$. A *face* of *A* is a subpoint configuration of *A* that has the form A_f for some $f \in V^*$, where

$$
A_f = (a_{e'} \in A : f(a_{e'}) \le f(a_e) \text{ for all } e \in E).
$$

Given $w \in \mathbb{R}^E$, the *lifted point configuration* A^w *in* $V \oplus \mathbb{R}$ *is* $A^w = (a_e \oplus w_e)_{e \in E}$ *. For* $f \in V^*$ we may consider the face *A w* $f_{\bigoplus 1}^w$ of A^w where $f \oplus 1 \in (V \oplus \mathbb{R})^*$ evaluates as

$$
(f \oplus 1)(v \oplus w) = f(v) + w.
$$

Point configurations of the form A_{f}^w $f_{\oplus 1}^w$ are also known as the *lower faces* of A^w . They should be thought of as those faces of A^w visible from points $v \oplus w$ with $w \ll 0$.

Now let $\pi\colon V\oplus\mathbb{R}\to V$ denote the natural projection. Then $\Delta_f = \pi(A^{w}_{f^c})$ $_{f\oplus 1}^w)$ is a subpoint configuration of *A* for any $f \in V^*$. The *regular subdivision* of *A* produced by w , denoted by $\mathsf{subd}_w(A)$, is the set of all subpoint configurations of the form Δ_f . These subpoint configurations are also known as the *cells* of the regular subdivision. The cells form a polyhedral subdivision of *A*, in particular, a subpoint configuration Γ ⊂ ∆*^f* is also in subd $_w(A)$ if and only if Γ is a face of Δ_f . In general, there may be points in A that are not contained in any cell of $\mathrm{subd}_w(A)$.

Given a regular subdivision Θ of *A*, define

$$
\tau(A,\Theta)^{\circ} = \{ w \in \mathbb{R}^{E} : \text{subd}_{w}(A) = \Theta \}, \quad \tau(A,\Theta) = \overline{\tau(A,\Theta)^{\circ}}.
$$

The sets $\tau(A,\Theta)^\circ$ are relatively open polyhedral cones in $\mathbb{R}^E.$ The cone $\tau(A,\Theta_2)$ is a face of the cone $\tau(A, \Theta_1)$ if and only if the subdivision Θ_1 is a refinement of the subdivision Θ2. The *secondary fan* of *A*, denoted by Sec *A*, is the complete polyhedral fan in **R***^E* whose cones are the $\tau(A, \Theta)$ as Θ ranges over all regular subdivisions of A.

It is clear that affinely equivalent *E*-labeled point configurations have the same secondary fan. Furthermore, the lineality space of Sec *A* is the set of those *w* for which the subdivision subd_{*w*}(*A*) is trivial, i.e. consists of the faces *A*. This happens if and only if there is an affine function *f* on \mathbb{R}^E such that $w_e = f(a_e)$ for all $e \in E$. We obtain

Proposition 2.1. *The lineality space of* Sec *A* is $\mathcal{L}(A)$ *.*

3 Initial ideals and Gröbner fans

We recall the construction of the Gröbner fan of a homogeneous ideal. See, e.g. [\[7](#page-11-0), Chapter 1] for details.

Consider the polynomial ring $S = \mathbb{C}[x_e]_{e \in E}$. A vector $w \in \mathbb{R}^E$ can be viewed as an **R**-grading on *S* that takes the value w_e on x_e . A polynomial $p \in S \setminus \{0\}$ is the sum of its homogeneous components with respect to this grading. The *initial form* of *p*, denoted by in_{*w*} *p*, is its (nonzero) homogeneous component of the least occurring grading. For an ideal *I* ⊂ *S* its *initial ideal* in_{*w*} *I* is the ideal spanned by all in_{*w*} *p* with *p* ∈ *I*.

For the remainder of the paper we fix an ideal $I \subset S$ that is homogeneous with respect to the standard grading $(1, \ldots, 1)$. We denote $R = S/I$ and consider the projective scheme Proj *R*.

Initial ideals play an important role in algebraic geometry by providing flat degenerations. Specifically, for any $w \in \mathbb{R}^E$ there is a flat degeneration of Proj *R* to Proj(*S*/ in_{*w*} *I*): a flat family over \mathbb{A}^1 with the fiber over 0 isomorphic to $\text{Proj}(S/\text{in}_w I)$ and all other fibers isomorphic to Proj *R*. Such a degeneration is known as an *initial degeneration* or a *Gröbner degeneration*.

Given an initial ideal *J* of *I*, define

$$
\tau(I,I)^{\circ} = \{ w \in \mathbb{R}^{E} : \text{in}_{w} I = J \}, \quad \tau(I,I) = \overline{\tau(I,I)^{\circ}}.
$$

The sets *τ*(*I*, *J*) ◦ are relatively open polyhedral cones in **R***^E* . The *Gröbner fan* of *I*, denoted by Gröb *I*, is the complete polyhedral fan in \mathbb{R}^E whose cones are the $\tau(I, I)$ as *J* ranges over all initial ideals of *I*.

Finally, the set of those *w* for which in*^w I* does not contain monomials is the *tropicalization* of *I*, denoted by Trop *I*. It is the support of a subfan of Gröb *I*.

4 From ideals to point configurations

The observations in Section [1](#page-2-0) let us associate a point configuration with the ideal *I*. The lineality space of Gröb *I*, which we denote $\mathcal{L}(I)$, consists of all those $w \in \mathbb{R}$ for which in_{*w*} *I* = *I*, i.e. *I* is homogeneous with respect to the grading *w*. In particular, $\mathcal{L}(I)$ contains $(1, \ldots, 1)$. We denote the point configuration $\mathcal{A}(\mathcal{L}(I))$ by $\mathcal{A}(I)$.

The torus $(\mathbb{R}^*)^E$ acts naturally on *S* by automorphisms: $t(x_e) = t_e x_e$ for $t \in (\mathbb{R}^*)^E$. Viewing \mathbb{R}^E as the tangent space of $(\mathbb{R}^*)^E$ at $(1,\ldots,1)$, the subspace $\mathcal{L}(I)$ may be seen as the tangent space of the subtorus $T \subset (\mathbb{R}^*)^E$ consisting of those *t* for which $t(I) = I$.

Furthermore, the point configuration $A(I)$ is naturally realized in the character lattice of *T*. The subspace $\mathcal{L}(I) \subset \mathbb{R}^{\overline{E}}$ is rational and $\mathbb{Z}^E \cap \mathcal{L}(I)$ is a lattice of rank equal to dim $\mathcal{L}(I)$; it is the cocharacter lattice of *T*. Let $M \subset \mathcal{L}(I)^*$ denote the dual lattice, i.e. the character lattice of *T*. The points of $A(I)$ lie in *M* — the point labeled by *e* is the character mapping $t \in T$ to t_e .

We now give our two main motivating examples.

Example 4.1. *Let* $A = (a_e)_{e \in E}$ *be an E-labeled point configuration in* \mathbb{R}^m *whose points lie in the lattice* **Z***m. Suppose I is the homogeneous toric ideal defined by A, i.e. the kernel of the homomorphism*

$$
S \to \mathbb{C}[t,z_1^{\pm 1},\ldots,z_m^{\pm 1}], \quad x_e \mapsto tz_1^{(a_e)_1}\ldots z_m^{(a_e)_m}.
$$

The linear subspace L(*I*) *consists of those points w from A by applying an affine function. In other words,* $\mathcal{L}(I) = \mathcal{L}(A)$ and Proposition [1.3](#page-3-0) *implies that* $\mathcal{A}(I)$ *is affinely equivalent to* A.

Since any toric (i.e. prime binomial) ideal is the kernel of a monomial map, we see that any toric ideal I can be identified with the toric ideal defined by the lattice point configuration A(*I*)*.*

Example 4.2. *Choose integers* $1 \leq k < n$ *and let E consist of integer tuples* (i_1, \ldots, i_k) *such that*

$$
1\leq i_1<\cdots
$$

The space \mathbb{C}^E can be identified with the exterior power $\wedge^k \mathbb{C}^n$, we consider the Plücker embedding *of the Grassmannian* $Gr(k, n) \hookrightarrow \mathbb{P}(\mathbb{C}^E)$. The ring S is the homogeneous coordinate ring of $\mathbb{P}(\mathbb{C}^E)$, and the Plücker ideal $I_{k,n}\subset S$ is the vanishing ideal of $\mathrm{Gr}(k,n)$. Then $\mathcal{L}(I_{k,n})\subset \mathbb{R}^E$ *is n-dimensional and is spanned by the vectors* w_1, \ldots, w_n *where* $(w_i)_e = 1$ *if the tuple* $e \in E$ *contains i and* $(w_i)_e = 0$ *otherwise.*

Denote the points in $A(I_{k,n})$ *by a_e* for $e \in E$. Consider the basis in $\mathcal{L}(I_{k,n})^*$ dual to w_1, \ldots, w_n . With respect to this dual basis, the vector $a_e \in \mathcal{L}(I_{k,n})^*$ is the indicator vector *of the tuple e. Thus,* $\mathcal{A}(I_{k,n})$ *consists of vertices of the hypersimplex* $\Delta(k,n)$ *.*

5 Bounds on the initial ideal

In this section we fix a point $w \in \mathbb{R}^E$ and study the initial ideal in_{*w*} *I*. We construct a lower bound on this ideal in terms of the regular subdivision subd_w $A(I)$ and an upper bound in terms of subdivision subd−*^w* A(*I*). As before, we write *a^e* to denote the element of $A(I)$ labeled by $e \in E$.

5.1 The lower bound

Let Θ denote the regular subdivision subd_{*w*} $A(I)$. As a first observation we note that $in_w(I) = I$ if and only if Θ is trivial as both conditions are equivalent to $w \in \mathcal{L}(I)$.

Now consider any cell ∆ of Θ. Denote **C**[∆] = **C**[*x^e*]*ae*∈∆, we view **C**[∆] as a subring of *S*. Let $I_\Lambda \subset \mathbb{C}[\Delta]$ be the image of *I* under the surjection $\rho_\Lambda : S \to \mathbb{C}[\Delta]$ that maps every *x*^{*e*} with *e* ∈ ∆ to itself and all other *x*^{*e*} to 0. Note that *I*[∆] is an ideal in $\mathbb{C}[\Delta]$ and is also a subspace of *S*. For two cells one of which is a face of the other, the respective ideals satisfy a particularly simple relation (note that the below need not hold when Δ is an arbitrary subpoint configurations of Γ).

Proposition 5.1. *For cells* $\Delta \subset \Gamma$ *of* Θ *one has* $I_{\Delta} = I_{\Gamma} \cap \mathbb{C}[\Delta]$ *.*

In particular, this lets us verify the following key fact only for maximal cells, i.e. those of dimension dim *A*.

Proposition 5.2. *If* Δ *is a cell of* Θ *, then* $I_{\Delta} \subset \text{in}_{w} I$ *.*

For a cell Δ of Θ let \tilde{I}_{Λ} denote the ideal in *S* generated by $I_{\Lambda} \subset \mathbb{C}[\Delta] \subset S$. We set

$$
I_w = \sum_{\Delta \text{ cell of } \Theta} \widetilde{I}_{\Delta}.
$$

In other words, *I^w* is the ideal in *S* generated by the subspaces *I*∆. By Proposition [5.1](#page-6-1) it suffices to sum over maximal cells ∆. Proposition [5.2](#page-6-2) gives the following lower bound on the initial ideal.

Theorem 5.3. *The initial ideal* in_{*w*} *I* contains I_w *.*

This theorem provides a surjection from S/I_w to $S/\text{in}_w I$, which geometrically can be interpreted as an upper bound on the initial degeneration.

Corollary 5.4. *There is a closed immersion* $\text{Proj}(S/\text{in}_{w}I) \hookrightarrow \text{Proj}(S/I_{w}).$

Example 5.5. *Let* $E = \{1, 2, 3, 4\}$ *and* $I = \langle x_1x_2 - x_3x_4 \rangle$ *. The subspace* $\mathcal{L}(I)$ *is cut out in* \mathbb{R}^E *by the equation* $w_1 + w_2 = w_3 + w_4$. To determine the point configuration $A(I)$ up to affine e quivalence, we may apply Proposition [1.4.](#page-3-1) The orthogonal projections $\pi_{\mathcal L(I)}(\varepsilon_i)$ are the vertices

of a square with the first and second point diagonally opposite each other. There are two nontrivial regular subdivisions of $\mathcal{A}(I)$ *. One of them is subd_w* $\mathcal{A}(I)$ *, induced by* $w_1 + w_2 < w_3 + w_4$ *. It has maximal two maximal cells:* $\Delta_3 = (a_1, a_2, a_3)$ *and* $\Delta_4 = (a_1, a_2, a_4)$ *. We see that in this case* I_{Δ_3} and I_{Δ_4} are the ideals generated by x_1x_2 in the respective rings. We obtain $I_w=\langle x_1x_2\rangle=0$ in_{*w*} *I.* The other nontrivial subdivision is given by w with $w_1 + w_2 > w_3 + w_4$ and also has *two maximal cells. In this case we similarly obtain* $I_w = in_w I = \langle x_3 x_4 \rangle$ *. Consequently, for the chosen ideal I the bound in Theorem [5.3](#page-6-0) is exact for any* $w \in \mathbb{R}^E$ *.*

Example 5.6. *A simple example of a proper inclusion* $I_w \subset \text{in}_w I$ *is given by* $E = \{1, 2\}$ *, the ideal* $I = \langle x_1x_2(x_1 - x_2) \rangle$ *and w with w*₁ $\neq w_2$ *. In this case* $\mathcal{A}(I)$ *consists of two coinciding points but* Θ *has only one cell consisting of a single point. This leads to* $I_w = 0$ *.*

We now consider the setting of Example [4.2](#page-5-0) in the special case $k = 2$, i.e. when *I* is the Plücker ideal defining $Gr(2, n)$. The following result can be deduced using the explicit description of Trop *I* given in [\[6](#page-11-6)] and the description of the corresponding subdivisions of ∆(2, *n*) given in [\[5\]](#page-11-7). It shows that the lower bound provided by Theorem [5.3](#page-6-0) can be exact in a nontrivial setting.

Theorem 5.7. *If* $I = I_{2,n}$ *is the Plücker ideal and w* \in Trop *I, then* $I_w = in_w I$ *.*

5.2 The upper bound

Let Θ^* denote the regular subdivision subd_{-*w*} $\mathcal{A}(I)$. In other words, Θ^* is the subdivision given by the *upper* faces of the lifted point configuration *A ^w*. For a cell ∆ of Θ[∗] we consider the ideal $I^{\Delta} = I \cap \mathbb{C}[\Delta]$ in $\mathbb{C}[\Delta]$, the latter ring defined as above. We give a series of statements concerning these ideals which are visibly dual to the statements in the previous subsection. For cells $\Delta \subset \Gamma$ of Θ^* let ρ_{Δ}^{Γ} ∆ denote the projection from **C**[Γ] to $\mathbb{C}[\Delta]$ taking $x_e \in \mathbb{C}[\Delta]$ to itself and all other variables to zero.

Proposition 5.8. *For cells* ∆ ⊂ Γ *of* Θ[∗] *one has ρ* Γ $\frac{\Gamma}{\Delta}(I^{\Gamma}) = I^{\Delta}.$

For a cell Δ of Θ^* , let \widetilde{I}^{Δ} denote the ideal in *S* generated by I^{Δ} and all x_e with $e \notin \Delta$. Proposition [5.8](#page-7-1) implies that $\tilde{I}^{\Gamma} \subset \tilde{I}^{\Delta}$ when $\Delta \subset \Gamma$. This, again, allows us to prove the next claim only for maximal cells ∆.

Proposition 5.9. *If* Δ *is a cell of* Θ^* *then* $\text{in}_w I \subset \widetilde{I}^{\Delta}$ *.*

We now set

$$
I^w = \bigcap_{\Delta \text{ cell of } \Theta^*} \widetilde{I}^{\Delta}.
$$

In fact, we have seen that it suffices to intersect over maximal ∆. Proposition [5.9](#page-7-2) gives an upper bound on the initial ideal.

Theorem 5.10. *The initial ideal* in_{*w*} *I is contained in* I^w *.*

The theorem provides a surjection from S/in_{w} *I* to S/I^{w} and its geometric counterpart is a lower bound on the initial degeneration.

Corollary 5.11. *There is a closed immerison of* $\text{Proj}(S/I^w)$ *into* $\text{Proj}(S/\text{in}_w I)$ *.*

Example 5.12. *Consider the setting of Example* [5.5.](#page-6-3) *If* $w_1 + w_2 > w_3 + w_4$, then subd_{-*w*} A(*I*) *has two maximal cells: the same two point configurations* Δ_3 *and* Δ_4 *. One sees that both* I^{Δ_3} and I^{Δ_4} are zero. However, $\widetilde{I}^{\Delta_3} = \langle x_4 \rangle$ and $\widetilde{I}^{\Delta_4} = \langle x_3 \rangle$ and, therefore, $I^w = \langle x_3 x_4 \rangle = \mathrm{in}_w I$. *Similarly, for* $w_1 + w_2 < w_3 + w_4$ *one has* $I^w = \langle x_1 x_2 \rangle = \text{in}_w I$. *Thus, for the chosen* I the *bound in Theorem [5.10](#page-7-0) is also exact for all w.*

In the above example *I* is toric. The restriction of the construction in this section to toric ideals is particularly well-behaved, this setting is discussed in [\[9\]](#page-11-3). As seen in Example [4.1,](#page-5-1) if *I* is toric, it can be identified with the toric ideal defined by the lattice point configuration A(*I*). From this we observe that *I* [∆] is the toric ideal defined by ∆ and Proj(*S*/e*I* [∆]) is the toric variety corresponding to ∆. Hence, Proj(*S*/*I ^w*) is a *semitoric variety* whose irreducible components are the toric varieties $\text{Proj}(S/\tilde{I}^{\Delta})$ for maximal cells ∆. Furthermore, it is precisely the reduction of the initial degeneration Proj(*S*/ in*^w I*).

Theorem 5.13 ([\[9,](#page-11-3) Theorem 3]). If I is toric, then I^w is the radical of $\text{in}_w I$.

6 Categorical limits

We present categorical counterparts of the results in the previous section. The quotients S/I_w and S/I^w arise as categorical limits, the surjections between these rings and $S/\text{ in }_wI$ are the respective mediating morphisms.

6.1 The colimit

Recall the setting and notations of Subsection [5.1.](#page-6-4) Denote $R_{\Delta} = S/I_{\Delta}$ for a cell Δ in Θ . Let $\mathcal{C}(\Theta)$ denote the category of cells in Θ with morphisms given by inclusions of cells. Proposition [5.1](#page-6-1) shows that we have embeddings *R*[∆] → *R*^Γ for inclusions ∆ ⊂ Γ. In other words, the rings R_{Λ} together with these embeddings form a diagram D of shape $\mathcal{C}(\Theta)$.

Next, for every ∆ we have a morphism *ψ*[∆] : *R*[∆] → *R^w* where *R^w* = *S*/*Iw*. Indeed, since *I*_{*w*} contains *I*_∆ we have *I*_∆ ⊂ *I*_{*w*} ∩ **C**[∆]. Therefore, *R*_∆ surjects onto **C**[∆]/(*I*_{*w*} ∩ **C**[∆]) while the latter embeds into R_w . We define ψ_{Δ} as the composition of these two maps. Together the ψ_{Δ} form a co-cone from $\mathcal D$ to R_w .

When every *a*_{*e*} lies in some cell of Θ our categorical interpretation is particularly simple. This is a natural case to consider: for example, this is true when $A(I)$ is the vertex set of a convex polytope, as is the case for the Plücker ideal and many others.

Theorem 6.1. If every a_e lies in some cell of Θ , then the co-cone from $\mathcal D$ to R_w formed by the *morphisms* ψ_{Δ} *is the colimit of* \mathcal{D} *. In short,* $R_w \cong \varinjlim_{\mathcal{C}(\Theta)} R_{\Delta}$ *.*

In the general case, an adjustment is needed. Consider those variables x_e for which *ae* is not contained in any cell of Θ. Let *S*⁰ be the ring of polynomials in all such *x^e* . We denote $\widehat{R}_\Delta = R_\Delta \otimes_{\mathbb{C}} S_0$. The embeddings $R_\Delta \to R_\Gamma$ provide embeddings $\widehat{R}_\Delta \to \widehat{R}_\Gamma$ which form a diagram $\widehat{\mathcal{D}}$. Furthermore, we have maps $\widehat{\psi}_{\Delta} \colon \widehat{R}_{\Delta} \to R_w$ given by $\widehat{\psi}_{\Delta}(p \otimes q) =$ *ψ*_Δ (p) *q*, which form a co-cone from *D* to *R_{<i>w*}</sub>.

Theorem 6.2. *The co-cone from* $\widehat{\mathcal{D}}$ *to* R_w *formed by the morphisms* $\widehat{\psi}_{\Delta}$ *is the colimit of* $\widehat{\mathcal{D}}$ *. In* $short$, $R_w \cong \varinjlim_{\mathcal{C}(\Theta)} \widehat{R}_\Delta$ *.*

Now, we also have a co-cone of morphisms ζ_{Δ} : $\widehat{R}_{\Delta} \to S/\operatorname{in}_{w} I$. Indeed, the surjection *S* → *S*/ in_{*w*} *I* provides a map S_0 → *S*/ in_{*w*} *I*. To obtain a map R_Λ → *S*/ in_{*w*} *I*, note that *I*[∆] ⊂ (in*^w I*) ∩ **C**[∆] in view of Proposition [5.2,](#page-6-2) and consider the composition

$$
R_{\Delta} \twoheadrightarrow \mathbb{C}[\Delta]/((\text{in}_{w} I) \cap \mathbb{C}[\Delta]) \hookrightarrow S/\text{in}_{w} I.
$$

We define *ξ*∆ as the tensor product of the two maps (of course, the first factor is not needed when every *a*_{*e*} lies in some cell of Θ). Theorem [6.2](#page-9-0) now implies the following.

Corollary 6.3. We have a unique morphism $\alpha: R_w \to S/\text{in}_w I$ such that $\alpha \circ \hat{\psi}_{\Delta} = \xi_{\Delta}$ for all *cells* ∆*. Furthermore, α coincides with the surjection provided by Theorem [5.3.](#page-6-0)*

6.2 The limit

We return to the setting and notations of Subsection [5.2.](#page-7-3) Remarkably, in this case the categorical interpretation is easier to define and does not require us to adjust for points not lying in any cell.

For a cell Δ of Θ^* denote the quotient $\mathbb{C}[\Delta]/I^{\Delta}$ by R^{Δ} . For an inclusion of cells $\Delta \subset \Gamma$, Proposition [5.8](#page-7-1) provides a surjection $R^{\Gamma} \to R^{\Delta}$. Together with these surjections the rings R^{Δ} form a diagram \mathcal{D}^* over the opposite category $\mathcal{C}(\Theta^*)^{\text{op}}$. We have $S/\widetilde{I}^{\Delta} = R^{\Delta}$ and $I^w \subset \tilde{I}^{\Delta}$ which provides surjections $\psi^{\Delta} \colon R^w \to \tilde{R}^{\Delta}$. These surjections form a cone from R^w to \mathcal{D}^* .

Theorem 6.4. The cone from R^w to \mathcal{D}^* formed by the morphisms ψ^{Δ} is the limit of \mathcal{D}^* . In short, $R^w \cong \varprojlim_{\mathcal{C}(\Theta^*)} R^{\Delta}$.

The surjections $\xi^\Delta\colon S/\mathop{\mathrm{in}}\nolimits_w I \to R^\Delta$ given by Proposition [5.9](#page-7-2) form a cone from $S/\mathop{\mathrm{in}}\nolimits_w I$ to \mathcal{D}^* . Proposition [6.4](#page-9-1) provides

Corollary 6.5. We have a unique morphism α : $S/\text{in}_w I \to R^w$ such that $\psi^{\Delta} \circ \alpha = \xi^{\Delta}$ for all *cells* ∆*. Furthermore, α coincides with the surjection provided by Theorem [5.10.](#page-7-0)*

7 Exactness of bounds and subfans

An immediate question raised by the above is for which *w* the bounds on the initial ideal provided by Theorems [5.3](#page-6-0) and [5.10](#page-7-0) are exact. We show that in both cases the set of such *w* has a surprisingly nice structure: it is the support of a subfan in the secondary fan.

First, let $\Omega(I)$ denote the set of $w \in \mathbb{R}^E$ for which $I_w = \text{in}_w I$. Since I_w is determined by subd_{*w*} $A(I)$, one sees that $\Omega(I)$ is union of relative interiors of cones in the common refinement of Gröb *I* and Sec $A(I)$. A stronger statement is also not hard to obtain: $\Omega(I)$ is a union of relative interiors of cones in Sec $\mathcal{A}(I)$. Indeed, suppose $I^w = \text{in}_w I$ and subd_w' $A(I)$ = subd_w $A(I)$. Then $I^w = I^{w'}$. However, in_{w'} I cannot be properly contained in $\text{in}_{w} I$ since both have the same Hilbert series as *I*. Hence, $I^{w'} = \text{in}_{w'} I$. A more involved argument can be used to further refine the claim.

Theorem 7.1. *The set* $\Omega(I)$ *is the support of a subfan of* Sec $\mathcal{A}(I)$ *.*

Example 7.2. Again, suppose that I is the Plücker ideal $I_{2,n}$. In this case, Sec $A(I)$ is the *secondary fan of* ∆(2, *n*)*. A regular subdivision of* ∆(2, *n*) *is* matroidal *if for every cell* ∆ *the tuples e with* $a_e \in \Delta$ *form the basis set of a matroid. The set of all w for which* subd_{*w*} $\mathcal{A}(I)$ *is matroidal is the Dressian Dr(2, n), it is the support of a subfan of Sec* $A(I)$ *. In the particular case of* $I_{2,n}$ *one has* $Dr(2,n) = Trop I$ *by results of* [\[6](#page-11-6)], so both have dimension $2n - 3$. Thus, *Theorem* [5.7](#page-7-4) *shows that* $Dr(2, n) \subset \Omega(I)$ *, however,* $\Omega(I)$ *can be substantially larger. For* $n = 4$ *it is not hard to check that* $\Omega(I) = \mathbb{R}^E$ *. For n* = 5, a computation in OSCAR [\[3\]](#page-11-8) shows that $\Omega(I)$ *is the union of 72 out of the 102 maximal cones of* Sec $\mathcal{A}(I)$ *.*

Next, let $\Omega^*(I)\subset \mathbb{R}^E$ be the set of all w for which $I^w=\text{in}_w(I).$ Let $-\operatorname{Sec}\mathcal{A}(I)$ denote the fan obtained from Sec $A(I)$ by reflection in the origin. It is similarly easy to see that $\Omega^*(I)$ is a union of relative interiors of cones in $-$ Sec $\mathcal{A}(I)$. And, again, a stronger statement can also be proved.

Theorem 7.3. *The set* $\Omega^*(I)$ *is the support of a subfan of* $-$ Sec $\mathcal{A}(I)$ *.*

Example 7.4. *Suppose I is toric. Theorem* [5.13](#page-8-0) *implies that* $w \in \Omega^*(I)$ *if and only if* $\text{in}_{w} I$ *is radical. However, the results in [\[7](#page-11-0), Chapter 8] show that* in*^w I is radical if and only if* −*w* ∈ *τ*(A(*I*), Θ) *for an* unimodular triangulation Θ*, i.e. a regular subdivision with every cell formed by the vertices of a unimodular simplex. We see that* Ω[∗] (*I*) *is the union of* −*τ*(A(*I*), Θ) *over all unimodular triangulation* Θ*.*

8 The very affine setting

In this section, we consider the case of very affine varieties. The distinguishing feature of this case is that the categorical statements can be phrased geometrically in terms of diagrams of schemes instead of rings.

Let $X = \text{Proj } R$ be the closed subscheme of $\mathbb{P}(\mathbb{C}^E)$ defined by the homogeneous ideal *I* ⊂ *S*. We assume that *X* meets the dense torus $({\bf C}^*)^E / {\bf C}^*$ of ${\bf P}({\bf C}^E)$ and denote by *X*[°] the scheme-theoretic intersection of *X* with this torus. More generally, given a face ∆ of Θ, we denote by *X* ◦ ∆ the scheme theoretic intersection of *X* with the coordinate subtorus $({\mathbb C}^*)^{\Delta}/{{\mathbb C}^*}$ of ${\mathbb P}({\mathbb C}^E)$.

We also assume $w \in \text{Trop } I$, this allows us to consider the initial degeneration $\text{in}_{w} X^{\circ}$ as a closed subscheme of $({\mathbb C}^*)^E/{\mathbb C}^*$ defined by the ideal in_{*w*} *I*.

Theorem 8.1. If $\Delta \subset \Gamma$ are faces of Θ , then X_{Δ}° and X_{Γ}° are nonempty, and the coordinate $surjection \ (\mathbb{C}^*)^{\Gamma}/\mathbb{C}^* \to (\mathbb{C}^*)^{\Delta}/\mathbb{C}^*$ *induces a morphism* $X^{\circ}_{\Gamma} \to X^{\circ}_{\Delta}$ *.*

This is largely a consequence of Proposition [5.1.](#page-6-1) Consequently, we may form the finite limit of affine schemes <u> $\varprojlim X_{\Delta}^{\circ}$ </u>.

Theorem 8.2. The coordinate surjections $({\bf C}^*)^E/{\bf C}^* \to ({\bf C}^*)^{\Delta}/{\bf C}^*$ induce maps $in_w X^{\circ} \to X_{\Delta}^{\circ}$. *Therefore we have a mediating morphism*

$$
\text{in}_{w} \, X^{\circ} \to \varprojlim X^{\circ}_{\Delta}.
$$

If each a^e ∈ A(*I*) *lies in some cell of* Θ*, then this morphism is a closed immersion.*

References

- [1] D. Corey. "Initial degenerations of Grassmannians". *Selecta Mathematica* **27** (2021), p. 57.
- [2] J. A. De Loera, J. Rambau, and F. Santos. *Triangulations*. Vol. 25. Algorithms and Computation in Mathematics. Structures for algorithms and applications. Springer-Verlag, Berlin, 2010.
- [3] W. Decker, C. Eder, C. Fieker, M. Horn, and M. Joswig, eds. *The Computer Algebra System OS-CAR: Algorithms and Examples*. 1st ed. Vol. 32. Algorithms and Computation in Mathematics. Springer, 2024.
- [4] I. Gelfand, M. Kapranov, and A. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants*. Modern Birkhäuser Classics. Boston: Birkhäuser, 2008.
- [5] M. Kapranov. "Chow quotients of Grassmannians I". *Advances in Soviet Mathematics* **16** (1993), 29–110.
- [6] D. Speyer and B. Sturmfels. "The Tropical Grassmannian". *Advances in Geometry* **4** (2004), pp. 389–411.
- [7] B. Sturmfels. *Gröbner Bases and Convex Polytopes*. Vol. 8. University Lecture Series. Providence: American Mathematical Society, 1995.
- [8] J. Tevelev. "Compactifications of Subvarieties of Tori". *American Journal of Mathematics* **129** (2007), 1087–1104.
- [9] C.-G. Zhu. "Degenerations of toric ideals and toric varieties". *Journal of Mathematical Analysis and Applications* **386** (2012), pp. 613–618.