

Indices and residues: from Poincaré-Hopf to Baum-Bott, and Marco Brunella

Maurício Corrêa and José Seade

*In memory of Marco Brunella (1964-2012),
whose mathematical legacy will endure forever.*

Abstract We study and discuss invariants of vector fields and holomorphic foliations that intertwine the theories of complex analytic singular varieties and singular holomorphic foliations on complex manifolds: two different settings with many points in common.

Contents

Introduction	1
1 Indices of vector fields on isolated complete intersection singularities	5
2 Singular Holomorphic foliations, Baum-Bott residues and indices	20
3 Indices for 1-dimensional holomorphic foliations on surfaces	28
4 Indices in higher dimensions and the Poincaré problem	32
5 Log Baum–Bott residues and Aleksandrov’s logarithmic index	38
6 Final remarks	41
References	43

Introduction

Holomorphic vector fields on complex manifolds determine holomorphic foliations by complex curves with singularities at the zeros -or singularities- of the vector field.

Maurício Corrêa

Università degli Studi di Bari, Via E. Orabona 4, I-70125, Bari, Italy, e-mail: mauricio.barros@uniba.it, mauriciomatufmg@gmail.com

José Seade

Instituto de Matemáticas, Unidad Cuernavaca, Universidad Nacional Autónoma de México. Ciudad Universitaria, 04510 Mexico City; e-mail: jseade@im.unam.mx

The leaves of this foliation are immersed Riemann surfaces. The most basic invariant of a vector field at an isolated singularity is its Poincaré-Hopf local index, and the Poincaré-Hopf index theorem can be regarded as saying that the Euler characteristic of the manifold, which is a global topological invariant, actually is localized at the singularities of the vector field. A myriad of important lines of research spring from this theorem, permeating all mathematics. Two of them are particularly relevant for us:

- The theory of characteristic classes of vector bundles and singular varieties. This is much related with indices of vector fields and frames; and
- The analogous theory for holomorphic singular foliations on manifolds.

Indeed, the foundational results by R. Bott [15] on indices and residues for holomorphic vector fields with singularities (not necessarily isolated) have profoundly influenced areas ranging from foliation theory to applications in enumerative algebraic geometry [54, 71]. This latter field was notably advanced by Kontsevich [71], who provided a rigorous framework for formulating and verifying predictions in the enumerative geometry of curves, as derived from Mirror Symmetry. In [11] Baum and Bott subsequently generalized Bott's theorem to meromorphic vector fields, and later extended these for higher-dimensional foliations [12]. Whence, we see that the indices and residues associated with foliations are of great significance, and the work of M. Brunella, T. Suwa, M. Soares and others turned this into a rich and fascinating area of mathematics where a lot is happening nowadays. See for instance [89] for an account on this subject.

On the other hand the astounding impact of characteristic classes in manifolds theory of course inspired the search for such invariants in the setting of singular varieties. The work of D. P. Sullivan, M.-H. Schwartz, R. MacPherson, W. Fulton, P. Baum and others laid down the foundations of the theory, and then the work of J.-P. Brasselet, J. Schürmann, L. Maxim, P. Aluffi and others are making this a rich and fascinating theory as well. And as noted for instance in [21, 22, 51, 29], the theory of indices of vector fields and 1-forms comes into the picture playing an important role.

In fact the study of indices and residues is of utmost importance in the local and global study of both, singular holomorphic foliations in complex manifolds and singular complex analytic varieties: two different theories with many points in common. The purpose of this work is to thread a path intertwining these two theories. We use as spearhead the GSV-index.

The first indices of vector fields and 1-forms on singular varieties in the literature appeared in the work of M. H. Schwartz [82] and R. MacPherson [76]. Yet, this was only for specific vector fields or 1-forms, aimed to defining Chern classes for singular varieties. The GSV index was introduced in [83, 62] for vector fields on isolated complex hypersurface singularities, and it began the theory of indices of vector fields and 1-forms on singular varieties.

The definition of the GSV index in 1987 was followed by the radial index (1991, and then again by different authors in 1997, 1998), the homological index (1998), indices for 1-forms (2001), the local Euler obstruction for vector fields (2004) and

for 1-forms (2005), the logarithmic index (2005) and several other related invariants. We briefly discuss these invariants in Section 1; we also comment on their relation with Chern classes for singular varieties, and we refer the reader to [22, 29] for more on this subject.

From the viewpoint of foliations and residues, in his remarkable paper [23], M. Brunella gave in 1997 an interpretation of the GSV index in the setting of holomorphic 1-dimensional foliations in complex surfaces, showing that this index is of great importance in the theory of holomorphic foliations. In fact several of the invariants that we study in the sequel spring from the Baum-Bott residues, so we briefly discuss these in Section 2. This section also serves as an introduction to the subject of singular holomorphic foliations and Pfaffian systems.

One of the main problems in this field is the famous Poincaré problem, which seeks to bound the degree of invariant varieties of holomorphic foliations in terms of the degree of the foliation. In [23], Brunella showed that the non-negativity of the total sum of the GSV index of the foliation along an invariant curve serves as an obstruction to affirmatively solving the Poincaré problem. He further proved in his book [27] that this invariant plays a significant role for the classification of holomorphic foliations in complex surfaces.

Section 3 has Brunella's article [23] as its core; we focus here on 1-dimensional holomorphic foliations on complex surfaces. We briefly discuss the Baum-Bott residues in this setting, and their relations with the Milnor number for foliations, the Camacho-Sad indices, the variations defined by Khanedani and Suwa, and the GSV index as defined by Brunella. We recall Brunella's theorem saying that if we have a holomorphic foliation \mathcal{F} on a complex surface X and S is a nondicritical separatrix at a point p then:

$$\text{GSV}(\mathcal{F}, S, p) \geq 0.$$

If \mathcal{F} further is a generalized curve (*i.e.* there are no saddle-nodes in its resolution) and S is maximal, union of all separatrices at p , then:

$$\text{GSV}(\mathcal{F}, S, p) = 0.$$

And the reciprocal from [36]: if $\text{GSV}(\mathcal{F}, S, p) = 0$, then \mathcal{F} is a generalized curve at p and S is the maximal separatrix.

Sections 4 and 5 look at higher dimensions. These are mostly based on work by Corrêa and Machado [43, 44] and Corrêa, Lourenço and Machado [39]. There are two previous articles that somehow paved the way. In [85], Seade and Suwa provide Baum-Bott type formulas for foliations by curves induced by global holomorphic vector fields on complex manifolds with a non-empty boundary. Also, a Bott-type formula for complex orbifolds is studied in [46], and formulas relating residues via good resolutions are obtained. To address the case of a projective variety Y with more complicated singularities, it is natural to consider a log resolution $(X, D) \rightarrow Y$ and to relate the residues of holomorphic vector fields on X to the Chern classes of the logarithmic tangent bundle $TX(-\log D)$. This is explained in Section 5.

In Section 4, using Aleksandrov's Decomposition Theorem we define, following [44], the GSV index for Pfaff systems ω of dimension k on complex manifolds X ,

relative to an invariant variety V in X which is a reduced local complete intersection of dimension $n - k$. This is done *à la* Brunella and associates an index to each codimension 1 component $S \subset V$ of the singular set of the Pfaff system induced by a twisted form $\omega \in H^0(X, \Omega_X^{n-k} \otimes \mathcal{L})$. It is proved that these indices localize the Chern class of the line bundle $\mathcal{L} \otimes \det(N_{V/X})^{-1}$ restricted to V . Then a method is given for calculating these GSV-indices, which can be compared with formulas by Suwa [90, Proposition 5.1] and Gómez-Mont [61]. This partially addresses Suwa's question in [90, Remark 5.3.(6)]. As an application, one has that the non-negativity of the GSV-index gives an obstruction to the solution of the Poincaré problem for Pfaff systems, in concordance with Brunella's observation for holomorphic foliations on surfaces. This further yields to a bound for the Poincaré problem similar to those previously established by Esteves and Cruz [48] and by Corrêa and Jardim [38]. Related results were previously obtained in [37, 87, 35, 26]. In particular, this recovers Soares' bound in the case where the invariant hypersurface V is smooth.

In section 4.1 we recall Suwa's interpretation in [90] of the GSV and virtual indices for vector fields as residues on complete intersection varieties with isolated singularities, and the fact that if $V = \{f_1 = \dots = f_k = 0\}$ is a local complete intersection in \mathbb{C}^n , with isolated singularity at 0, invariant by a foliation by curves \mathcal{F} singular at most at 0, then the index is a localization of the top Chern class:

$$\text{GSV}(\mathcal{F}, V, 0) = \text{Res}_{c_n}(\mathcal{F}, TU|_V; 0) = \text{Vir}(\mathcal{F}, V, 0).$$

Section 5 recalls Aleksandrov's logarithmic index $\text{Ind}_{\log}(\mathcal{F}, D, p)$ defined in [4], inspired by the work of Gómez-Mont, that measures the variation between the Poincaré–Hopf and the homological indices. Then, using this index, we address (following [43, 45]) the problem of establishing a global logarithmic residue theorem for meromorphic vector fields on compact complex manifolds.

One has that if \mathcal{F} is a one-dimensional foliation with isolated singularities and logarithmic along a normal crossings divisor D in a compact complex manifold X . Then

$$\int_X c_n(T_X(-\log D) - T\mathcal{F}) = \sum_{x \in \text{Sing}(\mathcal{F}) \cap (X \setminus D)} \mu_x(\mathcal{F}) + \sum_{x \in \text{Sing}(\mathcal{F}) \cap D} \text{Ind}_{\log}(\mathcal{F}, D, x),$$

where $\mu_x(\mathcal{F})$ is the Milnor number of \mathcal{F} at x . As an application, in [43], the authors prove that if \mathcal{F} is a one-dimensional foliation on \mathbb{P}^n , with n odd, and isolated singularities (non-degenerate) along a smooth divisor D , then $\text{Sing}(\mathcal{F}) \subset D$ if and only if Soares' bound [87] for the Poincaré problem is achieved, i.e, if and only if

$$\deg(D) = \deg(\mathcal{F}) + 1.$$

Recently, in [39], a logarithmic version of the Baum–Bott theorem was obtained for more general symmetric polynomials.

In the last section we briefly discuss relations with the adjunction formula and with other invariants, such as the Milnor and Tjurina numbers for foliations.

1 Indices of vector fields on isolated complete intersection singularities

Throughout this work, by a complete intersection singularity in \mathbb{C}^{n+k} we mean the germ (say at 0) of a geometric –non necessarily algebraic– complete intersection $(V, 0)$, defined by as many holomorphic equations as its codimension, say k . If V is defined by

$$f = (f_1, \dots, f_k) : (\mathbb{B}_\varepsilon, 0) \longrightarrow (\mathbb{C}^k, 0),$$

and the critical set of f meets V only at 0, then we say that $(V, 0)$ is an isolated complete intersection singularity, an ICIS for short.

A vector field ν on V is a continuous section of the bundle $T\mathbb{C}^{n+k}|_V$ that vanishes at 0 and for each $x \in V^* = V \setminus \{0\}$ the vector $\nu(x)$ is tangent to V^* . The vector field is holomorphic if the corresponding section is holomorphic. Notice that ν must vanish at 0 because this is an isolated singularity of V .

Notice that for each $x \in V^* = V \setminus \{0\}$, the tangent space $T_x V^*$ consists of all vectors in $T_x \mathbb{C}^{n+k}$ which are mapped to 0 by the differential of all the f_i :

$$T_x V^* = \{ \zeta \in T_x \mathbb{C}^{n+k} \mid df_i(x)(\zeta) = 0, \quad \forall i = 1, \dots, k \}.$$

Example 1. If V is defined in \mathbb{C}^{2n} by a map $f : (\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}, 0)$ and we set $V = f^{-1}(0)$, then the Hamiltonian vector field

$$\tilde{\zeta}(z_1, \dots, z_{2n}) = \left(-\frac{\partial f}{\partial z_2}, \frac{\partial f}{\partial z_1}, \dots, -\frac{\partial f}{\partial z_{2n}}, \frac{\partial f}{\partial z_{2n-1}} \right)$$

is tangent to V and it is zero only at the origin. Notice that this vector field is actually tangent to all the fibers $f^{-1}(t)$.

Example 2. Now let V be defined by the homogeneous polynomial map $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$ defined by $(z_1, \dots, z_n) \mapsto z_1^k + \dots + z_n^k$ for $k \geq 2$. Then the radial vector field $(z_1, \dots, z_n) \mapsto (z_1, \dots, z_n)$ is tangent to V and we will see later that, unlike the previous example, if $k \geq 3$ this can never be extended to the ambient space with an isolated singularity at 0 and being tangent to the fibers $f^{-1}(t)$. The case $k = 2$ is special (see Example 3).

1.1 The GSV index

Given an ICIS $(V, 0)$ as above, in \mathbb{C}^{n+k} , defined by $f = (f_1, \dots, f_k)$ we have that the (complex conjugate) gradient vector fields $\bar{\nabla} f_1, \dots, \bar{\nabla} f_k$ are linearly independent everywhere and normal to V (for the usual hermitian metric in \mathbb{C}^{n+k}). Hence if ν is a continuous vector field on V singular only at 0, then the set $\{\nu(z), \bar{\nabla} f_1, \dots, \bar{\nabla} f_k\}$ is a $(k+1)$ -frame at each point in $V^* := V \setminus \{0\}$, and up to homotopy it can be assumed to be orthonormal. Hence these vector fields define a continuous map from

V^* into the Stiefel manifold of complex orthonormal $(k+1)$ -frames in \mathbb{C}^{n+k} , denoted $W_k(n+k)$.

Let $\mathbf{K} = V \cap \mathbb{S}_\varepsilon$ be the link of 0 in V . It is an oriented, real manifold of dimension $(2n-1)$ and the above frame defines a continuous map

$$\phi_v = (v, \bar{\nabla} f_1, \dots, \bar{\nabla} f_k) : \mathbf{K} \longrightarrow W_{k+1}(n+k).$$

The Stiefel manifold $W_k(n+k)$ is diffeomorphic to the homogeneous space $U(n+k)/U(n)$ and therefore the homotopy sequence associated to this fibration implies that $W_k(n+k)$ is $(2n-2)$ -connected, while its homology in dimension $(2n-1)$ is isomorphic to \mathbb{Z} . Hence, if \mathbf{K} is connected, the map ϕ_v has a well defined degree $\deg(\phi_v) \in \mathbb{Z}$, defined by means of the induced homomorphism $H_{2n-1}(\mathbf{K}) \rightarrow H_{2n-1}(W_2(n+1))$ in the usual way.

Definition 1. Assume $n \geq 2$ or $n = 1$ and V is irreducible. Then **the GSV index** of v at $0 \in V$, $\text{Ind}_{\text{GSV}}(v; V, 0)$, is the degree of the above map ϕ_v .




Fig. 1 The GSV equals the PH index in a Milnor fiber.

Remark 1. Note that the link \mathbf{K} is always connected if $n \geq 2$. For $n = 1$ the link has one connected component for each irreducible component of the plane curve V and in this case one needs to take into account the intersection numbers of the branches. We refer to [22, 3.2] and to Section 3 for more about this case, which is specially important for the theory of holomorphic foliations.

We recall that one has a Milnor fibration associated to the map f , see [77, 66, 75], and the Milnor fiber F can be regarded as a compact $2n$ -manifold with boundary $\partial F = \mathbf{K}$. By the Ehresmann Fibration Lemma there is an ambient isotopy of the sphere \mathbb{S}_ε taking \mathbf{K} into the boundary ∂F , which can be extended to a collar of \mathbf{K} , which goes into a collar of ∂F in F . Hence v can be regarded as a non-singular vector field on ∂F .

We also recall that the Milnor fiber F has the homotopy type of a bouquet of spheres of middle dimension and therefore its Betti numbers are all 0 except $b_0 = 1$ and $b_n = \mu$ for some $\mu \geq 0$

Definition 2. The number $b_n = \mu$ is **the Milnor number** of V at 0.

Proposition 1. *The GSV index has the following properties:*

(1) *The GSV index of v at 0 equals the total Poincaré-Hopf index of v in the Milnor fiber:*

$$\text{GSV}(v; V, 0) = \text{PH}(v; F).$$

(2) *If v is everywhere transverse to \mathbf{K} , then*

$$\text{GSV}(v; V, 0) = 1 + (-1)^n \mu,$$

where n is the complex dimension of V and μ is the Milnor number.

The proof is easy and we refer to [22, 3.2] for details.

We notice that we know from [77] that if V is a hypersurface germ defined by a map f , then its Milnor number μ is the Poincaré-Hopf local index of the gradient vector field ∇f , and therefore it equals the intersection number,

$$\mu = \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1}, 0}}{\left(\frac{\partial f}{\partial z_0}, \dots, \frac{\partial f}{\partial z_n}\right)}. \quad (1)$$

For complete intersections the analogous expression is given by the Lê-Greuel formula [72, 63].

Example 3. Consider the radial vector field v_{rad} in Example 2 for $k = 2$. The statement (2) in the theorem above implies:

$$\text{GSV}(v_{rad}; V, 0) = 1 + (-1)^n \mu = 1 + (-1)^n.$$

Hence if n is odd, then the index is 0 and it is an exercise to show that in this case the vector field can be extended continuously to the Milnor fibers with no singularities.

When the vector field in question is holomorphic, the GSV-index has other interesting properties. We now discuss some of these. To begin, recall that if $\zeta = (\zeta_1, \dots, \zeta_n)$ is a holomorphic vector field in \mathbb{C}^n with an isolated singularity at 0, then its local Poincaré-Hopf index is the intersection number:

$$\text{PH}(v; 0) = \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1}, 0}}{(\zeta_1, \dots, \zeta_n)} \quad (2)$$

Recall that if ζ is a holomorphic vector field on a complex variety Z (for simplicity, say with isolated singularities), then the solutions of ζ are Riemann surfaces immersed in X , and one has an associated 1-dimensional holomorphic foliation with singularities at the zeroes of ζ .

Definition 3. Let $(V_i, 0)$, $i = 1, 2$, be germs of isolated hypersurface singularities defined by holomorphic map-germs $(\mathbb{C}^{n+1}, 0) \xrightarrow{f_i} (\mathbb{C}, 0)$, and let v_1, v_2 be germs of holomorphic vector fields on the V_i with an isolated singularity at 0. Then we say that:

1. The germs $(V_1, 0)$ and $(V_2, 0)$ are **topologically equivalent** if there exists an orientation preserving local homeomorphism h of \mathbb{C}^{n+1} , such that $f_2 = f_1 \circ h$. In this case we say that h is a topological equivalence between V_1 and V_2 .
2. The holomorphic vector fields v_1, v_2 are **topologically equivalent** if there exists a topological equivalence between V_1 and V_2 carrying the leaves of the foliation \mathcal{F}_1 of v_1 into the leaves corresponding to v_2 .

One has [62, Theorem 4.5]:

Theorem 1. *Let $(V, 0)$ denote the germ of an isolated hypersurface singularity in \mathbb{C}^{n+1} . Then, the GSV-index of a holomorphic vector field is a topological invariant. That is, if ζ is a holomorphic vector field on $(V, 0)$ and it is topologically equivalent to another holomorphic vector field ζ' on some other hypersurface germ, then ζ and ζ' have the same GSV-index.*

We recall that if \mathcal{F} is a 1-dimensional holomorphic foliation on a complex manifold, then \mathcal{F} is locally free and therefore if p is an isolated singularity of the foliation, then \mathcal{F} is locally generated by a holomorphic vector field. In analogy with equation 1 one has:

Definition 4. Let $\zeta = (\zeta_1, \dots, \zeta_n)$ be a holomorphic vector field tangent to \mathcal{F} in a neighborhood of the isolated singularity p . Then the **Milnor number** of \mathcal{F} at p is the Poincaré-Hopf local index of ζ at p , so it equals:

$$\mu(\mathcal{F}, p) := \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1}, 0}}{(\zeta_1, \dots, \zeta_n)}.$$

It is easy to prove (see [62]) that this invariant equals the PH-local index of any continuous vector field tangent to \mathcal{F} near p . One gets the following corollary to Theorem 1, which is Theorem A in [30].

Theorem 2. *The Milnor number of an isolated singularity of a 1-dimensional holomorphic foliation on a smooth complex manifold, is a topological invariant.*

We now remark that, as noticed in [13], given an isolated complex singularity germ $(V, 0)$ in some \mathbb{C}^m and a holomorphic vector field ζ on V with an isolated singularity at 0, there are infinitely many extensions of ζ to a holomorphic vector field in a neighborhood of 0 in \mathbb{C}^m . Yet, as noticed in [62, Section 3], if $(V, 0)$ is defined by a single equation f in \mathbb{C}^m , and if ζ is a holomorphic vector field on V with an isolated singularity at 0, then an extension $\tilde{\zeta}$ to the ambient space can have a singular locus of dimension at most 1, because cutting down by a hypersurface can reduce the dimension at most by 1.

One has the following Theorem 3.2 from [62].

Theorem 3. *Let $(V, 0)$ denote a hypersurface singularity germ in \mathbb{C}^{n+1} defined by a function f with at most an isolated singularity at 0 (which could be a smooth point in V). Let ζ be a holomorphic vector field on V with an isolated singularity at 0. Suppose there exists a holomorphic extension $\tilde{\zeta}$ of ζ to a neighborhood U of 0 in \mathbb{C}^{n+1} such that $\tilde{\zeta}$ is tangent to the fibers of the map f . Let Σ be the zero-locus of $\tilde{\zeta}$, which has dimension at most 1. Then:*

1. *The GSV-index of ζ in V is the intersection number of a Milnor fiber $F_t = f^{-1}(t) \cap U$ and Σ . That is:*

$$\text{GSV}(\zeta; V, 0) = \sum_{p \in U} \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1}, p}}{(\tilde{\zeta}_0, \dots, \tilde{\zeta}_n, f - t)},$$

where the $\tilde{\zeta}_0$ are the components of $\tilde{\zeta}$. Hence, in this case the GSV index is non-negative and it is 0 if and only if ζ has an isolated singularity at 0.

2. *Localized at 0 the formula becomes:*

$$\text{GSV}(\zeta; V, 0) = \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1}, 0}}{(\tilde{\zeta}_0, \dots, \tilde{\zeta}_n, f)} - \dim_{\mathbb{C}} \frac{(f) \cap (\tilde{\zeta}_0, \dots, \tilde{\zeta}_n, f)}{(f \tilde{\zeta}_0, \dots, f \tilde{\zeta}_n)}.$$

Notice that the number $\dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1}, 0}}{(\tilde{\zeta}_0, \dots, \tilde{\zeta}_n, f)}$ is a reminiscent of the Tjurina number for functions (see Section 6) and can be called the Tjurina number for vector fields. This appears again in the next section, in Gómez-Mont's formula for the homological index.

For instance, coming back to Example 2, the Milnor number is $(k - 1)^{n+1}$, by [77]. Hence, if n is odd, then the GSV index of the radial vector field is negative and therefore it can never be holomorphically extended being tangent to the Milnor fibers, not even with singularities.

1.2 The homological index

Theorem 3 gives an algebraic formula for the GSV index of holomorphic vector fields under very stringent conditions. Gómez-Mont's search for an algebraic formula in the vein of (2) for the GSV index, opened an area of research which is of interest on its own; this all springs from the definition of a homological index in [61]. This has given rise to a series of papers by Gómez-Mont and several co-authors (see [13, 58, 59, 60, 14]), including the real analytic setting where it extends to singular hypersurfaces the celebrated index formula of Eisenbud-Levine [53] and Khimshiashvili [69]. There is also a homological index for 1-forms in [52], and Aleksandrov's logarithmic index for vector fields and 1-forms [4, 6] that will be discussed in Section 5. In [93, 94] M. Zach uses variations of the homological index to study interesting generalizations of the Milnor number to the case of holomorphic maps on singular varieties.

The presentation below is essentially extracted from [22, Chapter 7] and it is a *résumé* of part of Gómez-Mont's work in [61].

We recall that as noted above, if $\zeta := (\zeta_1, \dots, \zeta_n)$ is a holomorphic vector field in \mathbb{C}^n with an isolated singularity at 0, then its Poincaré-Hopf local index is the intersection number:

$$\text{PH}(\zeta; 0) = \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^n, 0}}{(\zeta_1, \dots, \zeta_n)}.$$

We would like to have a similar expression for the GSV index. Yet we know from Example 3 that this index can be negative, even for holomorphic vector fields. Therefore an algebraic expression for the GSV index cannot simply be given by the dimension of some algebra. One may notice that for vector fields in \mathbb{C}^n one has the Koszul complex

$$0 \longrightarrow \Omega_{\mathbb{C}^n, 0}^n \longrightarrow \Omega_{\mathbb{C}^n, 0}^{n-1} \longrightarrow \cdots \longrightarrow \mathcal{O}_{\mathbb{C}^n, 0} \longrightarrow 0, \quad (3)$$

where the arrows are contraction by ζ . The Euler characteristic χ of this complex is $\text{PH}(\zeta; 0)$: this is the clue for defining the homological index, though there is an interesting difference. In the smooth case, all the homology of the Koszul complex vanishes except in top dimension, which contributes to χ with the PH index. In the singular case one has a similar complex but the homology groups do not vanish. The miracle is that “almost all of them” have the same dimension, so these cancel down when considering the Euler characteristic (see [61] for the precise statement).

Let $(V, 0) \subset (\mathbb{C}^m, 0)$ be a germ of a complex analytic variety of pure dimension n , which is regular on $V \setminus \{0\}$. So V is either regular at 0 or else it has an isolated singular point at the origin. A vector field ζ on $(V, 0)$ can always be defined as the restriction to V of a vector field \widehat{v} in the ambient space which is tangent to $V \setminus \{0\}$; ζ is holomorphic if \widehat{v} can be chosen to be holomorphic. So we may write $\zeta = (a_1, \dots, a_m)$ where the a_i are restriction to V of holomorphic functions on a neighborhood of 0 in $(\mathbb{C}^m, 0)$.

It is worth noting that given any space V as above, there are always holomorphic vector fields on V with an isolated singularity at 0. This non-trivial fact is a weak form of a stronger result ([13, p. 19]): in the space $\Theta(V, 0)$ of germs of holomorphic vector fields on V at 0, those having an isolated singularity form a connected, dense open subset $\Theta_0(V, 0)$. Essentially the same result implies also that every $\zeta \in \Theta_0(V, 0)$ can be extended to a germ of a holomorphic vector field in \mathbb{C}^m with an isolated singularity, though it can also be extended with a singular locus of dimension more than 0, a fact that may be useful for explicit computations (as in the sequel).

A (germ of a) holomorphic j -form on V at 0 means the restriction to V of a holomorphic j -form on a neighborhood of 0 in \mathbb{C}^m ; two such forms in \mathbb{C}^m are equivalent if their restrictions to V coincide on a neighborhood of 0 in V . We denote by $\Omega_{V, 0}^j$ the space of all such forms (germs); these are the Kähler differential forms on V at 0. So $\Omega_{V, 0}^0$ is the local structure ring $\mathcal{O}_{(V, 0)}$ of holomorphic functions on V at 0 and each $\Omega_{V, 0}^j$ is an $\Omega_{V, 0}^0$ -module. Notice that if the germ of V at 0 is determined by (f_1, \dots, f_k) then one has:

$$\Omega_{V,0}^j := \Omega_{\mathbb{C}^m,0}^j / (f_1 \Omega_{\mathbb{C}^m,0}^j + df_1 \wedge \Omega_{\mathbb{C}^m,0}^{j-1}, \dots, f_k \Omega_{\mathbb{C}^m,0}^j + df_k \wedge \Omega_{\mathbb{C}^m,0}^{j-1}), \quad (4)$$

where d is the exterior derivative.

Now, given a holomorphic vector field \widehat{v} at $0 \in \mathbb{C}^m$ with an isolated singularity at the origin, and a differential form $\omega \in \Omega_{\mathbb{C}^m,0}^j$, we can always contract ω by ζ in the usual way, thus getting a differential form $i_v(\omega) \in \Omega_{\mathbb{C}^m,0}^{j-1}$. If $\zeta = \widehat{v}|_V$ is tangent to V , then contraction is well defined at the level of differential forms on V at 0 and one gets a complex $(\Omega_{V,0}^\bullet, \nu)$:

$$0 \longrightarrow \Omega_{V,0}^n \longrightarrow \Omega_{V,0}^{n-1} \longrightarrow \cdots \longrightarrow \mathcal{O}_{V,0} \longrightarrow 0, \quad (5)$$

where the arrows are contraction by ζ and n is the dimension of V ; of course one also has differential forms of degree $> n$, but those forms do not play a significant role here. We consider the homology groups of this complex:

$$H_j(\Omega_{V,0}^\bullet, \nu) = \text{Ker}(\Omega_{V,0}^j \rightarrow \Omega_{V,0}^{j-1}) / \text{Im}(\Omega_{V,0}^{j+1} \rightarrow \Omega_{V,0}^j).$$

The first observation in [61] is that if V is regular at 0, so that its germ at 0 is that of \mathbb{C}^n at the origin, and if $\zeta = (a_1, \dots, a_n)$ has an isolated singularity at 0, then this is the usual Koszul complex. In that case, its homology groups vanish for $j > 0$, while

$$H_0(\Omega_{V,0}^\bullet, \nu) \cong \mathcal{O}_{\mathbb{C}^n,0} / (a_1, \dots, a_n).$$

In particular the complex is exact if $\zeta(0) \neq 0$. Since the contraction maps are $\mathcal{O}_{V,0}$ -module maps, this implies that if V has an isolated singularity at the origin, then the homology groups of this complex are concentrated at 0, and they are finite dimensional because the sheaves of Kähler differentials on V are coherent. Hence it makes sense to define:

Definition 5. The *homological index* $\text{Ind}_{\text{hom}}(\zeta, 0; V)$ of the holomorphic vector field ζ on $(V, 0)$ is the Euler characteristic of the above complex:

$$\text{Ind}_{\text{hom}}(\zeta, 0; V) = \sum_{i=0}^n (-1)^i h_i(\Omega_{V,0}^\bullet, \nu),$$

where $h_i(\Omega_{V,0}^\bullet, \nu)$ is the dimension of the corresponding homology group as a vector space over \mathbb{C} .

One has:

Theorem 7. *If the germ (V, p) is a complete intersection, then the GSV and the homological indices of every holomorphic local vector field on V coincide.*

This theorem was proved by Gómez-Mont in [61] when (V, p) is a hypersurface germ, and the general case is proved in [14]. For simplicity we restrict the discussion to the case where V is a hypersurface. In this case the proof of Theorem 7 in [61] has three main steps:

Step 1: Finding an algebraic formula for the homological index.

Step 2: Observe that the two indices coincide for one vector field if and only if they coincide for every vector field.

Step 3: Use the algebraic formula for the homological index and the basic properties of the GSV index, to prove that on every hypersurface germ there exists a tangent holomorphic vector field for which the two indices are 0.

Let us explain first the algebraic formulas for the homological index. Let $\mathbb{B} = \mathbb{B}^{n+1}$ be an open ball around the origin in \mathbb{C}^{n+1} . If $f : \mathbb{B}^{n+1} \rightarrow \mathbb{C}$ is a holomorphic function with $0 \in \mathbb{C}^{n+1}$ as its only critical point, then the 1-form $\sum_{i=1}^{n+1} \frac{\partial f}{\partial z_i} dz_i$ vanishes only at $0 \in \mathbb{C}^{n+1}$. Let \mathcal{I}_f be the Jacobian ideal $(\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_{n+1}}) \subset \mathcal{O}_{\mathbb{C}^{n+1},0}$ of f . Given a vector field ζ tangent to V , with a unique singularity at 0 , restriction of a vector field $\widehat{v} = (a_1, \dots, a_{n+1})$ on \mathbb{B} with a unique singularity at the origin, it is shown in [61, Theorem 2] that the homology groups $H_i(\Omega_{V,0}^\bullet, \zeta)$ of the above complex have dimensions:

$$\begin{aligned} h_0(\Omega_{V,0}^\bullet, \zeta) &= \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, a_1, \dots, a_{n+1}), \\ h_n(\Omega_{V,0}^\bullet, \zeta) &= \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, \mathcal{I}_f), \\ h_i(\Omega_{V,0}^\bullet, \zeta) &= \lambda, \quad \text{for } i = 1, \dots, n-1, \end{aligned}$$

where $0 \leq \lambda \leq \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, \mathcal{I}_f)$ is the integer:

$$\begin{aligned} \lambda &= \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, a_1, \dots, a_{n+1}) + \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/\left(\frac{df}{f}(\widehat{v}), a_1, \dots, a_{n+1}\right) \\ &\quad - \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(a_1, \dots, a_{n+1}), \end{aligned}$$

noticing that the tangency condition for \widehat{v} means that $\widehat{v}(df)$ is a multiple of f , so that $\frac{df}{f}(\widehat{v})$ is a holomorphic function on \mathbb{B} . As a consequence of these computations Gómez-Mont deduced the following expressions for the homological index (Theorem 1 in [61]).

Theorem 8. *Let $(V, 0)$ be an isolated hypersurface singularity (germ) of dimension n in \mathbb{C}^{n+1} , and let ζ be the restriction to V of a holomorphic vector field \widehat{v} on a neighborhood \mathbb{B} of 0 in \mathbb{C}^{n+1} , which has an isolated singularity at 0 and is tangent to V .*

(1) *If n is odd, then the homological index of ζ is:*

$$\text{Ind}_{\text{hom}}(\zeta, 0; V) = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, a_1, \dots, a_{n+1}) - \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, \mathcal{I}_f).$$

(2) *If n is even, then :*

$$\begin{aligned} \text{Ind}_{\text{hom}}(\zeta, 0; V) &= \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(a_1, \dots, a_{n+1}) \\ &\quad - \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/\left(\frac{df}{f}(\widehat{v}), a_1, \dots, a_{n+1}\right) + \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, \mathcal{I}_f). \end{aligned}$$

Notice that in the two formulas above, for even and odd dimensions, the following invariant appears:

$$\dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, a_1, \dots, a_{n+1}) .$$

This is reminiscent of the Tjurina number of holomorphic functions. The following name was coined in [33, p. 159]:

Definition 6. Let $\zeta = (a_1, \dots, a_{n+1})$ be a holomorphic vector field in \mathbb{C}^{n+1} be a holomorphic vector field in \mathbb{C}^{n+1} with an isolated zero at the origin. Then its **Tjurina (-Gómez-Mont)** number at 0 is:

$$\tau(\zeta) = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n+1},0}/(f, a_1, \dots, a_{n+1}) .$$

The next step in [61] for identifying the homological and the GSV indices for holomorphic vector fields on hypersurface germs is Step 2 above:

Lemma 1. *The difference*

$$\text{Ind}_{\text{hom}}(\zeta, 0; V) - \text{Ind}_{\text{GSV}}(\zeta, 0; V) = k ,$$

is a constant that depends only on V and not on the choice of vector field.

The proof of this lemma is an exercise using that both indices satisfy a certain “Law of conservation of the number”. This says roughly that if we perturb slightly a holomorphic vector field ζ at (V, p) , the new vector field ξ still has a singularity at p and new singularities arise at smooth points of V , then the index of ζ at p is the index of ξ at p plus the sum of the PH-indices of singularities of ξ at smooth points of V .

The final step for proving Theorem 7 consists in exhibiting a holomorphic vector on every hypersurface germ (V, p) for which the GSV and homological indices are both 0. When the dimension n of V is even, such an example in [61] is a Hamiltonian vector field as in Example 1. The case when n is odd is slightly more elaborate. In the two cases the vector field in question extends in an obvious way to the Milnor fibers with no singularities, and therefore the GSV index vanishes. Then Gómez-Mont uses Theorem 8 to prove that the homological index vanishes as well.

As a consequence we may state:

Corollary 1. *Let $(V, 0)$ be a hypersurface singularity in \mathbb{C}^{n+1} with an isolated singularity, and let $\zeta = (a_1, \dots, a_{n+1})$ be a holomorphic vector field tangent to V with an isolated singularity at 0. Then the homological index of ζ at 0, relative to V , equals the GSV index and if n is odd, this is the difference of their Tjurina numbers :*

$$\text{GSV}(\zeta, 0, V) = \tau(\zeta) - \tau(V) .$$

For n even this is:

$$\text{GSV}(\zeta, 0, V) = \text{PH}(\zeta, 0) + \tau(V) - \dim_{\mathbb{C}} \frac{\mathcal{O}_{\mathbb{C}^{n+1},0}}{\left(\frac{\zeta(f)}{f}, a_1, \dots, a_{n+1}\right)} .$$

Let us finish this section with two open questions:

Question 1: We know that the GSV index can be negative. We also know from [13] that given any isolated hypersurface germ $(V, 0)$ there are infinitely many holomorphic vector fields on V with an isolated singularity at 0, and those with the smallest homological (or GSV) index are dense. What is this index?

For example, if the singularity is quasi-homogeneous, this bound seems to be given by the index of the radial vector field, which equals the Euler characteristic of the Milnor fiber.

Question 2: Let X be a compact complex analytic variety with isolated singularities which are ICIS. Let ζ be a continuous vector field on X with isolated singularities at the singular points of X and perhaps at some other smooth points of X . Assume further that at each point in X_{sing} , ζ is holomorphic, and define its total homological index in the obvious way. Then it is an exercise to show that this is an invariant of X , independent of the vector field. What is this invariant? For instance, if X is a local complete intersection, this is the 0-degree Fulton class of X , by [86].

1.3 The virtual index

The GSV index inspired also a line of research in the vein of the Baum-Bott's residues, using Chern-Weil theory: this is the virtual index, defined by D. Lehmann, M. Soares and T. Suwa in [73] for vector fields on local complete intersections in complex manifolds. This was originally defined for holomorphic vector fields and it was later extended in [86] for continuous vector fields, which is useful for the study of characteristic classes on singular varieties (see Section 3). The virtual index is defined even if the vector field and the variety have non-isolated singularities; it attaches an index in \mathbb{Z} to each compact connected component of the singular set of the vector field. When the singularity is isolated, this coincides with the GSV index.

The starting point for defining the virtual index is recalling that for a continuous vector field v on a complex m -manifold X^m with an isolated singularity at a point, say 0, the local Poincaré-Hopf index can be regarded as being the localization at 0 of the top Chern class of X . This is explained in detail in [22] using Chern-Weil theory. In the isolated singularity case, which is the most relevant for this article, there is also the article [90] by T. Suwa, where he describes the localization in a particularly nice way, which is relevant for Section 4, where this is recalled (in 4.1).

Here we give an elementary intuitive explanation of the definition of the virtual index from the topological point of view. Let us explain.

The tangent bundle TX has Chern classes $c_1(X), \dots, c_m(X)$ in the cohomology ring $H^*(X; \mathbb{Z})$. If X is compact, cap product with the fundamental cycle $[X]$ maps these into homology classes in $H_*(X)$; these are called the homology Chern classes of X . The Gauss-Bonnet theorem tells us that the image in $H_0(X) \cong \mathbb{Z}$ of the top Chern class $c_m(X)$ is the Euler characteristic $\chi(X)$, which equals the total index of a vector field on X with isolated singularities, by the theorem of Poincaré-Hopf.

More generally, let v be a continuous vector field on X and assume v vanishes on some analytic subset S of X . We assume further that there is a regular neighborhood \mathcal{T} of S in X such that v is non-singular on $\mathcal{T} \setminus S$. Then V splits the Chern class $c_m(X)$ in two parts: one has support in the interior of \mathcal{T} and another in its exterior (see [22, Chapter 1] for details). By Alexander duality, these can be regarded as homology classes in $H_0(\mathcal{T})$ and $H_0(X \setminus \mathcal{T})$.

Definition 7. The contribution to $c_m(X)[X] = \chi(X)$ coming from $H_0(\mathcal{T}) \cong H_0(S)$ is the *localization of $c_m(X)[X]$ at S* .

If S is a point p , then the localization of $c_m(X)[X]$ at p is the corresponding local PH-index. We can always perturb v slightly so that it has isolated singularities in \mathcal{T} and the localization of $c_m(X)[X]$ at S is the total PH-index at S , see [22, Chapter 1].

We now want to adapt these ideas to vector fields on singular spaces. Let V be a subvariety of the complex manifold X^m defined by a regular section s of a holomorphic vector bundle E over X of rank k , with $1 \leq k < m$.

If s is transversal everywhere to the zero-section of E , then V is non-singular, the bundle $E|_V$ is isomorphic to the normal bundle of V and the Chern classes of V can be regarded as being those of the virtual bundle $TX|_V - E|_V$. Otherwise V is a singular local complete intersection and one defines its *virtual tangent bundle* by.

$$\tau V := TX|_V - E|_V .$$

The total Chern class of τV is defined in the usual way:

$$c_*(\tau V) = c_*(TX|_V) \cdot c_*(E|_V)^{-1} ,$$

where the term in the extreme right is the inverse of $c_*(E|_V)$ in the cohomology ring.

One gets the *total Fulton class* $c_*^{Fu}(V) = c_0^{Fu}(V) + \dots + c_n^{Fu}(V)$ by taking the cap product of $c_*(\tau V)$ with the fundamental cycle $[V]$. Fulton in [56] calls this the canonical class. The components $c_i^{Fu}(V) \in H_{2i}(V)$, for $i = 0, \dots, n$, are the *Fulton classes* of V .

We remark that in [56] Fulton actually defines these as classes of cycles in the Chow group and calls $c_*^{Fu}(V)$ the canonical class of V . His definition uses the Segre classes and it works for singular varieties in full generality (we refer to [22, 10, 18, 29] for thorough accounts, with different viewpoints, on Chern and their relations with singularity theory).

The idea now is to localize the zero-degree Fulton class (which corresponds to the top dimensional class in cohomology) using a tangent vector field, as we did on manifolds. The problem is that τV is a virtual bundle and the usual methods for localizing Chern classes via topology do not work.

Localization is done in [73, 20, 22] by means of Chern-Weil theory and it works equally well if we replace the isolated singularity p by a compact connected component S of the singular set. We refer to [22] for details.

Let us explain this construction in a special case which is particularly interesting and where it is much simpler. Assume that the regular section s that defines V can

be put in a flat family $\{s_t\}_{t \in \mathbb{C}}$ of holomorphic sections of E such that $s_0 = s$ and for $t \neq 0$ the section s_t is transversal to the zero section. Set $V_t := \{x \in X \mid s_t(x) = 0\}$.

We have a flat family of manifolds V_t degenerating to the special fiber V . Over each V_t one has the virtual bundle

$$TX|_{V_t} - E|_{V_t}.$$

For $t \neq 0$ the bundle $E|_{V_t}$ is isomorphic to the normal bundle of V_t , so the Chern classes of $TX|_{V_t} - E|_{V_t}$ are the Chern classes of the tangent bundle TV_t , and these can be regarded in homology. Denote these by $c_{*,t} \in H_*(V_t)$.

Now let $\tilde{\mathcal{T}}$ be a regular neighborhood of V in X , and let $\varepsilon_o > 0$ be small enough so that whenever $|t| < \varepsilon_o$, the manifold V_t is contained in $\tilde{\mathcal{T}}$. This exists by the compactness of V . Since $\tilde{\mathcal{T}}$ has V as a deformation retract, one has well defined homomorphisms:

$$H_*(V_t) \xrightarrow{\iota_*} H_*(\tilde{\mathcal{T}}) \xrightarrow{r_*} H_*(V),$$

induced by the inclusion of V_t in $\tilde{\mathcal{T}}$ and a retraction to V . Set $\mathcal{S} = r_* \circ \iota_*$. This is a special case of Verdier's specialization [91], and we know that the image $\mathcal{S}(c_{*,t}) \in H_*(V)$ are the Fulton classes of V . Let us summarize this discussion:

Proposition 2. *The Verdier specialization morphism \mathcal{S} carries the homology Chern classes of the general fiber V_t into the Fulton classes of the special fiber V .*

We now explain how to localize the top Fulton class of V at each connected component of its singular set V_{Sing} using a vector field. Intuitively this is easy: we use the vector field as we did on manifolds, to localize a contribution to the index at a component of the singular set of the vector field, and carry this contribution to the special fiber using the specialization morphism.

For simplicity we do it when the variety V , defined by a regular section of a holomorphic bundle E over X , only has isolated singularities, say p_1, \dots, p_r . The process in general is similar but a little more difficult technically.

For each p_i we may consider a conical neighborhood X_i of p_i in V and “replace it” by a local Milnor fiber $F_{i,t}$. We get a compact manifold \widehat{V}_t , and as $|t|$ tends to 0 this degenerates to V . As above, the homology Chern classes of V_t are mapped to the Fulton classes of V . Now, given a vector field on V with isolated singularities, we can take its restriction to the compact manifold with boundary $V \setminus (\bigcup_{i=1}^r X_i \cup)$, and extend it to all of \widehat{V}_t with isolated singularities on each local Milnor fiber $F_{i,t}$. This splits $c_n(\widehat{V}_t)[V_t]$ into a contribution coming from the smooth part $V \setminus (\bigcup_{i=1}^r X_i \cup)$, and a contribution coming from each $F_{i,t}$.

Definition 8. The virtual index of v at an isolated singularity p is the localization at p of the Fulton class $c_n(V) \in H_0(V) \cong \mathbb{Z}$ determined by v .

From the above construction we see that this invariant equals the total Poincaré-Hopf index of v at a local Milnor fiber when the singularity (V, p) can be put in a flat family as above. This condition is not necessary and the result is proved in general, either in [22, 20] via Chern-Weil theory, or in [29] topologically. We have:




Fig. 2 The virtual index is the localization at 0 of the PH index in a Milnor fiber.

Theorem 9. *The GSV and the virtual indices coincide for vector fields on ICIS germs.*

This theorem is discussed again in Section 4 from the viewpoint of Chern-Weil theory, following Suwa's viewpoint in [90] instead of the original one in [73] (or in [22]).

1.4 Radial index and the Milnor number

There are other indices of vector fields and 1-forms on singular varieties that play an important role in singularity theory. Two of these are the radial or Schwartz index, and the local Euler obstruction, which plays an important role for MacPherson's construction of Chern classes for singular varieties. This invariant has been extended to an index for vector fields and 1-forms on ICIS singularities. We refer to [22, 51, 29] for thorough accounts on this subject.

The radial (or Schwartz) index springs from the work of M. H. Schwartz in 1965, but she defined it only for a special class of vector fields, those obtained by radial extension. Her goal was defining Chern classes for singular varieties (see [22]). For instance, if p is an isolated singularity in an analytic space X , the only vector fields considered by M. H. Schwartz were the radial ones, *i.e.*, those transversal to each small sphere in the ambient space, centered at the singular point, and in that case the Schwartz index is 1. King and Trotman defined an index in general in [70], a 1991 preprint that unfortunately was not published till 2014, which essentially is the radial index. In the meantime, the same index was discovered simultaneously in [50] and [7]. This is defined in great generality, for real or complex analytic spaces, and we refer to [51] and [22, Ch. 2] for accounts on the subject. In the simplest case, where one considers an isolated singularity germ $(V, 0)$ and a continuous vector field on V with an isolated singularity at 0, this index is defined as:

$$\text{Rad}(v; 0, V) = 1 + d(v, v_{\text{rad}}),$$

where v_{rad} is a radial vector field on V (i.e. it is transversal to the intersection of V with every sufficiently small sphere, pointing outwards) and $d(v, v_{\text{rad}})$ is the difference between these two vector fields, as depicted in figure 3: this is the total index of a vector field ξ on a cylinder bounded by links $\mathbf{K}_{\varepsilon'} = V \cap \mathbb{S}_{\varepsilon'}$ and $\mathbf{K}_{\varepsilon} = V \cap \mathbb{S}_{\varepsilon}$ with $0 < \varepsilon' < \varepsilon$ and ξ restricts to v_{rad} on $\mathbf{K}_{\varepsilon'}$ and to v on \mathbf{K}_{ε} .




Fig. 3 The radial index

When V has non-isolated singularities, the radial index is defined similarly as above but there are two points of view. In [22, Ch. 2] one assigns an index to each connected component Σ of the singular set of a singular variety. For this one takes a regular neighborhood N_{Σ} of Σ with smooth boundary, and a smaller such neighborhood N'_{Σ} and use the cylinder bounded by these to define the difference $d(v, v_{\text{rad}})$, where v is the given vector field and v_{rad} is now the unit normal vector field on the boundary $\partial N'_{\Sigma}$, pointing outwards. In this case we define the radial index of v_{rad} to be the Euler characteristic $\chi(\Sigma)$ and the index of v is $\chi(S) + d(v, v_{\text{rad}})$.

In [50, 51] the definition is more refined as one must consider Whitney stratified vector fields with isolated singularities and one defines a radial index at each singularity of the vector field. The relation between the two definitions is as follows: the above radial index associated to Σ is the sum of the radial indices of the stratified vector field with isolated singularities contained in Σ . In either case the radial index at Σ actually is a class in $H_0(\Sigma; \mathbb{Z}) \cong \mathbb{Z}$, it can be regarded in $H_0(V; \mathbb{Z})$ via the inclusion, and it has the following important properties (see [22, 51]):

Theorem 10. 1. *Let $(V, 0)$ be an n -dimensional ICIS with Milnor number μ , and let v be a continuous vector field on V with an isolated singularity at 0 . Then their difference is the Milnor number of $(V, 0)$, independently of v :*

$$\mu = (-1)^n (\text{GSV}(v; 0, V) - \text{Rad}(v; 0, V)).$$

2. If V is a compact local complete intersection and v is a continuous vector field on V , singular at the singular set V_{sing} and at finitely many smooth points of V , then its total radial index (defined in the obvious way) is the Euler characteristic $\chi(V)$, which equals the 0-degree Schwartz-MacPherson class of V .

Remark 2 (Indices of 1-forms). The theory of indices of 1-forms on singular varieties is parallel to that for vector fields and it has been developed mostly by W. Ebeling and S. Gusein-Zade in a series of papers. We refer to their survey paper [51] for an account of this, and to [22, Ch. 9] for an introduction to the subject. Given an isolated singularity germ (V, p) in some \mathbb{C}^m and a holomorphic 1-form in a neighborhood of p , its restriction to V is a 1-form on V and its singularities are the points where its kernel is not transversal to V ; this includes p . There is a radial index for 1-forms with an isolated singularity at p , and if (V, p) is an ICIS, then there are also GSV and homological indices, and these coincide. The difference between these and the radial index is the local Milnor number. There is a local Euler obstruction as well.

Using 1-forms instead of vector fields has some advantages; perhaps a main one is that instead of imposing the tangency conditions one needs for vector fields, one needs a transversality condition for 1-forms. This allows, for instance, considering indices for collections of 1-forms. The indices we have envisaged so far morally correspond to the top Chern class, while the collections of 1-forms envisaged by Ebeling and Gusein-Zade correspond to different Chern numbers.

Remark 3 (Milnor classes). Classical Chern classes of complex manifolds can be defined as being the primary obstructions to constructing frames with appropriate indices at their singularities. These indices are natural extensions of the Poincaré-Hopf index. Similarly, for singular varieties equipped with a Whitney stratification, one may consider appropriate stratified frames and:

- One has a corresponding radial index for frames; the classes one gets are the Schwartz-MacPherson classes (see [22, Ch. 10]).
- If the variety is a local complete intersection in a manifold, one has a corresponding virtual index for frames, and the classes one gets are the Fulton classes (see [22, Ch. 10]).
- In the previous setting, the difference between the Fulton and the Schwartz-MacPherson classes are the Milnor classes of V .
- If V has only isolated singularities, the Milnor class in degree 0 is the sum of the local Milnor numbers (hence the name of Milnor classes). If V is a hypersurface with non-isolated singular set, the 0-degree Milnor class is Parusinski's generalized Milnor number.

Milnor classes arose first in the work of P. Aluffi [8, 9], though the actual name was coined later (see [29]).

2 Singular Holomorphic foliations, Baum-Bott residues and indices

A *regular* (smooth or non-singular) holomorphic foliation \mathcal{F} of dimension k on a complex manifold X is given by an open covering $\{U_\alpha\}$ of X , holomorphic submersions

$f_\alpha : U_\alpha \rightarrow \mathbb{C}^{n-k}$ and biholomorphisms

$$g_{\alpha\beta} : f_\beta(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n-k} \rightarrow f_\alpha(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n-k}$$

such that $f_\alpha = g_{\alpha\beta} \circ f_\beta$, whenever $U_\alpha \cap U_\beta \neq \emptyset$. That is (see figure 4), the diagramm

$$\begin{array}{ccc} U_\alpha \cap U_\beta & \xlongequal{\quad} & U_\alpha \cap U_\beta \\ \downarrow f_\beta & & \downarrow f_\alpha \\ f_\beta(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n-k} & \xrightarrow{g_{\alpha\beta}} & f_\alpha(U_\alpha \cap U_\beta) \subset \mathbb{C}^{n-k} \end{array}$$

commutes.




Fig. 4

If we set $f_\alpha = (f_\alpha^1, \dots, f_\alpha^{n-k})$, then the tangent of a fiber passing through $p \in U_\alpha$ is:

$$T_p(f_\alpha^{-1}(f_\alpha(p))) = \ker(df_\alpha)(p) = \{v \in T_p X; i_v(df_\alpha^1 \wedge \dots \wedge df_\alpha^{n-k})(p) = 0\}.$$

This means that the foliation is locally induced by the decomposable $(n-k)$ -form

$$df_\alpha^1 \wedge \dots \wedge df_\alpha^{n-k} \in H^0(U_\alpha, \Omega_{U_\alpha}^{n-k}),$$

and on $U_\alpha \cap U_\beta \neq \emptyset$ we have

$$df_\alpha^1 \wedge \cdots \wedge df_\alpha^{n-k} = \det(J(g_{\alpha\beta})) \cdot df_\beta^1 \wedge \cdots \wedge df_\beta^{n-k},$$

where $J(g_{\alpha\beta})$ is the jacobian matrix of the biholomorphic map $g_{\alpha\beta}$. Moreover, this gives us a holomorphic vector bundle of rank k defined by

$$T\mathcal{F} := \cup_\alpha \ker(df_\alpha) \subset TX$$

called the tangent bundle of \mathcal{F} . On the other hand, $T\mathcal{F}|_{U_\alpha}$ is generated by holomorphic vector fields $v_\alpha^1, \dots, v_\alpha^k$, such that $v_\alpha^i = \sum_{j=1}^k \Gamma_{ij}^k v_\alpha^j$, where Γ_{ij}^k are holomorphic functions on U_α . That is, the foliation is induced by the decomposable holomorphic k -vector field

$$v_\alpha^1 \wedge \cdots \wedge v_\alpha^k \in H^0(U_\alpha, \wedge^k TU_\alpha).$$

It follows from the Frobenius theorem that a holomorphic subvector bundle $\mathcal{E} \subset TX$ (called a distribution) of rank k is the tangent bundle of a foliation of dimension k if and only if $[\mathcal{E}, \mathcal{E}] \subset \mathcal{E}$. So, a holomorphic regular foliation induces an exact sequence of holomorphic vector bundles,

$$0 \longrightarrow T\mathcal{F} \xrightarrow{\phi} TX \xrightarrow{\pi} N\mathcal{F} \longrightarrow 0,$$

where the quotient bundle $N\mathcal{F}$ is called the normal bundle of \mathcal{F} . Notice that $\{\det(J(g_{\alpha\beta}))\}$ is a cocycle of $\wedge^{n-k} N\mathcal{F}$ the determinant of \mathcal{F} . Then, the foliation \mathcal{F} induces a nowhere vanishing holomorphic twisted $(n-k)$ -form,

$$\omega_{\mathcal{F}} \in H^0(X, \Omega_X^{n-k} \otimes \wedge^{n-k} N\mathcal{F}),$$

with values in the line bundle $\wedge^{n-k} N\mathcal{F}$, given by $\omega_{\mathcal{F}}|_{U_\alpha} = df_\alpha^1 \wedge \cdots \wedge df_\alpha^{n-k}$. Reciprocally, a nowhere vanishing twisted holomorphic $(n-k)$ -form $\omega \in H^0(X, \Omega_X^{n-k} \otimes L)$, with values in a line bundle L on X , induces a regular holomorphic foliation of dimension k , which is locally decomposable and integrable. That is, for all p there is an open neighborhood U and $\omega_1, \dots, \omega_{n-k} \in H^0(U, \Omega_U^{n-k})$, such that

$$\omega|_U = \omega_1 \wedge \cdots \wedge \omega_{n-k}$$

and

$$d\omega_i \wedge \omega_1 \wedge \cdots \wedge \omega_{n-k} = 0$$

for all $i = 1, \dots, n-k$. Then, by the Frobenius Theorem, there is a holomorphic map $f = (f^1, \dots, f^{n-k}) : U \rightarrow \mathbb{C}^{n-k}$, such that

$$\ker(\omega|_U) = \{v \in TU; i_v(\omega) = 0\} = \ker(df) = \ker(df^1 \wedge \cdots \wedge df^{n-k}) \subset TU.$$

Therefore, ω induces a holomorphic foliation \mathcal{F} of dimension k , with tangent bundle $T\mathcal{F} = \ker(\omega) \subset TX$. Moreover, for every $g \in H^0(X, \mathcal{O}_X^*)$ the form $\tilde{\omega} = g\omega$, yields the same foliation. In particular, if X is compact, then the foliation \mathcal{F} is represented by a point on the projective space

$$\mathbb{P}H^0(X, \Omega_X^{n-k} \otimes L).$$

In general, we can define.

Definition 9. A *regular Pfaff system* of dimension k on a *compact manifold* X is a point $[\omega] \in \mathbb{P}H^0(X, \Omega_X^{n-k} \otimes L)$, for some line bundle L , such that its singular set is $\text{Sing}(\omega) := \{\omega = 0\} = \emptyset$.

So, a regular Pfaff system ω is a regular foliation if and only if ω is *locally decomposable* and *integrable*.

2.1 Singular holomorphic foliations and Pfaffian systems

Let X be a complex manifold of dimension n . A singular holomorphic *distribution* \mathcal{F} , of dimension k , on X is given by an exact sequence

$$\mathcal{F} : 0 \longrightarrow T\mathcal{F} \xrightarrow{\phi} TX \xrightarrow{\pi} N\mathcal{F} \longrightarrow 0,$$

where $N\mathcal{F}$ is a torsion-free sheaf, called the *normal sheaf* of \mathcal{F} , and $T\mathcal{F}$ is a reflexive sheaf, of generic rank k , called the *tangent sheaf* of \mathcal{F} .

Definition 10. A *singular holomorphic foliation* of dimension k is a distribution \mathcal{F} , of dimension k , which is involutive, i.e. $[T\mathcal{F}, T\mathcal{F}] \subset T\mathcal{F}$. The *singular set* of the foliation \mathcal{F} is the singular set of its normal sheaf

$$\text{Sing}(\mathcal{F}) := \text{Sing}(N\mathcal{F}) = \{p \in X \mid (N\mathcal{F})_p \text{ is not locally free}\}.$$

Since $N\mathcal{F}$ is torsion-free, the singular set of \mathcal{F} has codimension ≥ 2 . The *singular scheme* of \mathcal{F} is defined as follows. Taking the maximal exterior power of the dual morphism $\phi^\vee : \Omega_X^1 \rightarrow T\mathcal{F}^*$ we obtain a morphism

$$\wedge^k \phi^\vee : \Omega_X^k \rightarrow \det(T\mathcal{F})^*$$

and an induced morphism $\Omega_X^k \otimes \det(T\mathcal{F}) \rightarrow \mathcal{O}_X$ whose image is an ideal sheaf \mathcal{I}_Z of a subscheme $Z \subset X$. Moreover, such a morphism induces a global section

$$v_{\mathcal{F}} \in H^0(X, \wedge^k TX \otimes \det(T\mathcal{F})^*)$$

which is called the *Pfaff field* induced by \mathcal{F} . This means that on a neighborhood U_p of a point p such that $T\mathcal{F}$ is locally free and generated by vector fields v_1, \dots, v_k , then $v_{\mathcal{F}}|_{U_p} = v_1 \wedge \dots \wedge v_k$ and $[v_i, v_j] = \sum_{k=1}^k \Gamma_{ij}^k v_k$, for all $i, j, k = 1, \dots, n$, where $\Gamma_{ij}^k \in \mathcal{O}_{U_p}$ and $v_i, v_j \in H^0(U_p, TU_p)$. Therefore, a singular holomorphic foliation \mathcal{F} of dimension k , induces a regular foliation \mathcal{F}^0 of dimension k on $X^0 := X - \text{Sing}(\mathcal{F})$.

The line bundle $K\mathcal{F} := \det(T\mathcal{F})^*$ is called the *canonical bundle* of \mathcal{F} . So, By the isomorphism $\wedge^k TX \simeq \Omega_X^{n-k} \otimes \det(TX)$ and the fact that $\det(N\mathcal{F}) \simeq \det(TX) \otimes K\mathcal{F}$ we conclude that the isomorphism

$$\wedge^k TX \otimes K\mathcal{F} \simeq \Omega_X^{n-k} \otimes \det(TX) \otimes K\mathcal{F} \simeq \Omega_X^{n-k} \otimes \det(N\mathcal{F})$$

says that the section $\nu_{\mathcal{F}}$ corresponds to a twisted $(n-k)$ -form

$$\omega_{\mathcal{F}} \in H^0(X, \Omega_X^{n-k} \otimes \det(N\mathcal{F}))$$

with values on $\det(N\mathcal{F})$ which yields a *Pfaff system*, of dimension $(n-k)$, associated with the corresponding regular foliation \mathcal{F}^0 on the regular part $X^0 = X - \text{Sing}(\mathcal{F})$.

So, in a neighborhood U_p of a point p where $N\mathcal{F}^*$ is locally free, we have

$$\omega_{\mathcal{F}}|_{U_p} = \omega_1 \wedge \cdots \wedge \omega_{n-k},$$

where $\omega_i \in H^0(U_p, \Omega_{U_p}^1)$ satisfies the integrability condition

$$d\omega_j \wedge \omega_1 \wedge \cdots \wedge \omega_{n-k} = 0$$

for all $j = 1, \dots, n-k$. Therefore,

$$\text{Sing}(\mathcal{F}) = \{\nu_{\mathcal{F}} = 0\} = \{\omega_{\mathcal{F}} = 0\}.$$

Definition 11. Let X be a compact complex manifold of dimension n . A *Pfaff system* of dimension k on X is an element

$$[\omega] \in \mathbb{P}H^0(X, \Omega_X^{n-k} \otimes \mathcal{L}),$$

where \mathcal{L} is a holomorphic line bundle on X , such that its singular set $\text{Sing}(\omega) := \{\omega = 0\}$ has codimension ≥ 2 .

Remark 4. Note that a holomorphic foliation of dimension k induces a Pfaff system of codimension $n-k$ that is generically decomposable. However, in general, a Pfaff system does not need to be generically decomposable. An example of a non-decomposable Pfaffian system is given by the symplectic 2-form in \mathbb{C}^{2m} . In canonical coordinates $(q_1, p_1, \dots, q_m, p_m) \in \mathbb{C}^{2m}$, this symplectic form is expressed as

$$\omega = \sum_{i=1}^m dq_i \wedge dp_i.$$

Example 4. A holomorphic foliation \mathcal{F} on \mathbb{P}^n of dimension k corresponds to a twisted $(n-k)$ -form $\omega \in H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^{n-k}(d+n-k+1))$, where $d \geq 0$ is called the *degree* of \mathcal{F} . We can interpret ω as a polynomial differential form

$$\omega = \sum_{1 \leq i_1 < \cdots < i_{n-k} \leq n} A_{i_1 \cdots i_{n-k}} dx_{i_1} \wedge \cdots \wedge dx_{i_{n-k}}$$

where the $A_{i_1 \cdots i_p}$ are homogeneous of degree $d+1$, and $\iota_R \omega = 0$, where

$$R = x_0 \frac{\partial}{\partial x_0} + \cdots + x_n \frac{\partial}{\partial x_n}$$

is the radial vector field. The local decomposability translates to the Plücker conditions

$$(i_v \omega) \wedge \omega = 0 \quad \text{for every } v \in \wedge^{n-k-1}(\mathbb{C}^{n+1}),$$

and the integrability condition becomes

$$(i_v \omega) \wedge d\omega = 0 \quad \text{for every } v \in \wedge^{n-k-1}(\mathbb{C}^{n+1}).$$

For more details, we refer the reader to [47, Section 1.3].

Example 5. Let $\psi : X \rightarrow Y$ be a holomorphic fibration, where Y is a complex manifold of dimension $n - k > 0$. Then ψ induces a holomorphic foliation \mathcal{F} on X of dimension k , whose tangent sheaf is the relative tangent sheaf of ψ , and whose singular set corresponds to the singularities of the fibration.

Definition 12. Let $\omega \in H^0(X, \Omega_X^{n-k} \otimes \mathcal{L})$ be a Pfaff system of dimension k on a complex manifold X . We say that an analytic subvariety $V \subset X$ is *invariant* by ω if $i^* \omega \equiv 0$, where $i : V \hookrightarrow X$ is the inclusion map. If \mathcal{F} is a holomorphic foliation of dimension k on X , then V is invariant by \mathcal{F} if it is invariant by its corresponding Pfaff system

$$\omega_{\mathcal{F}} \in H^0(X, \Omega_X^{n-k} \otimes \det(N\mathcal{F})).$$

Example 6. Consider the foliation \mathcal{F} in \mathbb{C}^2 induced by the holomorphic 1-form $\omega = x(5y^2 - 9) dx - y(5x^2 - 9) dy$. This foliation has 5 singular points given by the singular points of the invariant algebraic curve $C = \{(x^2 - y^2) \cdot (x^2 + 4y^2 - 9) = 0\}$. See figure 5.




Fig. 5

Example 7. In this example, we examine the topology, via the Milnor fibration, of a foliation on \mathbb{C}^2 induced by an exact holomorphic 1-form $\omega = df$, i.e., when the

foliation admits a holomorphic first integral. Suppose that the foliation is induced by the 1-form $d(x^3 - y^2)$ with an isolated singularity at 0. This foliation is tangent to the fibers of the polynomial function $f : \mathbb{C}^2 \rightarrow \mathbb{C}$, given by $f(x, y) = x^3 - y^2$, and it has the invariant (separatrix) curve $f^{-1}(0) := C$ with isolated singularity at 0. Pick the point $0 \in C$ and a small enough ϵ -ball $B_\epsilon(0)$ in \mathbb{C}^2 centered at the origin 0, with boundary the 3-sphere $S_\epsilon(x)$. Then by Milnor [77], $B_\epsilon(0) \cap C$ is homeomorphic to the cone on the real link $S_\epsilon(0) \cap C$. In this case, the link $S_\epsilon(0) \cap C \simeq \mathbb{S}^1$ is the trefoil knot. If we take a point $x_1 \in D_\delta(0) - \{0\} \subset \mathbb{C}$, with $|x_1| \ll \delta \ll \epsilon \leq 1$, then on the smooth curve $f^{-1}(x_1)$ we have also that $S_\epsilon(0) \cap f^{-1}(x_1)$ is the trefoil knot, while the leaves of the foliation contained within the ball, which are the Milnor fibers, have the homotopy of a bouquet of 2 circles, $\simeq \mathbb{S}^1 \vee \mathbb{S}^1$. See figure 6.

2.2 Baum–Bott residues

In this section we recall the Baum–Bott residue theorem for singular foliations on complex manifolds. A starting point is Bott’s vanishing theorem presented by R. Bott in the 1970 ICM in Nice (see [16]), where he discussed the state of the art in the problem of “foliating a manifold”. As he explains, this can be viewed in two steps:

Step 1: Given a (smooth or complex) n -manifold, construct a k -dimensional sub-bundle E of the tangent bundle TM ; and then,

Step 2: Given a subbundle E of TM as above, can it be deformed to an integrable one?

In the complex case we want E to be holomorphic.

Bott then gives a beautiful answer, which in the complex case can be stated as follows:

Theorem 11 (Bott’s vanishing theorem). *Let M be a compact complex analytic n -manifold which admits an integrable holomorphic subbundle E . Let $Q = TM/E$ be the quotient bundle, that one calls the normal bundle. Then the Chern ring TM/E , generated by the real Chern classes of TM/E , vanishes in dimensions greater than $2 \times \dim_{\mathbb{C}}(TM/E)$. That is,*

$$\text{Chern}^r(TM/E) = 0 \quad \forall r > \dim_{\mathbb{R}}(TM/E).$$

This provides topological obstructions for the integrability of a distribution on a manifold. It is worth saying that there is also previous work by Bott (e.g. [15]) where he relates the behavior of a holomorphic vector field at its zeroes to the characteristic numbers of complex manifolds and holomorphic bundles.

As an example (from [16]), consider a compact complex n -manifold M with vanishing Euler characteristic. This means that its top Chern class $c_n(M)$ vanishes as well, and also that by the theorem of Poincaré-Hopf, we can construct on M a nowhere vanishing C^∞ vector field v . This spans a smooth trivial 1-dimensional complex sub-bundle E of TM and one has a C^∞ ssplitting $TM \cong E \oplus Q$ where

$Q = TM/E$ is the normal bundle. Hence, since E is a trivial bundle, all the Chern numbers of M are degree n products of the Chern classes of Q , evaluated on the fundamental cycle $[M]$.

Now suppose that the vector field V actually is holomorphic, so it is integrable. Then one has a 1-dimensional holomorphic foliation \mathcal{E} with tangent bundle E . Then, by Bott's theorem, all degree n products of Chern classes of Q vanish. Hence, by the above observations, all the Chern numbers of M vanish, so we have:

Corollary 2. *If a compact complex manifold admits a nowhere vanishing holomorphic vector field, then all its Chern numbers are 0.*

Bott's theorem 11 was later extended by Baum-Bott to higher dimensional foliations, allowing also singularities. If there are no singularities, then the theorem states that the ring $\text{Chern}^r(Q)$, generated by the real Chern classes of the normal bundle, vanishes in dimensions greater than the real codimension of the foliation. In the presence of singularities, the theorem says that each such class, regarded in homology via Alexander duality, localizes as a sum of residues, which are homology classes of the singular set. More precisely:

Theorem 12 (Baum-Bott). [12] *Let \mathcal{F} be a holomorphic foliation of dimension k on a complex manifold X , and let φ be a homogeneous symmetric polynomial of degree d satisfying $n - k < d \leq n$. Let Z be a compact connected component of the singular set $\text{Sing}(\mathcal{F})$. Then, there exists a homology class $\text{Res}_\varphi(\mathcal{F}, Z) \in H_{2(n-d)}(Z; \mathbb{C})$ such that:*

- (i) $\text{Res}_\varphi(\mathcal{F}, Z)$ depends only on φ and on the local behavior of the leaves of \mathcal{F} near Z .
- (ii) Suppose that X is compact, and define $\text{Res}(\varphi, \mathcal{F}, Z) := \alpha_* \text{Res}_\varphi(\mathcal{F}, Z)$, where α_* is the composition of the maps

$$H_{2(n-d)}(Z; \mathbb{C}) \xrightarrow{i_*} H_{2(n-d)}(X; \mathbb{C}) \quad \text{and} \quad H_{2(n-d)}(X; \mathbb{C}) \xrightarrow{P} H^{2d}(X; \mathbb{C}),$$

where i_* is the induced map from the inclusion $i : Z \rightarrow X$, and P is the Poincaré duality map. Then

$$\varphi(N_{\mathcal{F}}) = \sum_Z \text{Res}(\varphi, \mathcal{F}, Z).$$

If the foliation has dimension one and isolated singularities, the residues can be expressed in terms of Grothendieck residues (see [11]). More precisely, in this case, for each $p \in \text{Sing}(\mathcal{F})$, an isolated singularity of \mathcal{F} , we have

$$\text{Res}(\varphi, \mathcal{F}, p) = \text{Res}_p \left[\begin{array}{c} \varphi(Jv) \\ v_1, \dots, v_n \end{array} \right], \quad (6)$$

where $v = (v_1, \dots, v_n)$ is a germ of a holomorphic vector field on (U, p) , serving as a local representative of \mathcal{F} , Jv is its Jacobian matrix, and

$$\text{Res}_p \left[\begin{array}{c} \varphi(Jv) \\ v_1, \dots, v_n \end{array} \right] = \left(\frac{1}{2\pi\sqrt{-1}} \right)^n \int_\Gamma \frac{\varphi(Jv)}{v_1, \dots, v_n},$$




Fig. 6

denotes the Grothendieck residue of $\varphi(J\nu)$ with respect to ν_1, \dots, ν_n at p , and Γ is the cycle $\{x \in U; |\nu_i(x)| = \epsilon_i; i = 1, \dots, n\}$.

In general, it is a difficult task to compute the Baum–Bott residues for foliations of higher dimension. In [92], Vishik proved Baum–Bott’s Theorem and provided a method to compute the residues, assuming that the foliation has a locally free tangent sheaf. In [17], F. Bracci and T. Suwa studied the behavior of Baum–Bott residues under smooth deformations.

Let us recall how to compute such residues in a particular case. Let $\text{Sing}_{n-k+1}(\mathcal{F}) \subset \text{Sing}(\mathcal{F})$ be the subset of components of pure dimension $n - k + 1$. Consider a symmetric homogeneous polynomial φ of degree $n - k + 1$. Let $Z \subset \text{Sing}_{n-k+1}(\mathcal{F})$ be an irreducible component. Take a generic point $p \in Z$ such that p is a point where Z is smooth and disjoint from the other irreducible components. Now, consider B_p a ball centered at p , of dimension $n - k + 1$ sufficiently small and transversal to Z in p . In [12], under a generic condition (Kupka condition [80]), Baum and Bott proved that

$$\text{Res}(\mathcal{F}, \varphi; Z) = \text{Res}_\varphi(\mathcal{F}|_{B_p}; p)[Z],$$

where $\text{Res}_\varphi(\mathcal{F}|_{B_p}; p)$ represents the Grothendieck residue at p of the one dimensional foliation $\mathcal{F}|_{B_p}$ on B_p and $[Z]$ denotes the fundamental cycle associated to Z .

In [42], the authors show that Baum–Bott’s generic hypothesis is not necessary, and we demonstrate that the above formula always holds as long as the singular set of the foliation has dimension less than or equal to $n - k + 1$. More recently, in [67] Kaufmann, Lärkäng, and Wulcan showed that the Baum–Bott residues can be expressed in terms of residue currents.

3 Indices for 1-dimensional holomorphic foliations on surfaces

In his remarkable paper [23], M. Brunella studied various invariants of holomorphic 1-dimensional foliations on complex surfaces that somehow spring from the Baum-Bott theory of residues. He then related all these invariants to establish interesting geometric properties of foliations. These results have been extended to higher dimensions and to other settings by various authors. This is the subject that we study in this section.

3.1 Baum-Bott-Brunella indices

In this subsection we look at the Milnor number of a foliation, already studied earlier (see 4), as well as at the Camacho-Sad index [32], the variation [68], defined by Khanedani-Suwa, and an invariant that Brunella called Baum-Bott invariant.

Let \mathcal{F} be a holomorphic foliation with isolated singularities on a complex surface X and let p be singular point of \mathcal{F} . From the previous section we know that there are two Baum-Bott residues associated to \mathcal{F} at p ; these correspond to the Chern numbers c_2 and c_1^2 . The one corresponding to c_2 , or to the symmetric function given by the determinant, has as residue at p the local Poincaré-Hopf index of a vector field generating \mathcal{F} locally. The local invariant we get is the local multiplicity, or Milnor number $\mu(\mathcal{F}, p)$ of the foliation defined in 4, which is a topological invariant. Following [23] we denote this invariant by $PH(\mathcal{F}, p) = \mu(\mathcal{F}, p)$.

The second Baum-Bott residue corresponds to the symmetric function given by the trace, or actually to the Chern number $c_1^2(N_{\mathcal{F}})$, where $N_{\mathcal{F}} \in H^1(X, \mathcal{O}^*)$ is the normal bundle of \mathcal{F} ; this bundle is defined even in the presence of singularities since \mathcal{F} is determined by a morphism of a line bundle into TM .

Following [23] we denote this by $BB(\mathcal{F}, p)$ and call it **the Baum-Bott invariant of \mathcal{F} at p** . This can be studied *à la* Godbillon-Vey. Let (z, w) be local coordinates centered at p and let $\omega = F(z, w)dw - G(z, w)dz$ be a holomorphic 1-form that generates \mathcal{F} locally. Let β be a complex valued smooth 1-form of type $(1, 0)$ on a punctured neighborhood $U^* = U \setminus \{p\}$ such that

$$d\omega = \beta \wedge \omega.$$

For instance one may take:

$$\beta = \frac{\frac{\partial F}{\partial z} + \frac{\partial G}{\partial w}}{|F|^2 + |G|^2} (\bar{F} dz + \bar{G} dw).$$

Then one has (see [23]):

$$BB(\mathcal{F}, p) = \frac{1}{(2\pi i)^2} \int_{\mathbb{S}_x^3} \beta \wedge d\beta,$$

where \mathbb{S}_ε^3 is the boundary sphere of a small ball around p with its natural orientation.

If the surface X is compact then one has a global result in the vein of the Poincaré-Hopf index theorem:

$$c_1^2(N_{\mathcal{F}}) = \sum_{p \in \text{sing}\mathcal{F}} \text{BB}(\mathcal{F}, p). \quad (7)$$

Now let S be a separatrix of \mathcal{F} at p . That is, a local holomorphic curve containing p which is invariant by \mathcal{F} . It is not assumed that this curve is neither smooth nor irreducible. The following invariant is defined in [68]:

Definition 13. The variation of \mathcal{F} at p relative to S is:

$$\text{Var}(\mathcal{F}, S, p) = \frac{1}{2\pi i} \int_{S \cap \mathbb{S}_\varepsilon} \beta,$$

with β as above.

This invariant has the following important global property.

Proposition 3 (Khanedani-Suwa). *Suppose S is now a compact holomorphic curve in X , invariant by the foliation, then:*

$$\sum_{p \in \text{Sing}(\mathcal{F}) \cap S} \text{Var}(\mathcal{F}, S, p) = c_1(N_{\mathcal{F}}) \cdot S.$$

Now we come to the Camacho-Sad index. This plays a key role for proving in [32] the separatrix theorem: every germ of a holomorphic vector field in \mathbb{C}^2 with an isolated singularity at the origin has at least one separatrix S . Yet, this index is not actually defined in \mathbb{C}^2 but up in a “desingularization” of the vector field, where one assumes that there is a smooth divisor C , invariant under the induced foliation, with isolated singularities. One attaches a CS-index to each such singularity, which actually is a localization of the Chern class of the normal bundle of C .

Let S be again a separatrix in X for the foliation \mathcal{F} and $p \in S$ a singularity of the foliation. Let ω be a local 1-form on X generating \mathcal{F} locally and f a local function that defines S near p , and we assume it is reduced. Then we know from [88] that there are functions g, ξ and a 1-form η on a neighborhood of p such that

$$g\omega = \xi df + f\eta$$

with $k \neq 0$ on $S^* = S \setminus \{p\}$. Notice that on S we have $g\omega = \xi df$ and $g \neq 0$ on S^* .

Definition 14. The Camacho-Sad index of \mathcal{F} at p relative to S is:

$$\text{CS}(\mathcal{F}, S, p) = -\frac{1}{2\pi i} \int_{S \cap \mathbb{S}_\varepsilon^3} \frac{1}{\xi} \eta.$$

One has:

Proposition 4.

$$\sum_{p \in \text{Sing}(\mathcal{F}) \cap S} \text{CS}(\mathcal{F}, S, p) = S \cdot S = c_1(N_S)[S].$$

Notice that all these formulas have a similar flavor, involving the geometry of the foliation to localize the Chern class of some normal bundle. This contrasts with the GSV index, which (by Proposition 9) is the localization of a Chern class of the virtual tangent bundle.

3.2 The GSV index for 1-dimensional holomorphic foliations

Given a foliation \mathcal{F} on a complex surface X as above, with a singularity at p and a separatrix S through p , S is locally generated by a holomorphic vector field v . Hence S can be regarded itself as being a hypersurface in X with an isolated singularity at p (in fact p can be a regular point of S), equipped with a vector field v . Thus one has the local GSV index of v in S at p , which we take as being its Poincaré-Hopf index in a Milnor fiber of a local defining function f for S (we take this as its definition even if S has several branches).

Brunella in [23] used the foliation to introduce an invariant $Z(\mathcal{F}, S, p)$ in the setting we now envisage, and proved that it coincides with the GSV index. Brunella then established surprising deep relationships between the Baum-Bott, Camacho-Sad, the variation and the GSV indices.

Let X be a compact complex surface, and let \mathcal{F} denote a one-dimensional holomorphic foliation on X . Let C be a reduced curve on X and x a point in C . When C is invariant under \mathcal{F} , we say that \mathcal{F} is *logarithmic along* C . Let $f = 0$ be a local equation for C in a neighborhood U_α of x , and let ω_α be a holomorphic 1-form that induces the foliation \mathcal{F} on U_α . Since \mathcal{F} is logarithmic along C , it follows from [81, 88] that there exist holomorphic functions g and ξ defined in a neighborhood of x that do not both vanish identically on C , such that

$$g \frac{\omega_\alpha}{f} = \xi \frac{df}{f} + \eta, \quad (8)$$

where η is an appropriate holomorphic 1-form.

Definition 15 (Brunella [23]). Let \mathcal{F} be a one-dimensional holomorphic foliation on a compact complex surface X that is logarithmic along a reduced curve $C \subset X$. Given $x \in C$, we define

$$Z(\mathcal{F}, C, x) = \sum_i \text{ord}_x \left(\frac{\xi}{g} |_{C_i} \right),$$

where $C_i \subset C$ are the irreducible components of C at x , and $\text{ord}_x \left(\frac{\xi}{g} |_{C_i} \right)$ denotes the order of vanishing of $\frac{\xi}{g} |_{C_i}$ at x .

First, we will see that Brunella's Z index coincides with the GSV index.

Proposition 5. [23, Page 532] *Let $p \in \text{Sing}(\mathcal{F}) \cap C$. Then*

$$Z(\mathcal{F}, C, p) = \text{GSV}(\mathcal{F}, C, p).$$

Proof. Let f be a holomorphic function on a neighborhood U of p and defining $C|_U = \{f = 0\}$. So, if ω induces \mathcal{F} on U we have

$$\omega = \frac{\xi}{g} df + \frac{f}{g} \eta.$$

Let $J : \Omega_U^1 \rightarrow TU$ denote the isomorphism from 1-forms to vector fields in a neighborhood U of p , defined by

$$\omega = P dx + Q dy \mapsto J(\omega) = Q \frac{\partial}{\partial x} - P \frac{\partial}{\partial y}.$$

The holomorphic vector field $J(\omega)$ defines the foliation \mathcal{F} and can be expressed as the sum of two meromorphic vector fields:

$$J(\omega) = \frac{\xi}{g} J(df) + \frac{f}{g} J(\eta) = v_1 + v_2,$$

with $v_1 = \frac{\xi}{g} J(df)$ and $v_2 = \frac{f}{g} J(\eta)$. Now, observe that

$$J(\omega)|_C = \left(\frac{\xi}{g} J(df) + \frac{f}{g} J(\eta) \right) |_C = \frac{\xi}{g} J(df)|_C = v_1|_C$$

and v_1 is tangent to $C_\epsilon = \{f = \epsilon\}$, for a small ϵ . The restriction $v_1|_{C_\epsilon}$ has poles at the points of $C_\epsilon \cap \{g = 0\}$ and zeros at the points of $C_\epsilon \cap \{\xi = 0\}$. Since the GSV index in C_ϵ is the Poincaré-Hopf index of $v_1|_{C_\epsilon}$, it is the difference between the number of zeroes and the number of poles of $v_1|_{C_\epsilon}$. Therefore,

$$\text{GSV}(\mathcal{F}, C, p) = \frac{1}{2\pi i} \int_{\partial C_\epsilon} \frac{d\xi}{\xi} - \int_{\partial C_\epsilon} \frac{dg}{g} = \frac{1}{2\pi i} \int_{\partial C} \frac{g}{\xi} d \left(\frac{\xi}{g} \right) = Z(\mathcal{F}, C, p).$$

This proves the proposition.

One has:

Proposition 6. [23, Proposition 5] *If C is any separatrix at p , then*

$$\text{Var}(\mathcal{F}, C, p) = \text{GSV}(\mathcal{F}, C, p) + \text{CS}(\mathcal{F}, C, p),$$

where $\text{Var}(\mathcal{F}, V, p)$ denotes the variational Khanedani-Suwa index of \mathcal{F} along V at p [68] and $\text{CS}(\mathcal{F}, C, p)$ is the Camacho-Sad index [32].

Remark 5. Recently, in [41], the authors generalized the variational index for currents invariant under foliations, with applications to the study of transcendental leaves of foliations and their accumulation on the singular set.

Theorem 13 (Brunella [23]). *Let \mathcal{F} be a one-dimensional holomorphic foliation on a compact complex surface X , logarithmic along a reduced curve $C \subset X$. Then*

$$\sum_{x \in \text{Sing}(\mathcal{F}) \cap C} \text{GSV}(\mathcal{F}, C, x) = N\mathcal{F} \cdot C - C \cdot C,$$

The GSV index is an integer and we already know that it can be negative. In fact, consider Example 2 and the foliation in \mathbb{C}^2 given by the homogeneous vector field $v = w^k \frac{\partial}{\partial z} - z^k \frac{\partial}{\partial w}$, whose leaves are the fibers of the map $f = z^{k+1} + w^{k+1}$. Then $S = f^{-1}(0)$ is a separatrix (with $k+1$ branches) and its GSV index is $1 - k^2$, which is negative whenever $k > 1$. Notice that this foliation has infinitely many separatrices.

Marco Brunella remarked that the non-negativity of the GSV index is the obstruction to the positive solution to ‘‘Poincaré problem’’, as we explain below in Section 4. The following results from [23] point in that direction and they are interesting on their own.

Recall that a separatrix S of \mathcal{F} at p is nondicritical if there is a sequence of blow-ups based at p , $\pi : \tilde{X} \rightarrow X$, such that π is a resolution of S , $\tilde{S} = \pi^{-1}(S)$ has only normal crossing singularities, and the divisor $\pi^{-1}(p)$ is \tilde{F} -invariant, where \tilde{F} is the induced foliation on \tilde{X} (see [30]). Then (see [23, Propositions 6 and 7]):

Theorem 14. *If S is a nondicritical separatrix at p then:*

$$\text{GSV}(\mathcal{F}, S, p) \geq 0.$$

If S further is a generalized curve (i.e. there are no saddle-nodes in its resolution) and S is maximal, union of all separatrices at p , then:

$$\text{GSV}(\mathcal{F}, S, p) = 0.$$

In [36], Cavalier and D. Lehmann proved that the converse is also true, i.e., if

$$\text{GSV}(\mathcal{F}, S, p) = 0,$$

then the foliation \mathcal{F} is a generalized curve at p , and S is the maximal separatrix.

4 Indices in higher dimensions and the Poincaré problem

As noted before, indices and residues play a significant role for the study of the famous Poincaré problem in dimensions 2 and higher. In this section we review some of these. The literature is vast and we will focus only on some of the works

most related with this article. Let us start by saying a few words about the problem itself.

In [79] H. Poincaré asked whether it is possible to decide if an algebraic differential equation in two variables is algebraically integrable, and he observed that in order to answer this question it is sufficient to know whether one can bound the degree of the generic leaves. He posed a question that in modern language can be stated as follows. Let \mathcal{F} be a holomorphic foliation in $\mathbb{P}_{\mathbb{C}}^2$ and C an irreducible curve which is invariant by \mathcal{F} : Is it possible to bound the degree of C in terms of the degree of \mathcal{F} ?

This problem has been studied by many authors in more generality. One may ask, for instance, if given a foliation \mathcal{F} on $\mathbb{P}_{\mathbb{C}}^n$ of degree d and a hypersurface V in $\mathbb{P}_{\mathbb{C}}^n$ invariant by \mathcal{F} , is it possible to bound the degree of V by a number which depends only on d ?

The answer in general is obviously negative, even for $n = 2$. For instance the foliation defined by the form $pwdz - qzdw$ has degree 1 and the invariant curve $z^p - w^q = 0$ has degree equal to the maximum of p, q . There are many more easy examples. Yet, there are many positive examples, and as explained in [26] the philosophy is that the failure of having a uniform bound is due to the existence of "bad singularities" of V or of \mathcal{F} .

As mentioned before, for $n = 2$, Brunella in [23] established a connection between the GSV-index and both the Khanedani-Suwa variational index [68] and the Camacho-Sad index [32], and he demonstrated that the non-negativity of the GSV-index acts as an obstruction to providing an affirmative answer to the *Poincaré problem* on compact complex surfaces.

In fact, more generally, the non-negativity of the GSV-index provides an obstruction to solving the generalized Poincaré problem for Pfaff systems on projective complex spaces as explained below.

In [44], a GSV-type index for varieties was introduced, in the spirit of Brunella, for varieties invariant under holomorphic Pfaff systems (which may not be locally decomposable) on projective manifolds.

Let U be a germ of an n -dimensional complex manifold. Let D be an analytic reduced hypersurface on U and consider its decomposition into irreducible components

$$D = D_1 \cup \cdots \cup D_{n-k},$$

and suppose that the analytic subvariety $V = D_1 \cap \cdots \cap D_{n-k}$ has pure codimension k . We assume that

$$V = \{z \in U : f_1(z) = \cdots = f_{n-k}(z) = 0\},$$

with $f_1, \dots, f_{n-k} \in \mathcal{O}_U$ and for each $i \in \{1, \dots, k\}$,

$$D_i = \{z \in U : f_i(z) = 0\}.$$

Since V is a reduced variety, then the k -form $df_1 \wedge \cdots \wedge df_{n-k}$ is not identically zero on each irreducible component of V . Following Aleksandrov [5], we denote by $\Omega_U^q(\hat{D}_i)$, for $q \geq 1$, the \mathcal{O}_U -module of meromorphic differential q -forms with

simple poles on the

$$\hat{D}_i = D_1 \cup \cdots \cup D_{i-1} \cup D_{i+1} \cup \cdots \cup D_{n-k},$$

for each $i = 1, \dots, n - k$. We have the following Aleksandrov's Decomposition Theorem, which is a generalization of Saito's residues [81].

Theorem 15 (Aleksandrov [5]). *Let $\omega \in \Omega_U^q(D)$ be a meromorphic q -form with simple poles on D . If for each $j = 1, \dots, k$,*

$$df_j \wedge \omega \in \sum_{i=1}^k \Omega_U^{q+1}(\hat{D}_i)$$

then, there exist a holomorphic function g , which is not identically zero on every irreducible component of V , a holomorphic $(q - k)$ -form $\xi \in \Omega_U^{q-k}$ and a meromorphic q -form $\eta \in \sum_{i=1}^k \Omega_U^q(\hat{D}_i)$ such that the following decomposition holds

$$g\omega = \frac{df_1}{f_1} \wedge \cdots \wedge \frac{df_k}{f_k} \wedge \xi + \eta. \quad (9)$$

We can apply this decomposition in our context.

Proposition 7. *Let $\omega \in H^0(X, \Omega_X^{n-k} \otimes \mathcal{L})$ be a Pfaff system of dimension k on a complex manifold X , and $V \subset X$ a reduced local complete intersection subvariety of dimension $n - k$ which is invariant by ω . Then for all local representations $\omega_\alpha = \omega|_{U_\alpha}$ of ω , and all local expressions of V in U_α*

$$V \cap U_\alpha = \{z \in U_\alpha : f_{\alpha,1}(z) = \cdots = f_{\alpha,n-k}(z) = 0\},$$

there are holomorphic functions $g_\alpha, \xi_\alpha \in \mathcal{O}(U_\alpha)$ such that

$$g_\alpha \omega_\alpha = \xi_\alpha df_{\alpha,1} \wedge \cdots \wedge df_{\alpha,n-k} + \eta_\alpha. \quad (10)$$

Moreover, g_α is not identically zero on every irreducible component of V and η_α is given by

$$\eta_\alpha = f_{\alpha,1} \eta_{\alpha,1} + \cdots + f_{\alpha,n-k} \eta_{\alpha,n-k},$$

where each $\eta_{\alpha,i} \in \Omega_{U_\alpha}^{n-k}$ is a holomorphic $(n - k)$ -form. In particular, the restriction

$$\frac{\xi_\alpha}{g_\alpha}|_V$$

is well defined.

In this context, we define the GSV index in Brunella's spirit, which coincides with the GSV index when V is a curve invariant under a foliation on a surface.

Definition 16. Under the assumptions of Proposition 7, suppose that $S \subset \text{Sing}(\omega) \cap V$ is a codimension one subvariety of V . The GSV-index of ω relative to V in S is defined by

$$\text{GSV}(\omega, V, S) := \sum_j \text{ord}_S \left(\frac{\xi_\alpha}{g_\alpha} |_{V_j} \right),$$

where the sum is taken over all irreducible components V_j of V and $\text{ord}_S \left(\frac{\xi_\alpha}{g_\alpha} |_{V_j} \right)$ denotes the order of vanishing of $\frac{\xi_\alpha}{g_\alpha} |_{V_j}$ along S .

The following result was proved in [44].

Theorem 16. *Let X be a projective manifold, and let $V \subset X$ be a reduced, locally complete intersection subvariety of dimension k , invariant under a Pfaff system of same dimension induced by a twisted form $\omega \in H^0(X, \Omega_X^{n-k} \otimes \mathcal{L})$. Then the following statements hold:*

- (a) *The integer number $\text{GSV}(\omega, V, S_i)$ depends only on the local representatives of ω , V , and S_i ;*
- (b) *If $\text{Sing}(\omega, V) := \text{Sing}(\omega) \cap V$ has codimension one in V , then the following formula is satisfied:*

$$\sum_i \text{GSV}(\omega, V, S_i) [S_i] = c_1([\mathcal{L} \otimes \det(N_{V/X})^{-1}]|_V) \frown [V],$$

where S_i denotes an irreducible component of $\text{Sing}(\omega, V)$, and $N_{V/X}$ is the normal sheaf of the subvariety V .

The following result offers a method for calculating the GSV-index for Pfaff systems, which can be compared with formulas in [90, Proposition 5.1] and [61]. Moreover, it partially addresses Suwa's question in [90, Remark 5.3.(6)].

Theorem 17. *With the setup as in Theorem 16, let $x \in V$, and let $\{f_1 = \dots = f_{n-k} = 0\}$ be a local equation for V in a neighborhood U of x . Consider*

$$\omega|_U = \sum_{|I|=n-k} a_I dZ_I, \quad \text{where } a_I \in \mathcal{O}(U),$$

as the holomorphic $(n-k)$ -form that induces the Pfaff system

$$\omega \in H^0(U, \Omega_U^{n-k} \otimes \mathcal{L})$$

on U . Then the following formula holds:

$$\text{GSV}(\omega, V, S_i) = \text{ord}_{S_i}(a_I|_V) - \text{ord}_{S_i}(\Delta_I|_V), \quad (11)$$

where Δ_I is the $(n-k) \times (n-k)$ minor of the Jacobian matrix $\text{Jac}(f_1, \dots, f_{n-k})$, corresponding to the multi-index I .

More precisely, let $\omega \in H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^k(d+n-k+1))$ be a holomorphic Pfaff system of dimension k and degree d . Let $V \subset \mathbb{P}^n$ be a reduced complete intersection variety of dimension k and multidegree (d_1, \dots, d_{n-k}) , invariant under ω . Suppose that $\text{Sing}(\omega, V)$ has codimension one in V . Then,

$$\sum_i \text{GSV}(\omega, V, S_i) \deg(S_i) = [d+n-k+1 - (d_1 + \dots + d_{n-k})] \cdot (d_1 \cdots d_{n-k}),$$

where S_i denotes an irreducible component of $\text{Sing}(\omega, V)$. Therefore, if $\text{GSV}(\omega, V, S_i) \geq 0$ for all i , we have

$$d_1 + \dots + d_{n-k} \leq d+n-k+1.$$

This bound is similar to those established by Esteves and Cruz [48] and by Corrêa and Jardim [38]. Related results were previously obtained by Cerveau and Lins Neto [37], Soares [87], Cavalier and Lehmann [35], Brunella and Esteves in [26]. In particular, Soares in [87] gave this bound for a foliation by curves:

$$d_1 + \dots + d_{n-1} \leq d+n,$$

whenever V is a smooth complete intersection invariant curve.

Example 8. Consider the Pfaff system $\omega \in H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}^{n-k}(d+n-k+1))$ given by

$$\omega = \sum_{0 \leq j \leq n-k} (-1)^j d_j f_j df_0 \wedge \cdots \wedge \widehat{df_j} \wedge \cdots \wedge df_{n-k},$$

where f_j is a homogeneous polynomial of degree d_j . We can see that

$$d_0 + d_1 + \dots + d_{n-k} = d+n-k+1.$$

Suppose that $\deg(f_0) = d_0 = 1$ and that $V = \{f_1 = \dots = f_{n-k} = 0\}$ is smooth. Then V is invariant by ω , and we have

$$d_1 + \dots + d_{n-k} = d+n-k.$$

4.1 GSV-indices for vector fields as residues

Suwa in [90] provided an alternative interpretation of the GSV and virtual indices for vector fields as residues on complete intersection varieties with isolated singularities. He considered the residue arising from the localization of the relevant Chern class of the ambient tangent bundle using a frame consisting of ν and additional vector fields. Suwa's definition is recalled as follows and does not depend on the choices made.

Let U be a neighborhood of the origin 0 in \mathbb{C}^m , and let V be a subvariety (reduced, but not necessarily irreducible) of pure dimension n in U . Assume that V contains 0 and that $V \setminus \{0\}$ is non-singular. We consider a closed ball B around 0 that is sufficiently small so that $R = B \cap V$ has a cone structure over $\partial R = L$, the link of V at 0 .

Let E be a C^∞ complex vector bundle of rank l on U , and let $\mathbf{s} = (s_1, \dots, s_r)$ be a C^∞ r -frame of E on a neighborhood V_0 of L in V , where $r = l - n + 1$. Let ∇_0 be an \mathbf{s} -trivial connection for $E|_{V_0}$, and let ∇_1 be a connection for E .

Definition 17. The residue of \mathbf{s} at 0 with respect to c_n is defined by

$$\text{Res}_{c_n}(\mathbf{s}, E|_V; 0) = \int_R c_n(\nabla_1) - \int_{\partial R} c_n(\nabla_0, \nabla_1).$$

We have an exact sequence

$$0 \longrightarrow TV_0 \longrightarrow TU|_{V_0} \xrightarrow{\pi} NV_0 \longrightarrow 0, \quad (12)$$

where TV_0 and TU denote the holomorphic tangent bundles of V_0 and U , respectively, and NV_0 is the normal bundle of V_0 in U . Suppose that V is a complete intersection defined by $f = (f_1, \dots, f_k)$ in U , where $k = m - n$. In a neighborhood of a regular point of f , we may choose (f_1, \dots, f_k) as part of the local coordinates on U , so that we have holomorphic vector fields $\frac{\partial}{\partial f_1}, \dots, \frac{\partial}{\partial f_k}$ away from the critical set of f . These vector fields are linearly independent and normal to the non-singular part of V , so that

$$\partial_f := \left(\pi \left(\frac{\partial}{\partial f_1} \Big|_{V_0} \right), \dots, \pi \left(\frac{\partial}{\partial f_k} \Big|_{V_0} \right) \right)$$

forms a frame for NV_0 . Suppose that we have a C^∞ nonsingular vector field v on V_0 .

Theorem 18. [90, Theorems 3.4 and 4.4] We have

$$\text{GSV}(v, 0) = \text{Res}_{c_n}((v, \partial_f), TU|_V; 0) = \text{Vir}(v, 0),$$

where $\text{Vir}(v, 0)$ is the virtual index of v at 0 .

Since the indices do not depend on the choices made, we have the following consequence.

Corollary 3. Let $V = \{f_1 = \dots = f_k = 0\}$ be a complete intersection, with isolated singularity at 0 , invariant by a foliation by curves \mathcal{F} singular at most at 0 . Then

$$\text{GSV}(\mathcal{F}, 0) = \text{Res}_{c_n}(\mathcal{F}, TU|_V; 0) = \text{Vir}(\mathcal{F}, 0).$$

Now, if $n = 2$, the formula 11 generalizes Suwa's formula for the GSV index. In fact, if $n = 2$, then $k = 1$, and V is a curve, and ω induces a foliation by curves. Let $x \in V$, and $\{f = 0\}$ be a local equation for V in a neighborhood U of x . Consider

$$\omega = a_1 dx - a_2 dy$$

and its dual vector field

$$v = a_2 \frac{\partial}{\partial x} + a_1 \frac{\partial}{\partial y}$$

Proposition 8. [90, Corollary 5.2] *Let $V = \{f = 0\}$ be a curve, with isolated singularity at 0, invariant by a foliation by curves \mathcal{F} singular at most at 0. Then*

$$Z(\omega, V, x) = \text{ord}_x(a_1|_V) - \text{ord}_x(\partial_x f|_V) = \text{ord}_x(a_2|_V) - \text{ord}_x(\partial_y f|_V).$$

An interesting application is that the GSV index provides an obstruction to the continuous extension of the normal bundle of the foliation from the regular part of the invariant variety. Indeed, let $(V, 0) \subset \mathbb{C}^m$ be an ICIS of dimension $n \geq 2$, and let v be a holomorphic vector field on V , singular only at 0. This vector field defines a one-dimensional holomorphic foliation \mathcal{F} on V singular at 0. On $V \setminus \{0\}$, we have the tangent bundle $T\mathcal{F}$ to the foliation, and we consider the smooth normal bundle $N\mathcal{F} = T(V \setminus \{0\})/T\mathcal{F}$.

Corollary 4. [84, Theorem 2] *Let $(V, 0)$ and \mathcal{F} be as above. Then the normal bundle to \mathcal{F} in $V \setminus \{0\}$ extends to 0 as a vector bundle (continuous or smooth) if and only if $\text{GSV}(\mathcal{F}, 0)$ is a multiple of $(n - 1)!$*

5 Log Baum–Bott residues and Aleksandrov’s logarithmic index

Using techniques from homological algebra, Aleksandrov [4] introduced an algebraic index, inspired by the work of Gómez-Mont, to measure the variation between the Poincaré–Hopf and homological indices. This is defined as follows: Let v be a holomorphic local vector field that induces foliation \mathcal{F} . The interior multiplication i_v then generates a complex of logarithmic differential forms:

$$0 \longrightarrow \Omega_{X,p}^n(\log \mathcal{D}) \xrightarrow{i_v} \Omega_{X,p}^{n-1}(\log \mathcal{D}) \xrightarrow{i_v} \cdots \xrightarrow{i_v} \Omega_{X,p}^1(\log \mathcal{D}) \xrightarrow{i_v} \mathcal{O}_{X,p} \longrightarrow 0.$$

Let p be an isolated singularity of the foliation \mathcal{F} . The i_v -homology groups of the complex $\Omega_{X,p}^\bullet(\log \mathcal{D})$ are finite-dimensional vector spaces. Accordingly, the Euler characteristic

$$\chi(\Omega_{X,p}^\bullet(\log \mathcal{D}), i_v) = \sum_{i=0}^n (-1)^i \dim H_i(\Omega_{X,p}^\bullet(\log \mathcal{D}), i_v)$$

of this complex of logarithmic differential forms is well-defined. Since this invariant does not depend on the choice of the local representative v of the foliation \mathcal{F} at p , the *logarithmic index* of \mathcal{F} at the point p is given by

$$\text{Ind}_{\log}(\mathcal{F}, \mathcal{D}, p) := \chi(\Omega_{X,p}^\bullet(\log \mathcal{D}), i_v).$$

In [43, 45], the authors addressed the problem of establishing a global logarithmic residue theorem for meromorphic vector fields on compact complex manifolds. They proved the following result under the assumption that the invariant divisors are of normal crossings type.

Theorem 19. *Let \mathcal{F} be a one-dimensional foliation with isolated singularities and logarithmic along a normal crossing divisor D in a compact complex manifold X . Then*

$$\int_X c_n(T_X(-\log D) - T\mathcal{F}) = \sum_{x \in \text{Sing}(\mathcal{F}) \cap (X \setminus D)} \mu_x(\mathcal{F}) + \sum_{x \in \text{Sing}(\mathcal{F}) \cap D} \text{Ind}_{\log}(\mathcal{F}, D, x),$$

where $\mu_x(\mathcal{F})$ is the Milnor number of \mathcal{F} at x .

In [43], the authors prove that if \mathcal{F} is a one-dimensional foliation on \mathbb{P}^n , with n odd, and isolated singularities (non-degenerate) along a smooth divisor D , then $\text{Sing}(\mathcal{F}) \subset D$ if and only if Soares' bound [87] for the Poincaré problem is achieved, i.e, if and only if

$$\deg(D) = \deg(\mathcal{F}) + 1.$$

More recently, in [39], the authors developed a Baum–Bott residue theory for foliations by logarithmic curves along free divisors:

Theorem 20. *Let \mathcal{F} be a one-dimensional holomorphic foliation on a complex manifold X , logarithmic along a free divisor $D \subset X$, and let φ be a homogeneous symmetric polynomial of degree n . If \mathcal{F} has only isolated singularities, then*

$$\int_X \varphi(T_X(-\log D) - T\mathcal{F}) = \sum_{x \in \text{Sing}(\mathcal{F}) \cap (X \setminus D)} \text{BB}_\varphi(v, x) + \sum_{x \in \text{Sing}(\mathcal{F}) \cap D} \text{Res}_\varphi^{\log}(\mathcal{F}, p),$$

where $\text{BB}_\varphi(v, x)$ is the Baum–Bott residue of \mathcal{F} at x with respect to φ . If, in addition, D has normal crossings at p , then

$$\text{Res}_\varphi^{\log}(\mathcal{F}, D, p) = (2\pi\sqrt{-1})^n \text{Res}_p \left[\begin{array}{c} \varphi(J_{\log} \vartheta) \\ v_1, \dots, v_n \end{array} \right].$$

Also, when $\varphi = \det$, we have

$$\text{Res}_{\det}^{\log}(\mathcal{F}, D, p) = (-1)^n (2\pi\sqrt{-1})^n \text{Ind}_{\log}(\mathcal{F}, D, p).$$

As an application of this result, we can derive a Baum–Bott-type formula for singular varieties. Consider $T_Y := \text{Hom}(\Omega_Y^1, \mathcal{O}_Y)$ the tangent sheaf of Y , where Ω_Y^1 represents the sheaf of Kähler differentials on Y . A holomorphic foliation on Y is a reflexive subsheaf $T\mathcal{F}$ of T_Y such that $[T\mathcal{F}, T\mathcal{F}] \subset T\mathcal{F}$. Then the following result holds.

Corollary 5. *Let Y be a compact normal variety, and let $T\mathcal{F} \subset T_Y$ be a one-dimensional holomorphic foliation with isolated singularities. Let φ be a homogeneous symmetric polynomial of degree $n = \dim(Y)$. If $\pi : (X, D) \rightarrow (Y, \emptyset)$ is a*

functorial log resolution of Y , then

$$\int_X \varphi(T_X(-\log D) - (T\pi^{-1}\mathcal{F})^*) = \sum_{x \in \text{Sing}(\mathcal{F}) \cap Y_{\text{reg}}} \text{BB}_\varphi(\mathcal{F}, x) + \sum_{S_\lambda \subset D \cap \text{Sing}(\pi^*\mathcal{F})} \text{Res}_\varphi^{\log}(\pi^*\mathcal{F}, S_\lambda),$$

where $\text{BB}_\varphi(\mathcal{F}, x)$ is the Baum–Bott index of \mathcal{F} at x with respect to φ . In particular, if $D \cap \text{Sing}(\pi^{-1}\mathcal{F})$ is isolated, then

$$\begin{aligned} \int_X \varphi(T_X(-\log D) - (T\pi^{-1}\mathcal{F})^*) &= \\ &= \sum_{x \in \text{Sing}(\mathcal{F}) \cap Y_{\text{reg}}} \text{BB}_\varphi(\mathcal{F}, x) + \sum_{x \in D \cap \text{Sing}(\pi^{-1}\mathcal{F})} \text{Res}_p \left[\begin{array}{c} \varphi(J_{\log}(\pi^{-1}\nu)) \\ \nu_1, \dots, \nu_n \end{array} \right]. \end{aligned}$$

If, in addition, $\pi^{-1}\mathcal{F}$ is induced by a global vector field ν and $\text{Sing}(\mathcal{F}) \subset \text{Sing}(Y)$, then the following Poincaré–Hopf type formula holds:

$$\chi(Y) = \sum_{x \in D \cap \text{Sing}(\pi^{-1}\mathcal{F})} \text{Res}_p \left[\begin{array}{c} \varphi(J_{\log}(\pi^{-1}\nu)) \\ \nu_1, \dots, \nu_n \end{array} \right] + \chi(\text{Sing}(Y)).$$

As a consequence of Theorem 20, we can show that the difference between the Baum–Bott and logarithmic indices can be expressed in terms of the GSV and Camacho–Sad indices. Before stating this result, let us first establish some notation. If $S \subset X$ is an analytic subset and \mathcal{F} is a foliation by curves in X , with isolated singularities, we denote by

$$\mathbf{R}(\mathcal{F}, S) = \sum_{x \in \text{Sing}(\mathcal{F}) \cap S} \mathbf{R}(\mathcal{F}, x),$$

where \mathbf{R} denotes some residue associated with \mathcal{F} along S . One gets (see [39, Corollary 1.5]):

Theorem 21. *Let \mathcal{F} be a one-dimensional holomorphic foliation on a compact complex surface X with a reduced invariant curve $S \subset X$. Then*

$$\text{BB}_{c_1^2}(\mathcal{F}, S) - \text{Res}_{c_1^2}^{\log}(\mathcal{F}, S) = 2\text{GSV}(\mathcal{F}, S) - \text{CS}(\mathcal{F}, S).$$

Hence, in particular, the non-negativity of

$$\text{BB}_{c_1^2}(\mathcal{F}, S) - \text{Res}_{c_1^2}^{\log}(\mathcal{F}, S) + \text{CS}(\mathcal{F}, S)$$

is an obstruction to an affirmative answer to the Poincaré problem. Moreover, if \mathcal{F} is a generalized curve along S , then

$$\text{Res}_{c_1^2}^{\log}(\mathcal{F}, S) = 2\text{Res}_{c_1^2}(\mathcal{F}, S).$$

Finally, a new obstruction to giving an affirmative answer to the Poincaré problem for foliations by curves on \mathbb{P}^n , n odd, is introduced as a log index as follows (see [39, Corollary 1.4]): let \mathcal{F} be a one-dimensional foliation, with isolated singularities, on \mathbb{P}^n with an invariant *free divisor* D . If n is odd and

$$\text{Res}_{\mathbb{C}^n}^{\log}(\mathcal{F}, \text{Sing}(\mathcal{F})) \geq 0$$

then

$$\deg(D) \leq \deg(\mathcal{F}) + n.$$

6 Final remarks

We have explored the intertwining amongst invariants for singular varieties and for singular foliations. There is a lot more in the literature, and in this section we briefly comment on some recent developments related to this work.

The Milnor number permeates this work and it is a topological invariant, fundamental in singularity theory. There are plenty of extensions and applications of this concept in the literature, see for instance [29] for an account on this.

The Tjurina number was introduced by Greuel in [64]; this is the dimension of the base space of a semi-universal deformation of an isolated hypersurface singularity. In fact one has the Milnor and the Tjurina algebras. For a map germ $(\mathbb{C}^n, o) \rightarrow (\mathbb{C}, 0)$ these are, respectively:

$$M_f = \frac{O_{\mathbb{C}^n, 0}}{\left(\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_n}\right)} \quad \text{and} \quad \mathcal{T}_f = \frac{O_{\mathbb{C}^n, 0}}{\left(f, \frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_n}\right)}.$$

Their dimensions over \mathbb{C} are the corresponding Milnor and Tjurina numbers, $\mu(f)$ and $\tau(f)$. There are deep relations between these two invariants (see for instance [65, 49]).

The Tjurina algebra for holomorphic vector fields, to our knowledge, appears first in Gómez-Mont's formula in [61] for the homological index (see 6), though he does not use this name. The actual name of Tjurina number for foliations was coined in [33, p. 159], where F. Cano, N. Corral and R. Mol use the intersection properties of polar curves for 1-dimensional foliations in complex surfaces, to study properties about non-dicritical singularities. In particular, they use the formula in [61] to give a polar interpretation of the GSV-index and use this to study the Poincaré problem for foliations in $\mathbb{P}_{\mathbb{C}}^2$.

In [55] the authors study relations between the Milnor and Tjurina numbers of singular foliations in the complex plane with respect to a balanced divisor of separatrices, an interesting concept introduced in [57]. In the non-dicritical case, where there are only finitely many separatrices of the foliation, a balanced divisor just means the union of all separatrices. For dicritical singularities this is a finite

set of separatrices which includes all the isolated separatrices and some separatrices from the ones associated to dicritical components. A balanced divisor provides a certain control of the algebraic multiplicity of the foliation and of its separatrices. The authors find in [55] important relations between various indices of a singular foliation, such as the Baum-Bott, Camacho-Sad, GSV variational indices and the Milnor and Tjurina numbers.

In [31], Camacho, Lins Neto and Sad posed the problem of determining whether non-trivial minimal sets exist for foliations on \mathbb{P}^2 . This question remains an open problem to this day. In [74], Lins Neto proved the non-existence of minimal sets for codimension one foliations on \mathbb{P}^n , with $n \geq 3$. Motivated by Lins Neto's work, Marco Brunella in [25] initiated the investigation of this problem in higher dimensions for codimension one holomorphic foliations with ample normal bundles. More precisely, Brunella proposed the following conjecture: *let \mathcal{F} be a codimension-one foliation on a projective manifold of dimension $n \geq 3$. If \mathcal{F} possesses an ample normal bundle, then every leaf of \mathcal{F} necessarily accumulates on its singular set.* In the remarkable work [28] by Brunella and Perrone, the authors, utilizing Baum-Bott residues, proved this conjecture for projective manifolds with Picard rank one. In [2], Adachi and Brinkschulte completely solved Brunella's conjecture by employing the localization of the first Atiyah class, following [1], of the determinant of the normal bundle of the foliation. A second proof of Brunella's conjecture was provided by Adachi, Biard, and Brinkschulte in [3], where the authors employed a residue formula for meromorphic connections. This approach was inspired by a residue formula proved by Pereira in [78]. Recently, in [41], the authors generalized the variational index for currents invariant under foliations, with applications to the study of transcendental leaves of foliations and their accumulation on the singular set. This result represents a higher-dimensional generalization of the result previously established by Brunella for foliations by curves, as proven in [24].

The adjunction formulas for foliations have played an important role in the classification of regular foliations on projective surfaces, as given by Brunella, and also in foliated birational geometry, see [27]. When the curve is invariant under the foliation, the difference $K_{\mathcal{F}} - K_C$ is the total sum of the GSV indices along C . More recently, Cascini and Spicer [34] obtained an adjunction formula for foliations on algebraic varieties, where a correction term serves as a generalization of Brunella's GSV index whenever the divisor is invariant under the foliation. We observe that Theorem 16 also provides an adjunction formula

$$K_{\mathcal{F}}|_V - K_V = \det(N_{\mathcal{F}})|_V + \det(N_{V/X})^{-1} = \sum_i \text{GSV}(\omega, V, S_i)[S_i],$$

since $\det(N_{\mathcal{F}})|_V = -K_X|_V + K_{\mathcal{F}}|_V$ and $\det(N_{V/X}) = -K_X|_V + K_V$.

In fact the GSV index was originally defined for vector fields on hypersurface singularities. The extension to complete intersections was given in [86] where this is used to prove an adjunction formula for local complete intersections.

References

1. M. Abate, F. Bracci, T. Suwa, and F. Tovena, *Localization of Atiyah classes*, Rev. Mat. Iberoam. **29** (2013), no. 2, 547–578. [42](#)
2. M. Adachi and J. Brinkschulte, *Dynamical aspects of foliations with ample normal bundle*, Indiana Univ. Math. J. **72** (2023), no. 5, 1931–1947. [42](#)
3. M. Adachi, S. Biard, and J. Brinkschulte, *A residue formula for meromorphic connections and applications to stable sets of foliations*, J. Geom. Anal., 33 (2023), 338. [42](#)
4. A. G. Aleksandrov, *The Index of Vector Fields and Logarithmic Differential Forms*. Funct. Anal. Appl. 39 (2005), 245–255. [4](#), [9](#), [38](#)
5. A.G. Aleksandrov, *Multidimensional residue theory and the logarithmic De Rham Complex*, Journal of Singularities, **5** (2012), 1–18 [33](#), [34](#)
6. A. G. Aleksandrov, *The index of differential forms on complete intersections*. Funct. Anal. Appl. 49 (2015), 1–14. [9](#)
7. M. Aguilar, J. Seade and A. Verjovsky, *Indices of vector fields and topological invariants of real analytic singularities*, Crelle’s, J. Reine u. Ange. Math. 504 (1998), 159–176. [17](#)
8. P. Aluffi, *MacPherson’s and Fulton’s Chern Classes of Hypersurfaces*. I.M.R.N. (1994), 455–465. [19](#)
9. P. Aluffi, *Chern classes for singular hypersurfaces*, Trans. Amer. Math. Soc. 351 (1999), 3989–4026. [19](#)
10. P. Aluffi, *Segre classes and invariants of singular varieties*. In “Handbook of Geometry and Topology of Singularities III”, ed. J. L. Cisneros Molina *et al*, Springer Verlag 2022, p. 419–491. [15](#)
11. P. Baum, R. Bott, *On the zeros of meromorphic vector fields*, Essay on Topology and Related Topics, Springer-Verlag, New York, 1970, 29–47. [2](#), [26](#)
12. P. Baum, R. Bott, *Singularities of holomorphic foliations*, J. Differential Geom. 7 (1972), 279–342. [2](#), [26](#), [27](#)
13. Ch. Bonatti and X. Gómez-Mont, *The index of a holomorphic vector field on a singular variety*, Astérisque 222 (1994), 9–35. [8](#), [9](#), [10](#), [14](#)
14. H.-C. G. von Bothmer, W. Ebeling and X. Gómez-Mont, *An algebraic formula for the index of a vector field on an isolated complete intersection singularity*, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 5, 1761–1783. [9](#), [11](#)
15. R. Bott, *A residue formula for holomorphic vector fields*, J. Differential Geom. 4 (1967), 311–332. [2](#), [25](#)
16. R. Bott, *On topological obstructions to integrability*. Actes Congr. Int. Math., Nice 1970, Tome 1, 27–36 (1971). [25](#)
17. F. Bracci, T. Suwa, *Perturbation of Baum-Bott residues*, Asian J. Math. 19, no. 5 (2015), 871–886. [27](#)
18. J.-P. Brasselet, *Characteristic Classes*. In “Handbook of Geometry and Topology of Singularities III”, ed. J. L. Cisneros Molina *et al*, Springer Verlag 2022, p. 303–417. [15](#)
19. J.-P. Brasselet, M.-H. Schwartz, *Sur les classes de Chern d’un ensemble analytique complexe*, Astérisque 82–83 (1981), 93–147.
20. J.-P. Brasselet, D. Lehmann, J. Seade and T. Suwa, *Milnor classes of local complete intersections*, Transactions A. M. S. 354 (2001), 1351–1371. [15](#), [16](#)
21. J.-P. Brasselet, D. T. Lê, J. Seade, *Euler obstruction and indices of vector fields*. Topology 39 (2000), 1193–1208. [2](#)
22. J.-P. Brasselet, J. Seade, T. Suwa. *Vector fields on singular varieties*. Lecture Notes in Mathematics 1987 (2009). Springer-Verlag, Berlin. [2](#), [3](#), [6](#), [7](#), [10](#), [14](#), [15](#), [16](#), [17](#), [18](#), [19](#)
23. M. Brunella, *Some remarks on indices of holomorphic fields*, Publicacions Matemàtiques, **41** (1997), 527–544. [3](#), [28](#), [30](#), [31](#), [32](#), [33](#)
24. M. Brunella, *Inexistence of invariant measures for generic rational differential equations in the complex domain*, Bol. Soc. Mat. Mexicana (3) **12** (1) (2006), 43–49. [42](#)
25. M. Brunella, *On the dynamics of codimension one holomorphic foliations with ample normal bundle*, Indiana Univ. Math. J., **57**(7) (2008), 3101–3113. [42](#)

26. M. Brunella, L. G. Mendes, *Bounding the degree of Solutions to Pfaff Equations*, Publicacions Matemàtiques, **44**, (2000), 593-604. [4](#), [33](#), [36](#)
27. M. Brunella, *Birational Geometry of Foliations*, IMPA Mathematical Publications, Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2004. [3](#), [42](#)
28. M. Brunella, C. Perrone, *Exceptional singularities of codimension one holomorphic foliations*, Publ. Mat. 55 (2011), 295–312. [42](#)
29. Callejas-Bedregal, R., Morgado, M. F. Z. and Seade, J., *Milnor number and Chern classes for singular varieties: an introduction*. In “Handbook of Geometry and Topology of Singularities III”, ed. J. L. Cisneros Molina *et al*, Springer Verlag 2022, p. 494-563. [2](#), [3](#), [15](#), [16](#), [17](#), [19](#), [41](#)
30. C. Camacho, A. Lins, P. Sad, *Topological invariants and equidesingularization for holomorphic vector fields*. J. Differ. Geom. 20, 143-174 (1984). [8](#), [32](#)
31. C. Camacho, A. Lins Neto, P. Sad, *Minimal sets of foliations on complex projective spaces*, Publications Mathématiques de l’Institut des Hautes Études Scientifiques, No. 68 (1988), 187-203. [42](#)
32. C. Camacho, P. Sad, *Invariant varieties through singularities of holomorphic vector fields*, Ann. of Math. **115** (1982), 579-595. nd topology of singularities III. Ed. J. L. Cisneros-Molina *et al*. Springer. 493-564 (2022). [28](#), [29](#), [31](#), [33](#)
33. F. Cano, N. Corral and R. Mol, *Local polar invariants for plane singular foliations*, Expo. Math. 37 (2019), 145-164 [13](#), [41](#)
34. P. Cascini, C. Spicer, *Foliation adjunction*, arXiv:2309.10697, 2024. [42](#)
35. V. Cavalier, D. Lehmann, *On the Poincaré inequality for one-dimensional foliations*, Compositio Mathematica, 142(2) (2006), 529-540. [4](#), [36](#)
36. V. Cavalier and D. Lehmann, *Localisation des résidus de Baum–Bott, courbes généralisées et K-théorie (I: feuilletages dans \mathbb{C}^2)*, Commentarii Mathematici Helvetici, **76** (2001), 665–683. [3](#), [32](#)
37. D. Cerveau and A. Lins Neto, *Holomorphic foliations in $\mathbb{P}_{\mathbb{C}}^2$ having an invariant algebraic curve*, Ann. Inst. Fourier 41 (1991), 883-903. [4](#), [36](#)
38. M. Corrêa Jr, M. Jardim, *Bounds for sectional genera of varieties invariant under Pfaff fields*, Illinois Journal of Mathematics, vol. **56** (2) (2012), 343-352. [4](#), [36](#)
39. M. Corrêa, F. Lourenço, and D. Machado, *Log Baum-Bott Residues for Foliations by Curves*, 2024. arXiv:2411.04015v1. [3](#), [4](#), [39](#), [40](#), [41](#)
40. M. Corrêa, *Analytic Varieties Invariant by Holomorphic Foliations and Pfaff Systems*, in: F. Cano, J.L. Cisneros-Molina, L. Dũng Tráng, J. Seade (eds) Handbook of Geometry and Topology of Singularities VI: Foliations, Springer, 2024.
41. M. Corrêa, A. Fernández-Pérez, and M. G. Soares, *Brunella–Khanedani–Suwa Variational Residues for Invariant Currents*, Journal of Singularities **23** (2021), 107–115. [32](#), [42](#)
42. M. Corrêa and F. Lourenço, *Determination of Baum-Bott residues of higher codimensional foliations*, Asian J. Math., 23 (3) (2019), 527–538. [27](#)
43. M. Corrêa, D. Machado, *Residue formulas for logarithmic foliations and applications*, Trans. Amer. Math. Soc., 371 (2019), 6403-6420. [3](#), [4](#), [39](#)
44. M. Corrêa and D. Machado, *GSV-index for holomorphic Pfaff systems*, Documenta Mathematica, vol. 25, pp. 1011–1027, 2020. [3](#), [33](#), [35](#)
45. M. Corrêa, D. Machado, *Global residue formula for logarithmic indices of one-dimensional foliations*, Ann. Math., 32 (2024), no. 4, 1095-1118. [4](#), [39](#)
46. M. Corrêa, A. M. Rodríguez, M. G. Soares, *A Bott type residues formula for complex orbifolds*, International Mathematics Research Notices, (10), 2889-2911, 2016. [3](#)
47. F. Cukierman and J. V. Pereira, *Stability of holomorphic foliations with split tangent sheaf*, Amer. J. Math., **130** (2008), no. 2, 413–439. [24](#)
48. J. D. A. Cruz, E. Esteves, *Regularity of subschemes invariant under Pfaff fields on projective spaces*. Commentarii Mathematici Helvetici **86** (2011), 947-965. [4](#), [36](#)
49. A. Dimca, G.-M. Greuel, *On 1-forms on isolated complete intersection on curve singularities*, J. of Singularities 18 (2018), 114-118. [41](#)
50. W. Ebeling and S. Gusein-Zade, *On the index of a vector field at an isolated singularity*. The Arnoldfest (Toronto, ON, 1997), 141-152, Fields Inst. Commun., 24, Amer. Math. Soc., Providence, RI, 1999. [17](#), [18](#)

51. W. Ebeling and S. Gusein-Zade, *Indices of vector fields and 1-forms*. In Handbook of geometry and topology of singularities III. Ed. J. L. Cisneros-Molina *et al.* Springer. 493-564 (2022). [2](#), [17](#), [18](#), [19](#)
52. W. Ebeling, G. Gusein-Zade, J. Seade. *Homological index for 1-forms and a Milnor number for isolated singularities*. International Journal of Maths. 15 (2004), no. 9, 895-905. [9](#)
53. D. Eisenbud and H. Levine, *An algebraic formula for the degree of a C^∞ map germ*, Ann. Math. (2) 106 (1977), no. 1, 19-44. [9](#)
54. G. Ellingsrud and S. Stromme, *Bott's formula and enumerative geometry*, J. Amer. Math. Soc. 9 (1996), no. 1, 175-193. [2](#)
55. A. Fernández-Pérez, E. R. García Barroso, N. Saravia-Molina. *On Milnor and Tjurina numbers of foliations*; arXiv:2112.14519. [41](#), [42](#)
56. W. Fulton, *Intersection Theory*, Springer-Verlag (1984). [15](#)
57. Y. Genzmer, *Rigidity for dicritical germ of foliation in \mathbb{C}^2* . Int. Math. Res. Notices 19 (2007), 14 p. [41](#)
58. L. Giraldo, X. Gómez-Mont and P. Mardesic, *On the complex formed by contracting differential forms with a vector field on a hypersurface singularity*, Bol. Soc. Mat. Mexicana (3) 7 (2001), no. 2, 211-221. [9](#)
59. L. Giraldo, X. Gómez-Mont and P. Mardesic, *A law of conservation of number for local Euler characteristics*, Complex manifolds and hyperbolic geometry (Guanajuato, 2001), 251-259, Contemp. Math., 311, Amer. Math. Soc., Providence, RI, 2002. [9](#)
60. L. Giraldo, X. Gómez-Mont and P. Mardesic, *On the index of vector fields tangent to hypersurfaces with non-isolated singularities*, J. London Math. Soc. (2) 65 (2002). [9](#)
61. X. Gómez-Mont. *An algebraic formula for the index of a vector field on a hypersurface with an isolated singularity*. J. Algebraic Geom. 7 (1998), 731-752. [4](#), [9](#), [10](#), [11](#), [12](#), [13](#), [35](#), [41](#)
62. X. Gómez-Mont, J. Seade, A. Verjovsky. *The index of a holomorphic flow with an isolated singularity*. Math. Ann. 291 (1991), 737-751. [2](#), [8](#)
63. G.-M. Greuel, *Der Gauß-Manin-Zusammenhang isolierter Singularitäten von vollständigen Durchschnitten*, Math. Ann. **214** (1975), 235-266. [7](#)
64. G.M. Greuel. *Dualität in der lokalen Kohomologie isolierter Singularitäten*. Math. Ann. 250 (2) (1980), 157-173. [41](#)
65. G.M. Greuel, Ch. Lossen, E. Shustin, *Introduction to Singularities and Deformations*. Springer Monographs in Mathematics, 2007. [41](#)
66. H. Hamm. *Lokale topologische Eigenschaften komplexer Räume*. Math. Ann., 191 (1971), 235-252. [6](#)
67. L. Kaufmann, R. Lärkäng, and E. Wulcan, *Baum-Bott residue currents*, arXiv preprint, arXiv:2302.08887, 2023. [27](#)
68. B. Khanedani, T. Suwa, *First variation of holomorphic forms and some applications*, Hokkaido Math. J. **26** (1997) , 323-335. [28](#), [29](#), [31](#), [33](#)
69. G.N. Khimshiashvili, *On the local degree of a smooth mapping*, Comm. Acad. Sci. Georgian SSR 85, No. 2 (1977), 309-312 (in Russian). [9](#)
70. H. King and D. Trotman, *Poincaré-Hopf theorems on stratified sets*, preprint 1996. (Published as "Poincaré-Hopf theorems on singular spaces", Proc. Lond. Math. Soc. (3) 108, No. 3, 682-703 (2014).) [17](#)
71. M. Kontsevich, *Enumeration of rational curves via torus actions*, The Moduli Space of Curves (Progress in Mathematics, Vol. 129), Birkhäuser, 1995, 335-368. [2](#)
72. D.T. Lê, *Computation of the Milnor number of an isolated singularity of a complete intersection*, Funct. Anal. Appl. **8** (1974), 45-49. [7](#)
73. D. Lehmann, M. Soares and T. Suwa, *On the index of a holomorphic vector field tangent to a singular variety*, Bol. Soc. Bras. Mat. 26 (1995), pp 183-199. [14](#), [15](#), [17](#)
74. A. Lins Neto, *A note on projective Levi flats and minimal sets of algebraic foliations*, Annales de l'Institut Fourier, Vol. 49, No. 4 (1999). [42](#)
75. E. Looijenga. *Isolated Singular Points on Complete Intersections*. Cambridge Univ. Press, Cambridge, London, New York, New Rochelle, Melbourne, Sydney, 1984. [6](#)
76. R. MacPherson, *Chern classes for singular varieties*, Ann. of Math **100** (1974), 423-432. [2](#)

77. J. W. Milnor. *Singular points of complex hypersurfaces*. Annals of Mathematics Studies, 61, Princeton University Press, Princeton, N.J., 1968. [6](#), [7](#), [9](#), [25](#)
78. J. V. Pereira, *Closed meromorphic 1-forms*, Handbook of Geometry and Topology of Singularities V: Foliations, 447–499. Springer, Cham (2024). [42](#)
79. H. Poincaré, *Sur l'integration algebrique des equations différentielles du premier ordre et du premier degré*, Rendiconti del Circolo Matematico di Palermo **5** (1891), 161-191; 11 (1897), 193–239. [33](#)
80. R. Rosas, *Transversely product singularities of foliations in projective spaces*, Comptes Rendus Mathématique, **361** (2023), no. G11, 1785–1787. [27](#)
81. K. Saito, *Theory of logarithmic differential forms and logarithmic vector fields*, J. Fac. Sci. Univ. Tokyo **27**(2) (1980), 265-291. [30](#), [34](#)
82. M.-H. Schwartz, *Classes caractéristiques définies par une stratification d'une variété analytique complexe*, C.R. Acad. Sci. Paris **260** (1965), 3262-3264, 3535-3537. [2](#)
83. J. Seade, *The index of a vector field on a complex surface with singularities*, in “The Lefschetz Centennial Conf.”, ed. A. Verjovsky, Contemp. Math. **58**, Part III, Amer. Math. Soc. (1987), 225-232. [2](#)
84. J. Seade, *Remarks on contact structures and vector fields on isolated complete intersection singularities*, The Proceedings of the IX Workshop on Real and Complex Singularities, edit. M. Saia *et al*, Contemporary Mathematics, AMS. 459, 2008, 195-205. [38](#)
85. J. Seade, T. Suwa. *A residue formula for the index of a holomorphic flow*. Mathematische Annalen 304 (1996), 621-634. [3](#)
86. J. Seade, T. Suwa. *An adjunction formula for local complete intersections*. Int. J. Maths. **9**, 1998, 759-768. [14](#), [42](#)
87. M. Soares, *The Poincaré problem for hypersurfaces invariant by one-dimensional foliations*, Invent. Math. **128** (1997), 495-500. [4](#), [36](#), [39](#)
88. T. Suwa, *Indices of holomorphic vector fields relative to invariant curves on surfaces*, Proc. Amer. Math. Soc. **123** (1995) , 2989-2997. [29](#), [30](#)
89. T. Suwa, *Indices of Vector Fields and Residues of Singular Holomorphic Foliations*, Actualités Mathématiques, Hermann Paris, 1998. [2](#)
90. T. Suwa, *GSV-Indices as Residues*, Journal of Singularities **9** (2014), 206-218. [4](#), [14](#), [17](#), [35](#), [36](#), [37](#), [38](#)
91. J.-L. Verdier. *Spécialisation des classes de Chern*. Astérisque 82-83 (1981), 149-159. [16](#)
92. M. S. Vishik, *Singularities of analytic foliations and characteristic classes*, Funct. Anal. Appl. **7** (1973), 1-15. [27](#)
93. M. Zach. *A generalization of Milnor's formula*. Math. Annalen **382**, 901-942. [9](#)
94. M. Zach. *Some Lê-Greuel type formulae on stratified spaces*. Preprint 2024, arXiv:2411.02682. [9](#)