

On large Sidon sets

Ingo Czerwinski* and Alexander Pott*

Abstract

A Sidon set M is a subset of \mathbb{F}_2^t such that the sum of four distinct elements of M is never 0. The goal is to find Sidon sets of large size. In this note we show that the graphs of almost perfect nonlinear (APN) functions with high linearity can be used to construct large Sidon sets. Thanks to recently constructed APN functions $\mathbb{F}_2^8 \rightarrow \mathbb{F}_2^8$ with high linearity, we can construct Sidon sets of size 192 in \mathbb{F}_2^{15} , where the largest sets so far had size 152. Using the inverse and the Dobbertin function also gives larger Sidon sets as previously known. Each of the new large Sidon sets M in \mathbb{F}_2^t yields a binary linear code with t check bits, minimum distance 5, and a length not known so far. Moreover, we improve the upper bound for the linearity of arbitrary APN functions.

Keywords Sidon sets, vectorial boolean functions, almost perfect nonlinear (APN) functions, linear binary codes.

Mathematics Subject Classification (2020) Primary: 11B13, 94D10. Secondary: 94B05.

1 Introduction

The following easy question seems to be difficult to answer: What is the largest number of elements in a set, in which an appropriate addition is defined, so that no sum appears twice?

In the early 1930s Sidon [19, 20] was the first who discussed such sets in the integers. Later Babai and Sós [1] generalised such sets to arbitrary groups and called them **Sidon sets**. In this note, we discuss only Sidon sets in \mathbb{F}_2^t , the t -dimensional vector space over the binary field \mathbb{F}_2 .

*Otto-von-Guericke-University Magdeburg, Faculty of Mathematics, Institute for Algebra and Geometry, 39106 Magdeburg, Germany, (ingo@czerwinski.eu, alexander.pott@ovgu.de)

Definition 1.1 ([1]). *A subset M of \mathbb{F}_2^t is called a Sidon set if $m_1 + m_2 \neq m_3 + m_4$ for all pairwise distinct $m_1, m_2, m_3, m_4 \in M$. Equivalently, the sum of any four distinct elements of M is different from 0.*

We emphasise that, by definition, every subset of a Sidon set is also a Sidon set.

The main research question on Sidon sets in \mathbb{F}_2^t is to find sets of large size. For dimensions $t \geq 11$, the maximum size is unknown. Therefore,

- either improve the lower bound by giving new constructions
- or improve the upper bound by new arguments.

The trivial upper bound (already mentioned in [1]) arises from the fact that the sums of two distinct elements of M have to be distinct and non-zero, hence

$$\binom{|M|}{2} = \frac{|M|(|M| - 1)}{2} \leq |\mathbb{F}_2^t \setminus \{0\}|.$$

So far, progress on the upper bound has only been made in terms of coding theory. This uses a one-to-one correspondence between sum-free Sidon sets and linear codes with minimum distance ≥ 5 . A set in \mathbb{F}_2^t is called *sum-free* if the sum of any three elements of this set is different from 0. If M is a sum-free Sidon set, then $M \cup \{0\}$ is Sidon but not sum-free anymore. An overview of the one-to-one correspondence can be found in [12].

The first improvement of the trivial bound was made in 1993 by Brouwer and Tolhuizen [4]. They improved the case t odd. The case t even was recently improved by the authors in [12]. We state here only the Brouwer-Tolhuizen version.

Proposition 1.2. *For any t , an upper bound for the maximum size of a Sidon set in \mathbb{F}_2^t is given by*

$$s_{max}(\mathbb{F}_2^t) \leq \begin{cases} 2^{\frac{t+1}{2}} - 2 & \text{for } t \text{ odd,} \\ \lfloor \sqrt{2^{t+1}} + 0.5 \rfloor & \text{for } t \text{ even,} \end{cases} \quad (1)$$

The improvement in [12] is that we may subtract 1 or 2 in the case that t is even, depending on some technical conditions.

For small dimensions $t \leq 14$, better upper bounds are known derived from coding-theory (see e.g Table 2 of [12] or Table 2 in this paper).

On the construction side, the following is known. In the case $t = 2n$ even, Sidon sets of size $2^n + 1$ are known. They are obtained from non-primitive BCH codes which are constructed as shortenings of codes with parameters $[2^n + 1, 2^n + 1 - 2n, 5]$ (see [15, page 586, 18.2 (ii)] or (C2) construction in [16]).

On large Sidon sets

Another construction is by Carlet, Mesnager and Picek [10, 11]. They have shown that the multiplicative subgroup M of size $2^n + 1$ in the finite field $\mathbb{F}_{2^{2n}}$ is a Sidon set in the additive group of the field. This set can also be constructed from a certain Goppa code, see [16]. If n is even, then M is Sidon and sum-free, and therefore $M \cup \{0\}$ Sidon too. We call constructions giving Sidon sets of such sizes for t even **classical**. These Sidon sets are of maximum size for $t = 4, 6, 8$. For $t = 10$ the construction leads to 33 but the largest possible Sidon set in dimension 10 has size 34.

The second important class of Sidon sets in even dimensions $t = 2n$ are Sidon sets of size 2^n and stems from almost perfect nonlinear functions. These functions are quite important from a cryptographic perspective.

Definition 1.3 ([17]). *A function $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ is called almost perfect nonlinear (APN) if for every $a \in \mathbb{F}_2^n \setminus \{0\}$ and $b \in \mathbb{F}_2^n$ the equation $F(x) + F(x+a) = b$ has at most 2 solutions in \mathbb{F}_2^n .*

Several classes of APN functions are known: there are some infinite families, see [9] for a summary, but also many sporadic constructions (which are quite important for us), see [2]. The relationship between APN functions and Sidon sets is the following:

Proposition 1.4. *A function $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ is APN if and only if its graph $G_F = \{(x, F(x)) : x \in \mathbb{F}_2^n\} \subseteq \mathbb{F}_2^n \times \mathbb{F}_2^n$ is a Sidon set in \mathbb{F}_2^{2n} .*

Note that the graph of an APN function is in \mathbb{F}_2^{2n} and has size 2^n which is slightly worse than the other constructions mentioned above. But they will play an important role in Section 4 to construct large Sidon sets in odd dimensions.

In the case of odd dimension $t = 2n - 1$, less is known how to construct large Sidon sets. There is only one construction which uses the shortening of an BCH code (see [15, page 586, 18.2 (vi)] or construction (C3) in [16]). It leads to Sidon sets in \mathbb{F}_2^{2n-1} of size $2^{n-1} + 2^{n/2}$ for n even and of size $2^{n-1} + 2^{\frac{n-1}{2}}$ for n odd. As in even dimension, we call constructions giving Sidon sets of such sizes for t odd **classical**. For $t = 7$, this construction gives a Sidon set of maximum possible size. For $t > 7$ and t odd, several sporadic constructions are known which give larger Sidon sets, see, for instance, Table 2.

Now, that we have seen the classical constructions of large Sidon sets and the APN functions as important examples for large Sidon sets, we will introduce a new construction of large Sidon sets in the next section. In its general form, it uses an existing Sidon set and intersects it with subspaces. In Section 3 we will then show that the Walsh spectrum and therefore the linearity of an APN function gives us information about the intersection of its graph with hyperplanes. In addition, we improve the upper bound for the linearity of an arbitrary APN

function. In Section 4 we then explicitly construct large Sidon sets from APN functions with high linearity, and in Section 5 we translate them into coding theory terms. Finally, in Section 6, we summarise all results and give an overview table about the maximum size of Sidon sets in small dimensions and related bounds and constructions.

2 Sidon sets in subspaces

The idea of constructing large Sidon sets is based on the fact that every subset of a Sidon set is also a Sidon set. If M is a Sidon set in \mathbb{F}_2^t and U an s -dimensional linear subspace of \mathbb{F}_2^t , then $M \cap U$ is a Sidon subset of \mathbb{F}_2^t and therefore a Sidon subset in the s -dimensional vector space U .

Also the intersection with affine subspaces can be used to construct Sidon sets, since with M also $M + b$ is Sidon for $b \in \mathbb{F}_2^t$. Hence, the intersection of M with an affine subspace $U + b$ leads to Sidon sets in U by intersecting $M + b$ with U (note $|M \cap (U + b)| = |(M + b) \cap U|$).

Our goal is to construct large Sidon sets using this procedure. In order to get a large Sidon set in this way, you need a subspace U which contains “many” elements. But one expects that large Sidon sets are “equally” distributed within \mathbb{F}_2^t , hence $|M \cap U| \approx |M|/2^{t-s}$ is the expectation. In other words: if we intersect a large Sidon set with a **hyperplane**, we expect that the size of the intersection is in the range of $|M|/2$. However, the upper bound reduces the size only by a factor $1/\sqrt{2}$ and we present examples where the intersection size of M with a hyperplane is in the range $|M|/\sqrt{2}$ (see Section 4).

Claude Carlet pointed out that the construction in Theorem 4.1 of [11] may be also reformulated as a construction of a large Sidon set as the intersection of the graph of an APN function with an appropriate subgroup (not a hyperplane).

3 Walsh Transform and Linearity

Let \mathbb{F}_2^n be equipped with a non-degenerate bilinear form $a \cdot b$.

Let $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^m$ be a function. The *Walsh transform* of F is the mapping $W_F: \mathbb{F}_2^n \times \mathbb{F}_2^m \rightarrow \mathbb{Z}$ defined by

$$W_F(a, b) = \sum_{x \in \mathbb{F}_2^n} (-1)^{a \cdot x + b \cdot F(x)}$$

and the *linearity* of F is defined by

$$\mathcal{L}(F) = \max_{a \in \mathbb{F}_2^n, b \in \mathbb{F}_2^m \setminus \{0\}} |W_F(a, b)|.$$

On large Sidon sets

In fact, the *Walsh transform* of F can also be formulated as a property of the Graph G_F inside its ambient space \mathbb{F}_2^{n+m} and can be described as follows:

Definition 3.1. *Let M be a subset of \mathbb{F}_2^t . The Walsh transform of M is the mapping $W_M: \mathbb{F}_2^t \rightarrow \mathbb{Z}$ defined by*

$$W_M(a) = \sum_{m \in M} (-1)^{a \cdot m}$$

and the linearity of M is defined by

$$\mathcal{L}(M) = \max_{a \in \mathbb{F}_2^t \setminus \{0\}} |W_M(a)|.$$

We remark that $W_M(a)$ is even if and only if $|M|$ is even.

Proposition 3.2. *Let $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^m$ be a function and G_F its graph, then $\mathcal{L}(F) = \mathcal{L}(G_F)$. In other words, the linearity of the function equals the linearity of its graph.*

Proof.

$$\begin{aligned} \mathcal{L}(F) &= \max_{a \in \mathbb{F}_2^n, b \in \mathbb{F}_2^m \setminus \{0\}} |W_F(a, b)| \\ &= \max_{a \in \mathbb{F}_2^n, b \in \mathbb{F}_2^m \setminus \{0\}} \left| \sum_{x \in \mathbb{F}_2^n} (-1)^{a \cdot x + b \cdot F(x)} \right| \\ &= \max_{(a, b) \in \mathbb{F}_2^n \times (\mathbb{F}_2^m \setminus \{0\})} \left| \sum_{x \in \mathbb{F}_2^n} (-1)^{(a, b) \cdot (x, F(x))} \right| \\ &= \max_{(a, b) \in \mathbb{F}_2^n \times \mathbb{F}_2^m \setminus (0, 0)} \left| \sum_{x \in \mathbb{F}_2^n} (-1)^{(a, b) \cdot (x, F(x))} \right| = \mathcal{L}(G_F). \end{aligned}$$

The last equality holds because of $\sum_{x \in \mathbb{F}_2^n} (-1)^{a \cdot x} = 0$ and therefore

$$\sum_{x \in \mathbb{F}_2^n} (-1)^{(a, 0) \cdot (x, F(x))} = 0$$

for $a \in \mathbb{F}_2^n \setminus \{0\}$. □

The following Proposition is well known.

Proposition 3.3. *Let M be a subset of \mathbb{F}_2^t , $a \in \mathbb{F}_2^t \setminus \{0\}$, $H_a := \{g \in \mathbb{F}_2^t : a \cdot g = 0\}$ and let $b \in \mathbb{F}_2^t \setminus H_a$. Then the Walsh coefficient $W_M(a)$ has different representations using intersection numbers with affine hyperplanes:*

$$\begin{aligned} W_M(a) &= |M \cap H_a| - |(M + b) \cap H_a| \\ &= |M| - 2 \cdot |(M + b) \cap H_a| \\ &= 2 \cdot |M \cap H_a| - |M| \end{aligned}$$

and

$$\begin{aligned} |M \cap H_a| &= \frac{|M| + W_M(a)}{2} \\ |(M + b) \cap H_a| &= \frac{|M| - W_M(a)}{2}. \end{aligned}$$

Proof.

$$\begin{aligned} W_M(a) &= \sum_{m \in M} (-1)^{a \cdot m} \\ &= \sum_{m \in M \cap H_a} 1 + \sum_{m \in (M+b) \cap H_a} -1 \\ &= |M \cap H_a| - |(M + b) \cap H_a| \end{aligned}$$

The remaining follows from $|M| = |M \cap H_a| + |(M + b) \cap H_a|$. \square

Corollary 3.4. *If M is subset of \mathbb{F}_2^t then there exists a hyperplane H and $b \in \mathbb{F}_2^t \setminus H$ such that*

$$|M \cap H| = \frac{|M| + \mathcal{L}(M)}{2}$$

or

$$|(M + b) \cap H| = \frac{|M| + \mathcal{L}(M)}{2}.$$

Proof. From the definition of $\mathcal{L}(M)$ it follows that $\mathcal{L}(M) = |W_M(a)|$ for some $a \in \mathbb{F}_2^t \setminus \{0\}$. Let $H_a = \{g \in \mathbb{F}_2^t : a \cdot g = 0\}$ and $b \in \mathbb{F}_2^t \setminus H_a$ as above. Proposition 3.3 shows the following: $|M \cap H_a| = \frac{|M| + W_M(a)}{2} = \frac{|M| + \mathcal{L}(M)}{2}$ if $W_M(a) > 0$, and $|(M + b) \cap H_a| = \frac{|M| - W_M(a)}{2} = \frac{|M| + \mathcal{L}(M)}{2}$ otherwise. \square

Corollary 3.5. *Let M be a Sidon set in \mathbb{F}_2^t . Then there exists a Sidon set in \mathbb{F}_2^{t-1} of size $\frac{|M| + \mathcal{L}(M)}{2}$.*

Proof. Use Corollary 3.4 and note that $M + b$ is Sidon too. \square

Corollary 3.6. *Let $F : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ be an APN function. Then there exists a Sidon set in \mathbb{F}_2^{2n-1} of size $2^{n-1} + \frac{\mathcal{L}(F)}{2}$.*

Proof. Direct consequence of Proposition 3.2 and Corollary 3.5. \square

In the next section, we will apply this corollary to several highly linear APN functions.

Not so much is known about an upper bound of the linearity of APN functions (see [8, Section 5.2]). An APN function $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ cannot have linearity 2^n , see [9]. For certain classes of APN functions smaller bounds are known. If F is a quadratic APN function then its linearity is $\leq 2^{n-1}$ (see [8]). Here quadratic means that all component functions have degree 2, or, in other words, the function $x \mapsto F(x+a) + F(x) + F(a) + F(0)$ is linear for $a \in \mathbb{F}_2^n$. We will give an interesting application of Corollary 3.6 to improve the bound on the linearity of arbitrary APN functions.

Assuming $\mathcal{L}(F) = 2^n - 2$ would lead to a Sidon set of size $2^{n-1} + 2^{n-1} - 1 = 2^n - 1$ in \mathbb{F}_2^{2n-1} due to Corollary 3.6. But this is not possible because of Proposition 1.2.

Corollary 3.7. *If $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ is APN then $\mathcal{L}(F) \leq 2^n - 4$.*

The connection between Sidon sets and APN functions also shows that no component function of an APN function can have weight 1, i.e. no component can have all its values 0 with only one exception. To see this, let $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ be a function and let $a \in \mathbb{F}_2^n \setminus \{0\}$. Then the *component function* $F_a: \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ of F is defined as $x \mapsto a \cdot F(x)$.

Corollary 3.8. *Let $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ be an APN function, $a \in \mathbb{F}_2^n \setminus \{0\}$, $\lambda \in \mathbb{F}_2$ and define $w_{a,\lambda} = |\{x \in \mathbb{F}_2^n : F_a(x) = \lambda\}|$. Then there exists a Sidon set of size $2^n - w_{a,\lambda}$ in \mathbb{F}_2^{2n-1} .*

Proof. The set $H_a = \{(x, y) \in \mathbb{F}_2^{2n} : (0, a) \cdot (x, y) = 0\}$ defines a hyperplane in \mathbb{F}_2^{2n} . Therefore the sets $G_F \cap H_a$ and $(G_F + b) \cap H_a$ are Sidon sets, where $b \in \mathbb{F}_2^{2n} \setminus H_a$. We have to choose the translate $G_F + b$ if $\lambda = 1$, compare with Proposition 3.3. \square

Corollary 3.9. *There is no APN function $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ with component function $x_1 x_2 \dots x_n$.*

Proof. Note that $x_1 x_2 \dots x_n$ gives a component function with $w_{a,1} = 1$, which results in a Sidon set of size $2^n - 1$ in \mathbb{F}_2^{2n-1} due to Corollary 3.8. But this Sidon set can not exist because of Proposition 1.2. \square

4 New construction

From Corollary 3.6 follows that it is sufficient to know the linearity of an APN function in order to know the best possible size when intersecting its graph with a hyperplane. Therefore we discuss in this section the known linearities of APN

functions. There are many good research papers and books which give a good overview about APN functions and their Walsh spectra, we refer to [6, 9, 18]

Let us begin with quadratic APN functions $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$. Several infinite families of quadratic APN functions as well as sporadic ones are known.

If n is odd, quadratic APN functions are almost bent, see [9]. That means that the linearity must be $2^{(n+1)/2}$. We may apply Corollary 3.6 to these APN functions and obtain Sidon sets of size $2^{n-1} + 2^{\frac{n-1}{2}}$ in \mathbb{F}_2^{2n-1} . This size matches the size of the classical constructions, hence we do not obtain Sidon sets larger than the known ones.

The situations is more interesting if n is even. The linearities of the known **infinite** families of quadratic APN functions for n even are $2^{n/2+1}$. Applying Corollary 3.6 results in Sidon sets of size $2^{n-1} + 2^{n/2}$. Again, this size matches the size of the classical constructions. Many **sporadic** functions for $n = 6$ and $n = 8$ are known, see [2, 5, 21, 22]. It is interesting that some of them have the best possible linearity 2^{n-1} (see Section 3). In the case $n = 6$ (see [5]) there is one quadratic function with linearity 32. Unfortunately, this yields only a Sidon set of size 48 in \mathbb{F}_2^{11} , which is not an improvement, but is equal to the best previously known example, see Table 2. In the case $n = 8$ we have examples with linearity 32 (the classical one), and also with linearity 64 and 128 (see [2]). If we use Corollary 3.6 for a function with linearity 128, we obtain the following remarkable improvement on the size of the largest known Sidon set (which was 152) in dimension 15:

Theorem 4.1. *There exists a Sidon set in \mathbb{F}_2^{15} of size 192.*

A maximal Sidon set in \mathbb{F}_2^{15} of size 192 is given in Table 1. In the table we use the standard integer representation of vectors in \mathbb{F}_2^t : the integer $\sum_{i=0}^{t-1} a_i 2^i$ in 2-adic representation “is” the vector (a_0, \dots, a_{t-1}) .

According to [2], there are four different APN function on 8 variables with linearity 128. Calculations show that each of the four functions leads to a Sidon set of size 192, which cannot be transformed with an affine permutation to one of the others. This has been also pointed out to us by Gabor Nagy.

Remark 4.2. *It might be possible that APN functions on \mathbb{F}_2^n with linearity 2^{n-1} exist for all even n . In that case we would have Sidon sets of size $2^{n-1} + 2^{n-2} = 3 \cdot 2^{n-2}$ in \mathbb{F}_2^{2n-1} .*

Next we discuss the inverse function $F(x) = x^{-1}$ on \mathbb{F}_{2^n} (here $F(0) = 0$). This is an APN function if and only if $n \geq 5$ is odd [9]. The Walsh coefficients are the elements $\equiv 0 \pmod{4}$ in the interval

$$[-2^{n/2+1} + 1, 2^{n/2+1} + 1],$$

see [14], and therefore:

On large Sidon sets

{	0,	1,	2,	4,	8,	16,	32,	64,
	128,	256,	423,	512,	732,	861,	1024,	1177,
	1228,	1335,	1350,	1481,	1594,	1934,	2048,	2415,
	2719,	2824,	3421,	3761,	3839,	3893,	3904,	4096,
	4233,	4446,	4726,	4834,	5013,	5199,	5322,	5366,
	5410,	5515,	5713,	6172,	6405,	6451,	6521,	6642,
	6672,	6862,	6888,	6914,	7358,	7681,	7909,	8040,
	8192,	8284,	8292,	8488,	8700,	8913,	9191,	9219,
	9398,	9817,	9875,	10262,	10378,	11071,	11153,	11647,
	11706,	11752,	11911,	11998,	12197,	12295,	12334,	12392,
	12484,	12555,	12646,	12927,	12965,	13110,	13682,	13858,
	14068,	14138,	14176,	14236,	14391,	14430,	14572,	14579,
	14808,	15005,	15008,	15205,	15238,	15583,	15843,	16008,
	16384,	16409,	16593,	16714,	17032,	17491,	17639,	17697,
	17741,	17942,	18157,	18201,	18261,	18342,	18423,	18531,
	18582,	18738,	18819,	18941,	19109,	19425,	19507,	19666,
	19728,	19779,	20100,	20225,	20384,	20565,	21605,	21933,
	21994,	22077,	22228,	22377,	22647,	22650,	22705,	23047,
	23486,	23495,	23582,	23680,	24332,	24346,	24443,	24444,
	24590,	24780,	24803,	25048,	25185,	25405,	25968,	25997,
	26016,	26553,	26949,	27003,	27092,	27310,	27613,	27821,
	28255,	28550,	28586,	29213,	29224,	29264,	29342,	29386,
	29429,	29463,	29688,	29786,	29926,	30412,	30663,	30900,
	31350,	31488,	31563,	31768,	31854,	31907,	32079,	32365 }

Table 1: Example of maximal Sidon set in \mathbb{F}_2^{15} of size 192.

Theorem 4.3. *The linearity of the inverse function equals to*

$$\lfloor 2^{n/2+1} + 1 \rfloor - (\lfloor 2^{n/2+1} + 1 \rfloor \bmod 4)$$

and leads to a Sidon set of size

$$2^{n-1} + \frac{1}{2} \left(\lfloor 2^{n/2+1} + 1 \rfloor - (\lfloor 2^{n/2+1} + 1 \rfloor \bmod 4) \right)$$

in \mathbb{F}_2^{2n-1} for odd $n \geq 5$.

Proof. The upper bound for the Walsh coefficients would give a linearity of

$$\lfloor 2^{n/2+1} + 1 \rfloor - (\lfloor 2^{n/2+1} + 1 \rfloor \bmod 4) \quad (2)$$

and the lower bound of

$$\begin{aligned} & | \lfloor -2^{n/2+1} + 1 \rfloor + 4 - (\lfloor -2^{n/2+1} + 1 \rfloor \bmod 4) | = \\ & \quad - \lfloor -2^{n/2+1} + 1 \rfloor - 4 + (\lfloor -2^{n/2+1} + 1 \rfloor \bmod 4). \end{aligned} \quad (3)$$

We define $\varepsilon = 2^{n/2+1} + 1 - \lfloor 2^{n/2+1} + 1 \rfloor$. Then

$$\lfloor -2^{n/2+1} + 1 \rfloor = -2^{n/2+1} + 1 - (1 - \varepsilon) = -2^{n/2+1} + \varepsilon.$$

Hence

$$\lfloor 2^{n/2+1} + 1 \rfloor + \lfloor -2^{n/2+1} + 1 \rfloor = 2^{n/2+1} + 1 - \varepsilon + -2^{n/2+1} + \varepsilon = 1 \quad (4)$$

and

$$(\lfloor 2^{n/2+1} + 1 \rfloor \bmod 4) \neq (\lfloor -2^{n/2+1} + 1 \rfloor \bmod 4). \quad (5)$$

Subtracting (3) from (2) and using (4) gives

$$5 - (\lfloor 2^{n/2+1} + 1 \rfloor \bmod 4) - (\lfloor -2^{n/2+1} + 1 \rfloor \bmod 4)$$

which is always ≥ 0 because of (5). This shows the linearity of the inverse. The existence of the Sidon set follows from Corollary 3.6. \square

Hence, the linearity is roughly $2^{n/2+1} = 2^{(n+2)/2}$, which is larger than the linearity of the almost bent functions (they have linearity $2^{(n+1)/2}$). This shows that the inverse function gives an improvement of the classical constructions. A few concrete values for the size of Sidon sets using the inverse function are contained in Table 2.

Let us finally discuss the Dobbertin functions [13]. It is an infinite family of APN functions $\mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$. The Walsh spectrum and the linearity is not known, but there is a conjecture about it, see [7]. The authors of that paper computed the Walsh spectrum for several cases ($n \leq 35$); in the range of our table Table 2, the cases $n = 5$ and $n = 10$ are important: The linearities are 12 ($n = 5$) and 80 ($n = 10$). If $n = 15$, the linearity is 576, a value which is larger than the linearity of the inverse function. But let us first note that the Dobbertin function $\mathbb{F}_2^{10} \rightarrow \mathbb{F}_2^{10}$ can be used to improve the best known lower bound on the size of Sidon sets in dimension $t = 19$, once more using Corollary 3.6:

Theorem 4.4. *There exists a Sidon set of size 552 in \mathbb{F}_2^{19} .*

Remark 4.5. *Dobbertin functions exist for all n divisible by 5. If the conjecture in [7] about the Walsh spectrum of these functions is true, then the general formula for the linearity of the Dobbertin function is $2^{3n/5} + 2^{2n/5}$ and leads to Sidon sets of size*

$$2^{n-1} + 2^{3n/5-1} + 2^{2n/5-1} \text{ in } \mathbb{F}_2^{2n-1}.$$

For the case n is divisible by 5, when the Dobbertin function is defined, this size is larger than the classical constructions.

5 Linear Codes

A (binary) *linear code* \mathcal{C} of length m and *dimension* k is a k -dimensional linear subspace \mathcal{C} of \mathbb{F}_2^m and $c \in \mathcal{C}$ is called a *code word* of \mathcal{C} . We consider all vectors to be row vectors. If the *minimum distance* of \mathcal{C} is d , that is the minimum number of non-zero entries of all non-zero code words of \mathcal{C} , then \mathcal{C} is called a $[m, k, d]$ -code. We call a $[m, m-t, d]$ -code, a linear code of length m , with t check bits and with minimum distance d . Such a $[m, m-t, d]$ -code \mathcal{C} can be defined as the kernel of a matrix \mathcal{H} with t rows, m columns and rank t which is called parity-check matrix of \mathcal{C} , i.e $\mathcal{C} = \{v \in \mathbb{F}_2^m : \mathcal{H} \cdot v^\top = 0\}$. The one-to-one correspondence between sets and linear codes states that a sum-free Sidon set M in \mathbb{F}_2^t with $|M| \geq t+1$ and $\dim\langle M \rangle = t$ (that is the dimension of the linear span of M) leads to a linear code \mathcal{C}_M of length $|M|$, with t check bits and with minimum distance ≥ 5 . Thereby the $|M|$ elements of M form the columns of the parity-check matrix of \mathcal{C}_M . Since every Sidon set can be transformed into a sum-free Sidon set by translation and removing one element, we obtain:

Proposition 5.1. *Every Sidon set M in \mathbb{F}_2^t leads to a $[m-1, m-t-1, \geq 5]$ -code.*

More details on the one-to-one correspondence can be found in [12]. In the following Theorem we summarise all the results of the previous section.

Theorem 5.2. (a) *There exists a $[191, 176, 5]$ -code.*

(b) *There exist $[m-1, m-2n-2, 5]$ -codes for*

$$m = 2^{n-1} + \frac{1}{2} (\lfloor 2^{n/2+1} + 1 \rfloor - (\lfloor 2^{n/2+1} + 1 \rfloor \bmod 4))$$

and odd $n \geq 5$.

(c) *There exist $[m-1, m-2n-2, 5]$ -codes for*

$$m = 2^{n-1} + 2^{3n/5-1} + 2^{2n/5-1}$$

and $n = 5, 10, 15, 20, 25, 30, 35$.

The $[191, 176, 5]$ -code from Theorem 5.2 (a) improves the best previously known length 151 of a linear code with 15 check bits and with minimum distance 5 (see [3]).

6 Summary and Table

In this work we constructed large Sidon sets in \mathbb{F}_2^{2n-1} in Section 4. We obtained these Sidon sets by intersecting the graph of a highly linear APN function $\mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ with affine hyperplanes.

The inverse function gives an infinite family of Sidon sets larger than the classical ones if n is odd. The known linearities of the Dobbertin functions give improvements.

In the case $n = 8$, sporadic examples of quadratic APN functions with high linearity 2^7 lead to Sidon sets of size 192 in \mathbb{F}_2^{15} , where the largest sets so far had size 152. If it is possible to find an infinite family of quadratic APN functions on \mathbb{F}_2^n with linearity 2^{n-1} we would have Sidon sets of size $3 \cdot 2^{n-2}$. This would give a considerable improvement of the best known Sidon size.

Recall that each of the new large Sidon set M in \mathbb{F}_2^t yields a linear code with t check bits, minimum distance 5, and a length not known so far (see Section 5).

Table 2 gives an overview about the maximum size of Sidon sets in small dimensions.

It is interesting that the connection between Sidon sets and APN functions which we discussed in this paper can also be used to improve the upper bound for the linearity of APN functions (Corollary 3.7).

Acknowledgement

We thank Claude Carlet and Gabor Nagy for useful comments on an early version of this paper. We are also grateful to Alexandr Polujan for fruitful discussions. Part of the research of the second author has been funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project No. 541511634 .

References

- [1] L. Babai and V. T. Sós, “Sidon sets in groups and induced subgraphs of Cayley graphs,” *European Journal of Combinatorics*, vol. 6, no. 2, pp. 101–114, 1985 (cited on pages 1, 2).

On large Sidon sets

	t	3	4	5	6	7	8	9	10	11	12	13	14	15
Best known upper bound		4	6	7	9	12	18	24	34	58	89	125	179	254
Best known size so far		4	6	7	9	12	18	24	34	48	66	82	129	152
Classical size		4	6	6	9	12	18	20	33	40	66	72	129	144
Dobbertin								22						
Inverse								22			74			
Quadratic, highest lin.						12			48					192
	t	16	17	18	19	20	21	22	23	24	25			
Best known upper bound		360	510	723	1022	1446	2046	2895	4094	5791	11583			
Best known size so far		258	272	513	544	1026	1056	2049	2112	4098	4160			
Classical size		258	272	513	544	1026	1056	2049	2112	4098	4160			
Dobbertin														552
Inverse			278					1068						4186
Quadratic, highest lin.														

Table 2: Sizes of Sidon sets in \mathbb{F}_2^t and related bounds and constructions.

- [2] C. Beierle and G. Leander, “New instances of quadratic APN functions,” *IEEE Transactions on Information Theory*, vol. 68, no. 1, pp. 670–678, 2022 (cited on pages 3, 8).
- [3] V. Bhargava, “The (151, 136) 10th-power residue code and its performance,” *Proceedings of the IEEE*, vol. 71, no. 5, pp. 683–685, 1983 (cited on page 12).
- [4] A. E. Brouwer and L. M. G. M. Tolhuizen, “A sharpening of the Johnson bound for binary linear codes and the nonexistence of linear codes with Preparata parameters,” *Designs, Codes and Cryptography*, vol. 3, no. 2, pp. 95–98, May 1993 (cited on page 2).
- [5] K. Browning, J. Dillon, R. Kibler, and M. McQuistan, “APN polynomials and related codes,” *Journal of Combinatorics, Information & System Sciences*, vol. 34, Jan. 2009 (cited on page 8).
- [6] L. Budaghyan, *Construction and analysis of cryptographic functions*. Springer, Cham, 2014, pp. viii+168 (cited on page 8).
- [7] L. Budaghyan, M. Calderini, C. Carlet, D. Davidova, and N. Kaleyski, “A note on the Walsh spectrum of Dobbertin APN functions,” in *Proceedings of SETA*, 2020 (cited on pages 10, 11).

- [8] C. Carlet, “Characterizations of the differential uniformity of vectorial functions by the Walsh transform,” *IEEE Transactions on Information Theory*, vol. 64, no. 9, pp. 6443–6453, 2018 (cited on page 7).
- [9] C. Carlet, *Boolean functions for cryptography and coding theory*. Cambridge University Press, New York, 2020, pp. xiv+562 (cited on pages 3, 7, 8).
- [10] C. Carlet and S. Mesnager, “On those multiplicative subgroups of $\mathbb{F}_{2^n}^*$ which are Sidon sets and/or sum-free sets,” *Journal of Algebraic Combinatorics*, vol. 55, no. 1, pp. 43–59, Feb. 2022 (cited on page 3).
- [11] C. Carlet and S. Picek, “On the exponents of APN power functions and Sidon sets, sum-free sets, and Dickson polynomials,” *Advances in Mathematics of Communications*, vol. 17, no. 6, pp. 1507–1525, 2023 (cited on pages 3, 4).
- [12] I. Czerwinski and A. Pott, “Sidon sets, sum-free sets and linear codes,” *Advances in Mathematics of Communications*, vol. 18, no. 2, pp. 549–566, 2024 (cited on pages 2, 11).
- [13] H. Dobbertin, “Almost perfect nonlinear power functions on $\text{GF}(2^n)$: A new case for n divisible by 5,” in *Finite Fields and Applications*, D. Jungnickel and H. Niederreiter, Eds., Berlin, Heidelberg: Springer Berlin Heidelberg, 2001, pp. 113–121 (cited on page 10).
- [14] G. Lachaud and J. Wolfmann, “The weights of the orthogonals of the extended quadratic binary Goppa codes,” *IEEE Transactions on Information Theory*, vol. 36, no. 3, pp. 686–692, 1990 (cited on page 8).
- [15] F. MacWilliams and N. Sloane, *The Theory of Error-Correcting Codes*, 2nd. North-holland Publishing Company, 1978 (cited on pages 2, 3).
- [16] G. P. Nagy, *Thin Sidon sets and the nonlinearity of vectorial Boolean functions*, 2022. arXiv: 2212.05887 [math.CO] (cited on pages 2, 3).
- [17] K. Nyberg and L. R. Knudsen, “Provable security against differential cryptanalysis,” in *Advances in Cryptology — CRYPTO’ 92*, E. F. Brickell, Ed., Berlin, Heidelberg: Springer Berlin Heidelberg, 1993, pp. 566–574 (cited on page 3).
- [18] A. Pott, “Almost perfect and planar functions,” *Designs, Codes and Cryptography*, vol. 78, no. 1, pp. 141–195, 2016 (cited on page 8).
- [19] S. Sidon, “Ein Satz über trigonometrische Polynome und seine Anwendung in der Theorie der Fourier-Reihen,” *Mathematische Annalen*, vol. 106, no. 1, pp. 536–539, Dec. 1932 (cited on page 1).
- [20] S. Sidon, “Über die Fourier Konstanten der Funktionen der Klasse L_p für $p > 1$,” *Acta Univ. Szeged Sect. Sci. Math*, vol. 7, pp. 175–176, 1935 (cited on page 1).

- [21] Y. Yu and L. Perrin, “Constructing more quadratic APN functions with the QAM method,” *Cryptogr. Commun.*, vol. 14, no. 6, pp. 1359–1369, 2022 (cited on page 8).
- [22] Y. Yu, M. Wang, and Y. Li, “A matrix approach for constructing quadratic APN functions,” *Des. Codes Cryptogr.*, vol. 73, no. 2, pp. 587–600, 2014 (cited on page 8).