

# SHARP BOUNDS FOR NEIGHBORHOOD DEGREE BASED INDICES OF GRAPHS

SANJU VAIDYA AND JEFF CHANG

ABSTRACT. In this paper, we will construct formulas and bounds for Neighborhood Degree-based indices of graphs and describe graphs that attain the bounds. Furthermore, we will establish a lower bound for the spectral radius of any graph.

## 1. INTRODUCTION

In the last forty years, many scientists have developed mathematical models for analyzing structures and properties of various chemical compounds. Graph theory plays a very important role in developing many types of models such as Quantitative Structure-Property Relationships (QSPR) models, and Quantitative Structure-Activity Relationships (QSAR) models ([3],[7], [11],[21]). In molecular graphs of chemical compounds, vertices correspond to atoms and edges correspond to the bonds between them. A topological index (connectivity index) is a type of a molecular descriptor that is based on the molecular graph of a chemical compound. In 1947 Harry Wiener introduced a topological index related to molecular branching. He correlated the indices with the boiling points of certain chemical compounds, alkanes. This inspired many mathematicians and chemists to develop more topological indices for molecular graphs.

In 1972, Zagreb indices were introduced by Gutman and Trijanastic [9]. They are based on the degrees of the vertices. They are very useful in modeling chemical and biological properties of chemical compounds, as shown by Klein, D. J. [11], Basak et al [3], and Todeschin et al [21]. Gutman et al [8] and Zhou et al [23] used Zagreb indices for analyzing bounds for Estrada index, which is based on the eigenvalues of the adjacency matrix.

Zagreb indices are generalized by many mathematicians and scientists. In 2019, Mondal et al [15] introduced the general neighborhood Zagreb index and proved its importance in predicting properties of chemical compounds. This index is based on the neighborhood degree of a vertex, which is the sum of the degrees of its adjacent vertices. In the last five years, many scientists used the neighborhood degree based indices to model various properties of chemical compounds which are used in medicines. For example, Mondal et al [17], investigated neighborhood degree based indices of some chemical compounds like remdesivir (GS-5734) which is useful for treatment of Covid-19 patients. In 2023, Balasubramanian and Chidambaram [2], used these indices for analysis of asthma drugs.

---

2010 *Mathematics Subject Classification.* Primary 05C07, 05C12, 05C35, 05C90.

The main research question is how to compute bounds and formulas of the indices stated above and find graphs which attain the bounds. In this paper, we will establish formulas for the general neighborhood Zagreb index. Additionally, we will find upper and lower bounds, and describe graphs which attain the bounds. Hu-Li-Shi-Xu-Gutman [10] investigated the zeroth-order general Randić index (which is the same as general Zagreb index) for molecular graphs in which the maximum vertex degree is at most 4. In Theorem 3.5 of this paper we will extend this further to the general neighborhood Zagreb index for graphs with maximum vertex degree more than 4.

In 2004, Yu-Lu-Tian [22] established a lower bound for the spectral radius of any graph, which is the largest eigenvalue of its adjacency matrix. They used the 2-degree of a vertex, which is the sum of the degrees of the adjacent vertices. So the 2-degree is the same as the neighborhood degree of a vertex. This is amazing! Using this and our lower bound for general neighborhood Zagreb index, we will find a lower bound for spectral radius of any graph in Section 4. Additionally, using generalization of Benakli, Halleck, and Kingan [4], we could define 2-distance neighborhood degree,  $\delta_2(v)$ , of any vertex  $v$  as the sum of the degrees of the vertices which are at distance 2 from the vertex  $v$ , which opens more research problems.

In Section 2 of this paper, we will review definitions of the first Zagreb index and general neighborhood Zagreb index. In Section 3 we will establish formulas of the general neighborhood Zagreb index for graphs. Additionally, we will find upper and lower bounds for the index and determine graphs which attain the bounds. In addition, we will establish sharp upper and lower bounds for the index for some special graphs. In Section 5 we will have discussion and conclusion.

## 2. REVIEW OF ZAGREB INDICES

We will use the notation and terminology introduced in Gutman et al [8] and Mondal et al [15].

In **molecular graphs** of chemical compounds, vertices correspond to individual atoms and edges correspond to the bonds between them. A **topological index** (connectivity index) is a type of a molecular descriptor that is based on the molecular graph of a chemical compound. The definition of the first Zagreb index, which is introduced by Gutman and Trijanastic [9], is as follows.

**Definition 2.1.** Let  $G$  be a graph with vertex set  $V(G)$  and  $d(u)$  be the degree of the vertex  $u \in V(G)$ . Then the first Zagreb index is defined as follows:

$$Z_G = M_1(G) = \sum_{u \in V(G)} d(u)^2.$$

In 2007 Gutman-Fortula-Marković-Glišić [8] proved that for a chemical tree (alkane) with  $n$  vertices  $Z_G = M_1(G) = 6n - 10 - 2n_2 - 2n_3$ , where  $n_i$  is the number of vertices of degree  $i$  for  $i = 2, 3$ . In Vaidya and Surendran [19], we established formulas of the first Zagreb index for molecular graphs of cycloalkanes, alkenes, and alkynes.

Zagreb indices have been generalized and studied for more than 30 years. In 2019, Mondal et al [15] introduced the following general neighborhood Zagreb index.

**Definition 2.2.** Let  $G$  be a graph with vertex set  $V(G)$ . For a vertex  $u \in V(G)$ , let  $\delta_G(u)$  be its neighborhood degree, which is the sum of the degrees of the vertices which are adjacent to the vertex  $u$ . Let  $\alpha$  be a nonzero real number such that  $\alpha \neq 1$ . Then the general neighborhood Zagreb index is defined as follows:

$$NM_\alpha(G) = \sum_{u \in V(G)} \delta_G(u)^\alpha.$$

### 3. FORMULAS AND BOUNDS FOR NEIGHBORHOOD DEGREE BASED INDICES

In this section we will establish some formulas for the general neighborhood Zagreb index. We will then find sharp upper and lower bounds for it using these formulas. First, we need the following Lemma which is proved in Vaidya-Chang [18].

**Lemma 3.1.** *Let  $p$  and  $q$  be any positive integers such that  $p < q$ . Let  $\alpha \in \mathbb{R}$  and  $\alpha \notin \{0, 1\}$ . Then we have the following.*

- (1) *If the real number  $\alpha < 0$  or  $\alpha > 1$ , then  $(p+i)^\alpha - p^\alpha - i\left(\frac{q^\alpha - p^\alpha}{q-p}\right) \leq 0$  for  $1 \leq i \leq q-p-1$  and if  $0 < \alpha < 1$ , then  $(p+i)^\alpha - p^\alpha - i\left(\frac{q^\alpha - p^\alpha}{q-p}\right) \geq 0$  for  $1 \leq i \leq q-p-1$ .*
- (2) *if the real number  $\alpha < 0$  or  $\alpha > 1$ , then  $(p+i)^\alpha - p^\alpha - i((p+1)^\alpha - p^\alpha) \geq 0$  for  $2 \leq i \leq q-p$  and for  $0 < \alpha < 1$ ,  $(p+i)^\alpha - p^\alpha - i((p+1)^\alpha - p^\alpha) \leq 0$  for  $2 \leq i \leq q-p$ .*

**Theorem 3.2.** *Let  $G$  be a graph with  $n$  vertices and  $m$  edges where  $n \geq 3$ . Let  $n_i$  denote the number of vertices of neighborhood degree  $i$ . Let  $\delta$  and  $\Delta$  respectively denote the minimum and the maximum neighborhood degree. Assume that  $\delta \neq \Delta$ . Let  $\alpha \in \mathbb{R}$  and  $\alpha \notin \{0, 1\}$ . Further denote  $s_\alpha = \frac{\Delta^\alpha - \delta^\alpha}{\Delta - \delta}$ . Then we have the following.*

- (1) *The general neighborhood Zagreb index*

$$NM_\alpha(G) = n\delta^\alpha + (M_1 - n\delta)s_\alpha + \sum_{i=1}^{\Delta-\delta-1} n_{\delta+i} [(\delta+i)^\alpha - \delta^\alpha - is_\alpha].$$

*Additionally, if  $\alpha < 0$  or  $\alpha > 1$  the coefficients  $(\delta+i)^\alpha - \delta^\alpha - is_\alpha \leq 0$  for  $1 \leq i \leq \Delta - \delta - 1$  and for  $0 < \alpha < 1$ , the coefficients  $(\delta+i)^\alpha - \delta^\alpha - is_\alpha \geq 0$  for  $1 \leq i \leq \Delta - \delta - 1$ .*

- (2) *The general neighborhood Zagreb index*

$$NM_\alpha(G) = n\delta^\alpha + (M_1 - n\delta)((\delta+1)^\alpha - \delta^\alpha) + \sum_{i=2}^{\Delta-\delta} n_{\delta+i} [(\delta+i)^\alpha - \delta^\alpha - i((\delta+1)^\alpha - \delta^\alpha)].$$

*Additionally, if  $\alpha < 0$  or  $\alpha > 1$ , the coefficients  $(\delta+i)^\alpha - \delta^\alpha - i((\delta+1)^\alpha - \delta^\alpha) \geq 0$  for  $2 \leq i \leq \Delta - \delta$  and for  $0 < \alpha < 1$ , the coefficients  $(\delta+i)^\alpha - \delta^\alpha - i((\delta+1)^\alpha - \delta^\alpha) \leq 0$  for  $2 \leq i \leq \Delta - \delta$ .*

*Proof.* We have  $\sum_{i=\delta}^{\Delta} in_i = M_1$  by part (i) of Lemma 1 of [16]. We also have  $n = \sum_{i=\delta}^{\Delta} n_i$ . Solving these equations, we will rewrite  $n_{\Delta}$  as follows.

$$n_{\Delta} = \frac{M_1 - n\delta - \sum_{i=\delta+1}^{\Delta-1} ((i - \delta)n_i)}{\Delta - \delta}.$$

Write  $NM_{\alpha}(G)$  in terms of  $n_i$ ,  $\delta \leq i \leq \Delta$ , and substitute  $n_{\delta}$  with  $n - \sum_{i=\delta+1}^{\Delta} n_i$ , we obtain

$$NM_{\alpha}(G) = \sum_{i=\delta}^{\Delta} i^{\alpha} n_i = \delta^{\alpha} (n - \sum_{i=\delta+1}^{\Delta} n_i) + \sum_{i=\delta+1}^{\Delta} i^{\alpha} n_i = n\delta^{\alpha} + \sum_{i=\delta+1}^{\Delta} n_i (i^{\alpha} - \delta^{\alpha}).$$

Further substitute  $n_{\Delta}$  into the sum and using Lemma 3.1, we get the result in part (1). Using the identities  $n_{\delta} + n_{\delta+1} = n - \sum_{i=\delta+2}^{\Delta} n_i$  and  $\delta n_{\delta} + (\delta + 1)n_{\delta+1} = M_1 - \sum_{i=\delta+2}^{\Delta} in_i$ , we solve for  $n_{\delta}$  and  $n_{\delta+1}$ . Substituting  $n_{\delta}$  and  $n_{\delta+1}$  into  $NM_{\alpha}(G) = \sum_{i=\delta}^{\Delta} i^{\alpha} n_i$ , and using Lemma 3.1 again, we get the result in part (2).  $\square$

The following corollary gives upper bounds or lower bounds for the general Zagreb index, depending on the values of  $\alpha$ . Both the bounds are sharp.

**Corollary 3.3.** *Let  $G$  be a graph with  $n$  vertices and  $m$  edges where  $n \geq 3$ . Let  $n_i$  denote the number of vertices of neighborhood degree  $i$ . Let  $\delta$  and  $\Delta$  respectively denote the minimum and the maximum neighborhood degree. Assume that  $\delta \neq \Delta$ . Let  $\alpha \in \mathbb{R}$  and  $\alpha \notin \{0, 1\}$ . Further denote  $s_{\alpha} = \frac{\Delta^{\alpha} - \delta^{\alpha}}{\Delta - \delta}$ . Then we have the following.*

- (1) *If  $\alpha < 0$  or  $\alpha > 1$ , the upper bound for the general neighborhood Zagreb index is  $NM_{\alpha}(G) \leq n\delta^{\alpha} + (M_1 - n\delta)s_{\alpha}$ .*
- (2) *If  $0 < \alpha < 1$ , it becomes the lower bound for the general first Zagreb index, that is,  $NM_{\alpha}(G) \geq n\delta^{\alpha} + (M_1 - n\delta)s_{\alpha}$ .*  
*Moreover, for parts (1) and (2), the equality occurs for any graph with vertices of neighborhood degrees  $\delta$  and  $\Delta$ .*
- (3) *If  $\alpha < 0$  or  $\alpha > 1$ , then the lower bound for the general neighborhood Zagreb index is  $NM_{\alpha}(G) \geq n\delta^{\alpha} + (M_1 - n\delta)((\delta + 1)^{\alpha} - \delta^{\alpha}) + n_{\Delta}(\Delta^{\alpha} - \delta^{\alpha} - (\Delta - \delta)((\delta + 1)^{\alpha} - \delta^{\alpha}))$ .*
- (4) *If  $0 < \alpha < 1$ , then the upper bound for the general neighborhood Zagreb index is  $NM_{\alpha}(G) \leq n\delta^{\alpha} + (M_1 - n\delta)((\delta + 1)^{\alpha} - \delta^{\alpha}) + n_{\Delta}(\Delta^{\alpha} - \delta^{\alpha} - (\Delta - \delta)((\delta + 1)^{\alpha} - \delta^{\alpha}))$ .*

*Moreover, for parts (3) and (4), the equality occurs if the graph  $G$  is a path.*

*Proof.* Follows from Theorem 3.2.  $\square$

**Example 3.4.** Here is an example. In Figure 1,  $\delta = 4$ ,  $\Delta = 10$ ,  $m = n = 12$  and all vertices have either neighborhood degree 4 or 10. It attains the bounds given in parts (1) and (2) of Corollary 3.3.

In Theorem (2.2) of [10], Hu-Li-Shi-Xu-Gutman established formulas for the maximum and the minimum zeroth-order general Randic index (which is same as general Zagreb index) for  $(n, m)$  molecular graphs if  $2m - n$  is congruent to 0, 1, or 2 modulo 3 and the maximum vertex degree is 4. In the following Theorem, we will extend this and find sharp




FIGURE 1.

bounds for the general neighborhood Zagreb index if  $M_1 - n\delta$  is congruent to any integer  $r$  modulo  $\Delta - \delta$  where  $0 \leq r \leq \Delta - \delta - 1$  and the maximum neighborhood degree  $\Delta \geq 3$ .

**Theorem 3.5.** *Let  $G$  be a graph with  $n$  vertices and  $m$  edges where  $n \geq 3$ . Let  $n_i$  denote the number of vertices of neighborhood degree  $i$ . Let  $\delta$  and  $\Delta$  respectively denote the minimum and the maximum neighborhood degree. Assume that  $\Delta - \delta \geq 2$  and  $M_1 - n\delta = q(\Delta - \delta) + r$ , where  $q$  is a positive integer and  $r$  is an integer such that  $0 \leq r \leq \Delta - \delta - 1$ . Let  $\alpha \in \mathbb{R}$  and  $\alpha \notin \{0, 1\}$ . Further denote  $s_\alpha = \frac{\Delta^\alpha - \delta^\alpha}{\Delta - \delta}$ . Then we have the following.*

- (1) *If  $r = 0$  and  $n_\Delta = q$ , then the graph  $G$  is a bi-neighborhood degree graph with vertices of neighborhood degree  $\delta$  and  $\Delta$ .*
- (2) *If  $r \geq 1$  and  $n_\Delta = q$ , then  $n_i = 0$  for  $\delta + r + 1 \leq i \leq \Delta - 1$  and  $n_{\delta+r} \leq 1$ .*
- (3) *If  $r \geq 1$  and  $\alpha < 0$  or  $\alpha > 1$  and  $n_{\delta+r} \neq 0$ , then the upper bound for the general neighborhood Zagreb index is  $NM_\alpha(G) \leq n\delta^\alpha + (M_1 - n\delta)s_\alpha + (\delta + r)^\alpha - \delta^\alpha - rs_\alpha$ .*
- (4) *If  $r \geq 1$  and  $0 < \alpha < 1$  and  $n_{\delta+r} \neq 0$ , it becomes the lower bound for the general neighborhood Zagreb index, that is,  $NM_\alpha(G) \geq n\delta^\alpha + (M_1 - n\delta)s_\alpha + (\delta + r)^\alpha - \delta^\alpha - rs_\alpha$ .*

*Additionally, in both parts (3) and (4) the equality occurs if the graph  $G$  has  $q$  vertices of neighborhood degree  $\Delta$ , a unique vertex of neighborhood degree  $\delta + r$ , and  $n - q - 1$  vertices of neighborhood degree  $\delta$ .*

*Proof.* We have  $\sum_{i=\delta}^{\Delta} in_i = M_1$  by part (i) of Lemma 1 of [16]. We also have  $n = \sum_{i=\delta}^{\Delta} n_i$ . By solving the equations, we get  $\sum_{i=\delta+1}^{\Delta} (i - \delta)n_i = M_1 - n\delta = q(\Delta - \delta) + r$ . If  $r = 0$  and  $n_\Delta = q$ , then  $n_i = 0$  for  $\delta + 1 \leq i \leq \Delta - 1$ . So we get (1). Also if  $r \geq 1$  and  $n_\Delta = q$ , then  $n_i = 0$  for  $\delta + r + 1 \leq i \leq \Delta - 1$  and  $n_{\delta+r} \leq 1$ . So we get (2). To prove (3), let  $r \geq 1$  and  $n_{\delta+r} \neq 0$ . By part (1) of Theorem 3.2 we have the general neighborhood Zagreb index  $NM_\alpha(G) = n\delta^\alpha + (M_1 - n\delta)s_\alpha + \sum_{i=1}^{\Delta-\delta-1} n_{\delta+i} [(\delta+i)^\alpha - \delta^\alpha - is_\alpha]$ . If  $\alpha < 0$  or  $\alpha > 1$ , then  $(\delta+i)^\alpha - \delta^\alpha - is_\alpha \leq 0$  for  $1 \leq i \leq \Delta - \delta - 1$ . So  $NM_\alpha(G) \leq n\delta^\alpha + (M_1 - n\delta)s_\alpha + n_{\delta+r} [(\delta+r)^\alpha - \delta^\alpha - (r)s_\alpha]$ .

Since  $n_{\delta+r} \neq 0$ , the result in part (3) follows. Similarly, the result in part (4) follows since for  $0 < \alpha < 1$ ,  $(\delta+i)^\alpha - \delta^\alpha - is_\alpha \geq 0$  for  $1 \leq i \leq \Delta - \delta - 1$ . In addition, by part (1) of Theorem (3.2), for both parts (3) and (4) the equality occurs if the graph  $G$  has  $q$  vertices




FIGURE 2.

of neighborhood degree  $\Delta$ , a unique vertex of neighborhood degree  $\delta + r$ , and  $n - q - 1$  vertices of neighborhood degree  $\delta$ .  $\square$

**Example 3.6.** For the graph in Figure 2, we have  $n = 5$ ,  $\delta = 3$ ,  $\Delta = 5$  and  $M_1 - n\delta = 7 = 3(\Delta - \delta) + 1$ . The graph has  $q = 3$  vertices of neighborhood degree  $\Delta = 5$ , a unique vertex of neighborhood degree  $\delta + r = 4$ , and the remaining  $n - q - 1$  vertices of neighborhood degree  $\delta = 3$ . It attains the bounds in parts (3) and (4) of Theorem 3.5.

#### 4. SPECTRAL RADIUS, NEIGHBORHOOD DEGREE, AND ITS GENERALIZATION

In 2004, Yu-Lu-Tian [22] established a lower bound for the spectral radius,  $\rho(G)$  of any graph  $G$ , which is the largest eigenvalue of its adjacency matrix. They used the 2-degree of a vertex, which is the sum of the degrees of the adjacent vertices. So the 2-degree is the same as the neighborhood degree of a vertex. They proved that the  $\rho^2 \geq \frac{NM_2(G)}{M_1}$ . Using this and part (2) of Theorem 3.2, we get the following Theorem.

**Theorem 4.1.** *Let  $G$  be a graph with  $n$  vertices and  $m$  edges where  $n \geq 3$ . Let  $\delta$  denote the minimum neighborhood degree. Then the spectral radius,  $\rho^2 \geq \frac{M_1(2\delta+1) - n\delta^2 - n\delta}{M_1}$ .*

*Proof.* Follows from Theorem 4 of Yu-Lu-Tian [22] and part 2 of Theorem 3.2.  $\square$

By using generalization of [4], we could define 2-distance neighborhood degree,  $\delta_2(v)$ , of any vertex  $v$  as the sum of the degrees of the vertices which are at distance 2 from the vertex  $v$ . Further we could define the general 2-distance neighborhood degree index as  $NM2_\alpha(G) = \sum_{u \in V(G)} \delta_2(u)^\alpha$ . In the following Theorem, we will establish some formulas for the 2-distance neighborhood degree index if the diameter of the graph is 2.

**Theorem 4.2.** *Let  $G$  be a graph of diameter 2 with  $n$  vertices and  $m$  edges where  $n \geq 3$ . Let  $n_i$  denote the number of vertices of 2-distance neighborhood degree  $i$ . Let  $d$  and  $D$  respectively denote the minimum and the maximum 2-distance neighborhood degree. Assume that  $d \neq 0$  and  $d \neq D$ . Let  $\alpha \in \mathbb{R}$  and  $\alpha \notin \{0, 1\}$ . Further denote  $S_\alpha = \frac{D^\alpha - d^\alpha}{D - d}$ . Then we have the following.*

(1) *The 2-distance neighborhood degree index*

$$NM2_\alpha(G) = nd^\alpha + (2m(n-1) - M_1 - nd)S_\alpha + \sum_{i=1}^{D-d-1} n_{d+i} [(d+i)^\alpha - d^\alpha - iS_\alpha].$$

(2) *The 2-distance neighborhood degree index*

$$NM2_{\alpha}(G) = nd^{\alpha} + (2m(n-1) - M_1 - nd)((d+1)^{\alpha} - d^{\alpha}) + \sum_{i=2}^{D-d} n_{d+i} [(d+i)^{\alpha} - d^{\alpha} - i((d+1)^{\alpha} - d^{\alpha})].$$

*Proof.* We have  $n = \sum_{i=d}^D n_i$  and  $\sum_{i=d}^D i n_i = 2m(n-1) - M_1$ . Solving these equations for  $n_d$  and  $n_D$  we get (1). Additionally, by solving the equations for  $n_d$  and  $n_{d+1}$ , we get (2).  $\square$

## 5. DISCUSSION AND CONCLUSION

In conclusion, we established formulas of the general neighborhood Zagreb index for graphs. Additionally, we found upper and lower bounds for the general neighborhood Zagreb index and determine graphs which attain the bounds. In addition, we established sharp upper and lower bounds for the general neighborhood Zagreb index for some special graphs. We also found lower bound for the spectral radius. Additionally, we discussed generalization of the neighborhood degree which leads to more research problems.

Since there are more than 100 topological indices, there are many open problems of finding bounds and formulas for them. They play a crucial role in modeling various properties of chemical compounds. This process of mathematical modeling is very useful in real life because it reduces cost and saves time in developing medicines. It is fascinating to see that neighborhood degree based indices are useful in finding bounds for spectral radius as well as predicting physico-chemical properties of chemical compounds.

## REFERENCES

1. Abdo, H., Dimitrov, D., & Gutman, I. (2017). On extremal trees with respect to the F-index. *Kuwait Journal of Science*, 44(3).
2. Balasubramaniyan, D., & Chidambaram, N. (2023). On some neighbourhood degree-based topological indices with QSPR analysis of asthma drugs. *The European Physical Journal Plus*, 138(9), 823.
3. Basak, S. C., Gute, B. D., & Grunwald, G. D. (1996). Estimation of the normal boiling points of haloalkanes using molecular similarity. *Croatica chemica acta*, 69(3), 1159-1173.
4. Benakli, N., Halleck, E., & Kingan, S. R. (2019). A characterization of 2-neighborhood degree list of diameter 2 graphs. *arXiv preprint arXiv:1909.08722*.
5. Došlić, T. (2008). Vertex-weighted Wiener polynomials for composite graphs. *Ars Math. Contemp*, 1(1), 66-80.
6. Furtula, B., & Gutman, I. (2015). A forgotten topological index. *Journal of mathematical chemistry*, 53(4), 1184-1190.
7. Gao, W., Farahani, M. R., & Shi, L. (2016). Forgotten topological index of some drug structures. *Acta medica mediterranea*, 32(1), 579-585.
8. Gutman, I., Furtula, B., Marković, V., & Glišić, B. (2007). Alkanes with greatest Estrada index. *Zeitschrift für Naturforschung A*, 62(9), 495-498.
9. Gutman, I., & Trinajstić, N. (1972). Graph theory and molecular orbitals. Total  $\varphi$ -electron energy of alternant hydrocarbons. *Chemical physics letters*, 17(4), 535-538.

10. Hu, Y., Li, X., Shi, Y., Xu, T., & Gutman, I. (2005). On molecular graphs with smallest and greatest zeroth-order general Randić index. *MATCH Commun. Math. Comput. Chem*, 54(2), 425-434.
11. Klein, D. J. (2002). Topological Indices and Related Descriptors in QSAR and QSPR Edited by James Devillers & Alexandru T. Balaban. Gordon and Breach Science Publishers: Singapore. 1999. 811 pp. 90-5699-239-2. *Journal of Chemical Information and Computer Sciences*, 42(6), 1507-1507.
12. Li, X., & Zhao, H. (2004). Trees with the first three smallest and largest generalized topological indices. *MATCH Commun. Math. Comput. Chem*, 50, 57-62.
13. Li, X., & Zheng, J. (2005). A unified approach to the extremal trees for different indices. *MATCH Commun. Math. Comput. Chem*, 54(1), 195-208.
14. Mansour, T., & Song, C. (2012). The  $a$  and  $(a,b)$ -Analogues of Zagreb Indices and Coindices of Graphs. *International Journal of Combinatorics*, 2012(1), 909285.
15. Mondal, S., De, N., & Pal, A. (2019). On some general neighborhood degree based topological indices. *International Journal of Applied Mathematics*, 32(6), 1037.
16. Mondal, S., De, N., & Pal, A. (2021). On neighborhood Zagreb index of product graphs. *Journal of molecular structure*, 1223, 129210.
17. Mondal, S., De, N., & Pal, A. (2022). Topological indices of some chemical structures applied for the treatment of COVID-19 patients. *Polycyclic Aromatic Compounds*, 42(4), 1220-1234.
18. Vaidya, S., & Chang, J. (2024). Sharp Bounds for Generalized Zagreb Indices of Graphs. *arXiv preprint arXiv:2409.06081*.
19. Vaidya, S., & Surendran, G. (2016). Bounds for topological indices of molecular graphs. *Congr. Numer.* 227, 51-64.
20. Soleimani, N., Nikmehr, M. J., & Tavallae, H. A. (2015). Computation of the different topological indices of nanostructures. *Journal of the national science foundation of Sri Lanka*, 43(2).
21. Todeschini, R., & Consonni, V. (2008). Handbook of molecular descriptors. John Wiley & Sons.
22. Yu, A., Lu, M., & Tian, F. (2004). On the spectral radius of graphs. *Linear algebra and its applications*, 387, 41-49.
23. Zhou, B. (2008). On Estrada index. *MATCH Commun. Math. Comput. Chem*, 60(2), 485-492.

MATH & CS DEPARTMENT, MERCY UNIVERSITY, 555 BROADWAY, DOBBS FERRY, NY 10522  
Email address: SVaidya@mercy.edu

MATH & CS DEPARTMENT, MERCY UNIVERSITY, 555 BROADWAY, DOBBS FERRY, NY 10522  
Email address: cchang4@mercy.edu