PENROSE TRANSFORMATION ON FLAG DOMAINS

KEFENG LIU AND YANG SHEN

ABSTRACT. Building on our recent work we construct the Penrose transformations of the cohomology groups of homogenous line bundles on flag domains $D = G_{\mathbb{R}}/T$ with $G_{\mathbb{R}}$ of Hermitian type. We give sufficient conditions for the injectivity of the Penrose transformations, and conditions under which the Penrose transformation of the automorphic cohomology groups on compact quotients of flag domains is an isomorphism. Finally we prove that the higher automorphic cohomology groups of certain homogeneous line bundles are isomorphic to the groups of automorphic forms on the Hermitian symmetric domain.

CONTENTS

0. INTRODUCTION

As noted in the preface of [\[1\]](#page-32-1), the Penrose transformation was originally introduced by Roger Penrose in the 1960s as an isomorphism between a sheaf cohomology group on a region of projective space and the solutions of a zerorest-mass field equation on a region of spacetime in mathematical physics. Since its introduction, numerous generalizations of the Penrose transformation have been studied in both mathematics and physics.

In recent works [\[4\]](#page-32-2), [\[5\]](#page-32-3), and [\[6\]](#page-32-4) by Carayol, as well as in [\[11\]](#page-33-0) and [\[12\]](#page-33-1) by Green, Griffiths, and Kerr, the Penrose transformation has been studied on flag domains for $SU(2, 1)$ and $Sp(4)$. In the context of Hodge theory and representation theory, the Penrose transformation on flag domains investigates the relationship between the cohomology groups of homogeneous line bundles on a classical flag domain D' and those on the non-classical flag domain D that is diffeomorphic to D' . Additionally, it also studies the relations between the corresponding automorphic cohomology groups on the compact quotients X and X' of D and D' respectively.

In our recent work [\[20\]](#page-33-2), we proved that any non-classical flag domain $D =$ $G_\mathbb{R}/V$ with $G_\mathbb{R}$ of Hermitian type is diffeomorphic to a classical flag domain $D',$

where $G_{\mathbb{R}}$ is a semisimple real Lie group and $V \subset G_{\mathbb{R}}$ is a compact subgroup containing a compact Cartan subgroup T . This represents the most general result on this topic, since any real Lie group $G_{\mathbb{R}}$ acting on a classical flag domain must be of Hermitian type. Furthermore, in [\[20\]](#page-33-2), we also showed that the compact, smooth quotient X of D admits no $\partial \overline{\partial}$ -structure compatible with the complex structure induced from D. This result is achieved by proving a conjecture of Green–Griffiths–Kerr concerning the vanishing of the automorphic cohomology groups of any non-trivial homogeneous vector bundles on X.

In contrast, it was proved in [\[13\]](#page-33-3) that the compact quotient X' of the classical flag domain D' is a projective manifold, and thus, it has an arithmetic structure on its automorphic cohomology group. Once the Penrose transformation is shown to be an isomorphism, the automorphic cohomology group on X will inherit the arithmetic structure from that on X' .

Therefore, in conjunction with our work [\[20\]](#page-33-2), the Penrose transformation on flag domains reveals unexpected properties. For further applications of the Penrose transformation, we refer the reader to [\[11\]](#page-33-0), Lecture 9, and [\[12\]](#page-33-1).

In this paper, we develop the Penrose transformation on general non-classical flag domains of the form $D = G_{\mathbb{R}}/T$ with $G_{\mathbb{R}}$ of Hermitian type, within the framework developed in [\[11\]](#page-33-0) and [\[12\]](#page-33-1). Our proof is based on overcoming several technical difficulties, with methods from deformation theory, geometric realizations of the representations of compact and non-compact real Lie groups, and geometry of homogenous manifolds.

Now let us introduce the main results of this paper.

Let D be a non-classical flag domain, and D' be the classical one which is diffeomorphic to D . There is an associated space W , called the corresponding space for D and D' . Moreover W is Stein with the basic diagram

where the fibers of π and π' are contractible. Then one can apply the EGW Theorem in [\[8\]](#page-32-5) to the homogeneous line bundles L_μ on D and $L_{\mu'}$ on D' to get

(2)
$$
H^*(D, L_{\mu}) = H^*_{DR}(\mathcal{W}, \Omega^{\bullet}_{\pi}(L_{\mu})),
$$

(3)
$$
H^*(D', L_{\mu'}) = H^*_{DR}(\mathcal{W}, \Omega^{\bullet}_{\pi'}(L_{\mu'})),
$$

where

$$
\Omega_{\pi}^{\bullet}(L_{\mu}) = \Omega_{\mathcal{W}}^{\bullet}/\pi^{*}\Omega_{D}^{\bullet} \otimes_{\mathcal{O}_{\mathcal{W}}} \pi^{*}L_{\mu}
$$

with the induced differential d_{π} , and the de Rham cohomology groups

$$
H^*_{DR}(\mathcal{W}, \Omega^\bullet_\pi(L_\mu))
$$

are the cohomology groups of global sections

 $\Gamma(\mathcal{W}, \Omega_{\pi}^{\bullet}(L_{\mu}); d_{\pi}).$

The notations for π' are similar.

From [\(2\)](#page-1-0) and [\(3\)](#page-1-0), the cohomology groups $H^*(D, L_\mu)$ and $H^*(D', L_{\mu'})$ can be related on W. More precisely we have the following definition of Penrose transformation on flag domains.

Lemma-Definition 0.1. Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of *Hermitian type and* D′ *be the classical flag domain that is diffeomorphic to* D*. The complex structures of* D *and* D′ *are given by*

$$
\mathcal{T}^{1,0}_o D \cong \mathfrak{n}_- = \mathfrak{k}_- \oplus \mathfrak{p}_-^1 \oplus \mathfrak{p}_-^2,
$$

$$
\mathcal{T}^{1,0}_o D' \cong \mathfrak{n}'_- = \mathfrak{k}_- \oplus \mathfrak{p}_+^1 \oplus \mathfrak{p}_-^2
$$

which only differ in the directions $\mathfrak{p}^1_\pm \subset \mathfrak{p}$.

Let μ and μ' be two weights such that $\mu + \rho = \mu' + \rho'$, where ρ (ρ' resp.) denotes *half of the sum of positive roots corresponding to* n[−] *(*n ′ [−] *resp.). Then the Penrose transformation*

$$
\mathcal{P}: H^0(D', L_{\mu'}) \to H^q(D, L_{\mu})
$$

is defined by

(4)
$$
H^{0}(D', L_{\mu'}) \xrightarrow{\mathcal{P}} H^{q}(D, L_{\mu})
$$

$$
\downarrow \cong \qquad \qquad \downarrow \cong
$$

$$
H^{0}_{DR}(\mathcal{W}, \Omega^{\bullet}_{\pi'}(L_{\mu'})) \xrightarrow{\omega^{\text{nc},1}} H^{q}_{DR}(\mathcal{W}, \Omega^{\bullet}_{\pi}(L_{\mu})),
$$

where $q = \dim_{\mathbb{C}} \mathfrak{p}^1_+$ and $\omega^{\text{nc},1} \in \wedge^q(\mathfrak{p}^1_+)^*$, which is considered as a left-invariant $differential\ q\hbox{-}form\ with\ values\ in\ L_{-2\rho_{\rm nc,1}}$ for $2\rho_{\rm nc,1}=\sum_{\beta\in\Delta^{{\rm nc},1}_+}\beta$ on $\mathcal W.$

Please see Section [1](#page-6-0) for the notations of the Lie algebras and the description of the complex structures of flag domains in terms of Lie algebras.

Let Δ be the root system for the Lie algebra of the complexification $G_{\mathbb{C}}$ of $G_{\mathbb{R}}$, and $\Delta_+^c, \Delta_+^{nc,1}, \Delta_+^{nc,2}$ be the sets of roots corresponding to $\mathfrak{k}_-, \mathfrak{p}_-^1, \mathfrak{p}_-^2$ respectively. The following theorem is the main result of this paper.

Theorem 0.2 (Theorem [3.5\)](#page-20-0)**.** *Let the notations and assumptions be as Lemma-*Definition [0.1.](#page-2-0) If for any $\beta \in \Delta_+^{\rm nc,1}$ there exists an $\alpha \in \Delta_+^{\rm c}$ such that

(5)
$$
\left(\alpha, \mu + \sum_{\beta' \in \Delta_{+}^{\mathrm{nc},1}} \beta' - \beta\right) = (\alpha, \mu' - \beta) < 0,
$$

then the Penrose transformation

$$
\mathcal{P}: H^0(D', L_{\mu'}) \to H^q(D, L_{\mu})
$$

given by [\(4\)](#page-2-1) *is injective.*

The main idea of proving Theorem [0.2](#page-2-2) can be sketched as follows. Let $[F]$ be the corresponding cohomological class of $F \in \Gamma(W, \Omega^{\bullet}_{\pi'}(L_{\mu'}))$, such that $[F\omega^{\text{nc},1}] = 0$. This means that there exists $\Psi \in \Gamma(W, \Omega^{q-1}_\pi(L_\mu))$ such that $F\omega^{\text{nc},1} =$ $d_{\pi}\Psi$. We need to show that $F=0$.

In the special cases of $SU(2, 1)$ and $Sp(4)$ where $q = 1$ and Ψ is a global section of L_{μ} on W, there is a natural space β and a holomorphic submersion $W \to \beta$, so that Ψ descends to a global section in $\Gamma(\mathcal{J}, L_\mu)$. Moreover, $\mathcal J$ is covered by the cycles isomorphic to $\mathbb{P}^1.$ Hence, for the weight μ in a certain range, one has the vanishing theorem

$$
\Gamma(\mathbb{P}^1, L_\mu|_{\mathbb{P}^1}) = 0,
$$

and hence $\Psi \in \Gamma(W, L_{\mu}) = 0$. This proves that $F\omega^{nc,1} = d_{\pi}\Psi = 0$, and hence $F = 0$. Please see [\[11\]](#page-33-0), Lecture 8, and [\[12\]](#page-33-1) for details.

In contrast to the cases of $SU(2,1)$ and $Sp(4)$, we generally have no similar space ∂ and $q = \#\Delta^{nc,1}_+ \geq 2$, which means that Ψ may have differential forms from \mathfrak{k}^*_+ and $(\mathfrak{p}^2_+)^*$. Instead, we turn to study the incidence space

$$
\mathcal{I} = \{(x, u) \in D \times \mathcal{U} : x \in Z_u\}
$$

with a holomorphic submersion $W \rightarrow J$ and prove the vanishing theorem on an open subset of $\mathcal{I}^{\circ} \subset \mathcal{I}$ in Theorem [2.7.](#page-14-0) Here, \mathcal{U} is the cycle space for the non-classical flag domain D. In the proof of Theorem [2.7,](#page-14-0) we use crucially the upper semi-continuity of the dimensions of a differential family of vector bundles proved by Kodaira and Spencer in [\[19\]](#page-33-4).

Let Ψ' be the projection of Ψ onto $\wedge^{q-1}(\mathfrak{p}^1_+)^*$. In the proof of Theorem [3.5,](#page-20-0) we use the locally trivial fibration structure $B_K/T_C \times V_\nu \times U$ of W, where $\{V_\nu \subset V_\nu\}$ $K_{\mathbb{C}}/B_K$ _{$\}_{\nu}$} is an open cover on which the homogenous vector bundles are trivialized and $U \subset \mathcal{U}$ is an open subset containing the base cycle $u_0 \simeq Z_0$ of D. Then we define Ξ from Ψ' such that Ξ is constant on $B_K/T_{\mathbb C}$, which is the fiber of $W \to \mathcal{I}$. Hence Ξ descends to a section on an open subset $\mathcal{I}^{\circ} = K_{\mathbb{C}}/B_K \times U$ of J containing the base cycle $K_{\mathbb{C}}/B_K \times \{u_0\}$, and Ξ is zero from the vanishing theorem on \mathcal{I}^o . Hence $\Xi = 0$ on the open subset $(K_{\mathbb{C}}/T_{\mathbb{C}}) \times U$ of W, and

$$
F\omega^{\mathrm{nc},1}|_{(K_{\mathbb{C}}/T_{\mathbb{C}})\times U}=\partial_{\mathfrak{p}^1_+}\Xi=0,
$$

where $\partial_{\mathfrak{p}^1_+}$ denotes the differentials of the coefficients of Ξ in the directions corresponding to p^1_+ . This implies that $F = 0$ on the open subset $(K_C/T_C) \times U$ of W, and $[F] = 0$ as the cohomological class in $\Gamma(W, \Omega_{\pi'}^{\bullet}(L_{\mu'}))$. This is the main idea of the proof of Theorem [0.2.](#page-2-2)

Next, we introduce the Penrose transformation on the compact quotients of flag domains.

Let $\Gamma \subset G_{\mathbb{R}}$ be a discrete, co-compact and neat subgroup. Then $X = \Gamma \backslash D$ and $X' = \Gamma \backslash D'$ are compact complex manifolds, and $W_{\Gamma} = \Gamma \backslash W$ is a complex manifold which is also Stein. Then the basic diagram [\(1\)](#page-1-1) becomes

Lemma-Definition 0.3. *Let the notations and assumptions be as Lemma-Definition [0.1.](#page-2-0)* Let $X = \Gamma \backslash D$ and $X' = \Gamma \backslash D'$ be the quotients under a discrete, co-compact *and neat subgroup* $\Gamma \subset G_{\mathbb{R}}$ *. Then the Penrose transformation on automorphic cohomology groups*

$$
\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})
$$

is defined by

(7)
$$
H^{0}(X', L_{\mu'}) \xrightarrow{\mathcal{P}} H^{q}(X, L_{\mu})
$$
\n
$$
\downarrow \cong
$$
\n
$$
H^{0}_{DR}(\mathcal{W}_{\Gamma}, \Omega^{\bullet}_{\pi'}(L_{\mu'})) \xrightarrow{\omega_{\Gamma}^{\text{nc},1}} H^{q}_{DR}(\mathcal{W}_{\Gamma}, \Omega^{\bullet}_{\pi}(L_{\mu})),
$$

where $\omega^{\mathrm{nc},1}_{\Gamma}$ $_{\Gamma}^{\rm nc,1}$ is the quotient under Γ of the left-invariant differential form $\omega^{\rm nc,1}$ given *in Lemma-Definition [0.1.](#page-2-0)*

Applying the proof of Theorem [0.2,](#page-2-2) we have the following theorem on the injectivity of the Penrose transformation on automorphic cohomology groups.

Theorem 0.4 (Theorem [4.2\)](#page-26-0)**.** *Let the notations be as Lemma-Definition [0.3.](#page-3-0) If for* any $\beta \in \Delta_+^{\rm nc,1}$ there exists an $\alpha \in \Delta_+^{\rm c}$ such that

(5)
$$
\left(\alpha, \mu + \sum_{\beta' \in \Delta^{nc,1}_+} \beta' - \beta\right) = (\alpha, \mu' - \beta) < 0,
$$

then the Penrose transformation on automorphic cohomology groups

$$
\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})
$$

given by Lemma-Definition [0.3](#page-3-0) is injective.

Hence in order that the Penrose transformation on the compact quotients is an isomorphism, we only need to check the dimensions. Thanks to Theorem 2.4 in [\[25\]](#page-33-5), we have that

$$
\dim_{\mathbb{C}} H^0(X', L_{\mu'}) = \dim_{\mathbb{C}} H^q(X, L_{\mu})
$$

depends only on the number determined by the weight $\mu + \rho = \mu' + \rho'$, provided that $\mu + \rho = \mu' + \rho'$ is regular and satisfies Property **W** of Williams in Lemma [4.5](#page-28-0) which is equivalent to conditions [\(9\)](#page-4-0) below.

Theorem 0.5 (Theorem [4.7\)](#page-29-0)**.** *Let the notations be as Lemma-Definition [0.3.](#page-3-0) Let* μ and μ' be two weights such that $\mu+\rho=\mu'+\rho'$ is regular and $\mu+\rho=\mu'+\rho'$ lies *in the Weyl chamber determined by the set of roots,*

(8)
$$
\Delta_{+}^{\rm c} \cup \Delta_{+}^{\rm nc,1} \cup \left(-\Delta_{+}^{\rm nc,2} \right) = \Delta_{+}^{\rm c} \cup \left(-\Delta_{+}^{\rm nc} \right).
$$

If conditions [\(5\)](#page-2-3) *are satisfied and moreover,*

(9)
$$
(\mu + 2\rho_{\text{nc},2}, \Delta_+^{\text{nc},1} \cup (-\Delta_+^{\text{nc},2})) = (\mu' + 2\rho'_{\text{nc}}, -\Delta_+^{\text{nc}}) > 0,
$$

then the Penrose transformation on automorphic cohomology groups

$$
\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})
$$

is an isomorphism.

Finally, we give an important example of homogenous line bundles satisfying the conditions in Theorem [0.5,](#page-4-1) which has applications to arithmetic geometry and number theory.

Let

$$
\mu'_{c} = -\sum_{\beta \in \Delta'^{\mathrm{nc}}_{+}} \beta = 2\rho_{\mathrm{nc},1} - 2\rho_{\mathrm{nc},2}.
$$

and k_0 the maximal positive integer such that $\mu_{c0}' \triangleq \mu_c'/k_0$ is still a weight. Let

$$
\mu'_{k} = k\mu'_{c0} = \frac{k}{k_0} (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}), \ \mu_{k} = \mu'_{k} - 2\rho_{\text{nc},1}.
$$

Then

$$
L_{\mu'_k} = \omega_{\mathbb{B}}^{\otimes k/k_0} \to D', L_{\mu_k} \to D
$$

are two line bundles whose cohomology groups are Penrose related, where $\omega_{\mathbb{B}}$ is the pull-back of the canonical bundle on $G_{\mathbb{R}}/K \cong \mathbb{B}$ via the holomorphic projection map

$$
p': D' \to G_{\mathbb{R}}/K.
$$

Theorem 0.6 (Theorem [4.8\)](#page-30-0)**.** *Let the notations be as above. Then there exists a positive integer* N *such that for* $k \geq N$ *, the Penrose transformation*

(10)
$$
\mathcal{P}: H^0(X', \omega_{\mathbb{B}}^{\otimes k/k_0}) \to H^q(X, L_{\mu_k})
$$

is an isomorphism. Therefore $H^q(X, L_{\mu_k})$ is isomorphic to the group

$$
H^0(\Gamma \backslash G_{\mathbb{R}}/K, \omega_{\mathbb{B}}^{\otimes k/k_0})
$$

of automorphic forms on $G_{\mathbb{R}}/K$.

The paper is organized as follows. In Section [1](#page-6-0) we introduce the basic notations of the geometry of flag domains and review our results on new complex structures of non-classical flag domains. In Section [2](#page-10-0) we introduce the corresponding space and the incidence variety for the flag domains and relate the cohomology groups of the homogenous line bundles on flag domains to the de Rham cohomology groups on the corresponding space. An important vanishing theorem is proved in this section, which is crucial to the proof of the injectivity of the Penrose transformation. In Section [3](#page-16-0) we define the Penrose transformation on flag domains and establish its injectivity under certain condition. In Section [4](#page-25-0) we introduce the Penrose transformation on the compact quotients of flag domains and prove that it is an isomorphism under the conditions given in Section [3](#page-16-0) and the conditions induced by Property **W** of Williams. We also apply our result to the groups of automorphic forms on Hermitian symmetric domains which can be identified with the higher automorphic cohomology groups of certain line bundles on the compact quotient of the non-classical flag domain.

1. PRELIMINARIES

In this section, we introduce the basic notions of flag domains and fix the notations in this paper. Additionally we recall our recent work [\[20\]](#page-33-2) on the new complex structures of non-classical flag domains, which is the foundation of this paper. References for this section are [\[9\]](#page-32-6), [\[15\]](#page-33-6) and [\[17\]](#page-33-7).

Let G be a reductive Q-algebraic group. Let $G_{\mathbb{R}}$ and $G_{\mathbb{C}}$ be the associated real and complex connected Lie groups respectively. A flag variety $\check{D} = G_{\mathbb{C}}/B$ is a complex homogenous manifold with $B \subset G_{\mathbb{C}}$ a Borel subgroup. A flag domain $D = G_{\mathbb{R}}/T$ is an open $G_{\mathbb{R}}$ -orbit in \tilde{D} , where the isotropy subgroup $T = G_{\mathbb{R}} \cap B$ is a compact Cartan subgroup of $G_{\mathbb{R}}$.

Let K be the maximal compact subgroup of $G_{\mathbb{R}}$ containing the compact Cartan subgroup T. The real Lie group $G_{\mathbb{R}}$ is said to be of Hermitian type if the Riemannian symmetric space $G_{\mathbb{R}}/K$ has a $G_{\mathbb{R}}$ -invariant complex structure. The flag domain D is called classical if $G_{\mathbb{R}}$ is of Hermitian type and the projection map

$$
p:\,D\to G_\mathbb{R}/K
$$

is a holomorphic map between complex manifolds. Otherwise, D is called nonclassical.

Hence non-classical flag domains include the cases that $G_{\mathbb{R}}$ is not of Hermitian type and that $G_{\mathbb{R}}$ is of Hermitian type and projection map p is not holomorphic. In this paper, we mainly consider the non-classical flag domain D with $G_{\mathbb{R}}$ of Hermitian type.

The complex structures of D and $G_{\mathbb{R}}/K$ can be described by the Lie algebras of the Lie groups mentioned above. For this, we fix some notations of the Lie algebras:

- Let $\mathfrak{h}_0 \subset \mathfrak{k}_0 \subset \mathfrak{g}_0$ be the Lie algebras of $T \subset K \subset G_{\mathbb{R}}$ respectively;
- Let $\mathfrak{b} \subset \mathfrak{g}$ be the Lie algebras of $B \subset G_{\mathbb{C}}$ respectively. Then $\mathfrak{g} = \mathfrak{g}_0 \otimes_{\mathbb{R}} \mathbb{C}$ is the complexification of \mathfrak{g}_0 and $\mathfrak{h}_0 = \mathfrak{b} \cap \mathfrak{g}_0$;
- Let $n_$ \subset g be the subalgebra such that

$$
T_o^{1,0}\check{D}=T_o^{1,0}D=\mathfrak{g}/\mathfrak{b}\cong \mathfrak{n}_-,
$$

where o is the base point in D .

In this paper, we also consider other real and complex homogeneous manifolds with base points denoted by σ whenever no ambiguity arises.

• Let $\mathfrak{h} \subset \mathfrak{k}$ be the complexications of $\mathfrak{h}_0 \subset \mathfrak{k}_0$ respectively. Then $\mathfrak{h} \subset \mathfrak{g}$ is the Cartan subalgebra, and there exist subspaces $\mathfrak{p}_0 \subset \mathfrak{g}_0$ and $\mathfrak{p} = \mathfrak{p}_0 \otimes_{\mathbb{R}} \mathbb{C} \subset \mathfrak{g}$ such that

(11)
$$
\mathfrak{g}_{\#} = \mathfrak{k}_{\#} \oplus \mathfrak{p}_{\#}, \text{ with } [\mathfrak{k}_{\#}, \mathfrak{k}_{\#}] \subset \mathfrak{k}_{\#}, [\mathfrak{k}_{\#}, \mathfrak{p}_{\#}] \subset \mathfrak{p}_{\#}, [\mathfrak{p}_{\#}, \mathfrak{p}_{\#}] \subset \mathfrak{k}_{\#},
$$

are Cartan decompositions on \mathfrak{g}_0 and \mathfrak{g} , where $\#$ is 0 or \emptyset ;

• Let $\mathfrak{n}_+ = \overline{\mathfrak{n}_-}$ and $\mathfrak{p}_\pm = \mathfrak{p} \cap \mathfrak{n}_\pm$, $\mathfrak{k}_\pm = \mathfrak{k} \cap \mathfrak{n}_\pm$.

We also denote by the complex Lie subgroups $T_{\mathbb{C}} \subset K_{\mathbb{C}} \subset G_{\mathbb{C}}$ corresponding to the complex Lie subalgebras $\mathfrak{h} \subset \mathfrak{k} \subset \mathfrak{g}$.

Since $\mathfrak{h} \subset \mathfrak{g}$ is the Cartan subalgebra, there exists a decomposition

(12)
$$
\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha},
$$

where $\Delta\subset \sqrt{-1}\mathfrak{h}_0^*$ is the root system of $\mathfrak g$ with respect to $\mathfrak h,$ and

$$
\mathfrak{g}_{\alpha} = \{ X \in \mathfrak{g} : [H, X] = \alpha(H)X, \forall h \in \mathfrak{h} \}
$$

is the root space which is one-dimensional with basis e_{α} for any $\alpha \in \Delta$.

Note that e_{α} , $\alpha \in \Delta$, can be considered as left-invariant vector fields on $G_{\mathbb{C}}$. Let $\{\omega^{\alpha} : \alpha \in \Delta\}$ be the dual of $\{e_{\alpha} : \alpha \in \Delta\}$. Then ω^{α} , $\alpha \in \Delta$, can be considered as left-invariant differential one-forms on $G_{\mathbb{C}}$.

Now we fix some notations for the root systems:

• Since $B \subset G_{\mathbb{C}}$ is Borel subgroup, there exists a set Δ_+ of positive roots such that

$$
\mathfrak{b}=\mathfrak{h}\oplus\bigoplus_{\alpha\in\Delta_+}\mathfrak{g}_{-\alpha},\,\mathfrak{n}_-=\bigoplus_{\alpha\in\Delta_+}\mathfrak{g}_\alpha;
$$

• Let $\Delta^c, \Delta^{\text{nc}} \subset \Delta$ be the sets of compact roots and noncompact roots respectively such that

$$
\mathfrak{k}=\mathfrak{h}\oplus\bigoplus_{\alpha\in\Delta^{\rm c}}\mathfrak{g}_\alpha,\,\mathfrak{p}=\bigoplus_{\beta\in\Delta^{\rm nc}}\mathfrak{g}_\beta.
$$

• Let $\Delta_{+}^{\rm c} = \Delta_{+} \cap \Delta^{\rm c}$ and $\Delta_{+}^{\rm nc} = \Delta_{+} \cap \Delta^{\rm nc}$. Then

$$
\mathfrak{k}_-=\bigoplus_{\alpha\in\Delta_+^\mathrm{c}}\mathfrak{g}_\alpha,\,\mathfrak{p}_-=\bigoplus_{\beta\in\Delta_+^\mathrm{nc}}\mathfrak{g}_\beta.
$$

With the above notations, we see that $n_$ or equivalently Δ_+ determines the complex structure on D via the integrable subbundle of the complexified differential tangent bundle,

$$
\mathrm{T}^{1,0}D = G_{\mathbb{R}} \times_T \mathfrak{n}_- \subset \mathrm{T}^{\mathbb{C}}D = G_{\mathbb{R}} \times_T \mathfrak{g}/\mathfrak{h},
$$

where the inclusion \subset is given by

(13)
$$
\mathfrak{n}_{-} \cong (\mathfrak{n}_{-} \oplus \mathfrak{h})/\mathfrak{h} \subset \mathfrak{g}/\mathfrak{h}.
$$

In the following, all the inclusions of integrable subbundles of the complex differential tangent bundles are given as [\(13\)](#page-7-0).

Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian type, i.e. there exists a $G_{\mathbb{R}}$ -invariant complex structure on the symmetric space $G_{\mathbb{R}}/K$. The $G_{\mathbb{R}}$ -invariant complex structure on $G_{\mathbb{R}}/K$ is given by an abelian subspace

$$
\mathfrak{p}_-'\subset \mathfrak{p},\,\mathfrak{p}_-'\oplus \mathfrak{p}_+'=\mathfrak{p}
$$

such that

$$
\mathrm{T}^{1,0}G_{\mathbb{R}}/K = G_{\mathbb{R}} \times_K \mathfrak{p}'_- \subset \mathrm{T}^{\mathbb{C}}G_{\mathbb{R}}/K = G_{\mathbb{R}} \times_K \mathfrak{g}/\mathfrak{k}.
$$

The following theorems are proved in [\[20\]](#page-33-2).

Theorem 1.1 (Theorem 5.8 of [\[20\]](#page-33-2)). Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain, where $G_{\mathbb{R}}$ is of Hermitian type with an abelian subspace $\mathfrak{p}'_-\subseteq \mathfrak{p}$ such that p'_⊕p'₊ = p. Then the flag domain D admits a new $G_\mathbb{R}$ -invariant complex structure, *given by*

(14)
$$
T^{1,0}D' = G_{\mathbb{R}} \times_T (\mathfrak{k}_- \oplus \mathfrak{p}'_-) \subseteq T^{\mathbb{C}}D
$$

where D′ *denotes the differentiable manifold* D *equipped with the new complex structure. Moreover,* $D' \subset \check{D}$ *is a classical flag domain with a holomorphic projection map*

$$
p': D' \to G_{\mathbb{R}}/K.
$$

And furthermore we have

Theorem 1.2 (Theorem 6.3 of [\[20\]](#page-33-2))**.** *Let the assumption be as in Theorem [1.1](#page-8-0) and let* Γ ⊆ G^R *be a co-compact and torsion-free discrete subgroup. Then we have two compact complex manifolds* $X = \Gamma \backslash D$ *and* $X' = \Gamma \backslash D'$ *which are diffeomorphic to each other, such that* X *is not in Fujiki class* C *while* X′ *is a projective manifold.*

In fact, in [\[20\]](#page-33-2), we proved that $X = \Gamma \backslash D$ admits no $\partial \overline{\partial}$ -structure compactible with the complex structure induced from D , by solving a conjecture of Green– Griffiths–Kerr about the vanishing theorems of automorphic cohomology of any non-trivial locally homogenous vector bundles on X.

In [\[7\]](#page-32-7), Carlson and Toledo proved that there exists no Kähler metric on X . In [\[14\]](#page-33-8), Griffiths, Robles and Toledo proved that the quotient $X = \Gamma \backslash D$, not necessarily compact or smooth, is not algebraic. Hence Theorem [1.2](#page-8-1) can be considered as a generalization of their results.

On the other hand, it is proved in [\[13\]](#page-33-3) that the compact quotient X' of the classical flag domain D' is a projective manifold. Thus, it is of interest to endow the automorphic cohomology groups on X with arithmetic structures inherited from those on X′ via the Penrose transformation discussed below.

Let us denote $n'_{-} = \mathfrak{k}_{-} \oplus \mathfrak{p}'_{-}$ and the set of positive roots corresponding to the complex structure of D' by Δ'_{+} , i.e.

$$
\Delta'_+ = \{ \alpha \in \Delta : \, e_\alpha \in \mathfrak{n}'_- \}.
$$

Since D is non-classical and D' is classical, the complex structures given by $\mathfrak{n}_$ and \mathfrak{n}'_- are different. Also note that \mathfrak{n}_- and \mathfrak{n}'_- only differ in \mathfrak{p}_- and \mathfrak{p}'_- and

$$
\mathfrak{p}_-\oplus\mathfrak{p}_+=\mathfrak{p}_-'\oplus\mathfrak{p}_+'=\mathfrak{p}.
$$

Hence if we let $\mathfrak{p}_-^2 = \mathfrak{p}_- \cap \mathfrak{p}_-'$ and $\mathfrak{p}_-^1 \subset \mathfrak{p}_-$ such that $\mathfrak{p}_- = \mathfrak{p}_-^1 \oplus \mathfrak{p}_-^2$, then

$$
\mathfrak{p}_-^\prime = \mathfrak{p}_+^1 \oplus \mathfrak{p}_-^2.
$$

In this paper, we write \mathfrak{s}_+ with subscript " $+$ " for the complex conjugate of the subspace \overline{s} _− with subscript "−" and vice versa.

In terms of roots, we have the union

$$
\Delta^{\text{nc}}_{+} = \Delta^{\text{nc},1}_{+} \cup \Delta^{\text{nc},2}_{+}
$$

where

$$
\Delta_+^{\text{nc},i} = \{ \beta : e_{\beta} \in \mathfrak{p}_{-}^i \}, i = 1, 2.
$$

10 KEFENG LIU AND YANG SHEN

For Δ_{+}^{\prime} , we have the union

$$
\Delta_+' = \Delta_+^{\prime c} \cup \Delta_+^{\prime nc}, \text{ where } \Delta_-^{\prime c} = \Delta_+^c, \, \Delta_+^{\prime nc} = \left(-\Delta_+^{nc,1}\right) \cup \Delta_+^{nc,2}.
$$

Remark 1.3. Since $[\mathfrak{k}, \mathfrak{p}'_{\pm}] \subset \mathfrak{p}'_{\pm}$, the complex structures on D' in Theorem [1.1](#page-8-0) can also be given by

$$
\mathrm{T}^{1,0}D' = G_{\mathbb{R}} \times_T (\mathfrak{k}'_- \oplus \mathfrak{p}'_-) \subseteq \mathrm{T}^{\mathbb{C}}D,
$$

where $f'_{-} \subset f$ is any Lie subalgebra. This complex structure may be useful in our further study of cup-products of automorphic cohomology groups on compact quotients of flag domains. Nevertheless, we will only consider the complex structure on D' with $\mathfrak{k}'_{-} = \mathfrak{k}_{-}$ for the sake of simplicity.

Lemma 1.4. For any
$$
\beta, \beta' \in \Delta'^{nc}_{+} = (-\Delta^{nc,1}_{+}) \cup \Delta^{nc,2}_{+}
$$
, we have that $(\beta, \beta') \geq 0$.

Proof. Otherwise we have that $(\beta, \beta') < 0$, then $\beta + \beta'$ is a weight in Δ^c , which implies that

$$
[e_{\beta}, e_{\beta'}] = N_{\beta, \beta'} e_{\beta + \beta'} \neq 0
$$

for e_{β} , $e_{\beta'} \in \mathfrak{p}'_{-}$, which contradicts to that \mathfrak{p}'_{-} is abelian.

Before closing this section, we introduce the following lemma for the proof of Lemma [3.1](#page-18-0) below.

Lemma 1.5. *Let the notations be as above. Then we have the following relations of the Lie brackets of the subspaces*

$$
\quad \ \ (15)\qquad \qquad [\mathfrak{k}_{-},\mathfrak{k}_{-}]\subset \mathfrak{k}_{-};
$$

(16)
$$
[\mathfrak{k}_{-}, \mathfrak{p}_{-}^{i}] \subset \mathfrak{p}_{-}^{i}, i = 1, 2;
$$

(17)
$$
[\mathfrak{p}_{-}^1, \mathfrak{p}_{-}^2] \subset [\mathfrak{p}_{-}, \mathfrak{p}_{-}] \subset \mathfrak{k}_{-};
$$

(18)
$$
[\mathfrak{p}_{-}^1, \mathfrak{p}_{+}^2] = 0, [\mathfrak{p}_{-}^i, \mathfrak{p}_{-}^i] = 0, i = 1, 2.
$$

The above relations also hold if we reverse the signs.

Proof. [\(15\)](#page-9-0) and [17](#page-9-0) follow directly from the properties of Cartan decomposition in [11](#page-6-1) and that n[−] is a subalgebra.

[\(18\)](#page-9-0) follows from that

$$
[\mathfrak{p}^1_-, \mathfrak{p}^2_+],\, [\mathfrak{p}^1_-, \mathfrak{p}^1_-]\subset [\mathfrak{p}'_+, \mathfrak{p}'_+]=0
$$

and that

$$
[\mathfrak{p}^2_-, \mathfrak{p}^2_-] \subset [\mathfrak{p}'_-, \mathfrak{p}'_-] = 0.
$$

Note that \mathfrak{p}'_- is preserved by \mathfrak{k}_\pm while \mathfrak{p}_- is only preserved by \mathfrak{k}_- . Hence

$$
[\mathfrak{k}_-,\mathfrak{p}_-^1]\subset [\mathfrak{k}_-,\mathfrak{p}_-]\subset \mathfrak{p}_-,\, [\mathfrak{k}_-,\mathfrak{p}_-^1]\subset [\mathfrak{k}_-,\mathfrak{p}_+']\subset \mathfrak{p}_+'
$$

which implies that

$$
[\mathfrak{k}_-,\mathfrak{p}_-^1]\subset \mathfrak{p}_-\cap \mathfrak{p}_+'=\mathfrak{p}_-^1.
$$

This proves [\(16\)](#page-9-0) with $i = 1$. A similar argument implies (16) with $i = 2$.

Remark 1.6. (1) Lemma [1.5](#page-9-1) has the counterpart relations of the corresponding subsets of roots. For example, [\(18\)](#page-9-0) implies that for any $\beta, \beta' \in \Delta^{nc,i}_+, \beta + \beta'$ is not a root, where $i = 1, 2$.

(2) For the sake of simplicity, we may assume that \mathfrak{p}^1 or $\Delta^{nc,1}_+$ is minimal in the sense that for any $\beta\in\Delta_+^{nc,1}$, there exists $\gamma\in\Delta_+^{nc,2}$ such that $\beta+\gamma\in\Delta_+^c$ is a root. If not, i.e. $\beta+\gamma$ is not a root for any $\gamma\in\Delta_+^{\mathrm{nc}},$ then we can put β into $\Delta_+^{\mathrm{nc},2}.$

2. CORRESPONDING SPACES

In this section, we introduce the corresponding space W for the flag domains D and D' . We relate the cohomology groups of the homogenous line bundles on D and D' to the de Rham cohomology groups on W, which is the foundation of the Penrose transformation. We also introduce the incidence variety J and prove the vanishing theorem on certain open subsets J^o of J , which is crucial to the proof of the injectivity of the Penrose transformation.

Let

$$
D = G_{\mathbb{R}}/T \subset \check{D} = G_{\mathbb{C}}/B
$$

be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian type. Then we have a natural projection map

(19)
$$
p: D = G_{\mathbb{R}}/V \to G_{\mathbb{R}}/K.
$$

Let $o \in D$ be a base point and

$$
\overline{o} = p(o) \in G_{\mathbb{R}}/K.
$$

The fiber

(20)
$$
Z_o \triangleq p^{-1}(\overline{o}) \simeq K/V \simeq K_{\mathbb{C}}/(K_{\mathbb{C}} \cap B)
$$

is a flag manifold which is an analytic subvariety of D . We call Z_o the base cycle for D.

We define the cycle space U by

(21) $\mathcal{U} =$ topological component of Z_o in $\{gZ_o : g \in G_{\mathbb{C}}, gZ_o \subset D\}$.

Note that for the classical flag domain D' , Z_o is also an analytic subvariety, while the cycle space \mathcal{U}' for D' is trivial in the sense that any $g \in G_{\mathbb{C}}$ such that $gZ_o \subset D'$ is an analytic subvariety if and only if $g \in G_{\mathbb{R}}$.

We collect some properties of the cycle space U as introduced in Lecture 6 of [\[11\]](#page-33-0).

Theorem 2.1. *(1) The cycle space* U *is Stein.*

(2) If D *is non-classical, then there is an open embedding*

$$
\mathfrak{U}\subset G_{\mathbb{C}}/K_{\mathbb{C}}\triangleq \check{\mathfrak{U}},
$$

where $K_{\mathbb{C}}$ *is the closed subgroup of* $G_{\mathbb{C}}$ *corresponding to* $\mathfrak{k} \subset \mathfrak{g}$ *.*

(3) If D is non-classical with $G_{\mathbb{R}}$ of Hermitian type, then there is a biholomor*phism*

$$
\mathcal{U}\cong\mathbb{B}\times\overline{\mathbb{B}}.
$$

where $\mathbb B$ *denotes the Hermitian symmetric space* $G_{\mathbb R}/K$ *and* $\overline{\mathbb B}$ *denotes the complex conjugate of* B*.*

Please see [\[9\]](#page-32-6) for the proofs of (1) and (2), and see [\[3\]](#page-32-8) or [9] for the proof of (3).

We define some closed subgroups of the complex Lie group $G_{\mathbb{C}}$ as follows:

$$
T_{\mathbb{C}} = \text{complexification of } T \text{ with Lie algebra } \mathfrak{h};
$$

$$
B_K = K_{\mathbb{C}} \cap B \text{ with Lie algebra } \mathfrak{b}_K = \mathfrak{h} \oplus \sum_{\alpha \in \Delta_{+}^{c}} \mathfrak{g}_{-\alpha},
$$

and define the corresponding complex homogenous spaces by

$$
\begin{array}{rcl}\n\check{\mathsf{W}} & = & G_{\mathbb{C}} / T_{\mathbb{C}}; \\
\check{\mathsf{J}} & = & G_{\mathbb{C}} / B_K.\n\end{array}
$$

Since $T_{\mathbb{C}} \subset B_K \subset B$ (or B') and $B_K \subset K_{\mathbb{C}}$, we have the following commutative diagram of projection maps of complex homogenous spaces

Recall that the open embedding $D\subset \check{D}$ induces the open embedding $\mathfrak{U}\subset \check{\mathfrak{U}}$ in [\(21\)](#page-10-1). We can also define the open subsets $W \subset \check{W}$ and $\mathcal{I} \subset \check{\mathcal{I}}$, such that each square box of the following diagram is Cartesian,

Then the projection maps in [\(22\)](#page-11-0) restrict to the following fibrations

We adopt the notations for fibrations from [\[11\]](#page-33-0) to facilitate reference, allowing readers to consult further details as needed.

Definition 2.2. *The space* W *is referred to as the enhanced flag variety, with* W *the correspondence space, and* I *is called the incidence variety.*

For $u \in \mathcal{U}$, we denote the corresponding analytic subvariety by $Z_u \subset D$. The incidence variety I can also be defined by

$$
\mathcal{I} = \{(x, u) \in D \times \mathcal{U} : x \in Z_u\}.
$$

Then the projection maps π_D and $\pi_\mathfrak{U}$ are the restrictions to J of the projections map $\text{pr}_1: D \times \mathcal{U} \to D$ and $\text{pr}_2: D \times \mathcal{U} \to \mathcal{U}$ respectively.

Proposition 2.3. *The space* \check{W} *is an affine algebraic variety and hence a Stein manifold; The open subset* W of \tilde{W} *is also Stein.*

Proof. Since all the Lie groups considered in this paper are linear algebraic groups, the complex Lie groups $G_{\mathbb{C}}$ and $T_{\mathbb{C}}$, which are the complexificaions of the compact real Lie groups, are reductive. Therefore the quotient $\check{W} = G_{\mathbb{C}}/T_{\mathbb{C}}$ is an affine algebraic variety, and hence Stein.

The open subset $W \subset \check{W}$ can also be defined as the fiber product $\mathcal{U} \times_{\check{U}} \check{W}$, which is a closed subset of $\mathfrak{U} \times \mathfrak{W}$. From Theorem [2.1,](#page-10-2) the product $\mathfrak{U} \times \mathfrak{W}$ of Stein manifolds is also Stein, and so is its closed submanifold W.

We also collect some basic facts about the fibers of the fibrations in [\(24\)](#page-11-1). Please see Lecture 7 in [\[11\]](#page-33-0) for details.

Proposition 2.4. (1) The fibers of $\pi_D : \mathcal{U} \to D$ and $\pi_J : \mathcal{W} \to \mathcal{I}$ are isomorphic t o the typical ones (i.e. f iber of the base point) $B/B_K \,\cong\, \mathfrak{p}_+$ and $B_K/H \,\cong\, \mathfrak{k}_+$ *respectively which are contractible affine algebraic varieties.*

(2) The fibers of π_u : J \rightarrow U *are isomorphic to the typical one* $Z_o \simeq K_{\rm C}/B_K$ *which is a projective variety;*

(3) The fibers of $\pi_{\text{I}} : \mathcal{W} \to \text{I}$ are isomorphic to the typical one $K_{\text{C}}/T_{\text{C}}$ which is *an affine algebraic variety.*

In order to relate the cohomology on D to that on W , we recall the theorem of EGW (Eastwood-Gindikin-Wong) in [\[8\]](#page-32-5) following Lecture 7 of [\[11\]](#page-33-0).

Let M, N be complex manifolds and $\pi : M \to N$ a holomorphic submersion. For $F \to N$ a holomorphic vector bundle, we let

- $\pi^{-1}F$ be the pullback to M of F as a sheaf of holomorphic sections of F;
- $\pi^* F = \pi^{-1} F \otimes_{\pi^{-1} \mathcal{O}_N} \mathcal{O}_M$.

Then $\pi^{-1}F \subset \pi^*F$ is identified with the sections of π^*F that are constant along the fibers of $M \rightarrow N$.

We define $\Omega_{\pi}^1 = \Omega_M^1 / \pi^* \Omega_N^1$ and $\Omega_{\pi}^q = \Lambda^q \Omega_{\pi}^1$ to be the sheaf over M of relative holomorphic q -forms, with the relative differential induced from d ,

$$
d_{\pi}: \Omega_{\pi}^q \to \Omega_{\pi}^{q+1}.
$$

Then we have the short exact sequence

$$
0 \to \pi^* \Omega^1_N \to \Omega^1_M \to \Omega^1_\pi \to 0,
$$

of sheaves of holomorphic forms.

Tensoring with π^*F , we have $\Omega^q_\pi(F) = \Omega^q_\pi \otimes_{\mathcal{O}_M} \pi^*F$ and the relative differential with values in F ,

$$
d_{\pi}: \Omega_{\pi}^q(F) \to \Omega_{\pi}^{q+1}(F).
$$

This gives the resolution of $\pi^{-1}F$

(25)
$$
0 \to \pi^{-1}F \to \pi^*F = \Omega^0_\pi(F) \xrightarrow{d_\pi} \Omega^1_\pi(F) \xrightarrow{d_\pi} \Omega^2_\pi(F) \to \cdots.
$$

Let $\mathbb{H}^*(M, \Omega^\bullet_\pi(F))$ denote the hypercohomology of the complex $\Omega^\bullet_\pi(F)$ of sheaves. Then [\(25\)](#page-13-0) implies that

$$
H^*(M, \pi^{-1}F) = \mathbb{H}^*(M, \Omega^{\bullet}_{\pi}(F)).
$$

Let

$$
H^*_{DR}(M,\Omega^\bullet_\pi(F))=H^*\left(\Gamma(M,\Omega^\bullet_\pi(F));d_\pi\right)
$$

be the de Rham cohomology groups, defined by the cohomology groups of the complex $(\Gamma(M, \Omega^\bullet_\pi(F)); d_\pi)$ of global holomorphic sections of $\Omega^\bullet_\pi(F)$.

Theorem 2.5 (EGW). Assume that M is Stein and the fibers of $M \rightarrow N$ are *contractible. Then*

$$
H^*(N, F) \cong H^*_{DR}(M, \Omega^{\bullet}_{\pi}(F)).
$$

Applying the theorem of EGW to the holomorphic submersion $\pi : W \to D$, we have that

(26)
$$
H^*(D, L_{\mu}) = H^*_{DR}(\mathcal{W}, \Omega^{\bullet}_{\pi}(L_{\mu})).
$$

Next we relate the de Rham cohomology groups on W to that on J . From the commutative diagram of holomorphic submersions

$$
W
$$
\n
$$
D \leftarrow \frac{W}{\pi D} \mathbf{J},
$$

we have the short exact sequence of sheaves on W,

$$
0 \to \pi_{\mathcal{I}}^*\Omega^1_{\pi_D} \to \Omega^1_{\pi} \to \Omega^1_{\pi_{\mathcal{I}}} \to 0,
$$

which defines a filtration F on Ω^\bullet_π by

$$
F^p \Omega^\bullet_\pi = \operatorname{Im}\{\pi^*_\jmath \Omega^p_{\pi_D} \otimes \Omega^{\bullet - p}_\pi \to \Omega^\bullet_\pi\}.
$$

The spectral sequence for F is

$$
E_0^{p,q} = \Gamma(\mathcal{W}, \text{Gr}_F^p \Omega_\pi^{p+q}(L_\mu))) = \Gamma(\mathcal{W}, \pi_j^* \Omega_{\pi_D}^p \otimes \Omega_{\pi_j}^q(L_\mu))
$$

\n
$$
d_0 = d_{\pi_0} : E_0^{p,q} \to E_0^{p,q+1}
$$

\n
$$
E_\infty^{p,q} = \text{Gr}_F^p H_{DR}^{p+q}(\mathcal{W}, \Omega_\pi^\bullet(L_\mu)).
$$

Then

$$
E_1^{p,q} = H^q_{DR}(\mathcal{W}, \pi_J^*\Omega^p_{\pi_D} \otimes \Omega^\bullet_{\pi_J}(L_\mu))_{d_{\pi_J}}.
$$

Here we write $H^*_{DR}(\cdots)_{d_{\pi_{\mathcal{I}}}}$ to emphasize the differential map. Again apply the theorem of EGW to the holomorphic submersion $\pi_{\mathcal{I}} : \mathcal{W} \to \mathcal{I}$ with sheaf $\Omega_{\pi_D}^p(L_\mu)$ to have that

$$
E_1^{p,q}=H^q_{DR}(\mathcal{W},\pi_J^*\Omega^p_{\pi_D}\otimes \Omega^\bullet_{\pi_J}(L_\mu))_{d_{\pi_J}}=H^q(\mathfrak{I},\Omega^p_{\pi_D}(L_\mu)).
$$

Hence

$$
E_1^{p,0} = \Gamma(1, \Omega_{\pi_D}^p(L_\mu)), d_1 = d_{\pi_D} : E_1^{p,0} \to E_1^{p+1,0},
$$

and

$$
E_2^{p,0} = H_{DR}^p(\mathcal{I}, \Omega_{\pi_D}^{\bullet}(L_\mu)).
$$

The quotient maps

$$
E_2^{p,0} \to E_3^{p,0} \to \cdots \to E_{\infty}^{p,0} = \mathrm{Gr}_F^p H_{DR}^p(\mathcal{W}, \Omega_\pi^{\bullet}(L_\mu))
$$

induce the surjective morphism $E_2^{p,0}\to E_\infty^{p,0}$, which is

$$
\pi_{\mathcal{I}}^* : H^p_{DR}(\mathcal{I}, \Omega_{\pi_D}^{\bullet}(L_\mu)) \to H^p_{DR}(\mathcal{W}, \pi_{\mathcal{I}}^* \Omega_{\pi_D}^{\bullet}(L_\mu)).
$$

In summary, we have proved the following.

Proposition 2.6. *The morphism*

$$
\pi_{\mathcal{I}}^*:\hspace{0.1 cm} H^*_{DR}(\mathcal{I}, \Omega_{\pi_D}^\bullet(L_\mu)) \rightarrow H^*_{DR}(\mathcal{W}, \pi_{\mathcal{I}}^*\Omega_{\pi_D}^\bullet(L_\mu))
$$

induced by $\pi_{\mathcal{I}}: \mathcal{W} \rightarrow \mathcal{I}$ *is surjective.*

Before closing this section, we prove the vanishing theorem on J for the injectivity of the Penrose transformation.

Theorem 2.7. Let L_{λ} be a homogenous line bundle on D with weight λ satisfying

$$
q_c(\lambda) \triangleq \#\{\alpha \in \Delta^c_+: (\lambda, \alpha) < 0\} \ge 1,
$$

i.e. there exists some $\alpha \in \Delta^c_+$ *such that* $(\lambda, \alpha) < 0$ *. Then*

(27)
$$
\Gamma(\mathbf{I}, \pi_D^* L_\lambda) = 0.
$$

Proof. **Step 1.** For any $G_{\mathbb{R}}$ -translation of the base cycle Z_o given in [\(20\)](#page-10-3), we have that

$$
\Gamma(gZ_o, L_\lambda|_{gZ_o}) = 0, \,\forall \, g \in G_\mathbb{R}.
$$

In fact, on the base cycle, we have

$$
L_\lambda|_{Z_o}=K\times_T\mathbb{C}_\lambda
$$

and hence the Bott–Borel–Weil theorem (c.f. [\[2\]](#page-32-9), [\[11\]](#page-33-0) or [\[22\]](#page-33-9)) implies that

$$
\Gamma(Z_o, L_\lambda|_{Z_o}) = H^0(Z_o, L_\lambda|_{Z_o}) = 0.
$$

On the $G_{\mathbb{R}}$ -translation gZ_o , the biholomorphic map $g : D \to D$ induces isomorphism

 $g: L_{\lambda}|_{Z_o} \to L_{\lambda}|_{gZ_o},$

which implies that

$$
\Gamma(gZ_o, L_\lambda|_{gZ_o}) = \Gamma(Z_o, L_\lambda|_{Z_o}) = 0.
$$

Step 2. There exists a nontrivial open subset $\mathcal{U}^{\circ} \subset \mathcal{U}$ containing the $G_{\mathbb{R}}$ translations of the base cycle Z_o such that

(28)
$$
\Gamma(Z_u, L_\lambda|_{Z_u}) = 0, \,\forall \, u \in \mathcal{U}^o.
$$

Let $u_0 \in \mathcal{U}$ such that $Z_{u_0} = gZ_o$ for some $g \in G_{\mathbb{R}}$. We can choose a connected open neighborhood U_{u_0} of u_0 in U, over which there exists an analytic family

$$
D \supset \mathcal{Z} \xrightarrow{\pi} U_{u_0}
$$

which is C^{∞} -trivial, such that $Z_u = \pi^{-1}(u)$ for any $u \in U_{u_0}$. Then the restriction

 $L_{\lambda}|_{\mathcal{Z}} \to \mathcal{Z}$

gives an family of line bundles

$$
(L_{\lambda}|_{\mathcal{Z}})|_{Z_u}=L_{\lambda}|_{Z_u}
$$

over U_{u_0} .

By Theorem 6 in [\[19\]](#page-33-4), we know that $\dim H^0(Z_u, L_\lambda|_{Z_u})$ is an upper semicontinuous function of $u \in U_{u_0}$, i.e. for any $u \in U_{u_0}$, one has

$$
\dim H^0(Z_u, L_\lambda|_{Z_u}) \le \dim H^0(Z_{u_0}, L_\lambda|_{Z_{u_0}}) = 0,
$$

which implies that $\Gamma(Z_u, L_\lambda|_{Z_u}) = 0$ for $u \in U_{u_0}$.

Let

$$
\mathcal{U}^o = \bigcup_{u = [gZ_o], g \in G_{\mathbb{R}}} U_u.
$$

Then \mathcal{U}^o satisfies [\(28\)](#page-15-0).

Step 3. Now we can prove the theorem.

Let

$$
\mathcal{I}^o = \{(x, u) \in D \times \mathcal{U}^o : x \in Z_u\} = \pi^{-1}_\mathcal{U}(\mathcal{U}^o).
$$

Then \mathcal{I}^o is an open subset of \mathcal{I} , and the restriction $\pi_D|_{\mathcal{I}^o}: \mathcal{I}^o \to D$ is still surjective, since for any $x \in D$ there exists $g \in G_{\mathbb{R}}$ such that gZ_o passes through x.

For any fixed $(x, u) \in \mathcal{I}^o$ with $x \in Z_u$, the subset

$$
\tilde{Z}_u = \{(y, u) \in \mathcal{I}^o : y \in Z_u\}
$$

is a compact subvariety of \mathcal{I}^o passing through (x, u) , which is isomorphic to $Z_u \subset \mathbb{R}^d$ D via π_D . Then

$$
\Gamma(\tilde{Z}_u, \pi_D^* L_\lambda|_{\tilde{Z}_u}) = \Gamma(Z_u, L_\lambda|_{Z_u}) = 0,
$$

by Step 2.

Since $\{\tilde{Z}_u : u \in \mathcal{U}^o\}$ covers \mathcal{I}^o and \mathcal{I}^o is a nontrivial open subset of \mathcal{I} , we have the conclusion [\(27\)](#page-14-1) of the theorem. \Box

Corollary 2.8. *Let the the assumption be as in Theorem [2.7.](#page-14-0) Let* U *be any small enough open subset of* U *containing the base point* u_o , *i.e.* the point representing *the base cycle* Z_o *. Then*

$$
\Gamma(\pi_{\mathfrak{U}}^{-1}(U), \pi_D^* L_\lambda|_{\pi_{\mathfrak{U}}^{-1}(U)}) = 0.
$$

Remark 2.9. (1) In Corollary [2.8,](#page-15-1) the open subset $U \subset \mathcal{U}$ can be chosen sufficiently small such that the vanishing theorem remains valid. This is because $\pi_u^{-1}(U)$ always contains and is covered by the cycles \tilde{Z}_u for $u \in U$. However we do not have the vanishing theorem on the open subset $V \subset J$, if V dose not contain any \tilde{Z}_u for $u \in \mathcal{U}$.

(2) In [\[14\]](#page-33-8), Griffiths, Robles and Toledo proved the cycle chain connectedness of the non-classical flag domains D , by proving that the incidence variety $\mathfrak I$ can be connected by integral curves of a subbundle $S \simeq \mathfrak{k}_- \oplus \mathfrak{p}_+$ of TJ. Now we can use the same method to show that J^o can be connected by integral curves of the restricted subbundle $S|_{\mathcal{I}^o}$, and conclude that any two points in D can be connected by a chain of cycles in \mathcal{U}^o . Here \mathcal{U}^o can be taken small enough containing the $G_{\mathbb{R}}$ -translations of the base cycles.

We can get more geometric properties of the non-classical flag domain from this observation in a forthcoming paper.

3. PENROSE TRANSFORMATION ON FLAG DOMAINS

In this section, we define the Penrose transformation on flag domains and establish its injectivity under specific conditions. Additionally, we present particular cases of weights that satisfy these conditions, so that the corresponding Penrose transformations are injective.

Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian type. Then Theorem [\(1.1\)](#page-8-0) implies that D is diffeomorphic to a classical flag domain D' . The complex structures of D and D' are given respectively by

$$
\mathcal{T}^{1,0}_o D \cong \mathfrak{n}_- = \mathfrak{k}_- \oplus \mathfrak{p}_-^1 \oplus \mathfrak{p}_-^2
$$

$$
\mathcal{T}^{1,0}_o D' \cong \mathfrak{n}'_- = \mathfrak{k}_- \oplus \mathfrak{p}_+^1 \oplus \mathfrak{p}_-^2.
$$

We define the projection map

$$
\pi_{D'} : \mathcal{I} \to D'
$$

by sending $(x, Z_u) \in \mathcal{I}$ to $x \in D'$, where $x \in D' \simeq D$ and $u = [Z_u] \in \mathcal{U}$. Then the fiber $\pi_{D'}^{-1}(x)$ is $\{u \in \mathcal{U} : x \in Z_u\}$. Note that there is only one $u \in \pi_{D'}^{-1}(x)$ such that Z_u is the compact analytic subvariety of D' , which is the $G_\mathbb{R}$ -translation of the base cycle Z_o . Nevertheless we still have that $\pi_{D'}$ remains a holomorphic map, as we have

$$
\mathrm{T}^{1,0}_o D' \cong \mathfrak{n}'_- \subset \mathfrak{k}_- \oplus \mathfrak{p}_- \oplus \mathfrak{p}_+ \cong \mathrm{T}^{1,0}_o \mathfrak{I}.
$$

Therefore we have the following commutative diagram of holomorphic fibrations

Since the fibers of π' are isomorphic to those of π , we can apply Theorem [2.5](#page-13-1) of EGW to the holomorphic submersion $\pi' : \mathcal{W} \to D'$ to have that

(30)
$$
H^*(D', L_{\mu'}) = H^*_{DR}(\mathcal{W}, \Omega^{\bullet}_{\pi'}(L_{\mu'})).
$$

The Penrose transformation relates the cohomology $H^*(D', L_{\mu'})$ to $H^*(D, L_{\mu})$ via the de Rham cohomology groups of the global sections on W of relative differential forms with values in $L_{\mu'}$ and L_{μ} .

Before giving the precise definition, we first introduce some basic lemmas from Lie algebra and geometry of homogenous manifolds.

In terms of root system, the complex structures on D and D' are given respectively by the set of positive roots

$$
\begin{array}{rcl}\n\Delta_+ &=& \Delta_+^c \cup \Delta_+^{nc,1} \cup \Delta_+^{nc,2} \\
\Delta_+' &=& \Delta_+^c \cup \left(-\Delta_+^{nc,1}\right) \cup \Delta_+^{nc,2}.\n\end{array}
$$

Let

$$
\begin{array}{lcl} \rho & = & \displaystyle \frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha = \frac{1}{2} \sum_{\alpha \in \Delta_+^c} \alpha + \frac{1}{2} \sum_{\alpha \in \Delta_+^{nc,1}} \alpha + \frac{1}{2} \sum_{\alpha \in \Delta_+^{nc,2}} \alpha \triangleq \rho_c + \rho_{\text{nc},1} + \rho_{\text{nc},2} \\ \rho' & = & \displaystyle \frac{1}{2} \sum_{\alpha \in \Delta_+'} \alpha = \rho_c - \rho_{\text{nc},1} + \rho_{\text{nc},2} \triangleq \rho_c + \rho_{\text{nc}}'. \end{array}
$$

For a weight λ we define

$$
q(\lambda) = #\{\alpha \in \Delta^c_+ : (\lambda, \alpha) < 0\} + #\{\alpha \in \Delta^{nc}_+ : (\lambda, \alpha) > 0\}
$$
\n
$$
q'(\lambda) = #\{\alpha \in \Delta'^c_+ : (\lambda, \alpha) < 0\} + #\{\alpha \in \Delta'^{nc}_+ : (\lambda, \alpha) > 0\}.
$$

From now on, we fix the positive integer

$$
q = \# \Delta_+^{\mathrm{nc},1},
$$

which is the number of non-compact roots whose signs are changed when we transform the complex structure on D to that on D' .

For convenience we write α , β and γ for the roots in $\Delta_{+}^{\rm c}$, $\Delta_{+}^{\rm nc,1}$ and $\Delta_{+}^{\rm nc,2}$ + respectively. Then

$$
\Delta_{+}^{c} = \{\alpha_1, \cdots, \alpha_d\},
$$

\n
$$
\Delta_{+}^{\text{nc},1} = \{\beta_1, \cdots, \beta_q\},
$$

\n
$$
\Delta_{+}^{\text{nc},2} = \{\gamma_1, \cdots, \gamma_r\},
$$

where $d = \dim_{\mathbb{C}} Z_o$ and $r = \dim_{\mathbb{C}} D - d - q$.

Let ω^{δ} be the dual to e_{δ} for $\delta \in \Delta$. Then ω^{δ} can be considered as a holomorphic left invariant differential form on $G_{\mathbb{C}}$, which satisfies that

$$
\omega^{\delta}(g.X) = -\delta(X)\omega^{\delta}(g), \,\forall X \in \mathfrak{h}, \,\forall g \in G_{\mathbb{C}}.
$$

Here $g.X$ denotes the right infinitesimal action of $\frak h$ on $G_{\mathbb C}.$ Hence, ω^δ descends to a holomorphic left-invariant differential 1-form with values in L_{δ} on $\dot{W} = G_{\mathbb{C}}/T_{\mathbb{C}}$ and then restricts to the open subset W, i.e.

$$
\omega^{\delta} \in \Gamma(\mathcal{W}, \Omega^1_{\mathcal{W}}(L_{\delta})).
$$

Lemma 3.1. Let $C_{\delta\delta'}^{\delta''}$, $\delta, \delta', \delta'' \in \Delta_+$, be the structure constants of the algebra $\mathfrak{n}_+ = \mathfrak{k}_+ \oplus \mathfrak{p}_+^1 \oplus \mathfrak{p}_+^2$ such that

$$
[e_{-\delta},e_{-\delta'}]=\sum_{\delta''\in\Delta_+}C_{\delta\delta'}^{\delta''}e_{-\delta''},\ C_{\delta\delta'}^{\delta''}=-C_{\delta'\delta}^{\delta''}.
$$

Then

(31)
$$
C^{\beta_j}_{\alpha_i \alpha_{i'}} = 0, C^{\gamma_k}_{\alpha_i \alpha_{i'}} = 0;
$$

(32)
$$
C^{\alpha_{i'}}_{\alpha_{i}\beta_{j}} = 0, C^{\gamma_{k}}_{\alpha_{i}\beta_{j}} = 0; C^{\alpha_{i'}}_{\alpha_{i}\gamma_{k}} = 0, C^{\beta_{j}}_{\alpha_{i}\gamma_{k}} = 0;
$$

(33)
$$
C_{\beta_j\gamma_k}^{\beta_{j'}}=0, C_{\beta_j\gamma_k}^{\gamma_{k'}}=0;
$$

(34)
$$
C_{\beta_j \beta_{j'}}^{\delta} = 0, C_{\gamma_k \gamma_{k'}}^{\delta} = 0, \forall \delta \in \Delta_+,
$$

for $1 \le i, i' \le d, 1 \le j, j' \le q, 1 \le k, k' \le r$.

Proof. Equation [\(31\)](#page-18-1), [\(32\)](#page-18-1),[\(33\)](#page-18-1),[\(34\)](#page-18-1) follow from [\(15\)](#page-9-0), [\(16\)](#page-9-0), [\(17\)](#page-9-0), [18](#page-9-0) in Lemma [1.5](#page-9-1) respectively.

In the following, we write $\omega \equiv_{\pi} \omega'$ for $\omega, \omega' \in \Gamma(\mathcal{W}, \Omega_{\pi}^{\bullet}(L_{\lambda}))$ if $\omega - \omega' = 0$ in $\Omega_{\pi}^{\bullet}(L_{\lambda}).$

Lemma 3.2. *Let*

$$
\omega^{-\alpha_i} \in \Gamma(\mathcal{W}, \Omega_\pi^1(L_{-\alpha_i})), \ \alpha_i \in \Delta_+^c, \ 1 \le i \le d;
$$

$$
\omega^{-\beta_j} \in \Gamma(\mathcal{W}, \Omega_\pi^1(L_{-\beta_j})), \ \alpha_j \in \Delta_+^{\text{nc},1}, \ 1 \le j \le q;
$$

$$
\omega^{-\gamma_k} \in \Gamma(\mathcal{W}, \Omega_\pi^1(L_{-\gamma_k})), \ \gamma_k \in \Delta_+^{\text{nc},2}, \ 1 \le k \le r.
$$

Then

$$
(35) \qquad d_{\pi}\omega^{-\alpha_i} \equiv_{\pi} -\frac{1}{2} \sum_{i',i''} C^{\alpha_i}_{\alpha_{i'}\alpha_{i''}} \omega^{-\alpha_{i'}} \wedge \omega^{-\alpha_{i''}} - \sum_{j,k} C^{\alpha_i}_{\beta_j\gamma_k} \omega^{-\beta_j} \wedge \omega^{-\gamma_k};
$$

(36)
$$
d_{\pi} \omega^{-\beta_j} \equiv_{\pi} - \sum_{i,j'} C^{\beta_j}_{\alpha_i \beta_{j'}} \omega^{-\alpha_i} \wedge \omega^{-\beta_{j'}};
$$

(37)
$$
d_{\pi} \omega^{-\gamma_k} \equiv_{\pi} - \sum_{i,k'} C^{\gamma_k}_{\alpha_i \gamma_{k'}} \omega^{-\alpha_i} \wedge \omega^{-\gamma_{k'}}.
$$

Proof. From (3.8) of [\[21\]](#page-33-10) (see also [\[13\]](#page-33-3) Page 260), we have that

$$
-2d_{\pi}\omega^{-\alpha_i} \equiv_{\pi} \sum_{\delta,\delta' \in \Delta_+} C^{\alpha_i}_{\delta\delta'}\omega^{-\delta} \wedge \omega^{-\delta'}
$$

\n
$$
= \sum_{i',i''} C^{\alpha_i}_{\alpha_i'\alpha_{i''}}\omega^{-\alpha_{i'}} \wedge \omega^{-\alpha_{i''}} + \sum_{i',j} C^{\alpha_i}_{\alpha_{i'}\beta_j}\omega^{-\alpha_{i'}} \wedge \omega^{-\beta_j} + \sum_{i',k} C^{\alpha_i}_{\alpha_{i'}\gamma_k}\omega^{-\alpha_{i'}} \wedge \omega^{-\gamma_k}
$$

\n
$$
+ \sum_{j,i'} C^{\alpha_i}_{\beta_j\alpha_{i'}}\omega^{-\beta_j} \wedge \omega^{-\alpha_{i'}} + \sum_{j,j'} C^{\alpha_i}_{\beta_j\beta_{j'}}\omega^{-\beta_j} \wedge \omega^{-\beta_{j'}} + \sum_{j,k} C^{\alpha_i}_{\beta_j\gamma_k}\omega^{-\beta_j} \wedge \omega^{-\gamma_k}
$$

\n
$$
+ \sum_{k,i'} C^{\alpha_i}_{\gamma_k\alpha_{i'}}\omega^{-\gamma_k} \wedge \omega^{-\alpha_{i'}} + \sum_{k,j} C^{\alpha_i}_{\gamma_k\beta_j}\omega^{-\gamma_k} \wedge \omega^{-\beta_j} + \sum_{k,k'} C^{\alpha_i}_{\gamma_k\gamma_{k'}}\omega^{-\gamma_k} \wedge \omega^{-\gamma_{k'}}.
$$

Then [\(35\)](#page-18-2) follows from Lemma [3.1](#page-18-0) and that $C^{\alpha_i}_{\beta_i}$ $\frac{\partial a_i}{\partial_j \gamma_k} = -C^{\alpha_i}_{\gamma_k}$ ι $\frac{\alpha_i}{\gamma_k\beta_j}$.

Similarly we can deduce [\(36\)](#page-18-2) and [37](#page-18-2) from Lemma [3.1](#page-18-0) and the anti-symmetry of the structure constants. \Box

Lemma 3.3. *Let*

$$
\omega^{\mathrm{nc},1} = \omega^{-\beta_1} \wedge \cdots \wedge \omega^{-\beta_q}
$$

be a left invariant differential form on $G_{\mathbb C}$. Then $\omega^{{\rm nc},1}$ descends to a left invariant *differential* q-form with values in $L_{-2\rho_{\text{net}}}$ on W, which is d_{π} -closed, i.e.

$$
\omega^{\text{nc},1} \in \mathcal{Z}_{d_{\pi}}(\mathcal{W}, \Omega_{\pi}^q(L_{-2\rho_{\text{nc},1}})).
$$

Proof. With the discussion before Lemma [3.1,](#page-18-0) we only need to prove that $d_{\pi}(\omega^{\mathrm{nc},1})=$ 0.

From [\(36\)](#page-18-2), we have that

$$
d_{\pi}\omega^{-\beta_j} \equiv_{\pi} -\sum_{i,j'} C^{\beta_j}_{\alpha_i\beta_{j'}} \omega^{-\alpha_i} \wedge \omega^{-\beta_{j'}},
$$

and the structure constants

$$
C_{\alpha_i\beta_{j'}}^{\beta_j} \neq 0
$$
 only if $j \neq j'$ and $\alpha + \beta_{j'} = \beta_j$.

Therefore

$$
d_{\pi}(\omega^{\text{nc},1}) = \sum_{1 \leq j \leq q} (-1)^{j-1} \omega^{-\beta_1} \wedge \cdots \wedge d_{\pi} \omega^{-\beta_j} \wedge \cdots \wedge \omega^{-\beta_q}
$$

\n
$$
\equiv_{\pi} \sum_{1 \leq j \leq q} (-1)^{j} \omega^{-\beta_1} \wedge \cdots \wedge \left(\sum_{i,j' \neq j} C_{\alpha_i \beta_{j'}}^{\beta_j} \omega^{-\alpha_i} \wedge \omega^{-\beta_{j'}}\right) \wedge \cdots \wedge \omega^{-\beta_q}
$$

\n
$$
= 0.
$$

Lemma-Definition 3.4. Let μ and μ' be two weights such that $\mu + \rho = \mu' + \rho'$. *Then the Penrose transformation*

$$
\mathcal{P}: H^0(D', L_{\mu'}) \to H^q(D, L_{\mu})
$$

is defined by

(38)
$$
H^{0}(D', L_{\mu'}) \xrightarrow{\mathcal{P}} H^{q}(D, L_{\mu})
$$
\n
$$
\downarrow \cong \qquad \qquad \downarrow \cong
$$
\n
$$
H^{0}_{DR}(\mathcal{W}, \Omega_{\pi'}^{\bullet}(L_{\mu'})) \xrightarrow{\omega^{\text{nc},1}} H^{q}_{DR}(\mathcal{W}, \Omega_{\pi}^{\bullet}(L_{\mu})).
$$

Proof. We need to check the well-definedness of P given as above. For any $\sigma \in H^0(D', L_{\mu'})$, let

$$
F_{\sigma} \in \mathcal{Z}_{d_{\pi'}}(\mathcal{W}, L_{\mu'}) = H_{DR}^0(\mathcal{W}, \Omega_{\pi'}^{\bullet}(L_{\mu'}))
$$

correspond to σ via the left vertical isomorphism in [\(38\)](#page-19-0).

For convenience, we will not distinguish the groups and their elements in the vertical isomorphisms in [\(38\)](#page-19-0).

Note that $\mu + \rho = \mu' + \rho'$ implies that $\mu = \mu' - 2\rho_{\text{nc},1}$. Hence Lemma [3.3](#page-19-1) implies that

$$
F_{\sigma}\omega^{\text{nc},1} \in \Gamma(\mathcal{W}, \Omega^q_{\pi}(L_{\mu})).
$$

We need to check that $d_{\pi}(F_{\sigma}\omega^{\mathrm{nc},1})=0.$

Since the complex structures of D and D' only differ in the directions corresponding to \mathfrak{p}^1_- , we have that

$$
(d_{\pi} - d_{\pi'})F_{\sigma} \equiv_{\pi} \sum_{j} a_j \omega^{-\beta_j}.
$$

Therefore from Lemma [3.3](#page-19-1) we have that

$$
d_{\pi}(F_{\sigma}\omega^{\text{nc},1}) = d_{\pi}(F_{\sigma}) \wedge \omega^{\text{nc},1} + F_{\sigma}d_{\pi}(\omega^{\text{nc},1})
$$

$$
\equiv_{\pi} (d_{\pi} - d_{\pi'}) (F_{\sigma}) \wedge \omega^{\text{nc},1}
$$

$$
\equiv_{\pi} \sum_{j} a_{j}\omega^{-\beta_{j}} \wedge \omega^{\text{nc},1} = 0.
$$

Hence

$$
F_{\sigma}\omega^{\mathrm{nc},1} \in \mathcal{Z}_{d_{\pi}}(\mathcal{W}, \Omega_{\pi}^q(L_{\mu})),
$$

and the Penrose transformation given by (38) is well-defined.

Theorem 3.5. Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian *type. Let* D′ *be the classical flag domain which is diffeomorphic to* D*.*

If μ *and* μ' *are two weights with* $\mu + \rho = \mu' + \rho'$ *, and satisfy that for any* $\beta \in \Delta^{nc,1}_+$ + there exists an $\alpha_\beta\in\Delta_+^\mathrm{c}$ such that

(39)
$$
(\alpha_{\beta}, \mu + 2\rho_{\text{nc},1} - \beta) = (\alpha_{\beta}, \mu' - \beta) < 0,
$$

then the Penrose transformation

$$
\mathcal{P}: H^0(D', L_{\mu'}) \to H^q(D, L_{\mu})
$$

given by [\(38\)](#page-19-0) *is injective.*

Proof. Let $F_{\sigma} \in Z_{d_{\pi'}}(\mathcal{W}, L_{\mu'})$ such that

$$
\mathcal{P}([F_{\sigma}])=[F_{\sigma}\omega^{\mathrm{nc},1}]=0
$$

in $H^q_{DR}(\mathcal{W}, \Omega^\bullet_\pi(L_\mu))$. Then there exists $\Psi \in \Gamma(\mathcal{W}, \Omega^{q-1}_\pi(L_\mu))$ such that

$$
F_{\sigma}\omega^{\mathrm{nc},1} = d_{\pi}\Psi.
$$

We need to show that $F_{\sigma} = 0$.

Since $F_{\sigma}\omega^{\text{nc},1}$ has only term in $\wedge^q(\mathfrak{p}_+^1)^* = \mathbb{C}\{\omega^{\text{nc},1}\}\,$, we see that

$$
[F_\sigma\omega^{\mathrm{nc},1}]\in H^q_{DR}(\mathcal{W},\pi^\ast_J\Omega^\bullet_{\pi_D}(L_\mu)).
$$

By Proposition [2.6,](#page-14-2) there exists $[\Phi] \in H^q_{DR}(\mathcal{I}, \Omega^{\bullet}_{\pi_D}(L_\mu))$ such that $\pi^*_\mathcal{I}[\Phi] = [F_{\sigma} \omega^{\text{nc},1}].$ Hence we can assume that F_{σ} is constant along the fibers of $\pi_{\mathcal{I}}: \mathcal{W} \to \mathcal{I}$.

Let $\Psi = \Psi' + \Psi''$, where

$$
\Psi' = \sum_{j} f_j \omega^{-\beta_1} \wedge \cdots \wedge \widehat{\omega^{-\beta_j}} \wedge \cdots \wedge \omega^{-\beta_q};
$$

$$
\Psi'' = \sum_{i} g_i \omega^{-\alpha_i} \wedge \cdots + \sum_{k} h_k \omega^{-\gamma_k} \wedge \cdots.
$$

Then Lemma [3.2](#page-18-3) implies that

(40)
\n
$$
F_{\sigma}\omega^{\text{nc},1} \equiv_{\pi} d_{\pi}\Psi' + d_{\pi}\Psi''
$$
\n
$$
\equiv_{\pi} \sum_{j} (-1)^{j-1} e_{-\beta_{j}}(f_{j}) \omega^{\text{nc},1}
$$
\n
$$
\pm \sum_{j \neq j'} f_{j} d_{\pi} \omega^{-\beta_{j'}} \wedge \omega^{-\beta_{1}} \wedge \cdots \widehat{\omega^{-\beta_{j'}}} \cdots \widehat{\omega^{-\beta_{j}}} \wedge \cdots
$$
\n
$$
+ \sum_{i,j} e_{-\alpha_{i}}(f_{j}) \omega^{-\alpha_{i}} \wedge \omega^{-\beta_{1}} \wedge \cdots \wedge \widehat{\omega^{-\beta_{j}}} \wedge \cdots
$$
\n
$$
+ \sum_{j,k} e_{-\gamma_{k}}(f_{j}) \omega^{-\gamma_{k}} \wedge \omega^{-\beta_{1}} \wedge \cdots \wedge \widehat{\omega^{-\beta_{j}}} \wedge \cdots + d_{\pi}\Psi''
$$

From [\(35\)](#page-18-2) and [\(37\)](#page-18-2), we see that $d_{\pi}\Psi''$ and

$$
\pm \sum_{j \neq j'} f_j d_{\pi} \omega^{-\beta_{j'}} \wedge \omega^{-\beta_1} \wedge \cdots \widehat{\omega^{-\beta_{j'}}} \cdots \widehat{\omega^{-\beta_j}} \wedge \cdots
$$

.

have no terms in $\wedge^q(\mathfrak{p}^1_+)^* = \mathbb{C}\{\omega^{\text{nc},1}\}\.$ Hence by comparing the types of both sides of [\(40\)](#page-21-0), we have that

(41)
$$
F_{\sigma}\omega^{\text{nc},1} = \sum_{j} (-1)^{j-1} e_{-\beta_j}(f_j) \omega^{\text{nc},1} \triangleq \partial_{\mathfrak{p}_+^1} \Psi',
$$

$$
\pm \sum_{j \neq j'} f_j d_{\pi}\omega^{-\beta_{j'}} \wedge \omega^{-\beta_1} \wedge \cdots \widehat{\omega^{-\beta_{j'}}} \cdots \widehat{\omega^{-\beta_j}} \wedge \cdots +
$$

$$
\sum_{i,j} e_{-\alpha_i}(f_j) \omega^{-\alpha_i} \wedge \cdots + \sum_{j,k} e_{-\gamma_k}(f_j) \omega^{-\gamma_k} \wedge \cdots + d_{\pi} \Psi'' = 0.
$$

Here we denote

$$
\partial_{\mathfrak{p}^1_+}(f\omega)=\sum_je_{-\beta_j}(f)\omega^{-\beta_j}\wedge\omega,
$$

which is globally defined on W.

Consider the fibrations in [\(24\)](#page-11-1)

where $\mathbb B$ denotes the Hermitian symmetric domain $G_{\mathbb R}/K$, which is isomorphic to a boundede domain B in \mathfrak{p}_- by the Harish-Chandra's embedding theorem, c.f. Lemma 7.11 in pages 390 – 391 in [\[15\]](#page-33-6).

Let $U = U_1 \times U_2 \subset U$ be an open subset of U with U_1 open subset of B and $U_2 = \overline{U_1} \subset \overline{\mathbb{B}}$. We take U small enough so that

$$
\pi_{\mathfrak{U}}^{-1}(U) \cong K_{\mathbb{C}}/B_K \times U, \pi'^{-1}(U) = \pi_{\mathfrak{I}}^{-1}\pi_{\mathfrak{U}}^{-1}(U) \cong K_{\mathbb{C}}/T_{\mathbb{C}} \times U
$$

and $\pi_{\mathfrak{U}}^{-1}(U) \subset \mathfrak{I}^o$ is covered by the compact analytic subvarieties $\tilde{Z}_u, \, u \in U,$ on which $\Gamma(\tilde Z_u,\pi_D^*L_{\mu_j})=0$ for line bundles $L_{\mu_j},\,1\leq j\leq q,$ which is to be defined below. Here \mathbb{J}^o and \tilde{Z}_u are given as in the proof of Theorem [2.7.](#page-14-0)

Then $\pi_{\mathcal{I}}$ is locally given by

$$
\pi_{\mathfrak{I}}|_{\pi'^{-1}(U)} = \pi_{B_K} \times \mathrm{id}_U : K_{\mathbb{C}}/T_{\mathbb{C}} \times U \to K_{\mathbb{C}}/B_K \times U,
$$

where $\pi_{B_K}: K_{\mathbb{C}}/T_{\mathbb{C}} \to K_{\mathbb{C}}/B_K$ is the projection map with contractible fibers $B_K/T_{\mathbb C}\simeq {\mathfrak k}_+.$

Note that all the homogenous vector bundles on the homogenous manifolds X can be trivialized simultaneously under an open cover of X. Let $\{V_{\nu}\}\$ be a cover of the flag variety $Z_o \cong K_\mathbb{C}/B_K$ under which all the homogenous vector bundles on $K_{\mathbb{C}}/B_K$ are trivialized. Then, since $B_K/T_{\mathbb{C}}$ is contractible, $\{B_K/T_{\mathbb{C}} \times V_{\nu}\}\)$ is a cover of $K_{\rm C}/T_{\rm C}$ under which all the homogenous vector bundles on $K_{\rm C}/T_{\rm C}$ are trivialized.

Let

$$
B_K/T_{\mathbb{C}} = \{w \in \mathfrak{k}_+ : w = (w_1, \dots, w_d)\},
$$

\n
$$
V_{\nu} = \{x_{\nu} \in \mathfrak{k}_- : x_{\nu} = (x_{\nu 1}, \dots, x_{\nu d})\},
$$

\n
$$
U_1 = \{(y, z) \in B : y = (y_1, \dots, y_q), z = (z_1, \dots, z_r)\} \subset \mathfrak{p}_- = \mathfrak{p}_-^1 \oplus \mathfrak{p}_-^2,
$$

be the coordinates on the corresponding spaces, with B a bounded subset of \mathbb{C}^{q+r} . Then

$$
\Psi'|_{B_K/T_\mathbb{C}\times V_\nu\times U}=\sum_j f_{j\nu}(w,x_\nu,y,\overline{y},z,\overline{z})\omega^{-\beta_1}\wedge\cdots\wedge\widehat{\omega^{-\beta_j}}\wedge\cdots\wedge\omega^{-\beta_q},
$$

where $w \in B_K/T_{\mathbb{C}}$, $x_v \in V_v$ and $(y, \overline{y}, z, \overline{z}) \in U$, and

$$
e_{-\alpha_i}(f_{j\nu})=\partial_{w_i}(f_{j\nu}),\ e_{-\beta_{j'}}(f_{j\nu})=\partial_{\overline{y}_{j'}}(f_{j\nu}),\ e_{-\gamma_k}(f_{j\nu})=\partial_{\overline{z}_k}(f_{j\nu}).
$$

Now we define $\Xi \in \Gamma(\pi^{\prime -1}(U), \Omega^{q-1}_\pi(L_\mu))$ by

$$
\Xi|_{B_K/T_{\mathbb{C}}\times V_{\nu}\times U}=\sum_j f_{j\nu}(0,x_{\nu},y,\overline{y},z,\overline{z})\omega^{-\beta_1}\wedge\cdots\wedge\widehat{\omega^{-\beta_j}}\wedge\cdots\wedge\omega^{-\beta_q}.
$$

Then [\(41\)](#page-21-1) implies that

$$
\partial_{\mathfrak{p}^1_+} \Xi|_{B_K/T_{\mathbb{C}} \times V_{\nu} \times U} = \sum_j (-1)^{j-1} \partial_{\overline{y}_j} (f_{j\nu}(0, x_{\nu}, y, \overline{y}, z, \overline{z}) \omega^{\text{nc},1}
$$

\n
$$
= F_{\sigma}(0, x_{\nu}, y, \overline{y}, z, \overline{z}) \omega^{\text{nc},1}|_{B_K/T_{\mathbb{C}} \times V_{\nu} \times U}
$$

\n
$$
= F_{\sigma}(w, x_{\nu}, y, \overline{y}, z, \overline{z}) \omega^{\text{nc},1}|_{B_K/T_{\mathbb{C}} \times V_{\nu} \times U},
$$

where the last equation follows from that F_σ is constant along the fibers of $\pi_\mathfrak{I}$: $W \to \mathcal{I}$. This implies that, when restricted to $\pi'^{-1}(U)$, we have

$$
\partial_{\mathfrak{p}^1_+} \Xi = F_\sigma \omega^{\mathrm{nc},1} |_{\pi'^{-1}(U)}.
$$

By construction, Ξ is constant in $w \in B_K/T_{\mathbb{C}}$, i.e. constant along the fibers of $\pi_{\mathcal{I}}: \, \mathcal{W} \to \mathcal{I}$, and hence descends to $\pi_{\mathcal{U}}^{-1}(U) \subset \mathcal{I}$,

$$
\Xi \in \bigoplus_{1 \le j \le q} \Gamma(\pi_{\mathfrak{U}}^{-1}(U), L_{\mu_j}),
$$

where $\mu_j = \mu + \beta_1 + \cdots + \beta_j + \cdots + \beta_q = \mu + 2\rho_{\text{nc},1} - \beta_j$.

By Borel-Weil-Bott theorem, condition [\(39\)](#page-20-1) implies that on the base cycle Z_o

$$
\Gamma(Z_o, L_{\mu_j}) = 0, 1 \le j \le q
$$

and so does on its small deformations $Z_u, \, u \, \in \, U.$ Since $\pi_{\mathrm{U}}^{-1}(U)$ is covered by the compact analytic subvarieties \tilde{Z}_u , $u \in U$, which are isomorphic to Z_u via the map π_D , we have that

$$
\Gamma(\pi_{\mathfrak{U}}^{-1}(U), L_{\mu_j}) = 0, 1 \le j \le q.
$$

Therefore $\Xi = 0$ and

$$
F_{\sigma}\omega^{\mathrm{nc},1}|_{\pi'^{-1}(U)}=\partial_{\mathfrak{p}^1_+}\Xi=0
$$

on the open subset $\pi^{-1}(U) \subset W$, which implies that $F_{\sigma} = 0$ on W. This proves the injectivity of the Penrose transformation.

We need to consider the case that the Penrose transformation is non-trivial, i.e.

$$
H^0(D', L_{\mu'}) \neq 0.
$$

A necessary condition for this is given as follows.

Theorem 3.6. Let the notations be as Theorem [3.5.](#page-20-0) If $H^0(D', L_{\mu'}) \neq 0$, then

 (42) $(\alpha) \geq 0, \forall \alpha \in \Delta^c_+.$

Proof. Assume on the contrary that there exists an $\alpha \in \Delta_+^{\rm c}$ such that

$$
(43) \qquad \qquad (\mu',\alpha)<0.
$$

Note that

$$
L_{\mu'}|_{Z_o}=K\times_T \mathbb{C}_{\mu'},
$$

where $Z_o = K/T \subset D'$ is the base cycle through the base point. Then from Borel-Weil-Bott theorem, we have that

$$
H^{0}(Z_{o}, L_{\mu'}|_{Z_{o}}) = H^{0}(Z_{o}, K \times_{T} \mathbb{C}_{\mu'}) = 0,
$$

under the assumption [\(43\)](#page-23-0).

For any $g \in G_{\mathbb{R}}$, $g: L_{\mu'}/Z_{\alpha} \to L_{\mu'}/Z_{\alpha}$ gives an isomorphism of line bundles, and hence

$$
H^0(gZ_o, L_{\mu'}|_{gZ_o}) = 0.
$$

Since D' is covered by the cycles gZ_o , $g \in G_{\mathbb{R}}$, we have that $H^0(D', L_{\mu'}) = 0$, which contradicts to the assumption of the theorem. Hence [\(42\)](#page-23-1) holds. \Box

From conditions [\(39\)](#page-20-1) and [\(42\)](#page-23-1) from Theorem [3.5](#page-20-0) and Theorem [3.6,](#page-23-2) it is reasonable to consider the weights μ, μ' satisfying that

(44)
$$
(\mu', \alpha) = 0, \forall \alpha \in \Delta^c_+.
$$

Then the injectivity condition [\(39\)](#page-20-1) of the Penrose transformation is reduced to the conclusion of Lemma [3.8](#page-24-0) below.

By the way, since D is non-classical, the equations

$$
(\lambda,\alpha)=0,\,\alpha\in\Delta_+^{\rm c}
$$

define a linear complex subspace of \mathfrak{h}^* of codimension ≥ 1 . Hence the set Σ of weights μ' satisfying [\(44\)](#page-24-1) is non-empty and once $\mu'_0 \in \Sigma$, then $k\mu'_0 \in \Sigma$ for $k \in \mathbb{Q}$ such that $k\mu_0'$ is a weight.

Theorem 3.7. Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian *type. Let* D′ *be the classical flag domain which is diffeomorphic to* D*. If* µ ′ *is weight such that*

$$
(\mu',\alpha)=0,\,\forall\,\alpha\in\Delta_+^{\rm c},
$$

and $\mu = \mu' - 2\rho_{\rm nc,1}$, then the Penrose transformation

$$
\mathcal{P}: H^0(D', L_{\mu'}) \to H^q(D, L_{\mu})
$$

given by [\(38\)](#page-19-0) *is injective.*

Proof. The theorem follows from Theorem [3.5](#page-20-0) and the following lemma. \Box

Lemma 3.8. *For any* $\beta \in \Delta^{nc,1}_+$ *, there exists an* $\alpha \in \Delta^c_+$ *such that* $(\beta, \alpha) > 0$ *.*

Proof. We first prove the lemma under the assumption that $G_{\mathbb{R}}$ is simple.

From Remark [1.6,](#page-10-4) we have that for any $\beta \in \Delta^{\text{nc},1}_+$, there exists $\gamma \in \Delta^{\text{nc},2}_+$ such that $\beta + \gamma \in \Delta_+^c$ is a root. Then

$$
(\beta, \beta + \gamma) = (\beta, \beta) + (\beta, \gamma).
$$

If $(\beta, \gamma) = 0$, then $(\beta, \beta + \gamma) > 0$. Otherwise, $(\beta, \gamma) < 0$, and

$$
\gamma + \beta, \cdots, \gamma + k\beta
$$
 are roots, where $k = -\frac{2(\beta, \gamma)}{(\beta, \beta)}$.

From Table 1 in Page 45 of [\[16\]](#page-33-11), the possible cases are $k = 1, 2, 3$.

If $k = 1, 3$, then $\gamma + k\beta \in \Delta^c_+$ and

$$
(\beta, \gamma + k\beta) = (\beta, \gamma) + k(\beta, \beta) = \frac{k}{2}(\beta, \beta) > 0.
$$

The remaining case is $k = 2$. Then $(\beta, \beta) = -(\beta, \gamma)$ and $2\beta + \gamma \in \Delta_+^{\text{nc},1}$. Since $\beta, \gamma \in \Delta_+$ and $(\beta, \gamma) < 0$, we can choose another simple roots of Δ starting from $\{\beta, \gamma, \dots\}$. Then according to the classifications of the Dynkin diagrams of the simple roots for Hermitian symmetric spaces in Lemma 4 of [\[23\]](#page-33-12), this corresponds to type B and type C, since the Dynkin diagram does not depend on the choice of the simple roots.

In the case of type C, Lemma 4 of [\[23\]](#page-33-12) implies that the non-compact roots are of the form

$$
\pm e_l \pm e_i, \ 1 \leq i \leq l-1,
$$

where $l = \text{rank } \mathfrak{g}$. But then $(\beta, \beta) = (\gamma, \gamma) = 2$ implies that

$$
k = -\frac{2(\beta, \gamma)}{(\beta, \beta)} = -\frac{2(\beta, \gamma)}{(\gamma, \gamma)} = 1,
$$

which is a contradiction.

Hence the case that $k = 2$ corresponds only to type B. Then Lemma 4 of [\[23\]](#page-33-12) implies that we can choose a set of simple roots from Δ^{nc} . In particular, the set $\{\beta, \gamma, \dots\}$ of simple roots chosen as above is a particular one, with the Dynkin diagram

$$
\circ \qquad \qquad \overbrace{\qquad \qquad \gamma'} \qquad \overbrace{\qquad \qquad }^{\circ \qquad \bullet \qquad \bullet \qquad \bullet \qquad \gamma}
$$

Now if $\Delta_{+}^{\text{nc},2} = \{\gamma\}$, then we can replace $\Delta_{+}^{\text{nc},1}$ with $\Delta_{+}^{\text{nc},2}$ so that $\gamma \in \Delta_{+}^{\text{nc},1}$ + and satisfies $(\gamma, \beta + \gamma) > 0$. Note that $(\gamma, \gamma) > (\beta, \beta)$. Otherwise, there exists $\gamma' \in \Delta^{\text{nc},2}_+$ such that $(\beta, \gamma') < 0$ with

$$
k' = -\frac{2(\beta, \gamma')}{(\beta, \beta)} = 1.
$$

Then the above argument implies that $\beta + \gamma' \in \Delta_+^c$ and $(\beta, \beta + \gamma') > 0$. This proves the lemma when $G_{\mathbb{R}}$ is simple.

In general, the flag domain

$$
D = D^1 \times \cdots \times D^n
$$

with $D^i = G_{\mathbb{R}}^i / T^i$ and $G_{\mathbb{R}}^i$ simple $1 \leq i \leq n$. Then we can take

$$
\Delta_+^{nc,1} = \Delta_+^{1nc,1} \cup \cdots \cup \Delta_+^{nc,1},
$$

where each $\Delta^{i\text{nc},1}_{+}$ satisfies the properties of the lemma. $\hfill \Box$

Remark 3.9. There should be a proof of the lemma without using the classification of the Hermitian symmetric spaces.

4. PENROSE TRANSFORMATION ON COMPACT QUOTIENTS OF FLAG DOMAINS

In this section, we introduce the Penrose transformation on the compact quotients of the flag domains and prove that it is an isomorphism under the conditions given in Section [3](#page-16-0) and the conditions induced by Property **W** of Williams. We also apply our result to the groups of automorphic forms on Hermitian symmetric domains, which can be identified with the higher automorphic cohomology groups of certain line bundles on the compact quotient of the non-classical flag domain.

Let $\Gamma \subset G_{\mathbb{R}}$ be a discrete, co-compact and neat subgroup. Then $X = \Gamma \backslash D$ and $X' = \Gamma \backslash D'$ are compact complex manifolds, and $\mathcal{I}_{\Gamma} = \Gamma \backslash \mathcal{I}$, $\mathcal{U}_{\Gamma} = \Gamma \backslash \mathcal{U}$ and $W_{\Gamma} = \Gamma \backslash W$ are complex manifolds. Moreover diagram [\(29\)](#page-16-1) induces the

following commutative diagram

The homogenous line bundles $L_{\mu} = G_{\mathbb{R}} \times_T \mathbb{C}_{\mu}$ and $L_{\mu'} = G_{\mathbb{R}} \times_T \mathbb{C}_{\mu'}$ on D and D' are $G_{\mathbb{R}}$ -invariant, and hence desecnd to the line bundles on X and X' respectively. We still denote them by L_{μ} and $L_{\mu'}$ for convenience.

The d_{π} -closed form

 $\omega^{\text{nc},1} = \omega^{-\beta_1} \wedge \cdots \wedge \omega^{-\beta_q} \in \Gamma(\mathcal{W}, \Omega_\pi^q(L_{-2\rho_{\text{nc},1}}))$

is left invariant, and descends to the form on W_{Γ} ,

$$
\omega_{\Gamma}^{\text{nc},1} \in \Gamma(\mathcal{W}_{\Gamma}, \Omega_{\pi}^q(L_{-2\rho_{\text{nc},1}})).
$$

In this section, we mainly consider the automorphic cohomology groups $H^*(X,L_\mu)$ and $H^*(X', L_{\mu'})$. From Page 236 of [\[11\]](#page-33-0), we know that $W_{\Gamma} = \Gamma \backslash W$ is also Stein. Applying Theorem [2.5](#page-13-1) of EGW again, we have that

$$
H^*(X, L_\mu) \cong H^*_{DR}(\mathcal{W}_{\Gamma}, \Omega^\bullet_\pi(L_\mu))
$$

$$
H^*(X', L_{\mu'}) \cong H^*_{DR}(\mathcal{W}_{\Gamma}, \Omega^\bullet_{\pi'}(L_{\mu'})).
$$

It is important to note that the arguments for the main results in Section [3](#page-16-0) are local. Consequently, we introduce the following definition of the Penrose transformation P on the compact quotients of flag domains and present the corresponding theorem establishing the injectivity of P.

Lemma-Definition 4.1. Let μ and μ' be two weights such that $\mu + \rho = \mu' + \rho'$. *Then the Penrose transformation on automorphic cohomology groups*

$$
\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})
$$

is defined by

(46)
$$
H^{0}(X', L_{\mu'}) \xrightarrow{\mathcal{P}} H^{q}(X, L_{\mu})
$$
\n
$$
\downarrow \cong
$$
\n
$$
H^{0}_{DR}(\mathcal{W}_{\Gamma}, \Omega^{\bullet}_{\pi'}(L_{\mu'})) \xrightarrow{\omega_{\Gamma}^{\text{nc},1}} H^{q}_{DR}(\mathcal{W}_{\Gamma}, \Omega^{\bullet}_{\pi}(L_{\mu})).
$$

We then have the following theorem.

Theorem 4.2. Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian *type. Let* D' *be the classical flag domain which is diffeomorphic to D. Let* $X = \Gamma \backslash D$ *and* $X' = \Gamma \backslash D'$ *be their corresponding quotients by a discrete, co-compact and neat subgroup* $\Gamma \subset G_{\mathbb{R}}$ *.*

If μ *and* μ' *are two weights with* $\mu + \rho = \mu' + \rho'$ *, and satisfy that for any* $\beta \in \Delta^{nc,1}_+$ + there exists $\alpha_\beta \in \Delta_+^{\rm c}$ such that

(39)
$$
(\alpha_{\beta}, \mu + 2\rho_{\text{nc},1} - \beta) = (\alpha_{\beta}, \mu' - \beta) < 0,
$$

Then the Penrose transformation on automorphic cohomology groups

 $\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})$

given by [\(46\)](#page-26-1) *is injective.*

Furthermore we have the following theorem.

Theorem 4.3. Let the notations be as Theorem [4.2.](#page-26-0) If μ and μ' are two weights *such that* $\mu + \rho = \mu' + \rho'$ and

$$
(\mu', \alpha) = 0, \,\forall \,\alpha \in \Delta^c_+.
$$

Then the Penrose transformation on automorphic cohomology groups

 $\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})$

given by [\(46\)](#page-26-1) *is injective.*

We are interested in the case that the Penrose transformation

(47)
$$
\mathcal{P}: H^0(X', L_{\mu'}) \xrightarrow{\cong} H^q(X, L_{\mu})
$$

is an isomorphism. Under Theorem [4.2,](#page-26-0) [\(47\)](#page-27-0) is reduced to proving that

(48)
$$
\dim_{\mathbb{C}} H^{0}(X', L_{\mu'}) = \dim_{\mathbb{C}} H^{q}(X, L_{\mu}).
$$

To get a natural condition for [\(48\)](#page-27-1), we introduce the work of Schmid, Griffiths, Williams and et al. on the geometric realizations of representations of semisimple real Lie groups. Good references for this survey are Schmid's paper [\[22\]](#page-33-9) and Lecture 5 and Lecture 9 of [\[11\]](#page-33-0). One is referred to [\[18\]](#page-33-13) for the basic notions in this area and more details.

First we fix some notations. Let $\widehat{G}_{\mathbb{R}}$ be the set of equivalence classes of irreducible unitary representations

 $\pi: G_{\mathbb{R}} \to \mathrm{Aut}(\tilde{V}_{\pi})$

of $G_\mathbb{R}$ on a Hilbert space \tilde{V}_π . Then the subspace $V_\pi~=~\tilde{V}_{\pi,K-\text{finite}}$ of $K\text{-finite}$ vectors in \tilde{V}_{π} is a Harish-Chandra module, which is called the Harish-Chandra module associated to the unitary representation $\tilde{V}_\pi.$ We also denote the unitary representation \tilde{V}_{π} and its associated Harish-Chandra module V_{π} by \tilde{V}_{ζ} and V_{ζ} respectively, if V_π has infinitesimal character $\chi_\zeta.$

The L^2 -cohomology group $H_{(2)}^k(D,L_\mu)$ of the homogeneous line bundle L_μ on D can be realized as

$$
H^k_{(2)}(D,L_\mu)=\int_{\widehat{G}_{\mathbb{R}}}\tilde{V_{\pi}}\widehat{\otimes} H^k(\mathfrak{n}_+,\tilde{V_{\pi}}^*)_{-\mu}
$$

via the Plancherel decomposition

$$
L^2(G_{\mathbb{R}}) = \int_{\widehat{G}_{\mathbb{R}}} \tilde{V}_{\pi} \widehat{\otimes} \tilde{V}_{\pi}^* d\pi,
$$

where $\tilde{V_{\pi}}$ * is the dual space of $\tilde{V_{\pi}}$, and $d\pi$ is the Plancherel measure which assigns positive mass to the discrete series representations. Here $H^k(\mathfrak{n}_+, \tilde{V_\pi})$ $(*)$ is the ncohomology and $H^k(\mathfrak{n}_+, \tilde{V_\pi})$ *)_{-µ} is the set of elements in $H^k(\mathfrak{n}_+,\tilde{V}_\pi)$ $\sum_{n=0}^{\infty}$ on which \mathfrak{h} acts by the weight $-\mu$.

The celebrated work of Schmid on the solution to the Langlands conjecture can be summarized as follows.

Theorem 4.4 (Schmid). Let L_{μ} be homogeneous line bundle on the flag domain $D = G_{\mathbb{R}}/T$ defined by a weight μ . Then we have the following realization of the *representation of* $G_{\mathbb{R}}$ *.*

(1) When $\mu + \rho$ *singular, i.e.* $\exists \alpha \in \Delta_+$ *s.t.* $(\mu + \rho, \alpha) = 0$ *, the L*²-cohomology group $H_{(2)}^{k}(D,L_{\mu})$ *is zero;*

(2) When $\mu + \rho$ *is regular, i.e.* $(\mu + \rho, \alpha) \neq 0$ *for any* $\alpha \in \Delta_+$ *, the L*²-cohomology group $H^k_{(2)}(D,L_\mu)$ is zero if $k\neq q(\mu+\rho)$, and $H^{q(\mu+\rho)}_{(2)}(D,L_\mu)$ is a discrete series *representation of* $G_{\mathbb{R}}$, whose associated Harish-Chandra module has infinitesimal *character* $\chi_{\mu+\rho}$ *;*

(3) When $\mu + \rho$ *belongs to the anti-dominant Weyl chamber, the cohomology* group $H^d(D, L_{\mu})$ is a Harish-Chandra module with infinitesimal character $\chi_{\mu+\rho}$, *where* $d = \dim_{\mathbb{C}} Z_o$.

The automorphic cohomology $H^k(X, L_\mu)$ on the compact quotient $X = \Gamma \backslash D$ can be realized as the n-cohomology:

$$
H^k(X, L_\mu) = \bigoplus_{\pi \in \widehat{G}_{\mathbb{R}}} H^k(\mathfrak{n}_+, V_\pi)_{-\mu}^{\oplus m_\pi(\Gamma)},
$$

where $m_{\pi}(\Gamma)$ is the multiplicity of the associated Harish-Chandra module V_{π} in $L^2(\Gamma \backslash G_{\mathbb{R}})$. Therefore $\dim_{\mathbb{C}} H^k(X, L_{\mu})$ is determined by

$$
\dim_{\mathbb{C}} H^k(\mathfrak{n}_+,V_\pi)^{\oplus m_\pi(\Gamma)}, \ \ \pi \in \widehat{G}_{\mathbb{R}}.
$$

For the n-cohomology, we have the following Williams Lemma from [\[24\]](#page-33-14) as introduced in Lecture 8 of [\[11\]](#page-33-0).

Lemma 4.5 (Williams Lemma). Let \tilde{V} be an irreducible unitary representation of $G_{\mathbb{R}}$ and V be its associated Harish-Chandra module. If μ is a weight satisfying

(1) $\mu + \rho$ *is regular*;

(2) Property W: For each $\beta \in \Delta^{nc}$ *with* $(\mu + \rho, \beta) > 0$ *, we have that*

$$
\left(\mu+\rho-\frac{1}{2}\sum_{\substack{\alpha\in\Delta\\\mu+\rho,\alpha>0}}\alpha,\,\beta\right)>0.
$$

Then $H^k(\mathfrak{n}_+, V)_{-\mu} \neq 0$ implies that

$$
\begin{cases} k=q(\mu+\rho)\\ \dim_{\mathbb{C}}H^k(\mathfrak{n}_+,V)_{-\mu}=1\\ \tilde{V}=\tilde{V}_{-(\mu+\rho)}, \end{cases}
$$

where $\tilde{V}_{-(\mu+\rho)}$ is a discrete series representation with infinitesimal character $\chi_{-(\mu+\rho)}.$

Another result we need is the following theorem.

Theorem 4.6 (Williams, Theorem 2.4 in [\[25\]](#page-33-5)). Let $X = \Gamma \backslash D$ ba a compact *quotient of the flag domain* $D = G_{\mathbb{R}}/T$ *and* L_{μ} *be a locally homogenous line bundle on X* defined by the weight μ . Assume that $\mu + \rho$ is regular and satisfies Property *W of Williams in Lemma [4.5.](#page-28-0) Then*

$$
\dim_{\mathbb{C}} H^{q(\mu+\rho)}(X, L_{\mu}) = m_{-(\mu+\rho)}(\Gamma),
$$

the multiplicity of $V_{-(\mu+\rho)}$ *in* $L^2(\Gamma \backslash G_{\mathbb{R}})$ *, and* $H^k(X, L_\mu) = 0$ *for* $k \neq q(\mu+\rho)$ *.*

With the above preparations we can prove the following theorem.

Theorem 4.7. Let $D = G_{\mathbb{R}}/T$ be a non-classical flag domain with $G_{\mathbb{R}}$ of Hermitian *type. Let* D' *be the classical flag domain which is diffeomorphic to* D *. Let* $X = \Gamma \backslash D$ *and* ^X′ = Γ\D′ *be their corresponding quotients by a discrete, co-compact and neat subgroup* $\Gamma \subset G_{\mathbb{R}}$ *.*

Let μ and μ' be two weights such that $\mu + \rho = \mu' + \rho'$ is regular and $\mu + \rho = \mu' + \rho'$ *lies in the Weyl chamber determined by the set of roots*

(49)
$$
\Delta_+^c \cup \Delta_+^{nc,1} \cup (-\Delta_+^{nc,2}) = \Delta_+^c \cup (-\Delta_+^{nc}) .
$$

If for any $\beta \in \Delta_+^{\rm nc,1}$ *there exists an* $\alpha_\beta \in \Delta_+^{\rm c}$ *such that*

(39)
$$
(\alpha_{\beta}, \mu + 2\rho_{\text{nc},1} - \beta) = (\alpha_{\beta}, \mu' - \beta) < 0,
$$

and moreover,

(50)
$$
(\mu + 2\rho_{\text{nc},2}, \Delta_+^{\text{nc},1} \cup (-\Delta_+^{\text{nc},2})) = (\mu' + 2\rho'_{\text{nc}}, -\Delta_+^{\text{nc}}) > 0,
$$

then the Penrose transformation

 $\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})$

given by [\(46\)](#page-26-1) *is an isomorphism.*

Proof. Since $\zeta = \mu + \rho = \mu' + \rho'$ is regular and lies in the Weyl chamber determined by the set of roots in [\(49\)](#page-29-1), we have that

(51)
$$
(\zeta, \alpha) > 0, \forall \alpha \in \Delta^c_+; (\zeta, \beta) > 0, \forall \beta \in \Delta^{nc,1}_+; (\zeta, \gamma) < 0, \forall \gamma \in \Delta^{nc,2}_+,
$$

which implies that

$$
q'(\mu' + \rho') = 0, \, q(\mu + \rho) = q,
$$

where $q = #\Delta_+^{\mathrm{nc},1}$.

From Theorem [4.2,](#page-26-0) the Penrose transformation

$$
\mathcal{P}: H^0(X', L_{\mu'}) \to H^q(X, L_{\mu})
$$

is injective. Hence we need to prove that

$$
\dim_{\mathbb{C}} H^0(X', L_{\mu'}) = \dim_{\mathbb{C}} H^q(X, L_{\mu}).
$$

For this we only need to check that conditions in [\(50\)](#page-29-2) is equivalent to Property **W** of Williams in Lemma [4.5,](#page-28-0) since then Theorem [4.6](#page-29-3) implies that

$$
\dim_{\mathbb{C}} H^0(X', L_{\mu'}) = m_{-(\mu' + \rho')}(\Gamma) = m_{-(\mu + \rho)}(\Gamma) = \dim_{\mathbb{C}} H^q(X, L_{\mu}).
$$

Note that under [\(51\)](#page-29-4) the conditions in Property **W**:

$$
\beta \in \Delta^{\text{nc}} \text{ with } (\mu + \rho, \beta) > 0 \Longleftrightarrow \beta \in \Delta_+^{\text{nc},1} \cup \left(-\Delta_+^{\text{nc},2} \right) = -\Delta_+^{\text{nc}}
$$

and

$$
\frac{1}{2} \sum_{\substack{\alpha \in \Delta \\ (\mu + \rho, \alpha) > 0}} \alpha = \frac{1}{2} \sum_{\alpha \in \Delta_+^c} \alpha + \frac{1}{2} \sum_{\beta \in \Delta_+^{\text{nc},1}} \beta - \frac{1}{2} \sum_{\gamma \in \Delta_+^{\text{nc},2}} \gamma
$$

$$
= \rho_c + \rho_{\text{nc},1} - \rho_{\text{nc},2} = \rho_c - \rho_{\text{nc}}'.
$$

Hence [\(50\)](#page-29-2) is equivalent to Property **W**.

Due to its significance in arithmetic geometry and number theory, we consider the homogenous line bundle $L_{\mu'_c}$ defined as the pull-back of the canonical bundle $\omega_{\mathbb{B}}$ via the holomorphic projection map

$$
p': D' \to G_{\mathbb{R}}/K \cong \mathbb{B},
$$

which is still denoted by $L_{\mu'_c}=\omega_{\mathbb B}\to D'.$ Since $\mathrm{T}^{1,0}_oG_{\mathbb R}/K\cong\mathfrak{p}'_-$, we have that

$$
\mu'_{c} = -\sum_{\beta \in \Delta'^{\text{nc}}_{+}} \beta = 2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}.
$$

Let k_0 the maximal positive integer such that $\mu'_{c0} \triangleq \mu'_{c}/k_0$ is still a weight. Then

$$
L_{\mu'_{c0}} = \omega_{\mathbb{B}}^{\otimes 1/k_0} \to D'
$$

is a well-defined line bundle on D' . Let

$$
\mu'_{k} = k\mu'_{c0} = \frac{k}{k_0} (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}), \ \mu_{k} = \mu'_{k} - 2\rho_{\text{nc},1}.
$$

Then

$$
L_{\mu'_k} = \omega_{\mathbb{B}}^{\otimes k/k_0} \to D', L_{\mu_k} \to D
$$

are two line bundles whose cohomology groups are Penrose related.

Theorem 4.8. *Let the notations be as above. Then there exists a positive integer* N such that for $k > N$, the Penrose transformation on automorphic cohomology *groups*

(52)
$$
\mathcal{P}: H^0(X', \omega_{\mathbb{B}}^{\otimes k/k_0}) \to H^q(X, L_{\mu_k})
$$

is an isomorphism. Therefore $H^q(X, L_{\mu_k})$ is isomorphic to the group

$$
H^0(\Gamma\backslash G_{\mathbb{R}}/K,\omega_{\mathbb{B}}^{\otimes k/k_0})
$$

of automorphic forms on $G_{\mathbb{R}}/K$.

Proof. The proof is divided into the following steps. **Step 1.** We first prove that

$$
(2\rho_{\mathop{\mathrm{nc}},1}-2\rho_{\mathop{\mathrm{nc}},2},\alpha)=-\sum_{\beta\in\Delta'^{\mathop{\mathrm{nc}}}_+}(\beta,\alpha)=0,\,\forall\,\alpha\in\Delta_+^c.
$$

Then $(\mu'_k, \alpha) = 0$, for any $\alpha \in \Delta^c_+$ and Theorem [4.3](#page-27-2) implies that the Penrose transformation [\(52\)](#page-30-1) is injective.

First note that

$$
[\mathfrak{k}_\pm,\mathfrak{p}_-']\subset \mathfrak{p}_-'\Longleftrightarrow \forall\,\alpha\in\Delta_+^c,\beta\in\Delta_+^{\prime\mathrm{nc}}, \beta\pm\alpha\in\Delta_+^{\prime\mathrm{nc}}\text{ or is not a root.}
$$

Now we fix any $\alpha \in \Delta_+^c$. If there exists $\beta \in \Delta_+^{\text{nc}}$ such that $(\beta, \alpha) \neq 0$, then we have either that $(\beta,\alpha)>0,$ which implies that

$$
\beta, \beta - \alpha, \cdots, \beta - k\alpha
$$
 are roots in Δ'^{nc}_+ , where $k = \frac{2(\beta, \alpha)}{(\alpha, \alpha)}$,

and

$$
(\beta + (\beta - \alpha) + \dots + (\beta - k\alpha), \alpha) = (k+1) (\beta, \alpha) - \frac{k(k+1)}{2} (\alpha, \alpha) = 0,
$$

or that $(\beta, \alpha) < 0$, which implies that

$$
\beta, \beta + \alpha, \cdots, \beta + k\alpha
$$
 are roots in Δ'^{nc}_{+} , where $k = \frac{-2(\beta, \alpha)}{(\alpha, \alpha)}$,

and

$$
(\beta + (\beta + \alpha) + \dots + (\beta + k\alpha), \alpha) = (k+1) (\beta, \alpha) + \frac{k(k+1)}{2} (\alpha, \alpha) = 0.
$$

Therefore

$$
\sum_{\substack{\beta \in \Delta'^{\mathrm{nc}}_{+} \\ (\beta, \alpha) \neq 0}} (\beta, \alpha) = 0
$$

and

$$
(2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}, \alpha) = -\sum_{\substack{\beta \in \Delta'^{\text{nc}}_{+} \\ (\beta,\alpha) = 0}} (\beta, \alpha) - \sum_{\substack{\beta \in \Delta'^{\text{nc}}_{+} \\ (\beta,\alpha) \neq 0}} (\beta, \alpha) = 0.
$$

Step 2. Lemma [1.4](#page-9-2) implies that

$$
(\beta, \beta') \ge 0, (\beta, \gamma) \le 0, (\gamma, \gamma') \ge 0, \forall \beta, \beta' \in \Delta^{\mathrm{nc}, 1}_{+}, \forall \gamma, \gamma' \in \Delta^{\mathrm{nc}, 2}_{+}.
$$

Hence

$$
(2\rho_{\mathrm{nc},1}-2\rho_{\mathrm{nc},2},\beta)=\left(\beta+\sum_{\beta'\in\Delta_+^{\mathrm{nc},1},\,\beta'\neq\beta}\beta'-\sum_{\gamma\in\Delta_+^{\mathrm{nc},2}}\gamma,\,\beta\right)>0,\,\forall\,\beta\in\Delta_+^{\mathrm{nc},1}
$$

and similarly

$$
(2\rho_{\mathrm{nc},1} - 2\rho_{\mathrm{nc},2}, \gamma) < 0, \,\forall \,\gamma \in \Delta_+^{\mathrm{nc},2}.
$$

Step 3. For

$$
\mu' + \rho' = \rho_c + \left(\frac{k}{k_0} - \frac{1}{2}\right) (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}),
$$

Step 1 and Step 2 imply that there exists a positive integer N_1 such that for $k > N_1$,

(53)
$$
(\mu' + \rho', \alpha) = (\rho_c, \alpha) > 0, \forall \alpha \in \Delta^c_+;
$$

(54)
$$
(\mu' + \rho', \beta) = \left(\frac{k}{k_0} - \frac{1}{2}\right) (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}, \beta) + (\rho_c, \beta) > 0, \forall \beta \in \Delta_+^{\text{nc},1};
$$

(55)
$$
(\mu' + \rho', \gamma) = \left(\frac{k}{k_0} - \frac{1}{2}\right) (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}, \gamma) + (\rho_c, \gamma) < 0, \forall \gamma \in \Delta_+^{\text{nc},2}.
$$

This implies that $\mu + \rho = \mu' + \rho'$ is regular and lies in the Weyl chamber determined by the set of roots in [\(49\)](#page-29-1).

Similarly, there exists a positive integer N with $N \geq N_1$ such that for $k \geq N$,

(56)
$$
(\mu' + 2\rho'_{\rm nc}, -\Delta'^{\rm nc}_{+}) = \left(\frac{k}{k_0} - 1\right) (2\rho_{\rm nc,1} - 2\rho_{\rm nc,2}, \Delta^{\rm nc,1}_{+} \cup \left(-\Delta^{\rm nc,2}_{+}\right)) > 0.
$$

Hence [\(50\)](#page-29-2) is satisfied.

Finally, by Theorem [4.7,](#page-29-0) Step 1 and Step 3 imply that the Penrose transformation [\(52\)](#page-30-1) is an isomorphism.

Remark 4.9. According to (53) , (54) , (55) and (56) , the minimum of N in Theorem [4.8](#page-30-0) is the least positive integer k such that

$$
\left(\frac{k}{k_0} - 1\right) (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}, \beta) + (\rho_c, \beta) > 0, \forall \beta \in \Delta_+^{\text{nc},1};
$$

$$
\left(\frac{k}{k_0} - 1\right) (2\rho_{\text{nc},1} - 2\rho_{\text{nc},2}, \gamma) + (\rho_c, \gamma) < 0, \forall \gamma \in \Delta_+^{\text{nc},2}.
$$

REFERENCES

- [1] R. J. Baston, and M. G. Eastwood, *The Penrose transform: its interaction with representation theory*, Courier Dover Publications, (2016).
- [2] R. Bott, Homogeneous vector bundles, *Annals of Math.*, **66** (1957), pp. 203–248.
- [3] D. Burns, S. Halverscheid, and R. Hind, The geoemtry of Grauert tubes and complexification of symmetric spaces, *Duke J. Math.*, **118** (2003), pp. 465–491.
- [4] H. Carayol, Limites dégénérées de séries discrètes, formes automorphes et variétés de Griffiths-Schmid: le cas du groupe U(2, 1), *Compos. Math.*, **111** (1998), pp. 51–88.
- [5] H. Carayol, Quelques relations entre les cohomologies des variétés de Shimura et celles de Griffiths-Schmid (cas du groupe SU(2, 1)), *Compos. Math.*, **121** (2000), pp. 305–335.
- [6] H. Carayol, Cohomologie automorphe et compactifications partielles de certaines variétés de Griffiths-Schmid, *Compos. Math.*, **141** (2005), pp. 1081–1102.
- [7] J. Carlson and D. Toledo, Compact quotients of non-classical domains are not Kähler, *Contemporary Mathematics,* **608** (2014), pp. 51–57.
- [8] M. Eastwood, S. Gindikin, and H. Wong, Holomorphic realizations of ∂-cohomology and constructions of representations, *J. Geom. Phys.*, **17**(3) (1995), pp. 231–244.
- [9] G. Fels, A. Huckleberry and J. A. Wolf, *Cycle Spaces of Flag Domains, Progress in Mathematics*, 245, Birkhäuser, (2006).
- [10] M. Green, P. Griffiths and M. Kerr, *Mumford-Tate Groups and Domains: Their Geometry and Arithmetic*, Annals of Math. Studies, **183**, Princeton University Press, Princeton, NJ, (2012).
- [11] M. Green, P. Griffiths and M. Kerr, *Hodge Theory, Complex Geometry, and Representation Theory*, Regional Conference Series in Mathematics, Volume **118**, American Mathematical Society, (2013).
- [12] M. Green, P. Griffiths and M. Kerr, Special values of automorphic cohomology classes, Vol. **231**, no. 1088, American Mathematical Society, (2014).
- [13] P. Griffiths and W. Schmid, Locally homogeneous complex manifolds, *Acta Math.*, **123** (1969), pp. 253–302.
- [14] P. Griffiths, C. Robles and D. Toledo, Quotients of non-classical flag domains are not algebraic, *Algebraic Geometry*, **1** (2014), pp. 1–13.
- [15] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces,* Academic Press, New York, (1978).
- [16] J. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag New York, (1972).
- [17] A. W. Knapp, *Lie groups beyond an introduction*, Progress in Mathematics, Vol. **140**, Boston: Birkhäuser, (1996).
- [18] A. W. Knapp, *Representation Theory of Semisimple Groups: An Overview Based on Examples*, Princeton Univ. Press, Princeton, NJ, (1986).
- [19] K. Kodaira and D. C. Spencer, On Deformations of Complex Analytic Structures, III, *Annals of Mathematics,* Second Series, **71**(1) (1960), pp. 43–76.
- [20] K. Liu and Y. Shen, Geometry of non-classical period domains, *[arXiv:2405.16536,](http://arxiv.org/abs/2405.16536)* (2024).
- [21] W. Schmid, Homogeneous complex manifolds and representations of semisimple Lie groups, Ph.D. dissertation, Univ. California, Berkeley (1967); reprinted in *Representation Theory and Harmonic Analysis on Semisimple Lie Groups*, Math. Surveys and Monogr. **31**, Amer. Math. Soc. (1989), pp. 223–286.
- [22] W. Schmid, Discrete series, *Proc. Symp. Pure Math.*, **61** (1997), pp. 83–113.
- [23] J. A. Wolf, On the Classification of Hermitian Symmetric Spaces, *Journal of Mathematics and Mechanics*, Vol. **13**, No. 3 (1964), pp. 489–495
- [24] F. Williams, Discrete series multiplicities in $L^2(\Gamma \backslash G)$ (II). Proof of Langlands conjecture, *Amer. J. Math.*, **107** (1985), pp. 367–376.
- [25] F. Williams, The n-cohomology of limits of discrete series, *J. Funct. Anal.*, **80** (1988), pp. 451–461.

MATHEMATICAL SCIENCES RESEARCH CENTER, CHONGQING UNIVERSITY OF TECHNOLOGY, CHONGQING 400054, CHINA;

DEPARTMENT OF MATHEMATICS,UNIVERSITY OF CALIFORNIA AT LOS ANGELES, LOS ANGELES, CA 90095-1555, USA

Email address: liu@math.ucla.edu

MATHEMATICAL SCIENCES RESEARCH CENTER, CHONGQING UNIVERSITY OF TECHNOLOGY, CHONGQING 400054, CHINA

Email address: syliuguang2007@163.com

