Quark Model Study of Doubly Heavy Ξ and Ω Baryons via Deep Neural Network and Hybrid Optimization

Zahra Ghalenovi^{1*}, Masoumeh Moazzen Sorkhi¹ and Amir Hossein Sovizi² ¹Department of Physics, Kosar University of Bojnord, Bojnourd 94156-15458, Iran ²Yadegar Emam, Shahre Rey Branch, Islamic Azad University, Tehran 18155-144, Iran

Abstract

In the past decades, research on the doubly heavy baryons developed rapidly in both experiments and theories. In this work we study the mass spectra and semileptonic decay widths of the doubly heavy Ξ and Ω baryons in the framework of the hypercentral constituent quark model. We solve the six-dimensional Schrödinger equation via deep learning and particle swarm optimization techniques to improve the speed and accuracy. Then, we predict the masses of the ground and excited states of single and doubly heavy baryons. Working close to the zero recoil point, we also study the $b \rightarrow c$ semileptonic decay widths and branching ratios of doubly heavy Ξ and Ω baryons. Our results are compared to the exist experimental data and other theoretical evaluations. Our predictions of mass spectrum and decay widths provide valuable information for the experiment searching for undiscovered heavy baryon states.

1 Introduction

The spectroscopy of doubly heavy baryons has become one of the hot topics in hadronic physics. During the last few years significant progress has been made in heavy baryon physics and many theoretical progresses have been achieved in doubly and triply heavy baryon spectroscopy [1, 2, 3]. The mass spectrum and width are two determinative quantities for theoretical and experimental researches and theoretical studies for the mass spectra and decay widths of heavy baryons are of special interest. The first observation of doubly heavy baryons was reported by the SELEX Collaboration for doubly heavy $M_{\Xi_{222}^{++}}$ baryon. They reported $M_{\Xi_{222}^{++}} = 3518 \pm 1.7$ MeV [4]. Later, the LHCb Collaboration reported the new baryon mass for the Ξ_{cc}^{++} state as $3621.40 \pm 0.72 \pm 0.14 \pm 0.27$ MeV and $3620.6 \pm 1.5 \pm 0.4 \pm 0.3$ MeV obtained in different decay modes[5, 6]. Recently, the LHCb collaboration has carried out new searches for the Ξ_{bc} and Ω_{bc} doubly heavy baryons with a data sample of proton-proton collisions, but these baryon states are not yet to be observed [7, 8, 9]. Although the Ξ_{bc} , Ω_{bc} , Ξ_{bb} and Ω_{bb} states has not been yet found in experiments, they are expected to be observed in the near future [10, 11]. The production of doubly heavy baryons can be done at the LHC or future high luminosity $e^+e^$ colliers. For the singly heavy baryons, mass spectra and some semileptonic decays have been measured in experiments [12, 13, 14, 15, 16, 17, 18, 19, 20, 21]. In comparison to the doubly charmed Ξ_{cc} baryon, searching for the other doubly heavy baryon states are more complicated. The search for these states is really hard for the experiment since higher energy and higher beam luminosity are required to produce them.

For the semileptonic decay widths of doubly heavy baryons no experimental data are also reported and only a limited number of theoretical calculations are available. As there are more doubly heavy baryons that may be obtained by experiment, proposing a theoretical model for their structures is

 $^{^{\}ast}z_ghalenovi@kub.ac.ir$

essential [22, 23, 24, 25, 26, 27, 28, 28, 29, 30, 31].

In recent years, machine learning has been the most popular computational approach for different fields of modern science and a significant raise of interest has been seen in that area. The machine learning techniques enable us to improve the performance and accuracy in theory and experiment.

The solution of three-body Schrödinger equation is complicated in hadron physics. The deep neural network (DNN) as a subfield of machine learning techniques provides a solution to this problem. Deep learning algorithms can identify the energy eigenvalues of the baryon systems. Use of neural networks for solving differential equations has some advantages [36, 37]. One point is that by increasing the dimensions of problem, the computational complexity does not increase considerably. The other advantage is that the solution of the problem is continuous over the domain of integration [32]. In this study, we present a mathematical model using the DNN combining with particle swarm optimization algorithm (PSO) which is fast and provides high accuracy, to estimate the masses of the single and doubly heavy baryons. Some earlier works studied the heavy hadron properties via neural networks and deep learning techniques [32, 33, 34, 35]. Ref. [32] solved the six-dimensional nonrelativistic Schrödinger equation via an artificial neural network and obtained the mass spectra of heavy Ω_b baryon states in a hypercentral approach. Ref. [33] used the neural networks and Gaussian processes to calculate the masses of the light baryons, charmonium and pentaquark states. The authors of Ref. [34] presented two approaches using the deep neural networks to predict the mass and width of the ordinary and exotic mesons. Ref. [35] solved five-body Schrödinger equation of hidden-charm pentaquark resonances by using the artificial neural network method and made predictions of parities for these states.

In this work, we study the single and doubly heavy baryon states in a hypercentral constituent quark model. Firstly, we introduce our model and simplify the three-body Schrödinger equation using the hypercentral model. Then, by applying the deep learning techniques, we evaluate the energy eigenvalues of the considered baryon systems. In our method, we also use the particle swarm optimization to get the best values of the energies. In the next step, we calculate the mass spectra of the ground and orbitally excited states of the single and doubly heavy baryons and compare our results to the exist experimental data and other predictions. The exist experimental data of the single heavy baryon masses help us to optimize our mass evaluations of the Ξ and Ω doubly heavy baryons. Working close to the zero recoil point, we simplify the form factors of $b \to c$ semileptonic decays and evaluate the semileptonic decay widths and branching ratios of the doubly heavy Ξ and Ω baryons. It helps us to get a deep understanding of their structure. We consider the $s = \frac{1}{2} \to s = \frac{1}{2}$ transitions including $\Xi_{bb} \to \Xi_{bc}, \Xi_{bc} \to \Xi_{cc}, \Omega_{bb} \to \Omega_{bc}, \text{ and } \Omega_{bc} \to \Omega_{cc}$ decays. The triplet doubly heavy baryons with $s = \frac{3}{2}$ are dominated by the strong or electromagnetic decays and due to the smallness of the weak coupling, their weak decays can not be observed in the experiments. Therefore, we ignore the calculations for the semileptonic decays of $s = \frac{3}{2}$ doubly heavy baryons.

The structure of this paper is in the following form. In Section 2 we introduce the three-body hypercentral quark model and present our potential model. In Section 3 the deep neural networks is briefly introduced. Our predictions for the ground and also orbitally excited states of the doubly heavy baryons are presented in Section 5. In Section 6 by introducing a universal function as the form factor of the considered transition, we calculate the semileptonic decay widths and corresponding branching fractions of doubly charm and bottom heavy baryons. A short summary is given in Section 7.

2 Phenomenological model

In order to obtain the heavy baryon masses we need to calculate the energy eigenvalues and also the eigenfunctions of the baryon states. We study the heavy baryons and therefore, the non-relativistic formalism is an appropriate method. We consider the baryon system as a bound state of three constituent quarks. Using the hypercentral constituent quark model [38, 39] we can simplify the three-body problem of the baryon states. The configurations of three quarks can be described by the Jacoobi

coordinates, ρ and λ ,

$$\vec{\rho} = \frac{1}{\sqrt{2}}(\vec{r_1} - \vec{r_2}), \qquad \vec{\lambda} = \frac{1}{\sqrt{6}}(\vec{r_1} + \vec{r_2} - 2\vec{r_3}) \tag{1}$$

such that

$$m_{\rho} = \frac{2m_1m_2}{m_1 + m_2}, \qquad m_{\lambda} = \frac{2m_3(m_1^2 + m_2^2 + m_1m_2)}{(m_1 + m_2)(m_1 + m_2 + m_3)}$$
(2)

Here m_1 , m_2 and m_3 are the constituent quark masses. We introduce the hyperspherical coordinates, hyperradius x and hyperangle ζ defined in terms of the Jacoobi coordinates ρ and λ by

$$x = \sqrt{\rho^2 + \lambda^2}, \qquad \xi = \arctan(\sqrt{\frac{\rho}{\lambda}}).$$
 (3)

The Hamiltonian of three-body baron system takes the form

$$H = \frac{p_{\rho}^2}{2m_{\rho}} + \frac{p_{\lambda}^2}{2m_{\lambda}} + V(x).$$
(4)

and the kinetic energy is given by $(\hbar = c = 1)$

$$-\left(\frac{\Delta_{\rho}}{2m_{\rho}} + \frac{\Delta_{\lambda}}{2m_{\lambda}}\right) = -\frac{1}{2\mu} \left(\frac{d^2}{dx^2} + \frac{5}{x}\frac{d}{dx} - \frac{L^2(\Omega_{\rho}, \Omega_{\lambda}, \xi)}{x^2}\right),\tag{5}$$

where $\mu = \frac{2m_{\rho}m_{\lambda}}{m_{\rho} + m_{\lambda}}$ is the reduced mass and Ω_{ρ} and Ω_{λ} are the angels of hyperspherical coordinates. The wave function of a three-body system can be defined in terms of the hyperspherical harmonic basis:

$$\Psi(\rho,\lambda) = \Sigma_{\gamma,l_{\rho},l_{\lambda}} N_{\gamma} \psi_{\nu,\gamma}(x) Y_{[\gamma],l_{\rho},l_{\lambda}}(\Omega_{\rho},\Omega_{\lambda},\xi).$$
(6)

where γ is the grand angular quantum number and ν counts the nodes of the wave function. The hyperradial wave function $\psi_{\nu,\gamma}(x)$ is obtained as a solution of the hyperradial equation [39]

$$\left[\frac{d^2}{dx^2} + \frac{5}{x}\frac{d}{dx} - \frac{\gamma(\gamma+4)}{x^2}\right]\psi_{\nu,\gamma}(x) = -2\mu[E_{\nu,\gamma} - V(x)]\psi_{\nu,\gamma}(x).$$
(7)

We consider the Killingbeck potential given by [43]

$$V(x) = \alpha x^2 + \beta x - \frac{\tau}{x},\tag{8}$$

here, x is the hyperradius and α , β and τ are constant. In the present work, we solve the Schrödinger equation 7 using deep learning technique combined with a PSO algorithm. We use a trial wave function as supposed in our previous works [40, 41]. In our study we consider the baryon states with zero nodes $\nu = 0$, and neglect the spin dependent interactions.

3 Deep Learning Model for Energy Evaluation

Neural network is a computational model motivated by the biological nervous system and is made up of computing units, called neurons. The first step in our framework involves the generation of a synthetic dataset by solving the time-independent Schrödinger equation and using the shooting method. The Schrödinger equation for the system under study is expressed as

$$\left[\frac{d^2}{dx^2} + \frac{5}{x}\frac{d}{dx} - \frac{\gamma(\gamma+4)}{x^2} - 2max^2 - 2mbx + \frac{2mc}{x}\right]\psi_{\gamma}(x) = -2mE_{\gamma}\psi_{\gamma}(x).$$
(9)

The goal is to compute E_{γ} energy eigenvalues such that the boundary conditions for $\psi_{\gamma}(x)$ are satisfied. Here deep learning model is designed to predict energy eigenvalues based on the features in dataset.

3.1 Model Architecture

The deep neural network architecture consists of three layers: an input layer, hidden layers, and output layer. These layers are connected to the next layer, however there is no connection among the neurons in the same layer. They are ideal tools to solve differential equations. In this study, we consider a feedforward neural network with one input layer, two hidden layers and one output layer. The output layer consists of a single neuron to predict the E_{γ} energy eigenvalues.

3.2 Training Procedure

The neural network is trained using the mean squared error (MSE) loss function:

$$MSE = \frac{1}{n} \sum_{i=1}^{n} (Y_{\text{true}} - Y_{\text{pred}})^2,$$
(10)

where n, Y_{true} and Y_{pred} are the number of data points, actual value for *i*-th data point and predicted value for the *i*-th data point. We employ the adaptive moment estimation optimizer for adaptive learning rate adjustments during training. The model is trained over a large number of epochs to ensure convergence. The dataset is split into 80% training and 20% validation subsets. The performance of the model can be evaluated by using the root mean squared error (RMSE) and MSE on the validation set to the prediction accuracy assessment.

4 Particle Swarm Optimization (PSO)

For the further refines of energy evaluations, we employ particle swarm optimization. PSO is a population-based optimization algorithm inspired by the social behavior of birds, and is effective in finding global optima in multi-dimensional search spaces. A swarm of particles is initialized randomly, where each particle represents a solution (energy value). The position of each particle is updated iteratively, and the fitness of each particle is evaluated using the trained deep learning model. The fitness function for each particle is based on the difference between the predicted energy value and the target energy value. The goal is to minimize this difference. The global best solution found by PSO algorithm represents the optimal energy estimations.

5 Doubly heavy baryon mass spectra

By using the obtained E_{γ} energies, we are able to evaluate the mass spectra of the baryon systems. The quark masses used in the present work are taken from our previous work [40]. We consider $m_q=320$ MeV, $m_s=440$ MeV, $m_c=1600$ MeV and $m_b=4670$ MeV.

Our calculated results of the masses of the ground and orbitally excited baryon states including *P*-wave and *D*-wave are listed in Tables 1, 2, 3 and 4 and compared with the exist experimental data [42] and the ones obtained from different theoretical models; non-relativistic quark model (NRQM), relativistic quark model (RQM), heavy diquark effective theory (EFT), heavy quark spin symmetry (HQSS) and Regge phenomenology (RPH) [43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53]. In our previous studies we have also considered the heavy baryon states in a quark model and solved the Schrödinger equation by the numerical method [43, 44]. Our evaluations for the ground and excited states of single charm and bottom baryon masses are listed in Tables 1 and 2. Our predictions for the ground and orbitally excited states of doubly heavy baryons are summarized in Tables 3 and 4. Our evaluated masses for the single heavy baryons are very close to the experimental data. We could obtain the experimentally measured mass of the Ξ_{cc}^{++} doubly heavy baron [6] for the *qcc* baryon state in Table 3.

State	Baryon	Our results	$\operatorname{Exp}[42]$	NRQM[43]	NRQM[45]
S-wave	Σ_c	2.453	2.455	2.459	2.452
	Ξ_c	2.463	2.465	2.504	2.473
	Ω_c	2.694	2.695	2.566	2.695
<i>P</i> -wave	Σ_c	2.791	2.792		2.794
	Ξ_c	2.789	2.791		2.726
	Ω_c	3.063	3.065		2.996
D-wave	Σ_c	3.088			3.058
	Ξ_c	3.113			2.960
	Ω_c	3.325			3.230

Table 1: Masses of the ground and orbitally excited states of single charm baryons (in GeV).

Table 2: Masses of the ground and orbitally excited states of single bottom baryons (in GeV).

State	Baryon	Our result	$\operatorname{Exp}[42]$	NRQM[43]	NRQM[46]	RPH[47]
	Σ_b	5.808	5.810	5.808	5.816	5813
S-wave	Ξ_b	5.925	5.797	5.848	5.793	5.793
	Ω_b	6.043	6.045	5.903	6.048	6.048
	Σ_b	6.096	6.098		6.105	6.098
P-wave	Ξ_b	6.213	6.227		6.133	6.080
	Ω_b	6.337	6.339		6.321	6.325
	Σ_b	6.384			6.363	6.369
D-wave	Ξ_b	6.352			6.355	6.354
	Ω_b	6.631			6.554	6.590

6 Semileptonic $B \rightarrow B' l \bar{\nu}$ decay widhts and branching fractions

The semileptonic decays caused by the weak force in which a hadron decays weakly into another hadron and a lepton and a neutrino are produced. To consider the semileptonic decays one needs to determine the form factors of the transitions. There are different methods to simplify the transition form factors [54, 56, 57, 58]. In the present work we study $B(1/2^+) \rightarrow B'(1/2^+)$ semileptonic decays of doubly heavy baryons for $b \rightarrow c$ transitions. The methodology for the decay widths follows the same pattern used in our previous work [40].

The transition matrix element for the semileptonic decay is

$$T = \frac{G_F}{\sqrt{2}} V_{cb} J_\mu \mathcal{L}^\mu \tag{11}$$

where G_F and V_{bc} refers to the Fermi Coupling constant and CKM matrix element. J_{μ} is the weak hadronic current and \mathcal{L}^{μ} is the leptonic current

$$J_{\mu} = V_{\mu} - A_{\mu} = \bar{\psi}^{c} \gamma_{\mu} (I - \gamma_{5}) \psi^{b}, \qquad \mathcal{L}^{\mu} = \bar{l} \gamma^{\mu} (1 - \gamma_{5}) \nu_{l}$$
(12)

where $\psi^{b(c)}$ is the charm or bottom quark field. $V_{\mu} \equiv \bar{\psi}^c \gamma_{\mu} \psi^b$ and $A_{\mu} \equiv \bar{\psi}^c \gamma_{\mu} \gamma_5 \psi^b$ are the vector and axial-vector weak currents, respectively. The hadron matrix elements is given in terms of the six form

Baryon	Content	Our results	NRQM[44]	NRQM[48]	RQM[49]	NRQM[50]
Ξ_{cc}	qcc	3.620	3.679	3.613	3.676	3.685
Ξ_{bc}	qbc	7.050	7.003	6.928	7.020	
Ξ_{bb}	qbb	10.200	10.325	10.198	10.340	10.314
Ω_{cc}	scc	3.750	3.830	3.712	3.815	3.832
Ω_{bc}	sbc	6.900	7.149	7.013	7.147	
Ω_{bb}	sbb	10.400	10.466	10.269	10.456	10.447

Table 3: Masses of the ground states of double heavy baryons (in GeV).

Table 4: Masses of the orbitally excited states of doubly heavy baryons (in GeV).

State	Baryon	Our results	NRQM[51]	EFT[52]	NRQM[53]
	Ξ_{cc}	3.928	3.853	4.028	3.855
	Ξ_{bc}	7.266	7.140		
	Ξ_{bb}	10.631	10.502	10.386	10.417
P-wave	Ω_{cc}	4.106	3.964	4.086	4.002
	Ω_{bc}	7.4457	7.375		
	Ω_{bb}	10.808	10.634	10.607	10.560
	Ξ_{cc}	4.237	4.026	4.321	
	Ξ_{bc}	7.575	7.307		
	Ξ_{bb}	10.939	10.658	10.585	
D-wave	Ω_{cc}	4.414	4.133	4.263	
	Ω_{bc}	7.754	7.807		
	Ω_{bb}	11.1177	10.783	10.723	

factors as fallows

$$H_{\mu} = \langle B'(p',s') | V_{\mu} - A_{\mu} | B(p,s) \rangle = \langle B'(p',s') | \bar{\psi}^{c} \gamma_{\mu} (I - \gamma_{5}) \psi^{b} | B(p,s) \rangle$$

= $\bar{u}'(p',s') \{ \gamma_{\mu} (F_{1}(\omega) - \gamma_{5}G_{1}(\omega)) + v_{\mu} (F_{2}(\omega) - \gamma_{5}G_{2}(\omega)) + v'_{\mu} (F_{3}(\omega) - \gamma_{5}G_{3}(\omega)) \} u(p,s)$ (13)

where $|B(p,s)\rangle$ and $|B'(p',s')\rangle$ refers to the initial and final baryons. u(p,s) and u'(p',s') are Dirac spinors, normalized as $\bar{u}u = 1$, and $v_{\mu} = p_{\mu}/m_B(v'_{\mu} = p'_{\mu}/m'_B)$ is the four velocity of the corresponding baryon. The differential decay rates are given in terms of the transversely Γ_T and longitudinally Γ_L polarized W's,

$$\frac{d\Gamma}{d\omega} = \frac{d\Gamma_L}{d\omega} + \frac{d\Gamma_T}{d\omega}.$$
(14)

The differential decay rates Γ_T and Γ_L for the $B^{1/2} \to B'^{1/2}$ transition are obtained in terms of the form factors [28, 29]

$$\frac{d\Gamma_T}{d\omega} = \frac{G_F^2 |V_{cb}|^2 M_{B'}^3}{12\pi^3} \sqrt{\omega^2 - 1} q^2 \{ (\omega - 1) |F_1(\omega)|^2 + (\omega + 1) |G_1(\omega)|^2 \},\tag{15}$$

and

$$\frac{d\Gamma_L}{d\omega} = \frac{G_F^2 |V_{cb}|^2 M_{B'}^3}{24\pi^3} \sqrt{\omega^2 - 1} \{(\omega - 1)|\mathcal{F}^V(\omega)|^2 + (\omega + 1)|\mathcal{F}^A(\omega)|^2\},\tag{16}$$

where

$$\mathcal{F}^{V,A}(\omega) = \left[(m_B \pm m_{B'}) F_1^{V,A}(\omega) + (1 \pm \omega) \left(m_{B'} F_2^{V,A}(\omega) + m_B F_3^{V,A}(\omega) \right) \right],$$

$$F_j^V \equiv F_j(\omega), \quad F_j^A \equiv G_j(\omega), \quad j = 1, 2, 3$$
(17)

Here the lepton masses are neglected. $\omega = v.v'$ is the velocity transfer and $q^2 = (p-p')^2 = m_B^2 + m_{B'}^2 - 2m_B m_{B'}\omega$ is the momentum transfer squared between the initial and final baryon. The form factors are dependent on the ω and q^2 . In the considered transitions ω ranges from $\omega = 1$ corresponding to zero recoil point of the final baryon to the maximum of $\omega_{max} = (m_B^2 + m_{B'}^2)/(2m_B m_{B'})$. Integrating over the parameter ω , we can obtain the total decay width

$$\Gamma = \int_{1}^{\omega_{max}} d\omega \frac{d\Gamma}{d\omega} = \int_{1}^{\omega_{max}} d\omega \left(\frac{d\Gamma_L}{d\omega} + \frac{d\Gamma_T}{d\omega}\right).$$
(18)

The form factors can be parametrized by different methods. Some earlier studies have simplified the transition form factors for the semileptonic decays of the baryons [54, 56, 57, 58]. Ref. [54] has studied the form factors of the doubly heavy baryon transitions using a Lorentz covariant constituent quark model. In the heavy quark limit and near to zero recoil point, the weak form factors can be simplified and expressed by a single IW function $\eta(\omega)$ [54, 59, 60]

$$F_1(\omega) = G_1(\omega) = \eta(\omega),$$

$$F_2(\omega) = F_3(\omega) = G_2(\omega) = G_3(\omega) = 0.$$
(19)

The universal function $\eta(\omega)$ is defined in terms of the kinematical parameter ω [54]

$$\eta(\omega) = exp\left(-3(\omega-1)\frac{m_{cc}^2}{\Lambda_B^2}\right)$$
(20)

with $m_{cc} = 2m_c$ for the $bc \to cc$ weak transitions. The cut-off parameter Λ_B is a parameter depends on the size of a baryon and its value ranges from $2.5 \leq \Lambda_B \leq 3.5$ GeV [54] where, the smaller of Λ_B gives smaller decay widths. In the case of the $bb \to bc$ transitions, m_{cc} should be replaced by m_{bb} and therefore, the IW functions for $bb \to bc$ and $bc \to cc$ transitions depends on the heavy flavor factors explicitly with no dependence on the light quark sector. The Λ_B dependence of η functions for $\Omega_{bb} \to \Omega_{bc}$ and $\Xi_{bb} \to \Xi_{bc}$ transitions with $\Lambda_B = 2.5$ GeV is shown in Fig. 1.

The ω dependence of the semileptonic decay rates of $\Xi_{bb} \to \Xi_{bc} \ell \bar{\nu}_{\ell}$, and $\Xi_{bc} \to \Xi_{cc} \ell \bar{\nu}_{\ell}$ transitions is shown in Figs. 2 and 3 respectively.

Regarding the equation 19 one can get the new relations for the differential decay rates

$$\frac{d\Gamma_T}{d\omega} = \frac{G_F^2 |V_{cb}|^2 M_{B'}^3}{6\pi^3} q^2 \omega \sqrt{\omega^2 - 1} \eta^2(\omega), \tag{21}$$

Table 5: Semileptonic decay widths of doubly heavy baryons in units of 10^{-14} GeV and their variations.

Decay	Our results	NRQM[40]	RCQM [54]	NRQM $[55]$	HQSS [26]	$LFQM^1$ [30]
$\Xi_{bc} \to \Xi_{cc} \ell \bar{\nu}_{\ell}$	$4.47^{+1.0}_{-1.20}$	4.39 ± 0.83	4.01 ± 1.21	5.07	$2.57^{+0.26}_{-0.03}$	4.50
$\Xi_{bb} \to \Xi_{bc} \ell \bar{\nu}_{\ell}$	$1.66^{+0.73}_{-0.64}$	1.75 ± 0.73	1.33 ± 0.61	1.63	$1.92^{+0.25}_{-0.05}$	3.30
$\Omega_{bc} \to \Omega_{cc} \ell \bar{\nu}_{\ell}$	$3.55^{+0.67}_{-0.84}$	4.70 ± 0.83	4.12 ± 1.10	5.39	$2.59^{+0.20}$	3.94
$\Omega_{bb} \to \Omega_{bc} \ell \bar{\nu}_{\ell}$	$2.01^{+0.99}_{-0.80}$	1.87 ± 0.76	1.92 ± 1.15	2.48	$2.14_{-0.02}^{+0.20}$	3.69



Figure 1: Behavior of η function versus Λ_B for $\Omega_{bb} \to \Omega_{bc} \ell \bar{\nu}_{\ell}$ and $\Xi_{bb} \to \Xi_{bc} \ell \bar{\nu}_{\ell}$ ($\ell = e \text{ or } \mu$) transitions ($\omega = \omega_{\max}$).



Figure 2: $\frac{d\Gamma_T}{d\omega}$, $\frac{d\Gamma_L}{d\omega}$ and $\frac{d\Gamma}{d\omega}$ semileptonic decay widths in units of 10^{-14} GeV for $\Xi_{bc} \to \Xi_{cc} \ell \bar{\nu}_{\ell}$ transition $(\ell = e \text{ or } \mu)$. The results for Ω_{bc} decay (not shown) are very similar.



Figure 3: $\frac{d\Gamma_L}{d\omega}$, $\frac{d\Gamma_L}{d\omega}$ and $\frac{d\Gamma}{d\omega}$ semileptonic decay widths in units of 10^{-14} GeV for $\Xi_{bb} \to \Xi_{bc} \ell \bar{\nu}_{\ell}$ transition ($\ell = e \text{ or } \mu$). The results for Ω_{bb} decay (not shown) are very similar.

The branching ratios of semileptonic decays of doubly heavy baryons can be obtained as following

$$\mathcal{B} = \Gamma \times \tau, \tag{22}$$

where τ is the lifetime of the initial baryon. We take the values of $\tau_{\Xi_{bb}} = 370 \times 10^{-15} s$, $\tau_{\Xi_{bc}} = 244 \times 10^{-15} s$ [61], $\tau_{\Omega_{bc}} = 220 \times 10^{-15} s$, and $\tau_{\Omega_{bb}} = 800 \times 10^{-15} s$ [62, 63] to evaluate the branching fractions.

Table 6: The calculated branching fractions of semileptonic decays of doubly heavy baryons.

Process	Our results	NRQM $[40]$	NRQM $[55]$	LFQM [30]
$\Xi_{bc} \to \Xi_{cc} \ell \bar{\nu}_{\ell}$	1.65×10^{-2}	$1.63 imes 10^{-2}$	1.11×10^{-2}	$1.67 imes10^{-2}$
$\Xi_{bb} \to \Xi_{bc} \ell \bar{\nu}_{\ell}$	0.61×10^{-2}	$0.98 imes 10^{-2}$	0.28×10^{-2}	1.86×10^{-2}
$\Omega_{bc} \to \Omega_{cc} \ell \bar{\nu}_{\ell}$	1.18×10^{-2}	$1.57 imes 10^{-2}$	$1.10 imes 10^{-2}$	$1.32 imes 10^{-2}$
$\Omega_{bb} \to \Omega_{bc} \ell \bar{\nu}_{\ell}$	2.45×10^{-2}	2.27×10^{-2}		4.49×10^{-2}

The calculated semileptonic decay widths of doubly heavy baryons and their variations are listed in Table 5. We take $\Lambda_B = 3$. The uncertainties in our predicted decay widths are due to the cut-off parameter Λ_B , which can take a value in the range $2.5 \leq \Lambda_B \leq 3.5$ GeV. We compare our results with those predicted by Refs. [26, 30, 40, 54, 55]. There is no experimental data for semileptonic decay widths of doubly heavy baryons. Our predicted decay widths are close to the ones obtained by Ref. [54]. For the $\Xi_{bc} \to \Xi_{cc}$, $\Xi_{bb} \to \Xi_{bc}$ and $\Omega_{bb} \to \Omega_{bc}$ transitions, our results are also in good agreement with those of our previous work [40]. The evaluated branching fractions of semileptonic decays are summarized in Table 6 and compared with those of other works [30, 40, 55]. We hope our results are helpful to extract the value of the CKM matrix element V_{cb} from near future experiments.

7 Summary

In the present work, we propose our phenomenological approach for estimation energy eigenvalues of three-body baryon systems by integrating the deep learning with particle swarm optimization methods.

By integrating deep learning with PSO, we have developed a powerful methodology for accurate energy prediction in three-body quantum systems. The deep learning model, trained on a large dataset generated using the shooting method, provides an initial estimate of the energy values while, PSO refines these predictions and leads to highly accuracy in calculations. This hybrid approach demonstrates the potential of combining machine learning and optimization techniques in solving challenging math problems in computational physics. By evaluation of baryon energies, we are able to predict the mass spectrum of the single and doubly heavy baryons which are not yet determined by experiment. The validity of our scheme for heavy baryon systems will be determined by consistency with the exist experimental data. Our evaluated masses are in a good agreement with the experiments and also with the predictions of other theoretical works. In our study the overall MSE, training MSE and validation MSE are 6.443×10^{-5} , 6.690×10^{-5} and 6.196×10^{-5} respectively. We also study the semileptonic decays of doubly heavy Ξ and Ω baryons. By introducing a simple form for the universal IW function, we investigate the semileptonic decay rates and branching fractions of the $bb \rightarrow bc$ and $bc \rightarrow cc$ transitions close to zero recoil limit. We present a comparison between our results and other available theoretical calculations, and find that the results are acceptable. Our predictions will be able to guide the future searches for the undiscovered single and doubly heavy baryons at LHCb, ATLAS, and CMS.

References

- [1] X. Z. Weng, X. L. Chen, and W. Z. Deng, Phys. Rev. D 97, 054008 (2018).
- [2] Z. Shah and A. K. Rai, Eur. Phys. J. C 77, 129 (2017).
- [3] H. Garcilazo, A. Valcarce, and J. Vijande, Phys. Rev. D 94, 074003 (2016).
- [4] M. Mattson et al. (SELEX Collaboration), Phys. Rev. Lett. 89, 112001 (2002).
- [5] R. Aaij et al. (LHCb Collaboration), Phys. Rev. Lett. **119**, 112001 (2017).
- [6] R. Aaij et al. (LHCb Collaboration), Phys. Rev. Lett. 121, 162002 (2018).
- [7] R. Aaij et al. (LHCb Collaboration), JHEP 11, 095 (2020).
- [8] R. Aaij et al. (LHCb Collaboration), Chin. Phys. C 45, 093002 (2021).
- [9] R. Aaij et al. (LHCb Collaboration), Chin. Phys. C 47, 093001 (2023).
- [10] A. Ali, A. Y. Parkhomenko, Q. Qin and W. Wang, Phys. Lett. B 782, 412 (2018).
- [11] A. Ali, Q. Qin and W. Wang, Phys. Lett. B 785, 605 (2018).
- [12] S. Navas et al. (Particle Data Group), Phys. Rev. D 110, 030001 (2024).
- [13] R. Aaij et al. [LHCb Collaboration], Phys. Rev. Lett. **118**, 182001 (2017).
- [14] J. Yelton et al. [Belle Collaboration], Phys. Rev. D 97, 051102 (2018).
- [15] B. Aubert et al. [BaBar Collaboration], Phys. Rev. D 78, 112003 (2008).
- [16] E. Solovieva et al. [Belle Collaboration], Phys. Lett. B 672, Phys. Lett B 672, 1-5 (2009).
- [17] M. Ablikim et al. (BESIII Collaboration), Phys. Lett. B 767, 42 (2017).
- [18] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **121**, 251801 (2018).
- [19] Y. B. Li et al. (Belle Collaboration), Phys. Rev. Lett. 127, 121803 (2021).
- [20] Y. B. Li et al. (Belle Collaboration), Phys. Rev. D 105, L091101 (2022).

- [21] R. Ammar et al. (CLEO Collaboration), Phys. Rev. Lett. 89, 171803 (2002).
- [22] Zhen-Yu Li, Guo-Liang Yu, Zhi-Gang Wang, Jian-Zhong Gu, and Hong-Tao Shen, Mod. Phys. Lett. A 38, 2350051 (2023).
- [23] Y. J. Shi, W. Wang and Z. X. Zhao, Eur. Phys. J. C 80, 398 (2020).
- [24] T. Gutsche, A. Mikhail, M.A. Ivanov, J. G. Körner, V. E. Lyubovitskij and Z. Tyulemissov, Phys. Rev. D 100, 114037 (2019).
- [25] Q.X. Yu, X.H. Guo, Nucl. Phys. B **947**, 114727 (2019).
- [26] E. Hernandez, J. Nieves and J. M. Verde-Velasco, Phys. Lett. B 663, 234 (2008).
- [27] V. E. Lyubovitskij, A. Faessler, T. Gutsche, M. A. Ivanov, and J. G. Körner, Prog. Part. Nucl. Phys. 50, 329 (2003).
- [28] C. Albertus, E. Hernandez and J. Nieves, Phys. Rev. D 71, 014012 (2005).
- [29] C. Albertus, E. Hernandez, J. Nieves and J. M. Verde-Velasco, Eur. Phys. J. A 32, 183 (2007), [Erratum]: Eur. Phys. J. A 36, 119 (2008).
- [30] W. Wang, F. S. Yu, Z.X. Zhao, Eur. Phys. J. C 77, 781 (2017).
- [31] Q. Qin, Y. J. Shi, W. Wang, Y. Guo-He, F. S. Yu and R. Zhu, Phys. Rev. D 103, 3 (2022).
- [32] H. Mutuk, Eur. Phys. J. A 56, 146 (2020).
- [33] Yarin Gal, Vishnu Jejjala, Damian Kaloni Mayorga Penac, Challenger, Mishraa, Int. J. Mod. Phys. A 37, 2250031 (2022).
- [34] M. Malekhosseini, S. Rostami, A. R. Olamaei, R. Ostovar and K. Azizi, Phys. Rev. D 110, 054011 (2024).
- [35] H. Mutuk, Chin. Phys. C 43, 093103 (2019).
- [36] N. Yadav, A. Yadav, M. Kumar, An Introduction to Neural Network Methods for Differential Equations, Springer in Applied Sciences and Technology, (2015).
- [37] D. R. Parisi, M. C. Mariani, M. A. Laborde, Chem. Eng. Process. 42, 715 (2003).
- [38] E. Santopinto, F. Iachello and M. M. Gainnini, Eur. Phys. J. A 1, 307 (1998);
- [39] M. M. Giannini, E. Santopinto and A. Vassallo, Prog. Part. Nucl. Phys. 50, 263 (2003); Eur Phys. J. A 12, 447 (2001).
- [40] Z. Ghalenovi, C. P. Shen and M. Moazzen Sorkhi, Phys. Lett. B 834, 137405 (2022).
- [41] Z. Ghalenovi and M. Moazzen Sorkhi, Eur. Phys. J. Plus **133**, 301 (2018).
- [42] S. Navas et al. (Particle Data Group), Phys. Rev. D 110, 030001 (2024).
- [43] Z. Ghalenovi, A. A. Rajabi, S. X. Qin and D. H. Rischke, Mod. Phys. Lett. A 29, 1450106 (2014).
- [44] Z. Ghalenovi and M. M. Sorkhi, Chin. Phys. C 47, 3 (2023).
- [45] Z. Shah, K. Thakkar, A. Kumar Rai and P. C. Vinodkumar, Eur. Phys. J. A 52, 313 (2016).
- [46] K. Thakkar, Z. Shah, A. Kumar Rai, P.C. Vinodkumar, Nucl. Phys. A 965, 57 (2017).
- [47] Ke-Wei Wei, B. Chen, N. Liu, Q-Qian Wang, Xin-Heng Guo, Phys. Rev. D 95, 116005 (2017).

- [48] C. Albertus, E. Hernández, and J. Nieves, Phys. Lett. B 683, 21 (2010).
- [49] W. Roberts and M. Pervin, Int. J. Mod. Phys. A 23, 2817 (2008).
- [50] T. Yoshida, E. Hiyama, A. Hosaka, M. Oka, and K. Sadato, Phys. Rev. D 92, 114029 (2015).
- [51] Z. Shah, K. Thakkar and A. K. Rai, Eur. Phys. J. C 76, 530 (2016).
- [52] J. Soto and J. T. Castellà, Phys. Rev. D 104, 074027 (2021).
- [53] E. Ortiz-Pacheco and R. Bijker, Phys. Rev. D 108, 054014 (2023).
- [54] A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Körner, and V. E. Lyubovitskij, Phys. Rev. D 80, 034025 (2009).
- [55] S. Rahmani, H. Hassanabadi, and H. Sobhani, Eur. Phys. J. C 80, 312 (2020).
- [56] H. Georgi and M. B. Wise, Phys. Lett. B 243, 279 (1990).
- [57] C. D. Carone, Phys. lett. B 253, 408 (1991).
- [58] J. M. Flynn and J. Neives, Phys. Rev. D 76, 017502 (2007) [Erratum: Phys. Rev. D 77, 099901 (2008).
- [59] N. Isgur and M. B. Wise, Nucl. Phys. B 348, 276 (1991).
- [60] D. Ebert, R. N. Faustov and V. O. Galkin, Phys. Rev. D 73, 094002 (2006).
- [61] M. Karliner and J. L. Rosner, Phys. Rev. D 90, 094007 (2014).
- [62] V. V. Kiselev and A. K. Likhoded, Phys. Usp. 172, 497 (2002).
- [63] V. V. Kiselev and A. K. Likhoded, Phys. Usp. 45, 455 (2002).