

# IMPROVEMENTS OF CERTAIN RESULTS OF THE CLASS $\mathcal{S}$ OF UNIVALENT FUNCTIONS

MILUTIN OBRADOVIĆ AND NIKOLA TUNESKI

*This paper is dedicated to the memory of our esteemed colleague, Derek K. Thomas, whose untimely passing occurred during the course of this work. We honor his legacy and the lasting impact of his achievements in the field of univalent functions.*

ABSTRACT. For  $f \in \mathcal{S}$ , the class univalent functions in the unit disk  $\mathbb{D}$  and given by  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  for  $z \in \mathbb{D}$ , we improve previous bounds for the second and third Hankel determinants in case when either  $a_2 = 0$ , or  $a_3 = 0$ . We also improve an upper bound for the coefficient difference  $|a_4| - |a_3|$  when  $f \in \mathcal{S}$ .

## 1. INTRODUCTION AND PRELIMINARIES

As usual, let  $\mathcal{A}$  be the class of functions  $f$  which are analytic in the open unit disc  $\mathbb{D} = \{z : |z| < 1\}$  of the form

$$(1) \quad f(z) = z + a_2 z^2 + a_3 z^3 + \dots,$$

and let  $\mathcal{S}$  be the subclass of  $\mathcal{A}$  consisting of functions that are univalent in  $\mathbb{D}$ .

In recent years a great deal of attention has been given to finding upper bounds for the modulus of the second and third Hankel determinants  $H_2(2)$  and  $H_3(1)$ , defined as follows whose elements are the coefficients of  $f \in \mathcal{S}$  (see e.g. [10]).

For  $f \in \mathcal{S}$

$$(2) \quad H_2(2) = a_2 a_4 - a_3^2$$

and

$$(3) \quad H_3(1) = a_3(a_2 a_4 - a_3^2) - a_4(a_4 - a_2 a_3) + a_5(a_3 - a_2^2).$$

Almost all results have concentrated on finding bounds for  $|H_2(2)|$  and  $|H_3(1)|$  for subclasses of  $\mathcal{S}$ , and only recently has a significant bound been found for the whole class  $\mathcal{S}$  and for both,  $|H_2(2)|$  and  $|H_3(1)|$  (see [7] and [8]). However finding exact sharp bounds remains an open problem.

In their paper [9] the authors gave the next results concerning coefficients bound and Hankel determinants  $|H_2(2)|$  and  $|H_3(1)|$ .

**Theorem 1.** *Let  $f \in \mathcal{S}$  and be given by (1) with  $a_2 = 0$ . Then*

---

2020 *Mathematics Subject Classification.* 30C45, 30C50, 30C55.

*Key words and phrases.* univalent functions, Grunsky coefficients, coefficient differences, Hankel determinant.

- (i)  $|a_3| \leq 1$ ,
- (ii)  $|a_4| \leq \frac{2}{3} = 0.666\dots$ ,
- (iii)  $|a_5| \leq \frac{5}{4} + \frac{1}{\sqrt{15}} = 1.508\dots$ ,
- (iv)  $|H_2(2)| \leq 1$ ,
- (v)  $|H_3(1)| \leq \frac{41}{20} = 2.05$ .

We would like to point out that there is a mistake in the estimate of  $|a_5|$  given in Theorem 2.1(iii) from [9]. The estimate given in the case (iii) in the theorem above is the correct one.

Similar results are given in [9] for the case when  $a_3 = 0$ .

**Theorem 2.** *Let  $f \in \mathcal{S}$  and be given by (1), with  $a_3 = 0$ . Then*

- (i)  $|a_2| \leq 1$ ,
- (ii)  $|a_4| \leq \frac{\sqrt{37} + 13}{12} = 1.59023\dots$ ,
- (iii)  $|a_5| \leq \frac{1}{4}\sqrt{\frac{757}{15}} + 1 = 2.77599\dots$ ,
- (iv)  $|H_2(2)| \leq 1.75088\dots$ ,
- (v)  $|H_3(1)| \leq 1.114596\dots$

A long standing problem in the theory of univalent functions is to find sharp upper and lower bounds for  $|a_{n+1}| - |a_n|$ , when  $f \in \mathcal{S}$ . Since the Keobe function has coefficients  $a_n = n$ , it is natural to conjecture that  $||a_{n+1}| - |a_n|| \leq 1$ . As early as 1933, this was shown to be false even when  $n = 2$ , when Fekete and Szegő [2] obtained the sharp bounds

$$-1 \leq |a_3| - |a_2| \leq \frac{3}{4} + e^{-\lambda_0}(2e^{-\lambda_0} - 1) = 1.029\dots,$$

where  $\lambda_0$  is the unique value of  $\lambda$  in  $0 < \lambda < 1$ , satisfying the equation  $4\lambda = e^\lambda$ . Hayman [4] showed that if  $f \in \mathcal{S}$ , then  $||a_{n+1}| - |a_n|| \leq C$ , where  $C$  is an absolute constant. The exact value of  $C$  is unknown, the best estimate to date being  $C = 3.61\dots$  (see Grinspan [3]), which because of the sharp estimate above when  $n = 2$ , cannot be reduced to 1.

In [9] the authors treated the difference coefficients  $|a_4| - |a_3|$  for  $f \in \mathcal{S}$ , which improves the previous cited result of Grinspan when  $n = 3$  as follows

**Theorem 3.** *Let  $f \in \mathcal{S}$  and be given by (1). Then*

$$|a_4| - |a_3| \leq 2.1033299\dots$$

For the proofs of the previously cited results, the authors mainly used the property of Grunsky coefficients given in the book of N. A. Lebedev ([5]). In the proofs of the results in this paper we also use the same tools but with different approach that brings improved estimates.

We proceed with the notations and results to be used.

Let  $f \in \mathcal{S}$  and let

$$\log \frac{f(t) - f(z)}{t - z} = \sum_{p,q=0}^{\infty} \omega_{p,q} t^p z^q,$$

where  $\omega_{p,q}$  are called Grunsky's coefficients with property  $\omega_{p,q} = \omega_{q,p}$ . For those coefficients in [1, 5] we can find the next Grunsky's inequality:

$$(4) \quad \sum_{q=1}^{\infty} q \left| \sum_{p=1}^{\infty} \omega_{p,q} x_p \right|^2 \leq \sum_{p=1}^{\infty} \frac{|x_p|^2}{p},$$

where  $x_p$  are arbitrary complex numbers such that last series converges.

Further, it is well-known that if  $f \in \mathcal{S}$  and has the form (1), then also

$$f_2(z) = \sqrt{f(z^2)} = z + c_3 + c_5 z^5 + \dots$$

belongs to the class  $\mathcal{S}$ . Then for the function  $f_2$  we have the appropriate Grunsky's coefficients of the form  $\omega_{2p-1,2q-1}^{(2)}$  and the inequality (4) gets the form

$$(5) \quad \sum_{q=1}^{\infty} (2q-1) \left| \sum_{p=1}^{\infty} \omega_{2p-1,2q-1}^{(2)} x_{2p-1} \right|^2 \leq \sum_{p=1}^{\infty} \frac{|x_{2p-1}|^2}{2p-1}.$$

As it has been shown in [5, p.57], if  $f$  is given by (1) then the coefficients  $a_2, a_3, a_4, a_5$  are expressed by Grunsky's coefficients  $\omega_{2p-1,2q-1}^{(2)}$  of the function  $f_2$  given by (3) in the following way (in the next text we omit upper index "(2)" in  $\omega_{2p-1,2q-1}^{(2)}$ ):

$$(6) \quad \begin{aligned} a_2 &= 2\omega_{11}, \\ a_3 &= 2\omega_{13} + 3\omega_{11}^2, \\ a_4 &= 2\omega_{33} + 8\omega_{11}\omega_{13} + \frac{10}{3}\omega_{11}^3, \\ a_5 &= 2\omega_{35} + 8\omega_{11}\omega_{33} + 5\omega_{13}^2 + 18\omega_{11}^2\omega_{13} + \frac{7}{3}\omega_{11}^4 \\ 0 &= 3\omega_{15} - 3\omega_{11}\omega_{13} + \omega_{11}^3 - 3\omega_{33} \\ 0 &= \omega_{17} - \omega_{35} - \omega_{11}\omega_{33} - \omega_{13}^2 + \frac{1}{3}\omega_{11}^4. \end{aligned}$$

We note that in [5] there exists a typing mistake for the coefficient  $a_5$ . Namely, instead of the term  $5\omega_{13}^2$ , there is  $5\omega_{15}^2$ .

Also, from (5) for  $x_{2p-1} = 0, p = 3, 4, \dots$ , we have

$$(7) \quad \begin{aligned} &|\omega_{11}x_1 + \omega_{31}x_3|^2 + 3|\omega_{13}x_1 + \omega_{33}x_3|^2 + |\omega_{15}x_1 + \omega_{35}x_3|^2 \\ &+ |\omega_{17}x_1 + \omega_{37}x_3|^2 \leq |x_1|^2 + \frac{|x_3|^2}{3}. \end{aligned}$$

From (7), having in mind that  $\omega_{31} = \omega_{13}$ , for  $x_1 = 1$  and  $x_3 = 0$  we have the next inequalities

$$\begin{aligned} |\omega_{11}|^2 &\leq 1, \\ |\omega_{11}|^2 + 3|\omega_{13}|^2 &\leq 1, \\ |\omega_{11}|^2 + 3|\omega_{13}|^2 + 5|\omega_{15}|^2 &\leq 1, \\ |\omega_{11}|^2 + 3|\omega_{13}|^2 + 5|\omega_{15}|^2 + 7|\omega_{17}|^2 &\leq 1. \end{aligned}$$

From the last inequalities we easily obtain

$$(8) \quad \begin{aligned} |\omega_{11}| &\leq 1, \\ |\omega_{13}| &\leq \frac{1}{\sqrt{3}}\sqrt{1-|\omega_{11}|^2}, \\ |\omega_{15}| &\leq \frac{1}{\sqrt{5}}\sqrt{1-|\omega_{11}|^2-3|\omega_{13}|^2}, \\ |\omega_{17}| &\leq \frac{1}{\sqrt{7}}\sqrt{1-|\omega_{11}|^2-3|\omega_{13}|^2-5|\omega_{15}|^2}. \end{aligned}$$

We note that we get the first inequality from (8) also using the fact  $|a_2| = |2\omega_{11}| \leq 2$  (see (6)).

## 2. MAIN RESULTS

We start with improvement of some results given in Theorem 1.

**Theorem 4.** *Let  $f \in \mathcal{S}$  and be given by (1) with  $a_2 = 0$ . Then*

- (i)  $|a_5| \leq \frac{3}{4} + \frac{1}{\sqrt{7}} = 1.12796\dots$ ,
- (ii)  $|H_3(1)| \leq 1.026\dots$

*Proof.*

- (i) The classical inequality  $|a_3 - a_2^2| \leq 1$  for  $f$  in  $\mathcal{S}$  when  $a_2 = 0$ , gives  $|a_3| \leq 1$ , which from (6) gives

$$(9) \quad |\omega_{13}| \leq \frac{1}{2}.$$

Since  $\omega_{11} = 0$  ( $\Leftrightarrow a_2 = 0$ ), from the relation for  $a_5$  in (6) and last relation in it, we obtain

$$(10) \quad |a_5| = |2\omega_{35} + 5\omega_{13}^2|$$

and

$$(11) \quad \omega_{35} = \omega_{17} - \omega_{13}^2.$$

Further, using the relations (10) and (11) we have

$$(12) \quad \begin{aligned} |a_5| &= |2\omega_{17} + 3\omega_{13}^2| \\ &\leq 2|\omega_{17}| + 3|\omega_{13}|^2 \\ &\leq \frac{2}{\sqrt{7}}\sqrt{1-3|\omega_{13}|^2} + 3|\omega_{13}|^2 \\ &\leq \frac{3}{4} + \frac{1}{\sqrt{7}} = 1.12796\dots, \end{aligned}$$

since by (6) ( $\omega_{11} = 0$ ),

$$(13) \quad |\omega_{17}| \leq \frac{1}{\sqrt{7}}\sqrt{1-3|\omega_{13}|^2-5|\omega_{15}|^2} \leq \frac{1}{\sqrt{7}}\sqrt{1-3|\omega_{13}|^2}$$

and  $|\omega_{13}| \leq \frac{1}{2}$  by (9).

(ii) When  $\omega_{11} = 0$ , the fifth relation in (6) gives  $\omega_{33} = \omega_{15}$ , and using (11), from (3) we have

$$\begin{aligned} |H_3(1)| &= |-8\omega_{13}^3 - 4\omega_{33}^2 + 2(2\omega_{35} + 5\omega_{13}^2)\omega_{13}| \\ &= |-8\omega_{13}^3 - 4\omega_{15}^2 + 4\omega_{13}(\omega_{17} - \omega_{13}^2) + 10\omega_{13}^3| \\ &= |-2\omega_{13}^3 - 4\omega_{15}^2 + 4\omega_{13}\omega_{17}| \\ &\leq 2|\omega_{13}|^3 + 4|\omega_{15}|^2 + 4|\omega_{13}||\omega_{17}| \\ &\leq 2|\omega_{13}|^3 + \frac{4}{5}(1 - 3|\omega_{13}|^2) + \frac{4}{\sqrt{7}}|\omega_{13}|\sqrt{1 - 3|\omega_{13}|^2} \\ &=: F_1(|\omega_{13}|), \end{aligned}$$

where

$$F_1(y) = 2y^3 + \frac{4}{5}(1 - 3y^2) + \frac{4}{\sqrt{7}}y\sqrt{1 - 3y^2}, \quad 0 \leq y \leq \frac{1}{2}.$$

Here we used the relations (10) and (13). Now, using the first derivative test we conclude that the function  $F_1$  attains its maximum for  $y_0 = 0.286667\dots$  with  $F_1(y_0) = 1.026\dots$

Finally, we note that the results (i) and (iv) in Theorem 1 are the best possible as the function  $f(z) = \frac{z}{1-z^2}$  shows.  $\square$

We next prove a similar result, this time assuming that  $a_3 = 0$ .

**Theorem 5.** *Let  $f \in \mathcal{S}$  and be given by (1), with  $a_3 = 0$ . Then*

- (i)  $|a_2| \leq 1$ ,
- (ii)  $|a_4| \leq \frac{1}{4}\sqrt{\frac{21}{5}} + \frac{5}{8} = 1.1373\dots$ ,
- (iii)  $|a_5| \leq 1.674896577\dots$ ,
- (iv)  $|H_2(2)| \leq 1.1373\dots$ ,
- (v)  $|H_3(1)| \leq 0.6647958756\dots$

*Proof.*

(i) We imitate the proof of Theorem 4(i) and from  $|a_3 - a_2^2| \leq 1$ ,  $a_3 = 0$ , we receive  $|a_2^2| \leq 1$ , i.e.,  $|a_2| \leq 1$ .

Further we will use that from (6) follows  $a_3 = 2\omega_{13} + 3\omega_{11}^2 = 0$ , and then

$$(14) \quad \omega_{13} = -\frac{3}{2}\omega_{11}^2 \quad \left( \Leftrightarrow \omega_{11}^2 = -\frac{2}{3}\omega_{13} \right),$$

and also, from  $|a_2| = |2\omega_{11}| \leq 1$ ,

$$|\omega_{11}| \leq \frac{1}{2} \quad \text{and} \quad |\omega_{13}| \leq \frac{3}{8} \quad (\text{by (14)}).$$

(ii) By using (6) and (14), we obtain

$$(15) \quad |a_4| = \left| 2\omega_{33} + 8\omega_{11} \left( -\frac{3}{2}\omega_{11}^2 \right) + \frac{10}{3}\omega_{11}^3 \right| = \left| 2\omega_{33} - \frac{26}{3}\omega_{11}^3 \right|.$$

On the other hand, using the fifth relation in (6), and (14), we have

$$(16) \quad \begin{aligned} \omega_{33} &= \omega_{15} - \omega_{11}\omega_{13} + \frac{1}{3}\omega_{11}^3 = \omega_{15} - \omega_{11} \left( -\frac{3}{2}\omega_{11}^2 \right) + \frac{1}{3}\omega_{11}^3 \\ &= \omega_{15} + \frac{11}{6}\omega_{11}^3. \end{aligned}$$

Combining (15) and (16) we obtain

$$(17) \quad \begin{aligned} |a_4| &= |2\omega_{15} - 5\omega_{11}^3| \leq 2|\omega_{15}| + 5|\omega_{11}|^3 \\ &\leq \frac{2}{\sqrt{5}} \sqrt{1 - |\omega_{11}|^2 - \frac{27}{4}|\omega_{11}|^4 + 5|\omega_{11}|^3} \\ &=: F_2(|\omega_{11}|), \end{aligned}$$

where

$$F_2(x) = \frac{2}{\sqrt{5}} \sqrt{1 - x^2 - \frac{27}{4}x^4 + 5x^3}, \quad 0 \leq x \leq \frac{1}{2},$$

and where we used (8) and (14) to obtain

$$(18) \quad |\omega_{15}| \leq \frac{1}{\sqrt{5}} \sqrt{1 - |\omega_{11}|^2 - \frac{27}{4}|\omega_{11}|^4}.$$

Finally,  $F_2(x)$  is strictly increasing function on the interval  $[0, 1/2]$ , attaining its maximal value  $\frac{1}{4}\sqrt{\frac{21}{5}} + \frac{5}{8} = 1.1373\dots$  for  $x = 1/2$ .

(iii) From the last relation in (6), using (14) and (16), after simple calculation we receive

$$(19) \quad \omega_{35} = \omega_{17} - \omega_{11}\omega_{15} - \frac{15}{4}\omega_{11}^4.$$

Using the relations (6) and (19), we get

$$(20) \quad \begin{aligned} |a_5| &= \left| 2\omega_{17} + 6\omega_{11}\omega_{15} - \frac{25}{4}\omega_{11}^4 \right| \\ &\leq 2|\omega_{17}| + 6|\omega_{11}||\omega_{15}| + \frac{25}{4}|\omega_{11}|^4 \\ &\leq \left( \frac{2}{\sqrt{7}} + \frac{6}{\sqrt{5}}|\omega_{11}| \right) \sqrt{1 - |\omega_{11}|^2 - \frac{27}{4}|\omega_{11}|^4 + \frac{25}{4}|\omega_{11}|^4} \\ &=: F_3(x), \end{aligned}$$

where

$$F_3(x) = \left( \frac{2}{\sqrt{7}} + \frac{6}{\sqrt{5}}x \right) \sqrt{1 - x^2 - \frac{27}{4}x^4 + \frac{25}{4}x^4}, \quad 0 \leq x \leq \frac{1}{2},$$

and where we used the estimate given in (18), the estimate from (8), and

$$(21) \quad \begin{aligned} |\omega_{17}| &\leq \frac{1}{\sqrt{7}} \sqrt{1 - |\omega_{11}|^2 - 3|\omega_{13}|^2 - 5|\omega_{15}|^2} \\ &\leq \frac{1}{\sqrt{7}} \sqrt{1 - |\omega_{11}|^2 - 3|\omega_{13}|^2}. \end{aligned}$$

Now, the first derivative test shows that on the interval  $[0, 1/2]$ , the function  $F_3$  has maximal value 1.674896577... attained for  $x = 0.43957885 \dots$

- (iv) Since  $a_3 = 0$  using (2) we have  $H_2(2) = a_2a_4$  and then from the relation (15) and estimation (18), we obtain

$$\begin{aligned} |H_2(2)| &= |a_2a_4| = |2\omega_{11}(2\omega_{15} - 5\omega_{11}^3)| \\ &\leq 4|\omega_{11}||\omega_{15}| + 10|\omega_{11}|^4 \\ &\leq \frac{4}{\sqrt{5}}|\omega_{11}|\sqrt{1 - |\omega_{11}|^2 - \frac{27}{4}|\omega_{11}|^4} + 10|\omega_{11}|^4 \\ &=: F_4(|\omega_{11}|), \end{aligned}$$

where

$$F_4(x) = \frac{4}{\sqrt{5}}x\sqrt{1 - x^2 - \frac{27}{4}x^4} + 10x^4, \quad 0 \leq x \leq \frac{1}{2}.$$

Again, the first derivative test shows that  $F_4$  is strictly increasing on the interval  $[0, 1/2]$  with maximal value 1.1373... attained for  $x = 1/2$ .

- (v) After applying  $a_3 = 0$  in the definition (3) we receive

$$(22) \quad H_3(1) = -a_4^2 - a_5a_2^2.$$

Recall that in (17) and (20) we found  $a_4 = 2\omega_{15} - 5\omega_{11}^3$  and  $a_5 = 2\omega_{17} + 6\omega_{11}\omega_{15} - \frac{25}{4}\omega_{11}^4$ , respectively. From these facts and (22), after some calculations we obtain

$$(23) \quad H_3(1) = -\omega_{15}^2 - 4\omega_{11}^3\omega_{15} - 8\omega_{11}^2\omega_{17}.$$

So, using (23) and estimates given in (18) and (21) we get

$$\begin{aligned} |H_3(1)| &= |\omega_{15}|^2 + 4|\omega_{11}|^3|\omega_{15}| + 8|\omega_{11}|^2|\omega_{17}| \\ &\leq \frac{1}{5} \left( 1 - |\omega_{11}|^2 - \frac{27}{4}|\omega_{11}|^4 \right) \\ &\quad + \left( \frac{4}{\sqrt{5}}|\omega_{11}|^3 + \frac{8}{\sqrt{7}}|\omega_{11}|^2 \right) \sqrt{1 - |\omega_{11}|^2 - \frac{27}{4}|\omega_{11}|^4} \\ &=: F_5(|\omega_{11}|), \end{aligned}$$

where

$$F_5(x) = \frac{1}{5} \left( 1 - x^2 - \frac{27}{4}x^4 \right) + \left( \frac{4}{\sqrt{5}}x^3 + \frac{8}{\sqrt{7}}x^2 \right) \sqrt{1 - x^2 - \frac{27}{4}x^4},$$

$0 \leq x \leq \frac{1}{2}$ . The first derivative test shows that the function  $F_5(x)$  when  $0 \leq x \leq 1/2$ , has maximal value 0.6647958756... attained for  $x = 0.458573 \dots$

□

\*\*\*\*\* DTS

**Theorem 6.** Let  $f \in \mathcal{S}$  and be given by (1).

- (i) Then  $|a_4| - |a_3| \leq 1.75185 \dots$
- (ii) If  $f$  is an odd function, then  $|a_5| - |a_3| \leq \frac{2}{\sqrt{7}} = 0.7559 \dots$

*Proof.*

(i) Using (6) and  $|\omega_{11}| \leq 1$ , we have

$$(24) \quad |a_4| - |a_3| \leq |a_4| - |\omega_{11}| |a_3| \leq |a_4 - \omega_{11} a_3| = 2 \left| \omega_{33} + 3\omega_{11}\omega_{13} + \frac{1}{6}\omega_{11}^3 \right|.$$

From the fifth relation in (6) we obtain

$$(25) \quad \omega_{33} = \omega_{15} - \omega_{11}\omega_{13} + \frac{1}{3}\omega_{11}^3,$$

and using the relations (24) and (25), after some simple calculations we have

$$\begin{aligned} |a_4| - |a_3| &\leq |2\omega_{15} + 4\omega_{11}\omega_{13} + \omega_{11}^3| \\ &\leq |\omega_{15}| + 4|\omega_{11}||\omega_{13}| + |\omega_{11}|^3 \\ &\leq \frac{2}{\sqrt{5}} \sqrt{1 - |\omega_{11}|^2 - 3|\omega_{13}|^2} + 4|\omega_{11}||\omega_{13}| + |\omega_{13}|^3 \\ &=: F_6(|\omega_{11}|, |\omega_{13}|), \end{aligned}$$

where

$$F_6(x, y) = \frac{2}{\sqrt{5}} \sqrt{1 - x^2 - 3y^2} + 4xy + x^3,$$

and  $0 \leq x \leq 1$ ,  $0 \leq y \leq \frac{1}{\sqrt{3}} \sqrt{1 - x^2}$ .

Now, we need to find maximum of the function  $F_6(x, y)$  when

$$(x, y) \in \left\{ (x, y) : 0 \leq x \leq 1, 0 \leq y \leq \frac{1}{\sqrt{3}} \sqrt{1 - x^2} \right\} := D_1.$$

We start the analysis in the interior of  $D_1$ . From

$$\frac{\partial F_6(x, y)}{\partial x} = -\frac{2x}{\sqrt{5}\sqrt{1 - x^2 - 3y^2}} + 3x^2 + 4y$$

and

$$\frac{\partial F_6(x, y)}{\partial y} = 4x - \frac{6y}{\sqrt{5}\sqrt{1 - x^2 - 3y^2}},$$

we receive that the stationary points (if existing) satisfy

$$3y \frac{\partial F_6(x, y)}{\partial x} - x \frac{\partial F_6(x, y)}{\partial y} = x^2(9y - 4) + 12y^2 = 0,$$

i.e., on  $D_1$ ,

$$x = \frac{2\sqrt{3}y}{\sqrt{4 - 9y}}.$$

Finally, substituting  $\frac{2\sqrt{3}y}{\sqrt{4 - 9y}}$  for  $x$  in  $\frac{\partial F_6(x, y)}{\partial x} = 0$ , and numerically solving the corresponding equation for  $y$ , we receive solution  $y_{01} = 0.2872\dots$ , and further  $x_{01} = \frac{2\sqrt{3}y_{01}}{\sqrt{4 - 9y_{01}}} = 0.83634\dots$ . It is easy to check that  $(x_{01}, y_{01}) \in D_1$  and that  $F_6(x_{01}, y_{01}) = 1.75185\dots$ .

On the edges of  $D_1$  we have:

- $F_6(0, y) = \frac{2}{\sqrt{5}} \sqrt{1 - 3y^2}$ , with maximal value for  $y \geq 0$ ,  $\frac{2}{\sqrt{5}} = 0.8944\dots$ ;
- $F_6(x, 0) = \frac{2}{\sqrt{5}} \sqrt{1 - x^2} + x^3$ , with maximal value for  $0 \leq x \leq 1$  equaling to  $1.13666\dots$  for  $x = 0.9494\dots$ ;
- $F_6\left(x, \frac{\sqrt{1 - x^2}}{\sqrt{3}}\right) = \frac{4}{\sqrt{3}} x \sqrt{1 - x^2} + x^3$ , with maximal value for  $0 \leq x \leq 1$  equaling to  $1.6496\dots$  for  $x = 0.8628\dots$



So, the function  $F_6$  attains its maximal value  $1.75185\dots$  in the interior point  $(x_{01}, y_{01})$  of its domain  $D_1$  and the conclusion of part (i) follows.

- (ii) Since  $f$  is odd,  $a_2 = 0$ , and then using  $|a_3 - a_2^2| \leq 1$ , we have  $|a_3| = |2\omega_{13}| \leq 1$ , i.e.,  $|\omega_{13}| \leq \frac{1}{2}$ . Further, from (6), (11) and (12), we receive

$$|a_5| = |2\omega_{17} + 3\omega_{13}^2|.$$

Also, since  $a_4 = 0$ , from (6) follows  $\omega_{33} = 0$  and  $\omega_{15} = 0$ . So, since  $|a_3| = |2\omega_{13}| \leq 1$ ,

$$\begin{aligned} |a_5| - |a_3| &\leq |a_5| - 2|\omega_{13}||a_3| = |2\omega_{17} + 3\omega_{13}^2| - 4|\omega_{13}|^2 \\ &\leq |(2\omega_{17} + 3\omega_{13}^2) - 4\omega_{13}^2| = |2\omega_{17} - \omega_{13}^2| \\ &\leq 2|\omega_{17}| + |\omega_{13}|^2 \leq \frac{2}{\sqrt{7}}\sqrt{1 - 3|\omega_{13}|^2} + |\omega_{13}|^2 \\ &\leq \frac{2}{\sqrt{7}}, \end{aligned}$$

where  $|\omega_{13}| \leq \frac{1}{2}$ .

□

**Remark 1.** *The estimate given in Theorem 6(ii) is an improvement of the result  $|a_5| - |a_3| < 1$  given in [6, p. 17].*

#### REFERENCES

1. P.L. Duren, *Univalent function*, Springer-Verlag, New York, 1983.
2. M. Fekete and G. Szegő, *Eine Bemerkung über ungerade schlichte Funktionen*, J. London Math. Soc. **8** (1933), 85–89.
3. A. Z. Grinspan, *The sharpening of the difference of the moduli of adjacent coefficients of schlicht functions*, In Some problems in modern function theory Proc. Conf. Modern Problems of Geometric Theory of Functions, Inst. Math., Acad. Sci. USSR, Novosibirsk, (Russian), Akad. Nauk SSSR Sibirsk. Otdel. Inst. Mat., Novosibirsk, (1976), 41–45.
4. W. K. Hayman, *On successive coefficients of univalent functions*, J. London Math. Soc., **38** (1963), 228–243.
5. N.A. Lebedev, *Area principle in the theory of univalent functions*, Published by "Nauka", Moscow, 1975 (in Russian).
6. A. Lecko, D. Partyka, *A generalized Fekete-Szegő functional and initial successive coefficients of univalent functions*, Bulletin des Sciences Mathématiques, 197 (2024), art. id. 103527.
7. M. Obradović and N. Tuneski, *Hankel determinants of second and third order for the class  $\mathcal{S}$  of univalent functions*, Math. Slovaca **71**(3), 649-654, 2021
8. Milutin Obradović and Nikola Tuneski, *An improvement fo the Hankel determinants of second and third order for the class  $\mathcal{S}$  of univalent functions*, .
9. M. Obradović, D.K. Thomas and N. Tuneski, *On the difference of coefficients of univalent functions*, Filomat **35**:11 (2021), 3653-3661.
10. D.K. Thomas, N. Tuneski, A. Vasudevarao, *Univalent Functions: A Primer, De Gruyter Studies in Mathematics* **69**, De Gruyter, Berlin, Boston, 2018.

DEPARTMENT OF MATHEMATICS, FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF BELGRADE,  
BULEVAR KRALJA ALEKSANDRA 73, 11000, BELGRADE, SERBIA  
Email address: obrad@grf.bg.ac.rs

DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF MECHANICAL ENGINEERING,  
SS. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE, KARPOŠ II B.B., 1000 SKOPJE, REPUBLIC OF  
NORTH MACEDONIA.  
Email address: nikola.tuneski@mf.edu.mk