

Sampling and Integration of Logconcave Functions by Algorithmic Diffusion

Yunbum Kook
Georgia Tech
yb.kook@gatech.edu

Santosh S. Vempala
Georgia Tech
vempala@gatech.edu

Abstract

We study the complexity of sampling, rounding, and integrating arbitrary logconcave functions. Our new approach provides the first complexity improvements in nearly two decades for general logconcave functions for all three problems, and matches the best-known complexities for the special case of uniform distributions on convex bodies. For the sampling problem, our output guarantees are significantly stronger than previously known, and lead to a streamlined analysis of statistical estimation based on dependent random samples.

Contents

1	Introduction	1
1.1	Results	4
1.2	Techniques and background	6
1.3	Preliminaries	12
2	Sampling from logconcave distributions	13
2.1	Reduction to an exponential distribution	14
2.2	Proximal sampler for the reduced exponential distribution	16
3	Warm-start generation via tilted Gaussian cooling	22
3.1	Tilted Gaussian Cooling	23
3.2	Proximal sampler for annealing distributions	24
3.3	Rényi-infinity sampling for logconcave distributions	31
4	Application 1: Rounding logconcave distributions	34
4.1	Isotropic rounding of a well-rounded distribution	36
4.2	Maintaining well-roundedness	39
4.3	Rounding the grounded distribution	42
5	Application 2: Integrating logconcave functions	43
5.1	Algorithm	44
5.2	Streamlined analysis	46
A	Functional inequalities and logconcave geometry	57

1 Introduction

Sampling and integration of logconcave functions are fundamental problems with numerous applications and important special cases such as uniform distributions over convex bodies and strongly logconcave densities. The study of these problems has led to many useful techniques. Both mathematically and algorithmically, general logconcave functions typically provide the “right” general abstraction. For example, many classical inequalities for convex bodies have natural extensions to logconcave functions (e.g., Grünbaum’s theorem, Brunn-Minkowski and Prékopa-Leindler, isotropic constant, etc.). The KLS hyperplane conjecture, first motivated by the analysis of the Ball Walk for sampling convex bodies, is in the setting of general logconcave densities. The current fastest algorithm for estimating the volume of a convex body crucially uses sampling from a sequence of logconcave densities, which is provably more efficient than using a sequence of uniform distributions. Sampling logconcave densities has many other applications as well, such as portfolio optimization, simulated annealing, Bayesian inference, differential privacy etc.

Sampling in high dimension is done by randomized algorithms based on Markov chains. These chains are set up to have a desired stationary distribution, which is relatively easy to ensure. For example, to sample uniformly, it suffices that the Markov chain is symmetric. Generally, to sample proportional to a desired function, it suffices to ensure the “detailed balance” condition. The main challenge is showing rapid mixing of the Markov chain, i.e., the convergence rate to the stationary distribution is bounded by a (small) polynomial in the dimension and other relevant parameters.

The traditional analysis of Markov chains for sampling high-dimensional distributions proceeds by analyzing the *conductance* of the Markov chain, the minimum conditional escape probability over all subsets of the state space of measure at most half. Bounding this is done by relating probabilistic (one-step distribution) distance to geometric distance, and then using a purely isoperimetric inequality for subsets of the support. Indeed, this approach led to several interesting questions and useful techniques, in particular the development of isoperimetric inequalities and the discovery of (nearly) tight bounds in many settings. To describe background and known results, let us first define the sampling problem (readers familiar with the problem can skip to §1.1).

Model and problems. We assume access to an integrable logconcave function $\exp(-V(x))$ in \mathbb{R}^n , via an evaluation oracle for a convex function $V : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$. We also assume that there exists a point x_0 and parameters r, R such that for the distribution π with density $d\pi \propto \exp(-V) dx$, a ball of radius r centered at x_0 is contained in the level set of π of measure $1/8^1$ and $\mathbb{E}_\pi[\|X - x_0\|^2] \leq R^2$. We refer to this as a *well-defined function oracle*², denoted by $\text{Eval}_{\mathcal{P}}(V)$ where \mathcal{P} indicates access to the actual values of parameters in \mathcal{P} (e.g., $\text{Eval}_{x_0, R}(V)$ presents both x_0 and R , while $\text{Eval}(V)$ does not provide any parameters).

Given this oracle, in this paper we consider three central problems: (1) sample from the distribution π , (2) find an affine transformation that places π in near-isotropic position, and (3) estimate the integral $\int e^{-V(x)} dx$ (i.e., the normalization constant of π). We measure the complexity in terms of the number of oracle calls and the total number of arithmetic operations.

Complexity of logconcave sampling. Sampling general logconcave functions, as an algorithmic problem, was first studied by Applegate and Kannan [AK91], who established an algorithm with complexity polynomial in the dimension assuming Lipschitzness of the function over its support. They did this by creating a sufficiently fine grid (so that the function changed by at most a small constant within each grid cell) and used a discrete “grid walk”. They sampled a grid point and

¹We use the standard convention of $1/8$; any constant bounded away from 1 would suffice. In fact, we will use an even weaker condition; see Problem 2.1 and Lemma 2.6.

²Without loss of generality, we will assume by scaling that $r = 1$.

used rejection sampling to output a point from the cube associated with the grid point. The output distribution is guaranteed to be within desired total variation (TV) distance from the target distribution. Following many developments, Lovász and Vempala improved the bound to scale as $n^2 R^2 / r^2$ [LV07, LV06a]. More specifically, [LV07, Theorem 2.2, 2.3] analyzed the Ball Walk and Hit-and-Run and proved a bound of

$$\frac{M^4}{\varepsilon^4} \frac{n^2 R^2}{r^2} \log \frac{M}{\varepsilon}$$

steps/queries to reach a distribution within ε TV-distance of the target starting from an M -warm distribution. [LV06a, Theorem 1.1] showed that Hit-and-Run achieves the same guarantee in

$$\frac{n^2 R^2}{r^2} \log^{\mathcal{O}(1)} \frac{nMR}{\varepsilon r}$$

steps. We note that in all previous work on general logconcave sampling the output guarantees are in TV-distance. It is often desirable to have stronger guarantees such as the KL or Rényi divergences.

Classical sampling algorithms. For target density proportional to e^{-V} , the Ball Walk with parameter δ proceeds as follows: at a point x with $e^{-V(x)} > 0$, sample a uniform random point y in the ball of radius δ centered at x ; go to y with probability $\min(1, \frac{e^{-V(y)}}{e^{-V(x)}}$), staying at x with the remaining probability. Hit-and-Run does not need a parameter: at current point x , pick a uniform random line ℓ through x , and go to a random point y along ℓ with probability proportional to $e^{-V(y)}$ (marginal along ℓ). For technical reasons, both of these walks (and almost all other walks) need to be *lazy*, i.e., they do nothing with probability $1/2$, and do the above step with probability $1/2$.

Sampling and isoperimetry. The complexity of sampling is intuitively tied closely to the isoperimetry of the target distribution. If the target has poor isoperimetry (roughly, a small measure surface can partition the support into two large measure subsets), then any “local” Markov chain will have difficulty moving from one large subset to its complement. There are two alternative views of isoperimetry — functional and geometric. We recap the definitions of isoperimetric constants.

Definition 1.1. We say that a probability measure π on \mathbb{R}^n satisfies a *Poincaré inequality* (PI) with parameter $C_{\text{PI}}(\pi)$ if for all smooth functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$,

$$\text{var}_\pi f \leq C_{\text{PI}}(\pi) \mathbb{E}_\pi[\|\nabla f\|^2], \tag{PI}$$

where $\text{var}_\pi f = \mathbb{E}_\pi[(f - \mathbb{E}_\pi f)^2]$.

The Poincaré inequality is implied by the generally stronger log-Sobolev inequality.

Definition 1.2. We say that a probability measure π on \mathbb{R}^n satisfies a *log-Sobolev inequality* (LSI) with parameter $C_{\text{LSI}}(\pi)$ if for all smooth functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$,

$$\text{Ent}_\pi(f^2) \leq 2C_{\text{LSI}}(\pi) \mathbb{E}_\pi[\|\nabla f\|^2], \tag{LSI}$$

where $\text{Ent}_\pi(f^2) := \mathbb{E}_\pi[f^2 \log f^2] - \mathbb{E}_\pi[f^2] \log \mathbb{E}_\pi[f^2]$.

For the geometric view, we define $\pi(\partial S) := \liminf_{\varepsilon \downarrow 0} \frac{\pi(S_\varepsilon) - \pi(S)}{\varepsilon}$, where $S_\varepsilon := \{x : d(x, S) \leq \varepsilon\}$.

Definition 1.3. The *Cheeger constant* $C_{\text{Ch}}(\pi)$ of a probability measure π on \mathbb{R}^n is defined as

$$C_{\text{Ch}}(\pi) = \inf_{S \subset \mathbb{R}^n, \pi(S) \leq \frac{1}{2}} \frac{\pi(\partial S)}{\pi(S)}.$$

Definition 1.4. The *log-Cheeger constant* $C_{\log\text{Ch}}(\pi)$ of a probability measure π on \mathbb{R}^n is defined as

$$C_{\log\text{Ch}}(\pi) = \inf_{S \subset \mathbb{R}^n, \pi(S) \leq \frac{1}{2}} \frac{\pi(\partial S)}{\pi(S) \sqrt{\log(1/\pi(S))}}.$$

It is known that for logconcave measures, $C_{\text{PI}}(\pi) = \Theta(C_{\text{Ch}}^{-2}(\pi))$ [Mil09] and $C_{\text{LSI}}(\pi) = \Theta(C_{\log\text{Ch}}^{-2}(\pi))$ [Led94]. Bounding these constants has been a major research topic for decades. In recent years, following many improvements, it has been shown that for isotropic logconcave measures, $C_{\text{PI}}(\pi) \lesssim \log n$ [Kla23] and for isotropic logconcave ones with support of diameter D , we have $C_{\text{LSI}}(\pi) \lesssim D$ [LV24]. The former is conjectured to be $\mathcal{O}(1)$ (the KLS conjecture), and the latter is the best possible.

The significance of these constants for algorithmic sampling became clear with the analysis of the **Ball Walk**, where the bound on its convergence from a warm start to a logconcave distribution depends directly on $C_{\text{Ch}}^{-2}(\pi)$ [KLS97] (this was the original motivation for the KLS conjecture) — the mixing time of the **Ball Walk** starting from an M -warm distribution to reach ε TV-distance is bounded by $\mathcal{O}(n^2 C_{\text{Ch}}^{-2} \text{poly}(\frac{M}{\varepsilon}))$. For the special case of uniformly sampling convex bodies, using the notion of a **Speedy Walk**, this can be improved to $\mathcal{O}(Mn^2 C_{\text{Ch}}^{-2} \log \frac{M}{\varepsilon})$, but using it in the analysis entails handling several technical difficulties. In principle, this should be extendable to general logconcave functions, albeit with formidable technical complications. We take a different approach.

Sampling, isoperimetry, and diffusion. The connection of isoperimetric constants to the convergence of continuous-time *diffusion* is a classical subject. The Langevin diffusion is a canonical stochastic differential equation (SDE) given by

$$dX_t = -\nabla V(X_t) dt + \sqrt{2} dB_t \quad X_0 \sim \nu_0,$$

where B_t is the standard Brownian process, and V is a smooth function. Then, under mild conditions, the law ν_t of X_t converges to the distribution with density ν proportional to e^{-V} in the limit. Moreover, it can be shown that

$$\chi^2(\nu_t \parallel \nu) \leq \exp\left(-\frac{2t}{C_{\text{PI}}(\nu)}\right) \chi^2(\nu_0 \parallel \nu), \quad \text{and} \quad \text{KL}(\nu_t \parallel \nu) \leq \exp\left(-\frac{2t}{C_{\text{LSI}}(\nu)}\right) \text{KL}(\nu_0 \parallel \nu).$$

A natural idea for a sampling algorithm is to discretize the diffusion equation. This discretized algorithm was shown to converge in $\text{poly}(n, \beta, C_{\text{PI}}(\nu))$ and $\text{poly}(n, \beta, C_{\text{LSI}}(\nu))$ iterations under **(PI)** [CEL⁺22] and **(LSI)** [VW19] respectively, assuming β -Lipschitzness of ∇V . Can this approach be extended to sampling general logconcave densities without the gradient Lipschitzness?

Algorithmic diffusion. By *algorithmic diffusion*, we mean the general idea of discretizing a diffusion process and proving guarantees on the discretization error and query complexity. Recently, [KVZ24] showed that such an algorithm (called **In-and-Out** there), which can be viewed as a **Proximal Sampler**, works for sampling from the uniform distribution over a convex body and recovers state-of-the-art complexity guarantees with substantially improved output guarantees (Rényi-divergences). The guarantee was extended to the Rényi-infinity (or pointwise) distance as well [KZ25]. A nice aspect of the analysis is that it shows convergence directly in terms of isoperimetric constants of target distributions. As a result, it provides a unifying point of view for analysis without requiring the use of technically sophisticated tools such as the **Speedy Walk** and s -conductance [LS93, KLS97, LV07]. We discuss the approach in more detail presently.

This brings us to the main motivation of the current paper: *Can we use algorithmic diffusion to obtain faster algorithms with stronger guarantees for sampling, rounding, and integrating general logconcave functions?*

Algorithm 1 Proximal Sampler PS_{exp}

Input: initial pt. $z_0 \sim \pi_0 \in \mathcal{P}(\mathbb{R}^{n+1})$, $\mathcal{K} = \{(x, t) : V(x) \leq nt\}$, $k \in \mathbb{N}$, threshold N , variance h .

Output: $z_{k+1} = (x_{k+1}, t_{k+1})$.

- 1: **for** $i = 0, \dots, k$ **do**
 - 2: Sample $y_{i+1} \sim \mathcal{N}(z_i, hI_{n+1})$.
 - 3: Sample $z_{i+1} \sim \mathcal{N}(y_{i+1} - hne_{n+1}, hI_{n+1})|_{\mathcal{K}}$.
 - 4: (↑) Repeat $z_{i+1} \sim \mathcal{N}(y_{i+1} - hne_{n+1}, hI_{n+1})$ until $z_{i+1} \in \mathcal{K}$. If $\#\text{attempts}_i \geq N$, declare **Failure**.
 - 5: **end for**
-

1.1 Results

In this paper, we propose new approaches for general logconcave sampling, rounding and integration, leading to the first improvements in the complexity of these problems in nearly two decades [LV06a]. Our methods crucially rely on a reduction from general logconcave sampling to *exponential sampling* in one higher dimension. With this reduction in hand, we develop a new framework for sampling and improve the complexity of four fundamental problems: (1) logconcave sampling from an $\mathcal{O}(1)$ -warm start, (2) warm-start generation, (3) isotropic rounding and (4) integration. For each problem, our improved complexity for general logconcave distributions matches the current best complexity for the uniform distribution over a convex body, a setting that has received much attention for more than three decades [DFK91, LS90, Lov91, LS93, KLS97, LV06b, CV18, KVZ24, KZ25]. We now present our main results, followed by a detailed discussion of the challenges and techniques in §1.2, and notations and preliminaries in §1.3.

Result 1: Logconcave sampling from a warm start. When a logconcave distribution $\pi^X \propto e^{-V}$ is presented with the evaluation oracle for V , we instead consider the “lifted” distribution

$$\pi(x, t) \propto \exp(-nt) \cdot \mathbb{1}[(x, t) : V(x) \leq nt]$$

in the augmented (x, t) -space, the X -marginal of which is exactly π^X (see Proposition 2.3). We quantify in §2 how this affects parameters pertinent to sampling (e.g., isoperimetric constants).

To sample from the exponential target π using a given \mathcal{R}_∞ warm-start, we use the Proximal Sampler (see Figure 1.1) with an evaluation oracle that returns nt conditioned on $V(x) \leq nt$. One iteration of the sampler is reversible with respect to the stationary distribution π , while it can be interpreted as a composition of two complimentary diffusion processes. Discarding the T -component of a sample, we are left with the X -component with law close to our desired target π^X . In this regard, our sampler is a natural generalization of In-and-Out [KVZ24], the Proximal Sampler for uniform sampling. Concretely, we establish mixing guarantees and query complexities of our sampler for the target logconcave distribution π^X in terms of its Poincaré constant (which in general satisfies $\|\text{Cov } \pi^X\| \leq C_{\text{PI}}(\pi^X) \lesssim \|\text{Cov } \pi^X\| \log n$ [Kla23]).

Theorem 1.5. *For any logconcave distribution π^X specified by a well-defined function oracle $\text{Eval}(V)$, for any given $\eta, \varepsilon \in (0, 1)$, $q \geq 2$, and π_0^X with $\mathcal{R}_\infty(\pi_0^X \| \pi^X) = \log M$, we can use the Proximal Sampler PS_{exp} (Algorithm 1) with suitable choices of parameters, so that with probability at least $1 - \eta$, we obtain a sample X such that $\mathcal{R}_q(\text{law } X \| \pi^X) \leq \varepsilon$, using $\tilde{\mathcal{O}}(qMn^2 (\|\text{Cov } \pi^X\| \vee 1) \text{polylog } \frac{1}{\eta\varepsilon})$ evaluation queries in expectation.*

See §2.2.1 for a more detailed description of the heat flow perspective and its benefits for sampling. Our specific setting of parameters can be found in Theorem 2.15. In comparison with the previous best complexity of $\tilde{\mathcal{O}}(n^2 \text{tr}(\text{Cov } \pi^X) \text{polylog } \frac{M}{\varepsilon})$, which is for TV-distance, the Proximal Sampler




Figure 1.1: Reduction to an exponential distribution and sampling via the Proximal Sampler PS_{exp} .

PS_{exp} achieves a provably better rate (since $\|\text{Cov } \pi^X\| \leq \text{tr}(\text{Cov } \pi^X)$) from an $\mathcal{O}(1)$ -warm start, and moreover does so in general \mathcal{R}_q -divergences. This mixing rate matches the previously known best rate for the uniform sampling by the Ball Walk [KLS97] as well as In-and-Out [KVZ24].

Result 2: Warm-start generation (sampling without a warm start). The Proximal Sampler PS_{exp} assumes access to a warm start in \mathcal{R}_∞ , and generating such a good warm-start is an important and challenging algorithmic problem in its own right. In §3, we propose Tilted Gaussian Cooling which generates an $\mathcal{O}(1)$ -warm start in the \mathcal{R}_∞ -divergence for any target logconcave distribution. This algorithm generalizes Gaussian Cooling [CV18], the known method for generating a warm-start for the uniform distribution over a convex body. The high-level idea is to follow a sequence $\{\mu_i\}_{i \in [m]}$ of distributions, where μ_1 is easy to sample from, μ_i and μ_{i+1} are close in some probability divergences, and μ_m is the desired target.

Our algorithm first reduces the original target to the exponential distribution and then follows annealing distributions of the form

$$\mu_{\sigma^2, \rho}(x, t) \propto \exp\left(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t\right) \cdot \mathbb{1}[\{V(x) \leq nt\} \cap \{\|x\| = \mathcal{O}(R), |t| = \mathcal{O}(1)\}],$$

with a carefully chosen schedule for updating the parameters σ^2 and ρ . Roughly speaking, we update the X -marginal through Gaussian Cooling and the T -marginal using an exponential *tilt*. In order to move across the annealing distributions, Tilted Gaussian Cooling requires an efficient sampler for the intermediate annealing distributions, ideally with guarantees in the \mathcal{R}_∞ -divergence in order to relay \mathcal{R}_∞ -warmness guarantees along the annealing scheme. We use the Proximal Sampler for these intermediate distributions with a \mathcal{R}_∞ -divergence guarantee, namely the Proximal Sampler PS_{ann} (Algorithm 3) in §3.2. In our computational model, this sampler started at a previous annealing distribution returns a sample with law μ satisfying $\mathcal{R}_\infty(\mu \parallel \mu_{\sigma^2, \rho}) \leq \varepsilon$, using $\tilde{\mathcal{O}}(n^2 \sigma^2 \text{polylog } R/\eta\varepsilon)$ evaluation queries in expectation (see Theorem 3.11).

By sampling these annealing distributions through PS_{ann} , Tilted Gaussian Cooling obtains an $\mathcal{O}(1)$ -warm start for π (and hence for the desired target π^X); then runs the Proximal Sampler PS_{exp} one final time to obtain a sample with \mathcal{R}_∞ -guarantees (this is a very strong notion of probability divergence that recovers all commonly used distances such as TV, KL, χ^2 , or \mathcal{R}_q).

Theorem 1.6. *For any logconcave distributions π^X specified by $\text{Eval}_{x_0, R}(V)$, for any given $\eta, \varepsilon \in (0, 1)$, Tilted Gaussian Cooling (Algorithm 2) with probability at least $1 - \eta$, returns a sample with law ν such that $\mathcal{R}_\infty(\nu \parallel \pi^X) \leq \varepsilon$, using $\tilde{\mathcal{O}}(n^2(R^2 \vee n) \text{polylog } 1/\eta\varepsilon)$ evaluation queries in expectation. Hence, if π^X is well-rounded (i.e., $R^2 \lesssim n$), then $\tilde{\mathcal{O}}(n^3 \text{polylog } 1/\eta\varepsilon)$ queries suffice.*

This improves the prior best complexity of $\tilde{\mathcal{O}}(n^3 R^2 \text{polylog } \frac{1}{\varepsilon})$ for general logconcave sampling (in TV-distance) from scratch by [LV06a, Corollary 1.2], and provides a much stronger \mathcal{R}_∞ -guarantee.

Moreover, this complexity matches the best-known complexity for uniform sampling with the TV-guarantee by [CV18] and with the \mathcal{R}_∞ -guarantee by [KZ25].

Result 3: Isotropic rounding of logconcave distributions. The above results on the complexity of logconcave sampling still have dependence on the second moment R^2 , so those are not fully polynomial in the problem parameters. In §4, we address this issue by initially running an algorithm for isotropic rounding which makes the covariance matrix of a given logconcave distribution near-isotropic (i.e., $\text{Cov } \pi^X \approx I_n$). After this rounding, $R^2 = \mathcal{O}(n)$, and thus the guarantee above turns into n^3 at the cost of a multiplicative factor of polylog R in the complexity. Isotropic rounding is also a useful tool for many other high-dimensional algorithms.

At a high level, just as in the warm-start generation, our algorithm follows a sequence $\{\mu_i\}$ of distributions while updating an affine map $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ along the way. An important subroutine is to convert a well-rounded distribution (i.e., with $\text{tr}(\text{Cov}) = \mathcal{O}(n)$) to one that is near-isotropic (i.e., with $\text{Cov} \approx I_n$). To this end, we generalize the approach taken in [JLLV21] for the uniform distribution — repeatedly [draw a few samples \rightarrow compute a crude estimate of the covariance \rightarrow identify skewed directions from the estimation and upscale them]. In doing so, we rely crucially on our sampler’s mixing rate in terms of $\|\text{Cov } \pi^X\|$ (instead of $\text{tr}(\text{Cov } \pi^X)$) in **Result 1** and its improved complexity for warm-start generation in **Result 2**.

Theorem 1.7. *For any logconcave distribution π^X specified by $\text{Eval}_{x_0, R}(V)$, there exists a randomized algorithm with query complexity of $\tilde{\mathcal{O}}(n^{3.5} \text{polylog } R)$ that finds an affine map F such that the pushforward of π^X via F is 1.01-isotropic with probability at least $1 - \mathcal{O}(n^{-1/2})$.*

Our complexity improves upon the previous best complexity of $\tilde{\mathcal{O}}(n^4 \text{polylog } R)$ for general logconcave distributions [LV06a, Theorem 6.1]. Moreover, our rate matches that of isotropic rounding for the uniform distribution [JLLV21, JLLV24].

Result 4: Integration of logconcave functions. Finally, we examine the complexity of integrating a logconcave function, a classical application of logconcave sampling. To this end, we extend an annealing approach in [CV18] to general logconcave integration in §5. Similar to Tilted Gaussian Cooling, we follow a sequence $\{f_i\}_{i \in [m]}$ of logconcave functions, where f_1 is easy to integrate, $\mu_i \propto f_i$ is an $\mathcal{O}(1)$ -warm start for $\mu_{i+1} \propto f_{i+1}$, the variance of f_{i+1}/f_i with respect to μ_i is small enough (e.g., $\chi^2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(m^{-1})$), and f_m is the target logconcave function.

Theorem 1.8. *For any $\varepsilon > 0$ and an integrable well-rounded logconcave function $f = e^{-V} : \mathbb{R}^n \rightarrow \mathbb{R}$ presented by $\text{Eval}_{x_0, R}(V)$, there exists an algorithm that with probability at least $3/4$, returns a $(1 + \varepsilon)$ -multiplicative approximation to the integral of f using $\tilde{\mathcal{O}}(n^3/\varepsilon^2)$ queries. For an arbitrary logconcave f given by a well-defined function oracle, the complexity is bounded by $\tilde{\mathcal{O}}(n^{3.5} \text{polylog } R + n^3/\varepsilon^2)$.*

This result improves the previous best complexity $\tilde{\mathcal{O}}(n^4/\varepsilon^2)$ of integrating logconcave functions [LV06a, Theorem 1.3]. Again, our improved complexity matches that for the uniform distribution (i.e., volume computation) [CV18, JLLV21, JLLV24]. We note that due to the strong guarantees of the new sampler, we can streamline the analysis of errors and dependence among samples used to estimate f_{i+1}/f_i , simplifying and strengthening earlier analyses [KLS97, LV06c, CV18].

1.2 Techniques and background

In proving our results — faster algorithms for the basic problems of sampling (with and without a warm start), rounding and integration — we do not take the most direct route. Instead we develop several techniques that appear to be interesting and we imagine will be useful in other contexts as well. These include the reversible heat-flow perspective for the design of polynomial-time algorithms

providing a direct connection to isoperimetric constants, sampling guarantees in the strong notion of \mathcal{R}_∞ -distance (which has other motivations such as differential privacy), relevant geometry of logconcave functions, and a streamlined analysis of estimation errors of dependent samples.

1.2.1 Going beyond the uniform distribution

The uniform distribution over a convex body is a special case of a logconcave distribution. The best-known polynomial complexity for uniform sampling is achieved by addressing three subproblems: (1) sampling from a warm start, (2) generating a warm start and (3) performing isotropic rounding. Extending the best-known guarantees and algorithms for each of these problems to general logconcave densities is the primary challenge addressed in this paper. In our proofs, we use general properties of logconcave functions and avoid any structural assumptions on the target density.

The first challenge is to establish a logconcave sampler with a mixing rate matching that of the best uniform samplers, such as the Ball Walk [KLS97] or In-and-Out [KVZ24]. These uniform samplers have a mixing rate of $n^2 \|\text{Cov } \pi^X\|$, while Hit-and-Run in [LV06a], previously the best for general logconcave distributions, has a rate of $n^2 \text{tr}(\text{Cov } \pi^X)$. The improved complexity bounds of these uniform samplers come from a better understanding of isoperimetric properties, such as the Cheeger and Poincaré constants for logconcave distributions, and this aspect is more direct for In-and-Out. Here we extend In-and-Out to general logconcave distributions. As we elaborate in §1.2.2, this requires implementing rejection sampling for the distributions of the form $\exp(-V(x) - \frac{1}{2h} \|x\|^2)$ (for a suitable h) using $\tilde{\mathcal{O}}(1)$ queries. However, with only logconcavity assumed for e^{-V} , determining a suitable proposal distribution and analyzing its query complexity present significant challenges.

Reduction to an exponential distribution. To extend methods from uniform to general logconcave distributions, we leverage a conceptual connection between logconcave sampling and convex optimization — sampling from $\exp(-V)$ is analogous to minimization of V . In optimization, we bound $V(x)$ with a new variable t , and add the convex constraint $V(x) \leq t$. Inspired by this, we consider an exponential distribution with density $\pi(x, t) \propto \exp(-nt) \cdot \mathbb{1}[(x, t) : V(x) \leq nt]$ on \mathbb{R}^{n+1} , with X -marginal $\pi^X \propto \exp(-V)$ (Proposition 2.3). The scaling of n in the exponential function is a natural choice as we will see later.

This reduction offers several advantages. First, the potential V becomes linear, opening up a possible extension of previous analyses. Second, it points to a clear path to generalizing ideas/guarantees devised for uniform sampling. Specifically, the conditional law $\pi^{X|T=t}$ of X given $T = t$ is the uniform distribution over the convex level set $L_{nt} = \{V(x) \leq nt\}$. Thus, π can be interpreted as an average of the uniform distribution over L_{nt} weighted by e^{-nt} , and this idea is crucial to our subsequent algorithms.

A similar exponential reduction was discussed conceptually in [KLSV22] and developed further in [KV24] for a sampling analogue of the interior-point method from optimization. Here, however, we apply this reduction in our general setup, without specific assumptions on the epigraph, and analyze structural properties of the reduced distribution. For example, we bound the mean and variance along the T -direction. This allows us to analyze how this reduction impacts key parameters of the original logconcave distribution, including the largest eigenvalue and trace of the covariance.

1.2.2 Logconcave sampling from a warm start

Sampling through diffusion. The proximal sampler [LST21] is essentially a Gibbs sampler from two conditional distributions. For a target distribution $\pi^X \propto \exp(-V)$, it introduces a new variable Y and considers the augmented distribution $\pi^{X,Y} \propto \exp(-V(x) - \frac{1}{2h} \|x - y\|^2)$ with parameter $h > 0$. One iteration involves two steps: (i) $y \sim \pi^{Y|X=x} = \mathcal{N}(x, hI_n)$ and (ii)

$x \sim \pi^{X|Y=y} \propto \exp(-V(\cdot) - \frac{1}{2h} \|\cdot - y\|^2)$. While step (i) is straightforward, step (ii) requires a nontrivial sampling procedure.

The complexity of the Proximal Sampler involves a mixing analysis (to determine the number of iterations for a desired accuracy to the target) and the query complexity of implementing step (ii). For the first part, [CCSW22] demonstrated that one iteration corresponds to simulating the heat flow for (i) and then a *time reversal* of the heat flow for (ii) (see §2.2.1 for details). This leads to exponential decay of the \mathcal{R}_q -divergence, with the decay rate dependent on the isoperimetry of the target distribution (e.g. (PI) and (LSI)). For the second part, prior approaches typically use rejection sampling or a different logconcave sampler (e.g., MALA, ULMC) under additional assumptions such as the smoothness of V or access to first-order and proximal oracles for V .

Sampling without smoothness. Previous studies of the Proximal Sampler focused on smooth unconstrained distributions, where the potential satisfies $\nabla^2 V \preceq \beta I_n$ for some $\beta < \infty$, leaving open the complexities of uniform or general logconcave sampling with hard constraints. [KVZ24] introduced In-and-Out, a version of the Proximal Sampler for uniform sampling over convex bodies. With $V(x) = (\mathbf{1}[x \in \mathcal{K}])^{-1}$ for a convex body \mathcal{K} , one iteration of In-and-Out draws $y \sim \mathcal{N}(x, hI_n)$ and then $x \sim \mathcal{N}(y, hI_n)|_{\mathcal{K}}$ (the Gaussian truncated to \mathcal{K}) using rejection sampling on the proposal $\mathcal{N}(y, hI_n)$. They introduce a *threshold* parameter on the number of rejection trials which ensures that the algorithm does not use too many queries.

This diffusion-based approach turns out to be stronger than the Ball Walk, the previous best uniform sampler with query complexity $\tilde{O}(n^2 \|\text{Cov } \pi^X\| \text{polylog } \frac{1}{\varepsilon})$ for obtaining an ε -close sample in TV-distance. In-and-Out provides \mathcal{R}_q -divergence guarantees with a matching rate of $n^2 \|\text{Cov } \pi^X\|$ for TV-distance, and its analysis is simpler than the Ball Walk. Since the latter’s analysis in [KLS97] goes through its biased version (called the Speedy Walk), it involves an understanding of an additional rejection step for making the biased distribution close to the uniform target, as well as the isoperimetric constant of the biased distribution. In contrast, In-and-Out achieves direct contraction towards the uniform target, with rate dependent on isoperimetric constants of the original target, not a biased one, and achieves stronger output guarantees. In-and-Out’s approach has also been extended to truncated Gaussian sampling [KZ25], leading to a \mathcal{R}_∞ -guarantee that improves upon the TV-guarantee of the Ball Walk [CV18].

Proximal sampler for general logconcave distributions. This prompts a natural question if we can extend In-and-Out beyond constant and quadratic potentials. Prior studies of the Proximal Sampler by [CCSW22, KVZ24] allow for an immediate mixing analysis through the isoperimetry of logconcave distributions. However, we also need a query complexity bound for sampling from $\pi^{X|Y=y} \propto \exp(-V(x) - \frac{1}{2h} \|x - y\|^2)$. While one might consider using Hit-and-Run [LV06a] for the second step, it requires *roundedness* of this distribution, and its current complexity is also unsatisfactory. Rejection sampling is another option, but without smoothness assumptions for general logconcave distributions, it is challenging to identify an appropriate proposal distribution and to bound the expected number of trials.

To handle an arbitrary convex potential V , we apply the exponential reduction, turning V into a convex constraint and work with the linear potential nt in one higher dimension. Extending In-and-Out to this exponential distribution, we propose the Proximal Sampler PS_{exp} (Algorithm 1). Since $C_{\text{PI}}(\pi) \lesssim (C_{\text{PI}}(\pi^X) + 1) \log n$ (Lemma 2.5), the mixing rate of PS_{exp} for π is close to that of the Proximal Sampler for π^X . Implementing the second step is now simpler due to the linear potential, allowing us to extend the analysis of In-and-Out for uniform and truncated Gaussian distributions to exponential distributions (Lemma 2.14). A new technical ingredient is bounding the rate of increase of $\int_{\mathcal{K}_\delta} e^{-nt} / \int_{\mathcal{K}} e^{-nt}$ as δ grows, where $\mathcal{K}_\delta = \{z \in \mathbb{R}^{n+1} : d(z, \mathcal{K}) \leq \delta\}$. We show that this growth rate is bounded by $e^{\delta n}$ (Lemma 2.6 and 2.13).

1.2.3 Warm-start generation

Prior work on warm-start generation [KLS97, LV06c, CV18] is based on using a sequence $\{\mu_i\}_{i \in [m]}$ of distributions, where μ_1 is easy to sample (e.g., uniform distribution over a unit ball), each μ_i is close to μ_{i+1} in probabilistic distance, and μ_m is the target. By moving along this sequence with a suitable sampler for intermediate annealing distributions, one can generate a warm start for the desired distribution. This approach is more efficient than trying to directly go from μ_1 to μ_m .

Warm-start generation for uniform distributions. The state-of-the-art algorithm for uniform distributions over convex bodies is **Gaussian Cooling** [CV18]. They set $\mu_i(x) \propto \exp(-\|x\|^2/(2\sigma_i^2))$, where σ_i^2 increases from n^{-1} to R^2 according to a suitable schedule. This approach ensures that μ_i is $\mathcal{O}(1)$ -warm with respect to μ_{i+1} , and uses the **Ball Walk** to sample from truncated Gaussians with TV-guarantees. Using a coupling argument, they showed that with high probability, this scheme outputs an ε -close sample to the uniform distribution in TV-distance, using $\tilde{\mathcal{O}}(n^2(R^2 \vee n) \text{polylog} \frac{1}{\varepsilon})$ membership queries. [KZ25] later improved this by replacing the **Ball Walk** with the **Proximal Sampler**, achieving \mathcal{R}_∞ -guarantees with the same complexity. As a result, **Gaussian Cooling** transfers \mathcal{R}_∞ -warmness across the sequence of distributions, and thus the complexity for uniform sampling from a convex body remains the same even for \mathcal{R}_∞ -divergence guarantees.

Going beyond uniform distributions. In [CV18], Cousins and Vempala raised the question of whether their annealing strategy can be extended to arbitrary logconcave distributions with complexity $\tilde{\mathcal{O}}(n^2(R^2 \vee n))$. A natural choice for annealing distributions is $\mu(x) \propto \exp(-V(x) - \frac{1}{2\sigma^2} \|x\|^2)$, which would still provide \mathcal{R}_∞ -closeness of consecutive distributions and allow for accelerated updates to σ^2 . However, prior samplers lack the necessary guarantees for these distributions, so we use the exponential reduction.

To generate a warm start for $\pi(x, t) \propto e^{-nt}|_{\mathcal{K}}$, it seems natural to consider an annealing distribution obtained by multiplying π by a Gaussian in “ (x, t) ” for a direct application of **Gaussian Cooling** in \mathbb{R}^{n+1} . However, due to different rates of changes in the quadratic term and linear term in t over an interval of length $\mathcal{O}(1)$, these two terms do not properly cancel each other, which implies that the warmness of μ_{m-1} with respect to $\mu_m = \pi$ is no longer $\mathcal{O}(1)$ -bounded.

We address this by introducing $\mu_{\sigma^2, \rho}(x, t) \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t) \cdot \mathbf{1}[(x, t) \in \bar{\mathcal{K}}]$ where $\sigma^2 \in (0, R^2]$, $\rho \in (0, n]$, and $\bar{\mathcal{K}} = \{V(x) \leq nt\} \cap \{\|x\| = \mathcal{O}(R), \|t\| = \mathcal{O}(1)\}$. Essentially, this runs **Gaussian Cooling** along the x -direction with an exponential tilt in the t -direction, ensuring that consecutive distributions are $\mathcal{O}(1)$ -close in \mathcal{R}_∞ . However, we need an efficient sampler for $\mu_{\sigma^2, \rho}$ with \mathcal{R}_∞ -guarantees to maintain \mathcal{R}_∞ -warmness across the annealing scheme.

Sampling from annealing distributions. Given the form of the annealing distribution (the potential is a combination of linear and quadratic terms), we use the **Proximal Sampler** to develop PS_{ann} (Algorithm 3). The query complexity for rejection sampling in the second step now can be derived from our analysis of PS_{exp} (Lemma 2.14) and the **Proximal Sampler** for truncated Gaussians [KZ25]. For mixing with \mathcal{R}_∞ guarantees, we apply a technique from [KZ25]. We obtain a mixing rate of $\mathcal{O}(h^{-1} C_{\text{LSI}}(\mu_{\sigma^2, \rho}))$ for the **Proximal Sampler** based on (LSI) (Proposition 2.7). This results in only a *doubly logarithmic* dependence on the initial warmness, implying convergence from any *feasible* start with an overhead of $\text{polylog}(n, R)$ (Lemma 3.5). This *uniform ergodicity* implies L^∞ -norm contraction of density toward the target [DMLM03] (see Lemma 3.4), leading to a \mathcal{R}_∞ -guarantee of PS_{ann} without significant overhead (Theorem 3.16).

In Lemma 3.3, we bound the LSI constant of the annealing distribution $\mu_{\sigma^2, \rho}$ by $\sigma^2 \vee 1$ via the Bakry-Émery criterion (Lemma A.1) and Holley–Stroock perturbation principle (Lemma A.2). For $\nu \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \frac{t^2}{2} - \rho t)|_{\bar{\mathcal{K}}}$, since the potential of ν is $\min(\sigma^{-2}, 1)$ -strongly convex, its LSI

without convex truncation is bounded by $\sigma^2 \vee 1$ through Bakry-Émery, and convex truncation to $\bar{\mathcal{K}}$ only helps in satisfying the criterion [BGL14]. Also, as $\sup t - \inf t = \Theta(1)$ over $\bar{\mathcal{K}}$, the ratio of $\mu_{\sigma^2, \rho}$ to ν is bounded below and above by $\Theta(1)$, so the perturbation principle ensures that $C_{\text{LSI}}(\mu_{\sigma^2, \rho}) \lesssim C_{\text{LSI}}(\nu) \leq \sigma^2 \vee 1$.

Gaussian cooling with exponential tilt. With the query complexity of PS_{ann} in mind, we design Tilted Gaussian Cooling (Algorithm 2) for warm-start generation for $\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}$. We run rejection sampling with proposal $\mathcal{N}(0, n^{-1}I_n) \otimes \text{Unif}(I_t)$ for some interval I_t of length $\Theta(1)$, and initial distribution $\exp(-\frac{n}{2}\|x\|^2)|_{\bar{\mathcal{K}}}$, which is $\mathcal{O}(1)$ -warm with respect to $\mu_1 = \exp(-\frac{n}{2}\|x\|^2 - t)|_{\bar{\mathcal{K}}}$. In Phase I, we update the two parameters according to $\sigma^2 \leftarrow \sigma^2(1 + n^{-1})$ and $\rho \leftarrow \rho(1 + n^{-1})$ while $\rho \leq n$ and $\sigma^2 \leq 1$. Since Phase I involves $\tilde{\mathcal{O}}(n)$ inner phases with complexities of sampling from each annealing distributions being $\tilde{\mathcal{O}}(n^2(\sigma^2 \vee 1)) = \tilde{\mathcal{O}}(n^2)$, the total complexity is $\tilde{\mathcal{O}}(n^3)$. In Phase II, we accelerate σ^2 -updates via $\sigma^2 \leftarrow \sigma^2(1 + \sigma^2/R^2)$ as in Gaussian Cooling. With $\tilde{\mathcal{O}}(R^2/\sigma^2)$ inner phases (for doubling of σ^2) and sampling complexity $\tilde{\mathcal{O}}(n^2\sigma^2)$ per inner phase, this has total complexity $\tilde{\mathcal{O}}(n^2R^2)$. At termination, PS_{exp} is run with $\mu_{R^2, n}$ as the initial distribution for target $\bar{\pi} \propto e^{-nt}|_{\bar{\mathcal{K}}}$, where these two are close in \mathcal{R}_{∞} . Using the LSI of $\bar{\pi}$ and the boosting scheme again, we can achieve an ε -close sample to π (not $\bar{\pi}$) in the \mathcal{R}_{∞} -divergence, using $\tilde{\mathcal{O}}(n^2(R^2 \vee n) \text{polylog} \frac{1}{\varepsilon})$ queries in total (Lemma 3.16).

1.2.4 Rounding

Rounding is the key to reducing the dependence on R from poly R to polylog R .

Isotropic rounding for uniform distributions. The previous best rounding algorithm for uniform distributions, proposed by [JLLV21], gradually isotropizes a sequence $\{\mu_i\}$ of distributions. For a convex body \mathcal{K} , they set $\mu_i = \text{Unif}(\mathcal{K} \cap B_{\delta^i}(0))$ for $\delta = 1 + n^{-1/2}$, increasing i while $\delta^i \leq R$. Their approach entails two important tasks: Outer loop: if $F(\mathcal{K} \cap B_r)$ is near-isotropic for an affine map F , then show that $F(\mathcal{K} \cap B_{\delta r})$ is well-rounded, and Inner loop: design an algorithm of n^3 -complexity that isotropizes a given well-rounded uniform distribution. The first task was accomplished by [JLLV24] through Paouris' lemma (i.e., exponential tail decay) and a universal property that the diameter of an isotropic convex body is bounded by $n + 1$.

The second task was addressed by repeating [draw samples \rightarrow compute crude covariance estimation \rightarrow upscale skewed directions of the covariance estimation]. They first run Gaussian Cooling to obtain a warm start for a uniform distribution μ from a convex body. Then, when the inner radius is r , the Ball Walk (or In-and-Out) is used to generate r^2 samples approximately distributed according to μ . These r^2 samples give a rough estimate of the covariance matrix $\bar{\Sigma}$ of μ such that $|\bar{\Sigma} - \Sigma| \lesssim nI_n$, where $\Sigma = \text{Cov} \mu$. Since the query complexity of these uniform samplers is $n^2\|\Sigma\|/r^2$, this procedure uses $n^2\|\Sigma\|$ queries in total. Then, it computes the eigenvalue/vectors of $\bar{\Sigma}$ and scales up (by a factor of 2) the subspace spanned by eigenvectors with eigenvalues less than n . One iteration of this process achieves two key properties: the largest eigenvalue of the covariance Σ increases by at most n additively while $r = \text{inrad} \mu$ almost doubles. Since the well-roundedness ensures that $\|\Sigma\| = \mathcal{O}(n)$ initially, the complexity of one iteration remains as $\tilde{\mathcal{O}}(n^3)$ throughout. Since there are $\sqrt{n} \log R$ outer iterations, the algorithm uses a total of $\tilde{\mathcal{O}}(n^{3.5} \text{polylog} R)$ queries.

Extension to general logconcave distributions. Our rounding algorithm essentially follows this approach, with several technical refinements. First of all, we define a *ground set* of a general logconcave distribution, namely the level set $\mathbf{L}_g := \{x : V(x) - \min V \leq 10n\}$. This ground set takes up most of measure due to the universal property in terms of the potential value (Lemma A.6). Focusing on the ground set is the first step toward a streamlined extension of the previous approach,

so we consider the *grounded distribution* $\nu^X := \pi^X|_{\mathbb{L}_g}$. Then, for $\nu_r^X := \nu^X|_{B_r(0)}$, we isotropize a sequence of distributions, given by $\nu_1^X \rightarrow \nu_\delta^X \rightarrow \nu_{\delta^2}^X \rightarrow \dots \rightarrow \nu_D^X \rightarrow \nu^X \rightarrow \pi^X$ for $D = \Theta(R)$.

To analyze the outer loop, we show that for an affine map F^X between \mathbb{R}^n , if $F_{\#}^X \nu_r^X$ is near-isotropic, then $(4F^X)_{\#} \nu_{\delta r}^X$ is well-rounded. Unfortunately, a varying density of $F_{\#}^X \nu_{\delta r}^X$ poses a daunting challenge in extending the previous proof in [JLLV24] to general logconcave distributions. Nonetheless, we can resolve this issue by working again with the exponential reduction (Lemma 4.12). This extension asks for a universal property that an isotropic grounded distribution has diameter of order $\mathcal{O}(n)$, similar to isotropic uniform distributions. We show this in Lemma 4.13. Transferring roundedness in the steps $\nu_D^X \rightarrow \nu^X$ and $\nu^X \rightarrow \pi^X$ is relatively straightforward by combining the change of measures and the reverse Hölder inequality for logconcave distributions.

For the inner loop, we can still apply the algorithm from [JLLV21] (or its streamlined version in [KZ24]) with only minor changes to constants. First, with the logconcave sampler PS_{exp} whose mixing rate depends on $\|\text{Cov } \pi^X\|$ rather than $\text{tr}(\text{Cov } \pi^X)$, the complexity analysis of the inner loop extends naturally to general logconcave distributions. Next, we note that the proofs for controlling $\|\Sigma_i\|$ and $\text{tr } \Sigma_i$ are identical to those for uniform distributions. The proof for the doubling of the inner radius at each iteration is nearly the same as in the uniform case, since the existence of a large ball due to isotropy is also a *universal* property of logconcave distributions (see Lemma A.10).

1.2.5 Integration

For integration, we use the stronger guarantees of our logconcave sampler, along with a version of the Tilted Gaussian Cooling scheme to obtain a cubic algorithm for *well-rounded* logconcave functions. For general logconcave functions, we use the rounding algorithm as a pre-processing step, then apply the integration algorithm to the near-isotropic distribution obtained after rounding.

Volume computation through annealing. Similar to sampling, prior volume algorithms also follow a sequence $\{f_i\}_{i \in [m]}$ of logconcave functions, moving across distributions $\mu_i \propto f_i$ using a logconcave sampler. The annealing scheme is designed in a way that f_1 is easy to integrate, μ_i is an $\mathcal{O}(1)$ -warm start for μ_{i+1} , the variance of the estimator $E_i = f_{i+1}/f_i$ with respect to μ_i is bounded by $m^{-1} \mathbb{E}_{\mu_i}[E_i^2]$ (i.e., $\chi^2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(m^{-1})$), and $f_{m+1} = f$ is the target logconcave function. Since $\mathbb{E}_{\mu_i} E_i = \int f_{i+1} / \int f_i$, accurate estimations of all E_i guarantee that the product $\int f_1 \times E_1 \cdots E_m$ is a good estimator of $\int f_1 \cdot \mathbb{E} E_1 \cdots \mathbb{E} E_m = \int f$.

The best-known algorithm, Gaussian Cooling [CV18], uses unnormalized Gaussian densities $f_i(x) = \exp(-\|x\|^2/2\sigma_i^2) \cdot \mathbb{1}_{\mathcal{K}}(x)$ for a convex body \mathcal{K} , along with an update schedule for σ_i^2 . As described earlier in §1.2.3, it uses the Ball Walk to sample from $\mu_i \propto f_i$ with TV-guarantees.

Extension to general logconcave functions. Our integration algorithm follows this approach, but once again in the lifted space. With PS_{ann} used for sampling, we follow a modified version of Tilted Gaussian Cooling for ease of analysis, particularly for variance control. We use $f_i(x, t) = \exp(-\|x\|^2/2\sigma_i^2 - \rho_i t)|_{\mathcal{K}}$ as the intermediate annealing functions, and ensure that $\mathcal{R}_{\infty}(\mu_i \parallel \mu_{i+1}) = \mathcal{O}(1)$ for efficient sampling from μ_{i+1} , and $\mathcal{R}_2(\mu_{i+1} \parallel \mu_i) = \exp(\mathcal{O}(m^{-1}))$ for $\text{var}_{\mu_i}(E_i/\mathbb{E}_{\mu_i} E_i) = \mathcal{O}(m^{-1})$. Since $\mathcal{O}(1)$ -warmness can be shown as in Tilted Gaussian Cooling, we elaborate on technical tools for variance control along with design of the algorithm. In Phase I, we go from $\exp(-n\|x\|^2/2)|_{\mathcal{K}}$ to $\exp(-\|x\|^2/2)|_{\mathcal{K}}$ with the update $\sigma^2 \leftarrow \sigma^2(1 + 1/n)$, where variance control is achieved by the logconcavity of $a \mapsto a^n \int h^a$ for a logconcave function h [KV06]. In Phase II, we move from $\exp(-\|x\|^2/2 - t)|_{\mathcal{K}}$ to $\exp(-\|x\|^2/2 - nt)|_{\mathcal{K}}$ with the update $\rho \leftarrow \rho(1 + 1/n)$, where variance control follows from the previous lemma and another lemma in [CV18]. Lastly in Phase III, we move from $\exp(-\|x\|^2/2 - nt)|_{\mathcal{K}}$ to $\exp(-nt)|_{\mathcal{K}}$ with the update $\sigma^2 \leftarrow \sigma^2(1 + \sigma^2/n)$, and we use the lemma in [CV18] again for variance control.

Streamlined statistical analysis. If all samples used are *independent*, and the estimators have moderate variance, namely $\text{var}_{\mu_i}(E_i/\mathbb{E}E_i) = \mathcal{O}(m^{-1})$, then $\text{var}(E_1 \cdots E_m) = (\prod(1 + m^{-1}) - 1) (\mathbb{E}E_i)^2 \approx \mathcal{O}(1) (\mathbb{E}[E_1 \cdots E_m])^2$, which implies concentration of the estimator around $\mathbb{E}[E_1 \cdots E_m] = \mathbb{E}E_1 \cdots \mathbb{E}E_m$ through Chebyshev’s inequality. However, samples given by (say) the Ball Walk in Gaussian Cooling are *approximately* distributed according to μ_i . In fact, samples drawn from μ_i and μ_{i+1} are *dependent*, since a sample from μ_i is used as a warm-start for μ_{i+1} . This means that the estimator $E_1 \cdots E_m$ is a *biased estimator* of $\mathbb{E}E_1 \cdots \mathbb{E}E_m (\neq \mathbb{E}[E_1 \cdots E_m])$. These two issues complicate statistical analysis in prior work on how close the estimator is to the integral of f , and require additional technical tools to address them. For the first issue, previous work used a coupling argument based on TV-distance (referred to as “divine intervention”) to account for the effects of approximate distributions (rather than exactly μ_i). For the second, they used the notion of α -mixing [Ros56] (referred to as “ μ -independence” therein) to bound the bias of the product estimator.

We simplify this statistical analysis substantially by using stronger guarantees of our sampler PS_{ann} . For the first issue, when $\bar{\mu}_i$ denotes the actual law of a sample X_i satisfying $\mathcal{R}_{\infty}(\bar{\mu}_i \parallel \mu_i) \leq \varepsilon$, we can notice that the probability of a bad event (any event) with respect to $\bar{\mu}_i$ (instead of μ_i) only increases by at most a multiplicative factor of $1 + \varepsilon$. Since the mixing time of PS_{ann} has a polylogarithmic dependence on $1/\varepsilon$, we can set ε polynomially small (i.e., $\varepsilon \leftarrow \varepsilon/\text{poly}(n, R)$), so we can enforce $\mathcal{R}_{\infty}(\otimes_i \bar{\mu}_i \parallel \otimes_i \mu_i) \leq \varepsilon$ without a huge overhead in the query complexity. For the second issue, we use notion of β -mixing (or the coefficient of absolute regularity) [KR60], which is stronger than α -mixing. This quantity basically measures the discrepancy between a joint distribution and the product of marginal distributions by $\|\text{law}(X_i, X_{i+1}) - \text{law } X_i \otimes \text{law } X_{i+1}\|_{\text{TV}} = \|\text{law}(X_i, X_{i+1}) - \bar{\mu}_i \otimes \bar{\mu}_{i+1}\|_{\text{TV}} \leq \beta$. Since the mixing of PS_{ann} via (LSI) ensures mixing from *any start* (Lemma 3.5), we can easily bound β by $\mathcal{O}(\varepsilon)$ (see (5.2) for details). Thus, when analyzing the probability of a bad event, we can replace $\otimes_i \bar{\mu}_i$ with $\text{law}(X_1, \dots, X_m)$ at the additive cost of $\mathcal{O}(m\varepsilon)$ in probability (we can replace $\varepsilon \leftarrow m\varepsilon$ once again).

1.3 Preliminaries

Let $\mathcal{P}(\mathbb{R}^n)$ be the family of probability measures (distributions) on \mathbb{R}^n that are absolutely continuous with respect to the Lebesgue measure. We use the same symbol for a distribution and density. For a set S and its indicator function $\mathbb{1}_S(x) = [x \in S]$, we use $\mu|_S$ to denote a distribution μ truncated to S (i.e., $\mu|_S \propto \mu \cdot \mathbb{1}_S$). For a measurable map $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $\mu \in \mathcal{P}(\mathbb{R}^n)$, the *pushforward* measure $F_{\#}\mu$ is defined as $F_{\#}\mu(A) = (\mu \circ F^{-1})(A)$ for a measurable set A . For $a, b \in \mathbb{R}$, we use $a \vee b$ and $a \wedge b$ to indicate their maximum and minimum, respectively. We use $B_r^n(x_0)$ to denote the n -dimensional ball of radius $r > 0$ centered at $x_0 \in \mathbb{R}^n$, dropping the superscript n if there is no confusion. We use $a = \tilde{\mathcal{O}}(b)$ to denote $a = \mathcal{O}(b \text{ polylog } b)$. For two symmetric matrices A, B , we use $|A| \preceq B$ to denote $-B \preceq A \preceq B$. Unless specified otherwise, for a vector v and a PSD matrix M , $\|v\|$ and $\|\Sigma\|$ refer to the ℓ_2 -norm of v and the operator norm of Σ , respectively. We use \mathbb{R} to denote the extended real number system $\mathbb{R} \cup \{\pm\infty\}$.

We recall notions of common probability divergences/distances between distributions.

Definition 1.9. For $\mu, \nu \in \mathcal{P}(\mathbb{R}^n)$, the *f-divergence* of μ towards ν with $\mu \ll \nu$ is defined as, for a convex function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ with $f(1) = 0$ and $f'(\infty) = \infty$,

$$D_f(\mu \parallel \nu) := \int f\left(\frac{d\mu}{d\nu}\right) d\nu.$$

For $q \in (1, \infty)$, the *KL-divergence* and χ^q -divergence correspond to $f(x) = x \log x$ and $x^q - 1$, respectively. The *q-Rényi divergence* is defined as

$$\mathcal{R}_q(\mu \parallel \nu) := \frac{1}{q-1} \log(\chi^q(\mu \parallel \nu) + 1).$$

The *Rényi-infinity divergence* is defined as

$$\mathcal{R}_\infty(\mu \parallel \nu) := \log \operatorname{ess\,sup}_\mu \frac{d\mu}{d\nu}.$$

A distribution μ is said to be *M-warm with respect to a distribution ν* if $\frac{\mu(S)}{\nu(S)} \leq M$ for any measurable subset S (i.e., μ is $\exp(\mathcal{R}_\infty(\mu \parallel \nu))$ -warm with respect to ν). The *total variation* (TV) distance for $\mu, \nu \in \mathcal{P}(\mathbb{R}^n)$ is defined by

$$\|\mu - \nu\|_{\text{TV}} := \frac{1}{2} \int |\mu(x) - \nu(x)| dx = \sup_{S \in \mathcal{F}} |\mu(S) - \nu(S)|,$$

where \mathcal{F} is the collection of all measurable subsets of \mathbb{R}^n .

We recall $\text{KL} = \lim_{q \downarrow 1} \mathcal{R}_q \leq \mathcal{R}_q \leq \mathcal{R}_{q'} \leq \mathcal{R}_\infty$ for $1 \leq q \leq q'$ and $2 \|\cdot\|_{\text{TV}}^2 \leq \text{KL} \leq \mathcal{R}_2 = \log(\chi^2 + 1) \leq \chi^2$. We refer readers to [VEH14] for basic properties of the Rényi-divergence (e.g., continuity/monotonicity in q). The data processing inequality is given in Lemma A.3 for reference.

2 Sampling from logconcave distributions

In this section, we study the complexity of sampling from a logconcave distribution under the setup stated in the introduction. We remark that the “roundness” (or regularity) of the distribution is specified through the existence of a ball in a level set of **constant measure**. We find it more convenient to work with the **values of the potential** (see Lemma 2.6). More precisely, we will focus on a particular level set, called the *ground set* of π , defined as

$$\mathsf{L}_{\pi, g} := \{x \in \mathbb{R}^n : V(x) - \min V \leq 10n\}. \quad (2.1)$$

Note that the ground set of π is *independent* of the normalization constant of π , which justifies using π in the subscript to indicate its uniqueness. We use $\mathsf{L}_{V, g}$ to reveal a specific choice of the potential V . We also define the notation: $V_0 := \min V$, and a level set $\mathsf{L}_{\pi, l} := \{x \in \mathbb{R}^n : V(x) - \min V \leq nl\}$. Hereafter, in $\text{Eval}_{\mathcal{P}}(V)$, the requirement of inclusion of $B_1(x_0)$ in the level set of π of measure $1/8$ is replaced by $B_1(x_0) \subset \mathsf{L}_{\pi, X, g}$. As we will see in Lemma 2.6, this is a weaker condition.

Problem 2.1 (Zeroth-order logconcave sampling from a warm start). Assume access to a well-defined function oracle $\text{Eval}(V)$ for convex $V : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$. Given target accuracy $\varepsilon > 0$ and M -warm initial distribution for π^X , how many evaluation queries do we need to obtain a sample X_* such that

$$D(\text{law } X_*, \pi^X) \leq \varepsilon \quad \text{for } D = \text{TV}, \text{KL}, \mathcal{R}_q, \text{ etc.}$$

Note that this *does not* require the knowledge of x_0, R , and $\min V$. Also, if the inner radius is r , then we can simply rescale the whole system by $x \mapsto r^{-1}x$.

We prove **Result 1** in this section:

Theorem 2.2 (Restatement of Theorem 1.5). *In the setting of Problem 2.1, there exists an algorithm that for given $\varepsilon, \eta \in (0, 1)$, $q \in [2, \infty)$, and M -warm initial distribution for π^X , with probability at least $1 - \eta$ returns a sample X_* satisfying $\mathcal{R}_q(\text{law } X_* \parallel \pi^X) \leq \varepsilon$, using $\tilde{\mathcal{O}}(qMn^2(\|\text{Cov } \pi^X\| \vee 1) \log^4 1/\eta\varepsilon)$ evaluation queries in expectation.*

See Theorem 2.15 for details. When $\|\text{Cov } \pi^X\| = \Omega(1)$ (the primary regime for algorithmic applications later), this complexity bound generalizes the well-known complexity bound of $\tilde{\mathcal{O}}(qMn^2 \|\text{Cov } \pi^X\| \text{polylog } 1/\eta\varepsilon)$ for uniform sampling from a convex body [KVZ24]. Also, if the inner radius is r , then the query complexity includes an additional multiplicative factor of r^{-2} .

2.1 Reduction to an exponential distribution

The logconcave sampling problem can be reduced to the following exponential sampling problem:

$$d\pi(x, t) \propto \exp(-nt) dx dt \quad \text{subject to } \mathcal{K} := \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : V(x) \leq nt\}. \quad (\text{exp-red})$$

We note that convexity of the constraint \mathcal{K} follows from that of V , and that the X -marginal of π exactly matches the desired target $\pi^X \propto \exp(-V)$.

Proposition 2.3. *The X -marginal π^X of π is proportional to $\exp(-V)$.*

Proof. We just integrate out t as follows:

$$\pi^X(x) = \int \pi(x, t) dt = \frac{\int \exp(-nt)|_{[V(x) \leq nt]} dt}{\int \exp(-nt)|_{[V(x) \leq nt]} dt dx} = \frac{\exp(-V(x))}{\int \exp(-V(x)) dx},$$

where the last equality follows from

$$\int \exp(-nt)|_{[V(x) \leq nt]} dt = \int_{V(x)/n}^{\infty} \exp(-nt) dt = \frac{1}{n} \int_{V(x)}^{\infty} \exp(-u) du = \frac{1}{n} \exp(-V(x)),$$

which completes the proof. \square

This implies that in order to sample from $\exp(-V)$, one can just focus on sampling $(x, t) \sim \pi$ and take x -part only. Precisely, once we obtain a sample $(X, T) \in \mathbb{R}^{n+1}$ such that $D(\text{law}(X, T) \parallel \pi) \leq \varepsilon$ for the f -divergence or Rényi divergence D , the data-processing inequality (DPI) (Lemma A.3) ensures that $D(\text{law } X \parallel \pi^X) \leq \varepsilon$. Therefore, the general logconcave sampling problem can be reduced to sampling from a particular form of the exponential distribution above. While it allows us to move to the simpler problem, for its mixing analysis, we will need to know the mean and variance along the new dimension as well as the operator norm of the covariance matrix of π .

Mean. For the mean $(\mu_x, \mu_t) := \mathbb{E}_\pi[(X, T)] \in \mathbb{R}^n \times \mathbb{R}$, we establish a useful bound on $\mu_t = \mathbb{E}_\pi T$.

Lemma 2.4. *In the setting of Problem 2.1, it holds that $0 \leq \mu_t - V_0/n \leq \sqrt{10}$.*

Proof. We directly compute $\mu_t = \mathbb{E}_{\pi T} T = \mathbb{E}_\pi T$ as follows:

$$\begin{aligned} \mu_t = \mathbb{E}_\pi T &= \frac{\int t e^{-nt}|_{[V(x) \leq nt]} dt dx}{\int e^{-nt}|_{[V(x) \leq nt]} dt dx} = \frac{\int \int_{V(x)/n}^{\infty} t e^{-nt} dt dx}{n^{-1} \int \exp(-V) dx} = \frac{\int \int_{V(x)}^{\infty} u e^{-u} du dx}{n \int \exp(-V) dx} \\ &= \frac{\int ([-u e^{-u}]_{V(x)}^{\infty} + \int_{V(x)}^{\infty} e^{-u} du) dx}{n \int \exp(-V) dx} = \frac{\int (V + 1) \exp(-V) dx}{n \int \exp(-V) dx} \\ &= n^{-1} \mathbb{E}_{\pi^X} (V + 1) = n^{-1} \mathbb{E}_\pi (V + 1). \end{aligned}$$

We now prove $\mathbb{E}_\pi (V - V_0) \leq \sqrt{10}n$. For $\pi^X = \exp(-W) / \int \exp(-W)$ with $W := V - V_0 \geq 0$, Lemma A.6 ensures that for $\beta \geq 2$,

$$\mathbb{P}_\pi(W(X) \geq \beta(n-1)) \leq (e^{1-\beta} \beta)^{n-1}.$$

Therefore, for $n \geq 2$ and the Gamma function Γ ,

$$\mathbb{E}_\pi W = \int_0^\infty \mathbb{P}_\pi(W(X) \geq u) du$$

$$\begin{aligned}
&= \int_0^{2^{(n-1)}} \mathbb{P}_\pi(W(X) \geq u) \, du + \int_{2^{(n-1)}}^\infty \mathbb{P}_\pi(W(X) \geq u) \, du \\
&\leq 2(n-1) + (n-1) \int_2^\infty \mathbb{P}_\pi(W(X) \geq \beta(n-1)) \, d\beta \\
&\leq 2(n-1) + (n-1) \int_0^\infty (e^{1-\beta}\beta)^{n-1} \, d\beta \\
&\leq n \left(2 \frac{n-1}{n} + \left(\frac{e}{n-1} \right)^{n-1} \frac{\Gamma(n)}{n} \right).
\end{aligned}$$

When $n = 1$, by using $\mathbb{P}_\pi(W(X) \geq t) \leq e^{-t}$ by [LV07, Lemma 5.6(a)], we also have $\mathbb{E}_\pi W \leq 1$. One can manually check that the final bound is at most $\sqrt{10}n$ for $n \leq 7$. When $n \geq 8$, since the digamma function $\psi(x) := \log \Gamma(x)$ satisfies $\psi(x) \leq \log x$, the second term in the parenthesis is decreasing in n , and thus $\mathbb{E}_\pi W \leq \sqrt{10}n$ as well. Substituting this bound back to the expression of μ_t , it follows that

$$\frac{V_0}{n} \leq \mu_t \leq \frac{V_0}{n} + \sqrt{10},$$

which completes the proof. \square

Covariance. Let us denote the covariance of π, π^X, π^T by $\Sigma := \mathbb{E}_\pi[((X, T) - (\mu_x, \mu_t))^{\otimes 2}]$, $\Sigma_x := \mathbb{E}_{\pi^X}[(X - \mu_x)^{\otimes 2}]$, $\Sigma_t := \mathbb{E}_{\pi^T}[(T - \mu_t)^2]$, respectively. We study two important quantities here — the largest eigenvalue (i.e., operator norm of Σ) and trace (i.e., second moment of π).

Lemma 2.5. *The variance in the t -direction is at most 160. Moreover, $\text{tr } \Sigma \leq \text{tr } \Sigma_x + 160$, and $\|\Sigma\| \leq 2(\|\Sigma_x\| + 160)$.*

Proof. Let us bound $\mathbb{E}_\pi[|T - \mu_t|^2] = \mathbb{E}_{\pi^T}[|T - \mu_t|^2]$. By the reverse Hölder (Lemma A.9),

$$\mathbb{E}_\pi[|T - \mu_t|^2] \leq \mathbb{E}_\pi[|T - V_0/n|^2] \leq 16 (\mathbb{E}_\pi|T - V_0/n|)^2 \leq 160.$$

Hence,

$$\text{tr } \Sigma = \mathbb{E}_\pi[\|X - \mu_x\|^2 + |T - \mu_t|^2] = \mathbb{E}_{\pi^X}[\|X - \mu_x\|^2] + \mathbb{E}_{\pi^T}[|T - \mu_t|^2] \leq R^2 + 160.$$

As for the largest eigenvalue, it follows that for a unit vector $v = (v_x, v_t) \in \mathbb{R}^n \times \mathbb{R}$,

$$\mathbb{E}_\pi[(v_x \cdot (X - \mu_x) + v_t (T - \mu_t))^2] \leq 2 \mathbb{E}_\pi[(v_x \cdot (X - \mu_x))^2] + 2 \mathbb{E}_\pi[(v_t (T - \mu_t))^2] \leq 2(\|\Sigma_x\| + 160).$$

Hence, $\|\Sigma\| \leq 2(\|\Sigma_x\| + 160)$. \square

We now show that the ‘measure-based’ regularity implies the ‘value-based’ one.

Lemma 2.6. *When a level set of $\pi^X \propto \exp(-V)$ of measure $\mathfrak{m} < 1$ contains $B_r(x_0)$ for $r > 0$ and $x_0 \in \mathbb{R}^n$, the level set $\mathsf{L}_{V,l} = \{V(x) - V_0 \leq \ell n\}$ with $\ell = 5 + 13 \log \frac{e}{1-\mathfrak{m}}$ contains $B_r(x_0)$.*

Proof. Consider the exponential reduction $\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}$ for $\mathcal{K} = \{(x, t) : V(x) \leq nt\}$. Since the variance in the t -direction is at most 160, Lemma A.7 leads to

$$\mathbb{P}_\pi(|T - \mu_t| \geq 13c) = \mathbb{P}_{\pi^T}(|T - \mu_t| \geq 13c) \leq \exp(-c + 1) \quad \text{for any } c > 0,$$

which implies $\mathbb{P}_\pi(T \geq 5 + V_0/n + 13c) \leq \mathbb{P}_{\pi^T}(T \geq \mu_t + 13c) \leq \exp(-c + 1)$. Hence, for $c = -\log \frac{e}{1-\mathfrak{m}}$, we have $\mathbb{P}_\pi(T - V_0/n \leq \ell) \geq \mathfrak{m}$. Since $\mathcal{K} \cap [t \leq V_0/n + \ell] = \mathcal{K} \cap (\mathsf{L}_{V,l} \times \mathbb{R})$, it follows from $\mathbb{P}_\pi(T \leq V_0/n + \ell) = \pi^X(\mathsf{L}_{V,l})$ that $\mathsf{L}_{V,l}$ contains $B_r(x_0)$. \square

2.2 Proximal sampler for the reduced exponential distribution

We now propose a sampling algorithm for this exponential distribution. For $z := (x, t) \in \mathbb{R}^n \times \mathbb{R}$ and parameter $h > 0$, we use the Proximal Sampler (PS) to sample from

$$\pi^{Z,Y}(z, y) \propto \exp(-\alpha^\top z - \frac{1}{2h} \|z - y\|^2)|_{\mathcal{K}} \quad \text{for } \alpha = ne_{n+1},$$

where its Z -marginal π^Z corresponds to our desired target $\pi^Z \propto \exp(-\alpha^\top z)|_{\mathcal{K}}$.

The Proximal Sampler for this target, called PS_{exp} (Algorithm 1), repeats the following two steps:

- [Forward] $y \sim \pi^{Y|Z=z} = \mathcal{N}(z, hI_{n+1})$.
- [Backward] $z \sim \pi^{Z|Y=y} \propto \exp(-\alpha^\top z - \frac{1}{2h} \|y - z\|^2)|_{\mathcal{K}} = \mathcal{N}(y - h\alpha, hI_{n+1})|_{\mathcal{K}}$ for $\alpha = ne_{n+1}$,

and the backward step is implemented by rejection sampling with proposal $\mathcal{N}(y - h\alpha, hI_{n+1})$.

As in previous work, the analysis can be divided into two parts: (1) mixing towards the target (§2.2.1), and (2) the query complexity of implementing the backward step (§2.2.2). The first part can be analyzed similarly to prior work on uniform sampling [KVZ24] or constrained Gaussian sampling [KZ25], while the second part introduces additional technical challenges.

2.2.1 Convergence rate

One iteration of the Proximal Sampler (one forward step + one backward step) for $\pi^X \propto \exp(-V)$ can be viewed as a composition of forward and backward heat flow, where it can be written as follows: for $\pi(x, y) \propto \exp(-V(x) - \frac{1}{2h} \|x - y\|^2)$, repeat (i) $y_{i+1} \sim \pi^{Y|X=x_i} = \mathcal{N}(x_i, hI_n)$ and (ii) $x_{i+1} \sim \pi^{X|Y=y_{i+1}}$. To make this perspective concrete, we borrow expositions from [KVZ24, p.4-5] and [Che24, §8.3]. For the laws of x_i and y_i denoted by μ_i^X and μ_i^Y , the forward step can be described as $\mu_{i+1}^Y = \int \pi^{Y|X=x} \mu_i^X(dx) = \mu_i^X * \mathcal{N}(0, hI_n)$. This step can be seen as simulating a Brownian motion with initial measure μ_i^X for time h :

$$dZ_t = dB_t \quad \text{with } Z_0 \sim \mu_i^X \implies Z_h \sim \mu_{i+1}^Y.$$

Let us denote $\mu P_t := \mu * \mathcal{N}(0, tI_n)$ for $\mu \in \mathcal{P}(\mathbb{R}^n)$ (i.e., $(P_t)_{t \geq 0}$ is the heat semigroup). Next, the backward step corresponds to $\mu_{i+1}^X = \int \pi^{X|Y=y} \mu_{i+1}^Y(dy)$, and it turns out that this process can also be represented by an SDE (called the backward heat flow): for a new Brownian motion B_t ,

$$dZ_t^\leftarrow = \nabla \log(\pi^X P_{h-t})(Z_t^\leftarrow) dt + dB_t \quad \text{for } t \in [0, h].$$

Its construction ensures that if $Z_0^\leftarrow \sim \delta_y$ (a point mass at y), then $Z_h^\leftarrow \sim \pi^{X|Y=y}$. Hence, if we initialize this SDE with $Z_0^\leftarrow \sim \mu_{i+1}^Y$, then the law of Z_h^\leftarrow corresponds to $\int \pi^{X|Y=y} \mu_{i+1}^Y(dy) = \mu_{i+1}^X$.

This perspective establishes a natural connection between mixing guarantees and constants of functional inequalities (e.g., (PI) and (LSI)) for a target distribution. [CCSW22] has already demonstrated this connection for the Proximal Sampler in the case of smooth, unconstrained distributions.

Proposition 2.7 ([CCSW22, Theorem 3 and 4]). *Assume that $\pi \in \mathcal{P}(\mathbb{R}^n)$ has a smooth density, and let P denote the Markov kernel of the Proximal Sampler. If π satisfies (LSI) with constant C_{LSI} , then for any $q \geq 1$ and distribution μ with smooth density and $\mu \ll \pi$,*

$$\mathcal{R}_q(\mu P \parallel \pi) \leq \frac{\mathcal{R}_q(\mu \parallel \pi)}{(1 + h/C_{\text{LSI}})^{2/q}}. \quad (2.2)$$

If π satisfies (PI) with constant C_{PI} , then

$$\chi^2(\mu P \parallel \pi) \leq \frac{\chi^2(\mu \parallel \pi)}{(1 + h/C_{\text{PI}})^2}. \quad (2.3)$$

Moreover, for all $q \geq 2$,

$$\mathcal{R}_q(\mu P \parallel \pi) \leq \begin{cases} \mathcal{R}_q(\mu \parallel \pi) - \frac{2 \log(1+h/C_{\text{PI}})}{q} & \text{if } \mathcal{R}_q(\mu \parallel \pi) \geq 1, \\ \frac{\mathcal{R}_q(\mu \parallel \pi)}{(1+h/C_{\text{PI}})^{2/q}} & \text{if } \mathcal{R}_q(\mu \parallel \pi) \leq 1. \end{cases} \quad (2.4)$$

Hence, its mixing times are roughly $qh^{-1}C_{\text{LSI}} \log \frac{\log \mathcal{R}_q}{\varepsilon}$ and $qh^{-1}C_{\text{PI}} \log \frac{\mathcal{R}_q}{\varepsilon}$ under (LSI) and (PI), respectively. We can extend this mixing result to constrained distributions under mild assumptions.

Lemma 2.8 ([KVZ24, Lemma 22]). *Let π be a probability measure, absolutely continuous with respect to the Lebesgue measure over \mathcal{K} . For $\{P_t\}_{t \geq 0}$ the heat semigroup, the forward and backward heat flow equations given by*

$$\begin{aligned} \partial_t \mu_t &= \frac{1}{2} \Delta \mu_t, \\ \partial_t \mu_t^\leftarrow &= -\operatorname{div}(\mu_t^\leftarrow \nabla \log(\pi P_{h-t})) + \frac{1}{2} \Delta \mu_t^\leftarrow \quad \text{with } \mu_0^\leftarrow = \mu_h, \end{aligned}$$

admit solutions on $(0, h]$, and the weak limit $\lim_{t \rightarrow h} \mu_t^\leftarrow = \mu_h^\leftarrow$ exists for any initial measure μ_0 with support³ contained in $\operatorname{supp} \pi$. Moreover, for any q -Rényi divergence with $q \in (1, \infty)$,

$$\mathcal{R}_q(\mu_h^\leftarrow \parallel \pi) \leq \lim_{t \downarrow 0} \mathcal{R}_q(\mu_{h-t}^\leftarrow \parallel \pi P_t).$$

Using this lemma, we now establish a mixing of the Proximal Sampler (for potentially constrained distributions) in χ^2 or \mathcal{R}_q for any $q \geq 2$ under (PI), and in \mathcal{R}_q for any $q \geq 1$ under (LSI).

Lemma 2.9. *Let P be the Markov kernel of the Proximal Sampler and μ be any initial distribution supported on $\mathcal{K} \subset \mathbb{R}^n$. Then, the contractions (2.2) under (LSI), and (2.3)-(2.4) under (PI) remain valid for $C_{\text{LSI}} = C_{\text{LSI}}(\pi^X)$ and $C_{\text{PI}} = C_{\text{PI}}(\pi^X)$.*

Proof. For any $\varepsilon > 0$, we have that $\mu_\varepsilon = \mu * \mathcal{N}(0, \varepsilon I_n)$ and $\pi_\varepsilon = \pi^X * \mathcal{N}(0, \varepsilon I_n)$ are C^∞ -smooth. Let D denote either χ^2 or \mathcal{R}_q , and define $C_\varepsilon := 1 + \frac{h-\varepsilon}{C_*(\pi^X)+\varepsilon} \leq 1 + \frac{h-\varepsilon}{C_*(\pi_\varepsilon)}$ for $* \in \{\text{PI}, \text{LSI}\}$, where the last inequality holds due to [Cha04, Corollary 13]:

$$C_{\text{PI}}(\pi^X * \mathcal{N}(0, \varepsilon I_n)) \leq C_{\text{PI}}(\pi^X) + \varepsilon \quad \& \quad C_{\text{LSI}}(\pi^X * \mathcal{N}(0, \varepsilon I_n)) \leq C_{\text{LSI}}(\pi^X) + \varepsilon.$$

Hence, by Proposition 2.7 with step size $h - \varepsilon$, we obtain that if $\mathcal{R}_q(\mu \parallel \pi^X) \leq 1$, for $\mu_0 = \mu$,

$$D(\mu_{h-\varepsilon}^\leftarrow \parallel \pi_\varepsilon) \leq C_\varepsilon^{-1} D(\mu_\varepsilon \parallel \pi_\varepsilon) \leq C_\varepsilon^{-1} D(\mu \parallel \pi^X),$$

where the last inequality follows from the DPI (Lemma A.3) for the f -divergence (χ^2) and Rényi divergence \mathcal{R}_q . By the lower semicontinuity of D (Lemma 2.8), sending $\varepsilon \rightarrow 0$ leads to

$$D(\mu P \parallel \pi^X) = D(\mu_h^\leftarrow \parallel \pi^X) \leq C_0^{-1} D(\mu \parallel \pi^X) = \frac{\chi^2(\mu \parallel \pi^X)}{(1 + h/C_{\text{PI}}(\pi^X))^2} \quad \text{or} \quad \frac{\mathcal{R}_q(\mu \parallel \pi^X)}{(1 + h/C_{\text{PI}}(\pi^X))^{2/q}}.$$

³The original result in [KVZ24] is proven only for a bounded support, but it can be extended to an unbounded support since the almost sure pointwise convergence of $\pi * \mathcal{N}(0, \varepsilon I_n)$ to π holds as $\varepsilon \rightarrow 0$ [Fol99, Theorem 8.15].

When $\mathcal{R}_q(\mu \parallel \pi^X) \geq 1$, a similar limiting argument shows that

$$\mathcal{R}_q(\mu P \parallel \pi^X) \leq \mathcal{R}_q(\mu_\epsilon \parallel \pi_\epsilon) - \frac{2 \log C_\epsilon}{q} \leq \mathcal{R}_q(\mu \parallel \pi^X) - \frac{2 \log C_0}{q},$$

where in the first inequality we used the lower-semicontinuity of \mathcal{R}_q , and in the second inequality used the DPI and then sent $\epsilon \rightarrow 0$. Contraction in \mathcal{R}_q for any $q \geq 1$ under (LSI) follows from the same argument above. \square

We can simply set $X \leftarrow Z = (X, T)$ and $Y \leftarrow Y$ above to obtain a mixing guarantee of the PS_{exp} in terms of $C_{\text{PI}}(\pi^Z)$, which satisfies $C_{\text{PI}}(\pi^Z) \lesssim (C_{\text{PI}}(\pi^X) + 1) \log n$ due to [Kla23] and Lemma 2.5.

Remark 2.10. In general, one cannot expect a mixing result of the PS_{exp} based on (LSI) even if the original distribution $\pi^X \propto \exp(-V)$ satisfies the LSI, as it does not hold in general for an exponential distribution. Hence, before sampling, an additional step such as truncation to a bounded convex set may be required. We will go through this procedure in §3 in order to invoke a mixing result via the LSI (see, for instance, Lemma 3.6).

2.2.2 Complexity of implementing the backward step

As opposed to the forward step, it is unclear how to sample from the backward distribution $\pi^{Z|Y=y} \propto \exp(-\alpha^\top z - \frac{1}{2h} \|y - z\|^2) |_{\mathcal{K}}$ for $\alpha = ne_{n+1}$. Due to

$$\alpha^\top z + \frac{1}{2h} \|z - y\|^2 = \frac{1}{2h} \|z - (y - h\alpha)\|^2 + \alpha^\top y - \frac{h}{2} \|\alpha\|^2,$$

the distribution can be written as

$$\pi^{Z|Y=y} \propto \exp(-\frac{1}{2h} \|z - (y - h\alpha)\|^2) |_{\mathcal{K}}.$$

Thus, we use rejection sampling with proposal $\mathcal{N}(y - h\alpha, hI_{n+1})$, accepting a proposed sample only if it lies within \mathcal{K} (which can be checked by using the evaluation oracle). The expected number of trials until the first acceptance is

$$\frac{1}{\ell(y)} := \frac{(2\pi h)^{(n+1)/2}}{\int_{\mathcal{K}} \exp(-\frac{1}{2h} \|z - (y - h\alpha)\|^2) dz}.$$

We show in this section that the expected number of wasted proposals is moderate under suitable choices of variance h and threshold N in the PS_{exp} . To this end, we first establish analytical and geometrical properties of $\pi^Y = \pi^Z * \mathcal{N}(0, hI_{n+1})$.

Density after taking the forward heat flow. We deduce a density of $\pi^Y = \pi^Z * \mathcal{N}(0, hI_{n+1})$, using the definition of convolution.

Lemma 2.11 (Density of π^Y). *For $\pi^Z \propto \exp(-\alpha^\top z) |_{\mathcal{K}}$, the density of $\pi^Y = \pi^Z * \mathcal{N}(0, hI_{n+1})$ is*

$$\pi^Y(y) = \frac{\int_{\mathcal{K}} \exp(-\frac{1}{2h} \|z - (y - h\alpha)\|^2) dz}{\int_{\mathcal{K}} \exp(-\alpha^\top z) dz \cdot (2\pi h)^{(n+1)/2}} \exp(-\alpha^\top y + \frac{1}{2} h \|\alpha\|^2) = \frac{\ell(y) \exp(-\alpha^\top y + \frac{1}{2} h \|\alpha\|^2)}{\int_{\mathcal{K}} \exp(-\alpha^\top z) dz}.$$

Proof. By the definition of the convolution,

$$\begin{aligned} \pi^Y(y) &= \frac{\int_{\mathcal{K}} \exp(-\alpha^\top z - \frac{1}{2h} \|z - y\|^2) dz}{\int_{\mathcal{K}} \exp(-\alpha^\top z) dz \cdot (2\pi h)^{(n+1)/2}} = \frac{\int_{\mathcal{K}} \exp(-\frac{1}{2h} \|z - (y - h\alpha)\|^2) dz}{\int_{\mathcal{K}} \exp(-\alpha^\top z) dz \cdot (2\pi h)^{(n+1)/2}} \exp(-\alpha^\top y + \frac{1}{2} h \|\alpha\|^2) \\ &= \frac{\ell(y)}{\int_{\mathcal{K}} \exp(-\alpha^\top z) dz} \exp(-\alpha^\top y + \frac{1}{2} h \|\alpha\|^2), \end{aligned}$$

which completes the proof. \square

We now identify an *effective domain* of π^Y , which takes up most of measure of π^Y . Below, $\mathcal{K}_\delta := \{x : d(x, \mathcal{K}) \leq \delta\}$ denotes the δ -blow up of \mathcal{K} .

Lemma 2.12. *In the setting of Problem 2.1, for $\tilde{\mathcal{K}} := \mathcal{K}_\delta + h\alpha$, if $\delta \geq 13hn$, then*

$$\pi^Y(\tilde{\mathcal{K}}^c) \leq \exp\left(-\frac{\delta^2}{2h} + 13\delta n\right).$$

Proof. Using the density formula of π^Y in Lemma 2.11,

$$\begin{aligned} \int_{\mathcal{K}} e^{-\alpha^\top z} dz \cdot \pi^Y(\tilde{\mathcal{K}}^c) &= \frac{1}{(2\pi h)^{(n+1)/2}} \int_{\tilde{\mathcal{K}}^c} \int_{\mathcal{K}} e^{-\frac{1}{2h} \|z-(y-h\alpha)\|^2} e^{-\alpha^\top y + \frac{1}{2} h\|\alpha\|^2} dz dy \\ &\stackrel{(i)}{=} \frac{1}{(2\pi h)^{(n+1)/2}} \int_{\mathcal{K}_\delta^c} \int_{\mathcal{K}} e^{-\frac{1}{2h} \|z-v\|^2} e^{-\alpha^\top v - \frac{1}{2} h\|\alpha\|^2} dz dv \\ &\stackrel{(ii)}{\leq} \frac{1}{(2\pi h)^{(n+1)/2}} \int_{\mathcal{K}_\delta^c} \int_{\mathcal{H}(v)} e^{-\frac{1}{2h} \|z-v\|^2} e^{-\alpha^\top v - \frac{1}{2} h\|\alpha\|^2} dz dv \\ &= \int_{\mathcal{K}_\delta^c} e^{-\alpha^\top v - \frac{1}{2} h\|\alpha\|^2} \left(\int_{d(v, \mathcal{K})}^{\infty} \frac{e^{-\frac{t^2}{2h}}}{\sqrt{2\pi h}} dt \right) dv =: (\#), \end{aligned}$$

where (i) follows from the change of variables by $v = y - h\alpha$, and in (ii) for $v \in \mathcal{K}_\delta^c$, $\mathcal{H}(v)$ denotes the supporting half-space at $\text{proj}_{\mathcal{K}}(v)$ containing \mathcal{K} , where $\text{proj}_{\mathcal{K}}(v)$ denotes the projection of v to \mathcal{K} . Denoting the CDF of the standard Gaussian by Φ , we have

$$\frac{1}{\sqrt{2\pi h}} \int_s^{\infty} e^{-\frac{t^2}{2h}} dt = 1 - \Phi(h^{-1/2}s) =: \Phi^c(h^{-1/2}s).$$

By the co-area formula and integration by parts, for the n -dimensional Hausdorff measure \mathcal{H}^n ,

$$\begin{aligned} (\#) &\stackrel{\text{coarea}}{=} e^{-\frac{1}{2}h\|\alpha\|^2} \int_\delta^\infty \Phi^c(h^{-1/2}s) \int_{\partial\mathcal{K}_s} e^{-\alpha^\top v} \mathcal{H}^n(dv) ds \\ &\stackrel{\text{IBP}}{=} e^{-\frac{1}{2}h\|\alpha\|^2} \left(\left[\Phi^c(h^{-1/2}s) \int_0^s \int_{\partial\mathcal{K}_t} e^{-\alpha^\top v} \mathcal{H}^n(dv) dt \right]_{s=\delta}^\infty + \int_\delta^\infty \frac{e^{-\frac{s^2}{2h}}}{\sqrt{2\pi h}} \int_0^s \int_{\partial\mathcal{K}_t} e^{-\alpha^\top v} \mathcal{H}^n(dv) dt ds \right) \\ &= e^{-\frac{1}{2}h\|\alpha\|^2} \left(\underbrace{\left[\Phi^c(h^{-1/2}s) \int_{\mathcal{K}_s \setminus \mathcal{K}} e^{-\alpha^\top v} dv \right]_{s=\delta}^\infty}_{=: \text{I}} + \int_\delta^\infty \frac{e^{-\frac{s^2}{2h}}}{\sqrt{2\pi h}} \int_{\mathcal{K}_s \setminus \mathcal{K}} e^{-\alpha^\top v} dv ds \right). \end{aligned} \quad (2.5)$$

As shown later in Lemma 2.13, this quantity is bounded by $e^{13sn} \int_{\mathcal{K}} \exp(-\alpha^\top v) dv$. From a standard bound on Φ^c [RW00, Theorem 20] given by

$$\Phi^c(h^{-1/2}s) \leq \frac{\exp(-\frac{s^2}{2h})}{\sqrt{2\pi}} \quad \text{for any } s > 0, \quad (2.6)$$

it follows that $\Phi^c(h^{-1/2}s) \int_{\mathcal{K}_s \setminus \mathcal{K}} e^{-\alpha^\top v} dv$ vanishes at $s = \infty$, and thus the term I can be bounded by 0.

Putting the bound in Lemma 2.13 into (2.5),

$$\int_{\mathcal{K}} e^{-\alpha^\top z} dz \cdot \pi^Y(\tilde{\mathcal{K}}^c) \leq \int_\delta^\infty \frac{1}{\sqrt{2\pi h}} e^{-\frac{s^2}{2h}} e^{13sn} ds \cdot \int_{\mathcal{K}} e^{-\alpha^\top z} dz.$$

Dividing both sides by $\int_{\mathcal{K}} e^{-\alpha^\top z} dz$ and invoking the standard bound on Φ^c again, we obtain

$$\pi^Y(\tilde{\mathcal{K}}^c) \leq \int_\delta^\infty \frac{1}{\sqrt{2\pi h}} e^{-\frac{s^2}{2h}} e^{13sn} ds = e^{52hn^2} \left(1 - \Phi\left(\frac{\delta - 13hn}{h^{1/2}}\right) \right)$$

$$\leq e^{52hn^2} \exp\left(-\frac{(\delta - 13hn)^2}{2h}\right) = \exp\left(-\frac{\delta^2}{2h} + 13\delta n\right),$$

which completes the proof. \square

This analytical property suggests taking $\delta \asymp 1/n$ and $h \asymp 1/n^2$. Thus, we set $h = \frac{c}{13^2 n^2}$ and $\delta = \frac{t}{13n}$ for parameters $c, t > 0$, under which $\delta \geq 13hn$ turns into $t \geq c$, and

$$\pi^Y(\tilde{\mathcal{K}}^c) \leq \exp\left(-\frac{t^2}{2c} + t\right).$$

We now prove an interesting result quantifying how fast $\int_{\mathcal{K}_s} \exp(-\alpha^\top v) dv$ increases in s .

Lemma 2.13. *In the setting of Problem 2.1,*

$$\frac{\int_{\mathcal{K}_s} \exp(-\alpha^\top v) dv}{\int_{\mathcal{K}} \exp(-\alpha^\top v) dv} \leq \exp(13sn).$$

Proof. Due to the assumption on the ground set $L_{V,g} = \{V(x) - V_0 \leq 10n\}$, it holds that $\mathcal{K} \cap [t = V_0/n + 10]$ contains $B_1^n(x_0)$. Then, the definition of \mathcal{K} ensures that \mathcal{K} contains $B_1^{n+1}(x_0, V_0/n + 11)$.

We now bound the quantity of interest: for $t_0 = 11 + V_0/n$,

$$\begin{aligned} \frac{\int_{\mathcal{K}_s} \exp(-\alpha^\top v) dv}{\int_{\mathcal{K}} \exp(-\alpha^\top v) dv} &= \frac{\int_{\mathcal{K}_s} \exp(-nt) dt dx}{\int_{\mathcal{K}} \exp(-nt) dt dx} = \frac{\int_{\mathcal{K}_s - t_0 e_{n+1}} \exp(-nt) dt dx}{\int_{\mathcal{K} - t_0 e_{n+1}} \exp(-nt) dt dx} \\ &\stackrel{(i)}{\leq} \frac{\int_{(1+s)(\mathcal{K} - t_0 e_{n+1})} \exp(-nt) dt dx}{\int_{\mathcal{K} - t_0 e_{n+1}} \exp(-nt) dt dx} \\ &\stackrel{(ii)}{=} (1+s)^{n+1} \frac{\int_{\mathcal{K} - t_0 e_{n+1}} \exp(-nt) \exp(-nst) dt dx}{\int_{\mathcal{K} - t_0 e_{n+1}} \exp(-nt) dt dx} \\ &\stackrel{(iii)}{\leq} e^{s(n+1)} e^{11sn} \frac{\int_{\mathcal{K} - t_0 e_{n+1}} \exp(-nt) dt dx}{\int_{\mathcal{K} - t_0 e_{n+1}} \exp(-nt) dt dx} \leq e^{13sn}, \end{aligned}$$

where (i) follows from $\mathcal{K}_s - t_0 e_{n+1} = (\mathcal{K} - t_0 e_{n+1})_s \subset (1+s)(\mathcal{K} - t_0 e_{n+1})$ due to $B_1(x_0, 0) \subset \mathcal{K} - t_0 e_{n+1}$, and in (ii) we used the change of variables by scaling the whole system by $(1+s)$, and (iii) follows from $t \geq -11$ in $\mathcal{K} - t_0 e_{n+1}$. \square

Per-step guarantees. Using the properties established above, we provide concrete choices of parameters, bounding a per-step query complexity for the backward step under a *warm start*.

Lemma 2.14. *In the setting of Problem 2.1, let μ be an M -warm initial distribution for π^Z . Given $k \in \mathbb{N}$ and $\eta \in (0, 1)$, set $S = \frac{16kM}{\eta} (\geq 16)$, $h = \frac{1}{n^2} \frac{(\log \log S)^2}{13^2 \log S}$ and $N = S \log^2 S$. Then, per iteration, the failure probability is at most η/k , and the expected number of queries is $\tilde{O}(M \log^2 \frac{k}{\eta})$.*

Proof. We note that $\mu * \mathcal{N}(0, hI_{n+1})$ is M -warm with respect to π^Y due to $d\mu/d\pi^X \leq M$. Hence, the failure probability per iteration is $\mathbb{E}_{\mu * \mathcal{N}(0, hI_{n+1})}[(1-\ell)^N] \leq M \mathbb{E}_{\pi^Y}[(1-\ell)^N]$. For $\tilde{\mathcal{K}} = \mathcal{K}_\delta + h\alpha$,

$$\begin{aligned} \mathbb{E}_{\pi^Y}[(1-\ell)^N] &= \int_{\tilde{\mathcal{K}}^c} \cdot + \int_{\tilde{\mathcal{K}} \cap [\ell \geq N^{-1} \log(3kM/\eta)]} \cdot + \int_{\tilde{\mathcal{K}} \cap [\ell \leq N^{-1} \log(3kM/\eta)]} \cdot \\ &\stackrel{(i)}{\leq} \pi^Y(\tilde{\mathcal{K}}^c) + \int_{[\ell \geq N^{-1} \log(3kM/\eta)]} \exp(-\ell N) d\pi^Y \end{aligned}$$

$$\begin{aligned}
& + \int_{\tilde{\mathcal{K}} \cap [\ell \leq N^{-1} \log(3kM/\eta)]} \frac{\ell(y) \exp(-\alpha^\top y + \frac{1}{2} h \|\alpha\|^2)}{\int_{\mathcal{K}} \exp(-\alpha^\top v) dv} dy \\
\stackrel{(ii)}{\leq} & \exp\left(-\frac{t^2}{2c} + t\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} \frac{\int_{\tilde{\mathcal{K}}} \exp(-\alpha^\top y + \frac{1}{2} h \|\alpha\|^2) dy}{\int_{\mathcal{K}} \exp(-\alpha^\top v) dv} \\
\stackrel{(iii)}{=} & \exp\left(-\frac{t^2}{2c} + t\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} \frac{\int_{\mathcal{K}_\delta} \exp(-\alpha^\top v - \frac{1}{2} h \|\alpha\|^2) dv}{\int_{\mathcal{K}} \exp(-\alpha^\top v) dv} \\
\stackrel{(iv)}{\leq} & \exp\left(-\frac{t^2}{2c} + t\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} e^{13\delta n} \\
\leq & \frac{\eta}{kM},
\end{aligned}$$

where in (i) we used the density formula of π^Y (Lemma 2.11), (ii) follows from Lemma 2.12, (iii) is due to the change of variables via $y = v + h\alpha$, in (iv) we used Lemma 2.13, and the last line follows from the choices of $c = \frac{(\log \log S)^2}{13^2 \log S}$, $t = \log \log S$, and $N = S \log^2 S$. Therefore, $\mathbb{E}_{\mu^* \mathcal{N}(0, hI_{n+1})}[(1 - \ell)^N] \leq \eta/k$. Note that under these choices of parameters, we have $t \geq c$ and so $\delta \geq 13hn$, as requested in Lemma 2.12.

We now bound the expected number of trials until the first acceptance. In a similar vein to above, the expected number is bounded by $M \mathbb{E}_{\pi^Y}[\ell^{-1} \wedge N]$, where

$$\begin{aligned}
\int_{\mathbb{R}^{n+1}} \frac{1}{\ell} \wedge N d\pi^Y & \leq \int_{\tilde{\mathcal{K}}} \frac{1}{\ell} d\pi^Y + N \pi^Y(\tilde{\mathcal{K}}^c) \leq \frac{\int_{\mathcal{K}_\delta} \exp(-\alpha^\top v - \frac{1}{2} h \|\alpha\|^2) dv}{\int_{\mathcal{K}} \exp(-\alpha^\top v) dv} + N \exp\left(-\frac{t^2}{2c} + t\right) \\
& \leq e^t + N \exp\left(-\frac{t^2}{2c} + t\right) \leq \log S + \log^2 S \\
& \leq 2 \log^2 \frac{16kM}{\eta}.
\end{aligned}$$

Therefore, from an M -warm start, the expected number of queries per iteration is $\tilde{\mathcal{O}}(M \log^2 \frac{k}{\eta})$. \square

Combining the results on the mixing and per-iteration complexity together, we can conclude the query complexity of sampling from a logconcave distribution, as claimed in Theorem 1.5.

Theorem 2.15. *In the setting of Problem 2.1, consider $\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}$ given in (exp-red). For any given $\eta, \varepsilon \in (0, 1)$, $q \geq 2$, $k \in \mathbb{N}$ defined below, the PS_{exp} with $h = (13^2 n^2 \log \frac{16kM}{\eta})^{-1}$, $N = \frac{16kM}{\eta} \log^2 \frac{16kM}{\eta}$, and initial distribution μ_0 which is M -warm with respect to π achieves $\mathcal{R}_q(\mu_k \parallel \pi) \leq \varepsilon$ (so $\mathcal{R}_q(\mu_k^X \parallel \pi^X) \leq \varepsilon$) after $k = \tilde{\mathcal{O}}(qn^2(\|\Sigma_x\| \vee 1) \log^2 \frac{M}{\eta\varepsilon})$ iterations, where μ_k is the law of k -th iterate, and Σ_x is the covariance matrix of $\pi^X \propto \exp(-V)$. With probability $1 - \eta$, the algorithm iterates k times successfully, using $\tilde{\mathcal{O}}(qMn^2(\|\Sigma_x\| \vee 1) \log^4 \frac{1}{\eta\varepsilon})$ evaluation queries in expectation. Moreover, an M -warm start for π^X can be used to generate an M -warm start for π .*

Proof. By Lemma 2.9 and Lemma 2.5, the PS_{exp} should iterate

$$k = \mathcal{O}(qh^{-1} C_{\text{PI}}(\pi) \log \frac{M}{\varepsilon}) = \mathcal{O}(qh^{-1}(\|\Sigma_x\| \vee 1) \log \frac{M}{\varepsilon}) \stackrel{\text{Remark 2.16}}{=} \tilde{\mathcal{O}}(qn^2(\|\Sigma_x\| \vee 1) \log^2 \frac{M}{\eta\varepsilon})$$

times to achieve ε -distance in \mathcal{R}_q . Under the choice of h and N as claimed, each iteration succeeds with probability $1 - k/\eta$ and uses $\tilde{\mathcal{O}}(M \log^2 k/\eta)$ queries in expectation by Lemma 2.14. Therefore, throughout k iterations, the algorithm succeeds with probability $1 - \eta$, using

$$\tilde{\mathcal{O}}(qMn^2(\|\Sigma_x\| \vee 1) \log^4 \frac{1}{\eta\varepsilon})$$

evaluation queries in expectation.

Next, an M -warm start μ^X for π^X can be lifted to an M -warm distribution on \mathbb{R}^{n+1} for π through the following procedure: generate $x \sim \mu^X$ and draw $t \sim \pi^{T|X=x} \propto e^{-nt} \cdot \mathbb{1}[t \geq V(x)/n]$, obtaining (x, t) . Note that $\pi^{T|X=x}$ can be sampled easily by drawing $u \sim \text{Unif}([0, 1])$ and taking $F^{-1}(u)$, where F is the CDF of $\pi^{T|X=x}$. Then, the density of the law of (x, t) is $\mu^X \cdot \pi^{T|X}$, and it follows from $\pi(x, t) = \pi^X \cdot \pi^{T|X}$ that

$$\frac{\mu^X \pi^{T|X}}{\pi} = \frac{\mu^X}{\pi^X} \leq M,$$

which completes the proof. \square

Remark 2.16 (Self-dependence of parameters). Upon substitution of the chosen h , the required number of iterations would be

$$k \gtrsim qn^2(\|\Sigma_x\| \vee 1) \log \frac{M}{\varepsilon} \log \frac{kM}{\eta},$$

where k shows up in both sides. When $c \geq 1$, since $x \geq c \log x$ holds for $x \geq 2c \log c$, the above bound on k can be turned into

$$k \gtrsim qn^2(\|\Sigma_x\| \vee 1) \log \frac{M}{\varepsilon} \log(qn^2(\|\Sigma_x\| \vee 1) \frac{M}{\eta} \log \frac{M}{\varepsilon}) = \tilde{\mathcal{O}}(qn^2(\|\Sigma_x\| \vee 1) \log \frac{M}{\varepsilon} \log \frac{M}{\eta}).$$

3 Warm-start generation via tilted Gaussian cooling

In the previous section, we assumed access to an M -warm distribution for $\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}$. Here we address the problem of generating a warm start (i.e., $M = \mathcal{O}(1)$).

Problem 3.1 (Warm-start generation). Given access to a well-defined function oracle $\text{Eval}_{x_0, R}(V)$ for convex $V : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, what is the complexity of generating an $\mathcal{O}(1)$ -warm start μ^X with respect to π^X (i.e., $\mathcal{R}_\infty(\mu^X \parallel \pi^X) = \mathcal{O}(1)$)?

Compared to Problem 2.1, this **requires** access to x_0 and R . We prove **Result 2** in this section:

Theorem 3.2 (Restatement of Theorem 1.6). *In the setting of Problem 3.1, there exists an algorithm that for given $\eta, \varepsilon \in (0, 1)$, with probability at least $1 - \eta$ returns a sample X_* satisfying $\mathcal{R}_\infty(\text{law } X_* \parallel \pi^X) \leq \varepsilon$, using $\tilde{\mathcal{O}}(n^2(R^2 \vee n) \text{polylog}^{1/\eta\varepsilon})$ evaluation queries in expectation. In particular, if π^X is well-rounded (i.e., $R^2 \asymp n$), then we need $\tilde{\mathcal{O}}(n^3 \text{polylog}^{1/\eta\varepsilon})$ queries in expectation.*

As in the previous section, we tackle this problem after reducing it to the exponential distribution through **(exp-red)**, and then truncate this distribution to a convex domain.

Convex truncation. Using Lemma A.7 with $\mathbb{E}_{\pi^X}[\|X - x_0\|^2] \leq R^2$, we have that for any $l \geq 1$,

$$\mathbb{P}_{\pi^X}(\|X - x_0\| \geq Rl) \leq \exp(-l + 1).$$

In addition, since $\mathbb{E}_{\pi^T}[|T - \mu_t|^2] = \mathbb{E}_\pi[|T - \mu_t|^2] \leq 160$ (Lemma 2.5), we also have

$$\mathbb{P}_{\pi^T}(|T - \mu_t| \geq 13l) \leq \exp(-l + 1) \quad \text{for any } l \geq 1.$$

Putting these two together and using $\mu_t \leq V_0/n + 5 \leq V(x_0)/n + 5$ (Lemma 2.4), for $\varepsilon > 0$, $l = \log \frac{2\varepsilon}{\varepsilon}$, and $B = \frac{V(x_0)}{n} + 5 + 13l$,

$$\mathbb{P}_\pi\{(B_{Rl}^n(x_0) \times (-\infty, B])^c\} \leq \mathbb{P}_\pi\{(B_{Rl}^n(x_0) \times \mathbb{R})^c\} + \mathbb{P}_\pi\{(\mathbb{R}^n \times [B, \infty))^c\}$$

$$= \mathbb{P}_{\pi^x}(\mathbb{R}^n \setminus B_{Rl}^n(x_0)) + \mathbb{P}_{\pi^T}(\mathbb{R} \setminus [B, \infty)) \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Thus, $\bar{\mathcal{K}} := \mathcal{K} \cap \{B_{Rl}(x_0) \times (-\infty, B]\}$ takes up $(1 - \varepsilon)$ measure of π , and $B_{Rl}(x_0) \times (V_0/n, B]$ could be thought of as the *essential domain* of π . Moreover, since $x_0 \in \mathbb{L}_{\pi^x, V}$ (i.e., $V(x_0) - V_0 \leq 10n$),

$$\frac{V(x_0)}{n} + 5 + 13l - \frac{V_0}{n} \leq 15 + 13l,$$

so the diameter of $\bar{\mathcal{K}}$ is $\mathcal{O}(lR)$. Going forward, we use $\bar{\pi}$ to denote $\pi|_{\bar{\mathcal{K}}} \propto \exp(-nt)|_{\bar{\mathcal{K}}}$.

3.1 Tilted Gaussian Cooling

We would like to find a sequence $\{\mu_i\}_{0 \leq i \leq m}$ of distributions, where μ_0 is easy to sample, μ_m is our target, and μ_i is $\mathcal{O}(1)$ -warm with respect to μ_{i+1} . Then, as μ_i is warm for μ_{i+1} , we can run an efficient sampler to go from μ_i to μ_{i+1} .

Gaussian Cooling, introduced in [CV18], can generate an $\mathcal{O}(1)$ -warm start for the uniform distribution over a convex body. However, extending this to general logconcave distributions has remained elusive. A straightforward extension to the exponential distribution $\bar{\pi}$ fails due to an exponential change in the value of e^{-nt} in $\bar{\mathcal{K}}$; roughly speaking, while t may vary by at most $\Theta(l)$, the density can become exponentially small relative to its maximum. This significant change in value prevents us from achieving $\mathcal{O}(1)$ -warmness, necessitating additional ideas to address this challenge.

We propose Tilted Gaussian Cooling (Algorithm 2), which handles the x -direction through the Gaussian Cooling algorithm and the t -direction through an exponential tilt. Its annealing distributions have two parameters σ^2 and ρ , given as

$$\mu_{\sigma^2, \rho}(x, t) \propto \exp\left(-\frac{1}{2\sigma^2} \|x - x_0\|^2 - \rho t\right)|_{\bar{\mathcal{K}}}.$$

We can translate the whole system by $(x, t) \mapsto (x - x_0, t - \frac{V(x_0)}{n} - 11)$ so that the shifted $\bar{\mathcal{K}}$ contains a unit ball centered at the origin (i.e., $x_0 = 0$), and t is within $[-21, 13l - 6]$ over $\bar{\mathcal{K}}$.

We develop in §3.2 a sampling algorithm PS_{ann} for these annealing distributions. Assuming this sampler is given, we provide an outline of Tilted Gaussian Cooling. For the annealing distribution $\mu_i := \mu_{\sigma_i^2, \rho_i}$, we use $\bar{\mu}_i$ to denote an approximate distribution close to μ_i produced by the sampler. Tilted Gaussian Cooling has two main annealing phases:

- Initialization ($\sigma^2 = n^{-1}$ and $\rho = l^{-1}$)
 - Initial distribution: $\mu_0, \bar{\mu}_0 \propto \exp(-\frac{n}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$ (run rejection sampling with proposal $\mathcal{N}(0, n^{-1}I_n) \otimes \text{Unif}([-21, 13l - 6])$)
 - Target distribution: $\mu_1 \propto \exp(-\frac{n}{2} \|x\|^2 - t/l)|_{\bar{\mathcal{K}}}$.
- Phase I ($n^{-1} \leq \sigma^2 \leq 1$ and $l^{-1} \leq \rho \leq n$)
 - Run the PS_{ann} with initial $\bar{\mu}_i$, target μ_{i+1} , and accuracy $\log 2$ in \mathcal{R}_∞ , where

$$\sigma_{i+1}^2 = \sigma_i^2 \left(1 + \frac{1}{n}\right) \quad \& \quad \rho_{i+1} = \rho_i \left(1 + \frac{1}{ln}\right).$$

- Phase II ($1 \leq \sigma^2 \leq l^2 R^2$)
 - Run the PS_{ann} with initial $\bar{\mu}_i$, target μ_{i+1} , and accuracy $\log 2$ in \mathcal{R}_∞ , where

$$\sigma_{i+1}^2 = \sigma_i^2 \left(1 + \frac{\sigma_i^2}{l^2 R^2}\right).$$

Algorithm 2 Tilted Gaussian Cooling

Input: convex $\mathcal{K} = \{(x, t) : V(x) \leq nt\}$, accuracy $\varepsilon > 0$, target dist. $\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}$ s.t. $B_1^n(x_0) \subset \mathcal{L}_{\pi^{x,g}}$ and $\mathbb{E}_{\pi}[\|X - x_0\|^2] \leq R^2$.

Output: a sample z with law approximately close to π .

- 1: Let $\bar{\mathcal{K}} = \mathcal{K} \cap (B_{Rl}^n(x_0) \times (-\infty, V(x_0)/n + 5 + 13l])$ with $l = \log \frac{6}{\varepsilon}$.
- 2: Draw $z_0 \sim \exp(-\frac{n}{2} \|x - x_0\|^2)|_{\bar{\mathcal{K}}}$ via rejection sampling with proposal $\mathcal{N}(x_0, n^{-1}I_n) \otimes \text{Unif}((V(x_0)/n - 10, V(x_0)/n + 5 + 13l])$.
- 3: Denote $\mu_{\sigma^2, \rho} := \exp(-\frac{1}{2\sigma^2} \|x - x_0\|^2 - \rho t)|_{\bar{\mathcal{K}}}$ for $\sigma^2 = n^{-1}$ and $\rho = l^{-1}$.
- 4: Get $z \sim \text{PS}_{\text{ann}}$ with initial z_0 , target $\mu_{\sigma^2, \rho}$, accuracy $\log 2$ (in \mathcal{R}_{∞}), and success prob. $\frac{\eta}{\widetilde{\mathcal{O}}(l^2 R^2 \vee n)}$.
- 5: **while** $\sigma^2 \in [n^{-1}, l^2 R^2]$ and $\rho \in [l^{-1}, n]$ **do**
- 6: Sample z from PS_{ann} with the same setup. Then update

$$\begin{aligned} \sigma^2 &\leftarrow \sigma^2 \left(1 + \frac{1}{n}\right) \quad \& \quad \rho \leftarrow \min(n, \rho \left(1 + \frac{1}{ln}\right)) \quad \text{when } \sigma^2 \in \left[\frac{1}{n}, 1\right], \rho \in \left[\frac{1}{l}, n\right] \\ \sigma^2 &\leftarrow \sigma^2 \left(1 + \frac{\sigma^2}{l^2 R^2}\right) \quad \text{when } \sigma^2 \in [1, l^2 R^2]. \end{aligned}$$

7: **end while**

- 8: Return $z \sim \text{PS}_{\text{exp}}$ with initial $\mu_{l^2 R^2, n}$, target $\pi|_{\bar{\mathcal{K}}}$, accuracy $\varepsilon/3$, and success prob. $\frac{\eta}{\widetilde{\mathcal{O}}(l^2 R^2 \vee n)}$.
-

- Termination ($\sigma^2 = l^2 R^2$)

– Run the PS_{exp} with initial $\mu_{l^2 R^2, n}(x, t)$ and target $\bar{\pi} \propto \exp(-nt)|_{\bar{\mathcal{K}}}$.

3.2 Proximal sampler for annealing distributions

We employ PS to sample from the annealing distributions $\{\mu_{\sigma^2, \rho}\}$. For $v := (x, t) \in \mathbb{R}^n \times \mathbb{R}$ and $w := (y, s) \in \mathbb{R}^n \times \mathbb{R}$, PS for this target and variance h , called PS_{ann} (Algorithm 3), can be implemented by repeating

- [Forward] $w \sim \mu^{W|V=v} = \mathcal{N}(v, hI_{n+1})$.
- [Backward] For $\tau := \frac{\sigma^2}{h + \sigma^2} < 1$, $y_{\tau} := \tau y$, and $h_{\tau} := \tau h$, run

$$v \sim \mu^{V|W=w} \propto \exp\left(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t - \frac{1}{2h} \|w - v\|^2\right)|_{\bar{\mathcal{K}}} \propto [\mathcal{N}(y_{\tau}, h_{\tau}I_n) \times \mathcal{N}(s - \rho h, h)]|_{\bar{\mathcal{K}}},$$

where the backward step is implemented by rejection sampling with the proposal $\mathcal{N}(y_{\tau}, h_{\tau}I_n) \times \mathcal{N}(s - \rho h, h)$. Note that PS_{exp} may be thought of as a special case of PS_{ann} with $\sigma^2 = \infty$. As in §2.2, we analyze its mixing and per-step complexity.

3.2.1 Convergence rate

For PS_{ann} , we use a mixing result based on (LSI), rather than (PI). As opposed to the exponential distribution, the annealing distributions have $C_{\text{LSI}}(\mu) < \infty$, so we can now invoke the (LSI)-mixing result in Lemma 2.9. Therefore, for $q \geq 1$, the kernel P of PS_{ann} , and initial distribution ν with $\nu \ll \mu$, it holds that

$$\mathcal{R}_q(\nu P \parallel \mu) \leq \frac{\mathcal{R}_q(\nu \parallel \mu)}{(1 + h/C_{\text{LSI}}(\mu))^{2/q}}. \quad (3.1)$$

Algorithm 3 PS_{ann}

Input: initial pt. $v_0 \sim \pi_0 \in \mathcal{P}(\mathbb{R}^{n+1})$, truncated $\bar{\mathcal{K}}$, $k \in \mathbb{N}$, threshold N , variances h , parameters $\sigma^2, \rho > 0$.

Output: $v_{k+1} = (x_{k+1}, t_{k+1})$.

- 1: Let $\tau = \frac{\sigma^2}{h+\sigma^2}$. Denote $v = (x, t)$ and $w = (y, s)$ in $\mathbb{R}^n \times \mathbb{R}$.
 - 2: **for** $i = 0, \dots, k$ **do**
 - 3: Sample $w_{i+1} \sim \mathcal{N}(v_i, hI_{n+1})$.
 - 4: Sample $v_{i+1} \sim \mathbf{G}(w_{i+1})|_{\bar{\mathcal{K}}}$ for $\mathbf{G}(w) := \mathcal{N}(\tau y, \tau h I_n) \times \mathcal{N}(s - \rho h, h)$.
 - 5: (↑) Repeat $v_{i+1} \sim \mathbf{G}(w_{i+1})$ until $v_{i+1} \in \bar{\mathcal{K}}$. If $\#\text{attempts}_i \geq N$, declare **Failure**.
 - 6: **end for**
-

We provide a bound on $C_{\text{LSI}}(\mu)$, using the Bakry-Émery criterion and bounded perturbation.

Lemma 3.3. $C_{\text{LSI}}(\mu) \lesssim \sigma^2 \vee l^2$ for $\mu(x, t) \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)|_{\bar{\mathcal{K}}}$.

Proof. Consider a probability measure defined by

$$\nu(x, t) \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \frac{1}{2l^2} t^2 - \rho t).$$

It is $\min(\sigma^{-2}, l^{-2})$ -strongly logconcave, so $C_{\text{LSI}}(\nu) \leq \sigma^2 \vee l^2$ by the Bakry-Émery criterion (Lemma A.1). It is well known that a convex truncation of strongly logconcave measures only helps in satisfying the criterion, so $C_{\text{LSI}}(\nu|_{\bar{\mathcal{K}}}) \leq \sigma^2 \vee l^2$ [Wan13, Theorem 3.3.2]. Since

$$\frac{d\mu}{d\nu|_{\bar{\mathcal{K}}}} = \mathbb{1}[(x, t) \in \bar{\mathcal{K}}] \exp\left(\frac{t^2}{2l^2}\right) \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \frac{1}{2l^2} t^2 - \rho t)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)},$$

and $t \in [-21, 13l - 6]$ over $\bar{\mathcal{K}}$, we have

$$\exp(-\Theta(1)) \leq \inf_{\bar{\mathcal{K}}} \frac{d\mu}{d\nu|_{\bar{\mathcal{K}}}} \leq \sup_{\bar{\mathcal{K}}} \frac{d\mu}{d\nu|_{\bar{\mathcal{K}}}} \leq \exp(\Theta(1)).$$

By the Holley–Stroock perturbation principle (Lemma A.2),

$$C_{\text{LSI}}(\mu) \leq \exp(\Theta(1)) C_{\text{LSI}}(\nu|_{\bar{\mathcal{K}}}) \lesssim \sigma^2 \vee l^2,$$

which completes the proof. \square

Combining the exponential contraction (3.1) and the bound on the LSI constant, we conclude that if PS_{ann} started at an initial distribution ν iterates $\mathcal{O}(qh^{-1}C_{\text{LSI}}(\mu) \log \frac{\mathcal{R}_q(\nu||\mu)}{\varepsilon}) = \mathcal{O}(qh^{-1}(\sigma^2 \vee l^2) \log \frac{\mathcal{R}_q(\nu||\mu)}{\varepsilon})$ times, then it achieves ε -distance within the target ν in \mathcal{R}_q -divergence.

We can strengthen this \mathcal{R}_q -divergence result to the stronger \mathcal{R}_∞ -divergence by following the boosting scheme used in [KZ25], which stems from a classical result in [DMLM03].

Lemma 3.4 ([KZ25, Theorem 14]). *Consider a Markov chain with initial distribution ν and kernel P reversible with respect to a stationary distribution μ . Then,*

$$\left\| \frac{d(\nu P^k)}{d\mu} - 1 \right\|_{L^\infty} \leq 2 \left\| \frac{d\nu}{d\mu} - 1 \right\|_{L^\infty} \text{ess sup}_x \|\delta_x P^k - \mu\|_{\text{TV}},$$

and in particular, $\mathcal{R}_\infty(\nu P^k || \mu) \leq 2 \left\| \frac{d\nu}{d\mu} - 1 \right\|_{L^\infty} \text{ess sup}_x \|\delta_x P^k - \mu\|_{\text{TV}}$.

In words, a uniform bound on the TV-distance from *any start* implies the Rényi-infinity bound. To use this result, we establish the uniform ergodicity of PS_{ann} .

Lemma 3.5. *For given $\varepsilon \in (0, 1)$, let P be the Markov kernel of PS_{ann} with variance h and target distribution μ . Then, $\sup_{v_0 \in \bar{\mathcal{K}}} \|\delta_{v_0} P^k - \mu\|_{\text{TV}} \leq \varepsilon$ for $k = \mathcal{O}(h^{-1}(\sigma^2 \vee l^2) \log \frac{n+h^{-1}R^2}{\varepsilon})$. If an initial distribution ν is M -warm with respect to μ , then $\mathcal{R}_\infty(\nu P^k \parallel \mu) \leq \varepsilon$ after $k = \mathcal{O}(h^{-1}(\sigma^2 \vee l^2) \log \frac{M(n+h^{-1}R^2)}{\varepsilon})$ iterations.*

Proof. We first bound the warmness of $\delta_{v_0} P$ for μ . By the definition of PS_{ann} , the forward step brings δ_{v_0} to $\mathcal{N}(v_0, hI_{n+1})$, and then the backward step $\pi^{V|W=(y,s)}(x, t) \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t - \frac{1}{2h} \|w - v\|^2)|_{\bar{\mathcal{K}}}$ brings it to, for $v = (x, t)$ and $w = (y, s)$

$$(\delta_{v_0} P)(x, t) = \mathbf{1}_{\bar{\mathcal{K}}}(x, t) \int \frac{\exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t - \frac{1}{2h} \|w - v\|^2) \exp(-\frac{1}{2h} \|w - v_0\|^2)}{(2\pi h)^{(n+1)/2} \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l - \frac{1}{2h} \|(y, s) - (z, l)\|^2) dz dl} dy ds.$$

For $D := \text{diam } \bar{\mathcal{K}} \lesssim Rl$, it follows from Young's inequality ($\|a - b\|^2 \leq 3\|a\|^2 + \frac{3}{2}\|b\|^2$) that

$$\frac{1}{2h} \|(y, s) - (z, l)\|^2 \leq \frac{3}{4h} \|(y, s)\|^2 + \frac{3}{2h} \|(z, l)\|^2 \leq \frac{3}{4h} \|(y, s)\|^2 + \frac{3D^2}{2h}.$$

Hence,

$$\delta_{v_0} P \leq \frac{\exp(\frac{3D^2}{2h}) \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)}{(2\pi h)^{\frac{n+1}{2}} \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l) dz dl} \int \exp(-\frac{1}{2h} \|w - v\|^2 - \frac{1}{2h} \|w - v_0\|^2 + \frac{3}{4h} \|w\|^2) dy ds.$$

The exponent of the integrand can be bounded as follows:

$$\begin{aligned} & -\frac{1}{2h} \|w - v\|^2 - \frac{1}{2h} \|w - v_0\|^2 + \frac{3}{4h} \|w\|^2 \\ &= -\frac{1}{4h} \|w - 2(v + v_0)\|^2 - \frac{1}{2h} \|v\|^2 - \frac{1}{2h} \|v_0\|^2 + \frac{1}{h} \|v + v_0\|^2 \\ &\leq -\frac{1}{4h} \|w - 2(v + v_0)\|^2 + \frac{4D^2}{h}, \end{aligned}$$

so it follows that

$$\delta_{v_0} P \leq \frac{2^{(n+1)/2} \exp(\frac{6D^2}{h}) \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l) dz dl}.$$

Dividing this by $\mu(x, t) \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)|_{\bar{\mathcal{K}}}$, we have

$$\frac{\delta_{v_0} P}{\mu} \leq 2^{\frac{n+1}{2}} \exp(6h^{-1}D^2),$$

which implies $\mathcal{R}_q(\delta_{v_0} P \parallel \mu) \lesssim n + h^{-1}D^2$. Thus, PS_{ann} can achieve $\mathcal{R}_q(\delta_{v_0} P^k \parallel \mu) \leq \varepsilon$ for $k = \mathcal{O}(qh^{-1}(\sigma^2 \vee l^2) \log \frac{n+h^{-1}D^2}{\varepsilon})$, and due to $2\|\cdot\|_{\text{TV}}^2 \leq \lim_{q \downarrow 1} \mathcal{R}_q$,

$$\sup_{v_0 \in \bar{\mathcal{K}}} \|\delta_{v_0} P^k - \mu\|_{\text{TV}} \leq \varepsilon.$$

For the second claim, replacing ε with ε/M and using Lemma 3.4 with $\|\frac{d\nu}{d\mu} - 1\|_{L_\infty} \leq M$, we have $\|\frac{d(\nu P^k)}{d\mu} - 1\|_{L_\infty} \leq \varepsilon$ and thus $\mathcal{R}_\infty(\nu P^k \parallel \mu) \leq \varepsilon$. \square

Similarly, we can establish a \mathcal{R}_∞ -guarantee for the exponential sampling considered in §2.2.

Lemma 3.6. *For given $\varepsilon \in (0, 1)$, let P be the Markov kernel of PS_{exp} with variance h and target distribution $\bar{\pi} \propto \exp(-nt)|_{\bar{\mathcal{K}}}$. Then, $\sup_{z_0 \in \bar{\mathcal{K}}} \|\delta_{z_0} P^k - \bar{\pi}\|_{\text{TV}} \leq \varepsilon$ for $k = \mathcal{O}(h^{-1} R^2 l^2 \log \frac{n+h^{-1}R^2}{\varepsilon})$. If an initial distribution ν is M -warm with respect to $\bar{\pi}$, then $\mathcal{R}_\infty(\nu P^k \parallel \bar{\pi}) \leq \varepsilon$ after $k = \mathcal{O}(h^{-1} R^2 l^2 \log \frac{M(n+h^{-1}R^2)}{\varepsilon})$ iterations.*

Proof. As before, for $z_0 \in \bar{\mathcal{K}}$, the composition of the forward and backward step brings δ_{z_0} to

$$(\delta_{z_0} P)(x, t) = \mathbf{1}_{\bar{\mathcal{K}}}(x, t) \int \frac{\exp(-nt - \frac{1}{2h} \|y - z\|^2) \exp(-\frac{1}{2h} \|y - z_0\|^2)}{(2\pi h)^{(n+1)/2} \int_{\bar{\mathcal{K}}} \exp(-nt - \frac{1}{2h} \|y - z\|^2) dz} dy.$$

Using $\frac{1}{2h} \|y - z\|^2 \leq \frac{3}{4h} \|y\|^2 + \frac{3}{2h} \|z\|^2 \leq \frac{3}{4h} \|y\|^2 + \frac{3D^2}{2h}$ (Young's inequality), for $D := \text{diam } \bar{\mathcal{K}} \lesssim Rl$,

$$\begin{aligned} \delta_{z_0} P &\leq \frac{\exp(\frac{3D^2}{2h}) \exp(-nt)}{(2\pi h)^{(n+1)/2} \int_{\bar{\mathcal{K}}} \exp(-nt) dx dt} \int \exp(-\frac{1}{2h} \|y - z\|^2 - \frac{1}{2h} \|y - z_0\|^2 + \frac{3}{4h} \|y\|^2) dy \\ &\leq \frac{2^{(n+1)/2} \exp(\frac{6D^2}{h}) \exp(-nt)}{\int_{\bar{\mathcal{K}}} \exp(-nt) dx dt}, \end{aligned}$$

where the last line follows from

$$\begin{aligned} &-\frac{1}{2h} \|y - z\|^2 - \frac{1}{2h} \|y - z_0\|^2 + \frac{3}{4h} \|y\|^2 \\ &= -\frac{1}{4h} \|y - 2(z + z_0)\|^2 - \frac{1}{2h} \|z\|^2 - \frac{1}{2h} \|z_0\|^2 + \frac{1}{h} \|z + z_0\|^2 \\ &\leq -\frac{1}{4h} \|y - 2(z + z_0)\|^2 + \frac{4D^2}{h}. \end{aligned}$$

Dividing $\delta_{z_0} P$ by $\bar{\pi}(x, t) \propto \exp(-nt)|_{\bar{\mathcal{K}}}$,

$$\frac{\delta_{z_0} P}{\bar{\pi}} \leq 2^{\frac{n+1}{2}} \exp(6h^{-1}D^2),$$

which implies $\mathcal{R}_q(\delta_{z_0} P \parallel \bar{\pi}) \lesssim \frac{q}{q-1} (n + h^{-1}D^2)$. Noting that the truncated distribution $\bar{\pi}$ has a bounded support, so $C_{\text{LSI}}(\bar{\pi}) = \mathcal{O}(D^2)$, the PS_{exp} can achieve $\mathcal{R}_q(\delta_{z_0} P^k \parallel \bar{\pi}) \leq \frac{\varepsilon}{M}$ after $k = \mathcal{O}(qh^{-1}D^2 \log \frac{M(n+h^{-1}D^2)}{\varepsilon})$ iterations, and

$$\sup_{z_0 \in \bar{\mathcal{K}}} \|\delta_{z_0} P^k - \bar{\pi}\|_{\text{TV}} \leq \frac{\varepsilon}{M}.$$

Hence, one can complete the proof using Lemma 3.4 with $\|\frac{d\nu}{d\bar{\pi}} - 1\|_{L^\infty} \leq M$. \square

3.2.2 Complexity of implementing the backward step

We recall the backward step: for $\tau = \frac{\sigma^2}{h+\sigma^2} < 1$, $y_\tau = \tau y$, and $h_\tau = \tau h$,

$$v \sim \mu^{V|W=w} \propto \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t - \frac{1}{2h} \|w - v\|^2)|_{\bar{\mathcal{K}}} \propto [\mathcal{N}(y_\tau, h_\tau I_n) \times \mathcal{N}(s - \rho h, h)]|_{\bar{\mathcal{K}}}.$$

Since we use the proposal $\mathcal{N}(y_\tau, h_\tau I_n) \times \mathcal{N}(s - \rho h, h)$ for rejection sampling, the success probability of each trial at $w = (y, v)$ is

$$\ell(w) := \frac{1}{(2\pi h_\tau)^n (2\pi h)^{1/2}} \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2h_\tau} \|x - y_\tau\|^2 - \frac{1}{2h} |t - (s - \rho h)|^2) dx dt.$$

Density after taking the forward step. As in Lemma 2.11, we deduce the density of $\mu^W = \mu^V * \mathcal{N}(0, hI_{n+1})$ by computing the convolution directly.

Lemma 3.7 (Density of μ^W). For $\mu^V \propto \exp(-\frac{\|x\|^2}{2\sigma^2} - \rho t)|_{\bar{\mathcal{K}}}$,

$$\begin{aligned} \mu^W(y, s) &= \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2h_\tau} \|x - y_\tau\|^2 - \frac{1}{2h} |t - (s - \rho h)|^2) dx dt}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t) dx dt \cdot (2\pi h)^{(n+1)/2}} \exp(-\frac{1}{2\tau\sigma^2} \|y_\tau\|^2) \exp(-\rho s + \frac{1}{2} \rho^2 h) \\ &= \frac{\tau^{n/2} \ell(y, s)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t) dx dt} \exp(-\frac{1}{2\tau\sigma^2} \|y_\tau\|^2) \exp(-\rho s + \frac{1}{2} \rho^2 h), \end{aligned}$$

We now claim that the following is the *effective domain* of μ^W :

$$\tilde{\mathcal{K}} = \begin{bmatrix} \tau^{-1} I_n & \\ & 1 \end{bmatrix} \bar{\mathcal{K}}_\delta + \begin{bmatrix} 0 \\ \rho h \end{bmatrix}. \quad (3.2)$$

Lemma 3.8. In the setting of Problem 3.1, for $\tilde{\mathcal{K}}$ in (3.2), if $\delta \geq 24hn$, then

$$\mu^W(\tilde{\mathcal{K}}^c) \leq \exp(-\frac{\delta^2}{2h} + 24\delta n + hn^2).$$

Proof. Using the density formula of μ^W in Lemma 3.7, for $v = (x, t)$, $w = (y, s)$, and $\bar{\mathcal{K}}_\delta^c := (\bar{\mathcal{K}}_\delta)^c$,

$$\begin{aligned} & \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t) dx dt \cdot \mu^W(\tilde{\mathcal{K}}^c) \\ &= \frac{1}{(2\pi h)^{\frac{n+1}{2}}} \int_{\bar{\mathcal{K}}^c} \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2h_\tau} \|x - y_\tau\|^2 - \frac{1}{2h} |t - (s - \rho h)|^2) \exp(-\frac{\|y_\tau\|^2}{2\tau\sigma^2} - \rho s + \frac{\rho^2 h}{2}) dv dw \\ &\stackrel{(i)}{=} \frac{\tau^{-n/2}}{(2\pi h_\tau)^{n/2} (2\pi h)^{1/2}} \int_{\bar{\mathcal{K}}_\delta^c} \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2h_\tau} \|x - y\|^2 - \frac{1}{2h} |t - s|^2) \exp(-\frac{\|y\|^2}{2\tau\sigma^2} - \rho s - \frac{\rho^2 h}{2}) dv dw \\ &\stackrel{(ii)}{\leq} \tau^{-n/2} \int_{\bar{\mathcal{K}}_\delta^c} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} \frac{1}{(2\pi h_\tau)^{n/2} (2\pi h)^{1/2}} \int_{\mathcal{H}(w)} \exp(-\frac{1}{2h_\tau} \|x - y\|^2 - \frac{1}{2h} |t - s|^2) dv dw \\ &\leq \tau^{-n/2} \int_{\bar{\mathcal{K}}_\delta^c} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} \left(\frac{1}{\sqrt{2\pi h}} \int_{d(w, \bar{\mathcal{K}})}^\infty \exp(-\frac{u^2}{2h}) du \right) dw =: (\#) \end{aligned}$$

where in (i) we used the change of variables via $y \leftarrow \tau y$ and $s \leftarrow s - \rho h$, and in (ii) $\mathcal{H}(w)$ denotes the supporting half-space at $\text{proj}_{\bar{\mathcal{K}}}(w)$ containing $\bar{\mathcal{K}}$ for given $w \in \bar{\mathcal{K}}_\delta^c$. By the co-area formula and integration by parts, for the n -dimensional Hausdorff measure \mathcal{H}^n ,

$$\begin{aligned} (\#) &\stackrel{\text{coarea}}{=} \tau^{-n/2} \int_\delta^\infty \Phi^c(h^{-1/2} u) \int_{\partial \bar{\mathcal{K}}_u} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} \mathcal{H}^n(dw) du \\ &\stackrel{\text{IBP}}{=} \tau^{-n/2} \left[\Phi^c(h^{-1/2} u) \int_0^u \int_{\partial \bar{\mathcal{K}}_l} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} \mathcal{H}^n(dw) dl \right]_{u=\delta}^\infty \\ &\quad + \tau^{-n/2} \int_\delta^\infty \frac{1}{\sqrt{2\pi h}} \exp(-\frac{u^2}{2h}) \int_0^u \int_{\partial \bar{\mathcal{K}}_l} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} \mathcal{H}^n(dw) dl du, \end{aligned}$$

and the double integral term is bounded as

$$\tau^{-n/2} \int_0^u \int_{\partial \bar{\mathcal{K}}_l} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} \mathcal{H}^n(dw) dl = \tau^{-n/2} \int_{\bar{\mathcal{K}}_u \setminus \bar{\mathcal{K}}} e^{-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s} dw$$

$$\stackrel{\text{Lemma 3.9}}{\leq} 2 \exp(hn^2 + 24un) \int_{\bar{\mathcal{K}}} e^{-\frac{1}{2\sigma^2} \|y\|^2 - \rho s} dy ds.$$

Using the tail bound on Φ^c in (2.6), $\tau^{-n/2} \Phi^c(h^{-1/2}u) \int_{\bar{\mathcal{K}}_u \setminus \bar{\mathcal{K}}} \exp(-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s) dw$ vanishes at $u = \infty$, and thus the first term of (#) can be bounded by 0. Hence,

$$\int_{\bar{\mathcal{K}}} e^{-\frac{1}{2\sigma^2} \|x\|^2 - \rho t} dx dt \cdot \mu^W(\tilde{\mathcal{K}}^c) \leq 2 \int_{\delta}^{\infty} \frac{1}{\sqrt{2\pi h}} \exp(-\frac{u^2}{2h}) \exp(hn^2 + 24un) du \int_{\bar{\mathcal{K}}} e^{-\frac{1}{2\sigma^2} \|x\|^2 - \rho t} dx dt.$$

Dividing both sides by $\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)$ and using the standard bound on Φ^c when $\delta \geq 23hn$,

$$\begin{aligned} \mu^W(\tilde{\mathcal{K}}^c) &\leq 2 \int_{\delta}^{\infty} \frac{1}{\sqrt{2\pi h}} e^{-\frac{s^2}{2h}} e^{hn^2 + 24sn} ds = 2e^{hn^2} \left(1 - \Phi\left(\frac{\delta - 24hn}{h^{1/2}}\right)\right) \\ &\leq \exp\left(-\frac{\delta^2}{2h} + 24\delta n + hn^2\right), \end{aligned}$$

which completes the proof. \square

This suggests taking $\delta \asymp 1/n$ and $h \asymp 1/n^2$ as in the exponential sampling. Precisely, under the choice of $h = \frac{c}{24^2 n^2}$ and $\delta = \frac{c/24+t}{n}$ for parameters $c, t > 0$, it follows that $\delta \geq 24hn$ and

$$\pi^Y(\tilde{\mathcal{K}}^c) \leq \exp\left(-\frac{t^2}{2c} + c\right).$$

We control how fast $\int_{\bar{\mathcal{K}}_u} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)$ increases in u .

Lemma 3.9. *In the setting of Lemma 3.8, for $\tau = \frac{\sigma^2}{h+\sigma^2} < 1$, $s > 0$, and $\rho \in (0, n]$,*

$$\tau^{-n/2} \int_{\bar{\mathcal{K}}_s} e^{-\frac{1}{2\tau\sigma^2} \|z\|^2 - \rho l} dz dl \leq 2 \exp(hn^2 + 24sn) \int_{\bar{\mathcal{K}}} e^{-\frac{1}{2\sigma^2} \|z\|^2 - \rho l} dz dl.$$

Proof. Consider when $\sigma^2 \leq 1/2n$. For $F : (z, l) \mapsto (\tau^{-1/2}z, l)$ between \mathbb{R}^{n+1} , and $\hat{\mathcal{K}} := \bar{\mathcal{K}} + \frac{V(x_0)}{n} - \frac{V_0}{n}$,

$$\begin{aligned} &\tau^{-n/2} \int_{\hat{\mathcal{K}}_s} \exp\left(-\frac{1}{2\tau\sigma^2} \|z\|^2 - \rho l\right) \\ &\stackrel{(i)}{\leq} \tau^{-n/2} \int_{(1+s)\hat{\mathcal{K}}} \exp\left(-\frac{1}{2\tau\sigma^2} \|z\|^2 - \rho l\right) = (1+s)^n \int_{F\hat{\mathcal{K}}} \exp\left(-\frac{(1+s)^2}{2\sigma^2} \|z\|^2 - \rho l(1+s)\right) \\ &\stackrel{(ii)}{\leq} e^{13sn} \int_{F\hat{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l\right) \\ &\stackrel{(iii)}{=} e^{13sn} \int_{\hat{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l\right) \times \frac{\mathbb{P}_{\mathcal{N}(0, \sigma^2 I_n) \times \text{Exp}}((X, T) \in F\hat{\mathcal{K}})}{\mathbb{P}_{\mathcal{N}(0, \sigma^2 I_n) \times \text{Exp}}((X, T) \in \hat{\mathcal{K}})} \\ &\stackrel{(iv)}{\leq} e^{13sn} \int_{\hat{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l\right) \times (\mathbb{P}_{\mathcal{N}(0, \sigma^2 I_n) \times \text{Exp}}(\|X\| \leq 1, T \geq 0))^{-1} \\ &\stackrel{(v)}{\leq} \frac{e^{13sn} e^{11sn}}{1 - d\sigma^2} \int_{\hat{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l\right) \leq 2e^{24sn} \int_{\hat{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l\right). \end{aligned}$$

In (i), we used $\hat{\mathcal{K}}_s \subset (1+s)\hat{\mathcal{K}}$ due to $B_1(0) \subset \hat{\mathcal{K}}$. (ii) follows from $l \geq -11$ in $F\hat{\mathcal{K}}$. In (iii), Exp denotes the 1-dim. exponential distribution with density proportional to $\exp(-\rho t)|_{[-t_0, \infty)}$ for $t_0 = \min_{\hat{\mathcal{K}}} t \geq -11$. (iv) and (v) follow from $B_1(0) \subset \hat{\mathcal{K}}$ and Chebyshev's inequality with $\rho \leq n$, respectively.

When $\sigma^2 \geq 1/2n$, starting from the (i)-line and using $\tau^{-n/2} \leq \exp(hn/2\sigma^2)$,

$$\begin{aligned} \tau^{-n/2} \int_{\tilde{\mathcal{K}}_s} \exp\left(-\frac{1}{2\tau\sigma^2} \|z\|^2 - \rho l\right) &\leq \exp\left(\frac{hn}{2\sigma^2} + sn\right) \int_{\tilde{\mathcal{K}}} \exp\left(-\frac{1}{2\tau\sigma^2} \|z\|^2 - \rho l(1+s)\right) \\ &\leq \exp(hn^2 + 12sn) \int_{\tilde{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma^2} \|z\|^2 - \rho l\right). \end{aligned}$$

We complete the proof by combining the two bounds in both cases. \square

Per-step guarantees. We provide the per-step query complexity and failure probability of implementing the backward step from a warm start.

Lemma 3.10. *In the setting of Problem 3.1, let ν be an M -warm initial distribution for $\mu^V = \mu_{\sigma^2, \rho}$ with $\sigma^2 > 0$ and $\rho \in (0, n]$. Given $k \in \mathbb{N}$ and $\eta \in (0, 1)$, set $Z = \frac{16kM}{\eta} (\geq 16)$, $h = \frac{1}{n^2} \frac{(\log \log Z)^2}{1200^2 \log Z}$ and $N = 2Z \log^2 Z$. Then, per iteration, the failure probability is at most η/k , and the expected number of queries is $\tilde{O}(M \log^2 \frac{k}{\eta})$.*

Proof. Since $\nu * \mathcal{N}(0, hI_{n+1})$ is M -warm with respect to μ^W due to $d\nu/d\mu^V \leq M$, the probability for the bad event per iteration is $\mathbb{E}_{\nu * \mathcal{N}(0, hI_{n+1})}[(1-\ell)^N] \leq M \mathbb{E}_{\mu^W}[(1-\ell)^N]$. For $\tilde{\mathcal{K}}$ given in (3.2),

$$\begin{aligned} \mathbb{E}_{\mu^W}[(1-\ell)^N] &= \int_{\tilde{\mathcal{K}}^c} \cdot + \int_{\tilde{\mathcal{K}} \cap [\ell \geq N^{-1} \log(3kM/\eta)]} \cdot + \int_{\tilde{\mathcal{K}} \cap [\ell \leq N^{-1} \log(3kM/\eta)]} \cdot \\ &\stackrel{(i)}{\leq} \mu^W(\tilde{\mathcal{K}}^c) + \int_{[\ell \geq N^{-1} \log(3kM/\eta)]} \exp(-\ell N) d\mu^W \\ &\quad + \int_{\tilde{\mathcal{K}} \cap [\ell \leq N^{-1} \log(3kM/\eta)]} \frac{\tau^{n/2} \ell(y, s) \exp(-\frac{1}{2\tau\sigma^2} \|y_\tau\|^2) \exp(-\rho s + \frac{1}{2} \rho^2 h)}{\int_{\tilde{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)} \\ &\stackrel{(ii)}{\leq} \exp\left(-\frac{t^2}{2c} + c\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} \frac{\int_{\tilde{\mathcal{K}}} \tau^{n/2} \exp(-\frac{1}{2\tau\sigma^2} \|y_\tau\|^2 - \rho s + \frac{1}{2} \rho^2 h)}{\int_{\tilde{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)} \\ &\stackrel{(iii)}{=} \exp\left(-\frac{t^2}{2c} + c\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} \frac{\tau^{-n/2} \int_{\tilde{\mathcal{K}}_\delta} \exp(-\frac{1}{2\tau\sigma^2} \|y\|^2 - \rho s - \frac{1}{2} \rho^2 h)}{\int_{\tilde{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t)} \\ &\stackrel{(iv)}{\leq} \exp\left(-\frac{t^2}{2c} + c\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} 2 \exp(hn^2 + 24\delta n) \\ &\stackrel{(v)}{\leq} \exp\left(-\frac{t^2}{4c}\right) + \frac{\eta}{3kM} + \frac{\log(3kM/\eta)}{N} 2 \exp(25t) \leq \frac{\eta}{kM}, \end{aligned}$$

where in (i) we used the density formula of μ^W in Lemma 3.7, (ii) follows from Lemma 3.8, (iii) is due to the change of variables, in (iv) we used Lemma 3.9, and (v) follows from the choices of $c = \frac{(\log \log Z)^2}{4 \cdot 24^2 \log Z}$, $t = \frac{1}{24} \log \log Z$, and $N = 2Z \log^2 Z$. Therefore, $\mathbb{E}_{\nu * \mathcal{N}(0, hI_{n+1})}[(1-\ell)^N] \leq \frac{\eta}{k}$.

The expected number of trials until the first acceptance is bounded by $M \mathbb{E}_{\mu^W}[\ell^{-1} \wedge N]$, where

$$\begin{aligned} \int_{\mathbb{R}^{n+1}} \frac{1}{\ell} \wedge N d\mu^W &\leq \int_{\tilde{\mathcal{K}}} \frac{1}{\ell} d\mu^W + N \mu^W(\tilde{\mathcal{K}}^c) \\ &\stackrel{(i)}{\leq} \frac{\int_{\tilde{\mathcal{K}}} \tau^{n/2} \exp(-\frac{1}{2\tau\sigma^2} \|y_\tau\|^2 - \rho s + \frac{1}{2} \rho^2 h) dy ds}{\int_{\tilde{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2} \|x\|^2 - \rho t) dx dt} + N \exp\left(-\frac{t^2}{2c} + c\right) \\ &\stackrel{(ii)}{\leq} 2 \exp(hn^2 + 24\delta n) + N \exp\left(-\frac{t^2}{2c} + c\right) \end{aligned}$$

$$\leq 4 \log^2 Z.$$

where (i) follows from Lemma 3.7 and Lemma 3.8, and in (ii) we repeat the arguments around (iii), (iv) above. Therefore, the expected number of queries per iteration is $\tilde{\mathcal{O}}(M \log^2 \frac{k}{\eta})$. \square

Combining the convergence rate and per-step guarantees of the PS_{ann} , we establish the query complexity of sampling from the annealing distribution from an M -warm start:

Theorem 3.11. *In the setting of Problem 3.1 without knowledge of x_0 and V_0 , for any given $\eta, \varepsilon \in (0, 1)$, $k \in \mathbb{N}$ defined below, PS_{ann} with $h = (1200^2 n^2 \log \frac{16kM}{\eta})^{-1}$, $N = \frac{32kM}{\eta} \log^2 \frac{16kM}{\eta}$, and initial distribution μ_0 which is M -warm with respect to $\mu := \mu_{\sigma^2, \rho}$ (where $\sigma > 0$ and $\rho \in (0, n]$) achieves $\mathcal{R}_{\infty}(\mu_k \parallel \mu) \leq \varepsilon$ after $k = \tilde{\mathcal{O}}(n^2(\sigma^2 \vee l^2) \log^2 \frac{MR}{\eta\varepsilon})$ iterations, where μ_k is the law of k -th iterate. With probability at least $1 - \eta$, the algorithm iterates k times successfully, using $\tilde{\mathcal{O}}(Mn^2(\sigma^2 \vee l^2) \log^4 \frac{R}{\eta\varepsilon})$ expected number of evaluation queries in total.*

Proof. By Lemma 3.5, PS_{ann} with given h should iterate $k = \tilde{\mathcal{O}}(n^2(\sigma^2 \vee l^2) \log^2 \frac{MR}{\eta\varepsilon})$ times to achieve ε -distance in \mathcal{R}_{∞} . Under the choice of h and N as claimed, each iteration succeeds with probability $1 - \eta/k$ and uses $\tilde{\mathcal{O}}(M \log^2 k/\eta)$ queries in expectation by Lemma 3.10. Therefore, throughout k iterations, the algorithm succeeds with probability $1 - \eta$, using $\tilde{\mathcal{O}}(Mn^2(\sigma^2 \vee l^2) \log^4 \frac{R}{\eta\varepsilon})$ queries. \square

3.3 Rényi-infinity sampling for logconcave distributions

Now that we have a sampler for the annealing distribution with \mathcal{R}_{∞} -guarantee, we analyze a query complexity of each phase in Tilted Gaussian Cooling.

Initialization. In the initialization where $\sigma^2 = 1/n$ and $\rho = 1/l$, we run PS_{ann} with initial distribution $\mu_0 = \exp(-\frac{n}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$ and target $\mu_1 \propto \exp(-\frac{n}{2} \|x\|^2 - t/l)|_{\bar{\mathcal{K}}}$. Recall that $l = \log \frac{2\varepsilon}{\varepsilon}$.

Lemma 3.12 (Initialization). *Rejection sampling for the initial $\mu_0 \propto \exp(-\frac{n}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$ takes $\mathcal{O}(l)$ queries in expectation. Given $\eta \in (0, 1)$, PS_{ann} with initial μ_0 , target $\mu_1 \propto \exp(-\frac{n}{2} \|x\|^2 - t/l)|_{\bar{\mathcal{K}}}$, $h^{-1} = \tilde{\Theta}(n^2 \log \frac{k}{\eta})$, $N = \tilde{\Theta}(k/\eta)$, and $k = \tilde{\Theta}(n^2 l^2 \log^4 \frac{MR}{\eta})$ iterates k times with probability at least $1 - \eta$, returning a sample with law $\bar{\mu}_1$ satisfying*

$$\text{ess sup}_{\mu_1} \left| \frac{d\bar{\mu}_1}{d\mu_1} - 1 \right| \leq 1,$$

where the expected number of evaluation queries used is $\tilde{\mathcal{O}}(n^2 l^2 \log^4 \frac{R}{\eta})$.

Proof. In the initialization through rejection sampling, the expected number of trials until the first acceptance is bounded by

$$\begin{aligned} \frac{\int_{\mathbb{R}^n} \exp(-\frac{n}{2} \|x\|^2) dx \cdot (15 + 13l)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2} \|x\|^2) dx dt} &\lesssim \frac{l \int_{\mathbb{R}^n} \exp(-\frac{n}{2} \|x\|^2) dx}{\int_{B_1(0)} \exp(-\frac{n}{2} \|x\|^2) dx} = l (\mathbb{P}_{\mathcal{N}(0, n^{-1}I_n)}(\|X\| \leq 1))^{-1} \\ &\stackrel{(i)}{\leq} l (1 - e^{-c\sqrt{n}})^{-1} = \mathcal{O}(l), \end{aligned}$$

where (i) follows from Paouris' lemma (Lemma A.8) for some universal constant c .

The warmness of μ_0 with respect to μ_1 is bounded as

$$\frac{d\mu_0}{d\mu_1} = \sup_{(x,t) \in \bar{\mathcal{K}}} \frac{\exp(-\frac{n}{2} \|x\|^2)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2} \|x\|^2) dx dt} \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2} \|x\|^2 - t/l) dx dt}{\exp(-\frac{n}{2} \|x\|^2 - t/l)}$$

$$\lesssim \sup_{\bar{\mathcal{K}}} \exp(t/l) \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2}\|x\|^2 - t/l) dx dt}{\int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2}\|x\|^2) dx} \lesssim 1.$$

Thus, PS_{ann} with initial μ_0 and target μ_1 can be run with target error of $\log 2$ in the \mathcal{R}_∞ -divergence, under a suitable choice of parameters given in Theorem 3.11. \square

Phase I. In the first phase ($\sigma^2 \in [\frac{1}{n}, 1]$ and $\rho \in [\frac{1}{l}, n]$), these parameters are updated according to

$$\sigma_{i+1}^2 = \sigma_i^2 \left(1 + \frac{1}{n}\right) \quad \& \quad \rho_{i+1} = \rho_i \left(1 + \frac{1}{ln}\right).$$

Note that PS_{ann} is initialized at $\bar{\mu}_i$ (not μ_i) for the target μ_{i+1} .

Lemma 3.13 (Phase I). *Given $\eta \in (0, 1)$, Phase I of Tilted Gaussian Cooling started at $\bar{\mu}_1$ succeeds with probability at least $1 - \eta$, returning a sample with law $\bar{\mu}_1$ satisfying*

$$\text{ess sup}_{\bar{\mathcal{K}}} \left| \frac{d\bar{\mu}_1}{d\mu_1} - 1 \right| \leq 1 \quad \text{for } \mu_1 \propto \exp(-\frac{1}{2}\|x\|^2 - nt)|_{\bar{\mathcal{K}}},$$

using PS_{ann} with suitable parameters. Phase I uses $\tilde{\mathcal{O}}(n^3 l^3 \log^5 \frac{R}{\eta})$ evaluation queries in expectation.

Proof. Since it takes at most ln many inner phases to double σ^2 and ρ , the total number of inner phases within Phase I is $m = \mathcal{O}(nl \log nl) = \tilde{\mathcal{O}}(nl)$.

The warmness between consecutive annealing distributions is bounded as

$$\begin{aligned} \frac{d\mu_i}{d\mu_{i+1}} &\leq \sup_{(x,t) \in \bar{\mathcal{K}}} \frac{\exp(-\frac{1}{2\sigma_i^2}\|x\|^2 - \rho_i t) \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma_{i+1}^2}\|x\|^2 - \rho_{i+1} t) dx dt}{\exp(-\frac{1}{2\sigma_{i+1}^2}\|x\|^2 - \rho_{i+1} t) \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma_i^2}\|x\|^2 - \rho_i t) dx dt} \\ &\leq \sup_{(x,t) \in \bar{\mathcal{K}}} \exp\left(\frac{\rho_i(t-t_0)}{ln}\right) \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma_{i+1}^2}\|x\|^2 - \rho_i(t-t_0)) dx dt}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma_i^2}\|x\|^2 - \rho_i(t-t_0)) dx dt} \\ &\stackrel{(i)}{\lesssim} \frac{(1 + \frac{1}{n})^{n/2} \int_{T\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma_i^2}\|x\|^2 - \rho_i(t-t_0)) dx dt}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma_i^2}\|x\|^2 - \rho_i(t-t_0)) dx dt} \leq \sqrt{e}, \end{aligned}$$

where in (i) we used $\rho_i \leq n$ and $t-t_0 \lesssim l$, and the change of variables via $T : (x, t) \mapsto ((1 + \frac{1}{n})^{-1/2}x, t)$.

Due to the design of the algorithm, PS_{ann} always ensures that $\text{ess sup}_{\mu_i} \left| \frac{d\bar{\mu}_i}{d\mu_i} - 1 \right| \leq 1$, and thus $\frac{d\bar{\mu}_i}{d\mu_{i+1}} \leq 2\sqrt{e}$. Hence, for each inner phase, PS_{ann} with target accuracy $\log 2$ in \mathcal{R}_∞ , target failure probability η/m , warmness $M = 2\sqrt{e}$, and suitable parameters succeeds with probability $1 - \eta/m$, returning a sample with law $\bar{\mu}_{i+1}$ such that

$$\text{ess sup}_{\bar{\mathcal{K}}} \left| \frac{d\bar{\mu}_{i+1}}{d\mu_{i+1}} - 1 \right| \leq 1,$$

and using $\tilde{\mathcal{O}}(n^2 l^2 \log^4 \frac{R}{\eta})$ many evaluation queries in expectation. Repeating this argument m many times, we complete the proof. \square

Phase II. In this second phase ($\sigma^2 \in [1, l^2 R^2]$), only σ^2 is updated according to

$$\sigma_{i+1}^2 = \sigma_i^2 \left(1 + \frac{\sigma_i^2}{l^2 R^2}\right).$$

Lemma 3.14 (Phase II). *Given $\eta \in (0, 1)$, Phase II of Tilted Gaussian Cooling started at $\bar{\mu}_1$ succeeds with probability at least $1 - \eta$, returning a sample with law $\bar{\mu}_1$ satisfying*

$$\text{ess sup}_{\bar{\mathcal{K}}} \left| \frac{d\bar{\mu}_1}{d\mu_1} - 1 \right| \leq 1 \quad \text{for } \mu_1 \propto \exp\left(-\frac{1}{2l^2R^2} \|x\|^2 - nt\right) \Big|_{\bar{\mathcal{K}}},$$

using PS_{ann} with suitable parameters. Phase II uses $\tilde{\mathcal{O}}(n^2R^2l^4 \log^4 \frac{1}{\eta})$ queries in expectation.

Proof. We note that for given σ^2 , it takes at most l^2R^2/σ^2 inner phases to double σ^2 , so the total number of inner phases within Phase II is $m = \tilde{\mathcal{O}}(l^2R^2)$.

The warmness between consecutive annealing distributions is bounded as

$$\begin{aligned} \frac{d\mu_i}{d\mu_{i+1}} &\leq \sup_{(x,t) \in \bar{\mathcal{K}}} \frac{\exp\left(-\frac{1}{2\sigma_i^2} \|x\|^2 - nt\right)}{\exp\left(-\frac{1}{2\sigma_{i+1}^2} \|x\|^2 - nt\right)} \frac{\int_{\bar{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma_{i+1}^2} \|x\|^2 - nt\right) dx dt}{\int_{\bar{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma_i^2} \|x\|^2 - nt\right) dx dt} \\ &\leq \frac{\int_{\bar{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma_{i+1}^2} \|x\|^2 - nt\right) dx dt}{\int_{\bar{\mathcal{K}}} \exp\left(-\frac{1}{2\sigma_i^2} \|x\|^2 - nt\right) dx dt} \leq \sqrt{e}, \end{aligned}$$

where the last line holds, since $\|x\|^2 \leq l^2R^2$ on $\bar{\mathcal{K}}$,

$$\exp\left(-\frac{1}{2\sigma_{i+1}^2} \|x\|^2\right) = \exp\left(-\frac{1}{2\sigma_i^2} \|x\|^2\right) \exp\left(\frac{1}{2(l^2R^2 + \sigma_i^2)} \|x\|^2\right) \leq \sqrt{e} \exp\left(-\frac{1}{2\sigma_i^2} \|x\|^2\right).$$

As in the analysis for Phase I, this implies that due to the design of the algorithm,

$$\frac{d\bar{\mu}_i}{d\mu_{i+1}} \leq 2\sqrt{e}.$$

Hence, for each inner phase, PS_{ann} with target accuracy $\log 2$ in \mathcal{R}_∞ , target failure probability η/m , warmness $M = 2\sqrt{e}$, and suitable parameters succeeds with probability $1 - \eta/m$, returning a sample with law $\bar{\mu}_{i+1}$ such that

$$\text{ess sup}_{\bar{\mathcal{K}}} \left| \frac{d\bar{\mu}_{i+1}}{d\mu_{i+1}} - 1 \right| \leq 1,$$

and using $\tilde{\mathcal{O}}(n^2(\sigma^2 \vee l^2) \log^4 \frac{R}{\eta})$ many evaluation queries in expectation. Each doubling part requires

$$\tilde{\mathcal{O}}\left(\frac{l^2R^2}{\sigma^2} \times n^2(\sigma^2 \vee l^2) \log^4 \frac{R}{\eta}\right) = \tilde{\mathcal{O}}\left(n^2R^2l^4 \log^4 \frac{1}{\eta}\right)$$

many queries in expectation. Therefore, the claim follows from multiplying this by $\log_2(lR)$. \square

Termination. In the last phase, PS_{exp} is run with initial distribution $\bar{\mu}_1$ and target $\bar{\pi} \propto e^{-nt} \Big|_{\bar{\mathcal{K}}}$.

Lemma 3.15 (Termination). *Given $\eta, \varepsilon \in (0, 1)$, PS_{exp} with initial distribution $\bar{\mu}_1$, $h^{-1} = \tilde{\Theta}(n^2 \log \frac{k}{\eta})$, $N = \tilde{\Theta}(k/\eta)$, and $k = \tilde{\Theta}(n^2R^2l^2 \log^4 \frac{M}{\eta\varepsilon})$ iterates k times with probability at least $1 - \eta$, returning a sample with law ν satisfying*

$$\text{ess sup}_{\bar{\pi}} \left| \frac{d\nu}{d\bar{\pi}} - 1 \right| \leq \varepsilon,$$

where the expected number of evaluation queries is $\tilde{\mathcal{O}}(n^2R^2l^2 \log^4 \frac{1}{\eta\varepsilon})$.

Proof. As $\|x\|^2 \leq l^2 R^2$ on $\bar{\mathcal{K}}$, the warmness between $\bar{\mu}_{\parallel}$ and $\bar{\pi}$ is bounded as

$$\frac{d\bar{\mu}_{\parallel}}{d\bar{\pi}} \leq \sup_{(x,t) \in \bar{\mathcal{K}}} \frac{\exp(-\frac{1}{2l^2 R^2} \|x\|^2 - nt)}{\exp(-nt)} \frac{\int_{\bar{\mathcal{K}}} \exp(-nt) dx dt}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2l^2 R^2} \|x\|^2 - nt) dx dt} \leq \sqrt{e},$$

and thus $\frac{d\bar{\mu}_{\parallel}}{d\bar{\pi}} \leq 2\sqrt{e}$. Therefore, by Lemma 3.6, PS_{exp} with initial $\bar{\mu}_{\parallel}$ and suitable parameters above outputs a sample with law ν such that

$$\text{ess sup}_{\bar{\mathcal{K}}} \left| \frac{d\nu}{d\bar{\pi}} - 1 \right| \leq \varepsilon,$$

using $\tilde{\mathcal{O}}(n^2 R^2 l^2 \log^4 \frac{1}{\eta \varepsilon})$ evaluation queries in expectation. \square

Final guarantee. As claimed in Theorem 1.5, we prove that Tilted Gaussian Cooling provides an $\mathcal{O}(1)$ -warm start for $\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}$, using $\tilde{\mathcal{O}}(n^2(R^2 \vee n))$ evaluation queries in expectation.

Theorem 3.16 (Warm-start generation). *In the setting of Problem 3.1, consider $\pi \propto \exp(-nt)|_{\mathcal{K}}$ given in (exp-red). For given $\eta, \varepsilon \in (0, 1)$, Tilted Gaussian Cooling succeeds with probability at least $1 - \eta$, returning a sample with law ν such that $\mathcal{R}_{\infty}(\nu \parallel \pi) \leq \varepsilon$, using $\tilde{\mathcal{O}}(n^2(R^2 \vee n) \log^8 1/\eta \varepsilon)$ evaluation queries in expectation. In particular, if π^X is well-rounded (i.e., $R^2 = \mathcal{O}(n)$), then we need $\tilde{\mathcal{O}}(n^3 \log^8 1/\eta \varepsilon)$ queries in expectation.*

Proof. Combining the four lemmas of this section with $\eta/4$ and $\varepsilon/3$ in place of η and ε , we conclude that by the union bound, the failure probability is at most η . Next, the choice of $l = \mathcal{O}(\log \frac{1}{\varepsilon})$ leads to

$$\sup_{\mathcal{K}} \frac{\bar{\pi}}{\pi} \leq \frac{1}{\mathbb{P}_{\pi}(Z \in \bar{\mathcal{K}})} \leq 1 + \frac{2\varepsilon}{3}.$$

Therefore, the claim follow from

$$\mathcal{R}_{\infty}(\nu \parallel \pi) \leq \mathcal{R}_{\infty}(\nu \parallel \bar{\pi}) + \mathcal{R}_{\infty}(\bar{\pi} \parallel \pi) \leq \frac{\varepsilon}{3} + \log(1 + \frac{2\varepsilon}{3}) \leq \varepsilon. \quad \square$$

4 Application 1: Rounding logconcave distributions

We have shown that the query complexity of generating a warm start (in fact, sampling from π^X with \mathcal{R}_{∞} -guarantee) is $\tilde{\mathcal{O}}(n^2(R^2 \vee n))$. However, R could be arbitrarily large so this query complexity is not actually polynomial in the problem parameters. Hence, prior work combines sampling algorithms with a *rounding* procedure, which makes π^X less skewed (e.g., $R^2 = \text{poly } n$). For instance, near-isotropic rounding makes $\text{Cov } \pi^X$ nearly the identity, under which $\mathbb{E}_{\pi^X}[\|X - \mathbb{E}X\|^2] \approx n$.

Problem 4.1 (Isotropic rounding). Assume access to a well-defined function oracle $\text{Eval}_{x_0, R}(V)$ for convex $V : \mathbb{R}^n \rightarrow \bar{\mathbb{R}}$. Given $s \geq 1$, what is the query complexity of finding an affine transformation $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $F_{\#} \pi^X$ satisfies

$$s^{-1} I_n \preceq \text{Cov } F_{\#} \pi^X = F[\text{Cov } \pi^X] F^{\top} \preceq s I_n.$$

We may assume that $\min V = 0$, since it can be found using the evaluation oracle when x_0 is given. This problem is equivalent to estimating the covariance of π^X accurately, since setting $F = [\text{Cov } \pi^X]^{-1/2}$ leads to $\text{Cov } F_{\#} \pi^X = I_n$. The prior best complexity of isotropic rounding is $\tilde{\mathcal{O}}(n^4)$ by [LV06a], and in this section we improve this to $\tilde{\mathcal{O}}(n^{3.5})$, as claimed in Theorem 1.7.

Theorem 4.2. *In the setting of Problem 4.1, there exists a randomized algorithm with query complexity of $\tilde{\mathcal{O}}(n^{3.5} \text{polylog } R)$ that finds an affine map $F^X : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $F^X \pi^X$ is 1.01-isotropic with probability at least $1 - \mathcal{O}(n^{-1/2})$.*

Setup. By translation, we assume throughout this section that $x_0 = 0$ and that $\mathcal{K} = \{(x, t) \in \mathbb{R}^{n+1} : V(x) - 11n \leq nt\}$ contains $B_1^{n+1}(0)$. We now introduce several notation for a concise description. In the first part of the algorithm, we mainly work with $d\nu^X \propto \exp(-V)|_{\mathbb{L}_{\pi^X, g}} dx$ truncated to the ground set $\mathbb{L}_{\pi^X, g}$, called the *grounded distribution*. One can think of it as a core of π^X in that (1) ν^X is 1.01-warm with respect to π^X (i.e., $\mathcal{R}_\infty(\nu^X \parallel \pi^X) \leq \log 1.01$), and (2) the values of V over the support of ν^X is within $10n$ from $\min V = 0$ (i.e., $\max_{\text{supp } \nu^X} V \leq 10n$). The first property immediately follows from

$$\sup_{\mathbb{L}_{\pi^X, g}} \frac{\nu^X}{\pi^X} \leq \frac{\int \exp(-V)}{\int_{\mathbb{L}_{\pi^X, g}} \exp(-V)} = (\mathbb{P}_{\pi^X}(X \in \mathbb{L}_{\pi^X, g}))^{-1} \leq 1.01, \quad (4.1)$$

where the last inequality is due to Lemma A.6 (with $\beta = 10$). We also denote by $\nu_r^X \propto \exp(-V)|_{\mathbb{L}_{\pi^X, g} \cap B_r(0)}$ the grounded distribution truncated to the ball $B_r(0)$. Hereafter, we use \mathbb{L}_g to indicate $\mathbb{L}_{\pi^X, g}$ if there is no confusion.

For each of π^X, ν^X, ν_r^X , its exponential counterpart can be written as, for the cylinders $C_g := \mathbb{L}_g \times \mathbb{R}$ and $C_r := B_r(0) \times \mathbb{R}$,

$$\pi(x, t) \propto \exp(-nt)|_{\mathcal{K}}, \quad \nu(x, t) \propto \exp(-nt)|_{\mathcal{K} \cap C_g}, \quad \nu_r(x, t) \propto \exp(-nt)|_{\mathcal{K} \cap C_g \cap C_r}.$$

We use $F^X : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ to denote an affine transformation and its embedding into \mathbb{R}^{n+1} defined by $F(x, t) := (F^X(x), t)$, respectively. Then, for $W := V \circ (F^X)^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}$,

$$\begin{aligned} F_{\#}\pi(x, t) &\propto \exp(-nt)|_{FK} \text{ where } FK = \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : W(x) \leq nt\}, \\ F_{\#}\nu(x, t) &\propto \exp(-nt)|_{FK \cap FC_g} \text{ where } FC_g = F^X \mathbb{L}_g \times \mathbb{R}, \\ F_{\#}\nu_r(x, t) &\propto \exp(-nt)|_{FK \cap FC_g \cap FC_r} \text{ where } FC_r = F^X B_r(0) \times \mathbb{R}. \end{aligned}$$

Note that $(F_{\#}\pi)^X = F_{\#}^X \pi^X \propto \exp(-W)$, $(F_{\#}\nu)^X = F_{\#}^X \nu^X \propto \exp(-W)|_{F^X \mathbb{L}_g}$, and $(F_{\#}\nu_r)^X = F_{\#}^X \nu_r^X \propto \exp(-W)|_{F^X(\mathbb{L}_g \cap B_r)}$. Lastly, it clearly holds that for $\mu \in \{\pi, \nu, \nu_r\}$,

$$\text{Cov } F_{\#}\mu = \begin{bmatrix} \text{Cov } F_{\#}^X \mu^X & * \\ * & \text{Cov } \mu^T \end{bmatrix}.$$

Roadmap to isotropy. At the high level, we gradually make π^X less skewed through an iterative process of sampling, estimating an approximate covariance matrix, and applying a suitable affine transformation. Assuming that Algorithm 4 (Isotropize) can solve Problem 4.1 when $R^2 = \mathcal{O}(n)$ by using $\tilde{\mathcal{O}}(n^3)$ many queries, we provide a brief overview of the entire rounding algorithm. Similar to our warm-start generation algorithm, for $\delta := 1 + n^{-1/2} > 1$ and $D := 2R \log 400$, this algorithm gradually isotropizes a sequence $\{\mu_i\}$ of distributions, $\nu_1^X \rightarrow \nu_\delta^X \rightarrow \nu_{\delta^2}^X \rightarrow \dots \rightarrow \nu_D^X \rightarrow \nu^X \rightarrow \pi^X$, passing along an affine map F^X such that if $F_{\#}^X \mu_i$ is near-isotropic, then $H_{\#}^X \mu_{i+1}$ (where H^X is a slight modification of F^X) satisfies regularity conditions:

- (LR) Lower regularity: the ground set of $H_{\#}^X \mu_{i+1}$ contains $B_1^n(x_0)$ for some x_0 .
- (UR) Upper regularity: $\mathbb{E}_{H_{\#}^X \mu_{i+1}} [\|X - x_0\|^2] \leq R^2 = \mathcal{O}(n)$.

This ensures that sampling from $H_{\#}^X \mu_{i+1}$ can be achieved with moderate complexity. We now provide a summary of the rounding algorithm:

1. Step 1: Make ν_D^X near-isotropic.

- (a) Run **Isotropize** on ν_1^X (which clearly satisfies (LR) and (UR) with $R^2 = \mathcal{O}(1)$) to obtain an affine map F^X such that $F_{\#}^X \nu_1^X$ is near-isotropic.
 - (b) Repeat while $r \leq D = \mathcal{O}(R)$: if $F_{\#}^X \nu_r^X$ is near-isotropic, then run **Isotropize** on $(4F^X)_{\#} \nu_{\delta r}^X$, and set $F^X \leftarrow 4F^X$ and $r \leftarrow \delta r$.
 - i. Justified by Lemma 4.12: If $F_{\#}^X \nu_r^X$ is near-isotropic, then $(4F^X)_{\#} \nu_{\delta r}^X$ satisfies (LR) and (UR) with $R^2 = \mathcal{O}(n)$.
2. Step 2: Given a near-isotropic $F_{\#}^X \nu_D^X$, make ν^X near-isotropic.
- (a) Run **Isotropize** on $(4F^X)_{\#} \nu^X$ to obtain a new map F^X that makes $F_{\#}^X \nu^X$ near-isotropic.
 - i. Justified by Lemma 4.14: If $F_{\#}^X \nu_D^X$ is near-isotropic, then $(4F^X)_{\#} \nu^X$ satisfies (LR) and (UR) with $R^2 = \mathcal{O}(n)$.
3. Step 3: Given a near-isotropic $F_{\#}^X \nu^X$, make π^X near-isotropic.
- (a) Draw $\tilde{\mathcal{O}}(n)$ samples from $(4F^X)_{\#} \pi^X$ to estimate its covariance accurately.
 - i. Justified by Lemma 4.15: If $F_{\#}^X \nu^X$ is near-isotropic, then $(4F^X)_{\#} \pi^X$ satisfies (LR) and $\|\text{Cov} F_{\#}^X \pi^X\| = \mathcal{O}(1)$.

Step 1 and 2 would use $\tilde{\mathcal{O}}(n^3 \times \sqrt{n} \log D) = \tilde{\mathcal{O}}(n^{3.5} \log D)$ queries, as **Isotropize** is called $\mathcal{O}(\sqrt{n} \log D)$ times in Step 1 and 2. In Step 3, as the PS_{exp} with an $\mathcal{O}(1)$ -warm start requires $\tilde{\mathcal{O}}(n^2)$ queries per sample, Step 3 uses $\tilde{\mathcal{O}}(n^3)$ queries in total. Therefore, the final complexity of this algorithm would be $\tilde{\mathcal{O}}(n^{3.5} \text{polylog } R/\eta)$ for a target failure probability η .

Remark 4.3 (Failure of samplers). Recall that the mixing time of PS_{exp} has a poly-logarithmic dependence on the target failure probability $\eta > 0$. Since the number k of iterations (of PS_{exp}) throughout the algorithm is polynomial in the problem parameters (e.g., n, R), we can, for instance, simply set $\eta = 1/(kn^{100})$ without affecting the final query complexity by more than polylogarithmic factors.

4.1 Isotropic rounding of a well-rounded distribution

In this section, we analyze the aforementioned algorithm **Isotropize**, which finds an affine map F^X making a well-rounded distribution near-isotropic. We state an exact setup for **Isotropize**.

Assumption 4.4. *Given access to an evaluation oracle for convex $U : \mathbb{R}^n \rightarrow \bar{\mathbb{R}}$ with $\min U \geq 0$, and distribution $d\mu^X \propto \exp(-U) dx$, assume that $B_1^n(x_0) \subset L_{\mu^X, g} = \{x \in \mathbb{R}^n : U(x) - \min U \leq 10n\}$ for known x_0 and $\mathbb{E}_{\mu^X}[\|X - x_0\|^2] \leq R^2 = C^2 n$ with a known constant $C \geq 1$, and that $U(x) \leq 10n$ for any $x \in \text{supp } \mu^X$ (i.e., $\text{supp } \mu^X = L_{\mu^X, g}$).*

We note that $\mu(x, t) \propto \exp(-nt)|_{\mathcal{S}}$ for $\mathcal{S} := \{(x, t) : U(x) - 11n \leq nt\}$ has an important property that the slice of $\text{supp } \mu$ at $t = -1$ recovers $\text{supp } \mu^X$, which simply follows from $\min U \geq \min V = 0$. Also, this setup can represent $F_{\#}^X \nu_r^X$ and $F_{\#}^X \nu^X$. For instance, $\mu^X = F_{\#}^X \nu_r^X$ can be obtained by setting $U = V \circ (F^X)^{-1} / \mathbf{1}[x \in F^X(L_g \cap B_r(0))]$.

Rounding algorithm, Isotropize. This algorithm generalizes the approach from [JLLV21], a rounding algorithm for well-rounded uniform distributions over a convex body, following a modified version of their algorithm studied in [KZ24] for streamlined analysis.

Algorithm 4 begins with Tilted Gaussian Cooling to generate an $\mathcal{O}(1)$ -warm start for μ , using n^3 queries (Theorem 3.16). The while-loops then proceeds as follows: for $r := \text{inrad } \mu^X$, the largest

Algorithm 4 Isotropize

Input: distribution $\mu \propto \exp(-nt)|_{\{(x,t):U(x)-11n \leq nt\}}$ over \mathbb{R}^{n+1} such that $B_1(0) \subset \text{supp } \mu^X = \mathcal{L}_{\mu^X, g}$ and $\mathbb{E}_{\mu^X}[\|X\|^2] \leq R^2 = \mathcal{O}(n)$.

Output: near-isotropic distribution $F_{\#}\mu_{\text{last}}$.

- 1: Run Tilted Gaussian Cooling to obtain $Z_0 = (X_0, T_0) \in \mathbb{R}^{n+1}$ with $\mathcal{R}_\infty(\text{law } Z_0 \parallel \mu) \leq \log 2$.
 - 2: Let $r_1 = 1$, $\mu_1 := \mu$, and $i = 1$.
 - 3: **while** $r_i^2 \lesssim n$ **do**
 - 4: For $k_i \asymp r_i^2 \text{polylog } n$, draw $\{Z_j = (X_j, T_j)\}_{j \in [k_i]} \leftarrow \text{PS}_{N_i}(\mu_i, \delta_{Z_0}, h_i)$ with $N_i \asymp r_i^{-2} n^3 \log n$ and $h_i \asymp r_i^2 (n^2 \log(r_i n))^{-1}$.
 - 5: Estimate $\bar{m}_i = \frac{1}{k_i} \sum_{j=1}^{k_i} X_j$ and $\bar{\Sigma}_i = \frac{1}{k_i} \sum_{j=1}^{k_i} (X_j - \bar{m}_i)^{\otimes 2}$.
 - 6: Compute $M_i = I_n + P_i$, where P_i is the orthogonal projection to the subspace spanned by eigenvectors of $\bar{\Sigma}_i$ with eigenvalue at most n . Define $M_i^*(x, t) := (M_i x, t)$.
 - 7: Set $\mu_{i+1} = (M_i^*)_{\#}\mu_i$, $X_0 \leftarrow M_i X_0$, $r_{i+1} = 2r_i(1 - 1/\log n)$, and $i \leftarrow i + 1$.
 - 8: **end while**
 - 9: Draw $\Theta(n \text{polylog } n)$ samples via $\text{PS}_N(\mu_{\text{last}}, \delta_{Z_0}, h)$ with $N \asymp n^2 \text{polylog } n$ and $h \asymp (n \log n)^{-1}$, and use their X -part to estimate the mean \bar{m} and covariance $\bar{\Sigma}$. For $F(x, t) := (\bar{\Sigma}^{-1/2}(x - \bar{m}), t)$, return $F_{\#}\mu_{\text{last}}$.
-

radius of a ball contained in $\text{supp } \mu^X = \mathcal{L}_{\mu^X, g}$, we **can** run the sampler PS_{exp} (with the $\mathcal{O}(1)$ -warm start and target failure probability $n^{-\mathcal{O}(1)}$) to generate r^2 samples distributed according to μ , since under Assumption 4.4, μ^X meets all conditions required by Problem 2.1. We then estimate an approximate covariance matrix $\bar{\Sigma}$ of μ^X (using these r^2 samples) such that $|\bar{\Sigma} - \Sigma| \lesssim nI_n$, where $\Sigma = \text{Cov } \mu^X$. This procedure requires $r^2 \times n^2 \|\Sigma\|/r^2 = n^2 \|\Sigma\|$ many queries in expectation. Next, we compute the eigenvalue/vectors of $\bar{\Sigma}$ and double the subspace spanned by the eigenvectors with eigenvalues less than n . One iteration of this inner loop achieves two key properties: (P1) the largest eigenvalue of the covariance Σ increases by at most n , and (P2) $r = \text{inrad } \mu^X$ almost doubles.

This algorithm repeats this inner loop (i.e., sampling–estimation–scaling) until r^2 reaches n . As r almost doubles every inner loop, the inner loop iterates at most $\mathcal{O}(\log n)$ times. Since the operator norm of the initial covariance matrix is $\mathcal{O}(n)$ due to Assumption 4.4, the operator norm will remain less than $\tilde{\mathcal{O}}(n)$ throughout the algorithm, and thus the query complexity of the inner loop is bounded by $\tilde{\mathcal{O}}(n^3)$.

Covariance estimation using dependent samples. We now specify how we can accurately estimate the target covariance Σ using r^2 many *dependent* samples drawn by the PS_{exp} . Letting P be the Markov kernel of the PS_{exp} with target $\mu(x, t) \propto \exp(-nt)|_{\mathcal{S}}$ and step size $h \asymp r^2/n^2$, we showed in Lemma 2.9 that PS has contraction in χ^2 -divergence, i.e., for any distribution $\nu \ll \mu$,

$$\chi^2(\nu P \parallel \mu) \leq \frac{\chi^2(\nu \parallel \mu)}{(1 + h/C_{\text{PI}}(\mu))^2},$$

which implies that the spectral gap of this Markov chain is at least $hC_{\text{PI}}^{-1} = \Omega(r^2 n^{-2} C_{\text{PI}}^{-1})$. Hence, a new Markov chain with kernel P^N for $N \asymp r^{-2} n^2 C_{\text{PI}}$ (i.e., composition of the PS_{exp} N -times) has a spectral gap at least 0.99 (arbitrarily close to 1). Hereafter, $\text{PS}_N(\mu, \nu, h)$ denotes a Markov chain with kernel P^N , initial distribution ν , and target distribution μ .

When the PS_{exp} samples from μ in Line 4, it takes a sample every N iteration; in other words, it draws $\Theta(r_i^2 \text{polylog } n)$ consecutive samples from the Markov chain with kernel P^N . Then, the following result allows us to estimate a covariance matrix of μ with a provable guarantee in Line 5 when using dependent samples under a spectral-gap condition:

Lemma 4.5 ([KZ24, Theorem 8]). *Let P be a reversible Markov chain on \mathbb{R}^n with stationary distribution μ and spectral gap λ . Let $(Z_i)_{i \in [k]}$ be a sequence of outputs given by P with initial distribution μ . If μ is logconcave and $\lambda \geq 0.99$, then the covariance estimator $\bar{\Sigma} = k^{-1} \sum_{i=1}^k (Z_i - \bar{\mathbf{m}})^{\otimes 2}$ for $\bar{\mathbf{m}} = k^{-1} \sum_{i=1}^k Z_i$ satisfies that for any $\varepsilon > 0$ and $\delta \in (0, n]$, with probability at least $1 - \mathcal{O}(n^{-1})$,*

$$|\bar{\Sigma} - \Sigma| \preceq \varepsilon \Sigma + \delta I_n \quad \text{for } \Sigma = \text{Cov } \mu,$$

so long as $k \asymp \frac{\text{tr } \Sigma}{\varepsilon \delta} \log^2 \frac{n \max(n, \text{tr } \Sigma)}{\delta} \log^2 n$.

We also rely on another result to justify the covariance estimation in Line 9, when computing the mean $\bar{\mathbf{m}}$ and covariance $\bar{\Sigma}$ of μ_{last} .

Lemma 4.6 ([KZ24, Corollary 10 and 33]). *In the setting of Lemma 4.5, $\bar{\Sigma}$ satisfies that for any $\varepsilon \in (0, 1)$, with probability at least $1 - \mathcal{O}(n^{-1})$,*

$$|\bar{\Sigma} - \Sigma| \preceq \varepsilon \Sigma \quad \text{and} \quad \|\bar{\mathbf{m}} - \mathbb{E}_\mu Z\|^2 \leq \varepsilon^2 \|\Sigma\|,$$

so long as $k \asymp \frac{n}{\varepsilon^2} \log^2 \frac{n}{\varepsilon} \log^2 n$.

We note that even when the chain starts from a non-stationary distribution ν , these statistical results still hold, with the bad probability increasing by a multiplicative factor of $\exp(\mathcal{R}_2(\nu \parallel \mu)) = 1 + \chi^2(\nu \parallel \mu)$ (see [KZ24, Lemma 29]). As PS_{exp} starts from an $\mathcal{O}(1)$ -warm start and $\mathcal{R}_2 \leq \mathcal{R}_\infty$, the bad probability only increases by a factor of $\mathcal{O}(1)$.

Analysis. We now examine the two key properties: (P1) the largest eigenvalue of the covariance of μ_i^X increases by at most $\tilde{\mathcal{O}}(n)$, and (P2) $r_i = \text{inrad } \mu_i^X$ nearly doubles every inner loop. [JLLV21] established these results regarding a rounding algorithm for the *uniform distribution* over a convex body. Notably, the proofs therein are independent of target distributions, allowing these results to hold for logconcave distributions as well. Here, we present an outline of these proofs, along with pointers to detailed arguments in previous works, following a streamlined analysis in [KZ24, §4.1].

We first state that PS_{exp} employed in Algorithm 4 has a spectral gap at least 0.99. Its proof is obvious from the discussion in the paragraph above.

Proposition 4.7 ([KZ24, Lemma 14]). *In Line 4 and 9, $\text{PS}_{N_i}(\mu_i, \cdot, h_i)$ with $N_i \asymp r_i^{-2} n^3 \log n$ and $h_i \asymp r_i^2 (n^2 \log(r_i n))^{-1}$ has a spectral gap at least 0.99.*

As for (P1), we need the following quantitative bounds on the operator norm and trace of $\Sigma_i^X = \text{Cov } \mu_i^X$ in each while loop of Algorithm 4. The original result was established by [JLLV21, Lemma 3.2], with a simpler proof shown in [KZ24, Lemma 16].

Lemma 4.8. *In Algorithm 4, the while-loop iterates at most $\mathcal{O}(\log n)$ times, during which $\text{tr } \Sigma_i^X \lesssim r_i^2 n$ and $\|\Sigma_i^X\| \lesssim n(1+i)$.*

The base case $i = 1$ holds since $\|\Sigma_1^X\| \leq \text{tr } \Sigma_1^X \leq R^2 = \mathcal{O}(n)$. Assuming the results hold for i as the induction hypothesis, one must need a guarantee on how close the covariance estimation in Line 5 is to the true covariance, in order to establish the claim for $i + 1$. The following lemma directly follows from Lemma 4.5 and Proposition 4.7.

Proposition 4.9 ([KZ24, Lemma 15]). *Each while-loop ensures that with probability at least $1 - \mathcal{O}(n^{-1})$, for $\Sigma_i^X = \text{Cov } \mu_i^X$ and $\Sigma_i = \text{Cov } \mu_i$,*

$$\frac{9}{10} \Sigma_i^X - \frac{n+1}{100} I_{n+1} \preceq \bar{\Sigma}_i \preceq \frac{11}{10} \Sigma_i^X + \frac{n+1}{100} I_{n+1} \quad \text{if } k_i \gtrsim \frac{\text{tr } \Sigma_i}{n} \text{polylog } n.$$

Since $\text{tr } \Sigma_i \leq \text{tr } \Sigma_i^X + \mathcal{O}(1)$ by Lemma 2.5 and $\text{tr } \Sigma_i^X \lesssim r_i^2 n$ from the induction hypothesis, the result above holds when $k_i \gtrsim r_i^2 \text{polylog } n$, thereby justifying the choice of k_i in Line 4. With this statistical estimation, the proof of Lemma 4.8 for $i + 1$ is exactly the same with [KZ24, Lemma 16].

Regarding (P2), we need minor adjustments to some constants in the proof of [KZ24, Lemma 17], whose original version (for the uniform distribution) appeared in [JLLV21, Lemma 3.2].

Lemma 4.10. *In Algorithm 4, under $r_{i+1} = 2(1 - 1/\log n)r_i$, each while-loop ensures $r_{i+1} \leq \text{inrad } \mu_{i+1}^X$.*

To demonstrate the existence of a ball of radius r_{i+1} centered at some point $c_{i+1} \in \text{supp } \mu_{i+1}^X$, the proof relies on two ellipsoids within $\text{supp } \mu_i^X$: a ball $B_{r_i}(c_i)$ from the induction hypothesis, and $\{\|x - \mathbf{m}_i\|_{(\Sigma_i^X)^{-1}} \leq e^{-2}\}$ for $\mathbf{m}_i = \mathbb{E}_{\mu_i^X} X$, whose existence follows from Lemma A.10. After multiplying $x \mapsto (I_n + P_i)x = M_i x$, for $c'_i := M_i c_i$ and $\mathbf{m}'_i := M_i \mathbf{m}_i$, the support of the pushforward $(M_i)_\# \mu_i^X = \mu_{i+1}^X$ contains two ellipsoids:

$$\mathcal{A} : \{(x - c'_i)^\top M_i^{-2}(x - c'_i) \leq r_i^2\}, \quad \text{and} \quad \mathcal{B} : \{(x - \mathbf{m}'_i)^\top (M_i \Sigma_i^X M_i)^{-1}(x - \mathbf{m}'_i) \leq e^{-2}\}.$$

Let \mathcal{S} be the subspace spanned by the eigenvectors of $\bar{\Sigma}_i$ with corresponding eigenvalues less than n . The transformation M_i scales up \mathcal{S} by a factor of two, ensuring that the \mathcal{A} -ellipsoid contains a $2r_i$ -ball centered at c'_i along \mathcal{S} . Moreover, as eigenvalues of $\bar{\Sigma}_i$ along \mathcal{S}^\perp are at least n , we can show that the \mathcal{B} -ellipsoid contains an $(n^{1/2}/10)$ -ball at \mathbf{m}'_i along \mathcal{S}^\perp . Using the convexity of $\text{supp } \mu_{i+1}^X$, one can show that the convex combination of these two ellipsoids leads to the existence of a ball of radius $2(1 - (\log n)^{-1})r_i$ at some point. We refer readers to [KZ24, Lemma 17] for details.

It is worth noting that $\mu_{i+1}^X \propto \exp(-W)$ for a new potential W satisfies that $W(x) - \min W = W(x) - \min U \leq W(x) \leq 10n$ for any $x \in \text{supp } \mu_{i+1}^X$ (i.e., $\text{supp } \mu_{i+1}^X = \mathcal{L}_{\mu_{i+1}^X, g}$), and this ground set contains a ball of radius r_{i+1} . Given that this aligns with the setup of Problem 2.1, we can proceed with the PS_{exp} to sample from μ_{i+1}^X , referring to the query complexity result in Theorem 2.15.

After the while-loop, in Line 9, we use $k \asymp n \text{polylog } n$ samples to estimate $\bar{\Sigma}$ and $\bar{\mathbf{m}}$, which by Lemma 4.6 satisfies that with probability at least $1 - \mathcal{O}(n^{-1})$,

$$(1 - 10^{-3}) \text{Cov } \mu_{\text{last}}^X \preceq \bar{\Sigma} \preceq (1 + 10^{-3}) \text{Cov } \mu_{\text{last}}^X.$$

Thus, for $F^X : x \mapsto \bar{\Sigma}^{-1/2}(x - \bar{\mathbf{m}})$, it holds that $F^X_\# \mu_{\text{last}}^X$ is 1.01-isotropic. Moreover, Lemma 4.6 implies that the mean of $F^X_\# \mu_{\text{last}}^X$ lies within a ball of radius 10^{-2} , i.e.,

$$\|\bar{\Sigma}^{-1/2}(\mathbf{m}_{\text{last}} - \bar{\mathbf{m}})\| \leq 0.01.$$

Combining these results, we record the final guarantee of Algorithm 4. A similar result for the rounding of uniform distributions can be found in [KZ24, Lemma 18].

Lemma 4.11 (Guarantee of Isotropize). *Assume that a distribution μ^X satisfies Assumption 4.4. Then, with probability at least $1 - \mathcal{O}(n^{-1/2})$, Algorithm 4 on the input μ returns a distribution $F_\# \mu_{\text{last}}$ using $\tilde{\mathcal{O}}(n^3)$ evaluation queries such that the X -marginal $F_\# \mu_{\text{last}}^X = (F_\# \mu_{\text{last}})^X$ is 1.01-isotropic, with its mean lying within a ball of radius 0.01.*

4.2 Maintaining well-roundedness

Recall that Step 1-(b) repeats the following while $r = \mathcal{O}(R)$: if $F_\# \nu_r^X$ is near-isotropic, then run **Isotropize** on $(4F^X)_\# \nu_{\delta r}^X$ for $\delta = 1 + n^{-1/2}$, and update $F^X \leftarrow 4F^X$ and $r \leftarrow \delta r$. To ensure this approach is valid, it must be the case that the input distribution $(4F^X)_\# \nu_{\delta r}^X$ satisfies Assumption 4.4 as well. We show that [JLLV24, Lemma 3.4], proven for the uniform distribution over a convex body, remains valid for any logconcave distribution.

Lemma 4.12. *Let $\delta = 1 + n^{-1/2}$ and $F^X : \mathbb{R}^n \rightarrow \mathbb{R}^n$ an invertible affine map. If $F_{\#}^X \nu_r^X$ is 1.01-isotropic with its mean lying within a ball of radius 0.01, then $(4F^X)_{\#} \nu_{\delta r}^X$ satisfies Assumption 4.4 (i.e., evaluation oracle, inclusion of $B_1(0)$, well-roundedness, and ground set equal to its support).*

Proof. By setting $U = V \circ (H^X)^{-1} / \mathbf{1}[x \in H^X(\mathbb{L}_g \cap B_{\delta r}(0))]$ with $H^X := 4F^X$, one can implement the evaluation oracle for the convex U (using the oracle for V) and recover $H_{\#}^X \nu_{\delta r}^X \propto \exp(-U)$. It also follows from the definition of U and \mathbb{L}_g that $U(x) \leq 10n$ on $\text{supp } H_{\#}^X \nu_{\delta r}^X$.

Let us now examine the inclusion of $B_1(0)$ in its ground set. When $F^X : x \mapsto Ax - \mu$ between \mathbb{R}^n , let us define an affine map $F_0^X : x \mapsto Ax - b$ from \mathbb{R}^n to \mathbb{R}^n which makes ν_r^X centered, noting that $\|b - \mu\| \leq 0.01$. As for the lower regularity, since $(F_0^X)_{\#} \nu_r^X$ is 1.01-isotropic with zero mean, by Lemma A.10, its support contains a ball of radius $(e\sqrt{1.01})^{-1} \geq 0.35$ centered at the origin. Hence, as $\|b - \mu\| \leq 0.01$, $\text{supp } F_{\#}^X \nu_r^X$ contains a ball of radius 0.3 centered at the origin, so does $\text{supp } F_{\#}^X \nu_{\delta r}^X$. Therefore, by scaling via $x \mapsto 4x$, we can ensure that $\text{supp } H_{\#}^X \nu_{\delta r}^X$ contains a unit ball $B_1(0)$. Due to $\min U \geq 0$ and $U(x) \leq 10n$ on $\text{supp } H_{\#}^X \nu_{\delta r}^X$, we clearly have $B_1(0) \subset \mathbb{L}_{H_{\#}^X \nu_{\delta r}^X, g}$.

We define an affine map $G_0^X : x \mapsto Ax - \delta b$ between \mathbb{R}^n . Observe that for a constant $c' \geq 2\delta$,

$$\begin{aligned} \mathbb{E}_{(G_0^X)_{\#} \nu_{\delta r}^X} \|X\| &= \int_0^{\infty} \mathbb{P}_{(G_0^X)_{\#} \nu_{\delta r}^X} (\|X\| \geq u) \, du \leq c' \sqrt{n} + \int_{c' \sqrt{n}}^{\infty} \mathbb{P}(\|X\| \geq u) \, du \\ &\leq c' \sqrt{n} + \delta \int_{c' \delta^{-1} \sqrt{n}}^{\infty} \mathbb{P}(\|X\| \geq \delta u) \, du. \end{aligned} \quad (4.2)$$

Let us bound the integrand in terms of $\mathbb{P}_{(F_0^X)_{\#} \nu_r^X} (\|X\| \geq u)$. For the embedded $G : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ defined by $G(x, t) = (G_0^X(x), t)$, we have

$$\text{supp } G_{\#} \nu_{\delta r} = G(\mathcal{K} \cap C_g \cap C_{\delta r}) = \begin{bmatrix} A & \\ & 1 \end{bmatrix} (\mathcal{K} \cap C_g \cap C_{\delta r}) - \begin{bmatrix} \delta b \\ 0 \end{bmatrix},$$

where $C_g = \mathbb{L}_g \times \mathbb{R}$ and $C_{\delta r} = B_{\delta r}(0) \times \mathbb{R}$. As \mathcal{K} and C_g contain the origin, and $\delta > 1$, we clearly have $\mathcal{K} \cap C_g \cap C_{\delta r} \subset \delta(\mathcal{K} \cap C_g \cap C_r)$, and thus

$$\begin{aligned} \text{supp } G_{\#} \nu_{\delta r} &\subseteq \delta \begin{bmatrix} A & \\ & 1 \end{bmatrix} (\mathcal{K} \cap C_g \cap C_r) - \begin{bmatrix} \delta b \\ 0 \end{bmatrix} = \delta \left(\begin{bmatrix} A & \\ & 1 \end{bmatrix} (\mathcal{K} \cap C_g \cap C_r) - \begin{bmatrix} b \\ 0 \end{bmatrix} \right) \\ &= \delta \text{supp } F_{\#} \nu_r, \end{aligned}$$

where $F : (x, t) \mapsto (F_0^X(x), t)$ is an affine map between \mathbb{R}^{n+1} . Moreover, it follows from $\delta C_{l\sqrt{n}} = C_{l\delta\sqrt{n}}$ for any $l > 0$ that

$$(\text{supp } G_{\#} \nu_{\delta r}) \setminus C_{l\delta\sqrt{n}} \subset [\delta \text{supp } F_{\#} \nu_r] \setminus [\delta C_{l\sqrt{n}}] = \delta [(\text{supp } F_{\#} \nu_r) \setminus C_{l\sqrt{n}}].$$

Integrating $\exp(-nt)$ over these regions and using $\inf_{\mathcal{K}} t \geq -11$ in (i) below,

$$\begin{aligned} \int_{(\text{supp } G_{\#} \nu_{\delta r}) \setminus C_{l\delta\sqrt{n}}} \exp(-nt) \, dx dt &\leq \int_{\delta [(\text{supp } F_{\#} \nu_r) \setminus C_{l\sqrt{n}}]} \exp(-nt) \, dx dt \\ &= \delta^{n+1} \int_{(\text{supp } F_{\#} \nu_r) \setminus C_{l\sqrt{n}}} \exp(-nt - t\sqrt{n}) \, dx dt \\ &\stackrel{(i)}{\leq} e^{11\sqrt{n}} \delta^{n+1} \int_{(\text{supp } F_{\#} \nu_r) \setminus C_{l\sqrt{n}}} \exp(-nt) \, dx dt \\ &= e^{11\sqrt{n}} \delta^{n+1} \cdot \mathbb{P}_{F_{\#} \nu_r} (\|X\| \geq l\sqrt{n}) \int_{\text{supp } F_{\#} \nu_r} \exp(-nt) \, dx dt. \end{aligned} \quad (4.3)$$

Since $(F_{\#}\nu_r)^X = (F_0^X)_{\#}\nu_r^X$ is an 1.01-isotropic centered logconcave distribution, a version of Paouris' theorem (Lemma A.8) implies that for some universal constant $c > 0$ and any $l \geq 2$,

$$\mathbb{P}_{F_{\#}\nu_r}(\|X\| \geq l\sqrt{n}) = \mathbb{P}_{(F_0^X)_{\#}\nu_r^X}(\|X\| \geq l\sqrt{n}) \leq \exp(-cl\sqrt{n}).$$

Dividing both sides of (4.3) by $\int_{\text{supp } G_{\#}\nu_{\delta r}} \exp(-nt) \, dxdt$, and substituting this Paouris bound,

$$\begin{aligned} \mathbb{P}_{(G_0^X)_{\#}\nu_{\delta r}^X}(\|X\| \geq l\delta\sqrt{n}) &= \mathbb{P}_{G_{\#}\nu_{\delta r}}(\|X\| \geq l\delta\sqrt{n}) \\ &\leq e^{11\sqrt{n}}\delta^{n+1} \exp(-cl\sqrt{n}) \frac{\int_{\text{supp } F_{\#}\nu_r} \exp(-nt) \, dxdt}{\int_{\text{supp } G_{\#}\nu_{\delta r}} \exp(-nt) \, dxdt} \\ &\leq e^{1+(12-cl)\sqrt{n}}. \end{aligned}$$

Putting this bound back to (4.2),

$$\begin{aligned} \mathbb{E}_{(G_0^X)_{\#}\nu_{\delta r}^X} \|X\| &\leq c'\sqrt{n} + \delta \int_{c'\delta^{-1}\sqrt{n}}^{\infty} \mathbb{P}_{(G_0^X)_{\#}\nu_{\delta r}^X}(\|X\| \geq \delta u) \, du \\ &\leq c'\sqrt{n} + e^{2+12\sqrt{n}} \int_{c'\delta^{-1}\sqrt{n}}^{\infty} e^{-cu} \, du \\ &= c'\sqrt{n} + c^{-1}e^{2+12\sqrt{n}}e^{-cc'\sqrt{n}/\delta}. \end{aligned}$$

By taking c' large enough, the second term can be made exponentially small. It then follows from the reverse Hölder (Lemma A.9) that

$$\mathbb{E}_{(G_0^X)_{\#}\nu_{\delta r}^X}[\|X\|^2] \lesssim (\mathbb{E}_{(G_0^X)_{\#}\nu_{\delta r}^X} \|X\|)^2 \lesssim n.$$

Thus, for $F^X(x) = Ax - \mu = G_0^X + \delta b - \mu$, we have

$$\begin{aligned} \mathbb{E}_{F_{\#}\nu_{\delta r}^X}[\|X\|^2] &\leq 2\mathbb{E}_{F_{\#}\nu_{\delta r}^X}[\|X - \delta b + \mu\|^2] + 2\|\delta b - \mu\|^2 = 2\mathbb{E}_{(G_0^X)_{\#}\nu_{\delta r}^X}[\|X\|^2] + 2\|\delta b - \mu\|^2 \\ &\lesssim n + \frac{\|b\|^2}{n} \lesssim n, \end{aligned}$$

where the last inequality follows from Lemma 4.13 below. Therefore, $\mathbb{E}_{H_{\#}\nu_{\delta r}^X}[\|X\|^2] = \mathcal{O}(n)$. \square

We now present a key geometric property concerning the diameter of the level set of logconcave distributions. We remark that this properly generalizes [LV07, Lemma 5.18], which says that if z is a maximum point of an isotropic logconcave density, then $\|z\| \leq n + 1$.

Lemma 4.13 (Diameter of level sets). *Suppose an isotropic logconcave density in \mathbb{R}^n is proportional to e^{-V} . Then, for $V_0 = \min V$ and any $c > 0$,*

$$\max_{x: V(x) - V_0 \leq cn} \|x\| \leq 8e^{c+1}n.$$

Proof. Pick any point x with $V(x) \leq V_0 + cn$. We may assume by rotation that $x = De_1$ for $D = \|x\| > 0$. Let us consider the marginal of the density along the direction e_1 , which is also isotropic with zero mean. We also consider the $(n-1)$ -dimensional slices perpendicular to e_1 at 0 and at D/n , calling them $\mathcal{S}(0)$ and $\mathcal{S}(D/n)$ respectively. We can define a bijective mapping $f: \mathcal{S}(D/n) \rightarrow \mathcal{S}(0)$ by $y \mapsto z = x + \frac{y-x}{1-n^{-1}}$; in words, $f(y)$ is the point $z \in \mathcal{S}(0)$ such that $y \in \overline{zx}$. Using the convexity of V with endpoints z and x ,

$$V(y) = V\left(\frac{n-1}{n}z + \frac{1}{n}x\right) \leq \frac{n-1}{n}V(z) + \frac{1}{n}V(x) \leq V(z) + c.$$

It then follows that

$$\begin{aligned} \int_{\mathcal{S}(D/n)} e^{-V(y)} \, dy &\geq e^{-c} \int_{\mathcal{S}(D/n)} e^{-V(f(y))} \, dy = \frac{e^{-c}}{|Df|} \int_{\mathcal{S}(0)} e^{-V(z)} \, dz \\ &= \left(1 - \frac{1}{n}\right)^{n-1} e^{-c} \int_{\mathcal{S}(0)} e^{-V(z)} \, dz \geq e^{-c-1} \int_{\mathcal{S}(0)} e^{-V(z)} \, dz. \end{aligned}$$

By Lemma A.11, the density of the marginal at 0 is at least $1/8$, so the density at D/n is at least $e^{-c-1}/8$. Since the value of the marginal density function is at least $\min(1/8, e^{-c-1}/8) = e^{-c-1}/8$ due to the convexity of V , the measure of the marginal in the range $[0, D/n]$ is at least

$$\frac{D}{8n} e^{-c-1}.$$

As this must be at most 1 (actually at most $1 - (1/e)$), we obtain that $D \leq 8e^{c+1} n$ as claimed. \square

4.3 Rounding the grounded distribution

We have justified Step 1 of the overall approach thus far. We now examine Step 2 and 3.

Step 2 ($\nu_D^X \rightarrow \nu^X$). Given an affine map F^X such that $F_{\#}^X \nu_D^X$ is 1.01-isotropic, and we show that $(4F^X)_{\#} \nu^X$ satisfies Assumption 4.4. The following result allows us to bound the complexity of Isotropize on $(4F^X)_{\#} \nu^X$ by $\tilde{\mathcal{O}}(n^3)$ as well.

Lemma 4.14. *Let $F^X : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an invertible affine map, and $D = 2R \log 400$. If $F_{\#}^X \nu_D^X$ is 1.01-isotropic with its mean located within 0.01 from the origin, then $(4F^X)_{\#} \nu_{\delta r}^X$ satisfies Assumption 4.4.*

Proof. Since ν^X is 1.01-warm with respect to π^X due to (4.1), $\mathbb{E}_{\nu^X}[\|X\|^2] \leq 1.01 \mathbb{E}_{\pi^X}[\|X\|^2] \leq (2R)^2$, so Lemma A.7 leads to

$$\mathbb{P}_{\nu^X}(\|X\| \geq 2tR) \leq e^{-t+1} \quad \text{for any } t \geq 1.$$

Thus, $\mathbb{P}_{\nu^X}(\|X\| \leq D) \geq 1 - \varepsilon$ for $D = 2R \log \frac{e}{\varepsilon}$, which implies

$$\frac{\nu_D^X}{\nu^X} = (\mathbb{P}_{\nu^X}(\text{supp } \nu_D^X))^{-1} = (\mathbb{P}_{\nu^X}(\|X\| \leq D))^{-1} \leq (1 - \varepsilon)^{-1}.$$

It means that $\mu_D := F_{\#}^X \nu_D^X$ is also $(1 - \varepsilon)^{-1}$ -warm with respect to $\mu := F_{\#}^X \nu^X$.

We first address the upper regularity (UR). For any unit vector $v \in \mathbb{R}^n$,

$$\begin{aligned} |\mathbb{E}_{\mu_D}|X \cdot v| - \mathbb{E}_{\mu}|X \cdot v| &= \left| \int |X \cdot v| (d\mu_D - d\mu) \right| \\ &\leq \sqrt{\mathbb{E}_{\mu}[|X \cdot v|^2]} \sqrt{\chi^2(\mu_D \parallel \mu)} \\ &\leq 4 \mathbb{E}_{\mu}|X \cdot v| \left(\frac{\mu_D}{\mu} - 1 \right), \end{aligned}$$

where the last line follows from the reverse Hölder (Lemma A.9). Rearranging terms, we obtain that

$$\begin{aligned} \mathbb{E}_{\mu}|X \cdot v| &\leq \frac{1 - \varepsilon}{1 - 5\varepsilon} \mathbb{E}_{\mu_D}|X \cdot v| \leq \frac{1 - \varepsilon}{1 - 5\varepsilon} (\mathbb{E}_{\mu_D}|(X - \mathbb{E}_{\mu_D} X) \cdot v| + 0.01) \\ &\leq \frac{1 - \varepsilon}{1 - 5\varepsilon} \times 1.02. \end{aligned}$$

Hence, by taking ε small enough, we can ensure that $\mathbb{E}_\mu |X \cdot v| \leq 1.05$ for any $v \in \mathbb{R}^n$, and this immediately implies that

$$\mathbb{E}_\mu [\|X\|^2] = \mathbb{E}_{F_{\#}^X \nu^X} [\|X\|^2] = \mathcal{O}(n).$$

As for the lower regularity (LR), we already showed in the proof of Lemma 4.12 that due to the isotropy, the support of $F_{\#}^X \nu_D^X$ contains a ball of radius 0.3 centered at the origin, so does that of $F_{\#}^X \nu^X$. Therefore, scaling by $x \mapsto 4x$, we can conclude that $B_1(0) \subseteq \text{supp}(4F^X)_{\#} \nu^X$ and that $\mathbb{E}_{(4F^X)_{\#} \nu^X} [\|X\|^2] = \mathcal{O}(n)$. The implementability of the evaluation oracle for the potential, as well as $\text{supp}(4F^X)_{\#} \nu^X = \mathcal{L}_{(4F^X)_{\#} \nu^X, g}$, can be proven similarly to Lemma 4.12. \square

Step 3 ($\nu^X \rightarrow \pi^X$). We show that if $F_{\#}^X \nu^X$ is 1.01-isotropic, then the largest eigenvalue of the covariance of $(4F^X)_{\#} \pi^X$ is $\mathcal{O}(1)$, and its ground set contains $B_1(0)$.

Lemma 4.15. *Let $F^X : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an invertible affine map such that $F_{\#}^X \nu^X$ is 1.01-isotropic with its mean located within 0.01 from the origin. Then, for the affine map $H^X = 4F^X$, the ground set $H^X \mathcal{L}_g$ of $H_{\#}^X \pi^X$ contains a unit ball $B_1(0)$, and $\|\text{Cov } H_{\#}^X \pi^X\| = \mathcal{O}(1)$.*

Proof. As in the proof of Lemma 4.14, for any unit vector $v \in \mathbb{R}^n$,

$$|\mathbb{E}_{F_{\#}^X \nu^X} |X \cdot v| - \mathbb{E}_{F_{\#}^X \pi^X} |X \cdot v|| \leq 4 \mathbb{E}_{F_{\#}^X \pi^X} |X \cdot v| \left(\frac{\nu^X}{\pi^X} - 1 \right) \leq 0.04 \mathbb{E}_{F_{\#}^X \pi^X} |X \cdot v|,$$

and thus $\mathbb{E}_{F_{\#}^X \pi^X} |X \cdot v| \leq 1.05 \mathbb{E}_{F_{\#}^X \nu^X} |X \cdot v|$. Using the reverse Hölder inequality in (i) below,

$$\begin{aligned} \mathbb{E}_{F_{\#}^X \pi^X} [(X - \mathbb{E}_{F_{\#}^X \pi^X} X) \cdot v]^2 &\leq \mathbb{E}_{F_{\#}^X \pi^X} [|X \cdot v|^2] \stackrel{(i)}{\leq} 16 (\mathbb{E}_{F_{\#}^X \pi^X} |X \cdot v|)^2 \leq 20 (\mathbb{E}_{F_{\#}^X \nu^X} |X \cdot v|)^2 \\ &\leq 20 (\mathbb{E}_{F_{\#}^X \nu^X} |(X - \mathbb{E}_{F_{\#}^X \nu^X} X) \cdot v| + 0.01)^2 \leq 50. \end{aligned}$$

Thus, the largest eigenvalue of $\text{Cov } H_{\#}^X \pi^X$ is $\mathcal{O}(1)$, and the ground set of $H_{\#}^X \pi^X$ contains $B_1(0)$. \square

We note that $\mathbb{E}_{H_{\#}^X \pi^X} [\|X\|^2] = \mathcal{O}(n)$, so we can first run Tilted Gaussian Cooling to generate an $\mathcal{O}(1)$ -warm start $Z_0 = (X_0, T_0)$ for $H_{\#} \pi$, using $\tilde{\mathcal{O}}(n^3)$ queries in expectation. Then, as in Isotropize, we run $\text{PS}_N(H_{\#} \pi, \delta_{Z_0}, h)$ with $N \asymp n^2 \text{polylog } n$ and $h \asymp (n^2 \text{polylog } n)^{-1}$, obtaining $k \asymp n \text{polylog } n$ many samples to compute the empirical covariance $\bar{\Sigma}$ for $\Sigma = \text{Cov } H_{\#}^X \pi^X$. It then follows from Lemma 4.6 that with probability at least $1 - \mathcal{O}(n^{-1})$,

$$0.99 \bar{\Sigma} \preceq \Sigma \preceq 1.01 \bar{\Sigma}.$$

Thus, $(\bar{\Sigma}^{-1/2} \circ H^X)_{\#} \pi^X$ becomes 1.01-isotropic. Since one iteration of PS_N uses $\tilde{\mathcal{O}}(N)$ queries, the total complexity of this procedure (in addition to Tilted Gaussian Cooling) would be $\tilde{\mathcal{O}}(n^3)$. Therefore, Step 3 also uses $\tilde{\mathcal{O}}(n^3)$ queries in expectation.

5 Application 2: Integrating logconcave functions

In this section, we propose a faster algorithm for integrating logconcave functions, leveraging our improved sampling guarantees and streamlining previous analysis in [KLS97, LV06c, CV18].

Problem 5.1 (Integration). Assume access to a well-defined function oracle $\text{Eval}_{x_0, R}(V)$ for convex $V : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$. Given $\varepsilon > 0$, what is the query complexity of computing a $(1 + \varepsilon)$ -multiplicative approximation F to the integral of $f = e^{-V}$ with probability at least $3/4$, i.e.,

$$(1 - \varepsilon) \int f \leq F \leq (1 + \varepsilon) \int f.$$

We can assume that $\min V = 0$ and that π^X is well-rounded due to the rounding algorithm in the previous section. Hereafter, we assume that $x_0 = 0$, $R^2 = C^2 n$ for a known constant $C > 0$, $\mathcal{K} = \{(x, t) : \mathbb{R}^n \times \mathbb{R} : V(x) - 11n \leq nt\}$, and that for $l = C^2 \vee \log \frac{8e}{\varepsilon}$,

$$\bar{\mathcal{K}} = \mathcal{K} \cap \{B_{Rl}(0) \times [-11, 4 + 13l]\}.$$

We recall that $\bar{\mathcal{K}}$ takes up $(1 - \frac{\varepsilon}{4})$ measure of π and denote $\bar{\pi} := \pi|_{\bar{\mathcal{K}}}$. In this section, we establish an improved complexity of integrating a logconcave function.

Theorem 5.2 (Restatement of Theorem 1.8). *For any integrable logconcave $f : \mathbb{R}^n \rightarrow \mathbb{R}$ given by a well-rounded function oracle and any $\varepsilon > 0$, there exists an algorithm that with probability at least $3/4$, returns a $(1 + \varepsilon)$ -multiplicative approximation to the integral of f using $\tilde{\mathcal{O}}(n^3/\varepsilon^2)$ queries. For an arbitrary logconcave f , the total query complexity is bounded by $\tilde{\mathcal{O}}(n^{3.5} \text{polylog } R + n^3/\varepsilon^2)$.*

5.1 Algorithm

The main idea for integration is also annealing, and will parallel the warm-start generation but in a more sophisticated way. We consider a sequence of distributions in \mathbb{R}^{n+1} starting with a simple logconcave function that is easy to integrate and ending with the target logconcave function, where two consecutive distributions μ_i and μ_{i+1} satisfy two conditions: (i) $\mathcal{R}_\infty(\mu_i \parallel \mu_{i+1}) = \mathcal{O}(1)$ for fast sampling from μ_{i+1} started at μ_i , and (ii) $\mathcal{R}_2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(n^{-1})$ or $\mathcal{O}(\sigma^2/C^2n)$ for small variance of estimators used in the algorithm.

This annealing scheme is used in [CV18] for the special case of Gaussians restricted to convex bodies. It needs an efficient sampler alongside with good dependence on the warm start and output guarantees. These were available at the time for Gaussians restricted to convex bodies. Our new sampler allows us to get similar strong guarantees for arbitrary logconcave distributions, so we will be able to extend the guarantees to logconcave integration by addressing additional technical difficulties.

Setup. Let m be the number of samples used for estimation, and p the number of total inner phases. Then, the integration algorithm (Algorithm 6) initializes m independent starts, generating a *thread* of samples denoted by $(Z_i^s = (X_i^s, T_i^s))_{0 \leq i \leq p}$ for $s \in [m]$. Here, the initial start Z_0^s is drawn from $\exp(-\frac{n}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$, and subsequent samples $(Z_i^s)_{i \in [p]}$ drawn through PS_{ann} are appended to the thread. Clearly, for each $i \in [p]$, the i -th samples Z_i^s across the threads are independent and have the same law (i.e., $Z_i^{s_1} \perp Z_i^{s_2}$ and law $Z_i^{s_1} = \text{law } Z_i^{s_2}$ for any $s_1 \neq s_2 \in [m]$). PS_{ann} with suitable parameters can ensure that the law of a sample Z_i^s , denoted by $\bar{\mu}_i = (\bar{\mu}_i^X, \bar{\mu}_i^T)$, is close to

$$\begin{aligned} \mu_0(x, t) &\propto \exp(-\frac{n}{2} \|x\|^2)|_{\bar{\mathcal{K}}} =: f_0(x, t), \\ \mu_i(x, t) &\propto \exp(-\frac{1}{2\sigma_i^2} \|x\|^2 - \rho_i t)|_{\bar{\mathcal{K}}} =: f_i(x, t) \quad \text{for } i \in [p], \end{aligned}$$

and we denote its integral (i.e., the normalization constant) by $F_i := \int f_i$.

Algorithm 5 Samp-Est ($Z_{i-1}, f_i, f_{i+1}, \epsilon$)

Input: initial point Z_{i-1} , target density f_i , upcoming density f_{i+1} , target accuracy ϵ .

Output: estimation E_i and samples Z_i approximately close to $\mu_{i+1} \propto f_{i+1}$.

- 1: For each $s \in [m]$, draw Z_i^s through PS_{ann} started at Z_{i-1}^s for target $\mu_i \propto f_i$ with accuracy ϵ .
- 2: Estimate

$$E_i = \frac{1}{m} \sum_{s=1}^m \frac{f_{i+1}(Z_i^s)}{f_i(Z_i^s)}.$$

- 3: Return E_i and $\{Z_i^s\}_{s \in [m]}$.
-

Estimation using drawn samples. Algorithm 6 calls Algorithm 5 for estimating $E_i := \frac{1}{m} \sum_{s=1}^m Y_i^s$, just the empirical average of Y_i^s defined as

$$Y_i^s := \frac{f_{i+1}(Z_i^s)}{f_i(Z_i^s)} \quad \text{for } i = 0, \dots, p, \text{ with convention } f_{p+1}(z) := \exp(-nt)|_{\bar{\mathcal{K}}}.$$

If each sample Z_i were to follow the *exact distribution* μ_i rather than the approximate one $\bar{\mu}_i$, then

$$\mathbb{E}E_i = \mathbb{E}Y_i = \int \frac{f_{i+1}(z)}{f_i(z)} \frac{f_i(z)}{\int f_i} dz = \frac{F_{i+1}}{F_i},$$

and thus

$$\underbrace{\int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2} \|x\|^2) dx dt}_{=: I_0} \times \mathbb{E}_{\mu_0} Y_0 \cdots \mathbb{E}_{\mu_p} Y_p = F_0 \frac{F_1}{F_0} \cdots \frac{F_{p+1}}{F_p} = \int_{\bar{\mathcal{K}}} \exp(-nt) dx dt.$$

Since $\mathbb{P}_\pi(X \in \bar{\mathcal{K}}) \geq 1 - \epsilon$ by construction, and $\int_{\bar{\mathcal{K}}} \exp(-nt) dx dt = \frac{1}{n} \int e^{-V}$, it implies that

$$(1 - \epsilon) \int e^{-V} \leq n I_0 \times \mathbb{E}Y_0 \cdots \mathbb{E}Y_p \leq \int e^{-V}. \quad (5.1)$$

While estimating $\mathbb{E}Y_i$ is straightforward since PS_{ann} yields as many samples from $\bar{\mu}_i \approx \mu_i$ as needed, we should also be able to estimate the initial integral $I_0 = \int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2} \|x\|^2) dx dt$.

The initial integral I_0 . For $I := [-11, 4 + 13l]$, we define $f_{-1}(x, t) := \exp(-\frac{n}{2} \|x\|^2) \cdot \mathbf{1}_I(t)$, and $\mu_{-1}(x, t) = \mathcal{N}(0, n^{-1}I_n) \otimes \text{Unif } I \propto f_{-1}(x, t)$. Using the estimate

$$Y_{-1}(x, t) := \frac{f_0(x, t)}{f_{-1}(x, t)} = \frac{\exp(-\frac{n}{2} \|x\|^2) \cdot \mathbf{1}_{\bar{\mathcal{K}}}(x, t)}{\exp(-\frac{n}{2} \|x\|^2) \cdot \mathbf{1}_I(t)} = \mathbf{1}_{\bar{\mathcal{K}}}(x, t) \quad \text{for } (x, t) \in \text{supp } \mu_{-1},$$

and its empirical average $E_{-1} := \frac{1}{m} \sum_{s=1}^m Y_{-1}^s$, we can approximate I_0 through $E_{-1} \int f_{-1}$ due to

$$E_{-1} \int f_{-1} \approx \mathbb{E}_{\mu_{-1}} Y_{-1} \times \int f_{-1} = \int \mathbf{1}_{\bar{\mathcal{K}}}(x, t) f_{-1}(x, t) dx dt = \int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2} \|x\|^2) dx dt = I_0.$$

Here, $\int f_{-1}$ can be computed by a formula

$$\int f_{-1} = \int \exp(-\frac{n}{2} \|x\|^2) dx \times (15 + 13l) = (2\pi n^{-1})^{n/2} (15 + 13l).$$

Overview of algorithm. We propose a new annealing scheme (Algorithm 6) to fulfill the two conditions on \mathcal{R}_2 and \mathcal{R}_∞ .

- **[Phase I]** While $\sigma^2 \in [n^{-1}, 1]$:
 - End-to-end: $\mu_{\text{start}} \propto \exp(-\frac{n}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$ to $\mu_{\text{end}} \propto \exp(-\frac{1}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$.
 - Update: $\sigma^2 \leftarrow \sigma^2 (1 + \frac{1}{n})$ and $\rho = 0$.
 - Guarantee: $\mathcal{R}_2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(n^{-1})$ & $\tilde{\mathcal{O}}(n)$ phases (Lemma 5.8).
- **[Phase II]** While $\rho \leq n$:
 - End-to-end: $\mu_{\text{start}} \propto \exp(-\frac{1}{2} \|x\|^2 - \frac{1}{28l} t)|_{\bar{\mathcal{K}}}$ to $\mu_{\text{end}} \propto \exp(-\frac{1}{2} \|x\|^2 - nt)|_{\bar{\mathcal{K}}}$.
 - Update: $[\sigma^2 \leftarrow \sigma^2 / (1 + \frac{1}{28nl})$ and $\rho \leftarrow \rho (1 + \frac{1}{28nl})]$ and $[\sigma^2 \leftarrow \sigma^2 (1 + \frac{1}{28nl})$ and $\rho \leftarrow \rho]$.⁴
 - Guarantee: $\mathcal{R}_2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(n^{-1})$ & $\tilde{\mathcal{O}}(nl)$ phases (Lemma 5.10).
- **[Phase III]** While $\sigma^2 \in [1, C^2n]$:
 - End-to-end: $\mu_{\text{start}} \propto \exp(-\frac{1}{2} \|x\|^2 - nt)|_{\bar{\mathcal{K}}}$ to $\mu_{\text{end}} \propto \exp(-\frac{1}{2C^2n} \|x\|^2 - nt)|_{\bar{\mathcal{K}}}$.
 - Update $\sigma^2 \leftarrow \sigma^2 (1 + \frac{\sigma^2}{C^2n})$ and $\rho = n$.
 - Guarantee: $\mathcal{R}_2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(\frac{\sigma^2}{C^2n})$ & $\tilde{\mathcal{O}}(\frac{C^2n}{\sigma^2})$ phases for doubling of σ^2 (see Lemma 5.11).

In summary, PS_{ann} started at $\bar{\mu}_{i-1}$ generates samples $Z \sim \bar{\mu}_i$, and estimate $f_{i+1}(Z)/f_i(Z)$. Meanwhile, we would like to ensure that (1) $\mathcal{R}_\infty(\mu_{i-1} \parallel \mu_i) = \mathcal{O}(1)$ when moving from $\bar{\mu}_{i-1}$ to $\bar{\mu}_i$, and (2) $\mathcal{R}_2(\mu_{i+1} \parallel \mu_i) = \mathcal{O}(n^{-1})$ in Phase I and II, and $\mathcal{O}(\frac{\sigma^2}{C^2n})$ in Phase III for variance control of f_{i+1}/f_i .

5.2 Streamlined analysis

We show that our estimate $E_{-1} \cdots E_p$ concentrates around $\mathbb{E}Y_{-1} \cdots \mathbb{E}Y_p$ with probability at least 0.9 (say).

5.2.1 Technical tools

Two sources of complications. Unfortunately, the samples Z_i in the algorithm follow an **approximate distribution** $\bar{\mu}_i$ (not exactly the target μ_i), so this entails a coupling characterization of TV-distance in prior work. Once this issue is properly addressed, readers may want to use a concentration of each E_i around $\mathbb{E}E_i$ based on $\text{var} E_i$, but this does not suit our purpose. For example, in Phase I, the (relative) variance $\text{var}(E_i/\mathbb{E}E_i)$ is bounded by $1/n$, so this leads to $E_i = (1 \pm \mathcal{O}(n^{-1/2})) \mathbb{E}E_i$ with high probability. Since there are $p = \tilde{\mathcal{O}}(n)$ many phases, the relative error of the product of E_i 's would blow up. Thus, one should rely on the concentration of the *product* $E_{-1} \cdots E_p$, not just combining that of each estimate E_i . To this end, we should be able to establish

$$\text{var}(E_{-1} \cdots E_p) \lesssim (\mathbb{E}[E_{-1} \cdots E_p])^2 \quad (\text{equivalently, } \mathbb{E}[E_{-1}^2 \cdots E_p^2] = (1 + \mathcal{O}(1)) (\mathbb{E}[E_{-1} \cdots E_p])^2).$$

Unfortunately, another complication comes into play due to **dependence of samples** *within* each thread, so one also needs $\mathbb{E}[E_{-1} \cdots E_p] = (1 \pm \mathcal{O}(1)) \mathbb{E}E_{-1} \cdots \mathbb{E}E_p$. Prior analysis of volume algorithms used the notion of α -mixing (called μ -independence therein) to address this technicality.

⁴In fact, we can combine two updates, but we separate them for ease of analysis.

Algorithm 6 Integral(f, ε)

Input: a logconcave function f in Problem 5.1, and target accuracy $\varepsilon > 0$.

Output: an estimation E for $\int f$.

1: $\diamond i = 0, l = C^2 \vee \log \frac{8\varepsilon}{\varepsilon}, (\sigma_0^2, \rho_0) = (\frac{1}{n}, 0), m = \frac{10^{11} l \log n l \log C^2 n}{\varepsilon^2}, \epsilon = \frac{\varepsilon^2}{10^{15} C^2 n l^2 \log^3(C^2 n l)}$. Denote

$$f_j(x, t) := \exp(-\frac{1}{2\sigma_j^2} \|x\|^2 - \rho_j t) \Big|_{\bar{\mathcal{K}}}.$$

2: Get $\{Z_{-1}^s\}_{s \in [m]} \sim \mu_{-1} = \mathcal{N}(0, \frac{1}{n} I_n) \otimes \text{Unif}([-11, 4+13l])$, and compute $E_{-1} = \frac{1}{m} \sum_{s=1}^m \mathbf{1}_{\bar{\mathcal{K}}}(Z_{-1}^s)$.

3: **while** $\sigma_i^2 \leq 1$ **do**

4: If $\sigma_i^2 = 1$, then go to next **while** after this iteration. Set

$$\begin{cases} \sigma_{i+1}^2 = 1 \wedge \sigma_i^2(1 + \frac{1}{n}) & \& \rho_{i+1} = 0 & \text{if } \sigma_i^2 < 1, \\ \sigma_{i+1}^2 = \sigma_i^2 & \& \rho_{i+1} = \frac{1}{28l} & \text{if } \sigma_i^2 = 1. \end{cases}$$

5: Run $(E_i, \{Z_i^s\}_{s \in [m]}) \leftarrow \text{Samp-Est}(\{Z_{i-1}^s\}_{s \in [m]}, f_i, f_{i+1}, \epsilon)$ and increment i .

6: **end while**

7: **while** $\rho_i < n$ **do**

8: Set $\beta = 1 + \frac{1}{28nl}$ if $\beta\rho_i < n$, and $\beta = n/\rho_i$ otherwise.

9: $(E_i, \{Z_i^s\}_{s \in [m]}) \leftarrow \text{Samp-Est}(\{Z_{i-1}^s\}_{s \in [m]}, f_i, f_{i+1}, \epsilon)$ for $\sigma_{i+1}^2 = \frac{\sigma_i^2}{\beta}, \rho_{i+1} = \beta\rho_i$. Increment i .

10: $(E_i, \{Z_i^s\}_{s \in [m]}) \leftarrow \text{Samp-Est}(\{Z_{i-1}^s\}_{s \in [m]}, f_i, f_{i+1}, \epsilon)$ for $\sigma_{i+1}^2 = \beta\sigma_i^2, \rho_{i+1} = \rho_i$. Increment i .

11: **end while**

12: **while** $\sigma_i^2 \leq C^2 n$ **do**

13: If $\sigma_i^2 = C^2 n$, then **leave** this loop after this iteration. Set

$$\begin{cases} \sigma_{i+1}^2 = C^2 n \wedge \sigma_i^2(1 + \frac{\sigma_i^2}{C^2 n}) & \& \rho_{i+1} = n & \text{if } \sigma_i^2 < C^2 n, \\ \sigma_{i+1}^2 = \infty & \& \rho_{i+1} = n & \text{if } \sigma_i^2 = C^2 n. \end{cases}$$

14: $(E_i, \{Z_i^s\}_{s \in [m]}) \leftarrow \text{Samp-Est}(\{Z_{i-1}^s\}_{s \in [m]}, f_i, f_{i+1}, \epsilon)$ and increment i .

15: **end while**

16: Return $E = n(2\pi n^{-1})^{n/2}(15+13l) \times E_{-1} E_0 \cdots E_i$ as the estimate of the integral of f .

Definition 5.3 (α -mixing, [Ros56]). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability triplet. Given two sub σ -algebras \mathcal{A} and \mathcal{B} , their α -mixing coefficient is defined by

$$\alpha(\mathcal{A}, \mathcal{B}) := \sup_{(A, B) \in \mathcal{A} \times \mathcal{B}} |\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)|.$$

For two random variables $X, Y : \Omega \rightarrow \mathbb{R}$, its α -mixing coefficient is defined by

$$\alpha(X, Y) = \sup_{(A, B) \in \sigma(X) \times \sigma(Y)} |\mathbb{P}(X \in A, Y \in B) - \mathbb{P}(X \in A)\mathbb{P}(Y \in B)|.$$

This notion naturally arises when bounding the covariance between X and Y through the following inequality due to [Dav68]: for $p, q, r \in \mathbb{R}_{>0}$ with $p^{-1} + q^{-1} + r^{-1} = 1$, and $\alpha = \alpha(X, Y)$,

$$|\text{Cov}(X, Y)| \leq 2\alpha^{1/p} \|X\|_q \|Y\|_r.$$

Previous volume-computation work used this inequality with $p = 1$ and $q = r = \infty$. Since the estimators Y_i are unbounded, they had to introduce several technical tools to handle this issue.

New tools. We provide new approaches that can handle these technicalities in a much cleaner way. As for the inexact distribution of samples (i.e., $\bar{\mu}_i$ instead of μ_i), PS_{ann} provides a provable guarantee on $\|\frac{d\bar{\mu}_i}{d\mu_i} - 1\|_{L^\infty}$, not just $\|\bar{\mu}_i - \mu_i\|_{\text{TV}}$, and this allows us to bypass a coupling argument. As for the dependence of samples, we use the notion of β -mixing (or the *coefficient of absolute regularity*) instead of α -mixing, which is viable also due to stronger guarantees of our samplers.

Definition 5.4 (β -mixing, [KR60]). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability triplet. Given two sub σ -algebras \mathcal{A} and \mathcal{B} , their β -mixing coefficient is defined by

$$\beta(\mathcal{A}, \mathcal{B}) := \|\mathbb{P}_{\sigma(\mathcal{A} \otimes \mathcal{B})} - \mathbb{P}_{\mathcal{A}} \otimes \mathbb{P}_{\mathcal{B}}\|_{\text{TV}}.$$

For two random variables $X, Y : \Omega \rightarrow \mathbb{R}$, it has an equivalent definition of

$$\beta(X, Y) = \|\text{law}(X, Y) - \text{law } X \otimes \text{law } Y\|_{\text{TV}}.$$

It holds that $2\alpha \leq \beta$ in general (see [Dou12, Proposition 1]), so β -mixing is stronger than α -mixing. When Y is generated by taking k steps of a Markov chain with kernel P started at X , it holds that (see [Dav74, Proposition 1])

$$\beta(X, Y) = \int_{\Omega} \|\delta_x P^k - \text{law } Y\|_{\text{TV}} (\text{law } X)(dx).$$

Recall that when PS_{ann} generates $Y \sim \bar{\mu}_{i+1}$ from $X \sim \bar{\mu}_i$, it follows from Lemma 3.5 that for $k = \tilde{O}(n^2(\sigma^2 \vee l^2) \text{polylog} \frac{R}{\varepsilon})$,

$$\sup_{x \in \mathcal{K}} \|\delta_x P^k - \mu_{i+1}\|_{\text{TV}} \leq \varepsilon,$$

and thus

$$\beta(X, Y) = \int_{\mathcal{K}} \|\delta_x P^k - \bar{\mu}_{i+1}\|_{\text{TV}} d\bar{\mu}_i(x) \leq \int_{\mathcal{K}} \|\delta_x P^k - \mu_{i+1}\|_{\text{TV}} d\bar{\mu}_i(x) + \|\mu_{i+1} - \bar{\mu}_{i+1}\|_{\text{TV}} \leq 2\varepsilon. \quad (5.2)$$

Therefore, $\|\text{law}(X, Y) - \text{law } X \otimes \text{law } Y\| \leq 2\varepsilon$, and it implies that we can “decouple” X and Y (i.e., use $\text{law } X \otimes \text{law } Y = \bar{\mu}_i \otimes \bar{\mu}_{i+1}$ instead of $\text{law}(X, Y)$) at the cost of small probability 2ε .

5.2.2 Sample and query complexities

We now analyze the variance of estimators and query complexities at initialization, Phase I, Phase II, and Phase III. We begin with a useful observation.

Lemma 5.5. For any $i = -1, 0, \dots, p$,

$$\text{var}_{\mu_i} \left(\frac{E_i}{\mathbb{E}_{\mu_i} E_i} \right) = \frac{1}{m} \text{var}_{\mu_i} \left(\frac{Y_i}{\mathbb{E}_{\mu_i} Y_i} \right) = \frac{1}{m} \text{var}_{\mu_i} \left(\frac{d\mu_{i+1}}{d\mu_i} \right) = \frac{1}{m} \chi^2(\mu_{i+1} \parallel \mu_i).$$

Proof. The first equality is obvious from independence of samples Z_i^s across threads $s \in [m]$. For the second, due to

$$\frac{Y_i}{\mathbb{E}_{\mu_i} Y_i} = \frac{f_{i+1}}{f_i} \frac{F_i}{F_{i+1}},$$

and $\mu_{i+1} \ll \mu_i$, we clearly have

$$\text{var}_{\mu_i} \left(\frac{Y_i}{\mathbb{E}_{\mu_i} Y_i} \right) = \text{var}_{\mu_i} \left(\frac{d\mu_{i+1}}{d\mu_i} \right),$$

and the third one follows from the definition of χ^2 . \square

We examine query and sample complexities for bounding the variance of estimator E_i by v (i.e., $\mathbb{E}_{\mu_i} [E_i^2] \leq (1 + v) (\mathbb{E}_{\mu_i} E_i)^2$).

Initialization. At initialization, Integral draws samples from μ_{-1} and μ_0 in Line 2 through rejection sampling. Below, we will set the target variance $v_0 = \frac{\epsilon^2}{10^9}$, which requires $m_0 \geq 10^{11}l/\epsilon^2$.

Lemma 5.6 (Initialization). *Given $v > 0$, in Line 2, Integral needs $m \geq \frac{100l}{v}$ samples to bound the relative variance $\text{var}(E/\mathbb{E}E)$ by v , using $\mathcal{O}(m)$ queries in expectation.*

Proof. For Line 2, we can show $\mathcal{R}_\infty(\mu_0 \parallel \mu_{-1}) = \log \mathcal{O}(l)$ as in the proof of Lemma 3.12:

$$\begin{aligned} \frac{d\mu_0}{d\mu_{-1}} &= \frac{\exp(-\frac{n}{2}\|x\|^2) \cdot \mathbf{1}_{\bar{\mathcal{K}}}(x, t) \int_{\mathbb{R}^n} \exp(-\frac{n}{2}\|x\|^2) dx \times |I|}{\exp(-\frac{n}{2}\|x\|^2) \cdot \mathbf{1}_I(t) \int_{\bar{\mathcal{K}}} \exp(-\frac{n}{2}\|x\|^2) dx dt} \\ &\leq (15 + 13l) \frac{\int_{\mathbb{R}^n} \exp(-\frac{n}{2}\|x\|^2) dx}{\int_{B_1^n(0) \times [0,1]} \exp(-\frac{n}{2}\|x\|^2) dx} \leq (15 + 13l) (\mathbb{P}_{\mathcal{N}(0, n^{-1}I_n)}(\|X\| \leq 1))^{-1} \\ &\lesssim l. \end{aligned}$$

Thus, by Lemma 5.5 and $\mathcal{R}_2 = \log(1 + \chi^2) \leq \mathcal{R}_\infty$,

$$\text{var}_{\mu_{-1}}\left(\frac{E_{-1}}{\mathbb{E}_{\mu_{-1}}E_{-1}}\right) = \frac{1}{m} \chi^2(\mu_0 \parallel \mu_{-1}) \leq \frac{100l}{m} \leq v.$$

Also, computing E_{-1} uses m many evaluation queries. \square

Phase I. We now analyze the variance of E_i under the update rule of $\sigma^2 \leftarrow \sigma^2(1 + \frac{1}{n})$. To this end, we recall a useful lemma from [KV06, Lemma 3.2]:

Lemma 5.7. *Let $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ be an integrable logconcave function. For $a > 0$ and $Z(a) = \int f(z)^a dz$, the function $a \mapsto a^n Z(a)$ is logconcave in a .*

Below, we will set the target variance $v_{1,1} = \frac{\epsilon^2}{10^9 n \log n}$ during $\sigma^2 < 1$ and $v_{1,2} = \frac{\epsilon^2}{10^9}$ when $\sigma^2 = 1$, each of which requires $m_{1,1} \geq \frac{3 \cdot 10^9 \log n}{\epsilon^2}$ and $m_{1,2} \geq \frac{2 \cdot 10^9}{\epsilon^2}$, respectively.

Lemma 5.8 (Phase I: Line 3). *Given $v > 0$ and $\epsilon > 0$, if $\sigma^2 < 1$, then Integral needs $m \geq \frac{3}{nv}$ samples to ensure $\text{var}_\mu(E/\mathbb{E}_\mu E) \leq v$. Also, when PS_{ann} draws m samples with law $\bar{\mu}$ satisfying $\mathcal{R}_\infty(\bar{\mu} \parallel \mu) \leq \epsilon$ in each inner phase, it requires $\tilde{\mathcal{O}}(mn^3 l^2 \log^4 \frac{R}{\epsilon})$ queries throughout Phase I. When $\sigma^2 = 1$, Integral needs $m \geq \frac{2}{v}$ samples and uses $\tilde{\mathcal{O}}(mn^2 l^2 \log^4 \frac{R}{\epsilon})$ queries.*

Proof. For $\alpha = 1/n$, let us denote

$$\mu_i(x, t) \propto \exp\left(-\frac{1}{2\sigma^2(1+\alpha)}\|x\|^2\right)\Big|_{\bar{\mathcal{K}}} \quad \& \quad \mu_{i+1}(x, t) \propto \exp\left(-\frac{1}{2\sigma^2}\|x\|^2\right)\Big|_{\bar{\mathcal{K}}}.$$

It then follows from the definition of L^2 -norm that

$$\left\| \frac{d\mu_{i+1}}{d\mu_i} \right\|_{L^2(\mu_i)}^2 = \frac{\int \exp(-\frac{1-\alpha}{2\sigma^2}\|x\|^2) \cdot \mathbf{1}_{\bar{\mathcal{K}}} \times \int_{\bar{\mathcal{K}}} \exp(-\frac{1+\alpha}{2\sigma^2}\|x\|^2) \cdot \mathbf{1}_{\bar{\mathcal{K}}}}{\left(\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2\sigma^2}\|x\|^2) \cdot \mathbf{1}_{\bar{\mathcal{K}}}\right)^2} = \frac{F(1-\alpha)F(1+\alpha)}{F(1)^2},$$

where $F(s) := \int \exp(-\frac{s}{2\sigma^2}\|x\|^2) \cdot \mathbf{1}_{\bar{\mathcal{K}}}$. Using logconcavity of $s^{n+1}F(s)$ in s (due to Lemma 5.7),

$$\frac{F(1-\alpha)F(1+\alpha)}{F(1)^2} \leq \frac{1}{(1-\alpha^2)^{n+1}} \leq 1 + \frac{3}{n}.$$

Since $1 + \chi^2(\mu_{i+1} \parallel \mu_i) = \|\mu_{i+1}/\mu_i\|_{L^2(\mu_i)}^2$, we can conclude that

$$\text{var}_{\mu_i}\left(\frac{E_i}{\mathbb{E}_{\mu_i}E_i}\right) = \frac{1}{m} \chi^2(\mu_{i+1} \parallel \mu_i) \leq \frac{3}{mn}.$$

Hence, it suffices to take $m \geq \frac{3}{nv}$ samples for estimation. Regarding the query complexity, there are $\mathcal{O}(n \log n)$ inner phases. Also, μ_{i-1} is always $\mathcal{O}(1)$ -warm with respect to μ_i , as in Lemma 3.13, thus PS_{ann} uses $\tilde{\mathcal{O}}(n^2 l^2 \log^4 \frac{R}{\epsilon})$ queries in expectation (to obtain one sample which is ϵ -close to μ_i in \mathcal{R}_∞). Therefore, Phase I requires $\tilde{\mathcal{O}}(mn^3 l^2 \log^4 \frac{R}{\epsilon})$ queries.

When $\sigma^2 = 1$, we let $\mu_i \propto \exp(-\frac{1}{2} \|x\|^2)|_{\bar{\mathcal{K}}}$ and $\mu_{i+1} \propto \exp(-\frac{1}{2} \|x\|^2 - \rho t)|_{\bar{\mathcal{K}}}$ for $\rho = \frac{1}{28l}$. Due to

$$\frac{d\mu_{i+1}}{d\mu_i} = \sup_{\bar{\mathcal{K}}} \frac{\exp(-\rho t) \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2} \|x\|^2)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2} \|x\|^2 - \rho t)} \leq \sup_{\bar{\mathcal{K}}} \exp(\rho(t - \min t)) \leq e,$$

it follows that for $m \geq 2/v$,

$$\text{var}_{\mu_i} \left(\frac{E_i}{\mathbb{E} E_i} \right) = \frac{1}{m} \chi^2(\mu_{i+1} \parallel \mu_i) \leq \frac{2}{m} \leq v.$$

Since μ_{i-1} is $\mathcal{O}(1)$ -warm with respect to μ_i , PS_{ann} uses $\tilde{\mathcal{O}}(mn^2 l^2 \log^4 \frac{R}{\epsilon})$ queries in expectation. \square

Phase II. Variance control in Phase II requires an additional lemma in [CV18, Lemma 7.8]. We present it with a streamlined proof.

Lemma 5.9. *Suppose a logconcave function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has support contained in $B_R(0)$. Let X be drawn from $\mu_{\sigma^2} \propto g(x, \sigma^2) := f(x) e^{-\|x\|^2/(2\sigma^2)}$ and $Y = g(X, \sigma^2(1+\alpha))/g(X, \sigma^2)$. For $\alpha \leq 1/2$,*

$$\text{var} \left(\frac{Y}{\mathbb{E} Y} \right) = \chi^2(\mu_{\sigma^2(1+\alpha)} \parallel \mu_{\sigma^2}) \leq \exp\left(\frac{2R^2\alpha^2}{\sigma^2}\right) - 1.$$

Proof. We can directly show that

$$\left\| \frac{d\mu_{\sigma^2(1+\alpha)}}{d\mu_{\sigma^2}} \right\|_{L^2}^2 = \frac{\int f(x) \exp(-\frac{1}{2\sigma^2} \|x\|^2) \times \int f(x) \exp(-\frac{1}{2\sigma^2} \frac{1+\alpha}{1+\alpha} \|x\|^2)}{(\int f(x) \exp(-\frac{1}{2\sigma^2(1+\alpha)} \|x\|^2))^2}.$$

Defining $g(s) := \log(\int f(x) \exp(-\frac{s}{2\sigma^2(1+\alpha)} \|x\|^2))$, we have

$$\mathcal{R}_2(\mu_{\sigma^2(1+\alpha)} \parallel \mu_{\sigma^2}) = g(1+\alpha) + g(1-\alpha) - 2g(1) = \int_0^\alpha (g'(1+t) - g'(1-t)) dt = \int_0^\alpha \int_{1-t}^{1+t} g''(q) dq dt. \quad (5.3)$$

For a logconcave distribution $\nu_q(x) \propto f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)$, we have

$$\begin{aligned} g''(q) &= -\frac{1}{2\sigma^2(1+\alpha)} \frac{d}{dq} \left(\frac{\int \|x\|^2 f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)}{\int f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)} \right) \\ &= \frac{1}{4\sigma^4(1+\alpha)^2} \left[\frac{\int \|x\|^4 f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)}{\int f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)} - \left(\frac{\int \|x\|^2 f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)}{\int f(x) \exp(-\frac{q}{2\sigma^2(1+\alpha)} \|x\|^2)} \right)^2 \right] \\ &= \frac{1}{4\sigma^4(1+\alpha)^2} \text{var}_{\nu_q}(\|X\|^2) \stackrel{\text{(PI)}}{\leq} \frac{1}{q\sigma^2(1+\alpha)} \mathbb{E}_{\nu_q}[\|X\|^2], \end{aligned}$$

where we used $C_{\text{PI}}(\nu_q) \leq \frac{\sigma^2(1+\alpha)}{q}$ (Bakry-Émery) in the last line. Due to $\sup_{\text{supp } \nu_q} \|X\| \leq R$,

$$g''(q) \leq \frac{R^2}{q\sigma^2}.$$

Putting this bound back to (5.3) and integrating,

$$\mathcal{R}_2(\mu_{\sigma^2(1+\alpha)} \parallel \mu_{\sigma^2}) \leq \frac{R^2}{\sigma^2} \int_0^\alpha \int_{1-t}^{1+t} \frac{1}{q} dq dt = \frac{R^2}{\sigma^2} \int_0^\alpha (\log(1+t) - \log(1-t)) dt \leq \frac{2R^2\alpha^2}{\sigma^2},$$

which completes the proof. \square

We now establish sample and query complexities under the two update rules: for $1 \leq \beta \leq 1 + \frac{1}{28nl}$, (i) $\sigma^2 \leftarrow \beta^{-1}\sigma^2$ and $\rho \leftarrow \beta\rho$, and (ii) $\sigma^2 \leftarrow \beta\sigma^2$. Below, we will set the target variance $v_2 = \frac{\epsilon^2}{2 \cdot 10^9 n \log n}$, which requires $m_2 \geq \frac{2 \cdot 10^9 \log nl}{\epsilon^2}$.

Lemma 5.10 (Phase II: Line 7). *Given $v > 0$ and $\epsilon > 0$, Integral needs $m \geq \frac{1}{nlv}$ samples to ensure $\text{var}_\mu(E/\mathbb{E}_\mu E) \leq v$. Also, when PS_{ann} draws m samples with law $\bar{\mu}$ satisfying $\mathcal{R}_\infty(\bar{\mu} \parallel \mu) \leq \epsilon$ in each inner phase, it requires $\tilde{\mathcal{O}}(mn^3 l^3 \log^4 \frac{R}{\epsilon})$ queries throughout Phase II.*

Proof. In the first update of $\sigma_{i+1}^2 = \sigma_i^2/\beta$ and $\rho_{i+1} = \beta\rho_i$, we can write $\mu_i(x, t) \propto \exp(-U(x, t))$ and $\mu_{i+1}(x, t) \propto \exp(-\beta U(x, t))$ for $U(x, t) = (\frac{1}{2} \|x\|^2 + \rho t)/\mathbf{1}_{\bar{\mathcal{K}}}(x, t)$, where $\frac{1}{28l} \leq \rho < n$. Then,

$$\left\| \frac{d\mu_{i+1}}{d\mu_i} \right\|_{L^2(\mu_i)}^2 = \frac{\int_{\mathbb{R}^{n+1}} \exp(-(2\beta - 1)U) \times \int_{\mathbb{R}^{n+1}} \exp(-U)}{(\int_{\mathbb{R}^{n+1}} \exp(-\beta U))^2}.$$

Using logconcavity of $s^{n+1} \int \exp(-sU)$ in s with endpoints $2\beta - 1$ and 1 (due to Lemma 5.7),

$$\frac{\int_{\mathbb{R}^{n+1}} \exp(-(2\beta - 1)U) \times \int_{\mathbb{R}^{n+1}} \exp(-U)}{(\int_{\mathbb{R}^{n+1}} \exp(-\beta U))^2} \leq \frac{\beta^{2n}}{(2\beta - 1)^n} \leq 1 + \frac{1}{nl}.$$

Since $1 + \chi^2(\mu_{i+1} \parallel \mu_i) = \|\mu_{i+1}/\mu_i\|_{L^2(\mu_i)}^2$, we can conclude that

$$\text{var}_{\mu_i} \left(\frac{E_i}{\mathbb{E}_{\mu_i} E_i} \right) = \frac{1}{m} \chi^2(\mu_{i+1} \parallel \mu_i) \leq \frac{1}{mnl},$$

so it suffices to take $m \geq \frac{1}{nlv}$ samples for estimation.

In the second update of $\sigma_{i+1}^2 = \beta\sigma_i^2 = 1$ and $\rho_{i+1} = \rho_i$, we can write $\mu_i^X \propto g(x) \exp(-\frac{\beta}{2} \|x\|^2)$ and $\mu_{i+1}^X \propto g(x) \exp(-\frac{1}{2} \|x\|^2)$ for some logconcave function g . Then, for $Y = \frac{\exp(-\|x\|^2/2)}{\exp(-\beta\|x\|^2/2)}$,

$$\begin{aligned} \text{var}_{\mu_i} \left(\frac{E_i}{\mathbb{E}_{\mu_i} E_i} \right) &= \frac{1}{m} \text{var}_{\mu_i} \left(\frac{Y}{\mathbb{E}_{\mu_i} Y} \right) = \frac{1}{m} \text{var}_{\mu_i} \left(\frac{d\mu_{i+1}^X}{d\mu_i^X} \right) = \frac{1}{m} \text{var}_{\mu_i^X} \left(\frac{d\mu_{i+1}^X}{d\mu_i^X} \right) \\ &\stackrel{(i)}{\leq} \frac{1}{m} \left(\exp\left(\frac{2C^2 n (\beta - 1)^2}{1/\beta}\right) - 1 \right) \leq \frac{1}{mnl}, \end{aligned} \quad (5.4)$$

where we used Lemma 5.9 in (i).

Regarding the query complexity, we show that μ_{i-1} is $\mathcal{O}(1)$ -warm with respect to μ_i . At the beginning of Phase II, for $t_0 := \inf_{\bar{\mathcal{K}}} t$,

$$\frac{d\mu_{i-1}}{d\mu_i} = \sup_{\bar{\mathcal{K}}} \frac{\exp(-\frac{1}{2} \|x\|^2)}{\exp(-\frac{1}{2} \|x\|^2 - \frac{1}{28l} t)} \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2} \|x\|^2 - \frac{1}{28l} t)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2} \|x\|^2)} \leq \sup_{\bar{\mathcal{K}}} \exp\left(\frac{1}{28l} (t - t_0)\right) \leq e.$$

In the first update,

$$\begin{aligned} \frac{d\mu_i}{d\mu_{i+1}} &= \sup_{\bar{\mathcal{K}}} \frac{\exp(-\frac{1}{2} \|x\|^2 - \rho t)}{\exp(-\frac{\beta}{2} \|x\|^2 - \beta\rho t)} \frac{\int_{\bar{\mathcal{K}}} \exp(-\frac{\beta}{2} \|x\|^2 - \beta\rho t)}{\int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2} \|x\|^2 - \rho t)} \leq \sup_{\bar{\mathcal{K}}} \frac{\exp(-\frac{1}{2} \|x\|^2 - \rho(t - t_0))}{\exp(-\frac{\beta}{2} \|x\|^2 - \beta\rho(t - t_0))} \\ &\leq \sup \exp\left(\frac{1}{56nl} \|x\|^2 + \frac{\rho}{28nl} (t - t_0)\right) \leq e^2. \end{aligned}$$

In the second update, the warmness can be shown as in Lemma 3.13. Putting these together, PS_{ann} uses $\tilde{\mathcal{O}}(n^2 l^2 \log^4 \frac{R}{\epsilon})$ queries per inner phase, and there are $\mathcal{O}(nl \log nl)$ inner phases in Phase II, so we need $\tilde{\mathcal{O}}(mn^3 l^3 \log^4 \frac{R}{\epsilon})$ queries throughout Phase II. \square

Phase III. We establish sample and query complexity under the update rule of $\sigma^2 \leftarrow \sigma^2 (1 + \frac{\sigma^2}{C^2 n})$. Since σ^2 increases from 1 to $C^2 n$, we can partition the whole phase by $\mathcal{O}(\log C^2 n)$ many doubling phases, where an initial σ^2 doubles in each doubling phase. Below, we will set the target variance $v_{3,1} = \frac{\epsilon^2 \sigma^2}{10^9 C^2 n \log C^2 n}$ during each doubling phase, where σ^2 is an initial value in the doubling phase. This requires $m_{3,1} \geq \frac{4 \cdot 10^9 \log C^2 n}{\epsilon^2}$. When $\sigma^2 = C^2 n$, we will set $v_{3,2} = \frac{\epsilon^2}{10^9}$, which require $m_{3,2} \geq \frac{2 \cdot 10^9}{\epsilon^2}$.

Lemma 5.11 (Phase III: Line 12). *Given $v > 0$ and $\epsilon > 0$, Integral needs $m \geq \frac{4\sigma^2}{C^2 n v}$ samples to ensure $\text{var}_\mu(E/\mathbb{E}_\mu E) \leq v$. Also, when PS_{ann} draws m samples with law $\bar{\mu}$ satisfying $\mathcal{R}_\infty(\bar{\mu} \parallel \mu) \leq \epsilon$ in each inner phase, it requires $\tilde{\mathcal{O}}(m C^2 n^3 l^2 \log^4 \frac{R}{\epsilon})$ queries during doubling of σ^2 . When $\sigma^2 = C^2 n$, Integral needs $m \geq \frac{2}{v}$ samples and uses $\tilde{\mathcal{O}}(m n^2 l^2 \log^4 \frac{R}{\epsilon})$ queries.*

Proof. Regarding the variance control, similar to (5.4), we can use Lemma 5.9 (with $R = C\sqrt{n}$ and $\alpha = \frac{\sigma^2}{C^2 n}$) to show that for $\sigma^2 \in [1, C^2 n]$,

$$\text{var}_{\mu_i} \left(\frac{E_i}{\mathbb{E} E_i} \right) = \frac{1}{m} \text{var} \left(\frac{Y}{\mathbb{E} Y} \right) \leq \frac{1}{m} \left(\exp \left(\frac{2C^2 n \alpha^2}{\sigma^2} \right) - 1 \right) \leq \frac{4\sigma^2}{C^2 m n}.$$

Hence, it suffices to take $m \geq \frac{4\sigma^2}{C^2 n v}$ samples for estimation. Next, we can show that μ_{i-1} is $\mathcal{O}(1)$ -warm with respect to μ_i during Phase III, as in the proof of Lemma 3.14. Since there are $\mathcal{O}(\frac{C^2 n}{\sigma^2})$ inner phases (for doubling of σ^2), and PS_{ann} requires $\tilde{\mathcal{O}}(n^2 \sigma^2 l^2 \log^4 \frac{R}{\epsilon})$ queries, Phase III uses $\tilde{\mathcal{O}}(m C^2 n^3 l^2 \log^4 \frac{R}{\epsilon})$ queries in expectation during each doubling phase.

When $\sigma^2 = C^2 n$, we let $\mu_i \propto \exp(-\frac{1}{2C^2 n} \|x\|^2 - nt)|_{\bar{\mathcal{K}}}$ and $\mu_{i+1} = \bar{\pi} \propto \exp(-nt)|_{\bar{\mathcal{K}}}$. Due to

$$\frac{d\mu_{i+1}}{d\mu_i} = \sup_{\bar{\mathcal{K}}} \frac{\exp(\frac{1}{2C^2 n} \|x\|^2) \int_{\bar{\mathcal{K}}} \exp(-\frac{1}{2C^2 n} \|x\|^2 - nt)}{\int_{\bar{\mathcal{K}}} \exp(-nt)} \leq \sqrt{e},$$

it follows that for $m \geq 2/v$,

$$\text{var}_{\mu_i} \left(\frac{E_i}{\mathbb{E} E_i} \right) = \frac{1}{m} \chi^2(\mu_{i+1} \parallel \mu_i) \leq \frac{2}{m} \leq v,$$

and to sample from μ_i , PS_{ann} uses $\tilde{\mathcal{O}}(m n^2 l^2 \log^4 \frac{R}{\epsilon})$ queries in expectation. \square

5.2.3 Final guarantee

With these bounds on the variance of estimators in **Integral**, we can bound the overall variance of the product of estimates using m samples. We then show that our final estimate in **Integral** is close to the integral $\int e^{-V}$ with high probability, streamlining previous analysis in [KLS97, LV06c, CV18].

Lemma 5.12. *If Integral runs with sample size $m = \epsilon^{-2}(10^{11} l \log n l \log C^2 n)$ and sample accuracy $\epsilon \leq \epsilon^2(10^{15} C^2 n l^2 \log^3(C^2 n l))^{-1}$, then it can compute a $(1 + \epsilon)$ -multiplicative approximation to $\int e^{-V}$ with probability at least 0.98.*

Proof. We note that the given sample size m satisfies all the required conditions (i.e., $m \geq m_0, m_{1,*}, m_2, m_{3,*}$) to attain a desired target variance in each phase.

We first assume that all samples $\{Z_i^s\}_{0 \leq i \leq p, s \in [m]}$ for estimations are independent (so $\{E_i\}_i$ are independent) and exactly distributed according to μ_i (not $\bar{\mu}_i$). Then, for $E = E_{-1} E_0 \cdots E_p$,

$$\text{var} E = \mathbb{E}[E_{-1}^2] \cdots \mathbb{E}[E_p^2] - (\mathbb{E} E_{-1})^2 \cdots (\mathbb{E} E_p)^2 = \prod_i (\text{var} E_i + (\mathbb{E} E_i)^2) - \prod_i (\mathbb{E} E_i)^2$$

$$\leq \prod_i (1 + v_i) (\mathbb{E}E_i)^2 - \prod_i (\mathbb{E}E_i)^2 = \left(\prod_i (1 + v_i) - 1 \right) \prod_i (\mathbb{E}E_i)^2 \leq \frac{\varepsilon^2}{10^8} \prod_i (\mathbb{E}E_i)^2,$$

where the last line follows from

$$\begin{aligned} \prod_i (1 + v_i) &\leq \exp\left(v_0 + v_{1,1} \times n \log n + v_{1,2} + v_2 \times 2nl \log nl + (v_{3,1} \times \frac{C^2 n}{\sigma^2}) \log C^2 n + v_{3,2}\right) \\ &\leq \exp\left(\frac{6\varepsilon^2}{10^9}\right) \leq 1 + \frac{\varepsilon^2}{10^8}. \end{aligned}$$

By Chebyshev's inequality,

$$\mathbb{P}_{(\otimes_i \mu_i)^{\otimes m}}(|E - \mathbb{E}E| \geq \frac{\varepsilon}{10^3} \mathbb{E}E) \leq 0.01.$$

Hence, (5.1) implies that with probability at least 0.99,

$$(1 - \varepsilon) \int f \leq E = n (2\pi n^{-1})^{n/2} (15 + 13l) \times E_{-1} E_0 \cdots E_i \leq (1 + \frac{\varepsilon}{10^3}) \int f. \quad (5.5)$$

We now take into two sources of complications: (1) inexact distributions of samples, and (2) dependence of samples within each thread. As for (1), since PS_{ann} ensures $\frac{\bar{\mu}_i}{\mu_i} \leq 1 + \varepsilon$, we have

$$\frac{\text{law } Z_1 \otimes \cdots \otimes \text{law } Z_p}{\mu_1 \otimes \cdots \otimes \mu_p} = \bigotimes_i \frac{\bar{\mu}_i}{\mu_i} \leq e^{2p\varepsilon},$$

and thus

$$\mathbb{P}_{(\otimes \bar{\mu}_i)^{\otimes m}}(|E - \mathbb{E}E| \geq \frac{\varepsilon}{10^3} \mathbb{E}E) \leq e^{2mp\varepsilon} \mathbb{P}_{\otimes \mu_i}(|E - \mathbb{E}E| \geq \frac{\varepsilon}{10^3} \mathbb{E}E) \leq 10^{-2} e^{2mp\varepsilon}.$$

Regarding (2), the triangle inequality results in

$$\begin{aligned} &\left\| \text{law}(Z_1, \dots, Z_p) - \bigotimes_{i=1}^p \text{law } Z_i \right\|_{\text{TV}} \leq \\ &\left\| \text{law}(Z_1, \dots, Z_p) - \text{law}(Z_1, \dots, Z_{p-1}) \otimes \text{law } Z_p \right\|_{\text{TV}} + \left\| \text{law}(Z_1, \dots, Z_{p-1}) \otimes \text{law } Z_p - \bigotimes_{i=1}^p \text{law } Z_i \right\|_{\text{TV}}. \end{aligned}$$

The first term equals $\left\| \text{law}(Z_{p-1}, Z_p) - \text{law } Z_{p-1} \otimes \text{law } Z_p \right\|_{\text{TV}}$, as $\{Z_i\}_i$ are generated by a Markov chain (see Lemma A.4 for details). Applying the DPI to the second term, we obtain that

$$\begin{aligned} &\left\| \text{law}(Z_1, \dots, Z_p) - \bigotimes_{i=1}^p \text{law } Z_i \right\|_{\text{TV}} \\ &= \left\| \text{law}(Z_{p-1}, Z_p) - \text{law } Z_{p-1} \otimes \text{law } Z_p \right\|_{\text{TV}} + \left\| \text{law}(Z_1, \dots, Z_{p-1}) - \bigotimes_{i=1}^{p-1} \text{law } Z_i \right\|_{\text{TV}}. \end{aligned}$$

As elaborated in (5.2), the first term is bounded by ε . Repeating the same argument on the second term, we have

$$\left\| \text{law}(Z_1, \dots, Z_p) - \bigotimes_{i=1}^p \text{law } Z_i \right\|_{\text{TV}} \leq 2p\varepsilon.$$

Using a similar argument based on the triangle inequality (see Lemma A.4 for details),

$$\left\| \text{law}(\{Z_i^s\}_{i,s}) - \left(\bigotimes_{i=1}^p \text{law } Z_i \right)^{\otimes m} \right\|_{\text{TV}} = \left\| (\text{law}(Z_1, \dots, Z_p))^{\otimes m} - \left(\bigotimes_{i=1}^p \text{law } Z_i \right)^{\otimes m} \right\|_{\text{TV}} \leq 2mp\epsilon.$$

Therefore, it holds from the definition of TV-distance that

$$\begin{aligned} \mathbb{P}_{(\text{law}(Z_1, \dots, Z_p))^{\otimes m}} (|E - \mathbb{E}E| \geq \frac{\epsilon}{10^3} \mathbb{E}E) &\leq \mathbb{P}_{(\otimes_i \bar{\mu}_i)^{\otimes m}} (|E - \mathbb{E}E| \geq \frac{\epsilon}{10^3} \mathbb{E}E) + 2mp\epsilon \\ &\leq 10^{-2} e^{2mp\epsilon} + 2mp\epsilon, \end{aligned}$$

and this is bounded by 0.02 when $\epsilon = (300mp)^{-1} \leq \epsilon^2 (10^{15} C^2 n l^2 \log^3(C^2 n l))^{-1}$. Therefore, (5.5) holds with probability at least 0.98. \square

We are now ready to prove the main result of this section, as claimed in Theorem 1.8.

Theorem 5.13. *In the setting of Problem 5.1, for any integrable logconcave $f : \mathbb{R}^n \rightarrow \mathbb{R}$ given by a well-rounded function oracle and any $\epsilon > 0$, with probability at least $3/4$, Integral (Algorithm 6) returns a $(1 + \epsilon)$ -multiplicative approximation to the integral of f using $\tilde{\mathcal{O}}(n^3/\epsilon^2)$ queries. For an arbitrary logconcave f , the total query complexity is bounded by $\tilde{\mathcal{O}}(n^{3.5} \text{polylog } R + n^3/\epsilon^2)$.*

Proof. By Lemma 5.12, we must have the sample size $m = \tilde{\mathcal{O}}(\frac{l \log n}{\epsilon^2})$ and sample accuracy $\epsilon = \mathcal{O}(\frac{\epsilon^2}{C^2 n l^2 \log^3(C^2 n l)})$. These conditions will decide query and sample complexities in each phase. At initialization (Lemma 5.6), under the choices of m and ϵ , we need

$$\text{Query: } \mathcal{O}(m) = \tilde{\mathcal{O}}\left(\frac{l \log n}{\epsilon^2}\right).$$

In Phase I (Lemma 5.8), we need

$$\begin{aligned} \sigma^2 < 1 : \text{Query: } &\tilde{\mathcal{O}}(mn^3 l^2 \log^4 \frac{R}{\epsilon}) = \tilde{\mathcal{O}}\left(\frac{n^3 l^4}{\epsilon^2}\right). \\ \sigma^2 = 1 : \text{Query: } &\tilde{\mathcal{O}}(mn^2 l^2 \log^4 \frac{R}{\epsilon}) = \tilde{\mathcal{O}}\left(\frac{n^2 l^3}{\epsilon^2}\right). \end{aligned}$$

In Phase II (Lemma 5.10), we need

$$\text{Query: } \tilde{\mathcal{O}}(mn^3 l^3 \log^4 \frac{R}{\epsilon}) = \tilde{\mathcal{O}}\left(\frac{n^3 l^4}{\epsilon^2}\right).$$

In Phase III (Lemma 5.11), we need

$$\text{Query: } \tilde{\mathcal{O}}(m C^2 n^3 l^2 \log^4 \frac{R}{\epsilon} \times \log C^2 n) = \tilde{\mathcal{O}}\left(\frac{C^2 n^3 l^3}{\epsilon^2}\right).$$

Putting these together with $l = C^2 \vee \log \frac{8\epsilon}{\epsilon}$, the query complexity of the algorithm is

$$\tilde{\mathcal{O}}\left(\frac{C^5 n^3}{\epsilon^2}\right).$$

For general logconcave f , we first use the rounding algorithm to bring f to well-rounded position, which takes $\tilde{\mathcal{O}}(n^{3.5})$ queries and then apply the above for well-rounded f . \square

Acknowledgement. We are grateful to Andre Wibisono for helpful discussions and to Sinho Chewi for pointing us to the Holley-Stroock principle and the reference [Wan13] used in the proof of Lemma 3.3. This work was supported in part by NSR Award CCF-2106444 and a Simons Investigator grant.

References

- [AK91] David Applegate and Ravi Kannan. Sampling and integration of near log-concave functions. In *Symposium on Theory of Computing (STOC)*, pages 156–163, 1991.
- [BGL14] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 103. Springer, 2014.
- [CCSW22] Yongxin Chen, Sinho Chewi, Adil Salim, and Andre Wibisono. Improved analysis for a proximal algorithm for sampling. In *Conference on Learning Theory (COLT)*, pages 2984–3014. PMLR, 2022.
- [CEL⁺22] Sinho Chewi, Murat A Erdogdu, Mufan Li, Ruoqi Shen, and Shunshi Zhang. Analysis of Langevin Monte Carlo from Poincaré to log-Sobolev. In *Conference on Learning Theory (COLT)*, pages 1–2. PMLR, 2022.
- [Cha04] Djilil Chafaï. Entropies, convexity, and functional inequalities, on Φ -entropies and Φ -sobolev inequalities. *Journal of Mathematics of Kyoto University*, 44(2):325–363, 2004.
- [Che24] Sinho Chewi. *Log-concave sampling*. Book draft available at <https://chewisinho.github.io>, 2024.
- [CV18] Ben Cousins and Santosh S Vempala. Gaussian Cooling and $O^*(n^3)$ algorithms for volume and Gaussian volume. *SIAM Journal on Computing (SICOMP)*, 47(3):1237–1273, 2018.
- [Dav68] Yu A Davydov. Convergence of distributions generated by stationary stochastic processes. *Theory of Probability & Its Applications*, 13(4):691–696, 1968.
- [Dav74] Yu A Davydov. Mixing conditions for Markov chains. *Theory of Probability & Its Applications*, 18(2):312–328, 1974.
- [DFK91] Martin Dyer, Alan Frieze, and Ravi Kannan. A random polynomial-time algorithm for approximating the volume of convex bodies. *Journal of the ACM (JACM)*, 38(1):1–17, 1991.
- [DMLM03] Pierre Del Moral, Michel Ledoux, and Laurent Miclo. On contraction properties of Markov kernels. *Probability Theory and Related Fields*, 126(3):395–420, 2003.
- [Dou12] Paul Doukhan. *Mixing: properties and examples*, volume 85. Springer Science & Business Media, 2012.
- [Fol99] Gerald B Folland. *Real analysis: modern techniques and their applications*, volume 40. John Wiley & Sons, 1999.
- [HS87] Richard Holley and Daniel W Stroock. Logarithmic Sobolev inequalities and stochastic Ising models. *Journal of Statistical Physics*, 46(5/6):1159–1194, 1987.
- [JLLV21] He Jia, Aditi Laddha, Yin Tat Lee, and Santosh S Vempala. Reducing isotropy and volume to KLS: an $O^*(n^3\psi^2)$ volume algorithm. In *Symposium on Theory of Computing (STOC)*, pages 961–974, 2021.

- [JLLV24] He Jia, Aditi Laddha, Yin Tat Lee, and Santosh S Vempala. Reducing isotropy and volume to KLS: Faster rounding and volume algorithms. *arXiv preprint arXiv:2008.02146*, 2024.
- [Kla23] Boáz Klartag. Logarithmic bounds for isoperimetry and slices of convex sets. *Ars Inveniendi Analytica*, 2023.
- [KLS97] Ravi Kannan, László Lovász, and Miklós Simonovits. Random walks and an $O^*(n^5)$ volume algorithm for convex bodies. *Random Structures & Algorithms (RS&A)*, 11(1):1–50, 1997.
- [KLSV22] Yunbum Kook, Yin Tat Lee, Ruoqi Shen, and Santosh S Vempala. Sampling with Riemannian Hamiltonian Monte Carlo in a constrained space. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 35, pages 31684–31696, 2022.
- [KR60] Andrei Nikolaevich Kolmogorov and Yu A Rozanov. On strong mixing conditions for stationary Gaussian processes. *Theory of Probability & Its Applications*, 5(2):204–208, 1960.
- [KV06] Adam Tauman Kalai and Santosh S Vempala. Simulated annealing for convex optimization. *Mathematics of Operations Research*, 31(2):253–266, 2006.
- [KV24] Yunbum Kook and Santosh S Vempala. Gaussian cooling and Dikin walks: The interior-point method for logconcave sampling. In *Conference on Learning Theory (COLT)*, pages 3137–3240. PMLR, 2024.
- [KVZ24] Yunbum Kook, Santosh S Vempala, and Matthew S Zhang. In-and-Out: Algorithmic diffusion for sampling convex bodies. In *To appear in Advances in Neural Information Processing Systems (NeurIPS)*, 2024.
- [KZ24] Yunbum Kook and Matthew S Zhang. Covariance estimation using Markov chain Monte Carlo. *arXiv preprint arXiv:2410.17147*, 2024.
- [KZ25] Yunbum Kook and Matthew S Zhang. Rényi-infinity constrained sampling with d^3 membership queries. In *To appear in Symposium on Discrete Algorithms (SODA)*, 2025.
- [Led94] Michel Ledoux. A simple analytic proof of an inequality by P. Buser. *Proceedings of the American Mathematical Society*, 121(3):951–959, 1994.
- [Lov91] László Lovász. *How to compute the volume?* DIMACS, Center for Discrete Mathematics and Theoretical Computer Science, 1991.
- [LS90] László Lovász and Miklós Simonovits. The mixing rate of Markov chains, an isoperimetric inequality, and computing the volume. In *Symposium on Foundations of Computer Science (FOCS)*, pages 346–354. IEEE, 1990.
- [LS93] László Lovász and Miklós Simonovits. Random walks in a convex body and an improved volume algorithm. *Random Structures & Algorithms (RS&A)*, 4(4):359–412, 1993.
- [LST21] Yin Tat Lee, Ruoqi Shen, and Kevin Tian. Structured logconcave sampling with a restricted Gaussian oracle. In *Conference on Learning Theory (COLT)*, pages 2993–3050. PMLR, 2021.

- [LV06a] László Lovász and Santosh S Vempala. Fast algorithms for logconcave functions: Sampling, rounding, integration and optimization. In *Symposium on Foundations of Computer Science (FOCS)*, pages 57–68. IEEE, 2006.
- [LV06b] László Lovász and Santosh S Vempala. Hit-and-run from a corner. *SIAM Journal on Computing (SICOMP)*, 35(4):985–1005, 2006.
- [LV06c] László Lovász and Santosh S Vempala. Simulated annealing in convex bodies and an $O^*(n^4)$ volume algorithm. *Journal of Computer and System Sciences (JCSS)*, 72(2):392–417, 2006.
- [LV07] László Lovász and Santosh S Vempala. The geometry of logconcave functions and sampling algorithms. *Random Structures & Algorithms (RS&A)*, 30(3):307–358, 2007.
- [LV24] Yin Tat Lee and Santosh S Vempala. Eldan’s stochastic localization and the KLS conjecture: Isoperimetry, concentration and mixing. *Annals of Mathematics*, 199(3):1043–1092, 2024.
- [Mil09] Emanuel Milman. On the role of convexity in isoperimetry, spectral gap and concentration. *Inventiones Mathematicae*, 177(1):1–43, 2009.
- [Pao06] Grigoris Paouris. Concentration of mass on convex bodies. *Geometric and Functional Analysis (GAFA)*, 16(5):1021–1049, 2006.
- [Ros56] Murray Rosenblatt. A central limit theorem and a strong mixing condition. *Proceedings of the National Academy of Sciences (PNAS)*, 42(1):43–47, 1956.
- [RW00] Mary Beth Ruskai and Elisabeth Werner. Study of a class of regularizations of $1/|x|$ using Gaussian integrals. *SIAM Journal on Mathematical Analysis (SIMA)*, 32(2):435–463, 2000.
- [VEH14] Tim Van Erven and Peter Harremoës. Rényi divergence and Kullback-Leibler divergence. *IEEE Transactions on Information Theory*, 60(7):3797–3820, 2014.
- [VW19] Santosh S Vempala and Andre Wibisono. Rapid convergence of the unadjusted Langevin algorithm: Isoperimetry suffices. *Advances in Neural Information Processing Systems (NeurIPS)*, 32, 2019.
- [Wan13] Feng-Yu Wang. *Analysis for diffusion processes on Riemannian manifolds*, volume 18. World Scientific, 2013.

A Functional inequalities and logconcave geometry

Functional inequalities.

Lemma A.1 (Bakry-Émery). *If the potential V is α -strongly convex, then a probability measure $\pi \propto \exp(-V)$ satisfies*

$$C_{\text{PI}}(\pi) \leq C_{\text{LSI}}(\pi) \leq \frac{1}{\alpha}.$$

Lemma A.2 (Bounded perturbation, [HS87]). *Suppose that $\pi \in \mathcal{P}(\mathbb{R}^n)$ satisfies (LSI) with constants $C_{\text{LSI}}(\pi)$. If $\mu \in \mathcal{P}(\mathbb{R}^n)$ satisfies $c \leq \frac{d\mu}{d\pi} \leq C$ for $c, C \in \mathbb{R}_{>0}$, then*

$$C_{\text{LSI}}(\mu) \leq \frac{C}{c} C_{\text{LSI}}(\pi).$$

Lemma A.3 (Data-processing inequality). *For $\mu, \nu \in \mathcal{P}(\mathbb{R}^n)$, Markov kernel P , f -divergence D_f , and $q \in [1, \infty]$, it holds that*

$$D_f(\mu P \parallel \nu P) \leq D_f(\mu \parallel \nu), \quad \text{and} \quad \mathcal{R}_q(\mu P \parallel \nu P) \leq \mathcal{R}_q(\mu \parallel \nu).$$

Lemma A.4 (Basic facts on TV). *When $\{Z_i\}_{i \in [p]}$ are generated by a Markov chain,*

$$\|\text{law}(Z_1, \dots, Z_p) - \text{law}(Z_1, \dots, Z_{p-1}) \otimes \text{law } Z_p\|_{\text{TV}} = \|\text{law}(Z_{p-1}, Z_p) - \text{law } Z_{p-1} \otimes \text{law } Z_p\|_{\text{TV}}.$$

Also, it holds that $\|\mu^{\otimes m} - \nu^{\otimes m}\|_{\text{TV}} \leq m \|\mu - \nu\|_{\text{TV}}$.

Proof. For the first claim, let us denote by $f_{1, \dots, p}$ the probability density of (Z_1, \dots, Z_p) . Then,

$$\begin{aligned} \|\text{law}(Z_1, \dots, Z_p) - \text{law}(Z_1, \dots, Z_{p-1}) \otimes \text{law } Z_p\|_{\text{TV}} &= \frac{1}{2} \int |f_{1, \dots, p}(x, y) - f_{1, \dots, p-1}(x) f_p(y)| \, dx dy \\ &= \frac{1}{2} \int |f_{p|1, \dots, p-1}(y|x) - f_p(y)| f_{1, \dots, p-1}(x) \, dx dy = \frac{1}{2} \int |f_{p|p-1}(y|x') - f_p(y)| f_{p-1}(x') \, dx' dy \\ &= \frac{1}{2} \int |f_{p-1, p}(x', y) - f_{p-1}(x') f_p(y)| \, dx' dy = \|\text{law}(Z_{p-1}, Z_p) - \text{law } Z_{p-1} \otimes \text{law } Z_p\|_{\text{TV}}. \end{aligned}$$

For the second claim, using the DPI for both terms in the RHS (or just the definition of TV),

$$\begin{aligned} \|\mu^{\otimes m} - \nu^{\otimes m}\|_{\text{TV}} &\leq \|\mu^{\otimes m-1} \otimes \mu - \nu^{\otimes m-1} \otimes \mu\|_{\text{TV}} + \|\nu^{\otimes m-1} \otimes \mu - \nu^{\otimes m-1} \otimes \nu\|_{\text{TV}} \\ &\leq \|\mu^{\otimes m-1} - \nu^{\otimes m-1}\|_{\text{TV}} + \|\mu - \nu\|_{\text{TV}}. \end{aligned}$$

Applying the same argument to the first term, we finish the proof. \square

Geometry of logconcave distributions. We recall the most basic property of logconcave functions and distributions:

Lemma A.5 (Dinghas, Leindler, Prékopa). *The product, minimum, and convolution of logconcave functions is logconcave. Any marginals of logconcave distributions are logconcave.*

The following explains why we can focus on the ground set $\mathsf{L}_{\pi, g}$ of a logconcave distribution π .

Lemma A.6 ([LV07, Lemma 5.16]). *Let π be a logconcave probability density on \mathbb{R}^n . For $\beta \geq 2$,*

$$\mathbb{P}_{\pi}(\pi(X) \leq e^{-\beta(n-1)} \max \pi) \leq (e^{1-\beta} \beta)^{n-1}.$$

We recall two useful tail bounds of logconcave distributions.

Lemma A.7 (Exponential decay, [LV07, Lemma 5.17]). *For logconcave $\pi \in \mathcal{P}(\mathbb{R}^n)$ with $\mathbb{E}_{\pi}[\|X\|^2] = R^2$, it holds that for any $t \geq 1$,*

$$\mathbb{P}_{\pi}(\|X\| \geq Rt) \leq e^{-t+1}.$$

Lemma A.8 (Paouris). *There exists a universal constant $c > 0$ such that for any s -isotropic centered logconcave $\pi \in \mathcal{P}(\mathbb{R}^n)$ and $t \geq 1$,*

$$\mathbb{P}_{\pi}(\|X\| \geq t\sqrt{sn}) \leq \exp(-ct\sqrt{n}).$$

Proof. For Σ the covariance matrix of π , we note that $\nu := (\Sigma^{-1/2})_{\#}\pi$ is isotropic logconcave. Then, for $E = \{X \in \mathbb{R}^n : \|X\|_{\Sigma^{-1}} \geq t\sqrt{n}\}$ and some universal constant $c > 0$,

$$\pi(E) = \int \mathbf{1}_E(x) \, d\pi(x) = \int \mathbf{1}_E(\Sigma^{1/2}y) \, d\nu(y) = \nu(\mathbb{R}^n \setminus B_{t\sqrt{n}}(0)) \leq \exp(-ct\sqrt{n}),$$

where the last line follows from Paouris' theorem [Pao06]. Due to $s^{-1}I_n \preceq \Sigma^{-1}$, we have $\mathbb{P}_{\pi}(\|X\| \geq t\sqrt{sn}) \leq \exp(-ct\sqrt{n})$, from which the claim easily follows. \square

The reverse direction of the Hölder inequality holds for logconcave distributions.

Lemma A.9 (Reverse Hölder, [LV07, Theorem 5.22]). *For logconcave $\pi \in \mathcal{P}(\mathbb{R}^n)$, it holds that $(\mathbb{E}_\pi[\|X\|^k])^{1/k} \leq 2k \mathbb{E}_\pi\|X\|$ for $k \geq 1$.*

The following folklore reveals a lower regularity of logconcave distributions:

Lemma A.10 ([LV07, Lemma 5.13]). *The support of any s -isotropic logconcave distribution on \mathbb{R}^n contains a ball of radius $1/\sqrt{se}$ centered at its mean.*

The following on one-dimensional isotropic logconcave density is crucial when proving Lemma 4.13.

Lemma A.11 ([LV07, Lemma 5.5 (b)]). *Let $f : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ be an isotropic logconcave density function. Then, $f(0) \geq 1/8$.*