
COMPUTABLE APPROXIMATIONS OF SEMICOMPUTABLE GRAPHS

VEDRAN ČAČIĆ, MATEA ČELAR, MARKO HORVAT, AND ZVONKO ILJAZOVIĆ

e-mail address: veki@math.hr, matea.celar@math.hr, mhorvat@math.hr, zilj@math.hr

University of Zagreb, Faculty of Science, Department of Mathematics

ABSTRACT. In this work, we study the computability of topological graphs, which are obtained by gluing arcs and rays together at their endpoints. We prove that every semicomputable graph in a computable metric space can be approximated, with arbitrary precision, by its computable subgraph with computable endpoints.

1. INTRODUCTION

Our work progresses the research programme, started by Miller [Mil02], that is trying to determine which conditions render semicomputable subsets of computable metric spaces computable [AH23, AH22, ČHI21, IS18, Kih12, IP18, BI14]. We consider (*topological*) *graphs* [Ilj20], i.e., disjoint unions of arcs and rays that can be glued together at their endpoints. Our main goal is to prove that every semicomputable graph can be approximated by a computable subgraph with computable endpoints. This subgraph differs from the original graph only near the uncomputable endpoints of the graph, where computable ones are found arbitrarily close and the corresponding edges made that much smaller.

The plan is to first prove the following: for a semicomputable set S in a computable metric space and a point $x \in S$ with a neighbourhood homeomorphic to \mathbb{R} , there is another neighbourhood of x which is a computable arc between two computable points in S . In our proof, we use a known result [IV17, Theorem 5.2] which states that for a semicomputable set S in a computable metric space, points in S which have a Euclidean neighbourhood also have a computable compact neighbourhood.

This first, auxiliary result allows us to shorten an edge of the graph ending in an uncomputable point so that it ends in a computable point. We then repeat this procedure to get a semicomputable subgraph with computable endpoints. To complete the proof that the resulting graph is indeed computable, we leverage another result from the literature [Ilj20, Theorem 5.2] which states that a topological pair of a graph and the set of all its endpoints has computable type. In particular, a semicomputable graph is computable if the set of all its endpoints is semicomputable; this generalizes the result [BI14, Theorem 7.5] that each semicomputable 1-manifold with semicomputable boundary and finitely many components is computable.

Note that replacing the setting of computable metric spaces with that of computable topological spaces would not lead to a more general result. Recently, it was shown [AH23,

Theorem 3.4] that every semicomputable set S in a computable topological space can be effectively embedded into the Hilbert cube, i.e. there is a computable homeomorphism f with a computable inverse such that $f(S)$ is semicomputable and for each computable subset T of the Hilbert cube, $f^{-1}(T)$ is computable.

2. BACKGROUND

2.1. Computable metric spaces. In this subsection we provide some basic facts about computable metric spaces. See [PER89, Wei00, Tur36, Wei93, BW99, BP03, Ilj09, IS18].

Let $k \in \mathbb{N} \setminus \{0\}$. A function $f : \mathbb{N}^k \rightarrow \mathbb{Q}$ is said to be **computable** if there exist computable (i.e. recursive) functions $a, b, c : \mathbb{N}^k \rightarrow \mathbb{N}$ such that

$$f(x) = (-1)^{c(x)} \cdot \frac{a(x)}{b(x) + 1}$$

for each $x \in \mathbb{N}^k$. A function $f : \mathbb{N}^k \rightarrow \mathbb{R}$ is said to be **computable** if there exists a computable function $F : \mathbb{N}^{k+1} \rightarrow \mathbb{Q}$ such that

$$|f(x) - F(x, i)| < 2^{-i}$$

for all $x \in \mathbb{N}^k$, $i \in \mathbb{N}$.

Let (X, d) be a metric space and let α be a sequence in X such that $\alpha(\mathbb{N})$ is a dense set in (X, d) . We say that (X, d, α) is a **computable metric space** if the function $(i, j) \mapsto d(\alpha_i, \alpha_j) : \mathbb{N}^2 \rightarrow \mathbb{R}$ is computable.

For example, if $n \in \mathbb{N} \setminus \{0\}$, d is the Euclidean metric on \mathbb{R}^n and $\alpha : \mathbb{N} \rightarrow \mathbb{Q}^n$ is an effective enumeration of \mathbb{Q}^n , then $(\mathbb{R}^n, d, \alpha)$ is a computable metric space. This space is called **computable Euclidean space**.

Let (X, d, α) be a computable metric space. A point $x \in X$ is said to be **computable** in (X, d, α) if there is a computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$d(x, \alpha_{f(k)}) < 2^{-k}$$

for all $k \in \mathbb{N}$.

From now on, let $(j, i) \mapsto (j)_i : \mathbb{N}^2 \rightarrow \mathbb{N}$ and $j \mapsto \bar{j} : \mathbb{N} \rightarrow \mathbb{N}$ be fixed computable functions such that

$$\{((j)_0, (j)_1, \dots, (j)_{\bar{j}}) \mid j \in \mathbb{N}\}$$

is the set of all nonempty finite sequences in \mathbb{N} .

For $j \in \mathbb{N}$ let

$$[j] := \{(j)_0, \dots, (j)_{\bar{j}}\}.$$

Then each nonempty finite subset of \mathbb{N} is equal to $[j]$ for some $j \in \mathbb{N}$.

Let (X, d) be a metric space, $A, B \subseteq X$ and $\varepsilon > 0$. We say that A and B are ε -**close**, and write $A \approx_\varepsilon B$, if

$$(\forall a \in A)(\exists b \in B)(d(a, b) < \varepsilon) \quad \text{and} \quad (\forall b \in B)(\exists a \in A)(d(a, b) < \varepsilon).$$

If A and B are nonempty compact sets in (X, d) , the number $\inf\{\varepsilon > 0 \mid A \approx_\varepsilon B\}$ is called the **Hausdorff distance** from A to B and it is denoted by $d_H(A, B)$.

It is not hard to check that, for $\varepsilon > 0$, we have $d_H(A, B) < \varepsilon$ if and only if $A \approx_\varepsilon B$.

Definition 2.1. Let (X, d, α) be a computable metric space. We say that a compact set $S \subseteq X$ is **computable** in (X, d, α) if $S = \emptyset$ or there is a computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$S \approx_{2^{-k}} \{\alpha_i \mid i \in [f(k)]\}, \quad \text{for all } k \in \mathbb{N}.$$

Let (X, d, α) be a computable metric space. Let $\tau_1, \tau_2 : \mathbb{N} \rightarrow \mathbb{N}$ be fixed computable functions such that $\{(\tau_1(i), \tau_2(i)) \mid i \in \mathbb{N}\} = \mathbb{N}^2$ and let $q : \mathbb{N} \rightarrow \mathbb{Q}$ be an effective enumeration of all positive rational numbers. For $i \in \mathbb{N}$ let

$$I_i := B(\alpha_{\tau_1(i)}, q_{\tau_2(i)}) \quad \text{and} \quad \hat{I}_i := \hat{B}(\alpha_{\tau_1(i)}, q_{\tau_2(i)}),$$

where $B(\alpha_{\tau_1(i)}, q_{\tau_2(i)})$ is the open ball in (X, d) with radius $\rho_i := q_{\tau_2(i)}$ centered at $\lambda_i := \alpha_{\tau_1(i)}$, and $\hat{B}(\lambda_i, \rho_i)$ is the corresponding closed ball.

For $j, l \in \mathbb{N}$ let

$$J_j := \bigcup_{i \in [j]} I_i, \quad \hat{J}_j := \bigcup_{i \in [j]} \hat{I}_i \quad \text{and} \quad J_{[l]} := \bigcup_{j \in [l]} J_j.$$

Then (J_j) and (\hat{J}_j) are, respectively, effective enumerations of all finite unions of rational open and closed balls.

Definition 2.2. Let (X, d, α) be a computable metric space.

- (i) A closed set S in (X, d) is said to be **computably enumerable** (c.e.) in (X, d, α) if the set $\{i \in \mathbb{N} \mid I_i \cap S \neq \emptyset\}$ is c.e.
- (ii) A compact set S in (X, d) is said to be **semicomputable** in (X, d, α) if the set $\{j \in \mathbb{N} \mid S \subseteq J_j\}$ is c.e.

It is not hard to see that these definitions do not depend on the choice of particular functions q, τ_1, τ_2 and $j \mapsto [j]$.

We write $K \subseteq_\varepsilon J_j$ if:

- $K \subseteq J_j$,
- for all $i \in [j]$, $I_i \cap K \neq \emptyset$, and
- for all $i \in [j]$, $\rho_i < \varepsilon$.

Note that $K \subseteq_\varepsilon J_j$ implies $K \approx_\varepsilon \{\lambda_i \mid i \in [j]\}$.

There is a characterization [Ilj13, Proposition 2.6] of a computable compact set in (X, d, α) :

$$S \text{ computable} \iff S \text{ c.e. and semicomputable.} \quad (2.1)$$

The notion of semicomputability can be generalized to noncompact sets [BI14, IS18].

Definition 2.3. Let (X, d, α) be a computable metric space.

A set $S \subseteq X$ is **semicomputable** in (X, d, α) if the following hold:

- (i) $S \cap B$ is a compact set for each closed ball B in (X, d) ;
- (ii) the set $\{(i, j) \in \mathbb{N}^2 \mid \hat{I}_i \cap S \subseteq J_j\}$ is c.e.

This notion extends the previously defined notion of semicomputability in the following sense: if S is a compact set in (X, d) , then S is semicomputable in the sense of the latter definition if and only if S is semicomputable in the sense of Definition 2.2(ii).

The notion of a computable set in a computable metric space can be now extended to noncompact sets in the obvious way: S is **computable** in (X, d, α) if S is semicomputable and c.e. in (X, d, α) .

Each semicomputable set S is co-c.e., which means that $X \setminus S = \bigcup_{i \in A} I_i$ for some c.e. set $A \subseteq \mathbb{N}$. On the other hand, a co-c.e. set need not be semicomputable, but the equivalence

$$S \text{ semicomputable} \iff S \text{ co-c.e.}$$

holds in some computable metric space, for example in computable Euclidean space $(\mathbb{R}^n, d, \alpha)$.

Definition 2.4. Let $i, j \in \mathbb{N}$.

- We say that I_i and I_j are **formally disjoint** if

$$d(\lambda_i, \lambda_j) > \rho_i + \rho_j.$$

We denote this by $I_i \diamond I_j$.

- We say that J_i and J_j are **formally disjoint** if $I_k \diamond I_l$ for all $k \in [i]$ and $l \in [j]$. We denote this by $J_i \diamond J_j$.

Note that $I_i \diamond I_j$ implies $\hat{I}_i \cap \hat{I}_j = \emptyset$. Similarly, $J_i \diamond J_j$ implies $\hat{J}_i \cap \hat{J}_j = \emptyset$.

Definition 2.5. Let $i, j \in \mathbb{N}$.

- We say that I_i is **formally contained** in I_j and we write $I_i \subseteq_{\forall} I_j$ if

$$d(\lambda_i, \lambda_j) + \rho_i < \rho_j.$$

- We say that J_i is **formally contained** in J_j and we write $J_i \subseteq_{\forall} J_j$ if

$$(\forall k \in [i])(\exists l \in [j])(I_k \subseteq_{\forall} I_l).$$

- We say that $J_{[i]}$ is **formally contained** in $J_{[j]}$ and we write $J_{[i]} \subseteq_{\forall} J_{[j]}$ if

$$(\forall k \in [i])(\exists l \in [j])(J_k \subseteq_{\forall} J_l).$$

Note that $I_i \subseteq_{\forall} I_j$ implies $\hat{I}_i \subseteq I_j$ and $J_i \subseteq_{\forall} J_j$ implies $\hat{J}_i \subseteq J_j$.

Note also that formal containment and formal disjointness are relations between numbers i and j (and not between the sets I_i and I_j or J_i and J_j), and that they are c.e.

Let (X, d, α) be a computable metric space. For every $j \in \mathbb{N}$ we define

$$\text{fdiam}(j) := \text{diam} \{ \lambda_u \mid u \in [j] \} + 2 \max \{ \rho_u \mid u \in [j] \}, \quad (2.2)$$

and call it the **formal diameter** of J_j . Again, this is formally a function of j , not of J_j .

It is easy to prove that the function $\text{fdiam} : \mathbb{N} \rightarrow \mathbb{R}$ is computable [Ilj09, Proposition 13].

2.2. Chains. Let (X, d) be a metric space. A finite sequence $\mathcal{C} = (C_0, \dots, C_n)$ of nonempty subsets of X is said to be a

- **quasi-chain** in X if, for all i, j , $C_i \cap C_j \neq \emptyset$ implies $|i - j| \leq 1$;
- **chain** in X if, for all i, j , $C_i \cap C_j \neq \emptyset$ is equivalent to $|i - j| \leq 1$;

Each set C_i is said to be the (i th) **link** of the (quasi-)chain \mathcal{C} . If $i < j$, we say that the link C_i **precedes** C_j , and if $i + 1 < j$, we say that C_i **strictly precedes** C_j .

A (quasi-)chain is said to be **open** in (X, d) if each of its links is open in (X, d) . Similarly, a (quasi-)chain is said to be **compact** if each of its links is compact.

Suppose S is a subset of X and $\mathcal{C} = (C_0, \dots, C_n)$ is a (quasi-)chain in X . We say that \mathcal{C} **covers** S if $S \subseteq C_0 \cup \dots \cup C_n$.

Suppose $\mathcal{A} = (A_0, \dots, A_n)$ is a finite sequence of nonempty bounded subsets of X . The **mesh** of \mathcal{A} is the number defined by

$$\text{mesh}(\mathcal{A}) := \max_{i=0}^n \text{diam } A_i.$$

If \mathcal{C} is a chain and $\text{mesh}(\mathcal{C}) < \varepsilon$, we say that \mathcal{C} is an ε -**chain**. We say that a continuum (i.e. a compact and connected metric space) X is **chainable** if there is an open ε -chain in X covering X for every $\varepsilon > 0$. For $a, b \in X$, we say that a continuum X is **chainable from a to b** if, for every $\varepsilon > 0$, there is an open ε -chain (U_0, \dots, U_n) in X covering X such that $a \in U_0$ and $b \in U_n$.

Let $\mathcal{C} = (C_0, \dots, C_n)$ and $\mathcal{D} = (D_0, \dots, D_m)$ be two (quasi-)chains in X . We say that \mathcal{D} **refines** \mathcal{C} if for each $i \in \{0, \dots, m\}$ there exists $j \in \{0, \dots, n\}$ such that $D_i \subseteq C_j$. If \mathcal{D} **refines** \mathcal{C} and $D_0 \subseteq C_0$ and $D_m \subseteq C_n$, we say that \mathcal{D} **strongly refines** \mathcal{C} .

Let (X, d, α) be a computable metric space and let $l \in \mathbb{N}$. We say that $(J_{j_0}, \dots, J_{j_m})$ is a **formal chain** in (X, d, α) if $J_{j_u} \diamond J_{j_v}$ for all $u, v \in \{0, \dots, m\}$ such that $|u - v| > 1$. Note that every formal chain is a quasi-chain.

2.3. Graphs. Let $n \in \mathbb{N} \setminus \{0\}$ and let $a, v \in \mathbb{R}^n$, $v \neq 0$. The set $\{a + tv \mid t \in [0, \infty)\}$ is called a **ray** in \mathbb{R}^n and we say that a is the (only) **endpoint** of this ray. Let K be a nonempty finite family of line segments and rays in \mathbb{R}^n such that

$$\begin{aligned} I, J \in K \text{ are such that} & \implies I \cap J = \{a\}, \text{ where } a \text{ is an} \\ I \neq J \text{ and } I \cap J \neq \emptyset & \text{ endpoint of both } I \text{ and } J. \end{aligned} \tag{2.3}$$

Then any topological space G homeomorphic to $\bigcup K$ is called a **graph**.

Let G be a graph and $x \in G$. We say that x is an **endpoint** of G if x has an open neighborhood in G which is homeomorphic to $[0, \infty)$ by a homeomorphism which maps x to 0.




Figure 1: A graph. Filled circles denote endpoints and arrows denote rays.

Suppose G is a graph and $f : \bigcup K \rightarrow G$ a homeomorphism, where K is the family from the definition of a graph. Then:

- $x \in G$ is an endpoint of G if and only if there exists $y \in \bigcup K$ such that $f(y) = x$ and such that y is an endpoint of a unique element of K ;
- G is compact if and only if K does not contain rays.

Note that our naming scheme differs slightly from [Ilj20]; what was previously called a *generalized graph* we call a *graph*, and what was previously called a *graph* is now a *compact graph*.

2.4. Auxiliary results. We will use the following results from the literature:

Lemma 2.6. [IP18, Lemma 3.8] *Let X be a set and let (C_0, \dots, C_m) and $(D_0, \dots, D_{m'})$ be two quasi-chains in X such that $(D_0, \dots, D_{m'})$ refines (C_0, \dots, C_m) . Suppose that $i, j, k \in \{0, \dots, m\}$ and $p, q \in \{0, \dots, m'\}$ are such that*

$$i < k < j, \quad p < q, \quad D_p \subseteq C_i \quad \text{and} \quad D_q \subseteq C_j.$$

Then there exists $r \in \{0, \dots, m'\}$ such that $p < r < q$ and $D_r \subseteq C_k$.

Lemma 2.7. [Ilj09, Lemma 41] *Let (X, d) be a metric space in which every closed ball is compact. Let $\mathcal{C}^k = (C_0^k, \dots, C_{m_k}^k)$, $k \in \mathbb{N}$ be a sequence of chains such that for all $k \in \mathbb{N}$, $(\text{Cl}(C_0^{k+1}), \dots, \text{Cl}(C_{m_{k+1}}^{k+1}))$ strongly refines $(C_0^k, \dots, C_{m_k}^k)$ and $\text{mesh}(\mathcal{C}^k) < 2^{-k}$. Let*

$$S = \bigcap_{k \in \mathbb{N}} \left(\text{Cl}(C_0^{k+1}) \cup \dots \cup \text{Cl}(C_{m_{k+1}}^{k+1}) \right).$$

Then S is a continuum, chainable from a to b , where $a \in \bigcap_{k \in \mathbb{N}} C_0^k$ and $b \in \bigcap_{k \in \mathbb{N}} C_{m_k}^k$.

Theorem 2.8. [IV17, Theorem 5.2] *Suppose (X, d, α) is a computable metric space, S a semicomputable set in this space and $x \in S$ a point which has a neighbourhood in S homeomorphic to \mathbb{R}^n for some $n \in \mathbb{N} \setminus \{0\}$. Then x has a computable compact neighbourhood in S .*

Theorem 2.9. [Ilj20, Theorem 6.5] *Let (X, d, α) be a computable metric space and let S be a semicomputable set in this space. Suppose S , as a subspace of (X, d) , is a graph such that the set E of all endpoints of S is semicomputable in (X, d, α) . Then S is computable in (X, d, α) .*

Proposition 2.10. [Ilj09, Proposition 12] *Let (X, d, α) be a computable metric space.*

- (i) *For all $j \in \mathbb{N}$, $\text{diam } \widehat{J}_j \leq \text{fdiam}(j)$.*
- (ii) *Let K and U be subsets of (X, d) such that K is nonempty and compact, U is open and $K \subseteq U$. Let $\varepsilon > 0$. Then there exists $j \in \mathbb{N}$ such that $K \subseteq J_j$, $\widehat{J}_j \subseteq U$ and $\text{fdiam}(j) < \text{diam } K + \varepsilon$.*

Lemma 2.11. *Let (X, d, α) be a computable metric space and let S be a semicomputable set in this space (not necessarily compact). Let $m \in \mathbb{N}$. Then $S \setminus J_m$ is a semicomputable set.*

Proof. Let $\Omega = \{(i, j) \in \mathbb{N}^2 \mid \widehat{I}_i \cap S \subseteq J_j\}$.

If B is a closed ball in (X, d) , then $(S \setminus J_m) \cap B$ is a closed set in (X, d) contained in $S \cap B$. Since $S \cap B$ is compact, $(S \setminus J_m) \cap B$ is compact too.

Let $i, j \in \mathbb{N}$. We have

$$\widehat{I}_i \cap (S \setminus J_m) \subseteq J_j \iff (\widehat{I}_i \cap S) \setminus J_m \subseteq J_j \iff \widehat{I}_i \cap S \subseteq J_j \cup J_m.$$

Choose a computable function $f : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that $J_a \cup J_b = J_{f(a,b)}$ for all $a, b \in \mathbb{N}$. Then

$$\widehat{I}_i \cap (S \setminus J_m) \subseteq J_j \iff \widehat{I}_i \cap S \subseteq J_{f(j,m)} \iff (i, f(j,m)) \in \Omega.$$

So the set of all $(i, j) \in \mathbb{N}^2$ such that $\widehat{I}_i \cap (S \setminus J_m) \subseteq J_j$ is the set of all $(i, j) \in \mathbb{N}^2$ such that $(i, f(j,m)) \in \Omega$, which is c.e. since Ω is c.e. and f is computable. \square

Corollary 2.12. *Let (X, d, α) be a computable metric space, let S be a semicomputable set in this space and let K and U be subsets of S such that K is compact, U is open in S and $K \subseteq U$. Then there exists a semicomputable compact set $S' \subseteq S$ such that $K \subseteq S' \subseteq U$.*

Proof. Since K is compact, it is fully contained in a rational closed ball with a sufficiently large diameter. Let $i \in \mathbb{N}$ be such that $K \subseteq L := \hat{I}_i \cap S$.

Consider the set $L \setminus U$. If it is empty, then $L \subseteq U$, so $S' := L$ is a semicomputable compact set such that $K \subseteq S' \subseteq U$.

Suppose $L \setminus U$ is not empty. Then it is a closed subset of a compact set L , therefore it is compact, and $L \setminus U \subseteq L \setminus K$. Since $L \setminus K$ is open in L , let V be an open set in (X, d) such that $V \cap L = L \setminus K$. Now, $L \setminus U$ and V are subsets of (X, d) such that $L \setminus U$ is nonempty and compact, V is open and $L \setminus U \subseteq V$. By Proposition 2.10(ii), there exists $j \in \mathbb{N}$ such that $L \setminus U \subseteq J_j \subseteq V$. Then $S' := L \setminus J_j$ is semicomputable compact and

$$L \setminus V \subseteq S' \subseteq L \setminus (L \setminus U).$$

Since

$$K = L \setminus (L \setminus K) = L \setminus (V \cap L) = L \setminus V \quad \text{and} \quad L \setminus (L \setminus U) \subseteq U,$$

we conclude that $K \subseteq S' \subseteq U$. □

Proposition 2.13. [BI14, Lemma 4.6] *Let (X, d, α) be a computable metric space and let A, B be disjoint nonempty compact sets in (X, d) . Then*

- (i) *For each $\varepsilon > 0$ there exists $j \in \mathbb{N}$ such that $A \subseteq_\varepsilon J_j$.*
- (ii) *There exists $\mu > 0$ such that for all $i, j \in \mathbb{N}$, if $A \subseteq_\mu J_i$ and $B \subseteq_\mu J_j$, then $J_i \diamond J_j$.*

Lemma 2.14. [BI14, Lemma 4.8] *Let (X, d, α) be a computable metric space. Let $A \subseteq X$, $j \in \mathbb{N}$ and $r > 0$ be such that $A \subseteq_r J_j$. Then*

$$\text{fdiam}(j) < 4r + \text{diam } A.$$

Proposition 2.15. [IP18, Proposition 4.13] *Let (X, d, α) be a computable metric space, $A \subseteq \mathbb{N}$ a c.e. set, K a nonempty compact set such that $K \subseteq \bigcup_{i \in A} I_i$. Then there exists $\mu > 0$ such that for all $i \in \mathbb{N}$, if $K \subseteq_\mu J_j$, then*

$$(\forall u \in [j])(\exists v \in A)(I_u \subseteq_\forall I_v).$$

Proposition 2.16. [IP18, Proposition 6.5.] *Let (X, d, α) be a computable metric space. Let $l, l', u, v \in \mathbb{N}$. The following statements hold:*

- (i) *if $J_u \subseteq_\forall J_v$, then $\text{Cl}(J_u) \subseteq J_v$;*
- (ii) *if $J_{[l]} \subseteq_\forall J_{[l']}$, then the finite sequence $(\text{Cl}(J_{(l)_0}), \dots, \text{Cl}(J_{(l)_{\bar{l}}}))$ refines the finite sequence $(J_{(l')_0}, \dots, J_{(l')_{\bar{l}'}})$.*

3. EXISTENCE OF A COMPUTABLE NEIGHBOURHOOD

The main result of this section is Theorem 3.3 which states that any point in a semicomputable set which has a neighbourhood N homeomorphic to \mathbb{R} also has a computable neighbourhood $N' \subseteq N$ which is an arc with computable endpoints.

Note that Theorem 2.8 has a similar statement: any point x in a semicomputable set which has a neighbourhood homeomorphic to \mathbb{R}^n also has a computable neighbourhood. However, it does not guarantee that this computable neighbourhood will be an arc with computable endpoints. Since computable points are dense in computable sets, there is a neighbourhood of x which is an arc with computable endpoints, but such neighbourhood is not necessarily computable (or even semicomputable).

In order to construct a neighbourhood which is simultaneously computable and an arc with computable endpoints, our approach is to look at the intersection of a sequence of chains whose links have strictly decreasing diameters.

We begin by listing some important auxiliary results.

Lemma 3.1. *Let (X, d, α) be a computable metric space, K a nonempty compact set, $a \in \mathbb{N}$ such that $K \subseteq J_a$. Then there exists $\mu > 0$ such that for all $j \in \mathbb{N}$, if $K \subseteq_\mu J_j$, then $J_j \subseteq_{\forall} J_a$.*

Proof. This follows immediately from Proposition 2.15 for $A = [a]$. \square

Lemma 3.2. *Let (X, d, α) be a computable metric space, (K_0, \dots, K_{n+1}) a compact quasi-chain in (X, d) , and $A \subseteq \mathbb{N}$ a finite set. Then for all $\varepsilon > 0$, there exist $p, l, q \in \mathbb{N}$ such that $\bar{l} = n - 1$ and*

- (a) $K_0 \subseteq_\varepsilon J_p$, $K_{n+1} \subseteq_\varepsilon J_q$ and $K_i \subseteq_\varepsilon J_{(l)_{i-1}}$ for all $i \in \{1, \dots, n\}$;
- (b) $(J_p, J_{(l)_0}, \dots, J_{(l)_{\bar{l}}}, J_q)$ is a formal chain;
- (c) for all $a \in A$, $K_0 \subseteq J_a$ implies $J_p \subseteq_{\forall} J_a$, $K_{n+1} \subseteq J_a$ implies $J_q \subseteq_{\forall} J_a$ and $K_i \subseteq J_a$ implies $J_{(l)_{i-1}} \subseteq_{\forall} J_a$, for all $i \in \{1, \dots, n\}$.

Proof. Fix $i, j \in \{0, \dots, n+1\}$ such that $|i - j| > 1$. Since (K_0, \dots, K_{n+1}) is a compact quasi-chain, K_i and K_j are disjoint nonempty compact sets in (X, d) . By Proposition 2.13(ii), there exists $\mu_{i,j} > 0$ such that for any $\alpha, \beta \in \mathbb{N}$,

$$K_i \subseteq_{\mu_{i,j}} J_\alpha \quad \text{and} \quad K_j \subseteq_{\mu_{i,j}} J_\beta \implies J_\alpha \diamond J_\beta. \quad (3.1)$$

Now fix $i \in \{0, \dots, n+1\}$ and $a \in A$ such that $K_i \subseteq J_a$. By Lemma 3.1, there exists $\mu_{i,a} > 0$ such that for any $\alpha \in \mathbb{N}$,

$$K_i \subseteq_{\mu_{i,a}} J_\alpha \implies J_\alpha \subseteq_{\forall} J_a. \quad (3.2)$$

Let

$$r := \min(\{\mu_{i,j} \mid i, j \in \{0, \dots, n+1\}, |i - j| > 1\} \cup \{\mu_{i,a} \mid i \in \{0, \dots, n+1\}, a \in A, K_i \subseteq J_a\} \cup \{\varepsilon\})$$

By Proposition 2.13(i), there exist $p, j_1, \dots, j_n, q \in \mathbb{N}$ such that

$$K_0 \subseteq_r J_p, K_1 \subseteq_r J_{j_1}, \dots, K_n \subseteq_r J_{j_n} \text{ and } K_{n+1} \subseteq_r J_q. \quad (3.3)$$

Let $l \in \mathbb{N}$ be such that $(j_1, \dots, j_n) = ((l)_0, \dots, (l)_{\bar{l}})$. Since $K \subseteq_{\lambda'} J_j$ implies $K \subseteq_\lambda J_j$ for any $\lambda' < \lambda$, we can now easily conclude that (a) follows from (3.3), (b) follows from (3.1), and (c) follows from (3.2). \square

Any topological space homeomorphic to $[0, 1]$ is called an **arc**. If A is an arc and $x \in A$ a point such that $A \setminus \{x\}$ is connected, then we say that x is an **endpoint** of A . Note that if $f : [0, 1] \rightarrow A$ is a homeomorphism, then $f(0)$ and $f(1)$ are the only endpoints of A .

Theorem 3.3. *Let S be a semicomputable set in a computable metric space (X, d, α) . Suppose a point $x \in S$ has an open neighborhood N in S which is homeomorphic to \mathbb{R} . Then there exist computable points $a, b \in N$ and a computable neighbourhood $N' \subseteq N$ of x in S which is an arc from a to b .*

Proof. Let N be an open neighborhood of x in S and let $f : \mathbb{R} \rightarrow N$ be a homeomorphism. Without loss of generality (we can always compose f with a translation by $f^{-1}(x)$), we may assume $f(0) = x$.

Consider the subsets $f([-3, 3])$ and $f(\langle -4, 4 \rangle)$ of S . The set $f(\langle -4, 4 \rangle)$ is a continuous image of an open set, so it is open in N and therefore open in S . The set $f([-3, 3])$ is a nonempty compact subset of $f(\langle -4, 4 \rangle)$. By Corollary 2.12, there exists a semicomputable compact set S' such that

$$f([-3, 3]) \subseteq S' \subseteq f(\langle -4, 4 \rangle). \quad (3.4)$$

Since f^{-1} is continuous and $f([-4, 4])$ is compact, we can choose $\varepsilon > 0$ such that

$$d(f(s), f(t)) < \varepsilon \implies |s - t| < \frac{1}{2}, \quad \text{for all } s, t \in [-4, 4]. \quad (3.5)$$

We may additionally assume $\varepsilon < 1$.

The set $f(\langle -3, 3 \rangle)$ is a neighbourhood of $f(-2)$ in S' homeomorphic to \mathbb{R} . By Theorem 2.8, there is a computable neighbourhood $N_{\tilde{a}}$ of $f(-2)$ in S' . Similarly, there is a computable neighbourhood $N_{\tilde{b}}$ of $f(2)$ in S' .

Since computable points are dense in computable sets, we can find computable points (see Figure 2)

$$\tilde{a} \in B(f(-2), \varepsilon) \cap N_{\tilde{a}} \quad \text{and} \quad \tilde{b} \in B(f(2), \varepsilon) \cap N_{\tilde{b}}. \quad (3.6)$$




Figure 2: The choice of points \tilde{a} and \tilde{b} from S' .

Note that $\tilde{a}, \tilde{b} \in S' \subseteq f(\langle -4, 4 \rangle)$, so $\tilde{a} = f(t_{\tilde{a}})$ and $\tilde{b} = f(t_{\tilde{b}})$ for some $t_{\tilde{a}}, t_{\tilde{b}} \in \langle -4, 4 \rangle$. Moreover, by (3.5), we have

$$t_{\tilde{a}} \in \langle -2.5, -1.5 \rangle \quad \text{and} \quad t_{\tilde{b}} \in \langle 1.5, 2.5 \rangle. \quad (3.7)$$

For $(p, l, q) \in \mathbb{N}^3$, we consider the following statements:

- (O1) $S' \subseteq J_p \cup J_{[l]} \cup J_q$;
- (O2) $(J_p, J_{(l)_0}, \dots, J_{(l)_l}, J_q)$ is a formal chain;
- (O3) $\tilde{a} \in J_p$;
- (O4) $\tilde{b} \in J_q$.

Each of these relations is c.e. (which follows from, respectively, the semicomputability of S' , the computable enumerability of formal disjointness, the computability of \tilde{a} and the computability of \tilde{b}), so the set

$$\Omega := \{l \in \mathbb{N} \mid (\exists p, q \in \mathbb{N})((\text{O1})\text{--}(\text{O4}) \text{ hold for } (p, l, q))\}$$

is also c.e. as a projection of their intersection.

Claim 1. *There exists $l \in \Omega$ such that $\text{fmesh}(l) < \frac{\varepsilon}{2}$, $d(\tilde{a}, J_{(l)_0}) < \frac{\varepsilon}{2}$ and $d(\tilde{b}, J_{(l)_l}) < \frac{\varepsilon}{2}$.*

Proof of Claim 1. Since f is uniformly continuous on $[-4, 4]$, we can find a subdivision $-4 = x_0 < x_1 < \dots < x_{n+1} = 4$ of $[-4, 4]$ such that for each $i \in \{0, \dots, n\}$,

$$\text{diam } f([x_i, x_{i+1}]) < \frac{\varepsilon}{4}. \quad (3.8)$$

Denote $C_i := f([x_i, x_{i+1}])$.

First, note that if $f(s) \in C_i$ and $f(t) \in C_j$, then

$$|j - i| > 2|t - s| - 1. \quad (3.9)$$

Indeed, suppose without loss of generality $s < t$ (the case $s = t$ is trivial); then $i \leq j$ and

$$t - s = (t - x_j) + (x_j - x_{j-1}) + \dots + (x_{i+1} - s),$$

where we have $j - i + 1$ summands. By (3.8) and (3.5), each one is less than $\frac{1}{2}$, so $|t - s| < \frac{1}{2}(|j - i| + 1)$, and (3.9) follows from this.

Let i be the index such that $t_{\tilde{a}} \in [x_i, x_{i+1})$ and let j be the index such that $t_{\tilde{b}} \in \langle x_j, x_{j+1} \rangle$. Then $\tilde{a} \in C_i$ and $\tilde{b} \in C_j$ and, by (3.7), we have

$$\begin{aligned} i - |i - 0| &> 2|t_{\tilde{a}} - (-4)| - 1 \geq 2|-2.5 + 4| - 1 = 2, \\ n - j &= |n - j| > 2|4 - t_{\tilde{b}}| - 1 \geq 2|4 - 2.5| - 1 = 2 \end{aligned}$$

and

$$|j - i| > 2|t_{\tilde{b}} - t_{\tilde{a}}| - 1 \geq 2|1.5 - (-1.5)| - 1 = 5.$$

Now we consider the compact quasi-chain

$$(C_0 \cup \dots \cup C_i, C_{i+1}, \dots, C_{j-1}, C_j \cup \dots \cup C_n).$$

By Lemma 3.2 (for $A = \emptyset$), we can find $p, l, q \in \mathbb{N}$ such that

$$C_0 \cup \dots \cup C_i \subseteq_{\frac{\varepsilon}{16}} J_p, \quad C_j \cup \dots \cup C_n \subseteq_{\frac{\varepsilon}{16}} J_q, \quad (3.10)$$

$$C_{i+1} \subseteq_{\frac{\varepsilon}{16}} J_{(l)_0}, \quad \dots, \quad C_{j-1} \subseteq_{\frac{\varepsilon}{16}} J_{(l)_i}. \quad (3.11)$$

and such that $(J_p, J_{(l)_0}, \dots, J_{(l)_i}, J_q)$ is a formal chain.

Now, we have $\tilde{a} \in C_i \subseteq J_p$, $\tilde{b} \in C_j \subseteq J_q$ and

$$S' \subseteq f([-4, 4]) = \bigcup_{i=0}^n C_i \subseteq J_p \cup J_{[l]} \cup J_q.$$

Therefore, (p, l, q) satisfies (O1)–(O4), so $l \in \Omega$. Moreover, since $\text{diam } C_k < \frac{\varepsilon}{4}$ for all $i < k < j$, (3.11) and Lemma 2.14 imply that for all $u \in [l]$,

$$\text{fdiam}(u) < 4 \cdot \frac{\varepsilon}{16} + \frac{\varepsilon}{4} = \frac{\varepsilon}{2},$$

so $\text{fmesh}(l) < \frac{\varepsilon}{2}$. Finally, since $f(x_{i+1}) \in C_i \cap C_{i+1} \subseteq C_i \cap J_{(l)_0}$, we have

$$d(\tilde{a}, J_{(l)_0}) \leq d(\tilde{a}, f(x_{i+1})) \leq \text{diam } C_i < \frac{\varepsilon}{2}$$

and, similarly, $d(\tilde{b}, J_{(l)_i}) < \frac{\varepsilon}{2}$. This concludes the proof of Claim 1.

Claim 2. If $l \in \Omega$, then

$$(J_{(l)_0} \cap S', \dots, J_{(l)_i} \cap S')$$

is an open chain in S' .

Proof of Claim 2. Let $p, q \in \mathbb{N}$ be such that (O1)–(O4) hold for (p, l, q) . We will prove that

$$\mathcal{C} := (J_p \cap S', J_{(l)_0} \cap S', \dots, J_{(l)_\bar{l}} \cap S', J_q \cap S')$$

is an open chain in S' .

The links of \mathcal{C} are clearly open in S' . By (O2), \mathcal{C} is a quasi-chain. It is sufficient to prove that neighbouring links of \mathcal{C} intersect.

By (3.7), (3.4) and (O1), we have

$$f([t_{\tilde{a}}, t_{\tilde{b}}]) \subseteq f([-3, 3]) \subseteq S' = \bigcup \mathcal{C}. \quad (3.12)$$

If $J_{(l)_{i-1}} \cap S'$ and $J_{(l)_i} \cap S'$ were disjoint for some $i > 0$, we would have

$$\begin{aligned} f([t_{\tilde{a}}, t_{\tilde{b}}]) \subseteq & ((J_p \cap S') \cup \dots \cup (J_{(l)_{i-1}} \cap S')) \cup \\ & \cup ((J_{(l)_i} \cap S') \cup \dots \cup (J_q \cap S')). \end{aligned}$$

However, this is impossible: because $f(t_{\tilde{a}}) \in J_p \cap S'$ and $f(t_{\tilde{b}}) \in J_q \cap S'$ by (O3) and (O4), the right-hand side is a disjoint union of nonempty open sets, but $f([t_{\tilde{a}}, t_{\tilde{b}}])$ is connected. The same argument shows that $J_p \cap S'$ intersects $J_{(l)_0} \cap S'$ and $J_{(l)_\bar{l}} \cap S'$ intersects $J_q \cap S'$. This concludes the proof of Claim 2.

For $(l, l') \in \mathbb{N}^2$, we consider the following statements:

- (G1) $l, l' \in \Omega$;
- (G2) $J_{[l']} \subseteq_{\forall} J_{[l]}$;
- (G3) $\text{fmesh}(l') < \frac{1}{2} \text{fmesh}(l)$;
- (G4) $J_{(l')_0} \subseteq_{\forall} J_{(l)_0}$;
- (G5) $J_{(l')_{\bar{l}'}} \subseteq_{\forall} J_{(l)_{\bar{l}}}$.

Denote by Γ the set of all $(l, l') \in \mathbb{N}^2$ which satisfy (G1)–(G5). By computable enumerability of Ω and \subseteq_{\forall} (as a relation on J -indices) and computability of fmesh , Γ is c.e.

Claim 3. *For each $l \in \Omega$, there exists $l' \in \Omega$ such that $(l, l') \in \Gamma$.*

Proof of Claim 3. Let $l \in \Omega$ and let $p, q \in \mathbb{N}$ be such that (O1)–(O4) hold for (p, l, q) .

By (O1), $\{J_p, J_{(l)_0}, \dots, J_{(l)_{\bar{l}}}, J_q\}$ is an open cover of S' . Let λ be a Lebesgue number of this cover.

Since J_p and J_q are open and $\tilde{a} \in J_p, \tilde{b} \in J_q$, there exist $r_1, r_2 > 0$ such that $B(\tilde{a}, r_1) \subseteq J_p$ and $B(\tilde{b}, r_2) \subseteq J_q$. Denote

$$r := \min \left\{ \lambda, r_1, r_2, \frac{1}{10} \text{fmesh}(l) \right\}.$$

Let $-4 = x_0 < x_1 < \dots < x_{n+1} = 4$ be a subdivision of $[-4, 4]$ such that for each $i \in \{0, \dots, n\}$

$$\text{diam } f([x_i, x_{i+1}]) < r. \quad (3.13)$$

Let $C_i := f([x_i, x_{i+1}]) \cap S', \forall i \in \{0, \dots, n\}$. Then (C_0, \dots, C_n) is a compact quasi-chain in S' which covers S' .

As in the proof of Claim 1, let i be the index such that $t_{\tilde{a}} \in [x_i, x_{i+1}]$ and let j be the index such that $t_{\tilde{b}} \in [x_j, x_{j+1}]$, so that $\tilde{a} \in C_i$ and $\tilde{b} \in C_j$. As before, we know that $i > 2$, $j < n - 2$ and $|i - j| > 5$.

Since $\tilde{a} \in C_i$, $\text{diam } C_i < r_1$ and $B(\tilde{a}, r_1) \subseteq J_p$ we have $C_i \subseteq J_p$. Similarly, $C_j \subseteq J_q$. By Lemma 2.6, there exists t such that $i < t < j$ and $C_t \subseteq J_{(l)_0}$. Let

$$v := \max \left\{ t \in \{i + 1, \dots, j - 1\} \mid C_t \subseteq J_{(l)_0} \right\}.$$

Again, by Lemma 2.6, there exists s such that $v \leq s < j$ and $C_s \subseteq J_{(l)\bar{i}}$. Let

$$w := \min \{s \in \{v, \dots, j-1\} \mid C_s \subseteq J_{(l)\bar{i}}\}.$$

We claim that (C_v, \dots, C_w) strongly refines $(J_{(l)_0}, \dots, J_{(l)\bar{i}})$. By definition, $C_v \subseteq J_{(l)_0}$ and $C_w \subseteq J_{(l)\bar{i}}$. Suppose $k \in \{v+1, \dots, w-1\}$. Then, since $\text{diam } C_k < \lambda$, C_k must be a subset of J_p, J_q or J_i for some $i \in [l]$. Suppose $C_k \subseteq J_p$. Since $C_w \subseteq J_{(l)\bar{i}}$, J_p strictly precedes $J_{(l)_0}$ and $J_{(l)_0}$ strictly precedes $J_{(l)\bar{i}}$, by Lemma 2.6 there exists u such that $k < u < w$ and $C_u \subseteq J_{(l)_0}$. However, this contradicts our choice of v because $v < k < u$. A similar argument shows that $C_k \not\subseteq J_q$, so C_k is a subset of J_i for some $i \in [l]$.

To summarize,

$$(C_0 \cup \dots \cup C_{v-1}, C_v, \dots, C_w, C_{w+1} \cup \dots \cup C_n)$$

is a compact quasi-chain in S' which covers S' and its subchain (C_v, \dots, C_w) strongly refines $(J_{(l)_0}, \dots, J_{(l)\bar{i}})$. By Lemma 3.2 (with $A = [l]$), we can find $p', l', q' \in \mathbb{N}$ such that

$$C_0 \cup \dots \cup C_{v-1} \subseteq_r J_{p'}, \quad C_{w+1} \cup \dots \cup C_n \subseteq_r J_{q'}, \quad (3.14)$$

$$C_v \subseteq_r J_{(l')_0}, \quad \dots, \quad C_w \subseteq_r J_{(l')\bar{i}}, \quad (3.15)$$

$$J_{(l')_0} \subseteq_{\forall} J_{(l)_0}, \quad J_{(l')\bar{i}} \subseteq_{\forall} J_{(l)\bar{i}} \quad \text{and} \quad J_{[l']} \subseteq_{\forall} J_{[l]} \quad (3.16)$$

and such that $(J_{p'}, J_{(l')_0}, \dots, J_{(l')\bar{i}}, J_{q'})$ is a formal chain.

Moreover, we have $\tilde{a} \in C_i \subseteq J_{p'}$, $\tilde{b} \in C_j \subseteq J_{q'}$ and

$$S' = f([-4, 4]) \cap S' = \bigcup_{i=1}^n C_i \subseteq J_{p'} \cup J_{[l']} \cup J_{q'}.$$

Therefore, (p', l', q') satisfies (O1)–(O4), which implies (G1); while (G2), (G4) and (G5) follow from (3.16). Finally, (3.13), (3.15) and Lemma 2.14 imply that

$$\text{fdiam}(k) < 4r + r \leq \frac{1}{2} \text{fmesh}(l), \quad \text{for each } k \in [l],$$

so (G3) also holds. This proves $(l, l') \in \Gamma$, so Claim 3 holds.

Claim 3 implies the existence of a partial recursive function $\psi : \Omega \rightarrow \mathbb{N}$ such that $(l, \psi(l)) \in \Gamma$ for all $l \in \Omega$. Let $l_0 \in \Omega$ be as in Claim 1 and define the sequence $(l_n)_{n \in \mathbb{N}}$ of natural numbers with

$$l_{n+1} := \psi(l_n), \quad \forall n \in \mathbb{N}.$$

Obviously, $(l_n)_{n \in \mathbb{N}}$ is recursive and $(l_n, l_{n+1}) \in \Gamma$ for all $n \in \mathbb{N}$.

Consider the sequence $(\mathcal{C}_n)_{n \in \mathbb{N}}$, where

$$\mathcal{C}_n := \left(J_{(l_n)_0} \cap S', \dots, J_{(l_n)\bar{i}_n} \cap S' \right), \quad \forall n \in \mathbb{N}. \quad (3.17)$$

By Claim 2, each \mathcal{C}_n is an open chain in S' . As a subspace of (X, d) , S' is a metric space in which every closed ball is compact (since S' itself is compact). Also, (G2) and Proposition 2.16(i) imply that

$$\left(\text{Cl}(J_{(l_{n+1})_0} \cap S'), \dots, \text{Cl}(J_{(l_{n+1})\bar{i}_{n+1}} \cap S') \right) \text{ strongly refines } \mathcal{C}_n \quad (3.18)$$

for each $n \in \mathbb{N}$. It follows from Claim 1 and (G3) that

$$\text{mesh}(\mathcal{C}_n) < 2^{-n}, \quad \forall n \in \mathbb{N}. \quad (3.19)$$

By Lemma 2.7,

$$N' := \bigcap_{n \in \mathbb{N}} \left(\bigcup_{j \in [l_{n+1}]} \text{Cl}(J_j \cap S') \right) \quad (3.20)$$

is a continuum in S' chainable from a to b , where

$$a \in \bigcap_{n \in \mathbb{N}} (J_{(l_n)_0} \cap S') \quad \text{and} \quad b \in \bigcap_{n \in \mathbb{N}} (J_{(l_n)_{l_n}} \cap S'). \quad (3.21)$$

Since

$$N' \subseteq S' \subseteq f(\langle -4, 4 \rangle) \subseteq N,$$

N' is a subset of N and a compact connected subset of $f(\langle -4, 4 \rangle)$, so it must be an arc. Since an arc can only be chainable from one of its endpoints to the other, a and b must be endpoints of N' .

Since $\text{fmesh}(l_n) < 2^{-n}$ for each $n \in \mathbb{N}$, (3.21) implies

$$d(a, \lambda_{(l_n)_0}) < 2^{-n} \quad \text{and} \quad d(b, \lambda_{(l_n)_{l_n}}) < 2^{-n}$$

for each $n \in \mathbb{N}$. It follows that a and b are computable points.

The function $n \mapsto \{\lambda_{(i)_0} \mid i \in [l_n]\}$ is computable. Let $n \in \mathbb{N}$. We claim that

$$N' \approx_{2^{-n}} \{\lambda_{(i)_0} \mid i \in [l_n]\}. \quad (3.22)$$

Let $y \in N'$. By (3.20) and (3.18), $y \in J_i \cap S'$ for some $i \in [l_n]$. Since $\text{fmesh}(l_n) < 2^{-n}$ and therefore $\text{fdiam}(i) < 2^{-n}$, it holds that

$$\text{diam}(J_i \cap S') \leq \text{diam}(J_i) \leq \text{fdiam}(i) < 2^{-n}.$$

Therefore, $d(y, \lambda_{(i)_0}) < 2^{-n}$.

Let $i \in [l_n]$. Since C_n is a chain which covers N' , its first link contains $a \in N'$ and its last link contains $b \in N'$, the connectedness of N' implies that its link $J_i \cap S'$ must intersect N' . Let $y \in N' \cap (J_i \cap S')$. Again, using $\text{fdiam}(i) < 2^{-n}$ we conclude that $d(y, \lambda_{(i)_0}) < 2^{-n}$.

It is not hard to conclude [Ilj13, Proposition 2.6] that there exists a computable function $\phi : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\{\lambda_{(i)_0} \mid i \in [l_n]\} = \{\alpha_i \mid i \in [\phi(k)]\},$$

so (3.22) implies that N' is a computable set in (X, d, α) . This concludes the proof of Theorem 3.3. \square

Lemma 3.4. *Let S be a semicomputable set in a computable metric space (X, d, α) . Suppose a point $x \in S$ has an open neighborhood N in S such that there exists a homeomorphism $f : [0, 1] \rightarrow N$ such that $f(0) = x$. Let $\varepsilon > 0$. Then there exists $a \in \langle 0, 1 \rangle$ such that $f(a)$ is a computable point, $S \setminus f([0, a])$ is a semicomputable set and $f([0, a]) \subseteq B(x, \varepsilon)$.*

Proof. Since f is continuous, there exists $t \in \langle 0, 1 \rangle$ such that

$$f([0, t]) \subseteq B(x, \varepsilon). \quad (3.23)$$

The set $f(\langle 0, 1 \rangle)$ is open in N and therefore open in S . Therefore $f(\langle 0, 1 \rangle)$ is an open neighborhood of $f(t)$ in S which is homeomorphic to \mathbb{R} and so Theorem 3.3 implies that there exists a computable neighborhood N' of $f(t)$ in S contained in $f(\langle 0, 1 \rangle)$ which is an arc with computable endpoints. Since f is a homeomorphism, $f^{-1}(N')$ is an arc in $\langle 0, 1 \rangle$, thus it is equal to $[a, b]$ for some $a, b \in \langle 0, 1 \rangle$, $a < t < b$ (see Figure 3).

Clearly, $f(a)$ is a computable point and

$$f([0, a]) \subseteq f([0, t]) \subseteq B(x, \varepsilon).$$

We claim that $S \setminus f([0, a])$ is a semicomputable set.




Figure 3: Points along an arc as in the proof of Lemma 3.4.
The highlighted arc is computable.

The set $f([0, b))$ is open in N and therefore open in S . Let U be an open set in (X, d) such that $f([0, b)) = S \cap U$. The set U is open, so it is a union of rational open balls. Since $f([0, a])$ is a compact set contained in U , there is a finite union of rational open balls J_m such that

$$f([0, a]) \subseteq J_m \subseteq U. \quad (3.24)$$

By Lemma 2.11, $S \setminus J_m$ is semicomputable. Since $f([a, b])$ is computable, it is also semicomputable, so $(S \setminus J_m) \cup f([a, b])$ is semicomputable.

Now $f(a)$ is a computable point and $f([0, a]) \subseteq B(x, \varepsilon)$. To prove that $S \setminus f([0, a])$ is semicomputable, we show that it is equal to $(S \setminus J_m) \cup f([a, b])$.

Indeed, since f is injective, $f([a, b])$ and $f([0, a])$ are disjoint, so $f([a, b]) \subseteq S \setminus f([0, a])$. It follows from (3.24) that $S \setminus J_m \subseteq S \setminus f([0, a])$. Therefore, $(S \setminus J_m) \cup f([a, b]) \subseteq S \setminus f([0, a])$.

Suppose $y \in S \setminus f([0, a])$. If $y \in J_m$, then by (3.24)

$$y \in U \cap S = f([0, b)) = f([0, a]) \cup f([a, b]),$$

which implies $y \in f([a, b])$. This proves $S \setminus f([0, a]) \subseteq (S \setminus J_m) \cup f([a, b])$. \square

4. COMPUTABLE APPROXIMATIONS OF SEMICOMPUTABLE SETS

4.1. Compact graphs. Let G be a compact graph. Then $G = \bigcup K$, where K is a nonempty finite family of (non-degenerate) line segments in \mathbb{R}^n such that the following holds:

$$\text{if } I, J \in K \text{ are such that } I \neq J \text{ and } I \cap J \neq \emptyset, \text{ then } I \cap J = \{a\}, \quad (4.1)$$

where a is an endpoint of both I and J .

Equivalently, a topological space G is a compact graph if and only if there exists a nonempty finite family \mathcal{A} of subspaces of G such that each $A \in \mathcal{A}$ is an arc, such that $G = \bigcup_{A \in \mathcal{A}} A$ and such that for all $A, B \in \mathcal{A}$ the following holds:

$$\text{if } A, B \in \mathcal{A} \text{ are such that } A \neq B \text{ and } A \cap B \neq \emptyset, \text{ then } A \cap B = \{a\}, \quad (4.2)$$

where a is an endpoint of both A and B [Ilj20, Remark 4.2]. If G is a compact graph and \mathcal{A} is a family with the described properties, then we say that \mathcal{A} **defines** G .

If G is a compact graph and if \mathcal{A} defines G , then $x \in G$ is an endpoint of G if and only if there exists a unique arc $A \in \mathcal{A}$ such that x is an endpoint of A [Ilj20]. In particular, the set of all endpoints of a compact graph G is finite.

If G is a compact graph and E the set of all its endpoints, then (G, E) has computable type [Ilj20]. Hence, if (X, d, α) is a computable metric space and S a semicomputable set in

this space which is, as a subspace of (X, d) , a compact graph, then S is computable if each endpoint of S is computable.

Suppose S is a semicomputable compact graph in a computable metric space (X, d, α) . If not all of its endpoints are computable, then S need not be computable (even a semicomputable arc in \mathbb{R} need not be computable [Mil02]). We want to show that such an S can be approximated by a computable subgraph T . Namely, for each uncomputable endpoint a of S , we cut off some small neighborhood of a in S and get a computable graph T with computable endpoints (see Figure 4). In general, endpoints of a computable graph T need not be computable, even when T is an arc [Mil02].




Figure 4: A semicomputable compact graph. Empty circles denote uncomputable points and filled circles denote computable points. The highlighted subset is a computable compact graph.

For simplicity, we will use the following notation: if A is an arc and $x, y \in A$ are such that $x \neq y$, then the unique subspace of A which is an arc whose endpoints are x and y will be denoted B by \overline{xy}^A .

Theorem 4.1. *Let (X, d, α) be a computable metric space and let S be a semicomputable compact graph in this space. Let $\{A_0, \dots, A_n\}$ be a family of arcs which defines S .*

Let Γ be the set of all endpoints of S which are uncomputable. Let $\varepsilon > 0$.

Then there exist arcs A'_0, \dots, A'_n such that for each $i \in \{0, \dots, n\}$ the following hold:

- (i) *if none of the endpoints of the arc A_i belong to Γ , then $A'_i = A_i$;*
- (ii) *if x and y are endpoints of A_i such that $x \in \Gamma$ and $y \notin \Gamma$, then $A'_i = \overline{zy}^{A_i}$, where $z \in A_i$ and $\overline{xz}^{A_i} \subseteq B(x, \varepsilon)$;*
- (iii) *if x and y are endpoints of A_i such that $x, y \in \Gamma$, then $A'_i = \overline{z_1 z_2}^{A_i}$, where $z_1, z_2 \in A_i$ and $\overline{xz_1}^{A_i} \subseteq B(x, \varepsilon)$, $\overline{z_2 y}^{A_i} \subseteq B(y, \varepsilon)$.*
- (iv) *the set $T = A'_0 \cup \dots \cup A'_n$ is a computable compact graph with computable endpoints.*

Proof. Let us define A'_0 in the following way. If none of the endpoints of A_0 belong to Γ , let $A'_0 = A_0$.

Suppose that $x \in \Gamma$ and $y \notin \Gamma$, where x and y are the endpoints of A_0 . There exists a homeomorphism $f : [0, 1] \rightarrow A_0$ such that $f(0) = x$ and $f(1) = y$. The set $f((0, 1))$ is open in S : its complement in S is equal to (note that $A_j \cap f((0, 1)) = \emptyset$ for $j \neq 0$ since A_j can intersect A_0 only in an endpoint, by the definition of a graph)

$$\{f(1)\} \cup \bigcup_{i \neq 0} A_i,$$

which is clearly a closed set in S .

Therefore, $f([0, 1))$ is an open neighbourhood of x in S . By Lemma 3.4, there exists $a \in \langle 0, 1 \rangle$ such that $f(a)$ is a computable point, $S \setminus f([0, a))$ is a semicomputable set and $f([0, a]) \subseteq B(x, \varepsilon)$.

Let $z = f(a) \in A_0$ and $A'_0 = \overline{zy}^{A_0}$. It holds that $\overline{xz}^{A_0} \subseteq B(x, \varepsilon)$. Now, the family of arcs $\{A'_0, A_1, \dots, A_n\}$ defines a compact graph T such that the set of all uncomputable endpoints of T is equal to $\Gamma \setminus \{x\}$. We also have

$$T = S \setminus f([0, a)),$$

so T is semicomputable.

Finally, if both endpoints x and y of A_0 belong to Γ , we proceed as in the previous case to obtain an arc of the form $\overline{xz_2}^{A_0}$, where z_2 is a computable point such that $z_2 \in A_0$, $\overline{z_2y}^{A_0} \subseteq B(y, \varepsilon)$ and such that $\overline{xz_2}^{A_0} \cup \bigcup_{i \neq 0} A_i$ is a semicomputable set. We then apply a similar procedure to the point x to obtain an arc of the form $\overline{z_1z_2}^{A_0}$, where z_1 is a computable point such that $z_1 \in \overline{xz_2}^{A_0}$, $\overline{xz_1}^{A_0} \subseteq B(x, \varepsilon)$ and such that $\overline{z_1z_2}^{A_0} \cup \bigcup_{i \neq 0} A_i$ is a semicomputable set. We define $A'_0 = \overline{z_1z_2}^{A_0}$.

As in the previous case, we have that the family $\{A'_0, A_1, \dots, A_n\}$ defines a semicomputable compact graph T such that the set of all uncomputable endpoints of T is equal to $\Gamma \setminus \{x, y\}$.

Now we define A'_1 in the same way and we get that the family $\{A'_0, A'_1, A_2, \dots, A_n\}$ defines a semicomputable compact graph T such that the set of all uncomputable endpoints of T is equal to $\{x \in \Gamma \mid x \text{ is an endpoint of } A_i \text{ for some } i \geq 2\}$.

In finitely many steps we get A'_0, \dots, A'_n which satisfy properties (i)-(iii) such that $\{A'_0, \dots, A'_n\}$ defines a semicomputable compact graph T which has no uncomputable endpoints. Hence, every endpoint of T is computable, so T is computable by Theorem 2.9. \square

An immediate consequence of Theorem 4.1 is the following fact.

Corollary 4.2. *Let (X, d, α) be a computable metric space and let S be a semicomputable compact graph in this space. Then for each $\varepsilon > 0$ there exists a computable compact graph T in (X, d, α) such that $T \subseteq S$, all endpoints of T are computable and $d_H(S, T) < \varepsilon$.*

4.2. Non-compact graphs. Having generalized notions of a semicomputable and computable set to non-compact sets, the question is under what conditions the implication

$$S \text{ semicomputable} \implies S \text{ computable} \tag{4.3}$$

holds. Beside the known conditions when S is compact, it is known that (4.3) holds when S is a graph such that each endpoint of S is computable [Ij20].

Any topological space homeomorphic to a ray is called a **topological ray**. If R is a topological ray and $a \in R$ such that $R \setminus \{a\}$ is connected, then we say that a is an **endpoint** of R . Note that each topological ray has a unique endpoint.

If G is a graph, then it follows easily from the definition that there exists a finite nonempty family \mathcal{A} of closed subspaces of G such that each element of \mathcal{A} is an arc or a topological ray, $G = \bigcup \mathcal{A}$ and for all $A, B \in \mathcal{A}$ the following holds:

$$\text{if } A, B \in \mathcal{A} \text{ are such that } A \neq B \text{ and } A \cap B \neq \emptyset, \text{ then } A \cap B = \{a\}, \tag{4.4}$$

where a is an endpoint of both A and B .

Conversely, let G be a topological space such that $G = \bigcup \mathcal{A}$, where \mathcal{A} is a finite nonempty family of closed subspaces of G such that each element of \mathcal{A} is an arc or a topological ray

and such that for all $A, B \in \mathcal{A}$ the implication (4.4) holds. We want to show that G is a graph.

Suppose $\mathcal{A} = \{A_0, \dots, A_m\}$. Similarly to [Ij20, Remark 4.2], we get that there exist $n \in \mathbb{N} \setminus \{0\}$ and finitely many subsets F_0, \dots, F_m of \mathbb{R}^n such that K_i is a line segment or a ray for each $i \in \{0, \dots, m\}$ and for each $i \in \{0, \dots, m\}$ there exists a homeomorphism $h_i : A_i \rightarrow K_i$ such that for all $i \neq j$ the following holds:

$$h_i(A_i \cap A_j) = h_j(A_i \cap A_j) = h_i(A_i) \cap h_j(A_j). \quad (4.5)$$

The fact that A_0, \dots, A_m are closed in G allows us to glue the maps h_0, \dots, h_m and to get a continuous bijection $f : G \rightarrow K_0 \cup \dots \cup K_m$. Each of the sets K_0, \dots, K_m is closed in \mathbb{R}^n , thus also in $K_0 \cup \dots \cup K_m$, and so f^{-1} is continuous too. Hence f is a homeomorphism. The equalities (4.5) imply that the family $\{K_0, \dots, K_m\}$ satisfies (2.3), which means that G is a graph.

If G is a graph and \mathcal{A} is a finite nonempty family of closed subspaces of G such that each element of \mathcal{A} is an arc or a topological ray and for all $A, B \in \mathcal{A}$ the implication (4.4) holds, then we say that \mathcal{A} **defines** G .

Let R be a topological ray and let $x, y \in R$, $x \neq y$. By \overline{xy}^R we denote a (unique) subspace of R which is an arc with endpoints x and y . Note the following: if a is an endpoint of R and $x \in R$, $x \neq a$, then $(R \setminus \overline{ax}^R) \cup \{x\}$ is a topological ray and a closed subspace of R .

The following theorem can be proved in the same way as Theorem 4.1.

Theorem 4.3. *Let (X, d, α) be a computable metric space and let S be a semicomputable graph in this space. Let $\{A_0, \dots, A_n\}$ be a family of arcs and topological rays which defines S . Let Γ be the set of all endpoints of S which are uncomputable. Let $\varepsilon > 0$. Then there exist arcs and topological rays A'_0, \dots, A'_n such that for each $i \in \{0, \dots, n\}$ the following hold:*

- (i) *if A_i is an arc whose endpoints do not belong to Γ , then $A'_i = A_i$;*
- (ii) *if A_i is an arc and x and y are endpoints of A_i such that $x \in \Gamma$ and $y \notin \Gamma$, then $A'_i = \overline{zy}^{A_i}$, where $z \in A_i$ and $\overline{xz}^{A_i} \subseteq B(x, \varepsilon)$;*
- (iii) *if A_i is an arc and x and y are endpoints of A_i such that $x, y \in \Gamma$, then $A'_i = \overline{z_1 z_2}^{A_i}$, where $z_1, z_2 \in A_i$ and $\overline{x z_1}^{A_i} \subseteq B(x, \varepsilon)$, $\overline{z_2 y}^{A_i} \subseteq B(y, \varepsilon)$.*
- (iv) *if A_i is a topological ray whose endpoint does not belong to Γ , then $A'_i = A_i$;*
- (v) *if A_i is a topological ray whose endpoint x belongs to Γ , then $A'_i = (A_i \setminus \overline{xz}^{A_i}) \cup \{z\}$, where $z \in A_i$ and $\overline{xz}^{A_i} \subseteq B(x, \varepsilon)$;*
- (vi) *the set $T = A'_0 \cup \dots \cup A'_n$ is a computable graph with computable endpoints.*

Although graphs in metric spaces are unbounded in general and therefore the Hausdorff distance does not make sense in this context, we do have the following immediate consequence of Theorem 4.3.

Corollary 4.4. *Let (X, d, α) be a computable metric space and let S be a semicomputable graph in this space. Then for each $\varepsilon > 0$ there exists a computable graph T in (X, d, α) such that $T \subseteq S$, all endpoints of T are computable and $S \approx_\varepsilon T$.*

4.3. 1-manifolds. A **1-manifold with boundary** is a second-countable Hausdorff space M such that each point of M has a neighborhood homeomorphic to $[0, \infty)$. The **boundary** ∂M of a such an M is defined as the set of all $x \in M$ such that x has a neighborhood in M homeomorphic to $[0, \infty)$ by a homeomorphism which maps x to 0.

If M is a 1-manifold with boundary, then each component of M is homeomorphic to one of these spaces [Sha11]:

$$\mathbb{R}, [0, \infty), \mathbb{S}^1, \text{ or } [0, 1].$$

We have the following: \mathbb{R} is the union of two rays in \mathbb{R} which intersect in a common endpoint, $[0, \infty)$ is a ray in \mathbb{R} , \mathbb{S}^1 is homeomorphic to the union of any three line segments in \mathbb{R}^2 each two of which intersect in a common endpoint, and $[0, 1]$ is a line segment in \mathbb{R} . In general, each component is a closed set. So if we additionally assume that M has finitely many components, it follows that M is a graph. Hence, each 1-manifold with boundary M with finitely many components is a graph and ∂M is the set of all endpoints of the graph M . We have the following consequence of Corollary 4.4.

Corollary 4.5. *Let (X, d, α) be a computable metric space and let M be a semicomputable 1-manifold in this space such that M has finitely many components. Then for each $\varepsilon > 0$ there exists a computable 1-manifold N in (X, d, α) such that $N \subseteq M$, each point of ∂N is computable and $M \approx_\varepsilon N$.*

REFERENCES

- [AH22] Djamel Eddine Amir and Mathieu Hoyrup. Computability of finite simplicial complexes. In *ICALP*, Paris, France, July 2022. URL: <https://inria.hal.science/hal-03564904>.
- [AH23] Djamel Eddine Amir and Mathieu Hoyrup. Strong computable type. *Computability*, 12(3):227–269, 2023.
- [BI14] K. Burnik and Z. Iljazović. Computability of 1-manifolds. *Logical Methods in Computer Science*, 10(2:8):1–28, 2014.
- [BP03] V. Brattka and G. Presser. Computability on subsets of metric spaces. *Theoretical Computer Science*, 305:43–76, 2003.
- [BW99] V. Brattka and K. Weihrauch. Computability on subsets of Euclidean space I: Closed and compact subsets. *Theoretical Computer Science*, 219:65–93, 1999.
- [ČHI21] Vedran Čačić, Marko Horvat, and Zvonko Iljazović. Computable subcontinua of semicomputable chainable hausdorff continua. *Theoretical Computer Science*, 892:155–169, 2021. URL: <https://www.sciencedirect.com/science/article/pii/S0304397521005478>, doi:10.1016/j.tcs.2021.09.018.
- [Ilj09] Z. Iljazović. Chainable and circularly chainable co-c.e. sets in computable metric spaces. *Journal of Universal Computer Science*, 15(6):1206–1235, 2009.
- [Ilj13] Z. Iljazović. Compact manifolds with computable boundaries. *Logical Methods in Computer Science*, 9(4:19):1–22, 2013.
- [Ilj20] Z. Iljazović. Computability of graphs. *Math. Log. Q.*, 66(1):51–64, 2020.
- [IP18] Z. Iljazović and B. Pažek. Computable intersection points. *Computability*, 7:57–99, 2018.
- [IS18] Z. Iljazović and I. Sušić. Semicomputable manifolds in computable topological spaces. *Journal of Complexity*, 45:83–114, 2018.
- [IV17] Zvonko Iljazović and Lucija Validžić. Computable neighbourhoods of points in semicomputable manifolds. *Annals of Pure and Applied Logic*, 168(4):840–859, 2017.
- [Kih12] T. Kihara. Incomputability of simply connected planar continua. *Computability*, 1(2):131–152, 2012.
- [Mil02] J. S. Miller. Effectiveness for embedded spheres and balls. *Electronic Notes in Theoretical Computer Science*, 66:127–138, 2002.
- [PER89] M.B. Pour-El and J.I. Richards. *Computability in Analysis and Physics*. Springer, Berlin, 1989.
- [Sha11] A.R. Shastri. *Elements of Differential Topology*. CRC Press, Taylor and Francis Group, 2011.
- [Tur36] A. M. Turing. On computable numbers, with an application to the entscheidungsproblem. *Proc. London Math. Soc.*, 42:230–265, 1936.
- [Wei93] K. Weihrauch. Computability on computable metric spaces. *Theoretical Computer Science*, 113:191–210, 1993.
- [Wei00] K. Weihrauch. *Computable Analysis*. Springer, Berlin, 2000.