# Differentially Private Learning Beyond the Classical Dimensionality Regime

Cynthia Dwork Harvard University dwork@seas.harvard.edu Pranay Tankala Harvard University pranay\_tankala@g.harvard.edu

Linjun Zhang Rutgers University linjun.zhang@rutgers.edu

November 20, 2024

#### Abstract

We initiate the study of differentially private learning in the proportional dimensionality regime, in which the number of data samples n and problem dimension d approach infinity at rates proportional to one another, meaning that  $d/n \to \delta$  as  $n \to \infty$  for an arbitrary, given constant  $\delta \in (0, \infty)$ . This setting is significantly more challenging than that of all prior theoretical work in high-dimensional differentially private learning, which, despite the name, has assumed that  $\delta = 0$  or is sufficiently small for problems of sample complexity O(d), a regime typically considered "low-dimensional" or "classical" by modern standards in high-dimensional statistics.

We provide sharp theoretical estimates of the error of several well-studied differentially private algorithms for robust linear regression and logistic regression, including output perturbation, objective perturbation, and noisy stochastic gradient descent, in the proportional dimensionality regime. The 1 + o(1) factor precision of our error estimates enables a far more nuanced understanding of the price of privacy of these algorithms than that afforded by existing, coarser analyses, which are essentially vacuous in the regime we consider.

We incorporate several probabilistic tools that have not previously been used to analyze differentially private learning algorithms, such as a modern Gaussian comparison inequality and recent universality laws with origins in statistical physics.

<sup>\*</sup>This work was supported in part by Simons Foundation Grant 733782, Cooperative Agreement CB20ADR0160001 with the United States Census Bureau, and NSF CAREER DMS-2340241.

# Contents

1	Introduction         1.1 Results Overview	<b>3</b> 5
	1.2 Technical Overview	11
	1.3 Related Work	13
<b>2</b>	Preliminaries	14
	2.1 Differential Privacy	14
	2.2 Convex Gaussian Minimax Theorem	16
	2.3 Universality	17
	2.4 Mathematical Miscellany	19
3	Objective Perturbation for Robust Linear Regression	22
	3.1 Special Case of the Utility Proof	28
	3.2 Proof of Theorem 3.7	29
	3.2.1 Step 1: Legendre Transform	30
	3.2.2 Step 2: CGMT Universality	31
	3.2.3 Step 3: CGMT Analysis	33
	3.2.4 Putting Steps 1, 2, and 3 Together	36
	3.3 Additional Proofs	38
4	Objective Perturbation for Logistic Regression	<b>39</b>
	4.1 Heuristic Derivation	41
	4.2 Proof of Theorem $4.2$	44
	4.2.1 Step 1: GFOM Universality	45
	4.2.2 Step 2: Legendre Transform	52
	4.2.3 Step 3: CGMT Analysis	54
	4.2.4 Putting Steps 1, 2, and 3 Together	57
5	Privacy of Objective Perturbation with Small $\lambda$	<b>59</b>
6	Output Perturbation	61
7	Noisy Stochastic Gradient Descent	64

# 1 Introduction

Over the last two decades, a groundbreaking line of research has demonstrated the limitations of classical statistical theory for modern high-dimensional learning problems. This vibrant area of research dates to the seminal works of [DMM09, DMM11, BM11a, BM11b, EKBB+13, Kar13]; more recent contributions may be found in [SC19, CS20, HS23, WZF24, Han24]. To give one example, Sur and Candès [SC19] showed that the classical theory for logistic regression can fail dramatically when applied in the proportional dimensionality regime, in which the number data samples n and problem dimension d approach infinity at rates proportional to one another, meaning that  $d/n \rightarrow \delta$  as  $n \rightarrow \infty$  for an arbitrary, given constant  $\delta \in (0, \infty)$ . In particular, while the classical theory predicts that the maximum likelihood estimator (MLE) is asymptotically unbiased with variance given by the inverse Fisher information, when, say,  $n \geq 5d$  or  $n \geq 10d$ , the MLE can exhibit nonzero bias, and greater variance than predicted by the classical theory. In a result we find particularly intriguing, they demonstrated a threshold for the dimensionality ratio d/n at which the very existence of the MLE undergoes a sharp *phase transition*.

Inspired by this body of work, we initiate the study of differentially private learning in the proportional dimensionality regime. As an illustrative example, consider the task of fitting a *d*-dimensional linear regression model to *n* labeled data samples  $(\boldsymbol{x}_i, y_i) \in \mathbb{R}^d \times \mathbb{R}$ . For simplicity, suppose that  $\boldsymbol{\beta}^* \in [-1, +1]^d$  denotes the unknown regression coefficients, that the feature vectors  $\boldsymbol{x}_i$  are subgaussian with covariance  $\frac{1}{d}\boldsymbol{I}_d$ , and that the labels  $y_i$  have constant variance given  $\boldsymbol{\beta}^*$  and  $\boldsymbol{x}_i$ . In this setting, it is known that there exists an  $(\varepsilon_{\text{DP}}, \delta_{\text{DP}})$ -differentially private algorithm [LKO22] such that for any sufficiently small  $\alpha > 0$ , the algorithm's output  $\hat{\boldsymbol{\beta}}$  has  $\operatorname{error}^1 \frac{1}{d} \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^*\|^2 \leq \alpha^2$  with probability 99%, and the algorithm only uses a sample of size

$$n \le O\left(\frac{d}{\alpha^2} + \frac{d + \log(1/\delta_{\rm DP})}{\alpha\varepsilon_{\rm DP}}\right) \cdot \operatorname{polylog}\left(\frac{1}{\alpha}\right).$$

This sample complexity upper bound is tight up to polylogarithmic factors [CWZ21], but nevertheless yields vacuous error bounds in the proportional regime. Indeed, even if we were to set  $\varepsilon_{\rm DP} = 1$ and  $\delta_{\rm DP} = 1/n^2$ , this upper bound would only guarantee that  $\alpha \leq O(1)$  because  $d/n = \Theta(1)$  in the proportional regime. In the absence of knowledge of the constant suppressed by the  $O(\cdot)$  notation, this bound on  $\alpha$  is no better than that of the trivial estimator  $\hat{\beta}_{\rm trivial} = 0$ , which has error

$$\frac{1}{d} \|\widehat{\boldsymbol{\beta}}_{\text{trivial}} - \boldsymbol{\beta}^{\star}\|^2 = \frac{1}{d} \|\boldsymbol{\beta}^{\star}\|^2 \le 1.$$

Therefore, the above-mentioned upper bound leaves unanswered the question of whether the algorithm in question exhibits a nontrivial privacy-utility tradeoff in the proportional regime (unless the constant  $\delta$  to which d/n converges is sufficiently small, a regime considered "low-dimensional" or "classical" by modern standards [Wai19]). Similar issues arise in other recent works on private linear regression, which assume either that  $n = \tilde{O}(d)$  or in some cases n = O(d) for a suppressed, unoptimized constant factor [Wan18, She19, VTJ22, BHH<sup>+</sup>24, BDE<sup>+</sup>24]. In this example of regression in the proportional regime, everything from the most trivial estimators to the most sophisticated algorithms achieves constant error, rather than error converging to zero. The relevant task then becomes understanding the precise values of these constants, which is of course not just of

<sup>&</sup>lt;sup>1</sup>The factor of 1/d in the estimation error is appropriate because the  $\boldsymbol{x}_i$  have covariance  $\frac{1}{d}\boldsymbol{I}_d$ . More generally, when the  $\boldsymbol{x}_i$  have covariance  $\boldsymbol{\Sigma} \in \mathbb{R}^{d \times d}$ , a natural notion of estimation error is  $\|\boldsymbol{\Sigma}^{1/2}(\boldsymbol{\beta} - \boldsymbol{\beta}^*)\|^2$ . A good rule of thumb is that in regression, the "scale" of  $\boldsymbol{\beta}$  ought to be inversely proportional to the "scale" of  $\boldsymbol{x}$ . All our statements regarding  $\|\boldsymbol{\beta}^*\| \leq O(\sqrt{d})$  and  $\|\boldsymbol{x}_i\| \leq O(1)$  and error metric  $\frac{1}{d}\|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|^2$  can be straightforwardly reformulated as statements regarding  $\|\boldsymbol{\beta}^*\| \leq O(1)$  and  $\|\boldsymbol{x}_i\| \leq O(\sqrt{d})$  and error metric  $\|\boldsymbol{\beta} - \boldsymbol{\beta}^*\|^2$ .

theoretical interest. Indeed, in genomic, neuroscience, and image data from various domains, the dimensionality d of the data is not only proportional to n, but often even larger than n, and so in all of these settings, we must go beyond the standard O(d/n) or  $O(\sqrt{d/n})$  convergence rates.

This example extends beyond linear regression, with similar challenges arising in other forms of regression and in areas such as heavy-tailed moment estimation [KSU20, LKKO21, HKM22, AKM<sup>+</sup>24], Gaussian parameter estimation [KLSU19, LKKO21, AL22, KMV22, KMS<sup>+</sup>22, AKT<sup>+</sup>23, HKMN23, BHS23, KDH23], high-dimensional hypothesis testing and selection [BKSW19, CKM<sup>+</sup>19, GKK<sup>+</sup>20, Nar22, PAA24], and empirical risk minimization [CMS11, KST12, BST14], which have all received sustained attention from the differentially private algorithm design community.

In this work, we carry out the first tight utility analyses of differentially private learning algorithms in the proportional dimensionality regime, where constants start to matter and order-wise bounds cease to be meaningful. We determine the utility, including the optimal constants, of three prominent differentially private estimation algorithms—output perturbation [DMNS06, CMS11], objective perturbation [CMS11, KST12], and noisy stochastic gradient descent (DP-SGD) [SCS13, BST14]—when applied to robust linear regression and logistic regression tasks. Our error bounds are tight up to a  $1 \pm n^{-\Omega(1)}$  factor for robust linear regression and a  $1 \pm e^{-(\log n)^{\Omega(1)}}$  factor for logistic regression. Our bounds depend in nuanced ways on the problem parameters, such as  $\varepsilon_{\rm DP}$ ,  $\delta_{\rm DP}$ , and the dimensionality ratio  $\delta$ . We summarize these results in Section 1.1 and preview our techniques in Section 1.2.

In the case of logistic regression, our main result for the (differentially private) objective perturbation algorithm closely parallels the breakthrough result of Sur and Candès [SC19] for the (non-private) maximum likelihood estimator. In that work, the authors supplanted the classical theory with a certain equation in  $\mathbb{R}^3$ , and the equation's solution characterizes the behavior of the MLE in the proportional regime (when it exists) more accurately than the Fisher information does. Our work similarly derives an equation in  $\mathbb{R}^3$ , now depending on the algorithm designer's chosen magnitude  $\nu > 0$  of noise for privacy, and its solution now characterizes the utility of the objective perturbation algorithm in the proportional regime, for a wide range of notions of utility. If one lets  $\nu \to 0$ , the objective perturbation algorithm's output coincides with the MLE (when it exists), and, as expected, our equations characterizing  $\hat{\beta}$  simplify to those of the non-private case.

**Contributions.** The contributions of this paper are summarized as follows:

- (1) We prove the first tight utility bounds for differentially private algorithms in the proportional dimensionality regime. We derive bounds for three commonly used algorithms: objective perturbation (Sections 3 and 4), output perturbation (Section 6), and DP-SGD (Section 7).
- (2) We introduce powerful techniques from the high-dimensional statistics literature that have not previously been used to analyze differentially private learning algorithms, which could be useful for many tasks in private high-dimensional statistics and machine learning beyond the ones we consider. The techniques include:
  - (a) The Convex Gaussian Minimax Theorem (CGMT), due to [Sto13, TOH15]. This theorem relates the value of a certain min-max optimization problem defined in terms of a Gaussian random matrix in  $\mathbb{R}^{n \times d}$  to the value of a simpler problem defined in terms of two Gaussian random vectors in  $\mathbb{R}^n$  and  $\mathbb{R}^d$ . Several of our utility analyses involve reformulating a differentially private algorithm as a min-max optimization problem, and then applying CGMT to achieve tight control of the algorithm's estimation error and, by duality, its residuals on the training data. Importantly, our proofs differ from prior CGMT

analyses of non-private estimators in several ways to account for the noise introduced for differential privacy.

- (b) Recent, rigorous formulations of the universality principle from statistical physics, which asserts that certain global properties of a dynamical system should be independent of its local details. We use universality to circumvent a weakness of CGMT: it requires Gaussian data. Gaussianity is incompatible with the private algorithms we consider, which all either require bounded data or enforce boundedness via feature- or gradient-clipping. Using universality, we extend our CGMT analyses to a broad class of non-Gaussian data distributions. Indeed, a major part of our proofs is verifying that the differentially private algorithms of interest to us indeed satisfy the various technical requirements needed to apply universality—specifically, the universality laws of [HS23, Han24], which were developed to study non-private estimators.
- (3) We prove an improved privacy bound for the objective perturbation algorithm (Section 5), the first to provide a DP guarantee even for arbitrarily small regularization strengths  $\lambda > 0$  and perturbation magnitudes  $\nu > 0$ . Combining our privacy bound with our tight utility bounds leads to the first meaningful *privacy-utility tradeoffs* for output perturbation, objective perturbation, and DP-SGD in the proportional dimensionality regime.

## 1.1 Results Overview

Our first technical result is a sharp utility analysis of the classic *objective perturbation* algorithm and the first such analysis to provide any guarantees at all in the proportional dimensionality regime. Objective perturbation is one of the most well-known differentially private empirical risk minimization algorithms, along with output perturbation and noisy stochastic gradient descent. Objective perturbation for risk minimization was introduced in the case of pure differential privacy by [CMS11], extended to the case of approximate differential privacy by [KST12], and studied further in several subsequent theoretical and empirical works [JT14, INS<sup>+</sup>19, NRVW20, RKW23].

For an informal description of the algorithm, suppose we are given a loss function  $\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y))$ and wish to select  $\boldsymbol{\beta} \in \mathbb{R}^d$  to minimize the average loss on a given data set of n labeled points  $(\boldsymbol{x}_1, y_1), \ldots, (\boldsymbol{x}_n, y_n) \in \mathbb{R}^d \times \mathbb{R}$ . The objective perturbation algorithm computes the minimizer  $\boldsymbol{\beta}$  of a *regularized* and *randomly perturbed* version of the standard objective function for empirical risk minimization:

$$\widehat{\boldsymbol{\beta}} = \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^d} \sum_{i=1}^n \ell(\boldsymbol{\beta}; (\boldsymbol{x}_i, y_i)) + \lambda \cdot \frac{1}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle.$$

The effect of the quadratic regularization term,  $\frac{1}{2} \|\boldsymbol{\beta}\|^2$ , is controlled by the *regularization strength*  $\lambda > 0$ . Similarly, the effect of the random linear perturbation term,  $\langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle$  for  $\boldsymbol{\xi} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{I}_d)$ , is controlled by the *perturbation strength*  $\nu > 0$  that ensures a certain level of differential privacy.

In [CMS11, KST12, BST14], the utility of  $\hat{\beta}$  is measured by its excess empirical risk:

$$\sum_{i=1}^{n} \ell(\widehat{\boldsymbol{\beta}}; (\boldsymbol{x}_i, y_i)) - \min_{\boldsymbol{\beta} \in \mathbb{R}^d} \sum_{i=1}^{n} \ell(\boldsymbol{\beta}; (\boldsymbol{x}_i, y_i)).$$

Essentially, this approach first compares  $\hat{\beta}$  to  $\hat{\beta}_{MLE} = \operatorname{argmin}_{\beta \in \mathbb{R}^d} \sum_{i=1}^n \ell(\beta; (x_i, y_i))$ , which can then be compared to a population version thereof, if desired. Another approach is to directly measure the utility of  $\hat{\beta}$  by its distance to a ground-truth vector  $\beta^* \in \mathbb{R}^d$  assumed to have generated the data. This alternate metric has also been used in many works on differentially private statistics,



Figure 1: The functions  $H_L$ ,  $H'_L$ , and  $H''_L$  with L = 1.

from the work of [DL09], to the more recent works on private Gaussian estimation starting with [KLSU19], and also on private regression [CWZ21, CWZ23].

One reason to study distance to  $\beta^*$  instead of excess empirical risk in the proportional dimensionality regime is offered by the work of [SC19], which showed that in this setting,  $\hat{\beta}_{MLE}$  is a remarkably bad point of reference even for tasks as straightforward as logistic regression—suffering from bias and inconsistency even for well-behaved data. (As a side note, we chose the subscript MLE because "maximum likelihood estimator" is an appropriate name in the common case where  $\ell$  computes the negative log likelihood of a statistical model with parameter  $\beta \in \mathbb{R}^d$ .) For this reason, in our work, we opt to directly measure the utility of  $\hat{\beta}$  by its estimation error with respect to  $\beta^*$ . Ultimately, our results will hold for a wide range of notions of distance to  $\beta^*$ .

Crucially, the regularization strength  $\lambda$  and the perturbation strength  $\nu$  determine both the privacy loss parameters ( $\varepsilon_{\rm DP}, \delta_{\rm DP}$ ) of  $\hat{\boldsymbol{\beta}}$  and also the utility, e.g.  $\frac{1}{d} \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2$ , as  $n \to \infty$  and  $d/n \to \delta$ . Although we analyze both relationships, it is the latter relationship that we precisely pin down in our work. For simplicity, we do so in the cases of robust linear regression and logistic regression, two fundamental and well-motivated learning tasks (in our later study of private gradient descent in Section 7, however, we will look at modified variants of these two tasks).

**Objective Perturbation for Robust Linear Regression.** Ordinary least squares linear regression is notoriously brittle in the presence of outliers in the data, and a popular method to improve outlier robustness is to replace squared loss with the *Huber loss* [Hub64], defined as

$$H_L(r) = \begin{cases} \frac{1}{2}r^2 & \text{if } |r| \le L\\ L|r| - \frac{1}{2}L^2 & \text{if } |r| \ge L \end{cases}$$

The Huber loss function coincides with the squared loss when the residual  $r = y - \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle$  is sufficiently small, but grows at a linear, rather than quadratic, rate when  $|r| \geq L$  for a specified constant L, which also happens to be the Lipschitz constant of the function. For ease of privacy analysis, we choose to study robust linear regression with Huber loss instead of traditional linear regression.

In our utility analysis of objective perturbation for robust linear regression (Theorem 1.1), we will consider data  $(\mathbf{X}, \mathbf{y}) \in \mathbb{R}^{n \times d} \times \mathbb{R}^n$  generated from a *linear model*, which means that there exists a ground-truth vector  $\boldsymbol{\beta}^* \in \mathbb{R}^d$  and suitably bounded error vector  $\boldsymbol{\varepsilon}^* \in \mathbb{R}^n$  independent of  $\mathbf{X}$  for which

$$y = X\beta^{\star} + \varepsilon^{\star}.$$

We will also assume that the feature matrix X is average-case in a certain sense, which is standard in the differentially private estimation literature. For concreteness, the reader can imagine that the entries are independent, random elements of  $\{-1/\sqrt{d}, +1/\sqrt{d}\}$ , but our results apply to various distributions with different conditions on the means and variances. We emphasize that privacy holds in a worst-case sense and requires no distributional assumptions. We state the informal theorem in terms of *zero-concentrated* differential privacy (zCDP) [DR16, BS16], which we define in Section 2.

**Theorem 1.1** (Robust Linear Regression with Objective Perturbation, Informal Version of Theorem 3.7). Let  $(\sigma^*, \tau^*)$  denote the solution to the following system of two scalar equations in two variables  $(\sigma, \tau)$ , which we write in terms of the algorithm's parameters  $\lambda, \nu > 0$ , the dimensionality ratio  $\delta \in (0, \infty)$ , dummy variables  $Z_1, Z_2 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$ , and  $\kappa^2 = \frac{1}{d} \|\boldsymbol{\beta}^*\|^2$  and  $\varepsilon^2 = \frac{1}{n} \|\boldsymbol{\varepsilon}^*\|^2$  as follows:

$$\sigma^{2} = \tau^{2} \left( \frac{1}{\delta} \mathbb{E} \left[ \min \left( L^{2}, \left( \frac{\sigma Z_{1} + \varepsilon Z_{2}}{1 + \tau} \right)^{2} \right) \right] + \lambda^{2} \kappa^{2} + \nu^{2} \right),$$
  
$$\tau = \frac{1}{\lambda \delta} \left( \delta - \frac{\tau}{1 + \tau} \Pr \left[ -L < \frac{\sigma Z_{1} + \varepsilon Z_{2}}{1 + \tau} < L \right] \right).$$

The output  $\hat{\beta}$  of the objective perturbation algorithm with Huber loss satisfies  $\rho_{\text{DP}}$ -zCDP for a constant  $\rho_{\text{DP}} = \rho_{\text{DP}}(L, \lambda, \nu)$  independent of n that is finite for any  $L, \lambda, \nu \in (0, \infty)$ . If the data are generated from a linear model as  $n \to \infty$  and  $d/n \to \delta$ , then w.h.p.,  $\hat{\beta}$  has estimation error

$$\frac{1}{d}\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2 = (\sigma^{\star})^2 \pm n^{-\Omega(1)}.$$

Theorem 1.1 is essentially a differentially private analogue of a result in Section 5.2 of [TAH18] for ridge regression with Gaussian features, which was later extended to subgaussian features by Theorem 3.12 of [HS23]. These works provide a pair of equations in  $(\sigma, \tau)$ , and the solution of this system can be used to characterize the behavior of the ridge regularized Huber regressor. Our pair of equations in Theorem 1.1 are quite similar, but we also capture the dependence on  $\nu$ , the magnitude of noise introduced for privacy by the objective perturbation algorithm. For intuition on where the two equations come from, we refer the reader to the technical overview in Section 1.2. We remark that as  $\nu \to 0$ , the equations of Theorem 1.1 recover those of [TAH18, HS23]. Thus, numerically comparing the solution to our equations with  $\nu > 0$  to those of prior work with  $\nu = 0$  allows us to precisely quantify the price of privacy of objective perturbation for robust linear regression. We carry out this comparison in Section 3.

Although the incorporation of  $\nu$  is the main quantitative difference between our theorem and those of [TAH18, HS23], our theorem also comes with a number of qualitative advantages that will become more apparent once we have presented its full version (Theorem 3.7). For example, while Theorem 3.12 of [HS23] assumes that  $\beta^*$  and  $\varepsilon^*$  are random with i.i.d. coordinates, our theorem does not even require them to be random (although, in the deterministic case, we do require certain mild conditions on their coordinates as  $n \to \infty$ ). Next, while [TAH18, HS23] only determine the limiting value of  $\frac{1}{d} \|\hat{\beta} - \beta^*\|^2$ , we also use  $\sigma^*$  and  $\tau^*$  to characterize a wide range of other scalar quantities of interest regarding the algorithm's output  $\hat{\beta}$ , both in relation to the ground-truth  $\beta^*$ and to the coefficients  $\boldsymbol{\xi}$  of the random linear perturbation term. Finally, the full version of our theorem studies not only  $\hat{\beta}$ , but also its predictions  $\boldsymbol{X}\hat{\beta}$  via duality.

To quickly build intuition for the distinct roles of  $\sigma^*$  and  $\tau^*$ , we remark that the full version of Theorem 1.1 will ultimately show that, in terms of the algorithm's perturbation vector  $\boldsymbol{\xi} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{I}_d)$  and an independent random vector  $\boldsymbol{z} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{I}_d)$ , we roughly have that

$$\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star} \approx -\tau^{\star} \lambda \boldsymbol{\beta}^{\star} - \tau^{\star} \nu \boldsymbol{\xi} + \sqrt{(\sigma^{\star})^2 - (\tau^{\star})^2 (\lambda^2 \kappa^2 + \nu^2)} \cdot \boldsymbol{z}.$$

By inspection, a larger  $\sigma^*$  leads to a larger coefficient on the z term, so  $\sigma^*$  is related to the variance of  $\hat{\beta} - \beta^*$ . Similarly, we see that  $\tau^*$  is related to the correlations of  $\hat{\beta}$  with both  $\boldsymbol{\xi}$  and  $\beta^*$ , the



Figure 2: The functions  $\rho$ ,  $\rho'$ ,  $\rho''$ , and  $\operatorname{prox}_{\gamma\rho}$  with  $\gamma = 3$ .

latter of which is related to the *bias* of  $\hat{\beta}$ . Formalizing this intuition will be easy to do given the full version of the theorem. Indeed, the bias, variance, and correlations are but a few of the "wide range" of scalar quantities to which we alluded in the previous paragraph.

Finally, we remark that the zCDP bound in Theorem 1.1, which we derive in Section 5, will be replaced in the full version (Theorem 3.7) with an even tighter privacy bound stated in terms of a family of ( $\varepsilon_{\text{DP}}, \delta_{\text{DP}}$ )-DP guarantees.

**Objective Perturbation for Logistic Regression.** In the case of binary classification, the response variable y lies in  $\{0,1\}^n$ , and a popular approach to understanding the relationship between y and X involves the *logistic loss*, which can be defined in terms of the convex function

$$\rho(t) = \log(1 + e^t).$$

Specifically, the logistic loss of  $\beta$  on a sample (x, y) is  $\rho(\langle x, \beta \rangle)$  if y = 0 and  $\rho(-\langle x, \beta \rangle)$  if y = 1.

In our utility analysis of objective perturbation for logistic regression (Theorem 1.2) we will consider data  $(\mathbf{X}, \mathbf{y}) \in \mathbb{R}^{n \times d} \times \{0, 1\}^n$  generated from a *logistic model*, in which there exists  $\boldsymbol{\beta}^{\star} \in \mathbb{R}^d$  with

$$\boldsymbol{y} \sim \text{Bernoulli}(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star})).$$

Here,  $\rho'(t) = 1/(1 + e^{-t})$  is the familiar sigmoid function that outputs values between 0 and 1.

The statement of Theorem 1.2, much like Theorem 1.1, will comprise a small number of scalar equations. This time, there are three equations instead of two, and we state them in terms of the *proximal operator* of the function  $\gamma \cdot \rho$  for a scalar  $\gamma > 0$ . Informally speaking, this function, denoted prox<sub> $\gamma\rho$ </sub>, takes as input a point  $s \in \mathbb{R}$  and outputs another point  $t \in \mathbb{R}$  that optimally balances two potentially competing considerations: minimization of the function  $\gamma \cdot \rho$  and proximity to the point s. It is helpful to know that as  $\gamma \to 0$ , the proximal operator roughly computes a gradient step of size  $\gamma$ , i.e. we have  $\operatorname{prox}_{\gamma\rho}(s) \approx s - \gamma \rho'(s)$ . For more detail, see Section 2 or [PB14].

**Theorem 1.2** (Logistic Regression with Objective Perturbation, Informal Version of Theorem 4.2). Let  $(\alpha^*, \sigma^*, \gamma^*)$  denote the solution to the following system of three scalar equations in three variables  $(\alpha, \sigma, \gamma)$ , which we write in terms of the algorithm's parameters  $\lambda, \nu > 0$ , the dimensionality ratio  $\delta \in (0, \infty)$ , dummy variables  $Z_1, Z_2 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$  and  $\kappa^2 = \frac{1}{d} \|\boldsymbol{\beta}^*\|^2$ , as follows:

$$\sigma^{2} = \gamma^{2} \left( \frac{1}{\delta} \mathbb{E} \Big[ 2\rho'(-\kappa Z_{1})\rho'(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_{1} + \sigma Z_{2}))^{2} \Big] + \nu^{2} \right),$$
  

$$\alpha = -\frac{1}{\delta} \mathbb{E} [2\rho''(-\kappa Z_{1})\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_{1} + \sigma Z_{2})],$$
  

$$\gamma = \frac{1}{\lambda\delta} \left( \delta - 1 + \mathbb{E} \left[ \frac{2\rho'(-\kappa Z_{1})}{1 + \gamma\rho''(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_{1} + \sigma Z_{2}))} \right] \right).$$

The output  $\hat{\beta}$  of the objective perturbation algorithm with logistic loss satisfies  $\rho_{\text{DP}}$ -zCDP for a constant  $\rho_{\text{DP}} = \rho_{\text{DP}}(\lambda, \nu)$  independent of n that is finite for any  $\lambda, \nu \in (0, \infty)$ . If the data are generated from a logistic model as  $n \to \infty$  and  $d/n \to \delta$ , then w.h.p.,  $\hat{\beta}$  has estimation error

$$\frac{1}{d}\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2 = (1 - \alpha^{\star})^2 \kappa^2 + (\sigma^{\star})^2 \pm e^{-(\log n)^{\Omega(1)}}.$$

Much like our results for robust linear regression, Theorem 1.2 is a differentially private analogue of existing, non-private results in the literature: this time the main result of Theorem 2 of [SAH19] for regularized logistic regression with Gaussian features, which was later extended to subgaussian features by Theorem 4.3 of [Han24]. These results are in turn generalizations of the landmark result of [SC19] on the behavior of the (non-regularized) MLE.

Once again, the main difference between the three equations in our Theorem 1.2 and those of prior work is our incorporation of the perturbation magnitude  $\nu$ . Taking  $\nu \to 0$ , we recover the system of equations of [SAH19], and taking both  $\nu \to 0$  and  $\lambda \to 0$ , we recover the equations of [SC19], up to some minor differences in notation and scaling conventions. Therefore, numerically comparing the solution to the equations of Theorem 1.2 with  $\nu > 0$  to those of prior work with  $\nu = 0$  allows us to precisely quantify the price of privacy in the case of objective perturbation for logistic regression. We carry out this comparison in Section 4.

Again, to quickly build intuition for the distinct roles of  $\alpha^*$ ,  $\sigma^*$ , and  $\gamma^*$ , we remark that the full version of Theorem 1.2 will ultimately show that, in terms of the algorithm's perturbation vector  $\boldsymbol{\xi} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{I}_d)$  and an independent random vector  $\boldsymbol{z} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{I}_d)$ , we roughly have that

$$\widehat{\boldsymbol{\beta}} \approx \alpha^{\star} \boldsymbol{\beta}^{\star} - \gamma^{\star} \boldsymbol{\nu} \boldsymbol{\xi} + \sqrt{(\sigma^{\star})^2 - (\gamma^{\star} \boldsymbol{\nu})^2} \cdot \boldsymbol{z}.$$

Consequently,  $\alpha^*$  is related to the *bias* of  $\hat{\beta}$  and  $\sigma^*$  is related to the variance of the debiased estimate  $\hat{\beta} - \alpha^* \beta^*$ . Moreover,  $\gamma^*$  is related to correlation of  $\hat{\beta}$  with  $\boldsymbol{\xi}$ . As in the case of robust linear regression, formalizing this intuition will be easy to do given the full version of the theorem. In the non-private case, a similar formulation exists involving only  $\beta^*$  and  $\boldsymbol{z}$  (the  $\boldsymbol{\xi}$  term disappears because  $\nu = 0$ ), where now  $\alpha^*$  and  $\sigma^*$  are defined in terms of the equations of [SC19] or [SAH19], depending on whether  $\lambda = 0$  or  $\lambda > 0$ .

Finally, we remark that the full version of our theorem (Theorem 4.2), will also have some qualitative advantages over prior results akin to those discussed for robust linear regression, and will also include a tighter privacy bounded stated in terms of ( $\varepsilon_{\text{DP}}, \delta_{\text{DP}}$ )-DP rather than  $\rho_{\text{DP}}$ -zCDP.

Privacy of Objective Perturbation with Small  $\lambda$ . Theorems 1.1 and 1.2 precisely determine the utility of objective perturbation as a function of the regularization strength  $\lambda$  and perturbation magnitude  $\nu$ , two parameters set by the algorithm designer, but these informal statements leave the algorithm's privacy loss parameter  $\rho_{\rm DP}(\lambda,\nu)$  unspecified. Ultimately, to precisely quantify the privacy-utility tradeoff of the algorithm, one needs explicit formulas for both the privacy loss and the utility as a function of  $\lambda, \nu$ . In Section 5, we determine, for any constant regularization strength  $\lambda > 0$  and any constant perturbation magnitude  $\nu > 0$ , privacy loss parameters independent of nfor which the algorithm satisfies ( $\varepsilon_{\rm DP}, \delta_{\rm DP}$ )-DP and  $\rho_{\rm DP}$ -zCDP even as  $n \to \infty$  and  $d/n \to \delta$ . Our proof builds on a closely related, recent result of [RKW23], which establishes privacy in the case where  $\lambda > s$  for a strictly positive constant s depending on the loss function's smoothness (e.g. s = 1 for robust linear regression, or s = 1/4 for logistic regression). We also show that the zCDP parameter  $\rho_{\rm DP}$  shrinks to 0 as  $\lambda$  and  $\nu$  increase. **Output Perturbation.** Our results on objective perturbation, which we laid out in the previous section, can easily be modified to characterize the utility of *output perturbation*, an even more foundational differentially private algorithm due to  $[DKM^+06]$ . Roughly speaking, [CMS11]suggest applying the output perturbation algorithm to empirical risk minimization by sampling  $\boldsymbol{\xi} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$  and then computing

$$\widehat{\boldsymbol{\beta}} = \left( \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^d} \sum_{i=1}^n \ell(\boldsymbol{\beta}; (\boldsymbol{x}_i, y_i)) + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 \right) + \nu \boldsymbol{\xi}.$$

As before, the behavior of the algorithm is controlled by the regularization strength  $\lambda > 0$  and perturbation strength  $\nu > 0$ , but here, the perturbation comes in the form of an additive noise term, rather than a random linear term in the objective. Extracting the  $\nu^2$  term from the systems of equations of Theorem 1.1 and Theorem 1.2 yields the following two corollaries, respectively.

**Corollary 1.3** (Linear Regression with Output Perturbation, Informal Version of Corollary 6.1). Let  $(\sigma^*, \tau^*)$  denote the solution to the system of equations from Theorem 1.1, replacing the first equation with:

$$\sigma^{2} = \tau^{2} \left( \frac{1}{\delta} \mathbb{E} \left[ \min \left( L^{2}, \left( \frac{\sigma Z_{1} + \varepsilon Z_{2}}{1 + \tau} \right)^{2} \right) \right] + \lambda^{2} \kappa^{2} \right).$$

Then, under the same conditions as Theorem 1.1, the estimation error of the perturbed output  $\hat{\beta}$  satisfies, w.h.p.,

$$\frac{1}{d}\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2 = (\sigma^{\star})^2 + \nu^2 \pm n^{-\Omega(1)}.$$

**Corollary 1.4** (Logistic Regression with Output Perturbation, Informal Version of Corollary 6.3). Let  $(\alpha^*, \sigma^*, \gamma^*)$  denote the solution to the system of equations from Theorem 1.2, replacing the first equation with

$$\sigma^{2} = \frac{\gamma^{2}}{\delta} \mathbb{E} \Big[ 2\rho'(-\kappa Z_{1})\rho'(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_{1} + \sigma Z_{2}))^{2} \Big].$$

Then, under the same conditions as Theorem 1.2, the estimation error of the perturbed output  $\hat{\beta}$  satisfies, w.h.p.,

$$\frac{1}{d} \|\widehat{\beta} - \beta^{\star}\|^2 = (1 - \alpha^{\star})^2 \kappa^2 + (\sigma^{\star})^2 + \nu^2 \pm e^{-(\log n)^{\Omega(1)}}$$

Noisy Stochastic Gradient Descent. The noisy stochastic gradient descent (a.k.a. DP-SGD) algorithm, which adds independent Gaussian noise to each iteration of SGD, is the cornerstone of modern, differentially private machine learning [ACG<sup>+</sup>16]. In Section 7, we present results for DP-SGD that parallel our results for objective perturbation (Theorems 1.1 and 1.2) and for output perturbation (Corollaries 1.3 and 1.4). These results for DP-SGD follow directly from a blackbox reduction to powerful, recent results in the non-private literature [GTM<sup>+</sup>24, Han24], and we validate them against simulated data. As such, we view this section more as a valuable point of reference than as a major technical contribution of our own. The results also come with some limitations. For example, certain technical requirements inherited from [GTM<sup>+</sup>24] prevent us from studying the standard formulations of robust linear regression and logistic regression, let alone the broad range of convex and non-convex optimization problems to which DP-SGD is typically applied. Also, the system of equations now has size  $O(T^2)$  for T steps of DP-SGD, where T must be held constant as  $n \to \infty$  and  $d/n \to \delta$ . Overcoming these obstacles is a promising direction for future work.

# 1.2 Technical Overview

In this section, we sketch the high-level idea behind our proofs of Theorems 1.1 and 1.2, which are our main results on objective perturbation in the proportional dimensionality regime. These proofs, which we carry out in Sections 3 and 4, respectively, comprise the bulk of our technical contribution. (As discussed in the previous section, our results for output perturbation are direct corollaries of Theorems 1.1 and 1.2, and our results for noisy stochastic gradient descent are similarly straightforward given prior results [ $GTM^+24$ , Han24] in the literature.)

Ultimately, for each of the three algorithms under consideration, we need two things:

• A privacy proof, which must hold for worst-case data points  $(\boldsymbol{x}_i, y_i)$ . These may be arbitrary points in  $B_R(\mathbf{0}) \times \mathbb{R}$  in the case of robust linear regression, or arbitrary points in  $B_R(\mathbf{0}) \times \{0, 1\}$ in the case of logistic regression, where  $B_R(\mathbf{0})$  denotes the ball of radius R centered at the origin in  $\mathbb{R}^d$ :

$$B_R(\mathbf{0}) = \{ \boldsymbol{x} \in \mathbb{R}^d : \|\boldsymbol{x}\| \le R \} \subseteq \mathbb{R}^d.$$

• A utility proof, which only needs to hold for data points  $(\boldsymbol{x}_i, y_i)$  sampled independently from an appropriate distribution over  $B_R(\mathbf{0}) \times \mathbb{R}$  in the case of robust linear regression, or over  $B_R(\mathbf{0}) \times \{0, 1\}$  in the case of logistic regression.

Our privacy proofs build on ones that exist in the differential privacy literature [BS16, BW18, RKW23], and we will hence inherit various boundedness requirements from those works. Specifically, to apply our privacy analysis to robust linear regression and logistic regression, for which the gradient of the loss function scales in magnitude with  $||\mathbf{x}||$ , we require the constraint that  $\mathbf{x}_1, \ldots, \mathbf{x}_n \in B_R(\mathbf{0})$ . This constraint can either be assumed as a priori knowledge, or enforced via a projection of unconstrained data in  $\mathbb{R}^d$  onto  $B_R(\mathbf{0})$ . Bounding the norm of each  $\mathbf{x}_i$  leads to a bound on the objective function's sensitivity, which ultimately yields our bound on the algorithm's privacy loss parameters ( $\varepsilon_{\text{DP}}, \delta_{\text{DP}}$ ) or  $\rho_{\text{DP}}$ .

Our utility proofs for objective perturbation are substantially more complex and comprise our main technical contribution. In these proofs, we initially focus on Gaussian features:

$$\boldsymbol{x}_1,\ldots,\boldsymbol{x}_n \stackrel{iid}{\sim} \mathcal{N}\left(\boldsymbol{0}, \frac{1}{d}\boldsymbol{I}_d\right).$$

The Gaussian distribution is not supported on  $B_R(\mathbf{0})$ , so we clearly must relax this assumption eventually. Nevertheless, Gaussianity is a fruitful starting point since it enables the use of Gaussian comparison inequalities. We focus on one such inequality called the *Convex Gaussian Minimax Theorem* (CGMT), which has seen very successful applications in the last decade to the kinds of high-dimensional regression problems of interest to us (for references, see the discussion of related works in Section 1.3). Roughly speaking, CGMT allows one to relate certain unwieldy optimization problems involving Gaussian random *matrices* to much simpler optimization problems involving Gaussian random vectors. We provide the formal statement of CGMT, along with a more detailed explanation of how it is used, in Section 2.

Since our aim is to apply CGMT, the first step of our proof is to reformulate the objective perturbation algorithm as min-max optimization of the form

$$\hat{\boldsymbol{\beta}} = \operatorname*{argmin}_{\boldsymbol{u} \in \mathcal{S}_{\boldsymbol{u}}} \max_{\boldsymbol{v} \in \mathcal{S}_{\boldsymbol{v}}} \underbrace{\langle \boldsymbol{X} \boldsymbol{u}, \boldsymbol{v} \rangle + \psi(\boldsymbol{u}, \boldsymbol{v})}_{Q_{\boldsymbol{u}, \boldsymbol{v}}}.$$

Here,  $S_{\boldsymbol{u}} \subseteq \mathbb{R}^d$  and  $S_{\boldsymbol{v}} \subseteq \mathbb{R}^n$  are constraint sets,  $\boldsymbol{X} = [\boldsymbol{x}_1 \cdots \boldsymbol{x}_n]^\top \in \mathbb{R}^{n \times d}$  denotes the feature matrix, which we also call the *design matrix*, and  $\psi : S_{\boldsymbol{u}} \times S_{\boldsymbol{v}} \to \mathbb{R}$  is a convex-concave function

that may depend on the ground-truth  $\beta^*$ , the regression errors  $\varepsilon^*$ , and the perturbation vector  $\boldsymbol{\xi}$ , but may *not* depend in any way on  $\boldsymbol{X}$ . We call  $\psi$  the *mean function* because  $\mathbb{E}[Q_{\boldsymbol{u},\boldsymbol{v}}] = \psi(\boldsymbol{u},\boldsymbol{v})$ , where the expectation is taken only over the randomness of  $\boldsymbol{X}$ .

At this point, we have reduced the task of understanding the behavior of  $\beta$  to the task of understanding the extrema of the random variables  $Q_{u,v}$  over  $u \in S_u$  and  $v \in S_v$ . In the cases of robust linear regression and logistic regression that we consider, the term  $\psi(u, v)$  will be very simple, so the main obstacle to optimizing  $Q_{u,v}$  by hand is the presence of the random matrix in the bilinear term  $\langle Xu, v \rangle$ .

Essentially, what CGMT enables us to do is replace the Gaussian random matrix  $\mathbf{X} \in \mathbb{R}^{n \times d}$ with two standard Gaussian random vectors  $\mathbf{g} \in \mathbb{R}^d$  and  $\mathbf{h} \in \mathbb{R}^n$ , after which solving the min-max optimization by hand becomes a straightforward matter of calculus. We take the derivatives of this *auxiliary optimization* with respect to  $\mathbf{u} \in \mathbb{R}^d$  and  $\mathbf{v} \in \mathbb{R}^n$  and set them equal to zero. Keeping in mind the constraint sets  $S_{\mathbf{u}}$  and  $S_{\mathbf{v}}$ , this yields a system of n + d equations in n + d variables characterizing the behavior of both this auxiliary optimizer and its corresponding dual variable. Ultimately, the guarantee of CGMT is that, over the randomness of  $\mathbf{X}$ , the solution to this auxiliary problem is indistinguishable from the solution  $\hat{\boldsymbol{\beta}}$  to the original problem, at least with respect to a wide range of *pseudo-Lipschitz* test functions.

To complete the proof in the case of Gaussian features, we show that, somewhat magically, this system of first-order optimality equations can be simplified to a fixed-size system of *scalar* equations (two equations in two variables in the case of robust linear regression, or three equations in three variables in the case of logistic regression) in the limit as  $n \to \infty$  with  $d/n \to \delta$ .

At this point, all that remains is to relax the Gaussianity assumption on X, which we emphasize is crucial for the analysis of any differentially private algorithm that assumes its input is bounded in  $B_R(\mathbf{0})$  or incorporates data clipping as a subroutine. To this end, we draw on recent, powerful theoretical results [HS23, Han24] that formalize the notion of *universality* from statistical physics. Informally speaking, universality is the idea that certain global properties of a system, such as the estimation error  $\frac{1}{d} \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2$ , should not depend too heavily on the local details of the system, such as the distribution of the entries of  $\boldsymbol{X}$ .

The first universality law we leverage is Corollary 2.6 of [HS23], which we call *CGMT universality*. This result tells us that even when X is not Gaussian, the distribution of the extrema of  $Q_{u,v}$  over  $u \in S_u$  and  $v \in S_v$  is essentially unchanged upon replacing X with a matrix X' with a few matching moments in the definition of  $Q_{u,v}$ . Taking X' = G to be Gaussian allows us to apply CGMT even to non-Gaussian X. This will ultimately allow us to consider X drawn from any isotropic distribution over  $B_R(\mathbf{0})$  with independent (not necessarily identically distributed!) entries. We believe our results can be extended to nonisotropic distributions, as well, which is a point we will revisit when we present the full versions of Theorems 1.1 and 1.2.

Applying CGMT universality requires several technical conditions on the original optimization problem to be verified. For our purposes, this roughly amounts to showing that the output  $\hat{\beta}$  of the objective perturbation algorithm satisfies  $\|\hat{\beta}\|_{\infty} \leq n^{o(1)}$ . We prove this in Lemma 3.14 using a leave-one-feature-out perturbation argument, a result that may be of independent interest.

Although CGMT universality as it appears in [HS23] is exactly what we need to analyze the objective perturbation with robust linear loss, frustratingly, it cannot be directly applied to logistic regression. The reason is that, in the case of the logistic loss, even the initial step of reformulating the perturbed objective as  $Q_{u,v} = \langle Xu, v \rangle + \psi(u, v)$  requires the assumption of Gaussianity to ensure the independence of the first and second terms, a requirement for CGMT. Consequently, we need a universality law that addresses the perturbed logistic objective function in its original form, before transforming it into  $Q_{u,v}$ . For this, we use the more recent universality law of [Han24], called generalized first-order method (GFOM) universality.

To apply GFOM universality, we first construct a sequence of " $\sigma$ -smoothed" iterates  $\beta_{\sigma}^{(t)}$  for  $t \in \mathbb{N}$  and  $\sigma > 0$  that start at  $\beta^{(0)} = \mathbf{0}$  but rapidly converge to  $\hat{\beta}$  as  $t \to \infty$  and  $\sigma \to 0$ . These iterates are *not* part of the objective perturbation algorithm, but rather exist solely for the purpose of analysis, as they fit the description of a "generalized first-order method" put forward in [CMW20]. What GFOM universality tells us is that the behavior of these iterates (and hence their limit  $\hat{\beta}$ ) is unchanged upon replacing X with a Gaussian matrix G with matching dimensions and a few matching moments. Our application of GFOM universality to the perturbed logistic objective is very similar to an argument of [Han24] carried out in the non-private case. The key differences arise in verifying that various technical conditions still hold in the presence of the random linear perturbation term introduced for privacy.

Ultimately, CGMT universality and GFOM universality allow us to reconcile the Gaussian design assumptions of CGMT analysis with the necessarily different data distributions encountered by differentially private algorithms, including output perturbation, objective perturbation, and noisy stochastic gradient descent. In terms of our final error bounds, CGMT universality, which we use for robust linear regression, incurs only an additive  $n^{-\Omega(1)}$  penalty. GFOM universality, which we use for logistic regression, incurs only an additive  $e^{-(\log n)^{\Omega(1)}}$  penalty.

# 1.3 Related Work

In recent years, a substantial body of work has focused on clarifying the capabilities and limitations of algorithms for differentially private regression [DL09, CMS11, KST12, Mir13, BST14, WFS15, She17, She19, LKO22, VTJ22, AJRV23, BHH<sup>+</sup>24, BDE<sup>+</sup>24]. The algorithms considered in these works vary along several axes, including time complexity, assumptions on the data, and the notion of privacy achieved, but to date, none have attempted to provide substantive theoretical guarantees in the proportional dimensionality regime as we do in this work. Beyond regression, there is an even larger body of work looking at other kinds of differentially private statistical problems, including but not limited to moment estimation for both light-tailed and heavy-tailed multivariate data [KLSU19, KSU20, LKKO21, HKM22, AL22, KMV22, KMS<sup>+</sup>22, AKT<sup>+</sup>23, HKMN23, BHS23, KDH23, AKM<sup>+</sup>24], learning multivariate mixture distributions [KSSU19, AAL21, AAL24], and high-dimensional hypothesis testing and selection [BKSW19, GKK<sup>+</sup>20, PAA24]. We suspect that a number of these problems might benefit from a treatment like ours in this new parameter regime.

One exception to the claim that our work is the first to consider the proportional regime is that there have indeed been works studying differentially private estimation in the regime where  $d \gg n$ under *sparsity* assumptions [SU17, CWZ21, ZWC<sup>+</sup>22, CWZ23, GH24, MJY24, KBY24], which we do not make. Such works are not comparable to ours, and under sparsity assumptions, the analogous challenge would be to analyze privacy-utility tradeoffs in the regime where n is proportional to, say,  $s \log(d/s)$ , where s is the sparsity level. Indeed, one can make a number of different arguments (e.g. via sample complexity, via  $\varepsilon$ -nets, or via the statistical dimension of [ALMT14]) that the effective "dimension" of some such problems is better described by  $O(s \log(d/s))$  than by O(d).

At a technical level, our proofs rely on certain probabilistic tools that, to our knowledge, have yet to appear in the DP literature, but have seen fruitful applications in the (non-private) modern statistics literature. For example, the Convex Gaussian Minimax Theorem that we use, though based on the classical Slepian's and Gordon's inequalities [Sle62, Gor85], was only presented in its current form and applied to modern statistical problems in the last decade [Sto13, TOH15, TAH18, DL20, DKT21, MM21, LGC<sup>+</sup>21, WWM21, LS22, ZZY22, HL22, MRSY23, CMW23]. Prior to this, the leading techniques for analysis in the proportional dimensionality regime included the *approximate message passing* (AMP) framework [DMM09, BM11a], which is arguably more complicated than CGMT, among other tools in random matrix analysis. The universality laws we use are from [HS23, Han24], but there are many works on this topic, including but not limited to [KM11, Kar13, BLM15, PH17, OT17, EK18, ASH19, MN17, DLS23, DSL24]. Privacy analyses we draw on include the works of [BS16, BW18, RW21, RKW23].

A final remark we make is that in the DP literature, the phrase "high-dimensional" typically refers to any problem in which d is large and the dependence of n on d is of interest, but in other communities, "high-dimensional" may refer to the proportional dimensionality regime, or the regime in which  $d \gg n$  under sparsity assumptions. In order to avoid confusion in our paper, which builds on prior work in multiple communities, we have opted to consistently use the phrase "proportional dimensionality" regime rather than "high-dimensional" regime when appropriate.

# 2 Preliminaries

Throughout this paper, when working with an infinite sequence indexed by  $n \in \mathbb{N}$ , which may be a sequence of scalars, vectors, matrices, functions, or subsets of  $\mathbb{R}^n$ , we will often refer to the sequence as though it were a single object of that type. For example, when referring to the problem dimension d, it should be understood that d is not merely a scalar, but rather a scalar sequence d(n) (equivalently,  $d_n$ ) indexed by n.

Consequently, by the proportional dimensionality regime, we mean the setting in which

$$\lim_{n \to \infty} \frac{d(n)}{n} = \delta,$$

where we denote the given dimensionality ratio by  $\delta \in (0, \infty)$ . In the proportional regime, any scalar sequence f(n) satisfying  $f(n) = \Theta(n^c)$  for a fixed constant c > 0 also satisfies  $f(n) = \Theta(d^c)$ , and vice versa. For consistency, when presented with such a notational choice, we shall always choose to express asymptotic bounds in terms of n, not d.

# 2.1 Differential Privacy

The algorithms we study in this paper all satisfy differential privacy (DP), the accepted definition of rigorous privacy protection for data analysis. We focus on the notion of *approximate*, or  $(\varepsilon, \delta)$ -differential privacy with respect to the *replace-one* notion of adjacency:

**Definition 2.1** ([DMNS06]). Given a universe  $\mathcal{X}$ , we say data sets  $\boldsymbol{x}, \boldsymbol{x}' \in \mathcal{X}^n$  are *adjacent* if  $x_i = x'_i$  except possibly in a single coordinate  $i \in [n]$ . We say a randomized algorithm  $\mathcal{A} : \mathcal{X}^n \to \mathcal{Y}$  satisfies  $(\varepsilon, \delta)$ -differential privacy if for all adjacent  $\boldsymbol{x}, \boldsymbol{x}' \in \mathcal{X}^n$  and subsets  $Y \subseteq \mathcal{Y}$ ,

$$\Pr[\mathcal{A}(\boldsymbol{x}) \in Y] \le e^{\varepsilon} \Pr[\mathcal{A}(\boldsymbol{x}') \in Y] + \delta.$$

We shall also require two other, related notions of differential privacy, known as *Rényi* and *zero-concentrated* DP, that have been used in major real-world deployments of DP. They are both defined in terms of the *Rényi divergence* of order  $\alpha > 1$  between two distributions P and Q over  $\mathcal{Y}$ :

$$D_{\alpha}(P||Q) = \frac{1}{\alpha - 1} \log \mathop{\mathbb{E}}_{y \sim P} \log \left(\frac{P(y)}{Q(y)}\right)^{\alpha - 1}$$

**Definition 2.2** (RDP [Mir17]). Given  $\alpha > 1$  and  $\varepsilon \ge 0$ , we say a randomized algorithm  $\mathcal{A} : \mathcal{X}^n \to \mathcal{Y}$  satisfies  $(\alpha, \varepsilon)$ -*Rényi DP* (RDP) if for all adjacent data sets  $\boldsymbol{x}, \boldsymbol{x}' \in \mathcal{X}^n$ ,

$$D_{\alpha}(\mathcal{A}(\boldsymbol{x}) \| \mathcal{A}(\boldsymbol{x}')) \leq \varepsilon.$$

Importantly, it is known that  $(\alpha, \varepsilon)$ -RDP implies  $\left(\varepsilon + \frac{\log(1/\delta)}{\alpha - 1}, \delta\right)$ -DP for any  $0 < \delta < 1$  [Mir17].

**Definition 2.3** (zCDP [BS16]). We say a randomized algorithm  $\mathcal{A} : \mathcal{X}^n \to \mathcal{Y}$  satisfies  $\rho$ -zeroconcentrated DP (zCDP) if  $\mathcal{A}$  satisfies  $(\alpha, \rho \alpha)$ -RDP for all  $\alpha > 1$ .

Outside of this section, we will use  $\varepsilon_{\text{DP}}$ ,  $\delta_{\text{DP}}$ ,  $\alpha_{\text{DP}}$ ,  $\rho_{\text{DP}}$  to refer to an algorithm's privacy parameters, rather than  $\varepsilon$ ,  $\delta$ ,  $\alpha$ ,  $\rho$ , in order to avoid confusion with other variable names. These include the regression errors  $\varepsilon$ , the dimensionality ratio  $\delta$ , the bias variable  $\alpha$ , and the function  $\rho$  used to define the logistic loss. For more background on differential privacy, including the motivation behind the definition, the protections it provides, interpretations of the privacy parameters, privacy composition theorems, and more, we refer the reader to [DR14, Vad17].

We collect here a handful of recent privacy analyses from the DP literature that we will repeatedly use in our paper. The first two results concern the *Gaussian mechanism*  $\mathcal{A}(\boldsymbol{x}) = f(\boldsymbol{x}) + \boldsymbol{z}$ where  $\boldsymbol{z} \sim \mathcal{N}(\boldsymbol{0}, \nu^2 \boldsymbol{I}_d)$  [DKM<sup>+</sup>06]. To state the result, which we will eventually use to analyze the output perturbation algorithm, recall that a function  $f : \mathcal{X}^n \to \mathbb{R}^d$  has  $\ell^2$  sensitivity at most  $\Delta$  if  $\|f(\boldsymbol{x}) - f(\boldsymbol{x}')\| \leq \Delta$  for all adjacent  $\boldsymbol{x}, \boldsymbol{x}' \in \mathcal{X}^n$ .

**Theorem 2.4** (Lemma 2.5 of [BS16]). If  $f : \mathcal{X}^n \to \mathbb{R}^d$  has  $\ell^2$  sensitivity at most  $\Delta$ , then the Gaussian mechanism w.r.t. f satisfies  $\rho$ -zCDP for  $\rho = \frac{\Delta^2}{2\nu^2}$ .

**Theorem 2.5** (Theorem 8 of [BW18]). If  $f : \mathcal{X}^n \to \mathbb{R}^d$  has  $\ell^2$  sensitivity at most  $\Delta$ , then for any  $\varepsilon \geq 0$  and  $\delta \in [0, 1]$ , the Gaussian mechanism w.r.t. f satisfies  $(\varepsilon, \delta)$ -DP if and only if

$$\delta \geq \operatorname{HockeyStick}\left(\varepsilon, \frac{\Delta}{\nu}\right) = \Phi\left(\frac{\Delta}{2\nu} - \frac{\varepsilon\nu}{\Delta}\right) - e^{\varepsilon}\Phi\left(-\frac{\Delta}{2\nu} - \frac{\varepsilon\nu}{\Delta}\right).$$

The name "HockeyStick" comes from the term *hockey stick divergence*, defined for  $\varepsilon \geq 0$  and distributions P and Q on  $\mathbb{R}$  as  $H_{e^{\varepsilon}}(P||Q) = \int_{-\infty}^{\infty} \max(0, P(x) - e^{\varepsilon}Q(x)) dx$ . As we have defined it, HockeyStick $(\varepsilon, \Delta/\nu)$  measures this divergence between  $P = \mathcal{N}(\Delta, \nu^2)$  and  $Q = \mathcal{N}(0, \nu^2)$ , which only depends on  $\Delta$  and  $\nu$  through their ratio  $\Delta/\nu$ .

The next result we state is a recent privacy analysis of objective perturbation from [RKW23]. Although we will not use this result directly in our paper, this result and its proof are the starting point from which our own privacy analysis in Section 5 proceeds. Indeed, the statement and proof of our privacy result will be very similar to that of [RKW23]. The key difference is that ours will apply to any regularization strength  $\lambda > 0$  and perturbation strength  $\nu > 0$ , but that of [RKW23] requires  $\lambda > s$  for some strictly positive s depending on the smoothness of the loss function.

The result of [RKW23], which we now state, concerns loss functions of the form  $\ell(\beta; (x, y)) = \ell_0(\langle x, \beta \rangle, y)$  for a function  $\ell_0 : \mathbb{R}^2 \to \mathbb{R}$ . Such loss functions are called *generalized linear model* (GLM) loss functions, and as the authors of [RKW23] emphasize, most privacy proofs for objective perturbation in the DP literature require this GLM assumption, even if they do not state it explicitly. We also remark that the result uses *add/remove-one* adjacency, in which a data set is viewed as a multiset (i.e. a function  $\mathcal{X} \to \{0\} \cup \mathbb{N}$ ), and two data sets x and x' are said to be adjacent if they differ by 1 in  $\ell^1$  norm. This is slightly different than the notion of adjacency we defined in Definition 2.1, but straightforward conversions between the two adjacency notions are possible.<sup>2</sup>

<sup>&</sup>lt;sup>2</sup>The statement of Theorem 2.6 is *not* identical to the statements of Theorems 3.1 and 3.2 in [RKW23] because we have corrected a few typos in their statement: first, in the  $(\varepsilon, \delta)$ -DP bound, we have added absolute value signs around the term  $\log(1 - s/\lambda)$  in the definition of  $\tilde{\varepsilon}$ . Similarly, we have added a factor of 2 in the denominator of  $L^2/2\nu^2$ . The absolute value signs and the factor of 2 are both present throughout the proof of Theorem 3.1 of [RKW23], and are only omitted in the theorem's statement.

**Theorem 2.6** (Theorems 3.1 and 3.2 of [RKW23]). Suppose that  $\ell_0 : \mathbb{R}^2 \to \mathbb{R}$  satisfies  $|\partial_1 \ell_0(\eta, y)| \leq L$  and  $0 \leq \partial_1^2 \ell_0(\eta, y) \leq s$  for some constants L, s > 0 and for all  $\eta, y \in \mathbb{R}$ . Then objective perturbation (Algorithm 1) with R = 1, any  $\lambda, \nu > 0$ , and the GLM loss function  $\ell(\beta; (x, y)) = \ell_0(\langle x, \beta \rangle, y)$  satisfies add/remove-one  $(\varepsilon, \delta)$ -differential privacy for any  $\varepsilon \geq 0$  and

$$\delta = \begin{cases} 2 \cdot \operatorname{HockeyStick}(\tilde{\varepsilon}, \frac{L}{\nu}) & \text{if } \hat{\varepsilon} \ge 0, \\ (1 - e^{\hat{\varepsilon}}) + 2e^{\hat{\varepsilon}} \cdot \operatorname{HockeyStick}\left(\frac{L^2}{2\nu^2}, \frac{L}{\nu}\right) & \text{otherwise}, \end{cases}$$

where we set  $\tilde{\varepsilon} = \varepsilon - |\log(1 - s/\lambda)|$  and  $\hat{\varepsilon} = \tilde{\varepsilon} - L^2/2\nu^2$ , provided that  $\lambda > s$ .

The algorithm also satisfies add/remove-one  $(\alpha, \varepsilon)$ -Rényi differential privacy for any  $\alpha > 1$  and

$$\varepsilon = -\log\left(1 - \frac{s}{\lambda}\right) + \frac{L^2}{2\nu^2} + \frac{1}{\alpha - 1}\log\mathbb{E}_{X \sim \mathcal{N}\left(0, \frac{L^2}{\nu^2}\right)}\left[e^{(\alpha - 1)|X|}\right].$$

The following lemma clarifies the conversion to which we alluded earlier from add/remove-one DP to replace-one DP, the version in Definitions 2.1 and 2.2.

**Lemma 2.7** (Size-2 Group Privacy). If a randomized algorithm  $\mathcal{A}$  satisfies add/remove-one  $(\varepsilon, \delta)$ -DP, then  $\mathcal{A}$  also satisfies replace-one  $(2\varepsilon, (1 + e^{\varepsilon})\delta)$ -DP. Similarly, if  $\mathcal{A}$  satisfies add/remove-one  $\rho$ -zCDP, then  $\mathcal{A}$  also satisfies replace-one  $4\rho$ -zCDP.

# 2.2 Convex Gaussian Minimax Theorem

In this section, we present the Convex Gaussian Minimax Theorem (CGMT), due to [Sto13, TOH15], which will play a key role in our analysis of objective perturbation.

As discussed in the technical overview in Section 1.2, CGMT allows us to better understand the optimizers of a certain class of min-max optimization problems that are defined in terms of a Gaussian random matrix  $\boldsymbol{G} \in \mathbb{R}^{n \times d}$ . The theorem relates the original problem to an *auxiliary* optimization problem in which the random matrix  $\boldsymbol{G}$  has been replaced with two random vectors  $\boldsymbol{g} \in \mathbb{R}^d$  and  $\boldsymbol{h} \in \mathbb{R}^n$ , whose dimensions match the number of columns and rows of  $\boldsymbol{G}$ , respectively. Remarkably, the theorem shows that the optimum value of the original problem, which is a scalarvalued function of  $\boldsymbol{G}$ , has a similar distribution to the optimum value of the auxiliary problem, which is a scalar-valued function of  $(\boldsymbol{g}, \boldsymbol{h})$  that is much easier to determine directly via calculus. The main assumption needed for CGMT is that the original min-max problem is convex-concave and depends bilinearly on  $\boldsymbol{G}$ .

At first glance, it may seem that knowing the optimum value of the original problem is not very useful. After all, in the proof sketch we outlined in Section 1.2, what we really wanted to understand was the *optimizer*  $\hat{\beta}$ , which is the location at which the optimum value is achieved. It turns out that by carefully choosing the constraint sets of the problem to which CGMT is applied, we can get our hands on a remarkable number of quantitative properties about  $\hat{\beta}$ . For example, suppose we would like to show that the estimation error of  $\hat{\beta}$  with respect to the ground-truth  $\beta^*$ is approximately  $(\sigma^*)^2$  in the limit. To do so, consider a small slack factor  $\varepsilon_n > 0$  and define the constraint set

$$\mathcal{S}_{\varepsilon_n} = \left\{ \boldsymbol{\beta} \in \mathbb{R}^d : \left| \frac{1}{d} \| \boldsymbol{\beta} - \boldsymbol{\beta}^{\star} \|^2 - (\sigma^{\star})^2 \right| \ge \varepsilon_n \right\}.$$

If we can show that, with high probability, the optimum value of the auxiliary problem over  $S_{\varepsilon_n}$  is strictly larger than the optimum value of the auxiliary problem over  $\mathbb{R}^d$ , then by CGMT, the same must be true of original problem. Since  $\hat{\beta}$  is the optimizer of the original problem, this would allow us to conclude that with high probability,  $\hat{\beta} \notin S_{\varepsilon_n}$ , so  $\frac{1}{d} \|\hat{\beta} - \beta^*\|^2 = (\sigma^*)^2 \pm \varepsilon_n$ . Nothing about the previous example is particularly specific to  $\ell^2$  estimation error, which suggests (correctly) that the argument can be carried out for a wide range of test functions. It turns out that we will be able to handle a broad class of *pseudo-Lipschitz* test functions. We will be able to handle slack factors on the order of  $\varepsilon_n = n^{-\Omega(1)}$  in the case of robust linear regression, or  $\varepsilon_n = e^{-(\log n)^{\Omega(1)}}$  in the case of logistic regression, both of which satisfy  $\varepsilon_n \to 0$  as  $n \to \infty$ .

**Theorem 2.8** (Convex Gaussian Minimax Theorem [Sto13, TOH15]). Given compact sets  $S_{\boldsymbol{u}} \subseteq \mathbb{R}^d$ and  $S_{\boldsymbol{v}} \subseteq \mathbb{R}^n$ , a continuous function  $\psi : S_{\boldsymbol{u}} \times S_{\boldsymbol{v}} \to \mathbb{R}$ , a matrix  $\boldsymbol{G} \in \mathbb{R}^{n \times d}$ , and vectors  $\boldsymbol{g} \in \mathbb{R}^d$ ,  $\boldsymbol{h} \in \mathbb{R}^n$ , consider the following two random variables indexed by  $\boldsymbol{u} \in S_{\boldsymbol{u}}$  and  $\boldsymbol{v} \in S_{\boldsymbol{v}}$ :

$$egin{aligned} Q_{oldsymbol{u},oldsymbol{v}} &= \langle oldsymbol{G}oldsymbol{u},oldsymbol{v} 
angle + \psi(oldsymbol{u},oldsymbol{v}) \ Q'_{oldsymbol{u},oldsymbol{v}} &= \|oldsymbol{u}\|\langleoldsymbol{h},oldsymbol{v}
angle + \|oldsymbol{v}\|\langleoldsymbol{g},oldsymbol{u}
angle + \psi(oldsymbol{u},oldsymbol{v}) \ \end{array}$$

We call  $Q_{u,v}$  and  $Q'_{u,v}$  the primary and auxiliary objective functions. If G, g, and h all have independent  $\mathcal{N}(0,\sigma^2)$  entries for some  $\sigma > 0$ , then:

(a) For any threshold  $t \in \mathbb{R}$ ,

$$\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}} < t\right] \leq 2\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}' < t\right].$$

(b) If  $S_u$  and  $S_v$  are convex and  $\psi$  is convex-concave on  $S_u \times S_v$ , then for any threshold  $t \in \mathbb{R}$ ,

$$\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}>t\right] \leq 2\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}'>t\right].$$

#### 2.3 Universality

Here, we state the two key universality laws from prior work that we will use in our analyses. The first result is Corollary 2.6 of [HS23], which we call CGMT universality. We have slightly rephrased the statement for clarity, at the cost of generality (e.g. by assuming that the mean function  $\psi$  is differentiable). We will eventually use this result in our analyses of objective perturbation and output perturbation for robust linear regression. As we discussed earlier in the technical overview of Section 1.2, this result allows us to extend CGMT analysis to non-Gaussian design matrices, as long as we have a few finite, matching moments.

**Theorem 2.9** (CGMT Universality: Corollary 2.6 of [HS23]). Consider random matrices  $A, B \in \mathbb{R}^{n \times d}$  with independent entries satisfying the following three moment conditions as  $n \to \infty$  with  $d = \Theta(n)$ :

- (1)  $\mathbb{E} \mathbf{A}_{ij} = \mathbb{E} \mathbf{B}_{ij} = 0$  for all  $i \in [n]$  and  $j \in [d]$ .
- (2)  $\mathbb{E} A_{ij}^2 = \mathbb{E} B_{ij}^2 = 1$  for all  $i \in [n]$  and  $j \in [d]$ .
- (3)  $\max_{i \in [n]} \max_{j \in [d]} \mathbb{E} |\mathbf{A}_{ij}|^3 + \mathbb{E} |\mathbf{B}_{ij}|^3 = O(1).$

Let  $S_{\boldsymbol{u}} \subseteq [-L_{\boldsymbol{u}}, L_{\boldsymbol{u}}]^d$  and  $S_{\boldsymbol{v}} \subseteq [-L_{\boldsymbol{v}}, L_{\boldsymbol{v}}]^n$  be measurable sets with  $L_{\boldsymbol{u}}, L_{\boldsymbol{v}} \ge 1$ , and set  $L = L_{\boldsymbol{u}} + L_{\boldsymbol{v}}$ . Let  $\psi : \mathbb{R}^d \times \mathbb{R}^n \to \mathbb{R}$  be a differentiable function with

$$\mathscr{M} = \max_{\boldsymbol{u} \in [-L,+L]^d} \max_{\boldsymbol{v} \in [-L,+L]^n} \|\nabla \psi(\boldsymbol{u},\boldsymbol{v})\|_1.$$

Finally, consider the following random variables indexed by  $u \in S_u$  and  $v \in S_v$ :

$$Q_{\boldsymbol{u},\boldsymbol{v}}^{\boldsymbol{A}} = \frac{1}{n^{3/2}} \langle \boldsymbol{A}\boldsymbol{u},\boldsymbol{v} \rangle + \psi(\boldsymbol{u},\boldsymbol{v}) \quad and \quad Q_{\boldsymbol{u},\boldsymbol{v}}^{\boldsymbol{B}} = \frac{1}{n^{3/2}} \langle \boldsymbol{B}\boldsymbol{u},\boldsymbol{v} \rangle + \psi(\boldsymbol{u},\boldsymbol{v}).$$

Then, for all thresholds  $t \in \mathbb{R}$ , all gaps g > 0, and all  $\omega \ge n$ ,

$$\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}^{\boldsymbol{A}} < t\right] - \Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}^{\boldsymbol{B}} < t+g\right] = O\left(\left(1+\frac{1}{g^3}\right)\left(\frac{\mathscr{M}}{\omega} + \frac{L^2\log^{2/3}(L\omega)}{n^{1/6}}\right)\right).$$

The second result we build on is Theorem 3.2 of [Han24], namely generalized first-order method (GFOM) universality. This result establishes universality for a broad class of iterative methods, called *first-order methods*, involving a random design matrix X. The result ensures that many *pseudo-Lipschitz* summaries of such iterates (see Section 2.4 for a definition of pseudo-Lipschitz) are essentially unchanged upon replacing X with a Gaussian matrix G of the same shape. As discussed in the technical overview of Section 1.2, this will prove useful for our analysis of objective perturbation and output perturbation for logistic regression, as well our DP-SGD results for both robust linear regression and logistic regression.

We have modified the statement slightly from the version that appeared in [Han24] to allow vector-valued iterates and different test functions per coordinate, changes that require only minimal, syntactic changes in the proof. Before stating the GFOM universality theorem, we give a formal definition of general first order methods:

**Definition 2.10.** Given  $\boldsymbol{u}^{(1)}, \ldots, \boldsymbol{u}^{(t)} \in \mathbb{R}^{m \times k}$ , let  $\boldsymbol{u}_i^{(s)} \in \mathbb{R}^k$  be the *i*<sup>th</sup> row of  $\boldsymbol{u}^{(s)}$ . Let  $\boldsymbol{u}^{(1:t)} \in \mathbb{R}^{t \times m \times k}$  be the tensor comprising  $\boldsymbol{u}^{(s)}$  for  $s \in [t]$ . Define  $\boldsymbol{u}_i^{(1:t)} \in \mathbb{R}^{t \times k}$  similarly. Given functions  $F_1, \ldots, F_m : \mathbb{R}^{t \times k} \to \mathbb{R}^k$ , let  $F(\boldsymbol{u}^{(1:t)}) \in \mathbb{R}^{m \times k}$  be the matrix with *i*<sup>th</sup> row  $F_i(\boldsymbol{u}_i^{(1:t)}) \in \mathbb{R}^k$ .

**Definition 2.11.** A general first order method (GFOM) consists of a matrix  $\boldsymbol{X} \in \mathbb{R}^{n \times d}$ , an initialization  $(\boldsymbol{u}^{(0)}, \boldsymbol{v}^{(0)}) \in \mathbb{R}^{d \times k} \times \mathbb{R}^{n \times \ell}$  with  $k, \ell = O(1)$  and  $d = \Theta(n)$ , and functions  $F_{1j}^{(t)}, F_{2j}^{(t)}$ :  $\mathbb{R}^{t \times k} \to \mathbb{R}^k$ ,  $G_{1i}^{(t)} : \mathbb{R}^{t \times \ell} \to \mathbb{R}^\ell$ , and  $G_{2i}^{(t)} : \mathbb{R}^{(t+1) \times \ell} \to \mathbb{R}^\ell$  for all  $t \in \mathbb{N}$ ,  $i \in [n]$ , and  $j \in [d]$ . Its iterates are

$$\boldsymbol{u}^{(t)} = \boldsymbol{X} F_1^{(t)}(\boldsymbol{v}^{(0:t-1)}) + G_1^{(t)}(\boldsymbol{u}^{(0:t-1)}) \in \mathbb{R}^{d \times k},$$
  
$$\boldsymbol{v}^{(t)} = \boldsymbol{X}^\top G_2^{(t)}(\boldsymbol{u}^{(0:t)}) + F_2^{(t)}(\boldsymbol{v}^{(0:t-1)}) \in \mathbb{R}^{n \times \ell}.$$
  
(2.12)

**Theorem 2.13** (GFOM Universality, Theorem 3.2 of [Han24]). Consider random matrices  $A, B \in \mathbb{R}^{n \times d}$  with independent entries satisfying the following conditions as  $n \to \infty$  with  $d = \Theta(n)$ :

- $\mathbb{E} \mathbf{A}_{ij} = \mathbb{E} \mathbf{B}_{ij} = 0$  for all  $i \in [n]$  and  $j \in [d]$ .
- $\mathbb{E} \mathbf{A}_{ij}^2 = \mathbb{E} \mathbf{B}_{ij}^2$  for all  $i \in [n]$  and  $j \in [d]$ .
- $\max_{i \in [n]} \max_{j \in [d]} \|\mathbf{A}_{ij}\|_{\psi_2} + \|\mathbf{B}_{ij}\|_{\psi_2} = O(n^{-1/2}).$

Also consider a GFOM satisfying the following condition for some  $\Lambda \geq 2$ , stated in terms of the Lipschitz norm  $||f||_{\text{Lip}} = \inf\{L \in \mathbb{R} : f \text{ is } L\text{-Lipschitz}\}\$  of a function f:

$$\max_{s \in [t]} \max_{q \in \{1,2\}} \max_{i \in [n]} \max_{j \in [d]} \|F_{qi}^{(t)}\|_{\text{Lip}} + \|G_{qj}^{(t)}\|_{\text{Lip}} + |F_{qi}^{(t)}(0)| + |G_{qj}^{(t)}(0)| \le \Lambda$$

Let  $\boldsymbol{u}^{(t)}(\boldsymbol{X})$  and  $\boldsymbol{v}^{(t)}(\boldsymbol{X})$  denote the output of this GFOM with matrix  $\boldsymbol{X}$ . Then, for any collection of  $\Lambda$ -pseudo-Lipschitz functions  $\psi_{1j} : \mathbb{R}^{t \times k} \to \mathbb{R}$  and  $\psi_{2i} : \mathbb{R}^{t \times \ell} \to \mathbb{R}$  of order p, for any  $q \in \mathbb{N}$ , there exists C > 0 such that

$$\mathbb{E} \left| \frac{1}{d} \sum_{j=1}^{d} \left( \psi_{1j}(\boldsymbol{u}_{j}^{(1:t)}(\boldsymbol{A})) - \psi_{1j}(\boldsymbol{u}_{j}^{(1:t)}(\boldsymbol{B})) \right) \right|^{q} + \mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} \left( \psi_{2i}(\boldsymbol{v}_{i}^{(1:t)}(\boldsymbol{A})) - \psi_{2i}(\boldsymbol{v}_{i}^{(1:t)}(\boldsymbol{B})) \right) \right|^{q} \\
\leq (\Lambda \log(n)(1 + \|\boldsymbol{u}^{(0)}\|_{\infty} + \|\boldsymbol{v}^{(0)}\|_{\infty}))^{Ct^{3}} n^{-1/(Ct^{3})}.$$

Note that in Theorem 2.13, if the initialization  $(\boldsymbol{u}^{(0)}, \boldsymbol{v}^{(0)})$  is random, then the conclusion of the theorem still holds even after conditioning on the value of  $(\boldsymbol{u}^{(0)}, \boldsymbol{v}^{(0)})$ .

#### 2.4 Mathematical Miscellany

**Basic Definitions.** We write vectors, matrices, and higher-order tensors in boldface. Given  $m \in \mathbb{N}$ , we write  $[m] = \{1, 2, \ldots, m\}$ . Given a vector  $\boldsymbol{x} \in \mathbb{R}^n$  and  $p \ge 1$ , we write  $\|\boldsymbol{x}\|_p = (\sum_{i=1}^n |x_i|^p)^{1/p}$  and  $\|\boldsymbol{x}\|_{\infty} = \max_{i \in [n]} |x_i|$  (by Hölder's inequality, if p < q, then  $\|\boldsymbol{x}\|_q \le \|\boldsymbol{x}\|_p \le n^{\frac{1}{p}-\frac{1}{q}}\|\boldsymbol{x}\|_q$ ). Given a matrix  $\boldsymbol{A} \in \mathbb{R}^{n \times m}$ , we define  $\|\boldsymbol{A}\|_{r \to s} = \max_{\|\boldsymbol{x}\|_r \le 1} \|\boldsymbol{A}\boldsymbol{x}\|_s$  and  $\|\boldsymbol{A}\|_p = \|\boldsymbol{A}\|_{p \to p}$ . By default, we let  $\|\boldsymbol{x}\| = \|\boldsymbol{x}\|_2$  and  $\|\boldsymbol{A}\| = \|\boldsymbol{A}\|_2$ . Given  $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^n$ , we write  $\boldsymbol{x} \odot \boldsymbol{y} = (x_1y_1, \ldots, x_ny_n) \in \mathbb{R}^n$  for their entrywise (a.k.a. Hadamard) product. Given a set  $\mathcal{S} \subseteq \mathbb{R}^n$ , a scalar  $c \in \mathbb{R}$  and a vector  $\boldsymbol{x} \in \mathbb{R}^n$ , we write  $\boldsymbol{x} + c \cdot \mathcal{S} = \{\boldsymbol{x} + c\boldsymbol{y} : \boldsymbol{y} \in \mathcal{S}\}$  for the rescaled and translated version of  $\mathcal{S}$ .

**Definition 2.14.**  $f : \mathbb{R}^m \to \mathbb{R}$  is *L*-pseudo-Lipschitz of order k if for all  $a, b \in \mathbb{R}^m$ ,

$$|f(a) - f(b)| \le L(1 + ||a|| + ||b||)^{k-1} ||a - b||.$$

We say f is *pseudo-Lipschitz* if there exist  $L \ge 0$  and  $k \in \mathbb{N}$  for which f is L-pseudo-Lipschitz of order k. We say f is L-Lipschitz if it is L-pseudo-Lipschitz of order 1. Let

$$||f||_{\text{Lip}} = \inf\{L \in \mathbb{R} : f \text{ is } L\text{-Lipschitz}\}.$$

The canonical example of an order-k pseudo-Lipschitz function is  $f(\mathbf{x}) = \|\mathbf{x}\|^k$ . Observe that if f is L-pseudo-Lipschitz of order k, there exists L' > 0 such that  $|f(\mathbf{x})| \leq L'(1 + \|\mathbf{x}\|^k)$ . We will often use the fact that  $(1 + \|\mathbf{a}\| + \|\mathbf{b}\|)^{k-1} \leq (3 \max\{1, \|\mathbf{a}\|, \|\mathbf{b}\|\})^{k-1} \leq O_k(1 + \|\mathbf{a}\|^{k-1} + \|\mathbf{b}\|^{k-1})$ .

**High-Dimensional Probability.** We collect some notation, definitions, and lemmas from probability that we will repeatedly use. For more detail, we refer the reader to [Ver18].

First, we say a sequence of events  $E_n$  holds with high probability (w.h.p.) if  $\Pr[E_n] \ge 1 - n^{-\Omega(1)}$ , or merely asymptotically almost surely (a.a.s.) if  $\Pr[E_n] \ge 1 - o(1)$ . We say a sequence of random variables  $x_1, x_2, \ldots \in \mathbb{R}$  converges in probability to the random variable  $x_0 \in \mathbb{R}$ , written  $x_n \xrightarrow{P} x_0$ , if for all c > 0, we have that a.a.s.,  $|x_n - x_0| \le c$ . Given  $\mathbf{p} \in [0, 1]^n$ , we will denote by Bernoulli( $\mathbf{p}$ ) the distribution over  $\{0, 1\}^n$  with independent, Bernoulli( $p_i$ ) coordinates. We will denote the standard Gaussian PDF and CDF by  $\varphi$  and  $\Phi$ , respectively. Given a vector  $\boldsymbol{\mu} \in \mathbb{R}^n$  and positive semidefinite matrix  $\boldsymbol{\Sigma} \in \mathbb{R}^{n \times n}$ , we will denote the multivariate Gaussian distribution with mean  $\boldsymbol{\mu}$  and covariance  $\boldsymbol{\Sigma}$  by  $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ . The key properties of the standard Gaussian distribution that we will use are its rotational invariance and Gaussian integration by parts, also known as Stein's lemma:

**Lemma 2.15** (Rotational Invariance). If  $z \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$  and  $\mathbf{Q} \in \mathbb{R}^{n \times n}$  is orthogonal  $(\mathbf{Q}^{-1} = \mathbf{Q}^{\top})$ , then  $\mathbf{Q}z \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ , as well.

**Lemma 2.16** (Stein's Lemma). If  $Z \sim \mathcal{N}(0,1)$  and  $g : \mathbb{R} \to \mathbb{R}$  is differentiable, then  $\mathbb{E}[g(Z)Z] = E[g'(Z)]$  if both expectations exist.

We also require the following notion of subgaussianity.

**Definition 2.17** (Subgaussian Norm). The subgaussian norm of a random variable  $X \in \mathbb{R}$  is

$$||X||_{\psi_2} = \inf \left\{ c > 0 \ \Big| \ \mathbb{E}[e^{(X/c)^2}] \le 2 \right\}.$$

We say X is subgaussian if  $||X||_{\psi_2} < \infty$ .

One can check that if  $X \sim \mathcal{N}(0,1)$ , then  $||X||_{\psi_2} = \Theta(1)$ . Moreover, we have have  $||X||_{\psi_2} = O(1)$  if and only if  $\Pr[|X| > t] < 2\exp(-\Omega(t^2))$ . Importantly, the subgaussian norm has the following relationship to projection onto a unit vector:

**Lemma 2.18.** If the random vector  $\mathbf{x} \in \mathbb{R}^d$  has independent, subgaussian components satisfying  $\|x_1\|_{\psi_2}, \ldots, \|x_d\|_{\psi_2} \leq \sigma$  and  $\mathbf{u} \in \mathbb{R}^n$  is independent of  $\mathbf{x}$  with  $\|\mathbf{u}\| = 1$ , then

$$\|\langle \boldsymbol{x}, \boldsymbol{u} \rangle\|_{\psi_2} = O(\sigma).$$

**Convex Analysis.** We now briefly review some concepts from convex analysis that we will frequently use. For the most basic definitions of convexity, as well as more detail on the concepts discussed here, we refer the reader to [Bub15, PB14].

**Definition 2.19** (Subdifferential). The *subdifferential* of a function  $f : \mathbb{R}^n \to \mathbb{R}$  at a point  $x \in \mathbb{R}^n$  is its set of *subgradients* g:

$$\partial f(\boldsymbol{x}) = \{ \boldsymbol{g} \in \mathbb{R}^n : \forall \boldsymbol{y} \in \mathbb{R}^n, f(\boldsymbol{y}) \ge f(\boldsymbol{x}) + \langle \boldsymbol{g}, \boldsymbol{y} - \boldsymbol{x} \rangle \}.$$

Minimizing a (strongly convex) function amounts to finding a point  $\boldsymbol{x}$  with a (nearly) vanishing subgradient  $\boldsymbol{g}$ :

**Lemma 2.20** (Convex Minimization). For any  $f : \mathbb{R}^n \to \mathbb{R}$ , the point  $\mathbf{x} \in \mathbb{R}^n$  is a global minimizer of f if and only if  $\mathbf{0} \in \partial f(\mathbf{x})$ . Relatedly, if f is m-strongly convex and  $\mathbf{g} \in \partial f(\mathbf{x})$ , then

$$f(\boldsymbol{x}) - \inf_{\boldsymbol{y} \in \mathbb{R}^n} f(\boldsymbol{y}) \le rac{1}{2m} \|\boldsymbol{g}\|^2.$$

We shall repeatedly use the minimax theorem, a cornerstone result of game theory.

**Lemma 2.21** (Sion's Minimax Theorem, [Sio58]). Suppose  $S_x \subseteq \mathbb{R}^d$  and  $S_y \subseteq \mathbb{R}^n$  are convex and the function  $f : \mathbb{R}^d \times \mathbb{R}^n \to \mathbb{R}$  satisfies the following two properties:

- The map  $\boldsymbol{x} \mapsto f(\boldsymbol{x}, \boldsymbol{y})$  is a closed, proper, convex function for all  $\boldsymbol{y} \in S_{\boldsymbol{y}}$ .
- The map  $\mathbf{y} \mapsto -f(\mathbf{x}, \mathbf{y})$  is a closed, proper, convex function for all  $\mathbf{x} \in \mathcal{S}_{\mathbf{x}}$ .

If either  $\mathcal{S}_{\boldsymbol{x}}$  or  $\mathcal{S}_{\boldsymbol{y}}$  is compact, then

$$\inf_{\boldsymbol{x}\in\mathcal{S}_{\boldsymbol{x}}}\sup_{\boldsymbol{y}\in\mathcal{S}_{\boldsymbol{y}}}f(\boldsymbol{x},\boldsymbol{y})=\sup_{\boldsymbol{y}\in\mathcal{S}_{\boldsymbol{y}}}\inf_{\boldsymbol{x}\in\mathcal{S}_{\boldsymbol{x}}}f(\boldsymbol{x},\boldsymbol{y})$$

We next introduce a bit of notation that will simplify the statements of our results.

**Definition 2.22** (Moreau Envelope, Proximal Operator). Given  $\tau \ge 0$ , the *Moreau envelope* of a closed, proper, convex function f is

$$e_f(\boldsymbol{x};\tau) = \begin{cases} \min_{\boldsymbol{y} \in \mathbb{R}^n} \left\{ \frac{1}{2\tau} \| \boldsymbol{y} - \boldsymbol{x} \|^2 + f(\boldsymbol{y}) \right\} & \text{if } \tau > 0, \\ f(\boldsymbol{x}) & \text{if } \tau = 0. \end{cases}$$

Its proximal operator is the point  $\boldsymbol{y}$  that minimizes  $e_f(\boldsymbol{x};\tau)$ :

$$\operatorname{prox}_{\tau f}(\boldsymbol{x}) = \operatorname{argmin}_{\boldsymbol{y} \in \mathbb{R}^n} \left\{ \frac{1}{2} \|\boldsymbol{y} - \boldsymbol{x}\|^2 + \tau f(\boldsymbol{y}) \right\}.$$

Here, the subscript  $\tau f$  refers to the product  $\tau \cdot f$ , i.e. the function f scaled by  $\tau$ . In the case that  $\tau = 1$ , we will simply write  $\operatorname{prox}_f(\boldsymbol{x})$ . When  $\tau = 0$ , so that  $\tau f$  vanishes everywhere, the function  $\operatorname{prox}_{\tau f}$  coincides with the identity function, i.e.  $\operatorname{prox}_0(\boldsymbol{x}) = \boldsymbol{x}$ .

Some remarks are in order. The proximal operator can be interpreted as a generalization of the projection operator. Indeed, suppose there is a nonempty, compact, convex set  $C \subseteq \mathbb{R}^d$  for which

$$f(\boldsymbol{x}) = egin{cases} 0 & ext{if } \boldsymbol{x} \in C \ +\infty & ext{if } \boldsymbol{x} \notin C \end{cases}$$

Then, f is a closed, proper, convex function and

$$\operatorname{prox}_f(\boldsymbol{x}) = \operatorname*{argmin}_{\boldsymbol{y} \in C} \| \boldsymbol{y} - \boldsymbol{x} \|.$$

Alternatively, the proximal operator can be interpreted as a *backward* gradient step with step size  $\tau$ . Indeed, if f is differentiable and  $\boldsymbol{y} = \text{prox}_{\tau f}(\boldsymbol{x})$ , then

$$\boldsymbol{y} = \boldsymbol{x} - \tau \nabla f(\boldsymbol{y}).$$

For general f, we have that y is the unique point such that

$$\boldsymbol{y} \in \boldsymbol{x} - \tau \partial f(\boldsymbol{y}),$$

where  $\partial f$  denotes the subgradient (see Definition 2.19). A priori, it is not apparent that this equation has a unique solution for  $\boldsymbol{y}$ , but this follows from the definition of  $\operatorname{prox}_f$  in Definition 2.22 and the strong convexity of  $\frac{1}{2} \|\boldsymbol{y} - \boldsymbol{x}\|^2$ . For comparison, the standard equation for a *forward* gradient step is

$$\boldsymbol{y} = \boldsymbol{x} - \tau \nabla f(\boldsymbol{x})$$

We remark that the proximal operator has a useful relationship to the convex conjugate operation, and that its derivatives can be expressed concisely:

**Definition 2.23** (Convex Conjugate). The *convex conjugate* of an extended real-valued function  $f : \mathbb{R}^n \to \mathbb{R} \cup \{\pm \infty\}$  is the convex function

$$f^{\star}(oldsymbol{y}) = \sup_{oldsymbol{x} \in \mathbb{R}^n} \langle oldsymbol{x}, oldsymbol{y} 
angle - f(oldsymbol{x})$$

If n = 1 and f is convex, then  $f^*$  is also called the Legendre transform of f, and  $(f^*)' = (f')^{-1}$ . Lemma 2.24 (Biconjugation). For a closed, proper, convex function f, we have  $f^{**} = f$ . Lemma 2.25 (Proximal Operator of Conjugate).  $\operatorname{prox}_{f^*}(x) = x - \operatorname{prox}_f(x)$ . Lemma 2.26 (Moreau Envelope Derivatives [SAH19]).

$$\nabla_{\boldsymbol{x}} e_f(\boldsymbol{x};\tau) = \frac{\boldsymbol{x} - \operatorname{prox}_{\tau f}(\boldsymbol{x})}{\tau}, \qquad \nabla_{\tau} e_f(\boldsymbol{x};\tau) = -\frac{1}{2} \left\| \frac{\boldsymbol{x} - \operatorname{prox}_{\tau f}(\boldsymbol{x})}{\tau} \right\|^2$$

#### Algorithm 1 Objective Perturbation

- 1: **input:** design matrix  $\boldsymbol{X} = [\boldsymbol{x}_1 \cdots \boldsymbol{x}_n]^\top \in \mathbb{R}^{n \times d}$  with  $\|\boldsymbol{x}_i\| \leq R$ , response vector  $\boldsymbol{y} \in \mathbb{R}^n$ , loss function  $\ell : \mathbb{R}^d \times \mathbb{R}^{d+1} \to \mathbb{R}$ , regularization strength  $\lambda > 0$ , perturbation strength  $\nu > 0$ .
- 2: Randomly sample the coefficients of the linear perturbation term:

$$\boldsymbol{\xi} \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_d)$$

3: Optimize the regularized and perturbed objective function:

$$\widehat{\boldsymbol{\beta}} = \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^d} \sum_{i=1}^n \ell(\boldsymbol{\beta}; (\boldsymbol{x}_i, y_i)) + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle$$

4: return  $\widehat{\boldsymbol{\beta}}$ 

# **3** Objective Perturbation for Robust Linear Regression

In this section, we precisely characterize the privacy-utility tradeoff for the well-known objective perturbation algorithm [CMS11, KST12, RKW23] when applied to the problem of robust linear regression. The version of the algorithm that we consider is described in Algorithm 1. Before stating the main result of this section (Theorem 3.7), we briefly review the setup for robust linear regression, comment on the assumptions that will be necessary for our utility analysis, and give insight into the meaning of the theorem's conclusion, which is significantly stronger than in the simplified statement Theorem 1.1 that we gave in the introduction.

**Robust Linear Regression Model.** We are given a dataset (X, y) of n samples. In the worst case,  $(x_i, y_i)$  are arbitrary points in  $\mathbb{R}^{d+1}$  with  $||x_i|| \leq R$ , but in the average case, there exists a ground-truth coefficient vector  $\beta^* \in \mathbb{R}^d$  and regression error vector  $\varepsilon^* \in \mathbb{R}^n$  independent of X such that

$$y = X \beta^{\star} + \varepsilon^{\star}.$$

To dampen the effect of outliers in the worst-case setting, we do not seek the least squares estimate, but rather a vector  $\hat{\beta}$  that minimizes the average *Huber loss* (robust linear loss) on the dataset:

**Definition 3.1.** The *L*-Lipschitz Huber loss is

$$H_L(r) = \begin{cases} \frac{1}{2}r^2 & \text{if } |r| \le L\\ L|r| - \frac{1}{2}L^2 & \text{if } |r| \ge L \end{cases}$$

We denote the *L*-truncation of  $r \in \mathbb{R}$  by

$$[r]_{L} = H'_{L}(r) = \begin{cases} -L & \text{if } r \leq -L, \\ r & \text{if } r \in [-L, +L], \\ +L & \text{if } r \geq L. \end{cases}$$

For  $\boldsymbol{r} \in \mathbb{R}^n$ , we define  $H_L(\boldsymbol{r}) = \sum_{i=1}^n H_L(r) \in \mathbb{R}$  and  $[\boldsymbol{r}]_L = \nabla H_L(\boldsymbol{r}) = ([r_1]_L, \dots, [r_n]_L) \in \mathbb{R}^n$ .

Assumptions. The main assumption for our utility analysis will be that the matrix X, which we call the *design matrix* or *feature matrix*, is subgaussian in the sense defined below.

**Definition 3.2** (Subgaussian Design). We say a random matrix  $X \in B_R(\mathbf{0})^n$  with  $d = \Theta(n)$  follows a subgaussian design if its entries  $x_{ij}$  are independent and satisfy the following conditions.

- (a) For all  $i \in [n]$  and  $j \in [d]$ , we have  $\mathbb{E}[x_{ij}] = 0$  and  $\mathbb{E}(x_{ij})^2 = 1/d$ .
- (b)  $\max_{i \in [n]} \max_{j \in [d]} ||x_{ij}||_{\psi_2} = O(1/\sqrt{n}).$

Note that any distribution supported on  $\left[-\frac{R}{\sqrt{d}}, +\frac{R}{\sqrt{d}}\right]^{n \times d} \subseteq B_R(\mathbf{0})^n$  with independent, variance 1/d entries automatically satisfies the conditions of Definition 3.2. For simplicity, we have required that the variance per coordinate is precisely 1/d, but this is merely a scaling convention, and our results extend in a straightforward manner to other isotropic distributions. In fact, given a recent line of work analyzing estimators in the proportional regime for design matrices with nonisotropic covariances or otherwise dependent entries [ZSC22, Fan22, BHX23, GTM<sup>+</sup>24], we suspect that the result of Theorem 3.7 can be further extended to nonisotropic X.

More interestingly, although we do impose some mild conditions on the ground-truth  $\beta^*$  and the regression errors  $\varepsilon^*$ , we do *not* require them to be random! This is important because several previous works indeed assume that  $\beta^*$  and  $\varepsilon^*$  are random with i.i.d. coordinates (e.g. [SC19, SAH19], or Theorem 3.12 of [HS23]). We will clarify the conditions on  $\beta^*$  and  $\varepsilon^*$  shortly once we have defined a certain notion of convergence.

Notion of Convergence. Theorem 3.7 will provide estimates for a large collection of scalar quantities that one might wish to compute on the coefficient vector output by the objective perturbation algorithm, along with non-asymptotic error bounds for those estimates that hold with high probability and decay at a  $n^{-\Omega(1)}$  rate. In order to state the theorem as concisely as possible, we introduce the notation  $\rightarrow$  to describe this type of convergence.<sup>3</sup>

**Definition 3.3** (Polynomial-Rate Pseudo-Lipschitz Convergence). Consider a random variable  $x_0 \in \mathbb{R}$  with finite moments of all orders and a vector  $\boldsymbol{x} \in \mathbb{R}^n$ . We write  $\boldsymbol{x} \rightsquigarrow x_0$ , if for all pseudo-Lipschitz functions  $f : \mathbb{R} \to \mathbb{R}$ ,

$$\exists c > 0 \quad \text{such that} \quad \Pr\left[\left|\frac{1}{n}\sum_{i=1}^{n}f(x_i) - \mathbb{E}[f(x_0)]\right| \le n^{-c}\right] \ge 1 - O(n^{-c}). \tag{3.4}$$

For  $\boldsymbol{x} \in \mathbb{R}^m$  with  $m = \Theta(n)$ , we write  $\boldsymbol{x} \rightsquigarrow x_0$  if (3.4) holds with m in place of n. Similarly, if  $\boldsymbol{x}_0 \in \mathbb{R}^{\ell}$  and  $\boldsymbol{X} = [\boldsymbol{x}_1 \cdots \boldsymbol{x}_m]^{\top} \in \mathbb{R}^{m \times \ell}$  for  $\ell = \Theta(1)$ , we write  $\boldsymbol{X} \rightsquigarrow \boldsymbol{x}_0$  if (3.4) holds for all pseudo-Lipschitz  $f : \mathbb{R}^{\ell} \to \mathbb{R}$  with m in place of n.

Observe that Definition 3.3 can be applied either to a sequence of deterministic vectors  $\boldsymbol{x} \in \mathbb{R}^m$ , or a sequence of random vectors  $\boldsymbol{x} \in \mathbb{R}^m$ . In the case of deterministic vectors, the condition that the absolute difference is bounded by  $n^{-c}$  with probability  $1 - O(n^{-c})$  becomes equivalent to the condition that the absolute difference is deterministically bounded by  $n^{-c}$ . Next, observe that Definition 3.3 is a strengthening of the familiar notion of *convergence in distribution*, applied to the empirical distribution with mass 1/m on each of the coordinates of  $\boldsymbol{x}$ . Convergence in distribution is weaker than Definition 3.3 because the former only requires (3.4) to hold for functions f that are

<sup>&</sup>lt;sup>3</sup>Although the  $\rightarrow$  abbreviation is our own, studying convergence with respect to pseudo-Lipschitz tests is certainly not new; it is considered, for example, in [BM11a, CMW20, Han24], among many other works.

both bounded and Lipschitz. Also, convergence in distribution only requires an o(1) convergence rate, but Definition 3.3 requires a much stronger  $n^{-\Omega(1)}$  convergence rate. The following lemmas provide intuition for the meaning of  $\mathbf{x} \rightsquigarrow x_0$ . We defer their proofs to Section 3.3.

**Lemma 3.5.** If  $x_0 \in \mathbb{R}$  has finite moments and  $x_1, \ldots, x_m \stackrel{iid}{\sim} x_0$ , then  $(x_1, \ldots, x_m) \rightsquigarrow x_0$ . **Lemma 3.6.** If  $x_0 \in \mathbb{R}$  has finite moments and  $\mathbf{x} \rightsquigarrow x_0$ , then for all c > 0, w.h.p.,  $\|\mathbf{x}\|_{\infty} \leq O(n^c)$ .<sup>4</sup>

In the following theorem, we will phrase not only the conclusion in terms of pseudo-Lipschitz convergence, but also the assumptions on  $\beta^*$  and  $\varepsilon^*$ . For a somewhat contrived example demonstrating the breadth of  $\beta^*$  to which our theorem can be applied, suppose the first 50% of the coordinates of  $\beta^*$  are deterministically equal to 0, the next 20% of the coordinates of  $\beta^*$  are sampled i.i.d. from  $\mathcal{N}(-3, 16)$ , and the final 30% deterministically comprise an arithmetic progression starting at -10 and ending at 10. In this example, we have that  $\beta^* \rightsquigarrow \beta_0^*$  a.s., where  $\beta_0^* \in \mathbb{R}$ follows a mixture distribution of a point mass at 0, the distribution  $\mathcal{N}(-3, 16)$ , and the uniform distribution on [-10, 10]. To verify the condition (3.4) with respect to, say,  $f(t) = t^2$ , one can check that w.h.p.,  $\frac{1}{d} \|\beta^*\|^2 = \mathbb{E}(\beta_0^*)^2 \pm O(n^{-1/2})$ , where

$$\mathbb{E}(\beta_0^{\star})^2 = 50\% \cdot 0 + 20\% \cdot (3^2 + 16) + 30\% \cdot \frac{1}{12}(10 - (-10))^2 = 15.$$

For an example of  $\beta^* \in \mathbb{R}^d$  that our theorem does *not* handle, consider  $\beta^* = (\sqrt{d}, 0, \dots, 0)$ . On the one hand, the empirical distribution of the coordinates of this  $\beta^*$  converges in distribution to the point mass at **0**, so if  $\beta^*$  were to converge with respect to pseudo-Lipschitz tests, it would also have to be toward the point mass at **0**. On the other hand, we have  $\frac{1}{d} \|\beta^*\|^2 = 1 \neq 0 \pm n^{-\Omega(1)}$ , so  $\beta^*$  fails to satisfy the condition (3.4) for  $f(t) = t^2$ .

We are now ready to state and prove the main theorem of this section.

**Theorem 3.7.** Let  $\hat{\beta}$  denote the output of Algorithm 1 with parameters  $R, \lambda, \nu > 0$  and instantiated with the L-Lipschitz Huber loss function:

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = H_L(y - \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle).$$

(a) (Privacy)  $\hat{\beta}$  satisfies  $(2\varepsilon_{\rm DP}, (1 + e^{\varepsilon_{\rm DP}})\delta_{\rm DP})$ -differential privacy for any  $\varepsilon_{\rm DP} \ge 0$  and

$$\delta_{\rm DP} = \begin{cases} 2 \cdot \text{HockeyStick}(\tilde{\varepsilon}_{\rm DP}, \frac{LR}{\nu}) & \text{if } \hat{\varepsilon}_{\rm DP} \ge 0, \\ (1 - e^{\hat{\varepsilon}_{\rm DP}}) + 2e^{\hat{\varepsilon}_{\rm DP}} \cdot \text{HockeyStick}\left(\frac{L^2R^2}{2\nu^2}, \frac{LR}{\nu}\right) & \text{otherwise,} \end{cases}$$

where we set  $\tilde{\varepsilon}_{\rm DP} = \varepsilon_{\rm DP} - \log(1 + R^2/\lambda)$  and  $\hat{\varepsilon}_{\rm DP} = \tilde{\varepsilon}_{\rm DP} - L^2 R^2/2\nu^2$ .

- (b) (Utility) Suppose the following hold for some  $\beta^* \in \mathbb{R}^d$  and  $\varepsilon^* \in \mathbb{R}^n$  as  $n \to \infty$  and  $d/n \to \delta$ :
  - (i)  $\mathbf{X} \in B_R(\mathbf{0})^n \subseteq \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y} = \mathbf{X} \boldsymbol{\beta}^* + \boldsymbol{\varepsilon}^*$ .
  - (ii) There exist random variables  $\beta_0^{\star}, \varepsilon_0^{\star} \in \mathbb{R}$  such that  $\beta^{\star} \rightsquigarrow \beta_0^{\star}$  and  $\varepsilon^{\star} \rightsquigarrow \varepsilon_0^{\star}$ .

Suppose there exist  $\sigma^*, \tau^* > 0$  solving the following system of two scalar equations in two variables  $(\sigma, \tau)$ , which we write in terms of a dummy variable  $Z \sim \mathcal{N}(0, 1)$  and  $\kappa^2 = \mathbb{E}(\beta_0^*)^2$  as

$$\sigma^2 = \tau^2 \left( \frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma Z + \varepsilon_0^{\star}}{1 + \tau} \right]_L^2 + \lambda^2 \kappa^2 + \nu^2 \right), \tag{3.8a}$$

$$\tau = \frac{1}{\lambda\delta} \left( \delta - \frac{\tau}{1+\tau} \Pr\left[ -L < \frac{\sigma Z + \varepsilon_0^*}{1+\tau} < L \right] \right).$$
(3.8b)

<sup>4</sup>Moreover, if  $\boldsymbol{x} \in \mathbb{R}^m$  is deterministic and  $\boldsymbol{x} \rightsquigarrow x_0$ , then  $\|\boldsymbol{x}\|_{\infty} \leq n^{o(1)}$  holds deterministically.

Then, in terms of  $(\sigma^*, \tau^*)$ , the estimation error  $\widehat{\beta} - \beta^*$  satisfies, for  $\xi_0 \sim \mathcal{N}(0, 1)$ ,

$$(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \, \widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}) \rightsquigarrow \left( \beta_{0}^{\star}, \, \xi_{0}, \, \tau^{\star} \left( \sqrt{\frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma^{\star} Z + \varepsilon_{0}^{\star}}{1 + \tau^{\star}} \right]_{L}^{2}} Z - \lambda \beta_{0}^{\star} - \nu \xi_{0} \right) \right).$$

Similarly, the L-truncated prediction error  $[\boldsymbol{y} - \boldsymbol{X}\widehat{\boldsymbol{eta}}]_L$  satisfies

$$(\boldsymbol{\varepsilon}^{\star}, [\boldsymbol{y} - \boldsymbol{X}\widehat{\boldsymbol{\beta}}]_L) \rightsquigarrow \left(\varepsilon_0^{\star}, \left[\frac{\sigma^{\star} Z + \varepsilon_0^{\star}}{1 + \tau^{\star}}\right]_L\right).$$

Several remarks about Theorem 3.7 are in order. First, we comment that part (a), whose proof we defer to Section 5, is a variant of Theorem 3.1 [RKW23] (for convenience, we included a statement of that prior result in Section 2). Essentially, part (a) establishes a link between any constant values of  $L, \lambda, \nu > 0$  and a family of  $(\varepsilon_{\text{DP}}, \delta_{\text{DP}})$ -DP privacy guarantees. In Section 5, we also give a single-parameter  $\rho_{\text{DP}}$ -zCDP privacy guarantee with a constant  $\rho_{\text{DP}} = \rho_{\text{DP}}(L, \lambda, \nu)$ , which is arguably easier to interpret but not as tight.

Next, we comment that part (b), whose proof will comprise the bulk of this section, yields several interesting corollaries about the behavior of  $\hat{\beta}$  when combined with Definition 3.3. For example, suppose we would like to know the bias of  $\hat{\beta}$ . If we take the order-2 pseudo-Lipschitz function  $f(\beta^*, u) = (\beta^*)^2 + \beta^* u$ , then Theorem 3.7(b) with Definition 3.3 tells us that the average value of f over the coordinates of  $(\beta^*, \hat{\beta} - \beta^*)$  can be approximated by the expected value of fover the randomness of  $(\beta_0^*, \tau^*(\sigma_v^* Z - \lambda \beta_0^* - \nu \xi_0))$  for an appropriate constant  $\sigma_v^* > 0$ . Since  $Z, \beta_0^*$ , and  $\xi_0$  are independent, we deduce that with high probability over the design matrix X and the randomness  $\boldsymbol{\xi}$  of the algorithm,

$$\frac{1}{d} \langle \widehat{\boldsymbol{\beta}}, \boldsymbol{\beta}^{\star} \rangle = \frac{1}{d} \sum_{j=1}^{d} f(\beta_{j}^{\star}, \widehat{\beta}_{j} - \beta_{j}^{\star}) = \mathbb{E}[(\beta_{0}^{\star})^{2} + \beta_{0}^{\star} \cdot \tau^{\star} (\sigma_{v}^{\star} Z - \lambda \beta_{0}^{\star} - \nu \xi_{0})] \pm n^{-\Omega(1)}$$
$$= (1 - \tau^{\star} \lambda) \kappa^{2} \pm n^{-\Omega(1)}.$$

It is apparent from equation (3.8b) that  $1 - \tau^* \lambda \in (0, 1)$ . Therefore, we deduce that  $\widehat{\boldsymbol{\beta}}$  is positively correlated with  $\boldsymbol{\beta}^*$ , but "shrunk" toward **0** by a factor of  $1 - \tau^* \lambda \pm n^{-\Omega(1)}$ . This is intuitively what we would expect given that the regularization term  $\frac{\lambda}{2} \|\boldsymbol{\beta}\|^2$  is centered at the origin.

Similarly, suppose we are interested in the mean squared error of  $\hat{\beta}$ . If we take the order-2 pseudo-Lipschitz function  $f(u) = u^2$ , then by Theorem 3.7(b) and Definition 3.3, with high probability,

$$\frac{1}{d} \|\widehat{\beta} - \beta^{\star}\|^{2} = \frac{1}{d} \sum_{j=1}^{d} f(\widehat{\beta}_{j} - \beta_{j}^{\star}) = \mathbb{E}(\tau^{\star}(\sigma_{v}^{\star}Z - \lambda\beta_{0}^{\star} - \nu\xi_{0}))^{2} \pm n^{-\Omega(1)}$$
$$= (\tau^{\star})^{2}((\sigma_{v}^{\star})^{2} + \lambda^{2}\kappa^{2} + \nu^{2}) \pm n^{-\Omega(1)}.$$

By equation (3.8a), the above expression is precisely  $(\sigma^*)^2 \pm n^{-\Omega(1)}$ , so we deduce that the root mean squared error of  $\beta^*$  is precisely  $\sigma^* \pm n^{-\Omega(1)}$ . Similar computations can be carried out in this manner to determine, for example, the correlation of  $\hat{\beta}$  with the perturbation vector  $\boldsymbol{\xi}$  or the fact that the residual error of the predictions  $\boldsymbol{X}\hat{\boldsymbol{\beta}}$  satisfies, w.h.p.,

$$\frac{1}{n} \left\| [\boldsymbol{y} - \boldsymbol{X} \widehat{\boldsymbol{\beta}}]_L \right\|^2 = \mathbb{E} \left[ \frac{\sigma^* Z + \varepsilon_0^*}{1 + \tau^*} \right]_L^2 \pm n^{-\Omega(1)}.$$



Figure 3: Theorem 3.7's predictions of the estimation error and truncated residuals of Algorithm 1 with Huber loss. Curves correspond to theoretical predictions, and dots correspond to the mean over 100 simulations of the algorithm on synthetic data with  $n \times d = 1000$ . In the left plots, the perturbation magnitude is  $\nu = 0$ , corresponding to the non-private case, but in the right plots,  $\nu = 1/5$ . In all plots, the Huber loss' Lipschitz constant is L = 10, the signal strength is  $\kappa = 1$ , and we consider  $\beta^* \sim \mathcal{N}(\mathbf{0}, \kappa^2 \mathbf{I}_d)$  and  $\varepsilon^* \sim \mathcal{N}(\mathbf{0}, (1/5)^2 \mathbf{I}_n)$ , along with  $\mathbf{X} \sim \frac{1}{\sqrt{d}} \text{Uniform}(\{-1, +1\}^{n \times d})$  and  $\mathbf{y} = \mathbf{X}\beta^* + \varepsilon^*$ .

The previous two examples both measure utility using squared  $\ell^2$  norm, but this was arbitrary. Indeed, by choosing different pseudo-Lipschitz test functions, we could just as easily have expressed many other possible utility metrics, including  $\frac{1}{d} \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|_{p}^{p}$  or  $\frac{1}{n} \|[\boldsymbol{y} - \boldsymbol{X}\widehat{\boldsymbol{\beta}}]_{L}\|_{p}^{p}$  for any  $p \geq 1$ , in terms of  $(\sigma^{\star}, \tau^{\star})$ . We validate these corollaries against synthetic data in Figure 3.

Next, we remark that equations (3.8a), (3.8b) can be simplified if one assumes approximately Gaussian errors. Indeed, if  $\varepsilon_0^* \sim \mathcal{N}(0, \sigma_{\varepsilon}^2)$  for some  $\sigma_{\varepsilon} > 0$ , then we have

$$\frac{\sigma Z + \varepsilon_0^*}{1 + \tau} \sim \mathcal{N}\left(0, \frac{\sigma^2 + \sigma_\varepsilon^2}{(1 + \tau)^2}\right).$$

Thus, we can express the variance of its *L*-truncation, as well as the probability it falls in the interval [-L, +L], using rescaled versions of the following formulas for the truncated Gaussian distribution: if  $Z \sim \mathcal{N}(0, 1)$  and s > 0, then in terms of the standard Gaussian PDF  $\varphi$  and CDF  $\Phi$ ,

$$\Pr[sZ \in [-L, +L]] = 2\Phi(L/s) - 1.$$
$$\mathbb{E}[(sZ)^2 \mid sZ \in [-L, +L]] = s^2 \left(1 - \frac{2(L/s)\varphi(L/s)}{2\Phi(L/s) - 1}\right).$$

**Simulations.** Before proceeding to the proof of Theorem 3.7, we include a non-rigorous and informal discussion of Figure 3, which validates the predictions of Theorem 3.7 against random, synthetic data. From a quantitative perspective, comparing the predicted error in the non-private case ( $\nu = 0$ ) to the differentially private case ( $\nu > 0$ ) allows us to make the first ever measurements of the *price of privacy* of objective perturbation in the proportional dimensionality regime.

However, there are also several interesting qualitative takeaways from Figure 3. For example, consider the plots on estimation error, which is closely related to the algorithm's test loss, or ability to generalize to out-of-sample data. In these plots, we see what appears to be a change in behavior around  $d/n \approx 1$  (equivalently, around  $n/(n+d) \approx 1/2$ ). On one side of this threshold, where n > d, the error of the objective perturbation algorithm is only mildly worse than in the non-private case, but on the other side, where n < d, there appears to be a dramatic deterioration, especially as  $\lambda \to 0$ . Interestingly, the "spike" in estimation error at n = d in the non-private case, closely related to the famous *double descent* phenomenon in modern machine learning, seems to go away in the differentially private case, or at least be of less importance. (Whether or not a "spike" occurs above the range of vertical axis values visualized in Figure 3 in somewhat irrelevant, in the sense that any estimator with error  $\frac{1}{d} \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^*\|^2$  above  $\kappa^2 = 1$  is already worse than the trivial estimator  $\hat{\boldsymbol{\beta}}_{\text{trivial}} = \mathbf{0}$ .)

On the other hand, we can tell a different story by looking at the clipped residuals of the algorithm's predictions on X, which is essentially a notion of training loss, or in-sample accuracy. Once again, there appears to be an interesting change in behavior around  $d/n \approx 1$ . In the non-private case, this is called the interpolation threshold, so as expected, on the side where n < d, the residual error is 0 as  $\lambda \to 0$ , but on the side where n > d, the residual error assumes a small positive value, even as  $\lambda \to 0$ . Strikingly, in the differentially private case, we observe a spike in residual error is, by definition, capped at  $L^2 = 100$ , but is nevertheless qualitatively similar to what we would expect of *test* loss, not training loss, in the non-private case.

One possible intuitive explanation for why differential privacy creates a spike in the training loss around n = d draws on a line of work on *adaptive data analysis* introduced by [DFH<sup>+</sup>15, BNS<sup>+</sup>16]. This work rigorously showed that any differentially private learning algorithm satisfying differential privacy *automatically* enjoys a certain resilience against overfitting and false discovery. With this in mind, it arguably makes sense that a training loss-related metric, when applied to a differentially private algorithm, might exhibit qualitative behaviors traditionally seen only when analyzing test loss.

# 3.1 Special Case of the Utility Proof

Before we begin the full utility proof, we will first prove the following special case. This special case assumes not only that we are given average-case features X, but also that the ground-truth coefficient vector  $\beta^*$  and error vector  $\varepsilon^*$  have i.i.d. coordinates. The conclusion of this special case consists of a single scalar, the limit of  $\frac{1}{d} || \hat{\beta} - \beta^* ||^2$ . Eventually, for the full proof of Theorem 3.7(b), we will remove these unnecessary i.i.d. assumptions on  $\beta^*$  and  $\varepsilon^*$  and determine the limit of both the estimation error  $\hat{\beta} - \beta^*$  and prediction error  $[\boldsymbol{y} - \boldsymbol{X}\hat{\beta}]_L$  with respect to all pseudo-Lipschitz test functions. However, we choose to first emphasize this special case because of its exceedingly simple proof given prior results in the high-dimensional statistics literature on non-private robust linear regression.

**Theorem 3.9.** Consider the setting of Theorem 3.7(b), but also assume that  $\beta_1^{\star}, \ldots, \beta_d^{\star} \stackrel{iid}{\sim} \beta_0^{\star}$  and  $\varepsilon_1^{\star}, \ldots, \varepsilon_n^{\star} \stackrel{iid}{\sim} \varepsilon_0^{\star}$  and that  $\varepsilon_0^{\star}$  is continuous. Define  $\sigma^{\star}$  as in Theorem 3.7. Then, as  $n \to \infty$ ,

$$\frac{1}{d} \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2 \xrightarrow{P} (\sigma^{\star})^2.$$

The i.i.d. assumption on  $\beta^*$  and  $\varepsilon^*$  is stronger than assuming  $\beta^* \rightsquigarrow \beta_0^*$  and  $\varepsilon^* \rightsquigarrow \varepsilon_0^*$ , which we saw in Lemma 3.5. Similarly, the conclusion of Theorem 3.9 is weaker than that of Theorem 3.7.

The idea behind Theorem 3.9 is a simple reduction to the non-private case. Indeed, the asymptotic behavior in the non-private case was derived by [TAH18] for Gaussian X and extended by [HS23] to subgaussian designs.

Proof of Theorem 3.9. In the absence of perturbation, which is the case when  $\nu = 0$ , Theorem 3.12 of [HS23] implies that under the stated assumptions, the scaled error of  $\hat{\beta}$  converges to the solution  $\sigma^*$  to the following system of two scalar equations in two variables  $(\sigma, \tau)$ . In terms of a dummy variable  $Z \sim \mathcal{N}(0, 1)$ , these equations are

$$\begin{split} \frac{1}{\delta} &-1 + \tau \lambda = \frac{1}{\delta} \mathbb{E}[\operatorname{prox}_{\tau H_L}'(\sigma Z + \varepsilon_0^{\star})], \\ &\sigma^2 = \frac{1}{\delta} \mathbb{E}[(\sigma Z + E - \operatorname{prox}_{\tau H_L}(\sigma Z + \varepsilon_0^{\star}))^2] + \lambda^2 \tau^2 \kappa^2. \end{split}$$

A simple calculation using the definition of the proximal operator (Definition 2.22) shows that for any  $\tau > 0$  and  $s \in \mathbb{R}$ ,

$$s - \operatorname{prox}_{\tau H_L}(s) = \tau \left[ \frac{s}{1+\tau} \right]_L$$

If  $|s| \neq (1 + \tau)L$ , then a similarly simple calculation shows that

$$1 - \operatorname{prox}_{\tau H_L}'(s) = \begin{cases} \frac{\tau}{1+\tau} & \text{if } |s| < (1+\tau)L, \\ 0 & \text{if } |s| > (1+\tau)L. \end{cases}$$

Substituting these formulas for  $s - \operatorname{prox}_{\tau H_L}(s)$  and  $1 - \operatorname{prox}'_{\tau H_L}(s)$  with  $s = \sigma Z + \varepsilon_0^*$  into the above system of equations yields the following simplification:

$$\sigma^{2} = \tau^{2} \left( \frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma Z + \varepsilon_{0}^{\star}}{1 + \tau} \right]_{L}^{2} + \lambda^{2} \kappa^{2} \right),$$

$$\tau = \frac{1}{\lambda \delta} \left( \delta - \frac{\tau}{1 + \tau} \Pr \left[ -L < \frac{\sigma Z + \varepsilon_{0}^{\star}}{1 + \tau} < L \right] \right).$$
(3.10)

This proves Theorem 3.9 in the non-private setting, where  $\nu = 0$ . To extend this result to the private setting, where  $\nu > 0$ , we carry out a straightforward reduction. To begin, we first rewrite the perturbed objective in Algorithm 1 by completing the square to combine the quadratic regularization term with the linear perturbation term. Indeed,

$$\widehat{oldsymbol{eta}} = \operatorname*{argmin}_{oldsymbol{eta} \in \mathbb{R}^d} H_L(oldsymbol{y} - oldsymbol{X}oldsymbol{eta}) + rac{\lambda}{2} \Big\| oldsymbol{eta} + rac{
u}{\lambda} oldsymbol{\xi} \Big\|^2.$$

Introducing the notation  $u = \beta - \beta^*$  and  $\hat{u} = \hat{\beta} - \beta^*$ , and recalling that  $\varepsilon^* = y - X\beta^*$ , this becomes

$$\widehat{oldsymbol{u}} = rgmin_{oldsymbol{u}\in\mathbb{R}^d} H_L(oldsymbol{arepsilon}^\star - oldsymbol{X}oldsymbol{u}) + rac{\lambda}{2} \Big\|oldsymbol{u} + \Big(eta^\star + rac{
u}{\lambda}oldsymbol{\xi}\Big)\Big\|^2$$

In this formulation, since X,  $\beta^*$ , and  $\varepsilon^*$  are independent, it is clear that  $\beta^*$  only influences the final, quadratic term of the optimization objective. Consequently,  $\hat{u}$  has the same value in the case of  $\nu > 0$  when the ground-truth vector is  $\beta^*$  as in the non-private case when ground-truth vector is

$$\boldsymbol{\beta}^{\star} + \frac{\nu}{\lambda} \boldsymbol{\xi}.$$

In equation (3.10), which we have already shown characterizes the limiting behavior in the nonprivate case, this amounts to defining  $\xi_0 \sim \mathcal{N}(0, 1)$  and replacing each occurrence of  $\mathbb{E}(\beta_0^*)^2 = \kappa^2$  with

$$\mathbb{E}\left(\beta_0^{\star} + \frac{\nu}{\lambda}\xi_0\right)^2 = \kappa^2 + \left(\frac{\nu}{\lambda}\right)^2$$

Doing so yields the system of equations from the statement of Theorem 3.7, which read

$$\sigma^{2} = \tau^{2} \left( \frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma Z + E}{1 + \tau} \right]_{L}^{2} + \lambda^{2} \kappa^{2} + \nu^{2} \right),$$
  
$$\tau = \frac{1}{\lambda \delta} \left( \delta - \frac{\tau}{1 + \tau} \Pr \left[ -L < \frac{\sigma Z + E}{1 + \tau} < L \right] \right).$$

Letting  $\sigma^*$  be the solution for  $\sigma$ , we conclude that

$$\frac{1}{\sqrt{d}} \|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\| \xrightarrow{P} \sigma^{\star}.$$

#### 3.2 Proof of Theorem 3.7

To prove Theorem 3.7 in its full generality, we can no longer apply a black-box reduction to prior work in the non-private case since this work assumes that  $\beta^*$  and  $\varepsilon^*$  are i.i.d. and does not fully determine the asymptotic behavior of  $\hat{\beta} - \beta^*$  and  $[\boldsymbol{y} - \boldsymbol{X}\hat{\beta}]_L$ . Instead, we will take a white-box approach, roughly following the proof strategy of [TAH18, HS23] and making several changes as necessary to accommodate the differences between their setting and ours. To reiterate, the main differences are the random linear perturbation term for differential privacy, our weakened assumptions on  $\beta^*$  and  $\varepsilon^*$ , and our conclusion that includes a description of  $[\boldsymbol{y} - \boldsymbol{X}\hat{\beta}]_L$  and involves a stronger notion of convergence. Consequently, the present proof is substantially more involved than the preceding one.

**Proof Overview.** Our proof proceeds in three steps, inspired by [HS23]:

- (1) Relate the perturbed Huber objective to a certain min-max optimization problem.
- (2) Use universality to relate this problem to another involving a Gaussian design matrix.
- (3) Use CGMT to analyze this min-max optimization with a Gaussian design matrix as  $n \to \infty$ .

For the first step, we will take the *Legendre transform* (a.k.a. convex conjugate) of the Huber loss function, which in our case will have the effect of isolating the randomness of  $\mathbf{X} \in \mathbb{R}^{n \times d}$  in a single, bilinear term.

For the second step, we will make critical use of Corollary 2.6 of [HS23], a powerful result that the authors call universality of the Gordon's max-min (min-max) cost optimum. The name "Gordon" comes from Gordon's inequality, a precursor to the convex Gaussian minimax theorem (CGMT) that pertains to the same families of Gaussian random variables and their extrema. For convenience, we have included a simplified rephrasing of this result, which we refer to as CGMT universality, as Theorem 2.9.

For the third step, we will use CGMT, roughly following the template in [TAH18], to determine the behavior of the estimation error  $\hat{\beta} - \beta^*$  as  $n \to \infty$ . Unlike [TAH18], our more detailed analysis will account for the random linear perturbation term, establish convergence in a stronger sense, and also explain the behavior of the residual error,  $y - X\hat{\beta}$ . For a statement of CGMT, see Theorem 2.8.

We remark that both steps 2 and 3 will be made more challenging by the fact that we do not assume that  $\beta^*$  and  $\varepsilon^*$  have i.i.d. coordinates.

#### 3.2.1 Step 1: Legendre Transform

The goal of this step is to prove the following lemma, which relates the output of Algorithm 1 with L-Lipschitz Huber loss to the min-max optimization of a certain random variable. Technically, the lemma holds in a worst-case sense, as neither the randomness of X and  $\xi$  nor any assumptions on  $\beta^*$  or  $\varepsilon^*$  are required for the proof. Eventually, the randomness of the optimization problem will arise from the randomness of the design matrix X.

**Lemma 3.11.** Let  $\hat{\beta}$  be the output of Algorithm 1 when instantiated with

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = H_L(y - \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle).$$

Fix  $\boldsymbol{\beta}^{\star} \in \mathbb{R}^d$  and define  $\boldsymbol{\varepsilon}^{\star} = \boldsymbol{y} - \boldsymbol{X} \boldsymbol{\beta}^{\star}$ . Given  $\boldsymbol{u} \in \mathbb{R}^d$  and  $\boldsymbol{v} \in [-L, +L]^n$ , define  $Q_{\boldsymbol{u}, \boldsymbol{v}}$  as:

$$Q_{\boldsymbol{u},\boldsymbol{v}} = -\langle \boldsymbol{X}\boldsymbol{u},\boldsymbol{v}\rangle + \psi(\boldsymbol{u},\boldsymbol{v}) \qquad for \qquad \psi(\boldsymbol{u},\boldsymbol{v}) = \frac{\lambda}{2} \|\boldsymbol{u}\|^2 + \langle \lambda \boldsymbol{\beta}^* + \nu \boldsymbol{\xi}, \boldsymbol{u} \rangle - \frac{1}{2} \|\boldsymbol{v}\|^2 + \langle \boldsymbol{\varepsilon}^*, \boldsymbol{v} \rangle.$$

Then,  $(\widehat{\boldsymbol{u}}, \widehat{\boldsymbol{v}}) = (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}, [\boldsymbol{y} - \boldsymbol{X}\widehat{\boldsymbol{\beta}}]_L)$  is the unique point in  $\mathbb{R}^d \times [-L, +L]^n$  satisfying

$$\max_{\boldsymbol{v}\in[-L,+L]^n} Q_{\widehat{\boldsymbol{u}},\boldsymbol{v}} = \min_{\boldsymbol{u}\in\mathbb{R}^d} Q_{\boldsymbol{u},\widehat{\boldsymbol{v}}}.$$

We call it the saddle point or Nash equilibrium of  $Q_{u,v}$ .

*Proof.* Algorithm 1 simply computes

$$\widehat{\boldsymbol{\beta}} = \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^d} H_L(\boldsymbol{y} - \boldsymbol{X}\boldsymbol{\beta}) + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle.$$

Making the substitution  $\boldsymbol{u} = \boldsymbol{\beta} - \boldsymbol{\beta}^{\star}$ , we rewrite this as

$$\widehat{oldsymbol{u}} = \operatorname*{argmin}_{oldsymbol{u} \in \mathbb{R}^d} H_L(oldsymbol{y} - oldsymbol{X}(oldsymbol{eta}^\star + oldsymbol{u})) + rac{\lambda}{2} \|oldsymbol{eta}^\star + oldsymbol{u}\|^2 + 
u \langle oldsymbol{\xi}, oldsymbol{eta}^\star + oldsymbol{u}
angle.$$

Expanding the quadratic term, recalling that  $\hat{u} = \hat{\beta} - \beta^*$  and  $\varepsilon^* = y - X\beta^*$ , and dropping terms that do not depend on u, we simplify this to

$$\widehat{\boldsymbol{u}} = \underset{\boldsymbol{u} \in \mathbb{R}^d}{\operatorname{argmin}} H_L(\boldsymbol{\varepsilon}^* - \boldsymbol{X}\boldsymbol{u}) + \frac{\lambda}{2} \|\boldsymbol{u}\|^2 + \langle \lambda \boldsymbol{\beta}^* + \nu \boldsymbol{\xi}, \boldsymbol{u} \rangle.$$
(3.12)

Next, we take the Legendre transform of  $H_L$  in equation (3.12). In our case, this simply amounts to applying the following identity regarding  $H_L$ , which is valid for all  $\mathbf{r} \in \mathbb{R}^n$ :

$$H_L(\boldsymbol{r}) = \max_{\boldsymbol{v} \in [-L,+L]^n} \langle \boldsymbol{r}, \boldsymbol{v} 
angle - rac{1}{2} \| \boldsymbol{v} \|^2.$$

The maximum is achieved iff  $\boldsymbol{v} = [\boldsymbol{r}]_L$ . Setting  $\boldsymbol{r} = \boldsymbol{\varepsilon}^{\star} - \boldsymbol{X} \boldsymbol{u}$ ,

$$\widehat{oldsymbol{u}} = rgmin_{oldsymbol{u} \in [-L,+L]^n} lpha_{oldsymbol{\varepsilon}^\star} - oldsymbol{X}oldsymbol{u}, oldsymbol{v} 
angle - rac{1}{2} \|oldsymbol{v}\|^2 + rac{\lambda}{2} \|oldsymbol{u}\|^2 + \langle \lambda oldsymbol{eta}^\star + 
u oldsymbol{\xi}, oldsymbol{u} 
angle.$$

Note that this objective function is simply  $Q_{u,v}$ . Given u, the maximum over v is achieved iff

$$oldsymbol{v} = [oldsymbol{r}]_L = [oldsymbol{\varepsilon}^\star - oldsymbol{X}oldsymbol{u}]_L$$

This coincides with  $\hat{\boldsymbol{v}} = [\boldsymbol{y} - \boldsymbol{X}\hat{\boldsymbol{\beta}}]_L$  when  $\boldsymbol{u} = \hat{\boldsymbol{u}}$ . The terms  $\frac{\lambda}{2} \|\boldsymbol{u}\|^2$  and  $-\frac{1}{2} \|\boldsymbol{v}\|^2$  ensure that  $Q_{\boldsymbol{u},\boldsymbol{v}}$  is  $\lambda$ -strongly convex in  $\boldsymbol{u}$  and 1-strongly concave in  $\boldsymbol{v}$ , so we conclude by the minimax theorem that  $(\hat{\boldsymbol{u}}, \hat{\boldsymbol{v}})$  is the unique saddle point of  $Q_{\boldsymbol{u},\boldsymbol{v}}$ .

# 3.2.2 Step 2: CGMT Universality

The goal of this step is to use CGMT universality (Theorem 2.9 in this paper, or Corollary 2.6 of [HS23]) to prove the following lemma, which relates the random variable  $Q_{u,v}$  to the random variable  $Q'_{u,v}$  that replaces -X with  $\frac{1}{\sqrt{d}}G$ , where G has the same shape as X but independent, standard Gaussian entries:

$$Q'_{\boldsymbol{u},\boldsymbol{v}} = \frac{1}{\sqrt{d}} \langle \boldsymbol{G} \boldsymbol{u}, \boldsymbol{v} \rangle + \psi(\boldsymbol{u}, \boldsymbol{v}).$$

Unlike X, whose rows have  $\ell^2$  norm bounded by R (as required for differential privacy), the entries of the matrix G are unbounded in the worst case. Despite this, the lemma shows that the random variable obtained by minimizing over u and maximizing over v has nearly the same CDF regardless of whether we start with  $Q_{u,v}$  or  $Q'_{u,v}$ . Note that the order of minimization over u and maximization over v below can be exchanged by the minimax theorem.

**Lemma 3.13.** If X follows a subgaussian design and  $y = X\beta^* + \varepsilon^*$  for  $\beta^* \rightsquigarrow \beta_0^*$  and  $\varepsilon^* \rightsquigarrow \varepsilon_0^*$ , then for all compact  $S_v \subseteq [-L, +L]^n$ , for all compact  $S_u \subseteq [-L_u, +L_u]^d$  with  $L_u \leq O(n^c)$  for a sufficiently small constant c > 0, and for all  $\tau \in \mathbb{R}$ ,

$$\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}'<\tau\right]\leq\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}<\tau+n^{1-\Omega(1)}\right]+n^{-\Omega(1)}$$

The above inequality also holds if we swap  $Q_{u,v}$  and  $Q'_{u,v}$ .

*Proof.* We calculate the gradient of  $\psi$ , as defined in Lemma 3.11:

$$\nabla \psi(\boldsymbol{u}, \boldsymbol{v}) = (\lambda \boldsymbol{u} + \lambda \boldsymbol{\beta}^{\star} + \nu \boldsymbol{\xi}, -\boldsymbol{v} + \boldsymbol{\varepsilon}^{\star}).$$

Since  $\beta^* \rightsquigarrow \beta_0^*$  and  $\varepsilon^* \rightsquigarrow \varepsilon_0^*$ , Lemma 3.6 implies that w.h.p., their  $\ell^{\infty}$  norms grow no faster than  $O(n^c)$ . The same is true of  $\boldsymbol{\xi}$  with high probability by Gaussianity, and for  $\boldsymbol{u}$  and  $\boldsymbol{v}$  by the assumptions on  $S_{\boldsymbol{u}}$  and  $S_{\boldsymbol{v}}$ , respectively. It follows that w.h.p.,

$$\|\nabla\psi(\boldsymbol{u},\boldsymbol{v})\|_1 = O(n^{1+c})$$

Rescaling the objective by  $\frac{1}{n}$  and applying Theorem 2.9 with  $\mathbf{A} = -\sqrt{d}\mathbf{X}$  and  $\mathbf{B} = \mathbf{G}$  (and vice versa) and  $g = n^{-1/21}$ , we conclude that

$$\Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}'<\tau\right] = \Pr\left[\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}<\tau\pm O(n^{20/21})\right] \pm O(n^{-\frac{1}{6}+\frac{1}{7}+2c+o(1)}).$$

The key nontrivial assumption of Lemma 3.13 is that the  $\ell^{\infty}$  diameter of the constraint set  $S_u$  satisfies  $L_u \leq O(n^c)$  for a sufficiently small c > 0. Therefore, to make effective use of this result, we must verify that for all c > 0, w.h.p.,

$$\|\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|_{\infty} = O(n^c).$$

By the triangle inequality, it suffices to check that  $\|\boldsymbol{\beta}^{\star}\|_{\infty} = O(n^c)$  and  $\|\boldsymbol{\hat{\beta}}\|_{\infty} = O(n^c)$ . For the former, Lemma 3.6 and  $\boldsymbol{\beta}^{\star} \rightsquigarrow \beta_0^{\star}$  imply that  $\|\boldsymbol{\beta}^{\star}\|_{\infty} = O(n^c)$  with high probability. For the latter, we prove the following lemma.

**Lemma 3.14.** Let  $\widehat{\boldsymbol{\beta}}$  be the output of Algorithm 1 with L-Lipschitz Huber loss  $\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = H_L(y - \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle)$ . Fix any  $\boldsymbol{\varepsilon}^{\star} \in \mathbb{R}^n$  and  $\boldsymbol{\beta}^{\star} \rightsquigarrow \beta_0^{\star}$ . If  $\boldsymbol{X}$  follows a subgaussian design and  $\boldsymbol{y} = \boldsymbol{X} \boldsymbol{\beta}^{\star} + \boldsymbol{\varepsilon}^{\star}$ , then for all c > 0, w.h.p. over  $\boldsymbol{X}$  and  $\boldsymbol{\xi}$ ,

$$\|\widehat{\boldsymbol{\beta}}\|_{\infty} \le O(n^c).$$

*Proof.* We use the leave-one-(feature)-out technique, as in [HS23], while accounting for the noise for differential privacy. First, denote the perturbed objective function by H:

$$H(\boldsymbol{\beta}) = H_L(\boldsymbol{y} - \boldsymbol{X}\boldsymbol{\beta}) + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle.$$

We wish to study the estimator  $\hat{\beta}^{(j)}$  obtained by omitting the  $j^{\text{th}}$  feature from all data points. By equation (3.12), in which  $\boldsymbol{u} = \boldsymbol{\beta} - \boldsymbol{\beta}^{\star}$  and  $\hat{\boldsymbol{u}} = \hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}$ , we see that dropping the  $j^{\text{th}}$  column from  $\boldsymbol{X}$  is equivalent to enforcing the constraint  $u_j = 0$ , or  $\beta_j = \beta_j^{\star}$ :

$$\widehat{\boldsymbol{\beta}}^{(j)} = \underset{\boldsymbol{\beta} \in \mathbb{R}^d}{\operatorname{argmin}} H(\boldsymbol{\beta}) \quad \text{s.t.} \quad \beta_j = \beta_j^\star.$$

To bound the  $j^{\text{th}}$  coordinate of  $\hat{\beta}$ , we will relate  $\hat{\beta}$  to  $\hat{\beta}^{(j)}$ , whose  $j^{\text{th}}$  coordinate is  $\beta_j^{\star}$  by definition. To this end, note that H is  $\lambda$ -strongly convex, so

$$\|\widehat{\boldsymbol{\beta}} - \widehat{\boldsymbol{\beta}}^{(j)}\| \le \frac{1}{\lambda} \|\nabla H(\widehat{\boldsymbol{\beta}}) - \nabla H(\widehat{\boldsymbol{\beta}}^{(j)})\|.$$

The distance  $\|\widehat{\beta} - \widehat{\beta}^{(j)}\|$  is at least  $|\widehat{\beta}_j - \widehat{\beta}_j^{(j)}|$ . Since  $\widehat{\beta}$  minimizes H, the gradient  $\nabla H(\widehat{\beta})$  vanishes. Similarly, the gradient  $\nabla H(\widehat{\beta}^{(j)})$  vanishes in all but its  $j^{\text{th}}$  coordinate, so the inequality simplifies to

$$|\widehat{\beta}_j - \beta_j^{\star}| \le \left| \frac{1}{\lambda} \nabla_j H(\widehat{\beta}^{(j)}) \right|.$$

All that remains is to bound the partial derivative on the right side. By direct calculation,

$$\nabla_j H(\boldsymbol{\beta}) = \langle [\boldsymbol{y} - \boldsymbol{X}\boldsymbol{\beta}]_L, \boldsymbol{X}_j \rangle + \lambda \beta_j + \nu \xi_j.$$

Here,  $X_j \in \mathbb{R}^n$  denotes the  $j^{\text{th}}$  column of X (not the  $j^{\text{th}}$  data point). Substituting  $\beta = \hat{\beta}^{(j)}$ , whose  $j^{\text{th}}$  coordinate is  $\beta_j^*$ , we see that

$$|\widehat{\beta}_j - \beta_j^{\star}| \leq \left|\frac{1}{\lambda} \nabla_j H(\widehat{\boldsymbol{\beta}}^{(j)})\right| = \left|\frac{1}{\lambda} \langle [\boldsymbol{y} - \boldsymbol{X} \widehat{\boldsymbol{\beta}}^{(j)}]_L, \boldsymbol{X}_j \rangle + \beta_j^{\star} + \frac{\nu}{\lambda} \xi_j \right|.$$

Observe that we can rewrite the truncated residual vector as  $[\boldsymbol{y} - \boldsymbol{X}\hat{\boldsymbol{\beta}}^{(j)}]_L = [\boldsymbol{\varepsilon}^{\star} - \boldsymbol{X}(\hat{\boldsymbol{\beta}}^{(j)} - \boldsymbol{\beta}^{\star})]_L$ , which, along with the fact that  $\hat{\boldsymbol{\beta}}^{(j)} - \boldsymbol{\beta}^{\star}$  vanishes in its  $j^{\text{th}}$  coordinate, makes it clear that it is independent of the vector  $\boldsymbol{X}_j$ . Since this truncated vector lies in  $[-L, +L]^n$  and is independent of  $\boldsymbol{X}_j$ , we have by subgaussianity (Lemma 2.18) and the triangle inequality that

$$\left\| |\widehat{\beta}_j - \beta_j^\star| - |\beta_j^\star| \right\|_{\psi_2} \le \frac{L\sqrt{n}}{\lambda} \|\boldsymbol{X}_j\|_{\psi_2} + \frac{\nu}{\lambda} \|\xi_j\|_{\psi_2}.$$

In a subgaussian design,  $\|\mathbf{X}_j\|_{\psi_2} = O(1/\sqrt{n})$ . Since  $\boldsymbol{\xi}$  is Gaussian,  $\|\boldsymbol{\xi}_j\|_{\psi_2} = O(1)$ . It follows that  $\||\widehat{\beta}_j - \beta_j^\star| - |\beta_j^\star|\|_{\psi_2} \le O(1)$  for each  $j \in [d]$ . Hence, by a union bound over all  $j \in [d]$ , as  $t \to \infty$ ,

$$\Pr\left[\max_{j\in[d]} \left| |\widehat{\beta}_j - \beta_j^{\star}| - |\beta_j^{\star}| \right| > t\right] \le n \cdot \exp(-\Omega(t^2)).$$

Setting  $t = \log n$  and recalling that  $\|\beta\|_{\infty} = \max_{j \in [d]} |\beta_j|$ , we see that with probability  $1 - n^{-\Omega(\log n)}$ ,

$$\|\widehat{\boldsymbol{\beta}}\|_{\infty} \le \log n + 2\|\boldsymbol{\beta}^{\star}\|_{\infty}.$$

To complete the proof, recall from Lemma 3.6 that for all c > 0, w.h.p.,  $\|\beta^{\star}\|_{\infty} = O(n^{c})$  as well.  $\Box$ 

### 3.2.3 Step 3: CGMT Analysis

In this step, we analyze the random variable  $Q'_{u,v}$  using CGMT (Theorem 2.8), roughly following the strategy of [TAH18].

In some ways, our proof is simpler than the proof in [TAH18]. This is partly because we focus on the special case of robust linear regression with  $\ell^2$  regularization, as opposed to a general loss function  $\mathcal{L}$  and general regularizer f. It is also simpler because our proof circumvents the introduction of several extraneous scalar- and vector-valued variables that are used in prior work.

In other ways, our proof is somewhat more complex than prior papers in this line of work. This is partly because we consider the random perturbation  $\boldsymbol{\xi}$  introduced for differential privacy, and partly because in order to make do with our weaker assumptions and still prove a stronger conclusion, we must take great care to bound the magnitude of various error terms.

To begin, we recall and analyze the *auxiliary* random variable for CGMT, defined as

$$Q_{oldsymbol{u},oldsymbol{v}}'' = rac{\|oldsymbol{u}\|}{\sqrt{d}} \langleoldsymbol{h},oldsymbol{v}
angle - rac{\|oldsymbol{v}\|}{\sqrt{d}} \langleoldsymbol{g},oldsymbol{u}
angle + \psi(oldsymbol{u},oldsymbol{v}) \qquad ext{for} \qquad oldsymbol{g} \sim \mathcal{N}(oldsymbol{0},oldsymbol{I}_d) ext{ and }oldsymbol{h} \sim \mathcal{N}(oldsymbol{0},oldsymbol{I}_n).$$

Eventually, we will use CGMT (Theorem 2.8) to relate  $Q_{u,v}$  via  $Q'_{u,v}$  to  $Q''_{u,v}$ .

**Lemma 3.15.** Define  $\sigma^*, \tau^* > 0$  as in Theorem 3.7(b), and consider the pair  $(\widetilde{u}, \widetilde{v})$  with

$$\widetilde{\boldsymbol{u}} = \tau^* \left( \sqrt{\frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma^* \boldsymbol{Z} + \varepsilon_0^*}{1 + \tau^*} \right]_L^2} \boldsymbol{g} - \lambda \boldsymbol{\beta}^* - \nu \boldsymbol{\xi} \right), \qquad \widetilde{\boldsymbol{v}} = \left[ \frac{\sigma^* \boldsymbol{h} + \boldsymbol{\varepsilon}^*}{1 + \tau^*} \right]_L.$$

Then, under the assumptions of Theorem 3.7(b), there exists a constant  $c^* \in \mathbb{R}$  such that w.h.p.,

- The function  $\boldsymbol{u} \mapsto Q_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}''$  is  $\lambda$ -strongly convex in  $\boldsymbol{u} \in \mathbb{R}^d$ ,
- The function  $\boldsymbol{v} \mapsto Q''_{\widetilde{\boldsymbol{u}},\boldsymbol{v}}$  is 1-strongly concave in  $\boldsymbol{v} \in [-L,+L]^n$ ,
- The pair  $(\widetilde{\boldsymbol{u}}, \widetilde{\boldsymbol{v}})$  satisfies

$$c^{\star}n - n^{1-\Omega(1)} \leq \min_{\boldsymbol{u} \in \mathbb{R}^d} Q_{\boldsymbol{u}, \widetilde{\boldsymbol{v}}}' \leq Q_{\widetilde{\boldsymbol{u}}, \widetilde{\boldsymbol{v}}}'' \leq \max_{\boldsymbol{v} \in [-L, +L]^n} Q_{\widetilde{\boldsymbol{u}}, \boldsymbol{v}}' \leq c^{\star}n + n^{1-\Omega(1)}.$$

We call  $(\widetilde{\boldsymbol{u}}, \widetilde{\boldsymbol{v}})$  an approximate saddle point of  $Q_{\boldsymbol{u},\boldsymbol{v}}''$ .

*Proof.* To show that  $(\tilde{u}, \tilde{v})$  is an approximate saddle point of  $Q''_{u,v}$ , it suffices to show that each of  $\tilde{u}$  and  $\tilde{v}$  is an *approximately best response* to the other, and that  $|Q''_{\tilde{u},\tilde{v}} - c^* n| \leq n^{1-\Omega(1)}$ . Formally, we say that  $\tilde{u}$  is an approximate best response to  $\tilde{v}$  if

$$Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}'' \leq \min_{\boldsymbol{u} \in \mathbb{R}^d} Q_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}'' + n^{1-\Omega(1)}.$$

Similarly, we say that  $\widetilde{v}$  is an approximate best response to  $\widetilde{u}$  if

$$Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}'' \ge \max_{\boldsymbol{v} \in [-L,+L]^n} Q_{\widetilde{\boldsymbol{u}},\boldsymbol{v}}'' - n^{1-\Omega(1)}$$

In order to prove these inequalities, we will check that certain derivatives approximately vanish.

(a) ( $\tilde{u}$  is an approximate best response to  $\tilde{v}$ ) We study the minimum over  $u \in \mathbb{R}^d$  of

$$Q_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}'' = \frac{\|\boldsymbol{u}\|}{\sqrt{d}} \langle \boldsymbol{h}, \widetilde{\boldsymbol{v}} \rangle - \frac{\|\widetilde{\boldsymbol{v}}\|}{\sqrt{d}} \langle \boldsymbol{g}, \boldsymbol{u} \rangle + \psi(\boldsymbol{u}, \widetilde{\boldsymbol{v}}).$$

If we can show that  $Q''_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}$  is a  $\lambda$ -strongly convex function of  $\boldsymbol{u}$ , then minimizing  $Q''_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}$  over  $\boldsymbol{u} \in \mathbb{R}^d$  reduces to finding a point  $\boldsymbol{u} \in \mathbb{R}^d$  at which the gradient  $\nabla_{\boldsymbol{u}}Q''_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}$  has small norm. To this end, observe that  $\psi$  contains the term  $\frac{\lambda}{2} ||\boldsymbol{u}||^2$ , so  $\psi(\boldsymbol{u},\widetilde{\boldsymbol{v}})$  is  $\lambda$ -strongly convex in  $\boldsymbol{u}$ . The term  $-\frac{\|\widetilde{\boldsymbol{v}}\|}{\sqrt{d}}\langle \boldsymbol{g}, \boldsymbol{u} \rangle$  is a linear function of  $\boldsymbol{u}$ , and hence convex. Finally, the term  $\frac{\|\boldsymbol{u}\|}{\sqrt{d}}\langle \boldsymbol{h},\widetilde{\boldsymbol{v}} \rangle$  will be convex, as well, if we can show that  $\langle \boldsymbol{h},\widetilde{\boldsymbol{v}} \rangle$  is positive with high probability.

Quantities like  $\langle \boldsymbol{h}, \tilde{\boldsymbol{v}} \rangle$ ,  $\langle \boldsymbol{g}, \tilde{\boldsymbol{u}} \rangle$ ,  $\|\tilde{\boldsymbol{v}}\|$ , and  $\|\tilde{\boldsymbol{u}}\|$  can be easily computed in the limit by combining the definitions of  $\tilde{\boldsymbol{u}}$  and  $\tilde{\boldsymbol{v}}$  in this lemma's statement with the assumptions on  $\boldsymbol{g}, \boldsymbol{h}, \boldsymbol{\beta}^{\star}, \boldsymbol{\varepsilon}^{\star}$ , and  $\boldsymbol{\xi}$ . By the definition of pseudo-Lipschitz convergence (Definition 3.3), w.h.p.,

$$\begin{split} \frac{\|\widetilde{\boldsymbol{u}}\|}{\sqrt{d}} &= \tau^{\star} \sqrt{(\sigma_{v}^{\star})^{2} + \lambda^{2} \kappa^{2} + \nu^{2}} \pm n^{-\Omega(1)}, \qquad \qquad \frac{1}{d} \langle \boldsymbol{g}, \widetilde{\boldsymbol{u}} \rangle = \tau^{\star} \sigma_{v}^{\star} \pm n^{-\Omega(1)}, \\ \frac{\|\widetilde{\boldsymbol{v}}\|}{\sqrt{d}} &= \sigma_{v}^{\star} \pm n^{-\Omega(1)}, \qquad \qquad \frac{1}{d} \langle \boldsymbol{h}, \widetilde{\boldsymbol{v}} \rangle = \frac{1}{\delta} \mathbb{E} \left[ \left[ \frac{\sigma^{\star} Z + \varepsilon_{0}^{\star}}{1 + \tau^{\star}} \right]_{L} Z \right] \pm n^{-\Omega(1)}. \end{split}$$

Above, for convenience, we have introduced the abbreviation

$$\sigma_v^{\star} = \sqrt{\frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma^{\star} Z + \varepsilon_0^{\star}}{1 + \tau^{\star}} \right]_L^2}.$$

Using equation (3.8a) (see Theorem 3.7(b)), we can simplify the expression for  $\|\tilde{\boldsymbol{u}}\|$  to

$$\frac{\|\widetilde{\boldsymbol{u}}\|}{\sqrt{d}} = \sigma^{\star} \pm n^{-\Omega(1)}$$

Using Stein's lemma followed by equation (3.8b), we can simplify the expression for  $\langle h, \tilde{v} \rangle$  to

$$\frac{1}{d}\langle \boldsymbol{h}, \widetilde{\boldsymbol{v}} \rangle = \frac{\sigma^{\star}}{\delta(1+\tau^{\star})} \Pr\left[-L < \frac{\sigma^{\star}Z + \varepsilon_{0}^{\star}}{1+\tau^{\star}} < L\right] \pm n^{-\Omega(1)} = \left(\frac{1}{\tau^{\star}} - \lambda^{\star}\right) \sigma^{\star} \pm n^{-\Omega(1)}.$$

By examining equation (3.8b), we see that  $1/\tau^* > \lambda^*$ , so the quantity above is positive with high probability, which establishes that  $Q''_{u,\tilde{v}}$  is  $\lambda$ -strongly convex in u with high probability. Therefore, all that remains is to evaluate its gradient at  $\tilde{u}$ . To this end, we calculate

$$\nabla_{\boldsymbol{u}} Q_{\boldsymbol{u},\boldsymbol{v}}'' = \frac{1}{\sqrt{d}} \frac{\boldsymbol{u}}{\|\boldsymbol{u}\|} \langle \boldsymbol{h}, \boldsymbol{v} \rangle - \frac{\|\boldsymbol{v}\|}{\sqrt{d}} \boldsymbol{g} + \lambda \boldsymbol{u} + \lambda \boldsymbol{\beta}^{\star} + \nu \boldsymbol{\xi}$$

Evaluating at  $(\tilde{u}, \tilde{v})$  and substituting our expressions for  $\langle h, \tilde{v} \rangle$ ,  $\langle g, \tilde{u} \rangle$ ,  $\|\tilde{v}\|$ , and  $\|\tilde{u}\|$  yields

$$\nabla_{\boldsymbol{u}} Q_{\boldsymbol{\widetilde{u}},\boldsymbol{\widetilde{v}}}'' = \left(\frac{1}{\tau^{\star}} \pm n^{-\Omega(1)}\right) \boldsymbol{\widetilde{u}} - (\sigma_{\boldsymbol{v}}^{\star} \pm n^{-\Omega(1)})\boldsymbol{g} + \lambda \boldsymbol{\beta}^{\star} + \nu \boldsymbol{\xi}$$

By substituting the definition of  $\widetilde{\boldsymbol{u}}$  in this lemma's statement and using the fact that for any constant c > 0, we have  $\|\widetilde{\boldsymbol{u}}\| + \|\boldsymbol{g}\| + \|\boldsymbol{\beta}^{\star}\| + \|\boldsymbol{\xi}\| \leq O(n^{\frac{1}{2}+c})$  w.h.p. (Lemma 3.6), we see that the above gradient clearly has  $\ell^2$  norm

$$\|\nabla_{\boldsymbol{u}} Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}''\| \le n^{\frac{1}{2} - \Omega(1)}$$

Thus, by  $\lambda$ -strong convexity, we have with high probability that

$$Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}'' \leq \min_{\boldsymbol{u} \in \mathbb{R}^d} Q_{\boldsymbol{u},\widetilde{\boldsymbol{v}}}' + n^{1 - \Omega(1)}$$

(b) ( $\tilde{v}$  is an approximate best response to  $\tilde{u}$ ) We study the max over  $v \in [-L, +L]^n$  of

$$Q_{\widetilde{oldsymbol{u}},oldsymbol{v}}''=rac{\|\widetilde{oldsymbol{u}}\|}{\sqrt{d}}\langleoldsymbol{h},oldsymbol{v}
angle-rac{\|oldsymbol{v}\|}{\sqrt{d}}\langleoldsymbol{g},\widetilde{oldsymbol{u}}
angle+\psi(\widetilde{oldsymbol{u}},oldsymbol{v}).$$

Analogously to the previous part, we first verify 1-strong concavity with respect to  $\boldsymbol{v}$ . To this end, observe that  $\psi$  contains the term  $-\frac{1}{2} \|\boldsymbol{v}\|^2$ , so  $\psi(\tilde{\boldsymbol{u}}, \boldsymbol{v})$  is 1-strongly concave in  $\boldsymbol{v}$ . The term  $\frac{\|\tilde{\boldsymbol{u}}\|}{\sqrt{d}} \langle \boldsymbol{h}, \boldsymbol{v} \rangle$  is a linear function of  $\boldsymbol{v}$ , and hence concave. Finally, the term  $-\frac{\|\boldsymbol{v}\|}{\sqrt{d}} \langle \boldsymbol{g}, \tilde{\boldsymbol{u}} \rangle$  is concave, as well, since we have already shown that  $\langle \boldsymbol{g}, \tilde{\boldsymbol{u}} \rangle$  is positive with high probability. Now, all that remains is to evaluate the gradient of  $Q''_{\tilde{\boldsymbol{u}}, \boldsymbol{v}}$  at  $\tilde{\boldsymbol{v}}$ . To this end, we calculate

$$\nabla_{\boldsymbol{v}} Q_{\boldsymbol{u},\boldsymbol{v}}'' = \frac{\|\boldsymbol{u}\|}{\sqrt{d}} \boldsymbol{h} - \frac{1}{\sqrt{d}} \frac{\boldsymbol{v}}{\|\boldsymbol{v}\|} \langle \boldsymbol{g}, \boldsymbol{u} \rangle + \boldsymbol{\varepsilon}^{\star}.$$

Evaluating at  $(\tilde{u}, \tilde{v})$  and substituting our expressions for  $\langle h, \tilde{v} \rangle$ ,  $\langle g, \tilde{u} \rangle$ ,  $\|\tilde{v}\|$ , and  $\|\tilde{u}\|$  yields

$$\nabla_{\boldsymbol{v}} Q_{\boldsymbol{\widetilde{u}},\boldsymbol{\widetilde{v}}}'' = (\sigma^{\star} \pm n^{-\Omega(1)})\boldsymbol{h} + \boldsymbol{\varepsilon}^{\star} - (1 + \tau^{\star} \pm n^{-\Omega(1)})\boldsymbol{\widetilde{v}}$$

By substituting the definition of  $\tilde{\boldsymbol{v}}$  and using the fact that for any constant c > 0, we have  $\|\tilde{\boldsymbol{v}}\| + \|\boldsymbol{h}\| + \|\boldsymbol{\varepsilon}^{\star}\| \le O(n^{\frac{1}{2}+c})$  w.h.p. (Lemma 3.6), we see that there exists a vector of  $\ell^2$  norm at most  $O(n^{\frac{1}{2}+c})$  in the set

$$\nabla_{\boldsymbol{v}} Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}'' - \partial \mathbf{1}_{[-L,+L]^n}(\widetilde{\boldsymbol{v}})$$

This set is a translation of the subdifferential set  $\partial \mathbf{1}_{[-L,+L]^n}(v) = \prod_{i=1}^n \partial \mathbf{1}_{[-L,+L]}(v_i)$ , where

$$\partial \mathbf{1}_{[-L,+L]}(v_i) = \begin{cases} \{0\} & \text{if } |v_i| < L, \\ (-\infty,0] & \text{if } v_i = -L, \\ [0,+\infty) & \text{if } v_i = L, \\ \varnothing & \text{if } |v_i| > L. \end{cases}$$

By 1-strong concavity on  $[-L, +L]^n$ , we have with high probability that

$$Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}'' \ge \max_{\boldsymbol{v} \in [-L,+L]^n} Q_{\widetilde{\boldsymbol{u}},\boldsymbol{v}}'' - n^{1-\Omega(1)}.$$

To conclude the proof, we remark that plugging our estimates for  $\langle \boldsymbol{h}, \tilde{\boldsymbol{v}} \rangle$ ,  $\langle \boldsymbol{g}, \tilde{\boldsymbol{u}} \rangle$ ,  $\|\tilde{\boldsymbol{v}}\|$ , and  $\|\tilde{\boldsymbol{u}}\|$  into the definition of  $Q''_{\boldsymbol{u},\boldsymbol{v}}$  similarly implies that there exists a constant  $c^* \in \mathbb{R}$  such that w.h.p.,

$$|Q_{\widetilde{\boldsymbol{u}},\widetilde{\boldsymbol{v}}}'' - c^* n| \le n^{1 - \Omega(1)}.$$

### 3.2.4 Putting Steps 1, 2, and 3 Together

Proof of Theorem 3.7. Note that part (a) follows immediately from Theorem 5.1, the observation that  $\rho'' : \mathbb{R} \to [0,1]$ , and a change of variables in the case that  $R \neq 1$ . The extra factors in  $(2\varepsilon_{\text{DP}}, (1 + e^{\varepsilon_{\text{DP}}})\delta_{\text{DP}})$  come from the conversion from add/remove-one DP to replace-one DP (see Lemma 2.7). Therefore, we focus on part (b). By Lemma 3.15, there exist constants  $c^* \in \mathbb{R}$  and  $c_{\text{cgmt}} > 0$  such that w.h.p,

$$c^*n - O(n^{1-c_{\rm cgmt}}) \le \min_{\boldsymbol{u} \in \mathbb{R}^d} Q_{\boldsymbol{u}, \widetilde{\boldsymbol{v}}}' \le Q_{\widetilde{\boldsymbol{u}}, \widetilde{\boldsymbol{v}}}'' \le \max_{\boldsymbol{v} \in [-L, +L]^n} Q_{\widetilde{\boldsymbol{u}}, \boldsymbol{v}}' \le c^*n + O(n^{1-c_{\rm cgmt}}).$$
(3.16)

By Lemma 3.6 and Lemma 3.14, for any arbitrarily small constant  $c_{\text{diam}} > 0$ , there exists an upper bound  $L_{\boldsymbol{u}} = O(n^{c_{\text{diam}}})$  such that w.h.p.,

$$\widetilde{\boldsymbol{u}}, \widehat{\boldsymbol{u}} \in [-L_{\boldsymbol{u}}, +L_{\boldsymbol{u}}]^d.$$

For brevity, set  $S_{\boldsymbol{u}} = [-L_{\boldsymbol{u}}, +L_{\boldsymbol{u}}]^d$  and  $S_{\boldsymbol{v}} = [-L, +L]^n$ . Then,  $\widetilde{\boldsymbol{u}} \in S_{\boldsymbol{u}}$  implies that w.h.p.,

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}'\leq c^{\star}n+O(n^{1-c_{\mathrm{cgmt}}}).$$

By CGMT (Theorem 2.8), we similarly have that w.h.p.,

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}' \leq c^{\star}n + O(n^{1-c_{\text{cgmt}}}).$$

By universality (Lemma 3.13), there is a constant  $c_{univ} \in (0, c_{cgmt}]$  such that w.h.p.,

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}} \le c^{\star}n + O(n^{1-c_{\mathrm{univ}}}).$$
(3.17)
By the definition of  $\tilde{u}$  in the statement of Lemma 3.15, along with our assumption that  $\beta^{\star} \rightsquigarrow \beta_0^{\star}$ ,

$$(\boldsymbol{\beta}^{\star},\boldsymbol{\xi},\widetilde{\boldsymbol{u}}) \rightsquigarrow \left(\beta_{0}^{\star},\ \xi_{0},\ \tau^{\star}\left(\sqrt{\frac{1}{\delta}\mathbb{E}\left[\frac{\sigma^{\star}Z+\varepsilon_{0}^{\star}}{1+\tau^{\star}}\right]_{L}^{2}}Z-\lambda\beta_{0}^{\star}-\nu\xi_{0}\right)\right).$$

For brevity, let  $u_0 \in \mathbb{R}$  denote the third random variable in the above triple. Then, the above assertion is that  $(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \widetilde{\boldsymbol{u}}) \rightsquigarrow (\beta_0^{\star}, \xi_0, u_0)$ , and we want to show that  $(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}) \rightsquigarrow (\beta_0^{\star}, \xi_0, u_0)$ , as well. To this end, fix an order-k pseudo-Lipschitz function  $f : \mathbb{R}^3 \to \mathbb{R}$ . By Definition 3.3, the above statement means that there exists a constant  $c_{pL} \in (0, c_{univ}]$  such that w.h.p.,

$$\left|\bar{f}(\boldsymbol{\beta}^{\star},\boldsymbol{\xi},\widetilde{\boldsymbol{u}}) - \mathbb{E}[f(\boldsymbol{\beta}_{0}^{\star},\boldsymbol{\xi}_{0},\boldsymbol{u}_{0})]\right| \leq O(n^{-c_{\mathrm{pL}}}),\tag{3.18}$$

where we have defined the function  $\overline{f} : \mathbb{R}^{d \times 3} \to \mathbb{R}$  by  $\overline{f}(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \boldsymbol{u}) = \frac{1}{d} \sum_{j=1}^{d} f(\beta_{j}^{\star}, \xi_{j}, u_{j})$ . Next, let  $c_{\text{slack}} > 0$  be any strictly positive constant satisfying the strict inequality  $c_{\text{slack}} < c_{\text{pL}}/2 - (k-1)c_{\text{diam}}$ . Note that such a choice of  $c_{\text{slack}}$  is always possible because, as we noted earlier, we can take the constant  $c_{\text{diam}}$  to be arbitrarily small. Next, define the open set

$$\mathcal{T}_{\boldsymbol{u}} = \left\{ \boldsymbol{u} \in \mathbb{R}^d : \left| \bar{f}(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \boldsymbol{u}) - \mathbb{E}[f(\boldsymbol{\beta}_0^{\star}, \xi_0, u_0)] \right| < n^{-c_{\text{slack}}} \right\}.$$
(3.19)

Observe that by Lemma 3.15, the auxiliary objective function is  $\lambda$ -strongly convex in  $\boldsymbol{u}$  when  $\boldsymbol{v} = \tilde{\boldsymbol{v}}$ . Also, since  $S_{\boldsymbol{u}} = [-L_{\boldsymbol{u}}, +L_{\boldsymbol{u}}]^d$ , the order-k pseudo-Lipschitz function  $\bar{f}$  is in fact Lipschitz continuous in  $\boldsymbol{u}$  over  $S_{\boldsymbol{u}}$  with Lipschitz constant

$$L_{\bar{f}} \leq \frac{1}{d} \cdot \sqrt{d} \cdot (1 + 2L_{\boldsymbol{u}})^{k-1} = O(n^{-\frac{1}{2} + (k-1)c_{\text{diam}}}).$$

Together,  $\Omega(1)$ -strong convexity,  $O(n^{-\frac{1}{2}+(k-1)c_{\text{diam}}})$ -Lipschitzness, (3.16), (3.18), and (3.19) imply that if we excise  $\mathcal{T}_{\boldsymbol{u}}$  from  $\mathcal{S}_{\boldsymbol{u}}$  to form a new compact constraint set  $\mathcal{S}_{\boldsymbol{u}} \setminus \mathcal{T}_{\boldsymbol{u}}$ , then w.h.p.,

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}\setminus\mathcal{T}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}''\geq c^{\star}n+\Omega\left(\left(\frac{n^{-c_{\mathrm{slack}}}}{n^{-\frac{1}{2}+(k-1)c_{\mathrm{diam}}}}\right)^{2}\right)-O(n^{1-c_{\mathrm{pL}}}).$$

The  $\Omega(\cdot)$  term on the right side simplifies to  $\Omega(n^{1-c_{\text{total}}})$  where  $c_{\text{total}} = 2c_{\text{slack}} + 2(k-1)c_{\text{diam}}$ . By our choice of  $c_{\text{slack}}$ , we have that  $\Omega(n^{1-c_{\text{total}}})$  dominates  $O(n^{1-c_{\text{pL}}})$ , so w.h.p.,

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}\setminus\mathcal{T}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}''\geq c^{\star}n+\Omega(n^{1-c_{\mathrm{total}}}).$$

Applying CGMT (Theorem 2.8) and universality (Lemma 3.13) as before (note that  $c_{\text{total}} < c_{\text{cgmt}}$  and  $c_{\text{total}} < c_{\text{univ}}$ ), we have w.h.p. that

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}\setminus\mathcal{T}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}\geq c^{\star}n+\Omega(n^{1-c_{\text{total}}}).$$
(3.20)

At this point, (3.17) and (3.20) imply that w.h.p.,

$$\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}\setminus\mathcal{T}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}>\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q_{\boldsymbol{u},\boldsymbol{v}}.$$

(note the strict inequality). Thus, the minimizer, which by Lemma 3.11 is  $\hat{u} = \hat{\beta} - \beta^* \in S_u$ , w.h.p. lies inside  $\mathcal{T}_u$ . Unpacking the definition of  $\mathcal{T}_u$  in (3.19), we conclude that

$$(\boldsymbol{\beta}^{\star},\boldsymbol{\xi},\widehat{\boldsymbol{\beta}}-\boldsymbol{\beta}^{\star}) \rightsquigarrow (\beta_0^{\star},\xi_0,u_0).$$

This completes the characterization of the estimation error of  $\hat{\beta}$  in part (b). For the truncated residual error, we now carry out an entirely analogous dual argument with the roles of  $\boldsymbol{u}$  and  $\boldsymbol{v}$  exchanged. As before, we start by noting that by CGMT (Theorem 2.8) and universality (Lemma 3.13), we have w.h.p. that

$$\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}Q_{\boldsymbol{u},\boldsymbol{v}}\geq c^{\star}n-O(n^{1-c_{\mathrm{univ}}}).$$

By the definition of  $\tilde{v}$  in the statement of Lemma 3.15, along with our assumption that  $\varepsilon^* \rightsquigarrow \varepsilon_0^*$ ,

$$(\boldsymbol{\varepsilon}^{\star}, \widetilde{\boldsymbol{v}}) \rightsquigarrow \left(\varepsilon_{0}^{\star}, \left[\frac{\sigma^{\star}Z + \varepsilon_{0}^{\star}}{1 + \tau^{\star}}\right]_{L}\right)$$

For brevity, let  $v_0 \in \mathbb{R}$  denote the second random variable in the above pair. Then, the above assertion is that  $(\boldsymbol{\varepsilon}^{\star}, \tilde{\boldsymbol{v}}) \rightsquigarrow (\varepsilon_0^{\star}, v_0)$ , and we want to show that  $(\boldsymbol{\varepsilon}^{\star}, [\boldsymbol{y} - \boldsymbol{X}\hat{\boldsymbol{\beta}}]_L) \rightsquigarrow (\varepsilon_0^{\star}, v_0)$  as well. To this end, as before, we take any order-k pseudo-Lipschitz function  $f : \mathbb{R}^2 \to \mathbb{R}$ , and excise from  $S_{\boldsymbol{v}}$  the open set  $\mathcal{T}_{\boldsymbol{v}}$  such that  $\boldsymbol{v} \in \mathcal{T}_{\boldsymbol{v}}$  iff the average value of f over the coordinates of  $(\boldsymbol{\varepsilon}^{\star}, \boldsymbol{v})$  differs from the expected value of f over the randomness of  $(\varepsilon_0^{\star}, v_0)$  by strictly less than  $n^{-c'_{\text{slack}}}$ , for a sufficiently small constant  $c'_{\text{slack}} > 0$ . Doing so yields (again by CGMT, universality, pseudo-Lipschitzness, and the strong concavity afforded by Lemma 3.15), w.h.p.,

$$\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}\setminus\mathcal{T}_{\boldsymbol{v}}}\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}Q_{\boldsymbol{u},\boldsymbol{v}}\leq c^{\star}n-\Omega(n^{1-c'_{\text{total}}}),$$

for an appropriate positive constant  $c'_{\text{total}} < c_{\text{univ}}$ . At this point, we have that w.h.p.,

$$\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}\setminus\mathcal{T}_{\boldsymbol{v}}}\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}Q_{\boldsymbol{u},\boldsymbol{v}}<\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}Q_{\boldsymbol{u},\boldsymbol{v}}$$

Thus, the maximizer, which by Lemma 3.11 is  $\hat{v} = [y - X\hat{\beta}]_L \in \mathcal{S}_v$ , w.h.p. lies inside  $\mathcal{T}_v$ , so

$$(\boldsymbol{\varepsilon}^{\star}, [\boldsymbol{y} - \boldsymbol{X}\widehat{\boldsymbol{\beta}}]_L) \rightsquigarrow \left(\varepsilon_0^{\star}, \left[\frac{\sigma^{\star} Z + \varepsilon_0^{\star}}{1 + \tau^{\star}}\right]_L\right).$$

#### 3.3 Additional Proofs

Proof of Lemma 3.5. If f is order-k pseudo-Lipschitz, then  $|f(x)| \leq L(1+|x|^k)$  for some L. Thus,  $f(x_0)$  has finite mean and variance as long as  $x_0$  has finite  $2k^{\text{th}}$  moment. By Chebyshev's inequality, independence of  $x_1, \ldots, x_m$ , and the fact that  $m = \Theta(n)$ ,

$$\Pr\left[\left|\frac{1}{m}\sum_{i=1}^{m}f(x_i) - \mathbb{E}[f(x_0)]\right| \ge n^{-\frac{1}{3}}\right] \le \frac{\operatorname{Var}(f(x_0))}{m(n^{-1/3})^2} = O(n^{-\frac{1}{3}}).$$

*Proof of Lemma 3.6.* For simplicity, suppose first that  $\boldsymbol{x}$  is deterministic. Then, by Definition 3.3,

$$\frac{1}{m} \sum_{i=1}^{m} |x_i|^k \le \mathbb{E} |x_0|^k + n^{-\Omega(1)}.$$
(3.21)

Consequently, for any threshold t > 0,

$$\frac{1}{m}\sum_{i=1}^{m} \mathbf{1}\left[|x_i|^k > t\right] \le \frac{\mathbb{E}|x_0|^k + n^{-\Omega(1)}}{t}.$$

Substituting  $t = m^2$  and rearranging terms,

$$\mathbf{1}\left[\max_{i\in[m]}|x_i| \le m^{\frac{2}{k}}\right] \ge 1 - \frac{\mathbb{E}|x_0|^k + n^{-\Omega(1)}}{m}.$$

Since  $m = \Theta(n)$ , the right hand side is strictly positive for large enough n. Therefore,

$$\max_{i\in[m]}|x_i|=O_k(n^{\frac{2}{k}})$$

Since this holds for all  $k \in \mathbb{N}$ , we conclude that

$$\|\boldsymbol{x}\|_{\infty} = \max_{i \in [m]} |x_i| = n^{o(1)}.$$

In the case that  $\boldsymbol{x}$  is random, then Definition 3.3 instead implies that (3.21) holds with high probability, and we conclude by the same argument that for any c > 0, w.h.p.,  $\|\boldsymbol{x}\|_{\infty} = O(n^c)$ .  $\Box$ 

# 4 Objective Perturbation for Logistic Regression

In this section, we study the privacy-utility tradeoff for objective perturbation, applied to the problem of *logistic* regression. The version of the algorithm that we consider is still that of Algorithm 1, but we will instantiate it with a different loss function  $\ell$ . Before stating the main result of this section (Theorem 4.2), we briefly review the setup for logistic regression and comment on some slight differences in the notion of convergence that will arise in the theorem's conclusion.

**Logistic Regression Model.** We are given a dataset  $(\mathbf{X}, \mathbf{y})$  of n samples. In the worst case,  $(\mathbf{x}_i, y_i)$  are arbitrary points in  $\mathbb{R}^{d+1}$  with  $\|\mathbf{x}_i\| \leq R$  and  $y \in \{0, 1\}$ , but in the average case, there is a ground-truth coefficient vector  $\boldsymbol{\beta}^* \in \mathbb{R}^d$  independent of  $\mathbf{X}$  such that

$$\boldsymbol{y}|(\boldsymbol{X},\boldsymbol{\beta}^{\star}) \sim \operatorname{Bernoulli}(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star})).$$

Here,  $\rho(t) = \log(1 + e^t)$ , so that  $\rho'(t) = 1/(1 + e^{-x})$  is the well-known sigmoid function. For  $t \in \mathbb{R}^n$ , we define  $\rho(t) = \sum_{i=1}^n \rho(t_i) \in \mathbb{R}$  and  $\rho'(t) = (\rho'(t_1), \ldots, \rho'(t_n)) \in \mathbb{R}^n$ . The logistic loss of a coefficient vector  $\beta$  on a data point (x, y) is

$$\rho(\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle) - y \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle = \rho(-(2y-1)\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle),$$

which simplifies to  $\rho(\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle)$  if y = 0 or  $\rho(-\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle)$  if y = 1.

Notion of Convergence. Since the universality results applicable to logistic regression in the existing statistics literature currently have slower convergence rates than those for robust linear regression, it will be convenient to introduce the following variant  $-\rightarrow$  of the  $\rightarrow$  notation we introduced in Section 3.

**Definition 4.1** (Slow Pseudo-Lipschitz Convergence). Consider a random variable  $x_0 \in \mathbb{R}$  with finite moments of all orders and a random vector  $\boldsymbol{x} \in \mathbb{R}^m$  with  $m = \Theta(n)$ . We write  $\boldsymbol{x} \dashrightarrow x_0$ , if for all  $f : \mathbb{R} \to \mathbb{R}$  that are either bounded and continuous, or pseudo-Lipschitz of order 2,

$$\exists c > 0 \quad \text{such that} \quad \Pr\left[\left|\frac{1}{m}\sum_{i=1}^{m} f(x_i) - \mathbb{E}[f(x_0)]\right| \le e^{-(\log n)^c}\right] \ge 1 - O(n^{-c}).$$

Define  $X \dashrightarrow x_0$  analogously for  $x_0 \in \mathbb{R}^{\ell}$ , as in Definition 3.3.

We are now ready to state and prove the main theorem of this section.

**Theorem 4.2.** Let  $\hat{\beta}$  denote the output of Algorithm 1 with parameters  $R, \lambda, \nu > 0$  and instantiated with the logistic loss function:

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = \rho(\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle) - y \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle.$$

(a) (Privacy)  $\hat{\boldsymbol{\beta}}$  satisfies  $(2\varepsilon_{\rm DP}, (1 + e^{\varepsilon_{\rm DP}})\delta_{\rm DP})$ -differential privacy for any  $\varepsilon_{\rm DP} \ge 0$  and

$$\delta_{\rm DP} = \begin{cases} 2 \cdot \text{HockeyStick}(\tilde{\varepsilon}_{\rm DP}, \frac{LR}{\nu}) & \text{if } \hat{\varepsilon}_{\rm DP} \ge 0\\ (1 - e^{\hat{\varepsilon}_{\rm DP}}) + 2e^{\hat{\varepsilon}_{\rm DP}} \cdot \text{HockeyStick}\left(\frac{L^2 R^2}{2\nu^2}, \frac{LR}{\nu}\right) & \text{otherwise,} \end{cases}$$

where we set  $\tilde{\varepsilon}_{\rm DP} = \varepsilon_{\rm DP} - \log(1 + R^2/4\lambda)$  and  $\hat{\varepsilon}_{\rm DP} = \tilde{\varepsilon}_{\rm DP} - L^2 R^2/2\nu^2$ .

- (b) (Utility) Suppose the following hold for some  $\beta^{\star} \in \mathbb{R}^d$  as  $n \to \infty$  and  $d/n \to \delta$ :
  - (i)  $\mathbf{X} \in B_R(\mathbf{0})^n \subseteq \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y} \sim \text{Bernoulli}(\rho'(\mathbf{X}\boldsymbol{\beta}^{\star}))$ .
  - (ii) There exists a random variable  $\beta_0^{\star} \in \mathbb{R}$  such that  $\beta^{\star} \rightsquigarrow \beta_0^{\star}$ .

Suppose there exist  $\sigma^*, \alpha^*, \gamma^* > 0$  solving the following system of three scalar equations in three variables  $(\sigma, \alpha, \gamma)$ , which we write in terms of dummy variables  $Z_1, Z_2 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$  and  $\kappa^2 = \mathbb{E}(\beta_0^*)^2$  as

$$\sigma^{2} = \gamma^{2} \left( \frac{1}{\delta} \mathbb{E} \Big[ 2\rho'(-\kappa Z_{1})\rho' \big( \operatorname{prox}_{\gamma\rho}(\kappa \alpha Z_{1} + \sigma Z_{2}) \big)^{2} \Big] + \nu^{2} \right), \tag{4.3a}$$

$$\alpha = -\frac{1}{\delta} \mathbb{E}[2\rho''(-\kappa Z_1) \operatorname{prox}_{\gamma\rho} \left(\kappa \alpha Z_1 + \sigma Z_2\right)], \tag{4.3b}$$

$$\gamma = \frac{1}{\lambda\delta} \left( \delta - 1 + \mathbb{E} \left[ \frac{2\rho'(-\kappa Z_1)}{1 + \gamma\rho''(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_1 + \sigma Z_2))} \right] \right).$$
(4.3c)

The estimation error  $\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}$  satisfies, for  $Z, \xi_0 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$ ,

$$(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \widehat{\boldsymbol{\beta}}) \dashrightarrow \left(\beta_0^{\star}, \xi_0, \alpha^{\star} \beta_0^{\star} + \sqrt{(\sigma^{\star})^2 - (\gamma^{\star} \nu)^2} Z - \gamma^{\star} \nu \xi_0\right)$$

The difference  $\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}) - \rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})$  satisfies, for  $Z_1, Z_2 \stackrel{iid}{\sim} \mathcal{N}(0,1)$  and  $y_0|Z_1 \sim \text{Bernoulli}(\rho'(\kappa Z_1))$ ,

$$(\boldsymbol{X}\boldsymbol{\beta}^{\star},\rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})) \dashrightarrow (\kappa Z_1, \alpha^{\star}\kappa Z_1 + \sigma^{\star}Z_2 + y_0 - \operatorname{prox}_{\gamma^{\star}\rho}(\alpha^{\star}\kappa Z_1 + \sigma^{\star}Z_2 + \gamma^{\star}y_0)).$$

As in the case of robust linear regression (Theorem 3.7), part (a) follows from the results of Section 5, and combining Theorem 4.2(b) with Definition 3.3 yields several interesting corollaries. For example, for the *bias* of  $\hat{\beta}$ , we see that w.h.p.,

$$\frac{1}{d}\langle \widehat{\boldsymbol{\beta}}, \boldsymbol{\beta}^{\star} \rangle = \mathbb{E}[(\alpha^{\star}\beta_{0}^{\star} + \sqrt{(\sigma^{\star})^{2} - (\gamma^{\star}\nu)^{2}}Z + \gamma^{\star}\nu\xi_{0}) \cdot \beta_{0}^{\star}] \pm e^{-(\log n)^{\Omega(1)}}$$
$$= \alpha^{\star}\kappa^{2} \pm e^{-(\log n)^{\Omega(1)}}.$$

Therefore, we deduce that  $\hat{\beta}$  is positively correlated with  $\beta^*$ , which is intuitively what we would expect, and this correlation is captured by  $\alpha^*$ . Similarly, for the variance of  $\hat{\beta}$ , we see that w.h.p.,

$$\frac{1}{d} \|\widehat{\boldsymbol{\beta}} - \alpha^{\star} \boldsymbol{\beta}^{\star}\|^{2} = \mathbb{E} \left( \sqrt{(\sigma^{\star})^{2} - (\gamma^{\star} \nu)^{2}} Z + \gamma^{\star} \nu \xi_{0} \right)^{2} \pm e^{-(\log n)^{\Omega(1)}}$$
$$= (\sigma^{\star})^{2} \pm e^{-(\log n)^{\Omega(1)}}.$$

In total, we see that the mean squared error of  $\widehat{\beta}$  is, w.h.p.,

$$\frac{1}{d} \|\widehat{\beta} - \beta^{\star}\|^2 = (1 - \alpha^{\star})^2 \kappa^2 + (\sigma^{\star})^2 \pm e^{-(\log n)^{\Omega(1)}}.$$

Similar calculations can be carried out for the difference  $\rho'(\mathbf{X}\boldsymbol{\beta}^{\star}) - \rho'(\mathbf{X}\widehat{\boldsymbol{\beta}})$ . We validate these corollaries against simulated data in Figure 4. For more commentary on how to interpret Theorem 4.2 and Figure 4, we refer the reader to the discussion surrounding (the very similarly-worded) Theorem 3.7 and Figure 3 on robust linear regression, in Section 3.

## 4.1 Heuristic Derivation

Before we begin the utility proof, we give a non-rigorous, heuristic derivation to motivate the three equations in the statement of Theorem 4.2(b). The derivation is based on Theorem 1 of [SAH19], which determines the asymptotic behavior of  $\hat{\beta}$  for regularized logistic regression, provided that  $\beta^*$  has i.i.d. coordinates and the regularizer is a deterministic function of the form  $\sum_{j=1}^{d} f(\beta_j)$ .

In order to apply Theorem 1 of [SAH19], one first assumes that the following system of six equations in six variables has a unique solution  $(\alpha^*, \sigma^*, \gamma^*, \theta^*, \tau^*, r^*)$ . These equations are stated in terms of  $Z, Z_1, Z_2 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$  and  $\beta^* \stackrel{iid}{\sim} \beta_0^*$ , as follows.<sup>5</sup>

$$\kappa^2 \alpha = \mathbb{E} \Big[ \beta_0^* \operatorname{prox}_{\lambda \delta \sigma \tau f} \Big( \sigma \tau \left( \theta \beta_0^* + r \sqrt{\delta} Z \right) \Big) \Big], \tag{4.4a}$$

$$\gamma = \frac{\sqrt{\delta}}{r} \mathbb{E} \Big[ Z \operatorname{prox}_{\lambda\delta\sigma\tau f} \Big( \sigma\tau \left( \theta\beta_0^\star + r\sqrt{\delta}Z \right) \Big) \Big], \tag{4.4b}$$

$$\kappa^2 \alpha^2 + \sigma^2 = \mathbb{E} \bigg[ \operatorname{prox}_{\lambda \delta \sigma \tau f} \Big( \sigma \tau \left( \theta \beta_0^\star + r \sqrt{\delta} Z \right) \Big)^2 \bigg], \tag{4.4c}$$

$$\gamma^2 = \frac{1}{r^2} \mathbb{E} \Big[ 2\rho'(-\kappa Z_1) \left( \kappa \alpha Z_1 + \sigma Z_2 - \operatorname{prox}_{\gamma \rho} (\kappa \alpha Z_1 + \sigma Z_2) \right)^2 \Big], \tag{4.4d}$$

$$\theta \gamma = -\mathbb{E} \left[ 2\rho''(-\kappa Z_1) \operatorname{prox}_{\gamma \rho} \left( \kappa \alpha Z_1 + \sigma Z_2 \right) \right], \tag{4.4e}$$

$$1 - \frac{\gamma}{\sigma\tau} = \mathbb{E}\left[\frac{2\rho'(-\kappa Z_1)}{1 + \gamma\rho''(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_1 + \sigma Z_2))}\right].$$
(4.4f)

The conclusion of [SAH19], informally speaking, is that the coordinates of  $\hat{\beta}$  behave like

$$\operatorname{prox}_{\lambda\delta\sigma^{\star}\tau^{\star}f}\left(\sigma^{\star}\tau^{\star}\left(\theta^{\star}\beta_{0}^{\star}+r^{\star}\sqrt{\delta}Z\right)\right).$$
(4.5)

Next, observe that logistic regression with objective perturbation can be recast as regularized logistic regression with a "randomized regularizer," as follows. First, sample  $\xi_1, \ldots, \xi_d \stackrel{iid}{\sim} \xi_0$  where  $\xi_0 \sim \mathcal{N}(0, 1)$  and define the functions  $f_0, \ldots, f_d : \mathbb{R} \to \mathbb{R}$  as

$$f_j(\beta) = \frac{\lambda}{2}\beta^2 + \nu\xi_j\beta.$$

Then, the objective perturbation algorithm's output  $\hat{\beta}$  can be written as

$$\widehat{\boldsymbol{\beta}} = \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^d} \rho(\boldsymbol{X}\boldsymbol{\beta}) - \langle \boldsymbol{y}, \boldsymbol{X}\boldsymbol{\beta} \rangle + \sum_{j=1}^d f_j(\beta_j).$$

<sup>&</sup>lt;sup>5</sup>We have replaced  $\delta$  with  $1/\delta$  and  $\lambda$  with  $\lambda\delta$  to account for differences in notation and scaling in [SAH19].



Figure 4: Theorem 4.2's predictions of the error of Algorithm 1 with logistic loss. Estimation error refers to  $\frac{1}{d} \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}\|^2$ . Difference of  $\rho'$  refers to  $\frac{1}{n} \|\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}) - \rho'(\boldsymbol{X}\hat{\boldsymbol{\beta}})\|^2$ , which is related to the residual vector  $\boldsymbol{y} - \rho'(\boldsymbol{X}\hat{\boldsymbol{\beta}})$  since  $\boldsymbol{y} \sim \text{Bernoulli}(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}))$ . Curves correspond to theoretical predictions, and dots correspond to the mean over 1000 simulations of the algorithm on synthetic data with  $n \times d = 1000$ . In the left plots, the perturbation magnitude is  $\nu = 0$ , but in the right plots,  $\nu = 1/5$ . In all plots, the signal strength is  $\kappa = 1$ , and we consider  $\boldsymbol{\beta}^{\star} \sim \mathcal{N}(\mathbf{0}, \kappa^2 \boldsymbol{I}_d)$ , along with  $\boldsymbol{X} \sim \frac{1}{\sqrt{d}} \text{Uniform}(\{-1,+1\}^{n \times d})$  and  $\boldsymbol{y} \sim \text{Bernoulli}(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}))$ .

Therefore, even though Theorem 1 of [SAH19] requires the coordinate-wise application of a single, deterministic regularizer f, and the above expression uses d distinct, "randomized regularizers"  $f_1, \ldots, f_d$ , one might guess that substituting  $f = f_0$  into the above system of six equations and taking the expectation over the randomness of  $f_0$  might nevertheless lead to the equations in Theorem 4.2(b), which we will later show correctly describe  $\hat{\beta}$ . Indeed, this is the case, which we now show. First, observe that

$$\operatorname{prox}_{tf_0}(x) = \frac{1}{1+t}x + \frac{t}{1+t}\left(-\frac{\nu}{\lambda}\xi_0\right) = \frac{x-t\cdot\frac{\nu}{\lambda}\xi_0}{1+t}.$$
(4.6)

Substituting (4.6) into (4.4b) yields

$$\gamma = \frac{\sqrt{\delta}}{r} \mathbb{E}\left[ Z \cdot \frac{\sigma \tau(\theta \beta_0^* + r\sqrt{\delta}Z) - \delta \sigma \tau \nu \xi_0}{1 + \lambda \delta \sigma \tau} \right]$$

If the random variables inside the expectation  $(Z, \beta_0^*, \text{ and } \xi_0)$  are all independent, we have  $\mathbb{E}[Z\beta_0^*] = \mathbb{E}[Z\xi_0] = 0$ . We also have  $\mathbb{E}[Z^2] = 1$ . Thus, the above equation simplifies to

$$\gamma = \frac{\delta \sigma \tau}{1 + \lambda \delta \sigma \tau}.$$
(4.7)

Next, substituting (4.6) into (4.4a) yields

$$\kappa^2 \alpha = \mathbb{E}\left[\beta_0^{\star} \cdot \frac{\sigma \tau(\theta \beta_0^{\star} + r\sqrt{\delta}Z) - \delta \sigma \tau \nu \xi_0}{1 + \lambda \delta \sigma \tau}\right]$$

Once again,  $\mathbb{E}[\beta_0^* Z] = \mathbb{E}[\beta_0^* \xi_0] = 0$  and we know  $\mathbb{E}(\beta_0^*)^2 = \kappa^2$ , so the above equation simplifies to

$$\kappa^2 \alpha = \frac{\sigma \tau \theta \kappa^2}{1 + \lambda \delta \sigma \tau}.$$

Using (4.7) to eliminate  $\tau$  further simplifies this to

$$\alpha = \frac{\gamma \theta}{\delta}.\tag{4.8}$$

Finally, substituting (4.6) into (4.4c) yields

$$\kappa^2 \alpha^2 + \sigma^2 = \mathbb{E}\left[\left(\frac{\sigma\tau(\theta\beta + r\sqrt{\delta}Z) - \delta\sigma\tau\nu\xi_0}{1 + \lambda\delta\sigma\tau}\right)^2\right]$$

Again using the fact that pairwise products of Z,  $\beta_0^{\star}$ , and  $\xi_0$  vanish in expectation, this becomes

$$\kappa^2 \alpha^2 + \sigma^2 = \left(\frac{\sigma\tau}{1 + \lambda\delta\sigma\tau}\right)^2 \left(\theta^2 \,\mathbb{E}(\beta_0^\star)^2 + r^2\delta \,\mathbb{E}[Z^2] + (\nu\delta)^2 \,\mathbb{E}[\xi_0^2]\right).$$

Eliminating  $\tau$  with (4.7) and  $\theta$  with (4.8) yields

$$\kappa^2 \alpha^2 + \sigma^2 = \left(\frac{\gamma}{\delta}\right)^2 \left( \left(\frac{\alpha\delta}{\gamma}\right)^2 \cdot \mathbb{E}(\beta_0^{\star})^2 + \frac{r^2}{\delta} \mathbb{E}[Z^2] + (\nu\delta)^2 \mathbb{E}[\xi_0^2] \right)$$

and substituting  $\mathbb{E}(\beta_0^{\star})^2 = \kappa^2$  and  $\mathbb{E}Z^2 = \mathbb{E}\xi_0^2 = 1$  finally yields

$$\sigma^2 = \frac{\gamma^2 r^2}{\delta} + \gamma^2 \nu^2. \tag{4.9}$$

At this point, equations (4.7), (4.8), and (4.9) can be summarized as

$$\theta = \frac{\alpha \delta}{\gamma}, \qquad \tau = \frac{\gamma}{\delta \sigma (1 - \lambda \gamma)}, \qquad r = \sqrt{\frac{\delta \sigma^2}{\gamma^2} - \delta \nu^2}.$$

Substituting these expressions into (4.4d), (4.4e), (4.4f) yields the system of three equations in Theorem 4.2(b). Therefore, we should expect that for the solution  $\alpha^*, \sigma^*, \gamma^* > 0$ , the expression (4.5) that captures the behavior of the coordinates of  $\hat{\beta}$  should reduce to

$$\alpha^*\beta_0^* + \sqrt{(\sigma^*)^2 - \gamma^2 \nu^2} Z + \gamma \nu \xi_0.$$

This completes our heuristic derivation, which we emphasize again was not rigorous because we did not satisfy the technical assumptions of [SAH19], including a single, deterministic regularizer  $f : \mathbb{R} \to \mathbb{R}$  applied coordinate-wise, and a ground-truth vector  $\beta^*$  with i.i.d. coordinates.

## 4.2 Proof of Theorem 4.2

The overall structure of our proof of Theorem 4.2 mirrors analyses for non-private logistic regression estimators that have appeared in the literature [SAH19, Han24], but we make several changes to account for the differences between their setting and ours. The main differences include our analysis of the random linear perturbation term introduced for differential privacy, our weakened assumptions on  $\beta^*$  ([SAH19] assumed  $\beta^*$  to be random with i.i.d. coordinates, but we do not), and our conclusion that includes a description of  $\rho'(\boldsymbol{X}\beta^*) - \rho'(\boldsymbol{X}\hat{\beta})$ . Our proof has three steps.

- (1) Use universality to relate the perturbed logistic objective to one with a Gaussian design.
- (2) Rewrite this modified problem as a min-max optimization amenable to CGMT.
- (3) Use CGMT to analyze this min-max optimization with a Gaussian design matrix as  $n \to \infty$ .

For the first step, we will rely on recent universality results of [Han24] that they use to study non-private, regularized logistic regression. In their most general framing, these universality results apply to any estimator that can be expressed as the output of a certain class of *first-order* algorithms, or a limit thereof. As we will show, the objective perturbation algorithm for logistic regression falls in this class. This allows us to relate the behavior of the algorithm's output  $\hat{\beta} \in \mathbb{R}^d$ (and the corresponding predictions  $\rho'(X\hat{\beta}) \in [0,1]^n$ ) in the setting amenable to differential privacy where X has bounded entries to the setting in which the entries of X are unbounded, but precisely follow a Gaussian law.

For the second step, as before, we will take the Legendre transform of the logistic loss function in order to isolate the randomness of the design matrix X in a single, bilinear term. Compared to the case of robust linear regression in Section 3.2, some extra care will be needed to ensure that this bilinear term is independent of the mean function  $\psi$ .

For the third step, we will use CGMT, roughly following the template in [SAH19], to derive the asymptotic behavior of the estimation error and prediction error of  $\hat{\beta}$ . Unlike [SAH19], our analysis will account for the random linear perturbation term, and also explain the behavior of  $\rho'(\mathbf{X}\boldsymbol{\beta}^{\star}) - \rho'(\mathbf{X}\hat{\boldsymbol{\beta}})$ , which is related to the residual error. As before, our analysis will be made more challenging by the fact that we do not assume that  $\boldsymbol{\beta}^{\star}$  has i.i.d. coordinates.

One will notice that the order of steps (1), (2), and (3) is different than in our analysis of robust linear regression in Section 3.2. The reason we must apply universality before transforming the problem into a min-max optimization amenable to CGMT is that, as we will see, this transformation *itself* requires the Gaussianity assumption. This is also why we cannot simply rely on the universality results of [HS23] (which require the problem to already be expressed in "CGMT form") and must instead use the more recent results of [Han24].

#### 4.2.1 Step 1: GFOM Universality

The goal of this step is to prove the following lemma, a differentially private analogue of Theorem 4.3 in [Han24]. The following lemma is a universality result that relates the output of objective perturbation (Algorithm 1) with the logistic loss function to its output when we replace X with  $\frac{1}{\sqrt{d}}G$ , where G has the same shape as X but independent, standard Gaussian entries. Unlike X, whose rows have  $\ell^2$  norm bounded by R (as required for differential privacy), the entries of the matrix G are unbounded in the worst case. Despite this, the lemma shows that the coordinate-wise average of any pseudo-Lipschitz test function, applied to either the estimate  $\hat{\beta}$  or its predictions  $\rho'(X\hat{\beta})$ , is roughly the same regardless of whether we start with X or  $\frac{1}{\sqrt{d}}G$ .

**Lemma 4.10.** Suppose  $\mathbf{X} \in \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y}|(\boldsymbol{\beta}^{\star}, \mathbf{X}) \sim \text{Bernoulli}(\boldsymbol{\rho}'(\mathbf{X}\boldsymbol{\beta}^{\star}))$ for some  $\boldsymbol{\beta}^{\star} \in \mathbb{R}^{d}$  satisfying  $\|\boldsymbol{\beta}^{\star}\| \leq O(n)$ . Let  $\mathbf{G} \in \mathbb{R}^{n \times d}$  have standard Gaussian entries, and let  $\hat{\boldsymbol{\beta}}(\mathbf{A})$  denote the output of Algorithm 1 with the logistic loss function and design matrix  $\mathbf{A} \in \{\mathbf{X}, \frac{1}{\sqrt{d}}\mathbf{G}\}$ . Then, for any collection of O(1)-pseudo-Lipschitz functions  $\psi_{1i} : \mathbb{R}^{2} \to \mathbb{R}$  and  $\psi_{2j} : \mathbb{R} \to \mathbb{R}$  of order 2 for  $i \in [n]$  and  $j \in [d]$ , we have that w.h.p.,<sup>6</sup>

$$\begin{aligned} &\left|\frac{1}{n}\sum_{i=1}^{n} \left(\psi_{1i}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{x}_{i}, \widehat{\boldsymbol{\beta}}(\boldsymbol{X}) \rangle) - \psi_{1i}\left(\left\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}}, \boldsymbol{\beta}^{\star} \right\rangle, \left\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}}, \widehat{\boldsymbol{\beta}}\left(\frac{\boldsymbol{G}}{\sqrt{d}}\right) \right\rangle\right)\right)\right| \\ &+ \left|\frac{1}{d}\sum_{j=1}^{d} \left(\psi_{2j}(\widehat{\beta}_{j}(\boldsymbol{X})) - \psi_{2j}\left(\widehat{\beta}_{j}\left(\frac{\boldsymbol{G}}{\sqrt{d}}\right)\right)\right)\right| \leq e^{-(\log n)^{\Omega(1)}}.\end{aligned}$$

Proof. As in the proof of Theorem 4.3 of [Han24] for non-private logistic regression, we will define a suitably smoothed sequence of iterates  $\beta^{(t)} \in \mathbb{R}^d$ , indexed by  $t \in \mathbb{N}$ , that converge to  $\hat{\beta}$  as  $t \to \infty$ . We will then conclude by Theorem 2.13, the universality of general first order methods (GFOMs). Unlike prior work, our proof will account for the noise introduced by objective perturbation for differential privacy. We also extend prior work by establishing universality for the logits  $X\hat{\beta}$  in addition to  $\hat{\beta}$ . Besides these changes, our proof and notation borrow heavily from that of Theorem 4.3 of [Han24], with a few simplifications due to our focus on the  $\ell^2$  regularizer.

To begin, we recast the logistic regression model in its *latent-variable* form. To this end, consider the distribution over  $\mathbb{R}$  with CDF  $\rho'$ , called the *logistic distribution*, and define  $\varepsilon_1^{\star}, \ldots, \varepsilon_n^{\star} \stackrel{iid}{\sim}$  Logistic. Then, the assumption that  $\boldsymbol{y} \sim$  Bernoulli $(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}))$  can be rephrased as

$$\boldsymbol{y} = \boldsymbol{1} [\boldsymbol{X}\boldsymbol{\beta}^{\star} + \boldsymbol{\varepsilon}^{\star} > 0].$$

Similarly, the loss value  $\ell(\boldsymbol{\beta}; (\boldsymbol{x}_i, y_i)) = \rho(\langle \boldsymbol{x}_i, \boldsymbol{\beta} \rangle) - y_i \langle \boldsymbol{x}_i, \boldsymbol{\beta} \rangle$  incurred on the *i*<sup>th</sup> data point can be rephrased as  $L(\langle \boldsymbol{x}_i, \boldsymbol{\beta} \rangle, \langle \boldsymbol{x}_i, \boldsymbol{\beta}^{\star} \rangle; \varepsilon_i^{\star})$ , where the function  $L : \mathbb{R}^3 \to \mathbb{R}$  is given by

$$L(x, y; \varepsilon) = \rho(-(2 \cdot \mathbf{1}[y + \varepsilon > 0] - 1)x).$$

<sup>&</sup>lt;sup>6</sup>Our proof will actually show that the bound holds both in expectation and with high probability.

Since we plan to take derivatives of L with respect to both its first and second inputs (call these operations  $\partial_1$  and  $\partial_2$ ), it will be convenient to consider the following smoothed version of L, parameterized by  $\sigma > 0$ . Letting  $\varphi : \mathbb{R} \to [0,1]$  be any infinitely differentiable, monotonically increasing function such that  $\varphi(x) = 0$  for all  $x \leq -1$  and  $\varphi(x) = 1$  for all  $x \geq 1$ , set  $\varphi_{\sigma}(x) = \varphi(x/\sigma)$  and  $L_{\sigma}(x, y; \varepsilon) = \rho(-(2\varphi_{\sigma}(y+\varepsilon)-1)x)$ . Similarly, set  $\varphi_0(x) = \mathbf{1}[x > 0]$  and  $L_0(x, y; \varepsilon) = L(x, y; \varepsilon)$ .

Next, consider the  $\sigma$ -smoothed version of the pertubed objective:

$$\mathcal{L}_{\sigma}(\boldsymbol{\beta}) = \sum_{i=1}^{n} L_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta} \rangle, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle; \varepsilon_{i}^{\star}) + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^{2} + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle.$$

Its gradient is

$$\nabla \mathcal{L}_{\sigma}(\boldsymbol{\beta}) = \sum_{i=1}^{n} \boldsymbol{x}_{i} \partial_{1} L_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta} \rangle, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle; \varepsilon_{i}^{\star}) + \lambda \boldsymbol{\beta} + \nu \boldsymbol{\xi},$$

where

$$\partial_1 L_{\sigma}(x,y;\varepsilon) = -(2\varphi_{\sigma}(y+\varepsilon)-1)\rho'(-(2\varphi_{\sigma}(y+\varepsilon)-1)x) \in [-1,1]$$
  
$$\partial_1^2 L_{\sigma}(x,y;\varepsilon) = (2\varphi_{\sigma}(y+\varepsilon)-1)^2 \rho''(-(2\varphi_{\sigma}(y+\varepsilon)-1)x) \in [0,1/4].$$

For some step size  $\eta$ , define the following gradient descent iterates on the  $\sigma$ -smoothed version of the perturbed objective:

$$\boldsymbol{\beta}_{\sigma}^{(0)} = \mathbf{0}, \qquad \boldsymbol{\beta}_{\sigma}^{(t)} = \boldsymbol{\beta}_{\sigma}^{(t-1)} - \eta \nabla \mathcal{L}_{\sigma}(\boldsymbol{\beta}_{\sigma}^{(t-1)}).$$

We now derive several useful properties of  $\beta_{\sigma}^{(t)}$ , such as its behavior as  $t \to \infty$ ,  $\sigma \to 0$  and upper bounds on  $\|\beta_{\sigma}^{(t)}\|$ ,  $\|\mathbf{X}\beta_{\sigma}^{(t)}\|_{\infty}$ . The argument for these upper bounds will be similar in spirit to the proof of Lemma 3.14 that we already saw and will, at various times, require bounding the effect of the noise introduced for differential privacy.

(a) (Behavior as  $t \to \infty$ ) In the limit, the iterates converge to  $\widehat{\beta}_{\sigma} = \operatorname{argmin}_{\beta \in \mathbb{R}^d} \mathcal{L}_{\sigma}(\beta)$ . To prove this, we subtract the equation  $\eta \nabla \mathcal{L}_{\sigma}(\widehat{\beta}_{\sigma}) = \mathbf{0}$  from the equation defining  $\beta_{\sigma}^{(t)}$ :

$$\boldsymbol{\beta}_{\sigma}^{(t)} - \widehat{\boldsymbol{\beta}}_{\sigma} = \left(\boldsymbol{\beta}_{\sigma}^{(t-1)} - \eta \nabla \mathcal{L}_{\sigma}(\boldsymbol{\beta}_{\sigma}^{(t-1)})\right) - \left(\widehat{\boldsymbol{\beta}}_{\sigma} - \eta \nabla \mathcal{L}_{\sigma}(\widehat{\boldsymbol{\beta}}_{\sigma})\right).$$

Observe that both invocations of  $\nabla \mathcal{L}_{\sigma}$  in the above equation contain an term of the form  $-\eta \nu \boldsymbol{\xi}$  due to objective perturbation, but these terms are canceled out by the subtraction. Thus, the remainder of this part of the proof is identical to the non-private case. We include it only for the sake of completeness:

We apply the mean value theorem to the  $\partial_1 L_{\sigma}$  terms in  $\nabla \mathcal{L}_{\sigma}$ , and use the fact that  $\partial_1^2 L_{\sigma}$  takes values in [0, 1/4] to obtain  $\mathbf{c} \in [0, 1/4]^n$  such that

$$\boldsymbol{\beta}_{\sigma}^{(t)} - \widehat{\boldsymbol{\beta}}_{\sigma} = \left( (1 - \eta \lambda) \boldsymbol{I} - \eta \sum_{i=1}^{n} c_i \boldsymbol{x}_i \boldsymbol{x}_i^{\top} \right) (\boldsymbol{\beta}_{\sigma}^{(t-1)} - \widehat{\boldsymbol{\beta}}_{\sigma}).$$

Since  $\boldsymbol{X}$  follows a subgaussian design, with probability  $1-e^{-\Omega(n)}$ , we have  $\boldsymbol{0} \leq \eta \sum_{i=1}^{n} c_i \boldsymbol{x}_i \boldsymbol{x}_i^{\top} \leq O(\eta) \boldsymbol{I}$ . Therefore, there exists some choice of step size  $\eta = \Theta(1/(1+\lambda))$  such that iterating the above equation for  $t = 1, 2, \ldots$  yields, with probability  $1 - e^{-\Omega(n)}$ ,

$$\|\boldsymbol{\beta}_{\sigma}^{(t)} - \widehat{\boldsymbol{\beta}}_{\sigma}\| \le e^{-\Omega(t)} \|\widehat{\boldsymbol{\beta}}_{\sigma}\|.$$

(b) (Behavior as  $\sigma \to 0$ ) We will analyze the behavior of the function  $\mathcal{L}_{\sigma}(\beta) - \mathcal{L}(\beta)$ . Clearly, the subtraction cancels out the objective perturbation term, so this part of the proof is also identical to the non-private case. We include it only for the sake of completeness:

The outputs of  $\varphi_{\sigma}$  and  $\varphi_0$  only differ on inputs smaller than  $\sigma$  in absolute value, and when they differ, they differ by at most 1. Using this observation, followed by Cauchy-Schwarz,

$$\begin{aligned} |\mathcal{L}_{\sigma}(\boldsymbol{\beta}) - \mathcal{L}_{0}(\boldsymbol{\beta})| &= \left| \sum_{i=1}^{n} (L_{\sigma} - L_{0}) (\langle \boldsymbol{x}_{i}, \boldsymbol{\beta} \rangle, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \varepsilon_{i}) \right| \\ &\leq 2 \sum_{i=1}^{n} |\langle \boldsymbol{x}_{i}, \boldsymbol{\beta} \rangle| \cdot |(\varphi_{\sigma} - \varphi_{0}) (\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle + \varepsilon_{i})| \\ &\leq 2 ||\boldsymbol{X}\boldsymbol{\beta}|| \left( \sum_{i=1}^{n} \mathbf{1} [\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle + \varepsilon_{i} \in [-\sigma, +\sigma]] \right)^{1/2}. \end{aligned}$$

Since the Logistic PDF takes values between 0 and 1/4, the terms of the parenthesized sum are independent Bernoulli random variables, each with parameter at most  $2\sigma \cdot 1/4$ . Consequently, standard concentration inequalities (e.g. Bernstein's) show that for any fixed constant D > 0, with probability  $1 - O(n^{-D})$ , the above sum is at most  $O(\sigma n + \log n)$ . We also have with probability  $1 - e^{-\Omega(n)}$  that  $\|\mathbf{X}\| = O(1)$ . Thus, with probability  $1 - O(n^{-D})$ ,

$$|\mathcal{L}_{\sigma}(\boldsymbol{\beta}) - \mathcal{L}_{0}(\boldsymbol{\beta})| \leq O((\sigma n + \log n)^{1/2} \|\boldsymbol{\beta}\|).$$

We apply this inequality to  $\hat{\beta}$  and  $\hat{\beta}_{\sigma}$ , which minimize  $\mathcal{L}$  and  $\mathcal{L}_{\sigma}$ , respectively. By the  $\lambda$ -strong convexity of these functions and Lemma 2.20, we deduce that with probability  $1 - O(n^{-D})$ ,

$$\|\widehat{\boldsymbol{\beta}}_{\sigma} - \widehat{\boldsymbol{\beta}}\| \le O((\sigma n + \log n)^{1/4} (\|\widehat{\boldsymbol{\beta}}\| + \|\widehat{\boldsymbol{\beta}}_{\sigma}\|)^{1/2}).$$

(c) (Worst-Case Upper Bounds on  $\|\beta_{\sigma}^{(t)}\|$ ,  $\|\widehat{\beta}_{\sigma}\|$ ) From the definition of  $\beta_{\sigma}^{(t)}$  and the fact that  $\partial_1 L$  takes values in [-1, +1],

$$\|\boldsymbol{\beta}_{\sigma}^{(t)}\| \le (1 - \eta\lambda) \|\boldsymbol{\beta}_{\sigma}^{(t-1)}\| + \eta \|\boldsymbol{X}^{\top}\|_{\infty \to 2} + \eta\nu \|\boldsymbol{\xi}\| = O(t \cdot (\|\boldsymbol{X}^{\top}\|_{\infty \to 2} + \|\boldsymbol{\xi}\|)).$$

The fact that  $\mathcal{L}_{\sigma}(\widehat{\beta}_{\sigma}) = \Omega(\|\widehat{\beta}_{\sigma}\|^2)$  is smaller than  $\mathcal{L}_{\sigma}(\mathbf{0}) = O(n)$  implies  $\|\widehat{\beta}_{\sigma}\| \leq O(\sqrt{n})$ .

(d) (High Probability Upper Bound on  $\|\beta_{\sigma}^{(t)}\|$ ,  $\|\widehat{\beta}_{\sigma}\|$ ) As in our analysis of  $t \to \infty$ , applying the mean value theorem to  $\partial_1 L_{\sigma}$  in the equation  $\nabla \mathcal{L}_{\sigma}(\widehat{\beta}_{\sigma}) = \mathbf{0}$  yields  $\mathbf{c} \in [0, 1/4]^n$  such that

$$\left(\lambda \boldsymbol{I} + \sum_{i=1}^{n} c_{i} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{\top}\right) \widehat{\boldsymbol{\beta}}_{\sigma} + \sum_{i=1}^{n} \boldsymbol{x}_{i} \partial_{1} L_{\sigma}(\boldsymbol{0}, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \varepsilon_{i}^{\star}) + \nu \boldsymbol{\xi} = \boldsymbol{0}$$

Therefore, with probability  $1 - e^{-\Omega(n)}$ ,

$$\|\widehat{\boldsymbol{\beta}}_{\sigma}\| \leq O\left(\left\|\sum_{i=1}^{n} \boldsymbol{x}_{i} \,\partial_{1} L_{\sigma}(\boldsymbol{0}, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \varepsilon_{i}^{\star})\right\| + \|\boldsymbol{\xi}\|\right).$$

The second term measures the norm of the random perturbation vector  $\boldsymbol{\xi}$ , and is clearly at most  $O(\sqrt{n \log n})$  with probability  $1 - O(n^{-D})$ , for any fixed constant D > 0. The same bound can be established for the first term, and the proof of this is identical to the non-private case. We include it only for the sake of completeness: Define the function  $F_{\sigma}(t) = \mathbb{E}[\varphi_{\sigma}(t+\varepsilon_0^{\star})]$ , where  $\varepsilon_0^{\star} \sim \text{Logistic.}$  Using the fact that  $\partial_1 L_{\sigma}(0, y, \varepsilon) = -(2\varphi_{\sigma}(y+\varepsilon)-1)$ , followed by the triangle inequality,

$$\begin{split} & \left\|\sum_{i=1}^{n} \boldsymbol{x}_{i} \partial_{1} L(\boldsymbol{0}, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \varepsilon_{i}^{\star})\right\|^{2} \\ &= \frac{1}{4} \left\|\sum_{i=1}^{n} \boldsymbol{x}_{i} (2\varphi_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle + \varepsilon_{i}^{\star}) - 1)\right\|^{2} \\ &\leq O\left(\underbrace{\left\|\sum_{i=1}^{n} \boldsymbol{x}_{i} (\varphi_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle + \varepsilon_{i}^{\star}) - F_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle))\right\|^{2}}_{S_{1}} + \underbrace{\left\|\sum_{i=1}^{n} \boldsymbol{x}_{i} (2F_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle) - 1)\right\|^{2}}_{S_{2}}\right). \end{split}$$

We bound the terms  $S_1$  and  $S_2$  separately. First,  $S_1 \leq O(n \log n)$  with probability  $1 - O(n^{-D})$ for any fixed constant D > 0 because for each fixed coordinate  $j \in [d]$ , the  $j^{\text{th}}$  coordinates of the summands across  $i \in [n]$  are centered, independent variables with subgaussian norm  $O(n^{-1/2})$ . To bound  $S_2$ , we use the fact that  $|F'_{\sigma}(t)| \leq ||\varphi'||_{\infty} \leq O(1)$  for any  $\sigma \geq 0$  and  $t \in \mathbb{R}$ . Thus, for each fixed coordinate  $j \in [d]$ , if we let  $\boldsymbol{x}_{i,(-j)} \in \mathbb{R}^d$  denote the version of  $\boldsymbol{x}_i$ with the  $j^{\text{th}}$  coordinate zeroed out, then there is some scalar  $\tau_j$  with  $|\tau_j| = O(1)$  such that

$$F_{\sigma}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle) = F_{\sigma}(\langle \boldsymbol{x}_{i,(-j)}, \boldsymbol{\beta}^{\star} \rangle) + \tau_{j} x_{ij} \beta_{j}^{\star}.$$

Consequently, expanding  $S_2$ ,

$$S_{2} \leq O\left(\sum_{j=1}^{d} \left(\sum_{i=1}^{n} x_{ij}\right)^{2} + \sum_{j=1}^{d} \left(\sum_{i=1}^{n} x_{ij} F_{\sigma}(\langle \boldsymbol{x}_{i,(-j)}, \boldsymbol{\beta}^{\star} \rangle)\right)^{2} + \sum_{j=1}^{d} \left(\sum_{i=1}^{n} x_{ij}^{2}\right)^{2} (\beta_{j}^{\star})^{2}\right).$$

Using standard subgaussian concentration inequalities on X, the fact that  $x_{ij}$  is independent of  $F_{\sigma}(\langle \boldsymbol{x}_{i,(-j)}, \boldsymbol{\beta}^{\star} \rangle)$ , and the fact that  $\|\boldsymbol{\beta}^{\star}\|_{2} = O(n)$ , we conclude that  $S_{2} \leq O(n \log n)$  with probability  $1 - O(n^{-D})$  for any constant D > 0.

In summary, we have shown that with probability  $1 - O(n^{-D})$ ,

$$\|\widehat{\boldsymbol{\beta}}_{\sigma}\| \leq O(\sqrt{n\log n}).$$

Combining this with the bound from part (a), with probability  $1 - O(n^{-D})$ ,

$$\|\boldsymbol{\beta}_{\sigma}^{(t)}\| \le O(\sqrt{n\log n}).$$

(e) (High Probability Upper Bound on  $\|\boldsymbol{X}\boldsymbol{\beta}_{\sigma}^{(t)}\|_{\infty}$ ) Fix an index  $k \in [n]$ , and let  $\mathcal{L}_{\sigma;[-k]}$  be the modified loss function that omits the term corresponding to the  $k^{\text{th}}$  sample  $\boldsymbol{x}_k$ . As before, define iterates

$$\boldsymbol{\beta}_{\sigma;[-k]}^{(0)} = \mathbf{0}, \qquad \boldsymbol{\beta}_{\sigma;[-k]}^{(t)} = \boldsymbol{\beta}_{\sigma;[-k]}^{(t-1)} - \eta \nabla \mathcal{L}_{\sigma;[-k]}(\boldsymbol{\beta}_{\sigma;[-k]}^{(t-1)}).$$

Clearly, these converge to  $\widehat{\beta}_{\sigma;[-k]} = \operatorname{argmin}_{\beta \in \mathbb{R}^d} \mathcal{L}_{\sigma;[-k]}(\beta)$  at the same rate that we proved  $\beta_{\sigma}^{(t)}$  converges to  $\widehat{\beta}_{\sigma}$ . We proceed by subtracting the equations defining  $\beta_{\sigma}^{(t)}$  and  $\beta_{\sigma;[-k]}^{(t)}$ , which cancels out the effect of linear objective perturbation. Therefore, the remainder of this part of the proof is identical to the non-private case. We include it only for the sake of completeness:

Similar to before, we see that there exist  $c \in [0, 1/4]^n$  and  $c_0 \in [-1, 1]$  such that

$$\boldsymbol{\beta}_{\sigma}^{(t)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t)} = \left( (1 - \eta \lambda) \boldsymbol{I} - \eta \sum_{i=1}^{n} c_i \boldsymbol{x}_i \boldsymbol{x}_i^{\mathsf{T}} \right) (\boldsymbol{\beta}_{\sigma}^{(t-1)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t-1)}) - \eta c_0 \boldsymbol{x}_k.$$

Notice that the final  $-\eta c_0 \boldsymbol{x}_k$  term arises because the gradient of  $\mathcal{L}_{\sigma}$  includes a term for the loss of  $(\boldsymbol{x}_k, y_k)$ , whereas  $\mathcal{L}_{\sigma;[-k]}$  does not. This implies that for appropriate  $\eta = \Theta(1)$ , with probability  $1 - e^{-\Omega(n)}$ ,

$$\|\boldsymbol{\beta}_{\sigma}^{(t)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t)}\| \le \frac{1}{2} \|\boldsymbol{\beta}_{\sigma}^{(t-1)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t-1)}\| + O(1).$$

Iterating the above equation for t = 1, 2, ... and taking a union bound over  $k \in [n]$  yields, with probability  $1 - e^{-\Omega(n)}$ ,

$$\|\boldsymbol{\beta}_{\sigma}^{(t)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t)}\| \le O(1).$$

A nearly identical argument yields, with probability  $1 - e^{-\Omega(n)}$ ,

$$\|\widehat{\boldsymbol{\beta}}_{\sigma} - \widehat{\boldsymbol{\beta}}_{\sigma;[-k]}\| \le O(1).$$

Next, by the triangle inequality,

$$\|\boldsymbol{X}\boldsymbol{\beta}_{\sigma}^{(t)}\|_{\infty} \leq \underbrace{\max_{k \in [n]} |\langle \boldsymbol{x}_{k}, \boldsymbol{\beta}_{\sigma;[-k]}^{(t)} \rangle|}_{S_{1}} + \underbrace{\max_{k \in [n]} |\langle \boldsymbol{x}_{k}, \boldsymbol{\beta}_{\sigma}^{(t)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t)} \rangle|}_{S_{2}}.$$

We bound  $S_1$  and  $S_2$  separately. First, for  $S_1$ , observe that  $\boldsymbol{x}_k$  and  $\boldsymbol{\beta}_{\sigma;[-k]}^{(t)}$  are independent by definition, and recall that we showed in part (a) that for any constant D > 0, with probability  $1 - O(n^{-D})$ , we have  $\|\boldsymbol{\beta}_{\sigma;[-k]}^{(t)}\| \leq O(\sqrt{n \log n})$ . Therefore, Lemma 2.18 implies that  $S_1 \leq O(\log n)$ . Next, for  $S_2$ , recall that we just proved  $\|\boldsymbol{\beta}_{\sigma}^{(t)} - \boldsymbol{\beta}_{\sigma;[-k]}^{(t)}\| \leq O(1)$ . Therefore, by Cauchy-Schwarz,  $S_2 \leq O(\sqrt{\log n})$ . Similar arguments hold with  $\hat{\boldsymbol{\beta}}_{\sigma;[-k]}$  in place of  $\boldsymbol{\beta}_{\sigma;[k]}^{(t)}$ . We conclude that for any constant D > 0, with probability  $1 - O(n^{-D})$ ,

$$\|\boldsymbol{X}\boldsymbol{\beta}_{\sigma}^{(t)}\|_{\infty} + \|\boldsymbol{X}\widehat{\boldsymbol{\beta}}_{\sigma}\|_{\infty} \leq O(\log n).$$

With our various bounds on the iterates  $\beta^{(t)}$  in hand, we now are ready to invoke GFOM universality (Theorem 2.13), which is the main step of the proof. We will construct the required GFOM using the following functions  $G_{\sigma,i} : \mathbb{R}^2 \to \mathbb{R}$  for  $\sigma \ge 0$  and  $i \in [n]$ :

$$G_{\sigma,i}(\boldsymbol{u}) = \partial_1 L_{\sigma}([u_1]_M, u_2; \varepsilon_i),$$

where  $M \leq O(\log n)$  is the bound on  $\|\boldsymbol{X}\boldsymbol{\beta}_{\sigma}^{(t)}\|_{\infty} + \|\boldsymbol{X}\widehat{\boldsymbol{\beta}}_{\sigma}\|_{\infty}$  from part (e). Let  $G_{\sigma} : \mathbb{R}^{m \times 2} \to \mathbb{R}$  be the separable function that applies  $G_{\sigma,i}$  to its  $i^{\text{th}}$  input. A direct calculation reveals that  $|\partial_1 \partial_2 L_{\sigma}(x, y; \varepsilon)| \leq O(|x|/\sigma)$ , from which it follows that

$$\sup_{\varepsilon \in \mathbb{R}^n} \max_{i \in [n]} \|G_{\sigma,i}\|_{\operatorname{Lip}} + |G_{\sigma,i}(\mathbf{0})| \le O\left(1 + \frac{M}{\sigma}\right).$$

Now define the GFOM iterates  $\boldsymbol{u}^{(t)} \in \mathbb{R}^{n \times 2}$  and  $\boldsymbol{v}^{(t)} \in \mathbb{R}^d$  by  $\boldsymbol{u}^{(0)} = \boldsymbol{0}$  and  $\boldsymbol{v}^{(0)} = \boldsymbol{0}$  and

$$\boldsymbol{u}^{(t)} = \boldsymbol{X} \left[ \boldsymbol{v}^{(t-1)} \mid \boldsymbol{\beta}^{\star} \right] \in \mathbb{R}^{n \times 2},$$
  
$$\boldsymbol{v}^{(t)} = \boldsymbol{X}^{\top} [-\eta G_{\sigma}(\boldsymbol{u}^{(t)})] + (1 - \eta \lambda) \boldsymbol{v}^{(t-1)} - \eta \nu \boldsymbol{\xi} \in \mathbb{R}^{d}.$$

Clearly, in the event  $E_{\sigma}$  that  $\|\boldsymbol{X}\boldsymbol{\beta}_{\sigma}^{(t)}\|_{\infty} \leq M$  at each step, the truncation  $[\cdot]_{M}$  in the definition of G never takes effect, so we have for all  $t \in \mathbb{N}$  that  $\boldsymbol{v}^{(t)}$  coincides with  $\boldsymbol{\beta}_{\sigma}^{(t)}$  and that the first column of  $\boldsymbol{u}^{(t)}$  coincides with  $\boldsymbol{X}\boldsymbol{\beta}_{\sigma}^{(t)}$ . By GFOM universality (Theorem 2.13), for any O(1)-pseudo-Lipschitz functions  $\psi_{1i}: \mathbb{R}^2 \to \mathbb{R}$  and  $\psi_{2j}: \mathbb{R} \to \mathbb{R}$  of order O(1) for  $i \in [n]$  and  $j \in [d]$ , there exists a constant C > 0 such that if  $\boldsymbol{G} \in \mathbb{R}^{n \times d}$  has independent standard Gaussian entries,

$$\mathbb{E}\left[\left|\frac{1}{n}\sum_{i=1}^{n}\left(\psi_{1i}(\langle \boldsymbol{x}_{i},\boldsymbol{\beta}^{\star}\rangle,\langle \boldsymbol{x}_{i},\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{X})\rangle)-\psi_{1i}\left(\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}},\boldsymbol{\beta}^{\star}\rangle,\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}},\boldsymbol{\beta}_{\sigma}^{(t)}\left(\frac{\boldsymbol{G}}{\sqrt{d}}\right)\rangle\right)\right)\right|\mathbf{1}_{E_{\sigma}}\right]\\+\mathbb{E}\left[\left|\frac{1}{d}\sum_{j=1}^{d}\left(\psi_{2j}(\beta_{\sigma,j}^{(t)}(\boldsymbol{X}))-\psi_{2j}\left(\beta_{\sigma,j}^{(t)}\left(\frac{\boldsymbol{G}}{\sqrt{d}}\right)\right)\right)\right|\mathbf{1}_{E_{\sigma}}\right]\leq(C\sigma^{-1}\log(n))^{Ct^{3}}n^{-1/(Ct^{3})}.$$

To conclude the proof, all that remains is to derive a version of the above inequality without the indicator functions  $\mathbf{1}_{E_{\sigma}}$ , and then consider the limit as  $t \to \infty$  and  $\sigma \to 0$ . The remainder of the proof is similar to that of the non-private case; the exception is that we often invoke the bounds from parts (a), (b), (c), (d), (e), whose proofs took into consideration the effect of objective perturbation at various points.

To remove the indicator functions, first recall that we showed in part (e) above that  $\Pr[E_{\sigma}] = 1 - O(n^{-D})$  for any constant D > 0. Thus, it suffices to show that even in the *worst* case, which includes the unlikely, probability  $O(n^{-D})$  event that  $E_{\sigma}$  does *not* occur, that all of the sums in the above display are bounded in expectation by  $O(n^c)$  for some constant c > 0 that does not depend on D. Indeed, taking D > c, this crude bound will lead to only a mild additive  $O(n^c \cdot n^{-D}) = n^{-\Omega(1)}$  increase to the GFOM universality error bound of  $(C\sigma^{-1}\log(n))^{Ct^3}n^{-1/(Ct^3)}$ .

To derive such a bound, observe that if the functions  $\psi_{1i}, \psi_{2j}$  are pseudo-Lispchitz of order 2, then for either  $A \in \{X, \frac{1}{\sqrt{d}}G\}$ ,

$$\mathbb{E}\left|\frac{1}{d}\sum_{j=1}^{d}\psi_{1j}(\langle \boldsymbol{a}_{i},\boldsymbol{\beta}^{\star}\rangle,\langle \boldsymbol{a}_{i},\boldsymbol{\beta}_{\sigma,j}^{(t)}(\boldsymbol{A})\rangle)\right| + \mathbb{E}\left|\frac{1}{d}\sum_{j=1}^{d}\psi_{2j}(\boldsymbol{\beta}_{\sigma,j}^{(t)}(\boldsymbol{A}))\right| \\
\leq O\left(1 + \frac{\mathbb{E}\|\boldsymbol{A}\boldsymbol{\beta}^{\star}\|^{2}}{n} + \frac{\mathbb{E}\|\boldsymbol{A}\cdot\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\|^{2}}{n} + \frac{\mathbb{E}\|\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\|^{2}}{n}\right).$$

By part (c), we have  $\|\boldsymbol{A}\boldsymbol{\beta}^{\star}\|^{2} + \|\boldsymbol{A}\cdot\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\|^{2} + \|\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\|^{2} \leq (tn\|\boldsymbol{A}\|\|\boldsymbol{\xi}\|)^{O(1)}$ , which is clearly  $n^{O(1)}$  in expectation, as desired.

We have thus removed the indicator functions  $\mathbf{1}_{E_{\sigma}}$ . Specifically, we have shown that for any collection of O(1)-pseudo-Lipschitz functions  $\psi_{1i}, \psi_{2j} : \mathbb{R} \to \mathbb{R}$  of order 2 and any constant D > 0, there exists a constant C > 0 such that with probability  $1 - Cn^{-D}$ :

$$\begin{split} & \mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} \left( \psi_{1i}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{X}) \rangle) - \psi_{1i} \left( \left\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}}, \boldsymbol{\beta}^{\star} \right\rangle, \left\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}}, \boldsymbol{\beta}_{\sigma}^{(t)} \left( \frac{\boldsymbol{G}}{\sqrt{d}} \right) \right\rangle \right) \right) \right| \\ & + \mathbb{E} \left| \frac{1}{d} \sum_{j=1}^{d} \left( \psi_{2j}(\beta_{\sigma,j}^{(t)}(\boldsymbol{X})) - \psi_{2j} \left( \beta_{\sigma,j}^{(t)} \left( \frac{\boldsymbol{G}}{\sqrt{d}} \right) \right) \right) \right| \leq (C\sigma^{-1}\log n)^{Ct^{3}} n^{-1/(Ct^{3})} + Cn^{-D}. \end{split}$$

Next, we consider what happens as  $t \to \infty$ . Indeed, using the definition of order-2 pseudo-

Lipschitzness, followed by Cauchy-Schwarz, for both  $A \in \{X, \frac{1}{\sqrt{d}}G\}$ ,

$$\begin{aligned} \left| \frac{1}{n} \sum_{i=1}^{n} \left( \psi_{1i}(\langle \boldsymbol{a}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{a}_{i}, \boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A}) \rangle) - \psi_{1i}(\langle \boldsymbol{a}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{a}_{i}, \widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A}) \rangle) \right) \right| \\ & \leq O\left( \frac{1}{n} \sum_{i=1}^{n} |\langle \boldsymbol{a}_{i}, \boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A}) - \widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A}) \rangle| \cdot (1 + |\langle \boldsymbol{a}_{i}, \boldsymbol{\beta}^{\star} \rangle| + |\langle \boldsymbol{a}_{i}, \boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A}) \rangle| + |\langle \boldsymbol{a}_{i}, \widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A}) \rangle|) \right) \\ & \leq O\left( \frac{1}{n} \cdot \|\boldsymbol{A}\| \cdot \|\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A}) - \widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A})\| \cdot (\sqrt{n} + \|\boldsymbol{A}\| \cdot (\|\boldsymbol{\beta}^{\star}\| + \|\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\| + \|\widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A})\|)) \right). \end{aligned}$$

Similarly, for either  $A \in \{X, \frac{1}{\sqrt{d}}G\}$ ,

$$\begin{aligned} \left| \frac{1}{d} \sum_{j=1}^{d} \left( \psi_{2j}(\beta_{\sigma,j}^{(t)}(\boldsymbol{A})) - \psi_{2j}(\widehat{\beta}_{\sigma,j}(\boldsymbol{A})) \right) \right| \\ & \leq O\left( \frac{1}{d} \sum_{j=1}^{d} |\beta_{\sigma,j}^{(t)}(\boldsymbol{A}) - \widehat{\beta}_{\sigma,j}(\boldsymbol{A})| \cdot (1 + |\beta_{\sigma,j}^{(t)}(\boldsymbol{A})| + |\widehat{\beta}_{\sigma,j}(\boldsymbol{A})|) \right) \\ & \leq O\left( \frac{1}{n} \cdot \|\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A}) - \widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A})\| \cdot (\sqrt{n} + \|\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\| + \|\widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A})\|) \right). \end{aligned}$$

As before, we consider separately the cases in which  $E_{\sigma}$  does and does not occur. In the case that  $E_{\sigma}$  does occur, our high-probability bound  $\|\beta_{\sigma}^{(t)}(\mathbf{A}) - \widehat{\beta}_{\sigma}(\mathbf{A})\| \le e^{-\Omega(t)} \|\widehat{\beta}_{\sigma}(\mathbf{A})\|$  from part (a) holds, as does our high probability bound  $\|\widehat{\beta}_{\sigma}(\mathbf{A})\| + \|\beta_{\sigma}^{(t)}(\mathbf{A})\| \le O(\sqrt{n \log n})$  from part (d), as well as the bound  $\|\mathbf{A}\| \le O(1)$ . In the case that  $E_{\sigma}$  does not occur, our crude bounds on all the aforementioned terms of order  $n^{O(1)}$  from part (c) still hold. Therefore, taking expectations of both sides of the above two displays, we see that for any constant D > 0, there exists a constant C > 0 such that

$$\mathbb{E}\left|\frac{1}{n}\sum_{i=1}^{n}\left(\psi_{1i}(\langle \boldsymbol{a}_{i},\boldsymbol{\beta}^{\star}\rangle,\langle \boldsymbol{a}_{i},\boldsymbol{\beta}_{\sigma}^{(t)}(\boldsymbol{A})\rangle)-\psi_{1i}(\langle \boldsymbol{a}_{i},\boldsymbol{\beta}^{\star}\rangle,\langle \boldsymbol{a}_{i},\widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A})\rangle)\right)\right| \\
+\mathbb{E}\left|\frac{1}{d}\sum_{j=1}^{d}\left(\psi_{2j}(\beta_{\sigma,j}^{(t)}(\boldsymbol{A}))-\psi_{2j}(\widehat{\boldsymbol{\beta}}_{\sigma,j}(\boldsymbol{A}))\right)\right| \leq e^{-t/C}(\log n)^{C}+Cn^{-D}.$$

Just as the above inequality uses the bound from part (a) (along with parts (c) (d), (e)) to determine the effect of taking  $t \to \infty$ , so too can we use the bound from part (b) (along with parts (c) (d), (e)) to determine the effect of taking  $\sigma \to 0$ :

$$\mathbb{E}\left|\frac{1}{n}\sum_{i=1}^{n} \left(\psi_{1i}(\langle \boldsymbol{a}_{i},\boldsymbol{\beta}^{\star}\rangle,\langle \boldsymbol{a}_{i},\widehat{\boldsymbol{\beta}}_{\sigma}(\boldsymbol{A})\rangle)-\psi_{1i}(\langle \boldsymbol{a}_{i},\boldsymbol{\beta}^{\star}\rangle,\langle \boldsymbol{a}_{i},\widehat{\boldsymbol{\beta}}(\boldsymbol{A})\rangle)\right)\right| \\ + \mathbb{E}\left|\frac{1}{d}\sum_{j=1}^{d} \left(\psi_{2j}(\widehat{\beta}_{\sigma,j}(\boldsymbol{A}))-\psi_{2j}(\widehat{\beta}_{j}(\boldsymbol{A}))\right)\right| \leq \left(\sigma+\frac{\log n}{n}\right)^{1/4} (\log n)^{C}+Cn^{-D}.$$

At this point, we have shown that for any constant D > 0, there exists a constant C > 0 such that

$$\begin{split} & \mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} \left( \psi_{1i}(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{x}_{i}, \widehat{\boldsymbol{\beta}}(\boldsymbol{X}) \rangle) - \psi_{1i} \left( \left\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}}, \boldsymbol{\beta}^{\star} \right\rangle, \left\langle \frac{\boldsymbol{g}_{i}}{\sqrt{d}}, \widehat{\boldsymbol{\beta}}\left( \frac{\boldsymbol{G}}{\sqrt{d}} \right) \right\rangle \right) \right) \right| \\ & + \mathbb{E} \left| \frac{1}{d} \sum_{j=1}^{d} \left( \psi_{2j}(\widehat{\beta}_{j}(\boldsymbol{X})) - \psi_{2j}\left( \widehat{\beta}_{j}\left( \frac{\boldsymbol{G}}{\sqrt{d}} \right) \right) \right) \right| \\ & \leq (C\sigma^{-1}\log n)^{Ct^{3}} n^{-1/(Ct^{3})} + e^{-t/C} (\log n)^{C} + \left( \sigma + \frac{\log n}{n} \right)^{1/4} (\log n)^{C} + Cn^{-D}. \end{split}$$

Taking  $t = (\log n)^{1/8}$  and  $\sigma = e^{-(\log n)^{1/8}}$  simplifies the entire bound to  $e^{-\Omega((\log n)^{1/8})}$ . Passing from a bound in expectation to a high-probability bound via Markov's inequality concludes the proof.

## 4.2.2 Step 2: Legendre Transform

The goal of this step is to prove the following lemma, which relates the output of Algorithm 1 with logistic loss to the min-max optimization of a certain random variable. As before, the lemma technically holds in a worst-case sense, and only later will the randomness of X become necessary.

In order to state the lemma, first recall from Section 2 that  $\rho^*$  denotes the convex conjugate of  $\rho$ . It is 4-strongly convex, and its negation is commonly known as the *binary entropy function* (in nats):

$$-\rho^{\star}(s) = \begin{cases} s \log\left(\frac{1}{s}\right) + (1-s) \log\left(\frac{1}{1-s}\right) & \text{if } 0 < s < 1, \\ 0 & \text{if } s = 0 \text{ or } s = 1. \end{cases}$$

**Lemma 4.11.** Fix any  $\beta^* \in \mathbb{R}^d$  and let  $\widehat{\beta}$  be the output of Algorithm 1 when instantiated with

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = \rho'(\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle) - y \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle.$$

Let  $Q_{\boldsymbol{\beta}, \boldsymbol{v}}$  be the following random variable indexed by  $\boldsymbol{\beta} \in \mathbb{R}^d$  and  $\boldsymbol{v} \in [0, 1]^n$ :

$$Q_{\boldsymbol{\beta},\boldsymbol{v}} = \langle \boldsymbol{X}\boldsymbol{\beta}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle - \rho^{\star}(\boldsymbol{v}).$$

Let  $\widehat{\boldsymbol{v}} = \rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})$ . Then,  $(\widehat{\boldsymbol{\beta}}, \widehat{\boldsymbol{v}})$  is the unique point in  $\mathbb{R}^d \times [0, 1]^n$  satisfying

$$\max_{\boldsymbol{v}\in[0,1]^n} Q_{\widehat{\boldsymbol{\beta}},\boldsymbol{v}} = \min_{\boldsymbol{\beta}\in\mathbb{R}^d} Q_{\boldsymbol{\beta},\widehat{\boldsymbol{v}}}.$$

We call it the saddle point or Nash equilibrium of  $Q_{\beta,v}$ .

*Proof.* Algorithm 1 simply computes

$$\widehat{\boldsymbol{\beta}} = \operatorname*{argmin}_{\boldsymbol{\beta} \in \mathbb{R}^d} \rho(\boldsymbol{X}\boldsymbol{\beta}) - \langle \boldsymbol{y}, \boldsymbol{X}\boldsymbol{\beta} \rangle + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle.$$

We shall take the *Legendre transform* of  $\rho$  in the above expression. In our case, this simply amounts to applying the following identity regarding  $\rho$ , which is valid for all  $t \in \mathbb{R}^n$ :

$$ho(oldsymbol{t}) = \max_{oldsymbol{v}\in[0,1]^n} ig\langle oldsymbol{t},oldsymbol{v}ig
angle - 
ho^\star(oldsymbol{v}).$$

The maximum is achieved iff  $\boldsymbol{v} = \rho'(\boldsymbol{t})$ . Setting  $\boldsymbol{t} = \boldsymbol{X}\boldsymbol{\beta}$ ,

$$\widehat{oldsymbol{eta}} = rgmin_{oldsymbol{eta} \in \mathbb{R}^d} \max_{oldsymbol{v} \in [0,1]^n} ig\langle oldsymbol{X} oldsymbol{eta}, oldsymbol{v} - oldsymbol{y} ig
angle - 
ho^{\star}(oldsymbol{v}) + rac{\lambda}{2} \|oldsymbol{eta}\|^2 + 
u \langle oldsymbol{\xi}, oldsymbol{u} 
angle.$$

Note that this objective function is simply  $Q_{\boldsymbol{u},\boldsymbol{v}}$ . Given  $\boldsymbol{\beta}$ , the maximum over  $\boldsymbol{v}$  is achieved iff  $\boldsymbol{v} = \rho'(\boldsymbol{X}\boldsymbol{\beta})$ , which coincides with  $\hat{\boldsymbol{v}}$  when  $\boldsymbol{\beta} = \hat{\boldsymbol{\beta}}$ . The terms  $\frac{\lambda}{2} \|\boldsymbol{\beta}\|^2$  and  $-\rho^*(\boldsymbol{v})$  ensure that  $Q_{\boldsymbol{\beta},\boldsymbol{v}}$  is  $\lambda$ -strongly convex in  $\boldsymbol{\beta} \in \mathbb{R}^d$  and 4-strongly concave in  $\boldsymbol{v} \in [0,1]^n$ , so we conclude by the minimax theorem that  $(\hat{\boldsymbol{\beta}}, \hat{\boldsymbol{v}})$  is the unique saddle point of  $Q_{\boldsymbol{\beta},\boldsymbol{v}}$ .

Next, we apply a trick from [SAH19] to ensure that the random bilinear term is independent of the remaining terms—a prerequisite for applying CGMT. In order to state the lemma, given a ground-truth coefficient vector  $\beta^* \in \mathbb{R}^d$ , define the linear subspaces

$$\operatorname{span}(\boldsymbol{\beta}^{\star}) = \{ t \boldsymbol{\beta}^{\star} : t \in \mathbb{R} \}, \qquad \operatorname{span}(\boldsymbol{\beta}^{\star})^{\perp} = \{ \boldsymbol{\beta} \in \mathbb{R}^{d} : \langle \boldsymbol{\beta}, \boldsymbol{\beta}^{\star} \rangle = 0 \}.$$

Given vectors  $\beta^{\parallel} \in \operatorname{span}(\beta^{\star})$  and  $\beta^{\perp} \in \operatorname{span}(\beta^{\star})^{\perp}$ , we will sometimes refer to their sum as

$$oldsymbol{eta} = oldsymbol{eta}^{\parallel} + oldsymbol{eta}^{\perp}$$
 ,

Conversely, given a vector  $\boldsymbol{\beta} \in \mathbb{R}^d$ , we will sometimes refer to its projections onto  $\operatorname{span}(\boldsymbol{\beta}^{\star})$  and  $\operatorname{span}(\boldsymbol{\beta}^{\star})^{\perp}$  by  $\boldsymbol{\beta}^{\parallel}$  and  $\boldsymbol{\beta}^{\perp}$ , respectively. Note that the following lemma relies on the rotational invariance provided by Gaussianity.

**Lemma 4.12.** Suppose that  $\mathbf{X} = \frac{1}{\sqrt{d}}\mathbf{G}$  with entries  $G_{ij} \stackrel{iid}{\sim} \mathcal{N}(0,1)$ . Let  $\mathbf{f} = \mathbf{G} \cdot \frac{\beta^*}{\|\beta^*\|}$ , let  $\mathbf{H}$  be an independent copy of  $\mathbf{G}$ , and let  $Q'_{\beta,v}$  be the following random variable indexed by  $\boldsymbol{\beta} \in \mathbb{R}^d$  and  $v \in [0,1]^n$ :

$$Q'_{\boldsymbol{\beta},\boldsymbol{v}} = \frac{1}{\sqrt{d}} \langle \boldsymbol{H} \boldsymbol{\beta}^{\perp}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{\|\boldsymbol{\beta}^{\parallel}\|}{\sqrt{d}} \langle \boldsymbol{f}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle - \rho^{\star}(\boldsymbol{v}).$$

Then, for any closed sets  $S_{\beta} \subseteq \mathbb{R}^d$  and  $S_{v} \subseteq [0,1]^n$ , the random variable  $\min_{\beta \in S_{\beta}} \max_{v \in S_{v}} Q_{\beta,v}$ has the same distribution as  $\min_{\beta \in S_{\beta}} \max_{v \in S_{v}} Q'_{\beta,v}$ . Moreover,  $f \sim \mathcal{N}(0, I_n)$  and

$$oldsymbol{y} |oldsymbol{f} \sim ext{Bernoulli}igg( 
ho'igg( rac{\|oldsymbol{eta}^{\star}\|}{\sqrt{d}}oldsymbol{f} igg) igg)$$

*Proof.* Substituting  $\mathbf{X} = \frac{1}{\sqrt{d}} \mathbf{G}$  and  $\boldsymbol{\beta} = \boldsymbol{\beta}^{\perp} + \boldsymbol{\beta}^{\parallel}$  into definition of  $Q_{\boldsymbol{\beta}, \boldsymbol{v}}$  yields

$$Q_{\boldsymbol{\beta},\boldsymbol{v}} = \frac{1}{\sqrt{d}} \langle \boldsymbol{G} \boldsymbol{\beta}^{\perp}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{1}{\sqrt{d}} \langle \boldsymbol{G} \boldsymbol{\beta}^{\parallel}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^2 + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle - \rho^{\star}(\boldsymbol{v}).$$

Observe that  $G\beta^{\parallel}$  and  $y \sim \text{Bernoulli}(\rho'(\frac{1}{\sqrt{d}}G\beta^{\star}))$  both depend on G only through the projections of its rows onto  $\text{span}(\beta^{\star})$ . By Gaussianity, these projections are independent of their projections onto  $\text{span}(\beta^{\star})^{\perp}$ , justifying the replacement of  $G\beta^{\perp}$  with  $H\beta^{\perp}$ . The facts that  $f \sim \mathcal{N}(0, I_n)$  and that  $G\beta^{\parallel} = \|\beta^{\parallel}\|f$  and  $G\beta^{\star} = \|\beta^{\star}\|f$  are also standard properties of multivariate Gaussians.  $\Box$ 

#### 4.2.3 Step 3: CGMT Analysis

In this step, we analyze the random variable  $Q'_{u,v}$  using CGMT (Theorem 2.8), roughly following the strategy of [SAH19].

As in the case of robust linear regression, our proof differs from past work in several ways. On the one hand, our proof is simpler, partly because it focuses on  $\ell^2$  regularization and partly because we avoid introducing several extraneous scalar- and vector-valued variables. On the other hand, our proof is somewhat more complex, partly due to our consideration of the perturbation vector  $\boldsymbol{\xi}$  introduced for differential privacy, and partly due to differences in the statement of our assumptions and conclusion, as well as our consideration of  $\rho'(\boldsymbol{X}\hat{\boldsymbol{\beta}})$  in addition to  $\hat{\boldsymbol{\beta}}$ .

To begin, we recall and analyze the auxiliary random variable to which CGMT (Theorem 2.8) pertains. We only define this random variable in the case that  $\boldsymbol{X} = \frac{1}{\sqrt{d}}\boldsymbol{G}$  and  $\boldsymbol{f} = \boldsymbol{G} \cdot \frac{\boldsymbol{\beta}^{\star}}{\|\boldsymbol{\beta}^{\star}\|}$ , in which case by Lemma 4.12 we have that  $\boldsymbol{f} \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_n)$  and

$$oldsymbol{y} \sim ext{Bernoulli}igg( 
ho'igg( rac{\|oldsymbol{eta}^{\star}\|}{\sqrt{d}}oldsymbol{f} igg) igg)$$

In terms of  $\boldsymbol{g} \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_d)$  and  $\boldsymbol{h} \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_n)$ , the auxiliary random variable is

$$Q_{\boldsymbol{\beta},\boldsymbol{v}}'' = \frac{\|\boldsymbol{\beta}^{\perp}\|}{\sqrt{d}} \langle \boldsymbol{h}, \boldsymbol{v} - \boldsymbol{y} \rangle - \frac{\|\boldsymbol{v} - \boldsymbol{h}\|}{\sqrt{d}} \langle \boldsymbol{g}, \boldsymbol{\beta}^{\perp} \rangle + \frac{\|\boldsymbol{\beta}^{\parallel}\|}{\sqrt{d}} \langle \boldsymbol{f}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{\lambda}{2} \|\boldsymbol{\beta}\|^{2} + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta} \rangle - \rho^{\star}(\boldsymbol{v}).$$

Eventually, we will use CGMT (Theorem 2.8) to relate  $Q''_{\beta,v}$  via  $Q'_{\beta,v}$  to  $Q_{\beta,v}$ .

**Lemma 4.13.** Define  $\alpha^*, \sigma^*, \gamma^* > 0$  as in Theorem 4.2(b), and consider the pair  $(\widetilde{\beta}, \widetilde{v})$  with

$$\begin{split} \widetilde{\boldsymbol{\beta}} &= \alpha^{\star} \boldsymbol{\beta}^{\star} + \sqrt{(\sigma^{\star})^{2} - (\gamma^{\star} \nu)^{2}} \boldsymbol{g} - \gamma^{\star} \nu \boldsymbol{\xi}, \\ \widetilde{\boldsymbol{v}} &= \boldsymbol{y} + \frac{1}{\gamma^{\star}} \left( \alpha^{\star} \kappa \boldsymbol{f} + \sigma^{\star} \boldsymbol{h} - \operatorname{prox}_{\gamma^{\star} \rho} (\alpha^{\star} \kappa \boldsymbol{f} + \sigma^{\star} \boldsymbol{h} + \gamma^{\star} \boldsymbol{y}) \right) \\ &= \boldsymbol{y} + (-1)^{\boldsymbol{y}} \odot \rho' (\operatorname{prox}_{\gamma^{\star} \rho} ((-1)^{\boldsymbol{y}} \odot (\alpha^{\star} \kappa \boldsymbol{f} + \sigma^{\star} \boldsymbol{h}))). \end{split}$$

Then, under the assumptions of Theorem 4.2(b), there exists a constant  $c^* \in \mathbb{R}$  such that w.h.p.

- The function  $\boldsymbol{\beta} \mapsto Q_{\boldsymbol{\beta}, \widetilde{\boldsymbol{v}}}''$  is  $\lambda$ -strongly convex in  $\boldsymbol{\beta} \in \mathbb{R}^d$ ,
- The function  $\boldsymbol{v} \mapsto Q''_{\widetilde{\boldsymbol{\beta}}, \boldsymbol{v}}$  is 4-strongly concave in  $\boldsymbol{v} \in [0, 1]^n$ ,
- The pair  $(\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}})$  satisfies

$$c^{\star}n - n^{1-\Omega(1)} \leq \min_{\boldsymbol{\beta} \in \mathbb{R}^d} Q_{\boldsymbol{\beta}, \widetilde{\boldsymbol{v}}}' \leq Q_{\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}}}' \leq \max_{\boldsymbol{v} \in [0, 1]^n} Q_{\widetilde{\boldsymbol{\beta}}, \boldsymbol{v}}' \leq c^{\star}n + n^{1-\Omega(1)}.$$

We call  $(\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}})$  an approximate saddle point of  $Q''_{\boldsymbol{\beta}, \boldsymbol{v}}$ .

Proof. We first remark that in this proof, we will repeatedly use the identities  $\operatorname{prox}_{\gamma\rho}(x+\gamma) = -\operatorname{prox}_{\rho}(-x)$  and  $\operatorname{prox}'_{\gamma\rho}(x) = 1/(1+\gamma\rho''(\operatorname{prox}_{\gamma\rho}(x)))$ , which are proved in [SAH19]. To show that  $(\widetilde{\beta}, \widetilde{v})$  is an approximate saddle point of  $Q''_{\beta,v}$ , it suffices to show that each of  $\widetilde{\beta}$  and  $\widetilde{v}$  is an approximately best response to the other, and that  $|Q''_{\widetilde{\beta},\widetilde{v}} - c^*n| \leq n^{1-\Omega(1)}$ . Formally, we say that  $\widetilde{\beta}$  is an approximate best response to  $\widetilde{v}$  if

$$Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}'' \leq \min_{\boldsymbol{u} \in \mathbb{R}^d} Q_{\boldsymbol{\beta},\widetilde{\boldsymbol{v}}}'' + n^{1-\Omega(1)}.$$

Similarly, we say that  $\widetilde{v}$  is an approximate best response to  $\widetilde{\beta}$  if

$$Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}'' \geq \max_{\boldsymbol{v} \in [0,1]^n} Q_{\widetilde{\boldsymbol{\beta}},\boldsymbol{v}}'' - n^{1-\Omega(1)}$$

In order to prove these inequalities, we will check that certain derivatives approximately vanish.

(a) ( $\tilde{\boldsymbol{\beta}}$  is an approximate best response to  $\tilde{\boldsymbol{v}}$ ) We first study  $\min_{\boldsymbol{\beta} \in \mathbb{R}^d} Q_{\boldsymbol{\beta}, \tilde{\boldsymbol{v}}}'$ . If we can show that  $Q_{\boldsymbol{\beta}, \tilde{\boldsymbol{v}}}'$  is a  $\lambda$ -strongly convex function of  $\boldsymbol{\beta}$ , then minimizing  $Q_{\boldsymbol{\beta}, \tilde{\boldsymbol{v}}}''$  over  $\boldsymbol{\beta} \in \mathbb{R}^d$  reduces to finding a point  $\boldsymbol{\beta} \in \mathbb{R}^d$  at which the gradient  $\nabla_{\boldsymbol{\beta}} Q_{\boldsymbol{\beta}, \tilde{\boldsymbol{v}}}''$  has small norm. To this end, observe that the term  $\frac{\lambda}{2} \|\boldsymbol{\beta}\|^2$  in  $Q_{\boldsymbol{\beta}, \tilde{\boldsymbol{v}}}''$  is  $\lambda$ -strongly convex in  $\boldsymbol{\beta}$ . The term  $-\frac{\|\tilde{\boldsymbol{v}}-\boldsymbol{y}\|}{\sqrt{d}} \langle \boldsymbol{g}, \boldsymbol{\beta}^{\perp} \rangle$  is a linear function of  $\boldsymbol{\beta}$ , and hence convex. Finally, the term  $\frac{\|\boldsymbol{\beta}^{\perp}\|}{\sqrt{d}} \langle \boldsymbol{h}, \tilde{\boldsymbol{v}} - \boldsymbol{y} \rangle$  will be convex, as well, if we can show that  $\langle \boldsymbol{h}, \tilde{\boldsymbol{v}} - \boldsymbol{y} \rangle$  is positive with high probability. Quantities like  $\langle \boldsymbol{h}, \tilde{\boldsymbol{v}} - \boldsymbol{y} \rangle$  can be easily computed in the limit by combining the definitions of  $\boldsymbol{\beta}$  and  $\tilde{\boldsymbol{v}}$  in this lemma's statement with the assumptions on  $\boldsymbol{f}, \boldsymbol{g}, \boldsymbol{h}, \boldsymbol{\beta}^*, \boldsymbol{y},$  and  $\boldsymbol{\xi}$ . Indeed, by the definition of pseudo-Lipschitz convergence (Definition 3.3), if  $f_0, g_0, h_0 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$  and  $y_0 | f_0 \sim \text{Bernoulli}(\rho'(\kappa f_0))$ , then w.h.p.,

$$\begin{split} \frac{1}{d} \langle \widetilde{\boldsymbol{\beta}}, \boldsymbol{\beta}^{\star} \rangle &= \frac{1}{\sqrt{d}} \| \widetilde{\boldsymbol{\beta}}^{\parallel} \| = \alpha^{\star} \kappa \pm n^{-\Omega(1)}, \\ &\quad \frac{1}{\sqrt{d}} \| \widetilde{\boldsymbol{\beta}}^{\perp} \| = \sigma^{\star} \pm n^{-\Omega(1)}, \\ &\quad \frac{1}{d} \langle \boldsymbol{g}, \widetilde{\boldsymbol{\beta}}^{\parallel} \rangle = 0 \pm n^{-\Omega(1)}, \\ &\quad \frac{1}{d} \langle \boldsymbol{g}, \widetilde{\boldsymbol{\beta}}^{\perp} \rangle = \sqrt{(\sigma^{\star})^2 - (\gamma^{\star} \nu)^2} \pm n^{-\Omega(1)}, \\ &\quad \frac{1}{\sqrt{d}} \| \widetilde{\boldsymbol{v}} - \boldsymbol{y} \| = \sqrt{\frac{1}{\delta}} \mathbb{E}[\rho'(\operatorname{prox}_{\gamma^{\star}\rho}((-1)^{y_0}(\alpha^{\star} \kappa f_0 + \sigma^{\star} h_0)))^2] \pm n^{-\Omega(1)}, \\ &\quad \frac{1}{d} \langle \boldsymbol{h}, \widetilde{\boldsymbol{v}} - \boldsymbol{y} \rangle = \frac{1}{\gamma^{\star}\delta} \left( \sigma^{\star} - \mathbb{E}[\operatorname{prox}_{\gamma^{\star}\rho}(\alpha^{\star} \kappa f_0 + \sigma^{\star} h_0 + \gamma^{\star} y_0) h_0] \right) \pm n^{-\Omega(1)}, \\ &\quad \frac{1}{d} \langle \boldsymbol{f}, \widetilde{\boldsymbol{v}} - \boldsymbol{y} \rangle = \frac{1}{\gamma^{\star}\delta} \left( \alpha^{\star} \kappa - \mathbb{E}[\operatorname{prox}_{\gamma^{\star}\rho}(\alpha^{\star} \kappa f_0 + \sigma^{\star} h_0 + \gamma^{\star} y_0) f_0] \right) \pm n^{-\Omega(1)}. \end{split}$$

We can simplify the expression for  $\|\tilde{\boldsymbol{v}} - \boldsymbol{y}\|$  using the fact that  $\Pr[y_0 = 0 \mid f_0] = \rho'(-\kappa f_0)$  and  $\Pr[y_0 = 1 \mid f_0] = \rho'(\kappa f_0)$ , followed by equation (4.3a) (see Theorem 4.2(b)):

$$\begin{aligned} \frac{1}{\sqrt{d}} \|\widetilde{\boldsymbol{v}} - \boldsymbol{y}\| &= \sqrt{\frac{1}{\delta} \mathbb{E}[2\rho'(-\kappa f_0)\rho'(\operatorname{prox}_{\gamma^*\rho}(\alpha^*\kappa f_0 + \sigma^*h_0))^2] \pm n^{-\Omega(1)}} \\ &= \sqrt{\left(\frac{\sigma^*}{\gamma^*}\right)^2 - \nu^2} \pm n^{-\Omega(1)}. \end{aligned}$$

Next, we can simplify the expression for  $\langle \boldsymbol{f}, \boldsymbol{\tilde{v}} - \boldsymbol{y} \rangle$  using Stein's lemma, followed by the formula for  $\operatorname{prox}_{\gamma^{\star}\rho}'$  given in Section 2, and finally equations (4.3b) and (4.3c):

$$\frac{1}{d} \langle \boldsymbol{f}, \widetilde{\boldsymbol{v}} - \boldsymbol{y} \rangle = \frac{\kappa}{\gamma^* \delta} \bigg( 1 - \mathbb{E}[2\rho'(-\kappa Z_1) \operatorname{prox}'_{\gamma^* \rho}(\alpha^* \kappa Z_1 + \sigma^* Z_2)] \\ + \frac{1}{\alpha} \mathbb{E}[2\rho''(-\kappa f_0) \operatorname{prox}_{\gamma^* \rho}(\alpha^* \kappa f_0 + \sigma^* h_0)] \bigg) \pm n^{-\Omega(1)} \\ = -\alpha^* \kappa \lambda \pm n^{-\Omega(1)}.$$

Similarly, we can simplify the expression for  $\langle \boldsymbol{h}, \tilde{\boldsymbol{v}} - \boldsymbol{y} \rangle$  using Stein's lemma, followed by the formula for  $\operatorname{prox}_{\gamma^{\star}\rho}'$  given in Section 2, and finally equation (4.3c):

$$\frac{1}{d} \langle \boldsymbol{h}, \widetilde{\boldsymbol{v}} - \boldsymbol{y} \rangle = \frac{\sigma^{\star}}{\gamma^{\star} \delta} \left( 1 - \mathbb{E}[2\rho'(-\kappa Z_1) \operatorname{prox}'_{\gamma^{\star} \rho}(\alpha^{\star} \kappa Z_1 + \sigma^{\star} Z_2)] \right) \pm n^{-\Omega(1)}$$
$$= \sigma^{\star} \left( \frac{1}{\gamma^{\star}} - \lambda \right) \pm n^{-\Omega(1)}.$$

Again by equation (4.3c), we see that  $1/\gamma^* > \lambda^*$ , so the quantity above is positive with high probability, which establishes that  $Q''_{\beta,\tilde{v}}$  is  $\lambda$ -strongly convex in  $\beta$  with high probability. Therefore, all that remains is to evaluate its gradient at  $\tilde{\beta}$ . To this end, we calculate

$$\nabla_{\boldsymbol{\beta}^{\perp}} Q_{\boldsymbol{\beta},\boldsymbol{v}}'' = \frac{1}{\sqrt{d}} \frac{\boldsymbol{\beta}^{\perp}}{\|\boldsymbol{\beta}^{\perp}\|} \langle \boldsymbol{h}, \boldsymbol{v} - \boldsymbol{y} \rangle - \frac{\|\boldsymbol{v} - \boldsymbol{y}\|}{\sqrt{d}} \boldsymbol{g}^{\perp} + \lambda \boldsymbol{\beta}^{\perp} + \nu \boldsymbol{\xi}^{\perp}.$$

Evaluating at  $(\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}})$  and substituting our expressions for  $\langle \boldsymbol{h}, \widetilde{\boldsymbol{v}} - \boldsymbol{y} \rangle$ ,  $\|\widetilde{\boldsymbol{\beta}}^{\perp}\|$ , and  $\|\widetilde{\boldsymbol{v}} - \boldsymbol{y}\|$  yields

$$\nabla_{\boldsymbol{\beta}^{\perp}} Q_{\boldsymbol{\widetilde{\beta}},\boldsymbol{\widetilde{v}}}'' = \left(\frac{1}{\gamma^{\star}} \pm n^{-\Omega(1)}\right) \boldsymbol{\widetilde{\beta}}^{\perp} - \left(\sqrt{\left(\frac{\sigma^{\star}}{\gamma^{\star}}\right)^2 - \nu^2} \pm n^{-\Omega(1)}\right) \boldsymbol{g}^{\perp} + \nu \boldsymbol{\xi}^{\perp}.$$

By substituting the definition of  $\tilde{\boldsymbol{\beta}}$  in this lemma's statement and using the fact that for any constant c > 0, we have  $\|\tilde{\boldsymbol{\beta}}\| + \|\boldsymbol{g}\| + \|\boldsymbol{\xi}\| \le O(n^{\frac{1}{2}+c})$  w.h.p. (Lemma 3.6), we see that the above gradient clearly has  $\ell^2$  norm

$$\|\nabla_{\boldsymbol{\beta}^{\perp}} Q_{\boldsymbol{\widetilde{\beta}},\boldsymbol{\widetilde{v}}}''\| \le n^{\frac{1}{2} - \Omega(1)}$$

Similarly, we calculate

$$abla_{oldsymbol{eta}} Q_{oldsymbol{eta},oldsymbol{v}} = rac{1}{\sqrt{d}} rac{oldsymbol{eta}^{\parallel}}{\|oldsymbol{eta}^{\parallel}\|} \langle oldsymbol{f},oldsymbol{v}-oldsymbol{y}
angle + \lambda oldsymbol{eta}^{\parallel}.$$

Evaluating at  $(\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}})$  and substituting our expressions for  $\langle \boldsymbol{f}, \widetilde{\boldsymbol{v}} - \boldsymbol{y} \rangle$  and  $\|\boldsymbol{\beta}^{\parallel}\|$  yields

$$\|\nabla_{\boldsymbol{\beta}^{\parallel}} Q_{\boldsymbol{\widetilde{\beta}},\boldsymbol{\widetilde{v}}}''\| \le n^{\frac{1}{2} - \Omega(1)}$$

Since we have shown that the derivatives of  $Q''_{\beta,\tilde{v}}$  with respect to  $\beta^{\parallel}$  and  $\beta^{\perp}$  both have norm  $n^{\frac{1}{2}-\Omega(1)}$ , by  $\lambda$ -strong convexity in  $\beta$ , we have with high probability that

$$Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}'' \leq \min_{\boldsymbol{\beta} \in \mathbb{R}^d} Q_{\boldsymbol{\beta},\widetilde{\boldsymbol{v}}}' + n^{1-\Omega(1)}$$

(b) ( $\tilde{\boldsymbol{v}}$  is an approximate best response to  $\tilde{\boldsymbol{\beta}}$ ) We study  $\max_{\boldsymbol{v}\in[0,1]^n} Q''_{\tilde{\boldsymbol{\beta}},\boldsymbol{v}}$ . Analogously to the previous part, we first verify 4-strong concavity with respect to  $\boldsymbol{v}$ . To this end, observe that the term  $-\rho^*(\boldsymbol{v})$  in  $Q''_{\tilde{\boldsymbol{\beta}},\boldsymbol{v}}$  is 4-strongly concave in  $\boldsymbol{v}$ . The term  $\frac{\|\tilde{\boldsymbol{\beta}}\|}{\sqrt{d}}\langle \boldsymbol{h}, \boldsymbol{v} - \boldsymbol{y} \rangle$  is a linear function of  $\boldsymbol{v}$ , and hence concave. Finally, the term  $-\frac{\|\boldsymbol{v}-\boldsymbol{y}\|}{\sqrt{d}}\langle \boldsymbol{g}, \tilde{\boldsymbol{\beta}} \rangle$  is concave, as well, since we have already shown that  $\langle \boldsymbol{g}, \tilde{\boldsymbol{\beta}} \rangle$  is positive with high probability. Now, all that remains is to evaluate the gradient of  $Q''_{\tilde{\boldsymbol{\beta}},\boldsymbol{v}}$  at  $\tilde{\boldsymbol{v}}$ . To this end, we calculate

$$\nabla_{\boldsymbol{v}} Q_{\boldsymbol{\beta},\boldsymbol{v}}'' = \frac{\|\boldsymbol{\beta}^{\perp}\|}{\sqrt{d}} \boldsymbol{h} - \frac{1}{\sqrt{d}} \frac{\boldsymbol{v} - \boldsymbol{y}}{\|\boldsymbol{v} - \boldsymbol{y}\|} \langle \boldsymbol{g}, \boldsymbol{\beta} \rangle + \frac{\|\boldsymbol{\beta}^{\parallel}\|}{\sqrt{d}} \boldsymbol{f} - (\rho^{\star})'(\boldsymbol{v}).$$

Evaluating at  $(\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}})$  and substituting our expressions for  $\|\boldsymbol{\beta}^{\perp}\|$ ,  $\|\widetilde{\boldsymbol{v}} - \boldsymbol{y}\|$ ,  $\langle \boldsymbol{g}, \widetilde{\boldsymbol{\beta}} \rangle$ ,  $\|\boldsymbol{\beta}^{\parallel}\|$  yields  $\nabla_{\boldsymbol{v}} Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}'' = (\alpha^{\star}\kappa \pm n^{-\Omega(1)})\boldsymbol{f} + (\sigma^{\star} \pm n^{-\Omega(1)})\boldsymbol{h} + (\gamma^{\star} \pm n^{-\Omega(1)})\boldsymbol{y} - ((\gamma^{\star} \pm n^{-\Omega(1)})\widetilde{\boldsymbol{v}} + (\rho^{\star})'(\widetilde{\boldsymbol{v}}))$ 

By substituting the definition of  $\tilde{v}$  in this lemma's statement and using standard properties about the relationship between  $\rho^*$  and  $\operatorname{prox}_{\gamma\rho}$  (see Section 2), we see that

$$\|\nabla_{\boldsymbol{v}}Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}''\| \le n^{\frac{1}{2}-\Omega(1)}$$

By 4-strong concavity, we have with high probability that

$$Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}'' \ge \max_{\boldsymbol{v} \in [0,1]^n} Q_{\widetilde{\boldsymbol{\beta}},\boldsymbol{v}}'' - n^{1-\Omega(1)}.$$

To conclude the proof, we remark that plugging our estimates into the definition of  $Q''_{\beta,v}$  similarly implies that there exists a constant  $c^* \in \mathbb{R}$  such that w.h.p.,

$$|Q_{\widetilde{\boldsymbol{\beta}},\widetilde{\boldsymbol{v}}}'' - c^* n| \le n^{1-\Omega(1)}.$$

#### 4.2.4 Putting Steps 1, 2, and 3 Together

Proof of Theorem 4.2. Note that part (a) follows immediately from Theorem 5.1, the observation that  $\rho'' : \mathbb{R} \to [0, 1/4]$ , and a change of variables in the case that  $R \neq 1$ . The extra factors in  $(2\varepsilon_{\text{DP}}, (1 + e^{\varepsilon_{\text{DP}}})\delta_{\text{DP}})$  come from the conversion from add/remove-one DP to replace-one DP (see Lemma 2.7). Therefore, we focus on part (b). By Lemma 4.13, there exist constants  $c^* \in \mathbb{R}$  and  $c_{\text{cgmt}} > 0$  such that w.h.p,

$$c^{\star}n - O(n^{1-c_{\rm cgmt}}) \leq \min_{\boldsymbol{\beta} \in \mathbb{R}^d} Q_{\boldsymbol{\beta}, \widetilde{\boldsymbol{v}}}' \leq Q_{\widetilde{\boldsymbol{\beta}}, \widetilde{\boldsymbol{v}}}' \leq \max_{\boldsymbol{v} \in [0,1]^n} Q_{\widetilde{\boldsymbol{\beta}}, \boldsymbol{v}}' \leq c^{\star}n + O(n^{1-c_{\rm cgmt}}).$$

By Lemma 3.6 and Lemma 3.14, for any arbitrarily small constant  $c_{\text{diam}} > 0$ , there exists an upper bound  $L_{\beta} = O(n^{c_{\text{diam}}})$  such that w.h.p.,

$$\widetilde{\boldsymbol{\beta}}, \widehat{\boldsymbol{\beta}} \in [-L_{\boldsymbol{\beta}}, +L_{\boldsymbol{\beta}}]^d.$$

For brevity, set  $S_{\beta} = [-L_{\beta}, +L_{\beta}]^d$  and  $S_{v} = [-1, +1]^n$ . Then,  $\widetilde{\beta} \in S_{\beta}$  implies that w.h.p.,

$$\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}\min_{\boldsymbol{\beta}\in\mathcal{S}_{\boldsymbol{\beta}}} Q_{\boldsymbol{\beta},\boldsymbol{v}}' \leq c^* n + O(n^{1-c_{\text{cgmt}}}).$$
(4.14)

Next, we would like to relate  $Q''_{\beta,v}$  to  $Q'_{\beta,v}$  via CGMT (Theorem 2.8). First, define the function

$$\psi(\boldsymbol{\beta}^{\perp},\boldsymbol{v}) = \min_{\boldsymbol{\beta}^{\parallel} \in (\mathcal{S}_{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\perp}) \cap \operatorname{span}(\boldsymbol{\beta}^{\star})} \frac{\|\boldsymbol{\beta}^{\parallel}\|}{\sqrt{d}} \langle \boldsymbol{f}, \boldsymbol{v} - \boldsymbol{y} \rangle + \frac{\lambda}{2} \|\boldsymbol{\beta}^{\perp} + \boldsymbol{\beta}^{\parallel}\|^{2} + \nu \langle \boldsymbol{\xi}, \boldsymbol{\beta}^{\perp} + \boldsymbol{\beta}^{\parallel} \rangle - \rho^{\star}(\boldsymbol{v}).$$

By Lemma 4.12, we can express  $Q'_{\beta,v}$  in the form amenable to CGMT using  $\psi$ :

$$\min_{\boldsymbol{\beta}\in\mathcal{S}_{\boldsymbol{\beta}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q'_{\boldsymbol{\beta},\boldsymbol{v}} = \max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}\min_{\boldsymbol{\beta}^{\perp}\in\mathcal{S}_{\boldsymbol{\beta}}\cap\mathrm{span}(\boldsymbol{\beta}^{\star})^{\perp}}\frac{1}{\sqrt{d}}\langle\boldsymbol{H}\boldsymbol{\beta}^{\perp},\boldsymbol{v}-\boldsymbol{y}\rangle + \psi(\boldsymbol{\beta}^{\perp},\boldsymbol{v}).$$

As per the discussion surrounding Lemma 4.12, the function  $\psi$  is independent of H (indeed, even though  $\boldsymbol{y}$  appears in the definition of the  $\psi$  function,  $\boldsymbol{y} \sim \text{Bernoulli}(\rho'(\frac{\|\boldsymbol{\beta}^*\|}{\sqrt{d}}\boldsymbol{f}))$  depends on the

design matrix only through f, not H). Thus, by (4.14) and CGMT (Theorem 2.8), we have that w.h.p.,

$$\min_{\boldsymbol{\beta}\in\mathcal{S}_{\boldsymbol{\beta}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q'_{\boldsymbol{\beta},\boldsymbol{v}} \leq c^{\star}n + O(n^{1-c_{\text{cgmt}}}).$$

By the definition of  $\widetilde{\beta}$  in the statement of Lemma 4.13, along with our assumption that  $\beta^* \rightsquigarrow \beta_0^*$ ,

$$(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \widetilde{\boldsymbol{\beta}}) \rightsquigarrow \left(\beta_0^{\star}, \, \xi_0, \, \alpha^{\star} \beta_0^{\star} + \sqrt{(\sigma^{\star})^2 - (\gamma^{\star} \nu)^2} Z + \gamma^{\star} \nu \xi_0\right)$$

For brevity, let  $\beta_0 \in \mathbb{R}$  denote the third random variable in the above triple. Then, the above assertion is that  $(\beta^*, \boldsymbol{\xi}, \tilde{\boldsymbol{\beta}}) \rightsquigarrow (\beta_0^*, \xi_0, \beta_0)$ , and we want to show that  $(\beta^*, \boldsymbol{\xi}, \hat{\boldsymbol{\beta}}) \rightsquigarrow (\beta_0^*, \xi_0, \beta_0)$ , as well. To this end, we use the same "excision" technique as we did in the proof of Theorem 3.7(b) in Section 3.2.4: fix an order-k pseudo-Lipschitz function  $f : \mathbb{R}^3 \to \mathbb{R}$ , and excise from  $S_{\boldsymbol{\beta}}$  the open set  $\mathcal{T}_{\boldsymbol{\beta}}$  where  $\boldsymbol{\beta} \in \mathcal{T}_{\boldsymbol{\beta}}$  iff the average value of f over the coordinates of  $(\boldsymbol{\beta}^*, \boldsymbol{\xi}, \boldsymbol{\beta})$  differs from the expected value of f over the randomness of  $(\beta_0^*, \xi_0, \beta_0)$  by strictly less than  $n^{-c'_{\text{slack}}}$ , for a sufficiently small constant  $c'_{\text{slack}}$ . Note that this excision adjusts the definition of the mean function  $\psi$ , as well. Then, applying CGMT and universality as before, along with the strong convexity afforded by Lemma 4.13, yields w.h.p.,

$$\min_{\boldsymbol{\beta}\in\mathcal{S}_{\boldsymbol{\beta}}\setminus\mathcal{T}_{\boldsymbol{\beta}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q'_{\boldsymbol{\beta},\boldsymbol{v}}\geq c^{\star}n+\Omega(n^{1-c_{\mathrm{total}}}),$$

for a constant  $0 < c_{\text{total}} < c_{\text{cgmt}}$ . Thus, w.h.p.,

$$\min_{\boldsymbol{\beta}\in\mathcal{S}_{\boldsymbol{\beta}}\setminus\mathcal{T}_{\boldsymbol{\beta}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q'_{\boldsymbol{\beta},\boldsymbol{v}}>\min_{\boldsymbol{\beta}\in\mathcal{S}_{\boldsymbol{\beta}}}\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}Q'_{\boldsymbol{\beta},\boldsymbol{v}}$$

In other words, if  $\mathbf{X} = \frac{1}{\sqrt{d}} \mathbf{G}$  is Gaussian, then w.h.p. the minimizer  $\hat{\boldsymbol{\beta}} \in S_{\boldsymbol{\beta}}$  lies inside  $\mathcal{T}_{\boldsymbol{\beta}}$ , meaning that  $(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \hat{\boldsymbol{\beta}}) \rightsquigarrow (\beta_0^{\star}, \xi_0, \beta_0)$ , as desired. This completes our characterization of the estimation error of  $\hat{\boldsymbol{\beta}}$  in the Gaussian case. For the difference  $\rho'(\mathbf{X}\boldsymbol{\beta}^{\star}) - \rho'(\mathbf{X}\hat{\boldsymbol{\beta}})$ , we carry out an entirely analogous dual argument with the roles of  $\boldsymbol{\beta}$  and  $\boldsymbol{v}$  exchanged. As before, we start by noting that by CGMT, we have w.h.p. that

$$\max_{\boldsymbol{v}\in\mathcal{S}_{\boldsymbol{v}}}\min_{\boldsymbol{u}\in\mathcal{S}_{\boldsymbol{u}}}Q'_{\boldsymbol{\beta},\boldsymbol{v}}\geq c^{\star}n-O(n^{1-c_{\mathrm{cgmt}}}).$$

By the definition of  $\tilde{\boldsymbol{v}}$  in the statement of Lemma 4.13, along with the fact that  $\boldsymbol{f} = \boldsymbol{G} \cdot \frac{\boldsymbol{\beta}^{\star}}{\|\boldsymbol{\beta}^{\star}\|}$ and  $\boldsymbol{\beta}^{\star} \rightsquigarrow \beta_0^{\star}$  with  $\mathbb{E}(\beta_0^{\star})^2 = \kappa^2$ , we have for dummy variables  $f_0, h_0 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$  and  $y_0|f_0 \sim$ Bernoullii $(\rho'(\kappa f_0))$  that

$$\left(\frac{\|\boldsymbol{\beta}^{\star}\|}{\sqrt{d}}\boldsymbol{f}, \widetilde{\boldsymbol{v}}\right) \rightsquigarrow \left(\kappa f_{0}, \ \alpha^{\star} \kappa f_{0} + \sigma^{\star} h_{0} + y_{0} - \operatorname{prox}_{\gamma^{\star} \rho}(\alpha^{\star} \kappa f_{0} + \sigma^{\star} h_{0} + \gamma^{\star} y_{0})\right)$$

For brevity, let  $v_0 \in \mathbb{R}$  denote the second random variable in the above pair. Then the above assertion is that  $(\frac{\|\boldsymbol{\beta}^{\star}\|}{\sqrt{d}}\boldsymbol{f}, \widetilde{\boldsymbol{v}}) \rightsquigarrow (\kappa f_0, v_0)$ , and we want to show that  $(\boldsymbol{X}\boldsymbol{\beta}^{\star}, \rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})) \rightsquigarrow (\kappa f_0, v_0)$ . Note that  $\frac{\|\boldsymbol{\beta}^{\star}\|}{\sqrt{d}}\boldsymbol{f} = \boldsymbol{X}\boldsymbol{\beta}^{\star}$  when  $\boldsymbol{X} = \frac{1}{\sqrt{d}}\boldsymbol{G}$ . Also note that by Lemma 4.11, we have  $\widehat{\boldsymbol{v}} = \rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})$ . Thus, applying the excision argument yet again to pass from  $(\frac{\|\boldsymbol{\beta}^{\star}\|}{\sqrt{d}}\boldsymbol{f}, \widetilde{\boldsymbol{v}})$  and  $(\frac{\|\boldsymbol{\beta}^{\star}\|}{\sqrt{d}}\boldsymbol{f}, \widehat{\boldsymbol{v}})$  yields  $(\boldsymbol{X}\boldsymbol{\beta}^{\star}, \rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})) \rightsquigarrow (\kappa f_0, v_0)$ , as desired. This concludes the proof in the Gaussian case, that  $\boldsymbol{X} = \frac{1}{\sqrt{d}}\boldsymbol{G}$ . For a general subgaussian design, we apply Lemma 4.10, along with the definition of slow pseudo-Lipschitz convergence (Definition 4.1), to conclude that

$$(\boldsymbol{\beta}^{\star},\boldsymbol{\xi},\widehat{\boldsymbol{\beta}}-\boldsymbol{\beta}^{\star}) \dashrightarrow (\beta_{0}^{\star},\,\xi_{0},\,\alpha^{\star}\beta_{0}^{\star}+\sqrt{(\sigma^{\star})^{2}-(\gamma^{\star}\nu)^{2}}Z+\gamma^{\star}\nu\xi_{0}),$$

and that for  $Z_1 = f_0$  and  $Z_2 = h_0$ ,

$$(\boldsymbol{X}\boldsymbol{\beta}^{\star},\,\rho'(\boldsymbol{X}\widehat{\boldsymbol{\beta}})) \dashrightarrow \left(\kappa Z_1,\,\,\alpha^{\star}\kappa Z_1 + \sigma^{\star}Z_2 + y_0 - \operatorname{prox}_{\gamma^{\star}\rho}(\alpha^{\star}\kappa Z_1 + \sigma^{\star}Z_2 + \gamma^{\star}y_0)\right). \qquad \Box$$

# 5 Privacy of Objective Perturbation with Small $\lambda$

In this section, we present our improved privacy proof for objective perturbation (Algorithm 1), from which we derived Theorems 3.7(a) and 4.2(a). The main result of this section, Theorem 5.1, is an extension of Theorem 2.6 (Theorems 3.1 and 3.2 of [RKW23]). While Theorem 5.1 will apply to all strictly positive regularization strengths  $\lambda > 0$  and perturbation magnitudes  $\nu > 0$ , Theorem 2.6 required that  $\lambda > s$  for some strictly positive s depending on the smoothness of the loss function. For example, s = 1 in the case of robust linear regression, and s = 1/4 in the case of logistic regression.

For this section only, we shall the refer to various privacy loss parameters of the objective perturbation algorithm by  $\varepsilon$ ,  $\delta$ ,  $\alpha$ , and  $\rho$ , rather than  $\varepsilon_{\text{DP}}$ ,  $\delta_{\text{DP}}$ ,  $\alpha_{\text{DP}}$ , and  $\rho_{\text{DP}}$ .

**Theorem 5.1.** Suppose that  $\ell_0 : \mathbb{R}^2 \to \mathbb{R}$  satisfies  $|\partial_1 \ell_0(\eta, y)| \leq L$  and  $0 \leq \partial_1^2 \ell_0(\eta, y) \leq s$  for some constants L, s > 0 and for all  $\eta, y \in \mathbb{R}$ . Then objective perturbation (Algorithm 1) with R = 1, any  $\lambda, \nu > 0$ , the GLM loss function  $\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = \ell_0(\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle, y)$ , and any strictly positive  $\lambda, \nu > 0$ , satisfies add/remove-one  $(\varepsilon, \delta)$ -differential privacy for any  $\varepsilon \geq 0$  and

$$\delta = \begin{cases} 2 \cdot \operatorname{HockeyStick}(\tilde{\varepsilon}, \frac{L}{\nu}) & \text{if } \hat{\varepsilon} \ge 0, \\ (1 - e^{\hat{\varepsilon}}) + 2e^{\hat{\varepsilon}} \cdot \operatorname{HockeyStick}\left(\frac{L^2}{2\nu^2}, \frac{L}{\nu}\right) & \text{otherwise,} \end{cases}$$

where we set  $\tilde{\varepsilon} = \varepsilon - \log(1 + s/\lambda)$  and  $\hat{\varepsilon} = \tilde{\varepsilon} - L^2/2\nu^2$ .

The algorithm also satisfies add/remove-one  $(\alpha, \varepsilon)$ -Rényi differential privacy for any  $\alpha > 1$  and

$$\varepsilon = \log\left(1 + \frac{s}{\lambda}\right) + \frac{L^2}{2\nu^2} + \frac{1}{\alpha - 1}\log\mathbb{E}_{X \sim \mathcal{N}\left(0, \frac{L^2}{\nu^2}\right)}\left[e^{(\alpha - 1)|X|}\right].$$

Before we prove Theorem 5.1, we make a couple comments. First, the only differences between the statement of this theorem and the statement of Theorem 2.6 are our removal of the assumption that  $\lambda > s$ , and, relatedly, our replacement of the quantity  $-\log(1 - s/\lambda)$  in both the approximate DP and RDP bounds with the strictly smaller quantity  $\log(1 + s/\lambda)$ . We also remark that the conclusion of Theorem 5.1 implies a  $\rho$ -zCDP bound with constant  $\rho$  for any  $\lambda, \nu > 0$ , justifying our informal commentary in Section 1.1:

**Corollary 5.2.** Under the same conditions as Theorem 5.1, Algorithm 1 satisfies  $\rho$ -zCDP for

$$\rho = \log\left(1 + \frac{s}{\lambda}\right) + \frac{L^2}{2\nu^2} + \sqrt{\frac{2}{\pi}} \cdot \frac{L}{\nu}.$$

*Proof.* By the definition of zCDP (Definition 2.2), we must show that the expression for  $\varepsilon$  in the RDP bound of Theorem 5.1 is at most  $\rho\alpha$ , for all  $\alpha > 1$ . Using a standard formula for the MGF of the *folded Gaussian distribution*, we see that if  $X \sim \mathcal{N}(0, \sigma^2)$ , then

$$\log \mathbb{E}[e^{t|X|}] = \frac{\sigma^2 t^2}{2} + \log(2\Phi(\sigma t)).$$

Substituting  $\sigma = L/\nu$  and  $t = \alpha - 1$  yields  $(\alpha, \varepsilon)$ -RDP for all  $\alpha > 1$  and

$$\varepsilon = \log\left(1 + \frac{s}{\lambda}\right) + \frac{L^2}{2\nu^2} + \left(\frac{L^2}{2\nu^2}(\alpha - 1) + \frac{\log(2\Phi(\frac{L}{\nu}(\alpha - 1)))}{\alpha - 1}\right)$$

Thus, the expression for  $\varepsilon$  has three terms: a constant term  $\log(1+s/\lambda)$ , a linear term  $(L^2/2\nu^2) \cdot \alpha$ , and third term  $\log(2\Phi(\frac{L}{\nu}(\alpha-1)))/(\alpha-1)$ , which one can check by hand strictly decreases from  $\sqrt{2/\pi}(L/\nu)$  to 0 as  $\alpha$  increases from 1 to  $\infty$ . We conclude that  $\varepsilon \leq \rho \alpha$  for the claimed value  $\rho$ .  $\Box$  *Proof of Theorem 5.1.* The main difference between this proof and the proof of Theorem 2.6 (Theorems 3.1 and 3.2 in [RKW23]) is our more careful bounding of a certain additive "change-of-variables" term in the privacy loss random variable. For clarity, we focus our attention on the part of the proof that changes.

To begin, let  $\widehat{\boldsymbol{\beta}} \in \mathbb{R}^d$  be the output of the algorithm when run on the data set  $(\boldsymbol{X}, \boldsymbol{y}) \in \mathbb{R}^{n \times d} \times \mathbb{R}^n$ . Let  $(\boldsymbol{X}', \boldsymbol{y}') \in \mathbb{R}^{(n \pm 1) \times d} \times \mathbb{R}^{(n \pm 1)}$  denote an add/remove-one adjacent data set. Because of the random perturbation term  $\boldsymbol{\xi} \in \mathbb{R}^d$  in the algorithm,  $\widehat{\boldsymbol{\beta}}$  is a random variable even for fixed  $\boldsymbol{X}$  and  $\boldsymbol{y}$ . Denote its density at  $\boldsymbol{\beta} \in \mathbb{R}^d$  by  $\mathsf{PDF}(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y}))$  and consider the privacy loss random variable

$$Z = \log\left(\frac{\mathsf{PDF}(\widehat{\boldsymbol{\beta}}; (\boldsymbol{X}, \boldsymbol{y}))}{\mathsf{PDF}(\widehat{\boldsymbol{\beta}}; (\boldsymbol{X}', \boldsymbol{y}'))}\right).$$

Note that both the numerator and the denominator involve  $\hat{\beta}$ , which is the output of the algorithm on  $(\mathbf{X}, \mathbf{y})$ , not  $(\mathbf{X}', \mathbf{y}')$ . In order to simplify Z, notice that there is a one-to-one correspondence between possible instantiations of the perturbation vector  $\boldsymbol{\xi}$  and possible outputs  $\boldsymbol{\beta}$  of the algorithm. Indeed, the algorithm outputs  $\boldsymbol{\beta}$  on input  $(\mathbf{X}, \mathbf{y})$  if and only if the chosen perturbation vector is

$$g(\boldsymbol{eta};(\boldsymbol{X},\boldsymbol{y})) = -rac{\lambda}{
u} \boldsymbol{eta} - rac{1}{
u} \sum_{i=1}^n \partial_1 \ell_0(\langle \boldsymbol{x}_i, \boldsymbol{eta} 
angle, y_i) \cdot \boldsymbol{x}_i.$$

Let  $J_g(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})) \in \mathbb{R}^{d \times d}$  denote the Jacobian matrix of the function g with respect to  $\boldsymbol{\beta}$ , and let  $\mathsf{PDF}_{\mathcal{N}}(\boldsymbol{z})$  denote the standard Gaussian density at  $\boldsymbol{z} \in \mathbb{R}^d$ . Using the change-of-variables formula, we can simplify the expression for Z to

$$Z = \underbrace{\log\left(\frac{\left|\det J_g(\widehat{\boldsymbol{\beta}}; (\boldsymbol{X}, \boldsymbol{y}))\right|}{\left|\det J_g(\widehat{\boldsymbol{\beta}}; (\boldsymbol{X}', \boldsymbol{y}'))\right|}\right)}_{(*)} + \underbrace{\log\left(\frac{\mathsf{PDF}_{\mathcal{N}}(g(\widehat{\boldsymbol{\beta}}; (\boldsymbol{X}, \boldsymbol{y})))}{\mathsf{PDF}_{\mathcal{N}}(g(\widehat{\boldsymbol{\beta}}; (\boldsymbol{X}', \boldsymbol{y}')))}\right)}_{(**)}.$$

To analyze (\*), the log determinant ratio, observe first that the function  $J_g$  satisfies

$$-J_g(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})) = \frac{\lambda}{\nu} \boldsymbol{I}_d + \frac{1}{\nu} \sum_{i=1}^n \partial_1^2 \ell_0(\langle \boldsymbol{x}_i, \boldsymbol{\beta} \rangle, y_i) \cdot \boldsymbol{x}_i \boldsymbol{x}_i^\top$$

Recall that the data sets  $(\mathbf{X}, \mathbf{y})$  and  $(\mathbf{X}', \mathbf{y}')$  differ in a single data point, which is present in one data set but not the other. Denote this point by  $(\mathbf{x}_0, y_0) \in \mathbb{R}^d \times \mathbb{R}$ , where  $\|\mathbf{x}_0\| \leq R = 1$ . Suppose first that  $(\mathbf{x}_0, y_0)$  is present in  $(\mathbf{X}, \mathbf{y})$  but not  $(\mathbf{X}', \mathbf{y}')$ . Then, we have that

$$-J_g(oldsymbol{eta};(oldsymbol{X},oldsymbol{y})) = -J_g(oldsymbol{eta};(oldsymbol{X}',oldsymbol{y}')) + rac{1}{
u}\partial_1^2\ell_0(\langleoldsymbol{x}_0,oldsymbol{eta}
angle,y_0)\cdotoldsymbol{x}_0oldsymbol{x}_0^{ op}.$$

Thus, by the matrix determinant lemma, we have that

$$\frac{\det(-J_g(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})))}{\det(-J_g(\boldsymbol{\beta}; (\boldsymbol{X}', \boldsymbol{y}')))} = 1 + \frac{1}{\nu} \partial_1^2 \ell_0(\langle \boldsymbol{x}_0, \boldsymbol{\beta} \rangle, y_0) \cdot \boldsymbol{x}_0^\top (-J_g(\boldsymbol{\beta}; (\boldsymbol{X}', \boldsymbol{y}')))^{-1} \boldsymbol{x}_0.$$

Observe that  $-J_g(\boldsymbol{\beta}; (\boldsymbol{X}', \boldsymbol{y}')) \succeq \frac{\lambda}{\nu} \boldsymbol{I}_d$ , that  $\|\boldsymbol{x}_0\| \leq 1$ , and that  $\partial_1^2 \ell(\langle \boldsymbol{x}_0, \boldsymbol{\beta} \rangle, y_0) \in [0, s]$ . Using these three bounds, it follows that

$$1 \leq \frac{\det(-J_g(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})))}{\det(-J_g(\boldsymbol{\beta}; (\boldsymbol{X}', \boldsymbol{y}')))} \leq 1 + \frac{s}{\lambda}.$$

### Algorithm 2 Output Perturbation

- 1: **input:** design matrix  $\boldsymbol{X} \in \mathbb{R}^{n \times d}$  with  $\|\boldsymbol{x}_i\| \leq R$ , response vector  $\boldsymbol{y} \in \mathbb{R}^n$ , loss function  $\ell : \mathbb{R}^d \times \mathbb{R}^{d+1} \to \mathbb{R}$ , regularization strength  $\lambda > 0$ , perturbation strength  $\nu > 0$ .
- 2: Randomly sample the additive perturbation term:

$$oldsymbol{\xi} \sim \mathcal{N}(oldsymbol{0}, oldsymbol{I}_d)$$

3: Optimize the regularized objective function:

$$\widetilde{oldsymbol{eta}} = rgmin_{oldsymbol{eta} \in \mathbb{R}^d} \sum_{i=1}^n \ell(oldsymbol{eta}; (oldsymbol{x}_i, y_i)) + rac{\lambda}{2} \|oldsymbol{eta}\|^2$$

4: Add the perturbation:

$$\widehat{\boldsymbol{\beta}} = \widetilde{\boldsymbol{\beta}} + \nu \boldsymbol{\xi}$$

5: return  $\hat{\beta}$ 

Similarly, in the case that  $(\boldsymbol{x}_0, y_0)$  is present in  $(\boldsymbol{X}', \boldsymbol{y}')$  but not  $(\boldsymbol{X}, \boldsymbol{y})$ , we have that

$$\left(1+\frac{s}{\lambda}\right)^{-1} \leq \frac{\det(-J_g(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})))}{\det(-J_g(\boldsymbol{\beta}; (\boldsymbol{X}', \boldsymbol{y}')))} \leq 1.$$

In either case, we have that  $|(*)| \leq \log(1 + s/\lambda)$  in absolute value, improving on the  $-\log(1 - s/\lambda)$  upper bound on |(\*)| from the proof of Theorem 2.6 (Theorems 3.1 and 3.2 of [RKW23]), which only held for  $\lambda > s$ . The remainder of the proof, which involves bounding (\*\*) and then analyzing various Rényi and hockey stick divergences, is identical to the proof of Theorem 2.6, and we omit it for brevity.

## 6 Output Perturbation

In this section, we consider two immediate corollaries of Theorem 3.7 and Theorem 4.2, respectively. These corollaries give us the privacy-utility tradeoffs for *output perturbation* (Algorithm 2), when applied to the problems of robust linear regression and logistic regression. To arrive at these corollaries, we simply observe that output perturbation can be viewed as objective perturbation with  $\nu = 0$  (i.e. no linear perturbation term), plus an additive Gaussian noise term to the output  $\hat{\beta}$ . Privacy follows immediately from Theorem 2.5 (the analytic Gaussian mechanism of [BW18]), along with the  $\ell^2$  sensitivity bound of  $2L/\lambda$  afforded by *L*-Lipschitzness and  $\lambda$ -strong convexity. In Figure 5, we validate the predictions of these corollaries against random, synthetic data, showing that output perturbation essentially incurs an additive increase in error that shifts the entire error curve of the non-private case "up" by  $\nu^2$ .

**Corollary 6.1** (Output Perturbation for Robust Linear Regression). Let  $\hat{\beta}$  denote the output of Algorithm 2 with parameters  $R, \lambda, \nu > 0$  and instantiated with the L-Lipschitz Huber loss function

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = H_L(y - \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle).$$

(a) (Privacy)  $\widehat{\beta}$  satisfies  $(\varepsilon_{\rm DP}, \delta_{\rm DP})$ -differential privacy for any  $\varepsilon_{\rm DP} \ge 0$  and

$$\delta_{\rm DP} = {\rm HockeyStick}\left(\varepsilon_{\rm DP}, \frac{2LR}{\lambda\nu}\right).$$

- (b) (Utility) Suppose the following hold for some  $\beta^{\star} \in \mathbb{R}^d$  and  $\varepsilon^{\star} \in \mathbb{R}^n$  as  $n \to \infty$  and  $d/n \to \delta$ :
  - (i)  $\mathbf{X} \in B_R(\mathbf{0})^n \subseteq \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y} = \mathbf{X} \boldsymbol{\beta}^\star + \boldsymbol{\varepsilon}^\star$ .
  - (ii) There exist random variables  $\beta_0^*, \varepsilon_0^* \in \mathbb{R}$  such that  $\beta^* \rightsquigarrow \beta_0^*$  and  $\varepsilon^* \rightsquigarrow \varepsilon_0^*$ .

Suppose there exist  $\sigma^*, \tau^* > 0$  solving the following system of two scalar equations in two variables  $(\sigma, \tau)$ , which we write in terms of a dummy variable  $Z \sim \mathcal{N}(0, 1)$  and  $\kappa^2 = \mathbb{E}(\beta_0^*)^2$  as

$$\sigma^2 = \tau^2 \left( \frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma Z + \varepsilon_0^*}{1 + \tau} \right]_L^2 + \lambda^2 \kappa^2 \right), \tag{6.2a}$$

$$\tau = \frac{1}{\lambda\delta} \left( \delta - \frac{\tau}{1+\tau} \Pr\left[ -L < \frac{\sigma Z + \varepsilon_0^*}{1+\tau} < L \right] \right).$$
(6.2b)

Then, in terms of  $(\sigma^*, \tau^*)$ , the estimation error  $\widehat{\beta} - \beta^*$  satisfies, for  $\xi_0 \sim \mathcal{N}(0, 1)$ ,

$$(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}) \rightsquigarrow \left( \beta_{0}^{\star}, \xi_{0}, \tau^{\star} \left( \sqrt{\frac{1}{\delta} \mathbb{E} \left[ \frac{\sigma^{\star} Z + \varepsilon_{0}^{\star}}{1 + \tau^{\star}} \right]_{L}^{2}} Z - \lambda \beta_{0}^{\star} \right) + \nu \xi_{0} \right)$$

**Corollary 6.3** (Output Perturbation for Logistic Regression). Let  $\hat{\beta}$  be the output of Algorithm 2 with parameters  $R, \lambda, \nu > 0$  and instantiated with the logistic loss function

$$\ell(oldsymbol{eta};(oldsymbol{x},y))=
ho(\langleoldsymbol{x},oldsymbol{eta}
angle)-y\langleoldsymbol{x},oldsymbol{eta}
angle.$$

(a) (Privacy)  $\hat{\beta}$  satisfies  $(\varepsilon_{\rm DP}, \delta_{\rm DP})$ -differential privacy for any  $\varepsilon_{\rm DP} \ge 0$  and

$$\delta_{\rm DP} = {\rm HockeyStick}\left(\varepsilon_{\rm DP}, \frac{2R}{\lambda\nu}\right)$$

- (b) (Utility) Suppose the following assumptions hold for some  $\beta^* \in \mathbb{R}^d$  as  $n \to \infty$  and  $d/n \to \delta$ :
  - (i)  $\mathbf{X} \in B_R(\mathbf{0})^n \subseteq \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y} \sim \text{Bernoulli}(\rho'(\mathbf{X}\boldsymbol{\beta}^{\star}))$ .
  - (ii) There exists a random variable  $\beta_0^{\star} \in \mathbb{R}$  with  $\kappa^2 = \mathbb{E}(\beta_0^{\star})^2$  such that  $\beta^{\star} \rightsquigarrow \beta_0^{\star}$ .

Note that  $\boldsymbol{\xi} \rightsquigarrow \xi_0$  for  $\xi_0 \sim \mathcal{N}(0, 1)$ .

Suppose there are unique  $\sigma^*, \alpha^*, \gamma^* > 0$  solving the following system of three scalar equations in three variables  $(\sigma, \alpha, \gamma)$ , which we write in terms of dummy variables  $Z_1, Z_2 \stackrel{iid}{\sim} \mathcal{N}(0, 1)$  as

$$\sigma^{2} = \frac{\gamma^{2}}{\delta} \mathbb{E} \Big[ 2\rho'(-\kappa Z_{1})\rho'(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_{1} + \sigma Z_{2}))^{2} \Big], \qquad (6.4a)$$

$$\alpha = -\frac{1}{\delta} \mathbb{E}[2\rho''(-\kappa Z_1) \operatorname{prox}_{\gamma\rho} (\kappa \alpha Z_1 + \sigma Z_2)], \qquad (6.4b)$$

$$\gamma = \frac{1}{\lambda\delta} \left( \delta - 1 + \mathbb{E} \left[ \frac{2\rho'(-\kappa Z_1)}{1 + \gamma\rho''(\operatorname{prox}_{\gamma\rho}(\kappa\alpha Z_1 + \sigma Z_2))} \right] \right).$$
(6.4c)

Then, the estimation error  $\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}^{\star}$  satisfies

 $(\boldsymbol{\beta}^{\star}, \boldsymbol{\xi}, \widehat{\boldsymbol{\beta}}) \dashrightarrow (\beta_0^{\star}, \xi_0, \alpha^{\star} \beta_0^{\star} + \sigma^{\star} Z + \nu \xi_0).$ 



(a) Algorithm 2 with Huber loss (L = 10) and  $\boldsymbol{\varepsilon}^{\star} \sim \mathcal{N}(\mathbf{0}, (1/5)^2 \boldsymbol{I}_n)$  and  $\boldsymbol{y} = \boldsymbol{X}\boldsymbol{\beta}^{\star} + \boldsymbol{\varepsilon}^{\star}$ .



(b) Algorithm 2 with logistic loss and  $\boldsymbol{y} \sim \text{Bernoulli}(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}))$ .

Figure 5: Predictions of Corollaries 6.1 and 6.3 on the estimation error of Algorithm 2. In all plots, curves correspond to theoretical predictions, and dots correspond to the mean over 100 simulations of the algorithm on synthetic data with  $n \times d = 1000$ . In the left plots, the perturbation magnitude is  $\nu = 0$ , but in the right plots,  $\nu = 1/2$ . In all plots, the signal strength is  $\kappa = 1$ , and we consider  $\beta^* \sim \mathcal{N}(\mathbf{0}, \kappa^2 \mathbf{I}_d)$ , along with  $\mathbf{X} \sim \frac{1}{\sqrt{d}}$ Uniform $(\{-1, +1\}^{n \times d})$ .

### Algorithm 3 Noisy Gradient Descent

- 1: **input:** design matrix  $\boldsymbol{X} = [\boldsymbol{x}_1 \cdots \boldsymbol{x}_n]^\top \in \mathbb{R}^{n \times d}$  with  $\|\boldsymbol{x}_i\| \leq R$ , response vector  $\boldsymbol{y} \in \mathbb{R}^n$ , loss function  $\ell : \mathbb{R}^d \times \mathbb{R}^{d+1} \to \mathbb{R}$ , step size  $\gamma > 0$ , noise magnitude  $\nu > 0$ .
- 2: Initialize  $\boldsymbol{\beta}^{(0)} = \mathbf{0}$
- 3: for t = 0, ..., T 1 do
- 4: Sample the  $t^{\text{th}}$  gradient perturbation vector:

$$\boldsymbol{\xi}^{(t)} \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_d)$$

5: Take a noisy gradient step:

$$\boldsymbol{\beta}^{(t+1)} = \boldsymbol{\beta}^{(t)} - \gamma \left( \sum_{i=1}^{n} \nabla_{\boldsymbol{\beta}} \, \ell(\boldsymbol{\beta}^{(t)}; (\boldsymbol{x}_i, y_i)) + \nu \boldsymbol{\xi}^{(t)} \right)$$

6: end for 7: return  $\widehat{\boldsymbol{\beta}} = \boldsymbol{\beta}^{(T)}$ 

## 7 Noisy Stochastic Gradient Descent

In this section, we present initial results for noisy stochastic gradient descent (DP-SGD) in the proportional regime that parallel our main results for objective perturbation (Theorems 3.7 and 4.2) and for output perturbation (Corollaries 6.1 and 6.3). The results in this section follow immediately from existing results in the non-private literature [ $GTM^+24$ , Han24], and as such, we consider the results of this section as a valuable point of reference for situating our other results, rather than a main contribution in their own right.

Due to certain separability requirements on the loss function in  $[\text{GTM}^+24]$ , which builds on [CMW20], as well as certain smoothness requirements in both of  $[\text{CMW20}, \text{GTM}^+24]$ , we cannot use these results in a black-box fashion to directly analyze the utility of DP-SGD on the robust linear regression and logistic regression tasks that we focused on in the other sections of this paper. Instead, we shift our attention to the following, non-standard, "conditional expectation" versions of the problems, which satisfy the necessary separability and smoothness requirements.

For robust linear regression, we previously assumed a labeled data set  $(\mathbf{X}, \mathbf{y})$  with  $\mathbf{y} = \mathbf{X}\beta^* + \varepsilon^*$ for suitably bounded  $\varepsilon^*$  and used the loss function  $H_L(\mathbf{y} - \mathbf{X}\beta)$ . We will now instead assume a labeled data set  $(\mathbf{X}, \mathbf{y})$  with  $\mathbf{y} = \mathbf{X}\beta^*$  and use the "conditional expectation" loss function

$$\ell_{\text{RobustLinearCE}}(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})) = \mathbb{E}_{\boldsymbol{\varepsilon}^{\star}}[H_L(\boldsymbol{y} + \boldsymbol{\varepsilon}^{\star} - \boldsymbol{X}\boldsymbol{\beta}) \mid \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{\beta}^{\star}]$$

For logistic regression, we previously assumed a labeled data set with  $\boldsymbol{y} \sim \text{Bernoulli}(\rho'(\boldsymbol{X}\boldsymbol{\beta}^{\star}))$ and used the loss function  $\rho(\boldsymbol{X}\boldsymbol{\beta}) - \langle \boldsymbol{y}, \boldsymbol{X}\boldsymbol{\beta} \rangle$ . We will now instead assume  $\boldsymbol{y} = \boldsymbol{X}\boldsymbol{\beta}^{\star}$  and use the "conditional expectation" loss function

$$\ell_{\text{LogisticCE}}(\boldsymbol{\beta}; (\boldsymbol{X}, \boldsymbol{y})) = \mathbb{E}_{\boldsymbol{\hat{y}} \sim \text{Bernoulli}(\boldsymbol{\rho}'(\boldsymbol{y}))}[\boldsymbol{\rho}(\boldsymbol{X}\boldsymbol{\beta}) - \langle \boldsymbol{\hat{y}}, \boldsymbol{X}\boldsymbol{\beta} \rangle \mid \boldsymbol{X}, \boldsymbol{\beta}, \boldsymbol{\beta}^{\star}].$$

These conditional expectation-based versions are admittedly unwieldy, but have the advantage of rigorously satisfying the assumptions needed to apply the results of [CMW20, GTM<sup>+</sup>24], because they are coordinate-wise separable and vary smoothly in  $X\beta$  and  $X\beta^*$ . For example, although [CMW20, GTM<sup>+</sup>24] both observe that their theorems happen to give correct predictions about the

behavior of (non-private) SGD for the standard formulation of logistic regression, they acknowledge that their theorems do not technically apply to this loss function because it takes as input  $\boldsymbol{y} = \mathbf{1}[\boldsymbol{X}\boldsymbol{\beta}^* + \boldsymbol{\varepsilon}^* > 0] \in \{0,1\}^n$  for  $\boldsymbol{\varepsilon}^* \sim \text{Logistic}$ , which is discontinuous with respect to  $\boldsymbol{X}\boldsymbol{\beta}^*$ .

For simplicity, we consider private, full batch gradient descent (Algorithm 3), although we could have just as easily analyzed the stochastic version for a mini-batch of size  $\Omega(n)$  using the same results from [GTM<sup>+</sup>24, Han24]. We validate the predictions of the following theorems against simulated data in Figure 6.

**Theorem 7.1** (DP-SGD for Robust Linear Regression, "Conditional Expectation" Version). Let  $\hat{\beta}$  be the output of T iterations of Algorithm 3 with parameters  $R, \gamma, \nu > 0$  and instantiated with the loss function

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = \mathbb{E}_{\varepsilon_0^{\star}} [H_L(y + \varepsilon_0^{\star} - \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle) \mid \boldsymbol{\beta}, (\boldsymbol{x}, y)]$$

for some fixed, continuous random variable  $\varepsilon_0^{\star} \in \mathbb{R}$ .

- (a) (**Privacy**)  $\hat{\boldsymbol{\beta}}$  satisfies  $\rho_{\text{DP}}$ -zCDP for  $\rho_{\text{DP}} = T \cdot \frac{2L^2 R^2}{\nu^2}$ .
- (b) (Utility) Suppose the following assumptions hold for some  $\beta^* \in \mathbb{R}^d$  as  $n \to \infty$  and  $d/n \to \delta$ :
  - (i)  $\mathbf{X} \in B_R(\mathbf{0})^n \subseteq \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y} = \mathbf{X} \boldsymbol{\beta}^{\star}$ .
  - (ii) There exists a random variable  $\beta_0^{\star} \in \mathbb{R}$  such that  $\beta^{\star} \rightsquigarrow \beta_0^{\star}$ .

Consider the following  $O(T^2)$  equations that recursively define, for all  $t, s \in \{0, 1, ..., T\}$ , the random variables  $\boldsymbol{\theta}^{(t)}, \boldsymbol{\eta}^{(t)}, \boldsymbol{u}^{(t)}, \boldsymbol{\omega}^{(t)} \in \mathbb{R}^2$  and  $\partial \boldsymbol{\eta}^{(t)} / \partial \boldsymbol{\omega}^{(s)} \in \mathbb{R}^{2 \times 2}$ , as well as the deterministic  $2 \times 2$  matrices  $\boldsymbol{R}_{\boldsymbol{\theta}}(t, s), \boldsymbol{R}_{\boldsymbol{g}}(t, s), \boldsymbol{\Gamma}^{(t)}, \boldsymbol{C}_{\boldsymbol{\theta}}(t, s), \boldsymbol{C}_{\boldsymbol{g}}(t, s) \in \mathbb{R}^{2 \times 2}$ .

The random variables are

$$\begin{split} \boldsymbol{\theta}^{(0)} &= \begin{bmatrix} 0\\ \beta_0^{\star} \end{bmatrix} \in \mathbb{R}^2, \\ \boldsymbol{\theta}^{(t+1)} &= (1 + \boldsymbol{\Gamma}^{(t)}) \boldsymbol{\theta}^{(t)} - \gamma \nu \begin{bmatrix} \xi_0^{(t)}\\ 0 \end{bmatrix} + \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{g}}(t,k) \boldsymbol{\theta}^{(k)} + \boldsymbol{u}^{(t)} \in \mathbb{R}^2, \\ \boldsymbol{\eta}^{(t)} &= -\gamma \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{\theta}}(t,k) \begin{bmatrix} \mathbb{E}_{\varepsilon_0^{\star}}[\eta_1^{(k)} - \eta_2^{(k)} - \varepsilon_0^{\star}]_L \\ 0 \end{bmatrix} + \boldsymbol{\omega}^{(t)} \in \mathbb{R}^2, \\ \frac{\partial \boldsymbol{\eta}^{(t)}}{\partial \boldsymbol{\omega}^{(s)}} &= \begin{cases} \mathbf{0} & \text{if } t < s, \\ I_2 & \text{if } t = s, \\ -\gamma \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{\theta}}(t,k) \operatorname{Pr}_{\varepsilon_0^{\star}} \left[ |\eta_1^{(k)} - \eta_2^{(k)} - \varepsilon_0^{\star}| < L \right] \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \frac{\partial \boldsymbol{\eta}^{(k)}}{\partial \boldsymbol{\omega}^{(s)}} & \text{if } t > s \end{cases} \right\} \in \mathbb{R}^{2 \times 2}, \end{split}$$

where  $\xi_0^{(0)}, \ldots, \xi_0^{(T)} \stackrel{iid}{\sim} \mathcal{N}(0, 1)$ , the random vector  $[(\boldsymbol{u}^{(0)})^\top | \cdots | (\boldsymbol{u}^{(t)})^\top]^\top \in \mathbb{R}^{2(t+1)}$  (for each  $t = 0, \ldots, T$ ) follows a centered multivariate Gaussian distribution with covariance matrix entries given by  $C_{\boldsymbol{g}}$ , and the random vector  $[(\boldsymbol{\omega}^{(0)})^\top | \cdots | (\boldsymbol{\omega}^{(t)})^\top]^\top \in \mathbb{R}^{2(t+1)}$  (for each  $t = 0, \ldots, T$ ) follows a centered multivariate Gaussian distribution with covariance matrix entries given by  $C_{\boldsymbol{\theta}}$ .

The deterministic  $2 \times 2$  matrices are

$$\begin{aligned} \boldsymbol{R}_{\boldsymbol{\theta}}(t+1,s) &= \begin{cases} \mathbf{0} & \text{if } t < s, \\ \mathbf{I}_{2} & \text{if } t = s, \\ (\mathbf{I}_{2} + \mathbf{\Gamma}^{(t)}) \mathbf{R}_{\boldsymbol{\theta}}(t,s) + \sum_{k=0}^{t-1} \mathbf{R}_{\boldsymbol{g}}(t,k) \mathbf{R}_{\boldsymbol{\theta}}(k,s) & \text{if } t > s \end{cases} \right\} \in \mathbb{R}^{2 \times 2} \\ \mathbf{R}_{\boldsymbol{g}}(t,s) &= -\frac{\gamma}{\delta} \mathbb{E} \left[ \Pr_{\varepsilon_{0}^{\star}} \left[ |\eta_{1}^{(t)} - \eta_{2}^{(t)} - \varepsilon_{0}^{\star}| < L \right] \left[ \begin{array}{c} 1 & -1 \\ 0 & 0 \end{array} \right] \frac{\partial \boldsymbol{\eta}^{(t)}}{\partial \boldsymbol{\omega}^{(s)}} \right] \in \mathbb{R}^{2 \times 2}, \\ \mathbf{\Gamma}^{(t)} &= -\frac{\gamma}{\delta} \mathbb{E} \left[ \Pr_{\varepsilon_{0}^{\star}} \left[ |\eta_{1}^{(t)} - \eta_{2}^{(t)} - \varepsilon_{0}^{\star}| < L \right] \left[ \begin{array}{c} 1 & -1 \\ 0 & 0 \end{array} \right] \right] \in \mathbb{R}^{2 \times 2}, \\ \mathbf{C}_{\boldsymbol{\theta}}(t,s) &= \mathbb{E} [\boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(s)})^{\top}] \in \mathbb{R}^{2 \times 2}, \\ \mathbf{C}_{\boldsymbol{g}}(t,s) &= \frac{\gamma^{2}}{\delta} \mathbb{E} \left[ \begin{array}{c} \mathbb{E}_{\varepsilon_{0}^{\star}} [\eta_{1}^{(t)} - \eta_{2}^{(t)} - \varepsilon_{0}^{\star}]_{L} \mathbb{E}_{\varepsilon_{0}^{\star}} [\eta_{1}^{(s)} - \eta_{2}^{(s)} - \varepsilon_{0}^{\star}]_{L} \begin{array}{c} 0 \\ 0 \end{array} \right] \in \mathbb{R}^{2 \times 2}. \end{aligned}$$

Then, for any pseudo-Lipschitz functions  $\psi : \mathbb{R}^{T+1} \to \mathbb{R}$  and  $\phi : \mathbb{R}^T \to R$ ,

$$\frac{1}{d} \sum_{j=1}^{d} \psi(\beta_{j}^{\star}, \beta_{j}^{(1)}, \dots, \beta_{j}^{(T)}) \xrightarrow{P} \mathbb{E}[\psi(\beta_{0}^{\star}, \theta_{1}^{(1)}, \dots, \theta_{1}^{(T)})],$$
$$\frac{1}{n} \sum_{i=1}^{n} \phi(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{(1)} \rangle, \dots, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{(T-1)} \rangle) \xrightarrow{P} \mathbb{E}[\phi(\omega^{(0)}, \eta_{1}^{(1)}, \dots, \eta_{1}^{(T)})].$$

*Proof.* Part (a) follows from standard properties of zCDP, such as Theorem 2.4. For part (b), consider the generalized first-order method initialized at  $v^{(0)} = [\mathbf{0} \mid \boldsymbol{\beta}^{\star}] \in \mathbb{R}^{d \times 2}$  and updated according to the following rule:

$$m{v}^{(t+1)} = h^{(t)}(m{v}^{(t)}; m{\xi}^{(t)}) + m{X}^{ op} g^{(t)}(m{r}^{(t)}) \in \mathbb{R}^{d imes 2},$$
  
 $m{r}^{(t)} = m{X} \sum_{k=0}^{t} m{v}^{(k)} \in \mathbb{R}^{n imes 2},$ 

where we have defined the functions  $g^{(t)} : \mathbb{R}^{n \times 2} \to \mathbb{R}^{n \times 2}$  and  $h^{(t)} : \mathbb{R}^{d \times 2} \to \mathbb{R}^{d \times 2}$  in terms of a dummy variable  $\varepsilon^{\star} = (\varepsilon_1^{\star}, \ldots, \varepsilon_n^{\star}) \stackrel{iid}{\sim} \varepsilon_0^{\star}$  as follows:

$$g^{(t)}([\boldsymbol{r}_1^{(t)} \mid \boldsymbol{r}_2^{(t)}]) = \left[-\gamma \mathop{\mathbb{E}}_{\boldsymbol{\varepsilon^\star}} [\boldsymbol{r}_1^{(t)} - \boldsymbol{r}_2^{(t)} - \boldsymbol{\varepsilon^\star}]_L \mid \boldsymbol{0}\right]$$

and

$$h^{(t)}([\boldsymbol{v}_1^{(t)} \mid \boldsymbol{v}_2^{(t)}]; \boldsymbol{\xi}^{(t)}) = \left[-\gamma \nu \boldsymbol{\xi}^{(t)} \mid \boldsymbol{0}\right].$$

It is straightforward to verify that  $\mathbf{v}^{(t)} = [\boldsymbol{\beta}^{(t)} \mid \boldsymbol{\beta}^{\star}] \in \mathbb{R}^{d \times 2}$ , where  $\boldsymbol{\beta}^{(t)}$  is the  $t^{\text{th}}$  iterate of DP-SGD (Algorithm 3). If  $\mathbf{X}$  were an isotropic Gaussian, our desired result would follow directly from Theorem 3.2 of [GTM<sup>+</sup>24] applied to this sequence, provided that we can verify their assumptions (A1), (A2), (A3.b) and (A4). Assumptions (A1) and (A2) are equivalent to our assumption (i) in part (b). The separability and smoothness of assumption (A3.b) are easily verified from the above expressions for  $g^{(t)}$  and  $h^{(t)}$ . For assumption (A4), observe that  $\boldsymbol{\beta}^{\star} \rightsquigarrow \boldsymbol{\beta}_{0}^{\star}$  implies that  $\frac{1}{d} \|\boldsymbol{\beta}^{\star}\|^{2}$  converges to a finite constant, namely  $\mathbb{E}(\boldsymbol{\beta}_{0}^{\star})^{2}$ , as  $n \to \infty$ . Substituting the functions g and h and the partial derivatives of g into the system of equations in their theorem, and then taking the limit as  $n \to \infty$  and  $d/n \to \delta$ , yields our claimed system of equations, which we have partially simplified. For  $\mathbf{X}$  following a general, subgaussian design, we simply apply Theorem 2.13 (GFOM Universality, Theorem 3.2 of [Han24]).

**Theorem 7.2** (DP-SGD for Logistic Regression, "Conditional Expectation" Version). Let  $\hat{\beta}$  be the output of T iterations of Algorithm 3 with parameters  $R, \gamma, \nu > 0$ , instantiated with the loss function

$$\ell(\boldsymbol{\beta}; (\boldsymbol{x}, y)) = \mathbb{E}_{\hat{y} \sim \text{Bernoulli}(\rho'(y))} \big[ \rho(\langle \boldsymbol{x}, \boldsymbol{\beta} \rangle) - \hat{y} \langle \boldsymbol{x}, \boldsymbol{\beta} \rangle \mid \boldsymbol{\beta}, (\boldsymbol{x}, y) \big]$$

- (a) (Privacy)  $\hat{\beta}$  satisfies  $\rho_{\rm DP}$ -zCDP for  $\rho_{\rm DP} = T \cdot \frac{2R^2}{\nu^2}$ .
- (b) (Utility) Suppose the following assumptions hold for some  $\beta^* \in \mathbb{R}^d$  as  $n \to \infty$  and  $d/n \to \delta$ :
  - (i)  $\mathbf{X} \in B_R(\mathbf{0})^n \subseteq \mathbb{R}^{n \times d}$  follows a subgaussian design and  $\mathbf{y} = \mathbf{X} \boldsymbol{\beta}^{\star}$ .
  - (ii) There exists a random variable  $\beta_0^{\star} \in \mathbb{R}$  such that  $\beta^{\star} \rightsquigarrow \beta_0^{\star}$ .

Consider the following  $O(T^2)$  equations that recursively define, for all  $t, s \in \{0, 1, ..., T\}$ , the random variables  $\boldsymbol{\theta}^{(t)}, \boldsymbol{\eta}^{(t)}, \boldsymbol{u}^{(t)}, \boldsymbol{\omega}^{(t)} \in \mathbb{R}^2$  and  $\partial \boldsymbol{\eta}^{(t)} / \partial \boldsymbol{\omega}^{(s)} \in \mathbb{R}^{2 \times 2}$ , as well as the deterministic  $2 \times 2$  matrices  $\boldsymbol{R}_{\boldsymbol{\theta}}(t, s), \boldsymbol{R}_{\boldsymbol{g}}(t, s), \boldsymbol{\Gamma}^{(t)}, \boldsymbol{C}_{\boldsymbol{\theta}}(t, s), \boldsymbol{C}_{\boldsymbol{g}}(t, s) \in \mathbb{R}^{2 \times 2}$ .

The random variables are

$$\begin{split} \boldsymbol{\theta}^{(0)} &= \begin{bmatrix} 0\\ \beta_0^{\star} \end{bmatrix} \in \mathbb{R}^2, \\ \boldsymbol{\theta}^{(t+1)} &= (1 + \boldsymbol{\Gamma}^{(t)}) \boldsymbol{\theta}^{(t)} - \gamma \nu \begin{bmatrix} \xi_0^{(t)}\\ 0 \end{bmatrix} + \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{g}}(t,k) \boldsymbol{\theta}^{(k)} + \boldsymbol{u}^{(t)} \in \mathbb{R}^2, \\ \boldsymbol{\eta}^{(t)} &= -\gamma \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{\theta}}(t,k) \begin{bmatrix} \rho'(\boldsymbol{\eta}_1^{(k)}) \rho'(-\boldsymbol{\eta}_2^{(k)}) - \rho'(-\boldsymbol{\eta}_1^{(k)}) \rho'(\boldsymbol{\eta}_2^{(k)}) \\ 0 \end{bmatrix} + \boldsymbol{\omega}^{(t)} \in \mathbb{R}^2, \\ \frac{\partial \boldsymbol{\eta}^{(t)}}{\partial \boldsymbol{\omega}^{(s)}} &= \begin{cases} \mathbf{0} & \text{if } t < s, \\ \mathbf{I}_2 & \text{if } t = s, \\ -\gamma \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{\theta}}(t,k) \begin{bmatrix} \rho''(\boldsymbol{\eta}_1^{(k)}) & -\rho''(\boldsymbol{\eta}_2^{(k)}) \\ 0 \end{bmatrix} \frac{\partial \boldsymbol{\eta}^{(k)}}{\partial \boldsymbol{\omega}^{(s)}} & \text{if } t > s \end{cases} \in \mathbb{R}^{2 \times 2}, \end{split}$$

where  $\xi_0^{(0)}, \ldots, \xi_0^{(T)} \stackrel{iid}{\sim} \mathcal{N}(0, 1)$ , the random vector  $[(\boldsymbol{u}^{(0)})^\top | \cdots | (\boldsymbol{u}^{(t)})^\top]^\top \in \mathbb{R}^{2(t+1)}$  (for each  $t = 0, \ldots, T$ ) follows a centered multivariate Gaussian distribution with covariance matrix entries given by  $C_{\boldsymbol{g}}$ , and the random vector  $[(\boldsymbol{\omega}^{(0)})^\top | \cdots | (\boldsymbol{\omega}^{(t)})^\top]^\top \in \mathbb{R}^{2(t+1)}$  (for each  $t = 0, \ldots, T$ ) follows a centered multivariate Gaussian distribution with covariance matrix entries given by  $C_{\boldsymbol{\theta}}$ .

The (deterministic / constant)  $2 \times 2$  matrices are

$$\begin{aligned} \boldsymbol{R}_{\boldsymbol{\theta}}(t+1,s) &= \begin{cases} \mathbf{0} & \text{if } t < s, \\ \mathbf{I}_{2} & \text{if } t = s, \\ (\mathbf{I}_{2} + \mathbf{\Gamma}^{(t)}) \boldsymbol{R}_{\boldsymbol{\theta}}(t,s) + \sum_{k=0}^{t-1} \boldsymbol{R}_{\boldsymbol{g}}(t,k) \boldsymbol{R}_{\boldsymbol{\theta}}(k,s) & \text{if } t > s \end{cases} \right\} \in \mathbb{R}^{2 \times 2} \\ \boldsymbol{R}_{\boldsymbol{g}}(t,s) &= -\frac{\gamma}{\delta} \mathbb{E} \left[ \begin{bmatrix} \rho^{\prime\prime}(\eta_{1}^{(t)}) & -\rho^{\prime\prime}(\eta_{2}^{(t)}) \\ 0 & 0 \end{bmatrix} \frac{\partial \boldsymbol{\eta}^{(t)}}{\partial \boldsymbol{\omega}^{(s)}} \right] \in \mathbb{R}^{2 \times 2}, \\ \mathbf{\Gamma}^{(t)} &= -\frac{\gamma}{\delta} \mathbb{E} \left[ \begin{array}{c} \rho^{\prime\prime}(\eta_{1}^{(t)}) & -\rho^{\prime\prime}(\eta_{2}^{(t)}) \\ 0 & 0 \end{bmatrix} \right] \in \mathbb{R}^{2 \times 2}, \\ \boldsymbol{C}_{\boldsymbol{\theta}}(t,s) &= \mathbb{E}[\boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(s)})^{\top}] \in \mathbb{R}^{2 \times 2}, \\ \boldsymbol{C}_{\boldsymbol{g}}(t,s) &= \frac{\gamma^{2}}{\delta} \mathbb{E} \left[ \begin{array}{c} (\rho^{\prime}(\eta_{1}^{(t)})\rho^{\prime}(-\eta_{2}^{(t)})-\rho^{\prime}(-\eta_{1}^{(t)})\rho^{\prime}(\eta_{2}^{(t)})) \\ \times (\rho^{\prime}(\eta_{1}^{(s)})\rho^{\prime}(-\eta_{2}^{(s)})-\rho^{\prime}(-\eta_{1}^{(s)})\rho^{\prime}(\eta_{2}^{(s)})) \\ 0 \end{bmatrix} \in \mathbb{R}^{2 \times 2}. \end{aligned}$$

Then, for any pseudo-Lipschitz functions  $\psi : \mathbb{R}^{T+1} \to \mathbb{R}$  and  $\phi : \mathbb{R}^T \to \mathbb{R}$ ,

$$\frac{1}{d} \sum_{j=1}^{d} \psi(\beta_{j}^{\star}, \beta_{j}^{(1)}, \dots, \beta_{j}^{(T)}) \xrightarrow{P} \mathbb{E}[\psi(\beta_{0}^{\star}, \theta_{1}^{(1)}, \dots, \theta_{1}^{(T)})],$$
$$\frac{1}{n} \sum_{i=1}^{n} \phi(\langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{\star} \rangle, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{(1)} \rangle, \dots, \langle \boldsymbol{x}_{i}, \boldsymbol{\beta}^{(T-1)} \rangle) \xrightarrow{P} \mathbb{E}[\phi(\omega^{(0)}, \eta_{1}^{(1)}, \dots, \eta_{1}^{(T)})].$$

The proof of Theorem 7.2 is essentially the same as that of Theorem 7.1, so we omit it.



(a) Algorithm 3 with Huber loss, "conditional expectation" version  $(L = 10, \varepsilon_0^* \sim \mathcal{N}(0, (1/5)^2))$ .



(b) Algorithm 3 with logistic loss, "conditional expectation" version.

Figure 6: Predictions of Theorems 7.1 and 7.2 on the estimation error of Algorithm 3. In all plots, curves correspond to theoretical predictions, and dots correspond to the mean over  $10^4$  simulations of the algorithm on synthetic data with  $n \times d = 1000$ . In the left plots, the noise magnitude is  $\nu = 0$ , but in the right plots,  $\nu = 1/10$ . In all plots, the signal strength is  $\kappa = 1$ , the step size is  $\gamma = 1/2 \cdot 1/(1 + \delta)$ , and we consider  $\beta^* \sim \mathcal{N}(\mathbf{0}, \kappa^2 \mathbf{I}_d)$ , along with  $\mathbf{X} \sim \frac{1}{\sqrt{d}} \text{Uniform}(\{-1, +1\}^{n \times d})$  and  $\mathbf{y} = \mathbf{X}\beta^*$ .

# References

- [AAL21] Ishaq Aden-Ali, Hassan Ashtiani, and Christopher Liaw. Privately learning mixtures of axis-aligned gaussians. In *Conference on Neural Information Processing Systems* (NeurIPS), 2021.
- [AAL24] Mohammad Afzali, Hassan Ashtiani, and Christopher Liaw. Mixtures of gaussians are privately learnable with a polynomial number of samples. In *International Conference* on Algorithmic Learning Theory (ALT), 2024.
- [ACG<sup>+</sup>16] Martin Abadi, Andy Chu, Ian Goodfellow, H. Brendan McMahan, Ilya Mironov, Kunal Talwar, and Li Zhang. Deep learning with differential privacy. In ACM Conference on Computer and Communications Security (CCS), 2016.
- [AJRV23] Kareem Amin, Matthew Joseph, Mónica Ribero, and Sergei Vassilvitskii. Easy differentially private linear regression. In *International Conference on Learning Representations (ICLR)*, 2023.
- [AKM<sup>+</sup>24] Sushant Agarwal, Gautam Kamath, Mahbod Majid, Argyris Mouzakis, Rose Silver, and Jonathan Ullman. Private mean estimation with person-level differential privacy. In ACM-SIAM Symposium on Discrete Algorithms (SODA), 2024.
- [AKT<sup>+</sup>23] Daniel Alabi, Pravesh K. Kothari, Pranay Tankala, Prayaag Venkat, and Fred Zhang. Privately estimating a gaussian: Efficient, robust, and optimal. In ACM Symposium on Theory of Computing (STOC), 2023.
- [AL22] Hassan Ashtiani and Christopher Liaw. Private and polynomial time algorithms for learning gaussians and beyond. In *Conference on Learning Theory (COLT)*, 2022.
- [ALMT14] Dennis Amelunxen, Martin Lotz, Michael B. McCoy, and Joel A. Tropp. Living on the edge: phase transitions in convex programs with random data. *Information and Inference: A Journal of the IMA*, 3(3):224–294, 06 2014.
- [ASH19] Ehsan Abbasi, Fariborz Salehi, and Babak Hassibi. Universality in learning from linear measurements. In *Conference on Neural Information Processing Systems (NeurIPS)*, 2019.
- [BDE<sup>+</sup>24] Gavin R Brown, Krishnamurthy Dj Dvijotham, Georgina Evans, Daogao Liu, Adam Smith, and Abhradeep Guha Thakurta. Private gradient descent for linear regression: Tighter error bounds and instance-specific uncertainty estimation. In International Conference on Machine Learning (ICML), 2024.
- [BHH<sup>+</sup>24] Gavin Brown, Jonathan Hayase, Samuel B. Hopkins, Weihao Kong, Xiyang Liu, Sewoong Oh, Juan C. Perdomo, and Adam Smith. Insufficient statistics perturbation: Stable estimators for private least squares. In *Conference on Learning Theory (COLT)*, 2024.
- [BHS23] Gavin Brown, Samuel Hopkins, and Adam Smith. Fast, sample-efficient, affineinvariant private mean and covariance estimation for subgaussian distributions. In *Conference on Learning Theory (COLT)*, 2023.
- [BHX23] Zhigang Bao, Qiyang Han, and Xiaocong Xu. A leave-one-out approach to approximate message passing, 2023.

- [BKSW19] Mark Bun, Gautam Kamath, Thomas Steinke, and Zhiwei Steven Wu. Private hypothesis selection. In *Conference on Neural Information Processing Systems (NeurIPS)*, 2019.
- [BLM15] Mohsen Bayati, Marc Lelarge, and Andrea Montanari. Universality in polytope phase transitions and message passing algorithms. *The Annals of Applied Probability*, 25(2):753 822, 2015.
- [BM11a] Mohsen Bayati and Andrea Montanari. The dynamics of message passing on dense graphs, with applications to compressed sensing. *IEEE Transactions on Information Theory*, 57(2):764–785, 2011.
- [BM11b] Mohsen Bayati and Andrea Montanari. The lasso risk for gaussian matrices. *IEEE Transactions on Information Theory*, 58(4):1997–2017, 2011.
- [BNS<sup>+</sup>16] Raef Bassily, Kobbi Nissim, Adam D. Smith, Thomas Steinke, Uri Stemmer, and Jonathan R. Ullman. Algorithmic stability for adaptive data analysis. In *ACM Symposium on Theory of Computing (STOC)*, 2016.
- [BS16] Mark Bun and Thomas Steinke. Concentrated differential privacy: Simplifications, extensions, and lower bounds. In Martin Hirt and Adam D. Smith, editors, *Theory of Cryptography Conference (TCC)*, 2016.
- [BST14] Raef Bassily, Adam D. Smith, and Abhradeep Thakurta. Private empirical risk minimization: Efficient algorithms and tight error bounds. In *IEEE Symposium on Foun*dations of Computer Science (FOCS), 2014.
- [Bub15] Sébastien Bubeck. Convex optimization: Algorithms and complexity. Foundations and Trends in Machine Learning, 8(3-4):231–357, 2015.
- [BW18] Borja Balle and Yu-Xiang Wang. Improving the gaussian mechanism for differential privacy: Analytical calibration and optimal denoising. In *International Conference on Machine Learning (ICML)*, 2018.
- [CKM<sup>+</sup>19] Clément L. Canonne, Gautam Kamath, Audra McMillan, Adam Smith, and Jonathan Ullman. The structure of optimal private tests for simple hypotheses. In ACM Symposium on Theory of Computing (STOC), 2019.
- [CMS11] Kamalika Chaudhuri, Claire Monteleoni, and Anand D. Sarwate. Differentially private empirical risk minimization. *Journal of Machine Learning Research*, 12(29):1069–1109, 2011.
- [CMW20] Michael Celentano, Andrea Montanari, and Yuchen Wu. The estimation error of general first order methods. In *Conference on Learning Theory (COLT)*, 2020.
- [CMW23] Michael Celentano, Andrea Montanari, and Yuting Wei. The Lasso with general Gaussian designs with applications to hypothesis testing. The Annals of Statistics, 51(5):2194 – 2220, 2023.
- [CS20] Emmanuel J. Candès and Pragya Sur. The phase transition for the existence of the maximum likelihood estimate in high-dimensional logistic regression. *The Annals of Statistics*, 48(1):27 42, 2020.

- [CWZ21] T. Tony Cai, Yichen Wang, and Linjun Zhang. The cost of privacy: Optimal rates of convergence for parameter estimation with differential privacy. The Annals of Statistics, 49(5):2825 – 2850, 2021.
- [CWZ23] T. Tony Cai, Yichen Wang, and Linjun Zhang. Score attack: A lower bound technique for optimal differentially private learning, 2023.
- [DFH<sup>+</sup>15] Cynthia Dwork, Vitaly Feldman, Moritz Hardt, Toniann Pitassi, Omer Reingold, and Aaron Leon Roth. Preserving statistical validity in adaptive data analysis. In ACM Symposium on Theory of Computing (STOC), 2015.
- [DKM<sup>+</sup>06] Cynthia Dwork, Krishnaram Kenthapadi, Frank McSherry, Ilya Mironov, and Moni Naor. Our data, ourselves: Privacy via distributed noise generation. In International Conference on the Theory and Applications of Cryptographic Techniques (EU-ROCRYPT), 2006.
- [DKT21] Zeyu Deng, Abla Kammoun, and Christos Thrampoulidis. A model of double descent for high-dimensional binary linear classification. *Information and Inference: A Journal* of the IMA, 11(2):435–495, 04 2021.
- [DL09] Cynthia Dwork and Jing Lei. Differential privacy and robust statistics. In ACM Symposium on Theory of Computing (STOC), 2009.
- [DL20] Oussama Dhifallah and Yue M. Lu. A precise performance analysis of learning with random features, 2020.
- [DLS23] Rishabh Dudeja, Yue M. Lu, and Subhabrata Sen. Universality of approximate message passing with semirandom matrices. *The Annals of Probability*, 51(5):1616 – 1683, 2023.
- [DMM09] David L Donoho, Arian Maleki, and Andrea Montanari. Message-passing algorithms for compressed sensing. Proceedings of the National Academy of Sciences, 106(45):18914–18919, 2009.
- [DMM11] David L Donoho, Arian Maleki, and Andrea Montanari. The noise-sensitivity phase transition in compressed sensing. *IEEE Transactions on Information Theory*, 57(10):6920–6941, 2011.
- [DMNS06] Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam D. Smith. Calibrating noise to sensitivity in private data analysis. In *Theory of Cryptography Conference* (*TCC*), 2006.
- [DR14] Cynthia Dwork and Aaron Roth. The algorithmic foundations of differential privacy. Foundations and Trends in Theoretical Computer Science, 9(3-4):211–407, 2014.
- [DR16] Cynthia Dwork and Guy N. Rothblum. Concentrated differential privacy, 2016.
- [DSL24] Rishabh Dudeja, Subhabrata Sen, and Yue M. Lu. Spectral universality in regularized linear regression with nearly deterministic sensing matrices. *IEEE Transactions on Information Theory*, 70(11):7923–7951, 2024.
- [EK18] Noureddine El Karoui. On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators. *Probability Theory and Related Fields*, 170(1):95–175, Feb 2018.
- [EKBB<sup>+</sup>13] Noureddine El Karoui, Derek Bean, Peter J Bickel, Chinghway Lim, and Bin Yu. On robust regression with high-dimensional predictors. *Proceedings of the National Academy of Sciences*, 110(36):14557–14562, 2013.
- [Fan22] Zhou Fan. Approximate Message Passing algorithms for rotationally invariant matrices. *The Annals of Statistics*, 50(1):197 224, 2022.
- [GH24] Kristian Georgiev and Samuel B. Hopkins. Privacy induces robustness: informationcomputation gaps and sparse mean estimation. In *Conference on Neural Information Processing Systems (NeurIPS)*, 2024.
- [GKK<sup>+</sup>20] Sivakanth Gopi, Gautam Kamath, Janardhan Kulkarni, Aleksandar Nikolov, Zhiwei Steven Wu, and Huanyu Zhang. Locally private hypothesis selection. In *Conference* on Learning Theory (COLT), 2020.
- [Gor85] Yehoram Gordon. Some inequalities for gaussian processes and applications. *Israel Journal of Mathematics*, 50(4):265–289, Dec 1985.
- [GTM<sup>+</sup>24] Cédric Gerbelot, Emanuele Troiani, Francesca Mignacco, Florent Krzakala, and Lenka Zdeborová. Rigorous dynamical mean-field theory for stochastic gradient descent methods. SIAM Journal on Mathematics of Data Science, 6(2):400–427, 2024.
- [Han24] Qiyang Han. Entrywise dynamics and universality of general first order methods, 2024.
- [HKM22] Samuel B. Hopkins, Gautam Kamath, and Mahbod Majid. Efficient mean estimation with pure differential privacy via a sum-of-squares exponential mechanism. In ACM Symposium on Theory of Computing (STOC), 2022.
- [HKMN23] Samuel B. Hopkins, Gautam Kamath, Mahbod Majid, and Shyam Narayanan. Robustness implies privacy in statistical estimation. In ACM Symposium on Theory of Computing (STOC), 2023.
- [HL22] Hong Hu and Yue M. Lu. Slope for sparse linear regression: Asymptotics and optimal regularization. *IEEE Transactions on Information Theory*, 68(11):7627–7664, 2022.
- [HS23] Qiyang Han and Yandi Shen. Universality of regularized regression estimators in high dimensions. *The Annals of Statistics*, 51(4):1799 1823, 2023.
- [Hub64] Peter J. Huber. Robust Estimation of a Location Parameter. The Annals of Mathematical Statistics, 35(1):73 – 101, 1964.
- [INS<sup>+</sup>19] Roger Iyengar, Joseph P. Near, Dawn Song, Om Thakkar, Abhradeep Thakurta, and Lun Wang. Towards practical differentially private convex optimization. In *IEEE Symposium on Security and Privacy*, 2019.
- [JT14] Prateek Jain and Abhradeep Guha Thakurta. (near) dimension independent risk bounds for differentially private learning. In *International Conference on Machine Learning (ICML)*, 2014.
- [Kar13] Noureddine El Karoui. Asymptotic behavior of unregularized and ridge-regularized high-dimensional robust regression estimators: rigorous results, 2013.
- [KBY24] Alexander Kent, Thomas B. Berrett, and Yi Yu. Rate optimality and phase transition for user-level local differential privacy, 2024.

- [KDH23] Rohith Kuditipudi, John Duchi, and Saminul Haque. A pretty fast algorithm for adaptive private mean estimation. In *Conference on Learning Theory (COLT)*, 2023.
- [KLSU19] Gautam Kamath, Jerry Li, Vikrant Singhal, and Jonathan Ullman. Privately learning high-dimensional distributions. In *Conference on Learning Theory (COLT)*, 2019.
- [KM11] Satish Babu Korada and Andrea Montanari. Applications of the lindeberg principle in communications and statistical learning. *IEEE Transactions on Information Theory*, 57(4):2440–2450, 2011.
- [KMS<sup>+</sup>22] Gautam Kamath, Argyris Mouzakis, Vikrant Singhal, Thomas Steinke, and Jonathan Ullman. A private and computationally-efficient estimator for unbounded gaussians. In Proceedings of Thirty Fifth Conference on Learning Theory (COLT), 2022.
- [KMV22] Pravesh Kothari, Pasin Manurangsi, and Ameya Velingker. Private robust estimation by stabilizing convex relaxations. In *Conference on Learning Theory (COLT)*, 2022.
- [KSSU19] Gautam Kamath, Or Sheffet, Vikrant Singhal, and Jonathan R. Ullman. Differentially private algorithms for learning mixtures of separated gaussians. In Conference on Neural Information Processing Systems (NeurIPS), 2019.
- [KST12] Daniel Kifer, Adam Smith, and Abhradeep Thakurta. Private convex empirical risk minimization and high-dimensional regression. In *Conference on Learning Theory* (COLT), 2012.
- [KSU20] Gautam Kamath, Vikrant Singhal, and Jonathan Ullman. Private mean estimation of heavy-tailed distributions. In *Conference on Learning Theory (COLT)*, 2020.
- [LGC<sup>+</sup>21] Bruno Loureiro, Cedric Gerbelot, Hugo Cui, Sebastian Goldt, Florent Krzakala, Marc Mezard, and Lenka Zdeborova. Learning curves of generic features maps for realistic datasets with a teacher-student model. In *Conference on Neural Information Process*ing Systems (NeurIPS), 2021.
- [LKKO21] Xiyang Liu, Weihao Kong, Sham M. Kakade, and Sewoong Oh. Robust and differentially private mean estimation. In Conference on Neural Information Processing Systems (NeurIPS), 2021.
- [LKO22] Xiyang Liu, Weihao Kong, and Sewoong Oh. Differential privacy and robust statistics in high dimensions. In *Conference on Learning Theory (COLT)*, 2022.
- [LS22] Tengyuan Liang and Pragya Sur. A precise high-dimensional asymptotic theory for boosting and minimum- $\ell_1$ -norm interpolated classifiers. The Annals of Statistics, 50(3):1669 1695, 2022.
- [Mir13] Darakhshan J. Mir. Differential privacy: An exploration of the privacy-utility landscape. PhD thesis, Rutgers University, 2013.
- [Mir17] Ilya Mironov. Rényi differential privacy. In *IEEE Computer Security Foundations* Symposium (CSF), 2017.
- [MJY24] Yuheng Ma, Ke Jia, and Hanfang Yang. Better locally private sparse estimation given multiple samples per user. In *International Conference on Machine Learning (ICML)*, 2024.

- [MM21] Léo Miolane and Andrea Montanari. The distribution of the Lasso: Uniform control over sparse balls and adaptive parameter tuning. *The Annals of Statistics*, 49(4):2313 2335, 2021.
- [MN17] Andrea Montanari and Phan-Minh Nguyen. Universality of the elastic net error. In 2017 IEEE International Symposium on Information Theory (ISIT), pages 2338–2342, 2017.
- [MRSY23] Andrea Montanari, Feng Ruan, Youngtak Sohn, and Jun Yan. The generalization error of max-margin linear classifiers: Benign overfitting and high dimensional asymptotics in the overparametrized regime, 2023.
- [Nar22] Shyam Narayanan. Private high-dimensional hypothesis testing. In *Conference on Learning Theory (COLT)*, 2022.
- [NRVW20] Seth Neel, Aaron Roth, Giuseppe Vietri, and Zhiwei Steven Wu. Oracle efficient private non-convex optimization. In International Conference on Machine Learning (ICML), 2020.
- [OT17] Samet Oymak and Joel A Tropp. Universality laws for randomized dimension reduction, with applications. Information and Inference: A Journal of the IMA, 7(3):337– 446, 11 2017.
- [PAA24] Alireza F. Pour, Hassan Ashtiani, and Shahab Asoodeh. Sample-optimal locally private hypothesis selection and the provable benefits of interactivity. In *Conference on Learning Theory (COLT)*, 2024.
- [PB14] Neal Parikh and Stephen P. Boyd. Proximal algorithms. Foundations and Trends in Optimization, 1(3):127–239, 2014.
- [PH17] Ashkan Panahi and Babak Hassibi. A universal analysis of large-scale regularized least squares solutions. In Conference on Neural Information Processing Systems (NeurIPS), 2017.
- [RKW23] Rachel Redberg, Antti Koskela, and Yu-Xiang Wang. Improving the privacy and practicality of objective perturbation for differentially private linear learners. In *Conference* on Neural Information Processing Systems (NeurIPS), 2023.
- [RW21] Rachel Redberg and Yu-Xiang Wang. Privately publishable per-instance privacy. In Conference on Neural Information Processing Systems (NeurIPS), 2021.
- [SAH19] Fariborz Salehi, Ehsan Abbasi, and Babak Hassibi. The impact of regularization on high-dimensional logistic regression. In Conference on Neural Information Processing Systems (NeurIPS), 2019.
- [SC19] Pragya Sur and Emmanuel J. Candès. A modern maximum-likelihood theory for high-dimensional logistic regression. Proceedings of the National Academy of Sciences, 116(29):14516-14525, 2019.
- [SCS13] Shuang Song, Kamalika Chaudhuri, and Anand D. Sarwate. Stochastic gradient descent with differentially private updates. In *IEEE Global Conference on Signal and Information Processing (GlobalSIP)*, 2013.

- [She17] Or Sheffet. Differentially private ordinary least squares. In International Conference on Machine Learning (ICML), 2017.
- [She19] Or Sheffet. Old techniques in differentially private linear regression. In International Conference on Algorithmic Learning Theory, 2019.
- [Sio58] Maurice Sion. On general minimax theorems. *Pacific Journal of Mathematics*, 8(1):171 176, 1958.
- [Sle62] David Slepian. The one-sided barrier problem for gaussian noise. The Bell System Technical Journal, 41(2):463–501, 1962.
- [Sto13] Mihailo Stojnic. A framework to characterize performance of lasso algorithms, 2013.
- [SU17] Thomas Steinke and Jonathan R. Ullman. Tight lower bounds for differentially private selection. In *IEEE Symposium on Foundations of Computer Science (FOCS)*, 2017.
- [TAH18] Christos Thrampoulidis, Ehsan Abbasi, and Babak Hassibi. Precise error analysis of regularized m-estimators in high dimensions. *IEEE Transactions on Information Theory*, 64(8):5592–5628, 2018.
- [TOH15] Christos Thrampoulidis, Samet Oymak, and Babak Hassibi. Regularized linear regression: A precise analysis of the estimation error. In *Conference on Learning Theory* (COLT), 2015.
- [Vad17] Salil P. Vadhan. The complexity of differential privacy. In Yehuda Lindell, editor, *Tutorials on the Foundations of Cryptography*, pages 347–450. Springer International Publishing, 2017.
- [Ver18] Roman Vershynin. High-Dimensional Probability: An Introduction with Applications in Data Science. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2018.
- [VTJ22] Prateek Varshney, Abhradeep Thakurta, and Prateek Jain. (nearly) optimal private linear regression for sub-gaussian data via adaptive clipping. In *Conference on Learning Theory (COLT)*, 2022.
- [Wai19] Martin J. Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2019.
- [Wan18] Yu-Xiang Wang. Revisiting differentially private linear regression: optimal and adaptive prediction & estimation in unbounded domain. In *Conference on Uncertainty in Artificial Intelligence (UAI)*, 2018.
- [WFS15] Yu-Xiang Wang, Stephen Fienberg, and Alex Smola. Privacy for free: Posterior sampling and stochastic gradient monte carlo. In *International Conference on Machine Learning (ICML)*, 2015.
- [WWM21] Shuaiwen Wang, Haolei Weng, and Arian Maleki. Does SLOPE outperform bridge regression? Information and Inference: A Journal of the IMA, 11(1):1–54, 11 2021.

- [WZF24] Tianhao Wang, Xinyi Zhong, and Zhou Fan. Universality of approximate message passing algorithms and tensor networks. *The Annals of Applied Probability*, 34(4):3943– 3994, 2024.
- [ZSC22] Qian Zhao, Pragya Sur, and Emmanuel J. Candès. The asymptotic distribution of the MLE in high-dimensional logistic models: Arbitrary covariance. *Bernoulli*, 28(3):1835 – 1861, 2022.
- [ZWC<sup>+</sup>22] Mingxun Zhou, Tianhao Wang, T-H. Hubert Chan, Giulia Fanti, and Elaine Shi. Locally differentially private sparse vector aggregation. In *IEEE Symposium on Security* and Privacy, 2022.
- [ZZY22] Xianyang Zhang, Huijuan Zhou, and Hanxuan Ye. A modern theory for highdimensional cox regression models, 2022.