

A cohomology-based Gromov-Hausdorff metric approach for quantifying molecular similarity

JunJie Wee^{1*}, Xue Gong^{2†}, Wilderich Tuschmann^{3‡}, and Kelin Xia^{2§}

¹Michigan State University, Department of Mathematics, East Lansing, MI 48824, USA

²Nanyang Technological University, Division of Mathematical Sciences, School of Physical and Mathematical Sciences, 637371, Singapore

³Karlsruher Institut für Technologie, Institut für Algebra und Geometrie, Arbeitsgruppe Differentialgeometrie, Karlsruhe, 76131, Germany

*weejunji@msu.edu

†xue.gong@ntu.edu.sg

‡wilderich.tuschmann@kit.edu

§xiakelin@ntu.edu.sg

ABSTRACT

We introduce, for the first time, a cohomology-based Gromov-Hausdorff ultrametric method to analyze 1-dimensional and higher-dimensional (co)homology groups, focusing on loops, voids, and higher-dimensional cavity structures in simplicial complexes, to address typical clustering questions arising in molecular data analysis. The Gromov-Hausdorff distance quantifies the dissimilarity between two metric spaces. In this framework, molecules are represented as simplicial complexes, and their cohomology vector spaces are computed to capture intrinsic topological invariants encoding loop and cavity structures. These vector spaces are equipped with a suitable distance measure, enabling the computation of the Gromov-Hausdorff ultrametric to evaluate structural dissimilarities. We demonstrate the methodology using organic-inorganic halide perovskite (OIHP) structures. The results highlight the effectiveness of this approach in clustering various molecular structures. By incorporating geometric information, our method provides deeper insights compared to traditional persistent homology techniques.

Introduction

In the last decade, concepts, methods, and techniques from topological and geometric data analysis^{1–4} have found profound applications, becoming indispensable tools in many areas of applied mathematics, such as image processing⁵, pattern recognition⁶, sensor network⁷, robotics⁸, computer graphics and vision⁹, cosmology¹⁰, medicine¹¹, computational and molecular biology¹². A central technique in Topological Data Analysis (TDA) is persistent homology, which provides a robust framework for capturing multi-scale topological features in data^{13,14}. However, while persistent homology is a powerful tool, it generally provides only limited information about the homology groups of the objects under study, often restricted to basic descriptors such as Betti numbers. In addition to topological tools, geometric theories and methods, such as discrete exterior calculus¹⁵, Laplace operators¹⁶, discrete Ricci curvatures^{17,18}, optimal transport¹⁹, parametric surfaces²⁰, and geometric flows²¹, have been widely applied in data analysis. These geometric approaches have often been shown to provide deeper insights than those achieved through purely topological methods.

A promising application of advanced geometric and topological techniques is molecular similarity analysis²², which is a fundamental concept in cheminformatics. Molecular similarity measures the resemblance between

molecules based on their structure, properties, or function, enabling applications in drug design²³, virtual screening²⁴, and reaction prediction²⁵. By assessing similarity, we can predict properties such as activity and toxicity, thereby accelerating the identification of promising compounds^{26,27}.

Existing methods for evaluating molecular similarity often rely on structural fingerprints, property-based metrics, or topological and 3D shape representations. For instance, structural fingerprints, such as Extended Connectivity Fingerprint (ECFP)²⁸ and MACCS keys²⁹, encode molecular features as binary sequences, which are typically compared using Tanimoto similarity³⁰. Property-based similarity involves comparing physicochemical attributes, such as molecular weight and polarity, using Euclidean or cosine distances³¹. Topological similarity, often evaluated through graph neural networks (GNNs)³², models molecular connectivity, while 3D similarity captures spatial geometry through metrics such as RMSD or Shape Tanimoto scores^{33,34}.

Recent advances in applied and computational topology and geometry offer powerful methods to assess molecular similarity. Persistent homology and its various refinements and extensions have proven invaluable for capturing molecular topological features in diverse chemical and biological applications^{35,36}. However, while effective in tracking the number of connected components, loops, and voids, persistent homology lacks the geometric information necessary to distinguish or identify individual features that arise during filtration. Previous studies have leveraged the connection between persistent homology and dendrograms^{37,38} to distinguish structures based on 0-dimensional homology groups, which correspond to connected components. These works employ geometric methods, such as the Gromov-Hausdorff distance, to measure the distance between two brain networks³⁹⁻⁴¹. However, there is still a need for methodologies that enable a more detailed and localized characterization of loops, voids, and higher-dimensional (co)homologies.

For the first time, we apply a cohomology-based Gromov-Hausdorff ultrametric method to study 1-dimensional and higher-dimensional (co)homology groups associated with loops, voids, and higher-dimensional cavity structures in simplicial complexes. The Gromov-Hausdorff distance (d_{GH})^{42,43} quantifies the dissimilarity between two metric spaces and possesses many desirable mathematical properties, making it widely used in differential geometry and applied algebraic topology for tasks such as shape matching, structural comparison, and characterization⁴⁴⁻⁴⁶. In the context of molecular structures A and B , modeled as simplicial complexes K_A and K_B , respectively, their cohomology groups give rise to the metric spaces $H^p(K_A)$ and $H^p(K_B)$. In theory, we can then compare their Gromov-Hausdorff distances. Although many lower bounds can be achieved in polynomial time^{46,47}, computing d_{GH} between arbitrary finite metric spaces is known to be NP-hard, as it leads to quadratic assignment problems⁴⁷. To address this challenge, we introduce a crucial second step: transforming $H^p(K_A)$ and $H^p(K_B)$ into ultrametric spaces, thus obtaining their respective dendrogram representations^{37,40,48}. This enables us to compute a more tractable ultrametric variation of the Gromov-Hausdorff distance, referred to as the Gromov-Hausdorff ultrametric (u_{GH})^{48,49}, which can be used to quantify molecular similarity.

The main contributions of this work are as follows:

1. We present the first workflow to utilize a Gromov-Hausdorff ultrametric approach based on 1-dimensional and higher-dimensional cohomology for assessing structural similarity between molecules, effectively capturing local topological features such as loops and voids that influence molecular properties.
2. We demonstrate the effectiveness of this workflow in clustering tasks using a dataset of organic-inorganic halide perovskites (OIHP).

The remainder of this paper is structured as follows: The next section introduces foundational concepts in algebraic topology, including simplicial complexes and the Hodge Laplacian, which are essential to our methodology. We then present our method for measuring structural similarity, followed by numerical experiments to validate our approach using OIHP material datasets. Finally, we discuss potential future

research directions.

Background

Simplicial complexes

In many applications, molecular data from biology, chemistry, and material science can be represented as graphs^{50,51}. In this framework, the vertices correspond to atoms, while the edges represent affinities between pairs of atoms. Simplicial complexes generalize graphs by capturing higher-order interactions through simplices, such as triangles and tetrahedra. In this and the next section, we aim to provide a brief introduction to the theoretical background necessary for performing computations on simplicial complexes. For a more comprehensive review, we direct readers to⁵²⁻⁵⁵.

If we let V be a finite set of vertices, then a p -simplex σ^p is a subset of V with $p + 1$ vertices. For example, a simplex of dimensions 0, 1, 2, and 3 can be viewed as a point, an edge, a triangle, and a tetrahedron, respectively. More precisely, a p -simplex $\sigma^p = \{v_0, v_1, v_2, \dots, v_p\}$ is defined as a convex hull formed by its $p + 1$ affinely independent points $v_0, v_1, v_2, \dots, v_p$ as follows:

$$\sigma^p = \left\{ \lambda_0 v_0 + \lambda_1 v_1 + \dots + \lambda_p v_p \mid \sum_{i=0}^p \lambda_i = 1; \forall i, 0 \leq \lambda_i \leq 1 \right\}.$$

A m -face of a p -simplex σ^p is defined as a convex hull formed by $m + 1$ vertices of σ^p , where $m < p$. If σ^m is a face of σ^p , denoted by $\sigma^m \subset \sigma^p$, then σ^p is also referred to as a *coface* of σ^m . The *upper degree* of a p -simplex σ^p , denoted by $\text{deg}(\sigma^p)$, is the number of $(p + 1)$ -simplices for which σ^p is a face. Two simplices σ_1^p and σ_2^p are *upper adjacent* and denoted by $\sigma_1^p \frown \sigma_2^p$ if they have a common coface. They are *lower adjacent* and denoted by $\sigma_1^p \smile \sigma_2^p$ if they share a common face. To facilitate computation, each simplex is assigned an orientation associated with the ordering of its vertices. If two p -simplices σ_1^p and σ_2^p are oriented similarly, then we write $\sigma_1^p \sim \sigma_2^p$. Conversely, we write $\sigma_1^p \not\sim \sigma_2^p$ if they have opposite orientations.

A p -dimensional simplicial complex K contains up to p -dimensional simplices and satisfies two conditions. First, any face of a simplex from K is also in K . Second, the intersection of any two simplices in K is either empty or a shared face. There are many common types of simplicial complexes such as Vietoris-Rips complex, Čech complex, Alpha complex, and clique complex.

The Combinatorial Hodge Laplacian

Computationally, combinatorial Hodge Laplacian or discrete Hodge Laplacian has been proposed^{52,54-60}. Essentially, this discrete version can be viewed as part of exterior calculus and discrete differential geometry. The concept of the combinatorial Hodge Laplacian is a part of Hodge theory, which has recently been applied to biomolecular data analysis⁶¹. Hodge Laplacian matrices of different dimensions can be constructed on a simplicial complex. A p -th dimensional Hodge Laplacian matrix characterizes topological connections between p -th simplices within a simplicial complex^{52,53,62}. Note that the graph Laplacian characterizes relations between vertices (0-simplices).

The p -th chain group $C_p(K)$ of a simplicial complex K over some field \mathbb{F} is a vector space over \mathbb{F} whose basis is the set of p -simplices of the simplicial complex K . Elements of $C_p(K)$ are called p -chains. The *dual* of $C_p(K)$, denoted by $C^p(K)$, is the set of all linear functionals on $C_p(K)$:

$$C^p(K) = \{ \phi : C_p(K) \rightarrow \mathbb{F} : \phi \text{ is linear} \}.$$

$C^p(K)$ is called the p -th *cochain group* and its elements are called p -*cochains*. Boundary operators are defined on both the chain and cochain groups. The *boundary map* ∂_p is a linear transformation which acts

on a p -simplex $\sigma^p = [u_0, u_1, \dots, u_p]$ as follows

$$\partial_p([u_0, u_1, \dots, u_p]) = \sum_{i=0}^p (-1)^i [u_0, \dots, u_{i-1}, u_{i+1}, \dots, u_p].$$

The *coboundary map* $\delta_p : C^p(K) \rightarrow C^{p+1}(K)$ is a linear transformation defined as follows: for a linear functional $\phi \in C^p(K)$ and a $p+1$ -simplex $\sigma^{p+1} = [u_0, u_1, \dots, u_{p+1}]$,

$$\delta_p(\phi)(\sigma^{p+1}) = \sum_{i=0}^{p+1} (-1)^i \phi([u_0, \dots, u_{i-1}, u_{i+1}, \dots, u_{p+1}]).$$

The boundary map gives rise to a *chain complex*, which is a sequence of chain groups connected by boundary maps as follows:

$$0 \rightarrow C_n(K) \rightarrow \dots \xrightarrow{\partial_{p+1}} C_p(K) \xrightarrow{\partial_p} C_{p-1}(K) \dots \xrightarrow{\partial_2} C_1(K) \xrightarrow{\partial_1} C_0(K) \rightarrow 0.$$

Similar to the boundary map giving rise to the chain complex, the coboundary operator gives rise to a *cochain complex*:

$$0 \leftarrow C^n(K) \leftarrow \dots \xleftarrow{\delta_p} C^p(K) \xleftarrow{\delta_{p-1}} C^{p-1}(K) \dots \xleftarrow{\delta_1} C^1(K) \xleftarrow{\delta_0} C^0(K) \leftarrow 0.$$

Since $C_p(K)$ and $C^p(K)$ are finite-dimensional, there exists unique matrix representations for ∂_p and δ_p . We have some useful relations regarding matrix representations of ∂_p and δ_p (A^T represents the transpose of a matrix A):

- For all $p \geq 0$, $\partial_{p+1}^T = \delta_p$,
- $\partial_p^T = \delta_p^*$,
- $\delta_p^T = \delta_p^*$.

Here, $\delta_p^* : C^{p+1}(K) \rightarrow C^p(K)$ is the *adjoint/transpose map* of δ_p where

$$\langle \delta_p(f), g \rangle = \langle f, \delta_p^*(g) \rangle,$$

for every $f \in C^p(K)$, $g \in C^{p+1}(K)$ and a suitable inner product $\langle \cdot, \cdot \rangle$ for $C^p(K)$ and $C^{p+1}(K)$. The adjoint of the boundary operator ∂_p , ∂_p^* is also defined analogously. Using the cochain group $C^p(K)$, we can define the p -th cocycle group Z^p and p -th coboundary group B^p as follows:

$$Z^p = \text{Ker}(\delta_p) = \{c \in C^p \mid \delta_p(c) = 0\},$$

$$B^p = \text{Im}(\delta_{p-1}) = \{c \in C^p \mid \exists d \in C^{p-1} : c = \delta_{p-1}(d)\}.$$

Then we have the p -th cohomology group $H^p = Z^p / B^p$.

The p -dimensional *combinatorial Hodge Laplacian* is the linear operator $\Delta_p : C^p(K) \rightarrow C^p(K)$ is defined as follows:

$$\Delta_p = \begin{cases} \delta_p^* \circ \delta_p + \delta_{p-1} \circ \delta_{p-1}^* & \text{if } p \geq 1, \\ \delta_p^* \circ \delta_p & \text{if } p = 0. \end{cases}$$

The case where $p = 0$ gives rise to the expression of the well-known graph Laplacian. Alternatively, the combinatorial Hodge Laplacian matrix can be expressed using the boundary matrix. The boundary operator

∂_p has a unique matrix representation. Given a simplicial complex K , the p -th boundary matrix \mathbf{B}_p is defined as,

$$(\mathbf{B}_p)_{ij} = \begin{cases} 1 & \text{if } \sigma_i^{p-1} \subset \sigma_j^p \text{ and } \sigma_i^{p-1} \sim \sigma_j^p, \\ -1 & \text{if } \sigma_i^{p-1} \subset \sigma_j^p \text{ and } \sigma_i^{p-1} \not\sim \sigma_j^p, \\ 0 & \text{if } \sigma_i^{p-1} \not\subset \sigma_j^p, \end{cases}$$

where σ_i^{p-1} is the i -th $(p-1)$ -simplex and σ_j^p is the j -th p -simplex.

Given that the highest order of the simplicial complex K is n , the p -th Hodge Laplacian (or combinatorial Laplacian) matrix \mathbf{L}_p of K is

$$\mathbf{L}_p = \begin{cases} \mathbf{B}_n^T \mathbf{B}_n & \text{if } p = n, \\ \mathbf{B}_p^T \mathbf{B}_p + \mathbf{B}_{p+1} \mathbf{B}_{p+1}^T & \text{if } 1 \leq p < n, \\ \mathbf{B}_1 \mathbf{B}_1^T & \text{if } p = 0. \end{cases}$$

Another way to define the Hodge Laplacian is through simplex relations. When $p = 0$,

$$(\mathbf{L}_0)_{ij} = \begin{cases} \deg(\sigma_i^0) & \text{if } i = j, \\ -1 & \text{if } i \neq j \text{ and } \sigma_i^0 \frown \sigma_j^0, \\ 0 & \text{if } i \neq j \text{ and } \sigma_i^0 \not\frown \sigma_j^0, \end{cases}$$

and \mathbf{L}_0 is exactly the same as the graph Laplacian matrix. When $p > 0$,

$$(\mathbf{L}_p)_{ij} = \begin{cases} \deg(\sigma_i^p) + p + 1 & \text{if } i = j, \\ 1 & \text{if } i \neq j, \sigma_i^p \not\sim \sigma_j^p, \sigma_i^p \smile \sigma_j^p \text{ and } \sigma_i^k \sim \sigma_j^p, \\ -1 & \text{if } i \neq j, \sigma_i^p \not\sim \sigma_j^p, \sigma_i^p \smile \sigma_j^p \text{ and } \sigma_i^k \not\sim \sigma_j^p, \\ 0 & \text{if } i \neq j, \sigma_i^p \frown \sigma_j^p \text{ or } \sigma_i^p \not\sim \sigma_j^p. \end{cases}$$

Mathematically, the eigenvalues of Hodge Laplacian matrices are independent of the choice of the orientation⁵⁷. The eigenspectrum of Hodge Laplacian matrices reveals topological information within simplicial complexes. For instance, the multiplicity of zero eigenvalues of \mathbf{L}_p corresponds to the Betti numbers β_p , reflecting the number of connected components (β_0), the number of cycles (β_1), the number of cavities (β_2), etc. Furthermore, the eigenvectors corresponding to the zero eigenvalues of \mathbf{L}_p , also referred to as cohomology generators, can be used to illustrate the p -dimensional cohomologies. Let n_p be the number of p -simplices in a simplicial complex K . Then, for any cohomology generator v of \mathbf{L}_p , we can arrange the entries of v to match the order of simplices in K . Specifically, the i -th entry v_i corresponds to the p -simplex σ_i^p in K . This allows us to visualize v on a simplicial complex.

The Gromov-Hausdorff Distance

The Gromov–Hausdorff distance measures the distance between two compact metric spaces^{42,43}. It measures the smallest distance at which two compact metric spaces can be considered “close”, which is particularly useful when the overall shape matching is more relevant than exact pointwise alignment. Given two metric spaces X and Y , the Gromov–Hausdorff distance looks for all possible isometric embeddings of X and Y into a common metric space Z and then calculates the Hausdorff distance⁶³ between these embeddings within Z . Formally, the Gromov-Hausdorff distance d_{GH} ^{42,43} between two compact metric spaces X and Y is defined as

$$d_{GH}(X, Y) = \inf d_H^Z(\varphi(X), \psi(Y)),$$

where the infimum is taken over all possible $Z \in \mathcal{M}$ and isometric embeddings $\varphi : X \hookrightarrow Z$ and $\psi : Y \hookrightarrow Z$, and d_H^Z denotes Hausdorff distance in Z .

Method

Distance Metrics for Cohomology Generators

For a given simplicial complex K , we denote the space consisting of all the p -dimensional cohomology generators arising from its Hodge Laplacian \mathbf{L}_p by $H^p(K)$. Subsequently, we want to construct a metric space from $H^p(K)$ by assigning distances between cohomology generators in $H^p(K)$. We will introduce three types of distance measures to construct the metric space: L_1 distance, cocycle distance, and Wasserstein distance.

Throughout this section, we assume that any two cohomology generators v and w from a Hodge Laplacian matrix \mathbf{L}_p must have a consistent order of entries. In other words, we can write

$$v = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_{n_p-1} \\ v_{n_p} \end{pmatrix} \begin{matrix} \sigma_1^p \\ \sigma_2^p \\ \vdots \\ \sigma_{n_p-1}^p \\ \sigma_{n_p}^p \end{matrix} \quad \text{and} \quad w = \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_{n_p-1} \\ w_{n_p} \end{pmatrix} \begin{matrix} \sigma_1^p \\ \sigma_2^p \\ \vdots \\ \sigma_{n_p-1}^p \\ \sigma_{n_p}^p \end{matrix},$$

where v_i and w_i are entries in v and w , respectively, corresponding to the simplex σ_i^p for all $1 \leq i \leq n_p$. Furthermore, we assume that any cohomology generator in $H^p(K)$ is normalized.

L_1 Distance

First, we define the L_1 distance between two cohomology generators as follows.

Definition 1 (L_1 distance). Let v and w be two cohomology generators from $H^p(K)$ where K is a simplicial complex. Then the L_1 distance between v and w is

$$\|v - w\|_1 = \sum_{i=1}^{n_p} |v_i - w_i|.$$

Note that in computation, the eigenvectors may have arbitrary signs. To eliminate the sign ambiguity introduced by the solver, we enforce that the first element of each cohomology generator is non-negative.

Cocycle Distance

As the L_1 norm of a cohomology generator tends to be larger when the cocycle contains more edges, it serves as a rough indicator of the size of the cocycle. Therefore, it is natural to consider another type of distance that measures the absolute difference between the L_1 norms of two cohomology generators. We refer to this as the *cocycle distance*.

Definition 2 (Cocycle distance). Let v and w be two cohomology generators in $H^p(K)$ where K is a simplicial complex. Their cocycle distance $d_s(v, w)$ is defined as the absolute difference in their L_1 norms:

$$d_s(v, w) = \left| \|v\|_1 - \|w\|_1 \right|.$$

For illustration, consider the BINOL-phosphoramidate **1_iv**, a BINOL (1,1'-bi-2-naphthol)-based phosphoric acid catalyst. We construct the atomic level simplicial complex representation of the molecule, where each vertex represents an atom, and generate the corresponding Hodge Laplacian matrices \mathbf{L}_0 and \mathbf{L}_1 .




Figure 1. Illustration of a catalyst 1_iv with its \mathbf{L}_0 and \mathbf{L}_1 cohomology generators. The \mathbf{L}_0 cohomology generator shows all the 0-simplices having equal value as it corresponds to the β_0 , which represents the connected components of the catalyst. The \mathbf{L}_1 matrix has 7 cohomology generators in total, each representing a unique cocycle within the catalyst.

The cohomology generators from \mathbf{L}_0 and \mathbf{L}_1 are depicted in Figure 1, where the color of each simplex σ_i^p is darker in blue if its associated $|v_i|$ in the cohomology generator is larger. From \mathbf{L}_0 , there is only one cohomology generator, representing a single connected component of the catalyst structure. Therefore, the cohomology generator in \mathbf{L}_0 assigns equal values to all vertices. On the other hand, \mathbf{L}_1 produces 7 cohomology generators highlighted in dark blue in Figure 1.

Each cohomology generator is a unique harmonic representative of a cohomology class. Loosely speaking, one can visualize all cycles using a combination of these cohomology generators and the boundaries of 2-simplices. In addition, the non-cohomology generators associated with non-zero eigenvalues reveal local and global clustering patterns. The Fiedler vector effectively clusters the vertices into two groups, colored in blue and red. For example, Figure 1 shows the Fiedler vector corresponding to the smallest non-zero eigenvalue of the Hodge Laplacian matrix, which clusters the vertices into two groups colored in blue and red.

Wasserstein Distance

Recall from the beginning of this section that any cohomology generator v in $H^p(K)$ is assumed to be normalized. Hence, given any two cohomology generators v and w in $H^p(K)$, we can square all the entries of v and w to obtain vectors v' and w' , whose entries sum to 1,

$$\text{i.e. } v' = \begin{pmatrix} v_1^2 \\ v_2^2 \\ \vdots \\ v_{n_p-1}^2 \\ v_{n_p}^2 \end{pmatrix} \begin{matrix} \sigma_1^p \\ \sigma_2^p \\ \vdots \\ \sigma_{n_p-1}^p \\ \sigma_{n_p}^p \end{matrix} \quad \text{and} \quad w' = \begin{pmatrix} w_1^2 \\ w_2^2 \\ \vdots \\ w_{n_p-1}^2 \\ w_{n_p}^2 \end{pmatrix} \begin{matrix} \sigma_1^p \\ \sigma_2^p \\ \vdots \\ \sigma_{n_p-1}^p \\ \sigma_{n_p}^p \end{matrix}. \quad (1)$$

Since the vectors v' and w' have entries that sums to 1, this allows us to treat the values in v' and w' as two probability measures m_1 and m_2 respectively. Now, we can define the pairwise distance between any two cohomology generators v and w to be the Wasserstein distance^{64, 65} between m_1 and m_2 .

Definition 3 (Wasserstein distance). The probability measures m_1 and m_2 are the probability distributions obtained from the entries of v' and w' as defined in (1). Let σ_i^p and σ_j^p be p -simplices in K . Then, $\xi(\sigma_i^p, \sigma_j^p)$

represents the amount of mass traveling from σ_i^p to σ_j^p . We require that the transportation from m_1 to m_2 is mass-preserving, i.e., $\sum_{\sigma_i^p \in K} \xi(\sigma_i^p, \sigma_j^p) = m_1(\sigma_i^p)$ and $\sum_{\sigma_j^p \in K} \xi(\sigma_i^p, \sigma_j^p) = m_2(\sigma_j^p)$. The Wasserstein distance between m_1 and m_2 , denoted by $W_1(m_1, m_2)$, is the minimum traveling distance that can be achieved, given by

$$W_1(m_1, m_2) = \inf_{\xi} \sum_{\sigma_i^p \in K} \sum_{\sigma_j^p \in K} d(\sigma_i^p, \sigma_j^p) \xi(\sigma_i^p, \sigma_j^p). \quad (2)$$

The distance between two simplices, denoted by $d(\sigma_i^p, \sigma_j^p)$, is defined as the minimum ℓ^2 distance between their corresponding vertices. Representing the sets of vertices in the simplices as $\sigma_i^p = \{x_1, x_2, \dots, x_p\}$ and $\sigma_j^p = \{u_1, u_2, \dots, u_p\}$, the distance is then given by

$$d(\sigma_i^p, \sigma_j^p) = \min_{\substack{1 \leq i \leq p \\ 1 \leq j \leq p}} \|\mathbf{x}_i - \mathbf{u}_j\|_2, \quad (3)$$

where \mathbf{x}_i and \mathbf{u}_j refer to the coordinates of the vertices x_i and u_j in \mathbb{R}^3 .

Cohomology-based Gromov–Hausdorff Ultrametric Approach for Structural Similarity

Given two molecular structures, A and B , we construct simplicial complexes, denoted by K_A and K_B respectively, where vertices represent atoms. There are various approaches for constructing the simplicial complex⁶⁶. For example, we can build a Vietoris–Rips complex such that a set of atoms is a simplex in K if every atom pair in the set has a ℓ^2 distance smaller than a specified filtration threshold. An alternative approach is to generate an Alpha complex, where a set of vertices forms a simplex when its filtration value is smaller than the threshold. For example, the filtration value of an atom pair is one-half of their ℓ^2 distance, and the filtration value for a triplet of atoms is the radius of the circle that passes through it.

After constructing the simplicial complex representation of a molecular structure, cohomology generators can be computed from its 1-dimensional Hodge Laplacian matrix \mathbf{L}_1 . Subsequently, we can compute a distance matrix comprised of pairwise distances between all cohomology generators from \mathbf{L}_1 . Three types of distance matrices can be derived for each molecular structure, corresponding to Definition 1, 2, or 3.

In theory, we can compare the Gromov-Hausdorff distances between the metric spaces formed by the cohomology generators using a selected distance measure. However, it has been proven that computing d_{GH} is NP-hard⁶⁷. Therefore, we will adopt a more tractable ultrametric variation of the Gromov-Hausdorff distance, called the Gromov-Hausdorff ultrametric, which can be computed in polynomial time⁴⁸.

An ultrametric space (X, d_X) is a metric space that satisfies the strong triangle inequality:

$$\forall x, x', x'' \in X, \text{ one has } d_X(x, x') \leq \max(d_X(x, x''), d_X(x'', x')).$$

The Gromov-Hausdorff ultrametric⁴⁹, denoted by u_{GH} , measures distances between compact ultrametric spaces X and Y as follows.

Definition 4 (The Gromov-Hausdorff ultrametric⁴⁹). The Gromov-Hausdorff ultrametric u_{GH} between *compact* ultrametric spaces X and Y is

$$u_{\text{GH}}(X, Y) := \inf d_H^Z(\varphi_X(X), \varphi_Y(Y)),$$

where the infimum is taken over all *ultrametric* spaces Z and isometric embeddings $\varphi_X : X \hookrightarrow Z$ and $\varphi_Y : Y \hookrightarrow Z$.

In our methodology, we transform metric spaces formed by the cohomology generators into an ultrametric space denoted by $H^1(K_A)$ and $H^1(K_B)$ using Algorithm 1 in⁴⁸. This transformation allows us to compute the Gromov-Hausdorff ultrametric $u_{\text{GH}}(H^1(K_A), H^1(K_B))$ between two ultrametric spaces associated with molecular structures A and B , thereby quantifying their structural similarity.




Figure 2. Illustration of 9 types of OIHP structures with formula MAPbX_3 , where MA refers to Methylammonium, Pb is lead and X is bromine (Br), chlorine (Cl) or iodine (I). (A) shows the three OIHP structures with bromine, chlorine, and iodine. (B) shows the three phases of OIHP structures, i.e. cubic, orthorhombic, and tetragonal. In total, we consider all 9 possible combinations. (C) Illustration of the construction of the Alpha complex from MAPbBr_3 structures in the cubic phase (left) with a filtration value of 3.5 Å (middle) and 4 Å (right), where the edges are colored by the absolute value of a cohomology generator, with red corresponding to a larger absolute value.

Results

Characterization of OIHP Material Structures

Organic-inorganic hybrid perovskite (OIHP) materials are favorable candidates for developing efficient and cost-effective solar cells. The stabilization of the OIHP structure is reliant on Van der Waals interactions and hydrogen bonding effects, which are closely associated with the distances between organic molecules and inorganic ions⁶⁸. Our u_{GH} -based features can be employed for characterizing OIHP material structures. To demonstrate this, we examine the clustering of Methylammonium lead halides (MAPbX_3 , X=Cl, Br, I). Three possible phases of MAPbX_3 are considered for each X-site atom, including the orthorhombic, tetragonal, and cubic phases as illustrated in Figure 2 (A)-(B).

We analyze 100 configurations from the molecular dynamics (MD) trajectories for each of the 9 OIHP structures, leading to 900 trajectories. Each MD trajectory arises from a molecular dynamics simulation to stabilize the initial configuration and consists of the 3D coordinates of atoms at a specific time. For each configuration, we construct an Alpha complex with 5 filtration thresholds of 3 Å, 3.5 Å, 4 Å, 5 Å, and 6 Å as shown in Figure 2 (C) and calculate its 1-dimensional co-homology generators and associated pairwise distances using L_1 distance as defined in Definition 1. We chose to construct alpha complexes, which are

based on the Delaunay triangulation of the points, because they better reflect the underlying geometry of atomic arrangement.

We consider the tasks of clustering 3 types of atoms for X-sites for each possible phase of MAPbX_3 , by calculating the u_{GH} between the 300 configurations of interest with the same phase for each of the 5 filtration values. This process generates a GH-based statistical feature vector with a length of $300 \times 5 = 1500$ for each configuration, allowing for differentiation among the three X-sites. In Figure 3 we compare the clustering result obtained using u_{GH} -based features with the ones achieved using the 3D coordinates of configurations. The clustering is performed utilizing t-SNE, with the x-axis and y-axis representing the two principal dimensions in the t-SNE model. Directly using the 3D coordinates proves ineffective in distinguishing between various X-site atoms. On the other hand, u_{GH} -based methods exhibit superior performance in clustering structures by X-site atoms.




Figure 3. The clustering of OIHP molecular configurations. Two clustering tasks are considered: (A) clustering of X-site atoms and (B) clustering of 9 types of OIHP structures. The t-SNE model is employed for clustering, where the x-axis and y-axis represent the two principal dimensions in the t-SNE model. We compare the clustering using 3D coordinates with those utilizing u_{GH} -based features with the L_1 distance. Using the 3D coordinates alone is insufficient for distinguishing between different X-site atoms and OIHP structures. Methods using u_{GH} -based features prove effective in both clustering tasks.

Discussion

In this work, we propose a new framework that employs the cohomology-based Gromov-Hausdorff ultrametric (u_{GH}) for quantifying structural similarity between molecular structures. In our workflow, we build simplicial complex representations of molecular structures and compute the u_{GH} between their respective cohomology

generator spaces. The cohomology generators effectively encode the local topological invariants, revealing cyclic patterns formed by edges. We illustrate the application of the cohomology-based Gromov-Hausdorff distance using organic-inorganic halide perovskites (OIHP) data. In our numerical experiments, the u_{GH} -based approach demonstrated effectiveness in clustering various structures while relying solely on the 3D coordinates proved to be insufficient.

We demonstrate the application of the cohomology-based u_{GH} approach in quantifying structural similarities. However, there are some limitations to our current work. For instance, we focused solely on the cohomology generators of the first-order Hodge Laplacian, which capture loop patterns. In the future, exploring non-cohomology generators may reveal clustering information, and higher-order Hodge Laplacians could unveil more complex structures such as cavities. Other potential future directions involve incorporating cohomology-based u_{GH} features into machine learning models for structure design and prediction. Additionally, our numerical experiments are limited to small molecules from chemistry and physics with hundreds of atoms. A promising avenue involves applying the proposed method to larger biological molecules with tens of thousands of molecules. For instance, it could be employed to quantify structural similarities between protein structures in drug design⁶⁹.

References

1. Carlsson, G. Topology and data. *Bull. Am. Math. Soc.* **46**, 255–308 (2009).
2. Wasserman, L. Topological data analysis. *Annu. Rev. Stat. Its Appl.* **5**, 501–532 (2018).
3. Le Roux, B. & Rouanet, H. *Geometric data analysis: from correspondence analysis to structured data analysis* (Springer Science & Business Media, 2004).
4. Kirby, M. *Geometric data analysis: an empirical approach to dimensionality reduction and the study of patterns* (John Wiley & Sons, Inc., 2000).
5. Klette, R. & Rosenfeld, A. *Digital geometry: Geometric methods for digital picture analysis* (Elsevier, 2004).
6. Carlsson, G. Topological pattern recognition for point cloud data. *Acta Numer.* **23**, 289–368 (2014).
7. Gao, J. & Guibas, L. Geometric algorithms for sensor networks. *Philos. Transactions Royal Soc. A: Math. Phys. Eng. Sci.* **370**, 27–51 (2012).
8. Bhattacharya, S. Search-based path planning with homotopy class constraints. *Proc. AAAI Conf. on Artif. Intell.* **24**, 1230–1237 (2010).
9. Skraba, P., Ovsjanikov, M., Chazal, F. & Guibas, L. Persistence-based segmentation of deformable shapes. In *2010 IEEE Computer Society Conference on Computer Vision and Pattern Recognition-Workshops*, 45–52 (IEEE, 2010).
10. Sousbie, T. The persistent cosmic web and its filamentary structure—i. theory and implementation. *Mon. Notices Royal Astron. Soc.* **414**, 350–383 (2011).
11. Skaf, Y. & Laubenbacher, R. Topological data analysis in biomedicine: A review. *J. Biomed. Informatics* **130**, 104082 (2022).
12. Yao, Y. *et al.* Topological methods for exploring low-density states in biomolecular folding pathways. *The J. Chem. Phys.* **130** (2009).
13. Edelsbrunner, Letscher & Zomorodian. Topological persistence and simplification. *Discret. & Comput. Geom.* **28**, 511–533 (2002).
14. Zomorodian, A. & Carlsson, G. Computing persistent homology. In *Proceedings of the twentieth annual symposium on Computational geometry*, 347–356 (2004).
15. Hirani, A. N. *Discrete exterior calculus* (California Institute of Technology, 2003).

16. Belkin, M. & Niyogi, P. Laplacian eigenmaps for dimensionality reduction and data representation. *Neural Comput.* **15**, 1373–1396 (2003).
17. Ollivier, Y. Ricci curvature of markov chains on metric spaces. *J. Funct. Analysis* **256**, 810–864 (2009).
18. Forman. Bochner's method for cell complexes and combinatorial ricci curvature. *Discret. & Comput. Geom.* **29**, 323–374 (2003).
19. Villani, C. *et al.* *Optimal transport: old and new*, vol. 338 (Springer, 2009).
20. Farin, G. *Curves and surfaces for computer-aided geometric design: a practical guide* (Elsevier, 2014).
21. Jin, M., Kim, J., Luo, F. & Gu, X. Discrete surface ricci flow. *IEEE Transactions on Vis. Comput. Graph.* **14**, 1030–1043 (2008).
22. Bender, A. & Glen, R. C. Molecular similarity: a key technique in molecular informatics. *Org. & Biomol. Chem.* **2**, 3204–3218 (2004).
23. Vilar, S. *et al.* Drug—drug interaction through molecular structure similarity analysis. *J. Am. Med. Informatics Assoc.* **19**, 1066–1074 (2012).
24. Eckert, H. & Bajorath, J. Molecular similarity analysis in virtual screening: foundations, limitations and novel approaches. *Drug Discov. Today* **12**, 225–233 (2007).
25. Arteca, G. A., Jammal, V. B., Mezey, P. G. & Mezey, P. G. Shape group studies of molecular similarity and regioselectivity in chemical reactions. *J. Comput. Chem.* **9**, 608–619 (1988).
26. Petrone, P. M. *et al.* Rethinking molecular similarity: comparing compounds on the basis of biological activity. *ACS Chem. Biol.* **7**, 1399–1409 (2012).
27. Basak, S. C., Grunwald, G. D., Host, G. E., Niemi, G. J. & Bradbury, S. P. A comparative study of molecular similarity, statistical, and neural methods for predicting toxic modes of action. *Environ. Toxicol. Chem. An Int. J.* **17**, 1056–1064 (1998).
28. Rogers, D. & Hahn, M. Extended-connectivity fingerprints. *J. Chem. Inf. Model.* **50**, 742–754 (2010).
29. Polton, D. Installation and operational experiences with maccs (molecular access system). *Online Rev.* **6**, 235–242 (1982).
30. Bajusz, D., Rácz, A. & Héberger, K. Why is Tanimoto index an appropriate choice for fingerprint-based similarity calculations? *J. Cheminformatics* **7**, 1–13 (2015).
31. Todeschini, R. & Consonni, V. *Handbook of molecular descriptors* (John Wiley & Sons, 2008).
32. Gilmer, J., Schoenholz, S. S., Riley, P. F., Vinyals, O. & Dahl, G. E. Neural message passing for quantum chemistry. In *International conference on machine learning*, 1263–1272 (PMLR, 2017).
33. Grant, J. A., Gallardo, M. A. & Pickup, B. T. A fast method of molecular shape comparison: A simple application of a gaussian description of molecular shape. *J. Comput. Chem.* **17**, 1653–1666 (1996).
34. Good, A. C. & Richards, W. G. Rapid evaluation of shape similarity using gaussian functions. *J. Chem. Inf. Comput. Sci.* **33**, 112–116 (1993).
35. Townsend, J., Micucci, C. P., Hymel, J. H., Maroulas, V. & Vogiatzis, K. D. Representation of molecular structures with persistent homology for machine learning applications in chemistry. *Nat. Commun.* **11**, 3230 (2020).
36. Xia, K. & Wei, G.-W. Persistent homology analysis of protein structure, flexibility, and folding. *Int. J. for Numer. Methods Biomed. Eng.* **30**, 814–844 (2014).
37. Carlsson, G. E., Mémoli, F. *et al.* Characterization, stability and convergence of hierarchical clustering methods. *J. Mach. Learn. Res.* **11**, 1425–1470 (2010).
38. Carlsson, G. & Mémoli, F. Persistent clustering and a theorem of j. kleinberg. *arXiv preprint arXiv:0808.2241* (2008).

39. Lee, H., Chung, M. K., Kang, H., Kim, B.-N. & Lee, D. S. Computing the shape of brain networks using graph filtration and gromov-hausdorff metric. In *Medical Image Computing and Computer-Assisted Intervention—MICCAI 2011: 14th International Conference, Toronto, Canada, September 18-22, 2011, Proceedings, Part II 14*, 302–309 (Springer, 2011).
40. Lee, H., Kang, H., Chung, M. K., Kim, B.-N. & Lee, D. S. Persistent brain network homology from the perspective of dendrogram. *IEEE Transactions on Med. Imaging* **31**, 2267–2277 (2012).
41. Chung, M. K., Lee, H., Solo, V., Davidson, R. J. & Pollak, S. D. Topological distances between brain networks. In *Connectomics in NeuroImaging: First International Workshop, CNI 2017, Held in Conjunction with MICCAI 2017, Quebec City, QC, Canada, September 14, 2017, Proceedings 1*, 161–170 (Springer, 2017).
42. Edwards, D. A. The structure of superspace. In *Studies in Topology*, 121–133 (Elsevier, 1975).
43. Gromov, M. Groups of polynomial growth and expanding maps (with an appendix by Jacques Tits). *Publ. Mathématiques de l’IHÉS* **53**, 53–78 (1981).
44. Memoli, F. On the use of Gromov-Hausdorff Distances for Shape Comparison. In Botsch, M., Pajarola, R., Chen, B. & Zwicker, M. (eds.) *Eurographics symposium on point-based graphics*, DOI: [10.2312/SPBG/SPBG07/081-090](https://doi.org/10.2312/SPBG/SPBG07/081-090) (The Eurographics Association, 2007).
45. Bronstein, A. M., Bronstein, M. M., Kimmel, R., Mahmoudi, M. & Sapiro, G. A gromov-Hausdorff framework with diffusion geometry for topologically-robust non-rigid shape matching. *Int. J. Comput. Vis.* **89**, 266–286 (2010).
46. Chazal, F., Cohen-Steiner, D., Guibas, L. J., Mémoi, F. & Oudot, S. Y. Gromov-Hausdorff stable signatures for shapes using persistence. *Comput. Graph. Forum* **28**, 1393–1403 (2009).
47. Mémoi, F. Some properties of Gromov–Hausdorff distances. *Discret. & Comput. Geom.* **48**, 416–440 (2012).
48. Mémoi, F., Smith, Z. & Wan, Z. The gromov-hausdorff distance between ultrametric spaces: its structure and computation. *J. Comput. Geom.* **14**, 78–143 (2023).
49. Zarichnyi, I. Gromov-Hausdorff ultrametric. *arXiv preprint math/0511437* (2005).
50. Bajorath, J. *Chemoinformatics: concepts, methods, and tools for drug discovery*, vol. 275 (Springer Science & Business Media, 2004).
51. Lo, Y. C., Rensi, S. E., Torng, W. & Altman, R. B. Machine learning in chemoinformatics and drug discovery. *Drug Discov. Today* **23**, 1538–1546 (2018).
52. Eckmann, B. Harmonische funktionen und randwertaufgaben in einem komplex. *Commentarii Math. Helvetici* **17**, 240–255 (1944).
53. Lim, L.-H. Hodge Laplacians on graphs. *SIAM Rev.* **62**, 685–715, DOI: [10.1137/18M1223101](https://doi.org/10.1137/18M1223101) (2020). <https://doi.org/10.1137/18M1223101>.
54. Shukla, S. & Yogeshwaran, D. Spectral gap bounds for the simplicial Laplacian and an application to random complexes. *J. Comb. Theory, Ser. A* **169**, 105134 (2020).
55. Torres, J. J. & Bianconi, G. Simplicial complexes: higher-order spectral dimension and dynamics. *J. Physics: Complex.* **1**, 015002 (2020).
56. Muhammad, A. & Egerstedt, M. Control using higher order Laplacians in network topologies. In *Proc. of 17th International Symposium on Mathematical Theory of Networks and Systems*, 1024–1038 (Citeseer, 2006).
57. Horak, D. & Jost, J. Spectra of combinatorial Laplace operators on simplicial complexes. *Adv. Math.* **244**, 303–336 (2013).
58. Barbarossa, S. & Sardellitti, S. Topological signal processing over simplicial complexes. *IEEE Transactions on Signal Process.* **68**, 2992–3007 (2020).

59. Mukherjee, S. & Steenbergen, J. Random walks on simplicial complexes and harmonics. *Random Struct. & Algorithms* **49**, 379–405 (2016).
60. Parzanchevski, O. & Rosenthal, R. Simplicial complexes: spectrum, homology and random walks. *Random Struct. & Algorithms* **50**, 225–261 (2017).
61. Wei, R. K. J., Wee, J., Laurent, V. E. & Xia, K. Hodge theory-based biomolecular data analysis. *Sci. Reports* **12**, 9699 (2022).
62. Schaub, M. T., Benson, A. R., Horn, P., Lippner, G. & Jadbabaie, A. Random walks on simplicial complexes and the normalized Hodge 1-Laplacian. *SIAM Rev.* **62**, 353–391, DOI: [10.1137/18M1201019](https://doi.org/10.1137/18M1201019) (2020). <https://doi.org/10.1137/18M1201019>.
63. Hausdorff, F. *Grundzüge der mengenlehre*, vol. 7 (von Veit, 1914).
64. Kolouri, S., Park, S. R., Thorpe, M., Slepcev, D. & Rohde, G. K. Optimal mass transport: Signal processing and machine-learning applications. *IEEE Signal Process. Mag.* **34**, 43–59 (2017).
65. Villani, C. *Topics in optimal transportation*, vol. 58 (American Mathematical Soc., 2021).
66. Carlsson, E., Carlsson, G. & De Silva, V. An algebraic topological method for feature identification. *Int. J. Comput. Geom. & Appl.* **16**, 291–314 (2006).
67. Schmiedl, F. Computational aspects of the Gromov–Hausdorff distance and its application in non-rigid shape matching. *Discret. & Comput. Geom.* **57**, 854–880 (2017).
68. Anand, D. V., Xu, Q., Wee, J., Xia, K. & Sum, T. C. Topological feature engineering for machine learning based halide perovskite materials design. *npj Comput. Mater.* **8**, 203 (2022).
69. Dean, P. M. *Molecular similarity in drug design* (Springer Science & Business Media, 2012).

Acknowledgements (not compulsory)

This work was supported in part by the Singapore Ministry of Education Academic Research fund Tier 1 grant RG16/23, Tier 2 grants MOE-T2EP20120-0010 and MOE-T2EP20221-0003. X.G. acknowledges support from NTU Presidential Postdoctoral Fellowship 023545-00001. The authors thank Yipeng Zhang for his contributions to the revision of this manuscript.

Author contributions statement

W.T. and K.X. designed the experiments. J.W. and X.G. conducted the experiments and analyzed the results. J.W. drafted the initial manuscript. X.G. revised the manuscript. All authors reviewed and contributed to the final manuscript.

Additional information

The authors declare that there are no competing interests related to this work.

Data availability

Code and data are available in GitHub repository <https://github.com/XueGong-git/pyGH>.